








T18Q1: Level 2 (L.G. 9)

Give the complete ionic equation for the reaction that occurs when aqueous solutions of lithium sulfide and copper (II) nitrate are mixed:



T18Q1: Solution

Give the complete ionic equation for the reaction that occurs when aqueous solutions of lithium sulfide and copper (II) nitrate are mixed:



Complete Ionic Equation:
contains the solids and ALL the ionic species



T18Q2: Level 2 (L.G. 9)

Give the net ionic equation for the reaction that occurs when aqueous solutions of barium nitrate and ammonium phosphate are mixed:

- A. $3\text{Ba}(\text{NO}_3)_2(\text{aq}) + 2(\text{NH}_4)_3\text{PO}_4(\text{aq}) \rightarrow \text{Ba}_3(\text{PO}_4)_2(\text{s}) + 6\text{NH}_4\text{NO}_3(\text{aq})$
- B. $3\text{Ba}(\text{NO}_3)_2(\text{aq}) + 2(\text{NH}_4)_3\text{PO}_4(\text{aq}) \rightarrow \text{Ba}_3(\text{PO}_4)_2(\text{aq}) + 6\text{NH}_4\text{NO}_3(\text{s})$
- C. $2\text{NO}_3^-(\text{aq}) + 6\text{NH}_4^+(\text{aq}) \rightarrow 6\text{NH}_4\text{NO}_3(\text{s})$
- D. $3\text{Ba}^{2+}(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) \rightarrow \text{Ba}_3(\text{PO}_4)_2(\text{s})$
- E. $2\text{NO}_3^-(\text{aq}) + 6\text{NH}_4^+(\text{aq}) + 2\text{PO}_4^{3-}(\text{aq}) \rightarrow \text{Ba}_3(\text{PO}_4)_2(\text{s}) + 6\text{NH}_4^+(\text{aq}) + \text{NO}_3^-(\text{aq})$

T18Q2: Solution

Give the net ionic equation for the reaction that occurs when aqueous solutions of barium nitrate and ammonium phosphate are mixed:



Net Ionic Equation:

Only the species that ACTUALLY react to form a solid!

All NH_4^+ and NO_3^- ion salts are soluble so any species with those ions will cancel out

T18Q3: Level 3 (L.G. 10)

Consider the decomposition of calcium carbonate to form calcium oxide and carbon dioxide. If the reaction produced 732 mL of CO_2 at 21°C and 77.1 cm Hg, how many grams of CaO are produced?

- A. 0.21 g**
- B. 1.73 g**
- C. 1.86 g**
- D. 3.10 g**
- E. 17.2 g**

T18Q3: Solution

Consider the decomposition of calcium carbonate to form calcium oxide and carbon dioxide. If the reaction produced 732 mL of CO_2 at 21°C and 77.1 cm Hg, how many grams of CaO are produced?

A. 0.21 g

B. 1.73 g

C. 1.86 g

D. 3.10 g

E. 17.2 g



2

$$PV = nRT$$

$$n = PV/RT$$

$$= (1.01 \text{ atm})(0.732 \text{ L}) / (0.082 \text{ L-atm/mol-K})(294 \text{ K})$$

$$= 0.031 \text{ moles}$$

$$77.1 \text{ cm Hg} = 1.01 \text{ atm}$$

$$21^\circ\text{C} = 294 \text{ K}$$

3

1:1 mole ratio

0.031 moles CaO

4

$$0.031 \text{ moles CaO} \times 56.08 \text{ g/mol}$$

$$= 1.73 \text{ grams}$$

T18Q4: Level 3 (L.G. 10)

Consider the reaction between 100 mL of 0.41 M iron(II) nitrate with 63 mL of 0.35M potassium phosphate. How many moles of precipitate are formed from this reaction?

- A. 0.011
- B. 0.014
- C. 0.022
- D. 0.041

T18Q4: Solution

Consider the reaction between 100 mL of 0.41 M iron(II) nitrate with 63 mL of 0.35M potassium phosphate. How many moles of precipitate are formed from this reaction?



A. 0.011

B. 0.014

C. 0.022

D. 0.041

2

$$M = \frac{\text{moles of solute}}{\text{volume of solution (L)}}$$

$$\text{moles} = M \times \text{vol (L)}$$

3 LR
calcs

$3\text{Fe}(\text{NO}_3)_2$	$2\text{K}_3\text{PO}_4$
$0.41 \text{ M} \times 0.100 \text{ L} = 0.041 \text{ mol}$	$0.35 \text{ M} \times 0.063 \text{ L} = 0.0221 \text{ mol}$
$0.041 \times (1/3)$ $= 0.014 \text{ mol Fe}_3(\text{PO}_4)_2$	$0.0221 \times (1/2)$ $= 0.011 \text{ mol Fe}_3(\text{PO}_4)_2$


T18Q5: Level 3 (L.G. 10)

What mass, in g, of AgCl is formed from the reaction of 75.0 mL of a 0.078 M $\text{AgC}_2\text{H}_3\text{O}_2$ solution with 55.0 mL of 0.109 M MgCl_2 solution?

- A. 0.860 g
- B. 1.72 g
- C. 2.56 g
- D. 3.20 g

T18Q5: Solution

What mass, in g, of AgCl is formed from the reaction of 75.0 mL of a 0.078 M $\text{AgC}_2\text{H}_3\text{O}_2$ solution with 55.0 mL of 0.109 M MgCl_2 solution?



A. 0.860 g

B. 1.72 g

C. 2.56 g

D. 3.20 g

2

$$M = \frac{\text{moles of solute}}{\text{volume of solution (L)}}$$

$$\text{moles} = M \times \text{vol (L)}$$



$$0.078 \text{ M} \times 0.075 \text{ L} = 0.006 \text{ mol}$$

$$0.109 \text{ M} \times 0.055 \text{ L} = 0.0060 \text{ mol}$$

LR
calcs

3

$$0.006 \times (2/2) = 0.006 \text{ mol AgCl}$$

$$0.006 \times (2/1) = 0.012 \text{ mol AgCl}$$

4

$$0.006 \text{ mol} \times (143.35 \text{ g} / 1 \text{ mol}) = 0.86 \text{ g}$$

T18Q6: Level 2 (L.G. 8)

What precipitate is most likely formed from a solution containing Ba^{+2} , Na^{+1} , OH^{-1} , and CO_3^{-2} ?

- A. NaOH
- B. BaCO_3
- C. Na_2CO_3
- D. Ba(OH)_2

T18Q6: Solution

What precipitate is most likely formed from a solution containing Ba^{+2} , Na^{+1} , OH^{-1} , and CO_3^{-2} ?

- A. NaOH → Soluble, bc Na is a group 1 metal
- B. BaCO_3
- C. Na_2CO_3 → Soluble, bc Na is a group 1 metal
- D. $\text{Ba}(\text{OH})_2$ → Slightly soluble, bc OH & Ba^{2+}

- 2. Alkali metal (Group 1A) salts and NH_4^+ are soluble.
- 3. F^- , S^{2-} , CO_3^{2-} , CrO_4^{2-} , PO_4^{3-} salts are insoluble, except for those containing Group 1A cations.
- 6. OH salts are insoluble (except for those containing Group 1A cations and NH_4^+ which are soluble and those containing Ca^{2+} , Sr^{2+} and Ba^{2+} which are slightly soluble).

T18Q7: Level 3 (L.G. 8)

How many of the following compounds are soluble in water?

$\text{Cu}(\text{OH})_2$ LiNO_3 NH_4Br K_2SO_4 PbCl_2 BaSO_4

- A. 0
- B. 1
- C. 2
- D. 3
- E. 4

T18Q7: Solution

How many of the following compounds are soluble in water?



A. 0

B. 1

C. 2

D. 3

E. 4

Soluble Soluble Soluble

1. Nitrate (NO_3^{-1}) and acetate ($\text{C}_2\text{H}_3\text{O}_2^{-1}$) salts are soluble.
2. Alkali metal (Group 1A) salts and NH_4^+ are soluble.
4. Cl^{-1} , Br^{-1} , and I^{-1} salts are soluble (except when combined with Ag^+ , Hg_2^{2+} , Pb^{2+}).
5. Sulfate salts are soluble (except when combined with Ag^+ , Hg_2^{2+} , Pb^{2+} , Ca^{2+} , Sr^{2+} and Ba^{2+}).
6. OH^{-1} salts are insoluble (except for those containing Group 1A cations and NH_4^+ which are soluble and those containing Ca^{2+} , Sr^{2+} and Ba^{2+} which are slightly soluble).




T18Q8: Level 1 (L.G. 6)

Choose the reaction that represents a combustion reaction:

- A. $\text{C}_6\text{H}_{12}\text{O}_2(\text{l}) + 8\text{O}_2(\text{g}) \rightarrow 6\text{CO}_2(\text{g}) + 6\text{H}_2\text{O}(\text{g})$
- B. $\text{Mg}(\text{s}) + \text{Cl}_2(\text{l}) \rightarrow \text{MgCl}_2(\text{aq})$
- C. $\text{C}_6\text{H}_{12}\text{O}_2(\text{l}) \rightarrow 6\text{C}(\text{s}) + 6\text{H}_2(\text{g}) + \text{O}_2(\text{g})$
- D. $\text{NaOH}(\text{aq}) + \text{CuCl}_2(\text{aq}) \rightarrow \text{NaCl}_2(\text{aq}) + \text{Cu}(\text{OH})_2(\text{s})$
- E. None of the above are combustion

T18Q8: Solution

Choose the reaction that represents a combustion reaction:



E. None of the above are combustion

A = Combustion (reaction with oxygen)

B = Synthesis (combining things)

C = Decomposition (breaking down)

D = Double displacement (swapping ions)

T18Q9: Level 2 (L.G. 9)

Give the complete ionic equation for the reaction that occurs when aqueous solutions of lithium sulfide and copper (II) nitrate are mixed:

