


Treatment Windows for Obesity-Induced Type II Diabetes with GLP-1 Agonist Drugs

By Eli Bullock-Papa

Quiz: What is something that looks cute but is extremely dangerous?



Quiz: What is something that looks cute but is extremely dangerous?




In this presentation:

1. What is diabetes
2. Overview of how it forms
3. A more complete model and why we think it's accurate
4. How the model has been used
5. What is missing from existing work and my research plan

What is Diabetes?

Diabetes Overview

Complex condition **diagnosed** by having **too much glucose in your blood**



Major Components Overview


Blood Glucose: sugar your body produces when you eat.

Insulin: a hormone produced to tell cell they should absorb glucose from the bloodstream.

Beta cells: cells in the pancreas that produce insulin.

Type 2 Diabetes

Type 2 diabetes



In type 2 diabetes, the pancreas makes some insulin but it is not working as well as it used to.

How does it
develop?

Diabetes Development and Obesity

Views have **changed over time** but this is the modern overview of **how we think Type 2 Diabetes (T2D) is developed in youth** (people under 40):

1. Due to the combination of diet and lack of exercise, a person starts to **gain excessive weight**, potentially leading to obesity.
2. In response to high blood sugar levels, the **pancreas produces more insulin** than normal to try to lower blood sugar levels.

Diabetes Development and Obesity

3. The body **begins to develop insulin resistance as a protective response**. It's theorized that this resistance helps shield the body's cells from excessive sugar intake and the stress of high insulin levels
4. The insulin resistance means that cells aren't taking up all of the blood glucose, and **blood sugar remains high** even with high insulin levels.
5. High blood sugar levels **keep beta cells producing insulin levels in overdrive**.
6. Over long stretches of time, this overwork causes **beta cells to begin dysfunctioning or even die**.
7. This plus continued insulin resistance means **blood glucose levels continue to spiral upwards**

A more complete
model

Model source:

[iScience](#). 2023 Nov 17; 26(11): 108324.

PMCID: PMC10665812


Published online 2023 Oct 23. doi: [10.1016/j.isci.2023.108324](https://doi.org/10.1016/j.isci.2023.108324)


PMID: [38026205](https://pubmed.ncbi.nlm.nih.gov/38026205/)


A data-driven computational model for obesity-driven diabetes onset and remission through weight loss


[Vehpi Yildirim](#),^{1,2,9,*} [Vivek M. Sheraton](#),^{2,3,4} [Ruud Brands](#),^{5,6} [Loes Crielaard](#),^{1,2} [Rick Quax](#),^{2,3} [Natal A.W. van Riel](#),^{7,8}
[Karien Stronks](#),^{1,2} [Mary Nicolaou](#),^{1,2} and [Peter M.A. Sloot](#)^{2,3}


► [Author information](#) ► [Article notes](#) ► [Copyright and License information](#) [PMC Disclaimer](#)















Glucose to β -cell Mass: The effect can be positive or negative depending on the circumstances. In this context, an optimal level of glucose is necessary to maintain β -cell mass. However, persistently high glucose levels (hyperglycemia) can lead to β -cell dysfunction and loss of mass.



































Remission



E_{GO}

IDGU



Liver



Fat Mass



FFA




Insulin

Diabetes

Inflammation to β -cell function: represents the adverse impact of inflammation on the pancreas's β -cells. Elevated levels of pro-inflammatory cytokines can trigger cellular stress responses, lead to programmed cell death, and interfere with insulin production and secretion processes, collectively contributing to the decline in the cells' ability to produce and regulate insulin effectively.

Ener
Inta




Is this model
accurate?

Why they think their model is accurate


- It incorporates what we know of as the most important processes in the formation of diabetes and T2D
- It reproduces experimental findings from the DIRECT study which compared weight intervention vs. diabetes treatment only. It reproduced:
 - Longer T2D duration for non-responders compared with responders
 - β -cell function increases in both groups but β -cell mass only increases in responders
 - Higher pancreas volume in non-diabetic controls
 - Greater increase in pancreas volumes of the responders vs non-responders

How did the paper
use this model?


Importance of Treatment Timelines



Bifurcation Analysis




Treatment efficacy based on BMI and beta cell mass



My contribution

Trends I think are relevant

1. Introduction of **GLP-1 agonist drugs like Semaglutide and Tirzepatide**. They have multiple effects that act potently on diabetes **and** obesity.
2. **Rises in rates of childhood obesity**. This means that by the time people are eligible for treatment with these GLP-1 agonists, **their window for effective treatment is already partly closed**



My research plan:

1. Get the model fully running on my computer using their provided python and xpp code posted on github.
2. Format trial data as necessary from the GLP-1 agonist drug trials - most importantly weight information of participants
3. Model different treatment timelines for ages at which to start treatment with the different drugs and their efficacy rates

Any Questions?