

Analyzing the choice of transportation

Elie Daher

July 3, 2017

Abstract

This document contains the problem of analysing the choice of transportation. This problem is from the chapter UTA Methods of the Book: Multiple Criteria Decision Analysis. This document was made during my internship at LAMSADE in the summer of 2017.

A DM wants to analyse the choice of transportation. The DM is interstered in the following criteria

1. price
2. time (min)
3. comfort (possibility to have a seat)

The evaluation of the previous criteria:

| Means of transportation | Price | Time | Comfort | Ranking of the DM |
|-------------------------|-------|------|---------|-------------------|
| RER | 3 | 10 | + | 1 |
| METRO (1) | 4 | 20 | ++ | 2 |
| METRO (2) | 2 | 20 | 0 | 2 |
| BUS | 6 | 40 | 0 | 3 |
| TAXI | 30 | 30 | +++ | 4 |

DM's preferences: $RER \succ Metro1 \approx Metro2 \succ Bus \succ Taxi$

1 Scale for each criteria

For each criteria, the interval $[g_i^*, g_{i*}]$ is cut into $(\alpha_i - 1)$ equal intervals. So in this case we have:

- Price $\rightarrow [30, 16, 2]$
- Time $\rightarrow [40, 30, 20, 10]$
- Comfort $\rightarrow [0, +, ++, +++]$

2 Marginal value by linear interpolation

The marginal value is calculated by a linear interpolation. In this case we have:

- $v[g(RER)] = 0.07v_1(16) + 0.93v_1(2) + v_2(10) + v_3(+)$
- $v[g(METRO1)] = 0.14v_1(16) + 0.86v_1(2) + v_2(20) + v_3(++)$
- $v[g(METRO2)] = v_1(2) + v_2(20) + v_3(0) = v_1(2) + v_2(20)$
- $v[g(BUS)] = 0.29v_1(16) + 0.71v_1(2) + v_2(40) + v_3(0) = 0.29v_1(16) + 0.71v_1(2)$
- $v[g(TAXI)] = v_1(30) + v_2(30) + v_3(+++) = v_2(30) + v_3(+++)$

3 Replace v_i with w_{ij}

- $v[g(RER)] = w_{11} + 0.93w_{12} + w_{21} + w_{22} + w_{23} + w_{31}$
- $v[g(METRO1)] = w_{11} + 0.86w_{12} + w_{21} + w_{22} + w_{31} + w_{32}$
- $v[g(METRO2)] = w_{11} + w_{12} + w_{21} + w_{22}$
- $v[g(BUS)] = w_{11} + 0.71w_{12}$
- $v[g(TAXI)] = w_{21} + w_{31} + w_{32} + w_{33}$

4 Difference between each pair of consecutive actions

- $\Delta(RER, METRO1) = 0.07w_{12} + w_{23} - w_{32} + \sigma_{RER} - \sigma_{METRO1} \geq \delta$
- $\Delta(METRO1, METRO2) = -0.14w_{12} + w_{31} + w_{32} + \sigma_{METRO1} - \sigma_{METRO2} = 0$
- $\Delta(METRO2, BUS) = 0.29w_{12} + w_{21} + w_{22} + \sigma_{METRO2} - \sigma_{BUS} \geq \delta$
- $\Delta(BUS, TAXI) = w_{11} + 0.71w_{12} - w_{21} - w_{31} - w_{32} - w_{33} + \sigma_{BUS} - \sigma_{TAXI} \geq \delta$

5 Linear Program

Main objectif: $[min]F = \sum_{a \in A_R} \sigma(a)$

subject to :

$$\begin{aligned} \Delta(RER, METRO1) &\geq \delta \\ \Delta(METRO1, METRO2) &= 0 \\ \Delta(METRO2, BUS) &\geq \delta \\ \Delta(BUS, TAXI) &\geq \delta \\ \sum_{i=1}^n u_i(g_i^*) &= 1 \end{aligned}$$

With $[min]F = \sum_{a \in A_R} \sigma(a)$ as the main objectif, we have the following linear program to solve:

| Desc | w_{11} | w_{12} | w_{21} | w_{22} | w_{23} | w_{31} | w_{32} | w_{33} | Result |
|-----------------------------------|----------|----------|----------|----------|----------|----------|----------|----------|---------------|
| $\Delta(RER, METRO1) \geq \delta$ | 0 | 0.07 | 0 | 0 | 1 | 0 | -1 | 0 | $\geq \delta$ |
| $\Delta(METRO1, METRO2) = 0$ | 0 | -0.14 | 0 | 0 | 0 | 1 | 1 | 0 | $= 0$ |
| $\Delta(METRO2, BUS) \geq \delta$ | 0 | 0.29 | 1 | 1 | 0 | 0 | 0 | 0 | $\geq \delta$ |
| $\Delta(BUS, TAXI) \geq \delta$ | 1 | 0.71 | -1 | 0 | 0 | -1 | -1 | -1 | $\geq \delta$ |
| $\sum_{i=1}^n u_i(g_i^*) = 1$ | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | $= 1$ |

So by using the `com.google.ortools` library, we can solve the Linear Program above with $\sigma = 0.05$. This Linear Program solution is coded in Java class `ChoiceTransportation`.

```
Minimize
Obj: +1 eRER_ +1 eRER__1 +1 eMETRO1_ +1 eMETRO1__2 +1 eMETRO2_ +1 eMETRO2__3 +1 e
Subject to
auto_c_000000000: +0.070000000000000001 w12 +1 w23 -1 w32 -1 eRER_ +1 eRER__1 +1 e
auto_c_000000001: -0.14 w12 +1 w31 +1 w32 -1 eMETRO1_ +1 eMETRO1__2 +1 eMETRO2_
auto_c_000000002: +0.29 w12 +1 w21 +1 w22 -1 eMETRO2_ +1 eMETRO2__3 +1 eBUS_ -1 e
auto_c_000000003: +1 w11 +0.71 w21 -1 w22 -1 w31 -1 w32 -1 w33 +1 eTAXI_ -1 eTAXI__
auto_c_000000004: +1 w11 +1 w12 +1 w21 +1 w22 +1 w23 +1 w31 +1 w32 +1 w33 = 1
```

An optimal solution has been found of the LP with $\sigma = 0.05$. The objective was accomplished with $[min]F = \sum_{a \in A_R} \sigma(a) = 0$ and $w_{11} = 0.5$, $w_{22} = 0.05$, $w_{23} = 0.05$, $w_{33} = 0.4$.

```
Problem solved in 286 milliseconds
Optimal objective value = 0.0
w31 = 0.0
w11 = 0.5
w22 = 0.05
w33 = 0.4
w21 = 0.0
w32 = 0.0
w12 = 0.0
w23 = 0.05
```