

Inequalities Witnessing Quantum Incompatibility in The Triangle Scenario

Thomas C. Fraser*

Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada

University of Waterloo, Waterloo, Ontario, Canada

(Dated: September 3, 2016)

Causal inference aims to identify which observable distributions are compatible with a given causal structure. Compatibility inequalities for the triangle scenario have been obtained and demonstrated to be violated by quantum configurations. Obtaining these inequalities has lead to the development of new computational techniques suitable for large causal structures with numerous outcomes. Numerical optimizations against these inequalities have found new quantum configurations that are incompatible with the triangle scenario, potentially suggesting new quantum resources qualitatively different from those known previously. **Todo (TC Fraser): Write a better abstract.**

* tcfrazer@tcfrazer.com

CONTENTS

I. Introduction	3
II. Definitions & Notation	3
A. Preliminary Definitions	6
III. Compatibility, Contextuality and the Marginal Problem	6
IV. The Triangle Scenario	8
V. The Fritz Distribution	9
VI. Summary of the Inflation Technique	10
VII. Certificate Inequalities	12
A. Casting the Marginal Problem as a Linear Program	12
B. Infeasibility Certificates	13
VIII. Covering Inequalities	14
A. Logical Covers	14
B. Extendable Covers	15
C. Complete Solution to The Marginal Problem	16
D. Extendable Cover Procedure	16
1. Optimizations	16
2. Avoiding Trivial Inequalities	16
E.	16
F. Irreducible Covers	16
G. Limited Factorization	17
H. Logical Implications & Inequalities	17
I. Implications in the Marginal Problem	17
J. Higher Order Marginal Implications	18
K. Hypergraph Transversals	18
L. Weighted Hypergraph Transversals	18
IX. Deriving Symmetric Inequalities	19
A. Causal Symmetry	19
B. Symmetric Marginal Polytope	20
C. Symmetric Logical Implications	21
D. Results	21
X. Non-linear Optimizations	21
A. Numerical Violations of The CHSH Inequality	22
XI. Conclusions	22
XII. Open Questions & Future Work	23
A. Exemplary Inequalities	23
B. Connections to Sheaf-Theoretic Treatment	23
C. Computationally Efficient Parametrization of the Unitary Group	23
D. Parametrization of Quantum States & Measurements	24
1. Quantum States	24
2. Measurements	25
3. Network Permutation Matrix	26
4. Degeneracy	27

I. INTRODUCTION

Todo (TC Fraser): Discuss why the marginal problem is important.

II. DEFINITIONS & NOTATION

Comment (TC Fraser): This entire section should be integrated into the body when logical to do so.

Definition 1. Following common notation used in [1], we use the notation $[k] = \{1, \dots, k\}$ to be a finite index set. Moreover, when referencing a finite set with k elements, we can write,

$$A_{[k]} \equiv \{A_i\}_{i \in [k]} = \{A_1, \dots, A_k\}$$

Definition 2. Each random variable v has a set of possible outcomes which are labeled by **outcome labels** O_v . The set of all outcomes themselves is its **outcome space** and is represented by the set of all functions from v to O_v ¹,

$$\text{Out}(v) \equiv \{f : \{v\} \rightarrow O_v\}$$

Each of these functions has a singular domain (v) and thus can be concisely written in a compact form $\{v \mapsto o\}$ where $o \in O_v$ represents some generic outcome label. Each function f should be interpreted as the *event* that v obtains the outcome o , as well as the function that takes v to o $\{v \mapsto f(v) = o\}$. Evidently $\text{Out}(v)$ is isomorphic to O_v , and in many cases the two can be considered equivalent; however we make this particular semantic distinction now in preparation of section IX.

Definition 3. As a natural generalization, when considering a larger set of random variables $V = \{v_i\}_{i \in [k]}$ we define the joint outcome space in an analogous way,

$$\text{Out}(V) \equiv \left\{ f : \{v_i\}_{i \in [k]} \rightarrow \{O_{v_i}\}_{i \in [k]} \mid \forall i \in [k] : f(v_i) \in O_{v_i} \right\} \quad (1)$$

Each outcome f can be compactly represented as a set of mappings over each element of V .

$$f = \{v_1 \mapsto f(v_1), \dots, v_k \mapsto f(v_k)\} = \{v_i \mapsto f(v_i)\}_{i \in [k]}$$

The domain of f refers to the set of random variables it evaluates and is denoted as $\text{Dom}(f)$. If $f \in \text{Out}(V)$, then $\text{Dom}(f) = V$.

Example 4. Interpreting $f \in \text{Out}(V)$ as an *event* or *outcome* on V , a probability distribution P_V over V will assign a non-negative probability $P_V(f)$ to f . Let $V = \{v_1, v_2\}$ with each $v \in V$ having 3 distinct outcomes $O_v = \{0, 1, 2\}$. Furthermore let $f \in \text{Out}(V)$ be the outcome $(v_1 \mapsto 2, v_2 \mapsto 1)$. The following class of notations will be considered interchangeable,

$$P_V(f) = P_{v_1, v_2}(f) = P(v_1 \mapsto f_1(v_1), v_2 \mapsto f_2(v_2)) = P(v_1 \mapsto 2, v_2 \mapsto 1) = P_{v_1 v_2}(21)$$

Each of which should be read as “the probability that v_1 gets outcome 2 AND v_2 gets outcome 1”.

Remark 5. In each of the above notations, P_V always represents a probability distribution over the *domain* of $f \in \text{Out}(V)$. In cases where the subscript is omitted, it should be assumed to be over the domain of f .

$$P_V(f) = P_{\text{Dom}(f)}(f) = P(f)$$

¹ Using functional language for outcome events is necessary for a sheaf-theoretic treatment of contextuality [2].

Definition 6. Let $V = \{v_1, \dots, v_k\}$ be a set of random variables and $W \subseteq V$ be a subset of V . Moreover, let $f \in \text{Out}(V)$ be a joint outcome over V ; assigning a unique label to each variable in V . The **restriction of f onto W** (denoted $f|_W \in \text{Out}(W)$) is the outcome of $\text{Out}(W)$ that *agrees* with each of f 's assignments for variables in W .

$$\forall v \in W : f(v) = f|_W(v)$$

Definition 7. Consider two sets of random variables V and W and two specific outcomes $f_V \in \text{Out}(V)$ and $f_W \in \text{Out}(W)$. The outcomes f_V and f_W are said to be **compatible** if they agree on their valuations of $V \cap W$. Compatibility will be denoted with ' \succsim '.

$$f_V \succsim f_W \iff f_V|_{V \cap W} = f_W|_{V \cap W}$$

Compatibility is a reflexive and symmetric, but *not* transitive. There are two cases when compatibility is transitive; whenever $V \cap W = \emptyset$ or $V = W$. Whenever $V \cap W = \emptyset$, f_V and f_W can be said to be *trivially* compatible. Moreover if $V = W$, compatibility becomes ordinary equivalence,

$$V = W \quad \text{and} \quad f_V \succsim f_W \iff f_V = f_W$$

Definition 8. Let $f_V \in \text{Out}(V)$ and $f_W \in \text{Out}(W)$ be outcomes of V and W respectively as in definition 7. If f_V and f_W are compatible, and $W \subseteq V$ then f_W is said to be **extendable** to f_V and is denoted ' \prec '.

$$f_W \prec f_V \iff f_W \succsim f_V \quad \text{and} \quad W \subseteq V \iff f_W = f_V|_W$$

The dual language is also appropriate: f_W is a **restriction** of f_V (denoted $f_V \succ f_W$) if and only if f_W is extendable to f_V . If $f_W \prec f_V$ then there exists a unique element $f_{V \setminus W} \in \text{Out}(V \setminus W)$ such that f_V can be reconstructed by gluing together $f_{V \setminus W}$ and f_W .

$$\forall v \in V : f_V(v) = \begin{cases} f_W(v) & v \in W \\ f_{V \setminus W}(v) & v \in V \setminus W \end{cases}$$

Remark 9. The notation ' $g \succsim h$ ' is used to indicate compatibility between $g \in \text{Out}(V)$ and $h \in \text{Out}(W)$ because of the following observation; g and h are compatible if and only if extensions of some outcome $f \in \text{Out}(V \cap W)$.

$$g \succsim h \iff \exists f \in \text{Out}(V \cap W) \mid g \succ f \prec h$$

In fact, such an f is uniquely defined,

$$\forall v \in V \cap W : f(v) = g(v) = h(v)$$

Definition 10. The set of all extendable outcomes of $f \in \text{Out}(W)$ into V is called the **extendable set of f into V** and can be written as,

$$\text{Ext}_V(f) \equiv \{g \in \text{Out}(V) \mid f \prec g\} \subseteq \text{Out}(V)$$

The extendable set of f into V is the subset of outcomes in $\text{Out}(V)$ that *agree* with f about valuations for variables in W .

Example 11. Let $W = \{a, b\}$ and $V = \{a, b, c\}$ be two sets of random variables. Clearly $W \subseteq V$; a prerequisite for extendability. Further impose that the outcome labels for each variable are the same: $O_a = O_b = O_c = \{0, 1, 2\}$. As an example, let $f = \{a \mapsto f(a) = 1, b \mapsto f(b) = 2\} \in \text{Out}(W)$; f is extendable to the outcome $g = \{a \mapsto g(a) = 1, b \mapsto g(b) = 2, c \mapsto g(c) = 1\} \in \text{Out}(V)$, and the extendable set of f into V is,

$$\text{Ext}_V(f) = \text{Ext}_{a,b,c}(a \mapsto 1, b \mapsto 2) = \{\{a \mapsto 1, b \mapsto 2, c \mapsto 0\}, \{a \mapsto 1, b \mapsto 2, c \mapsto 1\}, \{a \mapsto 1, b \mapsto 2, c \mapsto 2\}\}$$

Definition 12. A **graph** is an ordered tuple $(\mathcal{N}, \mathcal{E})$ of *nodes* and *edges* respectively where the nodes can represent any object and the edges are pairs of nodes. For convenience of notation, one defines an index set over the nodes denoted $\mathcal{I}_{\mathcal{N}}$.

$$\mathcal{N} = \{n_i \mid i \in \mathcal{I}_{\mathcal{N}}\} \quad \mathcal{E} = \{\{n_j, n_k\} \mid j, k \in \mathcal{I}_{\mathcal{N}}\}$$

Definition 13. A **directed graph** \mathcal{G} is an ordered tuple $(\mathcal{N}, \mathcal{E})$ of *nodes* and *edges* respectively where the nodes can represent any object and the edges are *ordered* pairs of nodes. For convenience of notation, one defines an index set over the nodes denoted $\mathcal{I}_{\mathcal{N}}$.

$$\mathcal{N} = \{n_i \mid i \in \mathcal{I}_{\mathcal{N}}\} \quad \mathcal{E} = \{n_j \rightarrow n_k \mid j, k \in \mathcal{I}_{\mathcal{N}}\}$$

Definition 14. The following definitions are common language in directed graph theory. Let $n, m \in \mathcal{N}$ be example nodes of the graph \mathcal{G} .

- The **parents of a node**: $\text{Pa}_{\mathcal{G}}(n) \equiv \{m \mid m \rightarrow n\}$
- The **children of a node**: $\text{Ch}_{\mathcal{G}}(n) \equiv \{m \mid n \rightarrow m\}$
- The **ancestry of a node**: $\text{An}_{\mathcal{G}}(n) \equiv \bigcup_{i \in \mathbb{N}} \text{Pa}_{\mathcal{G}}^i(n)$ where $\text{Pa}_{\mathcal{G}}^i(n) \equiv \text{Pa}_{\mathcal{G}}(\text{Pa}_{\mathcal{G}}^{i-1}(n))$ and $\text{Pa}_{\mathcal{G}}^0(n) = n$

All of these terms can be generalized to sets of nodes $N \subseteq \mathcal{N}$ through union over the elements,

- The **parents of a node set**: $\text{Pa}_{\mathcal{G}}(N) \equiv \bigcup_{n \in N} \text{Pa}_{\mathcal{G}}(n)$
- The **children of a node set**: $\text{Ch}_{\mathcal{G}}(N) \equiv \bigcup_{n \in N} \text{Ch}_{\mathcal{G}}(n)$
- The **ancestry of a node set**: $\text{An}_{\mathcal{G}}(N) \equiv \bigcup_{n \in N} \text{An}_{\mathcal{G}}(n)$

Moreover, an **induced subgraph** of \mathcal{G} due to a set of nodes $N \subseteq \mathcal{N}$ is the graph composed of N and all edges $e \in \mathcal{E}$ of the original graph that are contained in N .

$$\text{Sub}_{\mathcal{G}}(N) \equiv (N, \{e_i \mid i \in \mathcal{I}_{\mathcal{E}}, e_i \subseteq N\})$$

An **ancestral subgraph** of \mathcal{G} due to $N \subseteq \mathcal{N}$ is the induced subgraph due to the ancestry of N .

$$\text{AnSub}_{\mathcal{G}}(N) \equiv \text{Sub}_{\mathcal{G}}(\text{An}_{\mathcal{G}}(N))$$

Definition 15. A **directed acyclic graph** or **DAG** \mathcal{G} is an directed graph definition 13 with the additional property that no node n is in its set of **ancestors**.

$$\forall n \in \mathcal{N} : n \notin \bigcup_{i \in \mathbb{N}} \text{Pa}_{\mathcal{G}}^i(n)$$

Notice the difference between using the natural numbers \mathbb{N} to distinguish *ancestors* from *ancestry*.

Definition 16. A **hypergraph** denoted \mathcal{H} is an ordered tuple $(\mathcal{N}, \mathcal{E})$ of *nodes* and *hyperedges* respectively where the nodes can represent any object and the hyperedges are *subsets* of nodes. For convenience of notation, one defines an index set over the nodes and hyperedges of a hypergraph \mathcal{H} denoted $\mathcal{I}_{\mathcal{N}}$ and $\mathcal{I}_{\mathcal{E}}$ respectively.

$$\mathcal{H} = (\mathcal{N}, \mathcal{E}) \quad \mathcal{N} = \{n_i \mid i \in \mathcal{I}_{\mathcal{N}}\} \quad \mathcal{E} = \{e_i \mid i \in \mathcal{I}_{\mathcal{E}}, e_i \subseteq \mathcal{N}\}$$

Note that whenever the hyperedge or node index is arbitrary, it will be omitted. There is a dual correspondence between hyperedges $e \in \mathcal{E}$ and nodes $n \in \mathcal{N}$ in a Hypergraph. A hyperedge e is viewed as a set of nodes $\{n_i\}$, and a node n can be viewed as the set of hyperedges $\{e_i\}$ that contain it.

Definition 17. A **hypergraph transversal** (or edge hitting set) \mathcal{T} of a hypergraph \mathcal{H} is a set of nodes $\mathcal{T} \subseteq \mathcal{N}$ that have non-empty intersections with every hyperedge in \mathcal{E} .

$$\mathcal{T} = \{n_i \in \mathcal{N} \mid i \in \mathcal{I}_{\mathcal{T}}\} \quad \forall e \in \mathcal{E} : \mathcal{T} \cap e \neq \emptyset$$

Definition 18. A **necessary node** of a transversal \mathcal{T} is a node n such that $\mathcal{T} \setminus n$ is no longer a valid transversal. The set of all necessary nodes is denoted $\text{Nec}(\mathcal{T})$,

$$\text{Nec}(\mathcal{T}) = \{n \in \mathcal{T} \mid \exists e \in \mathcal{E} : (\mathcal{T} \setminus n) \cap e = \emptyset\}$$

An **unnecessary node** of a transversal \mathcal{T} is any node that is *not* necessary. The set of all unnecessary nodes is denoted $\text{UnNec}(\mathcal{T})$,

$$\text{UnNec}(\mathcal{T}) = \mathcal{T} \setminus \text{Nec}(\mathcal{T})$$

A **minimal hypergraph transversal** \mathcal{T} is any valid transversal of \mathcal{H} where every node n is *necessary*.

$$\mathcal{T} = \text{Nec}(\mathcal{T})$$

Definition 19. A **weighted hypergraph** $\mathcal{H}_{\mathcal{W}}$ is a regular hypergraph satisfying definition 16 equipped with a set of weights \mathcal{W} ascribed to each node such that a weighted hypergraph is written as a triplet $(\mathcal{W}, \mathcal{N}, \mathcal{E})$.

$$\mathcal{W} = \{w_i \mid i \in \mathcal{I}_{\mathcal{N}}, w_i \in \mathbb{R}\}$$

One would say that a particular node n_i carries weight w_i for each $i \in \mathcal{I}_{\mathcal{N}}$.

Definition 20. A **bounded transversal** of a weighted hypergraph $\mathcal{H}_{\mathcal{W}}$ is a transversal \mathcal{T} of the unweighted hypergraph \mathcal{H} and a real number t (denoted $\mathcal{T}_{\leq t}$) such that the sum of the node weights of the transversal is bounded by t .

$$\mathcal{T}_{\leq t} = \{n_i \mid i \in \mathcal{I}_{\mathcal{T}}\} \quad \text{s.t.} \quad \sum_{j \in \mathcal{I}_{\mathcal{T}}} w_j \leq t$$

One can define analogous **(strictly) upper/lower bounded transversals** by considering modifications of the notation: $\mathcal{T}_{< t}, \mathcal{T}_{\geq t}, \mathcal{T}_{> t}$.

Definition 21. A **causal structure** is simply a DAG with the extra classification of each node into one of two categories; the **latent nodes** and **observed nodes** denoted \mathcal{N}_L and \mathcal{N}_O . The latent nodes correspond to random variables that are either hidden through some fundamental process or cannot/will not be measured. The observed nodes are random variables that are measurable. Every node is either latent or observed and no node is both:

$$\mathcal{N}_L \cap \mathcal{N}_O = \emptyset \quad \mathcal{N}_L \cup \mathcal{N}_O = \mathcal{N}$$

Definition 22. The **product distribution** of two disjoint distributions P_V and P_W (where $V \cap W = \emptyset$) is denoted as usual with $P_V \times P_W$ and is defined as,

$$\forall f \in \text{Out}(V \cup W) : (P_V \times P_W)(f) \equiv P_V(f|_V) \cdot P_W(f|_W)$$

A product distribution of k mutually disjoint distributions is defined as,

$$\prod_{i \in [k]} P_{V_i} \equiv (P_{V_1} \times \cdots \times P_{V_k})$$

Definition 23. The **marginalization** of a distribution P_V to a distribution over $W \subseteq V$ is denoted $\sum_{V \setminus W} P_V$ and is defined such that,

$$\forall f \in \text{Out}(W) : \left(\sum_{V \setminus W} P_V \right)(f) \equiv \sum_{g \in \text{Ext}_V(f)} P_V(g)$$

A. Preliminary Definitions

Before discussing the present work, we

Todo (TC Fraser): How many definitions do I need to write??

III. COMPATIBILITY, CONTEXTUALITY AND THE MARGINAL PROBLEM

In order to determine if a given marginal distribution P_V or set of marginal distributions $\{P_{V_i}\}_{i \in [k]} \equiv \{P_{V_1}, \dots, P_{V_k}\}$ is compatible with a causal structure \mathcal{G} , one should first formalize what is meant by *compatible*.

Definition 24. A set of **causal parameters** for a particular causal structure \mathcal{G} is the specification of a conditional distribution for every node $n \in \mathcal{N}$ given it's parents in \mathcal{G} .

$$\left\{ P_{n|\text{Pa}_{\mathcal{G}}(n)} \mid n \in \mathcal{N} \right\}$$

Definition 25. A marginal distribution P_V is **compatible** with a causal structure \mathcal{G} (where it is assumed that $V \subseteq \mathcal{N}_O$) if there exists a *choice* of causal parameters $\left\{ P_{n|\text{Pa}_{\mathcal{G}}(n)} \mid n \in \mathcal{N} \right\}$ such that P_V can be *recovered* from the following series of operations:

1. First obtain a joint distribution over *all* nodes of the causal structure,

$$P_{\mathcal{N}} = \prod_{n \in \mathcal{N}} P_{n|\text{Pa}_{\mathcal{G}}(n)}$$

2. Then marginalize over the latent nodes of \mathcal{G} ,

$$P_{\mathcal{N}_O} = \sum_{\mathcal{N}_L} P_{\mathcal{N}}$$

3. Finally marginalize over the observed nodes not in V to obtain P_V ,

$$P_V = \sum_{\mathcal{N}_O \setminus V} P_{\mathcal{N}_O}$$

A set of marginal distributions $\{P_{V_i}\}_{i \in [k]}$ is compatible with \mathcal{G} if each distribution P_{V_i} can be made compatible by the *same* choice of causal parameters. A distribution P_V or set of distributions $\{P_{V_i}\}_{i \in [k]}$ is said to be **incompatible** with a causal structure if there *does not exist* a set of causal parameters with the above mentioned property.

Todo (TC Fraser): Source this?

Operations 2 and 3 of definition 25 are related to the *marginal problem*.

Definition 26. The Marginal Problem: Given a set of distributions $\{P_{V_i}\}_{i \in [k]} \equiv \{P_{V_1}, \dots, P_{V_k}\}$ where $V_i \subseteq \mathcal{J}$ for some set of random variables \mathcal{J} , does there exist a joint distribution $P_{\mathcal{J}}$ such that each given distribution P_{V_i} can be obtained from marginalizing $P_{\mathcal{J}}$?

$$\forall i \in [k] : P_{V_i} = \sum_{\mathcal{J} \setminus V_i} P_{\mathcal{J}}$$

Typically (although not strictly necessary²), \mathcal{J} is taken to mean the union of all V_i 's.

$$\mathcal{J} = V_1 \cup \dots \cup V_k = \bigcup_{i \in [k]} V_i \quad (2)$$

Todo (TC Fraser): Mention polytope perspective and existing methods

Definition 27. A reoccurring motif of these discussions will be the set of distributions $\{P_{V_i}\}_{i \in [k]} \equiv \{P_{V_1}, \dots, P_{V_k}\}$ mentioned in definition 26. In agreement with [1] we call the set of subsets $\mathcal{M} \equiv \{V_1, \dots, V_k\}$ the marginal contexts or the **maximal marginal scenario**³. An individual $V_i \in \mathcal{M}$ is a **marginal context**. The union of all contexts is denoted \mathcal{J} and is defined exactly as in eq. (2). Moreover, we will call the set of distributions a **marginal model** and denote it $P^{\mathcal{M}}$ provided that they are *compatible* **Todo (TC Fraser):** Mention quantum non-signalling and maybe $k \geq 3$:

$$\forall i \neq j \text{ if } V_i \cap V_j \neq \emptyset \text{ then } \sum_{V_i \setminus V_j} P_{V_i} = \sum_{V_j \setminus V_i} P_{V_j}$$

Following definition 2.3 in [1], a marginal model is said to be **contextual** if it *does not* admit a joint distribution $P_{\mathcal{J}}$ and **non-contextual** otherwise.

In addition, we elect to define the **marginal outcomes** $\text{Out}(\mathcal{M})$ to be the set of all outcomes belonging to outcomes $\text{Out}(V_i)$ of the marginal context⁴.

$$\text{Out}(\mathcal{M}) \equiv \prod_{i \in [k]} \text{Out}(V_i)$$

Each marginal outcome $m \in \text{Out}(\mathcal{M})$ belongs to a unique marginal outcome space $\text{Out}(V_i)$. To make reference to the marginal context V_i associated with m , recall remark 5: the context for m is precisely the domain of m ($V_i = \text{Dom}(m)$). Note that $\text{Out}(\mathcal{M})$ is a slight abuse of notation; $\text{Out}(\mathcal{M})$ is *not* a valid outcome space, analogous to the fact that $P^{\mathcal{M}}$ is *not* a probability distribution. Instead $\text{Out}(\mathcal{M})$ is a *flattened* collection of outcome spaces just as $P^{\mathcal{M}}$ is a collection of distributions. The **joint outcomes** $\text{Out}(\mathcal{J})$ are defined in the usual fashion of definition 2.

² Appending any additional variables to \mathcal{J} will not affect the existence of a joint distribution.

³ Rigorously speaking as defined by [1], a marginal scenario forms an *abstract simplicial complex* where it is required that $V' \in \mathcal{M}$ whenever $V' \subseteq V_i$ for some $i \in [k]$. “Maximal” refers to the restriction that $\forall i, j : V_i \not\subseteq V_j$.

⁴ Here ‘ \prod ’ refers to the *disjoint union* of outcome contexts; when $i \neq j$, $\text{Out}(V_i) \cap \text{Out}(V_j) = \emptyset$ since $V_i \neq V_j$.




FIG. 1. The casual structure of the Triangle Scenario. Three variables A, B, C are observable and illustrated as triangles, while X, Y, Z are latent variables illustrated as circles.




FIG. 2. An inflated causal structure of the Triangle Scenario fig. 1.

Definition 28. A **causal compatibility inequality** $\mathcal{I}_{\mathcal{M}}$ for a marginal scenario \mathcal{M} is a probabilistic inequality that is obeyed for every compatible marginal model $P^{\mathcal{M}}$. Whenever the marginal scenario \mathcal{M} in question is evident by context, the subscript of $\mathcal{I}_{\mathcal{M}}$ will be dropped leaving just \mathcal{I} .

Todo (TC Fraser): Discuss Compatibility, connection to cooperative games/resources, bell incompatibility? Todo (TC Fraser): Connection between contextuality and Compatibility via the marginal problem for causal parameters Todo (TC Fraser): Discuss what is meant by a ‘complete’ solution to the marginal problem Todo (TC Fraser): Maybe define the possibilistic marginal problem for later

IV. THE TRIANGLE SCENARIO

The triangle scenario \mathcal{T} is a causal structure composed of 3 parties A, B, C arranged in a triangular configuration while pair-wise sharing latent variables X, Y, Z . It has that has been studied extensively in existing literature (see [3, Fig. 1], [4, Fig. 6], [5, Fig. 8], [6, Fig. 8, Appendix E], [7, Fig. 3], [8, Fig. 1], ...) and is reproduced here in fig. 1 for convenience. At the time of publication for [5] it was noted that characterizing locality in \mathcal{T} remained an open problem and that identifying locality constraints in this configuration seemed challenging. In [6], \mathcal{T} was classified as an “interesting” causal structure.⁵ Additionally \mathcal{T} is presented along with a family of entropic inequalities that constrain the set of compatible distributions [6]. Furthermore in [7], Fritz demonstrated that \mathcal{T} is the *smallest* correlation scenario in which there exists quantum incompatible distributions by explicitly mentioning one (which we elect to call the *Fritz distribution* in section V) and poses an important question asking if other incompatible quantum distributions exist. Finally in [8], Wolfe et. al. make use of \mathcal{T} along with the Bell Scenario to demonstrate a novel technique for causal inference called the *inflation technique* (summarized in section VI). Wolfe et. al. validate the power and applicability of the inflation technique by proving incompatibility between \mathcal{T} and the W -type distribution; a quantum non-accessible distribution whose incompatibility is not witnessable by *any* entropic inequalities or any other known constraints [8]. Moreover, a family of polynomial, 2-outcome, compatibility inequalities are explicitly derived for an exemplary inflation of \mathcal{T} and 37 non-trivial representatives are printed in [8].

Identifying quantum-accessible distributions that are incompatible with the triangle scenario is of particular importance as it **Todo (TC Fraser): Discuss why we are doing this.**

As a preliminary search for quantum incompatibility in \mathcal{T} , we performed numerical optimizations using bipartite qubit states and 2-outcome POVM measurements (see appendix D for details) against the 37 compatibility inequalities printed in [8] as well as the entropic inequalities presented by [6]. Unfortunately, none of these inequalities could be

⁵ Indicating that conditional independence relations are not a sufficient characterization of compatibility for \mathcal{T} [6]. Note that in \mathcal{T} there are *zero* conditional independence relations over the observable nodes A, B, C .

violated. These early results suggest, although do not prove, that quantum non-locality is not present in \mathcal{T} for *two-outcome* measurements. We make *no claim* that this is a universal truth and simply take it as motivation to explore incompatibility in \mathcal{T} for a larger number of outcomes.

Specifically, we make use of the inflation technique for an inflated triangle scenario \mathcal{T}' depicted in fig. 2 and cast the *causal compatibility* problem as a *marginal problem*. We develop a new technique which we call the *extendable covering procedure* (ECP) to find a complete solution to the marginal problem that is suitable for large causal structures where the observable nodes are equipped with numerous outcomes. We use this technique to find numerous compatibility inequalities, some of which witness the incompatibility of the Fritz distribution. Before discussing the ECP, the Fritz distribution will be re-defined explicitly here, followed by a quick summary of the inflation technique of [8]. Following these summaries of existing work, we present the ECP and its connection to Hardy-type paradoxes [8–10], certificate inequalities and logical Bell inequalities [2]. Moreover, the ECP can be adapted to derive symmetric inequalities or inequalities bounding any subspace of marginal models. To conclude, we subject our found inequalities to non-linear optimizations and discuss the implications of our results.

V. THE FRITZ DISTRIBUTION

The **Fritz distribution** P_F is a quantum-accessible distribution known to be incompatible with the Triangle Scenario [7]. Explicitly, P_F is a three-party (A, B, C), four-outcome (1, 2, 3, 4) distribution that has form as follows:

$$\begin{aligned} P_F(111) = P_F(221) = P_F(412) = P_F(322) = P_F(233) = P_F(143) = P_F(344) = P_F(434) &= \frac{1}{32}(2 + \sqrt{2}) \\ P_F(121) = P_F(211) = P_F(422) = P_F(312) = P_F(243) = P_F(133) = P_F(334) = P_F(444) &= \frac{1}{32}(2 - \sqrt{2}) \end{aligned} \quad (3)$$

Here the notation $P_F(abc) = P_{ABC}(abc) = P(A = a, B = b, C = c)$ is used. The Fritz distribution P_F can be realized with the following quantum configuration:

$$\begin{aligned} \rho_{AB} &= |\Psi^+\rangle\langle\Psi^+| \quad \rho_{BC} = \rho_{CA} = |\Phi^+\rangle\langle\Phi^+| \\ M_A &= \{|0\psi_0\rangle\langle 0\psi_0|, |0\psi_\pi\rangle\langle 0\psi_\pi|, |1\psi_{-\pi/2}\rangle\langle 1\psi_{-\pi/2}|, |1\psi_{\pi/2}\rangle\langle 1\psi_{\pi/2}|\} \\ M_B &= \{|\psi_{\pi/4}0\rangle\langle\psi_{\pi/4}0|, |\psi_{5\pi/4}0\rangle\langle\psi_{5\pi/4}0|, |\psi_{3\pi/4}1\rangle\langle\psi_{3\pi/4}1|, |\psi_{-\pi/4}1\rangle\langle\psi_{-\pi/4}1|\} \\ M_C &= \{|00\rangle\langle 00|, |01\rangle\langle 01|, |10\rangle\langle 10|, |11\rangle\langle 11|\} \end{aligned} \quad (4)$$

Where for convenience of notation ψ_x is used to denote the superposition,

$$|\psi_x\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{ix}|1\rangle)$$

Additionally $|\Psi^+\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$ and $|\Phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ are two maximally entangled Bell states.

Fritz first proved it's incompatibility [7] by showing C acts a moderator to ensure measurement pseudo-settings for A and B are independent, satisfying non-broadcasting requirements for the standard Bell scenario. In fact, by coarse-graining outcomes for A and B and treating C as a measurement-setting moderator, P_F maximally violates the CHSH inequality. To illustrate this, begin with the CHSH inequality [11],

$$\langle AB|S_A = 1, S_B = 1\rangle + \langle AB|S_A = 1, S_B = 2\rangle + \langle AB|S_A = 2, S_B = 1\rangle - \langle AB|S_A = 2, S_B = 2\rangle \leq 2 \quad (5)$$

Where $\langle AB|S_A = i, S_B = j\rangle$ is the correlation between A and B given the measurement settings for A (B) is i (j) respectively. Next each of C 's outcomes become the conditioned settings in eq. (5),

$$\langle AB|C = 2\rangle + \langle AB|C = 3\rangle + \langle AB|C = 4\rangle - \langle AB|C = 1\rangle \leq 2 \quad (6)$$

Finally, specifying the two outcome coarse-graining as $\{1, 2, 3, 4\} \rightarrow \{(1, 4), (2, 3)\}$ gives a definition of correlation,

$$\langle AB\rangle \equiv 2\{P(11) + P(14) + P(41) + P(44) + P(22) + P(23) + P(32) + P(33)\} - 1$$

Which when applied to the Fritz distribution eq. (3) gives the following correlations:

$$\begin{aligned} \langle AB|C = 2\rangle &= 2\left\{\frac{1}{32}\left[2(2 + \sqrt{2}) + 2(2 - \sqrt{2})\right]\right\} - 1 = \frac{\sqrt{2}}{2} \\ \langle AB|C = 3\rangle &= \langle AB|C = 4\rangle = \frac{\sqrt{2}}{2} \quad \langle AB|C = 1\rangle = -\frac{\sqrt{2}}{2} \end{aligned}$$

Which when applied to eq. (6) gives the familiar Tsirelson violation of $2\sqrt{2} \not\leq 2$ [12].

Before continuing it is worth noting that eq. (3) is non-unique. Any distribution that is equal to eq. (3) via a permutation of outcomes or exchange of parties is also referred to as a Fritz distribution. Moreover, the quantum realization of eq. (4) is non-unique. In fact, eq. (4) is not the realization that Fritz originally had in mind. Nonetheless eq. (3) is taken as *the* Fritz distribution for concreteness throughout this paper.

In [7], Fritz posits the following question: *Find an example of non-classical quantum correlations in \mathcal{T} together with a proof of its non-classicality which does not hinge on Bell's Theorem.*

Todo (TC Fraser): Summarize Problem 2.17 in fritz BBT, make it more formal

VI. SUMMARY OF THE INFLATION TECHNIQUE

The causal inflation technique, first pioneered by Wolfe, Spekkens, and Fritz [8] and inspired by the *do calculus* and *twin networks* of Ref. [13], is a family of causal inference techniques that can be used to determine if a probability distribution is compatible or incompatible with a given causal structure. As a preliminary summary, the inflation technique begins by *augmenting* a causal structure with additional copies of its nodes, producing an *inflated* causal structure, and then exposes how causal inference tasks on the inflated causal structure can be used to make inferences on the original causal structure. Copies of the original nodes are distinguished by an additional subscript called the **copy-index**. For example node A of fig. 1 has copies A_1, A_2, A_3, A_4 in the inflation of fig. 2. Following the notions of Ref. [8], all such copies are deemed equivalent via a **copy-index equivalence** relation denoted ' \sim '. A copy-index is effectively arbitrary, so A' will refer to an arbitrary inflated copy of A .

$$A \sim A_1 \sim A' \not\sim B \sim B_1 \sim B'$$

We preemptively generalize the notion of copy-index equivalence to other mathematical objects like sets, graphs, and groups by saying that $X \sim Y$ if $X = Y$ upon removal of the copy-index. Equipped with the common graph-theoretic terminology and notation of definition 14, an inflation can be formally defined as follows:

Definition 29. An **inflation** of a causal structure \mathcal{G} is another causal structure \mathcal{G}' such that:

$$\forall n' \in \mathcal{N}' : \text{AnSub}_{\mathcal{G}'}(n') \sim \text{AnSub}_{\mathcal{G}}(n) \quad (7)$$

The motivation being that since the ancestry of each node n' in \mathcal{G}' plays the *exact same* role as the ancestry of its source copy n in \mathcal{G} , then every compatible joint distribution $P_{\mathcal{N}}$ on \mathcal{G} can be used to *create* compatible joint distributions on \mathcal{G}' . To do this, first notice that every compatible joint distribution $P_{\mathcal{N}}$ uniquely defines a set of causal parameters for \mathcal{G} .

$$\forall n \in \mathcal{N} : P_{n|\text{Pa}_{\mathcal{G}}(n)} = \sum_{\mathcal{N} \setminus n} P_{\mathcal{N}} \quad (8)$$

Now since eq. (7) enforces that $\text{Pa}_{\mathcal{G}'}(n') \sim \text{Pa}_{\mathcal{G}}(n)$, one can create a set of **inflated causal parameters** using eq. (8),

$$\forall n' \in \mathcal{N}' : P_{n'|\text{Pa}_{\mathcal{G}'}(n')} \equiv P_{n|\text{Pa}_{\mathcal{G}}(n)}$$

Which in turn uniquely defines a compatible joint distribution $P_{\mathcal{N}'}$ on \mathcal{G}' ⁶.

$$P_{\mathcal{N}'} = \prod_{n' \in \mathcal{N}'} P_{n'|\text{Pa}_{\mathcal{G}'}(n')}$$

This is known as the **weak inflation lemma**; compatible joint distributions over \mathcal{N} induce compatible joint distributions over \mathcal{N}' . Before generalizing to the inflation lemma, a careful observation needs to be made. For any pair of subsets of nodes $N \subseteq \mathcal{N}$ and $N' \subseteq \mathcal{N}'$ that are equivalent up to copy-index $N \sim N'$, if $\text{AnSub}_{\mathcal{G}}(N) \sim \text{AnSub}_{\mathcal{G}'}(N')$ then any compatible *marginal* distribution P_N over N induces a compatible marginal distribution $P_{N'}$ over N' . In fact since N' contains no duplicate nodes up to copy index (simply because $N \sim N'$ and N cannot contain duplicate nodes), $P_N = P_{N'}$. We refer such sets N' as the **injectable sets** of \mathcal{G}' and N the **images of the injectable sets** of \mathcal{G} .

$$\begin{aligned} \text{Inj}_{\mathcal{G}'}(\mathcal{G}') &\equiv \{N' \subseteq \mathcal{N}' \mid \exists N \subseteq \mathcal{N} : N \sim N'\} \\ \text{ImInj}_{\mathcal{G}}(\mathcal{G}') &\equiv \{N \subseteq \mathcal{N} \mid \exists N' \subseteq \mathcal{N}' : N \sim N'\} \end{aligned}$$

⁶ Of course not all compatible joint distributions on \mathcal{N}' are constructed in this way; all that is demonstrated is that all joint distributions constructed in this way are compatible.

Lemma 30. *The Inflation Lemma* *Given a particular inflation \mathcal{G}' of \mathcal{G} , if a marginal model $\{P_N \mid N \in \text{ImInj}_{\mathcal{G}}(\mathcal{G}')\}$ is compatible with \mathcal{G} then all marginal models $\{P_{N'} \mid N' \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')\}$ are compatible with \mathcal{G}' provided that $P_N = P_{N'}$ for all instances where $N \sim N'$. This is lemma 3 of [8].*

The inflation lemma is the most important result of the causal inflation technique [8]. The contrapositive version of lemma 30 is a powerful tool for determining compatibility. Any compatibility constraint on marginal models $\{P_{N'} \mid N' \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')\}$ of the inflated causal structure \mathcal{G}' correspond to valid compatibility constraints on marginal models $\{P_N \mid N \in \text{ImInj}_{\mathcal{G}}(\mathcal{G}')\}$ of the original causal structure. Corollary 5 of [8] proves this explicitly for incompatibility inequalities \mathcal{I} ; which the remainder of this work focuses on. Additionally, the inflation lemma holds when considering any subset of $\text{Inj}_{\mathcal{G}}(\mathcal{G}')$ (analogously $\text{ImInj}_{\mathcal{G}}(\mathcal{G}')$). Therefore, in situations where latent nodes are present in \mathcal{G} , one only needs to consider injectable sets that are composed of *observable nodes*.

In this work, we obtain inequalities that constrain the set of injectable marginal models $\{P_{N'} \mid N' \in \text{ImInj}_{\mathcal{G}}(\mathcal{G}')\}$ for the inflated Triangle Scenario of fig. 1 by considering the marginal problem (see section III) for the set of **maximally pre-injectable sets**⁷. A pre-injectable set V is a subset of \mathcal{N}' that can be decomposed into the disjoint union of injectable sets $V = \coprod_i N'_i \mid N'_i \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')$ that are mutually **ancestrally independent**,

$$\forall i, j : N'_i \perp N'_j \iff \text{An}_{\mathcal{G}'}(N'_i) \cap \text{An}_{\mathcal{G}'}(N'_j) = \emptyset$$

In doing so, any distribution over all nodes of a pre-injectable set $P_{\cup_i N'_i}$ will factorize according to graphical d -separation conditions [13],

$$P_V = P_{\cup_i N'_i} = \prod_i P_{N'_i}$$

This turns linear inequalities over the pre-injectable distributions into polynomial inequalities over the injectable distributions, allowing one to replace all such distributions with equivalent distributions over the original random variables of \mathcal{N} . $\text{PreInj}_{\mathcal{G}}(\mathcal{G}')$ will denote the set of all pre-injectable sets.

Focusing on the inflation \mathcal{T}' depicted in fig. 2, we obtained the injectable sets $\text{Inj}_{\mathcal{T}'}(\mathcal{T}')$ along with the maximally pre-injectable sets through a simple graphical procedure outlined in [8].

Maximal Pre-injectable Sets	Ancestral Independences
$\{A_1, B_1, C_1, A_4, B_4, C_4\}$	$\{A_1, B_1, C_1\} \perp \{A_4, B_4, C_4\}$
$\{A_1, B_2, C_3, A_4, B_3, C_2\}$	$\{A_1, B_2, C_3\} \perp \{A_4, B_3, C_2\}$
$\{A_2, B_3, C_1, A_3, B_2, C_4\}$	$\{A_2, B_3, C_1\} \perp \{A_3, B_2, C_4\}$
$\{A_2, B_4, C_3, A_3, B_1, C_2\}$	$\{A_2, B_4, C_3\} \perp \{A_3, B_1, C_2\}$
$\{A_1, B_3, C_4\}$	$\{A_1\} \perp \{B_3\} \perp \{C_4\}$
$\{A_1, B_4, C_2\}$	$\{A_1\} \perp \{B_4\} \perp \{C_2\}$
$\{A_2, B_1, C_4\}$	$\{A_2\} \perp \{B_1\} \perp \{C_4\}$
$\{A_2, B_2, C_2\}$	$\{A_2\} \perp \{B_2\} \perp \{C_2\}$
$\{A_3, B_3, C_3\}$	$\{A_3\} \perp \{B_3\} \perp \{C_3\}$
$\{A_3, B_4, C_1\}$	$\{A_3\} \perp \{B_4\} \perp \{C_1\}$
$\{A_4, B_1, C_3\}$	$\{A_4\} \perp \{B_1\} \perp \{C_3\}$
$\{A_4, B_2, C_1\}$	$\{A_4\} \perp \{B_2\} \perp \{C_1\}$

(9)

As can be counted, there are 12 maximally pre-injectable sets \mathcal{T}' which will be indexed 1 through 12 in the order seen above ($\{V_1, \dots, V_{12}\}$). The maximally pre-injectable sets $\{V_i\}_{i \in [12]}$ form a maximal marginal scenario \mathcal{M} over the observable nodes of \mathcal{T}' when equipped with outcome labels. Using lemma 30 inequalities $\mathcal{I}_{\text{PreInj}_{\mathcal{T}'}(\mathcal{T}')}$ constraining the set of contextual marginal models $P^{\mathcal{M}}$ can be deflated into polynomial inequalities over $\mathcal{I}_{\text{ImInj}_{\mathcal{T}'}(\mathcal{T}')}$ bounding compatible marginal models over the images of the injectable sets in \mathcal{T} .

$$\text{ImInj}_{\mathcal{T}}(\mathcal{T}') = \{\{A, B, C\}, \{A\}, \{B\}, \{C\}\}$$

⁷ The set of all pre-injectable sets forms a topological *simplicial complex*. ‘Maximal’ refers to the pre-injectable sets that are no proper subset of another.

VII. CERTIFICATE INEQUALITIES

A. Casting the Marginal Problem as a Linear Program

After obtaining the maximal pre-injectable sets associated with a particular inflation, one can write the marginal problem of definition 26 as a linear program. The key observation is that marginalization is a *linear* operator that can be performed via a matrix multiplication. To do this, we will define the *incidence matrix*.

Definition 31. The **incidence matrix** M for a marginal scenario $\mathcal{M} = \{V_1, \dots, V_k\}$ is a bit-wise matrix where the columns are indexed by *joint* outcomes $j \in \text{Out}(\mathcal{J})$ and the rows are indexed by *marginal* outcomes $m \in \text{Out}(\mathcal{M})$. The entries of M are populated whenever a row index is extendable to a column index.

$$M[m, j] = \delta_{m \prec j} \equiv \begin{cases} 1 & m \prec j \\ 0 & \text{otherwise} \end{cases}$$

Where $\delta_{m \prec j}$ is an *extendability indicator*. The incidence matrix has $|\text{Out}(\mathcal{J})|$ columns and $|\text{Out}(\mathcal{M})| = \sum_{i \in [k]} |\text{Out}(V_i)|$ rows. The number of non-zero entries of M is a simple expression,

$$\sum_{i \in [k]} |\text{Out}(V_i)| |\text{Out}(\mathcal{J} \setminus V_i)| = \sum_{i \in [k]} |\text{Out}(\mathcal{J})| = k |\text{Out}(\mathcal{J})|$$

This is due to the fact that each marginal context contributes a single non-zero entry to each column of M , resulting in $k |\text{Out}(\mathcal{J})|$ total non-zero entries.

Todo (TC Fraser): Computationally Efficient generation?

To illustrate this concretely, consider the following example:

Example 32. Let \mathcal{J} be 3 binary variables $\mathcal{J} = \{a, b, c\}$ and consider the marginal scenario $\mathcal{M} = \{\{a, c\}, \{b\}\}$. The incidence matrix becomes:

$$M = \begin{array}{c} \begin{matrix} (a,b,c) \mapsto \\ (a \mapsto 0, c \mapsto 0) \\ (a \mapsto 0, c \mapsto 1) \\ (a \mapsto 1, c \mapsto 0) \\ (a \mapsto 1, c \mapsto 1) \\ (b \mapsto 0) \\ (b \mapsto 1) \end{matrix} \end{array} \begin{pmatrix} \begin{matrix} (0,0,0) & (0,0,1) & (0,1,0) & (0,1,1) & (1,0,0) & (1,0,1) & (1,1,0) & (1,1,1) \end{matrix} \\ \begin{matrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \end{matrix} \end{pmatrix}$$

Example 33. The marginal problem for the inflated causal structure \mathcal{T}' depicted in fig. 2 concerns itself with the pre-injectable marginal scenario $\mathcal{M} = \text{PreInj}_{\mathcal{T}}(\mathcal{T}') = \{V_1, \dots, V_{12}\}$. Equipping the observed nodes of \mathcal{T}' with 4-element outcome labels, the incidence matrix M has 16,896 rows and 16,777,216 columns.

$$\begin{aligned} \# \text{ Rows} &= |\text{Out}(\mathcal{M})| = \sum_{i=1}^{12} |\text{Out}(V_i)| = \sum_{i=1}^{12} 4^{|V_i|} = 4 \cdot 4^6 + 8 \cdot 4^3 = 16,896 \\ \# \text{ Columns} &= |\text{Out}(\mathcal{J})| = 4^{|\mathcal{J}|} = 4^{12} = 16,777,216 \end{aligned}$$

With $12 \cdot 4^{12} = 201,326,592$ non-zero entries. For obvious reasons, no attempt is made to reproduce it here.

In order to describe how marginalization can be written as matrix multiplication $M \cdot x = b$, two more quantities need to be described:

Definition 34. The **joint distribution vector** $\mathcal{P}_{\mathcal{J}}$ for a probability distribution $P_{\mathcal{J}}$ is the vector whose entries are the positive, real-valued probabilities that $P_{\mathcal{J}}$ assigns to each joint outcome of $j \in \text{Out}(\mathcal{J})$. $\mathcal{P}_{\mathcal{J}}$ shares the same indices as the *column* indices of M .

$$\forall j \in \text{Out}(\mathcal{J}) : \mathcal{P}_{\mathcal{J}}^{\top}[j] = P_{\mathcal{J}}(j)$$

Definition 35. The **marginal distribution vector** $\mathcal{P}^{\mathcal{M}}$ for a marginal model $P^{\mathcal{M}} = \{P_{V_1}, \dots, P_{V_k}\}$ is the vector whose entries are probabilities over the set of marginal outcomes $\text{Out}(\mathcal{M})$. $\mathcal{P}^{\mathcal{M}}$ shares the same indices as the *row* indices of M .

$$\forall m \in \text{Out}(\mathcal{M}) : \mathcal{P}^{\mathcal{M}\top}[m] = P_{\text{Dom}(m)}(m)$$

Where $\text{Dom}(m) \in \mathcal{M}$ and $P_{\text{Dom}(m)} \in P^{\mathcal{M}}$ as per remark 5.

The marginal and joint distribution vectors are related via the incidence matrix M . Given a joint distribution vector $\mathcal{P}_{\mathcal{J}}$ one can obtain the marginal distribution vector $\mathcal{P}^{\mathcal{M}}$ by multiplying M by $\mathcal{P}_{\mathcal{J}}$.

$$\mathcal{P}^{\mathcal{M}} = M \cdot \mathcal{P}_{\mathcal{J}} \quad (10)$$

Remark 36. The particular ordering of the rows and columns of M carries no importance, but it *must* be consistent between M , $\mathcal{P}_{\mathcal{J}}$ and $\mathcal{P}^{\mathcal{M}}$. Throughout this work, a *canonical ordering* is used: the marginal contexts are written in the same order as listed in the marginal scenario \mathcal{M} and outcomes within a marginal context are enumerated in a *raveling* order; exemplified in example 32.

The marginal problem can now be rephrased in the language of the incidence matrix. Suppose one obtains a marginal distribution vector $\mathcal{P}^{\mathcal{M}}$. The marginal problem becomes equivalent to the question: *Does there exist a joint distribution vector $\mathcal{P}_{\mathcal{J}}$ such that eq. (10) holds?*

Definition 37. The Marginal Linear Program is the following linear program:

$$\begin{aligned} &\text{minimize: } \emptyset \cdot x \\ &\text{subject to: } x \succeq 0 \\ &\quad M \cdot x = \mathcal{P}^{\mathcal{M}} \end{aligned}$$

If this “optimization”⁸ is *feasible*, then there exists a vector x than can satisfy eq. (10) and is a valid joint distribution vector. Therefore feasibility implies that $P_{\mathcal{J}} = x$, solving the marginal problem with positive result. Moreover if the marginal linear program is *infeasible*, then there *does not* exist a joint distribution $P_{\mathcal{J}}$ over all random variables.

Definition 38. The Dual Marginal Linear Program is the dual of definition 37 formulated via a procedure similar to [14]:

$$\begin{aligned} &\text{minimize: } y \cdot \mathcal{P}^{\mathcal{M}} \\ &\text{subject to: } y \cdot M \succeq 0 \end{aligned}$$

Where y is a real valued vector with the same length as $\mathcal{P}^{\mathcal{M}}$.

B. Infeasibility Certificates

The dual marginal linear program also provides an answer to the marginal problem. To prove this, first notice that the dual problem is *never infeasible*; by choosing y to be trivially the null vector \emptyset of appropriate size, all constraints are satisfied. Secondly if $y \cdot M \succeq 0$ and $x \succeq 0$, then the following must hold if the primal is feasible:

$$y \cdot \mathcal{P}^{\mathcal{M}} = y \cdot M \cdot x \geq 0 \quad (11)$$

Therefore the *sign* of the dual value $d \equiv \min(y \cdot \mathcal{P}^{\mathcal{M}})$ solves the marginal problem. If $d < 0$ then eq. (11) is violated and therefore the marginal problem has negative result. Likewise if d satisfies eq. (11), then a joint distribution $P_{\mathcal{J}}$ exists. Before continuing, an important observation needs to be made. If $d \geq 0$, then it is exactly $d = 0$, due to the existence of the trivial $y = \emptyset$. This observation is an instance of the *Complementary Slackness Property* of [15]. **Comment (TC Fraser): Is this really the CSP?** Moreover, if $d < 0$, then it is unbounded $d = -\infty$. This latter point becomes clear upon recognizing that for any y such that $d < 0$, another y' can be constructed by multiplying y by a real constant α greater than one such that,

$$y' = \alpha y \mid \alpha > 1 \implies d' = \alpha d < d$$

Since a more negative d' can always be found, it must be that d is unbounded. This is a demonstration of the fundamental *Unboundedness Property* of [15]; if the dual is unbounded, then the primal is infeasible.

Comment (TC Fraser): Farkas's lemma here?

⁸ “Optimization” is presented in quotes here because the minimization objective is trivially always zero (\emptyset denotes the null vector of all zero entries). The primal value of the linear program is of no interest, all that matters is its *feasibility*.

Definition 39. An **infeasibility certificate** [16] is any vector y that satisfies the constraints of definition 38 and also permits violation of eq. (11) for some marginal distribution vector $\mathcal{P}^{\mathcal{M}}$.

$$y \in \mathbb{R}^{|\text{Out}(\mathcal{M})|} : y \cdot M \succeq 0, \quad y \cdot \mathcal{P}^{\mathcal{M}} < 0$$

Furthermore, any y satisfying $y \cdot M \succeq 0$ induces a **certificate inequality** that constraints the space of marginal distribution vectors which takes the symbolic form of eq. (11),

$$y \cdot \mathcal{P}^{\mathcal{M}} \geq 0$$

Where the entries of the certificate y act as coefficients for the entries of $\mathcal{P}^{\mathcal{M}}$.

Todo (TC Fraser): Discuss Infeasibility Certificates basis **Todo (TC Fraser): Discuss the certificate inequalities we found.**

$$\mathcal{I}_{\text{Mosek-Cert}}, \mathcal{I}_{\text{Cert}}$$

VIII. COVERING INEQUALITIES

Section VII discusses how one can obtain a valid compatibility inequality that witnesses incompatibility for a particular marginal model $P^{\mathcal{M}}$ by writing the marginal problem as a linear program. It was demonstrated that every incompatible distribution is witness-able by some certificate inequality. However, when a particular marginal model $P^{\mathcal{M}}$ is not known, a linear optimization offers no utility. **Todo (TC Fraser): Motivate why certificates are not enough, want many solutions, want logical foundation**

In this section, we will first summarize a method that can be used to obtain a complete solution to the *possibilistic* marginal problem and indirectly offer a *partial* solution to the *probabilistic* marginal problem. Doing so corresponds to enumerating all

A. Logical Covers

In this section, we summarize the familiar notion of a logical cover. Suppose one has a particular marginal outcome $a \in \text{Out}(\mathcal{M})$ and a marginal model $P^{\mathcal{M}}$ that assigns the probability $P(a)$ to the outcome a . If the marginal model $P^{\mathcal{M}}$ is to admit a joint distribution $P_{\mathcal{J}}$, then it must be that $P(a)$ is equal to a marginalization over a 's extendable set into \mathcal{J} .

$$P(a) = \sum_{j \in \text{Ext}_{\mathcal{J}}(a)} P_{\mathcal{J}}(j) \quad (12)$$

This corresponds to row a of the incidence matrix M . Next consider a subset of marginal outcomes $C \subseteq \text{Out}(\mathcal{M})$. Summing over the associated probabilities $\{P(c)\}_{c \in C}$ and applying eq. (12) yields,

$$\sum_{c \in C} P(c) = \sum_{c \in C} \sum_{j \in \text{Ext}_{\mathcal{J}}(c)} P_{\mathcal{J}}(j) \quad (13)$$

If for all joint distributions $P_{\mathcal{J}}$ it can be demonstrated that the following holds,

$$\sum_{c \in C} \sum_{j \in \text{Ext}_{\mathcal{J}}(c)} P_{\mathcal{J}}(j) \geq \sum_{j \in \text{Ext}_{\mathcal{J}}(a)} P_{\mathcal{J}}(j) \quad (14)$$

Then combining eqs. (12) to (14) yields a valid compatibility inequality for \mathcal{M} .

$$P(a) \leq \sum_{c \in C} P(c) \quad (15)$$

In order for eq. (14) to hold for *all* joint distributions $P_{\mathcal{J}}$, it must hold on an algebraic level; each term on the right-hand side of eq. (14) also appears on the left-hand side of eq. (14).⁹ Therefore, identifying all compatibility inequalities that can be written in the form of eq. (15) corresponds to identifying all sets $C \subseteq \text{Out}(\mathcal{M})$ such that their joint distribution terms algebraically *cover* the joint distribution terms for $a \in \text{Out}(\mathcal{M})$.

The notion of forming a

⁹ If there were a joint outcome j^* present on the right-hand side of eq. (14) but not the left, then eq. (14) would be violated by the deterministic distribution $P_{\mathcal{J}}(j^*) = 1$

B. Extendable Covers

We will now generalize the notion of a logical cover by considering more exotic antecedents corresponding to multiple marginal outcomes $A \subseteq \text{Out}(\mathcal{M})$ along with weights Γ_A for each outcome. Afterwards we demonstrate that these generalized covers also admit non-trivial probabilistic compatibility inequalities.

Definition 40. A **multiplicity set** or **m-set** Γ_S for a marginal scenario \mathcal{M} is a function $\Gamma_S : \text{Out}(\mathcal{M}) \rightarrow \mathbb{Z}_{\geq 0}$ that assigns to each marginal outcome $m \in \text{Out}(\mathcal{M})$ a *multiplicity* $\Gamma_S(m) \in \mathbb{Z}_{\geq 0}$. The subscript S is a subset of marginal outcome $S \subseteq \text{Out}(\mathcal{M})$ indicating which marginal outcomes receive non-zero multiplicity.

$$\forall s \in S : \Gamma_S(s) \in \mathbb{N} \quad \forall m \in \text{Out}(\mathcal{M}) \setminus S : \Gamma_S(m) = 0$$

We define the **extension of an m-set** Γ_S into \mathcal{J} to be the m-set Γ_{E_S} which assigns multiplicities to *joint* outcomes $j \in \text{Out}(\mathcal{J})$. The extension of S denoted E_S is the set of all joint outcomes that are extensions of marginal outcomes $s \in S$.

$$E_S = \bigcup_{s \in S} \text{Ext}_{\mathcal{J}}(s)$$

While the multiplicity of each element $j \in E_S$ corresponds to the sum over multiplicities $\Gamma_S(s)$ where $j \in \text{Ext}_{\mathcal{J}}(s)$.

$$\forall j \in E_S : \Gamma_{E_S}(j) = \sum_{s \in S} \Gamma_S(s) \delta_{s \prec j} \quad (16)$$

With all other joint outcomes receiving a zero multiplicity $\forall j \notin E_S : \Gamma_{E_S}(j) = 0$.

Definition 41. An m-set Γ_C is an **extendable cover** of an m-set Γ_A for a marginal scenario \mathcal{M} if and only if the following two conditions hold¹⁰:

$$E_A \subseteq E_C \quad (17)$$

$$\forall j \in E_A : \Gamma_{E_A}(j) \leq \Gamma_{E_C}(j) \quad (18)$$

Identifying extendable covers is of great importance because of the following theorem.

Theorem 42. *If the m-set Γ_C is an extendable cover of the m-set Γ_A for a marginal scenario \mathcal{M} , then the following is a valid compatibility $\mathcal{I}_{\mathcal{M}}$ for the marginal scenario \mathcal{M} .*

$$\sum_{a \in A} \Gamma_A(a) P(a) \leq \sum_{c \in C} \Gamma_C(c) P(c) \quad (19)$$

Proof. To prove that eq. (19) is a valid compatibility inequality $\mathcal{I}_{\mathcal{M}}$ for all marginal models $P^{\mathcal{M}}$, we will demonstrate that eq. (19) follows from the assumption that a joint distribution $P_{\mathcal{J}}$ exists.

For each m-set Γ_S write the weighted sum over the probabilities $\{P(s)\}_{s \in S}$ in terms of the joint distribution $P_{\mathcal{J}}$ using eq. (16).

$$\begin{aligned} \sum_{s \in S} \Gamma_S(s) P(s) &= \sum_{s \in S} \Gamma_S(s) \sum_{j \in \text{Ext}_{\mathcal{J}}(s)} P_{\mathcal{J}}(j) \\ &= \sum_{s \in S} \Gamma_S(s) \sum_{j \in \text{Out}(\mathcal{J})} \delta_{s \prec j} P_{\mathcal{J}}(j) \\ &= \sum_{j \in \text{Out}(\mathcal{J})} \left[\sum_{s \in S} \Gamma_S(s) \delta_{s \prec j} \right] P_{\mathcal{J}}(j) \\ &= \sum_{j \in E_S} \Gamma_{E_S}(j) P_{\mathcal{J}}(j) \end{aligned} \quad (20)$$

¹⁰ In fact eq. (18) implies eq. (17) since $\forall j \notin E_A : \Gamma_{E_A}(j) = 0$.

First examine the substitution $S \rightarrow C$,

$$\sum_{c \in C} \Gamma_C(c) P(c) = \sum_{j \in E_C} \Gamma_{E_C}(j) P_{\mathcal{J}}(j)$$

By eq. (17) $E_A \subseteq E_C$,

$$\sum_{c \in C} \Gamma_C(c) P(c) = \sum_{j \in E_A} \Gamma_{E_C}(j) P_{\mathcal{J}}(j) + \sum_{j \in E_C \setminus E_A} \Gamma_{E_C}(j) P_{\mathcal{J}}(j)$$

Notice that the sum over $j \in E_C \setminus E_A$ is entirely positive since $\Gamma_{E_C}(j) \in \mathbb{N}$ and $P_{\mathcal{J}}(j) \geq 0$. Moreover eq. (18) gives $\Gamma_{E_C}(j) \geq \Gamma_{E_A}(j)$. Combining these observations along with eq. (20) for $S \rightarrow A$, one arrives at the inequality,

$$\sum_{c \in C} \Gamma_C(c) P(c) \geq \sum_{j \in E_A} \Gamma_{E_A}(j) P_{\mathcal{J}}(j) = \sum_{a \in A} \Gamma_A(a) P(a)$$

Which is equivalent to eq. (19); every non-contextual marginal model $P^{\mathcal{M}}$ must satisfy eq. (19). \square

C. Complete Solution to The Marginal Problem

Theorem 43.

D. Extendable Cover Procedure

The **extendable cover procedure** begins with an antecedent m-set Γ_A and iteratively produces consequent m-sets Γ_C that are extendable covers of Γ_A . To facilitate this, we define a **partial consequent m-set** to be any marginal m-set Γ_S that *has yet to* become an extendable cover of Γ_A . Also define the null m-set Γ_{\emptyset} to be the m-set assigning zero multiplicity to each marginal outcome.

Algorithm 1 The Extendable Cover Procedure (ECP)

$$\Gamma_S^0 = \Gamma_{\emptyset}$$

Todo (TC Fraser): Trivial nodes Todo (TC Fraser): Duplication under Deflation Todo (TC Fraser): Memory Todo (TC Fraser): Parallelization Todo (TC Fraser): Simplicial Complex

1. Optimizations

2. Avoiding Trivial Inequalities

A **trivial compatibility inequality** $\mathcal{I}_{\mathcal{M}}$ is an inequality that is satisfied by all marginal models $P^{\mathcal{M}}$, not just compatible ones. Trivial compatibility inequalities are of no importance Since the set of all extendable cover inequalities
 Todo (TC Fraser):

E.

F. Irreducible Covers

Todo (TC Fraser): Basis certificates

G. Limited Factorization

H. Logical Implications & Inequalities

Following the work conducted by Mansfield and Fritz [10], we consider a possibilistic implications of the form,

$$a \implies c_1 \vee \cdots \vee c_m = \bigvee_{i \in [m]} c_i \quad (21)$$

Where a and each of the c_i 's are simply events or outcomes of a particular set of variables. The letter ' a ' is chosen for the event a since it takes the place of the logical **antecedent** of eq. (21). Likewise, the letter ' c ' is chosen to represent logical **consequents**. We refer to the set of all c_i 's simply as $C = \{c_i \mid i \in [m]\}$. The implication eq. (21) can be read as *whenever a occurs, at least one element of c also occurs*.

It is possible to turn possibilistic implications into probabilistic inequalities by recognizing that the logical implication of eq. (21) induces the inequality,

$$P(a) \leq P\left(\bigvee_{i=1}^m c_i\right)$$

Furthermore utilizing Boole's inequality,

$$P\left(\bigvee_{i=1}^m c_i\right) \leq \sum_{i=1}^m P(c_i) \quad (22)$$

Gives,

$$P(a) \leq \sum_{i=1}^m P(c_i) \quad (23)$$

Such that whenever the inequality eq. (23) is violated, the implication in eq. (21) is violated as well. Note that the converse is *not* true; if the inequality eq. (23) holds true, it is still possible for there to be a violation of eq. (21).

Remark 44. An important result of Boole's inequality is that eq. (22) becomes an exact *equality* whenever elements of C are pairwise disjoint. Therefore, finding a set C of pairwise disjoint events satisfying eq. (21) will give rise to tighter inequalities.

I. Implications in the Marginal Problem

The question then remains, *how does non-contextuality give rise to implications of the form of eq. (21)?* Begin by making the principle assumption that a joint distribution *does* exist for a marginal model $P^{\mathcal{M}}$. This assumption induces logical *tautologies* that take the form of eq. (21), which we call **marginal implications**, using the following train of logic.

1. Suppose a particular marginal outcome $a \in O_{\mathcal{M}}$ happens to occur.
2. Since a joint distribution exists, then a refers to some *incomplete* knowledge about a *joint* event $j = o[\mathcal{J}] \in O_{\mathcal{J}}$ that actually occurred. More precisely, this set of possible j 's is the extendable set of a in $O_{\mathcal{J}}$.

$$j \in \text{Ext}_{\mathcal{J}}(a)$$

3. Therefore whenever a occurs, one of the elements j of $\text{Ext}_{\mathcal{J}}(a)$ *has* to occur.

$$a \implies \bigvee_{j \in \text{Ext}_{\mathcal{J}}(a)} j \quad (24)$$

4. Now suppose we could obtain a set of marginal outcomes $C = \{c_1, \dots, c_m\}$ each different from a such that for every j , $\text{Ext}(c \rightarrow j)$ for at least one element $c \in C$. If such a C can be found, then the possibility of at least one j occurring implies the possibility of at least one c occurring.

$$\bigvee_{j \in \text{Ext}_{\mathcal{J}}(a)} j \implies c_1 \vee \cdots \vee c_m = \bigvee_{i \in [m]} c_i \quad (25)$$

5. Combing eq. (24) with eq. (25), one obtains eq. (21).

Todo (TC Fraser): Talk about why this is called hardy paradox

Finding all marginal implications for a chosen marginal outcome a corresponds to finding a set of marginal outcomes $C = \{c_1, \dots, c_m\}$ whose extendable sets *cover* $\text{Ext}_{\mathcal{J}}(a)$. Formally this corresponds to a **set covering** problem **Todo (TC Fraser): cite** which we elected to cast as the equivalent **hypergraph transversal problem**.

Todo (TC Fraser): Mention sufficient solution to the possibilistic marginal problem **Todo (TC Fraser): Illustrate how it can distinguish more than possibilistic differences**

Given an antecedent a , we can construct a hypergraph \mathcal{H}_a called the **marginal hypergraph** with nodes \mathcal{N}_a and \mathcal{E}_a . The nodes of this hypergraph are the subset of marginal outcomes $\mathcal{N}_a \subseteq O^{\mathcal{M}}$ *compatible* with a . The edges are labeled by outcomes in $\text{Ext}_{\mathcal{J}}(a) \subseteq O_{\mathcal{J}}$, namely the set of j 's, and contain all nodes compatible with j .

$$\mathcal{N}_a = \{n \in O^{\mathcal{M}} \mid \text{Com}(n, a)\} \quad (26)$$

$$\mathcal{E}_a = \{\{n \in O^{\mathcal{M}} \mid \text{Com}(n, j)\}, j \in \text{Ext}_{\mathcal{J}}(a)\} \quad (27)$$

Remark 45. Note that every node is contained in some edge. To prove this, consider the definition of compatibility between outcomes. If every n is compatible with a , then there exists some outcome $j \in \text{Ext}_{\mathcal{J}}(A)$ such that $\text{Ext}(n \rightarrow j)$. Therefore since $\text{Ext}(n \rightarrow j)$, $\text{Com}(n, j)$; satisfying the central condition of eq. (27).

J. Higher Order Marginal Implications

Todo (TC Fraser): Discuss (m, n) - type implications and the non-triviality **Todo (TC Fraser): Link to logical bell inequalities/completeness or not?**

K. Hypergraph Transversals

Section VIII demonstrates how finding possibilistic marginal implications can be casted as a hypergraph transversal problem. This section aims to summarize the general idea behind existing algorithms and also discusses some of the inequalities found using these algorithms. **Todo (TC Fraser): References here**

Definition 46. A **hypergraph transversal generation** is any algorithm that correctly generates the complete set of all minimal transversals of \mathcal{H} .

Definition 47. Any hypergraph with strictly non-empty edges $\forall e \neq \emptyset$ will always admit the **trivial transversal** \mathcal{T}^* where all nodes are considered as members $\mathcal{T}^* = \mathcal{N}$. Any hypergraph with empty edges will be called a **degenerate hypergraph** as it admits no transversals¹¹.

Remark 45 guarantees that all marginal hypergraphs are non-degenerate. There are two distinct approaches to hypergraph transversal generation: *top-down* and *bottom-up*.

Definition 48. A **top-down** hypergraph transversal generation refers to any algorithm that begins with the trivial transversal \mathcal{T}^* and iteratively removes unnecessary nodes from \mathcal{T}^* .

Definition 49. A **bottom-up** hypergraph transversal generation refers to any algorithm that begins with the empty set \emptyset and iteratively adds nodes.

One should select a top-down method if the typical size of minimal transversals $|\mathcal{T}|$ is comparable to the number of nodes $|\mathcal{N}|$, otherwise a bottom-up method will perform better. For our purposes we implemented a deep-first transversal algorithm similar to Recalling remark 45

L. Weighted Hypergraph Transversals

In section VII is was demonstrated that the Fritz distribution is witness-able via a certificate inequality. It is also possible to witness **Todo (TC Fraser): Discuss the Inequalities Derived/ Trivial and non-trivial**

Todo (TC Fraser): Weighted transversals and Optimizations **Todo (TC Fraser): Seeding inequalities (huge advantage here)**

¹¹ Most authors require that all hypergraphs be non-degenerate [17].

IX. DERIVING SYMMETRIC INEQUALITIES

Symmetric compatibility inequalities are useful for a number of reasons. First, Bancal et. al. [18] discuss computational advantages in considering symmetric versions of marginal polytopes mentioned in section III; the number of extremal points typically grows exponentially in \mathcal{J} , but only polynomial for the symmetric polytope. They also note a number of interesting inequalities (such as CHSH [11]) can be written in a way that is symmetric under the exchange of parties, demonstrating that quantum-non-trivial inequalities can be recovered from facets of a symmetric polytope. Second, numerical optimizations against inequalities invariant under exchange of parties will lead to one of two interesting cases: either the extremal distribution is invariant under exchange of parties or it is not. The latter case generates a family of incompatible distributions obtained by exchanging parties of the found extremal distribution¹². In this section we discuss how to achieve computational advantage and its application to the inequality techniques mentioned in section VII and section VIII H.

A. Causal Symmetry

The aim of this section is to formalize what types of symmetries are present in a particular causal structure \mathcal{G} . The group of these causal symmetries will define, for each compatibility inequality $\mathcal{I}_{\mathcal{M}}$, a family of inequalities that are also valid incompatibility inequalities.

First consider the permutation group acting on a set of nodes $N \subseteq \mathcal{N}$ denoted $\text{Perm}(N)$ to be the set of all bijective maps φ from N to N .

$$\text{Perm}(N) = \{\varphi \mid \varphi : N \rightarrow N\}$$

The action of $\varphi \in \text{Perm}(N)$ on a causal structure \mathcal{G} is defined via an extension of φ to the corresponding element in $\text{Perm}(\mathcal{N})$ that leaves nodes $n \in \mathcal{N} \setminus N$ invariant,

$$\begin{aligned}\varphi[\mathcal{G}] &\equiv (\varphi[\mathcal{N}], \varphi[\mathcal{E}]) \\ \varphi[\mathcal{N}] &\equiv \{\varphi[n] \mid n \in \mathcal{N}\} = \mathcal{N} \\ \varphi[\mathcal{E}] &\equiv \{\varphi[e] \mid e \in \mathcal{E}\}\end{aligned}$$

To motivate a more general treatment of such symmetries, consider the Triangle Scenario of fig. 1. Due to the rotational and reflective symmetries of the Triangle Scenario, any permutation $\varphi \in \text{Perm}(\mathcal{N}_O)$ of the observable nodes $\mathcal{N}_O = \{A, B, C\}$ creates a new causal structure $\varphi[\mathcal{G}]$ that is *equivalent* to \mathcal{G} up to a relabeling of *latent* nodes. By construction, any valid compatibility inequality $\mathcal{I}_{\mathcal{M}}$ for the marginal scenario $\mathcal{M} = \{V_1, \dots, V_k\}$ is *independent* of the labeling of latent nodes. Therefore applying φ to all distributions in $\mathcal{I}_{\mathcal{M}}$ yields $\varphi[\mathcal{I}_{\mathcal{M}}] \equiv \varphi[\mathcal{I}]_{\varphi[\mathcal{M}]}$, another valid compatibility inequality for the permuted marginal scenario $\varphi[\mathcal{M}]$ defined as,

$$\varphi[\mathcal{M}] = \{\varphi[V_1], \dots, \varphi[V_k]\}$$

More generically, any permutation φ that preserves the graphical structure of \mathcal{G} can be applied to $\mathcal{I}_{\mathcal{M}}$ to generate new and valid compatibility inequalities. Specifically this alludes to the **graph automorphism group** $\text{Aut}(\mathcal{G})$ of \mathcal{G} .

$$\text{Aut}(\mathcal{G}) \equiv \{\varphi \in \text{Perm}(\mathcal{N}) \mid \varphi[\mathcal{G}] = \mathcal{G}\}$$

In general, $\text{Aut}(\mathcal{G})$ could include elements φ that map latent nodes to observable nodes¹³. This behaviour is *undesired* for causal inference where the latent nodes will never appear in a marginal scenario. Instead, the subgroup of $\text{Aut}(\mathcal{G})$ that never maps observable nodes to latent nodes is the **set-wise stabilizer** of $\text{Aut}(\mathcal{G})$ for \mathcal{N}_O and will be expressed with a subscript.

Definition 50. The **causal symmetry group** for a causal structure \mathcal{G} is the graph automorphism subgroup that stabilizes the observable nodes.

$$\text{Aut}_{\mathcal{N}_O}(\mathcal{G}) = \{\varphi \in \text{Perm}(\mathcal{N}) \mid \varphi[\mathcal{G}] = \mathcal{G}, \forall n \in \mathcal{N}_O : \varphi[n] \in \mathcal{N}_O\}$$

Given any compatibility inequality $\mathcal{I}_{\mathcal{M}}$, each element φ of the causal symmetry group defines a new valid compatibility inequality $\varphi[\mathcal{I}_{\mathcal{M}}] = \varphi[\mathcal{I}]_{\varphi[\mathcal{M}]}$.

¹² If the extremal distribution is not invariant under exchange of parties, there is indication that the space of accessible distributions is non-convex.

¹³ In practice however, it is rare to be considering a causal structure \mathcal{G} where some latent node $n_L \in \mathcal{N}_L$ takes on a role indistinguishable from some observable node $n_O \in \mathcal{N}_O$, so this won't be an issue.

The inflation technique discussed in section VI allows one to derive compatibility inequalities for a causal structure \mathcal{G} by considering the marginal problem over the pre-injectable sets of \mathcal{G}' denoted $\text{PreInj}_{\mathcal{G}}(\mathcal{G}') = \{V_1, \dots, V_k\}$. It is important to recall that due to lemma 30, inequalities \mathcal{I}' for \mathcal{G}' are *only* transferable to inequalities \mathcal{I} for \mathcal{G} if \mathcal{I}' is in terms of distributions over the pre-injectable sets $\mathcal{I}'_{\text{PreInj}_{\mathcal{G}}(\mathcal{G})}$. As a consequence of this observation, we need to consider a subgroup of the causal symmetry group that preserves the pre-injectable sets,

$$\varphi[\text{PreInj}_{\mathcal{G}}(\mathcal{G}')] = \text{PreInj}_{\mathcal{G}}(\mathcal{G}') \quad (28)$$

Definition 51. The **restricted causal symmetry group** Φ for a causal structure \mathcal{G} and a marginal scenario \mathcal{M} is the graph automorphism subgroup that stabilizes the marginal scenario \mathcal{M} .

$$\Phi \equiv \text{Aut}_{\mathcal{M}}(\mathcal{G}) = \{\varphi \in \text{Perm}(\mathcal{N}) \mid \varphi[\mathcal{G}] = \mathcal{G}, \forall V \in \mathcal{M} : \varphi[V] \in \mathcal{M}\}$$

Given any compatibility inequality $\mathcal{I}_{\mathcal{M}}$ each element φ of the causal symmetry group defines a new valid compatibility inequality $\varphi[\mathcal{I}_{\mathcal{M}}] = \mathcal{I}_{\mathcal{M}}$ over the *same* marginal context \mathcal{M} .

The restricted causal symmetry group for the pre-injectable sets of an inflation \mathcal{G}' is the ideal symmetry group for use with the inflation technique. If one obtains a particular inequality $\mathcal{I}'_{\text{PreInj}_{\mathcal{G}}(\mathcal{G})}$ as well as the restricted causal symmetry group Φ , an entire family of valid compatibility inequalities are also obtained.

$$\left\{ \varphi[\mathcal{I}']_{\text{PreInj}_{\mathcal{G}}(\mathcal{G}')} \mid \varphi \in \Phi \right\} \text{ where } \Phi = \text{Aut}_{\text{PreInj}_{\mathcal{G}}(\mathcal{G}')}(\mathcal{G}')$$

We have obtained Φ for the inflated triangle scenario fig. 2. Φ is an order 48 group with the following 4 generators:

φ_1	φ_2	φ_3	φ_4
$A_1 \rightarrow A_4$	$A_1 \rightarrow A_1$	$A_1 \rightarrow C_1$	$A_1 \rightarrow A_1$
$A_2 \rightarrow A_3$	$A_2 \rightarrow A_3$	$A_2 \rightarrow C_2$	$A_2 \rightarrow A_2$
$A_3 \rightarrow A_2$	$A_3 \rightarrow A_2$	$A_3 \rightarrow C_3$	$A_3 \rightarrow A_3$
$A_4 \rightarrow A_1$	$A_4 \rightarrow A_4$	$A_4 \rightarrow C_4$	$A_4 \rightarrow A_4$
$B_1 \rightarrow B_4$	$B_1 \rightarrow C_1$	$B_1 \rightarrow A_1$	$B_1 \rightarrow B_2$
$B_2 \rightarrow B_3$	$B_2 \rightarrow C_3$	$B_2 \rightarrow A_2$	$B_2 \rightarrow B_1$
$B_3 \rightarrow B_2$	$B_3 \rightarrow C_2$	$B_3 \rightarrow A_3$	$B_3 \rightarrow B_4$
$B_4 \rightarrow B_1$	$B_4 \rightarrow C_4$	$B_4 \rightarrow A_4$	$B_4 \rightarrow B_3$
$C_1 \rightarrow C_4$	$C_1 \rightarrow B_1$	$C_1 \rightarrow B_1$	$C_1 \rightarrow C_3$
$C_2 \rightarrow C_3$	$C_2 \rightarrow B_3$	$C_2 \rightarrow B_2$	$C_2 \rightarrow C_4$
$C_3 \rightarrow C_2$	$C_3 \rightarrow B_2$	$C_3 \rightarrow B_3$	$C_3 \rightarrow C_1$
$C_4 \rightarrow C_1$	$C_4 \rightarrow B_4$	$C_4 \rightarrow B_4$	$C_4 \rightarrow C_2$

It is easy to verify that $\varphi_1, \varphi_2, \varphi_3, \varphi_4$ are all automorphisms of the inflated Triangle Scenario \mathcal{G}' of fig. 2. Moreover they stabilize the observable nodes, and the map pre-injectable sets of eq. (9) to the pre-injectable sets. Finally, in the Triangle Scenario we have that,

$$\Phi = \text{Aut}_{\text{PreInj}_{\mathcal{G}}(\mathcal{G}')}(\mathcal{G}') \sim \text{Aut}_{\mathcal{N}_O}(\mathcal{G})$$

To see this, φ_1 and φ_4 become the identity element \mathbb{I} in $\text{Aut}_{\mathcal{N}_O}(\mathcal{G})$ upon removal of the copy-index, leaving φ_2 to generate reflections and φ_3 to generate rotations.

B. Symmetric Marginal Polytope

Up until this point, we have discussed how new inequalities can be created from known ones by exploiting causal symmetry. Although very useful, we divert our attention to finding *symmetric* inequalities themselves.

$$\forall \varphi \in \Phi : \varphi[\mathcal{I}_{\mathcal{M}}] = \mathcal{I}_{\mathcal{M}}$$

In order to do this, we must first define how elements $\varphi \in \Phi$ act on marginal outcomes $O_{\mathcal{M}}$.

Definition 52. Let $\mathcal{J} = \{v_1, \dots, v_k\}$ be a collection of random variables and let $\varphi \in \text{Perm}(\mathcal{J})$. \mathcal{J} is said to be **label invariant with respect to φ** if the outcome labels for each variable $v \in \mathcal{J}$ are unchanged under the action of φ .

$$\forall v \in \mathcal{J} : O_v = O_{\varphi[v]}$$

Label invariance is a prerequisite for any kind of symmetry among random variables.

Definition 53. Let $\mathcal{J} = (v_1, \dots, v_k)$ be a collection of random variables along with a specific outcome $m \in \text{Out}(V)$ where $V \subseteq \mathcal{J}$. The action of $\varphi \in \text{Perm}(\mathcal{J})$ on m is a new outcome $\varphi[m] \in \text{Out}(\varphi[V])$ over $\varphi[V] \subseteq \mathcal{J}$ defined as follows,

$$\forall v \in V : (\varphi[m])(\varphi[v]) \equiv m(v)$$

The idea being that φ modifies m in such a way that permutes its domain while leaving its range invariant.

Example 54. Consider the outcome $m = \{A \mapsto 0, B \mapsto 1, C \mapsto 2\} \in \text{Out}(\{A, B, C\})$ where all variables have 4 possible outcomes $O_A = O_B = O_C = \{0, 1, 2, 3\}$ and the permutation $\varphi = \{A \rightarrow D, B \rightarrow C, C \rightarrow B, D \rightarrow A\}$. The action of φ on m is as follows,

$$\begin{aligned} \varphi[m] &= \varphi[\{A \mapsto m(A), B \mapsto m(B), C \mapsto m(C)\}] \\ &= \{\varphi[A] \mapsto m(A), \varphi[B] \mapsto m(B), \varphi[C] \mapsto m(C)\} \\ &= \{D \mapsto m(A), C \mapsto m(B), B \mapsto m(C)\} \\ &= \{D \mapsto 0, C \mapsto 1, B \mapsto 2\} \\ &= \{B \mapsto 2, C \mapsto 1, D \mapsto 0\} \end{aligned}$$

Evidently, $\varphi[m] \in \text{Out}(\{B, C, D\})$.

Through repeated action of $\varphi \in \Phi$ on marginal outcomes $m \in \text{Out}(\mathcal{M})$ and joint outcomes $j \in \text{Out}(\mathcal{J})$, one can define group orbits of Φ in $\text{Out}(\mathcal{M})$ and $\text{Out}(\mathcal{J})$.

$$\begin{aligned} \text{Orb}_{\Phi}(m) &\equiv \{\varphi[m] \mid \varphi \in \Phi\} \\ \text{Orb}_{\Phi}(j) &\equiv \{\varphi[j] \mid \varphi \in \Phi\} \end{aligned}$$

Definition 55. The **symmetric incidence matrix** M_{Φ} for a marginal scenario \mathcal{M} and a particular restricted causal symmetry Φ is a contracted version of the incidence matrix M for \mathcal{M} . Each row of M_{Φ} corresponds to a marginal orbit $\text{Orb}_{\Phi}(m)$. Analogously each column of M_{Φ} corresponds to a joint orbit $\text{Orb}_{\Phi}(j)$. The entries of M_{Φ} are integers and correspond to summing over the rows and columns of M that belong to each orbit.

$$M_{\Phi}[\text{Orb}_{\Phi}(m), \text{Orb}_{\Phi}(j)] = \sum_{\substack{j' \in \text{Orb}_{\Phi}(j) \\ m' \in \text{Orb}_{\Phi}(m)}} M[m', j']$$

C. Symmetric Logical Implications

D. Results

Todo (TC Fraser): Identify the desired symmetry group
 Todo (TC Fraser): How we obtained the desired symmetry group
 Todo (TC Fraser): Group orbits to symmetric marginal description matrix
 Todo (TC Fraser): Infeasibility on symmetric marginal problem
 Todo (TC Fraser): Hardy Transversals can't work on the symmetric marginal problem
 Todo (TC Fraser): Symmetrizing non-symmetric inequalities through avoiding orbits
 Todo (TC Fraser): higher order transversals on mutually impossible events

X. NON-LINEAR OPTIMIZATIONS

Todo (TC Fraser): Mention outlook on the importance of classification and determining interesting inequalities [18]
 Compatibility inequalities for a given causal structure are fantastic for finding incompatible distributions. In the inflation technique, this is no exception. Parameterizing a space distributions using a set of real-valued parameters λ , enables us to perform numerical optimizations against these inequalities in hopes that a particular set of parameters λ is able to generate an incompatible distribution P . To illustrate this generic procedure and its reliability, we will first examine the popular CHSH inequality.




FIG. 3. Convex optimizations against $\mathcal{I}_{\text{CHSH}}$ recover algebraic violation of 4.




FIG. 4. Quantum optimizations against $\mathcal{I}_{\text{CHSH}}$ recover maximum violation of $2\sqrt{2}$.

A. Numerical Violations of The CHSH Inequality

The CHSH inequality [11] can be viewed as a causal compatibility inequality for the iconic Bell Scenario (Fig. 19 of [19], Fig. 11 of [8], Fig. 1 a) of [20], etc.) corresponding to Bell's notion of local causality [19]. It constrains the set of 2-outcome bipartite distributions over local binary measurement settings for each party $P_{AB|S_AS_B} \equiv \{P_{AB|00}, P_{AB|01}, P_{AB|10}, P_{AB|11}\}$. Numerical optimization *should* obtain the algebraic violation associated with the PR-Box correlations [21]. Maintaining full generality, we simply need to parameterize these 4 distributions using eq. (E2), each requiring 4 real-valued parameters. We define the optimization target for the CHSH inequality to be the left-hand-side of eq. (5),

$$\mathcal{I}_{\text{CHSH}} = \langle AB|11 \rangle + \langle AB|12 \rangle + \langle AB|21 \rangle - \langle AB|22 \rangle$$

Figure 3 demonstrates this optimization for 5 random seed parameters λ^0 , each converging to the expected value of 4. Analogously, [12]

Todo (TC Fraser): Demonstrate Quantum, Convexity Todo (TC Fraser): Why Inequalities are great for optimizations Todo (TC Fraser): Non-linearity Todo (TC Fraser): Techniques Used Todo (TC Fraser): Finding maximum violation of CHSH easily Todo (TC Fraser): Unreliance when number of parameters increases Todo (TC Fraser): Issues with local minimum Todo (TC Fraser): Using initial conditions close to fritz, obtain greater violation Todo (TC Fraser): Greater violation shares possibilistic structure of fritz and violates CHSH under definition Todo (TC Fraser): Not realizable with maximally entangled qubit states Todo (TC Fraser): Not realizable with separable measurements Todo (TC Fraser): Many non-trivial inequalities to be tested Todo (TC Fraser): inequality \rightarrow dist \rightarrow inequality evolution

XI. CONCLUSIONS

Todo (TC Fraser): Inflation technique allows one to witness fritz incompatibility Todo (TC Fraser): Linear optimization induces certificates which are incompatibility witnesses Todo (TC Fraser): There are quantum distributions in the Triangle Scenario that are incompatible and different from fritz in terms of entanglement but not possibilistic structure

XII. OPEN QUESTIONS & FUTURE WORK

The marginal problem has been well studied in various contexts [1, 8, 22, ...]. Reference [1] contains an excellent summary of its connection to other fields of research including knowledge integration, database theory and coalition games in game theory. **Todo (TC Fraser): Lots of stuff**

Appendix A: Exemplary Inequalities

Appendix B: Connections to Sheaf-Theoretic Treatment

Appendix C: Computationally Efficient Parametrization of the Unitary Group

Spengler, Huber and Hiesmayr [23] suggest the parameterization of the unitary group $\mathcal{U}(d)$ using a $d \times d$ -matrix of real-valued parameters $\lambda_{n,m}$,

$$U = \left[\prod_{m=1}^{d-1} \left(\prod_{n=m+1}^d \exp(iP_n \lambda_{n,m}) \exp(i\sigma_{m,n} \lambda_{m,n}) \right) \right] \cdot \left[\prod_{l=1}^d \exp(iP_l \lambda_{l,l}) \right] \quad (\text{C1})$$

Where P_l are one-dimensional projective operators,

$$P_l = |l\rangle\langle l| \quad (\text{C2})$$

and the $\sigma_{m,n}$ are generalized anti-symmetric σ -matrices,

$$\sigma_{m,n} = -i|m\rangle\langle n| + i|n\rangle\langle m|$$

Where $1 \leq m < n \leq d$. Spengler et. al. proved the validity of eq. (C1) in Ref. [23].

For the sake of reference, let us label the matrix exponential terms in eq. (C1) in a manner that corresponds to their affect on an orthonormal basis $\{|1\rangle, \dots, |d\rangle\}$.

$$\begin{aligned} GP_l &= \exp(iP_l \lambda_{l,l}) \\ RP_{n,m} &= \exp(iP_n \lambda_{n,m}) \\ R_{m,n} &= \exp(i\sigma_{m,n} \lambda_{m,n}) \end{aligned} \quad (\text{C3})$$

It is possible to remove the reliance on matrix exponential operations in eq. (C1) by utilizing the explicit form of the exponential terms in eq. (C3). As a first step, recognize the defining property of the projective operators eq. (C2),

$$P_l^k = (|l\rangle\langle l|)^k = |l\rangle\langle l| = P_l$$

This greatly simplifies the global phase terms GP_l ,

$$GP_l = \exp(iP_l \lambda_{l,l}) = \sum_{k=0}^{\infty} \frac{(iP_l \lambda_{l,l})^k}{k!} = \mathbb{I} + \sum_{k=1}^{\infty} \frac{(i\lambda_{l,l})^k}{k!} P_l^k = \mathbb{I} + P_l \left[\sum_{k=1}^{\infty} \frac{(i\lambda_{l,l})^k}{k!} \right] = \mathbb{I} + P_l (e^{i\lambda_{l,l}} - 1) \quad (\text{C4})$$

Analogously for the relative phase terms $RP_{n,m}$,

$$RP_{n,m} = \dots = \mathbb{I} + P_n (e^{i\lambda_{n,m}} - 1) \quad (\text{C5})$$

Finally, the rotation terms $R_{m,n}$ can also be simplified by examining powers of $i\sigma_{m,n}$,

$$R_{m,n} = \exp(i\sigma_{m,n} \lambda_{m,n}) = \sum_{k=0}^{\infty} \frac{(|m\rangle\langle n| - |n\rangle\langle m|)^k \lambda_{m,n}^k}{k!}$$

One can verify that the following properties hold,

$$\begin{aligned} &(|m\rangle\langle n| - |n\rangle\langle m|)^0 = \mathbb{I} \\ \forall k \in \mathbb{N}, k \neq 0 : &(|m\rangle\langle n| - |n\rangle\langle m|)^{2k} = (-1)^k (|m\rangle\langle m| + |n\rangle\langle n|) \\ \forall k \in \mathbb{N} : &(|m\rangle\langle n| - |n\rangle\langle m|)^{2k+1} = (-1)^k (|m\rangle\langle n| - |n\rangle\langle m|) \end{aligned}$$

Revealing the simplified form of $R_{m,n}$,

$$R_{m,n} = \mathbb{I} + (|m\rangle\langle m| + |n\rangle\langle n|) \sum_{j=1}^{\infty} (-1)^j \frac{\lambda_{n,m}^{2j}}{(2j)!} + (|m\rangle\langle n| - |n\rangle\langle m|) \sum_{j=0}^{\infty} (-1)^j \frac{\lambda_{n,m}^{2j+1}}{(2j+1)!}$$

$$R_{m,n} = \mathbb{I} + (|m\rangle\langle m| + |n\rangle\langle n|)(\cos \lambda_{n,m} - 1) + (|m\rangle\langle n| - |n\rangle\langle m|) \sin \lambda_{n,m} \quad (\text{C6})$$

By combining the optimizations of eqs. (C5) to (C4) together we arrive at an equivalent form for eq. (C1) that is computational more efficient.

$$U = \left[\prod_{m=1}^{d-1} \left(\prod_{n=m+1}^d R P_{n,m} R_{m,n} \right) \right] \cdot \left[\prod_{l=1}^d G P_l \right] \quad (\text{C7})$$

In quantum mechanics, the global phase of a state $|\psi\rangle \in \mathcal{H}^n$ is a *redundant* parameter. Parameterizing unitaries using eq. (C7) is especially attractive since the global phase terms $G P_l$ can be dropped, allowing one to parameterize all unitaries in $\mathcal{U}(d)$ up to this degeneracy [23]¹⁴.

$$U_{/GP_l} = \left[\prod_{m=1}^{d-1} \left(\prod_{n=m+1}^d R P_{n,m} R_{m,n} \right) \right] \quad (\text{C8})$$

Todo (TC Fraser): Explanation of Computational Complexity $\mathcal{O}(d^3)$ vs. $\mathcal{O}(1)$ using [24] Todo (TC Fraser): Pre-Caching for Fixed dimension d Todo (TC Fraser): Talk about inverse via haar measure

Appendix D: Parametrization of Quantum States & Measurements

Throughout section X, we utilize a variety of parameterizations of quantum states and measurements in order to generate quantum-accessible probability distributions. There are numerous techniques that can be used when parameterizing quantum states and measurements [23, 25–28] with applications Todo (TC Fraser): Finish this sentence. For our purposes, we need to parameterize the space of quantum-accessible distributions P_Q that are *realized* on the Triangle Scenario. We have implemented P_Q under the following description.

$$P_{ABC}(abc) = \text{Tr}[\Omega^\top \rho_{AB} \otimes \rho_{BC} \otimes \rho_{CA} \Omega M_{A,a} \otimes M_{B,b} \otimes M_{C,c}] \quad (\text{D1})$$

1. Quantum States

The bipartite states $(\rho_{AB}, \rho_{BC}, \rho_{CA})$ of eq. (D1) were taken to be two-qubit density matrices acting on $\mathcal{H}^2 \otimes \mathcal{H}^2$.¹⁵ The space of all such states corresponds to the space of all 4×4 positive semi-definite hermitian matrices with unitary trace. Throughout this section, we refer to these bipartite states simply as ρ unless otherwise indicated. There are three distinct techniques that we have considered.

Taking inspiration from [28], we can parameterize all such density matrices ρ using **Cholesky Parametrization** [29]. The Cholesky decomposition allows one to write any hermitian positive semi-definite matrix ρ in terms of a lower (or upper) triangular matrix T using $\rho = T^\dagger T$. Our Cholesky parameterization consists of assigning 16 real-valued parameters λ to the entries of T and generating a unitary trace ρ similar to eq. (4.4) of [28].

$$\rho = \frac{T^\dagger T}{\text{Tr}(T^\dagger T)} \quad T = \begin{bmatrix} \lambda_1 & 0 & 0 & 0 \\ \lambda_2 + i\lambda_3 & \lambda_4 & 0 & 0 \\ \lambda_5 + i\lambda_6 & \lambda_7 + i\lambda_8 & \lambda_9 & 0 \\ \lambda_{10} + i\lambda_{11} & \lambda_{12} + i\lambda_{13} & \lambda_{14} + i\lambda_{15} & \lambda_{16} \end{bmatrix} \quad (\text{D2})$$

Our deviation from exclusively using eq. (D2) is two-fold. First, eq. (D2) is degenerate in that the normalization indicates only $16 - 1 = 15$ parameters are required for fully generic parameterization of all such states ρ . Removing

¹⁴ In our implementation, we accomplish this by explicitly setting $\lambda_{l,l} = 0$ in eq. (C4)

¹⁵ We also considered qutrit \mathcal{H}^3 qutrit \mathcal{H}^4 states. However for 6 d -dimensional \mathcal{H}^d states, the joint density matrix ρ acts on $(\mathcal{H}^d)^{\otimes 6}$ making it a (d^6, d^6) matrix with d^{12} entries. Computationally only $d = 2$ was feasible for our optimization tasks.

this degeneracy is possible although difficult. Second, the parameters λ_i carry no physical meaning associated with the state ρ , unlike our next parameterization.

In Spengler, Huber and Hiesmayr's work [23], they discuss how to parameterization density matrices ρ acting on \mathcal{H}^d of rank k through it's spectral decomposition,

$$\rho = \sum_{i=1}^k p_i |\psi_i\rangle\langle\psi_i| \quad p_i \geq 0, \sum_i p_i = 1, k \leq d \quad (\text{D3})$$

Where any orthonormal basis $\{|\psi_i\rangle\}$ of \mathcal{H}^d can be transformed into a computational basis $\{|i\rangle\}$ by a unitary $U \in \mathcal{U}(d)$ such that $|\psi_i\rangle = U|i\rangle$. We refer to eq. (D3) as the **Spengler Parametrization**. Without loss of generality we parameterize all full-rank ($k = d$) matrices by simultaneously parameterizing the $d = 4$ eigenvalues p_i of eq. (D3) using eq. (E1) and the unitary group $\mathcal{U}(4)$ up to global phase equivalence using eq. (C8). Parameterizing ρ using the Spengler parameterization requires $3 + 12 = 15$ parameters; admitting no degeneracies.

Finally in cases where we wish to restrict ourselves to *pure* bipartite states $\rho = |\psi\rangle\langle\psi|$, we have the luxury to use a **Schmidt Parametrization**. This is accomplished via a Schmidt decomposition $|\psi_{AB}\rangle = \sum_i \sigma_i |i_A\rangle \otimes |i_B\rangle$ where normalization demands that $\sum_i \sigma_i^2 = 1$, $\{|i_A\rangle\}$ and $\{|i_B\rangle\}$ are orthonormal bases for \mathcal{H}_A and \mathcal{H}_B respectively [30]. Additionally for qubit sources we can *choose* our orthonormal bases to be the computational basis $\{|0\rangle, |1\rangle\}$ and write,

$$|\psi\rangle = \cos^2(\lambda_1)|0\rangle \otimes |0\rangle + \sin^2(\lambda_1)|1\rangle \otimes |1\rangle$$

Where only 1 real-valued parameter $\lambda = \{\lambda_1\}$ is required to parameterize all pure states up to local unitaries. Pure states are also attractive due to their computational advantage in computing eq. (D1). If each state ρ is decomposable into $|\psi\rangle\langle\psi|$, then eq. (D1) can be written as,

$$P_{ABC}(abc) = \langle\psi_{AB}\psi_{BC}\psi_{CA}|\Omega M_{A,a} \otimes M_{B,b} \otimes M_{C,c}\Omega^\top|\psi_{AB}\psi_{BC}\psi_{CA}\rangle \quad (\text{D4})$$

Avoiding the expensive matrix multiplications of eq. (D1).

2. Measurements

With full generality, we consider a measurement M to be a **projective-operator valued measure (POVM)** represented by a set of hermitian, positive semi-definite operators $\{M_i\}_{i=1,\dots,k}$ ¹⁶ acting on \mathcal{H}^d summing to the identity,

$$\forall |\phi\rangle \in \mathcal{H}^d : \langle\phi|M_i|\phi\rangle \geq 0 \quad \sum_{i=1}^k M_i = \mathbb{I}_{\mathcal{H}^d} \quad (\text{D5})$$

When considering $k = 2$ outcome measurements acting on \mathcal{H}^4 we parameterize the first POVM element M_1 by using a Cholesky parameterization similar to eq. (D2) without normalizing for trace. Afterwards, M_2 is fully determined by eq. (D5).

$$M_1 = T^\dagger T \quad M_2 = \mathbb{I} - T^\dagger T \quad (\text{D6})$$

However, in order for M_2 to be positive semi-definite, the largest eigenvalue of M_1 has to be less than 1. To see this is a necessary and sufficient constraint, first expand out eq. (D6),

$$\langle\phi|M_2|\phi\rangle = \langle\phi|\mathbb{I} - M_1|\phi\rangle = |\phi|^2 - \langle\phi|\left(\sum_{i=1}^d m_1^{(i)}|m_1^{(i)}\rangle\langle m_1^{(i)}|\right)|\phi\rangle$$

Next write a generic $|\phi\rangle \in \mathcal{H}^d$ in terms of a linear combination of the eigenvectors of M_1 ¹⁷.

$$\langle\phi|M_2|\phi\rangle = \sum_j \left|\langle\phi|m_1^{(j)}\rangle\right|^2 - \sum_i m_1^{(i)} \left|\langle\phi|m_1^{(i)}\rangle\right|^2 = \sum_i \left(1 - m_1^{(i)}\right) \left|\langle\phi|m_1^{(i)}\rangle\right|^2 \quad (\text{D7})$$

¹⁶ The i -th element of M is referenced using a subscript M_i . The set of measurement elements for a particular party X will be written M_X . When both party and element are to be referenced, we write $M_{X,i}$.

¹⁷ The eigenvectors of M_1 form an orthonormal basis for \mathcal{H}^d because M_1 is Hermitian

Since $|\phi\rangle$ is arbitrary, for each i set $|\phi\rangle = |m_1^{(i)}\rangle$ to see that each eigenvalue of M_1 needs to be less than 1.

$$\langle m_1^{(i)} | M_2 | m_1^{(i)} \rangle = (1 - m_1^{(i)}) \geq 0 \implies m_1^{(i)} \leq 1 \quad (\text{D8})$$

By eq. (D7) is not difficult to see that eq. (D8) is a sufficient condition. During optimization, eq. (D8) can either be enforced passively as a constraint or directly by normalizing M_1 by its largest eigenvalue $\max(m_1^{(i)})$ whenever necessary.

When generalizing the above parameterization to more than 2 outcomes, only necessary conditions were found. Generating k -outcome POVM measurements is doable using rejection sampling techniques such as those used in [31] however a valid parameterization with little to no degeneracy was not found. Upon making this observation, a necessary departure to **projective-valued measures (PVMs)** is warranted¹⁸. With loss of generality, consider the set of PVMs M satisfying eq. (D5) in addition to orthogonal and projective properties,

$$M_i M_j = \delta_{ij} M_i \quad M_i = |m_i\rangle\langle m_i| \quad (\text{D9})$$

Parameterizing M for k -outcome measurements corresponds to parameterizing the set of all k -th order orthonormal sub-bases of \mathcal{H}^d . First note that any such basis $\{|\psi_1\rangle, \dots, |\psi_k\rangle\}$ can be transformed into the computational basis $\{|1\rangle, \dots, |k\rangle\}$ by a unitary denoted $U \in \mathcal{U}(d)$,

$$U|\psi_i\rangle = |i\rangle$$

With this observation we just need to parameterize the set of all unitaries $\mathcal{U}(d)$,

$$M = \{U|i\rangle\langle i|U^\dagger\}_{i \in 1, \dots, k}$$

Specifically, the projective property each M_i means that the global phase of U is completely arbitrary; one only needs to consider parameterizing unitaries up to global phase eq. (C8). This method was inspired by the *measurement seeding* method of Pál and Vértesi's [33] iterative optimization technique.

Analogously to eq. (D4), projective measurements offer considerable computational advantage as eq. (D1) can be rewritten as,

$$P_{ABC}(abc) = \langle m_{A,a} m_{B,b} m_{C,c} | \Omega^\top \rho_{AB} \otimes \rho_{BC} \otimes \rho_{CA} \Omega | m_{A,a} m_{B,b} m_{C,c} \rangle \quad (\text{D10})$$

3. Network Permutation Matrix

Finally, we introduce the **network permutation matrix** Ω for the Triangle Scenario of fig. 1. For bipartite qubit states, Ω becomes a 64×64 bit-wise matrix that acts on the measurements M and is depicted in fig. 5. To illuminate its necessity, consider eq. (D1) without Ω .

$$\begin{aligned} P_{ABC}(abc) &\stackrel{?}{=} \text{Tr}[(\rho_{AB} \otimes \rho_{BC} \otimes \rho_{CA})(M_A^a \otimes M_B^b \otimes M_C^c)] \\ &= \text{Tr}[(\rho_{AB} M_A^a) \otimes (\rho_{BC} M_B^b) \otimes (\rho_{CA} M_C^c)] \\ &= \text{Tr}(\rho_{AB} M_A^a) \text{Tr}(\rho_{BC} M_B^b) \text{Tr}(\rho_{CA} M_C^c) \\ &= P_{A|\rho_{AB}}(a) P_{B|\rho_{BC}}(b) P_{C|\rho_{CA}}(c) \end{aligned}$$

On an operational level, this corresponds to A making a measurement on *both* subsystems of ρ_{AB} and *not* on any component of ρ_{CA} . This is analogously troubling for B and C as well. The network permutation matrix Ω corresponds to *aligning* the underlying 6-qubit joint state ρ with the joint measurement M . To understand its effect, consider its effect on 6-qubit pure state $|q_1\rangle \otimes \dots \otimes |q_6\rangle = |q_1 q_2 q_3 q_4 q_5 q_6\rangle$ where $\forall i : |q_i\rangle \in \mathcal{H}^2$.

$$\Omega |q_1 q_2 q_3 q_4 q_5 q_6\rangle = |q_2 q_3 q_4 q_5 q_6 q_1\rangle$$

Ω acts as a *partial transpose* on $(\mathcal{H}^2)^{\otimes 6}$ by shifting the underlying tensor structure one subsystem to the “left”. It is uniquely defined by its action on all 2^6 orthonormal basis elements of $(\mathcal{H}^2)^{\otimes 6}$,

$$\Omega \equiv \sum_{|q_i\rangle \in \{|0\rangle, |1\rangle\}} |q_2 q_3 q_4 q_5 q_6 q_1\rangle \langle q_1 q_2 q_3 q_4 q_5 q_6|$$

¹⁸ Strictly speaking, when the number of outcomes (k) *matches* the Hilbert space dimension (d), eq. (D5) implies eq. (D9) by completeness. When considering 4 outcome measurements on bipartite qubit states in \mathcal{H}^4 , PVMs are completely general. Moreover, Naimark's dilation theorem guarantees that PVMs acting on \mathcal{H}^q can *emulate* the behaviour of any POVM acting on \mathcal{H}^d provided that q is sufficiently larger than d [32].




FIG. 5. The network permutation matrix Ω for $(\mathcal{H}^2)^{\otimes 6}$ realized on the triangle scenario. Black represents a value of 1 and 0 otherwise.

4. Degeneracy

Todo (TC Fraser): Discuss local unitary degeneracy

Appendix E: Convex Parametrization of Finite Probability Distributions

As discussed in section X, there is a need to parameterize the family of all probability distributions P_V over a given set of variables $V = (v_1, \dots, v_{|V|})$. If the cardinality of O_V is finite, then this is computationally feasible. The space of probability distributions over $n = |O_V|$ distinct outcomes forms a $n - 1$ dimensional convex polytope naturally embedded in $\mathbb{R}_{\geq 0}^n$ [34] that is parameterizable by $n - 1$ real value parameters; normalization $\sum_{o[V] \in O_V} P_V(o[V]) = 1$ accounts for the ‘-1’. An example of a non-degenerate parameterization of P_V consists of $n - 1$ parameters $\lambda = (\lambda_1, \dots, \lambda_{n-1})$, $\lambda_i \in [0, \pi/2]$ which generate the n probability values p_j using hyperspherical coordinates [23, 26],

$$p_j = \cos^2 \lambda_j \prod_{i=1}^{j-1} \sin^2 \lambda_i \quad \forall j \in 1, \dots, n-1$$

$$p_n = \prod_{i=1}^{n-1} \sin^2 \lambda_i$$
(E1)

Furthermore due to the periodicity of the parameter space λ , eq. (E1) can be used for either constrained or unconstrained optimization problems. For continuity reasons, unconstrained optimizations are performed whenever possible.

Although non-degenerate, this parameterization suffers from uniformity; a randomly sampled vector of parameters λ *does not* translate to a randomly sampled probability P_V . An easy-to-implement, degenerate parameterization of P_V can be constructed by simply beginning with n real parameters $\lambda = (\lambda_1, \dots, \lambda_n)$, then making them positive and normalized by their sum¹⁹.

$$p_j = \frac{|\lambda_j|}{\sum_{i=1}^n |\lambda_i|} \quad \forall j \in 1, \dots, n$$
(E2)

¹⁹ Strictly speaking, eq. (E2) *also* suffers from non-uniformity; being biased toward uniform probability distributions P_V . Todo (TC Fraser): Discuss rejection sampling simplex algorithms

For various convex optimization tasks sensitive to initial conditions outlined section X, the latter parameterization of eq. (E2) generally performed better than the former eq. (E1).

-
- [1] Tobias Fritz and Rafael Chaves, “Entropic inequalities and marginal problems,” (2011), [10.1109/TIT.2012.2222863](#), [arXiv:1112.4788](#).
 - [2] Samson Abramsky and Adam Brandenburger, “The sheaf-theoretic structure of non-locality and contextuality,” (2011), [10.1088/1367-2630/13/11/113036](#), [arXiv:1102.0264](#).
 - [3] Bastian Steudel and Nihat Ay, “Information-theoretic inference of common ancestors,” (2010), [arXiv:1010.5720](#).
 - [4] Rafael Chaves, Lukas Luft, and David Gross, “Causal structures from entropic information: geometry and novel scenarios,” *New Journal of Physics* **16**, 043001 (2014).
 - [5] Cyril Branciard, Denis Rosset, Nicolas Gisin, and Stefano Pironio, “Bilocal versus nonbilocal correlations in entanglement-swapping experiments,” *Phys. Rev. A* **85** (2012), [10.1103/physreva.85.032119](#).
 - [6] Joe Henson, Raymond Lal, and Matthew F Pusey, “Theory-independent limits on correlations from generalized bayesian networks,” *New Journal of Physics* **16**, 113043 (2014), [arXiv:1405.2572](#).
 - [7] Tobias Fritz, “Beyond bell’s theorem: Correlation scenarios,” (2012), [10.1088/1367-2630/14/10/103001](#), [arXiv:1206.5115](#).
 - [8] Elie Wolfe, Robert W. Spekkens, and Tobias Fritz, “The inflation technique for causal inference with latent variables,” (2016).
 - [9] Yeong-Cherng Liang, Robert W. Spekkens, and Howard M. Wiseman, “Specker’s parable of the overprotective seer: A road to contextuality, nonlocality and complementarity,” *Physics Reports* **506**, 1–39 (2011).
 - [10] Shane Mansfield and Tobias Fritz, “Hardy’s non-locality paradox and possibilistic conditions for non-locality,” (2011), [10.1007/s10701-012-9640-1](#), [arXiv:1105.1819](#).
 - [11] John F. Clauser, Michael A. Horne, Abner Shimony, and Richard A. Holt, “Proposed experiment to test local hidden-variable theories,” *Phys. Rev. Lett.* **23**, 880–884 (1969).
 - [12] B. S. Cirel’son, “Quantum generalizations of bell’s inequality,” *Lett Math Phys* **4**, 93–100 (1980).
 - [13] Judea Pearl, *Causality: Models, Reasoning, and Inference* (Cambridge University Press, 2009).
 - [14] Sébastien Lahaie, “How to take the dual of a linear program,” (2008).
 - [15] Bradley Hax, and Magnanti, *Applied Mathematical Programming* (Addison-Wesley, 1977) Chap. 4, pp. 143–144.
 - [16] Erling D. Andersen, *Computational Optimization and Applications* **20**, 171–183 (2001).
 - [17] Dimitris J. Kavvadias and Elias C. Stavropoulos, “An efficient algorithm for the transversal hypergraph generation,” *JGAA* **9**, 239–264 (2005).
 - [18] Jean-Daniel Bancal, Nicolas Gisin, and Stefano Pironio, “Looking for symmetric bell inequalities,” *Journal of Physics A: Mathematical and Theoretical* **43**, 385303 (2010).
 - [19] Christopher J. Wood and Robert W. Spekkens, “The lesson of causal discovery algorithms for quantum correlations: Causal explanations of bell-inequality violations require fine-tuning,” (2012), [10.1088/1367-2630/17/3/033002](#), [arXiv:1208.4119](#).
 - [20] Armin Tavakoli, “Quantum correlations in connected multipartite bell experiments,” (2015), [10.1088/1751-8113/49/14/145304](#), [arXiv:1509.08491](#).
 - [21] D. Rohrlich and S. Popescu, “Nonlocality as an axiom for quantum theory,” (1995), [arXiv:quant-ph/9508009](#).
 - [22] N. N. Vorob’ev, “Consistent families of measures and their extensions,” *Theory of Probability & Its Applications* **7**, 147–163 (1962).
 - [23] Christoph Spengler, Marcus Huber, and Beatrix C. Hiesmayr, “A composite parameterization of unitary groups, density matrices and subspaces,” (2010), [10.1088/1751-8113/43/38/385306](#), [arXiv:1004.5252](#).
 - [24] Cleve Moler and Charles Van Loan, “Nineteen dubious ways to compute the exponential of a matrix, twenty-five years later,” *SIAM Rev.* **45**, 3–49 (2003).
 - [25] Dénes Petz and László Ruppert, “Optimal quantum-state tomography with known parameters,” (2015), [10.1088/1751-8113/45/8/085306](#), [arXiv:1511.06666](#).
 - [26] S. R. Hedemann, “Hyperspherical parameterization of unitary matrices,” (2013), [arXiv:1303.5904](#).
 - [27] Kazuyuki Fujii, Kunio Funahashi, and Takayuki Kobayashi, “Jarlskog’s parametrization of unitary matrices and qudit theory,” (2005), [arXiv:quant-ph/0508006](#).
 - [28] Daniel F. V. James, Paul G. Kwiat, William J. Munro, and Andrew G. White, “Measurement of qubits,” *Phys. Rev. A* **64** (2001), [10.1103/physreva.64.052312](#).
 - [29] Markus Grasmair, “On the existence of a cholesky factorization,” (2014).
 - [30] Michael A. Nielsen and Isaac L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition* (Cambridge University Press, 2011) Chap. The Schmidt decomposition and purifications, p. 109.
 - [31] Dénes Petz and László Ruppert, “Optimal quantum-state tomography with known parameters,” *Journal of Physics A: Mathematical and Theoretical* **45**, 085306 (2012).
 - [32] Asher Peres, *Quantum Theory: Concepts and Methods (Fundamental Theories of Physics)* (Springer, 1995) Chap. 9.6, p. 285.
 - [33] Károly F. Pál and Tamás Vértesi, “Maximal violation of a bipartite three-setting, two-outcome bell inequality using infinite-dimensional quantum systems,” *Phys. Rev. A* **82** (2010), [10.1103/physreva.82.022116](#).
 - [34] Nicolas Brunner, Daniel Cavalcanti, Stefano Pironio, Valerio Scarani, and Stephanie Wehner, “Bell nonlocality,” (2013),

[10.1103/RevModPhys.86.419](#), [arXiv:1303.2849](#).