

Causal Compatibility Inequalities Admitting of Quantum Violations in the Triangle Scenario

T.C. Fraser^{1,2}


¹Perimeter Institute for Theoretical Physics
Ontario, Canada

²University of Waterloo
Ontario, Canada


Quantum Networks Workshop, IIP, Nov. 2016

1. Thank International Institute for Physics (IIP) for support
2. Thank Perimeter Institute for Theoretical Physics (PI) for support

The Triangle Scenario (TS)



└ The Triangle Scenario (TS)



1. Research project's goal was to derive new causal compatibility inequalities that distinguish quantum correlations from classical correlations
2. Triangle Scenario has been studied extensively before
3. In Branciard et al. 2012 it was noted that characterizing locality in TS remained an open problem and that identifying compatibility constraints in this configuration **seemed challenging**
4. In Fritz 2012, Fritz demonstrated that TS is the **smallest** correlation scenario in which there exists quantum incompatible distributions (proof without inequalities)
5. In Henson, Lal, and Pusey 2014, TS was classified as an **interesting** causal structure: conditional independence relations are not a sufficient characterization of compatibility (there are none)
6. Several other authors (see Steudel and Ay 2010, Chaves, Luft, and Gross 2014, Wolfe, Spekkens, and Fritz 2016, ...) have investigated TS without achieving research objective

- 1 Report the discovery of such inequalities
- 2 Explain how these inequalities were obtained
- 3 Symmetric inequalities
- 4 Greater violations
- 5 Different kinds of non-locality?

└ This Talk

- Report the discovery of such inequalities
- Explain how these inequalities were obtained
- Symmetric inequalities
- Greater violations
- Different kinds of non-locality?

1. The purpose of this talk is to present these new-found inequalities and explain how they were obtained
2. Talk about how symmetries play two roles in the search for quantum non-locality
3. Demonstrate how these inequalities are robust to noise
4. Finding maximally violating distributions

Example Inequality

- Quick example/preview:

$$\begin{aligned} &P(110)P(223) + P(110)P(233) + P(110)P(323) + P(110)P(333) \leq \\ &2P(020)P(213) + 2P(023)P(210) + 2P(023)P(310) + 2P(030)P(213) + \\ &2P(033)P(210) + 2P(033)P(310) + 2P(120)P(213) + 2P(123)P(210) + \\ &2P(123)P(310) + 2P(130)P(213) + 2P(132)P(311) + 2P(133)P(210) + \\ &\quad + \cdots \text{ 324 more terms } \cdots + \\ &P(320)P(323) + P(320)P(333) + P(323)P(330) + P(330)P(333) \end{aligned}$$

- $P(abc)$ shorthand for $P_{ABC}(abc)$
- Four outcomes for each A, B, C
- Polynomial in P_{ABC} , marginals $P_{AB}, P_{BC}, P_{AC}, P_A, P_B, P_C$

Triangle Inequalities

└ Example Inequality

■ Quick example/preview:

$$\begin{aligned}
&P(110)P(223) + P(110)P(233) + P(110)P(323) + P(110)P(333) \leq \\
&2P(020)P(213) + 2P(023)P(210) + 2P(023)P(310) + 2P(030)P(213) + \\
&2P(033)P(210) + 2P(033)P(310) + 2P(120)P(213) + 2P(123)P(210) + \\
&2P(123)P(310) + 2P(130)P(213) + 2P(132)P(311) + 2P(133)P(210) + \\
&\quad + \dots + 324 \text{ more terms} \dots + \\
&P(320)P(323) + P(320)P(333) + P(323)P(330) + P(330)P(333)
\end{aligned}$$

■ $P(abc)$ shorthand for $P_{ABC}(abc)$ ■ Four outcomes for each A, B, C ■ Polynomial in P_{ABC} , marginals $P_{AB}, P_{AC}, P_{BC}, P_A, P_B, P_C$

1. As a quick example or preview of what is to come, here is a an example inequality admits quantum violations in the triangle scenario
2. Some features of note: inequality is polynomial in P_{ABC} and its marginals

Question: Which marginal models $P^{\mathcal{M}}$ are **compatible** with a causal structure \mathcal{G} ?

- **Marginal model** $P^{\mathcal{M}}$ is collection of probability distributions

$$P^{\mathcal{M}} = \{P_{V_1}, \dots, P_{V_k}\}$$

- **Marginal scenario** $\mathcal{M} = \{V_1, \dots, V_k\}$

$$V \in \mathcal{M}, V' \subseteq V \implies V' \in \mathcal{M}$$

- **Joint random variables** $\mathcal{J} = \bigcup_i V_i = \{v_1, \dots, v_n\}$
- **Causal Structure** $\mathcal{G} = (\mathcal{N}, \mathcal{E})$ is a directed acyclic graph (DAG)
- Nodes classified into **latent nodes** \mathcal{N}_L and **observed nodes** \mathcal{N}_O

└ Notation

Question: Which marginal models P^M are compatible with a causal structure \mathcal{G} ?

- **Marginal model** P^M is collection of probability distributions

$$P^M = \{P_{V_1}, \dots, P_{V_k}\}$$

- **Marginal scenario** $\mathcal{M} = \{V_1, \dots, V_k\}$

$$V \in \mathcal{M}, V' \subseteq V \implies V' \in \mathcal{M}$$

- **Joint random variables** $\mathcal{J} = \bigcup_i V_i = \{v_1, \dots, v_n\}$

- **Causal Structure** $\mathcal{G} = (\mathcal{N}, \mathcal{E})$ is a directed acyclic graph (DAG)

- Nodes classified into **latent nodes** N_L and **observed nodes** N_O

1. Before continuing, I will define exactly what I mean by causal compatibility
2. Causal compatibility refers to the compatibility between causal structures and marginal models
3. Marginal model is collection of probability distributions over sets of random variables
4. Marginal scenario refers to the those sets of random variables
5. The complete set of random variables are the joint random variables

Let $n, m \in \mathcal{N}$ be nodes of the graph \mathcal{G} .

- **parents of n** : $\text{Pa}_{\mathcal{G}}(n) \equiv \{m \mid m \rightarrow n\}$
- **children of n** : $\text{Ch}_{\mathcal{G}}(n) \equiv \{m \mid n \rightarrow m\}$
- **ancestry of n** : $\text{An}_{\mathcal{G}}(n) \equiv \bigcup_{i \in \mathbb{W}} \text{Pa}_{\mathcal{G}}^i(n)$

$$\text{Pa}_{\mathcal{G}}^0(n) = n \quad \text{Pa}_{\mathcal{G}}^i(n) \equiv \text{Pa}_{\mathcal{G}}(\text{Pa}_{\mathcal{G}}^{i-1}(n))$$

Notation extends to sets of nodes $N \subseteq \mathcal{N}$,

- **parents of N** : $\text{Pa}_{\mathcal{G}}(N) \equiv \bigcup_{n \in N} \text{Pa}_{\mathcal{G}}(n)$
- **children of N** : $\text{Ch}_{\mathcal{G}}(N) \equiv \bigcup_{n \in N} \text{Ch}_{\mathcal{G}}(n)$
- **ancestry of N** : $\text{An}_{\mathcal{G}}(N) \equiv \bigcup_{n \in N} \text{An}_{\mathcal{G}}(n)$

An **induced subgraph** of $\mathcal{G} = (\mathcal{N}, \mathcal{E})$ due to $N \subseteq \mathcal{N}$

$$\text{Sub}_{\mathcal{G}}(N) = (N, \{e \in \mathcal{E} \mid e \subseteq N\})$$

Question: Which marginal models $P^{\mathcal{M}}$ are **compatible** with a causal structure \mathcal{G} ?

Answer: $P^{\mathcal{M}}$ is compatible with \mathcal{G} if there exists a set of **casual parameters**

$$\left\{ P_{n|\text{Pa}_{\mathcal{G}}(n)} \mid n \in \mathcal{N} \right\}$$

Such that for each $V \in \mathcal{M}$, P_V can be recovered:

1 $P_{\mathcal{N}} = \prod_{n \in \mathcal{N}} P_{n|\text{Pa}_{\mathcal{G}}(n)}$

2 $P_V = \sum_{\mathcal{N} \setminus V} P_{\mathcal{N}}$

Inequality: A **casual compatibility inequality** I is an inequality over $P^{\mathcal{M}}$ that is satisfied by all compatible $P^{\mathcal{M}}$

Two necessary components to compatibility:

- 1 **Marginal problem:** $\forall V \in \mathcal{M} : P_V = \sum_{\mathcal{N} \setminus V} P_{\mathcal{N}}$
 - Is the marginal model contextual or non-contextual?
 - 3 distinct ways to tackle this problem
 - 1 Convex hull, Polytope projection, Fourier-Motzkin
 - 2 Possibilistic Hardy Inequalities (Hypergraph transversals)
 - 3 Linear Program Feasibility/Infeasibility
- 2 **Markov Separation:** $P_{\mathcal{N}} = \prod_{n \in \mathcal{N}} P_{n|\text{Pa}_{\mathcal{G}}(n)}$
 - Much harder to determine since latent nodes \mathcal{N}_O have unspecified behaviour
 - It is possible to turn Markov Separation problem into a Marginal problem (at least partially)

Developed by Wolfe, Spekkens, and Fritz
Wolfe, Spekkens, and Fritz 2016

Definition

An **inflation** of a causal structure \mathcal{G} is another causal structure \mathcal{G}' such that:

$$\forall n' \in \mathcal{N}', n' \sim n \in \mathcal{N} : \text{AnSub}_{\mathcal{G}'}(n') \sim \text{AnSub}_{\mathcal{G}}(n)$$


Where $\text{AnSub}_{\mathcal{G}}(n)$ denotes the ancestral sub-graph of n in \mathcal{G}

$$\text{AnSub}_{\mathcal{G}}(n) = \text{Sub}_{\mathcal{G}}(\text{An}_{\mathcal{G}}(n))$$


And ' \sim ' is a **copy-index** equivalence relation

$$A_1 \sim A_2 \sim A \not\sim B_1 \sim B_2 \sim B$$


Some Inflations of the Triangle Scenario




The Triangle Scenario



Spiral Inflation



Cut Inflation



Large Inflation

Inflation Lemma

If one has obtained \mathcal{G} , inflation \mathcal{G}' and *compatible* marginal distribution P_N where $N \subseteq \mathcal{N}$, then:

- 1 There exists causal parameters $\{P_{n|\text{Pa}_{\mathcal{G}}(n)} \mid n \in \mathcal{N}\}$ such that

$$P_N = \prod_{n \in N} P_{n|\text{Pa}_{\mathcal{G}}(n)}$$

- 2 $\text{AnSub}_{\mathcal{G}'}(n') \sim \text{AnSub}_{\mathcal{G}}(n) \implies \text{Pa}_{\mathcal{G}'}(n') \sim \text{Pa}_{\mathcal{G}}(n)$

- 3 Construct **inflated causal parameters**

$$\forall n' \in \mathcal{N}' : P_{n'|\text{Pa}_{\mathcal{G}'}(n')} \equiv P_{n|\text{Pa}_{\mathcal{G}}(n)}$$

- 4 Obtain *compatible* marginal distributions over any $N' \subseteq \mathcal{N}'$

$$P_{N'} = \prod_{n' \in N'} P_{n'|\text{Pa}_{\mathcal{G}'}(n')}$$

- Inflation procedure holds for any $N \in \mathcal{N}, N' \in \mathcal{N}'$ where $N \sim N'$
- Define **injectable sets of \mathcal{G}'** and **images of the injectable of \mathcal{G}**

$$\text{Inj}_{\mathcal{G}}(\mathcal{G}') \equiv \{N' \subseteq \mathcal{N}' \mid \exists N \subseteq \mathcal{N} : N \sim N'\}$$

$$\text{ImInj}_{\mathcal{G}}(\mathcal{G}') \equiv \{N \subseteq \mathcal{N} \mid \exists N' \subseteq \mathcal{N}' : N \sim N'\}$$

- For $N' \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')$ there is a *unique* $N \subseteq \mathcal{N}$ such that $N \sim N'$
- For $N \in \text{ImInj}_{\mathcal{G}}(\mathcal{G})$ there can *exist many* $N' \subseteq \mathcal{N}'$ such that $N \sim N'$


Lemma

The Inflation Lemma: Wolfe, Spekkens, and Fritz 2016, lemma 3
Given a particular inflation \mathcal{G}' of \mathcal{G} , if a marginal model $\{P_N \mid N \in \text{ImInj}_{\mathcal{G}}(\mathcal{G}')\}$ is compatible with \mathcal{G} then all marginal models $\{P_{N'} \mid N' \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')\}$ are compatible with \mathcal{G}' provided that $P_N = P_{N'}$ for all instances where $N \sim N'$.

Corollary

Any causal compatibility inequality I' constraining the injectable sets $\text{Inj}_{\mathcal{G}}(\mathcal{G}')$ can be *deflated* into a causal compatibility inequality I constraining the images of the injectable sets $\text{ImInj}_{\mathcal{G}}(\mathcal{G}')$.

- Deflation only holds when inequality constrains probabilities
 $P_{N'}, N' \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')$
- **Linear inequality** for \mathcal{G}'



- **Polynomial inequality** for $\mathcal{G}!$


- d -separation relations + inflation = polynomial inequalities over \mathcal{G}
- Restrict focus to sets N' that are partitioned into N'_1, N'_2 d -separated by empty set \emptyset
- A **pre-injectable set** N' :

$$N' = \coprod_i N'_i \quad \forall i : N'_i \in \text{Inj}_{\mathcal{G}}(\mathcal{G}')$$

$$\forall i, j : N'_i \perp N'_j \iff \text{An}_{\mathcal{G}'}(N'_i) \cap \text{An}_{\mathcal{G}'}(N'_j) = \emptyset$$

- Only need to consider **maximal pre-injectable sets** $\text{PreInj}_{\mathcal{G}}(\mathcal{G}')$

Pre-injectable Sets of Large Inflation



- Has 12 maximal pre-injectable sets (to follow)

Large Inflation Pre-injectable Sets

Maximal Pre-injectable Sets

$\{A_1, B_1, C_1, A_4, B_4, C_4\}$

$\{A_1, B_2, C_3, A_4, B_3, C_2\}$

$\{A_2, B_3, C_1, A_3, B_2, C_4\}$

$\{A_2, B_4, C_3, A_3, B_1, C_2\}$

$\{A_1, B_3, C_4\}$

$\{A_1, B_4, C_2\}$

$\{A_2, B_1, C_4\}$

$\{A_2, B_2, C_2\}$

$\{A_3, B_3, C_3\}$

$\{A_3, B_4, C_1\}$

$\{A_4, B_1, C_3\}$

$\{A_4, B_2, C_1\}$

Ancestral Independences

$\{A_1, B_1, C_1\} \perp \{A_4, B_4, C_4\}$

$\{A_1, B_2, C_3\} \perp \{A_4, B_3, C_2\}$

$\{A_2, B_3, C_1\} \perp \{A_3, B_2, C_4\}$

$\{A_2, B_4, C_3\} \perp \{A_3, B_1, C_2\}$

$\{A_1\} \perp \{B_3\} \perp \{C_4\}$

$\{A_1\} \perp \{B_4\} \perp \{C_2\}$

$\{A_2\} \perp \{B_1\} \perp \{C_4\}$

$\{A_2\} \perp \{B_2\} \perp \{C_2\}$

$\{A_3\} \perp \{B_3\} \perp \{C_3\}$

$\{A_3\} \perp \{B_4\} \perp \{C_1\}$

$\{A_4\} \perp \{B_1\} \perp \{C_3\}$

$\{A_4\} \perp \{B_2\} \perp \{C_1\}$

Deriving Inequalities

- Inflation facilitates turning linear, inflated inequalities into polynomial deflated ones
- **Question:** How to derive compatibility inequalities for \mathcal{G}' ?
- **Answer:** Use your favorite technique for deriving compatibility inequalities:
 - Entropic inequalities
 - Finite outcome inequalities
- Here we solve the marginal problem for $\mathcal{M} = \text{PreInj}_{\mathcal{G}}(\mathcal{G}')$
- First some notation and formalism

Outcomes and Events

Definition

Each variable v has finite set of **outcomes** O_v .

Each set of variables V has finite set of **events** $\mathcal{E}(V)$:

$$\mathcal{E}(V) \equiv \{s : V \rightarrow O_V \mid \forall v \in V, s(v) \in O_v\}$$

Definition

The set of events over the joint variables $\mathcal{E}(\mathcal{J})$ are termed the **joint events**.

Definition

The set of events over the marginal contexts are the **marginal events**

$$\mathcal{E}(\mathcal{M}) \equiv \coprod_{V \in \mathcal{M}} \mathcal{E}(V)$$

Definition

The **joint distribution vector** $\mathcal{P}^{\mathcal{J}}$

$$\mathcal{P}_j^{\mathcal{J}} = P_{\mathcal{J}}(j) \quad \forall j \in \mathcal{E}(\mathcal{J})$$

Definition

The **marginal distribution vector** $\mathcal{P}^{\mathcal{M}}$

$$\mathcal{P}_m^{\mathcal{M}} = P_{\mathcal{D}(m)}(m) \quad \forall m \in \mathcal{E}(\mathcal{M}), \mathcal{D}(m) \in \mathcal{M}$$

Can now write complete marginal problem as matrix multiplication:

$$\forall V \in \mathcal{M} : P_V = \sum_{\mathcal{J} \setminus V} P_{\mathcal{J}} \iff \mathcal{P}^{\mathcal{M}} = M \cdot \mathcal{P}^{\mathcal{J}}$$

Incidence Matrix

- **Incidence matrix** M is a bit-wise matrix
- Row-indexed by marginal events $m \in \mathcal{E}(\mathcal{M})$
- Column-indexed by joint events $j \in \mathcal{E}(\mathcal{J})$

$$M_{m,j} = \begin{cases} 1 & m = j|_{\mathcal{D}(m)} \\ 0 & \text{otherwise} \end{cases}$$

$$\# \text{Columns} = |\mathcal{E}(\mathcal{J})| = \prod_{v \in \mathcal{J}} |O_v|$$

$$\# \text{Rows} = |\mathcal{E}(\mathcal{M})| = \sum_{V \in \mathcal{M}} \prod_{v \in V} |O_v|$$

Example

Let \mathcal{J} be 3 binary variables $\mathcal{J} = \{A, B, C\}$ and \mathcal{M} be the marginal scenario $\mathcal{M} = \{\{A, B\}, \{B, C\}, \{A, C\}\}$. The incidence matrix becomes:

$$M = \begin{array}{l} (A,B,C) = \\ (A=0,B=0) \\ (A=0,B=1) \\ (A=1,B=0) \\ (A=1,B=1) \\ (B=0,C=0) \\ (B=0,C=1) \\ (B=1,C=0) \\ (B=1,C=1) \\ (A=0,C=0) \\ (A=0,C=1) \\ (A=1,C=0) \\ (A=1,C=1) \end{array} \begin{pmatrix} (0,0,0) & (0,0,1) & (0,1,0) & (0,1,1) & (1,0,0) & (1,0,1) & (1,1,0) & (1,1,1) \\ \mathbf{1} & \mathbf{1} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \mathbf{1} & \mathbf{1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \mathbf{1} & \mathbf{1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \mathbf{1} & \mathbf{1} \\ \mathbf{1} & 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 \\ 0 & \mathbf{1} & 0 & 0 & 0 & \mathbf{1} & 0 & 0 \\ 0 & 0 & \mathbf{1} & 0 & 0 & 0 & \mathbf{1} & 0 \\ 0 & 0 & 0 & \mathbf{1} & 0 & 0 & 0 & \mathbf{1} \\ \mathbf{1} & 0 & \mathbf{1} & 0 & 0 & 0 & 0 & 0 \\ 0 & \mathbf{1} & 0 & \mathbf{1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \mathbf{1} & 0 & \mathbf{1} & 0 \\ 0 & 0 & 0 & 0 & 0 & \mathbf{1} & 0 & \mathbf{1} \end{pmatrix}$$

Marginal Linear Program

- Obtain inequalities from incidence matrix M and known incompatible distribution $\mathcal{P}^{\mathcal{M}}$

Marginal LP:

minimize: $\emptyset \cdot x$

subject to: $x \succeq 0$

$$M \cdot x = \mathcal{P}^{\mathcal{M}}$$

Dual Marginal LP:

minimize: $y \cdot \mathcal{P}^{\mathcal{M}}$

subject to: $y \cdot M \succeq 0$

- y is an **infeasibility certificate**:

$$y \cdot M \cdot x = y \cdot \mathcal{P}^{\mathcal{M}} \geq 0$$

- **Infeasibility inequality**: $y \cdot \mathcal{P}^{\mathcal{M}} \geq 0$
- Most linear programming toolkits return certificates (*Mosek*, *Gurobi*, *CPLEX*, *cvxr/cvxopt*.)

└ Marginal Linear Program

■ Obtain inequalities from incidence matrix M and known incompatible distribution \mathcal{P}^M

Marginal LP:

minimize $0 \cdot x$

subject to: $x \geq 0$

$$M \cdot x = \mathcal{P}^M$$

Dual Marginal LP:

minimize $y \cdot \mathcal{P}^M$

subject to: $y \cdot M \geq 0$

■ y is an infeasibility certificate:

$$y \cdot M \cdot x = y \cdot \mathcal{P}^M \geq 0$$

■ Infeasibility inequality: $y \cdot \mathcal{P}^M \geq 0$


■ Most linear programming toolkits return certificates (Mosek, Gurobi, CPLEX, cvx/conopt.)

1. It is possible to derive non-contextuality inequalities from M and a known incompatible distribution \mathcal{P}^M
2. Note the null objective; the minimization objective is trivially always zero
3. The primal value of the linear program is of no interest, all that matters is its *feasibility*.
4. If *feasible*, then there exists a vector x that is a valid joint distribution vector \mathcal{P}^J .
5. Feasibility implies that $\mathcal{P}^J = x$.
6. Infeasibility implies contextuality


Known Incompatible Distribution in The Triangle Scenario

Fritz 2012 provides quantum-accessible distribution incompatible with TS

$$P_{ABC}(abc) = \text{Tr}[\Pi^\top \rho_{AB} \otimes \rho_{BC} \otimes \rho_{CA} \Pi M_{A,a} \otimes M_{B,b} \otimes M_{C,c}]$$



Fritz Distribution



$$\text{Yellow} = \frac{1}{32} (2 - \sqrt{2}) \quad \text{Black} = \frac{1}{32} (2 + \sqrt{2})$$

$$P_F(000) = P_F(110) = P_F(301) = P_F(211) = P_F(122) = P_F(032) = P_F(233) = P_F(323) = \frac{1}{32}(2 + \sqrt{2})$$

$$P_F(010) = P_F(100) = P_F(311) = P_F(201) = P_F(132) = P_F(022) = P_F(223) = P_F(333) = \frac{1}{32}(2 - \sqrt{2})$$

- C 's outcome acts as measurement “setting” for A , B ;
independent of ρ_{AB}
- Correlation coarse graining $\{0, 1, 2, 3\} \rightarrow \{(0, 3), (1, 2)\}$

$$\langle AB|C = 0, 1, 2 \rangle = \frac{1}{\sqrt{2}} \quad \langle AB|C = 3 \rangle = -\frac{1}{\sqrt{2}}$$

- Gives CHSH violation

$$\langle AB|C = 0 \rangle + \langle AB|C = 1 \rangle + \langle AB|C = 2 \rangle - \langle AB|C = 3 \rangle = 2\sqrt{2} \not\leq 2$$

Quantum Implementation of Fritz Distribution [Optional]

■ States:

$$\rho_{AB} = |\Psi^+\rangle\langle\Psi^+| \quad \rho_{BC} = \rho_{CA} = |\Phi^+\rangle\langle\Phi^+|$$

■ Maximally entangled Bell states:

$$|\Psi^+\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle) \quad |\Phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$$

■ Measurements:

$$\begin{aligned} M_A &= \left\{ |0\psi_0\rangle\langle 0\psi_0|, |0\psi_\pi\rangle\langle 0\psi_\pi|, |1\psi_{-\pi/2}\rangle\langle 1\psi_{-\pi/2}|, |1\psi_{\pi/2}\rangle\langle 1\psi_{\pi/2}| \right\} \\ M_B &= \left\{ |\psi_{\pi/4}0\rangle\langle\psi_{\pi/4}0|, |\psi_{5\pi/4}0\rangle\langle\psi_{5\pi/4}0|, |\psi_{3\pi/4}1\rangle\langle\psi_{3\pi/4}1|, |\psi_{-\pi/4}1\rangle\langle\psi_{-\pi/4}1| \right\} \\ M_C &= \{ |00\rangle\langle 00|, |01\rangle\langle 01|, |10\rangle\langle 10|, |11\rangle\langle 11| \} \end{aligned}$$

■ Shorthand: $|\psi_x\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{ix}|1\rangle)$

- Proof contingent on perfect correlation between C and pseudo-settings
- Proof not robust to noise

Problem (2.17 in Fritz 2012)

Find an example of non-classical quantum correlations in TS together with a proof of its non-classicality which does not hinge on Bell's Theorem.

- “...would be helpful to have inequalities...”
- Possible to find inequalities violated by P_F using the Large inflation of the TS

- Joint variables are all of the observable nodes $\mathcal{N}'_O = \mathcal{J}$

$$\mathcal{J} = \{A_1, A_2, A_3, A_4, B_1, B_2, B_3, B_4, C_1, C_2, C_3, C_4\}$$

- Marginal scenario is composed of pre-injectable sets
 $\mathcal{M} = \text{PreInj}_{\mathcal{G}}(\mathcal{G}')$
- Inequalities violated by Fritz distribution are inherently 4-outcome
- Incidence matrix M is **very large** $\sim 2.25\text{Gb}$
 - #Columns = $4^{12} = 16,777,216$
 - #Rows = $4 \times 4^6 + 8 \times 4^3 = 16,896$
 - #Non-zero Entries = 201,326,592

Triangle Scenario Inequality

- Desirable to find compatibility inequality I such that

$$\forall \varphi \in \text{Perm}(A, B, C) : \varphi[I] = I$$

- Compatibility is independent of variable labels
 $I, \mathcal{G} \rightarrow \varphi[I], \varphi[\mathcal{G}]$
- Need $\varphi[\mathcal{G}] = \mathcal{G}$ to find new $\varphi[I]$

Definition

The **causal symmetry group** of causal structure \mathcal{G} :

$$\text{Aut}(\mathcal{G}) = \{\varphi \in \text{Perm}(\mathcal{N}) \mid \varphi[\mathcal{G}] = \mathcal{G}\}$$

Strictly speaking, one needs to preserve observable nodes:

$$\text{Aut}_{\mathcal{N}_O}(\mathcal{G}) = \{\varphi \in \text{Aut}(\mathcal{G}) \mid \varphi[\mathcal{N}_O] = \mathcal{N}_O\}$$

└ Causal Symmetry

- Desirable to find compatibility inequality I such that

$$\forall \varphi \in \text{Perm}(A, B, C) : \varphi[I] = I$$

- Compatibility is independent of variable labels
 $I, G \rightarrow \varphi[I], \varphi[G]$
- Need $\varphi[G] = G$ to find new $\varphi[I]$

Definition

The *causal symmetry group* of causal structure G :

$$\text{Aut}(G) = \{\varphi \in \text{Perm}(N) \mid \varphi[G] = G\}$$

Strictly speaking, one needs to preserve observable nodes:

$$\text{Aut}_{N_O}(G) = \{\varphi \in \text{Aut}(G) \mid \varphi[N_O] = N_O\}$$

1. Fritz distribution is incompatible with Triangle scenario because party C plays the role of measurement settings for both A and B
2. In order to find quantum distributions different from P_F in the Triangle Scenario, it is therefore desirable to find a proof of its incompatibility (i.e. inequality) that is symmetric under exchange of parties
3. Surprisingly, it is possible to do so!
4. Here is how.
5. First, we will formally define the symmetry group in question
6. Causal symmetry group is the group of automorphisms on causal structure

- Causal symmetry group for \mathcal{G}' is no good!
- Not possible to deflate inequality if it's not in terms of injectable sets

Definition

The **restricted causal symmetry group** Φ of \mathcal{G}' :

$$\Phi = \text{Aut}_{\text{PreInj}_{\mathcal{G}}}(\mathcal{G}')$$

Restricted Causal Symmetry of Large Inflation

- Φ for the large inflation is an order 48 group with 4 generators

φ_1	φ_2	φ_3	φ_4
$A_1 \rightarrow A_4$	$A_1 \rightarrow A_1$	$A_1 \rightarrow C_1$	$A_1 \rightarrow A_1$
$A_2 \rightarrow A_3$	$A_2 \rightarrow A_3$	$A_2 \rightarrow C_2$	$A_2 \rightarrow A_2$
$A_3 \rightarrow A_2$	$A_3 \rightarrow A_2$	$A_3 \rightarrow C_3$	$A_3 \rightarrow A_3$
$A_4 \rightarrow A_1$	$A_4 \rightarrow A_4$	$A_4 \rightarrow C_4$	$A_4 \rightarrow A_4$
$B_1 \rightarrow B_4$	$B_1 \rightarrow C_1$	$B_1 \rightarrow A_1$	$B_1 \rightarrow B_2$
$B_2 \rightarrow B_3$	$B_2 \rightarrow C_3$	$B_2 \rightarrow A_2$	$B_2 \rightarrow B_1$
$B_3 \rightarrow B_2$	$B_3 \rightarrow C_2$	$B_3 \rightarrow A_3$	$B_3 \rightarrow B_4$
$B_4 \rightarrow B_1$	$B_4 \rightarrow C_4$	$B_4 \rightarrow A_4$	$B_4 \rightarrow B_3$
$C_1 \rightarrow C_4$	$C_1 \rightarrow B_1$	$C_1 \rightarrow B_1$	$C_1 \rightarrow C_3$
$C_2 \rightarrow C_3$	$C_2 \rightarrow B_3$	$C_2 \rightarrow B_2$	$C_2 \rightarrow C_4$
$C_3 \rightarrow C_2$	$C_3 \rightarrow B_2$	$C_3 \rightarrow B_3$	$C_3 \rightarrow C_1$
$C_4 \rightarrow C_1$	$C_4 \rightarrow B_4$	$C_4 \rightarrow B_4$	$C_4 \rightarrow C_2$

Symmetric Incidence

- Group orbits through repeated action of $\varphi \in \Phi$ on $m \in \mathcal{E}(\mathcal{M})$ and $j \in \mathcal{E}(\mathcal{J})$

$$\Phi[m] \equiv \{\varphi[m] \mid \varphi \in \Phi\}$$

$$\Phi[j] \equiv \{\varphi[j] \mid \varphi \in \Phi\}$$

- Construct **symmetric incidence matrix** $\Phi[M]$

$$\Phi[M]_{\Phi[m], \Phi[j]} = \sum_{m' \in \Phi[m]} \sum_{j' \in \Phi[j]} M_{m', j'}$$

$$\Phi[M] = \Lambda_{\Phi[\mathcal{E}(\mathcal{M})]} \cdot M \cdot \Lambda_{\Phi[\mathcal{E}(\mathcal{J})]}$$

- $\Phi[M]$ not a bit-wise matrix like M
- For large inflation M is $16,896 \times 16,777,216$
- For large inflation $\Phi[M]$ is $450 \times 358,120$

Party Symmetric Inequality

$$2[P(001)P(333)]_3 + 2[P(010)P(323)]_3 + 6[P(000)P(323)]_3 + 6[P(000)P(333)]_1$$


$$\leq$$

$$\begin{aligned} & 12[P(031)P(302)]_6 + 12[P(033)P(303)]_6 + 12[P(103)P(130)]_6 + 12[P(203)P(230)]_6 + 12[P(203)P(330)]_6 + 2[P(001)P(320)]_6 + 2[P(002)P(221)]_3 + 2[P(003)P(211)]_6 + \\ & 2[P(003)P(331)]_3 + 2[P(011)P(211)]_3 + 2[P(012)P(322)]_6 + 2[P(013)P(313)]_6 + 2[P(013)P(332)]_6 + 2[P(020)P(111)]_3 + 2[P(020)P(211)]_6 + 2[P(021)P(212)]_6 + \\ & 2[P(022)P(211)]_3 + 2[P(022)P(212)]_6 + 2[P(022)P(322)]_3 + 2[P(023)P(232)]_6 + 2[P(030)P(212)]_3 + 2[P(031)P(231)]_6 + 2[P(032)P(331)]_6 + 2[P(033)P(333)]_3 + \\ & 2[P(101)P(131)]_3 + 2[P(101)P(132)]_6 + 2[P(102)P(131)]_6 + 2[P(102)P(132)]_6 + 2[P(102)P(133)]_6 + 2[P(110)P(133)]_6 + 2[P(110)P(212)]_6 + 2[P(110)P(222)]_3 + \\ & 2[P(110)P(223)]_3 + 2[P(112)P(331)]_3 + 2[P(120)P(122)]_6 + 2[P(121)P(201)]_6 + 2[P(122)P(200)]_3 + 2[P(122)P(202)]_6 + 2[P(122)P(210)]_6 + 2[P(122)P(300)]_3 + \\ & 2[P(130)P(232)]_6 + 2[P(130)P(233)]_6 + 2[P(131)P(201)]_6 + 2[P(131)P(202)]_3 + 2[P(131)P(313)]_3 + 2[P(133)P(200)]_3 + 2[P(133)P(201)]_6 + 2[P(133)P(211)]_3 + \\ & 2[P(133)P(212)]_6 + 2[P(133)P(300)]_3 + 2[P(202)P(231)]_6 + 2[P(210)P(222)]_6 + 2[P(220)P(222)]_3 + 2[P(220)P(313)]_6 + 2[P(221)P(313)]_6 + 2[P(222)P(331)]_3 + \\ & 2[P(223)P(331)]_3 + 2[P(230)P(312)]_6 + 2[P(231)P(313)]_6 + 2[P(232)P(320)]_6 + 2[P(302)P(322)]_6 + 2[P(320)P(323)]_6 + 2[P(330)P(332)]_3 + 3[P(000)P(003)]_3 + \\ & 3[P(010)P(301)]_6 + 4[P(001)P(131)]_6 + 4[P(002)P(020)]_6 + 4[P(002)P(133)]_6 + 4[P(002)P(323)]_6 + 4[P(010)P(123)]_6 + 4[P(013)P(212)]_6 + 4[P(013)P(312)]_6 + \\ & 4[P(023)P(221)]_6 + 4[P(023)P(222)]_6 + 4[P(023)P(322)]_6 + 4[P(031)P(211)]_6 + 4[P(032)P(321)]_6 + 4[P(100)P(123)]_6 + 4[P(100)P(232)]_6 + 4[P(100)P(313)]_6 + \\ & 4[P(112)P(310)]_6 + 4[P(122)P(203)]_6 + 4[P(122)P(302)]_6 + 4[P(130)P(222)]_6 + 4[P(130)P(223)]_6 + 4[P(222)P(310)]_6 + 4[P(223)P(320)]_6 + 4[P(231)P(301)]_6 + \\ & 4[P(312)P(330)]_6 + 6[P(001)P(031)]_6 + 6[P(001)P(033)]_6 + 6[P(002)P(300)]_6 + 6[P(002)P(330)]_3 + 6[P(003)P(032)]_6 + 6[P(003)P(131)]_6 + 6[P(003)P(132)]_6 + \\ & 6[P(011)P(300)]_3 + 6[P(011)P(320)]_6 + 6[P(012)P(200)]_6 + 6[P(012)P(301)]_6 + 6[P(013)P(030)]_6 + 6[P(013)P(110)]_6 + 6[P(013)P(120)]_6 + 6[P(013)P(303)]_6 + \\ & 6[P(020)P(102)]_6 + 6[P(020)P(103)]_6 + 6[P(020)P(123)]_6 + 6[P(020)P(202)]_3 + 6[P(020)P(203)]_6 + 6[P(020)P(311)]_6 + 6[P(020)P(322)]_6 + 6[P(020)P(330)]_6 + \\ & 6[P(022)P(303)]_6 + 6[P(030)P(033)]_6 + 6[P(030)P(101)]_3 + 6[P(030)P(133)]_6 + 6[P(030)P(202)]_3 + 6[P(030)P(303)]_3 + 6[P(030)P(332)]_6 + 6[P(031)P(203)]_6 + \\ & 6[P(032)P(310)]_6 + 6[P(033)P(101)]_6 + 6[P(033)P(130)]_6 + 6[P(033)P(200)]_3 + 6[P(033)P(212)]_6 + 6[P(033)P(220)]_6 + 6[P(033)P(222)]_3 + 6[P(033)P(230)]_6 + \\ & 6[P(033)P(322)]_3 + 6[P(100)P(203)]_6 + 6[P(101)P(130)]_6 + 6[P(103)P(310)]_6 + 6[P(113)P(130)]_6 + 6[P(113)P(230)]_6 + 6[P(113)P(330)]_3 + 6[P(122)P(330)]_6 + \\ & 6[P(130)P(313)]_6 + 6[P(132)P(303)]_6 + 6[P(133)P(303)]_6 + 6[P(133)P(320)]_6 + 6[P(200)P(203)]_6 + 6[P(201)P(230)]_6 + 6[P(203)P(231)]_6 + 6[P(223)P(300)]_6 + \\ & 8[P(003)P(320)]_6 + 8[P(032)P(300)]_6 \end{aligned}$$

Parameterizing Quantum Distributions

For our purposes, we need to parameterize the space of quantum-accessible distributions that are *realized* on the Triangle Scenario

$$P_{ABC}(abc) = \text{Tr}[\Pi^\top \rho_{AB} \otimes \rho_{BC} \otimes \rho_{CA} \Pi M_{A,a} \otimes M_{B,b} \otimes M_{C,c}]$$



- Attempt to find new non-classical distributions
- Objective function $f(\lambda) \in \mathbb{R}$
 - 1 Real-valued parameters $\lambda = (\lambda_0, \dots, \lambda_n)$
 - 2 Quantum states/measurements $\rho_{AB}, \rho_{BC}, \rho_{CA}, M_A, M_B, M_C$
 - 3 Distribution P_{ABC}
 - 4 Plug into inequality I in homogeneous form $I(P_{ABC}) \geq 0$
 - 5 Output is objective value $I(P_{ABC})$
- Numerical minimization of $f(\lambda)$

$$f(\lambda_{(k+1)}) = \lambda_{(k)} - \gamma_{(k)} \nabla f(\lambda_{(k)})$$

- Non-convex, non-linear, smooth/continuous
- Gradient Descent, BFGS Method, Nelder-Mead simplex method
- Stochastic methods: simulated annealing, basin-hopping

Parameterizing Unitary Group

- Spengler, Huber and Heismayr Spengler, Huber, and Hiesmayr 2010 demonstrate a parameterization of $\mathcal{U}(d)$ where the parameters are organized in a $d \times d$ -matrix of real values $\lambda_{n,m}$

$$U = \left[\prod_{m=1}^{d-1} \left(\prod_{n=m+1}^d R_{m,n} R P_{n,m} \right) \right] \cdot \left[\prod_{l=1}^d G P_l \right]$$

- Global Phase Terms: $G P_l = \exp(i P_l \lambda_{l,l})$
- Relative Phase Terms: $R P_{n,m} = \exp(i P_n \lambda_{n,m})$
- Rotation Terms: $R_{m,n} = \exp(i \sigma_{m,n} \lambda_{m,n})$
- Projection Operators: $P_l = |l\rangle\langle l|$
- Anti-symmetric σ -matrices: $\sigma_{m,n} = -i|m\rangle\langle n| + i|n\rangle\langle m|$
- Parameters $\lambda_{n,m} \in [0, 2\pi]$

Parameterizing Unitary Group Cont'd

- Each parameter $\lambda_{n,m}$ has physical interpretation
- Degeneracies are easily eliminated such as global phase

$$\forall l = 1, \dots, d : \lambda_{l,l} = 0 \implies GP_l = \mathbb{1}$$

- Parameterize $U \in \mathcal{U}(d)$ up to global phase denoted $\tilde{U} \in \mathcal{U}(d)$
- Computationally efficient

$$GP_l = \mathbb{1} + P_l(e^{i\lambda_{l,l}} - 1)$$

$$RP_{n,m} = \mathbb{1} + P_n(e^{i\lambda_{n,m}} - 1)$$

$$\begin{aligned} R_{m,n} = \mathbb{1} &+ (|m\rangle\langle m| + |n\rangle\langle n|)(\cos \lambda_{n,m} - 1) \\ &+ (|m\rangle\langle n| - |n\rangle\langle m|) \sin \lambda_{n,m} \end{aligned}$$

Parameterizing States

- Each latent resource $\rho \in (\rho_{AB}, \rho_{BC}, \rho_{CA})$ modeled as bipartite qubit state acting on $\mathcal{H}^{d/2} \otimes \mathcal{H}^{d/2}$
- $d \times d$ positive semi-definite (PSD) hermitian matrices with unitary trace
- **Cholesky Parametrization** allows one to write any hermitian PSD as $\rho = T^\dagger T$
- For $d = 4$:

$$T = \begin{bmatrix} \lambda_1 & 0 & 0 & 0 \\ \lambda_2 + i\lambda_3 & \lambda_4 & 0 & 0 \\ \lambda_5 + i\lambda_6 & \lambda_7 + i\lambda_8 & \lambda_9 & 0 \\ \lambda_{10} + i\lambda_{11} & \lambda_{12} + i\lambda_{13} & \lambda_{14} + i\lambda_{15} & \lambda_{16} \end{bmatrix}$$

- d^2 real-valued parameters
- Normalized $\rho = T^\dagger T / \text{Tr}(T^\dagger T)$ adds degeneracy

Parameterizing States Cont'd

- **SHH parameterization** Spengler, Huber, and Hiesmayr 2010
exploits spectral decomposition; for rank $k \leq d$ density matrix

$$\rho = \sum_{i=1}^k p_i |\psi_i\rangle \langle \psi_i| \quad p_i \geq 0, \sum_i p_i = 1$$

- Orthonormal k -element sub-basis $\{|\psi_i\rangle\}$ of \mathcal{H}^d can be transformed into computational basis $\{|i\rangle\}$ by unitary $U \in \mathcal{U}(d)$ such that $|\psi_i\rangle = U|i\rangle$
- Freedom to choice k
- Parameterize ρ through $\{p_i\}$ and \tilde{U}_k

$$\tilde{U}_k = \prod_{m=1}^k \left(\prod_{n=m+1}^d R_{m,n} R P_{n,m} \right)$$

- real-value parameters $d^2 - (d - k)^2 - k$ for \tilde{U}_k , $k - 1$ for $\{p_i\}$
(no degeneracy)

Parameterizing POVMs

- Each party (A, B, C) is assigned a **projective-operator valued measure (POVM)** (M_A, M_B, M_C)

$$\forall |\psi\rangle \in \mathcal{H}^d : \langle \psi | M_\chi | \psi \rangle \geq 0 \quad M_\chi = M_\chi^\dagger$$

- n -outcome measurement

$$M_\chi = \{M_{\chi,1}, \dots, M_{\chi,n}\} \quad \sum_{i=1}^n M_{\chi,i} = \mathbb{1}$$

- For $n = 2$ outcomes, a parameterization exists by constraining the eigenvalues of $M_{\chi,i}$; for $n > 2$ not aware of anything
- Warrants consideration of **projective-valued measures (PVMs)** (for $n = d$ this is without loss of generality)

Parameterizing POVMs

- Each party (A, B, C) is assigned a **projective-operator valued measure (POVM)** (M_A, M_B, M_C)

$$\forall |\psi\rangle \in \mathcal{H}^d : \langle \psi | M_k | \psi \rangle \geq 0 \quad M_k = M_k^\dagger$$

- n-outcome measurement**

$$M_k = \{M_{k,1}, \dots, M_{k,n}\} \quad \sum_{i=1}^n M_{k,i} = 1$$

- For $n = 2$ outcomes, a parameterization exists by constraining the eigenvalues of $M_{k,i}$ for $n > 2$ not aware of anything
- Warrants consideration of **projective-valued measures (PVMs)** (for $n = d$ this is without loss of generality)

1. Naimark's Dilation Theorem

Parameterizing PVMs

- Each party (A, B, C) is assigned n -outcome (M_A, M_B, M_C) such that,

$$M_{\chi,i}M_{\chi,j} = \delta_{ij}M_{\chi,i} \quad M_{\chi,i} = |m_{\chi,i}\rangle\langle m_{\chi,i}|$$

- Inspired by Pál and Vértesi 2010, parameterizing PVMs means parameterizing a n -element sub-basis $\{|m_{\chi,i}\rangle\}$
- Use unitary transformation again

$$\{|m_{\chi,1}\rangle, \dots, |m_{\chi,n}\rangle\} = \{U|1\rangle, \dots, U|n\rangle\}$$


- Global phase and remaining basis irrelevant: \tilde{U}_n requires $n(2d - n - 1)$ real-valued parameters
- PVMs are computationally more efficient

$$P_{ABC}(abc) = \langle m_{A,a}m_{B,b}m_{C,c} | \Pi^\top \rho_{AB} \otimes \rho_{BC} \otimes \rho_{CA} \Pi | m_{A,a}m_{B,b}m_{C,c} \rangle$$

Network Permutation Matrix

- States and measurements in the Triangle Scenario are not aligned
- Without Π , P_{ABC} would be separable
- Required to align B 's measurement over $\text{Tr}_{A,C}(\rho_{AB} \otimes \rho_{BC})$
- Π is a $2^6 \times 2^6$ matrix
- Shifts one qubit to the left

$$\Pi \equiv \sum_{|q_i\rangle \in \{|0\rangle, |1\rangle\}} |q_2 q_3 q_4 q_5 q_6 q_1\rangle \langle q_1 q_2 q_3 q_4 q_5 q_6|$$



- Full rank ρ, M gives 81 free parameters

$$|\lambda| = 3 \cdot (12 + 3) + 3 \cdot 12 = 81$$

- Parameterization of quantum distributions still degenerate
- Noisy seed (Gaussian noise):


$$\lambda_{(0)} = \lambda_{(F)} + \delta\lambda \quad \delta\lambda_i \sim \mathcal{N}(\mu = 0, \sigma^2 = (2\pi 10^x)^2)$$

- Uniform seed:

$$\lambda_{(0),i} \sim \mathcal{U}([0, 2\pi])$$


$$\mathcal{U}_\varepsilon(P) = (1 + \varepsilon)P + \varepsilon\mathcal{U}$$

Uniform Distribution




$$\text{Yellow square} = \frac{1}{64}$$


Fritz Local Minima




Fritz Local Minima Zoomed




Max Entangled vs. Max Violating (???) [Optional]



Max Entangled vs. Max Violating (???) [Optional]



Max Entangled vs. Max Violating



Maximally Violating Distributions

- Able to out-perform violation provided by Fritz distribution
- Maximally-violating states are not maximally-entangled; similar to detection loop-hole example of Methot and Scarani 2006
- Violation very sensitive to the initial parameters $\lambda_{(0)}$
- Both symmetric and asymmetric inequalities exhibit same qualitative features

- New causal compatibility inequalities have been found for the TS
- Inflation technique capable of producing inequalities with quantum/classical witnesses
- Proof of non-classicality is robust to noise
- Fritz witness-able by party symmetric inequalities
- Maximally violating distributions are different than Fritz but also similar
- Further research is necessary

Post-doc Opportunities At Perimeter

- Branciard, Cyril et al. (2012). “Bilocal versus nonbilocal correlations in entanglement-swapping experiments”. In: *Phys. Rev. A* 85.3.
- Chaves, Rafael, Lukas Luft, and David Gross (2014). “Causal structures from entropic information: geometry and novel scenarios”. In: *New Journal of Physics* 16.4, p. 043001.
- Fritz, Tobias (2012). “Beyond Bell’s Theorem: Correlation Scenarios”. In: eprint: [arXiv:1206.5115](https://arxiv.org/abs/1206.5115).
- Henson, Joe, Raymond Lal, and Matthew F Pusey (2014). “Theory-independent limits on correlations from generalized Bayesian networks”. In: *New Journal of Physics* 16.11, p. 113043. eprint: [arXiv:1405.2572](https://arxiv.org/abs/1405.2572).
- Methot, A. A. and V. Scarani (2006). “An anomaly of non-locality”. In: eprint: [arXiv:quant-ph/0601210](https://arxiv.org/abs/quant-ph/0601210).
- Pál, Károly F. and Tamás Vértesi (2010). “Maximal violation of a bipartite three-setting, two-outcome Bell inequality using infinite-dimensional quantum systems”. In: *Phys. Rev. A* 82.2.

- Spengler, Christoph, Marcus Huber, and Beatrix C. Hiesmayr (2010). “A composite parameterization of unitary groups, density matrices and subspaces”. In: [eprint: arXiv:1004.5252](#).
- Steudel, Bastian and Nihat Ay (2010). *Information-theoretic inference of common ancestors*. [eprint: arXiv:1010.5720](#).
- Wolfe, Elie, Robert W. Spekkens, and Tobias Fritz (2016). *The Inflation Technique for Causal Inference with Latent Variables*. [eprint: arXiv:1609.00672](#).