

Reviewed Preprint

v1 • September 5, 2025

Not revised

Reviewed Preprint

v2 • March 30, 2026

Revised by authors

✉ For correspondence:

daan@mbg.au.dk

* These authors contribute equally

Competing interests: No competing interests declared

Funding: See [page 39](#)

Reviewing editor: Aya Ito-Ishida, RIKEN Center for Brain Science, Japan

© 2025, Authier et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Pathogenic O-GlcNAc dyshomeostasis is associated with cortical malformations and hyperactivity

Florence Authier^{1,*}, Asad Jan^{2,*}, Islam Faress¹, Christian Stald Skoven³, Iria Esperon-Abril¹, Shagana Tharmakulasingam Balasubramaniam¹, Kévin-Sébastien Coquelin¹, Jens R Nyengaard², Carsten Scavenius¹, Benedetta Attianese¹, Oscar G Sevillano-Quispe¹, Simon Fristed Eskildsen³, Jesper Skovhus Thomsen⁴, Brian Hansen³, Daan MF van Aalten^{1,5} ✉

¹Section for Neurobiology and DANDRITE, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark • ²Core Center for Molecular Morphology, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark • ³CFIN, Aarhus Hospital, Aarhus, Denmark • ⁴Department of Biomedicine, Aarhus University, Aarhus, Denmark • ⁵Division of Cell and Developmental Biology, School of Life Sciences, University of Dundee, Dundee, United Kingdom

eLife Assessment

This is an **important** study that takes a key step towards understanding developmental disorders linked to mutations in the O-GlcNAc transferase enzyme by generating a mouse model harboring the C921Y mutation. While the mechanisms remain open, the study thoroughly examines behavioral and anatomical differences in these mice and provides **convincing** evidence for behavioral hyperactivity and learning/memory deficits, as well as phenotypic differences in skull and brain formation. This study will be of interest to those studying neurodevelopmental disorders and associated mechanisms.

<https://doi.org/10.7554/eLife.107170.2.sa3>

Abstract

Missense variants in the O-GlcNAc transferase (*OGT*) gene have recently been shown to segregate with a syndromic form of intellectual disability (OGT-ID), underscoring the importance of protein O-GlcNAcylation in brain function. However, the underlying pathophysiological mechanisms linking ID to potential OGT malfunction—whether developmental, neurophysiological, or both—remain unclear. Here, we present comprehensive analyses encompassing behaviour and brain architecture of a rodent model carrying the pathogenic C921Y OGT-ID variant. These mice show a range of behavioural deficits, including hyperactivity, impulsivity, and associative learning phenotypes. Structural studies, using micro-computed tomography and magnetic resonance imaging, revealed reduced skull size, microcephaly, reduced cortical thickness and hypoplastic corpus callosum. Detailed histological analyses revealed dysplastic changes in the neocortex, predominantly affecting the superficial layers of cingulate cortex. Mechanistically, quantitative proteomic analyses revealed O-GlcNAc dyshomeostasis associated with distinct perturbed molecular pathways involved in brain development. Taken together, these data reveal neurodevelopmental defects associated with O-GlcNAc dyshomeostasis and provide a platform for dissecting mechanism and treatments of OGT-ID.

Introduction

Intellectual disability (ID) is a prevalent neurodevelopmental condition estimated to affect 1-3% of the global population and is characterized by a significant impairment in cognitive function, along with compromised adaptive and social behaviour¹. Developmental anomalies in the human

cerebral cortex, particularly the neocortex, are common causes of intellectual disability (ID)^{2,3}. A recently reported syndromic form of ID is associated with missense variants in the *OGT* gene, which resides on chromosome Xq13.1, and encodes the O-linked N-acetylglucosamine transferase (OGT). This enzyme facilitates the covalent attachment of O-linked N-acetylglucosamine (O-GlcNAc) to the hydroxyl groups of serine and threonine (Ser/Thr) residues on nuclear, cytoplasmic and mitochondrial proteins^{4,5}. O-GlcNAcylation is a dynamic and highly conserved modification governed solely by two enzymes, OGT for attachment^{6,7} and O-GlcNAcase (OGA) for removal⁸. This sets O-GlcNAcylation apart from other post-translational modifications, such as phosphorylation and ubiquitination, that rely on hundreds of “writer” and “eraser” enzymes, associated with specific sets of “reader” proteins^{9,10}. O-GlcNAcylation regulates cellular functions, such as transcriptional activation¹¹, gene expression¹², stress response¹³ and proteostasis¹¹, in response to physiological changes. Structural studies of OGT have demonstrated that the enzyme is composed of an N-terminal tetratricopeptide repeat (TPR) domain, involved in substrate recognition and binding of OGT interactors^{9,10}, and a catalytic glycosyltransferase domain. In addition to catalysing O-GlcNAcylation, the catalytic domain is involved in the proteolytic activation of the transcriptional coregulator Host Cell Factor 1 (HCF-1), itself an ID-associated protein^{14,15}.

OGT and OGA are notably abundant in the brain, with particularly enrichment in hippocampal formation and related structures¹⁶. Thus, O-GlcNAcylation plays a critical role in neuronal survival, development, and synaptic function. For instance, pan-neuronal knockout of OGT results in severe neurodevelopmental defects and early neurodegeneration^{17,18}. Moreover, deletion of OGT in the dopaminergic neurons of substantia nigra or peripheral neurons in the dorsal root ganglion induces widespread apoptosis^{19,20}. At the synapse, OGT and OGA display distinct localization patterns: OGA is absent from the post-synaptic density, while OGT is distributed evenly between pre- and post-synaptic compartments in excitatory synapses^{21,22}. In this context, aynaptosome mass spectrometry (MS) data have revealed that nearly 20% of synaptic proteins, including ankyrin G, CaMKIV, and GluA2, are modified by O-GlcNAc, which dynamically regulates excitability and transmission^{23,24}.

To date, 17 pathogenic missense variants in the *OGT* gene have been identified as causal for a newly described syndromic form of ID, termed OGT-ID, also known as O-GlcNAc Transferase Congenital Disorder of Glycosylation (OGT-CDG)^{25–29}. This syndrome is frequently comorbid with neurological and psychiatric disorders such as epilepsy and autism spectrum disorder (ASD), both of which are associated with focal cortical dysplasia (FCD)^{30–32}. In addition to ID and maladaptive behaviour, OGT-ID patients exhibit a plethora of muscular, facial and neurological abnormalities, such as hypotonia, craniofacial dysmorphism, microcephaly, fifth finger clinodactyly, and developmental delay^{25,26,33}. Missense and exon-skipping variants in OGT have been found across both functional domains (i.e. TPR and catalytic), with affected patients exhibiting similar clinical features. These observations hint towards pathogenic mechanisms whereby the OGT variants affect neurodevelopmental processes and/or brain function leading to intellectual disability. Nevertheless, understanding the potential impact of OGT-ID variants on brain development has remained challenging, primarily due to the lack of viable vertebrate models since germline knockout (KO) of *Ogt* in mice leads to embryonic lethality³⁴.

Among OGT-ID variants, the catalytically impaired C921Y variant has been the most extensively studied. It was identified in three affected brothers born to a healthy, non-consanguineous couple in Denmark. The affected individuals exhibit developmental delay, autistic features, dysmorphic traits, osteoporosis and seizures³⁵. Functional studies have shown that the OGT^{C921Y} variant has decreased glycosyltransferase activity, both *in vitro* and in mouse embryonic stem cells (mESCs). In OGT^{C921Y} mESCs, the mutation reduces O-GlcNAcylation and decreases expression of stem cell markers Oct4, Sox2, and alkaline phosphatase (ALP)³², suggesting that OGT plays a critical role in embryonic stem cell self-renewal and pluripotency³². In *Drosophila melanogaster* models, the OGT^{C921Y} mutation has been shown to reduce O-GlcNAcylation during development³⁶, a defect that can be rescued by genetic or pharmacological inhibition of OGA. This mutation disrupts larval neuromuscular junction development and shortens sleep bout duration, with both effects being partially reversible through OGA inhibition. These findings suggest that certain aspects of OGT-ID

pathology are developmental in origin, while others may be reversible. Although these studies indicate that early differentiation and development are affected, a vertebrate model is required to dissect the developmental and brain-wide effects of OGT-ID variants, as the precise mechanisms remain unknown.

Here, we use an OGT^{C921Y} mouse model of OGT-ID to reveal a range of behavioural deficits, including hyperactivity, impulsivity, and associative learning phenotypes. Structural studies, using micro-computed tomography and magnetic resonance imaging, revealed reduced skull size, microcephaly, reduced cortical thickness and hypoplastic corpus callosum. Detailed histological analyses uncovered nodular cortical dysplasia, predominantly affecting the superficial layers of cingulate cortex. Mechanistically, quantitative proteomic analyses suggested perturbed regulation of distinct perturbed molecular pathways involved in brain development in the presence of O-GlcNAc dyshomeostasis. Taken together, these data reveal neurodevelopmental defects associated with O-GlcNAc dyshomeostasis and provide a platform for dissecting mechanism and treatments of OGT-ID.

Materials & Methods

Animal husbandry

The OGT^{C921Y} line was previously generated and reported³⁷ and was maintained under C57BL/6J background (Janvier, France). Animal cohorts were obtained from crossing of male OGT^{WT} with female OGT^{C921Y/+}. Only male mice were used in all experiments. Animals were housed in digitally ventilated cages (Tecniplast, Italy) with water and food available *ad libitum* and 12/12 h light/dark cycles in the Skou animal facility of Aarhus University. All animal studies and breeding were performed in accordance with the ARRIVE guidelines and the European Communities Council Directive (2010/EU) and were approved by the Danish Animal Experiments Inspectorate (Dyreforsøgstilsynet), under Breeding license 2022-15-0202-00135 and Project licenses: 2023-15-0201-01426 and 2020-15-0201-00421.

Non-invasive monitoring of spontaneous activity in home cages (DVC)

Patterns of locomotion and spontaneous activity were continuously monitored over 24 hours in home cages through a specialized digitally ventilated cages (DVC) platform (Tecniplast, Italy) between the age of 30 days and 105 days (termination). This platform is based on electrical capacitance sensing technology, with incorporates a sensor board equipped with an integrated circuit comprising 12 electrodes directly beneath the floor of the cages³⁸. The DVC circuit measures changes in the electrical capacitance signal from each electrode in response to the movement of a water-filled body (animal) close to or away from a given electrode. The measurements, performed approximately 4 times per second, are remotely relayed to the centralized DVC analytics platform (Tecniplast, Italy). In this web-based interface, time-stamped data for each cage can be visualized using in-built tools (e.g. daily rhythms, cumulative activity/locomotion index aggregated per minute/hour/day, bedding status, light or dark period activity, heatmaps etc.). In the default setup, the DVC analytics web-interface plots the animal locomotion index as arbitrary units normalized between 0% and 100%, representing the overall activity performed in the cage by the animals, i.e., the signal is measured for each cage and not each animal.

Behaviour assessment

Prior to behavioural testing, all animals were handled daily by the experimenter for one week. All handling took place in the experimental room with an ambient light setting of 25-30 Lux. All animals were housed in groups of 2 to 3 individuals. Animals were 10 weeks old at the start of behavioural testing, reaching 14 weeks old at the end. Before any behavioural test, mice were placed in the experiment room for at least 30 min prior to testing. The data analyses and video

quantification were performed blindly with respect to the genotype. The sample size was determined based on previous experience and validated using Post-hoc Power Calculation using ClinCalc online tool (<https://clincalc.com/stats/Power.aspx> )

Locomotor behaviour and sensorimotor coordination

Open Field: The activity of mice in an open field maze was recorded using ANY-maze video tracking software (Stoelting Europe, Ireland). Individual animals were placed in a 33.5 x 33.5 x 39 cm opaque box and were allowed to explore for 10 min over three consecutive days. Time spent in the centre and periphery and time moving in the periphery were analysed to investigate parameters of locomotion and anxiety.

Rotarod: Coordination skills were assessed using rotarod apparatus. Mice were placed on a Ugo Basile NB 80534 rotarod with an increase in speed from 4 to 40 rpm and an acceleration time of 2 min (40 rpm/min is reached after 2 min) for maximum 5 min. Latency to fall from the rotarod and speed were measured.

Static rods: Mice were placed at the extremity of the rod with head facing the void. Time spent by the mouse to perform a t-turn and to reach the goal platform were recorded. If the animal fell <5 s (presumably due to a misplacement), the mouse was tested again. After three consecutive fails or an up-side down of a mouse, a 120 s score was reported. Five rod diameters (35, 28, 22, 15 and 10 mm) were used for each animal.

Pole test: Mice were placed close to the top of a pole (40 cm) with the head of the mouse facing up. Times to perform a t-turn and reach the ground were recorded. The test was performed three times for each mouse. If the mouse succeeded the three attempts, a score of 3 was reported. For each slide and fail to perform the test within the 120 s cut off, an additional 1 point was incremented in the total score.

Cognitive performance

Novel Object Recognition: Novelty associated short- and long-term memory were assessed in an open-field maze. During the familiarisation phase mice are free to explore the maze containing two identical familiar objects for 10 min. After 90 min for short-term and 24 h for long-term memory, one of the familiar objects is replaced by a novel object and mice are free to explore the maze for 10 min. Time spent exploring objects and number of explorations was measured until reaching a total exploration time of 20 s for each mouse. *Spontaneous alternation:* The spontaneous alternation test was performed in a T-maze and was used to assess spatial working memory. Mice were free to choose left or right arms for 7 trials. Each arm entry was recorded to calculate the percentage of spontaneous alternation corresponding to the number of correct Left-Right (L/R) or Right-Left (R/L) sequences. Mice that were able to remember which arms they had entered most recently would choose a different one to explore.

Aversive conditioning and recall: The apparatus consisted of an open-top cage (24 × 20 × 30 cm) with metal floor bars, placed inside a soundproof cubicle (55 × 60 × 57 cm) (Ugo Basile, Italy). Three minutes after being placed in the conditioning chamber, the mice were conditioned using four tone-foot shock (CS-US) pairings (n = 9 per group). Each pairing consisted of a 25 s, 7 kHz tone (CS), followed by a 25 s gap (trace period), and then a 1.5 s foot shock (US) at 0.5 mA. After the conditioning session, the animals were isolated for 10–15 min before being returned to their home cage with their littermates. Long-term memory recall was assessed in a novel context 24 h after conditioning. Following a 3 min acclimatisation period in the novel context, the mice were exposed to four CS presentations without foot shocks. The intertrial interval ranged from 120 to 180 s for both the conditioning and testing sessions. The behavioural responses were recorded using a top-mounted camera and freezing and locomotor activity were automatically scored using ANY-maze software (Stoelting Europe, Ireland). The freezing percentage represents the time the mouse spent freezing during the CS presentation or the total duration of the time bin.

Anxiety behaviour

Elevated-plus maze: Anxiety-like behaviour was assessed using the elevated plus maze paradigm. Mice were placed in the centre of the cross-shaped maze comprised of two open arms (125 Lux) and two closed arms (25 Lux) and allowed to explore the maze for 5 min. Time, distance and number of entries in each of the open and closed arms were analysed as an approximation for anxiety.

Dark-light paradigm: Light-like anxiety behaviour was assessed using the dark-light paradigm at 180 Lux. Mice were placed in an arena containing both dark and light compartments. At the start of the test, mice were placed in the dark compartment and were free to explore both compartments for 10 min. Time, distance and number of entries in the light compartment were analysed to investigate the level of anxiety.

Compulsive behaviour

Marbles: Before the test, 20 glass marbles (1.6 cm in diameter) were placed on the top of the bedding (in five rows of four marbles) in a cage with 5 cm bedding. At the end of the test, the number of buried marbles (covered by at least 75% of bedding) was counted.

Digging and self-grooming: Mice were placed in a cage with 5 cm (digging) or 1 cm (self-grooming) bedding for 3 min. Number of digging/self-grooming/rearing events and time spent digging/self-grooming/rearing were recorded. *Nesting:* Mice were placed in a cage with 1 cm bedding in the presence of a cotton pad as nesting material for 1 h. Cotton pads were weighed before and 24 h after testing to allow drying. Percentage of material removed was quantified.

Litter burrowing: Mice were placed in a cage in the presence of a PVC tube filled with 120 g of litter bedding for 30 min. Percentage of litter material removed was quantified.

Brain perfusion for structural analyses

Brains were perfusion fixed and prepared for in-skull MRI (OGT^{WT} n = 15 and OGT^{C921Y} n = 16). Perfusion fixation was performed after the mice had been anesthetized by an intraperitoneal injection of Euthanimal (250 mg/kg, Alfasan, 088672). Then, the brain was fixed by transcardiac perfusion at 125 mmHg, to be close to physiologic brain perfusion as previously reported³⁹ using 25 mL heparinized (Heparin, 0.2 mL/100 mL, 5000 IU/mL, Pan Pharma, 482480) Natriumchlorid (9 mg/mL, B Braun, 5/389885/0417) for 3 min 30 s, followed by 25 mL of buffered 10% formalin solution (VWR Qpath Chemicals, 11699404) for 3 min 30 s. After decapitation, the mandible and extracranial tissue were removed from the skull to avoid susceptibility artifacts from air bubbles trapped in fur and cavities during imaging. Hereafter, the in-skull brains were stored in 10% formalin solution for at least one week prior imaging. One WT mouse was excluded due to misperfusion.

Magnetic resonance imaging (MRI) and image analysis

Before imaging, the fixed brain samples were washed in PBS for at least 24 h to increase MRI signal by removal of excess fixative⁴⁰. For imaging, the samples were subsequently mounted in a 15 mL centrifuge tube filled with a perfluorocarbon based liquid (Fluorinert, 3M, PN: FC-770) as is standard^{41–46}.

MRI data collection: MRI scans were acquired on a 9.4 T preclinical system (BioSpec 94/20, Bruker Biospin, Ettlingen, Germany) using a bore-mounted 25 mm quadrature transmit-receive coil. To avoid sample vibrations, the tube containing the sample was secured in a custom polyethylene foam cylinder inside the coil. In-house 3D-printed sample holders ensured consistent positioning of the samples throughout experiments. High-resolution B0 maps were acquired before each sequence, allowing shimming using Bruker's MAPSHIM. Both DKI data and structural data were acquired for each sample.

Diffusion kurtosis analysis: DKI data was collected using an 8-segmented diffusion-weighted spin-echo EPI sequence with a $150 \times 150 \mu\text{m}$ in-plane resolution and $250 \mu\text{m}$ slice thickness (60 slices for whole-brain coverage). Five unweighted volumes were acquired for signal normalization followed by 30 isotropically distributed encoding directions at each of three non-zero b-values (0.5, 1.0, 2.0 $\text{ms}/\mu\text{m}^2$). Additional scan parameters were time between diffusion gradients (Δ) = 15 ms, diffusion gradients duration (δ) = 6 ms, 20 averages, effective echo time (TE) = 27.7 ms, repetition time (TR) = 3500 ms, bandwidth = 278 kHz, resulting in a DKI scan time of 19h26m40s per animal. In addition, a rapid acquisition with relaxation enhancement (RARE) sequence with a $50 \times 50 \mu\text{m}$ in-plane resolution and $250 \mu\text{m}$ slice thickness was performed. Here, the 60 slices were positioned identically to the DKI data to allow for precise multi-atlas segmentation (MAS, details below) and ROI-specific extraction of DKI parameters. The scan parameters used were effective TE = 10.5 ms, TR = 3000 ms, 30 averages, and RARE factor = 2, with a scan time of 2h28m30s per animal. For the volumetric analysis, we acquired data with an isotropic resolution of $50 \mu\text{m}$ using a 3D fast low-angle shot (FLASH) sequence. Scan parameters for this were: TR = 88.5 ms, TE = 13.7 ms, matrix size = $360 \times 198 \times 300$, FOV = $18 \times 9.9 \times 15 \text{ mm}$, and 4 averages, resulting in a scan time of 6h31m21s per animal. For all scan types, data quality was ensured by visual inspection and samples rescanned if needed to ensure consistently high data quality for subsequent analyses. For diffusion kurtosis (DKI) analysis all DKI data from all samples were pre-processed in MATLAB (MathWorks Inc., v. 2022a) for noise floor correction, denoising and Gibbs ringing removal as described in^{46,47}. After pre-processing, DKI data analysis was performed using inhouse MATLAB scripts as previously described^{47,48} yielding metrics of mean water diffusivity (MD), tissue anisotropy (FA) and the mean kurtosis (MK, an index of tissue microstructure) in each voxel⁴⁷.

Multi-atlas segmentation: To systematically extract regional information from the mouse brain, a multi-atlas segmentation (MAS)^{46,49,50} was performed using 10 ex vivo NeAt templates from C57/BL6J mice (60,61). This followed procedures as described previously⁴⁶. Thus, high-resolution labelled images were obtained and then down sampled to match the in-plane resolution of the DKI data allowing extraction of DKI metrics from anatomically well-defined regions of interest (ROIs). Regional voxel values of DKI metrics were filtered for outliers (defined as values exceeding 3 times the median absolute deviation) and used for statistical analysis.

Volumetric analysis: The high-resolution FLASH images were processed using an in-house pipeline⁴⁶ applying B1 inhomogeneity correction⁵³, denoising⁵⁴, and intensity normalization. Spatial alignment with a high-resolution template of the C57BL/6J mouse⁵⁵ was done by manually initializing a linear registration⁵⁶ followed by a non-linear registration⁵⁷. Neuroanatomical labels from the C57BL/6J mouse atlas were subsequently transformed and resampled to data native space using the calculated deformation fields and affine transformations for calculation of individual regional volumes. From this, absolute brain volume and regional volumes were calculated. For all brains, relative regional volume (RRV; region size as % of total brain volume) was calculated to account for total brain size variation.

Statistics: To investigate group differences, permutation tests were performed for either total brain volume (1M permutations), regional volumes (40 regions, 100k permutations) or extracted voxelwise DKI metrics for each region (20 regions, 100k permutations). Uncorrected significance is reported for p values < 0.05 and indicated by asterisks (*). Pound symbols (#) indicate significance below $\alpha = 0.05$ divided by total number regions tested for group mean (green) and group median (red), respectively. Section signs (§) indicate significance ($p < 0.05$) with p values adjusted for false discovery rate (Benjamini-Hochsberg)⁵⁸.

Cortical thickness: Cortical thickness was calculated as previously described⁵⁹. Briefly, Laplace's equation, with fixed boundary conditions for each of the inner and outer surfaces, was solved. For this, the inner and outer surfaces of the cortex were defined based on the anatomical atlas and transformed to the given mouse. For each point on the cortical surface, the length of a streamline connecting the inside and outside surfaces was used to define the thickness. Cortical thickness was averaged within the bilateral frontal, occipital, and parieto-temporal lobes as well as the entorhinal cortex. Statistical maps of group differences in cortical thickness were generated by fitting a general linear model at each surface vertex (SurfStat,

<http://www.math.mcgill.ca/keith/surfstat/> [47]). Given the multiple comparisons performed, statistical maps were family-wise error (FWE) corrected using random field theory [38] with $\alpha = 0.001$ as cluster defining threshold. All statistical maps were thresholded at $p = 0.05$ (uncorrected and corrected).

Micro Computed Tomography (MicroCT)

Following MRI, the in-skulls brain samples were imaged using MicroCT scanning (vivaCT 80, Scanco Medical AG, Brüttisellen Switzerland). Skulls of 20 week old mice were placed in the scanner and imaged using 500 projections over 180°, an isotropic voxel size of 39 μm , X-ray voltage of 55 kVp, X-ray current of

105 μA , and an average time of 200 ms. Images were reconstructed and converted to DICOM files that were exported for subsequent analysis in 3D slicer (<http://www.slicer.org>) [48] and rendered in 3D. Skulls were isolated applying a threshold that separates bone from soft tissue. Segmented skulls and extracted endocasts were exported as 3D models. Furthermore, the coordinates of 45 surface landmarks were registered in a semi-automated fashion: a random skull was chosen as template model and landmarked manually, and these landmarks were automatically applied to the rest of the dataset using the ALPACA module in 3D Slicer. Additionally, the PseudoLMGenerator module was used to obtain a dense network of surface landmarks consisting of 768 points from the skulls and used as input for Principal Components Analysis (PCA). The image processing was performed with the operator blinded for the group distribution. For 3D visualization of differences between WT and OGT^{C921Y} mice by heat maps, models for the average shapes of OGT^{WT} and OGT^{C921Y} skulls were generated from the General Procrustes Analysis (GPA) module, aligned, superimposed, and the model-to-model distance was measured. Skull shape and size differences were assessed by measuring distances between the 45 surface landmarks and performing Euclidean distances matrix analysis. A previously collected cohort of 8 weeks old mice skulls were also analysed with this pipeline⁶⁰. One WT skull was excluded from analysis due to imaging artefact.

Histology and immunofluorescence (IF) microscopy

Following MicroCT scanning, formalin fixed and paraffin embedded brain sections (10 μm thickness) were obtained from the male WT and OGT^{C921Y} mice ($n = 6$ per group). To better characterize the radial and tangential cytoarchitecture in the neocortex, as well as to assess the bilaterality of any incidental findings in brain structure, one hemisphere was cut in sagittal orientation while the opposite hemisphere was cut in coronal orientation. Sections were deparaffinised and stained with haematoxylin and eosin, as described⁶¹. Serial sections were stained with cresyl violet for assessing the cellular arrangement of neurons in cortical layers, and separately with Luxol Fast blue for visualizing white matter distribution, including the arrangement of large tracts⁶¹. Then, high resolution views were obtained with an Olympus VS120 digital slide scanner equipped for bright field imaging. Slide scans were imported into Qupath (v. 0.5.1)⁶² and regions of interest (ROI) were outlined manually guided by the Mouse Brain Atlas (Paxinos and Franklin's The Mouse Brain in Stereotaxic Coordinates, 4th Edition). ROI were segmented using the Qupath cell detection on the haematoxylin channel.

Cellular identity was further verified by immunofluorescence (IF) microscopy using the following primary antibodies: anti-NeuN (1:500; ABN78; Sigma-Aldrich) and anti-glial fibrillary acidic protein- GFAP (1:500; ab68428; abcam). For this purpose, sections were deparaffinised and incubated in a blocking buffer comprising 5% normal goat serum in Tris-buffered saline for 1 h at room temperature. Then, the sections were incubated overnight at 4°C with the primary antibodies diluted in phosphate-buffered saline containing 0.3% Triton-X and 0.5% bovine serum albumin- BSA: IF detection was performed by fluorophore conjugated secondary antibodies: Alexa-Fluor488 goat anti-rabbit (1:1000; A-11034; Thermo Fisher) and Alexa-Fluor594 goat anti-mouse (1:1000; A-11005; Thermo Fisher). Image acquisition was performed on a Leica DM6000 upright microscope equipped with a fluorescence light source. Multiplex IHC analyses were performed in Discovery Ultra automated staining system (Roche Diagnostics) after

deparaffinization and heat-induced antigen retrieval using DISCOVERY CC1 buffer (Roche Diagnostics catalogue # 06414575001) for 32 minutes at 95 °C. Then, sections were sequentially incubated with the following primary antibodies: anti-NeuN (Rabbit polyclonal, Sigma-Aldrich catalogue # ABN78; dilution, 1:500), anti-Olig2 (clone: EP112; Rabbit monoclonal, Roche Diagnostics catalogue # 07667973001; dilution, 1:500), anti-Reelin (Goat polyclonal, Novus Biologicals catalogue # AF3820, dilution: 1:100) and anti- Pou3F2/BRN2 (Rabbit polyclonal, Proteintech catalogue # 14596-1-AP; dilution, 1:250). Chromogenic detection was performed using Discovery Omnimap Horseradish peroxidase (HRP)-conjugated secondary antibodies with following chromogens: DISCOVERY Teal HRP (for NeuN; Roche Diagnostics catalogue # 08254338001), DISCOVERY ChromoMap DAB (for Olig2: Roche Diagnostics catalogue # 05266645001), DISCOVERY Purple (for Reelin; Roche Diagnostics catalogue # 07053983001) and DISCOVERY yellow HRP (for POU3F2; Roche Diagnostics catalogue # 8502641001). In order to avoid cross-reaction between antibodies, denaturation between chromogenic detection of markers was performed using ULTRA CC2 buffer (Roche Diagnostics catalogue # 05424542001) at 100 °C for 24 minutes. High resolution views were obtained with an Olympus VS120 digital slide scanner in bright field imaging mode.

Tissue collection and dissociation for biochemical analyses

The prefrontal cortex was rapidly isolated from whole 16 week old male mouse brain (n = 3 per group), snap frozen and stored at -80 °C until processing. Tissues were disrupted in phosphate-buffered saline (PBS) two times at 5000 rpm for 30 s with 10 s break using a Precellys® 24 Touch homogenizer (Bertin Technologies). Homogenates were split in half for further protein and RNA extractions.

Mass Spectrometry (MS) and data analysis

Protein extracts from prefrontal cortex tissues were prepared for mass spectrometry using S-Trap micro spin columns (Protifi), including three washes with 50% CHCl₃/50% MeOH. Trypsin digestion (Proteomics grade, Sigma-Aldrich) was performed for 16 h at 37 °C. The resulting peptides were lyophilized and dissolved in 0.5% formic acid. LC-MS/MS was conducted using an EASY-nLC 1200 system (Thermo Scientific) connected to an Orbitrap Eclipse Tribrid Mass Spectrometer (Thermo Scientific) with a 2 cm trap column (100 µm i.d.) and a 15 cm analytical column (75 µm i.d.), both packed in-house with ReproSil-Pur C18-AQ 1.9 µm resin (Dr. Maisch GmbH). Peptides were eluted at 250 nl/min using an 80 min gradient from 5% to 44% phase B (0.1% formic acid and 80% acetonitrile), followed by a 30 s gradient to 100% phase B and 5 min at 100% B. Protein identification and quantification were performed using Proteome Discoverer 2.5 (Thermo Scientific). Data were searched against the mouse reference proteome (uniprot.org) using the Sequest search engine with the following parameters: MS error tolerance of 10 ppm, MS/MS error tolerance of 0.02 Da, trypsin as the protease with two missed cleavages, and carbamidomethylation as a fixed modification. Variable modifications included HexNAc (ST) and oxidation (M). Label-free quantification was based on precursor ions using unique peptides quantified in at least 2 out of 3 replicates. Peptide intensities were normalized to total peptide intensity and scaled using the average of all samples. Protein ratios were based on summed peptide abundances with imputation using replicate-based resampling. Significantly regulated proteins were identified using ANOVA, with adjustments for multiple testing.

Western immunoblotting

Brain homogenates were lysed using 10x RIPA buffer (Cell Signaling) as previously described³⁷. For MS, 50 mL of lysates were stored at -80 °C until further processing. For western blot, the rest of the lysates were centrifuged at 14,000 rpm for 20 min at 4 °C, and the protein concentration was determined with Pierce™ BCA Protein Assay kit (Thermo Scientific, 23227). Proteins (20 µg) were separated on precast 4-12% NuPAGE Bis-Tris Acrylamide gels (Invitrogen) and transferred to nitrocellulose membrane. Membranes were incubated with primary antibodies in 5% bovine serum albumin in Tris-buffered saline buffer with 0.1% Tween-20 overnight at 4 °C. Anti-OGA (1:1000 dilution; HPA036141; Sigma), anti-O-GlcNAc (RL2) (1:1000 dilution; NB300-524, Novus

Biologicals), anti-OGT (F-12) (1:1000 dilution; sc-74546; Santa Cruz), mouse anti-actin (1:5000 dilution; A5441; Merck) antibodies were used. Next, the membranes were incubated with IR680/800-labeled secondary antibodies at room temperature for 1 h. Blots were imaged using a Li-Cor Odyssey infrared imaging system (Li-Cor), and signals were quantified using Emperia software (Li-Cor). Results were normalized to the mean of each corresponding WT replicates set and represented as a fold change relative to WT.

RT-qPCR

Total RNA was purified from brain homogenates using RNAeasy Kit (Qiagen) as previously described³⁷. The threshold-crossing value was normalized to internal control transcripts (*18S*, *Actb*, and *Pgk1*). Results were normalized to the mean of each corresponding WT replicate set and represented as a fold change relative to WT.

Statistics

Statistical analyses were performed with Prism 9 (Graph Pad) unless specified otherwise. D'Agostino & Pearson, Shapiro–Wilk, and Kolmogorov-Smirnov normality tests were performed to verify normality. For data that fulfilled normality requirements, unpaired *t* tests were used for pairwise comparisons of WT and OGT^{C921Y} data or Two-way Anova for multiple comparisons were used. For data sets that did not fulfil normality, Mann-Whitney tests were used for pairwise comparisons.

Results

OGT^{C921Y} mice display postnatal growth development delay and hyperactivity

In patients, OGT-ID variants associate with balance difficulties, ataxia and hypotonia, suggesting possible locomotor defects in addition to ID and behavioural deficits⁶³. To investigate whether the OGT^{C921Y} variant phenocopies such deficits in mice, we first assessed a range of behavioural traits, including locomotion, anxiety, compulsivity, learning and memory in a mouse line carrying this variant, the generation of which has recently been reported³⁷. As part of our overall behavioural screening, animal body weight was monitored weekly. During the initial period after weaning (3 to 8 weeks old), no difference in body weight was observed between OGT^{WT} and OGT^{C921Y} mice suggesting similar developmental trajectories between both genotypes. However, we observed statistically significant lower gain in body weight in the OGT^{C921Y} mice from 9 to 20 weeks old compared to the WT mice (Time x Genotype $p = <0.0001$, 2-way ANOVA) (Fig. 1A), suggesting that OGT^{C921Y} mice show postnatal growth development delay.

Shortly after weaning, we placed the animals in digitally ventilated cages to continuously monitor the patterns of spontaneous activity in a non-invasive manner³⁸. For this purpose, we tracked animal activity longitudinally, starting at postnatal day 30 until 105 days of age (data collected over 75 consecutive days; $n = 10$ per group, housed in five separate cages). These experiments revealed hyperactive patterns of home cage activity by the OGT^{C921Y} mice, compared to the WT littermates, as early as day 35 ($p = <0.0001$, 2-way ANOVA) (Fig. 1B). Intriguingly, this period (postnatal 9-20 weeks) also marks the stage reflecting the observed postnatal growth development delay in the OGT^{C921Y} mice (Fig. 1A). Evaluation of daily rhythms and home cage activity during light/dark periods revealed that the OGT^{C921Y} mice exhibited frequent periods of hyperactivity during the dark phase (night time, active time for rodents), compared to the WT (Fig. 1C, D). However, the resting time (light phase, day time) was similar between the genotypes (Fig. 1C, D) ruling out potential disturbances in circadian rhythms, which otherwise could affect their normal day/night patterns of home cage activity.

We next investigated general locomotor activity in an open field arena. Mice were placed in open arenas and allowed to explore for 10 min on three consecutive days to evaluate both exploration of, and habituation to, an unfamiliar environment. Over the total three day period, OGT^{C921Y} mice




Figure 1. OGT^{C921Y} mice show postnatal growth development delay and increased spontaneous activity.

Significance is shown as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$. (A) Body weight of male OGT^{WT} ($n=15$) and OGT^{C921Y} ($n=16$) mice from 3 weeks to 20 weeks old. 2-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (B) Line chart representing the cumulative locomotion index (x-axis, days) for 75 days. Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons (OGT^{WT}, $n = 10$ housed in 5 cages; OGT^{C921Y}, $n = 10$ housed in 5 cages). (C) Line chart representing the cumulative locomotion index with light (6 am – 6 pm) and dark (6 pm – 6 am) periods in DVC cages over a longitudinal period up to 75 days (x-axis showing time in hours, 24-h format). The arrows indicate increased nocturnal activity by OGT^{C921Y} mice compared to WT littermates. (D) Heat maps representation of the cumulative locomotion index with light and dark periods in DVC cages for 75 days (x-axis showing time in hours over 24-hour period each day, y-axis showing days with Day 31 as the starting point on top). The arrow indicates increased nocturnal activity by OGT^{C921Y} mice compared to WT littermates. (E) Representative tracking plot over three consecutive days of a male OGT^{WT} and male OGT^{C921Y} mice in the open field arena. (F) Distance travelled over three consecutive days of male OGT^{WT} ($n = 15$) and male OGT^{C921Y} ($n = 16$) mice in the open field arena. 2-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (G) Maximal speed displayed by male OGT^{WT} ($n = 15$) and male OGT^{C921Y} ($n = 16$) mice in the open field arena. 2-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons.

showed an increase in activity in the arena as reflected by an increase in total distance travelled (Day 1 $p = 0.0011$; Day 2 $p = <0.0001$; Day 3 $p = <0.0001$, 2-way ANOVA) and maximum speed (Day 1 $p = 0.0013$; Day 2 $p = 0.001$; Day 3 $p = <0.0001$, 2-way ANOVA) compared to WT mice (Fig. 1E, F, G). To assess whether this hyperactivity was due to higher anxiety levels in OGT^{C921Y} mice, we evaluated the fraction of time spent in the periphery versus centre of the arena. Over the three days of testing, OGT^{C921Y} mice demonstrated thigmotaxic behaviour and increased distance travelled in both periphery (Day 1 $p = 0.0004$; Day 2 $p = <0.0001$; Day 3 $p = <0.0001$, 2-way ANOVA) and centre areas (Day 1 $p = 0.014$; Day 2 $p = 0.0012$; Day 3 $p = 0.0193$, 2-way ANOVA), but no preference for the periphery of the arena compared to WT, as shown by similar time spent in the periphery (Day 1 $p = 0.1771$; Day 2 $p = 0.2657$; Day 3 $p = 0.8885$, 2-way ANOVA) and centre areas (Day 1 $p = 0.1785$; Day 2 $p = 0.2543$; Day 3 $p = 0.8849$, 2-way ANOVA) between both genotypes (Fig. S1A-D). During Elevated Plus Maze (EPM), a test to assess elevated and open space anxiety, OGT^{C921Y} mice covered more distance ($p = <0.0001$, t -test) and performed higher number of entries in the centre ($p = <0.0001$, t -test) resulting in an increase in open/close time ratio ($p = 0.0063$, t -test) (Fig. S1E-H). Similarly, OGT^{C921Y} mice travelled more distance (0-5 min $p = 0.009$; 5-10 min $p = <0.0001$, 2-way ANOVA) and displayed frequent light-dark transitions compared to WT mice during the dark-light paradigm test (0-5 min $p = 0.001$; 5-10 min $p = 0.3374$, 2-way ANOVA), with no difference in time spent in the light compartment between both genotypes (0-5 min $p = 0.0926$; 5-10 min $p = 0.0608$, 2-way ANOVA) (Fig. S1I-L). These findings potentially rule out anxiety-like behaviour in both the OGT^{C921Y} and the WT mice.

We also assessed motor skills, including sensorimotor coordination and balance. No differences were observed in mean speed ($p = 0.9056$, t -test) or time spent on the rotarod ($p = 0.814$, t -test) between WT and OGT^{C921Y} mice (Fig. S2A, B). In the pole test, both OGT^{C921Y} and WT mice were able to perform a t-turn and reach the ground with similar scores ($p = 0.1523$, Mann-Whitney test) (Fig. S2C). During static rod tests, OGT^{C921Y} mice spent more time than their WT littermates to complete the required t-turn (35 mm $p = 0.0018$; 28 mm $p = 0.0003$; 22 mm $p = 0.1201$; 15 mm $p = 0.0005$; 10 mm $p = <0.0001$, 2-way ANOVA) although all mice reached the goal platform (Fig. 2A, B, C). Taken together these data rule out gross defects in motor coordination and balance in the OGT^{C921Y} animals, while revealing a degree of hyperactivity and stereotypy.

OGT^{C921Y} mice exhibit reduced digging behaviour

In order to further characterize the stereotypic phenotypes that could reflect ASD features observed in some OGT-ID patients⁶³, we assessed compulsive and repetitive behaviour. The OGT^{C921Y} showed a reduced number of buried marbles in a marble burying test ($p = 0.0002$, t -test) (Fig. 2D, E) and a reduction in litter displacement in a burrowing test ($p = 0.0107$, t -test) (Fig. 2F, G) compared to WT mice. To investigate whether the reduction in burying/burrowing activity was due to reduced digging behaviour, WT and OGT^{C921Y} were individually placed in cages containing 5 cm deep litter. The number of digging events and time spent digging during 3 min were measured. The OGT^{C921Y} mice exhibit both reduced number of digging events ($p = 0.0004$, t -test) and digging time ($p = 0.0003$, t -test) compared to WT mice (Fig. 2H, I). These findings were not due to lower activity of the OGT^{C921Y} mice, but rather to an increase in rearing events ($p = 0.1978$, t -test) and time spent rearing ($p = 0.0317$, t -test) compared to WT mice (Fig. 2J, K). The number of self-grooming events ($p = 0.7421$, t -test) and time spent self-grooming ($p = 0.3785$, t -test) were similar between the two genotypes (Fig. S2D, E). In addition, OGT^{C921Y} mice removed the same amount of material as WT mice during the nesting test ($p = 0.9774$, t -test) (Fig. S2F).

OGT^{C921Y} mice manifest features of impulsive behaviour and enhanced long term memory recall

We next investigated the effect of the OGT^{C921Y} variant on memory. First, we evaluated spontaneous alternation using a T-maze test as a measure of working spatial memory. The mice were placed in a T-shape maze and allowed to choose freely between right (R) and left (L) arms for 7 successive trials (Fig. 3A). Both OGT^{C921Y} and WT mice show similar percentage of




Figure 2. Motor, digging and burying activity.

Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. Student t test was used for statistics. (A) Representative image of experimental set-up for the static rods test. (B) Time to t-turn of male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during the static rods test. (C) Delta time defined as (Total time - time to turn) of male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during the static rods test. (D) Representative images of the marble test at 0 min (T₀) and 30 min (T₃₀). (E) Number of buried marbles by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during the marble test. (F) Representative image of the litter burrowing test. (G) Percentage of litter removed by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during the litter burrowing test. (H) Number of digging events by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during 3 min observation. (I) Time spent digging by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during 3 min observation. (J) Number of rearing events by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during 3 min observation. (K) Time spent rearing by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) during 3 min observation.

spontaneous alternation ($p = 0.2347$, t -test) (Fig. 3B [↗](#)), suggesting intact short-term spatial working memory in the OGT^{C921Y} mice. Nevertheless, the OGT^{C921Y} mice display a significantly reduced latency of choice ($p = 0.0175$, Mann-Whitney test) compared to WT mice (Fig. 3C [↗](#)), suggesting possible impulsive behaviour.

We also investigated novelty-associated memory using the Novel Object Recognition (NOR) paradigm. We used two different sets of objects to assess both short (90 min) and long (24 h) term memory. During the familiarisation phase, the mice were placed in an open field arena with two identical objects for 10 min. After 90 min or 24 h, mice were placed in the same arena containing one familiar object and one novel object (Fig. 3D [↗](#)). Time spent exploring each object was recorded allowing 20 s total exploration of both objects. Discrimination indexes were approached by calculating $(t_{\text{novel}} - t_{\text{familiar}}) / t_{\text{total}}$. After a 90 min interval, OGT^{C921Y} mice showed a similar discrimination index as WT mice ($p = 0.1482$, t -test), suggesting normal short term memory function in this assay (Fig. 3E [↗](#)). However, while both WT and OGT^{C921Y} show discrimination indices greater than 0.2 after 24 h, OGT^{C921Y} mice showed significantly greater discrimination index compared to WT mice ($p = 0.0434$, t -test), potentially indicating enhanced long term memory recall (Fig. 3F [↗](#)).

OGT^{C921Y} mice exhibit impaired plasticity during associative learning

To assess learning and memory function, we subjected OGT^{C921Y} mice and WT littermates to aversive Pavlovian trace conditioning - an established hippocampus- and cortex-dependent learning and memory paradigm where mice learn to associate a neutral tone with an aversive foot shock in a non-contiguous manner. Learning involves the presentation of a tone as the conditioned stimulus (CS), followed by a trace period, after which a foot shock is delivered as the unconditioned stimulus (US) (Fig. 3G [↗](#)). We used freezing, defined as the complete cessation of movement (except for breathing), as the conditioned response (CR) and a surrogate measure of short-term adaptation and aversive learning. Freezing behaviour was monitored during the acquisition phase and again 24 h later during a recall session (Fig. 3G [↗](#)). We observed that OGT^{C921Y} mice exhibited significant learning deficits, as indicated by a delay in freezing behaviour compared to WT littermates (Time x Genotype $p = <0.0001$, 2-way ANOVA) (Fig. 3H [↗](#)). This learning deficit cannot be attributed to somatosensory anomalies, as both genotypes displayed comparable responses to the US, as indicated by locomotor activity evoked by the foot shock (Time x Genotype $p = 0.3323$, 2-way ANOVA) (Fig. 3I [↗](#)). Furthermore, mutant mice reached the same CR levels as WT mice by the fourth CS-tone presentation (Fig. 3H [↗](#)), suggesting unimpaired auditory function. During the 24 h recall, there were no differences in freezing behaviour in OGT^{C921Y} mice compared to WT mice (Time x Genotype $p = 0.5337$, 2-way ANOVA) (Fig. 3J [↗](#)), suggesting an impaired plasticity during the acquisition phase. Taken together, these data suggest that OGT^{C921Y} exhibit impaired plasticity during associative learning.

OGT^{C921Y} mice exhibit reduced skull size and shape deformation

OGT-ID is often associated with microcephaly or craniofacial deformities, which were also recapitulated in our initial report using a small cohort ($n < 5$) of 2 month old OGT^{C921Y} mice³⁷. Here we aimed for a comprehensive skull shape analyses using micro-computed tomography (μ CT) in larger cohorts of older (5 month) mice (WT, $n = 14$; OGT^{C921Y}, $n = 16$). Average skull shapes were superimposed for OGT^{WT} and OGT^{C921Y} mice and skull shape differences between the two groups were analysed. The OGT^{C921Y} skulls were smaller in both the anterior and posterior areas, whereas the top of the skull appeared more curved compared to the WT skulls (Fig. 4A [↗](#)). We performed similar analyses on the previously reported μ CT data collected from 2 month old animals³⁷ to detect possible temporal aspects to these defects. The 2 month old OGT^{C921Y} mice show similar but more prominent shape differences than the 5 month old OGT^{C921Y} mice, suggesting an age-dependent convergence of skull shapes (Fig. 4A [↗](#)). Next, surface landmarks were used to perform Principal Component Analysis (PCA) allowing comparison of skull shape




Figure 3. OGT^{C921Y} show impulsivity, enhanced long term memory and delayed aversive associated learning.

Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. (A) Schematic of the T-maze paradigm. Mice are free to explore both arms for 7 additional trials. Each choice and latency were recorded. (B) Percentage of correct alternation (L-R/R-L sequences) of OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) during the T-maze test. Student *t* test was used for statistics. (C) Latency of arm entry of OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) during the T-maze test. Student *t* test was used for statistics. (D) Representative image of the Novel Object Recognition (NOR) test. During the familiarization phase, mice are free to explore an arena with two identical objects. During the test phase, mice were free to explore the same arena where one of the familiar objects has been replaced by a novel object to assess short (90 min) and long (24 h) memory. (E) Discrimination index of OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) during the short term (90 min) NOR test. Student *t* test was used for statistics. (F) Discrimination index of OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) during the long term (24h) NOR test. Student *t* test was used for statistics. (G) Diagram showing the behavioural paradigm (top), tone, conditioned stimulus (CS), trace period, and unconditioned stimulus, foot shock (US) lengths (bottom). (H) Freezing behaviour of OGT^{WT} and OGT^{C921Y} (n = 9 per group) during the acquisition session of the aversive conditioning. 2-way ANOVA time x genotype interaction $p < 0.0001$, $F(11,176) = 5.572$. (I) Locomotor activity of OGT^{WT} and OGT^{C921Y} (n = 9 per group) evoked by the US. The data represent the 3 s bin following the foot shock. Two-way ANOVA time x genotype interaction $p = 0.3323$, $F(3,48) = 1.166$. (J) Freezing behaviour of OGT^{WT} and OGT^{C921Y} (n = 9 per group) in the recall session where the CS were presented without US. The trace period refers to the 25 s following the CS presentation. Two-way ANOVA time x genotype interaction $p = 0.5337$, $F(11,176) = 0.9082$.

independently of size differences (Fig. S3B). Skulls from 5 month old OGT^{WT} and OGT^{C921Y} mice were separated along the first principal component (PC1) with a higher dispersion in the OGT^{C921Y} group suggesting heterogeneity in skull shape deformation phenotype penetrance. More than 63 % of the OGT^{C921Y} skulls display a positive PC1 value while 93 % of the WT skulls display a negative PC1 value. The skull of the OGT^{C921Y} mice has a shorter and rounder shape than the average WT mice (Fig. 4B). We performed Euclidean Distance Matrix Analysis (EDMA) using 45 relevant landmarks from the Richtsmeier laboratory resource (<https://getahead.la.psu.edu>) to identify the key regions contributing to the shape differences (Fig. S3A). At 5 months of age, 55% of the distances were at least 2% shorter in the OGT^{C921Y} skulls than in WT skulls, with the largest differences affecting the cranial base length (Fig. S3C). Similarly, in skulls from 2 month old animals, distances that are shorter than 5% in the OGT^{C921Y} skulls were predominantly located at the base of the skull (Fig. S3C). The main bone contributing to the cranial base length is the sphenoid, and shortening in the sphenoid bone determines the curvature of the cranial vault postnatally⁶⁴. This may indicate that defects in the postnatal development of the skull could cause the observed phenotype in the OGT^{C921Y} mice. Similar skull shape deformities have been reported in mouse models of Fragile X syndrome⁶⁵ and skeletal defect syndromes, such as osteogenesis imperfecta⁶⁶.

At both ages (i.e., 2 and 5 months old), the OGT^{C921Y} mice, the reduction in cranial base distances and overall reduction in skull size were obvious (2 months $p = 0.0016$; 5 months $p = 0.0101$, 2-way ANOVA) (Fig. 4C). From 3D reconstruction of the skulls, we determined the endocranial cavity volume as an indirect measure of the brain volume. At both ages tested, OGT^{C921Y} mice had a smaller internal cranial volume compared to their WT littermates (2 months $p = 0.013$; 5 months $p = <0.0001$, 2-way ANOVA) (Fig. 4D), suggesting a smaller brain volume. Taken together, these findings indicate that the OGT^{C921Y} mice exhibit antero-posterior skull growth defects leading to reduced skull size and shape deformation, suggesting dysmorphic features and reduced brain size.

OGT^{C921Y} display reduced cortical thickness and hypoplastic changes in brain structures

Neocortex hypoplasia, changes in cortical thickness and white matter integrity defects have been reported in mouse models of neurodevelopment disorders including ASD⁶⁷, CHARGE⁶⁸ and Rett⁶⁹ syndromes. To assess whether the reduction in brain size is general or localized to specific brain regions, we employed magnetic resonance imaging (MRI). Specifically, we performed volumetry analysis of anatomically well-defined brain regions based on high-resolution MRIs of 50 μm isotropic resolution, covering both the grey and white matter. The OGT^{C921Y} mice showed a reduced total brain volume compared to WT (Fig. 5A). Regional volumetric analyses showed a reduction in absolute volumes in most brain regions (Fig. S4). When expressed as regional relative volume (RRV) variation, the OGT^{C921Y} mice showed significantly reduced (group median) volumes of several brain regions including the frontal lobe of the cerebral cortex, corpus callosum, basal forebrain, globus pallidus, internal capsule and stria terminalis (Fig. 5C, S5). In contrast, RRV was increased in the lateral septum, medulla, hippocampus and hypothalamus regions of the OGT^{C921Y} mice compared to WT mice (Fig. 5C, S5). Cortical thickness analysis revealed significantly thinner cortex bilaterally in posterior regions in the OGT^{C921Y} mice compared to WT mice, with differences on the order of 100 μm (Fig. 5B).

We next assessed potential local brain microstructural defects using diffusion kurtosis imaging (DKI)^{70–73}, which provides sensitive indices of water mobility (mean diffusivity, MD) and diffusion directionality (fractional anisotropy, FA) in tissue as well as markers of tissue complexity (mean kurtosis, MK). Collectively, the MRI markers employed here are known to be sensitive to subtle tissue alterations in both rodent^{41,42,48,74–76} and human brain^{77–80}. We observed no difference in these DKI metrics between OGT^{C921Y} and WT mice in any of the 20 automatically segmented regions including neocortex (Figs. S6–9). Taking together, these MRI data suggest that OGT^{C921Y} show preserved brain microstructure at the resolution (150 μm x 150 μm x 250 μm) of the DKI experiments yet reduced cortical thickness.




Figure 4. Micro-computed tomography of OGT-ID mouse skulls indicates a reduced endocranial volume and shorter skull length.

Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. (A) Heatmaps obtained of the average OGT^{C921Y} skull ($n = 16$) to the average OGT^{WT} skull ($n = 14$). Red and blue regions indicate that the average OGT^{C921Y} skull is smaller or larger respectively than the average OGT^{WT} in those areas. (B) PCA biplot of PC1 and PC2 of the surface landmarks of 20 weeks old male OGT^{WT} and OGT^{C921Y} skulls (top). Percentages in the axis indicate the explained variance. Deformations in PC1 showing shape differences across this axis are shown (bottom). (C) Volume of the skulls of 8 weeks and 20 weeks old male OGT^{WT} and OGT^{C921Y} mice. Student t test was used for statistics. (D) Volume of the extracted endocranial cavity of 8 weeks and 20 weeks old male OGT^{WT} and OGT^{C921Y} mice. Student t test was used for statistics.




Figure 5. OGT^{C921Y} mice show changes in regional brain volume and reduced cortical thickness.

Volumetry and cortical thickness from high resolution T1-weighted Magnetic Resonance (MR) images. (A) Total brain volume of 20 weeks old male OGT^{WT} and OGT^{C921Y} mice. Groupwise total brain volume where each dot represents one subject. The horizontal lines correspond to group extrema and median. Asterisks (*) mark significance ($p < 0.05$) based on permutation tests (1M permutations) of either the mean (green asterisk) or median (red asterisk) of the two groups. (B) Heatmaps of group differences in cortical thickness between male OGT^{WT} and OGT^{C921Y} mice. Maps are effect sizes (top row), p-values of significant differences both uncorrected for multiple comparisons (middle row) and corrected with family-wise error (FWE). $N = 15$ per genotype. One mutant was not perfused correctly and removed from analysis. (C) Regional brain volumes represented as percentage of whole brain volume of male OGT^{WT} and OGT^{C921Y} mice. Asterisks (*) indicate uncorrected significance ($p < 0.05$), based on permutation tests (100k permutations for each region). Pound symbols (#) indicate significance below alpha (0.05) divided by total number regions tested (40) for mean (green) and median (red), respectively. Section signs (§) indicate significance ($p < 0.05$) with p-values adjusted for false discovery rate (Benjamini-Hochsberg, BH). Y-axes are scaled to individual ROIs to highlight group variation and difference.

OGT^{C921Y} mice exhibit dysmorphic features in superficial cortical layers, predominantly affecting the cingulate

Previous studies indicate that congenital dysplasia in cortical organization manifests as a range of malformations including disrupted cortical laminar organization, neuronal heterotopia in the subcortical white matter, misplaced neurons in cortical lamina I, clustering of neurons in the grey matter, and the presence of dysmorphic neurons^{81–83}. To evaluate cortical cytoarchitecture at the microscopic level and for detecting any features suggestive of focal cortical dysplasia (FCD) in OGT^{C921Y} mice, we subjected the brain sections to histological analyses using H&E (for cells), cresyl violet (for neurons), luxol fast (for white matter) staining (Fig. S10 [↗](#)). In the H&E analyses, we observed an overall normal 6-layered cortical organization across the regions involved in primary sensorimotor modalities, including the somatomotor cortex M1/M2, primary somatosensory cortex S1/S2, auditory cortex and primary visual cortex V1 (Fig. S11 [↗](#)). Intriguingly, part of the cingulate (retrosplenial Area 29/Area 30, according to Paxinos and Franklin) found within the paramedian portion (sagittal section, interaural 0.36) showed FCD in 5 out of 6 animals in the OGT^{C921Y} cohort (Fig. 6A–C [↗](#), compared to WT). The most conspicuous microscopic finding suggestive of focal cortical dysplasia was seen in the form of pseudosulcus formation in the cingulate, such that the superficial cortical layers (I–III) appeared to be displaced inwards (Fig. 6B–C [↗](#), black arrow), and resembled polymicrogyria similar to those observed in other congenital brain malformations^{84,85}. In two extreme cases, nodular arrangements of cells within the superficial layers (II/III bordering IV), with a central halo containing loosely arranged eosinophilic tissue, were observed (Fig. 6B–C [↗](#), yellow arrow). Having ruled out gross structural alterations (except in the cingulate cortex), we next assessed cortical cell density, which would reveal hypo or hyperproliferative neurodevelopmental anomalies. Total cell density analyses in the neocortical regions (H&E stained serial sections, reflecting both neuronal and non-neuronal cells), did not indicate drastic differences between the two groups in the major cortical regions examined (Fig. 6D [↗](#)). Staining analyses using cresyl violet indicated that the nodular collections in the layers II/III bordering IV were predominantly neuronal cell bodies (Fig. 6E [↗](#)). In luxol fast stained sections, these malformations contained collections of ectopic white matter, which populated the center of nodular malformations (Fig. 6F [↗](#)) and/or found as patchy deposits in the vicinity of superficial layers (II–IV) (Fig. 6C [↗](#)). To further establish the cell identity in the cortical malformations, we performed immunofluorescence analyses using neuronal nuclei marker (NeuN) and astroglial marker glial acidic fibrillary protein (GFAP). These analyses further corroborated the notion that the foci of FDC in the cingulate of OGT^{C921Y} mice were predominantly populated by neuronal cells and were devoid of astrocytes, which otherwise would indicate reactive astrogliosis seen in states of brain trauma (Fig. 6G–H [↗](#), compare WT in Fig. 6G [↗](#) to OGT^{C921Y} in Fig. 6H [↗](#)). Furthermore, normal localization of astroglial cells in the subcortical white matter was evident in the WT and OGT^{C921Y} cohorts (Fig. 6G–H [↗](#), merged images in the panels on right). In parallel, multiplex IHC analyses confirmed that the nodular cortical malformations predominantly affected layers II–V, as indicated by the distribution of neuronal markers Reelin and POU class 3 homeobox 2 (POU3F2), and were populated by neurons and oligodendrocytes in the vicinity (Fig. S12 [↗](#)). Taken in conjunction with the MRI (reduced cortical thickness), the cortical dysplasia revealed by the histology analyses suggest a neurodevelopmental component to OGT-ID affecting the cortical superficial layers, which in OGT^{C921Y} mice predominantly affects the cingulate.

Quantitative proteomics analyses revealed distinctly perturbed molecular pathways in prefrontal cortex of the OGT^{C921Y} mice

In order to decipher the molecular phenotypes that could be associated with the cortical defects in the OGT-ID mutant strain, we performed proteome analysis on the prefrontal cortex (PFC) of male OGT^{C921Y} and WT mice using label-free quantification. A total of 5488 proteins were identified with a 1 % false discovery rate which we then ranked as top 50 most affected (and potentially revealing) factors. This ranking was based on defined cut-off criteria for both the upregulated (p value ≤ 0.05 ; log-2 fold change ≥ 1.45) and down-regulated proteins (p value ≤ 0.05 , log-2 fold




Figure 6. Pseudosulcus formation, cortical dysplasia and ectopic white matter in the cingulate cortex of OGT^{C921Y} mice.

(A-B) Representative low magnification (10X) sagittal views of WT (in A) and OGT^{C921Y} mice (in B), showing cortical malformations in the cingulate cortex of the latter by H&E staining. The black arrow points to instances of pseudosulcus formation and the yellow arrow points to instances of cortical dysplasia observed in the layers II-IV (scale bar = 500 μ m). (C) Representative high magnification (40X) views extracted from the insets in A except one marked with *, which is depicting nodular arrangement of cells in a brain section from a different OGT^{C921Y} animal (scale bar = 100 μ m). (D) Regional cell density assessed in 10X views (H&E staining) of two-serial sections from WT and OGT^{C921Y} mice. Error bars indicate Mean $\pm$ SEM (n = 6 / group; Mann-Whitney test, none significant). Regions examined include: (coronal sections) primary motor cortex M1/M2, primary somatosensory cortex S1/S2, auditory cortex (A); (sagittal sections) visual cortex V1 and cingulate cortex (C). Also see Fig. S10 depicting coordinates and region demarcations (dashed rectangles) used in the cell density analyses. (E-F) Representative low magnification (10X) sagittal views showing cresyl violet staining for neuronal cell bodies (in E) and myelin staining by luxol fast (in F). Notice the instances of ectopic white matter (red arrows) in the cingulate cortex of OGT^{C921Y} mice (scale bar = 200 μ m). in 6F.

change ≤ -1.33). Then we performed pathways enrichment and downstream analyses using the built-in tools of the STRING database (Fig. 7A, B). To begin with, the expression of OGA was significantly downregulated in the cortex of OGT^{C921Y} (Fig. 7B) in the mass spectrometry analyses, as well as in western blotting analyses of the cortex ($p = 0.0223$, t -test) (Fig. S13A, C) as we have reported previously⁶⁰. Although no significant changes in the expression of OGT were observed in OGT^{C921Y} cortex ($p = 0.788$, t -test) (Fig. S13A, D), there was a significant increase in OGT/OGA protein ratio in OGT^{C921Y} mice ($p = <0.0001$, t -test) (Fig. S13E), which is driven by the reduction in OGA levels to compensate for reduced OGT activity. Despite this compensatory mechanism, global O-GlcNAcylation of proteins in the brain was drastically impaired in OGT^{C921Y} brain compared to WT ($p = 0.0018$, t -test) (Fig. S13O, Q). The perturbed regulation of *Ogt/Oga* ratio (Oga mRNA $p = 0.0087$; Ogt mRNA $p = 0.0894$; Ogt/Oga ratio $p = <0.0001$, t -test) was further confirmed at the transcriptional level by RT-PCR (Fig. S14A, B, C). Of note, similar O-GlcNAc dyshomeostasis at protein/mRNA levels was observed in other brain regions analysed except for OGT protein levels that were found reduced in both hippocampus and cerebellum (Fig. S14), suggesting brain region specific effect of the OGT^{C921Y} mutation. In the prefrontal cortex region, gene ontology analyses in the STRING database pointed to significant upregulation in the pathways regulating cellular processes (GO:0009987), cellular metabolic pathways (GO:0044237), nervous system development (GO:0007399), protein catabolic process (GO:0030163) and lysosomal transport (GO:0007041) (Fig. 7C). Among the top up regulated proteins identified in our dataset (Fig. 7A), thirty-five proteins belong to neurodevelopmental pathways, including proteins involved in neuronal migration (PLXND1, FAT3, ASTN2, NEUROD1), neurogenesis (NEUROD1, RBBP5, RBBP6) and synaptic function (CLCN3, SORCS3, AP3B1).

Intriguingly, we also identified significant downregulation in the pathways regulating small molecule metabolic process (GO: 0044281), carbohydrate derivative biosynthetic process (GO: 1901137), cellular respiration (GO: 0045333), ATP metabolic process (GO: 00046034) and mitochondrial acetyl-CoA biosynthetic process from pyruvate (GO:0061732) (Fig. 7D). Moreover, the perturbed proteomic signatures in the PFC corresponded to distinct Monarch enrichment profiles⁸⁶ in the STRING database including abnormal CNS myelination (HP:0011400), agenesis of corpus callosum (HP: 0001274), upper motor neuron dysfunction (HP: 0002493) and abnormal muscle physiology and function (HP: 0011804; HP: 0003808; HP: 0001252; HP: 0001319; HP 0003394) (Fig. S13A, B).

Taken together, these data suggest that perturbations in protein O-GlcNAcylation in brain of OGT^{C921Y} are associated with distinct alterations in molecular pathways that potentially impact brain development.

Discussion

Although several OGT variants linked to ID have been reported, the mechanisms underlying the disorder remain unknown. Potential mechanisms that have been proposed⁶³ (Fig. 8A) include loss of O-GlcNAcylation on OGT substrates important for brain function and development^{28,37,63}, OGT aggregation due to reduced OGT stability⁶³, impaired OGT interactome⁸⁷, misprocessing of HCF1¹⁵, or loss of OGA^{27,28,37}, a common feature observed in models of OGT-ID. Lastly, O-GlcNAc dyshomeostasis has been recently suggested as a common mechanism in OGT-ID variants and has been proposed as a biomarker to identify new pathogenic variants using a stem cell reporter line⁸⁸. Furthermore, these hypotheses are not mutually exclusive and can individually or in combination account for the cognitive dysfunction due to impaired brain development, defects in synaptogenesis/synaptic pruning and/or neural transmission impairment (Fig. 8B). Vertebrate models are a key step towards deciphering the mechanisms linked to the disease, and here we have described a detailed characterization of a mouse model of one OGT-ID variant (OGT^{C921Y}).

The OGT^{C921Y} variant is found in a Danish family with three affected male siblings that inherited the variant through their affected mother. The male siblings exhibit dysmorphic features, ID with poor language skills and late onset epilepsy. Autistic features including repetitive mannerisms, and hyperactivity have also been reported³². Mild hyperactivity has also been reported in a patient carrying another OGT-ID variant⁸⁹. The OGT^{C921Y} mice recapitulate features of impaired cognitive




Figure 7. Quantitative proteomics analyses revealed distinctly perturbed molecular pathways in prefrontal cortex of the OGT^{C921Y} mice.

(A) List of Top 50 proteins up-regulated (cut off criteria p value 0.05, log 2 change 1.45). (B) List of Top 50 proteins down-regulated (cut off criteria p value 0.05, log 2 change -1.33). (C) Up-regulated biological pathways enrichment in the STRING database. (D) Down-regulated biological pathways enrichment in the STRING database.




Figure 8. Graphical summary of the neurodevelopmental, structural and behaviour defects identified in the OGT^{C921Y} mice.

This Figure was created using BioRender.com [Sevillano quispe, O. \(2026\) https://BioRender.com/n211133](https://BioRender.com/n211133)

function, impulsive behaviour and increased spontaneous activity (or hyperactivity) both in their home cage and in the open field arena. These phenotypes (hyperactivity and impulsivity) are often observed in mouse models recapitulating human ID and autism disorders. For instance, *Fmr1*^{Ko} mice exhibit hyperactivity in open field assays and increased exploratory behaviour as seen by an increased number of light-dark transitions in the dark-light paradigm similarly to our observations in the OGT^{C921Y} mice^{90–92}. Moreover, hyperactivity and impulsivity behaviours present in a haploinsufficiency mouse model of *Kdm6b* were rescued with methylphenidate, a neurostimulant used to alleviate attention deficit and hyperactivity in individuals with ADHD⁹³. In addition, the stereotypy and increased rearing activity observed in the OGT^{C921Y} mice suggest locomotor compulsive behaviour similar to that reported in the individuals carrying the C921Y variant³². We have observed that, unlike the WT mice, the locomotor activity of the OGT^{C921Y} mice remained unaffected during repeated measurements in the open field test, which could suggest an impairment in spatial habituation learning. Interestingly, defects in habituation learning were also observed in *Drosophila* models carrying OGT-ID variants during a light-off jump habituation test⁹⁴. As habituation corresponds to the simplest form of learning⁹⁵, these data hint towards some learning and memory defects in OGT^{C921Y} mice. This is further indicated by a delay in freezing behaviour during acquisition compared to WT littermates during the aversive Pavlovian paradigm, a well-established hippocampus- and cortex-dependent learning and memory.

Reduction of brain size has been reported in OGT-ID patients, including microcephaly and brain atrophy²⁹. In the present study, both the shape and morphology of the skulls of the OGT^{C921Y} mice were affected. The morphological alteration of the skulls reduced the endocast volume (as an estimate of brain volume) significantly in both 2 month old and 5 month old OGT^{C921Y} mice compared to their WT littermates. MRI brain structural analyses indicate that redistribution of regional volumes rather than overall reduction underpins the lower total brain volume in the OGT^{C921Y} mice. The frontal lobes of the cerebral cortex were prominently affected and associated with bilateral reduced cortical thickness suggesting overall reduction of cortical grey matter. Reduced brain volume, in particular the prefrontal cortex region, has also been associated with several neurodevelopment disorders in human including ADHD^{96,97} and associated rodent models⁹⁸. Moreover, reduced brain volume and grey matter density in cerebellum and cortical regions are observed in adolescents with ASD associated with intellectual deficits⁹⁶.

While the MRI data suggest an overall reduction in the size of the prefrontal cortex, histological analyses revealed FCD and pseudosulci formation in areas of the cingulate (retrosplenial) cortex in the OGT^{C921Y} mice with patchy distribution of ectopic white matter in the vicinity of these cortical malformations. These features are reminiscent of cortical polymicrogyria seen in some types of CDG disorders such as muscular dystrophy-dystroglycanopathy (MDDG) syndromes^{84,85}, and in some inborn errors of metabolism characterized by defective neural migration⁹⁹. Given that OGT^{C921Y} mice exhibit considerable reduction in cortical protein O-GlcNAcylation, it is plausible to postulate that these forms of cortical dysplasia (pseudosulci) arise as neurodevelopmental sequelae to the perturbed O-GlcNAcome. FCD has been linked to cognitive impairment in certain X-linked intellectual disabilities (XLIDs), such as Fragile X syndrome¹⁰⁰. FCD arises from localized malformations in the cortex caused by disruptions in neuronal proliferation, differentiation, and migration during corticogenesis^{101–103}. However, some studies suggest that similar disruptions in the organization of cortical layers may also arise due to disorders of post-migrational development of neural progenitors and not solely due to defects in proliferation or migration¹⁰⁴. Therefore, it remains to be determined whether the features of cortical malformation (microcephaly and disturbed cytoarchitecture in cingulate) in the OGT^{C921Y} mice pertain to these categories. Behavioural and neuroimaging studies indicate that distinct regions within the cingulate participate in complex cognitive tasks involving affective response and decision making^{105,106}. Meta-analyses of neuroimaging studies in human subjects with psychiatric disease (major depressive disorder, bipolar disorder, schizophrenia, anxiety and addiction) purport grey matter loss in the dorsal anterior cingulate as a frequent finding^{107,108}. Taken together, these suggest a link between the cortical structure defects and the behavioural deficits observed in the OGT^{C921Y}

mice. However, further systemic interventions (e.g. optogenetic manipulations) will be needed to investigate whether the behavioural phenotypes of OGT^{C921Y} mice solely arise from cortical dysplasia in the cingulate cortex.

There is a significant dearth of information on possible cortical malformations in patients with OGT-ID. This is largely due to the lack of clinical data available as brain imaging has only been reported in a few cases, or that the defects are beyond the limits of detection/resolution in routine clinical MRI scans. Nevertheless, defects in white matter composition have been reported in patients with OGT-ID including thin corpus callosum and periventricular leukomalacia²⁹. Both defects have previously been associated with learning difficulties and ID^{96,109}. Changes in white matter structures have also been found in neuroimaging analysis of neurodevelopment disorders such as Fragile-X¹¹⁰, developmental delay^{111,112} and ADHD¹¹³. Together, these observations suggest that changes in white matter structures could underline learning and behavioural deficits in neurodevelopment disorders. Interestingly, the OGT^{C921Y} mice also showed reduced volumes of several white matter structures including corpus callosum, internal capsule, cerebral peduncle, corticospinal tracts and stria terminalis. These observations could guide future studies in investigating whether OGT-ID variants cause impaired myelination processes leading to cognitive impairment.


Our proteomics analyses revealed a number of perturbed molecular pathways in the prefrontal cortex of the OGT^{C921Y} mice. Interestingly, some of the deregulated proteins are associated with brain malformation including progressive microcephaly (ARSA, CHMP1A, QARS1)^{114–116}, cortical dysplasia (RALGAPB, PACS2)^{90,9}, myelination defects (ARSA, CHMP1A, GABRA2, HEXA, RMND1)^{114,117–121} and agenesis of the corpus callosum (EFNB1, TUBA8, CLCN3, PLXND1)^{122–125}, providing potential candidates underlying the brain defects identified in the OGT^{C921Y} mice. In addition, the RHO GTPase cycle pathway (HAS-9012999) was significantly upregulated from the Reactome database. RHO GTPase family are involved in cell migration, division and polarity and play crucial roles in neurodevelopment¹²⁶. While a detailed description of all the top 50 upregulated proteins is beyond the scope of this Discussion, the ubiquitin ligase RB binding protein 6 (RBBP6) promotes ubiquitination of the Y box protein 1 (YBX1) followed by its degradation by the proteasome. YBX1 is a nucleic acid binding protein required for forebrain specification, cell proliferation and neuronal differentiation through the suppression of RNA polymerase II mediated transcription¹²⁷. Interestingly, while RBBP6 was found up regulated, its target YBX1 was identified as significantly down regulated in our dataset, suggesting a role of the RBBP6/YBX1 axis in the brain structural defects observed in the OGT^{C921Y} mice. Future studies at early timepoints during brain development will be needed to explore these candidate conveyors of OGT-ID.

Whereas structural brain defects suggest a neurodevelopment origin, it is as yet unknown whether the associated cortical dysfunction originates from structural defects during neurodevelopment or neurophysiological defects due to the continuous presence of the catalytically impaired OGT^{C921Y} variant in neurons, their synapses and glia. Non-invasive monitoring of the OGT^{C921Y} mice activity patterns (in DVC) showed that spontaneous hyperactivity was not detected from weaning but continuously progressed from approximately 7 weeks of age, coinciding with a delay in postnatal growth. With reasonable caution, this is reminiscent of clinical findings in patients with OGT-ID who generally start developing symptoms in the first two years. From a translational perspective, these features hint towards a progressive postnatal appearance of behaviour and morphological deficits in the OGT^{C921Y} mice, thus potentially offering a therapeutic window for future interventions. It is hoped that cognitive impairment due to prefrontal cortical dysfunction can partly be rescued postnatally even in presence of brain structural defects, with notable examples from research in models of Fragile X and Rett's neurodevelopmental syndromes^{69,97,100}.

Conclusions

In conclusion, we report that O-GlcNAc dyshomeostasis in brains of OGT^{C921Y} mice is associated with distinct behavioural phenotypes reflecting hyperactivity, impulsivity and learning deficits. These phenotypes were accompanied by features consistent with perturbed neurodevelopment including skull deformation, microcephaly and focal cortical dysplasia in the cingulate cortex (Fig. 8B). Moreover, the glimpse offered by changes in the neocortical proteome of OGT^{C921Y} mice and the observed O-GlcNAc dyshomeostasis will guide future studies in unravelling the pathophysiology of the disorder, and hold promise for the development of novel therapeutic interventions in OGT-ID.

Supplementary material




Supplementary Figure S1. OGT^{C921Y} mice show similar anxiety-like behaviour and increased exploration

Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. (A) Distance travelled in periphery by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the open field arena. Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (B) Time spent in periphery by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the open field arena. Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (C) Distance travelled in centre by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the open field arena. Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (D) Time spent in centre by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the open field arena. Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (E) Representative tracking plot of male OGT^{WT} and OGT^{C921Y} mice in the elevated plus maze (EPM). (F) Total distance travelled of male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the EPM. Student t test was used for statistics. (G) Number of centre entries of male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the EPM. Student t test was used for statistics. (H) Ratio between time spent in open and close arms by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice in the EPM. Student t test was used for statistics. (I) Representative tracking plot of male OGT^{WT} and OGT^{C921Y} mice in the light compartment during the dark-light paradigm (D/L). (J) Total distance travelled of male

OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) mice in the light compartment during the dark-light paradigm (D/L). Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (K) Number of centre entries of male OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) mice in the light compartment during the dark-light paradigm (D/L). Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons. (L) Time spent of male OGT^{WT} (n = 15) and OGT^{C921Y} (n = 16) mice in the light compartment during the dark-light paradigm (D/L). Two-way ANOVA (Alpha, 0.05) followed by Tukey's column comparisons.


Supplementary Figure S2. OGT^{C921Y} mice show similar motor skills


Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. (A) Mean score reached by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice on the rotarod. (B) Mean time spent by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice on the rotarod. (C) Score reached by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice during the pole test. (D) Number of self-grooming events male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice during 3 min observation. (E) Time spent self-grooming by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice during 3 min observation. (F) Percentage of material removed from a cotton pad by male OGT^{WT} ($n = 15$) and OGT^{C921Y} ($n = 16$) mice during 3 min during the nesting test.



Supplementary Figure S3. OGT^{C921Y} mice show reduced distances in the cranial base.

(A) Lateral and superior views of a representative microCT 3D reconstruction of a mouse skull presenting the 45 landmarks used for Euclidean distance matrix analysis (EDMA) (left) and surface markups used for Principal Components Analysis and average shapes generation (right). (B) Lateral and inferior views of representative 3D reconstructions of mouse skulls presenting the linear distances that are at least 5% shorter in 2 and 5 months old OGT^{C921Y} skulls ($p < 0.01$, two-tailed unpaired t-test used)






Supplementary Figure S4. Volumetry from segmentation of T1-weighted structural images (FLASH) from Magnetic Resonance Imaging (MRI) illustrated as absolute volumes.

Asterisks (*) indicate uncorrected significance ($p < 0.05$), based on permutation tests (100k permutations for each region). Pound symbols (#) indicate significance below alpha (0.05) divided by total number regions tested (40) for mean (green) and median (red), respectively. Section signs (§) indicate significance ($p < 0.05$) with p values adjusted for false discovery rate (Benjamini-Hochsberg, BH). Y-axes are scaled to individual ROIs to highlight group variation and difference.




Supplementary Figure S5. Relative volumetry from T1 segmentation.

Volumetry from segmentation of T1-weighted structural images (FLASH) from Magnetic Resonance Imaging (MRI) illustrated as regional relative volumes (divided by individual total brain volume). Asterisks (*) indicate uncorrected significance ($p < 0.05$), based on permutation tests (100k permutations for each region). Pound symbols (#) indicate significance below alpha (0.05) divided by total number regions tested (40) for mean (green) and median (red), respectively. Section signs (§) indicate significance ($p < 0.05$) with p values adjusted for false discovery rate (Benjamini-Hochsberg, BH). Y-axes are scaled to individual ROIs to highlight group variation and difference.




Supplementary Figure S6. OGT^{C921Y} mice show conserved DKI metrics for Neocortex


(A) Middle coronal section of subject:207 illustrating the different calculated DKI metrics: Mean Diffusivity (MD), Fractional Anisotropy (FA) and Mean Kurtosis (MK). (B) Extracted DKI metrics (MD, FA and MK) for Neocortex of all subjects in each group for neocortex of all mice.




Supplementary Figure S7. Subject-wise median of extracted Fractional Anisotropy (FA) values in 20 selected regions of interest (ROI).



Supplementary Figure S8. Subject-wise median of extracted Mean Diffusivity (MD) values in 20 selected regions of interest (ROI).




Supplementary Figure S9. Subject-wise median of extracted Mean Kurtosis (MK) values in 20 selected regions of interest (ROI).



Supplementary Figure S10. Panoramic views of the H&E stained sagittal and coronal brain sections depicting neuroanatomical landmarks used for measurements of cortical thickness and cell density.

(A) Representative images showing coronal panoramic views of the brain sections used for the measurements of cortical cell density. The sections were used to sample regions in the cortex at the level of Bregma 0.14 and Bregma -2.06. Regional cell counts were obtained from primary motor cortex M1/M2 and primary somatosensory cortex S1/S2, and from the auditory cortex. (B) Representative images showing sagittal panoramic views of the brain sections used for the measurements of cortical cell density. The sections were used to sample medial (interaural, 0.36, in C) and lateral (interaural, 2.28, in D) regions of the cortex. Regional cell counts were obtained from cingulate cortex and visual cortex V. Abbreviations: Aud/AuV, auditory cortex (in B); CPu, caudate putamen; superior colliculus. Neuroanatomical annotations are based on the Paxinos and Franklin's Mouse Brain in Stereotaxic Coordinates, Elsevier Publishing, 4th Edition.



Supplementary Figure S11. Low magnification (10X) views of the H&E stained sagittal and coronal brain sections used for assessing cortical organization.

Representative images showing low magnification (10X) views of select cortical regions from OGT^{WT} (in A) and OGT^{C921Y} (in B) mice. Areas shown include primary motor cortex M1/M2, primary somatosensory cortex S1/S2, auditory cortex and visual cortex V1. Notice a mild degree of cortical dysplasia in layers II-III in the auditory cortex of OGT^{C921Y} mice (yellow arrow). The images were obtained from coronal sections (Bregma, 0.14, for M1/M2 and S1/S2; Bregma, -2.06, auditory cortex) and sagittal sections (Interaural 2.28, visual cortex V1). Scale bar = 200 μ m. Also see Fig. S10. Neuroanatomical annotations are based on the Paxinos and Franklin's Mouse Brain in Stereotaxic Coordinates, Elsevier Publishing, 4th Edition.





Figure S12. Histological characterization of cortical malformations by multiplex IHC.

(A-B) Representative low magnification (20X) sagittal views of WT (A) and OGT^{C921Y} mice (B), showing cellular populations in the vicinity of nodular cortical malformation in the cingulate cortex of the latter using multiplex IHC chromogenic detection (NeuN in teal, Olig2 in brown, Reelin in purple and POU3F2 in yellow)- scale bar = 200 μ m. Numerals I-VI in S12a-d indicate neocortical layers. (C-D) Representative high magnification (40X) views extracted from the insets in A (wild type) and B (OGT^{C921Y}) respectively. Notice the undulating pattern of NeuN in layers II/III (black arrow, in B) and mixed cellular populations in the vicinity of cortical malformations in layers II-V of an OGT^{C921Y} animal-scale bar = 100 μ m.




Supplementary Figure S13. *OGT^{C921Y}* mice show altered OGT/OGA ratio and global reduction in O-GlcNAc levels in brain regions.

Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ ($n = 3$ per group). Student t test was used for statistics. (A) Western blot of OGA and OGT level in male prefrontal cortex and hippocampus of *OGT^{WT}* and *OGT^{C921Y}* mice. Actin antibodies were used as loading control. (B) Western blot of OGA and OGT level in male striatum and cerebellum of *OGT^{WT}* and *OGT^{C921Y}* mice. Actin antibodies were used as loading control. Quantification of OGA (C), OGT (D) and OGT/OGA ratio (E) proteins levels in prefrontal cortex from the Western blot in panel A. Quantification of OGA (F), OGT (G) and OGT/OGA ratio (H) proteins levels in striatum from the Western blot in panel B. Quantification of OGA (I), OGT (J) and OGT/OGA ratio (K) proteins levels in hippocampus from the Western blot in panel A. Quantification of OGA (L), OGT (M) and OGT/OGA ratio (N) proteins levels in prefrontal cortex from the Western blot in panel B. (O) Western blot of O-GlcNAc levels in prefrontal cortex and hippocampus of male *OGT^{WT}* and *OGT^{C921Y}* mice. Actin antibodies were used as loading control. (P) Western blot of O-GlcNAc levels in striatum and cerebellum of male *OGT^{WT}* and *OGT^{C921Y}* mice. Actin antibodies were used as loading control. Quantification of O-GlcNAc levels in prefrontal cortex (Q), hippocampus, striatum (S) and cerebellum (T) from the Western blot in panel O and P.).


Supplementary Figure S14. OGT^{C921Y} mice show altered *Ogt/Oga* mRNA ratio levels in brain regions.

Significance is shown as * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ ($n = 3$ per group). Student t test was used for statistics. Quantification of Oga (A), Ogt (B) and Ogt/Oga ratio (C) mRNA levels in prefrontal cortex Quantification of Oga (D), Ogt (E) and Ogt/Oga ratio (F) mRNA levels in hippocampus. Quantification of Oga (G), Ogt (H) and Ogt/Oga ratio (I) mRNA levels in striatum. Quantification of Oga (J), Ogt (K) and Ogt/Oga ratio (L) mRNA levels in cerebellum.



Supplementary Figure S15. Perturbed proteomic signatures in the prefrontal cortex corresponded to distinct Monarch enrichment profiles in the OGT^{C921Y} mice.

(A) Up-regulated human phenotypes enrichment in the Monarch database. (B) Down-regulated human phenotypes enrichment in the Monarch database.



Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD074604. Behaviour and histology source data files have been deposited at <https://osf.io/b5zqr/overview>.

Acknowledgements

The authors would like to thank Trine Mikkelesen (JRN lab) for the assistance with the histology workflow, Kristian Graff (Department of Molecular Biology and Genetics, Aarhus University), for the assistance in behaviour equipment build-up and Kamilla Zahll Hornbek (Department of Biomedicine, Aarhus University) for breeding and animal care assistance.

Additional information

Funding

This work was funded by a Wellcome Trust Investigator Award (110061), a Novo Nordisk Fonden Laureate award (NNF21OC0065969) and a Villum Fonden Investigator (00054496) to D.M.F.v.A. Supported in part by the Danish Research Institute of Translational Neuroscience – DANDRITE of the Nordic-EMBL Partnership for Molecular Medicine and Lundbeckfonden. The Novo Nordisk Foundation is gratefully acknowledged for funding the Scanco μ CT equipment as a part of the Aarhus X-ray Imaging Alliance (AXIA).

Author contributions

F.A. and D.M.F.v.A conceived the study; F.A., I.F., C.S.S., S.T.B., K.S.C., B.A., J.S.T. performed experiments; C.S. performed mass spectrometry; F.A., I.F., C.S.S., K.S.C., I.E.A., S.F.E., B.H. analysed data; A.J. and J.R.N. performed histology and image analyses; O.G.S.Q. provided illustration used in Figure 8; F.A., A.J. and D.M.F.v.A. interpreted the data and wrote the manuscript with input from all authors.

Funding

Funder	Grant reference number	Author
Wellcome Trust (WT)	https://doi.org/10.35802/110061	Daan MF van Aalten
Novo Nordisk Fonden (NNF)	NNF21OC0065969	Daan MF van Aalten
Villum Fonden (Villum Foundation)	00054496	Daan MF van Aalten

Author ORCID iDs

Christian Stald Skoven: <https://orcid.org/0000-0002-0639-6718>
Jesper Skovhus Thomsen: <https://orcid.org/0000-0001-9386-6679>
Daan MF van Aalten: <https://orcid.org/0000-0002-1499-6908>

References

1. **Tassé MJ**, Luckasson R, Schalock RL (2016) The Relation Between Intellectual Functioning and Adaptive Behavior in the Diagnosis of Intellectual Disability. *Intellect Dev Disabil* **54**:381-390 <https://doi.org/10.1352/1934-9556-54.6.381> | PubMed
2. **Barkovich AJ**, Guerrini R, Kuzniecky RI, Jackson GD, Dobyns WB (2012) A developmental and genetic classification for malformations of cortical development: update 2012. *Brain J Neurol* **135**:1348-1369 <https://doi.org/10.1093/brain/aws019> | PubMed

3. Kurabayashi N, Fujii K, Otake Y, et al. (2023) Neocortical neuronal production and maturation defects in the TcMAC21 mouse model of Down syndrome. *iScience* **26**:108379 <https://doi.org/10.1016/j.isci.2023.108379> | PubMed
4. Torres CR, Hart GW (1984) Topography and polypeptide distribution of terminal N-acetylglucosamine residues on the surfaces of intact lymphocytes. Evidence for O-linked GlcNAc. *J Biol Chem* **259**:3308-3317 [https://doi.org/10.1016/s0021-9258\(17\)43295-9](https://doi.org/10.1016/s0021-9258(17)43295-9) | PubMed
5. Wulff-Fuentes E, Berendt RR, Massman L, et al. (2021) The human O-GlcNAc database and meta-analysis. *Sci Data* **8**:25 <https://doi.org/10.1038/s41597-021-00810-4> | PubMed
6. Gao Y, Wells L, Comer FI, Parker GJ, Hart GW (2001) Dynamic O-glycosylation of nuclear and cytosolic proteins: cloning and characterization of a neutral, cytosolic beta-N-acetylglucosaminidase from human brain. *J Biol Chem* **276**:9838-9845 <https://doi.org/10.1074/jbc.M010420200> | PubMed
7. Fujiki R, Hashiba W, Sekine H, et al. (2011) GlcNAcylation of histone H2B facilitates its monoubiquitination. *Nature* **480**:557-560 <https://doi.org/10.1038/nature10656> | PubMed
8. Park SY, Kim HS, Kim NH, et al. (2010) Snail1 is stabilized by O-GlcNAc modification in hyperglycaemic condition. *EMBO J* **29**:3787-3796 <https://doi.org/10.1038/emboj.2010.254> | PubMed
9. Levine ZG, Fan C, Melicher MS, Orman M, Benjamin T, Walker S (2018) O-GlcNAc Transferase Recognizes Protein Substrates Using an Asparagine Ladder in the Tetratricopeptide Repeat (TPR) Superhelix. *J Am Chem Soc* **140**:3510-3513 <https://doi.org/10.1021/jacs.7b13546> | PubMed
10. Meek RW, Blaza JN, Busmann JA, Alteen MG, Vocadlo DJ, Davies GJ (2021) Cryo-EM structure provides insights into the dimer arrangement of the O-linked β -N-acetylglucosamine transferase OGT. *Nat Commun* **12**:6508 <https://doi.org/10.1038/s41467-021-26796-6> | PubMed
11. Fujiki R, Hashiba W, Sekine H, et al. (2011) GlcNAcylation of histone H2B facilitates its monoubiquitination. *Nature* **480**:557-560 <https://doi.org/10.1038/nature10656> | PubMed
12. Constable S, Lim JM, Vaidyanathan K, Wells L (2017) O-GlcNAc transferase regulates transcriptional activity of human Oct4. *Glycobiology* **27**:927-937 <https://doi.org/10.1093/glycob/cwx055> | PubMed
13. Dikic I, Wakatsuki S, Walters KJ (2009) Ubiquitin-binding domains - from structures to functions. *Nat Rev Mol Cell Biol* **10**:659-671 <https://doi.org/10.1038/nrm2767> | PubMed
14. Capotosti F, Guernier S, Lammers F, et al. (2011) O-GlcNAc Transferase Catalyzes Site-Specific Proteolysis of HCF-1. *Cell* **144**:376-388 <https://doi.org/10.1016/j.cell.2010.12.030> | PubMed
15. Lazarus MB, Jiang J, Kapuria V, et al. (2013) HCF-1 Is Cleaved in the Active Site of O-GlcNAc Transferase. *Science* **342**:1235-1239 <https://doi.org/10.1126/science.1243990> | PubMed
16. O'Donnell N, Zachara NE, Hart GW, Marth JD (2004) Ogt-Dependent X-Chromosome-Linked Protein Glycosylation Is a Requisite Modification in Somatic Cell Function and Embryo Viability. *Mol Cell Biol* **24**:1680 <https://doi.org/10.1128/MCB.24.4.1680-1690.2004> | PubMed
17. Lee BE, Kim HY, Kim HJ, et al. (2020) O-GlcNAcylation regulates dopamine neuron function, survival and degeneration in Parkinson disease. *Brain J Neurol* **143**:3699-3716 <https://doi.org/10.1093/brain/awaa320> | PubMed
18. Su C, Schwarz TL (2017) O-GlcNAc Transferase Is Essential for Sensory Neuron Survival and Maintenance. *J Neurosci* **37**:2125 <https://doi.org/10.1523/JNEUROSCI.3384-16.2017> | PubMed
19. Cole RN, Hart GW (2001) Cytosolic O-glycosylation is abundant in nerve terminals. *J Neurochem* **79**:1080-1089 <https://doi.org/10.1046/j.1471-4159.2001.00655.x> | PubMed
20. Lagerl f O, Hart GW, Haganir RL (2017) O-GlcNAc transferase regulates excitatory synapse maturity. *Proc Natl Acad Sci U S A* **114**:1684-1689 <https://doi.org/10.1073/PNAS.1621367114> | PubMed
21. Vosseller K, Trinidad JC, Chalkley RJ, et al. (2006) O-linked N-acetylglucosamine proteomics of postsynaptic density preparations using lectin weak affinity chromatography and mass spectrometry. *Mol Cell Proteomics MCP* **5**:923-934 <https://doi.org/10.1074/mcp.T500040-MCP200> | PubMed

22. **Trinidad JC**, Barkan DT, Gullledge BF, et al. (2012) Global identification and characterization of both O-GlcNAcylation and phosphorylation at the murine synapse. *Mol Cell Proteomics MCP* **11**:215-229 <https://doi.org/10.1074/mcp.O112.018366> | PubMed
23. **Vaidyanathan K**, Niranjana T, Selvan N, et al. (2017) Identification and characterization of a missense mutation in the O-linked β -N-acetylglucosamine (O-GlcNAc) transferase gene that segregates with X-linked intellectual disability. *J Biol Chem* **292**:8948-8963 <https://doi.org/10.1074/jbc.M116.771030> | PubMed
24. **Willems AP**, Gundogdu M, Kempers MJE, et al. (2017) Mutations in N-acetylglucosamine (O-GlcNAc) transferase in patients with X-linked intellectual disability. *J Biol Chem* **292**:12621-12631 <https://doi.org/10.1074/jbc.M117.790097> | PubMed
25. **Vaidyanathan K**, Niranjana T, Selvan N, et al. (2017) Identification and characterization of a missense mutation in the O-linked β -N-acetylglucosamine (O-GlcNAc) transferase gene that segregates with X-linked intellectual disability. *J Biol Chem* **292**:8948-8963 <https://doi.org/10.1074/jbc.M116.771030> | PubMed
26. **Willems AP**, Gundogdu M, Kempers MJE, et al. (2017) Mutations in N-acetylglucosamine (O-GlcNAc) transferase in patients with X-linked intellectual disability. *J Biol Chem* **292**:12621-12631 <https://doi.org/10.1074/jbc.M117.790097> | PubMed
27. **Pravata VM**, Muha V, Gundogdu M, et al. (2019) Catalytic deficiency of O-GlcNAc transferase leads to X-linked intellectual disability. *Proc Natl Acad Sci* **116**:14961-14970 <https://doi.org/10.1073/pnas.1900065116> | PubMed
28. **Pravata VM**, Gundogdu M, Bartual SG, et al. (2020) A missense mutation in the catalytic domain of O-GlcNAc transferase links perturbations in protein O-GlcNAcylation to X-linked intellectual disability. *FEBS Lett* **594**:717-727 <https://doi.org/10.1002/1873-3468.13640> | PubMed
29. **Pravata VM**, Omelková M, Stavridis MP, et al. (2020) An intellectual disability syndrome with single-nucleotide variants in O-GlcNAc transferase. *Eur J Hum Genet* **28**:706-714 <https://doi.org/10.1038/s41431-020-0589-9> | PubMed
30. **Taylor DC**, Falconer MA, Bruton CJ, Corsellis JAN (1971) Focal dysplasia of the cerebral cortex in epilepsy. *J Neurol Neurosurg Psychiatry* **34**:369-387 <https://doi.org/10.1136/jnnp.34.4.369> | PubMed
31. **Casanova MF**, El-Baz AS, Kamat SS, et al. (2013) Focal cortical dysplasias in autism spectrum disorders. *Acta Neuropathol Commun* **1** <https://doi.org/10.1186/2051-5960-1-67> | PubMed
32. **Omelková M**, Fenger CD, Murray M, et al. (2023) An O-GlcNAc transferase pathogenic variant linked to intellectual disability affects pluripotent stem cell self-renewal. *Dis Model Mech* **16**:dmm049132 <https://doi.org/10.1242/dmm.049132> | PubMed
33. **Selvan N**, George S, Serajee FJ, et al. (2018) O-GlcNAc transferase missense mutations linked to X-linked intellectual disability deregulate genes involved in cell fate determination and signaling. *J Biol Chem* **293**:10810-10824 <https://doi.org/10.1074/jbc.RA118.002583> | PubMed
34. **Shafi R**, Iyer SPN, Ellies LG, et al. (2000) The O-GlcNAc transferase gene resides on the X chromosome and is essential for embryonic stem cell viability and mouse ontogeny. *Proc Natl Acad Sci* **97**:5735-5739 <https://doi.org/10.1073/pnas.100471497> | PubMed
35. **Omelková M**, Fenger CD, Murray M, et al. (2023) An O-GlcNAc transferase pathogenic variant linked to intellectual disability affects pluripotent stem cell self-renewal. *Dis Model Mech* **16**:dmm049132 <https://doi.org/10.1242/dmm.049132> | PubMed
36. **Czajewski I**, Swain B, Xu J, McDowall L, Ferenbach AT, Van Aalten DM (2024) Rescuable sleep and synaptogenesis phenotypes in a Drosophila model of O-GlcNAc transferase intellectual disability. *eLife* **13**:e90376 <https://doi.org/10.7554/eLife.90376> | PubMed
37. **Authier F**, Ondruskova N, Ferenbach AT, McNeilly AD, van Aalten DMF (2024) Neurodevelopmental defects in a mouse model of O-GlcNAc transferase intellectual disability. *Dis Model Mech* **17**:dmm050671 <https://doi.org/10.1242/dmm.050671> | PubMed

38. Piilgaard L, Rose L, Justinussen JL, et al. (2023) Non-invasive detection of narcolepsy type I phenotypical features and disease progression by continuous home-cage monitoring of activity in two mouse models: the HCRT-KO and DTA model. *Sleep* **46**:zsad144 <https://doi.org/10.1093/sleep/zsad144> | PubMed
39. Schwarzmaier SM, Knarr MRO, Hu S, Ertürk A, Hellal F, Plesnila N (2022) Perfusion pressure determines vascular integrity and histomorphological quality following perfusion fixation of the brain. *J Neurosci Methods* **372**:109493 <https://doi.org/10.1016/j.jneumeth.2022.109493> | PubMed
40. Shepherd TM, Thelwall PE, Stanis GJ, Blackband SJ (2009) Aldehyde Fixative Solutions Alter the Water Relaxation and Diffusion Properties of Nervous Tissue. *Magn Reson Med Off J Soc Magn Reson Med Soc Magn Reson Med* **62**:26-34 <https://doi.org/10.1002/mrm.21977> | PubMed
41. Khan AR, Chuhutin A, Wiborg O, et al. (2016) Biophysical modeling of high field diffusion MRI demonstrates micro-structural aberration in chronic mild stress rat brain. *NeuroImage* **142**:421-430 <https://doi.org/10.1016/j.neuroimage.2016.07.001> | PubMed
42. Chuhutin A, Hansen B, Włodarczyk A, Owens T, Shemesh N, Jespersen SN (2020) Diffusion Kurtosis Imaging maps neural damage in the EAE model of multiple sclerosis. *NeuroImage* **208** <https://doi.org/10.1016/j.neuroimage.2019.116406> | PubMed
43. Bay V, Kjølbj BF, Iversen NK, et al. (2018) Stroke infarct volume estimation in fixed tissue: Comparison of diffusion kurtosis imaging to diffusion weighted imaging and histology in a rodent MCAO model. *PLoS One* **13**:e0196161 <https://doi.org/10.1371/journal.pone.0196161> | PubMed
44. Jespersen SN, Bjarkam CR, Nyengaard JR, et al. (2010) Neurite density from magnetic resonance diffusion measurements at ultrahigh field: comparison with light microscopy and electron microscopy. *NeuroImage* **49**:205-216 <https://doi.org/10.1016/j.neuroimage.2009.08.053> | PubMed
45. Vestergaard-Poulsen P, Wegener G, Hansen B, et al. (2011) Diffusion-weighted MRI and quantitative biophysical modeling of hippocampal neurite loss in chronic stress. *PLoS One* **6**:e20653 <https://doi.org/10.1371/journal.pone.0020653> | PubMed
46. Knopper RW, Skoven CS, Eskildsen SF, Østergaard L, Hansen B (2024) The effects of locus coeruleus ablation on mouse brain volume and microstructure evaluated by high-field MRI. *Front Cell Neurosci* **18** <https://doi.org/10.3389/fncel.2024.1498133> | PubMed
47. Hansen B, Khan AR, Shemesh N, et al. (2017) White matter biomarkers from fast protocols using axially symmetric diffusion kurtosis imaging. *NMR Biomed* **30**:e3741 <https://doi.org/10.1002/nbm.3741> | PubMed
48. Ardalan M, Chumak T, Quist A, et al. (2022) Reelin cells and sex-dependent synaptopathology in autism following postnatal immune activation. *Br J Pharmacol* **179**:4400-4422 <https://doi.org/10.1111/bph.15859> | PubMed
49. Ma D, Cardoso MJ, Modat M, et al. (2014) Automatic structural parcellation of mouse brain MRI using multi-atlas label fusion. *PLoS One* **9**:e86576 <https://doi.org/10.1371/journal.pone.0086576> | PubMed
50. Ma D, Holmes HE, Cardoso MJ, et al. (2019) Study the Longitudinal in vivo and Cross-Sectional ex vivo Brain Volume Difference for Disease Progression and Treatment Effect on Mouse Model of Tauopathy Using Automated MRI Structural Parcellation. *Front Neurosci* **13** <https://doi.org/10.3389/fnins.2019.00011> | PubMed
51. Ma Y, Hof PR, Grant SC, et al. (2005) A three-dimensional digital atlas database of the adult C57BL/6J mouse brain by magnetic resonance microscopy. *Neuroscience* **135**:1203-1215 <https://doi.org/10.1016/j.neuroscience.2005.07.014> | PubMed
52. Ma Y, Smith D, Hof PR, et al. (2008) In Vivo 3D Digital Atlas Database of the Adult C57BL/6J Mouse Brain by Magnetic Resonance Microscopy. *Front Neuroanat* **2** <https://doi.org/10.3389/neuro.05.001.2008> | PubMed
53. Sled JG, Zijdenbos AP, Evans AC (1998) A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE Trans Med Imaging* **17**:87-97 <https://doi.org/10.1109/42.668698> | PubMed

54. **Coupe P**, Yger P, Prima S, Hellier P, Kervrann C, Barillot C (2008) An optimized blockwise nonlocal means denoising filter for 3-D magnetic resonance images. *IEEE Trans Med Imaging* **27**:425-441 <https://doi.org/10.1109/TMI.2007.906087> | PubMed
55. **Dorr AE**, Lerch JP, Spring S, Kabani N, Henkelman RM (2008) High resolution three-dimensional brain atlas using an average magnetic resonance image of 40 adult C57Bl/6J mice. *NeuroImage* **42**:60-69 <https://doi.org/10.1016/j.neuroimage.2008.03.037> | PubMed
56. **Collins DL**, Neelin P, Peters TM, Evans AC (1994) Automatic 3D intersubject registration of MR volumetric data in standardized Talairach space. *J Comput Assist Tomogr* **18**:192-205 <https://doi.org/10.1097/00004728-199403000-00005> | PubMed
57. **Avants BB**, Tustison NJ, Stauffer M, Song G, Wu B, Gee JC (2014) The Insight ToolKit image registration framework. *Front Neuroinformatics* **8** <https://doi.org/10.3389/fninf.2014.00044> | PubMed
58. **Benjamini Y**, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol* **57**:289-300 <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
59. **Lerch JP**, Carroll JB, Dorr A, et al. (2008) Cortical thickness measured from MRI in the YAC128 mouse model of Huntington's disease. *NeuroImage* **41**:243-251 <https://doi.org/10.1016/j.neuroimage.2008.02.019> | PubMed
60. **Authier F**, Ondruskova N, Ferenbach AT, McNeilly AD, van Aalten DMF (2024) Neurodevelopmental defects in a mouse model of O-GlcNAc transferase intellectual disability. *Dis Model Mech* **17**:dmm050671 <https://doi.org/10.1242/dmm.050671> | PubMed
61. **Robson K** (2005) Neuropathology Techniques. *Neuropathol Appl Neurobiol* **31**:204-205 <https://doi.org/10.1111/j.1365-2990.2005.00585.x>
62. **Bankhead P**, Loughrey MB, Fernández JA, et al. (2017) QuPath: Open source software for digital pathology image analysis. *Sci Rep* **7**:16878 <https://doi.org/10.1038/s41598-017-17204-5> | PubMed
63. **Pravata VM**, Omelková M, Stavridis MP, et al. (2020) An intellectual disability syndrome with single-nucleotide variants in O-GlcNAc transferase. *Eur J Hum Genet. Published online* <https://doi.org/10.1038/s41431-020-0589-9> | PubMed
64. **Adasooriya D**, Kyeong M, Cho SW (2023) Sphenoid Bone Determines the Curvature of the Cranial Vault in Postnatal Skull Development in C57BL/6 Mice. *J Bone Metab* **30**:93-101 <https://doi.org/10.11005/jbm.2023.30.1.93> | PubMed
65. **Heulens I**, Suttie M, Postnov A, et al. (2013) Craniofacial characteristics of fragile X syndrome in mouse and man. *Eur J Hum Genet* **21**:816-823 <https://doi.org/10.1038/ejhg.2012.265> | PubMed
66. **Husain TS**, Moore JC, Huston LA, et al. (2024) Neurocranial growth in the OIM mouse model of osteogenesis imperfecta. *Anat Rec* **307**:581-591 <https://doi.org/10.1002/ar.25307> | PubMed
67. **Fenlon LR**, Liu S, Gobius I, et al. (2015) Formation of functional areas in the cerebral cortex is disrupted in a mouse model of autism spectrum disorder. *Neural Develop* **10** <https://doi.org/10.1186/s13064-015-0033-y> | PubMed
68. **Donovan APA**, Rosko L, Ellegood J, et al. (2023) Pervasive cortical and white matter anomalies in a mouse model for CHARGE syndrome. *J Anat* **243**:51-65 <https://doi.org/10.1111/joa.13856> | PubMed
69. **Allemang-Grand R**, Ellegood J, Spencer Noakes L, et al. (2017) Neuroanatomy in mouse models of Rett syndrome is related to the severity of Mecp2 mutation and behavioral phenotypes. *Mol Autism* **8**:32 <https://doi.org/10.1186/s13229-017-0138-8> | PubMed
70. **Jensen JH**, Helpert JA, Ramani A, Lu H, Kaczynski K (2005) Diffusional kurtosis imaging: the quantification of non-gaussian water diffusion by means of magnetic resonance imaging. *Magn Reson Med* **53**:1432-1440 <https://doi.org/10.1002/mrm.20508> | PubMed
71. **Hansen B**, Lund TE, Sangill R, Jespersen SN (2013) Experimentally and computationally fast method for estimation of a mean kurtosis. *Magn Reson Med* **69**:1754-1760 <https://doi.org/10.1002/mrm.24743> | PubMed

72. Hansen B, Lund TE, Sangill R, Stubbe E, Finsterbusch J, Jespersen SN (2016) Experimental considerations for fast kurtosis imaging. *Magn Reson Med* **76**:1455-1468 <https://doi.org/10.1002/mrm.26055> | PubMed
73. Hansen B, Jespersen SN (2017) Recent Developments in Fast Kurtosis Imaging. *Front Phys* **5** <https://doi.org/10.3389/fphy.2017.00040>
74. Qvist P, Eskildsen SF, Hansen B, et al. (2018) Brain volumetric alterations accompanied with loss of striatal medium-sized spiny neurons and cortical parvalbumin expressing interneurons in Brd1+/- mice. *Sci Rep* **8**:16486 <https://doi.org/10.1038/s41598-018-34729-5> | PubMed
75. Joseph JV, Blaavand MS, Cai H, et al. (2023) STING activation counters glioblastoma by vascular alteration and immune surveillance. *Cancer Lett* **579** <https://doi.org/10.1016/j.canlet.2023.216480> | PubMed
76. Lindhardt TB, Skoven CS, Bordoni L, Østergaard L, Liang Z, Hansen B (2024) Anesthesia-related brain microstructure modulations detected by diffusion magnetic resonance imaging. *NMR Biomed* **37**:e5033 <https://doi.org/10.1002/nbm.5033> | PubMed
77. Vukovic N, Hansen B, Lund TE, Jespersen S, Shtyrov Y (2021) Rapid microstructural plasticity in the cortical semantic network following a short language learning session. *PLoS Biol* **19**:e3001290 <https://doi.org/10.1371/journal.pbio.3001290> | PubMed
78. Tietze A, Hansen MB, Østergaard L, et al. (2015) Mean Diffusional Kurtosis in Patients with Glioma: Initial Results with a Fast Imaging Method in a Clinical Setting. *AJNR Am J Neuroradiol* **36**:1472-1478 <https://doi.org/10.3174/ajnr.A4311> | PubMed
79. Lohmeier J, Radbruch H, Brenner W, et al. (2024) Detection of recurrent high-grade glioma using microstructure characteristics of distinct metabolic compartments in a multimodal and integrative 18F-FET PET/fast-DKI approach. *Eur Radiol* **34**:2487-2499 <https://doi.org/10.1007/s00330-023-10141-0> | PubMed
80. Thaler C, Kyselyova AA, Faizy TD, et al. (2021) Heterogeneity of multiple sclerosis lesions in fast diffusional kurtosis imaging. *PloS One* **16**:e0245844 <https://doi.org/10.1371/journal.pone.0245844> | PubMed
81. Fauser S, Huppertz HJ, Bast T, et al. (2006) Clinical characteristics in focal cortical dysplasia: a retrospective evaluation in a series of 120 patients. *Brain J Neurol* **129**:1907-1916 <https://doi.org/10.1093/brain/awl133> | PubMed
82. Tassi L, Colombo N, Garbelli R, et al. (2002) Focal cortical dysplasia: neuropathological subtypes, EEG, neuroimaging and surgical outcome. *Brain J Neurol* **125**:1719-1732 <https://doi.org/10.1093/brain/awf175> | PubMed
83. Blümcke I, Thom M, Aronica E, et al. (2011) The clinicopathologic spectrum of focal cortical dysplasias: a consensus classification proposed by an ad hoc Task Force of the ILAE Diagnostic Methods Commission. *Epilepsia* **52**:158-174 <https://doi.org/10.1111/j.1528-1167.2010.02777.x> | PubMed
84. Paprocka J, Jezela-Stanek A, Tyłki-Szymańska A, Grunewald S (2021) Congenital Disorders of Glycosylation from a Neurological Perspective. *Brain Sci* **11**:88 <https://doi.org/10.3390/brainsci11010088> | PubMed
85. Squier W, Jansen A (2014) Polymicrogyria: pathology, fetal origins and mechanisms. *Acta Neuropathol Commun* **2** <https://doi.org/10.1186/s40478-014-0080-3> | PubMed
86. Putman TE, Schaper K, Matentzoglou N, et al. (2024) The Monarch Initiative in 2024: an analytic platform integrating phenotypes, genes and diseases across species. *Nucleic Acids Res* **52**:D938-D949 <https://doi.org/10.1093/nar/gkad1082> | PubMed
87. Mayfield JM, Hitefield NL, Czajewski I, et al. (2024) O-GlcNAc transferase congenital disorder of glycosylation (OGT-CDG): Potential mechanistic targets revealed by evaluating the OGT interactome. *J Biol Chem* **300**:107599 <https://doi.org/10.1016/j.jbc.2024.107599> | PubMed

88. Yuan H, Mitchell CW, Ferenbach AT, et al. (2024) Exploiting O-GlcNAc dyshomeostasis to screen O-GlcNAc transferase intellectual disability variants. *Stem Cell Rep* **19**:102380 <https://doi.org/10.1016/j.stemcr.2024.11.010> | PubMed
89. Pravata VM, Gundogdu M, Bartual SG, et al. (2019) A missense mutation in the catalytic domain of O-GlcNAc transferase links perturbations in protein O-GlcNAcylation to X-linked intellectual disability. *FEBS Lett* <https://doi.org/10.1002/1873-3468.13640> | PubMed
90. Ding Q, Sethna F, Wang H (2014) Behavioral analysis of male and female Fmr1 knockout mice on C57BL/6 background. *Behav Brain Res* **0** <https://doi.org/10.1016/j.BBR.2014.05.046> | PubMed
91. Peier AM, McIlwain KL, Kenneson A, Warren ST, Paylor R, Nelson DL (2000) (Over)correction of FMR1 deficiency with YAC transgenics: behavioral and physical features. *Hum Mol Genet* **9**:1145-1159 <https://doi.org/10.1093/HMG/9.8.1145> | PubMed
92. Huynh TN, Shah M, Koo SY, Faraud KS, Santini E, Klann E (2015) eIF4E/Fmr1 Double Mutant Mice Display Cognitive Impairment in Addition to ASD-like Behaviors. *Neurobiol Dis* **83** <https://doi.org/10.1016/j.NBD.2015.08.016> | PubMed
93. Gao Y, Aljazi MB, He J (2022) Kdm6b Haploinsufficiency Causes ASD/ADHD-Like Behavioral Deficits in Mice. *Front Behav Neurosci* **16** <https://doi.org/10.3389/FNBEH.2022.905783> | PubMed
94. Fencikova M, Muha V, Mariappa D, et al. (2022) Intellectual disability-associated disruption of O-GlcNAc cycling impairs habituation learning in Drosophila. *PLoS Genet* **18**:e1010159 <https://doi.org/10.1371/JOURNAL.PGEN.1010159> | PubMed
95. Rankin CH, Abrams T, Barry RJ, et al. (2009) Habituation Revisited: An Updated and Revised Description of the Behavioral Characteristics of Habituation. *Neurobiol Learn Mem* **92**:135 <https://doi.org/10.1016/j.NLM.2008.09.012> | PubMed
96. Spencer MD, Moorhead TWJ, Lymer GKS, et al. (2006) Structural correlates of intellectual impairment and autistic features in adolescents. *NeuroImage* **33**:1136-1144 <https://doi.org/10.1016/j.neuroimage.2006.08.011> | PubMed
97. Kolk SM, Rakic P (2022) Development of prefrontal cortex. *Neuropsychopharmacology* **47**:41-57 <https://doi.org/10.1038/s41386-021-01137-9> | PubMed
98. Russell VA, Sagvolden T, Johansen EB (2005) Animal models of attention-deficit hyperactivity disorder. *Behav Brain Funct* **1** <https://doi.org/10.1186/1744-9081-1-9> | PubMed
99. Schiller S, Rosewich H, Grünwald S, Gärtner J (2020) Inborn errors of metabolism leading to neuronal migration defects. *J Inherit Metab Dis* **43**:145-155 <https://doi.org/10.1002/jimd.12194> | PubMed
100. Reynolds CD, Nolan SO, Smith GD, et al. (2020) Increased Expression of Fragile X Mental Retardation Protein in Malformative Lesions of Patients with Focal Cortical Dysplasia. *Neuroreport* **31**:1036-1041 <https://doi.org/10.1097/WNR.0000000000001517> | PubMed
101. Raymond AA, Fish DR, Sisodiya SM, Alsanjari N, Stevens JM, Shorvon SD (1995) Abnormalities of gyration, heterotopias, tuberous sclerosis, focal cortical dysplasia, microdysgenesis, dysembryoplastic neuroepithelial tumour and dysgenesis of the archicortex in epilepsy. Clinical, EEG and neuroimaging features in 100 adult patients. *Brain J Neurol* **118**:629-660 <https://doi.org/10.1093/brain/118.3.629> | PubMed
102. Barkovich AJ, Kuzniecky RI, Dobyns WB, Jackson GD, Becker LE, Evrard P (1996) A classification scheme for malformations of cortical development. *Neuropediatrics* **27**:59-63 <https://doi.org/10.1055/s-2007-973750> | PubMed
103. Barkovich AJ, Kuzniecky RI, Jackson GD, Guerrini R, Dobyns WB (2005) A developmental and genetic classification for malformations of cortical development. *Neurology* **65**:1873-1887 <https://doi.org/10.1212/01.wnl.0000183747.05269.2d> | PubMed
104. Judkins AR, Martinez D, Ferreira P, Dobyns WB, Golden JA (2011) Polymicrogyria includes fusion of the molecular layer and decreased neuronal populations but normal cortical laminar organization. *J Neuropathol Exp Neurol* **70**:438-443 <https://doi.org/10.1097/NEN.0b013e31821ccf1c> | PubMed

105. Leech R, Sharp DJ (2014) The role of the posterior cingulate cortex in cognition and disease. *Brain* **137**:12-32 <https://doi.org/10.1093/brain/awt162> | PubMed
106. Kolling N, Wittmann MK, Behrens TEJ, Boorman ED, Mars RB, Rushworth MFS (2016) Value, search, persistence and model updating in anterior cingulate cortex. *Nat Neurosci* **19**:1280-1285 <https://doi.org/10.1038/nn.4382> | PubMed
107. Goodkind M, Eickhoff SB, Oathes DJ, et al. (2015) Identification of a common neurobiological substrate for mental illness. *JAMA Psychiatry* **72**:305-315 <https://doi.org/10.1001/jamapsychiatry.2014.2206> | PubMed
108. Opel N, Goltermann J, Hermesdorf M, Berger K, Baune BT, Dannlowski U (2020) Cross-Disorder Analysis of Brain Structural Abnormalities in Six Major Psychiatric Disorders: A Secondary Analysis of Mega- and Meta-analytical Findings From the ENIGMA Consortium. *Biol Psychiatry* **88**:678-686 <https://doi.org/10.1016/j.biopsych.2020.04.027> | PubMed
109. Choi JY, Rha D, Park ES (2016) The Effects of the Severity of Periventricular Leukomalacia on the Neuropsychological Outcomes of Preterm Children. *J Child Neurol* **31**:603-612 <https://doi.org/10.1177/0883073815604229> | PubMed
110. Barnea-Goraly N, Eliez S, Hedeus M, et al. (2003) White matter tract alterations in fragile X syndrome: Preliminary evidence from diffusion tensor imaging. *Am J Med Genet B Neuropsychiatr Genet* **118B**:81-88 <https://doi.org/10.1002/ajmg.b.10035> | PubMed
111. Cascio C, Styner M, Smith RG, et al. (2006) Reduced relationship to cortical white matter volume revealed by tractography-based segmentation of the corpus callosum in young children with developmental delay. *Am J Psychiatry* **163**:2157-2163 <https://doi.org/10.1176/ajp.2006.163.12.2157> | PubMed
112. Pujol J, López-Sala A, Sebastián-Gallés N, et al. (2004) Delayed myelination in children with developmental delay detected by volumetric MRI. *NeuroImage* **22**:897-903 <https://doi.org/10.1016/j.neuroimage.2004.01.029> | PubMed
113. Ishii K, et al. (2015) Neuronal Heterotopias Affect the Activities of Distant Brain Areas and Lead to Behavioral Deficits. *J Neurosci Off J Soc Neurosci* **35**:12432-12445 <https://doi.org/10.1523/JNEUROSCI.3648-14.2015> | PubMed
114. Doherty K, Frazier SB, Clark M, et al. (2019) A closer look at ARSA activity in a patient with metachromatic leukodystrophy. *Mol Genet Metab Rep* **19**:100460 <https://doi.org/10.1016/j.ymgmr.2019.100460> | PubMed
115. Coulter ME, Dorobantu CM, Lodewijk GA, et al. (2018) The ESCRT-III Protein CHMP1A Mediates Secretion of Sonic Hedgehog on a Distinctive Subtype of Extracellular Vesicles. *Cell Rep* **24**:973-986.e8. <https://doi.org/10.1016/j.celrep.2018.06.100> | PubMed
116. Chan DL, Rudinger-Thirion J, Frugier M, et al. (2022) A case of QARS1 associated epileptic encephalopathy and review of epilepsy in aminoacyl-tRNA synthetase disorders. *Brain Dev* **44**:142-147 <https://doi.org/10.1016/j.braindev.2021.10.009> | PubMed
117. He T, Sun H, Xu B, et al. (2023) Novel bi-allelic variants of CHMP1A contribute to pontocerebellar hypoplasia type 8: additional clinical and genetic evidence. *Front Neurol* **14**:1228218 <https://doi.org/10.3389/fneur.2023.1228218> | PubMed
118. Feng Y, Wei ZH, Liu C, et al. (2022) Genetic variations in GABA metabolism and epilepsy. *Seizure Eur J Epilepsy* **101**:22-29 <https://doi.org/10.1016/j.seizure.2022.07.007> | PubMed
119. Lv Y, Ji L, Dai H, et al. (2024) Identification of key regulatory genes involved in myelination after spinal cord injury by GSEA analysis. *Exp Neurol* **382**:114966 <https://doi.org/10.1016/j.expneurol.2024.114966> | PubMed
120. Gupta A, Colmenero I, Ragge NK, et al. (2016) Compound heterozygous RMND1 gene variants associated with chronic kidney disease, dilated cardiomyopathy and neurological involvement: a case report. *BMC Res Notes* **9** <https://doi.org/10.1186/s13104-016-2131-2> | PubMed

121. Cesani M, Lorioli L, Grossi S, et al. (2016) Mutation Update of ARSA and PSAP Genes Causing Metachromatic Leukodystrophy. *Hum Mutat* **37**:16-27 <https://doi.org/10.1002/humu.22919> | PubMed
122. Gürsoy S, Hazan F, Öztürk T, Çolak R, Çalkavur Ş (2021) Evaluation of Sporadic and Familial Cases with Craniofrontonasal Syndrome: A Wide Clinical Spectrum and Identification of a Novel EFN1 Gene Mutation. *Mol Syndromol* **12**:269-278 <https://doi.org/10.1159/000515697> | PubMed
123. Abdollahi MR, Morrison E, Sirey T, et al. (2009) Mutation of the Variant α -Tubulin TUBA8 Results in Polymicrogyria with Optic Nerve Hypoplasia. *Am J Hum Genet* **85**:737-744 <https://doi.org/10.1016/j.ajhg.2009.10.007> | PubMed
124. Duncan AR, Polovitskaya MM, Gaitán-Peñas H, et al. (2021) Unique variants in CLCN3, encoding an endosomal anion/proton exchanger, underlie a spectrum of neurodevelopmental disorders. *Am J Hum Genet* **108**:1450-1465 <https://doi.org/10.1016/j.ajhg.2021.06.003> | PubMed
125. Tomas-Roca L, Tsaalbi-Shtylik A, Jansen JG, et al. (2015) De novo mutations in PLXND1 and REV3L cause Möbius syndrome. *Nat Commun* **6**:7199 <https://doi.org/10.1038/ncomms8199> | PubMed
126. Govek EE, Newey SE, Van Aelst L (2005) The role of the Rho GTPases in neuronal development. *Genes Dev* **19**:1-49 <https://doi.org/10.1101/gad.1256405> | PubMed
127. Evans MK, Matsui Y, Xu B, et al. (2020) Ybx1 fine-tunes PRC2 activities to control embryonic brain development. *Nat Commun* **11**:4060 <https://doi.org/10.1038/s41467-020-17878-y> | PubMed

Peer reviews

Reviewer #1 (Public review):

This study established a C921Y OGT-ID mouse model, systematically demonstrating in mammals the pathological link between O-GlcNAc metabolic imbalance and neurodevelopmental disorders (cortical malformation, microcephaly) as well as behavioral abnormalities (hyperactivity, impulsivity, learning/memory deficits). Researchers comprehensively assessed the model phenotype through integrated multi-level analysis methods, including long-term behavioral monitoring, high-resolution brain structural imaging (micro-CT and MRI), histopathology, and quantitative proteomics.

The core strength of this study lies in its multimodal experimental design. The evidence chain spanning in vivo behavior, brain structure, and molecular characteristics demonstrates high consistency and correlation. Of particular note is the combination of non-invasive behavioral tracking with quantitative neuroimaging techniques, providing objective validation for the observed phenotypes. The findings support the authors' core conclusion: O-GlcNAc homeostasis imbalance correlates with neurodevelopmental deficits, including structural abnormalities in specific brain regions and altered cognitive behaviors. Furthermore, this model reproduces certain clinical features observed in human patients.

Nevertheless, several avenues remain open for further exploration. For instance, sample sizes in certain omics analyses remain relatively small, and investigations into downstream molecular mechanisms are still confined to the level of correlation-direct causal validation through genetic or pharmacological interventions is still required. Furthermore, as this model focuses on a single recurrent mutation, the generalizability of its findings to other OGT-ID variants remains to be verified.

It provides the first actionable vertebrate model for neurodevelopmental disorders with unclear mechanisms, filling a critical gap in this field. The multidimensional research methods established in the paper-such as the digital behavioral phenotyping workflow-also offer valuable references for related disease studies.

<https://doi.org/10.7554/eLife.107170.2.sa2>

Reviewer #2 (Public review):

Summary:

The authors are trying to understand why certain mutants of O-GlcNAc transferase (OGT) appear to cause developmental disorders in humans. As an important step towards that goal, the authors generated a mouse model with one of these mutations that disrupts OGT activity. They then go on to test these mice for behavioral differences, finding that the mutant mice exhibit some signs of hyperactivity and differences in learning and memory. They then examine alterations to the structure of the brain and skull, and again find changes in the mutant mice that have been associated with developmental disorders. Finally, they identify proteins that are up or down regulated between the two mice as potential mechanisms to explain the observations.

Strengths:

The major strength of this manuscript is the creation of this mouse model, as a key step in beginning to understand how OGT mutants cause developmental disorders. This line will prove important for not only the authors but other investigators as well, enabling the testing of various hypotheses and potentially treatments. The experiments are also rigorously performed and the conclusions are well supported by the data.

Weaknesses:

The only weakness is a lack of mechanistic insight. However, this certainly may come in the future through more targeted experimentation using this mouse model. I do not recommend that these experiments need to be performed in this manuscript.

Comments on revisions:

The authors have addressed all of my suggestions proactively.

<https://doi.org/10.7554/eLife.107170.2.sa1>

Author Response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public review):

This study established a C921Y OGT-ID mouse model, systematically demonstrating in mammals the pathological link between O-GlcNAc metabolic imbalance and neurodevelopmental disorders (cortical malformation, microcephaly) as well as behavioral abnormalities (hyperactivity, impulsivity, learning/memory deficits). However, critical flaws in the current findings require resolution to ensure scientific rigor.

The most concerning finding appears in Figure S12. While Supplementary Figure S12 demonstrates decreased OGA expression without significant OGT level changes in C921Y mutants via Western blot/qPCR, previous reports (Florence Authier, et al., Dis Model Mech. 2023) described OGT downregulation in Western blot and an increase in qPCR in the same models. The opposite OGT expression outcomes in supposedly identical mouse models directly challenge the model's reliability. This discrepancy raises serious concerns about either the experimental execution or the interpretation of results. The authors must revalidate the data with rigorous controls or provide a molecular biology-based explanation.

We thank the reviewer for their time and effort in improving the quality of our manuscript.

We would like to point out that the results presented in the previous Fig. S12 (now Fig. S13) are from different ages of the mice and restricted to the prefrontal cortex, compared to the previous report (Florence Authier, et al., Dis Model Mech. 2023) where we showed OGT and OGA mRNA/protein expression in total brain homogenates. In this previous study, we observed a significant reduction in OGT protein levels while OGT mRNA levels were significantly increased in the brains of 3 months old mutant C921Y compared to WT controls. However, in our current study (Figure S12, now S13), OGA and OGT mRNA/protein expression have been a) restricted to the pre-frontal cortex and b) are from 4 months old male mice. Therefore, a direct comparison of findings from total brain vs. prefrontal cortex would be speculative. In our present work, OGT protein levels are not changed in the pre-frontal cortex, while OGT mRNA levels are increased (similarly to the total brain data), albeit not significantly.

It is plausible that the different levels of OGT protein expression in total brain (previous study) and prefrontal cortex (current study) potentially reflect regional differences in the regulation of OGT protein levels/stability, since OGT mRNA levels are increased in both cases. This notion is also supported by additional analyses in three other brain regions (hippocampus, striatum and cerebellum) and these data are now included in Figures S13 and S14.

A few additional comments to the author may be helpful to improve the study.

Major

(1) While this study systematically validated multi-dimensional phenotypes (including neuroanatomical abnormalities and behavioral deficits) in OGT C921Y mutant mice, there is a lack of relevant mechanisms and intervention experiments. For example, the absence of targeted intervention studies on key signaling pathways prevents verification of whether proteomics-identified molecular changes directly drive phenotypic manifestations.

We agree with the reviewer that the suggested experiments would further strengthen our work. However, the extensive nature of the suggested studies would result in considerable delay in sharing this work with the scientific and patient communities. Nevertheless, we appreciate the reviewers' comment and will continue to work along these lines, and report in a follow up manuscript in the future.

(2) Although MRI detected nodular dysplasia and heterotopia in the cingulate cortex, the cellular basis remains undefined. Spatiotemporal immunofluorescence analysis using neuronal (NeuN), astrocytic (GFAP), and synaptic (Synaptophysin) markers is recommended to identify affected cell populations (e.g., radial glial migration defects or intermediate progenitor differentiation abnormalities).

Following the reviewers' suggestion, we have performed additional analyses to identify the cellular composition of the observed nodular dysplasia using neuronal and glial markers. These new analyses indicate that the nodular collections in the layers II/III were predominantly neurons, for example see cresyl violet (Fig. 6E). Moreover, we have also performed immunofluorescence imaging using NeuN and GFAP (Fig. 6G-H), which reflect that the dystrophic collections are predominantly neurons. To further corroborate these findings, we have also performed multiplex IHC analyses, presented in Fig. S12, which indicate that: i) the nodular cortical malformations were populated by neurons and oligodendrocytes and ii) predominantly affected layers II-V, as reflected by the distribution of neuronal markers Reelin and POU class 3 homeobox 2 (POU3F2), and collectively (Fig. 6 and Fig. S12) reflect neuronal disorganisation due to migration defects rather than differentiation defects. We

appreciate the reviewers' suggestion to perform spatiotemporal analyses of these cellular features; however, tissue from defined stages of development is not available.

(3) While proteomics revealed dysregulation in pathways including Wnt/ β -catenin and mTOR signaling, two critical issues remain unresolved: a) O-GlcNAc glycoproteomic alterations remain unexamined; b) The causal relationship between pathway changes and O-GlcNAc imbalance lacks validation. It is recommended to use co-immunoprecipitation or glycosylation sequencing to confirm whether the relevant proteins undergo O-GlcNAc modification changes, identify specific modification sites, and verify their interactions with OGT.

We agree with the referee that these experiments would further strengthen the work. However, we respectfully point out that the inference that altered proteins must themselves be O-GlcNAc modified is not necessarily correct. For instance, O-GlcNAcylation of unknown protein kinase X, E3 ligase/DUB, Y or transcription factor Z could indirectly affect these pathways/proteins. Nevertheless, we have performed further experiments to explore whether Wnt/ β -catenin and mTOR signalling are functionally affected, as pointed out by the referee. In the qPCR analyses, we did not observe significant changes in expression of Wnt target genes (*Cdkn1a*, *Ccnd1*, *Myc*, *Ramp3*, *Tfrc*), neither in protein levels of key proteins involved in Wnt/ β -catenin (non-phosphorylated β -catenin) and mTOR (phosphorylated rpS6) signalling by western blots (data not shown). These results suggest that both pathways are not functionally deregulated in prefrontal cortex of adult OGT^{C921Y} mice to a significant extent.

(4) Given that OGT-ID neuropathology likely originates embryonically, we recommend serial analyses from E14.5 to P7 to examine cellular dynamics during critical corticogenesis phases.

We appreciate the reviewers' suggestion to perform spatiotemporal analyses of these cellular dynamics; however, tissue from defined stages of development is not available. As stated above, we want to share our current findings with the scientific and patient communities in a timely manner, and the suggested experiments could form the foundation of a follow up study in the future.

(5) The interpretation of Figure 8A constitutes overinterpretation. Current data fail to conclusively demonstrate impairment of OGT's protein interaction network and lack direct evidence supporting the proposed mechanisms of HCF1 misprocessing or OGA loss.

Thank you for the comment. To avoid misleading the readers, we have removed panel A from the previous version of Figure 8 and updated the version of record.

Reviewer #2 (Public review):

Summary:

The authors are trying to understand why certain mutants of O-GlcNAc transferase (OGT) appear to cause developmental disorders in humans. As an important step towards that goal, the authors generated a mouse model with one of these mutations that disrupts OGT activity. They then go on to test these mice for behavioral differences, finding that the mutant mice exhibit some signs of hyperactivity and differences in learning and memory. They then examine alterations to the structure of the brain and skull and again find changes in the mutant mice that have been associated with developmental disorders. Finally, they identify proteins that are up- or down-regulated between the two mice as potential mechanisms to explain the observations.

Strengths:

The major strength of this manuscript is the creation of this mouse model, as a key step in beginning to understand how OGT mutants cause developmental disorders. This line will prove important for not only the authors but other investigators as well, enabling the testing of various hypotheses and potentially treatments. The experiments are also rigorously performed, and the conclusions are well supported by the data.

Weaknesses:

The only weakness identified is a lack of mechanistic insight. However, this certainly may come in the future through more targeted experimentation using this mouse model.

We agree with the reviewer that the suggested experiments would further strengthen our work. However, the extensive nature of the suggested studies would result in considerable delay in sharing this work with the scientific and patient communities. Nevertheless, we appreciate the reviewers' comment and will continue to work along these lines, and report in a follow up manuscript in the future.

Recommendations for the authors:

Editor's note:

Should you choose to revise your manuscript, if you have not already done so, please include full statistical reporting including exact p-values wherever possible alongside the summary statistics (test statistic and df) and, where appropriate, 95% confidence intervals. These should be reported for all key questions and not only when the p-value is less than 0.05 in the main manuscript.

Statistics including exact p-values have been included in the main text for all key questions where appropriate.

Reviewer #1 (Recommendations for the authors):

(1) In Figure 1F, the y-axis labels and scale values are partially obscured by graphical elements, compromising accurate interpretation of the data range.

Panel 1F has been adjusted to make the y-axis label visible.

(2) Regarding the histological analyses in Figure 6, the current H&E staining and Luxol Fast Blue myelin staining results lack age-matched wild-type control samples processed in parallel, which undermines experimental comparability. To enhance methodological rigor, control group staining results should be displayed adjacent to each experimental group image.

The original Figure 6 already contained comparison between WT and OGT^{C921Y} tissues. The Figure has been updated with additional data from the WT and C921Y mutant groups shown side by side.

Reviewer #2 (Recommendations for the authors):

(1) I believe that Figures S1 and S2 were switched during the submission. The legends are correct, so the authors should just be careful with the order when they upload the final versions.

Figures S1 and S2 have been re-ordered.

(2) On page 18, the authors state, "Although no significant changes in the expression of OGT were observed in OGTC921Y cortex (Figure S12A, C), there was a significant increase in OGT/OGA protein ratio in OGTC921Y mice (Fig. S12D). As a functional consequence,

global O-GlcNAcylation of proteins in the brain was drastically impaired in the OGTC921Y brain compared to WT (Figure S12E, F).

To me, this statement suggests that the incorrect ratio of OGT to OGA is responsible for the altered O-GlcNAc levels. I think this is missing important information. The authors are, I'm sure, aware that OGT and OGA expression is linked to O-GlcNAc levels. I think it would be better to describe the situation here as the tissue attempting to respond to lower OGT activity by lowering OGA levels. However, the tissue is not fully successful, resulting in lower overall O-GlcNAc levels as seen by RL2. If the difference were only driven by the OGT/OGA ratio, one would expect increased O-GlcNAc levels due to decreased OGA. I think it is important to point out more details here for non-expert readers.

Thank you for the insightful comment, we have included these aspects in the revised text, please see page 20.

(3) I am a little surprised that the authors did not explore differences in O-GlcNAc-modified proteins through a more targeted enrichment of these proteins for analysis of potential modification differences, in addition to just changes in protein abundance.

We agree that these experiments would further strengthen the work. However, it is not known yet whether OGT-CDG is caused by loss of O-GlcNAc modification on specific proteins or due to as yet to decipher mechanisms (e.g. OGT interactome, HCF1 processing, feedback on OGA levels) which we are not able to confirm in the current manuscript. Therefore, as a starting point, we have performed whole proteome analysis to establish candidate hypothesis which could lead to discovering cellular and molecular mechanisms underlying OGT-CDG. Lastly, we appreciate the reviewers' comment and will continue to work along these lines, and report in a follow up manuscript in the future.

<https://doi.org/10.7554/eLife.107170.2.sa0>