

Homeodomain proteins hierarchically specify neuronal diversity and synaptic connectivity

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

December 18, 2023 (this version)

Reviewed preprint version 1

October 2, 2023

Sent for peer review

July 12, 2023

Posted to bioRxiv

June 15, 2023

Chundi Xu , Tyler B. Ramos, Edward M. Rogers, Michael B. Reiser, Chris Q. Doe

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon; Eugene, OR 97403, USA •
Janelia Research Campus, Howard Hughes Medical Institute, 19700 Helix Drive, Ashburn, VA, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

How our brain generates diverse neuron types that assemble into precise neural circuits remains unclear. Using *Drosophila* lamina neuron types (L1-L5), we show that the primary homeodomain transcription factor (HDTF) Brain-specific homeobox (Bsh) is initiated in progenitors and maintained in L4/L5 neurons to adulthood. Bsh activates secondary HDTFs Ap (L4) and Pdm3 (L5) and specifies L4/L5 neuronal fates while repressing the HDTF Zfh1 to prevent ectopic L1/L3 fates (control: L1-L5; Bsh-knockdown: L1-L3), thereby generating lamina neuronal diversity for normal visual sensitivity. Subsequently, in L4 neurons, Bsh and Ap function in a feed-forward loop to activate the synapse recognition molecule DIP- β , thereby bridging neuronal fate decision to synaptic connectivity. Expression of a Bsh:Dam, specifically in L4, reveals Bsh binding to the *DIP- β* locus and additional candidate L4 functional identity genes. We propose that HDTFs function hierarchically to coordinate neuronal molecular identity, circuit formation, and function. Hierarchical HDTFs may represent a conserved mechanism for linking neuronal diversity to circuit assembly and function.

eLife assessment

This paper, offering insights into the mechanisms of neuronal cell type diversification, provides **important** findings that have theoretical or practical implications beyond a single subfield. The data are **compelling** and provide evidence that features methods, data and analyses that are more rigorous than the current state-of-the-art.

Introduction

Our ability to perceive and respond to the world requires a diverse array of neuron types characterized initially by transcription factor (TF) combinatorial codes, followed by neuron type-specific functional attributes such as cell surface molecules, neurotransmitters, and ion channels. It has been well-documented how initial neuronal diversity is generated: in both *Drosophila* and mouse, spatial and temporal factors act combinatorially to generate molecularly distinct newborn neurons (Bayraktar and Doe, 2013 [DOI](#); Doe, 2017 [DOI](#); Erclik et al., 2017 [DOI](#); Holguera and Desplan,

2018 [\[1\]](#); Sen et al., 2019 [\[2\]](#)). Yet most spatial and temporal factors are only transiently present in newborn neurons; therefore, another mechanism is required to bridge initial fate to mature features such as connectivity, neurotransmitters, and ion channels. It remains poorly understood how the initial fate decision of newborn neurons leads to the functional identity of mature neurons. Work from the Hobert lab in *Caenorhabditis elegans* found that each adult neuron type expresses a unique combination of HDTFs, which have been called terminal selectors (Hobert, 2021 [\[3\]](#); Reilly et al., 2020 [\[4\]](#)). Terminal selector HDTFs not only drive the expression of neuron functional identity genes but also activate pan-neuronal genes (Hobert, 2021 [\[3\]](#); Howell et al., 2015 [\[5\]](#); Kratsios et al., 2015 [\[6\]](#); Stefanakis et al., 2015 [\[7\]](#)). Loss of terminal selector HDTFs frequently results in altered neuronal identity and function (Arlotta and Hobert, 2015 [\[8\]](#); Cros and Hobert, 2022 [\[9\]](#); Reilly et al., 2022 [\[10\]](#)).

Although a great deal is known about HDTFs in the specification of neuron-type specific morphology and synaptic connectivity in *C. elegans* (Chen et al., 2012 [\[11\]](#); Cubelos et al., 2010 [\[12\]](#); De Marco Garcia and Jessell, 2008 [\[13\]](#); Friedrich et al., 2016 [\[14\]](#); Hasegawa et al., 2013 [\[15\]](#); Ozel et al., 2022 [\[16\]](#); Sakkou et al., 2007 [\[17\]](#); Santiago and Bashaw, 2014 [\[18\]](#); Thor and Thomas, 1997 [\[19\]](#)), it remains unknown the extent to which this model is generalizable to other organisms. Here we use the *Drosophila* lamina, the first ganglion in the optic lobe, to test the hypothesis that HDTFs couple the initial fate decision to later circuit formation and functional aspects of the neuron. The *Drosophila* lamina has only five intrinsic neuron types (L1-L5), which are analogous to bipolar cells in the vertebrate (Sanes and Zipursky, 2010 [\[20\]](#)). During late larval and early pupal stages, lamina progenitor cells (LPCs) give rise to L1-L5 neurons (Fernandes et al., 2017 [\[21\]](#); Huang et al., 1998 [\[22\]](#)). The core motion detection neurons L1-L3 receive direct synaptic inputs from photoreceptors and mediate visual motion detection, whereas L4 and L5 receive synaptic inputs from L2 and L1, respectively, and their function is currently unclear (Meinertzhagen and O'Neil, 1991 [\[23\]](#); Rivera-Alba et al., 2011 [\[24\]](#); Silies et al., 2013 [\[25\]](#); Tuthill et al., 2013 [\[26\]](#)). The cell bodies of each lamina neuron type are localized in a layer-specific manner (Tan et al., 2015 [\[27\]](#)). L2/L3 cell bodies are intermingled in the most distal layer, while L1, L4, and L5 form distinct layers progressively more proximal (**Figure 1** [\[28\]](#)). Each lamina neuron type expresses unique TF markers. L1, L2, and L3 neurons express Svp, Bab2, and Erm, respectively (Tan et al., 2015 [\[27\]](#)). L4 and L5 neurons express the HDTFs Bsh/Ap and Bsh/Pdm3, respectively (Hasegawa et al., 2013 [\[15\]](#); Tan et al., 2015 [\[27\]](#)). Work from Hasegawa et al has shown that in *bsh* mutants, L4 adopts L3-like morphology, L5 becomes glia, and Ap mRNA level is reduced (Hasegawa et al., 2013 [\[15\]](#)). This suggests that the HDTF Bsh is important for L4/L5 neuron type specification.

Here, we show that HDTFs function hierarchically in coupling neuronal fate specification to circuit assembly. First, a primary (earlier-initiated) HDTF Bsh activates two secondary (later-initiated) HDTFs Ap (L4) and Pdm3 (L5) and specifies L4 and L5 neuronal fates. Next, we show that the primary HDTF Bsh generates lamina neuronal diversity (control: L1-L5; Bsh-knockdown: L1-L3) required for normal visual sensitivity. Third, in L4 neurons, Bsh and Ap function in a feedforward loop to activate the synapse recognition molecule (DIP- β), thereby bridging neuronal fate decision to synaptic connectivity. Fourth, Bsh in L4 neurons directly binds at the DIP- β locus and other L4 functional identity genes. Our work may provide a conserved mechanism of HDTFs hierarchically linking neuronal diversity to circuit assembly.

Results

Sequential expression of HDTFs during lamina neurogenesis

L4 and L5 neurons are generated by a subset of LPCs during late larval and early pupal stages (Fernandes et al., 2017 [\[21\]](#); Huang et al., 1998 [\[22\]](#)), but it is unknown exactly when Bsh, Ap and Pdm3 are initially expressed. To address this question, we identified the Tailless (Tll) TF as a novel marker for LPCs (**Figure 1A-C** [\[29\]](#)). Indeed, Tll and the neuronal marker Elav have precise




Figure 1.

Sequential initiation of HDTFs during lamina neurogenesis.

(A-A'') Tll is identified as an LPC marker, expressed complementary to Elav; Dac labels both Tll+ LPCs and Elav+ neurons. LF: lamina furrow. Here and below, scale bar: 10 μ m, $n \geq 5$ brains.

(B-B'') Tll+ cells are localized both within the lamina columns and before the columns. Lamina columns (white dash circle) are outlined by the photoreceptor axons, which are labeled by Chaoptin. $n \geq 5$ brains. LF: lamina furrow.

(C-C'') Bsh is expressed in Tll+ Elav-LPCs (white dash circle) as well as in Elav+ L4 and L5 neurons.

(D-D'') Ap is expressed in L4 neurons. Newborn L4 neurons are Bsh+ Elav+ Ap- (yellow line circle), while older L4 neurons are Bsh+ Elav+ Ap+ (white line circle).

(E-E'') Pdm3 is expressed in L5 neurons. Newborn L5 neurons are Bsh+ Pdm3- (yellow line circle), while older L5 neurons are Bsh+ Pdm3+ (white line circle).

(F-F'') Bsh, Ap, and Pdm3 expressions are maintained in adults.

(G) Schematic of lamina neuron development in early pupa.

(H) Summary.

complementary high-level expression, validating Tll as an LPC marker (**Figure 1A-C**), summarized in **1G**). Importantly, we found that Bsh was first detected in a subset of Tll+ LPCs (**Figure 1C**). In contrast, Ap and Pdm3 were first detected much later in L4 and L5 neurons, respectively (**Figure 1D-E**). Ap and Pdm3 were never detected in LPCs or newborn L4 or L5 neurons (**Figure 1D-E**), showing that Bsh is expressed prior to Ap and Pdm3. Expression of Bsh, Ap, and Pdm3 was maintained in L4/L5 neurons into the adult (**Figure 1F**), consistent with a potential role as terminal selectors in maintaining neuron identity (Deneris and Hobert, 2014; Serrano-Saiz et al., 2018). We conclude that Bsh expression is initiated in LPCs, while Ap and Pdm3 are initiated in neurons. Due to its earlier expression, we refer to Bsh as a primary HDTF, and due to their later expression, we refer to Ap and Pdm3 as secondary HDTFs (**Figure 1H**).

Bsh activates Ap/Pdm3 expression and specifies L4/L5 neuronal fates

Bsh is expressed prior to Ap and Pdm3, raising the possibility that Bsh activates Ap and Pdm3 expression. We used R27G05-Gal4 to express Bsh-RNAi in LPCs. To confirm whether R27G05-Gal4 is turned on in LPCs, we used R27G05-Gal4>UAS-myristoylated-GFP, and indeed, GFP can be detected in all LPCs and lamina neurons until ~66 hours after pupa formation (APF) (**Figure 2—figure supplement 1A, D, E**). To test whether R27G05-Gal4 can be turned on in lamina neurons, we took advantage of tubP-Gal80[ts] to temporally control R27G05-Gal4>UAS-myristoylated-GFP. We found that GFP can be detected in most lamina neurons when tubP-Gal80[ts] was inactivated from the beginning of lamina neurogenesis. In contrast, GFP is absent in most lamina neurons when tubP-Gal80[ts] was inactivated from the end of lamina neurogenesis, suggesting that R27G05-Gal4 is functionally an LPC-Gal4 transgene (**Figure 2—figure supplement 1B, C**). As expected, Bsh-knockdown (KD) (R27G05-Gal4>UAS-Bsh-RNAi) eliminates Bsh expression in LPCs and newborn neurons (**Figure 2A-F**). Bsh remains undetectable in the lamina neurons in the adult, despite the lack of RNAi expression, which indicates that Bsh expression is unable to reinitiate in neurons if lost in LPCs (**Figure 2—figure supplement 2A'-B'**). Importantly, Bsh-KD resulted in a nearly complete loss of Ap and Pdm3 expression (**Figure 2G-J**); these neurons are not dying, as the number of Elav+ lamina neurons is unchanged (**Figure 2K**). Note that ectopic Ap expression in L5 is caused by the R27G05-Gal4 line, probably due to its genome insertion site, but this does not affect our conclusion that Bsh is required for Ap and Pdm3 expression. Consistent with the Bsh-KD phenotype, we found similar results using a Bsh Crispr/Cas9 knockout (KO) (**Figure 2—figure supplement 2E-I**). Taken together, we conclude that the primary HDTF Bsh is required to drive the expression of the secondary HDTFs Ap and Pdm3 and specify L4/L5 neuronal fates (**Figure 2L**).

To determine if Bsh is continuously required for Ap expression in L4, we first performed Bsh knockdown specifically in L4 neurons using L4-split Gal4 (31C06-AD, 34G07-DBD)>UAS-Bsh-RNAi. Interestingly, this fails to knock down Bsh, suggesting L4-split Gal4 depends on Bsh expression. Next, we tried a Bsh Crispr/Cas9 knock out (Bsh-KO: 31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs) to remove Bsh expression beginning in L4 neurons, and it indeed led to a significant decrease in Bsh+ neurons between 2d-3d APF (**Figure 2—figure supplement 3**). Despite the loss of Bsh expression in most L4 neurons, we observed no loss of Ap expression and no derepression of other lamina neuron markers (**Figure 2—figure supplement 4**). These results show (a) that all known lamina neuron markers are independent of Bsh regulation in neurons and (b) that Ap may undergo positive autoregulation after its initiation, rendering it independent of Bsh. Autoregulation is a common feature of HDTFs (Leyva-Díaz and Hobert, 2019). We conclude that Bsh is not required to maintain Ap expression or repress other lamina neuron markers in L4 neurons.

Bsh suppresses L1/L3 neuronal fates

We next asked whether L4/L5 neurons are transformed into another neuronal type following Bsh-KD (R27G05-Gal4>UAS-Bsh-RNAi). Above, we showed that the number of Elav+ lamina neurons remains unchanged in Bsh-KD, indicating that L4/L5 have assumed a different cell fate (**Figure 2K**). To test for ectopic generation of another lamina neuron type, we assayed the expression of Svp (L1), Bab2 (L2), and Erm (L3). We found that Bsh-KD led to ectopic expression of the L1 and L3 markers Svp and Erm in the positions normally occupied by L4/L5 cell bodies (**Figure 3A-E**). Notably, we never saw cell bodies co-expressing Erm and Svp, excluding the possibility of an ectopic hybrid neuronal fate in Bsh-KD. In contrast, the L2 marker Bab2 was unaffected by Bsh-KD (**Figure 3F-H**). Together, our data suggest that the absence of Bsh may generate ectopic L1 and L3 neuron types at the expense of L4 and L5.

To confirm L1 and L3 neurons are ectopically generated at the expense of L4 and L5 in Bsh-KD, we used Bsh-LexA>LexAop-GFP to trace L5 neurons (**Figure 3—figure supplement 1A-A'**). Indeed, control GFP+ neurons turn on L5 marker Pdm3 soon after their birth (**Figures 3I-I'**). In contrast, Bsh-KD generated GFP+ neurons that turn on L1 marker Svp instead of L5 marker Pdm3 (**Figure 3I-L**). Interestingly, some of the GFP+Svp+ neurons had cell bodies displaced from the L5 layer into the L1 layer (**Figure 3J**), which suggests that L1 neurons may actively seek out their appropriate settling layer. Furthermore, we also observed a transformation of L5 to L1 in neuronal morphology. In control, GFP+ neurons, which trace L5 neurons, have very few dendrites in the lamina neuropil (**Figure 3M**). In contrast, Bsh-KD resulted in GFP+ neurons elaborating L1-like dendrite arbors - bushy dendrites throughout the lamina (**Figure 3N, O**; summarized in **3P**). Because Bsh-KD generates L1 neuron type at the expense of L5, the ectopic L3 neurons must be generated at the expense of L4, which is confirmed by our accompanying work (Xu et al., 2023). This is also consistent with a previous report showing the transformation of L4 to L3 morphology in *bsh* mutant clones, although, unlike our results, they observed an L5>glial fate change using L5-Gal4 to trace L5 in *bsh* clones (Hasegawa et al., 2013). The difference in results (where we see L5>L1 and they see L5>glia) is likely due to the unfaithful expression of L5-Gal4 in *bsh* mutants where L5 neuron type is missing. Together, we conclude that the absence of Bsh generates ectopic L1 and L3 neuron types at the expense of L5 and L4, respectively.

Bsh-KD in LPCs results in a loss of Ap expression and ectopic L1/L3 marker expression. To exclude the possibility that Bsh represses L1/L3 fates through Ap, we knocked down Ap expression from their time of birth using an LPC-Gal4 line (R27G05-Gal4>Ap-RNAi). As expected, Ap-KD eliminates Ap expression in L4 neurons (**Figure 3—figure supplement 2**). Furthermore, Ap remained undetectable in lamina neurons in the adult, which indicates that Ap expression is unlikely able to reinitiate if normal initiation is lost (**Figure 3—figure supplement 2**). Importantly, loss of Ap did not affect Bsh expression in L4 and did not lead to ectopic expression of other lamina neuron markers (**Figure 3—figure supplement 3**), suggesting that Bsh but not Ap is required to repress L1/L3 neuronal fates. Taken together, we conclude that the primary HDTF Bsh, but not the secondary HDTF Ap, promotes L4/L5 neuronal fates and suppresses L1/L3 fates, thereby generating lamina neuronal diversity (**Figure 3P**).

Bsh represses Zfh1 to suppress L1/L3 neuronal fates

How does the primary HDTF Bsh repress L1 and L3 neuronal fates? We hypothesize that Bsh might repress an unidentified HDTF, which is shared by L1 and L3 and required to generate L1 and L3 neuron types. To find this HDTF, we screened published RNA-seq data (Tan et al., 2015) and found that the HDTF Zfh1 was present in all LPCs before becoming restricted to L1 and L3 neurons (**Figure 4A-D**). To determine if Zfh1 is required for L1 and L3 neuronal fates, we used RNAi to perform Zfh1-KD in LPCs (R27G05-Gal4>Zfh1-RNAi). As expected, Zfh1-KD significantly decreased Zfh1 nuclear levels in all LPCs and neurons (**Figure 4D-E'**, **Figure 4—figure supplement 1A-D**) and resulted in a loss of Svp+ L1 and Erm+ L3 neurons (**Figure 4D-F**), consistent with a role in specifying L1 and L3 neuronal identity. In contrast, Zfh1-KD did not increase the total




Figure 3.

Bsh suppresses L1/L3 neuronal fate.

(A-E) Bsh-KD in LPCs results in the ectopic expression of the L1 marker Svp and L3 marker Erm in L4/L5 cell body layers (circled). (C) Schematic of lamina neuron development from 1day pupae to adult. (D and E) Quantification of Erm and Svp expression. Here and below, scale bar, 10 μ m.

(F-H) Bsh-KD in LPCs does not produce ectopic Bab2-positive neurons or glia in the L5 layer (circled). $n \geq 5$ brains. Genotype: R27G05-Gal4>UAS-Bsh-RNAi. (H) Quantification of Bab2 expression.

(I-L) Bsh-KD in LPCs results in ectopic Svp+ L1 neurons at the expense of Pdm3+ L5 neurons. Bsh GFP+ neurons marked with yellow arrowheads show L1 marker Svp expression in Bsh-KD while L5 marker Pdm3 expression is in control. Genotype: Bsh-LexA>LexAop-GFP. (K and L) Quantification of Pdm3 and Svp expression.

(M-O) Bsh-KD transforms L5 neuron morphology to L1-like neuronal morphology. (M) Control L5 neurons have very few dendrites in the lamina neuropil. Genotype: R27G05-Gal4, Bsh-LexA>LexAop-GFP. (N) Bsh-KD transforms L5 neuron morphology to L1-like neuronal morphology. Genotype: R27G05-Gal4>UAS-Bsh-RNAi; Bsh-LexA>LexAop-GFP. (O) Control L1 neurons show bushy dendrites throughout the lamina. Genotype: svp-Gal4, R27G05-FLP>UAS-FRT-stop-FRT-myrGFP.

(P) Summary.

Data are presented as mean $\pm$ SEM. Each dot represents each brain. $n=5$ brains in (D), (E), (H), (K), and (L). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, ns=not significant, unpaired t-test.

number of Ap/Pdm3+ neurons (**Figure 4—figure supplement 1E-G**), showing that *Zfh1* does not repress L4/L5 neuronal identity. We observed that *Zfh1*-KD resulted in fewer lamina neurons overall, including a reduction of Bsh+ L4 and L5 neurons (**Figure 4—figure supplement 1E'-K**); this suggests a potential role for *Zfh1* in LPCs in regulating lamina neurogenesis. Importantly, Bsh-KD (R27G05-Gal4>Bsh-RNAi) resulted in ectopic expression of *Zfh1* in the L4/L5 cell body layers and in GFP+ neurons (normally L5) (**Figure 4G-I**, **Figure 4—figure supplement 1L-N**). We propose that Bsh and *Zfh1* are both primary HDTFs (specifying L4/L5 and L1/L3, respectively) and that Bsh represses *Zfh1* to suppress L1/L3 neuronal fates (**Figure 4J**).

Bsh:DamID reveals Bsh direct binding to L4 identity genes and pan-neuronal genes

In *C. elegans*, terminal selectors show non-redundant control of neuronal identity genes and redundant control of pan-neuronal genes. To see if the same mechanism is used by the HDTF Bsh, we profiled Bsh direct targets with precise spatial and temporal control: only in L4 neurons at the time of synapse formation (46-76h APF). To do this, we used Targeted DamID (Aughey et al., 2021), which can be used to identify Bsh DNA-binding sites across the genome (**Figure 5A**). We generated a Bsh:Dam transgenic fly line according to published methods (Aughey et al., 2021) and expressed it specifically in L4 neurons using the L4-Gal4 transgene R31C06-Gal4 during synapse formation (**Figure 5—figure supplement 1A-B"**). We verified that the Bsh:Dam fusion protein was functional by rescuing Ap and Pdm3 expression (**Figure 5—figure supplement 1C-E**). We performed three biological replicates which had high reproducibility (**Figure 5B**).

Next, we determined which Bsh-bound genomic targets showed enriched transcription in L4 neurons during synapse formation using recently published L4 scRNA sequencing data from the same stage (Jain et al., 2022). There are 958 genes that are significantly transcribed in L4 at 48h or 60h APF. Among them, 421 genes show Bsh:Dam binding peaks, while 537 genes do not show Bsh:Dam binding peaks (**Figure 5C**). Genes having Bsh:Dam binding peaks include numerous candidate L4 identity genes: ion channels, synaptic organizers, cytoskeleton regulators, synaptic recognition molecules, neuropeptide/receptor, neurotransmitter/receptor, and pan-neuronal genes (**Figure 5D**, Supplementary File 1). Genes expressed in L4 but not having Bsh:Dam binding peaks include long non-coding RNA, mitochondrial genes, ribosomal protein, heat shock protein, ATP synthase, and others (Supplementary File 2).

We previously showed that *DIP-β*, a cell surface protein of the immunoglobulin superfamily, is specifically expressed in L4 neurites in the proximal lamina, and is required for proper L4 circuit formation (Xu et al., 2019). Here we found Bsh binding peaks in L4 neurons within the first intron of the *DIP-β* gene (**Figure 5D**). Interestingly, the Ecdysone receptor (EcR), which controls the temporal expression of *DIP-β* in L4 neurons, also has a DNA-binding motif in the *DIP-β* first intron (Jain et al., 2022), suggesting Bsh and the EcR pathway may cooperate to achieve the proper spatial (Bsh) and temporal (EcR) expression pattern of the *DIP-β* synapse recognition molecule. Taken together, consistent with the work of terminal selector in *C. elegans*, we found evidence that Bsh:Dam shows direct binding to L4 identity genes -- including *DIP-β* -- as well as pan-neuronal genes (summarized in **Figure 5E**).

Although our experiment focused on Bsh targets during the stages of synapse formation, we checked for Bsh binding at the HDTF loci that might be bound by Bsh during the earlier stages of neuronal identity specification. We found that Bsh:Dam showed a Bsh binding peak at the *ap* locus, suggesting that Bsh may redundantly maintain Ap expression in L4 neurons through the stage of synapse formation (along with potential Ap positive autoregulation) (**Figure 5—figure supplement 1F**). In contrast, Bsh:Dam did not show binding peaks at *pdm3* or *zfh1* loci, which might be due to the inaccessibility of *pdm3* or *zfh1* loci in L4 neurons. Indeed, Dam (open




Figure 4.

Bsh represses Zfh1 to suppress L1/L3 neuronal fate.

(A-A'') Zfh1 is expressed in all LPCs and some lamina neurons at 12h APF. GFP labels all lamina cells. Elav labels lamina neurons. The yellow dash circle outlines LPCs and the white dash circle outlines lamina neurons. Genotype: R27G05-Gal4>UAS-myrGFP. Here and below, scale bar, 10 μ m. n $\geq$ 5 brains.

(B-B'') Zfh1 is expressed in Svp+ L1 and Erm+ L3 neurons at 1-2d APF; Svp and Erm are never coexpressed. The white dashed circle outlines L1 and L3 neurons and the yellow line indicates the rough boundary between L1 and L3 cell bodies.

(C) Schematic of lamina neuron development from 1 day pupae to adult.

(D-F) Zfh1-KD in LPCs results in a loss of Svp+ L1 and Erm+ L3 neurons. Quantification: the number of Erm+ or Svp+ cell bodies in a single optical slice. Genotype: R27G05-Gal4>UAS-zfh1-RNAi.

(G-I) Bsh-KD in LPCs results in ectopic Zfh1 in L4/L5 layers. White circles label Bsh+ cell bodies in L4 layer in control. L5 layer, white outline. Quantification: the percentage of Bsh+ or Zfh1+ neurons in L5 layer. Genotype: R27G05-Gal4>UAS-bsh-RNAi.

(J) Summary.

Data are presented as mean $\pm$ SEM. Each dot represents each brain. n=6 brains in (F) and n=5 brains in (I). ***P<0.001, unpaired t-test.




Figure 5.

Bsh:DamID reveals Bsh direct binding to L4 identity genes and pan-neuronal genes.

(A) Schematic for TaDa method.

(B) Both Dam and Bsh:Dam show high Pearson correlation coefficients among their three biological replicates, with lower Pearson correlation coefficients between Dam and Bsh:Dam. A heat map is generated using the multiBamSummary and plotCorrelation functions of deepTools.

(C) Bsh:Dam peaks in L4 (46h-76h APF) called by find_peaks and L4 scRNA-seq data (48h APF; 60h APF) (Jain et al., 2022) are combined (see method).

(D) Bsh:Dam shows strong signals in L4 identity genes and pan-neuronal genes. The Dam alone signal indicates the open chromatin region in L4 neurons. The y axes of Bsh:Dam signal represent log₂(Bsh:Dam/Dam) scores. Bsh peaks in L4 neurons were generated using find_peaks (FDR<0.01; min_quant=0.9) and peaks2genes.

(E) Summary

chromatin) (Aughey et al., 2018) does not show a peak at *pdm3* or *zfh1* loci, suggesting that *pdm3* or *zfh1* loci are not accessible in L4 neurons during synaptogenesis (Figure 5—figure supplement 1F).

Bsh and Ap form a coherent feed-forward loop to activate DIP-β

Here we ask whether Bsh or Ap are required for the expression of DIP-β in L4 neurons. We found that Bsh-KO (31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs) only in L4 neurons resulted in a strong decrease in DIP-β levels (Figure 6A-D). Furthermore, using the STaR method (Chen et al., 2014; Xu et al., 2019), we found that L4 primary dendrite length and presynaptic Bruchpilot (Brp) puncta in the proximal lamina were both decreased following Bsh-KO (Figure 6E-I); this is distinct from the DIP-β-KD phenotype (Xu et al., 2019). We conclude that Bsh is required to activate DIP-β expression in L4 and regulate L4 morphology and connectivity. Because Bsh-KO has a distinct phenotype from DIP-β-KD, it is likely that Bsh has distinct targets in addition to DIP-β.

To test whether Ap controls DIP-β expression in L4, we knocked down Ap in L4 neurons (R27G05-Gal4>Ap-RNAi) and observed a strong decrease in DIP-β levels in L4 neurons (Figure 6J-M). Note that L4-split Gal4 (31C06-AD, 34G07-DBD)>UAS-Ap-RNAi cannot knock down Ap in L4 neurons due to the dependence of L4-split Gal4 on Ap expression. We next combined the STaR method with Ap-shRNA (see Materials and Methods), resulting in a loss of Ap expression in L4 at 2d APF and a strong decrease in DIP-β levels in L4 neurons (Figure 6—figure supplement 1). Importantly, Ap-KD in L4 neurons increased Brp puncta in the distal/proximal lamina and increased the length of L4 primary dendrites, which is similar to DIP-β-KD phenotype (Xu et al., 2019) (Figure 6N-R). We conclude that Ap is required to activate DIP-β expression in L4 and regulate L4 morphology and connectivity mainly through DIP-β.

A coherent feed-forward motif is A activates B, followed by A and B, both activating C (Mangan and Alon, 2003). This is what we observe for Bsh, Ap, and DIP-β. Bsh activates Ap in L4 neurons soon after their birth (Figure 2G, H), and Bsh and Ap are both required to activate DIP-β (Figure 6A-D, J-M). Importantly, loss of Bsh in L4 neurons decreases DIP-β levels (Figure 6A-D) without altering Ap expression (Figure 2—figure supplement 4F). Similarly, loss of Ap has no effect on Bsh (Figure 3—figure supplement 3A-B'''), yet it decreases DIP-β levels (Figure 6J-M). In conclusion, we have defined a coherent feed-forward loop in which Bsh activates Ap, and then both are independently required to promote expression of the synapse recognition gene DIP-β, thereby bridging neuronal fate decision to synaptic connectivity (Figure 6S).

Bsh is required for normal visual behavior

Bsh is required to specify L4/L5 neuronal fate and generate lamina neuronal diversity (control: L1-L5; Bsh-KD: L1-L3) (Figure 3), raising the hypothesis that lack of Bsh may compromise lamina function. To test this, we used an apparatus (the Fly Vision Box) that integrates multiple assays, including visual motion (Zhu et al., 2009), phototaxis (Benzer, 1967), and spectral preference (Gao et al., 2008) in flies walking in transparent tubes (Isaacson et al., 2023) (Figure 7A; see Methods). We used the LPC-specific driver R27G05-Gal4 to express Bsh RNAi in LPCs. We found that Bsh-KD in LPCs resulted in a lack of Bsh in the adult lamina (Isaacson et al., 2023) (Figure 7—figure supplement 1). Flies with Bsh-KD in LPCs, where L4/L5 neurons are transformed into L1/L3 neurons, showed a reduced response to a high-speed stimulus, suggesting weakened sensitivity to visual motion (Figure 7B). Previous work found that L4 function is not required for motion detection when silencing L4 or L5 neuron activity alone (Silies et al., 2013; Tuthill et al., 2013), which suggests that L4 and L5 acting together might be required for normal sensitivity to visual motion. When tested with both UV and green LEDs, the Bsh-KD flies had reduced phototaxis to both dim and bright lights, suggesting less sensitivity to both UV and green lights. Interestingly, the Bsh-KD flies exhibited larger responses toward bright UV illumination in the spectral preference assay (Figure 7C, D). This apparent attraction to bright UV light may result from more weakened sensitivity to green light, which may be expected since L4 and L5 are




Figure 6.

Bsh and Ap form a coherent feed-forward loop to activate DIP-β.

(A-D) Bsh Crispr KO in postmitotic L4 neurons results in loss of DIP-β expression in the proximal lamina neuropil (arrow) at 3d APF. DIP-β expression is detected using an anti-DIP-β antibody. The signal in the distal lamina is from non-lamina neurons, probably LawF. Significantly reduced DIP-β fluorescence intensity is observed in the proximal lamina (75%–100% distance, marked by red bar (C, D). *** $P < 0.001$, unpaired t-test, $n = 8$ brains, each line represents each brain, scale bar, 10 μm .

Genotype: 31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs.

(E-I) Bsh Crispr KO in L4 neurons results in a decrease of primary dendrite length and proximal synapse number in postmitotic L4 neurons of 1d adults. Here and below, white dash lines indicate the lamina neuropil and yellow lines show the boundary between the distal and proximal lamina. The average number of Brp puncta in L4 neurons present within the distal or proximal halves of lamina cartridges. * $P < 0.05$, *** $P < 0.001$, ns= not significant, unpaired t-test, $n = 100$ cartridges, $n = 5$ brains, each dot represents one cartridge, data are presented as mean $\pm$ SEM. Genotype: 31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs, UAS-myrGFP, UAS-RSR, 79C23-S-GS-rst-stop-rst-smFPV5-2a-GAL4.

(J-M) Ap RNAi KD in postmitotic L4 neurons results in loss of DIP-β expression in the proximal lamina neuropil (arrow) at 3d APF. The signal in the distal lamina is from non-lamina neurons, probably LawF. Significantly reduced DIP-β fluorescence intensity is observed in the proximal lamina (75%–100% distance, marked by red bar (L, M). *** $P < 0.001$, unpaired t-test, $n = 8$ brains, each line represents each brain, scale bar, 10 μm . Genotype: R27G05-Gal4>UAS-ApRNAi.

(N-R) Ap-KD in L4 neurons results in an increase of primary dendrite length and proximal synapse number in postmitotic L4 neurons in 1d adults. The average number of Brp puncta in L4 neurons present within the distal or proximal halves of lamina cartridges. * $P < 0.05$, *** $P < 0.001$, ns= not significant, unpaired t-test, $n = 100$ cartridges, $n = 5$ brains, each dot represents one cartridge, data are presented as mean $\pm$ SEM. Genotype: 31C06-AD, 34G07-DBD>UAS-RSR, 79C23-S-GS-rst-stop-rst-smFPV5-2a-GAL4, UAS-Ap-shRNA, UAS-myrGFP.

(S) Summary.

indirect R1-6 targets (Takemura et al., 2015 [↗](#)). Taken together, we conclude that the primary HDTF Bsh is required to generate lamina neuronal diversity and normal visual behavior (**Figure 7E** [↗](#)).

Discussion

HDTFs are evolutionarily conserved factors in specifying neuron-type specific structure and function (Hobert, 2021 [↗](#); Hobert and Kratsios, 2019 [↗](#); Kitt et al., 2022 [↗](#)). In *C. elegans*, some HDTFs function as terminal selectors, controlling the expression of all neuronal identity genes and diversifying neuronal subtypes, while other HDTFs act downstream of terminal selectors to activate a subset of identity genes (Gordon and Hobert, 2015 [↗](#); Hobert, 2016 [↗](#)). Here, we show that the Bsh primary HDTF functions for L4/L5 fate specification by promoting expression of the Ap and Pdm3 secondary HDTFs and suppressing the HDTF Zfh1 to inhibit ectopic L1/L3 fate, thereby generating lamina neuronal diversity. In L4, Bsh and Ap act in a feed-forward loop to drive the expression of synapse recognition molecule DIP- β , thereby bridging neuronal fate decision to synaptic connectivity (**Figure 7F** [↗](#)). Our DamID data provides support for several hundred Bsh direct binding targets that also show enriched expression in L4 neurons; these Bsh targets include predicted and known L4 identity genes as well as pan-neuronal genes, similar to the regulatory logic first observed in *C. elegans* (Hobert, 2021 [↗](#); Stefanakis et al., 2015 [↗](#)) (**Figure 5C** [↗](#)). HDTFs are widely expressed in the nervous system in flies, worms, and mammals. By characterizing primary and secondary HDTFs according to their initiation order, we may decode conserved mechanisms for generating diverse neuron types with precise circuits assembly.

How can a single primary HDTF Bsh activate two different secondary HDTFs and specify two distinct neuron fates: L4 and L5 (**Figure 7F** [↗](#))? In our accompanying work (Xu et al., 2023 [↗](#)), we show that Notch signaling is activated in newborn L4 but not in L5. This is not due to an asymmetric partition of a Notch pathway component between sister neurons, as is common in most regions of the brain (Li et al., 2013 [↗](#); Mark et al., 2021 [↗](#)), but rather due to L4 being exposed to Delta ligand in the adjacent L1 neurons; L5 is not in contact with the Delta+ L1 neurons and thus does not have active Notch signaling. We show that while Notch signaling and Bsh expression are mutually independent, Notch is necessary and sufficient for Bsh to specify L4 fate over L5. The Notch^{ON} L4, compared to Notch^{OFF} L5, has a distinct open chromatin landscape which allows Bsh to bind distinct genomic loci, leading to L4-specific identity gene transcription. We propose that Notch signaling and HDTF function are integrated to diversify neuronal types.

We used DamID (this work) and a scRNAseq dataset (Jain et al., 2022 [↗](#)) to identify genomic loci containing both Bsh direct binding sites and L4-enriched expression. Genes that have Bsh:Dam binding peaks but are not detected in L4 scRNA sequencing data at 48h or 60h APF might be due to the following reasons: they are transcribed later, at 60h – 76h APF; the algorithm (find_peaks; peaks2genes) that we used to detect Bsh:Dam peaks and call the corresponding genes is not 100% accurate; some regulatory regions are outside the stringent ± 1 kb association with genes; Bsh may act as transcription repressor; TFs generally act combinatorially as opposed to alone and that many required specific cooperative partner TFs to also be bound at an enhancer for gene activation; and scRNAseq data is not 100% accurate for representing gene transcription (**Figure 5C** [↗](#), Supplementary File 3).

Does the primary HDTF Bsh control all L4 neuronal identity genes? It seems likely, as Bsh:Dam shows binding to L4-transcribed genes that could regulate L4 neuronal structure and function, including the functionally validated synapse recognition molecule DIP- β . Furthermore, we found Bsh and Ap form a feed-forward loop to control DIP- β expression in L4 neurons. Similarly, in *C. elegans*, terminal selectors UNC-86 and PAG-3 form a feed-forward loop with HDTF CEH-14 to control the expression of neuropeptide FLP-10, NLP-1 and NLP-15 in BDU neurons (Gordon and Hobert, 2015 [↗](#)), suggesting an evolutionarily conserved approach, using feed-forward loops, for




Figure 7.

Bsh+ L4/L5 are required for normal visual sensitivity.

(A) Schematic of the Fly Vision Box.

(B) Bsh-KD adult flies show reduced responses to a high-speed stimulus. Left: stimulus with different temporal frequency; right: stimulus with 5Hz temporal frequency, but with the indicated contrast level.

(C) Bsh-KD adult flies show reduced phototaxis to both dim and bright lights. Relative intensity was used: 15 and 200 for dim and bright green respectively; 20 and 100 for dim and bright UV respectively.

(D) Bsh-KD adult flies show larger responses towards bright UV illumination. For lower UV levels, flies walk towards the green LED, but walk towards the UV LED at higher UV levels. Data are presented as mean $\pm$ SD, with individual data points representing the mean value of all flies in each tube. For control (mcherry-RNAi), $n=18$ groups of flies (tubes), for Bsh-KD, $n=16$ groups of flies, run across 3 different experiments. Each group is 11-13 male flies. * $P < 0.05$, unpaired, two-sample t-test controlled for false discovery rate.

(E) Model. Left: In wild type, Zfh1+ LPCs give rise to L1 and L3 neurons, whereas Zfh1+Bsh+ LPCs give rise to L4 and L5 neurons. The lineage relationship between these neurons is unknown. Right: Bsh KD results in a transformation of L4/L5 into L1/L3 which may reveal a simpler, ancestral pattern of lamina neurons that contains the core visual system processing arrangement.

(F) Summary.

terminal selectors to activate neuronal identity genes. An important future direction would be testing whether Bsh controls the expression of all L4 identity genes via acting with Ap in a feed-forward loop. One intriguing approach would be profiling the Ap genome-binding targets in L4 during the synapse formation window and characterizing the unique and sharing genome-binding targets of Bsh and Ap in L4 neurons. Further, it would be interesting to test whether the primary HDTF Bsh functions with Ap to maintain neuron type-specific morphology, connectivity, and function properties in adults.

Newborn neurons are molecularly distinct prior to establishing their characteristic morphological or functional attributes. We discovered that the primary HDTF Bsh is specifically expressed in newborn L4 and L5 neurons and is required to specify L4 and L5 fates, suggesting that identifying differentially expressed factors in newborn neurons is essential to decoding neuron type specification. We note that primary and secondary TFs may be HDTFs as well as non-HDTFs. For example, the primary HDTF Zfh1 is required to activate Svp in L1 and Erm in L3, neither of which are HDTF, though Erm has a significant function in L3 axon targeting (Peng et al., 2018 [DOI](#)). This suggests that the primary HDTF can activate non-HDTFs to initiate neuron identity features. Recent work in *Drosophila* medulla found that a unique combination of TFs (a mix of HDTFs and non-HDTFs) is required to control neuron identity features (Özel et al., 2022 [DOI](#)). It would be important to dissect whether there is hierarchical expression and function within these TF combinations and to test whether HDTFs activate non-HDTFs.

Evolution can drive a coordinated increase in neuronal diversity and functional complexity. We hypothesize that there was an evolutionary path promoting increased neuronal diversity by the addition of primary HDTF Bsh expression. This is based on our finding that the loss of a single HDTF (Bsh) results in reduced lamina neuron diversity (only L1-3), which may represent a simpler ancestral brain. A similar observation was described in *C. elegans* where the loss of a single terminal selector caused two different neuron types to become identical, which was speculated to be the ancestral ground state (Arlotta and Hobert, 2015 [DOI](#); Cros and Hobert, 2022 [DOI](#); Reilly et al., 2022 [DOI](#)), suggesting phylogenetically conserved principles observed in highly distinct species. An interesting possibility is that evolutionarily primitive insects, such as silverfish (Truman and Riddiford, 1999 [DOI](#)), lack Bsh expression and L4/L5 neurons, retaining only the core motion detection L1-L3 neurons. Our findings provide a testable model whereby neural circuits evolve more complexity by adding the expression of a primary HDTF (Figure 7E [DOI](#)).

Acknowledgements

We thank Stefan Abreo for technical assistance; Claude Desplan, Lawrence Zipursky, Makoto Sato, Markus Affolter, Richard Mann, Jing Peng, Cheng-Yu Lee, James Skeath, Cheng-Ting Chien for antibodies; Emily Nielson for the Fly Vision Box schematic; Claire Managan and Adam Taylor (Janelia Research Campus) for technical assistance and data processing; Lihi Zelnik-Manor and Pietro Perona (Caltech) for the original fly-tracking code; Ayanthi Bhattacharya and Sonia Sen for advice on DamID; and Claude Desplan, Vilaiwan Fernandes, Kristen Lee, Emily Heckman, Peter Newstein and Sarah Ackerman for comments on the manuscript. Stocks obtained from the Bloomington *Drosophila* Stock Center were used in this study.

Author contributions

CX: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Resources; Writing—original draft; Writing—review & editing; Visualization; Project administration

TBR: Validation; Formal analysis; Investigation; Resources; Writing—review & editing

EMR: Resources; Writing—review & editing

MBR: Methodology; Software; Formal analysis; Data curation; Writing—review & editing

CQD: Methodology; Resources; Writing—review & editing; Visualization; Supervision; Project administration; Funding acquisition

Declaration of interests

The authors declare no competing financial or non-financial interests.

Data Availability Statement

All resources will be provided upon request. DamID data in this publication have been deposited in NCBI's GEO and are accessible through GEO Series accession number GSE246726.

Figure legends

Materials and Methods




Figure 2—figure supplement 1.

R27G05-GAL4 is turned on in all LPCs and turned off in lamina neurons.

(A-A''') R27G05-Gal4 drives GFP expression in all LPCs and lamina neurons (white dash circle) which are labeled by Dac at 96 hours after larval hatching (ALH). Hth labels LPCs next to lamina furrow (Piñeiro et al., 2014). Here and below, scale bar, 10 μ m.

(B, C) R27G05-Gal4 is turned off in lamina neurons. Activating R27G05-Gal4 at 29°C at the beginning of wandering L3 for 20 hours (by inactivating Gal80ts) reveals continued expression of GFP; in contrast, activating R27G05-Gal4 at 29°C at 1d APF for 20 hours reveals little GFP, showing that R27G05 is not expressed in pupal stages containing postmitotic lamina neurons. Genotype: R27G05-Gal4>UAS-myrGFP, tubP-Gal80[ts].

(D, E) Although R27G05-Gal4 is turned off by 1d APF, GFP persists in lamina neuron cell bodies until 66h APF and in their neurites until 75h APF. Genotype: R27G05-Gal4>UAS-myrGFP.




Figure 2—figure supplement 2.

Bsh-KD or Bsh-KO in LPCs affects L4/L5 neuronal fate.

(A-D) Bsh-KD in LPCs removes most L4 (Bsh/Ap) and L5 (Bsh/Pdm3) neuron markers of 3-day adult flies. The Ap expression in L5 is caused by the Gal4 driver line but is irrelevant here. (C and D) The number of Pdm3+ (C) or Ap+/Pdm3- (D) cell bodies in a single slice in control (A) and Bsh-KD (B). Here and below, scale bar, 10 μ m, white circle outlines the region of L4 and L5 cell bodies. (E-I) Bsh-KO in LPCs (27G05-Gal4>UAS-Cas9, UAS-Bsh-sgRNAs) decreases Ap and Pdm3 expression in the lamina of 3-4-day old pupa. (G-I) The number of Bsh+ (G), Pdm3+ (H), or Ap+/Pdm3- (I) cell bodies in a single slice in control (E) and Bsh-KO (F).

Data are presented as mean $\pm$ SEM. Each dot represents each brain. n=6 brains in (C) and (D), n=5 brains in (G), (H), and (I). Each dot represents each brain. ***P<0.001, unpaired t-test.




Figure 2—figure supplement 3.

Bsh-KD in L4 neurons results in loss of detectable Bsh between 2d and 3d APF.

(A-C) Bsh expression is detected at 2d APF following Bsh-KO in L4 neurons. Genotype: 31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs.

(D-F) Bsh expression is absent at 3d APF following Bsh-KD in L4 neurons. Genotype: 31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs. GFP labels L4 neurons. White arrowhead indicates L4 cell body lacking Bsh expression.

Data are presented as mean $\pm$ SEM. ns= not significant, *** P <0.001, unpaired t-test, n = 6 brains; each dot represents one brain. Scale bar, 10 μ m.




Figure 2—figure supplement 4.

Bsh is not required in L4 neurons to maintain L4 neuronal fate.

(A-B'') L1 marker Svp and L2 marker Bab2 are not detected in L4 cell bodies when knocking out Bsh in L4 (31C06-AD, 34G07-DBD>UAS-Cas9, UAS-Bsh-sgRNAs). Here and below, L4 is labeled by GFP (31C06-AD, 34G07-DBD> UAS-myrGFP). White outline, L4 cell body. Scale bar, 10 μ m. (C-D'') L3 marker Erm is not detected in L4 cell bodies when knocking out Bsh in L4.

(E-F'') Bsh becomes absent in most L4 while Ap expression remains normal when knocking out Bsh L4.

(G-H'') L5 marker Pdm3 is not detected in L4 cell bodies when knocking out Bsh in L4.

(I) The percentage of each lamina neuron marker in L4 neurons (GFP+). ***P<0.001, ns= not significant, unpaired t-test, n= 5 or 6 brains, each dot represents each brain, data are presented as mean $\pm$ SEM.




Figure 3—figure supplement 1.

Loss of Bsh in LPCs transforms L5 neurons into L1 neurons.

(A-A'') Bsh-LexA drives LexAop-myrGFP expression in L5 neurons but not L4 at 20h APF. Older GFP cell bodies express L5 marker Pdm3. Ap labels L4 neurons. The white circle outlines GFP+ cells. scale bar, 10 μ m.




Figure 3—figure supplement 2.

Ap-KD eliminates Ap initiation and maintenance in L4 neurons.

(A-C) Ap initiation in L4 neurons is eliminated at 19-20h APF when knocking down Ap (27G05-GAL4>UAS-Ap-RNAi). The top Bsh+ labels L4 and the bottom Bsh+ labels L5. Ap and Pdm3 are expressed in older L4 and older L5 neurons respectively. The Ap expression in L5 neurons is caused by the Gal4 driver line but is irrelevant here. Scale bar, 10 μ m. (C) The percentage of Ap expression in L4 neurons (top Bsh+ row) in control (A) and Ap-KD (B).

(D-F) Ap expression remains undetectable in adults though Bsh expression remains normal. Bsh is unlikely able to reinitiate Ap expression if normal Ap initiation is lost. Scale bar, 10 μ m. (F) The percentage of Ap expression (Ap+/Pdm3-) in L4 neurons (Bsh+/Pdm3-) in control (D) and Ap-KD (E).

Data are presented as mean $\pm$ SEM. Each dot represents each brain. $n=6$ brains in (C) and (F). *** $P<0.001$, unpaired t-test.




Figure 3—figure supplement 3.

Ap is not required to suppress other lamina neuron markers in L4 neurons.

(A-H'') L1 (Svp, Zfh1), L2 (Bab2), L3 (Erm, Zfh1), and L5 (Pdm3) markers are not expressed in L4 neurons (top Bsh+ row) at 3-4d APF when knocking down Ap (27G05-GAL4>UAS-Ap-RNAi). Scale bar, 10 μ m.

(I) The percentage of each lamina neuron marker in L4 neurons (top row of Bsh+). *** $P < 0.001$, ns = not significant, unpaired t-test, $n = 6$ brains, each dot represents each brain, data are presented as mean $\pm$ SEM.




Figure 4—figure supplement 1.

Zfh1 expression and epistasis with Bsh.

(A-D) Zfh1-KD in LPCs significantly reduces Zfh1 expression in all LPCs and lamina neurons at 12h APF. The white line circle outlines LPCs. The yellow line circle outlines lamina neurons. White dash circle outline Zfh1+ glia. Scale bar here and below, 10 μ m. Genotype: R27G05-Gal4>UAS-Zfh1-RNAi. Quantification: Zfh1 signal of three cell bodies from each brain and normalized Zfh1 signal by setting the highest Zfh1 signal as 100.

(E-H) The total number of Ap+ and Pdm3+ cell bodies is decreased when knocking down Zfh1 in LPCs. The number of Bsh+ L4 and L5 neurons is decreased when knocking down Zfh1 in LPCs. Scale bar, 10 μ m. (G) The total number of Ap+ and Pdm3+ cell bodies in a single slice. (H) The number of Bsh+ L4 and L5 neurons in a single slice. Genotype: R27G05-Gal4>UAS-Zfh1-RNAi.

(I-K) The number of lamina neurons is decreased when knocking down Zfh1 in LPCs. Scale bar, 10 μ m. (K) The number of Elav+ cells in a single slice. Genotype: R27G05-Gal4>UAS-Zfh1-RNAi. (L-N) Zfh1 is ectopically expressed in GFP+ neurons. GFP (Bsh-LexA>LexAop-myrGFP) labels LPCs and L5 neurons in control. The white line circle outlines GFP+ neurons. GFP+ LPCs are TII+ (yellow arrowhead). Scale bar, 10 μ m. (N) The percentage of Zfh1+ cells in GFP+ neurons (Elav+) in control (L) and Bsh-KD (M). We divided GFP+ neurons into three domains with equal width from newborn to older neurons.

Data are presented as mean $\pm$ SEM. Each dot represents a brain. n=5 brains per genotype. ** P <0.01, *** P <0.001, unpaired t-test.




Figure 5—figure supplement 1.

UAS-Bsh:Dam expressed by 31C06-Gal4 is functional and specifically expressed in L4 neurons; Bsh:Dam shows a binding peak in *ap* locus but not *pdm3* or *zfh1* in L4 neurons.

(A) Schematic of the system used to restrict Bsh:Dam expression spatially to L4 (using the 31C06-Gal4 driver) and temporally to 46-76h APF (using the temperature-sensitive Gal4 inhibitor, Gal80ts; red T-bar). Note the high expression of GFP from the first ORF (thick green arrow), and low expression of Bsh or Bsh:Dam following the readthrough of the stop codons prior to the second ORF (thin green arrow).

(B-B'') 31C06-Gal4, Gal80ts, UAS-Bsh:Dam is expressed in Ap⁺ L4 neurons when conditionally expressed from 46h-76h APF. Scale bar, 10 μ m.

(C-E) Bsh:Dam retains Bsh function and can partially rescue Ap and Pdm3 expressions in Bsh-null mutant flies (Bsh-null mutant; R27G05-Gal4>UAS-myGFP-Bsh:Dam). Note that Bsh:Dam expression from the second ORF is extremely low to avoid toxicity and is not detected by immunostaining. The white circle outlines the cell body area of lamina neurons. (E) Quantification of the number of Ap⁺ or Pdm3⁺ or Bsh⁺ cell bodies within z stack of 2.88 μ m in control (C-C') and Bsh:Dam (D-D'). Scale bar, 10 μ m. Data are presented as mean $\pm$ SEM. Each dot represents a brain. Control: n=5 brains; Bsh:Dam: n=4 brains. ns= not significant, **P<0.001, unpaired t-test.

(F) Bsh:Dam shows strong signals in *ap* gene but not *pdm3* or *zfh1* gene in L4 neurons during the synapse formation window. The Dam alone signal indicates the open chromatin region in L4 neurons. The y axes of Bsh:Dam signal represent log2 (Bsh:Dam/Dam) scores. Bsh peaks in L4 neurons were generated using find_peaks (FDR<0.01; min_quant=0.9) and peaks2genes.




Figure 6—figure supplement 1.

DIP-β expression is disrupted when knocking down Ap in L4 neurons.

(A-C) Ap-KD (31C06-AD, 34G07-DBD>UAS-RSR, 79C23-S-GS-rst-stop-rst-smFPV5-2a-GAL4, UAS-Ap-shRNA; see methods for an explanation of this genotype) significantly reduced Ap expression in L4 neurons at 2d APF. The white line circle outlines lamina neuron cell bodies. Scale bar, 10 μ m. (C) Quantification of Ap signal in the cell bodies of L4 neurons in control (A) and Ap-KD (B). We averaged the Ap signal of three cell bodies from each brain and normalized the Ap signal by setting the highest Ap signal as 100. ***P<0.001, unpaired t-test, n= 6 brains, each dot represents each brain, data are presented as mean $\pm$ SEM.

(D-F') DIP-β expression is disrupted in L4 neurons in Ap-KD (31C06-AD, 34G07-DBD>UAS-RSR and 79C23-S-GS-rst-stop-rst-smFPV5-2a-GAL4) and UAS-Ap-shRNA. Note that the DIP-β signal in the proximal lamina belongs to L4 in control (arrow, L4), whereas the DIP-β signal in the distal lamina belongs to unknown cells. Scale bar, 10 μ m.

(E-G) Quantification of DIP-β fluorescence intensity along the long axis of lamina cartridges (see white lines in (D) and (F)) from the distal dash line to the proximal dash line. Significantly reduced fluorescence intensity is observed in the proximal lamina (75%–100% distance, marked by the red bar in (E) and (G)) of Ap-KD flies compared to the control. ***P<0.001, unpaired t-test, n=5 brains, each line represents each brain.




Figure 7—figure supplement 1.

Bsh-KD in LPCs results in loss of Bsh in the adult fly.

(A-A') Control flies have Bsh⁺ neurons in the adult L4 neurons (arrow). Genotype: w⁺, R27G05-Gal4>UAS-mcherry-RNAi, tubP-Gal80[ts]. Neuropil marker Brp (blue), Bsh neurons (green). Scale bar, 10 μm.

(B-B') Bsh-KD in LPCs results in the loss of Bsh⁺ neurons in adult L4 neurons. Neuropil marker Brp (blue), there are no Bsh⁺ neurons, although there is background staining (green). Scale bar, 10 μm. Genotype: w⁺, R27G05-Gal4>UAS-Bsh-RNAi, tubP-Gal80[ts].

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (<i>D. melanogaster</i>)	10xUAS-IVS-myristoylated-GFP	Bloomington Drosophila Stock Center	RRID: BDSC_32199	w[1118]; P{y[+7.7] w[+mC]=10XUAS-IVS-myr::GFP}su(Hw) attP5
strain, strain background (<i>D. melanogaster</i>)	<i>R27G05GAL4</i>	Bloomington Drosophila Stock Center	RRID: BDSC_48073	w[1118]; P{y[+7.7] w[+mC]=GMR27G05-GAL4}attP2
strain, strain background (<i>D. melanogaster</i>)	UAS-Bsh-RNAi	Bloomington Drosophila Stock Center	RRID: BDSC_29336	y[1] v[1]; P{y[+7.7] v[+t1.8]=TRiP.JF02498}attP2
strain, strain background (<i>D. melanogaster</i>)	Bsh-LexA	Bloomington Drosophila Stock Center	RRID: BDSC_52834	w[1118]; P{y[+7.7] w[+mC]=GMR64B07-lexA}attP40
strain, strain background (<i>D. melanogaster</i>)	13xLexAop-IVS-myr::GFP	Bloomington Drosophila Stock Center	RRID: BDSC_32211	y[1] w[*] P{y[+7.7] w[+mC]=13XLexAop2-IVS-myr::GFP}su(Hw) attP8

strain, strain background (<i>D. melanogaster</i>)	UAS-Cas9 (attp40)	Bloomington Drosophila Stock Center	RRID: BDSC_58985	P{ry[+t7.2]=hsFLP}12, y[1] w[*]; P{y[+t7.7]w[+mC]=UAS-Cas9.P2}attP40
strain, strain background (<i>D. melanogaster</i>)	UAS-Cas9 (attp2)	Bloomington Drosophila Stock Center	RRID: BDSC_58986	P{ry[+t7.2]=hsFLP}12, y[1] w[*]; P{y[+t7.7]w[+mC]=UAS-Cas9.P2}attP2/TM6B, Tb[1]
strain, strain background (<i>D. melanogaster</i>)	UAS-Bsh-sgRNA	Vienna Drosophila Resource Center	VDRC 341537	P{ry[+t7.2]=hsFLP}1, y[1] w[1118]; P{y[+t7.7]w[+mC]=HD_CF000611}attP40/CyO-GFP
strain, strain background (<i>D. melanogaster</i>)	UAS-Ap-RNAi	Bloomington Drosophila Stock Center	RRID: BDSC_41673	y[1] sc[*] v[1] sev[21]; P{y[+t7.7]v[+t1.8]=TRiP.HMS02207}attP2
strain, strain background (<i>D. melanogaster</i>)	20xUAS-RSR.PEST	Bloomington Drosophila Stock Center	RRID: BDSC_55795	w[1118]; P{y[+t7.7]w[+mC]=20XUAS-RSR.PEST}attP2
strain, strain background (<i>D. melanogaster</i>)	UAS-Ap-shRNA	Vienna Drosophila Resource Center	VDRC 330463	P{VSH330463}attP40
strain, strain background (<i>D. melanogaster</i>)	UAS-Bsh-HA	Bloomington Drosophila Stock Center	RRID: BDSC_83310	y[1] w[1118]; PBac{y[+mDint2]w[+mC]=UAS-bsh.ORF.3xHA.GW}VK00018/CyO

<i>melanogaster</i>)				
strain, strain background (D. <i>melanogaster</i>)	tubP- GAL80[ts]	Bloomington Drosophila Stock Center	RRID: BDSC_7017	w[*]; P{w[+mC]=tubP- GAL80[ts]}2/TM2
strain, strain background (D. <i>melanogaster</i>)	UAS-Zfh1- RNAi	Vienna Drosophila Resource Center	VDRC 103205	P{KK109931}VIE -260B
strain, strain background (D. <i>melanogaster</i>)	Brp-rst- stop-rst- smFPV5- 2a-GAL4	Jing Peng (Harvard Medical School)		W; Bl/CyO-GFP; Brp-rst-stop-rst- smFPV5-2A- Gal4/tm6b
strain, strain background (D. <i>melanogaster</i>)	31C06AD (III), 34G07DBD (III)	Gift from Janelia Research Campus (Tuthill et al., 2013b)		w; UAS- FLP/CyO; 31c06A, 34G07DBD/tm6b
strain, strain background (D. <i>melanogaster</i>)	31C06- Gal4, UAS- myristoylat ed- tdTomato	Gift from Lawrence Zipursky		;Bl/CyO; 31c06- Gal4, UAS- myristoylated- tdTomato/tm6b
strain, strain background (D. <i>melanogaster</i>)	VALIUM2 0-mCherry	Bloomington Drosophila Stock Center	RRID: BDSC_35785	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=VALIU M20- mCherry}attP2

strain, strain background (<i>D. melanogaster</i>)	Bsh-ORF-3XHA (86Fb)	FlyORF Webshop	Cat#F000054	M{UAS-bsh. ORF. 3xHA. GW}ZH-86Fb
strain, strain background (<i>D. melanogaster</i>)	flyORF-TaDa	Bloomington Drosophila Stock Center	RRID: BDSC_91637	w[1118]; M{RFP[3xP3.PB]w[+mC]=FlyORF-TaDa}ZH-86Fb
strain, strain background (<i>D. melanogaster</i>)	hs-FlpD5; FlyORF-TaDa	Bloomington Drosophila Stock Center	RRID: BDSC_91638	w[1118]; P{y[+t7.7]w[+mC]=hs-FLPD5}attP40; M{RFP[3xP3.PB]w[+mC]=FlyORF-TaDa}ZH-86Fb
strain, strain background (<i>D. melanogaster</i>)	Bsh-null mutant	Gift from Makoto Sato		
strain, strain background (<i>D. melanogaster</i>)	Bsh-TaDa	This paper		w; +/CyO; UAS-GFP-Bsh-DAM/tm6b; See Generating Bsh-TaDa fly line in methods
Antibody	Chicken polyclonal	Abcam	Cat#ab13970, RRID_300798	Anti-GFP (1:1000)
Antibody	Rabbit polyclonal	Gift from Claude Desplan (Özel et al., 2021)		Anti-Bsh (1:1000)

Antibody	Guinea pig polyclonal	Gift from Lawrence Zipursky (Tan et al., 2015) & Makoto Soto (Hasegawa et al., 2011)		Anti-Bsh (1:1000)
Antibody	Rabbit polyclonal	Gift from Markus Affolter (Bieli et al., 2015)		Anti-Apterous (1:1000)
Antibody	Rat monoclonal	Gift from Cheng-Ting Chien (Chen et al., 2012)		Anti-Pdm3 (1:200)
Antibody	Rabbit polyclonal	Gift from Cheng-Yu Lee (Janssens et al., 2014)		Anti-Erm (1:100)
Antibody	Rat monoclonal	Gift from Jing Peng (Santiago et al., 2021)		Anti-Erm (1:70)
Antibody	Mouse monoclonal	Developmental Studies Hybridoma Bank	Cat#Seven-up D2D3, RRID_2618079	Anti-Svp (1:10)
Antibody	Rabbit polyclonal	Gift from James Skeath (Tian et al., 2004)		Anti-Zfh1 (1:1000)
Antibody	Rabbit polyclonal	Asian Distribution Center for Segmentation Antibodies	Code#812	Anti-Tailless (1:200)

Antibody	Mouse monoclonal	Developmental Studies Hybridoma Bank	Cat#mAbdac1-1, RRID: AB_579773	Anti-Dac (1:100)
Antibody	Mouse monoclonal	Developmental Studies Hybridoma Bank	Cat#Elav-9F8A9, RRID: AB_528217	Anti-Elav (1:200)
Antibody	Mouse monoclonal	Developmental Studies Hybridoma Bank	Cat#Rat-Elav-7E8A10 anti-elav, RRID: AB_528218	Anti-Elav (1:50)
Antibody	Mouse monoclonal	Developmental Studies Hybridoma Bank	Cat#24B10, RRID: AB_528161	Anti-Chaoptin (1:20)
Antibody	Auinea pig polyclonal	Gift from Matthew Pecot Lab (Xu et al., 2019)		Anti-DIP- β (1:300)
Antibody	Mouse monoclonal	Bio-Rad Laboratories	Cat#MCA1360A647, RRID: AB_770156	Anti-V5-TAG:Alexa Fluor® 647 (1:300)
Antibody	Rat monoclonal	Sigma Aldrich	Cat#12158167001, RRID: AB_390915	Anti-HA (1:100)
Antibody	Guinea pig polyclonal	Gift from Richard Mann (Casares and Mann, 1998)		Anti-Hth (1:2000)
Antibody	Mouse monoclonal	Developmental Studies Hybridoma Bank	Cat#nc-82, RRID: AB_2314866	Anti-Brp (1:50)

Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#712-545-153, RRID: AB_2340684	Alexa Fluor 488 anti-rat (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#703-545-155, RRID: AB_2340375	Alexa Fluor 488 anti-chicken (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#706-545-148, RRID: AB_2340472	Alexa Fluor 488 anti-guinea pig (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#711-545-152, RRID: AB_2313584	Alexa Fluor 488 anti-rabbit (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#715-545-150, RRID: AB_2340846	Alexa Fluor 488 anti-mouse (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#715-295-151, RRID: AB_2340832	Rhodamine Red-X anti-mouse (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#712-295-153, RRID: AB_2340676	Rhodamine Red-X anti-rat (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#711-295-152, RRID: AB_2340613	Rhodamine Red-X anti-rabbit (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#706-295-148, RRID: AB_2340468	Rhodamine Red-X donkey anti-guinea pig (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#711-605-152, RRID: AB_2492288	Alexa Fluor 647 donkey anti-rabbit (1:400)

Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#715-605-151, RRID: AB_2340863	Alexa Fluor 647 donkey anti-mouse (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#706-605-148, RRID: AB_2340476	Alexa Fluor 647 anti-guinea pig (1:400)
Antibody	Donkey polyclonal	Jackson ImmunoResearch Lab	Cat#712-605-153, RRID: AB_2340694	Alexa Fluor 647 anti-rat (1:400)
sequence-based reagent	Oligonucleotide	Integrated DNA technologies		DamID Adaptor (top strand): CTAATACGACT CACTATAGGGC AGCGTGGTCGC GGCCGAGGA
sequence-based reagent	Oligonucleotide	Integrated DNA technologies		DamID Adaptor (bottom strand): TCCTCGGCCG
sequence-based reagent	Oligonucleotide	Integrated DNA technologies		DamID Primer for PCR: GGTCGCGGCCGA GGATC
commercial assay or kit	QIAamp DNA Micro Kit	Qiagen	Cat#56304	
commercial assay or kit	PCR Purification Kit	Qiagen	Cat#28104	
chemical compound, drug	EDTA	Sigma-Aldrich	Cat#E6758	

chemical compound, drug	DpnI and CutSmart buffer	NEB	Cat#R0176S	
chemical compound, drug	DpnII and DpnII buffer	NEB	Cat#R0543S	
chemical compound, drug	MyTaq HS DNA Polymerase	Bioline	Cat#BIO-21112	
chemical compound, drug	AlwI	NEB	Cat#R0513S	
chemical compound, drug	RNase A (DNase free)	Roche	Cat#11119915001	
chemical compound, drug	T4 DNA ligase and 10x buffer	NEB	Cat#M0202S	
software, algorithm	Fiji	(Schindelin et al., 2012)	https://imagej.nih.gov/ij/download.html	
software, algorithm	FastQC (v0.11.9)	The Babraham Bioinformatics group	https://www.bioinformatics.babraham.ac.uk/projects/download.html#fastqc	
software, algorithm	MATlab	Mathworks	https://www.mathworks.com/products/matlab.html	
software, algorithm	damidseq_pipeline	(Marshall and Brand, 2015)	https://owenjm.github.io/damidseq_pipeline/	

software, algorithm	Bowtie2 (v2.4.5)	(Langmead and Salzberg, 2012)	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml	
software, algorithm	IGV (v.2.13.2)	(Robinson et al., 2011)	https://software.broadinstitute.org/software/igv/download	
software, algorithm	SAMtools (v1.15.1)	(Li et al., 2009)	http://www.htslib.org/download/	
software, algorithm	deepTools (v3.5.1)	(Ramírez et al., 2016)	https://deeptools.readthedocs.io/en/develop/content/installation.html	
software, algorithm	Find_peaks	(Marshall et al., 2016)	https://github.com/owenjm/find_peaks	

Contact for reagent and resource sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact Chundi Xu (cxu3@uoregon.edu) and Chris Doe (cdoe@uoregon.edu).

Experimental model and subject details

All flies were reared at 25°C on standard cornmeal fly food, unless otherwise stated. For all RNAi and shRNA knockdown experiments, crosses are kept at 25°C and their progeny are kept at 28.5°C with 16:8 hours light-dark cycle from the embryo stage until dissection. For all Gal80ts experiments, crosses are kept at 18°C and progenies are kept at 29°C at the desired time.

Method details

Animal collections

For the Bsh-misexpression experiment, crosses were reared at 25°C in collection bottles fitted with 3.0% agar apple juice caps containing plain yeast paste. Embryos were then collected on 3.0% agar apple juice caps with plain yeast paste for 4 hours. The collected embryos were moved to 18°C until 72 hours after larval hatching (ALH). The larvae at 72h ALH were moved to 29°C until 58 hours after pupal formation.

For the experiment R27G05-GAL4>UAS-myrGFP, tubP-GAL80[ts], the progeny is kept in 18°C from embryo and moved to 29.2°C at early wandering L3 or 1d APF for 20 hours.

For the behavioral experiments, the progeny is kept in 18°C from embryo and moved to 29°C with 16:8 hours light-dark cycle from the early larval stage until behavioral tests. Male flies at 2-5days after the eclosion at 29°C were used for the Fly Vision Box experiments.

Immunohistochemistry

Fly brains were dissected in Schneider's medium and fixed in 4% paraformaldehyde in phosphate buffered saline (PBS) for 25 min. After fixation, brains were quickly washed with PBS with 0.5% Triton X-100 (PBT) and incubated in PBT for at least 2 hr at room temperature. Next, samples were incubated in blocking buffer (10% normal donkey serum, 0.5% Triton X-100 in PBS) overnight at 4°C. Brains were then incubated in primary antibody (diluted in blocking buffer) at 4°C for at least two nights. Following primary antibody incubation, brains were washed with PBT. Next, brains were incubated in secondary antibody (diluted in blocking buffer) at 4°C for at least one day. Following secondary antibody incubation, brains were washed with PBT. Finally, brains were mounted in SlowFade Gold antifade reagent (Thermo Fisher Scientific, Waltham, MA). Images were acquired using a Zeiss 800 confocal and processed with Image J and Adobe Photoshop.

Knocking down HDTF in neurons

Genotype for knocking down Ap specifically in L4 neurons: 31C06-AD, 34G07-DBD>UAS-RSR:PEST, 79C23-S-GS-RSRT-Stop-RSRT-smFP:V5-2a-GAL4, UAS-Ap-shRNA. The genetic element 79C23-S is a bacterial artificial chromosome that encodes the *Brp* gene (Chen et al., 2014 [DOI](#)). 31C06-AD, 34G07-DBD drives the expression of R recombinase (RSR) in L4 neurons and RSR removes the stop codon from 79C23-S-GS-RSRT-Stop-RSRT-smFP:V5-2a-GAL4. Therefore Brp:smFP:V5-2a-GAL4 is transcribed and translated in L4 neurons into two proteins, Brp:smFP:V5 and GAL4. GAL4 together with 31C06-AD, 34G07-DBD drives the expression of Ap-shRNA to knockdown Ap. There is no continuous Ap-shRNA expression when using 31C06-AD, 34G07-DBD to drive expression of Ap-shRNA directly because 31C06-AD, 34G07-DBD depends on Ap.

Generating Bsh-TaDa fly line

Bsh-TaDa fly line was generated using the FlyORF-TaDa system described in (Aughey et al., 2021 [DOI](#)). Homozygous *hs-FlpD5*; FlyORF-TaDa virgin females were crossed to males from Bsh-ORF-3xHA line. Progeny (larval stage) were heat shocked at 37°C for 60 min, once per day. After eclosion, F1 male flies were crossed to MKRS/TM6B virgin females. F2 males and virgin females with the correct eye phenotype (*w*⁻; 3xP3-dsRed2⁺) were crossed to establish a balanced stock.

TaDa in L4 neurons at the time of synapse formation

Homozygous *tubP-GAL80[ts]*; 31C06-Gal4, UAS-myristoylated-tdTomato males were crossed to homozygous virgin females (FlyORF-TaDa line for Dam; Bsh-TaDa line for Bsh:Dam). Crosses were reared at 18°C. To perform TaDa in L4 neurons during synapse formation window, we collected pupae with the age of 46h APF and moved them to 29°C to activate 31C06-Gal4 for 24 hours (Figure 5—figure supplement 1 [DOI](#)). Then lamina were collected (age equivalent at 25°C: 76h APF) in cold PBS within one hour and stored at -20°C immediately until sufficient lamina were collected – for each group, about 70 lamina from 35 pupae. The TaDa experimental pipeline was followed according to (Marshall et al., 2016 [DOI](#)) with a few modifications. Briefly, DNA was extracted using a QIAamp DNA Micro Kit (Qiagen), digested with DpnI (NEB) overnight, and cleaned up using Qiagen PCR purification columns. DamID adaptors were ligated using T4 DNA ligase (NEB) followed by DpnII (NEB) digestion for 2 h and PCR amplification using MyTaq HS DNA polymerase (Bioline). The samples were sequenced on the NovaSeq at 118 base pairs and 27-33 million single end reads per sample.

Bioinformatic analysis

The TaDa sequencing data was analyzed as described previously (Sen et al., 2019 [DOI](#)). Briefly, each file was assessed for quality using FastQC (v0.11.9). The damidseq pipeline was run to generate Dam bedgraph files, Log2 ratio bedgraph files (Bsh:Dam/Dam), Dam bam files and Bsh:Dam bam files as described previously (Marshall and Brand, 2015 [DOI](#)). The bedgraph files were used for data visualization on IGV (v2.13.2) (Robinson et al., 2011 [DOI](#)). The Log2 ratio bedgraph files

(Bsh:Dam/Dam) were used for calling Bsh peaks in L4 neurons using `find_peaks` (FDR<0.01; `min_quant=0.9`) and the generated peak files were used for calling genes using `peaks2genes` (https://github.com/owenjm/find_peaks). The gene list of Bsh peaks (score>2.5) in L4 neurons were then combined with L4 scRNAseq data (Jain et al., 2022) (normalization number>2). The bam files of Dam and Bsh:Dam were used to create sorted bam files and indexed bam files (bam.bai) using SAMtools (v1.15.1) (Li et al., 2009). Sorted bam files and indexed bam files were then computed using the `multiBamSummary` and `plotCorrelation` functions of deepTools (v3.5.1) (Ramírez et al., 2016) for correlation coefficients between biological replicates.

Behavioral experiments

The Fly Vision Box apparatus was developed at the Janelia Research Campus as a high-throughput assay integrating several tests of visually guided behavior for flies in tubes. Groups of flies (usually 10-15) are placed in the clear acrylic tubes, inside a temperature-controlled box. The box contains 6 tubes, each with a strip of green LEDs (patterns of 4 pixels on / 4 pixels off moving at the indicated temporal frequency or contrast) lining one wall, for the motion vision experiments, and a single green and single UV LED on each end of each corridor, for the phototaxis and spectral-preference assays. During the spectral preference task, a green LED is illuminated (at level 10) at the end of each tube, while at the opposite end of the tubes a UV LED is illuminated at increasing brightness levels. The tubes are capped with a polished acrylic plug that is transparent. 4 small (pager) motors are mounted on the corners of the box to provide a mechanical startle in between trials. A camera mounted above the box records fly movements at 25 frames/s. The camera is fitted with an Infrared-passing filter and the tubes are suspended above an Infrared backlight.

In visual motion response assay, stimulus with different temporal frequency (0, 0.5, 2, 5, 10, 20, 42 HZ) or 5Hz temporal frequency, but with the indicated contrast level were given. The flies walk against the direction of motion, quantified with a Direction Index. The phototaxis behavior measures the movement of flies towards UV or green LED with the indicated (relative) intensity level at the end of the tubes. The Direction Index is shown integrated over each trial length. In spectral preference task, for lower UV levels, flies walk towards the green LED, but walk towards the UV LED at higher UV levels.

A full experiment lasts ~30 minutes and consists of 44 conditions, presented in short blocks, during which a series of conditions (e.g. the 4 LED settings for the phototaxis assay) are presented twice (for Phototaxis) or 4 times each for the other tests, where on consecutive trials the stimulus is presented with opposite direction (for motion) or at the opposite ends of the tubes (for phototaxis and spectral preference). Each trial begins immediately after a 0.5 s mechanical startle by the motors.

The box is heated to 34°C. The heating and motorized startle ensure that flies are active throughout and respond to these stimuli.

Quantification and Statistical Analysis for behavioral experiments

Videos are tracked offline with custom code written in MATLAB. After tracking, the trajectories were analyzed on a per-frame basis to see whether flies were walking in one direction (either following the moving patterns or walking towards one end-cap LED) or the other. The behavior is summarized as a 'Direction Index' which is simply the difference between flies walking in one direction and the other (by convention the direction against the motion, or towards the illuminated LED, is the positive one) divided by the total number of flies in the tube. For example, with 12 flies in a tube, during one frame, 7 are moving in the positive direction and 4 are moving in the negative direction (and 1 is not moving), then for this frame, $DI = (7-4)/12 = 0.25$.

For the summarized plots in **Figure 7**, the Direction Index is averaged over the entire 10 second duration of each motion stimulus trial. For the phototaxis and spectral preference experiments, the DI is integrated (accumulated) across time and the data recorded for each trial is the peak (negative or positive) of this curve. For the 10 or 15 second trials, the maximum possible value is 10 or 15 ($10, 15 \text{ s} \times 1$). Individual data points in **Figure 7** represent the mean or peak DI metric summarizing all the flies in each tube, and the summary data are the mean and standard deviation across tubes.

The data summarized are part of a larger series of experimental conditions that include several other tasks that are largely redundant with those presented. Nevertheless, the flies experienced 44 different conditions, and so all data were used for a False Discovery Rate controlling procedure (Benjamini and Hochberg, 1995) with $q = 0.05$. The data from conditions not shown do not contain additional tests with statistically significant differences between the 2 tested genotypes.

Quantification of DIP- β fluorescence signal

Using image J, we quantified DIP- β signal in control and knockdown brains by measuring fluorescence signal along the long axis of lamina cartridges (see white lines in **Figure 6A**) from the distal dash line to the proximal dash line (3 cartridges per brain). Signal intensity values and cartridge lengths were converted to percentages by setting the highest intensity within each cartridge as 100% intensity and the full length of the cartridge as 100% distance. Statistical analysis using unpaired t tests was performed after setting uniform intervals (using the spline function on MATLAB) of 0.01% distance. We presented 20-100% as 0-100% to focus on L4 signal (the first 20% is DIP- β signal in LaWF2).

Quantification of Brp puncta in the distal and proximal regions of lamina cartridges

Using confocal microscopy, we generated z-stacks of the lamina down the long axis of lamina cartridges. Within each z-stack (i.e. each optic lobe) 20 cartridges in the center of the lamina were identified and the number of Brp puncta in their distal halves was counted. The top (distal edge; top dash line in **Figure 6E**) and bottom (proximal edge; bottom dash line in **Figure 6E**) of each cartridge was determined by the first section below L4 cell body and last section containing L4 neuron processes (myrtd::TOM), respectively. The midpoint of each cartridge was then identified as the section in between the top and bottom sections. Brp puncta were counted in the sections distal to the midpoint of each cartridge as distal Brp puncta and in the sections proximal to the midpoint of each cartridge as proximal Brp puncta. Genotypes were scored in a blind manner.

Quantification of L4 proximal neurite length

In the same cartridges as those chosen to quantify the Brp puncta, the number of z-stacks that contained L4 proximal neurites were calculated as a percentage of the entire long axis, and represented the values presented for proximal neurite length.

Statistical Analysis

Statistics were performed using a combination of Microsoft Excel, MATLAB (MathWorks) and Prism (GraphPad) software. Unpaired t-test was used, unless otherwise noted. Data are presented as mean $\pm$ SEM unless otherwise noted. A 95% confidence interval was used to define the level of significance. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, ns=not significant. All other relevant statistical information can be found in the figure legends.

References

- Arlotta P, Hobert O (2015) **Homeotic Transformations of Neuronal Cell Identities** *Trends Neurosci* **38**:751–762 <https://doi.org/10.1016/j.tins.2015.10.005>
- Aughey GN, Delandre C, McMullen JPD, Southall TD, Marshall OJ (2021) **FlyORF-TaDa allows rapid generation of new lines for in vivo cell-type-specific profiling of protein–DNA interactions in *Drosophila melanogaster*** *G3 GenesGenomesGenetics* **11** <https://doi.org/10.1093/g3journal/jkaa005>
- Aughey GN, Estacio Gomez A, Thomson J, Yin H, Southall TD (2018) **CATaDa reveals global remodelling of chromatin accessibility during stem cell differentiation in vivo** *eLife* **7** <https://doi.org/10.7554/eLife.32341>
- Bayraktar OA, Doe CQ (2013) **Combinatorial temporal patterning in progenitors expands neural diversity** *Nature* **498**:449–455 <https://doi.org/10.1038/nature12266>
- Benjamini Y, Hochberg Y (1995) **Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing** *J R Stat Soc Ser B Methodol* **57**:289–300 <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Benzer S (1967) **BEHAVIORAL MUTANTS OF *Drosophila* ISOLATED BY COUNTERCURRENT DISTRIBUTION** *Proc Natl Acad Sci* **58**:1112–1119 <https://doi.org/10.1073/pnas.58.3.1112>
- Casares F, Mann RS (1998) **Control of antennal versus leg development in *Drosophila*** *Nature* **392**:723–726 <https://doi.org/10.1038/33706>
- Chen C-K, Chen W-Y, Chien C-T (2012) **The POU-domain protein Pdm3 regulates axonal targeting of R neurons in the *Drosophila* ellipsoid body** *Dev Neurobiol* **72**:1422–1432 <https://doi.org/10.1002/dneu.22003>
- Chen Y, Akin O, Nern A, Tsui CYK, Pecot MY, Zipursky SL (2014) **Cell-type-Specific Labeling of Synapses In Vivo through Synaptic Tagging with Recombination** *Neuron* **81**:280–293 <https://doi.org/10.1016/j.neuron.2013.12.021>
- Cros C, Hobert O (2022) ***Caenorhabditis elegans* sine oculis/SIX-type homeobox genes act as homeotic switches to define neuronal subtype identities** *Proc Natl Acad Sci* **119** <https://doi.org/10.1073/pnas.2206817119>
- Cubelos B *et al.* (2010) **Cux1 and Cux2 Regulate Dendritic Branching, Spine Morphology, and Synapses of the Upper Layer Neurons of the Cortex** *Neuron* **66**:523–535 <https://doi.org/10.1016/j.neuron.2010.04.038>
- De Marco Garcia NV, Jessell TM. (2008) **Early Motor Neuron Pool Identity and Muscle Nerve Trajectory Defined by Postmitotic Restrictions in Nkx6.1 Activity** *Neuron* **57**:217–231 <https://doi.org/10.1016/j.neuron.2007.11.033>
- Deneris ES, Hobert O (2014) **Maintenance of postmitotic neuronal cell identity** *Nat Neurosci* **17**:899–907 <https://doi.org/10.1038/nn.3731>

Doe CQ (2017) **Temporal Patterning in the Drosophila CNS** *Annu Rev Cell Dev Biol* **33**:219–240 <https://doi.org/10.1146/annurev-cellbio-111315-125210>

Erclik T *et al.* (2017) **Integration of temporal and spatial patterning generates neural diversity** *Nature* **541**:365–370 <https://doi.org/10.1038/nature20794>

Fernandes VM, Chen Z, Rossi AM, Zipfel J, Desplan C (2017) **Glia relay differentiation cues to coordinate neuronal development in Drosophila** *Science* **357**:886–891 <https://doi.org/10.1126/science.aan3174>

Friedrich J, Sorge S, Bujupi F, Eichenlaub MP, Schulz NG, Wittbrodt J, Lohmann I (2016) **Hox Function Is Required for the Development and Maintenance of the Drosophila Feeding Motor Unit** *Cell Rep* **14**:850–860 <https://doi.org/10.1016/j.celrep.2015.12.077>

Gao S *et al.* (2008) **The Neural Substrate of Spectral Preference in Drosophila** *Neuron* **60**:328–342 <https://doi.org/10.1016/j.neuron.2008.08.010>

Gordon PM, Hobert O (2015) **A Competition Mechanism for a Homeotic Neuron Identity Transformation in C. elegans** *Dev Cell* **34**:206–219 <https://doi.org/10.1016/j.devcel.2015.04.023>

Hasegawa E, Kaido M, Takayama R, Sato M (2013) **Brain-specific-homeobox is required for the specification of neuronal types in the Drosophila optic lobe** *Dev Biol* **377**:90–99 <https://doi.org/10.1016/j.ydbio.2013.02.012>

Hobert O (2021) **Homeobox genes and the specification of neuronal identity** *Nat Rev Neurosci* **22**:627–636 <https://doi.org/10.1038/s41583-021-00497-x>

Hobert O (2016) **Terminal Selectors of Neuronal Identity** *Current Topics in Developmental Biology* :455–475 <https://doi.org/10.1016/bs.ctdb.2015.12.007>

Hobert O, Kratsios P (2019) **Neuronal identity control by terminal selectors in worms, flies, and chordates** *Curr Opin Neurobiol* **56**:97–105 <https://doi.org/10.1016/j.conb.2018.12.006>

Holguera I, Desplan C (2018) **Neuronal specification in space and time** *Science* **362**:176–180 <https://doi.org/10.1126/science.aas9435>

Howell K, White JG, Hobert O (2015) **Spatiotemporal control of a novel synaptic organizer molecule** *Nature* **523**:83–87 <https://doi.org/10.1038/nature14545>

Huang Z, Shilo BZ, Kunes S (1998) **A retinal axon fascicle uses Spitz, an EGF receptor ligand, to construct a synaptic cartridge in the brain of Drosophila** *Cell* **95**:693–703 [https://doi.org/10.1016/S0092-8674\(00\)81639-6](https://doi.org/10.1016/S0092-8674(00)81639-6)

Isaacson MD *et al.* (2023) **Isaacson MD, Eliason JLM, Nern A, Rogers EM, Lott GK, Tabachnik T, Rowell WJ, Edwards AW, Korff WL, Rubin GM, Branson K, Reiser MB. 2023. Small-field visual projection neurons detect translational optic flow and support walking control. doi:10.1101/2023.06.21.546024 Small-field visual projection neurons detect translational optic flow and support walking control** <https://doi.org/10.1101/2023.06.21.546024>

Jain S, Lin Y, Kurmangaliyev YZ, Valdes-Aleman J, LoCascio SA, Mirshahidi P, Parrington B, Zipursky SL (2022) **A global timing mechanism regulates cell-type-specific wiring programmes** *Nature* :1–7 <https://doi.org/10.1038/s41586-022-04418-5>

- Janssens DH, Komori H, Grbac D, Chen K, Koe CT, Wang H, Lee C-Y (2014) **Earmuff restricts progenitor cell potential by attenuating the competence to respond to self-renewal factors** *Dev Camb Engl* **141**:1036–1046 <https://doi.org/10.1242/dev.106534>
- Kitt MM, Tabuchi N, Spencer WC, Robinson HL, Zhang XL, Eastman BA, Lobur KJ, Silver J, Mei L, Deneris ES (2022) **An adult-stage transcriptional program for survival of serotonergic connectivity** *Cell Rep* **39** <https://doi.org/10.1016/j.celrep.2022.110711>
- Kratsios P, Pinan-Lucarré B, Kerk SY, Weinreb A, Bessereau J-L, Hobert O (2015) **Transcriptional Coordination of Synaptogenesis and Neurotransmitter Signaling** *Curr Biol* **25**:1282–1295 <https://doi.org/10.1016/j.cub.2015.03.028>
- Langmead B, Salzberg SL (2012) **Fast gapped-read alignment with Bowtie 2** *Nat Methods* **9**:357–359 <https://doi.org/10.1038/nmeth.1923>
- Leyva-Díaz E, Hobert O (2019) **Transcription factor autoregulation is required for acquisition and maintenance of neuronal identity** *Development* **146** <https://doi.org/10.1242/dev.177378>
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup (2009) **The Sequence Alignment/Map format and SAMtools** *Bioinforma Oxf Engl* **25**:2078–2079 <https://doi.org/10.1093/bioinformatics/btp352>
- Li X, Erclik T, Bertet C, Chen Z, Voutev R, Venkatesh S, Morante J, Celik A, Desplan C (2013) **Temporal patterning of Drosophila medulla neuroblasts controls neural fates** *Nature* **498**:456–462 <https://doi.org/10.1038/nature12319>
- Mangan S, Alon U (2003) **Structure and function of the feed-forward loop network motif** *Proc Natl Acad Sci* **100**:11980–11985 <https://doi.org/10.1073/pnas.2133841100>
- Mark B, Lai S-L, Zarin AA, Manning L, Pollington HQ, Litwin-Kumar A, Cardona A, Truman JW, Doe CQ (2021) **A developmental framework linking neurogenesis and circuit formation in the Drosophila CNS** *eLife* **10** <https://doi.org/10.7554/eLife.67510>
- Marshall OJ, Brand AH. (2015) **damidseq_pipeline: an automated pipeline for processing DamID sequencing datasets** *Bioinformatics* **31**:3371–3373 <https://doi.org/10.1093/bioinformatics/btv386>
- Marshall OJ, Southall TD, Cheetham SW, Brand AH (2016) **Cell-type-specific profiling of protein– DNA interactions without cell isolation using targeted DamID with next-generation sequencing** *Nat Protoc* **11**:1586–1598 <https://doi.org/10.1038/nprot.2016.084>
- Meinertzhagen IA, O’Neil SD (1991) **Synaptic organization of columnar elements in the lamina of the wild type in Drosophila melanogaster** *J Comp Neurol* **305**:232–263 <https://doi.org/10.1002/cne.903050206>
- Özel MN, Gibbs CS, Holguera I, Soliman M, Bonneau R, Desplan C (2022) **Coordinated control of neuronal differentiation and wiring by sustained transcription factors** *Science* **378** <https://doi.org/10.1126/science.add1884>
- Peng J, Santiago IJ, Ahn C, Gur B, Tsui CK, Su Z, Xu C, Karakhanyan A, Silies M, Pecot MY (2018) **Drosophila Fezf coordinates laminar-specific connectivity through cell-intrinsic and cell-extrinsic mechanisms** *eLife* **7** <https://doi.org/10.7554/eLife.33962>

- Piñeiro C, Lopes CS, Casares F (2014) **A conserved transcriptional network regulates lamina development in the Drosophila visual system** *Dev Camb* **141**:2838–2847 <https://doi.org/10.1242/dev.108670>
- Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke T (2016) **deepTools2: a next generation web server for deep-sequencing data analysis** *Nucleic Acids Res* **44**:W160–W165 <https://doi.org/10.1093/nar/gkw257>
- Reilly MB, Cros C, Varol E, Yemini E, Hobert O (2020) **Unique homeobox codes delineate all the neuron classes of C. elegans** *Nature* **584**:595–601 <https://doi.org/10.1038/s41586-020-2618-9>
- Reilly MB *et al.* (2022) **Widespread employment of conserved C. elegans homeobox genes in neuronal identity specification** *PLOS Genet* **18** <https://doi.org/10.1371/journal.pgen.1010372>
- Rivera-Alba M, Vitaladevuni SN, Mishchenko Y, Lu Z, Takemura S, Scheffer L, Meinertzhagen IA, Chklovskii DB, de Polavieja GG. (2011) **Wiring Economy and Volume Exclusion Determine Neuronal Placement in the Drosophila Brain** *Curr Biol* **21**:2000–2005 <https://doi.org/10.1016/j.cub.2011.10.022>
- Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP (2011) **Integrative genomics viewer** *Nat Biotechnol* **29**:24–26 <https://doi.org/10.1038/nbt.1754>
- Sakkou M, Wiedmer P, Anlag K, Hamm A, Seuntjens E, Ettwiller L, Tschöp MH, Treier M (2007) **A Role for Brain-Specific Homeobox Factor Bsx in the Control of Hyperphagia and Locomotory Behavior** *Cell Metab* **5**:450–463 <https://doi.org/10.1016/j.cmet.2007.05.007>
- Sanes JR, Zipursky SL (2010) **Design Principles of Insect and Vertebrate Visual Systems** *Neuron* **66**:15–36 <https://doi.org/10.1016/j.neuron.2010.01.018>
- Santiago C, Bashaw GJ (2014) **Transcription factors and effectors that regulate neuronal morphology** *Development* **141**:4667–4680 <https://doi.org/10.1242/dev.110817>
- Santiago IJ *et al.* (2021) **Drosophila Fezf functions as a transcriptional repressor to direct layer-specific synaptic connectivity in the fly visual system** *Proc Natl Acad Sci* **118** <https://doi.org/10.1073/pnas.2025530118>
- Schindelin J *et al.* (2012) **Fiji: an open-source platform for biological-image analysis** *Nat Methods* **9**:676–682 <https://doi.org/10.1038/nmeth.2019>
- Sen SQ, Chanchani S, Southall TD, Doe CQ (2019) **Neuroblast-specific open chromatin allows the temporal transcription factor, Hunchback, to bind neuroblast-specific loci** *eLife* **8** <https://doi.org/10.7554/eLife.44036>
- Serrano-Saiz E, Leyva-Díaz E, De La Cruz E, Hobert O. (2018) **BRN3-type POU Homeobox Genes Maintain the Identity of Mature Postmitotic Neurons in Nematodes and Mice** *Curr Biol* **28**:2813–2823 <https://doi.org/10.1016/j.cub.2018.06.045>
- Silies M, Gohl DM, Fisher YE, Freifeld L, Clark DA, Clandinin TR (2013) **Modular Use of Peripheral Input Channels Tunes Motion-Detecting Circuitry** *Neuron* **79**:111–127 <https://doi.org/10.1016/j.neuron.2013.04.029>
- Stefanakis N, Carrera I, Hobert O (2015) **Regulatory Logic of Pan-Neuronal Gene Expression in C. elegans** *Neuron* **87**:733–750 <https://doi.org/10.1016/j.neuron.2015.07.031>

- Takemura Shin-ya *et al.* (2015) **Synaptic circuits and their variations within different columns in the visual system of *Drosophila*** *Proc Natl Acad Sci* **112**:13711–13716 <https://doi.org/10.1073/pnas.1509820112>
- Tan L *et al.* (2015) **Ig Superfamily Ligand and Receptor Pairs Expressed in Synaptic Partners in *Drosophila*** *Cell* **163**:1756–1769 <https://doi.org/10.1016/j.cell.2015.11.021>
- Thor S, Thomas JB (1997) **The *Drosophila* islet Gene Governs Axon Pathfinding and Neurotransmitter Identity** *Neuron* **18**:397–409 [https://doi.org/10.1016/S0896-6273\(00\)81241-6](https://doi.org/10.1016/S0896-6273(00)81241-6)
- Tian X, Hansen D, Schedl T, Skeath JB (2004) **Epsin potentiates Notch pathway activity in *Drosophila* and *C. elegans*** *Dev Camb Engl* **131**:5807–5815 <https://doi.org/10.1242/dev.01459>
- Truman JW, Riddiford LM (1999) **The origins of insect metamorphosis** *Nature* **401**:447–452 <https://doi.org/10.1038/46737>
- Tuthill JC, Nern A, Holtz SL, Rubin GM, Reiser MB (2013) **Contributions of the 12 Neuron Classes in the Fly Lamina to Motion Vision** *Neuron* **79**:128–140 <https://doi.org/10.1016/j.neuron.2013.05.024>
- Xu C, Ramos TB, Marshall O, Doe CQ (2023) **Xu C, Ramos TB, Marshall O, Doe CQ. 2023. Notch signaling and Bsh homeodomain activity are integrated to diversify *Drosophila* lamina neuron types.** doi:<https://www.biorxiv.org/content/10.1101/2023.06.15.545141v1> *Notch signaling and Bsh homeodomain activity are integrated to diversify *Drosophila* lamina neuron types*
- Xu C *et al.* (2019) **Control of Synaptic Specificity by Establishing a Relative Preference for Synaptic Partners** *Neuron* **103**:865–877 <https://doi.org/10.1016/j.neuron.2019.06.006>
- Zhu Y, Nern A, Zipursky SL, Frye MA (2009) **Peripheral Visual Circuits Functionally Segregate Motion and Phototaxis Behaviors in the Fly** *Curr Biol* **19**:613–619 <https://doi.org/10.1016/j.cub.2009.02.053>

Article and author information

Chundi Xu

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon; Eugene, OR 97403, USA

For correspondence: cxu3@uoregon.edu

ORCID iD: [0000-0002-1056-8893](https://orcid.org/0000-0002-1056-8893)

Tyler B. Ramos

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon; Eugene, OR 97403, USA

Edward M. Rogers

Janelia Research Campus, Howard Hughes Medical Institute, 19700 Helix Drive, Ashburn, VA, USA

Michael B. Reiser

Janelia Research Campus, Howard Hughes Medical Institute, 19700 Helix Drive, Ashburn, VA, USA

ORCID iD: [0000-0002-4108-4517](https://orcid.org/0000-0002-4108-4517)

Chris Q. Doe

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon; Eugene, OR 97403, USA

Copyright

© 2023, Xu et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Hugo Bellen

Baylor College of Medicine, United States of America

Senior Editor

K VijayRaghavan

National Centre for Biological Sciences, Tata Institute of Fundamental Research, India

Reviewer #1 (Public Review):

In this very strong and interesting paper the authors present a convincing series of experiments that reveal molecular mechanism of neuronal cell type diversification in the nervous system of *Drosophila*. The authors show that a homeodomain transcription factor, Bsh, fulfills several critical functions - repressing an alternative fate and inducing downstream homeodomain transcription factors with whom Bsh may collaborate to induce L4 and L5 fates (the author's accompanying paper reveals how Bsh can induce two distinct fates). The authors make elegant use of powerful genetic tools and an arsenal of satisfying cell identity markers.

I believe that this is an important study because it provides some fundamental insights into the conservation of neuronal diversification programs. It is very satisfying to see that similar organizational principles apply in different organism to generate cell type diversity. The authors should also be commended for contextualizing their work very well, giving a broad, scholarly background to the problem of neuronal cell type diversification.

My one suggestion for the authors is to perhaps address in the Discussion (or experimentally address if they wish) how they reconcile that Bsh is on the one hand: (a) continuously expressed in L4/L4, (b) binding directly to a cohort of terminal effectors that are also continuously expressed but then, on the other hand, is not required for their maintaining L4 fate? A few questions: Is Bsh only NOT required for maintaining Ap expression or is it also NOT required for maintaining other terminal markers of L4? The former could be easily explained - Bsh simply kicks off Ap, Ap then autoregulates, but Bsh and Ap then continuously activate terminal effector genes. The second scenario would require a little more complex mechanism: Bsh binding of targets (with Notch) may open chromatin, but then once that's done, Bsh is no longer needed and Ap alone can continue to express genes. I feel that the

authors should be at least discussing this. The postmitotic Bsh removal experiment in which they only checked Ap and depression of other markers is a little unsatisfying without further discussion (or experiments, such as testing terminal L4 markers). I hasten to add that this comment does not take away from my overall appreciation for the depth and quality of the data and the importance of their conclusions.

Reviewer #2 (Public Review):

Summary:

In this paper, the authors explore the role of the Homeodomain Transcription Factor Bsh in the specification of Lamina neuronal types in the optic lobe of *Drosophila*. Using the framework of terminal selector genes and compelling data, they investigate whether the same factor that establishes early cell identity is responsible for the acquisition of terminal features of the neuron (i.e., cell connectivity and synaptogenesis).

The authors convincingly describe the sequential expression and activity of Bsh, termed here as 'primary HDTF', and of Ap in L4 or Pdm3 in L5 as 'secondary HDTFs' during the specification of these two neurons. The study demonstrates the requirement of Bsh to activate either Ap and Pdm3, and therefore to generate the L4 and L5 fates. Moreover, the authors show that in the absence of Bsh, L4 and L5 fates are transformed into a L1 or L3-like fates.

Finally, the authors used DamID and Bsh:DamID to profile the open chromatin signature and the Bsh binding sites in L4 neurons at the synaptogenesis stage. This allows the identification of putative Bsh target genes in L4, many of which were also found to be upregulated in L4 in a previous single-cell transcriptomic analysis. Among these genes, the paper focuses on Dip- β , a known regulator of L4 connectivity. They demonstrate that both Bsh and Ap are required for Dip- β , forming a feed-forward loop. Indeed, the loss of Bsh causes abnormal L4 synaptogenesis and therefore defects in several visual behaviors.

The authors also propose the intriguing hypothesis that the expression of Bsh expanded the diversity of Lamina neurons from a 3 cell-type state to the current 5 cell-type state in the optic lobe.

Strengths:

Overall, this work presents a beautiful practical example of the framework of terminal selectors: Bsh acts hierarchically with Ap or Pdm3 to establish the L4 or L5 cell fates and, at least in L4, participates in the expression of terminal features of the neuron (i.e., synaptogenesis through Dip- β regulation).

The hierarchical interactions among Bsh and the activation of Ap and Pdm3 expression in L4 and L5, respectively, are well established experimentally. Using different genetic drivers, the authors show a window of competence during L4 neuron specification during which Bsh activates Ap expression. Later, as the neuron matures, Ap becomes independent of Bsh. This allows the authors to propose a coherent and well-supported model in which Bsh acts as a 'primary' selector that activates the expression of L4-specific (Ap) and L5-specific (Pdm3) 'secondary' selector genes, that together establish neuronal fate.

Importantly, the authors describe a striking cell fate change when Bsh is knocked down from L4/L5 progenitor cells. In such case, L1 and L3 neurons are generated at the expense of L4 and L5. The paper demonstrates that Bsh in L4/L5 represses *Zfh1*, which in turn acts as the primary selector for L1/L3 fates. These results point to a model where the acquisition of Bsh during evolution might have provided the grounds for the generation of new cell types, L4 and L5, expanding lamina neuronal diversity for a more refined visual behaviors in flies. This is an intriguing and novel hypothesis that should be tested from an evo-devo standpoint, for instance by identifying a species when L4 and L5 do not exist and/or Bsh is not expressed in L neurons.

To gain insight into how Bsh regulates neuronal fate and terminal features, the authors have profiled the open chromatin landscape and Bsh binding sites in L4 neurons at mid-pupation using the DamID technique. The paper describes a number of genes that have Bsh binding peaks in their regulatory regions and that are differentially expressed in L4 neurons, based on available scRNAseq data. Although the manuscript does not explore this candidate list in depth, many of these genes belong to classes that might explain terminal features of L4 neurons, such as neurotransmitter identity, neuropeptides or cytoskeletal regulators. Interestingly, one of these upregulated genes with a Bsh peak is Dip- β , an immunoglobulin superfamily protein that has been described by previous work from the author's lab to be relevant to establish L4 proper connectivity. This work proves that Bsh and Ap work in a feed-forward loop to regulate Dip- β expression, and therefore to establish normal L4 synapses. Furthermore, Bsh loss of function in L4 causes impairs visual behaviors.

Weaknesses:

- The last paragraph of the introduction is written using rhetorical questions and does not read well. I suggest rewriting it in a more conventional direct style to improve readability.
- A significant concern is the way in which information is conveyed in the Figures. Throughout the paper, understanding of the experimental results is hindered by the lack of information in the Figure headers. Specifically, the genetic driver used for each panel should be adequately noted, together with the age of the brain and the experimental condition. For example, R27G05-Gal4 drives early expression in LPCs and L4/L5, while the 31C06-AD, 34G07-DBD Split-Gal4 combination drives expression in older L4 neurons, and the use of one or the other to drive Bsh-KD has dramatic differences in Ap expression. The indication of the driver used in each panel will facilitate the reader's grasp of the experimental results.
- Bsh role in L4/L5 cell fate:
 - o It is not clear whether Tll+/Bsh+ LPCs are the precursors of L4/L5. Morphologically, these cells sit very close to L5, but are much more distant from L4.
 - o Somatic CRISPR knockout of Bsh seems to have a weaker phenotype than the knockdown using RNAi. However, in several experiments down the line, the authors use CRISPR-KO rather than RNAi to knock down Bsh activity: it should be explained why the authors made this decision. Alternatively, a null mutant could be used to consolidate the loss of function phenotype, although this is not strictly necessary given that the RNAi is highly efficient and almost completely abolishes Bsh protein.
 - o Line 102: Rephrase "R27G05-Gal4 is expressed in all LPCs and turned off in lamina neurons" to "is turned off as lamina neurons mature", as it is kept on for a significant amount of time after the neurons have already been specified.
 - o Line 121: "(a) that all known lamina neuron markers become independent of Bsh regulation in neurons" is not an accurate statement, as the markers tested were not shown to be dependent on Bsh in the first place.
 - o Lines 129-134: Make explicit that the LPC-Gal4 was used in this experiment. This is especially important here, as these results are opposite to the Bsh Loss of Function in L4 neurons described in the previous section. This will help clarify the window of competence in which Bsh establishes L4/L5 neuronal identities through ap/pdm3 expression.
- DamID and Bsh binding profile:
 - o Figure 5 - figure supplement 1C-E: The genotype of the Control in (C) has to be described within the panel. As it is, it can be confused with a wild type brain, when it is in fact a Bsh-KO mutant.
 - o It is not clear how L4-specific Differentially Expressed Genes were found. Are these genes DEG between Lamina neurons types, or are they upregulated genes with respect to all neuronal clusters? If the latter is the case, it could explain the discrepancy between scRNAseq DEGs and Bsh peaks in L4 neurons.

- Dip- β regulation:
 - Line 234: It is not clear why CRISPR KO is used in this case, when Bsh-RNAi presents a stronger phenotype.
 - Figure 6N-R shows results using LPC-Gal4. It is not clear why this driver was used, as it makes a less accurate comparison with the other panels in the figure, which use L4-Split-Gal4. This discrepancy should be acknowledged and explained, or the experiment repeated with L4-Split-Gal4>Ap-RNAi.
 - Line 271: It is also possible that L4 activity is dispensable for motion detection and only L5 is required.
- Discussion: It is necessary to de-emphasize the relevance of HDTFs, or at least acknowledge that other, non-homeodomain TFs, can act as selector genes to determine neuronal identity. By restricting the discussion to HDTFs, it is not mentioned that other classes of TFs could follow the same Primary-Secondary selector activation logic.

Author Response

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public Review):

In this very strong and interesting paper the authors present a convincing series of experiments that reveal molecular mechanism of neuronal cell type diversification in the nervous system of Drosophila. The authors show that a homeodomain transcription factor, Bsh, fulfills several critical functions - repressing an alternative fate and inducing downstream homeodomain transcription factors with whom Bsh may collaborate to induce L4 and L5 fates (the author's accompanying paper reveals how Bsh can induce two distinct fates). The authors make elegant use of powerful genetic tools and an arsenal of satisfying cell identity markers.

Thanks!

I believe that this is an important study because it provides some fundamental insights into the conservation of neuronal diversification programs. It is very satisfying to see that similar organizational principles apply in different organisms to generate cell type diversity. The authors should also be commended for contextualizing their work very well, giving a broad, scholarly background to the problem of neuronal cell type diversification.

Thanks!

My one suggestion for the authors is to perhaps address in the Discussion (or experimentally address if they wish) how they reconcile that Bsh is on the one hand: (a) continuously expressed in L4/L4, (b) binding directly to a cohort of terminal effectors that are also continuously expressed but then, on the other hand, is not required for their maintaining L4 fate? A few questions: Is Bsh only NOT required for maintaining Ap expression or is it also NOT required for maintaining other terminal markers of L4? The former could be easily explained - Bsh simply kicks off Ap, Ap then autoregulates, but Bsh and Ap then continuously activate terminal effector genes. The second scenario would require a little more complex mechanism: Bsh binding of targets (with Notch) may open chromatin, but then once that's done, Bsh is no longer needed and Ap alone can continue to express genes. I feel that the authors should be at least discussing this. The postmitotic Bsh removal experiment in which they only checked Ap and depression of other markers is a little unsatisfying without further discussion (or experiments, such as

testing terminal L4 markers). I hasten to add that this comment does not take away from my overall appreciation for the depth and quality of the data and the importance of their conclusions.

Great suggestions, we will discuss these two hypotheses as requested.

Bsh initiates Ap expression in L4 neurons which then maintain Ap expression independently of Bsh expression, likely through Ap autoregulation. During the synaptogenesis window, Ap expression becomes independent from Bsh expression, but Bsh and Ap are both still required to activate the synapse recognition molecule DIP-beta. Additionally, Bsh also shows putative binding to other L4 identity genes, e.g., those required for neurotransmitter choice, and electrophysiological properties, suggesting Bsh may initiate L4 identity genes as a suite of genes. The mechanism of maintaining identity features (e.g., morphology, synaptic connectivity, and functional properties) in the adult remains poorly understood. It is a great question whether primary HDTF Bsh maintains the expression of L4 identity genes in the adult. To test this, in our next project, we will specifically knock out Bsh in L4 neurons of the adult fly and examine the effect on L4 morphology, connectivity, and function properties.

Reviewer #2 (Public Review):

Summary:

In this paper, the authors explore the role of the Homeodomain Transcription Factor Bsh in the specification of Lamina neuronal types in the optic lobe of Drosophila. Using the framework of terminal selector genes and compelling data, they investigate whether the same factor that establishes early cell identity is responsible for the acquisition of terminal features of the neuron (i.e., cell connectivity and synaptogenesis).

Thanks for the positive words!

The authors convincingly describe the sequential expression and activity of Bsh, termed here as 'primary HDTF', and of Ap in L4 or Pdm3 in L5 as 'secondary HDTFs' during the specification of these two neurons. The study demonstrates the requirement of Bsh to activate either Ap and Pdm3, and therefore to generate the L4 and L5 fates. Moreover, the authors show that in the absence of Bsh, L4 and L5 fates are transformed into a L1 or L3-like fates.

Thanks!

Finally, the authors used DamID and Bsh:DamID to profile the open chromatin signature and the Bsh binding sites in L4 neurons at the synaptogenesis stage. This allows the identification of putative Bsh target genes in L4, many of which were also found to be upregulated in L4 in a previous single-cell transcriptomic analysis. Among these genes, the paper focuses on Dip-β, a known regulator of L4 connectivity. They demonstrate that both Bsh and Ap are required for Dip-β, forming a feed-forward loop. Indeed, the loss of Bsh causes abnormal L4 synaptogenesis and therefore defects in several visual behaviors. The authors also propose the intriguing hypothesis that the expression of Bsh expanded the diversity of Lamina neurons from a 3 cell-type state to the current 5 cell-type state in the optic lobe.

Thanks for the excellent summary of our findings!

Strengths:

Overall, this work presents a beautiful practical example of the framework of terminal selectors: Bsh acts hierarchically with Ap or Pdm3 to establish the L4 or L5 cell fates and, at least in L4, participates in the expression of terminal features of the neuron (i.e., synaptogenesis through Dip- β regulation).

Thanks!

The hierarchical interactions among Bsh and the activation of Ap and Pdm3 expression in L4 and L5, respectively, are well established experimentally. Using different genetic drivers, the authors show a window of competence during L4 neuron specification during which Bsh activates Ap expression. Later, as the neuron matures, Ap becomes independent of Bsh. This allows the authors to propose a coherent and well-supported model in which Bsh acts as a 'primary' selector that activates the expression of L4-specific (Ap) and L5-specific (Pdm3) 'secondary' selector genes, that together establish neuronal fate.

Thanks again!

Importantly, the authors describe a striking cell fate change when Bsh is knocked down from L4/L5 progenitor cells. In such cases, L1 and L3 neurons are generated at the expense of L4 and L5. The paper demonstrates that Bsh in L4/L5 represses Zfh1, which in turn acts as the primary selector for L1/L3 fates. These results point to a model where the acquisition of Bsh during evolution might have provided the grounds for the generation of new cell types, L4 and L5, expanding lamina neuronal diversity for a more refined visual behaviors in flies. This is an intriguing and novel hypothesis that should be tested from an evo-devo standpoint, for instance by identifying a species when L4 and L5 do not exist and/or Bsh is not expressed in L neurons.

Thanks for the appreciation of our findings!

To gain insight into how Bsh regulates neuronal fate and terminal features, the authors have profiled the open chromatin landscape and Bsh binding sites in L4 neurons at mid-pupation using the DamID technique. The paper describes a number of genes that have Bsh binding peaks in their regulatory regions and that are differentially expressed in L4 neurons, based on available scRNAseq data. Although the manuscript does not explore this candidate list in depth, many of these genes belong to classes that might explain terminal features of L4 neurons, such as neurotransmitter identity, neuropeptides or cytoskeletal regulators. Interestingly, one of these upregulated genes with a Bsh peak is Dip- β , an immunoglobulin superfamily protein that has been described by previous work from the author's lab to be relevant to establish L4 proper connectivity. This work proves that Bsh and Ap work in a feed-forward loop to regulate Dip- β expression, and therefore to establish normal L4 synapses. Furthermore, Bsh loss of function in L4 causes impaired visual behaviors.

Thanks for the excellent summary of our findings.

Weaknesses:

- The last paragraph of the introduction is written using rhetorical questions and does not read well. I suggest rewriting it in a more conventional direct style to improve readability.

We agree and have updated the text as suggested.

• A significant concern is the way in which information is conveyed in the Figures. Throughout the paper, understanding of the experimental results is hindered by the lack of information in the Figure headers. Specifically, the genetic driver used for each panel should be adequately noted, together with the age of the brain and the experimental condition. For example, R27G05-Gal4 drives early expression in LPCs and L4/L5, while the 31C06-AD, 34G07-DBD Split-Gal4 combination drives expression in older L4 neurons, and the use of one or the other to drive Bsh-KD has dramatic differences in Ap expression. The indication of the driver used in each panel will facilitate the reader's grasp of the experimental results.

We agree and have updated the figure annotation.

• Bsh role in L4/L5 cell fate: o It is not clear whether Tll+/Bsh+ LPCs are the precursors of L4/L5. Morphologically, these cells sit very close to L5, but are much more distant from L4.

Our current data show L4 and L5 neurons are generated by different LPCs. However, currently, we don't have tools to demonstrate which subset of LPCs generate which lamina neuron type. We are currently working on a follow-up manuscript on LPC heterogeneity, but those experiments have just barely been started.

• Somatic CRISPR knockout of Bsh seems to have a weaker phenotype than the knockdown using RNAi. However, in several experiments down the line, the authors use CRISPR-KO rather than RNAi to knock down Bsh activity: it should be explained why the authors made this decision. Alternatively, a null mutant could be used to consolidate the loss of function phenotype, although this is not strictly necessary given that the RNAi is highly efficient and almost completely abolishes Bsh protein.

The reason we chose CRISPR-KO (L4-specific Gal4, uas-Cas9, and uas-Bsh-sgRNAs) is that it effectively removed Bsh expression from the majority of L4 neurons. However, it failed to knock down Bsh in L4 neurons using L4-split Gal4 and Bsh-RNAi because L4-split Gal4 expression depends on Bsh. We have updated this explanation in the text.

• Line 102: Rephrase "R27G05-Gal4 is expressed in all LPCs and turned off in lamina neurons" to "is turned off as lamina neurons mature", as it is kept on for a significant amount of time after the neurons have already been specified.

Thanks; we have made that change.

• Line 121: "(a) that all known lamina neuron markers become independent of Bsh regulation in neurons" is not an accurate statement, as the markers tested were not shown to be dependent on Bsh in the first place.

Good point. We have rephrased it as "that all known lamina neuron markers are independent of Bsh regulation in neurons".

• Lines 129-134: Make explicit that the LPC-Gal4 was used in this experiment. This is especially important here, as these results are opposite to the Bsh Loss of Function in L4 neurons described in the previous section. This will help clarify the window of competence in which Bsh establishes L4/L5 neuronal identities through ap/pdm3 expression.

Thanks! We have updated Gal4 information in the text for every manipulation.

- *DamID and Bsh binding profile:*
- *Figure 5 - figure supplement 1C-E: The genotype of the Control in (C) has to be described within the panel. As it is, it can be confused with a wild type brain, when it is in fact a Bsh-KO mutant.*

Great point! Thank you for catching this and we have updated it.

- *It Is not clear how L4-specific Differentially Expressed Genes were found. Are these genes DEG between Lamina neurons types, or are they upregulated genes with respect to all neuronal clusters? If the latter is the case, it could explain the discrepancy between scRNAseq DEGs and Bsh peaks in L4 neurons.*

We did not use “L4-specific Differentially Expressed Genes”. Instead, we used all genes that are significantly transcribed in L4 neurons (line 209-213).

- *Dip-β regulation:*
- *Line 234: It is not clear why CRISPR KO is used in this case, when Bsh-RNAi presents a stronger phenotype.*

As we explained above, the reason we chose CRISPR-KO (L4-specific Gal4, uas-Cas9, and uas-BshsgRNAs) is that it effectively removed Bsh expression from the majority of L4 neurons. However, it failed to knock down Bsh in L4 neurons using L4-split Gal4 and Bsh-RNAi because L4-split Gal4 expression depends on Bsh. We have updated this explanation in the text.

- *Figure 6N-R shows results using LPC-Gal4. It is not clear why this driver was used, as it makes a less accurate comparison with the other panels in the figure, which use L4-Split-Gal4. This discrepancy should be acknowledged and explained, or the experiment repeated with L4-Split-Gal4>Ap-RNAi.*

I think you mean 6J-M shows results using LPC-Gal4. We first tried L4-Split-Gal4>Ap-RNAi but it failed to knock down Ap because L4-Split-Gal4 expression depends on Ap. We have added this to the text.

- *Line 271: It is also possible that L4 activity is dispensable for motion detection and only L5 is required.*

Thanks! Work from Tuthill et al, 2013 showed that L5 is not required for any motion detection. We have included this citation in the text.

- *Discussion: It is necessary to de-emphasize the relevance of HDTFs, or at least acknowledge that other, non-homeodomain TFs, can act as selector genes to determine neuronal identity. By restricting the discussion to HDTFs, it is not mentioned that other classes of TFs could follow the same PrimarySecondary selector activation logic.*

That is a great point, thank you! We have included this in the discussion.