

A genome-wide association study implicates the olfactory system in *Drosophila melanogaster* diapause-associated lifespan extension and fecundity

Sreesankar Easwaran, Denise J. Montell 

Molecular, Cellular, and Developmental Biology Department University of California Santa Barbara, CA 93106

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The effects of environmental stress on animal life are gaining importance with climate change. Diapause is a dormancy program that occurs in response to an adverse environment, followed by resumption of development and reproduction upon the return of favorable conditions. Diapause is a complex trait, so we leveraged the *Drosophila* genetic reference panel (DGRP) lines and conducted a Genome-Wide Association Study (GWAS) to characterize the genetic basis of diapause. We assessed post-diapause and non-diapause fecundity across all DGRP lines. GWAS revealed 546 genetic variants, encompassing single nucleotide polymorphisms, insertions and deletions associated with post-diapause fecundity. We identified 291 candidate diapause-associated genes, 40 of which had previously been associated with diapause. 89 of the candidates were associated with more than one SNP. Gene network analysis indicated that the diapause-associated genes were primarily linked to neuronal and reproductive system development. Similarly, comparison with results from other fly GWAS revealed the greatest overlap with olfactory-behavior-associated and fecundity-and-lifespan-associated genes. An RNAi screen of the top candidates identified two neuronal genes, *Dip-γ* and *Scribbler*, to be required during recovery for post-diapause fecundity. We complemented the genetic analysis with a test of which neurons are required for successful diapause. We found that although amputation of the antenna had little to no effect on non-diapause lifespan, it reduced diapause lifespan and postdiapause fecundity. We further show that olfactory receptor neurons and temperature-sensing neurons are required for successful recovery from diapause. Our results provide insights into the molecular, cellular, and genetic basis of adult reproductive diapause in *Drosophila*.

eLife assessment

This **useful** study shows how genetic variation is associated with fecundity following a period of reproductive diapause in female *Drosophila*. The work identifies the olfactory system as central to successful diapause with associated changes in longevity and fecundity. While the genetic screening and methods used are **solid**, the approach to assessing diapause is **incomplete** and could benefit from additional orthogonal experiments.

<https://doi.org/10.7554/eLife.98142.1.sa3>

Introduction

To endure adverse environments, many organisms from worms and flies to mammals initiate a dormancy program called diapause, triggered by cues like unfavorable temperatures, drought, or starvation (Tatar and Yin 2001 [↗](#); Hand et al. 2016 [↗](#); Deng et al. 2018 [↗](#); Hussein et al. 2022 [↗](#); Easwaran and Montell 2023 [↗](#)). By enhancing survival, diapause can support both beneficial insects like pollinators, and harmful insects such as agricultural pests and disease vectors. Elucidating diapause mechanisms may suggest ways to control harmful insects and support beneficial organisms. Furthermore, unraveling diapause genetics holds promise for understanding stem cell resilience (Easwaran et al. 2022 [↗](#)), aging (Fontana et al. 2010 [↗](#); Hutfilz 2022 [↗](#)), and even cancer dormancy (Lin and Zhu 2021 [↗](#); Easwaran and Montell 2023 [↗](#)).

Successful diapause occurs optimally at different developmental stages depending on the organism. In *Drosophila* species, the optimal stage is the newly-eclosed adult. During diapause, feeding, activity levels, and mating behaviors change, metabolism slows, reproduction arrests, stress resilience increases, and lifespan is lengthened (Kubrak et al. 2014 [↗](#)).

The genetic basis of adult reproductive diapause in *Drosophila* is likely complex because the diapause program involves sensing environmental cues and responding by reprogramming behavior, metabolism, reproduction, and aging, continuously monitoring environmental conditions, and reactivating development and/or reproduction when conditions become favorable. Historically, studies of *Drosophila* diapause have focused primarily on entry into diapause by measuring arrest of egg production, specifically at the stage of yolk accumulation referred to as vitellogenesis (Saunders et al. 1989 [↗](#), 1990 [↗](#)). However, it has become clear that diapause is a comprehensive program that changes virtually every aspect of the fly's life from behavior and to metabolism, reproduction, and lifespan (Tatar and Yin 2001 [↗](#); Kubrak et al. 2014 [↗](#); Denlinger 2023 [↗](#)). Prior studies have identified temperature and day-length as relevant environmental cues and key hormonal regulators such as insulin, juvenile hormone (JH), and 20-hydroxyecdysone (20E), which coordinate metabolism and reproduction (Hutfilz 2022 [↗](#); Denlinger 2023 [↗](#)). Recent studies identify specific circadian neurons that sense changes in temperature and relay that information to the reproductive system (Meiselman et al. 2022 [↗](#); Hidalgo et al. 2023 [↗](#)). Yet, much remains to be learned about this fascinating life history trait (Denlinger 2023 [↗](#)).

One approach to analyzing complex traits and mapping quantitative trait loci (QTL) is to conduct a Genome-Wide Association Study (GWAS) (Huang et al. 2014 [↗](#); Morozova et al. 2015 [↗](#); Shorter et al. 2015 [↗](#)). The *Drosophila* Genome Reference Panel (DGRP) comprises fully sequenced, highly inbred lines of *Drosophila melanogaster* and serves as a valuable tool for understanding genotype-phenotype relationships in *Drosophila* (Mackay et al. 2012 [↗](#); Morozova et al. 2015 [↗](#); Mackay and Huang 2018 [↗](#)).

We used the DGRP to carry out a *Drosophila* GWAS and identify genes and gene classes associated with successful diapause. We developed a novel method for assessing successful diapause, by evaluating not only the arrest of egg development or the recovery of egg maturation but a more stringent criterion: the ability of post-diapause flies to produce viable adult progeny. Of the 291 diapause-associated genes we identified, ~40 have previously been implicated in diapause in flies or another organism. The classes of genes most over-represented in the diapause-associated set are genes that regulate neuronal development and gonad development. We show that *Gal4*-mediated RNAi knockdown is effective at 10°C, and we identify two neuronal genes required during recovery for post-diapause fecundity. We further complement this genetic analysis by identifying neurons in the fly antenna required for successful diapause. Our studies implicate the olfactory system as critically important in successful diapause.

Results

Quantifying successful diapause across the DGRP lines

Key features of *Drosophila* diapause include ovarian arrest and post-diapause recovery of fertility (the ability to produce any progeny) and fecundity (the number of progeny produced). Whereas the majority of studies of *Drosophila* diapause focus on arrest of oogenesis, we quantified diapause by assessing the ability of newly-eclosed flies to undergo 35 days of diapause, recover, and produce viable progeny, which is a more stringent test of success (Fig. 1A [↗](#)).

DGRP lines were allowed to develop under non-diapause conditions (25°C and 12:12 L:D). Virgin females were collected and transferred to diapausing conditions (10°C and 8:16 L:D) for 5 weeks. Subsequently, flies were shifted to 18°C for one day, followed by incubation at 25°C in a fresh vial of fly food for recovery. We evaluated the ability of post-diapause flies to reproduce by mating individual females with 2 Canton-S (CS) male control (non-diapausing) flies for 4 days. Afterward, parents were removed, and progeny were allowed to develop for 12 more days. We then measured post-diapause fecundity by counting the number of adult progeny that eclosed. A minimum of 20 female flies per DGRP line were tested. Non-diapause fecundity was measured by setting crosses immediately upon collecting virgin flies. The average non-diapause and post-diapause 4-day fecundity of DGRP lines are shown in Figure 1B [↗](#), arranged in ascending order of non-diapause fecundity. In general, most lines showed lower fecundity post-diapause compared to newly-eclosed flies in optimal conditions. However, there were interesting exceptions. For example, lines 309, 738, and 853 exhibited better fecundity post-diapause compared even to young flies in optimal conditions (Supplemental Table S1). The DGRP lines exhibit variability in fecundity both post-diapause and in non-diapause conditions [Fig. 1B [↗](#) and (Durham et al. 2014 [↗](#))]. Therefore, we normalized post-diapause to non-diapause fecundity (Fig. 1C [↗](#) and Supplemental Table S1).

The lines that showed poor non-diapause fecundity that improved after diapause (e.g. 362, 737, 822) suggested that there might be a genetic tradeoff between the two. However, the overall correlation (Fig. 1D [↗](#)) between non-diapause and post-diapause fecundity was 32.3% ($R^2=0.3228$, Pearson's correlation coefficient, $r=0.5682$), suggesting a positive rather than negative correlation overall between post-diapause and non-diapause fecundity. The frequency distributions of 4-day fecundity for the non-diapause (Fig. 1E [↗](#)), and normalized fecundity (Fig. 1F [↗](#)) conditions are shown. Although fecundity is lower post-diapause than in young flies, even in Canton-S (CS) controls, it is higher than for flies maintained in non-diapause conditions for 35 days (Fig. 1G [↗](#) and Supplemental Table S2).

GWAS for diapause using the DGRP tool

The normalized fecundity scores (post-diapause 4-day fecundity/non-diapause control 4-day fecundity) served as the basis for conducting a GWAS using the DGRP2 web tool (Mackay et al. 2012 [↗](#); Huang et al. 2014 [↗](#)). A total of 546 genetic variants, encompassing single nucleotide




Figure 1.

Quantification of diapause in *Drosophila* Genetic Reference Panel (DGRP) lines measured as the ratio of post-diapause to non-diapause fecundity.

(A) Schematic of experimental workflow. (B) Average number of progenies produced (fecundity) in a 4-day individual female fly mating experiment of DGRP lines either as non-diapausing (yellow) or after a 35-day post-diapause (blue) virgin flies. Each dot represents the average fecundity and the line above represents the standard error. DGRP line numbers are indicated wherever the post-diapause fecundity exceeds the non-diapause fecundity. (C) Normalized post-diapause fecundity [average of (individual post-diapause fecundity/mean non-diapause fecundity)] of each DGRP line. (D) Correlation of post-diapause to non-diapause fecundity. Pearson's $r=0.5682$, $r^2=0.3228$. (E-F) Frequency distribution of DGRP lines fecundity under non-diapause (E) and of the normalized post-diapause fecundity (F). (G) Average 4-day fecundity of single female flies, each crossed with 2 young Canton S male flies, aged for 1, 35, or 42 days in non-diapause conditions or kept in diapause conditions followed by recovery. 1-way ANOVA and Tukey's multiple comparison test, compact letter display shows comparisons. n is the number of individual female fly fecundity measured.

polymorphisms (SNPs) and insertions/deletions (indels), were identified as associated with diapause fecundity (**Fig. 2A-B** and Supplemental Table S3). When a variant is located within 1kb up- or down-stream of an annotated gene, or within the gene, it is considered potentially associated with that gene. We thus identified 291 candidate diapause-associated fecundity genes. Notably, 40 out of the 291 genes had previously been reported to be associated with diapause (Supplemental Table S4) either through functional analysis [e.g., insulin receptor (InR) and *couch potato* (*cpo*)] (Schmidt et al. 2008; Zhang and Denlinger 2011; Kankare et al. 2012; Sim and Denlinger 2013; Kubrak et al. 2014), due to changes in gene expression [e.g., expanded (ex) and Laminin A (LanA)] (Zhao et al. 2016), or associated with cold tolerance [β -Tub97EF (Myachina et al. 2017)] (Supplemental Table S4). Eighty-nine genes were associated with more than one diapause-associated variant (e.g. at least two different SNPs) (**Fig. 2C**).

Network analysis of diapause-associated genes

Using the gene list obtained from the GWAS, we performed a gene network analysis using the GeneMania application within Cytoscape (Shannon et al. 2003; Warde-Farley et al. 2010). This application facilitates the generation of network predictions based on a combination of known physical and genetic interactions, co-localization, co-expression, shared protein domains, pathway data, and predicted functional relationships between genes. Additionally, it identifies subnetworks of genes related by functional Gene Ontology (GO). Enriched subnetworks are identified by dividing the number of genes from the input set by the total number of genes associated with a specific gene ontology. The subnetwork analysis is presented in **Figure 2D**. Two primary classes of GO terms are significantly enriched in the diapause GWAS set compared to the genome as a whole: nervous system development and the reproductive system/gonad development. These two categories are consistent with the idea that changes in the environment are sensed by the nervous system and the information is relayed to the reproductive system.

We also compared the list of diapause-associated genes to results of other *Drosophila* GWAS (**Fig. 3**, Supplemental Table S5). Intriguingly, the most significant overlap was with genes identified as associated with olfactory behavior (Horváth and Kalinka 2018), which has not previously been described as important for diapause. The second most significant overlap was with genes identified in non-diapause fecundity and lifespan (Durham et al. 2014). Less significant overlaps were found with genes associated with circadian rhythm (Harbison et al. 2019), chill coma recovery (Mackay et al. 2012), development time associated with lead toxicity (Zhou et al. 2016), alcohol sensitivity (Morozova et al. 2015), starvation resistance (Mackay et al. 2012), mean food intake (Garlapow et al. 2015), sleep (Harbison et al. 2013), neurotoxin methylmercury (MeHg) tolerance (Montgomery et al. 2014), brain wiring regulators (Li et al. 2020), and death due to traumatic brain injury (Katzenberger et al. 2015). Overlaps with genes associated with courtship, viability in lead toxicity (Zhou et al. 2016), effects of genetic architecture and Wolbachia status on starvation (Huang et al. 2014), aggression (Shorter et al. 2015), ER stress (Chow et al. 2013), leg development (Grubbs et al. 2013), startle response (Mackay et al. 2012), and embryo development time (Horváth et al. 2016) were not statistically significant. In summary, both analyses implicated the nervous system, and potentially the olfactory system in particular, in the regulation of post-diapause fecundity.

RNAi analysis of GWAS candidates identifies neural genes required for recovery

Genes with multiple SNPs are good candidates for influencing diapause traits. To assess the functional significance of the top candidates, we conducted an RNAi screen of the genes with multiple associated SNPs. First, we evaluated the effectiveness of *Gal4* under diapausing conditions. As a control, we crossed *Mat- α -tub-Gal4*, which drives expression in the female germline to *UASp-F-tractin.tdTomato* to assess the effectiveness of *Gal4*-mediated expression at




Figure 2.

Genome-wide association of *Drosophila* diapause.

(A) Manhattan plot for genome-wide association distribution. The position of each point along the y-axis indicates $-\log_{10}(P\text{-value})$ of association of a SNP, insertion, or deletion. Points above the blue line have a P value $< 1e^{-5}$. The red line represents Bonferroni corrected P value = $4.8e^{-8}$. (B) Q-Q plot of P values from the DGRP single variant GWAS with the red line representing expected P-value and observed P values deviating (black dots) from expected. (C) Numbers of genetic variants and candidate genes associated with diapause according to the GWAS. (D) Subnetworks from the Cytoscape analysis showing q-value (using the Benjamini-Hochberg procedure) for each subnetwork identified.




Figure 3.

Common genes to diapause-GWAS hits and other behavior-associated genes.

(A-T) Venn diagrams illustrate the intersection of genes associated with diapause identified through Genome-Wide Association Studies (diapause-GWAS) with genes from other behavior-related gene lists obtained from various studies. The diapause-GWAS gene set is represented as the first set throughout the figure, while subsequent sets represent different behavior-related gene lists identified in separate studies. The percentage of common genes compared to the total genes from different respective behavior-associated gene lists are provided for each Venn diagram. p-values of overlap to the diapause gene list determined by Fisher's exact tests are also provided. Venn diagrams are arranged in the order of p-values.

10°C. We compared tdTomato expression in flies maintained at different temperatures for 3 weeks. Compared to 25°C (Fig. 4A) and 18°C (Fig. 4B), flies at 10°C exhibited as high or higher expression of tdTomato (Fig. 4C).

To evaluate the effectiveness of *Gal4*-mediated RNAi at 10°C, we used *Mat-α-tub-Gal4* to drive *zpg* (*zero population growth*, aka *INX4*) RNAi and assessed the extent of Zpg knockdown by immunostaining with an anti-Zpg antibody. We used *Mat-α-tub-Gal4* because Zpg is expressed in the germline, and we chose to target Zpg because of the availability of the anti-Zpg antibody. Zpg is expressed from the earliest stages of germline development in the germarium (Fig. 4E, E'). *Mat-α-tub-Gal4* is not expressed detectably in the germarium but is expressed in early egg chambers (Fig. 4C). So, to quantify the knockdown efficiency, we measured the level of Zpg in stage 3 egg chambers relative to the germarium staining, as an internal control. Remarkably, comparison of *Mat-α-tub-Gal4>zpg* RNAi flies kept at different temperatures for 3 weeks revealed that *Gal4*-mediated RNAi was as effective at 10°C, as it was at 18°C and 25°C (Fig. 4D-H).

The top 15 GWAS hits for which multiple RNAi lines were available were selected for RNAi screening (Supplemental Table S6). We used *tub-Gal4* because it is widely expressed, including in neurons. We measured non-diapause and post-diapause fecundity. For the five RNAi lines that proved lethal, we used *tubulin-Gal80^{ts}* to suppress the RNAi expression during development and diapause and then shifted the flies to 30 °C to inactivate the Gal80 and drive the RNAi during recovery (Fig. 4I), and measured post-diapause fecundity. In this way we could test specifically for a function in post-diapause recovery (Supplementary Table S5). RNAi against two genes, *Defective-proboscis-extension-response interacting protein-γ* (Dip-γ) and *Scribbler* (*sbb*) significantly reduced post-diapause fecundity (Fig. 4J).

To determine whether Dip-γ and *sbb* are required specifically in neurons we crossed *nSyb-Gal4*, a pan-neuronal driver, to validated *UAS-Dip-γ* RNAi and *UAS-sbb* RNAi lines (Davis et al. 2014; Shimozono et al. 2019). Neuron-specific knockdown of Dip-γ (Fig. 4K, Supplemental Table S7) caused as severe a defect in post-diapause fecundity as ubiquitous RNAi, in contrast to glial knockdown using *Repo-Gal4* (Fig. 4L, Supplemental Table S7). Pan neuronal RNAi of *sbb* (Fig. 4K) also significantly inhibited post-diapause fecundity more than glial RNAi (Fig. 4L). We conclude that Dip-γ and *sbb* are required in neurons for successful diapause, consistent with the enrichment of this gene class in the diapause GWAS.

Post-diapause fecundity requires neurons in the antenna

While we were able to show significant functional effects of Dip-γ and *sbb*, GWAS studies by their nature identify many genes with small effects, which are too small to detect individually. Furthermore, the overlap with genes associated with olfactory behavior led us to complement the genetic analysis by testing whether the antenna and neurons within it are required for successful diapause. We removed the antenna from CS flies and measured post-diapause fecundity. As a control for the surgery, we removed the arista, which is an appendage from the antenna. Removal of the antenna but not the arista reduced post-diapause fecundity compared to unmanipulated controls (Fig. 5A and Supplemental Table S9). Removal of the antenna but not the arista also slightly, but statistically significantly, reduced the number of germline stem cells post-diapause (Fig. 5B). These results suggest that sensory cells in the antenna are important for successful diapause.

Another important diapause trait is lifespan extension. So we tested the effect of removing the antenna on post-diapause and non-diapause lifespan (Fig. 5C). Antenna removal had almost no effect on non-diapause lifespan (median survival 72 days w/o antenna vs 75 days with antenna) (Fig. 5C). In contrast, antennaless flies exhibited a substantial reduction in diapause lifespan (median survival 95 days) compared to flies that had intact antennae (median survival 142 days) (Fig. 5C), confirming the importance of the antenna for successful diapause.




Figure 4.

RNAi-mediated loss of function study to identify genes involved in diapause.

(A-C) *Mat- α -tub-Gal4* driving expression of *UASp-F-tractin.tdTomato* (red) at the indicated temperatures. Scale bars are 100 μ m. (D) Quantification of *zpg* RNAi knockdown in Stage-3 egg chambers normalized to the level of Zpg in the germarium. (1-way ANOVA and Tukey's multiple comparison test, compact letter display shows comparisons). The numbers (n) of Stage 3 egg chambers quantified are shown. (E-H) Representative images of egg chambers stained with anti-Zpg antibody (green) from either control (no-knockdown), (E, E') or knockdown of *zpg* (F-H') driven by *Mat- α -tub-Gal4* at different temperatures. (E'-H') are higher magnification, single channel views of the ovarioles shown in (E-H). Scale bars are 100 μ m for (E-H) and 20 μ m in (E'-H'). All flies in (A-H') were kept at respective temperatures for 3 weeks. (I) Experimental design for RNAi knockdown specifically during recovery for the experiment shown in (J). The temperature-sensitive Gal80 repressor of Gal4 prevented RNAi expression during development and diapause. Incubation at 30°C during recovery inactivates Gal80, allowing Gal4-mediated RNAi knockdown. (J) Ubiquitous knockdown of *Dip-y* or *sbb* with *tub Gal4* specifically during recovery as shown in (I) significantly reduces post-diapause/non-diapause fecundity compared to the control (*tubGal80^{TS}; tubGal4 > Ctrl* RNAi #9331). (K) Pan-neuronal RNAi knockdown of *Dip-y* and *sbb* with *nSybGal4* significantly reduces post-diapause/non-diapause fecundity compared to the control (*nSyb Gal4 > Ctrl* RNAi #54037). (L) Glia-specific knockdown of *Dip-y* or *sbb* with *Repo Gal4* causes little or no reduction in post-diapause/non-diapause fecundity (Control- *Repo Gal4 > Ctrl* RNAi #54037). In (J-L), 1-way ANOVA and Tukey's multiple comparison test, compact letter display shows comparisons. n is the number of individual female flies tested.




Figure 5.

Neural control of diapause.

(A) Effect of antenna removal on recovery of fecundity post-diapause. Arista removal was used as a control for the surgery. *n* is the number of individual female flies tested. 1-way ANOVA and Tukey's multiple comparison test, compact letter display shows comparisons. (B) Effect of antenna removal on GSC recovery after 5 weeks of diapause. *n* is the number of germaria counted (there are typically 2-3 GSCs/germarium). 1-way ANOVA and Tukey's multiple comparison test with compact letter display to show comparisons. (C) Role of antenna in lifespan extension in diapause. Control and antennalless flies were maintained at 25°C for 2 weeks post-surgery to allow for wound healing. Control flies were maintained at 25°C and diapause flies were moved to 10°C. Median survival for flies with intact antenna in diapause (+A_Diap) - 142 days; antennalless flies in diapause (-A_Diap) - 95 days; Control with intact antenna in optimal conditions (+A_Ctrl) - 75 days; and antennalless flies in optimal conditions (-A_Ctrl) - 72 days. Survival curves are compared pairwise using the Log-rank (Mantel-Cox) test and χ^2 values are: Diap +/- antenna = 86, Ctrl +/- antenna = 11.7, Diap +antenna vs Ctrl +antenna = 158.3, and Diap -antenna vs Ctrl -antenna = 105.8. (D-E) Effects on diapause of inactivating neuronal transmission (by driving tetanus toxin using UAS-TeTxLC.tdt, in odor responsive neurons using the indicated *Gal4* lines. Orco is the co-receptor for all of the odorant receptors (D). (E) Ir8a is a co-receptor involved in organic acid detection. Ir25a is a co-receptor involved in chemo- and thermo-sensation. Ir76b is involved in detection of various amines and salt. Ir84a is involved in detection of phenylacetic acid and male courtship behavior. Hot Cells are heat-sensitive cells in the arista.

Drosophila antennae are involved in various sensory modalities, including olfaction. To identify which neurons are required for diapause, we suppressed neuronal activity in subsets of cells by expressing the tetanus toxin light chain protein (TNT), which selectively cleaves the neuronal isoform of fly synaptobrevin. Orco is the virtually universal co-receptor for all of the odorant receptors of the OR class (Montell 2021 [↗](#)), so we used Orco-*Gal4* to drive expression. We found that blocking neuronal transmission in the Orco-expressing neurons decreased the post-diapause/non-diapause fecundity ratio (Fig. 5D [↗](#), Supplemental Table S9), further implicating odor perception in successful diapause (see discussion).

Ionotropic receptors (IRs) are another class of odorant receptor (Montell 2021 [↗](#)). Ir25a is broadly expressed, and inhibiting Ir25a-expressing neurons by expressing TNT reduced the post-diapause/non-diapause fecundity ratio compared to *Gal4s* expressed in other IR-expressing cells such as Ir8a and Ir84a, which are involved in odor perception, and Ir76b, which is involved in contact chemosensation (Fig. 5E [↗](#) and Supplemental Table S9). We also found that inhibiting neurons expressing Ir21a, mutations in which disrupt warm and cool avoidance behaviors (Budelli et al. 2019 [↗](#)), dramatically impaired recovery from diapause. Blocking Ir21a neuronal transmission prevented flies from exiting diapause, as evidenced by 100% mortality of the post-diapause female flies moved to recovery conditions (n= 25). Expressing TNT in the “hot cell” neurons (Ni et al. 2013 [↗](#)), which respond to heating, using *HC-Gal4*, also significantly reduced the post-diapause fecundity (Fig. 5E [↗](#) and Supplemental Table S9).

Discussion

Genome wide analysis of the effect of diapause on fecundity

Drosophila diapause is a fascinating life history trait that confers resilience in stressful environments and extends reproductive potential and organismal longevity. Our GWAS study identified nearly 300 candidate genes associated with an important but understudied feature of diapause: the ability to recover and reproduce successfully post-diapause. The most striking finding is that genes associated with neural development are highly over-represented in the diapause set. We further confirmed that at least two such genes, *Dip-γ* and *sbb*, are essential for post-diapause fecundity.

When we started this project, little was known about the neural control of diapause. Consistent with our finding reported here that neural genes are enriched in the diapause GWAS, and our recent finding that circadian activity and sleep are dramatically altered in diapause (Meyerhof et al. 2024 [↗](#)), two recent studies have reported that low temperature affects two subsets of circadian neurons, DN3s, and sLNvs, which in turn impact reproductive arrest (Meiselman et al. 2022 [↗](#); Hidalgo et al. 2023 [↗](#)). sLNv neurons secrete the neuropeptide Pigment Dispersing Factor (PDF) onto insulin producing cells (IPCs), which secrete insulin-like peptides (ILPs), which in turn promote JH production. At 10°C, PDF mRNA and protein levels are reduced in sLNv neurons, reducing JH production, which arrests vitellogenesis. In parallel, cool-temperature-induced reduction in DN3 activity disinhibits cholinergic ASTC-R2 neurons, which in turn inhibit vitellogenic egg chamber development. These neurons do not synapse onto the IPCs, and the mechanism by which the ASTC-R2 neurons affect oogenesis is unknown. Thus, there are at least two parallel mechanisms by which cool temperatures affect neural activity and contribute to reproductive arrest. It is not yet clear how cool temperatures reduce PDF mRNA or how PDF mRNA and protein levels rebound during recovery.

We found that the neuronal transcription factor *Sbb* is required post-diapause for recovery, so it is interesting to speculate that *Sbb* could affect PDF mRNA abundance. *Sbb* functions predominantly as a transcriptional repressor, so it could promote PDF expression in recovery by inhibiting expression or activity of a repressor of PDF. Alternatively, PDF reduces levels of the transcriptional

co-activator Eyes Absent (Eya) in the IPCs, so reduced PDF leads to elevated Eya protein levels during diapause. It is possible that Sbb represses Eya transcription to facilitate recovery post-diapause. A third possibility is that Sbb functions in the DN3-ASTC-R2 pathway. We have also found that circadian rhythms and sleep behavior are dramatically altered in flies at 10°C (Meyerhof et al. 2024 [↗](#)). It will be of interest to determine if Dip-γ and *sbb* are required for those behavioral effects at low temperatures, in addition to the effects on post-diapause fecundity described here.

Olfactory neurons are required for post-diapause fecundity

We complemented our genetic analysis with an investigation of the cells that contribute to diapause. The overlap between olfactory-behavior-associated genes and diapause-associated genes inspired us to test if the antenna is required, and we found that the antenna is required both for post-diapause fecundity and for lifespan extension. We conclude that cells in the antenna transmit important information for successful diapause. Furthermore, inhibiting olfactory receptor neurons by expressing TNT with Orco-Gal4 caused a similar effect on post-diapause fecundity. This is interesting because Meiselman et al found that the antenna was not required for accumulation of mature eggs post-diapause (Meiselman et al. 2022 [↗](#)). Our measurement of fecundity required not only that mature eggs developed in the ovary but also that they were fertilized, laid, and could develop into adults. So, what is required for production of viable progeny in addition to mature eggs? Since both feeding and mating are suppressed during diapause, the results suggest that the antenna may be required to promote these behaviors, consistent with the known role of the antenna in providing sensory information important for flies to seek food and mates. The antenna may be dispensable for relieving the vitellogenesis block in the ovary but may specifically be required for animals to find sufficient food and mates post-diapause to support maximal fecundity.

Consistent with the possible role of the olfactory system, in a screen of 505 gene trap lines, Tunstall et al. identified 16 genes that were expressed in ORNs, three of which are present in our diapause-associated gene set: *tai*, *cpo*, and *dpp* (Tunstall et al. 2012 [↗](#)). *Cpo* is a known diapause-associated gene (Kankare et al. 2012 [↗](#)), and we show here that *dpp* is also required for post-diapause fecundity. Tunstall et al showed that *tai* is required for normal olfactory-mediated attraction to food.

We further explored whether IR-expressing ORNs might also be required by inhibiting subsets of IR-expressing cells with TNT. The most remarkable finding was that expressing TNT with *Ir21a-Gal4* resulted in post-diapause death, indicating that *Ir21a-Gal4*-expressing cells are essential for successful exit from dormancy. *Ir21a* is an ionotropic receptor expressed in cells sensitive to cooling (cool cells). Animals mutant for *Ir21a* fail to detect cooling and also fail to avoid both warm and cool temperatures (Budelli et al. 2019 [↗](#)). That cells required for temperature sensation are essential for diapause is perhaps not surprising, however it is at first glance puzzling that inhibiting the *Ir21a*-expressing cells affects diapause because *Ir21a*-expressing cells are located in the arista, which when amputated did not affect post-diapause fecundity in our experiments. Similarly puzzling is the observation that inhibition of the “hot cells,” which are also located in the arista in close proximity to the cool cells, also impairs post-diapause fecundity. One possible explanation for these results could be that cool cells and hot cells inhibit one another such that eliminating cool cell activity actually disinhibits hot cells and vice versa. In fact, Budelli et al (Budelli et al. 2019 [↗](#)) found hot cell spiking in *Ir21a* mutants as well as defects in avoidance of both warm and cool temperatures. If inhibiting cooling-sensitive cells hyperactivates warming-sensitive cells and vice versa, then eliminating both cell types by removal of the arista might be less harmful than inactivating individual cell types.

In summary, we report the first GWAS of diapause, using the stringent assay of the ability to enter diapause, survive, exit diapause, and produce fertile offspring. This analysis has revealed a key role for the olfactory system, as well as specific genes and cells required for diapause lifespan

extension and post-diapause fecundity. Additionally, the 291 candidate diapause-associated genes identified here provide many intriguing candidates for future study.

Methods

Drosophila stocks

The majority of the experiments were carried out using the inbred, sequenced lines of the *Drosophila* Genetic Reference Panel (DGRP) (Huang et al. 2014 [DOI](#)). The RNAi lines and *Gal4* stocks were from BDSC and VDRC (GD44077, SH330386, 27049, GD10268, GD40698, 32034, KK101659, GD36166, 40889, 27508, GD991, GD992, GD21052, KK100412, 54811, GD51936, KK106911, 44661, 53315, 54037, 57840, KK104551, GD17767, 63013, 56867, 80461, KK104056, 35785, KK60102, 9331, 5138, 7019, 23292, 41731, 41728, 51311, 41750, 28838, 35607, 7063, 58989, 51635, 64349, and 7415). *Hot cell-Gal4* (Gallio et al. 2011 [DOI](#)) is a gift from Zuker lab and *Ir21a-Gal4* (Ni et al. 2016 [DOI](#)) is a gift from Craig Montell lab.

Fecundity assay

For the non-diapause fecundity assessment, a minimum of 15 newly-eclosed, not more than 6-hour-old virgin females of each strain were collected under mild CO₂ anesthesia. These females were individually placed into fly food vials containing cornmeal, molasses, agar medium, and yeast. In individual female fly crosses, one female from the DGRP line and two male flies of the same genotype (Canton S) were added to each vial.

In the post-diapause group, a minimum of 20 newly eclosed, not more than 6-hour-old virgin female flies of each strain were collected and promptly transferred to a cold room under diapause conditions (10°C and 8L:16D) for 5 weeks. After this diapause induction period, the flies were shifted to 18°C and 12L:12D for 1 day for temperature acclimation. Subsequently, the flies were transferred to new vials with fresh food medium dusted with yeast and placed at 25°C and 12L:12D for an additional day. Similar to the non-diapause crosses, individual female fly crosses were set up. The flies were allowed to mate and lay eggs for 4 days at 25°C under a normal photoperiod of 12L:12D. After 4 days, the parental flies were discarded, and the resulting progeny were counted on the 16th day from the initial mating start date. The progeny count was limited to the number of adult flies and any black pupae produced by a single female during these 16 days. If female flies failed to survive diapause, the line was given the lowest score of zero, indicating an inability to diapause. Those who died during the 4-day cross period were not considered for scoring. In cases of low replicates due to deaths during diapause, recovery, or cross, the process was repeated to obtain sufficient replicates. The fecundity of flies can vary with the age of fly food, dryness of fly food, age of male flies used to assess the female fecundity, crowding in the bottle, etc. To control all these variables, we have always used fresh fly food (1-week-old) and 1-week-old male flies to reduce the variation. Also, we control the number of flies in collection bottles to 30-40 - about 25-30 females and 10 males and flipped every 3-4 days to a new bottle. We have also observed the fly vials regularly to prevent dryness of food and added water whenever necessary.

Similarly, assessments of non-diapause and post-diapause fecundity were conducted for the RNAi experiments. The post-diapause fecundity was normalized by dividing individual female fly fecundity by average non-diapause fecundity of the same genotype.

Scoring of fecundity

For both the non-diapause and the post-diapause, the fecundity of each strain was calculated based on the average number of progeny produced across all replicates, excluding cases in which the female and/or both males died during the 4-day mating window. To accurately assess the diapause capacity of each strain, we normalized it by dividing the individual post-diapause

fecundity by the average of non-diapause fecundity for each DGRP line. The average of this normalized post-diapause fecundity were then utilized in our GWAS analysis pipeline (Mackay et al. 2012 [\[1\]](#)). This normalization eliminated the basal difference in fecundity among DGRP lines, allowing us to focus on the variability in diapause.

Data analysis

GWAS was carried out using DGRP2 website. We conducted gene ontology enrichment and network analyses based on the top variants ($P < 1e10^{-5}$) associated with the mean post-diapause fecundity/non-diapause fecundity score using the GeneMANIA application in Cytoscape (Shannon et al. 2003 [\[2\]](#); Warde-Farley et al. 2010 [\[3\]](#)). Set Comparison Appyter was used for comparing the diapause-GWAS gene list to other behavior gene lists using 13500 as the background gene list and 0.05 as the significance level to calculate the p-value (Clarke et al. 2021 [\[4\]](#)).

Immunostaining

Immunofluorescence was performed using standard procedures. Briefly, adult female ovarioles were carefully dissected in phosphate-buffered saline (PBS) using a bent tungsten needle, pulling on the stalk region of older egg chambers to minimize damage to the germarium. Ovarioles prepared for immunostaining were fixed in 4% paraformaldehyde (PFA) in PBS for 20 minutes. To prevent sample sticking and facilitate settling, 20 μ l of PBST (PBS+0.2% Triton X100) was added to the fixing solution. After fixation, ovarioles were washed three times for 10 minutes each in PBST and then incubated with primary antibodies at 4°C overnight. The following morning, ovarioles were washed twice for 15 minutes in PBST before incubation with secondary antibodies for at least 2 hours at room temperature. After the removal of secondary antibodies, the samples were washed three times in PBST for 10 minutes each. Hoechst was used as a nuclear stain and added to the second wash solution. The ovarioles were subsequently mounted in Vectashield and stored at 4°C until imaging. All antibody dilutions were prepared in PBST. Due to difficulties in settling diapause samples in the solutions, a 5-minute hold step was introduced on a stand before removing solutions from the tube at each step, consistently followed for all controls as well. The anti-Zpg antibody (rabbit polyclonal) was used at a 1:20000 dilution (Smendziuk et al. 2015 [\[5\]](#)). The antibodies used for identifying Germline Stem Cells (GSCs) are Hts (1B1) and Vasa (DSHB) (Easwaran et al. 2022 [\[6\]](#)).

Lifespan analysis

Virgin female flies were collected and moved to the conditions to assess their lifespan. In the antenna removed fly lifespan measurement, before moving to the diapausing conditions we gave 14 days at optimal conditions (25°C, 12L:12D photoperiod) to heal the wound caused by amputation of the antenna. Flies were kept in food vials with a density of 20 or fewer flies per vial. Deaths were censored daily and all vials were flipped every other day for lifespan measurement at optimal condition (25°C, 12L:12D photoperiod) and once every month for lifespan measurement at diapause condition (10°C, 16L:8D photoperiod) to prevent desiccation and drying up of fly food. The Survival plot was created in Prism using the Kaplan-Meier survival analysis. Survival curves are compared pairwise by the Log-rank (Mantel-Cox) test to obtain the χ^2 - values.

Supplemental Table S1. Non-diapause and post-diapause fecundity in DGRP lines.

Quantification of fecundity of DGRP lines in non-diapausing and post-diapausing conditions.

Supplemental Table S2. Fecundity of Canton S control flies scored at different ages and conditions.

Supplemental Table S3. Gene variants associated with post-diapause fecundity identified by GWAS. Results using the DGRP2 tool to analyze the ratio of post-diapause to non-diapause fecundity.

Supplemental Table S4. Candidate diapause-associated genes identified by GWAS. Genes within 1kb upstream or downstream of the gene variants associated with diapause are candidate diapause-associated genes. The number of different associations are provided along with references for genes previously associated with diapause.

Supplemental Table S5. Comparison of diapause-GWAS gene list to other behavior-associated gene lists. Diapause-GWAS gene lists are compared to other behavior-associated gene lists from different studies (DOIs are included in each tab from where gene lists are obtained). Common genes identified, Venn diagram, and p-values are added in each tab of the Excel sheet.

Supplemental Table S6. RNAi screen of top 15 GWAS-associated genes. Fecundity assessment of RNAi-mediated knockdown of different genes from the GWAS list. Non-diapause and post-diapause fecundity were measured and post-diapause fecundity was normalized by dividing individual post-diapause female fly fecundity by average non-diapause fecundity.

Supplemental Table S7. Tissue specific RNAi of *Dip-γ* and *sbb*. Fecundity assessment of pan neuronal and glial specific RNAi-mediated knockdown of *Dip-γ* and *sbb*.

Supplemental Table S8. Role of antennae in ability of flies to undergo diapause. Fecundity assay of Canton S control flies with antennae removed along with controls.

Supplemental Table S9. Identification of specific cell types involved in diapause in fly antennae. Fecundity assay used to identify diapause modifying cell types in fly antennae by specifically blocking neuronal transmission through the expression of tetanus toxin.

Competing interests

The authors declare no competing interests.

Acknowledgements

We thank Dominique Houston, Mackenzie Kui, Yishi Xu, and Alyssa Chow for technical assistance and members of the lab for discussions. We thank Maddalina Nano for proofreading the manuscript. This work was supported by NIH grant R01AG36907 to D.J.M.

References

Budelli G *et al.* (2019) **Ionotropic receptors specify the morphogenesis of phasic sensors controlling rapid thermal preference in drosophila** *Neuron* **101**:738–747

Chow CY, Wolfner MF, Clark AG (2013) **Using natural variation in *Drosophila* to discover previously unknown endoplasmic reticulum stress genes** *Proc Natl Acad Sci USA* **110**:9013–9018

Clarke DJB *et al.* (2021) **Appyters: Turning Jupyter Notebooks into data-driven web apps** *Patterns (N Y)* **2**

- Davis SM, Thomas AL, Nomie KJ, Huang L, Dierick HA (2014) **Tailless and Atrophia control *Drosophila* aggression by regulating neuropeptide signalling in the pars intercerebralis** *Nat Commun* **5**
- Deng L, Li C, Chen L, Liu Y, Hou R, Zhou X (2018) **Research advances on embryonic diapause in mammals** *Anim Reprod Sci* **198**:1–10
- Denlinger DL (2023) **Insect diapause: from a rich history to an exciting future** *J Exp Biol* **226**
- Durham MF, Magwire MM, Stone EA, Leips J (2014) **Genome-wide analysis in *Drosophila* reveals age-specific effects of SNPs on fitness traits** *Nat Commun* **5**
- Easwaran S, Montell DJ (2023) **The molecular mechanisms of diapause and diapause-like reversible arrest** *Biochem Soc Trans* **51**:1847–1856
- Easwaran S, Van Ligten M, Kui M, Montell DJ (2022) **Enhanced germline stem cell longevity in *Drosophila* diapause** *Nat Commun* **13**
- Fontana L, Partridge L, Longo VD (2010) **Extending healthy life span--from yeast to humans** *Science* **328**:321–326
- Gallio M, Ofstad TA, Macpherson LJ, Wang JW, Zuker CS (2011) **The coding of temperature in the *Drosophila* brain** *Cell* **144**:614–624
- Garlapow ME, Huang W, Yarboro MT, Peterson KR, Mackay TFC (2015) **Quantitative Genetics of Food Intake in *Drosophila melanogaster*** *PLoS ONE* **10**
- Grubbs N, Leach M, Su X, Petrisko T, Rosario JB, Mahaffey JW (2013) **New components of *Drosophila* leg development identified through genome wide association studies** *PLoS ONE* **8**
- Hand SC, Denlinger DL, Podrabsky JE, Roy R (2016) **Mechanisms of animal diapause: recent developments from nematodes, crustaceans, insects, and fish** *Am J Physiol Regul Integr Comp Physiol* **310**:R1193–211
- Harbison ST, Kumar S, Huang W, McCoy LJ, Smith KR, Mackay TFC (2019) **Genome-Wide Association Study of Circadian Behavior in *Drosophila melanogaster*** *Behav Genet* **49**:60–82
- Harbison ST, McCoy LJ, Mackay TFC (2013) **Genome-wide association study of sleep in *Drosophila melanogaster*** *BMC Genomics* **14**
- Hidalgo S, Anguiano M, Tabuloc CA, Chiu JC (2023) **Seasonal cues act through the circadian clock and pigment-dispersing factor to control EYES ABSENT and downstream physiological changes** *Curr Biol* **33**:675–687
- Horváth B, Betancourt AJ, Kalinka AT (2016) **A novel method for quantifying the rate of embryogenesis uncovers considerable genetic variation for the duration of embryonic development in *Drosophila melanogaster*** *BMC Evol Biol* **16**
- Horváth B, Kalinka AT (2018) **The genetics of egg retention and fertilization success in *Drosophila*: One step closer to understanding the transition from facultative to obligate viviparity** *Evolution* **72**:318–336

Huang W *et al.* (2014) **Natural variation in genome architecture among 205 *Drosophila melanogaster* Genetic Reference Panel lines** *Genome Res* **24**:1193–1208

Hussein AM, Balachandar N, Mathieu J, Ruohola-Baker H (2022) **Molecular Regulators of Embryonic Diapause and Cancer Diapause-like State** *Cells* **11**

Hutfilz C (2022) **Endocrine regulation of lifespan in insect diapause** *Front Physiol* **13**

Kankare M, Salminen TS, Lampinen H, Hoikkala A (2012) **Sequence variation in couch potato and its effects on life-history traits in a northern malt fly, *Drosophila montana*** *J Insect Physiol* **58**:256–264

Katzenberger RJ, Chtarbanova S, Rimkus SA, Fischer JA, Kaur G, Seppala JM, Swanson LC, Zajac JE, Ganetzky B, Wassarman DA (2015) **Death following traumatic brain injury in *Drosophila* is associated with intestinal barrier dysfunction** *eLife* **4**

Kubrak OI, Kučerová L, Theopold U, Nässel DR (2014) **The sleeping beauty: how reproductive diapause affects hormone signaling, metabolism, immune response and somatic maintenance in *Drosophila melanogaster*** *PLoS ONE* **9**

Lin Y-H, Zhu H (2021) **A malignant case of arrested development: cancer cell dormancy mimics embryonic diapause** *Cancer Cell* **39**:142–144

Li J *et al.* (2020) **Cell-Surface Proteomic Profiling in the Fly Brain Uncovers Wiring Regulators** *Cell* **180**:373–386

Mackay TFC, Huang W (2018) **Charting the genotype-phenotype map: lessons from the *Drosophila melanogaster* Genetic Reference Panel** *Wiley Interdiscip Rev Dev Biol* **7**

Mackay TFC *et al.* (2012) **The *Drosophila melanogaster* Genetic Reference Panel** *Nature* **482**:173–178

Meiselman MR, Alpert MH, Cui X, Shea J, Gregg I, Gallio M, Yapici N (2022) **Recovery from cold-induced reproductive dormancy is regulated by temperature-dependent AstC signaling** *Curr Biol* **32**:1362–1375

Meyerhof GT, Easwaran S, Bontempo A, Montell C, Montell DJ (2024) **Altered circadian rhythm, sleep, and rhodopsin 7-dependent shade preference during diapause in *Drosophila melanogaster*** *PNAS*

Montell C (2021) ***Drosophila* sensory receptors-a set of molecular Swiss Army Knives** *Genetics* **217**:1–34

Montgomery SL, Vorobjeikina D, Huang W, Mackay TFC, Anholt RRH, Rand MD (2014) **Genome-wide association analysis of tolerance to methylmercury toxicity in *Drosophila* implicates myogenic and neuromuscular developmental pathways** *PLoS ONE* **9**

Morozova TV, Huang W, Pray VA, Whitham T, Anholt RRH, Mackay TFC (2015) **Polymorphisms in early neurodevelopmental genes affect natural variation in alcohol sensitivity in adult *Drosophila*** *BMC Genomics* **16**

Myachina F, Bosshardt F, Bischof J, Kirschmann M, Lehner CF (2017) ***Drosophila* β -Tubulin 97EF is upregulated at low temperature and stabilizes microtubules** *Development* **144**:4573–4587

- Ni L, Bronk P, Chang EC, Lowell AM, Flam JO, Panzano VC, Theobald DL, Griffith LC, Garrity PA (2013) **A gustatory receptor paralogue controls rapid warmth avoidance in *Drosophila*** *Nature* **500**:580–584
- Ni L, Klein M, Svec KV, Budelli G, Chang EC, Ferrer AJ, Benton R, Samuel AD, Garrity PA (2016) **The Ionotropic Receptors IR21a and IR25a mediate cool sensing in *Drosophila*** *eLife* **5**
- Saunders DS, Henrich VC, Gilbert LI (1989) **Induction of diapause in *Drosophila melanogaster*: photoperiodic regulation and the impact of arrhythmic clock mutations on time measurement** *Proc Natl Acad Sci USA* **86**:3748–3752
- Saunders DS, Richard DS, Applebaum SW, Ma M, Gilbert LI (1990) **Photoperiodic diapause in *Drosophila melanogaster* involves a block to the juvenile hormone regulation of ovarian maturation** *Gen Comp Endocrinol* **79**:174–184
- Schmidt PS, Zhu C-T, Das J, Batavia M, Yang L, Eanes WF (2008) **An amino acid polymorphism in the couch potato gene forms the basis for climatic adaptation in *Drosophila melanogaster*** *Proc Natl Acad Sci USA* **105**:16207–16211
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003) **Cytoscape: a software environment for integrated models of biomolecular interaction networks** *Genome Res* **13**:2498–2504
- Shimozono M, Osaka J, Kato Y, Araki T, Kawamura H, Takechi H, Hakeda-Suzuki S, Suzuki T (2019) **Cell surface molecule, Klingon, mediates the refinement of synaptic specificity in the *Drosophila* visual system** *Genes Cells* **24**:496–510
- Shorter J, Couch C, Huang W, Carbone MA, Peiffer J, Anholt RRH, Mackay TFC (2015) **Genetic architecture of natural variation in *Drosophila melanogaster* aggressive behavior** *Proc Natl Acad Sci USA* **112**:E3555–63
- Sim C, Denlinger DL (2013) **Insulin signaling and the regulation of insect diapause** *Front Physiol* **4**
- Smendziuk CM, Messenberg A, Vogl AW, Tanentzapf G (2015) **Bi-directional gap junction-mediated soma-germline communication is essential for spermatogenesis** *Development* **142**:2598–2609
- Tatar M, Yin C (2001) **Slow aging during insect reproductive diapause: why butterflies, grasshoppers and flies are like worms** *Exp Gerontol* **36**:723–738
- Tunstall NE, Herr A, de Bruyne M, Warr CG (2012) **A screen for genes expressed in the olfactory organs of *Drosophila melanogaster* identifies genes involved in olfactory behaviour** *PLoS ONE* **7**
- Warde-Farley D *et al.* (2010) **The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function** *Nucleic Acids Res* **38**:W214–20
- Zhang Q, Denlinger DL (2011) **Elevated couch potato transcripts associated with adult diapause in the mosquito *Culex pipiens*** *J Insect Physiol* **57**:620–627

Zhao X, Bergland AO, Behrman EL, Gregory BD, Petrov DA, Schmidt PS (2016) **Global Transcriptional Profiling of Diapause and Climatic Adaptation in *Drosophila melanogaster*** *Mol Biol Evol* **33**:707–720

Zhou S, Morozova TV, Hussain YN, Luoma SE, McCoy L, Yamamoto A, Mackay TFC, Anholt RRH (2016) **The Genetic Basis for Variation in Sensitivity to Lead Toxicity in *Drosophila melanogaster*** *Environ Health Perspect* **124**:1062–1070

Editors

Reviewing Editor

Jiwon Shim

Hanyang University, Seoul, Korea, the Republic of

Senior Editor

Utpal Banerjee

University of California, Los Angeles, Los Angeles, United States of America

Reviewer #1 (Public Review):

Summary:

The paper begins with phenotyping the DGRP for post-diapause fecundity, which is used to map genes and variants associated with fecundity. There are overlaps with genes mapped in other studies and also functional enrichment of pathways including most surprisingly neuronal pathways. This somewhat explains the strong overlap with traits such as olfactory behaviors and circadian rhythm. The authors then go on to test genes by knocking them down effectively at 10 degrees. Two genes, *Dip-gamma* and *sbb*, are identified as significantly associated with post-diapause fecundity, and they also find the effects to be specific to neurons. They further show that the neurons in the antenna but not the arista are required for the effects of *Dip-gamma* and *sbb*. They show that removing the antenna has a diapause-specific lifespan-extending effect, which is quite interesting. Finally, ionotropic receptor neurons are shown to be required for the diapause-associated effects.

Strengths and Weaknesses:

Overall I find the experiments rigorously done and interpretations sound. I have no further suggestions except an ANOVA to estimate the heritability of the post-diapause fecundity trait, which is routinely done in the DGRP and offers a global parameter regarding how reliable phenotyping is. A minor point is I cannot find how many DGRP lines are used.

<https://doi.org/10.7554/eLife.98142.1.sa2>

Reviewer #2 (Public Review):

Summary

In this study, Easwaran and Montell investigated the molecular, cellular, and genetic basis of adult reproductive diapause in *Drosophila* using the *Drosophila* Genetic Reference Panel (DGRP). Their GWAS revealed genes associated with variation in post-diapause fecundity across the DGRP and performed RNAi screens on these candidate genes. They also analyzed the functional implications of these genes, highlighting the role of genes involved in neural and germline development. In addition, in conjunction with other GWAS results, they noted the importance of the olfactory system within the nervous system, which was supported by

genetic experiments. Overall, their solid research uncovered new aspects of adult diapause regulation and provided a useful reference for future studies in this field.

Strengths:

The authors used whole-genome sequenced DGRP to identify genes and regulatory mechanisms involved in adult diapause. The first *Drosophila* GWAS of diapause successfully uncovered many QTL underlying post-diapause fecundity variations across DGRP lines. Gene network analysis and comparative GWAS led them to reveal a key role for the olfactory system in diapause lifespan extension and post-diapause fecundity.

Weaknesses:

(1) I suspect that there may be variation in survivorship after long-term exposure to cold conditions (10°C, 35 days), which could also be quantified and mapped using genome-wide association studies (GWAS). Since blocking *Ir21a* neuronal transmission prevented flies from exiting diapause, it is possible that natural genetic variation could have a similar effect, influencing the success rate of exiting diapause and post-diapause mortality. If there is variation in this trait, could it affect post-diapause fecundity? I am concerned that this could be a confounding factor in the analysis of post-diapause fecundity. However, I also believe that understanding phenotypic variation in this trait itself could be significant in regulating adult diapause.

(2) On p.10, the authors conclude that "*Dip-γ* and *sbb* are required in neurons for successful diapause, consistent with the enrichment of this gene class in the diapause GWAS." While I acknowledge that the results support their neuronal functions, I remain unconvinced that these genes are required for "successful diapause". According to the RNAi scheme (Figure 4I), *Dip-γ* and *sbb* are downregulated only during the post-diapause period, but still show a significant effect, comparable to that seen in the *nSyb* Gal4 RNAi lines (Figure 4K). In addition, two other RNAi lines (SH330386, 80461) that did not show lethality did not affect post-diapause fecundity. Notably, RNAi (27049, KK104056) substantially reduced non-diapause fecundity, suggesting impairment of these genes affects fecundity in general regardless of diapause experience. Therefore, the reduced post-diapause fecundity observed may be a result of this broader effect on fecundity, particularly in a more "sensitized" state during the post-diapause period, rather than a direct regulation of adult diapause by these genes.

(3) The authors characterized 546 genetic variants and 291 genes associated with phenotypic variation across DGRP lines but did not prioritize them by significance. They did prioritize candidate genes with multiple associated variants (p.9 "Genes with multiple SNPs are good candidates for influencing diapause traits."), but this is not a valid argument, likely due to a misunderstanding of LD among variants in the same gene. A gene with one highly significantly associated variant may be more likely to be the causal gene in a QTL than a gene with many weakly associated variants in LD. I recommend taking significance into account in the analysis.

<https://doi.org/10.7554/eLife.98142.1.sa1>

Reviewer #3 (Public Review):

Summary:

Drosophila melanogaster of North America overwinters in a state of reproductive diapause. The authors aimed to measure 'successful' *D. melanogaster* reproductive diapause and reveal loci that impact this quantitative trait. In practice, the authors quantified the number of eggs

produced by a female after she exited 35 days of diapause. The authors claim that genes involved with olfaction in part contribute to some of the variation in this trait.

Strengths:

The work used the power platform of the fly DRGP/GWAS. The work tried to verify some of the candidate loci with targeted gene manipulations.

Weaknesses:

Some context is needed. Previous work from 2001 established that *D. melanogaster* reproductive diapause in the laboratory suspends adult aging but reduces post-diapause fecundity. The work from 2001 showed the extent fecundity is reduced is proportional to diapause duration. As well, the 2001 data showed short diapause periods used in the current submission reduce fecundity only in the first days following diapause termination; after this time fecundity is greater in the post-diapause females than in the non-diapause controls.

In this context, the submission fails to offer a meaningful concept for what constitutes 'successful diapause'. There is no biological rationale or relationship to the known patterns of post-diapause fecundity. The phenotype is biologically ambiguous.

I have a serious concern about the antenna-removal design. These flies were placed on cool/short days two weeks after surgery. Adults at this time will not enter diapause, which must be induced soon after eclosion. Two-week-old adults will respond to cool temperatures by 'slowing down', but they will continue to age on a time scale of day-degrees. This is why the control group shows age-dependent mortality, which would not be seen in truly diapaused adults. Loss of antennae increases the age-dependent mortality of these cold adults, but this result does not reflect an impact on diapause.

- Appraisal of whether the authors achieved their aims, and whether the results support their conclusions.

The work falls well short of its aim because the concept of 'successful diapause' is not biologically established. The paper studies post-diapause fecundity, and we don't know what that means. The loci identified in this analysis segregate for a minimally constructed phenotype. The results and conclusions are orthogonal.

- The likely impact of the work on the field, and the utility of the methods and data to the community.

The work will have little likely impact. Its phenotype and operational methods are weakly developed. It lacks insight based on the primary literature on post-diapause. The community of insect diapause investigators are not likely to use the data or conclusions to understand beneficial or pest insects, or the impact of a changing climate on how they over-winter.

<https://doi.org/10.7554/eLife.98142.1.sa0>

Author response:

eLife assessment

*This useful study shows how genetic variation is associated with fecundity following a period of reproductive diapause in female *Drosophila*. The work identifies the olfactory system as central to successful diapause with associated changes in longevity and fecundity. While the genetic screening and methods used are solid, the approach to assessing diapause is incomplete and could benefit from additional orthogonal experiments.*

Response: We agree that, as with most studies, additional follow-up work will be informative.

Public Reviews:

Reviewer #1 (Public Review):

Summary:

The paper begins with phenotyping the DGRP for post-diapause fecundity, which is used to map genes and variants associated with fecundity. There are overlaps with genes mapped in other studies and also functional enrichment of pathways including most surprisingly neuronal pathways. This somewhat explains the strong overlap with traits such as olfactory behaviors and circadian rhythm. The authors then go on to test genes by knocking them down effectively at 10 degrees. Two genes, Dip-gamma and sbb, are identified as significantly associated with post-diapause fecundity, and they also find the effects to be specific to neurons. They further show that the neurons in the antenna but not the arista are required for the effects of Dip-gamma and sbb. They show that removing the antenna has a diapause-specific lifespan-extending effect, which is quite interesting. Finally, ionotropic receptor neurons are shown to be required for the diapause-associated effects.

Strengths and Weaknesses:

Overall I find the experiments rigorously done and interpretations sound. I have no further suggestions except an ANOVA to estimate the heritability of the post-diapause fecundity trait, which is routinely done in the DGRP and offers a global parameter regarding how reliable phenotyping is. A minor point is I cannot find how many DGRP lines are used.

Response: Thank you for the suggestions. We screened 193 lines and we will add that information to the methods.

Additionally, we will add the heritability estimate of the post-diapause fecundity trait.

Reviewer #2 (Public Review):

Summary

In this study, Easwaran and Montell investigated the molecular, cellular, and genetic basis of adult reproductive diapause in Drosophila using the Drosophila Genetic Reference Panel (DGRP). Their GWAS revealed genes associated with variation in post-diapause fecundity across the DGRP and performed RNAi screens on these candidate genes. They also analyzed the functional implications of these genes, highlighting the role of genes involved in neural and germline development. In addition, in conjunction with other GWAS results, they noted the importance of the olfactory system within the nervous system, which was supported by genetic experiments. Overall, their solid research uncovered new aspects of adult diapause regulation and provided a useful reference for future studies in this field.

Strengths:

The authors used whole-genome sequenced DGRP to identify genes and regulatory mechanisms involved in adult diapause. The first Drosophila GWAS of diapause successfully uncovered many QTL underlying post-diapause fecundity variations across DGRP lines. Gene network analysis and comparative GWAS led them to reveal a key role for the olfactory system in diapause lifespan extension and post-diapause fecundity.

Weaknesses:

(1) I suspect that there may be variation in survivorship after long-term exposure to cold conditions (10°C, 35 days), which could also be quantified and mapped using genome-wide association studies (GWAS). Since blocking *Ir21a* neuronal transmission prevented flies from exiting diapause, it is possible that natural genetic variation could have a similar effect, influencing the success rate of exiting diapause and post-diapause mortality. If there is variation in this trait, could it affect post-diapause fecundity? I am concerned that this could be a confounding factor in the analysis of post-diapause fecundity. However, I also believe that understanding phenotypic variation in this trait itself could be significant in regulating adult diapause.

Response: We agree that it is possible that the ability to endure cool temperatures per se may influence post-diapause fecundity. However, cool temperature is the essential diapause-inducing condition in *Drosophila*, so it is not obvious how to separate those effects experimentally, and we agree that phenotypic variation in the cool-sensitivity trait itself could be significant in regulating diapause.

(2) On p.10, the authors conclude that "*Dip-γ* and *sbb* are required in neurons for successful diapause, consistent with the enrichment of this gene class in the diapause GWAS." While I acknowledge that the results support their neuronal functions, I remain unconvinced that these genes are required for "successful diapause". According to the RNAi scheme (Figure 4I), *Dip-γ* and *sbb* are downregulated only during the post-diapause period, but still show a significant effect, comparable to that seen in the *nSyb Gal4* RNAi lines (Figure 4K).

Response: Our definition of successful diapause is the ability to produce viable adult progeny post-diapause, which requires that the flies enter, maintain, and exit diapause, alive and fertile. We will restate our conclusion to say that *Dip-γ* and *sbb* are required for post-diapause fecundity.

In addition, two other RNAi lines (SH330386, 80461) that did not show lethality did not affect post-diapause fecundity.

Response: We interpret those results to mean that those RNAi lines were not effective since *Dip-γ* and *sbb* are known to be essential.

Notably, RNAi (27049, KK104056) substantially reduced non-diapause fecundity, suggesting impairment of these genes affects fecundity in general regardless of diapause experience. Therefore, the reduced post-diapause fecundity observed may be a result of this broader effect on fecundity, particularly in a more "sensitized" state during the post-diapause period, rather than a direct regulation of adult diapause by these genes.

Response: Ubiquitous expression of RNAi lines #27049 or #KK104056 was lethal, so we included the tubGAL80ts repressor to prevent RNAi from taking effect during development. Flies had to be shifted to 30 °C to inactivate the repressor and thereby activate the RNAi. At 30 °C, fecundity of the controls (GFP RNAi lines #9331, KK60102) were also lower (average non-diapause fecundity = 12 and 19 respectively) and similar to #27049 or #KK104056. We also assessed the knockdown using Repo GAL4 and nSyb GAL4 and did not find a significant difference/decline in the non diapause fecundity for #27049 and #KK104056 as compared to a nonspecific RNAi control (#54037).

(3) The authors characterized 546 genetic variants and 291 genes associated with phenotypic variation across DGRP lines but did not prioritize them by significance. They

did prioritize candidate genes with multiple associated variants (p.9 "Genes with multiple SNPs are good candidates for influencing diapause traits."), but this is not a valid argument, likely due to a misunderstanding of LD among variants in the same gene. A gene with one highly significantly associated variant may be more likely to be the causal gene in a QTL than a gene with many weakly associated variants in LD. I recommend taking significance into account in the analysis.

We agree with the reviewer, and in Supplemental Table S3 we list top-associated SNPs in order from the lowest (most significant) p-value. Most of the top-associated genes from this analysis were uncharacterized CG numbers for which there were insufficient tools available for validation purposes. Nevertheless, there is overlap amongst the highly significant genes by p-value and those with multiple SNPs. Amongst the top 15 genes with multiple associated SNPs CG18636 & CR15280 ranked 3rd by p-value, CG7759 ranked 4th, CG42732 ranked 10th, and Drip ranked 30th (all above the conservative Bonferroni threshold of 4.8×10^{-8}) while three Sbb-associated SNPs also appear in Table 3 above the standard 5×10^{-5} threshold.

Reviewer #3 (Public Review):

Summary:

Drosophila melanogaster of North America overwinters in a state of reproductive diapause. The authors aimed to measure 'successful' D. melanogaster reproductive diapause and reveal loci that impact this quantitative trait. In practice, the authors quantified the number of eggs produced by a female after she exited 35 days of diapause. The authors claim that genes involved with olfaction in part contribute to some of the variation in this trait.

Strengths:

The work used the power platform of the fly DRGP/GWAS. The work tried to verify some of the candidate loci with targeted gene manipulations.

Weaknesses:

Some context is needed. Previous work from 2001 established that D. melanogaster reproductive diapause in the laboratory suspends adult aging but reduces post-diapause fecundity. The work from 2001 showed the extent fecundity is reduced is proportional to diapause duration. As well, the 2001 data showed short diapause periods used in the current submission reduce fecundity only in the first days following diapause termination; after this time fecundity is greater in the post-diapause females than in the non-diapause controls.

Response: The 2001 paper by Tatar et al. reports the number of eggs laid after 3, 6, or 9 weeks in diapause conditions. Thus the diapause conditions used in this study (35 days or 5 weeks) are neither short nor long, rather intermediate. Does the reviewer have a specific concern?

In this context, the submission fails to offer a meaningful concept for what constitutes 'successful diapause'. There is no biological rationale or relationship to the known patterns of post-diapause fecundity. The phenotype is biologically ambiguous.

Response: We have unambiguously defined successful diapause as the ability to produce viable adult progeny post-diapause. Other groups have measured % of flies that arrest ovarian development or % of post-diapause flies with mature eggs in the ovary, or # eggs laid post-diapause; however we suggest that # of viable adult progeny produced post-diapause is more meaningful than the other measurements from the point of view of perpetuating the species.

I have a serious concern about the antenna-removal design. These flies were placed on cool/short days two weeks after surgery. Adults at this time will not enter diapause, which must be induced soon after eclosion. Two-week-old adults will respond to cool temperatures by 'slowing down', but they will continue to age on a time scale of day-degrees. This is why the control group shows age-dependent mortality, which would not be seen in truly diapaused adults. Loss of antennae increases the age-dependent mortality of these cold adults, but this result does not reflect an impact on diapause.

Response: The reviewer has a point. We carried out the lifespan study under two different conditions: either by removing the antenna and moving the flies directly to 10 °C or by removing the antenna and allowing a “wound healing” period prior to moving the flies to 10 °C (out of concern that the flies might have died quickly because wound healing may be impaired at 10 °C). In both cases, lifespan was shortened. We will add a discussion of the technical limitations of this experiment.

• *Appraisal of whether the authors achieved their aims, and whether the results support their conclusions.*

The work falls well short of its aim because the concept of 'successful diapause' is not biologically established. The paper studies post-diapause fecundity, and we don't know what that means. The loci identified in this analysis segregate for a minimally constructed phenotype. The results and conclusions are orthogonal.

Response: It is unclear to us why the reviewer has such a negative opinion of measuring post-diapause fecundity, specifically the ability to produce viable progeny post-diapause. The value of this measurement seems obvious from the point of view of perpetuating the species.

• *The likely impact of the work on the field, and the utility of the methods and data to the community.*

The work will have little likely impact. Its phenotype and operational methods are weakly developed. It lacks insight based on the primary literature on post-diapause. The community of insect diapause investigators are not likely to use the data or conclusions to understand beneficial or pest insects, or the impact of a changing climate on how they over-winter.

Response: The reviewer has not explained why his/her opinion is so negative.

<https://doi.org/10.7554/eLife.98142.1.sa4>