

POMC neurons control fertility through differential signaling of MC4R in Kisspeptin neurons

Reviewed Preprint

v2 • January 24, 2025

Revised by authors

Reviewed Preprint

v1 • September 11, 2024

Rajae Talbi, Todd L Stincic, Kaitlin Ferrari, Choi Ji Hae, Karol Walec, Elizabeth Medve, Achi Gerutshang, Silvia León, Elizabeth A McCarthy, Oline K Rønnekleiv, Martin J Kelly , Víctor M Navarro 

Harvard Medical School, Boston, United States • Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States • Division of Neuroscience, Oregon National Primate Research Center, Beaverton, United States • Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, United States • Harvard Program in Neuroscience, Boston, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

The study presents **compelling** evidence that the melanocortin system originating in the arcuate nucleus of the hypothalamus plays a crucial role in puberty onset, representing a significant advance in our understanding of reproductive biology. The research employs innovative approaches and benefits from the combined expertise of two respected laboratories, enhancing the robustness of the findings. Given the potential impact on human health and the strength of the evidence presented, this **fundamental** work will likely influence the field substantially and may inform future clinical applications.

<https://doi.org/10.7554/eLife.100722.2.sa4>

Abstract

Inactivating mutations in the melanocortin 4 receptor (*MC4R*) gene cause monogenic obesity. Interestingly, female patients also display various degrees of reproductive disorders, in line with the subfertile phenotype of *MC4RKO* female mice. However, the cellular mechanisms by which *MC4R* regulates reproduction are unknown. *Kiss1* neurons directly stimulate gonadotropin-releasing hormone (GnRH) release through two distinct populations; the *Kiss1*^{ARH} neurons, controlling GnRH pulses, and the sexually dimorphic *Kiss1*^{AVPV/PeN} neurons controlling the preovulatory LH surge. Here, we show that *Mc4r* expressed in *Kiss1* neurons regulates fertility in females. *In vivo*, deletion of *Mc4r* from *Kiss1* neurons in female mice replicates the reproductive impairments of *MC4RKO* mice without inducing obesity. Conversely, reinsertion of *Mc4r* in *Kiss1* neurons of *MC4R* null mice restores estrous cyclicity and LH pulsatility without reducing their obese phenotype. *In vitro*, we dissect the specific action of *MC4R* on *Kiss1*^{ARH} vs *Kiss1*^{AVPV/PeN} neurons and show that *MC4R* activation excites *Kiss1*^{ARH} neurons through direct synaptic actions. In contrast, *Kiss1*^{AVPV/PeN} neurons are normally inhibited by *MC4R* activation except under elevated estradiol levels, thus facilitating the activation of *Kiss1*^{AVPV/PeN} neurons to induce the LH surge driving ovulation

in females. Our findings demonstrate that POMC^{ARH} neurons acting through MC4R, directly regulate reproductive function in females by stimulating the “pulse generator” activity of Kiss1^{ARH} neurons and restricting the activation of Kiss1^{AVPV/PeN} neurons to the time of the estradiol-dependent LH surge, and thus unveil a novel pathway of the metabolic regulation of fertility by the melanocortin system.

Introduction

Obesity rates have skyrocketed in Western societies in the last decades, resulting in an alarming rise in comorbidities that place a significant burden on healthcare systems. The increase in obesity correlates with a decrease in fertility rates, leading to conception challenges that are experienced by approximately 15% of couples in the United States currently (1 [↗](#)).

The melanocortin 4 receptor (MC4R) binds i) α-melanocyte stimulating hormone (αMSH), an agonist product of the pro-opiomelanocortin (*Pomc*) gene, and ii) the inverse agonist, agouti-related peptide (AgRP), to regulate food intake and energy expenditure (2 [↗](#), 3 [↗](#)). While the role of MC4R on food intake is largely mediated by neurons located in the paraventricular nucleus of the hypothalamus (PVN) (4 [↗](#)), its expression in the brain is widespread (5 [↗](#)) with the specific function of the different MC4R-expressing neurons in areas beyond the PVN remaining to be fully explored. Inactivating mutations in *MC4R* are a leading cause of monogenic obesity in humans (6 [↗](#)) and cause excessive obesity and hyperphagia in mice (7 [↗](#)). Scant evidence in humans shows an association between *MC4R* mutations and higher incidence of hypogonadotropic hypogonadism (HH) (8 [↗](#)), alterations in the timing of puberty onset (9 [↗](#)), and polycystic ovary syndrome (PCOS) (10 [↗](#)); however, conflicting reports exist (6 [↗](#)). Therefore, the role of MC4R signaling in reproductive function in humans remains controversial despite the clear association found in mice. Supporting this role of MC4R, *Mc4r* null mice display an array of reproductive abnormalities that largely affects females, characterized by irregular estrous cycles, disrupted luteinizing hormone (LH) secretion, reduced corpora lutea and reduced fertility (11 [↗](#)–13 [↗](#)). Further evidence from mice demonstrates that MC4R agonists robustly increase LH release in a kisspeptin-dependent manner (14 [↗](#)).

Kisspeptin is the most potent gonadotropin releasing hormone (GnRH) secretagogue known to date and it is mainly produced in two distinct neuronal populations. Kiss1 neurons of the arcuate nucleus of the hypothalamus (Kiss1^{ARH}), present in both sexes, control the pulsatile (tonic) release of GnRH, and sex steroids attenuate their release of kisspeptin. Kiss1 neurons of the anteroventral periventricular continuum area (Kiss1^{AVPV/PeN}), are predominantly present in females, and are responsible for generating the preovulatory GnRH/LH surge essential for ovulation (15 [↗](#)). Despite these critical roles of both populations of Kiss1 neurons for reproduction, the underlying mechanisms that determine how each Kiss1 population responds differently to sex steroids to regulate the GnRH tonic vs surge release remains unresolved.

In rodents, compelling evidence indicates a close interaction between Kiss1 neurons and the melanocortin system: 1) fibers from POMC neurons in the arcuate nucleus (POMC^{ARH}) project to and juxtapose Kiss1^{ARH} neurons (14 [↗](#)); 2) the melanocortin signaling through MC4R contributes to the permissive role of leptin on puberty onset (14 [↗](#), 16 [↗](#), 17 [↗](#)); and 3) MC4R expression on both Kiss1^{ARH} (18 [↗](#)–20 [↗](#)) and Kiss1^{AVPV/PeN} (18 [↗](#), 21 [↗](#)). Altogether, this evidence suggests a clear role for MC4R in Kiss1 neurons, in the control of reproductive maturation and fertility.

In this study, we investigated the contribution of MC4R signaling versus obesity *per se* in the etiology of the reproductive impairments observed in *Mc4r* null mice, which could explain similar impairments observed in humans. Using genetic mouse models with specific deletion or reinsertion of *Mc4r* in Kiss1 neurons, we show that MC4R action in Kiss1 neurons is necessary for

normal reproductive function in female mice. Whole cell voltage clamp recordings evidenced an excitatory action of MC4R on Kiss1^{ARH} neurons and an estradiol-dependent inhibitory action on Kiss1^{AVPV/PeN} neurons, with important implications for the timing of the preovulatory LH surge.

Results

Mc4r within Kiss1 neurons determines the timing of puberty onset in females

The expression of *Mc4r* within Kiss1^{ARH} and Kiss1^{AVPV/PeN} neurons has already been described elsewhere (18–21), suggesting a role for MC4R in the regulation of fertility. To further investigate the specific role of MC4R in Kiss1 neurons, we generated a Kiss1-specific MC4R knockout mouse model (Kiss1-MC4RKO). The specific deletion of *Mc4r* from Kiss1 neurons, as well as the absence of global recombination, were confirmed through RNAscope. While Kiss1 neurons of Kiss1-MC4RKO mice lack *Mc4r* transcript compared to their control littermates (Figure S1A), *Mc4r* was detectable in the PVN of all mice, which is a major site of *Mc4r* expression in the brain in the regulation of metabolism, therefore supporting the specific deletion of *Mc4r* only within Kiss1 neurons (Figure S1B).

Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening (VO) and first estrus (FE). Kiss1-MC4RKO females showed a significant advancement in the age of VO ($P=0.0150$, Kiss1-MC4RKO: 24.67 ± 0.39 vs controls: 26.32 ± 0.53) and FE ($P=0.0341$, Kiss1-MC4RKO: 32.63 ± 1.4 vs controls: 37.84 ± 1.90) compared to their littermate controls (Figure 1A, B). Body weight of Kiss1-MC4RKO females at the age of puberty onset was similar between groups ($P=0.2596$, Kiss1-MC4RKO: 12.41 ± 0.30 vs controls: 12.89 ± 0.26) (Figure 1C). These data suggest that melanocortin signaling on Kiss1 neurons participates in the timing of puberty onset in females. Since the role of the melanocortin system on puberty onset is largely unexplored, we evaluated the expression profile of the main components of this system in the hypothalami of WT female mice at postnatal days (PND) 10, 15, 22 and 30. Interestingly, expressions of *Agrp*, *Mc3r* and *Mc4r* were significantly lower at the time of puberty onset (PND30) compared to earlier developmental ages (Figure 1D). This supports our *in vivo* data and suggests that a decrease in the hypothalamic melanocortin tone drives puberty onset, in line with the advancement in the age of VO and FE observed in Kiss1-MC4RKO female mice (Figure 1A, B).

MC4R in Kiss1 neurons is required for female reproduction

Because Kiss1-MC4RKO females showed altered puberty onset, we further investigated their reproductive phenotype. Interestingly, Kiss1-MC4RKO female mice displayed normal BW throughout the time of the study (up to PND150) (Figure 2A); however, they presented with irregular estrous cycles with predominantly more time spent in diestrus ($P=0.0001$) and less time spent in estrus ($P<0.0001$) than their control littermates (Figure 2B, C). The assessment of the pulsatile release of LH every 10 min over 180 min revealed no change in the total number of pulses between groups but a significant increase in basal LH release ($P=0.04$) in Kiss1-MC4RKO female mice (0.42 ± 0.03) compared to controls (0.34 ± 0.02) (Figure 2D–H). The analysis of the gene expression of the “KNDy” systems in the ARH, which control the GnRH pulse generator (15), revealed normal expression levels of *Kiss1*, *Tac2* and *Tacr3*, but significantly lower expression of *Pdyn* ($P=0.0067$, Kiss1-MC4RKO: 0.78 ± 0.038 vs controls: 1.00 ± 0.019) (Figure 2I–L). The conserved expression of *Kiss1*, *Tac2* and *Tacr3* correlates with the preserved LH pulse frequency and amplitude in Kiss1-MC4RKO mice, while the lower inhibitory tone of dynorphin (Figure 2I) correlates with the higher basal LH levels (Figure 2E). However, the mRNA expression of these genes, alone, does not necessarily reflect changes of their activities. To assess the contribution of MC4R signaling in Kiss1 neurons to the induction of ovulation through the preovulatory LH surge,




Figure 1.

***Mc4r* expressed in Kiss1 neurons determines the timing of puberty onset.**

Kiss1-MC4RKO females display advanced puberty onset, assessed by daily monitoring of vaginal opening (**A**) and first estrus (**B**), as documented by cumulative percent and mean age of animals at vaginal opening (**A**) and first estrus (**B**) in Kiss1-MC4RKO females (n=24) compared to WT littermates (n=19). *p<0.05 by Student's t-test. Data presented as the mean± SEM. (**C**) Female Kiss1-MC4RKO (n=19) have normal body weight at the time of puberty onset compared to their WT littermates (n=15). (**D**) Ontogeny expression of melanocortin genes (*AgRP*, *Pomc*, *Mc4r* and *Mc3r*) in the ARH of WT female mice at different postnatal pre-pubertal and pubertal ages: P10, P15, P22 and P30, normalized to the housekeeping gene *Hprt*. (n values: females at P10 (n=6), P15 (n=6), P22 (n=6), and P30 (n=5)). Groups with different letters are significantly different (p<0.05), as determined by one-way ANOVA followed by the Student-Newman-Keuls test. Data presented as the mean± SEM.

Kiss1-MC4RKO females and control littermates were submitted to an LH surge induction protocol that showed the LH surge was significantly blunted in the Kiss1-MC4RKO females ($P=0.0091$), while the protocol clearly evoked the expected afternoon rise of LH in control mice (**Figure 2M**). In line with these findings, the ovaries of Kiss1-MC4RKO females displayed fewer corpora lutea, markers of recent ovulation ($P=0.0054$, Kiss1-MC4RKO: 0.80 ± 0.37 vs controls: 2.80 ± 0.37), in addition to increased cystic follicles ($P=0.0337$, Kiss1-MC4RKO: 1.60 ± 0.40 vs controls: 0.40 ± 0.24) (**Figure 2N-P**), which correlate with decreased fertility as observed by the extended time to deliver pups (*i.e.* longer time to get pregnant) ($P=0.0030$, Kiss1-MC4RKO: 30.30 ± 4.60 vs controls: 20.88 ± 0.22), and fewer pups per litter ($P=0.0095$, Kiss1-MC4RKO: 6.66 ± 0.55 vs controls: 8.44 ± 0.24) (**Figure 2S**, *T*).

The increase in serum LH levels and decreased ovulation observed in the Kiss1-MC4RKO females is reminiscent of polycystic ovary syndrome (PCOS) mouse models (22, 23). Thus, we investigated whether Kiss1-MC4RKO females display a PCOS-like phenotype. We analyzed circulating levels of testosterone (T) and anti-müllerian hormone (AMH), which are frequently elevated in PCOS models (24). The Kiss1-MC4RKO females expressed normal T and AMH levels compared to control mice in diestrus (**Figure 2Q**, *R*). Thus, we can exclude a PCOS-like reproductive phenotype mediated by the lack of melanocortin signaling on Kiss1 neurons.

Re-insertion of MC4R in Kiss1 neurons of MC4RKO mice improves reproductive function

Kiss1-MC4RKO females displayed reproductive abnormalities resembling those described in MC4RKO females (11–13). Thus, we hypothesized that the reproductive defects described for the MC4RKO mice would be, at least in part, due to the absence of MC4R signaling in Kiss1 neurons. To further investigate this hypothesis, we generated mice that do not express *Mc4r* anywhere (MC4R-LoxTB, *i.e.* MC4RKO) or that express *Mc4r* only in Kiss1 neurons (*Kiss1-cre:MC4R-loxTB*). The specific re-insertion of *Mc4r* within Kiss1 neurons in the *Kiss1-cre:MC4R-loxTB* mice was confirmed through RNAscope (**Figure S2A**). *Mc4r* expression was not detected in the PVN of these mice (**Figure S2B**), and it was only detected in Kiss1 neurons in the *Kiss1-cre:MC4R-loxTB* mice. Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening (VO) and first estrus (FE). MC4R-LoxTB and *Kiss1-cre:MC4R-loxTB* female mice displayed normal timing of puberty onset compared to their control littermates, as assessed by VO ($P=0.104$, MC4R-LoxTB: 30.86 ± 1.90 vs *Kiss1-cre:MC4R-loxTB*: 30.09 ± 1.00 vs controls: 27.57 ± 0.81), and FE ($P=0.8472$, MC4R-LoxTB: 35.14 ± 2.46 vs *Kiss1-cre:MC4R-loxTB*: 34.55 ± 1.64 vs controls: 35.80 ± 0.84) (**Figure 3A**), despite displaying significantly higher body weight at the time of puberty onset ($P=0.0001$, MC4R-LoxTB: 15.49 ± 0.50 vs *Kiss1-cre:MC4R-loxTB*: 15.54 ± 0.40 vs Controls: 13.29 ± 0.30) (**Figure 3B**). As expected, MC4R-LoxTB females (MC4RKO) displayed increased body weight (**Figure 3C**) and an array of reproductive impairments that included: irregular estrous cycles with significantly more days in diestrus ($P=0.0016$) and fewer days in estrus ($P=0.0317$) compared to controls (**Figure 3D**, *E*), significantly higher serum LH levels characterized by higher basal ($P=0.0882$, MC4R-LoxTB: 0.48 ± 0.02 vs Controls: 0.37 ± 0.06) and amplitude levels per LH pulse ($P=0.0102$, MC4R-LoxTB: 0.730 ± 0.009 vs Controls: 0.58 ± 0.07) (**Figure 3F-J**), and fewer corpora lutea ($P=0.0366$, MC4R-LoxTB: 0.80 ± 0.80 vs Controls: 4.20 ± 1.15) (**Figure 3K**, *L*), recapitulating the same reproductive phenotype observed in Kiss1-MC4RKO mice despite the differences in BW. Re-introduction of MC4R into Kiss1 neurons in *Kiss1-cre:MC4R-loxTB* mice completely recovered estrous cyclicity, which was similar to controls (diestrus: $P=0.1932$, proestrus: $P=0.8262$, estrus: $P=0.0547$, compared to controls) (**Figure 3D**, *E*) despite *Kiss1-cre:MC4R-loxTB* mice showing the same degree of obesity as MC4R-LoxTB mice (**Figure 3C**), indicating that obesity *per se* was not mediating the irregular estrous cycles in MC4RKO mice. As indicated above, MC4R-LoxTB mice display higher overall circulating LH levels, and this feature was also recovered by the re-insertion of MC4R in Kiss1 neurons of *Kiss1-cre:MC4R-loxTB* mice (basal LH: 0.32 ± 0.04 , LH amplitude: 0.48 ± 0.04 , LH pulses/180min: 2.40 ± 0.67 and AUC: 70.62 ± 7.15 compared to controls) (**Figure 3G-J**), further confirming that the melanocortin action on Kiss1 neurons is required for the normal control of gonadotropin release. Despite these significant improvements in reproductive function,




Figure 2.

Deletion of MC4R from Kiss1 neurons impairs fertility in Kiss1-MC4RKO females.

(A) Female Kiss1-MC4RKO (n=19) have normal body weight from weaning and until post-natal day (PND) 150 compared to their WT littermates (n=15). (B) Representative examples of estrous cycles in Kiss1-MC4RKO and control females (n=5/group) assessed by daily vaginal cytology for 15 days. (C) Kiss1-MC4RKO females displayed irregular estrous cycles with a longer time in diestrus and a shorter time in estrus compared to control females. *** $P < 0.001$ Two-way ANOVA. (D) Pattern of LH pulsatility was analyzed in gonad intact Kiss1-MC4RKO (n=6) and control (n=7) females. # Represents LH pulses. (E) LH total secretory mass assessed by area under the curve (AUC), (F) LH pulse amplitude, (G) Basal LH and (H) Total number of pulses/180 min were analyzed. * $p < 0.05$ by Student's t-test. The expression of the KNDy genes *Pdyn* (I), *Kiss1* (J), *Tac2* (K) and *Tacr3* (L), was assessed in the ARH of adult Kiss1-MC4RKO (n=6) and control (n=3) females. ** $p < 0.01$ by Student's t-test. Data presented as the mean $\pm$ SEM. (M) Kiss1-MC4RKO (n=6) and control (n=5) females were subjected to an LH surge induction protocol. LH samples were collected in the morning (AM [8 a.m.]) and evening (PM [7 p.m.]) after lights off. ** $P < 0.01$, Two-way ANOVA followed by Bonferroni *post hoc* test. (N) Ovarian histology shows a decrease in the number of corpora lutea (CL, O) and increase in the number of cystic follicles (CF, P) of Kiss1-MC4RKO compared to controls (n=5/group). # represents cystic follicles. Serum levels of Testosterone (Q) and AMH (R) in adult gonad intact Kiss1-MC4RKO and control females (n=5/group). Student t test for unpaired samples. Data presented as the mean $\pm$ SEM. Kiss1-MC4RKO females display impaired fertility characterized by increased time to deliver pups (S) and decreased number of pups per litter (T), (n=9/group). * $p < 0.05$ and ** $p < 0.01$ by Student's t-test. Data presented as the mean $\pm$ SEM.

Kiss1-cre:MC4R-loxTB mice presented fewer corpora lutea than controls (0.80 ± 0.80), similar to MC4R-LoxTB mice, suggesting that an ovulatory impairment still persists (Figure 3K, L). While tonic LH release and estrous cyclicity are predominantly controlled by Kiss1^{ARH} neurons, ovulation is mediated by the induction of the preovulatory LH surge by Kiss1^{AVPV/PeN} neurons. It is possible that the metabolic signals derived from their obese phenotype, and/or the absence of the direct action of MC4R in GnRH neurons (16, 25) prevents the complete recovery of ovulation in *Kiss1-cre:MC4R-loxTB* mice. Both genetic models displayed significantly lower T (MC4R-LoxTB: 28.50 ± 6.20 vs *Kiss1-cre:MC4R-loxTB*: 31.22 ± 3.90 vs Controls: 62.84 ± 16.53) and AMH (MC4R-LoxTB: 150.6 ± 22.0 vs *Kiss1-cre:MC4R-loxTB*: 146.9 ± 10.8 vs Controls: 321.8 ± 15.7) levels than control mice in diestrus (Figure 3M, N). Therefore, we can exclude, once again, a PCOS-like reproductive phenotype mediated by the lack of melanocortin signaling.

Kiss1^{ARH} neurons are excited by MC4R agonists

Based on the expression of *Mc4r* in Kiss1 neurons and the reproductive impairment found after *Mc4r* deletion within Kiss1 neurons (Figure 2), we hypothesized that Kiss1^{ARH} neurons are direct targets of melanocortins and would respond to MC4R activation. Initially we did whole-cell, voltage clamp recordings in Kiss1^{ARH} neurons (Figure 4A). *Kiss1-Cre* x Ai32 or *Kiss1-Cre* AAV-injected mice underwent ovariectomies (OVX) and were given estradiol (E2) replacement (see Methods). We targeted fluorescent cells for recording which were isolated synaptically by bathing the slices in tetrodotoxin (TTX, 1 μ M). Focal application of the high-affinity melanocortin receptor agonist melanotan II (MTII, ~250 nM) elicited a small inward current in half of the isolated Kiss1^{ARH} cells (10/21) (Figure 4B). Next, we examined if the E2 state affected MTII response by recording from Kiss1^{ARH} neurons from OVX females and found two thirds (6/9 cells) responded, but there was no significant difference in the average current (Figure 4C). Finally, it must be noted that at this concentration MTII is a non-selective MCR agonist, so we followed up using a perfusion of the MC4R-selective agonist THIQ (100 nM) to determine if activation of MC4Rs alone was sufficient to invoke a postsynaptic response in Kiss1^{ARH} neurons from OVX+E2 females. Similar to MTII, THIQ was able to elicit an excitatory inward current in Kiss1^{ARH} neurons (4/5 cells, -7.4 ± 0.5 pA), indicating MC4R activation is sufficient (Figure 4D).

POMC neurons synapse directly with Kiss1^{ARH} neurons

Based on the pharmacological activation of MCRs, we wanted to address whether POMC^{ARH} neurons are the source of melanocortins to excite Kiss1^{ARH} neurons directly. To answer this question, we injected an AAV-EF1 α -DIO-ChR2:mCherry vector into the ARH of adult *Pomc-Cre* female mice (Figure 4E) (24). After 2-4 weeks we did whole cell recordings from putative Kiss1^{ARH} neurons and looked for a response to high frequency optogenetic stimulation of POMC fibers in OVX + E2-treated females (Figure 4F). We were able to confirm that these were Kiss1 neurons based on the presence of a persistent sodium current and/or *post hoc* identification by scRT-PCR (40/77) cells, Figure 4G). While this current is more prevalent in the AVPV Kiss1 population, Kiss1 neurons are the only ARH neurons to display this electrophysiological “fingerprint” of a pronounced INaP paired with a high capacitance and a low input resistance (26). Additionally, while recording from putative Kiss1^{ARH} neurons, we optogenetically stimulated POMC fibers at low frequency (1 Hz, 5 ms, 470 nm light) and recorded fast postsynaptic inward currents. The fast kinetics of the EPSC are also a tell-tale sign of an ionotropic glutamatergic response (27). Previously, CNQX was sufficient to block similar excitatory ESPCs in other postsynaptic targets of POMC neurons (28). While the consistent latency from optogenetic stimulus and current response was indicative of a single synaptic delay, we wanted to establish that it was a direct synaptic connection, as we have previously shown for output of POMC neurons (28) and Kiss1 neurons (29–31). First, we abrogated the optogenetic response with the addition of TTX (1 μ M) to the bath, we were able to rescue the postsynaptic glutamate response with the addition of the K⁺ channel blockers 4-aminopyridine (4-AP; 0.5 mM)




Figure 3.

Re-insertion of MC4R in Kiss1 neurons restores estrous cyclicity and LH pulsatility in *Kiss1-cre:MC4R-LoxTB* females.

(A) *Kiss1-cre:MC4R-LoxTB* (n=11) and *MC4R-LoxTB* (MC4RKO) (n=7) displayed normal puberty onset as compared to their control littermates (n=14), as documented by cumulative percent and mean age of animals at vaginal opening and first estrus. (B) *Kiss1-cre:MC4R-LoxTB* and *MC4R-LoxTB* both had significantly higher body weights at the time of puberty onset compared to their controls. $^{***}P < 0.01$, $^{***}P < 0.001$ by one way ANOVA. (C) *MC4R-LoxTB* (n=7) and *Kiss1-cre:MC4R-LoxTB* (n=11) females displayed significantly higher body weight than their littermates (n=10) from post-natal day 30 onwards, $^{*}p < 0.05$ and $^{**}P < 0.01$. Data presented as the mean $\pm$ (SEM). (D) Representative examples of estrous cycles of *MC4R-LoxTB* (n=4), *Kiss1-cre:MC4R-LoxTB* (n=5) and control littermate mice, (n=4) assessed by daily monitoring of vaginal smears for 15 days. (D/M: diestrus/metestrus, P: proestrus, E: estrus). (E) *MC4R-LoxTB* females displayed irregular estrous cycles, presenting longer time in diestrus and shorter time in estrus compared to control females, while *Kiss1-cre:MC4R-LoxTB* females displayed regular estrous cyclicity, similar to controls. $^{*}p < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$, $^{****}P < 0.0001$, Two-way ANOVA. Data presented as the mean $\pm$ SEM. (F) Pattern of LH pulsatility analyzed in gonad intact *MC4R-LoxTB* (n=4), *Kiss1-cre:MC4R-LoxTB* (n=5) and control littermates females (n=5). LH samples were collected every 10 min for 180 minutes; # represents LH pulses. (G) LH total secretory mass, (H) LH pulse amplitude, (I) basal LH, and (J) total number of pulses/180 min were assessed. $^{*}p < 0.05$, $^{**}P < 0.01$ by one way ANOVA. (K) Representative samples of ovarian histology from *MC4R-LoxTB*, *Kiss1-cre:MC4R-LoxTB* and control females (n=5/group); CL: Corpora lutea. Data are presented as the mean $\pm$ SEM. (L) Ovarian histology showed a significant decrease in the number of corpora lutea in the *MC4RKO* and *Kiss1-cre:MC4R-LoxTB* compared to controls. Groups with different letters are significantly different. Serum levels of (M) Testosterone and (N) AMH in adult gonad intact females *MC4R-LoxTB*, *Kiss1-cre:MC4R-LoxTB* and their control littermates. $^{*}p < 0.05$, $^{****}P < 0.0001$, One Way ANOVA. Data are presented as the mean $\pm$ SEM.




Figure 4.

Kiss1^{ARH} neurons are activated by melanocortin agonists and respond to optogenetic stimulation of POMC neurons.

(A) Slices were taken from *Kiss1-cre* injected (confocal image) brains to target fluorescent cells (arrowhead) for recording (white outline over electrode). (B) Whole cell voltage clamp recording of Kiss1^{ARH} neurons following the direct application of the high affinity melanocortin receptor agonist melanotan II (MTII, 250 nM) was added directly to the bath and the response was compared between slices from OVX and OVX+E females. (C) While the average inward current was slightly higher in the OVX+E state, this was not a significant difference (Student's t-test for unpaired samples, $p > 0.05$). (D) The MC4R-selective agonist THIQ (100 nM) was perfused, and excitatory inward currents were generated in Kiss1^{ARH} neurons. (E) Images are of AAV-driven labeling of POMC cells as seen through the confocal. (F) Using the AAV-driven expression of channelrhodopsin in adult *Pomc-cre* mice, high frequency stimulation elicited a slow inward current in Kiss1 neurons. (G) The identity of cells was confirmed through the presence of a persistent sodium current (see (26)) and/or with RT-PCR of harvested cytoplasm showing *Kiss1* expression. (Gel: MM=molecular marker, TC=tissue controls). (H) A direct synaptic projection from POMC to Kiss1^{ARH} neuron (*posthoc* identified) was confirmed using the "rescue" protocol: 1, baseline glutamatergic responses were initially generated (black trace); 2, then action potentials were eliminated by blocking voltage-gated sodium channels with TTX and the postsynaptic response (red trace); 3, blockade of potassium channels facilitated calcium entry into the terminal through ChR2 to release synaptic vesicles, "rescuing" the postsynaptic glutamate response (green trace). (I) Glutamatergic responses were also often observed, particularly in the ventral ARH. As seen when targeting low input resistance neurons (i.e., Kiss1^{ARH}) with low frequency optogenetic stimulation (5 ms pulse, 50 ms inter-spike interval), the first response in OVX + vehicle females was larger relative to the second response in OVX + E2-treated female mice. Representative traces are the average of 30 sweeps. (J) The averaged paired-pulse ratio was lower in recordings from estradiol-treated female mice, indicating an increased release probability (Students t-test $p < 0.05$). (K) High frequency optogenetic stimulation elicited a small inward current. (L) In a different cell, a high frequency inward current was noted before constant perfusion of SHU9119 for 15 minutes (break between traces). When the stimulation protocol was repeated, no inward current was elicited. (M) The majority of high frequency responses were inward and only twice was an inhibitory outward current recorded in identified Kiss1^{ARH} neurons. Perfusion of the non-selective opioid receptor antagonist naloxone reversed the current, eliminating MCR-activation as the mechanism.

and tetraethyl ammonium (TEA; 7.5 mM) (**Figure 4H**). K⁺ channel block enables the calcium influx from ChR2 activation in the presynaptic terminals to be sufficient to restore vesicle fusion. Therefore, there appears to be no intervening synapses between POMC and the downstream Kiss1^{ARH} neurons.

Previously we found that E2 increases glutamate release from POMC neurons by increasing *Vglut2* (28). Therefore, we compared the postsynaptic glutamatergic responses following two optogenetic stimuli (50 ms interval between light flashes) in Kiss1^{ARH} neurons from OVX + vehicle versus OVX + E2-treated female mice (**Figure 4I**). As anticipated, we found that treatment with E2 increased the probability of glutamate release from POMC neurons onto Kiss1^{ARH} neurons based on the significant decrease in the paired pulse ratio of the two stimuli (32) (**Figure 4J**). Although in the present study we used an *in vivo* treatment paradigm, we know from previous studies that this augmentation of glutamate release can happen quite rapidly after a brief exposure to E2 *in vitro* (within 15 min) (28). Therefore, Kiss1^{ARH} neurons are excited by the glutamatergic input from POMC neurons in an E2-dependent manner.

Next, we tested for a melanocortin response using a high frequency stimulation protocol known to elicit peptide release (33). Only cells displaying a glutamatergic response were tested. Indeed, high frequency stimulation evoked an excitatory inward current (**Figure 4K**) that was blocked by the selective MC3/4R antagonist SHU 9119 (**Figure 4L**). However, the number of high frequency responses seen in Kiss1 neurons was unexpectedly infrequent and small (4/16, 25%, mean inward current: -1.4±3.1 pA) compared to the efficacy of pharmacological MCR activation. E2-treatment enhances *Pomc* expression and β-endorphin labeling (34) (35), but the effect, if any, on αMSH release or the MCRs in Kiss1 neurons is unknown. We suspected that the high frequency stimulation was eliciting co-release of αMSH and β-endorphin which would exert opposite effects on a postsynaptic cell. In order to isolate melanocortin signaling we pretreated slices with the non-selective opioid antagonist naloxone (1 μM) and found an increase in the likelihood and magnitude of a response to high frequency stimulation (9/23, 40%, mean inward current: -7.6±2.9 pA) (**Figure 4M**).

Together these optogenetic findings reinforced our pharmacological results showing that Kiss1^{ARH} neurons are excited by MCR agonists. Additional studies would need to be done to identify the cation conductance that is affected by the MC4R signaling cascade in Kiss1^{ARH} neurons, but clearly we have established that there are excitatory glutamatergic and peptidergic inputs from POMC to Kiss1^{ARH} neurons.

Kiss1^{AVPV/PeN} neurons are inhibited by MC4R agonists

Having established a direct, excitatory projection from POMC to Kiss1^{ARH} neurons, we next examined POMC inputs to Kiss1^{AVPV/PeN} neurons. First, we used immunocytochemistry to label fibers expressing αMSH to demonstrate that POMC fibers densely innervated the AVPV/PeN area (**Figure 5A**). Next, we recorded from AVPV neurons in slices taken from OVX+E2, POMC-Cre mice injected with AAV1-DIO-YFP:ChR2 into the ARH (**Figure 5B**). We targeted cells along the ventricle in the AVPV and PeN, surrounded by YFP terminals, and Kiss1^{AVPV/PeN} neurons were identified based on the expression of several endogenous conductances (*I_{NaP}*, T-type calcium current, h-current) that are unique to these neurons (26, 36). We recorded a direct glutamatergic synaptic response following optogenetic stimulation in six neurons, which was further verified through pharmacological “rescue” of the synaptic response in the majority (5/6) neurons tested (**Figure 5C**), indicating that POMC neurons make direct monosynaptic connections with Kiss1^{AVPV/PeN} neurons.

We next hypothesized that Kiss1^{AVPV/PeN} neurons would be excited by exogenous application of melanocortin agonists because the receptor is typically Gs-coupled (37, 38). We did whole-cell, voltage clamp recordings from Kiss1^{AVPV/PeN-Cre::Ai32} neurons, which were isolated




Figure 5.

Kiss1^{AVPV/PeN} neurons are inhibited by MC4R agonists in an E2-dependent manner.

(A) Immunohistochemistry showing robust labeling of α -MSH fibers in the AVPV/PeN region in OVX+E2 WT female mice. (B) Low and high power bright-field images taken during electrophysiology recording in OVX+E2 POMC-Cre mice expressing YFP:ChR2 in the ARH following AAV injection. (left) Low power image shows the location of recorded cells. (right) Higher power image of area from white box inset displaying POMC fibers innervating the area. (C) Fibers surrounded putative Kiss1^{AVPV} neurons (expressing I_{NaP} , IT and I_h) and optogenetic stimuli was able to elicit postsynaptic currents that were eliminated with tetrodotoxin (1 μ M), but “rescued” with addition of K^+ channel blockers (4-AP and TEA) which indicates a monosynaptic connection between POMC^{ARH} and Kiss1^{AVPV} neurons. (D) Whole-cell voltage clamp recordings were made in ChR2-YFP positive cells in brain slices taken from Kiss1-cre: Ai32 female mice. Bath application of melanotan II (MTII, 500 nM) generated an inhibitory outward current in a Kiss1^{AVPV/PeN} neuron from OVX female. As the cell was synaptically isolated using bath-applied tetrodotoxin (TTX, 1 μ M) this represents a direct effect. Washout of MTII while still in TTX quickly led to a return to baseline RMP. (E) An IV relationship was plotted using voltage steps before and after MTII administration. The crossing at -80 mV ($\sim$ EK⁺) indicates that the opening of K^+ channels underlie the MC4R inhibition of Kiss1^{AVPV} neurons. (F) In a subset of recordings from OVX brain slices, the selective membrane estrogen receptor (Gq-mER) agonist STX (10 nM) was added to the bath for $\sim$ 10 minutes prior to addition of MTII. STX pretreatment resulted in either a greatly attenuated outward current or even an inward current as shown in this example. (G) The mean outward current was significantly higher when recording in Kiss1^{AVPV/PeN} neurons from brain slices from vehicle-treated, OVX females compared to E2-treated, OVX females or acute STX-treated brain slices from OVX females (One-way ANOVA $F_{(2,19)}=12.32$, $p<0.001$; Holm-Sidak *posthoc* comparisons found significant differences between all groups; * $p<0.05$, *** $p<0.001$). (H) There was no difference in the frequency of mEPSCs after MTII, calculated as a percent of baseline between groups. (I) However, there was a significant difference in the mEPSC amplitude (calculated as a percent of baseline) between all groups: main effect $F_{(2,19)}=10.58$, $p<0.001$. *Posthoc* comparisons using Holm-Sidak found OVX+STX to be different from both OVX ($p<0.001$) and OVX+E ($p<0.05$).

synaptically by bathing the neurons in TTX (1 μ M), from OVX females. For these experiments, we tested the response to the high affinity melanocortin receptor agonist MTII (500 nM). Surprisingly, we consistently measured an outward (inhibitory) current that could be reversed on washout of MTII (**Figure 5D**). This outward current was associated with an increase in a K^+ conductance based on the current-voltage plot (*i.e.*, the outward current reversed at $\sim E_{K^+}$, **Figure 5E**). Also, the inwardly-rectifying I/V could indicate that MC4R is coupled to activation of inwardly-rectifying K^+ channels as has been previously reported for MC4R signaling in pre-autonomic parasympathetic neurons in the brainstem (39). We hypothesized that the coupling could be modulated by E2 via a Gq -coupled membrane estrogen receptor (Gq-mER) as we have previously demonstrated in POMC neurons (40–42). For these experiments, we again utilized OVX females and targeted Kiss1^{AVPV/PeN}-Cre::Ai32 neurons. Once in the whole-cell voltage-clamp configuration, we perfused the slices with STX (10 nM), a selective ligand for the putative Gq-mER (42), and tested the response to the MTII. Indeed, the outward current (inhibitory) response to MTII was completely abrogated and even reversed by STX (**Figure 5F**, G). The short-term (bath) treatment with STX ensured that there was no desensitization of the Gq-mER with longer-term (*in vivo*) treatment with E2. Therefore, estrogen receptor activation can rapidly uncouple (*i.e.*, desensitize) the MC4R inhibitory response in Kiss1^{AVPV/PeN} neurons. We would predict that the intracellular signaling cascade for the heterologous desensitization is similar to what we have elucidated in POMC neurons (42), but this will need to be determined in future experiments. We next investigated the response in vehicle-treated, OVX females. Indeed, Kiss1^{AVPV/PeN} neurons were even more inhibited by the same MTII exposure (**Figure 5G**). Therefore, in contrast to Kiss1^{ARH} neurons, the MC4R appears to be coupled to opening of K^+ channels in Kiss1^{AVPV/PeN} neurons. To further establish whether the MTII was having pre- or postsynaptic effects, we measured the glutamatergic mEPSCs before and after melanocortin receptor activation. We found there was no significant effect on the frequency (**Figure 5H**), but there was an effect on the amplitude (**Figure 5I**). This further supports a postsynaptic locus of STX's effects. However, *posthoc* comparisons found STX increased amplitude compared to both OVX and OVX+E. This result hints at a membrane delimited effect as STX is a much more potent agonist for Gq-mER than E2 (43).

Discussion

Our findings show that melanocortins act directly on Kiss1 neurons to contribute to the metabolic regulation of fertility, in line with previous reports showing that α MSH stimulates LH release in a kisspeptin-dependent manner (14) and that Kiss1 neurons express the melanocortin receptor MC4R (18–21). Several studies in humans have linked *Mc4r* mutations to reproductive abnormalities, including precocious puberty (9), PCOS (10), and hypogonadism (8). However, a direct association between *MC4R* mutations and reproductive function, independent from the obese condition of these patients, has not been identified (6). In the current study, we show that the reproductive impairments observed in MC4R deficient mice, which replicate many of the conditions described in humans, are largely mediated by the direct action of melanocortins via MC4R on Kiss1 neurons and not to their obese phenotype. This is because the ablation of MC4R from Kiss1 neurons largely replicated the reproductive impairments observed in MC4RKO female mice without inducing obesity, and the selective reinsertion of MC4R into Kiss1 neurons of MC4RKO mice significantly improved their reproductive function without changing their obese phenotype. Strikingly, puberty onset was advanced in Kiss1-MC4RKO females. Our findings revealed a low melanocortin tone in the hypothalamus of WT females during pubertal development, characterized by lower levels of *Agrp*, *Mc4r* and *Mc3r* expression at the time of puberty onset ($\sim$ PND30). This developmental decline in the expression of melanocortin receptors is in line with the advanced timing of puberty onset in females when MC4R is congenitally ablated from Kiss1 neurons. A recent study using whole body MC3RKO mice showed a trend to delayed puberty onset (19), suggesting a larger role of melanocortins in the timing of puberty onset. Here, we show that in females the combination of MC3R and MC4R action may be necessary for

pubertal development. Interestingly, the MC4R-LoxTB and *Kiss1-cre:MC4R-LoxTB* mice do not show an advancement in puberty onset as we observed in *Kiss1-MC4RKO* mice; however, these mice are obese at the time of puberty onset already, which may affect the timing of sexual maturation independently of MC4R action in *Kiss1* neurons. Of note, the findings presented in this study do not rule out the existence of additional sites of action of MC4R in other neuronal populations to regulate reproduction, *e.g.*, GnRH neurons, which also express MC4R (16, 25). Indeed, the lack of full recovery of the reproductive function in *Kiss1-cre:MC4R-LoxTB* mice suggests that MC4R in *Kiss1* neurons is necessary but not sufficient to achieve full reproductive capabilities.

In addition to early puberty onset, *Kiss1-MC4RKO* females displayed an impaired preovulatory LH surge driving ovulation and had fewer corpora lutea in their ovaries compared to control littermates, further supporting a role of MC4R in regulating ovulation. In adulthood, the reproductive phenotype observed in *Kiss1-MC4RKO* and *MC4R-loxTB* female mice (increased LH, irregular estrous cycles, oligo-ovulation, increased cystic follicles) correlates with the phenotype observed in PCOS mouse models (22, 23). In fact, an association between *MC4R* mutations and PCOS has been reported (10). However, our mouse models failed to display higher levels of circulating androgens or AMH, two of the hallmarks of PCOS (24), suggesting that melanocortin signaling is unlikely to be a main contributing factor in the development of this syndrome.

Our data revealing early puberty onset and impaired LH pulse and surge driving ovulation support a direct role of MC4R on the two primary *Kiss1* populations regulating these functions (ARH and AVPV/PeN). We further investigated this and found that MC4R signaling excites *Kiss1*^{ARH} neurons and inhibits *Kiss1*^{AVPV/PeN} neurons in an estradiol-dependent manner. In *Kiss1*^{ARH} neurons, bath applied or optogenetically-evoked release of melanocortin agonists induce a direct excitatory inward current. Our studies using whole cell voltage clamp in female mice were able to detect a melanocortin-mediated inward current in *Kiss1*^{ARH} neurons that would increase excitability. Previously, an MCR-activation effect was not detected in cell-attached recordings of diestrus females (14, 44). However, cell-attached recordings do not measure a direct response in isolated *Kiss1*^{ARH} neurons but rather the summation of multiple synaptic inputs to *Kiss1*^{ARH} neurons. Most importantly, we did not see a difference in the excitatory response in synaptically isolated *Kiss1*^{ARH} neurons from E2-treated versus vehicle-treated, ovariectomized females. This emphasizes the importance of doing whole-cell recording from isolated *Kiss1* neurons. The modulatory actions of POMC^{ARH} neuronal projections to *Kiss1*^{ARH} neurons can also occur through a) the excitatory action of glutamate in a process that is facilitated in the presence of estradiol, in line with previous publications (33, 45), and b) the inhibitory action of β -endorphin, suggesting that POMC^{ARH} neurons are able to exert a precise regulatory role of the GnRH pulse generator, *i.e.* *Kiss1*^{ARH} neurons, in response to metabolic challenges. In the AVPV/PeN, MC4R agonists inhibited *Kiss1*^{AVPV/PeN} neurons in an estradiol-dependent manner, which utilized a G α_q -coupled membrane estrogen receptor (STX receptor) to attenuate the inhibitory tone of MC4R in these neurons. Rapid estrogen signaling may act to ease transitions between states. Membrane delimited E2 actions can quickly attenuate or enhance coupling between receptors and signaling cascades. These effects will precede E2-driven changes in gene expression that produce more stable alterations in signaling. This combination of mechanisms will reduce any lag between rises in serum E2 and physiological effects. Considering the abbreviated mouse reproductive cycle, parallel mechanisms acting on different timescales are particularly important.

Our pharmacology data clearly demonstrates the estradiol-dependent action of MC4R in *Kiss1*^{AVPV/PeN} neurons, suggesting that similar to *Kiss1*^{ARH} neurons, this population is regulated by melanocortins. The effect of MC4R agonists on *Kiss1* neurons in the presence of TTX, suggests a direct synaptic effect without the need of channelrhodopsin-2(ChR2)-assisted circuit mapping. Nonetheless, to further demonstrate this interaction, we provide evidence of profound innervation of α MSH fibers and the presence of synaptic contact between POMC and *Kiss1*^{AVPV/PeN} neurons through the optogenetic stimulation of a glutamatergic response in *Kiss1*^{AVPV/PeN} neurons after

low frequency stimulation of POMC terminals. We intentionally avoided high frequency stimulation of POMC terminals to prevent the co-release of β -endorphins (e.g., **Figure 4M**) and their possible additive effect on the inhibitory action of MC4R. Future studies will be aimed at the characterization of the endogenous opioid pathways in the induction of the LH surge.

To date, two instances of MC4R-mediated inhibition have been described in the central nervous system: MC4R can $G_{i/o}$ couple to open K^+ (Kir6.2) channels in parasympathetic preganglionic neurons in the brainstem (39), and MC4R can directly couple to K^+ (Kv7.1) channels in the hypothalamic paraventricular nucleus neurons (46). Interestingly, our findings describe an inhibitory role of MC4R in the reproductive neuroendocrine axis. This steroid state dependent effect is particularly relevant because in the context of reproduction, the two populations of Kiss1 neurons are strikingly different in their response to estradiol, where peptide expression in Kiss1^{ARH} and Kiss1^{AVPV/PeN} neurons are inhibited or stimulated, respectively, in order to mount the negative versus positive feedback of sex steroids (15). This differential regulation allows for the episodic versus surge release of GnRH; however, the cellular mechanisms underlying these opposing roles of Kiss1 neurons to the same stimulus, *i.e.* circulating E2 levels, remain unknown. Although there were no differences in the magnitude of the electrophysiological responses to MTII in Kiss1^{ARH} neurons in vehicle- versus E2-treated, ovariectomized females, we cannot rule out that E2 affects other systems activated by the MC4R signaling cascade as recently reported in MC4R expressing neurons of the ventromedial nucleus of the hypothalamus (VMH) (47). Interestingly, puberty onset was advanced in Kiss1-MC4RKO females. This data suggests that the inhibitory tone of MC4R signaling on Kiss1^{AVPV/PeN} neurons might also be involved in the timing of puberty onset, which in turn suggests a role for this female-specific population of Kiss1 neurons in sexual maturation. Furthermore, our findings suggest that metabolic cues, through the regulation of the melanocortin output onto Kiss1^{AVPV/PeN} neurons, are essential for the timing and magnitude of the GnRH/LH surge. This study suggests that the melanocortin-kisspeptin pathway is one of the multiple pathways essential for the metabolic regulation of the HPG axis. Future studies are warranted to fully elucidate synaptic input and postsynaptic cascades in the POMC-Kiss1^{AVPV/PeN} circuit.

Altogether, our data reveal a differential regulatory action of MC4R in the neural control of GnRH/LH release, participating in both the surge (E2-treated, ovariectomized female) and the pulse-like (ovariectomized female) modes of LH release through the cellular regulation of Kiss1 neurons, which translate into our findings on the conditional knockout of *Mc4r* from Kiss1 neurons (**Figure 6**). These findings are important because the reproductive abnormalities often attributed to obesity in MC4R deficient patients may not be caused by the excess in body weight but, at least in part, by a deficiency in MC4R signaling directly at the level of Kiss1 neurons that affects predominantly the ability to mount a preovulatory LH surge.

Materials and methods

In-vivo experimental procedures

Generation of Kiss1-MC4RKO, Kiss1-cre:MC4R-LoxTB and MC4R-LoxTB Transgenic Mice

Kiss1-MC4RKO mice were generated by crossing *Kiss1-cre* knock-in mice and *MC4R*^{lox/lox} mice. *Kiss1-cre* mice (RRID:MGI:6278139) were obtained from Dr. Richard Palmiter (University of Washington, Seattle, WA) (49) and *Mc4r*^{lox/lox} (RRID:IMSR_JAX:023720) were a gift from Dr. Brad Lowell (Beth Israel Deaconess Hospital, Boston, MA) (4). These mice were crossed to generate Kiss1-MC4RKO mice lacking *Mc4r* expression selectively from Kiss1 neurons (*Kiss1-cre*^{+/+}; *MC4R*^{lox/lox}) and their control littermates (*MC4R*^{lox/lox}). To generate *Kiss1-cre:MC4R-LoxTB* and




Figure 6.

Working model of the neuronal input from POMC ARH to Kiss1 ARH and AVPV populations.

POMC^{ARH} neurons excite Kiss1^{ARH} neurons via glutamate and αMSH release, which activate glutamatergic ionotropic receptors and MC4R, respectively. In low E2 state, αMSH fibers project to the AVPV/PeN, and MC4R inhibit Kiss1^{AVPV/PeN} neurons via opening a K⁺ channel. However, in high E2 state, E2 enhances the glutamatergic excitation of both POMC and Kiss1 ARH neurons. E2-treatment rapidly abrogates (uncouples) the inhibitory MC4R signaling in Kiss1^{AVPV/PeN} neurons, allowing greater excitatory actions of Kiss1^{ARH} neurons via enhanced glutamate release (26 [DOI](#)).

MC4R-LoxTB mice, *MC4R-LoxTB* mice were purchased from The Jackson Laboratory (*MC4R-LoxTB*; catalog no. 006414). These mice present with a loxp-flanked transcriptional blocking (*LoxTB*) sequence preventing normal endogenous gene transcription and translation from the endogenous locus. Homozygous *MC4R-loxTB/loxTB* mice are devoid of functional *MC4R* mRNA (*MC4RKO* mice), while the presence of Cre recombinase on the *Kiss1* promotor in *Kiss1-cre* mice will result in the removal of the transcription blocker and subsequent expression of *MC4R* in tissue-specific sites (i.e. *Kiss1* neurons) therefore resulting in the generation of *Kiss1-cre:MC4R-LoxTB* (with *Mc4r* expression restored in *Kiss1* neurons), their obese control littermates *MC4R-LoxTB* mice (*MC4RKO*), and their WT controls (*MC4R*^{+/+}). To rule out early embryonic recombination of the *MC4R*^{loxTB/loxTB} or *MC4R*^{lox/lox} alleles, we ran PCR assays on tail DNA designed to detect wild-type allele (*MC4R*^{+/+}), undeleted lox (*MC4R*^{lox/lox}) or loxTB alleles (*MC4R*^{loxTB/loxTB}). Genotyping was confirmed by sending tissue to Transnetyx, Inc., for testing by real-time polymerase chain reaction. Mice were housed in Harvard Medical School Animal Resources facilities where they were fed standard mouse chow (Teklad F6 Rodent Diet 8664) and were given *ad libitum* access to tap water under constant conditions of temperature (22–24°C) and light (12 hr light [07:00]/dark [19:00] cycle).

RNAscope in situ hybridization

To validate the *Kiss1-MC4RKO*, *Kiss1-cre:MC4R-LoxTB* and *MC4R-LoxTB* (*MC4RKO*) mouse models and investigate the co-expression of *Kiss1* and *Mc4r* mRNA in these mice, dual fluorescence ISH was performed using RNAscope (ACD, Multiplex Fluorescent v.2) according to the manufacturer's protocol using the following probes: *Mc4r* (319181-C2) and *Kiss1* (500141-C1). Brains (n=4/group) from WT OVX (for expression in the ARH) and OVX+E2 (for expression in the AVPV/PeN) mice and OVX *Kiss1-MC4RKO*, *Kiss1-cre:MC4R-LoxTB* and *MC4RKO* were removed fresh frozen on dry ice, and then stored at -80°C until sectioned. Five sets of 20 µm sections in the coronal plane were cut on a cryostat, from the diagonal band of Broca to the mammillary bodies, thaw mounted onto SuperFrost Plus slides (VWR Scientific) and stored at -80°C until use. A single set was used for ISH experiment (adjacent sections 100 µm apart). Images were taken at 20x magnification of the sections containing AVPV, PeN, and the three rostro-to-caudal levels of the ARH, and *Kiss1* neurons expressing (*Kiss1-cre:MC4R-LoxTB* mice) or lacking (*Kiss1-MC4RKO* mice) *Mc4r* were identified using ImageJ.

Reproductive maturation of *Kiss1-MC4RKO*, *Kiss1-cre:MC4R-LoxTB* and *MC4R-LoxTB* mice

To assess the reproductive phenotype of mice with selective reinsertion of *Mc4r* in *Kiss1* neurons (*Kiss1-cre:MC4R-LoxTB*, n=11), selective deletion of *Mc4r* from *Kiss1* neurons (*Kiss1-MC4RKO*, n=24), global deletion of *Mc4r* (*MC4R-LoxTB*, n=7); and their control *MC4R*^{lox/lox} littermates (n=14), mice were weaned at post-natal day (PND) 21 and were monitored daily for puberty onset. Females were monitored daily for vaginal opening (VO, indicative of the complete canalization of the vaginal cavity) and for first estrus (first day with cornified cells determined by daily morning vaginal cytology) after the day of VO. Body weight (BW) was measured at the day of puberty onset to determine if changes in puberty onset could be due to differences in BW. Estrous cyclicity was monitored in females by daily vaginal cytology, for a period of 15 days, in 6-month-old mice and their respective control littermates (n=5/group). Cytology samples were obtained every morning (9 am), placed on a glass slide and stained with hematoxylin and eosin for determination of the estrous cycle stage under the microscope.

Fecundity test of *Kiss1-MC4RKO* females

Adult 6-month-old *Kiss1-MC4RKO* and control littermate female mice (n=9/group) were placed with adult WT males proven to father litters for 90 days and time to deliver pups and number of pups per litter were monitored.

Characterization of the estradiol-induced luteinizing hormone surge

Kiss1-MC4RKO (n=6) and control MC4R^{lox/lox} littermate (n=5) adult female mice were subjected to bilateral ovariectomy (OVX) via abdominal incision under light isoflurane anesthesia. Immediately after OVX, capsules filled with E2 (1 ug/20g BW) were implanted subcutaneously via a small mid-scapular incision on the back. Five days later, mice were subcutaneously injected in the morning with estradiol benzoate (1 ug/20g BW) to produce elevated proestrus-like E2 levels (preovulatory LH surge) on the following day (50 [4](#)). Blood samples were collected at 8am and 7pm; LH levels were stored at -80°C until measured via LH ELISA.

Ovarian histology and hormone measurements

Bilateral ovariectomy of 6-month-old Kiss1-MC4RKO (n=4), *Kiss1-cre*:MC4R-LoxTB, MC4R-LoxTB and their control littermates (n=5/group) was performed under light isoflurane anesthesia. Briefly, the ventral skin was shaved and cleaned, and one small abdominal incision was made. Once the ovaries were identified and excised, the muscle incision was sutured, and the skin was closed with surgical clips. Ovaries were stored in Bouin's fixative, sectioned, and stained with hematoxylin and eosin at the Harvard Histopathology Core. Corporal lutea were counted in the middle section of each ovary. Serum samples were also collected for analysis of testosterone and anti-müllerian hormone (AMH) levels in these mice. These hormones levels were measured at the University of Virginia Ligand Assay core with the Mouse & Rat Testosterone ELISA assay (reportable average range 10-1600 ng/dL; sensitivity of 10 ng/dL); AMH ELISA assay (reportable average range 0.2 – 15 ng/mL; sensitivity of 0.2 ng/mL).

LH pulsatile secretion profile in gonad intact Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB female mice

To assess the profile of LH pulses secretion, adult 6 months old gonad intact Kiss1-MC4RKO females (n=6), their control Mc4r^{lox/lox} littermates (n=7); and *Kiss1-cre*:MC4R-LoxTB (n=5/group), MC4R-LoxTB and their control WT females (n=4/group) were handled daily for 3 weeks to allow acclimation to sampling conditions prior to the experiment. Pulsatile measurements of LH secretion were assessed in diestrus by repeated blood collection through a single incision at the tip of the tail. The tail was cleaned with saline and 4 µL of blood was taken at each time point from the cut tail with a pipette. We collected sequential blood samples every 10 minutes over a 180-minute sampling period. Samples were immediately frozen on dry ice and stored at -80 °C until analyzed with LH ELISA as previously described (51 [4](#)). The functional sensitivity of the ELISA assay was 0.0039 ng/ml with a CV% of 3.3%.

LH pulses analysis

LH pulses in mice were analyzed using a custom-made MATLAB-based algorithm. The MATLAB code includes a loop that determines LH pulses as any LH peak: (i) whose height is 20% greater than the heights of the 2 previous values; (ii) 10% greater than the height of the following value; and (iii) the peak at the second time interval needs to be 20% greater than the single value that comes before it to be considered a pulse, as we previously described (52 [4](#)).

LH pulsatility was assessed by measuring: (1) the total secretory mass, assessed by area under the curve (AUC); (2) the LH pulse amplitude, calculated by averaging the 4 highest LH values in the samples collection period for each animal; (3) the basal LH, calculated by averaging the 4 lowest LH values in the samples collection period for each animal; and (4) the total number of pulses throughout the 180 minutes sampling period.

Immunohistochemistry

Animals and Treatment

Coronal brain blocks (2 mm each) from adult female C57BL/6 mice (n=4) were fixed by immersion in 4% paraformaldehyde for ~ 8 hours, cryoprotected in 30% sucrose solution, frozen in isopentane at -55°C, sectioned coronally on a cryostat at 20 µm, and thaw-mounted onto Superfrost Plus slides (Fisher Scientific, Pittsburgh, PA). The 20 µm sections were stored at -20°C until used for immunocytochemistry.

Immunocytochemistry

The sections were rinsed in PB (0.1M phosphate buffer, pH 7.4) for at least 30 min. Next, sections were incubated with normal serum corresponding to the host for the secondary antiserum (5 % normal serum with 0.3% Triton-X100 in PBS for 30 min), rinsed in PB and then incubated for ~45 h at 4° C with a rabbit polyclonal antiserum against α-MSH (1:2,500). The specificity of this antiserum has been documented (53). After rinsing, sections were first incubated for 2-3 hours at room temperature with biotinylated donkey anti rabbit gamma globulin (IgG; 1:500) and next with streptavidin-Alexa 488 (1:2500). Both the primary and secondary antibodies were diluted in tris-(hydroxymethyl) aminomethane (0.5 %, Jackson ImmunoResearch, Philadelphia, PA) in PB containing 0.7 % seaweed gelatin (Jackson ImmunoResearch, Philadelphia, PA) and 0.5 % Triton X-100 and 3% bovine serum albumin (BSA; Jackson ImmunoResearch, Philadelphia, PA) adjusted to pH 7.6. Following a final rinse overnight, slides were coverslipped with gelvatol containing the anti-fading agent, 1,4-diazabicyclo(2,2)octane (DABCO; Cold Spring Harbor Protocols, 2006).

Imaging

Photomicrographs of labeling were initially acquired using a Nikon E800 fluorescent microscope (Eclipse E800; Nikon Instruments, Melville, NY) equipped with a fiber illuminator (Intensilight C-HGFI; Nikon Instruments) and a high-definition digital microscope camera head (DS-Fi1; Nikon Instruments) interfaced with a PC-based camera controller (DS-U3; Nikon Instruments).

Real time quantitative PCR

i) To investigate the changes in the expression of the melanocortin genes *Agrp*, *Pomc*, *Mc3r* and *Mc4r* in the MBH during development in prepubertal and pubertal WT females at ages P10 (n=6), P15 (n=6), P22 (n=6), and P30 (n=5), and ii) evaluate the gene expression profile of *Pdyn*, *Kiss1*, *Tac2*, *Tacr3*, in the ARH of adult intact (in diestrus) female Kiss1-MC4RKO (n=6) and their control MC4R^{lox/lox} littermates (n=3), the brains were removed and rapidly embedded in Tissue-Tek, frozen in -30°C isopentane solution and stored at -80°C until use. Frozen tissue punches were recovered through MBH with a 1 mm diameter canula (54). These tissue punches encompassed the whole MBH from the WT females (figure 1D) and the ARH from the Kiss1-MC4RKO and their control females (Figure 2I). The tissues were homogenized, and total RNA was isolated using TRIzol reagent (Invitrogen) followed by chloroform/isopropanol extraction. RNA purity and concentration were measured via an absorbance spectrophotometer (260/280 nm > 1.8; NanoDrop 1000, Thermo Fisher Scientific). Total RNA (1 µg) was reverse transcribed to cDNA using random hexamers (High-Capacity cDNA Synthesis Kit, Life Technologies). Quantitative real-time PCR assays were performed using SYBR Green RT-qPCR master mix (Applied Biosystems) and analyzed using ABI Prism 7000 SDS software (Applied Biosystems). The cycling conditions were: 2 minutes incubation at 95°C (hot start), 45 amplification cycles (95°C for 30 seconds, 60°C for 30 seconds, and 45 seconds at 75°C, with fluorescence detection at the end of each cycle), followed by melting curve of the amplified products obtained by ramped increase of the temperature from 55°C to 95°C to confirm the presence of single amplification product per reaction. PCR specificity was verified by melting curve analysis and agarose gel electrophoresis. Each sample was run in duplicate to

obtain an average cycle threshold (CT) value and relative expression of each target gene was determined using the comparative Ct method (55). The data were normalized to Hypoxanthine Guanine Phosphoribosyltransferase (*Hprt*) expression levels in each sample. Results were expressed as fold differences in relative gene expression with respect to i) P10 for melanocortin genes expression analysis during development in WT mice, and ii) controls for the *KNDy* genes expression in female *Kiss1*-MC4RKO mice. The primers used are listed in **Table 1**.

Gene	Gene length (pb)	Accession #	Primers	Location (nt)	Sequence
<i>Hprt</i>	352	NM_013556.2	Hprt-F	704-728	CCTGCTGGATTACATTAAAGCGCTG
			Hprt-R	377-401	GTCAAGGGCATATCCAACAACAAAC
<i>Agrp</i>	136	NM_001271806.1	Agrp-F	466-488	GCCTCAAGAAGACAACCTGCAGAC
			Agrp-R	580-601	AAGCAGGACTCGTGCAGCCTTA
<i>Pomc</i>	138	NM_001278584.1	Pomc-F	187-208	CCATAGATGTGTGGAGCTGGTG
			Pomc-R	303-324	CACCTCCGTTGCCAGGAAACAC
<i>Mc4r</i>	101	NM_016977.4	Mc4r-F	552-570	CCCGGACGGAGGATGCTAT
			Mc4r-R	632-652	TCGCCACGATCACTAGAATGT
<i>Pdyn</i>	200	NM_018863.4	Pdyn-F	45-64	ACAGGGGGAGACTCTCATCT
			Pdyn-R	223-244	GGGGATGAATGACCTGCTTACT
<i>Kiss1</i>	129	AF472576.1	Kiss1-F	147-166	GCTGCTGCTTCTCCTCTGTG
			Kiss1-R	256-275	TCTGCATACCGCGATTCTT
<i>Tac2</i>	234	NM_001199971.1	Tac2- F	238-257	GCTCCACAGCTTTGTCCTTC
			Tac2- R	452-471	GCTAGCCTTGCTCAGCACTT
<i>Tacr3</i>	159	NM_021382.6	Tacr3-F	759-779	GCCATTGCAGTGGACAGGTAT
			Tacr3-R	898-917	ACGGCCTGGCATGACTTTTA

Data analysis

Statistical data are expressed as means $\pm$ SEM, where n represents the number of animals in each study group. The significance of differences between groups was evaluated using unpaired two-tailed Student's t test, or a one- or two-way ANOVA test (with *post hoc* comparisons). Significance level was set at $p < 0.05$. All analyses were performed with GraphPad Prism Software, Inc (San Diego, CA).

In-vitro experimental procedures

Animals

All animal procedures described in the electrophysiology studies were performed in accordance with institutional guidelines based on National Institutes of Health standards and approved by the Institutional Animal Care and Use Committees at Oregon Health and Science University and Appalachian State University. *Kiss1-cre* (v2) mice (49) were crossed with Ai32 (56) or C57B6J mice. *Pomc-Cre* mice (RRID:IMSR_JAX:005965) (57) were crossed with wildtype C57B6J (RRID:IMSR_JAX:000664) mice. The Ai32 cross was not used with *Pomc-Cre* mice because the gene is transiently expressed during development in some cells fated to be Kiss1 or AgRP cells (58) and early recombinant events lead to persistent expression of the ChR2-mCh fusion protein in non-POMC cells. However, we have previously shown that AAV-driven expression in adult *Pomc-*

cre animals is restricted to β -endorphin labeled cells in the ARH (45 [↗](#)), avoiding this non-specificity issue. All colonies were maintained onsite under controlled temperature (21–23 °C) and photoperiod (12:12-h light-dark cycle 0600 to 1800) while receiving ad libitum food (5L0D; LabDiet, St. Louis, MO) and water access. Following surgeries, mice received a s.c. dose of 4–5 mg/kg carprofen (Rimadyl; Pfizer Animal Health, New York, NY) and given at least 1 week of recovery.

Ovariectomies

Seven to ten days prior to each experiment, ovaries were removed as described previously while under isoflurane anesthesia (45 [↗](#)). Two days before experiments females received either an injection of sesame oil (50 μ l, sc; Sigma-Aldrich, St. Louis, MO) or a priming dose (0.25 μ g/50 μ l sesame oil, sc) of E2 benzoate (Sigma-Aldrich, St. Louis, MO) in the morning. On the following day, oil or a high (1.5 μ g) dose of E2 benzoate, which generates a LH surge, was administered (59 [↗](#)). Circulating levels of E2 were verified by the uterine weights (<25 mg for OVX and >95 mg for OVX E2-treated) at the time of euthanasia (59 [↗](#)).

AAV Delivery

Bilateral ARH injections of AAV1-Ef1a-DIO-ChR2:mCherry (RRID:Addgene_20297) or AAV1-Ef1a-DIO-ChR2:YFP (RRID:Addgene_20298) were performed on adult *Kiss1-cre* mice or *Pomc-cre* mice on a stereotaxic frame under isoflurane anesthesia. ARH injection coordinates were anteroposterior (AP): −1.10 mm, mediolateral (ML): $\pm$ 0.30 mm, dorsoventral (DL): −5.80 mm (surface of brain z = 0.0 mm); 400 nl of the AAV (2.0×10^{12} particles/ml) was injected (100 nl/min) into each position. Mice were given carprofen for analgesia and allowed to recover for at least two weeks before euthanasia.

Visualized whole-cell patch recordings

Hypothalamic coronal brain slices (240 μ m) were made from female mice, using a Leica VT1000S vibratome, in ice cold cutting solution bubbled with O_2/CO_2 (95%/5%). Slices were then transferred to a holding chamber with artificial cerebrospinal fluid bubbled with the same gas mix and allowed to recover for at least 1 hour. For recordings, slices were placed in a perfusion chamber and visualized with an Olympus BX51W1 using either differential infrared contrast or oblique illumination. *Kiss1* neurons in the ARH or AVPV/PeN were targeted for electrophysiological recordings as done previously and described below (60 [↗](#), 61 [↗](#)). Final concentration was calculated based on the drug in the known volume of the bath. Perfused drugs (TTX, naloxone, and STX) were constantly circulated and given at least 10 minutes to reach maximal effect. Focal application of drugs (MTII and THIQ) was done with the pump stopped (10–20 minutes). 0.3 to 1 μ l was directly added to the known volume of the bath to achieve the desired final concentration. This approach enables precise timing of recordings and reduces the risk of receptor desensitization.

Solutions/drugs

Standard vibratome slicing, external, and internal recording solutions were utilized as previously described (60 [↗](#), 61 [↗](#)). Tetrodotoxin was purchased from Alomone Labs (Jerusalem, Israel), Melanotan II and α MSH from Tocris (Minneapolis, MN). TEA, 4-AP, 17 β -estradiol benzoate, and Naloxone were purchased from Millipore-Sigma. STX was produced by AAPharmaSyn, LLC (Ann Arbor, MI) under contract.

Electrophysiology data analysis

Electrophysiological data were analyzed using Clampfit 10/11 (Molecular Devices) and Prism 7/10 (Dotmatics). All values are expressed as Mean $\pm$ SEM. Comparisons between two groups were made using un-paired Student's t-test or between multiple groups using an ANOVA (with *post hoc*

comparisons) with p-values < 0.05 considered significant. When variances differed significantly, Mann-Whitney U test was used instead.

Targeting of Kiss1 neurons for electrophysiological recordings

For non-optogenetic experiments, brain slices were taken from AAV injected Kiss1-Cre AAV or Kiss1xAi32 female mice. Ai32 mice (RRID:IMSR_JAX:024109, C57BL/6 background) carry the floxed ChR2 (H134R)-EYFP gene in their Gt(ROSA)26Sor Locus (56 [DOI](#)), allowing its expression in a Cre-dependent manner. Due to concerns of nonspecific expression (62 [DOI](#)), we previously validated this model using single cell RT-PCR and documented that *Kiss1* mRNA was detectable in 99% of individually harvested eYFP cells (n=126). In addition, we have used both AAV injected Kiss1-Cre AAV or Kiss1xAi32 female mice and found no differences in electrophysiological results (60 [DOI](#)). We used AAV injected POMC-Cre female mice and avoided small soma, high input resistance (>800 MΩ), ventrally located cells that are typically NPY/AgRP neurons. Instead, we targeted larger, more dorsomedial neurons in the ARH while avoiding fluorescent (POMC) cells. Unlike ARH Kiss1 neurons, POMC neurons do not make reciprocal projections, so uninfected POMC neurons do not display monosynaptic EPSCs in response to optogenetic stimulation. Next, we used a ramp IV protocol to probe for the presence of a persistent sodium current (I_{NaP}). While this current is more prevalent in the AVPV Kiss1 population, Kiss1 neurons are the only ARH neurons to display this electrophysiological “fingerprint” of a pronounced I_{NaP} paired with a high capacitance and low input resistance (26 [DOI](#)). Finally, we also harvested the cytosol at the end of all recordings and measured the expression of *Kiss1*. Only cells that expressed a persistent sodium current and/or expressed *Kiss1* mRNA were included in final analysis (n = 40 ARH & 6 AVPV).

Acknowledgements

The authors would like to recognize the excellent technical expertise of Ms. Martha A. Bosch (tissue preparation, immunohistochemical procedure and single cell RT-PCR of recorded neurons). In addition, we thank Dr. Rona Carroll for her assistance at Harvard Medical School and Mr. Cole Martinson, a student worker in the Ronnekleiv/ Kelly laboratories, for his assistance with genotyping and care of mouse colonies at OHSU. The University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core is supported by the Eunice Kennedy Shriver NICHD/NIH (NCTRI) Grant P50-HD28934. This work was supported by PHS MPI grant DK68098 to MJK and OKR; HD090151, HD099084 and DK133760 to VMN, and The Charles A. King Trust Postdoctoral Research Fellowship Award, The Lalor Foundation Postdoctoral Fellowship Award, the Women’s brain initiative Fellowship Award, the ROSA SCORE pilot grant (supported by NIH Research Grant U54 AG062322 funded by The National Institute on Aging (NIA) and Office of Research on Women’s Health (ORWH)) and the IBRO-ISN Research Fellowship Award to RT, and startup funds from Appalachian State University to TLS.

Additional information

Authors contribution

RT, TLS, MJK and VMN conceived and designed the study; RT, TLS, KF, CJH, EM, AG, KW, SL, EAM, OKR performed the experiments. RT, TLS, OKR, MJK and VMN analyzed data. RT, TLS and VMN wrote the manuscript with input from all the authors.

Supplementary figures

Supplemental figure 1 [↗](#)

Supplemental figure 2 [↗](#)

References

1. Health U. 2020) **Infertility** <http://obgyn.ucla.edu/infertility>
2. Andermann ML, Lowell BB 2017) **Toward a Wiring Diagram Understanding of Appetite Control** *Neuron* **95**:757–78
3. Cone RD 2006) **Studies on the physiological functions of the melanocortin system** *Endocr Rev* **27**:736–49
4. Shah BP, Vong L, Olson DP, Koda S, Krashes MJ, Ye C, et al. 2014) **MC4R-expressing glutamatergic neurons in the paraventricular hypothalamus regulate feeding and are synaptically connected to the parabrachial nucleus** *Proceedings of the National Academy of Sciences of the United States of America* **111**:13193–8
5. Wang K, Mao W, Zhang X, Zhao Y, Fan K, Pan D, et al. 2020) **Neuroanatomy of melanocortin-4 receptor pathway in the mouse brain** *Open Life Sci* **15**:580–7
6. Farooqi IS, Keogh JM, Yeo GS, Lank EJ, Cheetham T, O’Rahilly S 2003) **Clinical spectrum of obesity and mutations in the melanocortin 4 receptor gene** *The New England journal of medicine* **348**:1085–95
7. Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, et al. 2005) **Divergence of melanocortin pathways in the control of food intake and energy expenditure** *Cell* **123**:493–505
8. Hainerová IA, Zamrazilová H, Sedláčková D, Hainer V 2011) **Hypogonadotropic hypogonadism in a homozygous MC4R mutation carrier and the effect of sibutramine treatment on body weight and obesity-related health risks** *Obes Facts* **4**:324–8
9. Doulla M, McIntyre AD, Hegele RA, Gallego PH 2014) **A novel MC4R mutation associated with childhood-onset obesity: A case report** *Paediatr Child Health* **19**:515–8
10. Batarfi AA, Filimban N, Bajouh OS, Dallol A, Chaudhary AG, Bakhshab S 2019) **MC4R variants rs12970134 and rs17782313 are associated with obese polycystic ovary syndrome patients in the Western region of Saudi Arabia** *BMC Med Genet* **20**
11. Chen X, Huang L, Tan HY, Li H, Wan Y, Cowley M, et al. 2017) **Deficient melanocortin-4 receptor causes abnormal reproductive neuroendocrine profile in female mice** *Reproduction (Cambridge, England)* **153**:267–76
12. Sandrock M, Schulz A, Merkwitz C, Schoneberg T, Spanel-Borowski K, Ricken A 2009) **Reduction in corpora lutea number in obese melanocortin-4-receptor-deficient mice** *Reprod Biol Endocrinol* **7**
13. Cui L, Tan C, Huang L, Wang W, Huang Z, Geng F, et al. 2022) **Dapagliflozin partially restores reproductive function in MC4R KO obese female mice** *The Journal of endocrinology* **254**:65–76
14. Manfredi-Lozano M, Roa J, Ruiz-Pino F, Piet R, Garcia-Galiano D, Pineda R, et al. 2016) **Defining a novel leptin-melanocortin-kisspeptin pathway involved in the metabolic control of**

puberty *Mol Metab* **5**:844–57

15. Goodman RL, Herbison AE, Lehman MN, Navarro VM (2022) **Neuroendocrine control of gonadotropin-releasing hormone: Pulsatile and surge modes of secretion** *Journal of neuroendocrinology* **34**
16. Israel DD, Sheffer-Babila S, de Luca C, Jo YH, Liu SM, Xia Q, et al. (2012) **Effects of leptin and melanocortin signaling interactions on pubertal development and reproduction** *Endocrinology* **153**:2408–19
17. Manfredi-Lozano M, Roa J, Tena-Sempere M (2018) **Connecting metabolism and gonadal function: Novel central neuropeptide pathways involved in the metabolic control of puberty and fertility** *Front Neuroendocrinol* **48**:37–49
18. Cravo RM, Margatho LO, Osborne-Lawrence S, Donato J, Atkin S, Bookout AL, et al. (2011) **Characterization of Kiss1 neurons using transgenic mouse models** *Neuroscience* **173**:37–56
19. Lam BYH, Williamson A, Finer S, Day FR, Tadross JA, Gonçalves Soares A, et al. (2021) **MC3R links nutritional state to childhood growth and the timing of puberty** *Nature* **599**
20. Villa PA, Ruggiero-Ruff RE, Jamieson BB, Campbell RE, Coss D (2024) **Obesity alters POMC and kisspeptin neuron crosstalk leading to reduced luteinizing hormone in male mice** *The Journal of Neuroscience*
21. Stephens SBZ, Kauffman AS (2021) **Estrogen Regulation of the Molecular Phenotype and Active Translatome of AVPV Kisspeptin Neurons** *Endocrinology* **162**
22. Moore AM, Prescott M, Campbell RE (2013) **Estradiol negative and positive feedback in a prenatal androgen-induced mouse model of polycystic ovarian syndrome** *Endocrinology* **154**:796–806
23. McCarthy EA, Dischino D, Maguire C, Leon S, Talbi R, Cheung E, et al. (2022) **Inhibiting Kiss1 Neurons With Kappa Opioid Receptor Agonists to Treat Polycystic Ovary Syndrome and Vasomotor Symptoms** *J Clin Endocrinol Metab* **107**:e328–e47
24. Dewailly D, Barbotin AL, Dumont A, Catteau-Jonard S, Robin G (2020) **Role of Anti-Müllerian Hormone in the Pathogenesis of Polycystic Ovary Syndrome** *Frontiers in endocrinology* **11**
25. Roa J, Herbison AE (2012) **Direct regulation of GnRH neuron excitability by arcuate nucleus POMC and NPY neuron neuropeptides in female mice** *Endocrinology* **153**:5587–99
26. Zhang C, Bosch MA, Qiu J, Ronnekleiv OK, Kelly MJ (2015) **17beta-Estradiol increases persistent Na(+) current and excitability of AVPV/PeN Kiss1 neurons in female mice** *Mol Endocrinol* **29**:518–27
27. Clements JD, Westbrook GL (1991) **Activation kinetics reveal the number of glutamate and glycine binding sites on the N-methyl-D-aspartate receptor** *Neuron* **7**:605–13
28. Stincic TL, Grachev P, Bosch MA, Rønnekleiv OK, Kelly MJ (2018) **Estradiol drives the anorexigenic activity of proopiomelanocortin neurons in female mice** *eNeuro* **5**:e0103–18
29. Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, et al. (2016) **High-frequency stimulation-induced peptide release synchronizes arcuate kisspeptin neurons and excited GnRH neurons** *eLife* **5**

30. Qiu J, Rivera HM, Bosch MA, Padilla SL, Stincic TL, Palmiter RD, et al. 2018) **Estrogenic-dependent glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits in females** *eLife* **7**
31. Stincic TL, Qiu J, Connors AM, Kelly MJ 2021) **Rønnekleiv OK. Arcuate and Preoptic Kisspeptin neurons exhibit differential projections to hypothalamic nuclei and exert opposite postsynaptic effects on hypothalamic paraventricular and dorsomedial nuclei in the female mouse** *eneuro*
32. Herman MA, Ackermann F, Trimbuch T, Rosenmund C 2014) **Vesicular glutamate transporter expression level affects synaptic vesicle release probability at hippocampal synapses in culture** *The Journal of Neuroscience* **34**:11781–91
33. Qiu J, Rivera HM, Bosch MA, Padilla SL, Stincic TL, Palmiter RD, et al. 2018) **Estrogenic-dependent glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits in females** *Elife* **7**
34. Thornton JE, Loose MD, Kelly MJ, Rønnekleiv OK 1994) **Effects of estrogen on the number of neurons expressing β -endorphin in the medial basal hypothalamus of the female guinea pig** *The Journal of comparative neurology* **341**:68–77
35. Petersen SL, Keller ML, Carder SA, McCrone S 1993) **Differential Effects of Estrogen and Progesterone on Levels of POMC mRNA Levels in the Arcuate Nucleus: Relationship to the Timing of LH Surge Release** *Journal of Neuroendocrinology* **5**:643–8
36. Zhang C, Tonsfeldt KJ, Qiu J, Bosch MA, Kobayashi K, Steiner RA, et al. 2013) **Molecular mechanisms that drive estradiol-dependent burst firing of Kiss1 neurons in the rostral periventricular preoptic area** *American journal of physiology Endocrinology and metabolism* **305**:E1384–97
37. Mountjoy KG, Mortrud MT, Low MJ, Simerly RB, Cone RD 1994) **Localization of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain** *Molecular Endocrinology* **8**:1298–308
38. Yu J, Gimenez LE, Hernandez CC, Wu Y, Wein AH, Han GW, et al. 2020) **Determination of the melanocortin-4 receptor structure identifies Ca^{2+} as a cofactor for ligand binding** *Science* **368**:428–33
39. Sohn JW, Harris LE, Berglund ED, Liu T, Vong L, Lowell BB, et al. 2013) **Melanocortin 4 receptors reciprocally regulate sympathetic and parasympathetic preganglionic neurons** *Cell* **152**:612–9
40. Kelly MJ, Loose MD, Ronnekleiv OK 1992) **Estrogen suppresses mu-opioid- and GABAB-mediated hyperpolarization of hypothalamic arcuate neurons** *The Journal of neuroscience: the official journal of the Society for Neuroscience* **12**:2745–50
41. Lagrange AH, Ronnekleiv OK, Kelly MJ 1994) **The potency of mu-opioid hyperpolarization of hypothalamic arcuate neurons is rapidly attenuated by 17 beta-estradiol** *The Journal of neuroscience: the official journal of the Society for Neuroscience* **14**:6196–204
42. Qiu J, Bosch MA, Tobias SC, Grandy DK, Scanlan TS, Ronnekleiv OK, et al. 2003) **Rapid signaling of estrogen in hypothalamic neurons involves a novel G-protein-coupled estrogen receptor that activates protein kinase C** *The Journal of neuroscience: the official journal of the Society for Neuroscience* **23**:9529–40

43. Qiu J, Bosch MA, Tobias SC, Krust A, Graham SM, Murphy SJ, et al. 2006) **A G-protein-coupled estrogen receptor is involved in hypothalamic control of energy homeostasis** *The Journal of neuroscience: the official journal of the Society for Neuroscience* **26**:5649–55
44. Hessler XL Sabine, Herbison Allan E. 2020) **Direct inhibition of arcuate kisspeptin neurones by neuropeptide Y in the male and female mouse** *Journal of Neuroendocrinology*
45. Stincic TL, Grachev P, Bosch MA, Ronnekleiv OK, Kelly MJ 2018) **Estradiol Drives the Anorexigenic Activity of Proopiomelanocortin Neurons in Female Mice** *eNeuro* **5**
46. Ghamari-Langroudi M, Digby GJ, Sebag JA, Millhauser GL, Palomino R, Matthews R, et al. 2015) **G-protein-independent coupling of MC4R to Kir7.1 in hypothalamic neurons** *Nature* **520**:94–8
47. Krause WC, Rodriguez R, Gegenhuber B, Matharu N, Rodriguez AN, Padilla-Roger AM, et al. 2021) **Oestrogen engages brain MC4R signalling to drive physical activity in female mice** *Nature* **599**:131–5
48. Brix N, Ernst A, Lauridsen LLB, Parner E, Stovring H, Olsen J, et al. 2019) **Timing of puberty in boys and girls: A population-based study** *Paediatr Perinat Epidemiol* **33**:70–8
49. Padilla SL, Johnson CW, Barker FD, Patterson MA, Palmiter RD 2018) **A Neural Circuit Underlying the Generation of Hot Flushes** *Cell Rep* **24**:271–7
50. Dror T, Franks J, Kauffman AS 2013) **Analysis of multiple positive feedback paradigms demonstrates a complete absence of LH surges and GnRH activation in mice lacking kisspeptin signaling** *Biology of reproduction* **88**
51. Steyn FJ, Wan Y, Clarkson J, Veldhuis JD, Herbison AE, Chen C 2013) **Development of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult male mice** *Endocrinology* **154**:4939–45
52. Talbi R, Ferrari K, Choi JH, Gerutshang A, McCarthy EA, Dischino D, et al. 2021) **Characterization of the Action of Tachykinin Signaling on Pulsatile LH Secretion in Male Mice** *Endocrinology* **162**
53. Dave JR, Rubinstein N, Eskay RL 1985) **Evidence that b-endorphin binds to specific receptors in rat peripheral tissues and stimulates the adenylate cyclase-adenosine 3',5'-monophosphate system** *Endocrinology* **117**:1389–96
54. Naulé L, Robert V, Parmentier C, Martini M, Keller M, Cohen-Solal M, et al. 2015) **Delayed pubertal onset and prepubertal Kiss1 expression in female mice lacking central oestrogen receptor beta** *Hum Mol Genet* **24**:7326–38
55. Pfaffl MW 2001) **A new mathematical model for relative quantification in real-time RT-PCR** *Nucleic Acids Res* **29**
56. Madisen L, Mao T, Koch H, Zhuo JM, Berenyi A, Fujisawa S, et al. 2012) **A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing** *Nature Neuroscience* **15**:793–802
57. Balthasar N, Coppari R, McMinn J, Liu SM, Lee CE, Tang V, et al. 2004) **Leptin receptor signaling in POMC neurons is required for normal body weight homeostasis** *Neuron* **42**:983–91

58. Padilla SL, Carmody JS, Zeltser LM 2010) **Pomc-expressing progenitors give rise to antagonistic neuronal populations in hypothalamic feeding circuits** *Nature Medicine* **16**:403–5
59. Bosch MA, Tonsfeldt KJ, Ronnekleiv OK 2013) **mRNA expression of ion channels in GnRH neurons: subtype-specific regulation by 17beta-estradiol** *Molecular and cellular endocrinology* **367**:85–97
60. Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, et al. 2016) **High-frequency stimulation-induced peptide release synchronizes arcuate kisspeptin neurons and excites GnRH neurons** *Elife* **5**
61. Stincic TL, Qiu J, Connors AM, Kelly MJ, Ronnekleiv OK 2021) **Arcuate and Preoptic Kisspeptin Neurons Exhibit Differential Projections to Hypothalamic Nuclei and Exert Opposite Postsynaptic Effects on Hypothalamic Paraventricular and Dorsomedial Nuclei in the Female Mouse** *eNeuro* **8**
62. Qiu J, Stincic TL, Bosch MA, Connors AM, Kaech Petrie S, Ronnekleiv OK, et al. 2021) **Deletion of Stim1 in Hypothalamic Arcuate Nucleus Kiss1 Neurons Potentiates Synchronous GCaMP Activity and Protects against Diet-Induced Obesity** *The Journal of neuroscience: the official journal of the Society for Neuroscience* **41**:9688–701

Author information

Rajae Talbi*

Harvard Medical School, Boston, United States, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

ORCID iD: [0000-0001-7158-6246](https://orcid.org/0000-0001-7158-6246)

*These authors contributed equally

Todd L Stincic^{†*}

Division of Neuroscience, Oregon National Primate Research Center, Beaverton, United States

[†]Present address: Department of Biology, Appalachian State University, Boone.

*These authors contributed equally

Kaitlin Ferrari

Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Choi Ji Hae

Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Karol Walec

Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Elizabeth Medve

Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Achi Gerutshang

Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Silvia León

Harvard Medical School, Boston, United States, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Elizabeth A McCarthy

Harvard Medical School, Boston, United States, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States

Oline K Rønnekleiv

Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, United States, Division of Neuroscience, Oregon National Primate Research Center, Beaverton, United States

Martin J Kelly

Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, United States, Division of Neuroscience, Oregon National Primate Research Center, Beaverton, United States

For correspondence: kellym@ohsu.edu

Víctor M Navarro

Harvard Medical School, Boston, United States, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, United States, Harvard Program in Neuroscience, Boston, United States

For correspondence: vnavarro@bwh.harvard.edu

Editors

Reviewing Editor

Ashley Webb

Buck Institute for Research on Aging, Novato, United States of America

Senior Editor

Ma-Li Wong

State University of New York Upstate Medical University, Syracuse, United States of America

Reviewer #1 (Public review):

Summary:

The authors investigate the role of the melanocortin system in puberty onset. They conclude that POMC neurons within the arcuate nucleus of the hypothalamus provide important but differing input to kisspeptin neurons in the arcuate or rostral hypothalamus.

Strengths:

Innovative and novel
Technically sound
Well-designed
Thorough

Weaknesses:

There were no major weaknesses identified.

<https://doi.org/10.7554/eLife.100722.2.sa3>

Reviewer #2 (Public review):

Summary:

This interesting manuscript describes a study investigating the role of MC4R signalling on kisspeptin neurons. The initial question is a good one. Infertility associated with MC4 mutations in humans has typically been ascribed to the consequent obesity and impaired metabolic regulation. Whether there is a direct role for MC4 in regulating the HPG axis has not been thoroughly examined. Here, the researchers have assembled an elegant combination of targeted loss of function and gain of function in vivo experiments, specifically targeting MC4 expression in kisspeptin neurons. This excellent experimental design should provide compelling evidence for whether melanocortin signalling directly affects arcuate kisspeptin neurons to support normal reproductive function. There were definite effects on reproductive function (irregular estrous cycle, reduced magnitude of LH surge induced by exogenous estradiol). However, the magnitude of these responses and the overall effect on fertility were relatively minor. The mice lacking MC4R in kisspeptin neurons remained fertile despite these irregularities. The second part of the manuscript describes a series of electrophysiological studies evaluating the pharmacological effects of melanocortin signalling in kisspeptin cells in ex-vivo brain slices. These studies characterised interesting differential actions of melanocortins in two different populations of kisspeptin neurons. Collectively, the study provides some novel insights into how direct actions of melanocortin signalling via the MC4 receptor in kisspeptin neurons contribute to the metabolic regulation of the reproductive system. Importantly, however, it is clear that other mechanisms are also at play.

Strengths:

The loss of function/gain of function experiments provides a conceptually simple but hugely informative experimental design. This is the key strength of the current paper - especially the knock-in study that showed improved reproductive function even in the presence of ongoing obesity. This is a very convincing result that documents that reproductive deficits in MC4R knockout animals (and humans with deleterious MC4R gene variants) can be ascribed to impaired signalling in the hypothalamic kisspeptin neurons and not necessarily caused as a consequence of obesity. As concluded by the authors: "reproductive impairments observed in MC4R deficient mice, which replicate many of the conditions described in humans, are

largely mediated by the direct action of melanocortins via MC4R on Kiss1 neurons and not to their obese phenotype." This is important, as it might change how such fertility problems are treated.

I would like to see the validation experiments for the genetic manipulation studies given greater prominence in the manuscript because they are critical to interpretation. Presently, only single unquantified images are shown, and a much more comprehensive analysis should be provided.

Weaknesses:

(1) Given that mice lacking MC4R in kisspeptin neurons remained fertile despite some reproductive irregularities, this can be described as a contributing pathway, but other mechanisms must also be involved in conveying metabolic information to the reproductive system. This is now appropriately covered in the discussion.

(2) The mechanistic studies evaluating melanocortin signalling in kisspeptin neurons were all completed in ovariectomised animals (with and without exogenous hormones) that do not experience cyclical hormone changes. Such cyclical changes are fundamental to how these neurons function in vivo and may dynamically alter how they respond to hormones and neuropeptides. Eliminating this variable makes interpretation difficult, but the authors have justified this as a reductionist approach to evaluate estradiol actions specifically. However, this does not reflect the actual complexity of reproductive function.

For example, the authors focus on a reduced LH response to exogenous estradiol in ovariectomised mice as evidence that there might be a sub-optimal preovulatory LH surge. However, the preovulatory LH surge (in intact animals) was not measured.

They have not assessed why some follicles ovulated, but most did not. They have focused on the possibility that the ovulation signal (LH surge) was insufficient rather than asking why some follicles responded and others did not. This suggests some issue with follicular development, likely due to changes in gonadotropin secretion during the cycle and not simply due to an insufficient LH surge.

<https://doi.org/10.7554/eLife.100722.2.sa2>

Reviewer #3 (Public review):

The manuscript by Talbi R et al. generated transgenic mice to assess the reproduction function of MC4R in Kiss1 neurons in vivo and used electrophysiology to test how MC4R activation regulated Kiss1 neuronal firing in ARH and AVPV/PeN. This timely study is highly significant in neuroendocrinology research for the following reasons.

(1) The authors' findings are significant in the field of reproductive research. Despite the known presence of MC4R signaling in Kiss1 neurons, the exact mechanisms of how MC4R signaling regulates different Kiss1 neuronal populations in the context of sex hormone fluctuations are not entirely understood. The authors reported that knocking out *Mc4r* from Kiss1 neurons replicates the reproductive impairment of MC4RKO mice, and *Mc4r* expression in Kiss1 neurons in the MC4R null background partially restored the reproductive impairment. MC4R activation excites Kiss1 ARH neurons and inhibits Kiss1 AVPV/PeN neurons (except for elevated estradiol).

(2) Reproduction dysfunction is one of obesity comorbidities. MC4R loss-of-function mutations cause obesity phenotype and impaired reproduction. However, it is hard to determine the causality. The authors carefully measured the body weight of the different mouse models (Figure 1C, Figure 2A, Figure 3B). For example, the Kiss1-MC4RKO females

showed no body weight difference at puberty onset. This clearly demonstrated the direct function of MC4R signaling in reproduction but was not a consequence of excessive adiposity.

(3) Gene expression findings in the "KNDy" system align with the reproduction phenotype.

(4) The electrophysiology results reported in this manuscript are innovative and provide more details of MC4R activation and Kiss1 neuronal activation.

Overall, the authors have presented sufficient background in a clear, logical, and organized structure, clearly stated the key question to be addressed, used the appropriate methodology, produced significant and innovative main findings, and made a justified conclusion.

Comments on revisions:

The authors have addressed my comments.

<https://doi.org/10.7554/eLife.100722.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public reviews:

We are grateful to the reviewers and the editorial team for their feedback and thorough revisions of our paper. We also appreciate their acknowledgement that this study represents a significant advancement in the field of reproductive neuroendocrinology and offers insights on the contribution of obesity vs melanocortin signaling in women's fertility. In the revised version, we will provide a more detailed clarification of the data and methodology and adhere to the reviewers' suggestions.

Please find below our answers to specific concerns in the public review:

Given the fact that mice lacking MC4R in Kiss1 neurons remained fertile despite some reproductive irregularities, the overall tone and some of the conclusions of the manuscript (e.g., from the abstract: "... Mc4r expressed in Kiss1 neurons is required for fertility in females") were overstated. Perhaps this can be described as a contributing pathway, but other mechanisms must also be involved in conveying metabolic information to the reproductive system.

We will tone down these statements throughout the manuscript to indicate that MC4R in Kiss1 neurons plays a role in the metabolic control of fertility (rather than "...is required for fertility")

The mechanistic studies evaluating melanocortin signalling in Kiss1 neurons were all completed in ovariectomised animals (with and without exogenous hormones) that do not experience cyclical hormone changes. Such cyclical changes are fundamental to how these neurons function in vivo and may dynamically alter the way they respond to neuropeptides. Therefore, eliminating this variable makes interpretation difficult.

Mice lack true follicular and luteal phases and therefore it is impossible to separate estrogen-mediated changes from progesterone-mediated changes (e.g., in a proestrous female). Therefore, we use an ovariectomized female model in which we can generate a LH surge with an E2-replacement regimen [1]. This model enables us to focus on estrogen effects,

exclude progesterone effects, and minimize variability. Inclusion of cycling females would make interpretation much more difficult.

(1) Bosch et al., 2013 Mol & Cell Endo; <https://doi.org/10.1016/j.mce.2012.12.021>

Use of the POMC-Cre to target ontogenetic inputs to Kiss1 neurons might have targeted a wider population of cells than intended.

POMC is transiently expressed during embryonic development in a portion of cells fated to be Kiss1 or NPY/AgRP neurons [1-2]. Therefore, this is a valid concern when crossing with a floxed mouse. However, use of AAVs in adult animals avoids this issue and leads to specific expression in POMC neurons [3]. This POMC-Cre mouse has been used extensively with AAVs to drive specific expression in POMC neurons by other laboratories [4-7]. Therefore, we are confident that our optogenetic studies have narrowly targeted POMC inputs.

(1) Padilla et al., 2010 Nat Med; <https://doi.org/10.1038/nm.2126>

(2) Lam et al., 2017 Mol Metab; <https://doi.org/10.1016/j.molmet.2017.02.007>

(3) Stincic et al., 2018 eNeuro; <https://doi.org/10.1523/eneuro.0103-18.2018>

(4) Fenselau et al., 2017 Nat Neuro; <https://doi.org/10.1038/nn.4442>

(5) Rau & Hentges, 2019 J Neuro; <https://doi.org/10.1523/jneurosci.3193-18.2019>

(6) Fortin et al., 2021 Nutrients; <https://doi.org/10.3390/nu13051642>

(7) Villa et al., 2024 J Neuro; <https://doi.org/10.1523/jneurosci.0222-24.2024>

Recommendations for Authors

We thank the reviewers and the editorial team for their comments and thorough revisions of our paper. We have now addressed their comments and edited the manuscript accordingly:

Reviewer #1 (Recommendations For The Authors):

L80 - This is an awkward sentence; it isn't an inverse agonist of the AgRP; this may read better just to say that the inverse agonist, AgRP.

Thank you for this comment. This has now been changed in the text (L80).

L86 - This text reads as if mice have an inherent obesity issue.

This has also now been addressed in the text (L86).

L131 - The numbers of digits past the decimal point should match for both mean and SEM.

This has also now been addressed throughout the text.

Figure 1D: Revise the bar graphs with distinct SEM bars, as these data are not generated within the same mice.

The graphs are now changed, and they include distinct SEM and individual data points.

Figure 2I-L - An n of 3 for controls is pretty minimal, though the clustering of data points is tight.

We thank the reviewer for this comment, and we emphasize that while we agree that an n=3 for controls is minimal, the mRNA level values of this group are close, therefore the clustering of the data points is tight. We are happy to provide the raw data value for these groups if the reviewer wishes to.

L159 - The role of reduced dynorphin mRNA is pretty speculative with regard to basal levels of LH, especially since no other indices of LH secretion were affected. It should also be recognized that mRNA levels do not always equate to activity.

We agree with the reviewer that our explanation of the role of the reduced dynorphin with regards to the elevated basal LH is speculative, however, we only report that the higher LH levels correlates with the lower expression of the Pdyn gene expression, which is in line with the well documented role of Dynorphin on inhibiting LH secretion. We also recognize that mRNA levels don't necessarily reflect activity. We have now added this statement to the text (L159).

L164 - Given the ovary data, it seems that the increase seen in KO mice isn't quite sufficient, but is it known how much of a surge is necessary for ovulation in mice?

We agree with the reviewer's comment that the LH surge in Kiss1MC4RKO group is not enough to consistently induce ovulation, which is supported by the decrease in the numbers of corpora lutea data (Figure 2, O).

According to literature, an LH surge in the female mice is estimated by a LH value >4 ng/ml (Bahougne et al., 2020). According to this rule, our data show that only two females out of six had LH surge in the KO group, while four females out of five had LH surge in the control group.

L211 - According to the figure, LH pulses were not recovered and remained similar to KO levels. Looking at the LH secretory patterns presented, it seems like the pulse frequency data should be interpreted with some caution, given that some of the pulses identified are tenuous at best.

We agree that the LH pulses identified by our software (criteria described in the methods) are variable in shape (LH pulses are difficult to detect clearly in gonad intact females) and did not differ in number between groups; however, the reinsertion of Mc4r within Kiss1 neurons restored LH basal levels, amplitude and total secretory mass, which are clear indicatives of a significant improvement in the ability of these mice to release LH.

L218 - Is there a reason why the surge was not looked at in these groups?

Ovarian histology is the best indicator of ovulation. In these mice, corpora lutea were absent, indicating impaired ovulation, thus, we did not consider performing an LH surge protocol was necessary.

L244 - This would also fit with previous findings in sheep that not all Kiss neurons express MC receptors

We agree with this comment.

L329 - Given the rapidity of its actions, how would this membrane ER function during a normal surge?

Rapid estrogen signaling can act to ease transitions between states. Membrane delimited E2 actions can quickly attenuate or enhance coupling between receptors and signaling cascades. These effects will precede E2-driven changes in gene expression that produce more stable alterations in signaling. This combination of mechanisms will reduce any lag between rises in serum E2 and physiological effects. Considering the abbreviated mouse reproductive cycle, parallel mechanisms acting on different timescales are particularly important.

L365 - I'm a little confused as to how this particular work sheds light on a role for MC3R. Is the relative distribution of the two isoforms within Kiss neurons known?

In the present study, we report that hypothalamic Mc3r expression decreases leading up to the age of puberty onset (p30), in line with the profile of expression of Mc4r and a recent publication involving Mc3r in puberty onset (Lam et al., 2021), suggesting that both receptors may be involved in the control of reproductive function, potentially through the direct regulation of Kiss1 neurons as characterized in our present study.

L422 - While I understand the nature of this statement, the receptor may simply reflect the activity of what binds to it, i.e., AgRP vs. alpha-MSH, suggesting that maybe the prepubertal period is more AgRP-dominated.

We agree with this statement, and this needs to be further investigated.

L495 - Reinsertion of Mc4R in Kiss1 neurons

Thank you for this comment. This is now corrected in the text (L501).

L524 - Bilateral ovariectomy of 6-month

Thank you for this comment. This is now corrected in the text (L530).

L538 - Is it known what stage of the cycle these mice were in when samples were collected?

Yes, the samples were collected in diestrus. This is now mentioned in the text (L548)

L556 - Pulse amplitude is usually measured relative to the preceding nadir.

The method that we have been consistently using in our lab is the average of the 4 highest LH values in the samples collection period for each animal. We have found this to be consistent and representative of the overall amplitude (McCarthy et al., 2021; Talbi et al., 2021).

L594 - This is a little confusing - the whole MBH would contain the ARH, but only the ARH was collected from the KO mice. If the whole MBH, dynorphin and Tac3, and Tac3 are expressed outside of the ARC, making interpretation of changes specifically within the ARH is difficult.

Here (L592), we describe two different experiments, as mentioned by i) and ii).

For experiment 1 (i): MBH was used in the WT mice at ages P10, P15, P22 and P30 to investigate the expression of the melanocortin genes (Agrp, Pomc, Mc3r and Mc4r).

For experiment 2 (ii): In both KO and control groups, only the micro-dissected ARH was used to investigate genes expressions of Pdyn, Kiss1, Tac2, Tacr3.

Reviewer #2 (Recommendations For The Authors):

The validation experiments for the various manipulations are currently presented in the supplementary data. Still, in my opinion, these are critically important for interpreting the data, and it should be considered to present these more comprehensively in the main body of the manuscript. In Figure S1, it seems that the exposure of the two images is not the same, with a higher background in the control. Has this image been adjusted to highlight the staining, while the other has not? It looks like there remains a low level of expression still present in at least some of the KO cells - this may reflect difficulties using RNAscope (with its extreme amplification) to detect the absence of a signal, or it could also be that the knockout is incomplete. A percentage of cells still express MC4R. I think this should be acknowledged or discussed.

We thank the reviewer for the feedback. While we agree that the validation of the mouse model is critical, we would like to keep it in the supplemental data.

We also agree that the exposure looks different between the KO and WT controls, and we thank the reviewer for this comment. The quality of the photograph decreased when transferring to the manuscript. This has now been improved in the revised figure.

As for the MC4R expression in some of the KO cells, we believe that MC4R is expressed in non Kiss1 cells as shown in the merged figure. Therefore, we believe that the Knockout of Mc4r in Kiss1 neurons is complete in these mice.

The clear difference from the PVN's lack of effect is convincing and indicates that a specific knockout has been achieved. Is equivalent data also available for the AVPV population of cells that are examined later in the manuscript? Do those Kiss1 neurons also express the MC4R? The same question applies to the knock-in experiment: Was the expression of MC4R also driven in the AVPV population using this approach

Yes, Kiss1 neurons in the AVPV also express MC4R as indicated in this study, and thus Mc4r is removed/reinserted in the AVPV as well in this mouse model.

The quantitative RT-qPCR data on developmental changes in metabolic signaling molecules are really peripheral to the paper's main question. Relative to the validation experiments (as discussed above), I think these are less important data and could be placed into a supplementary figure. The discussion of these data becomes problematic, e.g., on line 359, the changes are described as "a low melanocortin tone..." but this seems problematic when referring to reduced expression of AgRP, an inverse agonist at the MC4R. If you are going to present these data, individual data points should be shown. Similarly, the question about whether this is a PCOS-like phenotype is perhaps worth asking. Still, the simple assessment of T and AMH could also be reported in a sentence without necessarily showing the data (or placing it in a supplementary figure). Better to focus on the key question - which is the role of MC4R signaling in Kiss1 neurons.

We understand this reviewer's concerns, however, due to the impact of MC4R signaling (particularly in the context of AgRP) on puberty, we strongly believe that the reader will benefit from expression profile across ages so we will respectfully disagree and keep in the main figure.

Per this reviewer's comment, we have now added individual data points to Figure 1D.

We also agree with the reviewer that the T and AMH data are not in the main scope of the paper, but since we uncovered a PCOS-like phenotype in female mice with specific deletion of

Mc4r from Kiss1 neurons, it is important to keep these data in the main figure to show that the phenotype does not fully resemble a PCOS model.

Having praised the experimental design, I think it is fair to acknowledge that the reproductive data from these experiments remain difficult to interpret. I understand that it is difficult to illustrate estrous cycles, but the "quantitative" data on percentages of time spent in any one stage are not as informative as seeing the actual individual patterns in Figure 2B. Were all of the animals consistently like the one illustrated, with persistent diestrus and only occasional evidence of ovulation?

We agree that Figure 2C may be difficult to interpret but it is the best way to capture the all the data points for each group.

All the 5 Kiss1MC4RKO females had persistent diestrus phases with only one or two estrus phases over 15 days (except for one female who had 4 estrous days), compared to control females who had 7 to 9 days of estrous, as shown in the graph (except for one female who had 5 days of estrus over 15 days period).

Given that LH pulses appear to be normal, does this, in fact, suggest an ovarian problem? Is that possible? Are MC4R and Kiss1 co-expressed in the ovary? Or do you think this suggests an ovulation problem, perhaps driven by the impaired LH surge?

This reviewer is correct in that our findings suggest a central defect in ovulation based on the deficit observed in the preovulatory LH surge. Thus, it is possible to have normal LH pulses, which are driven by one population of Kiss1 neurons (ARH) and the LH surge, driven by a distinct population of Kiss1 neurons (AVPV).

Similarly, the response to the "LH surge induction protocol" is impaired (why not look at endogenous LH surges?). It seems that ovulation should be an all-or-none phenomenon in that if the LH surge is sufficient to induce ovulation, then all available follicles would be ovulated. If it is not, then no follicles will be ovulated. Why fewer follicles are ovulated in the gene-targeted animals seems more likely to be due to impaired follicular development rather than a subthreshold LH surge. So, this again points back to the ovary. Or perhaps we need a more thorough assessment of the pattern of LH pulses throughout the cycles in these animals.

An LH surge induction protocol allows us to submit all female mice to the same conditions and expect a similar response, which is then optimal to compare with animals with an expected ovulation deficit, as it eliminates external factors. We disagree in that ovulation is an all-or-none phenomenon because in mice numerous follicles mature at the same time and thus a decrease in the number of ovulated oocytes may be significant between groups even if the animals are not completely infertile.

Collectively, my assessment of these data is that there are effects on reproduction, but they are actually relatively subtle. There were abnormal cycles and impaired LH surge in response to exogenous estrogen. But the animals are not actually infertile, so can ovulate and express normal reproductive behavior. So while there is a role for MC4R signalling in Kiss1 neurons, it may be a contributing modulatory role rather than a major regulatory mechanism. I think the tone of the descriptions should reflect this. I like the way it is framed in some parts of the discussion ("reproductive impairments...mediated by MC4R in Kiss1 neurons and not by their obese phenotype"), but the overall significance of this is overstated in some places, such as the abstract and in other parts of the discussion ("this population is tightly controlled by melanocortins").

As mentioned in previous responses, ovulation in mice is not all-or nothing, so while the mice can reproduce, the disruption in the central mechanisms that control ovulation and irregular estrous cycles are a significant advancement in the field with strong translational potential to species where only one oocyte is usually ovulated, like in humans, where reproductive disorders in MC4R patients had been attributed to the obesity phenotype rather than to a central action of MC4R (as the reviewer captured in their comment). This is one of the main findings of this study.

The overstatement has been now addressed throughout the text.

For in vitro studies, all mice were ovariectomized and given estradiol "replacement." What was the rationale for this? Wouldn't this suppress the basal activity of these neurons? Then it appears that some of the animals were studied as ovariectomised (for an unspecified time but apparently ">7 days", without hormone replacement. The basal activity of these cells would be dramatically different. I think these artificial manipulations make these data quite difficult to interpret. How does this reflect the situation in a normal (or abnormal) estrous cycle? My understanding is that the brain slice approach already compromises the ability of this population of cells to function as a coordinated network (i.e., coordinated episodes of activity that are seen in vivo have not been observed in vitro in brain slices). Ovariectomizing and providing exogenous hormones also removes the additional regulatory elements of the cyclical changes in hormone inputs, so the cells may or may not behave like they would in vivo. Perhaps the authors could justify their choice of experimental model.

We have clarified that the mice were ovariectomized for 7-10 days. A group of 3 mice are OVXed at once and then used on subsequent days a week later. This delay is both for the recovery of the animal and to allow for “washout” of endogenous ovarian hormones. For optogenetic studies, we were not measuring basal activity. Rather, we prioritized the ability to detect a postsynaptic response. While E2 decreases the networked activity of Kiss1- ARH neurons, the Hcn channels, calcium channels, and Vglut2 expression are all increased. This leads to increased excitability and more glutamate release. Mice lack true follicular and luteal phases and therefore it is impossible to separate estrogen-mediated changes from progesterone-mediated changes (e.g., in a proestrous female). Therefore, we use an ovariectomized female model in which we can generate a LH surge with an E2-replacement regimen (Bosch et al., J Mol Cell Endocrinology 2013). This model enables us to focus on estrogen effects, exclude progesterone effects, and minimize variability. Finally, we have documented that Kiss1^{ARH} neurons retain the synchronization of their neuronal firing in the hypothalamic slice preparation (Qiu et al., eLife 2016).

Figure 4E shows neurons' staining after expressing a Cre-dependent channel rhodopsin vector into POMC-Cre mice. The number of labelled cells looks markedly larger than expected for adult POMC neurons. Was the specificity of this approach to neurons expressing POMC checked? I understand that the POMC-Cre mice have been criticised for ectopic expression of Cre during development in other populations of neurons in the arcuate nucleus that does not express POMC, such as the AgRP neurons (e.g., PMID: 22166984). Is it possible that this is not a problem in adult animals? Has that been validated in these animals? The description of the method suggests that it is acknowledged that some of the expression driven in these animals might be in AgRP neurons. Still, optogenetic activation of these cells will include all cells expressing Cre at the time of AAV administration.

POMC is transiently expressed during embryonic development in a portion of cells fated to be Kiss1 or NPY/AgRP neurons. Therefore, this is a valid concern when crossing with a floxed mouse. However, use of AAVs in adult animals avoids this issue and leads to specific

expression in POMC neurons. This POMC-Cre mouse has been used extensively with AAVs to drive specific expression in POMC neurons by other laboratories (Padilla et al., Nat Med 2010; Lam et al., Mol Metab 2017; Stincic et al., eNeuro 2018 eNeuro; Fenselau et al., Nat Neuro 2017). We have previously shown that AAV-driven mCherry expression is limited to cells labeled with a beta-endorphin antibody (Stincic et al., 2018 eNeuro). Therefore, we are confident that our optogenetic studies have narrowly targeted POMC inputs.

Some additional explanation of the electrophysiology result may be required. For example, on Line 292, I'm confused by Fig 4M. Why is the response to 20Hz stimulation different in this cell (compared to the one in 4L) before administering naloxone? What proportion of cells showed this opposite response? On line 307: Is 5 cells sufficient for testing the POMC inputs onto AVPV and PeN Kiss1 neurons? How many slices/animals are included in collecting these 5 cells? The rapid action of STX illustrates the ability to modulate the response to MTII, but I am struggling to understand the implications of this in a physiological context. Suppose this response is desensitized by longer-term treatment with E2 (as indicated in the manuscript). Is it relevant to normal regulation during the cycle (particularly in the AVPV, where the key regulatory step seems to be the prolonged exposure to high estradiol as part of the preovulatory signals leading up to the LH surge)?

As stated in the text, E2 has been shown to increase POMC expression and beta-Endorphin immunostaining. We do not know the effects of E2 on aMSH expression and release. E2 also tends to attenuate the coupling between inhibitory postsynaptic metabotropic (Gi,o-coupled) receptors and signaling cascades. So, there is likely a combination of pre- and post-synaptic mechanisms contributing to these responses. However, the focus of the current studies was on the predominant melanocortin signaling and, as such, we chose to eliminate the influence of opioid signaling. We have added two more cells to this group, both of which were successfully rescued for a total of 5 of 6 cells (6 slices, 5 animals). Between the labeling of b-endorphin fibers and high rate of rescue, we do believe that this is sufficient evidence to support a direct POMC input to Kiss1^{AVPV/PeN} neurons.

Line 52: "Here, we show that Mc4r expressed in Kiss1 neurons is required for fertility in females." The knockout animals remain fertile, so this conclusion needs to be re-worded.

Thank you for this comment. This has now been changed (L52).

Line 80: "The melanocortin 4 receptor (MC4R) binds α -melanocyte stimulating hormone (aMSH), an agonist product of the pro-opiomelanocortin (Pomc) gene, and the inverse agonist of the agouti-related peptide (AgRP) to regulate food intake and energy expenditure" Is this the correct wording? I think it should be stated that AgRP is an inverse agonist at the MC4R, not that aMSH is the inverse agonist of AgRP. Re-work this sentence.

Thank you for this comment. This has now been changed (L79-80).

Line 88: "... however, conflicting reports exist". Describe what these conflicting reports show. Many MC4 variants ("mutations") are expressed in humans, but few will fully inactivate signalling like the mouse knockout.

We thank the reviewer for this comment. By conflicting data, we refer to the studies that report no reproductive impairments in women with MC4R mutations. Either because the metabolic impairments (obesity, hyperphagia, hyperinsulinemia, hyperleptinemia, etc) are so strong that the focus is skewed to these issues, without a full reproductive assessment in these women, or simply because the reviewer mentioned, not all MC4R mutations fully

inactivate its signaling in humans - as opposed to mouse models where reproductive disruption has been described previously in full body MC4RKOs.

Line 91: "...that largely affects females". Is this a genuine sex difference, or are reproductive deficits simply more overt in female rodents? I think the Coss paper (reference 19 in the manuscript) showed a greater effect of diet-induced obesity in males than in females.

We believe that sex differences exist with regards to the role of MC4R in the regulation of fertility, as we show that most of this effect is mediated by MC4R signaling in Kiss1 AVPV neurons, a neuronal population that is specific to the female brain.

As far as we can tell, the Coss paper (Villa et al., 2024) has only tested males but not females. Moreover, they investigated the effect of diet induced obesity in mice on their fertility (specifically LH secretion), while in this study we are specifically looking at the deletion of MC4R from Kiss1 neurons, and these mice were not obese (Figure 2A). While both these conditions induce impaired fertility, the mechanisms and signaling pathways are different (our mice lack MC4R signaling while the obese mice have a decrease in MC4R expression but the signaling is still functional).

Line 392: also Hessler et al. PMID: 32337804.

This reference is now added to the text (Line 393).

Line 433. The discussion of how advanced puberty onset (seen in the Kiss1-specific KO animals) might be caused by MC4R signalling in AVPV Kiss1 neurons, which are sexually dimorphic, which might explain sex differences in puberty timing in mammals seems extremely speculative and based on limited data. More targeted experiments would be needed to address this, and I think this speculation should be removed here.

This speculation has now been removed from the text.

Line 438: "Furthermore, our findings suggest that metabolic cues, through the regulation of the melanocortin output onto Kiss1AVPV/PeN neurons, are essential for the timing and magnitude of the GnRH/LH surge." Again, I think this is overstating the present data, which has only looked at an artificial hormone administration regime. The animals are fertile and, thus, must be able to mount a sufficient LH surge. The major effect, in fact, seems to be on their cycle, perhaps leading to impaired follicular development. Please acknowledge that this will be one of the multiple pathways by which metabolic information is fed into the HPG axis.

In addition to the effect on their cycles as mentioned by the reviewer, the Kiss1MC4RKO females also display impaired fertility (Figure 2, S-T) and fewer corpora lutea which is in line with the impaired mounting of LH surge (Figure 2, M). Even if the LH surge is induced by the hormone administration protocol, it only reflects the natural ability of the HPG axis to mount the surge, as this regimen is only there to mimic the endogenous hormonal changes leading to LH surge and therefore ovulation, in a controlled manner. Nonetheless, we agree with this reviewer that this is not the sole mechanism by which metabolism regulates reproductive function and this has been emphasized in the paper. (line 443)

Reviewer #3 (Recommendations For The Authors):

The decreased melanocortin tone drives puberty onset (Figure 1D), and this is correlative. The transgenic animals' hypothalamic expression of Agrp, Pomc, Mc4r, and Mc3r can be

measured to strengthen the claim. Hprt expression should be demonstrated, as this housekeeping gene was used as a common denominator.

We thank the reviewer for this comment. While we think that indeed, measuring Agrp, Pomc, Mc4r, and Mc3r gene expressions in the transgenic mice will strengthen our claim and give more insights into the melanocortins tone during pubertal maturation, this is unfortunately not feasible as it will involve generating a lot of mice (at least n=40 pups for an n=5/group, KO and control littermates, females only -which will require setting up lots of breeding pairs-) during different ages throughout puberty.

As for the gene expression of Hprt, because we have 6 mice per age, 4 ages total, every gene (Agrp, Pomc, Mc4r, Mc3r) was run in a separate plate with Hprt as its own housekeeping gene. Samples were run in duplicates for each Hprt and melanocortin genes in a 96 well = 48 wells for Hprt and 48 wells for each of the melanocortin genes. Therefore, it won't be possible to represent one Hprt expression for all the four genes, however every gene was normalized to the Hprt gene expression that was ran in the same plate).

In Figures 4 and 5, dot plots can be used (as opposed to the bar graphs) to better reflect the individual data points.

Figures 4 and 5 have been revised to include individual data points.

The electrophysiology experiment requires more details in the method section. In addition to the publication cited, a brief recap of the methodology used in this paper, such as the focal application of MTII (Figure 4B), is also needed.

We have added more details to the Methods.

<https://doi.org/10.7554/eLife.100722.2.sa0>