

Decoding Phase Separation of Prion-Like Domains through Data-Driven Scaling Laws

Reviewed Preprint

v2 • January 6, 2025

Revised by authors

Reviewed Preprint

v1 • July 22, 2024

M Julia Maristany, Anne Aguirre Gonzalez, Jorge R Espinosa, Jan Huertas, Rosana Colleparado-Guevara, Jerelle A Joseph 

Department of Physics, University of Cambridge, Cambridge, UK • Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK • Universidad Complutense de Madrid, Madrid, Spain • Department of Genetics, University of Cambridge, Cambridge, UK • Department of Chemical and Biological Engineering, Princeton University, Princeton, USA • Omenn-Darling Bioengineering Institute, Princeton University, Princeton, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

The authors performed extensive coarse-grained molecular dynamics simulations of 140 different prion-like domain variants to interrogate how specific amino acid substitutions determine the driving forces for phase separation. The analyses are **solid**, and the derived predictive scaling laws can aid in identifying potential phase-separating regions in uncharacterized proteins. Overall, this is a **valuable** contribution to the field of biomolecular condensates. It exemplifies how data-driven methodologies can uncover new insights into complex biological phenomena.

<https://doi.org/10.7554/eLife.99068.2.sa4>

Abstract

Proteins containing prion-like low complexity domains (PLDs) are common drivers of the formation of biomolecular condensates and are prone to misregulation due to amino acid mutations. Here, we exploit the accuracy of our residue-resolution coarse-grained model, Mpipi, to quantify the impact of amino acid mutations on the stability of 140 PLD mutants from six proteins (hnRNPA1, TDP43, FUS, EWSR1, RBM14, and TIA1). Our simulations reveal the existence of scaling laws that quantify the range of change in the critical solution temperature of PLDs as a function of the number and type of amino acid sequence mutations. These rules are consistent with the physicochemical properties of the mutations and extend across the entire family tested, suggesting that scaling laws can be used as tools to predict changes in the stability of PLD condensates. Our work offers a quantitative lens into how the emergent behavior of PLD solutions varies in response to physicochemical changes of single PLD molecules.

Introduction

Biomolecular condensates are highly multicomponent systems with remarkable specificity: they exhibit non-random molecular compositions and, concomitantly, contain distinct physicochemical environments defined by the collective behavior of their components [1, 2]. Phase separation has emerged as a leading mechanism to account for the formation of biomolecular condensates. In this thermodynamic process, segregative (density-driven) and associative (connectivity-driven) transitions are coupled to produce protein- and nucleic acid-rich phases suspended in the cellular cytoplasm or nucleoplasm [3]. Within this framework, intrinsically disordered proteins/regions (IDP/IDRs) have emerged as strong contributors to the multivalency and phase separation capacity of many naturally occurring proteins [1, 4–12]. Moreover, IDRs characterized by amino acid sequences of low complexity and enriched in aromatic residues, such as the prion-like low-complexity domains (PLDs) of RNA-binding proteins [8, 13–16], are prone to form condensates that behave like viscous fluids. Furthermore, the PLDs of various naturally occurring proteins, such as Fused in Sarcoma (FUS), Trans-activation response DNA-binding protein 43 (TDP-43), and the heterogeneous nuclear ribonucleoprotein A1 (hnRNPA1) have been reported to transition from liquid-like condensates into dynamically arrested solids and glasses over time [17, 18]. Such arrested states have been postulated as precursors of pathological protein aggregates and amyloids [19]. Understanding the molecular factors that regulate the ability of PLDs to phase separate is of crucial importance, as these are conjectured to also facilitate their aggregation [20].

Biomolecular condensates are networked fluids sustained by multivalent macromolecules that interconnect with more than two partners simultaneously [1, 2, 21–25]. Thus, the factors that increase the molecular valency, give rise to more densely connected liquid networks, and more stable condensates [1, 23, 26–29]. For PLDs, multivalency has been shown to be strongly promoted by the presence of aromatic residues, which act as “stickers” that enable associative π – π interactions, and also cation– π contacts with arginine and lysine [8, 9, 30–36]. Indeed, the range of coexistence of PLD condensates has been shown to change most significantly in response to mutations involving aromatic residues [9, 37]. Quantitative experimental phase diagrams for a set of hnRNPA1-IDR variants in combination with mean-field simulations demonstrated that tyrosine drives stronger associative interactions than phenylalanine, arginine is very context-specific, and lysine destabilizes PLD–PLD interactions [33]. More importantly, such study demonstrates that, at least for hnRNPA1, there exist quantitative rules that explain the changes in thermodynamic parameters of condensates in response to precise amino-acid sequence mutations.

In this work, we investigate whether there are scaling laws that can predict how the critical parameters of a broader collection of PLDs change in response to amino acid sequence mutations. Scaling laws are mathematical relationships that describe how the properties of a system change for a given variable, such as its size or composition [38]. In the context of polymer-like systems such as IDPs, scaling laws have offered invaluable insight into vastly different aspects of their physical behavior: for example, they were instrumental in determining that the relationship between viscosity and radius of gyration in polymer solutions is described by a power law between the two quantities [39]; demonstrating that aging scaling during polymer collapse is a universal property in homopolymers [40]; determining that in single polymer systems, their end-to-end distance is directly proportional to their length [41]; showing that foldable sequences with more compact unfolded states are recent products of protein evolution [42]. In the context of protein condensates, we hypothesized that scaling laws, if found, may allow us to quantify how condensates respond to different intrinsic perturbations—such as amino acid mutations or presence/absence of different post-translational modifications, changes in protein length, increased/decreased flexibility—or external stimuli—such as changes to the pH of the

medium or the ionic environment. Thus, we focused on determining if such laws exist for PLDs by exploring six different RNA-binding proteins (i.e., hnRNPA1, TDP43, FUS, EWSR1, RBM14, and TIA1; **Fig. 1** [A,B](#)), linked to the etiology of neurodegenerative disorders through pathology and genetics [[43](#)–[53](#)].

Leveraging the accuracy and efficiency of a modern sequence-dependent coarse-grained model [[32](#)], we quantify the phase diagrams (temperature-vs-density plane) of large set of mutants spanning all six PLDs mentioned above. We performed Direct Coexistence molecular dynamics (MD) simulations (**Fig. 1** [C](#)), exploring simultaneously the protein-rich and protein-depleted phases of the condensate for each mutant, in the temperature–concentration phase space. Our massive set of simulations encompasses 140 PLD mutants, each sampled at 8 to 12 different temperatures. Accounting for independent replicas to ensure proper sampling and convergence, the dataset presented here comprises the results of 2000 independent Direct Coexistence simulations (**Fig. 1** [D](#)), totaling 1.2 milliseconds of accumulated sampling.

Our work puts forward a data-driven scaling-law framework for the study of PLD phase behavior. We reveal that the changes in the critical solution temperatures of PLDs, as a function of the number and type of amino acid sequence mutations (e.g., aromatic, positively charged, etc), can indeed be quantified by scaling laws, which are highly conserved across the family of PLDs we investigate. The addition of aromatic residues yields a positive linear correlation between mutation fraction and critical temperature changes, consistent with the dominant role of aromatic residues in PLD phase separation. We find that protein length dictates scaling effects when aromatic residues are mutated into uncharged, non-aromatic amino acids. We also quantify the effects of arginine mutations on condensate stability, where removing arginines decreases the critical temperature of the condensate—revealing an inverse relationship between the fraction of arginine deletions and critical temperatures. Finally, polar amino acids emerge as subtle modulators of the phase behavior of PLDs. By examining the impact of different amino acid substitutions on the critical solution temperature, our results provide insights into the underlying molecular mechanisms of protein condensation. The scaling laws presented in this study serve two purposes: firstly, to verify the generality of the molecular grammar that governs the phase behavior across various PLD proteins, and secondly, to quantify the extent to which the critical solution temperature of a PLD protein solution will change in response to specific amino acid sequence changes. We anticipate that scaling laws for other families of proteins with very different compositional biases (e.g., transcription factors, nucleoporins, or molecular chaperones) may exist, but could be distinct from those we report here for PLDs. Overall, our work suggests that scaling laws represent useful predictors of the phase behavior of protein families.

Results

Generating and Analyzing an Extensive Dataset of Critical Solution Temperatures for Prion-like Domain Variants

The phase behavior of PLDs and their modulation with amino acid sequence mutations has been extensively investigated with a variety of experimental techniques and with theory and simulation methods for systems including the hnRNPA1 PLD [[33](#), [54](#)], the FUS PLD [[8](#), [49](#), [51](#), [55](#)–[63](#)], and the TDP-43 PLD [[45](#), [46](#), [50](#), [64](#), [65](#)], among many others. Here, we perform residue-resolution coarse-grained Molecular Dynamics (MD) simulations (see **Fig. 1** [C](#), with further details in Methods) with our Mpipi model [[32](#)] to quantitatively compare the critical parameters of a total of 140 mutants of the PLDs of TDP-43, FUS, EWSR1, hnRNPA1, RBM14, and TIA1 (**Fig. 1** [A](#)) under consistent solution conditions. From our simulations, we compute temperature-vs-density phase diagrams and estimate the critical solution temperature for each system. The PLDs of the proteins we investigate have similar composition biases—i.e., are highly rich in neutral, polar, and aromatic residues—and present less than 5% of total charged amino




Figure 1.

Data generation and representation of dataset for probing scaling laws of PLD phase behavior

(A) Representation of the prion-like domains (grey-shaded regions) of each considered protein, alongside the associated number of variants simulated, totaling a set of 140 sequences. **(B)** Composition of the wild-type sequence of the PLDs simulated, in terms of the percentage of neutral amino acids (A, C, I, L, M, P, S, T, and V; no net charge at pH 7 and no π electrons in the side chain), glycine (G), negative (D, E), positive (K, H, R), neutral- π (N and Q; no net charge at pH 7 with π electrons in the side chain) and aromatic (F, W, Y) residues. **(C)** Top panel: Representation of the simulation model: each amino acid of the protein is represented by a single bead, bonded interactions are modeled with springs, and non-bonded interactions are modeled with a combination of the Wang-Frenkel potential and a Coulomb term with Debye-Hückel screening. Middle panel: The Direct Coexistence simulation method, in the slab geometry, is employed, where both the protein-rich and protein-depleted phases are simulated simultaneously. Bottom panel: Finally, to obtain the data point in the concentration-temperature phase space, at a given temperature the density profile is computed and the concentrations of the rich and depleted phases are obtained via a suitable fitting (Methods). **(D)** Representation of the entire computational data set. Orange data points represent variants where charged or aromatic residues were mutated (i.e., stickers), while cyan data points represent all other types of mutations studied (i.e., spacers).

acids in their sequence (**Fig. 1** [B](#)). Thus, our simulation results allow us to assess if there exists a code—i.e., a simple set of scaling laws—that can predict how the critical parameters of the family of PLDs will scale in response to amino acid sequence mutations.

Our residue-resolution model Mpipi (see Methods) is ideally suited to investigate the phase behavior of PLDs because it achieves the correct balance of the dominant π - π and cation- π interactions, which drive the phase behavior of PLDs. Indeed, the Mpipi model recapitulates the quantitative experimental phase diagrams of hnRNPA1 variants by Bremer and colleagues [33] with remarkable accuracy [32]. It is worth noting that the Mpipi model was not parameterized based on the experimental dataset of Bremer and colleagues, but rather the model parameters were developed independently by combining atomistic umbrella sampling MD simulations of amino acid pairs with the bioinformatics data of Vernon and colleagues for π -driven contacts [35]. Using Flory–Huggins–Staverman theory, critical temperatures for phase separation were extracted from the experimental data of Bremer et al., which were then used to validate the Mpipi model [32].

The amino acid sequences of all the proteins we investigated (Supporting Information) were designed based on five main considerations. (1) The set of mutants should serve to probe the effects of both the strong modulators of PLD phase separation—namely, aromatic and arginine interactions—and the subtle modulators—that is, polar and neutral amino acids, previously identified experimentally [8, 9, 33]. (2) To ensure that the observed changes in the critical solution temperatures can be attributed unequivocally to specific amino-acid-type changes, we mutated only one amino acid type at a time. However, multiple mutations of the same amino acid type were allowed. (3) Mutations were distributed homogeneously across the entire protein. (4) Deleted amino acids (e.g. aromatic deletions) were replaced by weakly interacting amino acids (i.e. serines, threonines, glycines, and alanines). For each PLD protein, the identity of the weakly interacting residues used as replacements were chosen to preserve the percentages of the various types of weakly interacting amino acids present originally in the unmodified PLD. (5) Achieve a data set that spans a wide array of critical solution temperatures, as shown in **Fig. 1 D**.

Validation of the Mpipi Model for Phase behavior of Prion-like Domains

The availability of quantitative experimental binodals for the hnRNPA1-IDR and various amino acid sequence variants [33] has allowed us to test the accuracy of the Mpipi model to describe the phase diagrams of PLD solutions [32]. Here, we make such a test more stringent by extending the hnRNPA1-IDR validation set from 9 PLD variants to 21, selecting mutations spanning changes in aromatic, charged, polar, and neutral residues. The amino acid sequences of all the hnRNPA1-IDR variants considered are described in the Supporting Information and are denoted using the intuitive nomenclature established by Bremer and colleagues: variants that mutate n residues of type X into m residues of type Z are labeled as $-nX + mZ$, and so forth. For more details on the nomenclature, see Methods.

For each hnRNPA1-IDR variant, we perform Direct Coexistence MD simulations, compute critical temperatures from the simulations, and compare them with the experimental critical temperatures (**Fig. 2 A–D**). We use the fitting methodology outlined in Methods to estimate the critical solution temperatures from the experimental data.

When comparing with experimental critical temperatures, our model achieves a Pearson correlation coefficient higher than 0.93 for all types of mutations analyzed. Additionally, for all cases, the root-mean-square deviation (**D**) between the experimental and simulated critical temperatures is ≤ 14 Kelvin. Our validation set, which expands the range of proteins variants originally tested [32], highlights that the Mpipi potential can effectively capture the thermodynamic behavior of a wide range of hnRNPA1-PLD variants, and suggests that Mpipi is




Figure 2.

Mpipi model quantitatively captures phase behavior for a large set of hnRNPA1 PLD mutants.

Mutations are divided into A aromatic, B Arg/Lys, C charged, and D Gly/Ser sequence variants of the PLD of the protein hnRNPA1, and comparison with experimental results. The hnRNPA1 PLD was used to validate the accuracy of the model, by extracting critical temperatures from experimental phase diagrams of a large set of these variants [33]. To assess how well the Mpipi model performed for PLDs, we compared critical temperatures obtained from simulations with the corresponding ones extracted from the experimental data for those variants that were available (depicted in the top right of each binodal, where r is the Pearson coefficient and D is the root mean square deviation between simulated and experimental values). For the variants where the experimental critical temperature could not be determined, we make a comparison between the critical temperature with the saturation concentration, since for these systems we expect them to be inversely correlated. The naming system for the variants is discussed in detail in the text.

adequate for proteins with similar sequence compositions, as in the set of proteins analyzed in this study. In recent work by others [66], Mpipi was tested against experimental radius of gyration data for 137 disordered proteins and the model produced highly accurate results, which further suggests the applicability of the approach to a broad range of sequences.

Large-Scale Data Generation on Critical Parameters of PLD Phase Separation

The next step of the study consists of generating a large amount of data, needed to report the critical temperatures of the PLD condensates. In the Direct Coexistence method, the diluted phase and the condensate are simulated in the same rectangular box separated by an interface. The phase diagrams we compute explore the plane of temperature (vertical axis) versus density (horizontal axis). For each phase diagram, we run a set of about 20 Direct Coexistence simulations—each simulation explores the phase behavior of the system at different constant temperatures. For a given value of the temperature, if two phases are detected in our simulations, we measure a “density profile”: the density of the proteins as a function of the long side of the simulation box (Bottom panel in Fig. 1 C). We then plot the two coexistence densities for each temperature we simulate and build the coexistence curve (also known as binodal), Fig. 3. The coexistence curve shows the values of temperature and protein density for which phase separation will occur. The region above the coexistence curve is the “one-phase region” where the temperature is too high for the enthalpic gain coming from the associative protein–protein interactions to overcome the entropic loss upon phase separation. The region below the coexistence curve, the “two-phase coexistence region” represents lower temperatures for which the enthalpic gain due to protein–protein interactions is now sufficiently high to favor phase separation into a condensed (protein-enriched) and a diluted (protein-depleted) liquid phase. The maximum in the coexistence curve is known as the critical point: the upper critical solution temperature (herein, critical solution temperature) and corresponding critical density. For temperatures higher than the critical solution temperature, phase separation is no longer observed.

The protein solutions that we investigate with our coarse-grained simulations must be sufficiently large, in terms of the number of copies of the same protein, to minimize finite size effects—for instance, to adequately sample bulk phenomena without dominating interfacial effects. At the same time, our simulations must be small enough to ensure computational convergence in a feasible timescale. Following extensive testing of computational convergence for systems of different sizes and MD trajectories of different time lengths, we determined that a system composed of 64 protein replicas, with trajectories of 200 ns ensured rigorous convergence and reproducible results, while taking into careful consideration the trade-off between accuracy and computational demands. The size of the simulation boxes were also optimized, again to ensure finite size effects did not affect the predictions (See Methods and Materials, and Supporting Information).

As a result, we generated a dataset of critical parameters of over an order of magnitude higher than previously reported studies [9, 33, 67–70]. This extensive dataset (Fig. 3) not only offers new insights into phase separation phenomena but also sets a new benchmark for data-driven research in this field.

A Data-Driven Scaling Law Approach for Studying Phase Behavior of Prion-like Domains

Scaling laws can reveal key relationships between system variables, such as composition and critical parameters, providing a useful theory to predict and interpret the regulation of complex biological phenomena such as phase separation. In the context of PLD phase behavior, scaling laws would provide a quantitative theory to analyze how alterations in amino acid composition affect the critical solution parameters—a crucial aspect for understanding protein behavior in




Figure 3.

Complete set of 140 binodals, in the temperature versus density phase space, generated via largscale molecular dynamics simulations.

Binodals are grouped according to the mutation type (Aromatic, Arg/Lys, Gln/Asn (Polar), Ala/Ser and Thr/Gly/Ser (Neutral)). Each set is further grouped based on the PLD in question.

biological systems. Here, our objective is to discern which scaling laws can effectively capture the variations of PLD critical parameters as a function of sequence mutations. To accomplish this, we employ a systematic approach of varying the amino acid sequence of the proteins considered, to determine how the phase behavior of PLDs changes with amino acid compositional perturbations.

For each type of mutation we analyze, we define and test three types of scaling laws. These scaling laws illustrate a different relationship between critical solution temperature changes and the number of amino acid mutations. The first and simplest scaling law is a linear relationship between these two quantities,

$$\Delta T_c = S_{X_1 \rightarrow X_2} N_{X_1 \rightarrow X_2}, \quad (1)$$

where ΔT_c is the variation on the critical temperature between the mutated protein and the corresponding unmodified reference sequence, X_i is an amino acid (e.g., Y, F, R, and so forth) and $N_{X_1 \rightarrow X_2}$ is the number of X_1 residues that have been mutated to X_2 . Thus, $S_{X_1 \rightarrow X_2}$ in this case quantifies the variation in the critical solution temperature of the protein solution for a mutation of one X_1 residue to X_2 . If the data is well represented by this first scaling law, it would imply that the effects of amino acid mutations on PLD phase behavior depend exclusively on the total number of mutations, but are not affected by other PLD parameters, such as the protein length. Thus, the first scaling law (Eq. 1) is the simplest possible model we test here for how incremental mutations may collectively influence the phase behavior of PLDs.

Next, we define a second type of scaling law, which hypothesizes that the changes to the critical solution temperature upon mutation is proportional to the fraction of residues that are mutated, rather than their raw number. That is,

$$\Delta T_c = S_{X_1 \rightarrow X_2} \frac{N_{X_1 \rightarrow X_2}}{L} = S_{X_1 \rightarrow X_2} f_{X_1 \rightarrow X_2}, \quad (2)$$

where, L represents the number of total amino acids in the PLD, or protein length, and $f_{X_1 \rightarrow X_2} = \frac{N_{X_1 \rightarrow X_2}}{L}$ is the fraction of X residues that have been mutated to X_1 . Thus, this second scaling law hypothesizes that the effect of mutations on the critical solution temperature is more pronounced in shorter proteins (where $N_{X_1 \rightarrow X_2}$ mutations constitute a larger fraction of the protein) and less so in longer proteins (where $N_{X_1 \rightarrow X_2}$ mutations represent a smaller fraction of the total number of amino acids). The expectation is that because longer PLDs have higher valencies than shorter ones, the density of inter-molecular connections they form inside condensates is more robust towards perturbations than that of shorter PLDs [71].

Finally, we define a third scaling law,

$$\Delta T_c = S_{X_1 \rightarrow X_2} \frac{N_{X_1 \rightarrow X_2}}{\sqrt{L}} \quad (3)$$

where the change in critical solution temperature is hypothesized to be directly proportional to the number of mutations ($N_{X_1 \rightarrow X_2}$) but inversely linear to $\sqrt{L}$. Like our second scaling law, this third relationship also presumes that longer PLDs form condensates with critical parameters that are less affected by mutations than shorter PLDs. However, by considering $\sqrt{L}$, instead of L as in Eq. 2, the expectation is that the buffering effects of protein length are more modest than those stated by Eq. 2. Notably, the amino acid composition of PLDs can be understood by the stickers-and-spacers framework [72, 73]. Aromatic and arginine residues have been proposed as stickers, contributing to the stability of the liquid network via associative interactions, yet some groups argue that all residues matter equally. Our data suggests that while a strict black-and-white delineation might not be appropriate, certain residues such as Tyr or Arg demonstrably have a

larger impact on T_c when mutated. Conversely, other residues exhibit smaller effects, acting more like spacers interspersed among the stickers. Eq. 3 suggests that as a protein lengthens, the number of residues that can establish sufficiently strong inter-protein interactions grows too; yet, it grows much more conservatively than L because a fraction of L is made of spacers.

With these scaling laws as ansatz, we set to uncover trends in the dataset we built from our MD simulations of 140 variants of different PLDs. Adopting these laws as foundational hypotheses, we explore the relationship between alterations in composition and changes in critical solution temperatures, offering insights into the molecular drivers of PLD phase separation.

Aromatic Mutations Exhibit a Linear Correlation Between Mutation Fraction and Critical Solution Temperatures

Aromatic amino acids, namely tyrosine (Tyr, Y), phenylalanine (Phe, F), and tryptophan (Trp, W), are widely regarded as essential contributors to phase separation of PLDs, due to their ability to form π - π and cation- π contacts [9, 30, 31, 33, 35, 36, 74–76]. Among aromatics, experiments and simulations agree that Tyr is a stronger contributor to the stability of condensates than Phe [8, 31, 33, 60], and that both the composition and the patterning of aromatic residues in PLDs have an impact on the condensate critical parameters and its material state [18, 68, 77, 78].

Aromatic residues are highly abundant in PLDs, with the fraction of aromatic residues in the PLD variants analyzed in this study ranging from 0.19 to 0.43 (Fig. 1B). Furthermore, because different aromatics are unequal contributors to PLD phase behavior, we analyzed the impact of mutations for Tyr and Phe and Trp separately.

To quantify the impact of mutating Tyr to Phe in shifting the critical solution temperatures of our set of PLDs—i.e., determining which of our proposed scaling laws best describes the data—we designed 17 protein mutants that varied the Tyr to Phe ratio in the family of PLDs and computed their phase diagrams (Fig. 4A). Our simulations reveal that Tyr is stronger than Phe as a driving force for phase separation for the entire PLD family considered, a finding that replicates the results of Bremer *et al.* [33] for hnRNPA1 and confirms that it extends to other PLD proteins. By analyzing the change of critical solution temperature with respect to the unmodified PLD (or wild type, WT) against the number of mutations, our results reveal a linear relationship between the number of Tyr to Phe mutations and the change in the critical temperature of the system; our first scaling law, $S_{Y \rightarrow F}$, Eq. 4. Here, ΔT_c represents the temperature difference between the mutated protein and its wild-type sequence, and $N_{Y \rightarrow F}$ represents the number of Tyr residues that were mutated to Phe in the sequence.

$$S_{Y \rightarrow F} = \frac{\Delta T_c}{N_{Y \rightarrow F}} = -0.40 \pm 0.04 \quad (4)$$

Note that, by definition $S_{Y \rightarrow F} = -S_{F \rightarrow Y}$, since $N_{Y \rightarrow F}$ is defined as positive for Tyr to Phe mutations and negative for Phe to Tyr mutations, or, in other words, $N_{Y \rightarrow F} = -N_{F \rightarrow Y}$. For our data set, we find $S_{Y \rightarrow F}$ to be -0.40 ± 0.04 , where the error given is the standard error. The way to interpret this result is that a single mutation of a Tyr residue to a Phe residue in a given PLD protein would result in a -0.4 K variation in the critical solution temperature for phase separation. Such a modest variation due to a single-point mutation is smaller than the typical error associated with estimating the absolute value of the critical solution temperature itself using residue-resolution coarse-grained Direct Coexistence simulations (approximately 3 K). Unambiguously measuring the value of the critical solution temperature with sub-Kelvin accuracy poses a considerable challenge also in experiments [79, 80]. However, we can uncover statistically significant trends by analyzing the behavior of the critical solution temperature in a large set of PLDs with diverse

numbers of mutations rather than a single case. In particular, our analysis reveals that the impact of Tyr to Phe mutations is modest and the effect of multiple mutations can be described adequately as cumulative.

In addition to these analyses, the role of Tyr/Phe to Trp mutations, albeit Trp being a rare residue in prion domains, was investigated in TIA1 due to the presence of Trp in the wild type of this PLD, **Fig. 4B**. The simplest scaling law that describes this mutation (**Equation 5**) is fundamentally equal to the law describing Tyr to Phe mutations. However, a notable distinction is that the $S_{F/Y \rightarrow W}$ scaling constant for our data is approximately 4.3 ± 0.1 , which is one order of magnitude greater than $S_{Y \rightarrow F}$, consistent with the much stronger π - π contacts established by Trp. It is important to note that, in this case, similarly to Try to Phe mutations, multiple Ansatz align closely with the observed data, which underscores the fact that, in the context of these mutations, the size of the side chains does not significantly alter phase behavior, as the predominance of aromatic perturbations exerts a dominating effect.

$$S_{F/Y \rightarrow W} = \frac{\Delta T_c}{N_{F/Y \rightarrow W}} = 4.3 \pm 0.1 \quad (5)$$

Across the whole family of PLDs analyzed, our data reveals that changes in aromatic amino acid identity impact the critical solution temperatures modestly (< 1 K per mutations involving F and Y) to strongly (4 K for mutations involving Trp) and independently of the protein length. Such a result is consistent with the mutations in aromatic identity preserving the total number of associative interactions that the PLDs can establish, and hence, the density of molecular connections in the condensed liquid network. That is, aromatic mutations are expected to influence the energetics and lifetimes of the multivalent interactions among PLDs, but not the number of intermolecular contacts they form.

Strong Critical Temperature Reduction Due To Aromatic Deletions is Buffered By Protein Length

We next examine the role of replacing aromatic residues (Tyr and Phe) with uncharged, non-aromatic residues—such as Gly, Ser, Gln, Ala, and Asn—which are prevalent in PLDs. The mutations were designed to preserve amino acid sequence compositional biases by replacing Tyr and Phe with uncharged, non-aromatic residues in the same proportions as they appear in the wild type of each sequence, i.e., maintaining the relative amounts of each type of residue found in the sequence. All amino acid sequences are available in the Supporting Information.

Our simulations show that aromatic deletions strongly reduce the critical solution temperature of the PLD solutions, **Fig. 4C**. The best scaling law from our ansatz to describe the data is **Eq. 3**, which involves the square root of the PLD length. That is, $S_{Y/F \rightarrow X}$, where X represents any uncharged, non-aromatic residue.

$$S_{Y/F \rightarrow X} = \frac{\Delta T_c \sqrt{L}}{N_{Y/F \rightarrow X}} = -56 \pm 5 \quad (6)$$

Here, once again, $N_{Y/F \rightarrow X}$ is the number of mutations, and L is the length of the PLD. For our data set, the value of the scaling constant $S_{Y/F \rightarrow X}$ is -56 ± 5 . To interpret this result, let us consider a model PLD of two hundred amino acids ($L = 200$). If we mutate one Tyr to an uncharged, non-aromatic amino acid, the change in the critical temperature—computed via the formula $S_{Y/F \rightarrow X} \times \frac{N_{Y \rightarrow X}}{\sqrt{L}}$ —is approximately -4 K. Such a value is consistent with the dominance of π - π and cation- π contacts in PLD phase separation reported experimentally [9, 30, 31, 33, 35, 36, 74–76]. As stated above, aromatic residues behave as associate hubs within PLDs condensates. When these aromatic residues are replaced with uncharged non-aromatic amino acids the density of connections network of molecular connections in the liquid is strongly




Figure 4.

Mutations in aromatic amino acids have strong effects on the critical solution temperature of PLDs.

A. Tyr to Phe mutations. The number of Tyr mutated to Phe (x-axis) versus the change in critical temperature, (y-axis), computed as the critical temperature of the variant minus that of the corresponding PLD wild-type sequence. The trendline defines the simplest, dominating scaling law for this mutation. The R^2 value of 0.88 shows the agreement of our data to the scaling law defined by Equation 4. **B.** Tyr or Phe to Trp mutations. In this case, the overall trendline, defined by equation 5 has an R^2 of 0.99. **C.** Analysis of the variants involving mutations of aromatic residues to uncharged, non-aromatic amino acids, a.k.a. aromatic deletions. In the y-axis, the change in the critical temperature of the variant to that of the wild type sequence, and in the x-axis, a renormalized measure of mutations: the fraction of aromatic residues mutated times the $\sqrt{L}$, to account for the competing physical interactions between PLDs. Error bars indicate the standard error associated with replicas of the simulation.

perturbed, as we have shown earlier [81]. To observe this in more detail, we computed select contact maps (see **Fig. 2**) – notably, mutations that removed aromatic amino acids strongly altered the primary sites of interactions.

Our results reveal that the length of the PLDs strongly influences how the critical parameters of PLD solutions change in response to aromatic deletions. The effect of mutations is stronger in shorter proteins because each aromatic deletion represents a higher percentage of the total number of associative residues in the PLD, than in longer proteins. Longer proteins have higher valencies and redundant “stickers” that facilitate a network rearrangement.

Our results highlight the delicate balance between amino acid composition and protein length in determining the phase behavior of PLDs, providing a deeper understanding of the molecular basis of protein phase transitions.

Arginine Mutations Drastically Alter Condensate Stability

The distribution and balance of charged amino acids—i.e., aspartic acid (Asp, D), glutamic acid (Glu, E), histidine (His, H), arginine (Arg, R), and lysine (Lys, K)—within an IDR affects its solubility, conformational ensemble, and the range of stability of the ensuing condensates it might form. PLDs are not particularly rich in charged residues: in the wild-type PLDs of our data set, the fraction of charged residue ranges from 0.02 to 0.1 (**Fig. 1B**), and the fraction of positively charged amino acids is narrowly distributed around 0.015, except for hnRNPA1, which has a value of 0.09 (**Fig. 1B**). However, despite their low prevalence in PLDs, charged residues, in particular Arg, have been shown to be important contributors to the stability of a wide-range of biomolecular condensates [30, 33, 82–84].

While both Arg and Lys are positively charged amino acids at physiological pH, it has been demonstrated that they have unequal roles in the stability of biomolecular condensates [30, 33, 60, 83, 85]. Both residues can form charge–charge and cation– π associative interactions, but the case of Arg is unique: the sp^2 -hybridized planar guanidinium group of Arg confers it a special capacity to form strong hybrid π – π /cation– π contacts with the π -electrons of aromatic rings [31]. Thus, to quantify the impact of Arg in the range of stability of PLD condensates, we designed mutants that replaced Arg with Lys and vice-versa (**Fig. 5**).

The best fitting scaling law associated with mutating Arg to Lys is inversely dependent on the fraction of mutations, rather than the number of residues mutated (**Eq. 7**). The resulting scaling constant for this particular mutation, i.e., $S_{R \rightarrow K}$, has a value of -1300 ± 100 , indicating that for a typical PLD of $L = 200$, the mutation of a single Arg into Lys results in a reduction of the critical temperature of the system by 6.5 K. Hence, replacing Arg with Lys is a destabilizing mutation of PLD condensates in agreement with the wide-body of experimental evidence [30, 33, 60, 83, 85].

$$S_{R \rightarrow K} = \frac{\Delta T_c}{f_{R \rightarrow K}} = -1300 \pm 100 \quad (7)$$

The length dependency of this particular scaling law suggests that the impact of Arg to Lys mutations on the phase behavior of PLDs primarily depends on the relative concentration of these mutations (mutation fraction). Additionally, it proposes that longer sequences may exhibit increased resilience against Arg to Lys mutations. This resilience is likely attributed to the greater separation of charge within the longer sequences.

To further analyze the effect of the charge of the PLDs in this set of mutants, in **Fig. 5** we include a scalebar showing the net charge per residue (NCPR). Mutants that present higher NCPR present the biggest decrease in their phase-separating propensities, which is consistent with the




Figure 5.

Critical effects of Arg mutations in the phase transition temperature of the condensates

(A) Analysis of the critical solution temperature of PLDs with Arg to Lys mutations. The number of Arg to Lys mutations divided by PLD length (x-axis) versus the change in the critical solution temperature, that is, $T_c(\text{variant}) - T_c(\text{WT})$ (y-axis). The trendline, representing **Equation 7**, fits the data with an $R^2 = 0.94$, and defines the stability measure of this perturbation.

(B) Analysis of the critical solution temperature of PLDs with Arg mutations to uncharged, non-aromatic amino acids that maintain the WT compositional percentages (mainly Gly, Ser, and Ala. For more details see Supporting Information). Arg deletion fraction (x-axis) versus $T_c(\text{variant}) - T_c(\text{WT})$ (y-axis). The trendline, which fits the data with $R^2 = 0.96$, defines the stability measure, represented in **Equation 8**.

expectation that condensates with a neutral overall charge are more energetically stable [33, 86, 87].

When we replace Arg with uncharged, non-aromatic residues, the resulting scaling law also depends inversely on the fraction of mutations, with a value of -640 ± 50 , amounting to a reduction of 3.2 K in T_c per Arg deletion for a model PLD of length $L = 200$.

$$S_{R \rightarrow X} = \frac{\Delta T_c}{f_{R \rightarrow X}} = -640 \pm 50 \quad (8)$$

Therefore, our results reveal that it is almost twice more destabilizing to mutate Arg to Lys than to replace Arg with any uncharged, non-aromatic amino acid, namely, Gly, Ser or Ala. Mutating Arg to an uncharged non-aromatic amino acid is less destabilizing than mutating it to Lys likely because the charge-reduced variants decrease the cation–cation repulsion of the PLDs, with respect to the Arg-rich or Lys-rich systems. Furthermore, mutating Arg to Lys decreases the enthalpic gain for condensate formation not only via the replacement of stronger Arg-based associative interactions for weaker Lys-based ones but also by increasing the overall solubility of the Lys-rich PLD [36, 88], given that the hydration free energy of Arg is less favorable than that of Lys [31, 36, 82].

Polar Amino Acids Subtly Modulate PLD Phase behavior

Uncharged, non-aromatic amino acids, including Gly, Ala, proline (P, Pro), Ser, and Thr, are commonly found in PLDs. Because they generally form weaker residue–residue interactions and present smaller excluded volumes than aromatics, they are classified as spacers that contribute to maintaining the solubility of proteins and material states of condensates [68, 72]. These amino acids can also have other functions in proteins, and the role they play as spacers depends on their chemical make-up, but also on the specific context in which they are found [33, 37, 89].

Here, we investigate the role of glutamine (Gln, Q) and asparagine (Asp, N). Both Gln and Asn are polar amino acids, hence their charge distribution is more asymmetric, presenting a significant dipole moment. Consequently, they can establish transient ion–dipole, dipole–dipole and π –dipole interactions with other residues. While the Mpipi model does not account explicitly for polarity, it is worth noting that it was parametrized based on bioinformatics data, which implicitly incorporates this aspect. Thus, our study allows for the observation of emergent behaviors that stem from this inherent characteristic.

We designed mutations that vary the relative amounts of Gln and Asn, and examined trends in their phase-separating propensity. Increasing the number of Gln per unit length by mutating Asn to Gln increases the ability of the system to phase separate, likely due to Gln having a higher dipole moment than Asn, which is approximated in our model via marginally stronger non-ionic interactions formed by Gln over Asn (Fig. 6A).

$$S_{N \rightarrow Q} = \frac{\Delta T_c}{f_{N \rightarrow Q}} = 50 \pm 5 \quad (9)$$

The value of the scaling constant $S_{N \rightarrow Q}$ is 50 ± 5 , which is significantly lower than that of mutations of residues with a net charge or aromatic deletions (rounding 1300, for example, for arginine to lysine mutations). This result indicates that mutations of Asn to Gln or vice versa have a smaller impact in the critical parameters than aromatic deletions or mutations of charged residues.

Finally, we also explored the role of Ser, Gly, Thr, and Ala on the critical parameters of the condensate. While Ser and Thr are also polar amino acids, their dipole moment is significantly smaller than that of Gln and Asp. Consistently, the scaling constants involving mutations Ser and




Figure 6.

Polar and neutral amino acid mutations show subtle modulation of the critical solution temperatures of PLDs
A) Analysis of the critical solution temperature of PLDs with Asn to Gln mutations.

The number of Gln to Asn mutations renormalized by dividing by PLD length (x-axis) versus the temperature change $T_c(\text{variant}) - T_c(\text{WT})$ (y-axis). The trendline, which fits the data with an $R^2 = 0.84$, fits the scaling law **Equation 9**. (B) Analysis of the critical solution temperature of PLDs with Gly/Ser to Thr mutations (x-axis) versus the temperature change $T_c(\text{variant}) - T_c(\text{WT})$ (y-axis). The trendlines, representing **Equations 10** and 11 fit the data with an $R^2 = 0.7$. (C) The number of Ala to Ser mutations (x-axis) versus the temperature change $T_c(\text{variant}) - T_c(\text{WT})$. The trendline, **Equation 12**, fits the data with an $R^2 = 0.77$

Thr mutations—i.e., $S_{S \rightarrow T}$ —are considerably smaller (**Fig. 6** B and C) overall. A comparable pattern is noted for $S_{G \rightarrow T}$, with G expected to exhibit significantly lower polarity compared to Gln and Asp.

$$S_{S \rightarrow T} = \frac{\Delta T_c}{N_{S \rightarrow T}} = -0.09 \pm 0.03 \quad (10)$$

$$S_{G \rightarrow T} = \frac{\Delta T_c}{N_{G \rightarrow T}} = -0.08 \pm 0.02 \quad (11)$$

Regarding Ala, we also tested its relative role with respect to Ser.

$$S_{A \rightarrow S} = \frac{\Delta T_c}{N_{A \rightarrow S}} = 0.16 \pm 0.05 \quad (12)$$

Amino acids with larger side chains, such as Thr, have a larger excluded volume than those with smaller side chains, such as Gly or Ser. The corresponding value of the scaling constants $S_{S \rightarrow T}$ and $S_{G \rightarrow T}$ suggests that in the case of Thr mutations, mutations of neutral, non-aromatic amino acids to those with larger excluded volumes inhibit phase separation of the PLD by lowering the corresponding critical temperature, while mutations to those amino acids with smaller excluded volumes have an opposite effect. This is likely due to increased flexibility of the protein backbone, allowing the chain to adopt multiple conformations that promote multivalent contacts. Importantly, these trends cannot be anticipated by inspecting the parameters of the Mpipi model; instead, they emerge from examining the behavior of the collective system.

On the other hand, for mutations involving Ser, Gly, and Ala—all small amino acids with minor side chain size variations—the impact on phase separation seems to stem predominantly from changes in solvation. Solvation plays a key role in phase separation since it can help balance the driving forces of this process [90, 91]. While hydrophobic regions of the protein, with low solvation free energies, tend to form aggregates to reduce interactions with the solvent environment; hydrophilic regions promote interactions with water. Whether these regions inhibit or promote phase separation is likely due to a balance of opposing effects. On the one hand, hydrophilic regions can form energetically favorable contacts with solvent, promoting dispersion and suppressing phase separation. However, when these regions are recruited to condensates the release of water molecules can help to off-set the entropic cost of sequestering hydrophobic regions within a protein droplet, thereby facilitating phase separation.

Specifically, mutations from Gly to Ser (not shown) increase the size of the side chain while also enhancing solvation, while mutations from Ala to Ser do not significantly change the size of the side chain but significantly increase the protein solvation. In the first case, the critical parameters of phase separation remain virtually unchanged, only presenting a very minor decrease. In the case of Ala to Ser mutations, phase separation is slightly enhanced. Both scaling laws $S_{G \rightarrow S}$ and $S_{A \rightarrow S}$ reflect such an effect, and underlie the context-dependent nature of these mutations.

Overall, mutations of unchanged, non-aromatic amino acids seem to have an impact on phase separation due to two competing effects: a change in side chain size and a change in solvation. However, it is important to note that the magnitude of the scaling laws that govern these mutations is minimal, and thus, likely extremely sensitive to small fluctuations such as changes in the system and solution conditions. Notably, these findings are consistent with experiments that report small changes in phase separation propensity for mutations involving neutral, non-aromatic residues [8, 33].

Beyond composition: patterning considerations

Beyond composition, the relative positioning of amino acids in the sequence, i.e. sequence pattern, also plays a role in protein folding, function and, in this case, phase behavior [9, 86, 92, 93]. Due to the sequence heterogeneity of PLDs, and, more generally, IDPs, it is particularly complex to define adequate patterning measures [87, 94–96].

In this work, we have taken advantage of our large data set to analyze the impact of amino acid sequence patterning by adopting a patterning order parameter [9, 86, 97], σ_{aro} , that has been designed to describe patterning of aromatic residues in such IDPs.

To compute σ_{aro} , we divide the amino acid sequence into peptide segments of equal length, compute a measure of sequence aromatic asymmetry for each segment, and finally sum the values of such measure over all the segments. Consistent with literature [9, 86, 97], we take l to be 5 and 6. σ_{aro} is then given by:

$$\sigma_{\text{aro}} = \left(\sum_{\text{segments}} \frac{\sigma_i}{\sigma_{\text{max},i}} \right) \frac{N_{\text{stickers}}}{L}, \quad (13)$$

where L is the total number of amino acids in the protein domain, N_{stickers} the number of sticker residues—i.e. charged or aromatic residues—and σ_i has the functional form

$$\sigma_i = \left(\left(2 \frac{N_{\text{aro},i}}{l_i} - 1 \right)^2 - \sigma_T \right)^2. \quad (14)$$

Here, $N_{\text{aro},i}$ is the number of aromatic residues in the i -th segment, l_i the segment length, and σ_T is a function of the total number of aromatic residues in the sequence N_{aro} :

$$\sigma_T = \left(2 \frac{N_{\text{aro}}}{L} - 1 \right)^2 \quad (15)$$

In turn, $\sigma_{\text{max},i}$ is the measure of aromatic amino acid distribution for an artificial sequence with all the aromatic amino acids concentrated in part of the chain.

Notably, our results, presented in **Fig. 7**, indicate that the sequences with the most dispersed aromatic residues (i.e., low values of σ_{aro}) phase separate more readily. Dispersed residues result in greater effective valency of a PLD, leading to a higher probability of forming a percolated network structure to sustain the condensate. Additionally, this result demonstrates that the patterning order parameter, σ_{aro} , can be used to predict the phase-separating propensity of a PLDs as the function of aromatic residue distribution. This result is consistent with previous studies that demonstrate the importance of aromatic residue patterning on the phase behaviour of PLDs [9, 98].

Discussion

Computer simulations have advanced our ability to map emergent properties of protein solutions to the chemical make-up of their individual biomolecules. For instance, modern residue-resolution protein coarse-grained models [32, 74, 99–103] can now be used to quantify the modulation of protein phase behavior as a result of amino acid sequence mutations of their native proteins with near-quantitative agreement with experiments [32, 101, 102]. In this study, we exploit the accuracy and efficiency of our protein model, Mpipi [32], to compute and analyze the temperature-vs-density phase diagrams of a large set of PLD proteins—140 variants of the




Figure 7.

Prion-like domains featuring sequences with dispersed aromatic residues show greater propensities for phase separation.

A Representation of the aromatic amino acids present in the wild type sequences of each PLD considered B Location of the positively charged (red) and negatively charged (blue) amino acids present in the wild type sequences of each PLD considered C Analysis of the critical solution temperature versus the σ order parameter, defined in Equation 13. Those groups of variants with the same σ value show their average critical temperature highlighted, and softer values for other data points. For lower values of σ , the critical temperature present in the condensates is higher, and vice versa, indicating that those variants with a more homogeneous distribution of aromatic amino acids have condensates more stable at higher temperatures.

proteins hnRNPA1, TDP43, FUS, EWSR1, RBM14, and TIA1. Collectively, our set of phase diagrams reveal conserved scaling laws that quantitatively predict the change in the critical solution temperatures of single-component PLD solutions as a function of the number and type of amino acid mutations of the PLDs.

To facilitate the interpretation of the scaling laws derived in our study, we have organized the data in a table that illustrates the effect of single point mutations on the critical solution temperatures of the family of PLDs, as a function of the PLD length.

Consistent with experimental evidence and computational predictions, interactions involving aromatic residues—namely, π - π and cation- π —are dominant in stabilizing condensates formed by PLDs [8, 32, 33]. Notably, our simulations reveal the largest decrease in the critical solution temperature of a PLD condensate occurs when an aromatic residue is replaced by an uncharged, non-aromatic amino acid. Our scaling law for such aromatic deletions ($56 \text{ K}/\sqrt{L}$) predicts that the destabilizing effect of replacing an aromatic with a non-aromatic residue is linearly amplified by the number of such mutations and screened by the square root of the length of the protein ($\sqrt{L}$), indicating a disruption in the percolated network of interactions sustaining the condensate. Condensates made of longer PLDs are more robust against aromatic deletions because, by construction, each aromatic residue represents a smaller percentage of the total valency of the PLD in question. In other words, longer PLDs are higher valency molecules than their shorter counterparts and can, thus, form a densely connected percolating condensed-liquid network, even when a few of their aromatic stickers have been deleted.

Our findings also highlight the crucial role of Arg in the critical parameters of PLD condensates. Specifically, our scaling law for Arg deletions ($640 \text{ K}/L$) predicts that the destabilizing effect of an Arg mutation to an uncharged, non-aromatic residue is inversely proportional to the length of the protein (L), as well as linear with respect to the number of mutations. More specifically, it indicates that, for a PLD of the length of 100 amino acids, the change in the critical temperature will be of the order of 6.4 K. While formulaically similar to the scaling law for aromatic deletions, a notable difference is a change in the dependence on the PLD length. In the case of Arg deletions, the effect of mutations is screened by L rather than $\sqrt{L}$, indicating a stronger screening dependence on the PLD length. This indicates that for long PLDs, the destabilizing effects due to Arg deletions are diminished when compared to aromatic deletions, while in shorter PLDs this trend is reversed with each Arg deletion exerting a more significant destabilizing effect than the aromatic deletions.

These results underscore once again how for PLDs cation- π and π - π interactions are the main drivers of phase separation [9, 30, 31, 33, 35, 36, 74–76]. While it may seem counterintuitive for Arg deletions to have a lesser effect in the critical parameters of longer PLDs than their aromatic counterparts (as Arg residues are known to form strong cation- π interactions [104, 105]) the explanation lies in the type of systems in question, which are particularly rich in aromatic residues. Longer PLDs have, as a result of their higher valency, a larger network of possible pairwise interactions, which can mitigate the destabilizing effect of Arg deletions. In the case of shorter PLDs, each Arg residue represents a larger proportion of the total valency, and thus its deletion has a proportionally larger impact on the critical parameters. In contrast, aromatic residues contribute to the stability of the condensate through both cation- π and π - π interactions, which are predicted to be weaker than their cation- π analogues [31, 32], but more numerous in PLDs. In a longer PLD, the deletion of an aromatic residue results in the loss of more potential interactions, creating a larger destabilizing effect than the deletion of an Arg residue. However, in shorter PLDs, due to their lower total valency, the loss of potential associative interactions that result from the deletion of an aromatic residue is lower. Hence, the destabilizing effect resulting from this mutation is comparatively less than for the Arg deletion.

Our scaling laws have also allowed us to quantify the destabilizing effect of Arg to Lys mutations in the phase separation of PLDs. Concretely, mutations of Arg residues to Lys result in an overall decrease in the critical temperature of phase separation, two-fold in comparison to the impact of mutating Arg into an alternate uncharged non-aromatic residue. Consequently, the scaling law considerably intensifies for mutations from Arg to Lys, nearly doubling the effects of each mutation in the critical temperature of the condensate (See **Table 1**). This implies that each mutation of Arg to Lys exerts a substantially more destabilizing effect than mutations of Arg to other uncharged, non-aromatic residues, consistent with experimental evidence [30, 33, 83, 106] and highlighting the significant, modulatory role Lys plays in the phase behavior of PLDs.

Uncharged, non-aromatic residues such as Ser, Gly, Asn, Gln, and Thr are found to play a marginal role in PLD condensate stability [33, 37, 68, 72]. Notably, PLD condensates are highly stable in the face of Gly, Ser, Thr and Ala mutations. This behavior is quantified by their respective scaling laws, with values for the scaling constants for mutations from Gly to Thr, Ser to Thr and Ala to Ser at -0.09 K, -0.08 K and 0.16 K respectively. These numerical values indicate that, for a change of one degree in the critical temperature, approximately 10 amino acids of this set would need to undergo mutation. Notably, these scaling laws do not display a significant length dependency. This observation aligns with the idea that these mutations primarily affect the condensate properties through modulation in excluded volume and solvation, rather than by interrupting specific interaction networks.

In contrast, the condensates display heightened sensitivity towards Asn and Gln mutations, underscored by the Asn to Gln scaling of 50 K/L. This law is also inversely length dependent, indicating that mutations on these amino acids affect interaction networks and that mutations Asn to Gln result in a heightened ability of the system to phase separate. We hypothesize that, in addition to differences in solvation volume, the polarity of each amino acid plays a role by forming charge–dipole and dipole–dipole interactions. Both Gln and Asn are polar amino acids, with Gln having a longer side chain than Asn, which could potentially interact with a larger number of nearby atoms. Furthermore, the additional methylene group ($-\text{CH}_2-$) may slightly alter the electron distribution around the amide group, which can lead to the formation of more stable dipole–dipole contacts. It is important to note that our computational model, Mpipi, is a charge-fixed model, and hence does not directly capture the effects of polarization. However, the lack of physical details in the Mpipi model is compensated by the parameters having been derived from bioinformatics data.

While the Mpipi model has proven to be a powerful tool in predicting phase behaviour of PLDs, it is vitally important to acknowledge its limitations. Notably, the model does not account for temperature-dependent dependencies related to solvent effects, such as hydrophobic interactions and charge-mediated interactions facilitated by desolvation penalties, which implies that the model might not accurately predict the effects of mutations in particular sequences, such as sequences high in proline or aliphatic residues, where the interaction landscape is temperature sensitive. Additionally, the Mpipi model is not predictive of folding, or secondary structure. An assumption we made was that the mutations introduced do not significantly alter the secondary structure of each PLD, which is a necessary simplification for the scope of this study.

Collectively, the definition of a set of scaling laws acts as a unifying framework for the study of the critical parameters of PLD phase separation, by allowing for direct comparison of individual sequence effects. Such laws are cumulative, meaning that the effects of successive mutations can be predicted by the addition of the appropriate laws, as long as the resulting mutation keeps within PLD composition and patterning biases.

L	$Y \rightarrow F$	$F/Y \rightarrow W$	$F/Y \rightarrow X$	$R \rightarrow K$	$R \rightarrow X$
100	-0.4 ± 0.04	4.3 ± 0.1	-5.6 ± 0.5	-13 ± 1	-6.4 ± 0.5
200	-0.4 ± 0.04	4.3 ± 0.1	-3.9 ± 0.4	-6.5 ± 0.5	-3.2 ± 0.3
300	-0.4 ± 0.04	4.3 ± 0.1	-3.2 ± 0.3	-4.3 ± 0.3	-2.1 ± 0.2

L	$N \rightarrow Q$	$S \rightarrow T$	$G \rightarrow T$	$A \rightarrow S$
100	0.5 ± 0.05	-0.09 ± 0.03	-0.08 ± 0.02	0.16 ± 0.05
200	0.25 ± 0.03	-0.09 ± 0.03	-0.08 ± 0.02	0.16 ± 0.05
300	0.17 ± 0.02	-0.09 ± 0.03	-0.08 ± 0.02	0.16 ± 0.05

Table 1.

Predicted effect of single point mutations on the critical temperature, in Kelvin, of PLDs, as a function of their length, L

Our work explores a new framework to describe, quantitatively, the phase behavior of a large set of protein variants via the identification of scaling laws. Our scaling laws can predict changes in the stability of condensates based on amino acid sequence mutations. This code not only allows for testing the effects of composition bias on PLD phase separation but also serves as a predictive tool, allowing for a better selection of mutations to achieve desired changes in different condensates. This study is focused on PLDs, but we hypothesize similar scaling behavior in other IDPs, with the laws having a dependence on the protein length that reflects the nature of the underlying interaction networks of the system. Our findings have important implications for understanding PLD condensation and for designing modifications to counter their aberrant aggregation, by setting the basis for characterizing and quantifying the molecular determinants of PLD phase behavior.

Methods and materials

Design of the dataset

To design sequence variants to test the phase behavior of prion-like domains, we use in-silico targeted mutagenesis to introduce specific changes to the amino acid sequence of the PLD of interest. These changes involve substituting different amino acids, while carefully maintaining patterning and compositional biases, to test the role of given amino acids in the phase-separating propensities of the respective PLDs.

The nomenclature employed is consistent with that of Bremer and colleagues [33]. Namely, if n amino acids of type X are removed from the sequence and replaced with n amino acids of type Y , the variant is labeled $-nX + nY$. If instead, n amino acids of type X are removed from the sequence and replaced with amino acids (often different types) in proportion to the compositional bias of the PLD, the variant is simply labeled $-nX$.

To begin, we select a family of PLDs composed of hNRNPA1, RBM14, FUS, TDP43, TIA1 and EWSR1 and perform mutations on each sequence to generate a set of variants. Accordingly, we assess whether the effects of sequence mutations in PLD phase behaviors hold for the whole family of PLDs. Furthermore, our data set allows us to develop a set of rules that can aid in understanding and predicting PLD phase behavior. The amino acid sequences of the PLDs, and their corresponding variants, investigated herein are provided in the Supporting Information.

In total, we compute phase diagrams of 140 PLD variants via molecular dynamics (MD). This task required a large amount of CPU hours, over a 1.25 million in total. Given these demands, efficient utilization of high-performance computing (HPC) resources and proper parallelization of the computational tasks are paramount to ensuring optimal performance. To this end, we compute MD trajectories using the LAMMPS (Large-scale Atomic/Molecular Massively Parallel Simulator) software [107], exploiting its efficient parallelization algorithms. Another significant computational challenge involves the large size of data generated. Each single run produced a significant amount of trajectory, backup, and analysis data, and a large number of files. On average, each run produced over 50 files and a minimum of 5 GB of data. To efficiently handle this, we implement a robust, tiered, data storage and retrieval system, ensuring that the vast amounts of data is organized through a combination of indexing and partitioning, enhancing search efficiency and accessibility. We leveraged the Lustre filesystem for its high performance and scalability, making it the ideal choice for managing and accessing large volumes of active data required by our computational jobs. Custom scripts are employed for automated data processing, enabling us to extract meaningful insights from the complex datasets. All scripts are available in our GitHub repository.

Critical Parameter Determination Through MD

We used MD simulations to determine the full binodals (i.e., the boundaries that delimit the two-phase region from the one-phase region) for the PLDs. Specifically, we perform MD simulations in the canonical ensemble – i.e., fixed volume (V), temperature (T), and number of particles (N). For each binodal, a series of independent NVT simulations are performed at different temperatures and the density of each of the coexisting phases are determined. Thus, this approach yields binodals in the temperature–density phase space.

Direct Coexistence Simulations

Specifically, we use Direct Coexistence simulations, which enables the simultaneous simulation of both high- and low-density phases of a mixture, separated by an interface, within a single simulation box. Additionally, these simulations are performed with periodic boundary conditions in each dimension.

To initiate the NVT Direct Coexistence simulations, we start with one copy of a PLD in a cubic simulation cell and replicate it to the target number of chains (64 in total). We then iteratively scaled the simulation box to compress the chains into a high density direct coexistence simulation (approx. 0.8–1 g/cm³) [100]. Here, we use a Langevin thermostat [108] (at the target temperature, with a damping time of 40 ps) for temperature control. Keeping the length of the box in the X and Y dimensions fixed, we then elongate the Z dimension of the box. We then simulate each system at the target temperature for over 200 ns (see Convergence test in the SI). From each simulation we compute the density profile in the Z direction and determine the density of the dilute and condensed phases. These data are then used to produce the binodals.

Mpapi model

All the simulations are run using the Mpapi coarse-grained force-field [32] with an implicit solvent. In the Mpapi model electrostatic, cation– π and π – π interactions are carefully balanced to capture the phase behavior of PLDs. Notably, Mpapi achieves quantitative agreement with experiments in terms of prediction of critical solution temperatures of such systems. In Mpapi, each amino acid is represented by a single unique bead – i.e., of corresponding mass, molecular diameter (σ), charge (q), and an energy scale that reflects the relative frequency of planar π – π contacts (ϵ) [32].

The potential energy in the Mpapi model is computed as follows:

$$E_{\text{mpapi}} = E_{\text{bond}} + E_{\text{elec}} + E_{\text{pair}}. \quad (16)$$

Beads are connected via harmonic springs

$$E_{\text{bond}} = \sum_i \frac{1}{2} k (r_i - r_i^0)^2. \quad (17)$$

Here k is the bond force constant (19.1 kcal mol^{−1} Å^{−2}) and r_i^0 is the equilibrium bond length, which is set to 3.81 Å.

The electrostatic contribution is modeled via a Coulomb interaction term, with Debye–Hückel screening.

$$E_{\text{elec}} = \sum_{i,j} \frac{q_i q_j}{4\pi\epsilon_r \epsilon_0 r_{ij}} \exp(-\kappa r_{ij}), \quad (18)$$

where q_i and q_j are the charges of the bead i and j , respectively. ϵ_0 is the electric constant and $\epsilon_r = 80$ is the relative dielectric constant of water. $\kappa = 0.126 \text{ \AA}^{-1}$ is the inverse Debye screening length, which corresponds to monovalent salt concentration of 150 mM.

Finally, short-ranged non-bonded interactions are modeled with the Wang–Frenkel potential

$$E_{\text{pair}} = \sum_{i,j} \phi_{i,j} = \sum_{i,j} \epsilon_{i,j} \alpha_{i,j} \left[\left(\frac{\sigma_{i,j}}{r} \right)^{2\mu_{i,j}} - 1 \right] \left[\left(\frac{\sigma_{i,j}}{r} \right)^{2\nu_{i,j}} - 1 \right]^{2\nu_{i,j}} \quad (19)$$

where E_{pair} is the potential energy between all pairs of amino acids, $\phi_{i,j}$ is the potential energy between the pairs of amino acids i and j , $\epsilon_{i,j}$ is an energy parameter that dictates the strength of interaction of that particular pair, $\sigma_{i,j}$ is average the Van der Waals radii of the pair that dictate the position of the energy minima, r is the distance between center of masses and $\nu_{i,j}$ and $\mu_{i,j}$ are fitted exponential parameters that determine the shape of the energy curve. Full details of the Mpipi model are described in 32.

Data fitting

To estimate the critical temperatures of the binodals computed in Section , we use the law of coexistence densities

$$(\rho_{\text{high}}(T) - \rho_{\text{low}}(T))^{3.06} = d \left(1 - \frac{T}{T_c} \right) \quad (20)$$

The constant 3.06 in the equation is a dimensionless empirical factor that was derived from simulations of the 3D Ising model. The critical densities are computed by assuming that the law of rectilinear diameters holds, namely

$$\rho_{\text{high}}(T) + \rho_{\text{low}}(T) = 2\rho_c + 2A(T - T_c) \quad (21)$$

where $\rho_{\text{high}}(T)$, $\rho_{\text{low}}(T)$ and ρ_c are the densities of the high-density and low-density phases and the critical density, respectively, T_c is the critical temperature, and d and A are fitting parameters.

Finite-size scaling

In direct coexistence simulations in the slab geometry, finite-size effects can arise from either the cross-sectional area (i.e., size of the interface) or from the long axis (i.e., fraction bulk condensate). To ensure our calculations are robust with respect to system size, we perform finite-size scaling at fixed system number density: (1) varying the size of the cross section and (2) varying the length of the long axis. Using these tests, we confirm that the phase diagrams computed for systems of varying sizes (27, 64, 125 and 216 chains) exhibit no significant difference in predicted densities (see [Figure 2](#) in Supporting Information). Our scaling analysis revealed that 64 chains in a box of dimensions 125×125×1000 is the minimum size that ensures proteins do not interact with their periodic images and that the condensed phase is large enough to represent a true bulk phase.

Acknowledgements

We thank Pin Yu Chew for useful comments on the manuscript. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 803326 to R.C.G.). MJM would like to acknowledge the Winton Programme for Physics of Sustainability for doctoral funding. AAG is funded by the ERC (grant agreement No 803326). JRE acknowledges funding from the Ramon y Cajal fellowship (RYC2021-030937-I) and the Spanish National Agency for Research under the grant PID2022-136919NA-C33. This work was supported by the Engineering and Physical Sciences Research

Council (EPSRC) [grant number EP/X02332X/1 to JH] under the UK Research and Innovation (UKRI) Post-doctoral Fellowships Guarantee Scheme [project TF-CHROM-LLPS]. JAJ acknowledges research support from departmental start-up funds provided by the Department of Chemical and Biological Engineering and the Omenn–Darling Bioengineering Institute at Princeton University. This project made use of time on HPC granted via the UK High-End Computing Consortium for Biomolecular Simulation, HECBioSim (<http://hecbiosim.ac.uk> ) , supported by EPSRC (grant no. EP/R029407/1).

Code and data availability

The source code for running the Mpipi model in LAMMPS, example input scripts, as well as instructions for reproducing the data through a demo can be found at the GitHub repository: https://github.com/JuliMaristany/PLD_Scaling_Laws  . The data supporting the findings of this manuscript will be available via a figshare repository.

Full amino acid sequences

RBM14

WT(Q96PK6)	1	PMTAQAASYRAQPSVSLGAPYRGQLASPSQSAASSLGPPYGAQPSASALSSYGGQAAAASSLNSYGAQGSLLASYG
-20Y	1	G.S.Q.G.S.
-10Y	1	G.S.Q.G.S.
-5Y	1	S.S.S.S.
-30Y+30F	1	F.F.F.F.F.
-20Y+20F	1	F.F.F.F.F.
-10Y+10F	1	F.F.F.F.F.
-5Y+5F	1	F.F.F.F.F.
-10T+10S	1	S.S.S.S.
-10T+10G	1	G.G.G.G.
-5T+5S	1	S.S.S.S.
-5T+5G	1	G.G.G.G.
-20Q+20N	1	N.N.N.N.
-10Q+20N	1	N.N.N.N.
-5Q+5N	1	N.N.N.N.
-4N+4Q	1	Q.Q.Q.Q.
WT(Q96PK6)	79	NQPSYGAQAASSYGVRAAASSYNTQGAASSLGSYGAQAASYGQAASSLAYGAQAASYNAPASASNAQSAAPYAAQ
-20Y	79	G.Q.Q.G.S.G.Q.S.
-10Y	79	Q.Q.S.S.G.G.G.
-5Y	79	G.G.S.S.S.S.
-30Y+30F	79	F.F.F.F.F.F.F.
-20Y+20F	79	F.F.F.F.F.F.F.
-10Y+10F	79	F.F.F.F.F.F.F.
-5Y+5F	79	F.F.F.F.F.F.F.
-10T+10S	79	S.S.S.S.S.
-10T+10G	79	G.G.G.G.G.
-5T+5S	79	S.S.S.S.S.
-5T+5G	79	G.G.G.G.G.
-20Q+20N	79	N.N.N.N.N.
-10Q+20N	79	N.N.N.N.N.
-5Q+5N	79	N.N.N.N.N.
-4N+4Q	79	Q.Q.Q.Q.Q.
WT(Q96PK6)	157	QAASYSQPAAYVAQPATAAAYASQPAAYAAQATTPMAGSYGAQPVVQTQLNSYGAQASMG LSGSYGAQSAAAATG SY
-20Y	157	Q.S.Q.S.S.Q.S.
-10Y	157	S.S.Q.S.S.S.
-5Y	157	S.S.Q.Q.S.
-30Y+30F	157	F.F.F.F.F.F.F.
-20Y+20F	157	F.F.F.F.F.F.F.
-10Y+10F	157	F.F.F.F.F.F.F.
-5Y+5F	157	F.F.F.F.F.F.F.
-10T+10S	157	S.S.S.S.S.
-10T+10G	157	G.G.G.G.G.
-5T+5S	157	S.S.S.S.S.
-5T+5G	157	G.G.G.G.G.
-20Q+20N	157	N.N.N.N.N.
-10Q+20N	157	N.N.N.N.N.
-5Q+5N	157	N.N.N.N.N.
-4N+4Q	157	Q.Q.Q.Q.Q.
WT(Q96PK6)	235	GAAAAYGAQPSATLAAPYRTQSSASLAASYAAQHPQAAASYRGQPGNAYDGAGQPSAAYLSMSQGAVANANSTPPPY
-20Y	235	S.Q.Q.S.S.Q.S.
-10Y	235	S.S.S.S.S.
-5Y	235	S.S.S.S.
-30Y+30F	235	F.F.F.F.F.F.
-20Y+20F	235	F.F.F.F.F.F.
-10Y+10F	235	F.F.F.F.F.F.
-5Y+5F	235	F.F.F.F.F.F.
-10T+10S	235	S.S.S.S.S.
-10T+10G	235	G.G.G.G.G.
-5T+5S	235	S.S.S.S.S.
-5T+5G	235	G.G.G.G.G.
-20Q+20N	235	N.N.N.N.N.
-10Q+20N	235	N.N.N.N.N.
-5Q+5N	235	N.N.N.N.N.
-4N+4Q	235	Q.Q.Q.Q.Q.

FUS

(P35637)	1	M A S N D Y T Q Q A T Q S Y G A Y P T Q P G Q G Y S Q Q S S Q P Y G Q Q S Y S G Y S Q S T D T S G Y G Q S S Y S S Y G Q S
-28Y+28F	1	 F F . . F F F F F
-14Y+14F	1	 F F F F F F
-7Y+7F	1	 F . F F
-14Y	1	 Q G S Q G
-7Y	1	. Q . G
-10T+10S	1	 S . . . S S . S . . . S
-10T+10G	1	 G . . . G G . G . . . G
-5T+5S	1	 S S . S
-5T+5G	1	 G G . G
-10G+10T	1	. T
-10S+10T	1	 T . T
-20Q+20N	1	 N . . . N N N N N N
-10Q+10N	1	 N N N N
-5Q+5N	1	 N . N
-4N+4Q	1	. .
(P35637)	62	Q N T G Y G T Q S T P Q G Y G S T G G Y G S S Q S S Q S S Y G Q Q S S Y P G Y G Q Q P A P S S T S G S Y G S S S Q S S S Y
-28Y+28F	62	 F F F F F F F F
-14Y+14F	62	 F F F F F
-7Y+7F	62	. F . F
-14Y	62	 S . Q G S
-7Y	62	 S . Q
-10T+10S	62	. . . S . . . S . . . S . . . S . S
-10T+10G	62	. . . G . . . G . . . G . . . G . G
-5T+5S	62	 S S . S
-5T+5G	62	 G G . G
-10G+10T	62	. T . T
-10S+10T	62	. T T T
-20Q+20N	62	 N N N N N N N
-10Q+10N	62	. N N N
-5Q+5N	62	. N N
-4N+4Q	62	. Q .
(P35637)	123	G Q P Q S G S Y S Q Q P S Y G G Q Q Q S Y G Q Q Q S Y N P P Q G Y G Q Q N Q Y N S S S G G G G G G G G G G N Y G Q D Q S S
-28Y+28F	123	 F F F F F F F F
-14Y+14F	123	 F F F F F
-7Y+7F	123	. F . F
-14Y	123	 Q S G
-7Y	123	 S . G
-10T+10S	123	. .
-10T+10G	123	. .
-5T+5S	123	. .
-5T+5G	123	. .
-10G+10T	123	. T . T
-10S+10T	123	. T T T
-20Q+20N	123	. . . N N N N N N N N
-10Q+10N	123	. . . N N N N N
-5Q+5N	123	. N . N
-4N+4Q	123	. Q . . . Q Q
(P35637)	184	M S S G G G S G G G Y G N Q D Q S G G G G S G G Y G Q Q D R G G R G R G G S G G G G G G G G G G Y N R S S G
-28Y+28F	184	 F F F
-14Y+14F	184	. F .
-7Y+7F	184	. .
-14Y	184	 G . G
-7Y	184	 G .
-10T+10S	184	. .
-10T+10G	184	. .
-5T+5S	184	. .
-5T+5G	184	. .
-10G+10T	184	 T T T
-10S+10T	184	. . . T .
-20Q+20N	184	. .
-10Q+10N	184	. .
-5Q+5N	184	. .
-4N+4Q	184	. Q

EWSR1

Dataset representation

WT(Q01844)	1	MASTDYSTYSQAAQQGY SAYTAQPTQGYAQTQTAYGQQSYGTYGQPTDVSYTQAQTTATYGGTAYATSYGQP
-20Y	1	T.....Q.....S.....G.....T.....Q.....S.....T.....Q.....
-10Y	1	T.....Q.....S.....G.....T.....Q.....S.....T.....Q.....
-5Y	1	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-30Y+30F	1	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-20Y+20F	1	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-10Y+10F	1	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-5Y+5F	1	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-30T+30S	1	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-30T+30G	1	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-20T+20S	1	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-10T+10S	1	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-20T+20G	1	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-10T+10G	1	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-30S+30T	1	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20G+20T	1	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10G+10T	1	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20S+20T	1	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10S+10T	1	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20Q+20N	1	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-10Q+10N	1	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-5Q+5N	1	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-4N+4Q	1	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
WT(Q01844)	74	PTGYTTPAPQAYSQPVQGYGTGAYDTTATVTTTQASYYAQSYGTQPAYPAYCQPPAATAPTRPDQGNKPT
-20Y	74	T.....G.....Q.....Q.....S.....S.....S.....S.....S.....S.....
-10Y	74	Q.....S.....G.....G.....G.....G.....G.....G.....G.....G.....
-5Y	74	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-30Y+30F	74	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-20Y+20F	74	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-10Y+10F	74	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-5Y+5F	74	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-30T+30S	74	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-30T+30G	74	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-20T+20S	74	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-10T+10S	74	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-20T+20G	74	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-10T+10G	74	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-30S+30T	74	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20G+20T	74	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10G+10T	74	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20S+20T	74	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10S+10T	74	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20Q+20N	74	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-10Q+10N	74	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-5Q+5N	74	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-4N+4Q	74	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
WT(Q01844)	147	ETSQPQSSTGGYNQPSLGYGQSNYSYPQVPGSYPMQPVAPPSPYPTSYSSSTQPTSYDQSSYSQQNTYGGPSS
-20Y	147	Q.....Q.....T.....S.....S.....S.....S.....S.....S.....
-10Y	147	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-5Y	147	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-30Y+30F	147	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-20Y+20F	147	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-10Y+10F	147	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-5Y+5F	147	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-30T+30S	147	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-30T+30G	147	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-20T+20S	147	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-10T+10S	147	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-20T+20G	147	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-10T+10G	147	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-30S+30T	147	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20G+20T	147	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10G+10T	147	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20S+20T	147	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10S+10T	147	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20Q+20N	147	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-10Q+10N	147	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-5Q+5N	147	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-4N+4Q	147	Q.....Q.....Q.....Q.....Q.....Q.....Q.....Q.....Q.....Q.....
WT(Q01844)	220	YGGQSSYGGQSSYGGQPPPTSYPPQTGSYSQAPSQYSQQSSSYGGQSSFRQDHPSSMGVYGG
-20Y	220	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-10Y	220	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-5Y	220	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-30Y+30F	220	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-20Y+20F	220	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-10Y+10F	220	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-5Y+5F	220	F.....F.....F.....F.....F.....F.....F.....F.....F.....F.....
-30T+30S	220	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-30T+30G	220	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-20T+20S	220	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-10T+10S	220	S.....S.....S.....S.....S.....S.....S.....S.....S.....S.....
-20T+20G	220	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-10T+10G	220	G.....G.....G.....G.....G.....G.....G.....G.....G.....G.....
-30S+30T	220	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20G+20T	220	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10G+10T	220	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20S+20T	220	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-10S+10T	220	T.....T.....T.....T.....T.....T.....T.....T.....T.....T.....
-20Q+20N	220	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-10Q+10N	220	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-5Q+5N	220	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....
-4N+4Q	220	N.....N.....N.....N.....N.....N.....N.....N.....N.....N.....

Appendix 1

Convergence tests

Molecular Dynamic simulations are inherently sensitive to both the length of simulated time and system scaling factor such as the number of particles present in the system and the size of the simulation box. In this study, we systematically tested all these aspects to ensure proper convergence of the results.

The system was placed in the middle of the simulation box, and the length of X and Y was the same as the length of the condensate, while the box was elongated in Z, to about ten times the size of X, Y (see 1). Importantly, the cross-section is large enough to avoid protein self-interactions through the periodic boundary conditions (e.g., cross-section sizes are at least twice the protein radius of gyration in condensate bulk conditions).

We tested for time convergence by plotting the obtained density of the protein wild types after increasing simulation times, from 50 ns to 350 ns. On simulated timesteps above 100 ns, the density of the condensate exhibited no significant differences. We thus concluded that, to optimize speed-up, 200 ns provided sufficient sampling for each point of the binodal.

Finally, we decided to keep the number of particles at a constant value of 64, which we found to be a good balance between minimizing finite-size effects and optimizing simulation speed (see [Figure 2](#)). We tested this by varying the corresponding number of protein chains of the FUS WT as a model protein from 27 to 216 and found that in these systems there was not a significant difference in the simulated density. Additionally, with 64 protein chains, we could ensure a bulk large enough to sustain the condition that the cross-section of the box should be higher than twice the radii of gyration of the protein chain.

Appendix 2

Normalised and comparative contact maps

Further study of the differences between variants at the chemical level was carried out by computing the inter-molecular contacts per residue inside a phase-separated droplet. For each variant, the contact matrix was computed at a temperature around ten percent below the T_c . The cutoff to define a pairwise distance as a contact was dynamically calculated for each pair, as 1.2 times the average Van der Waals radius of both pair residues. Given residues i and j belong to different molecules, the contact maps were computed as shown below:

$$c_{i,j} = \sum_{i,j} \chi_{ij} \quad (22)$$

$$\chi_{ij} = \begin{cases} 1 & \text{if } |r_i - r_j| \leq r_{ij}^{\text{cutoff}} \\ 0 & \text{otherwise} \end{cases} \quad (23)$$

To get a more accurate picture of the main interactions driving phase separation for each variant, we normalised the contact maps to account for the occurrence of each residue in the sequence. This allowed us to account for the disparities in residue distributions in the sequences, and extract the relevant residue-residue interactions. For each residue-residue pair, the normalisation was performed as follows:

$$C = \begin{bmatrix} \alpha_{M,M} & \alpha_{M,G} & \dots & \alpha_{M,I} \\ \alpha_{G,M} & \alpha_{G,G} & \dots & \alpha_{G,I} \\ \dots & \dots & \dots & \dots \\ \alpha_{I,M} & \alpha_{I,G} & \dots & \alpha_{I,I} \end{bmatrix} \quad (24)$$

$$\alpha_{i,j} = \frac{c_{i,j}}{n_i \cdot n_j} \quad (25)$$

where n_i is the number of i residues in the IDP sequence. The resulting contact matrix is min–max normalised:

$$\beta_{i,j} = \frac{\alpha_{i,j} - \min C}{\max C - \min C} \quad (26)$$

$$C_{\text{norm}} = \begin{bmatrix} \beta_{M,M} & \beta_{M,G} & \cdots & \beta_{M,I} \\ \beta_{G,M} & \beta_{G,G} & \cdots & \beta_{G,I} \\ \cdots & \cdots & \cdots & \cdots \\ \beta_{I,M} & \beta_{I,G} & \cdots & \beta_{I,I} \end{bmatrix} \quad (27)$$

The comparative contact maps (**Fig. 1** and **2**), on the other hand, are computed as the difference in contacts of a protein variant respective to its wild type, as:

$$C_{\text{comparative}} = C_{\text{variant}} - C_{\text{WT}} \quad (28)$$


which is finally mean-normalised.

$$\gamma_{i,j} = \frac{\alpha_{i,j} - \overline{C}_{\text{comparative}}}{\max C_{\text{comparative}} - \min C_{\text{comparative}}} \quad (29)$$

$$C_{\text{comparative, normed}} = \begin{bmatrix} \gamma_{M,M} & \gamma_{M,G} & \cdots & \gamma_{M,I} \\ \gamma_{G,M} & \gamma_{G,G} & \cdots & \gamma_{G,I} \\ \cdots & \cdots & \cdots & \cdots \\ \gamma_{I,M} & \gamma_{I,G} & \cdots & \gamma_{I,I} \end{bmatrix} \quad (30)$$

Appendix 0 figure 8.


Representation of the entire computational data set, divided on mutations on 'sticker' or 'spacer' residues. Orange data points, on the left, represent variants where charged or aromatic residues were mutated (i.e., stickers), while cyan data points, on the right, represent all other types of mutations studied (i.e., spacers)




Appendix 1 figure 1.

Convergence tests for the WT sequence of the PLD of FUS.

The density profiles of different and independent simulations are plotted across the perpendicular axis to the condensate interfaces. Each simulation was performed for different timescales (as specified in the legend) to check for proper convergence.






Appendix 1 figure 2.

Convergence tests for the WT sequence of the PLD of FUS.


The density profiles of different and independent simulations are plotted across the perpendicular axis to the condensate interfaces. Each simulation was performed for different system sizes (aka, number of chains and corresponding box size, as specified in the legend) to check for proper convergence. To the right, a representation of the systems of increasing number of chains we employed to check for convergence. In each case, the overall density of the box is kept constant, while the cross-section and length is varied in order to maintain the density constraint. For 27 chains, the box dimensions are 109×109×765 Å; for 64 chains 145×145×1021 Å; for 125 chains 182×182×1276 Å and for the final case of 216 218×218×1531 Å.




Appendix 2 figure 1.

Normalised contact maps of different *R to K* variants of PLDs: A) TDP B) FUS, C) hnRNPA1

A)



B)



Appendix 2 figure 2.

Normalised contact maps of different aromatic variants of PLDs: A) FUS B) hnRNPA1.

References

- [1] Banani Salman F., Lee Hyun O., Hyman Anthony A., Rosen Michael K. (2017) **Biomolecular condensates: Organizers of cellular biochemistry** *Nature Reviews Molecular Cell Biology* **18**:285–298
- [2] Li Pulong *et al.* (2012) **Phase transitions in the assembly of multivalent signalling proteins** *Nature* **483**:336–340
- [3] Mittag Tanja, Pappu Rohit V (2022) **A conceptual framework for understanding phase separation and ad-dressing open questions and challenges** *Molecular cell* **82**:2201–2214
- [4] Shin Yongdae, Brangwynne Clifford P. (2017) **Liquid phase condensation in cell physiology and disease** *Science* **357**
- [5] Gomes Edward, Shorter James (2019) **The molecular language of membraneless organelles** *Journal of Biological Chemistry* **294**:7115–7127
- [6] Nott Timothy J *et al.* (2015) **Phase transition of a disordered nuage protein generates environmentally responsive membraneless organelles** *Molecular cell* **57**:936–947
- [7] Smith Jarrett, Calidas Deepika, Schmidt Helen, Lu Tu, Rasoloson Dominique, Seydoux Geraldine (2016) **Spatial patterning of p granules by rna-induced phase separation of the intrinsically-disordered protein meg-3** *Elife* **5**
- [8] Wang Jie *et al.* (2018) **A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding Proteins** *Cell* **174**:688–699
- [9] Martin Erik W, Holehouse Alex S, Peran Ivan, Farag Mina, Incicco J Jeremias, Bremer Anne, Grace Christy R, Soranno Andrea, Pappu Rohit V, Mittag Tanja (2020) **Valence and patterning of aromatic residues determine the phase behavior of prion-like domains** *Science* **367**:694–699
- [10] Martin Erik W, Holehouse Alex S (2020) **Intrinsically disordered protein regions and phase separation: sequence determinants of assembly or lack thereof** *Emerging topics in life sciences* **4**
- [11] Posey Ammon E, Holehouse Alex S, Pappu Rohit V, Rhoades Elizabeth (2018) **Phase separation of intrinsically disordered proteins** *Methods in Enzymology* :1–30
- [12] Elbaum-Gar?nkle Shana, Kim Younghoon, Szczepaniak Krzysztof, Chen Carlos Chih-Hsiung, Eckmann Christian R, Myong Sua, Brangwynne Clifford P (2015) **he disordered p granule protein laf-1 drives phase separation into droplets with tunable viscosity and dynamics** *Proceedings of the National Academy of Sciences* **112**:7189–7194
- [13] Franzmann Titus M *et al.* (2018) **Phase separation of a yeast prion protein promotes cellular fitness** *Science* **359**
- [14] Maharana Shovamayee *et al.* (2018) **Rna buffers the phase separation behavior of prion-like rna binding proteins** *Science* **360**:918–921

- [15] Gotor Nieves Lorenzo, Armaos Alexandros, Calloni Giulia, Burgas Marc Torrent, Vabulas R Martin, De Groot Natalia Sanchez, Tartaglia Gian Gaetano (2020) **Rna-binding and prion domains: the yin and yang of phase separation** *Nucleic acids research* **48**:9491–9504
- [16] Molliex Amandine, Temirov Jamshid, Lee Jihun, Coughlin Maura, Kanagaraj Anderson P., Kim Hong Joo, Mittag Tanja, Taylor J. Paul (2015) **Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and Drives Pathological Fibrillization** *Cell* **163**:123–133
- [17] Jawerth Louise *et al.* (2020) **Protein condensates as aging maxwell ?uids** *Science* **370**:1317–1323
- [18] Alshareedah Ibraheem, Borchers Wade M, Cohen Samuel R, Farag Mina, Singh Anurag, Bremer Anne, Pappu Rohit V, Mittag Tanja, Banerjee Priya R (2023) **Sequence-encoded grammars determine material properties and physical aging of protein condensates** *bioRxiv* :2023–4
- [19] Linsenmeier Miriam, Hondele Maria, Grigolato Fulvio, Secchi Eleonora, Weis Karsten, Arosio Paolo (2022) **Dynamic arrest and aging of biomolecular condensates are modulated by low-complexity domains, rna and biochemical activity** *Nature Communications* **13**
- [20] Sprunger Macy L, Jackrel Meredith E (2021) **Prion-like proteins in phase separation and their link to disease** *Biomolecules* **11**
- [21] Banani SF (2016) **Compositional control of phase-separated cell. bodies** *Cell* **166**:651–663
- [22] Wang Jie *et al.* (2018) **A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding Proteins** *Cell* **174**:688–699
- [23] Choi Jeong Mo, Holehouse Alex S., Pappu Rohit V. (2020) **Physical Principles Underlying the Complex Biology of Intracellular Phase Transitions** *Annual Review of Biophysics* **49**:107–133
- [24] Choi Jeong-Mo, Hyman Anthony A, Pappu Rohit V (2020) **Generalized models for bond percolation transitions of associative polymers** *Physical Review E* **102**
- [25] Harmon Tyler S, Holehouse Alex S, Rosen Michael K, Pappu Rohit V (2017) **Intrinsically disordered linkers determine the interplay between phase separation and gelation in multivalent proteins** *elife* **6**
- [26] Iserman Christiane *et al.* (2020) **Condensation of ded1p promotes a translational switch from housekeeping to stress protein production** *Cell* **181**:818–831
- [27] Adame-Arana Omar, Weber Christoph A, Zaburdaev Vasily, Prost Jacques, Jülicher Frank (2020) **Liquid phase separation controlled by ph** *Biophysical Journal* **119**:1590–1605
- [28] Munder Matthias Christoph *et al.* (2016) **A ph-driven transition of the cytoplasm from a ? uid-to a solid-like state promotes entry into dormancy** *eLife* **5**
- [29] Cinar Hasan, Fetahaj Zamira, Cinar Süleyman, Vernon Robert M., Chan Hue Sun, Winter Roland H. A. (2019) **Temperature, hydrostatic pressure, and osmolyte effects on liquid–liquid phase separation in protein condensates: Physical chemistry and biological implications** *Chemistry – A European Journal* **25**:13049–13069
- [30] Fisher Rachel S, Elbaum-Gar?nkle Shana (2020) **Tunable multiphase dynamics of arginine and lysine liquid condensates** *Nature communications* **11**

- [31] Krainer Georg *et al.* (2021) **Reentrant liquid condensate phase of proteins is stabilized by hydrophobic and non-ionic interactions** *Nature Communications* **12**:1–14
- [32] Joseph Jerelle A., Reinhardt Aleks, Aguirre Anne, Chew Pin Yu, Russell Kieran O., Espinosa Jorge R., Garaizar Adiran, Collepardo-Guevara Rosana (2021) **Physics-driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy** *Nature Computational Science* **1**:732–743
- [33] Bremer Anne, Farag Mina, Borchers Wade M., Peran Ivan, Martin Erik W., Pappu Rohit V., Mittag Tanja (2022) **Deciphering how naturally occurring sequence features impact the phase behaviors of disordered prion-like domains** *Nature chemistry* **14**
- [34] Brady Jacob P *et al.* (2017) **Structural and hydrodynamic properties of an intrinsically disordered region of a germ cell-specific protein on phase separation** *Proceedings of the National Academy of Sciences* **114**:E8194–E8203
- [35] Vernon RM (2018) **Pi-Pi contacts are an overlooked protein feature relevant to phase separation** *eLife* **7**
- [36] Fossat Martin J, Zeng Xiangze, Pappu Rohit V (2021) **Uncovering differences in hydration free energies and structures for model compound mimics of charged side chains of amino acids** *The Journal of Physical Chemistry B* **125**:4148–4161
- [37] Martin Erik W, Mittag Tanja (2018) **Relationship of sequence and phase separation in protein low-complexity regions** *Biochemistry* **57**:2478–2487
- [38] Ball R. C. (1991) **Scaling in Polymer Physics** Boston, MA: Springer US :483–490
- [39] Dunstan Dave E. (2019) **The viscosity-radius relationship for concentrated polymer solutions** *Scientific Reports* **9**
- [40] Majumder Suman, Christiansen Henrik, Janke Wolfhard (2020) **Understanding nonequilibrium scaling laws governing collapse of a polymer** *The European Physical Journal B* **93**:1–19
- [41] Rubinstein Michael, Colby Ralph H, *et al.* (2003) **Polymer physics** New York: Oxford university press
- [42] Hofmann Hagen, Soranno Andrea, Borgia Alessandro, Gast Klaus, Nettels Daniel, Schuler Benjamin (2012) **Polymer scaling laws of unfolded and intrinsically disordered proteins quantified with single-molecule spectroscopy** *Proceedings of the National Academy of Sciences* **109**:16155–16160
- [43] Clarke Joseph P, Thibault Patricia A, Salapa Hannah E, Levin Michael C (2021) **A comprehensive analysis of the role of hnrnp a1 function and dysfunction in the pathogenesis of neurodegenerative disease** *Frontiers in Molecular Biosciences* **8**
- [44] Kim Hong Joo *et al.* (2013) **Mutations in prion-like domains in hn-rnpa2b1 and hnrnpa1 cause multisystem proteinopathy and als** *Nature* **495**:467–473
- [45] Wang I-Fan, Wu Lien-Szu, Shen CK James (2008) **Tdp-43: an emerging new player in neurodegenerative diseases** *Trends in molecular medicine* **14**:479–485

- [46] Jo Myungjin, Lee Shinrye, Jeon Yu-Mi, Kim Seyeon, Kwon Younghwi, Kim Hyung-Jun (2020) **The role of tdp-43 propagation in neurodegenerative diseases: integrating insights from clinical and experimental studies** *Experimental & molecular medicine* **52**:1652–1662
- [47] Ash Peter EA *et al.* (2021) **Tia1 potentiates tau phase separation and promotes generation of toxic oligomeric tau** *Proceedings of the National Academy of Sciences* **118**
- [48] Apicco Daniel J *et al.* (2018) **Reducing the rna binding protein tia1 protects against tau-mediated neurodegeneration in vivo** *Nature neuroscience* **21**:72–80
- [49] Deng Hao, Gao Kai, Jankovic Joseph (2014) **The role of fus gene variants in neurodegenerative diseases** *Nature Reviews Neurology* **10**:337–348
- [50] Clotilde Lagier-Tourenne and Don W Cleveland (2009) **Rethinking als: The fus about tdp-43** *Cell* **136**:1001–1004
- [51] Couthouis Julien *et al.* (2012) **Evaluating the role of the fus/tls-related gene ewsr1 in amyotrophic lateral sclerosis** *Human molecular genetics* **21**:2899–2911
- [52] Lee Junghee, Nguyen Phuong T, Shim Hyun Soo, Hyeon Seung Jae, Im Hyeonjoo, Choi Mi-Hyun, Chung Sooyoung, Kowall Neil W, Lee Sean Bong, Ryu Hoon (2019) **Ewsr1, a multifunctional protein, regulates cellular function and aging via genetic and epigenetic pathways** *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* **1865**:1938–1945
- [53] Li Yun R, King Oliver D, Shorter James, Gitler Aaron D (2013) **Stress granules as crucibles of als pathogenesis** *Journal of cell biology* **201**:361–372
- [54] Farag Mina, Borchers Wade M., Bremer Anne, Mittag Tanja, Pappu Rohit V. (2023) **Phase separation in mixtures of prion-like low complexity domains is driven by the interplay of homotypic and heterotypic interactions** *bioRxiv*
- [55] Reber Stefan *et al.* (2021) **The phase separation-dependent FUS interactome reveals nuclear and cytoplasmic function of liquid-liquid phase separation** *Nucleic Acids Research* **49**:7713–7731
- [56] Feng Zhe, Chen Xudong, Wu Xiandeng, Zhang Mingjie (2019) **Formation of biological condensates via phase separation: Characteristics, analytical methods, and physiological implications** *Journal of Biological Chemistry* **294**:14823–14835
- [57] Welsh Timothy J *et al.* (2022) **Surface electrostatics govern the emulsion stability of biomolecular condensates** *Nano letters* **22**:612–621
- [58] Luo Feng, Gui Xinrui, Zhou Heng, Gu Jing, Li Yichen, Liu Xiangyu, Zhao Minglei, Li Dan, Li Xueming, Liu Cong (2018) **Atomic structures of FUS LC domain segments reveal bases for reversible amyloid ?bril formation** *Nat. Struct. Mol. Biol* **25**:341–346
- [59] Murray Dylan T, Kato Masato, Lin Yi, Thurber Kent R, Hung Ivan, McKnight Steven L, Tycko Robert (2017) **Structure of FUS protein ?brils and its relevance to self-assembly and phase separation of low-complexity domains** *Cell* **171**:615–627
- [60] Qamar Seema *et al.* (2018) **Fus phase separation is modulated by a molecular chaperone and methylation of arginine cation- π interactions** *Cell* **173**:720–734

- [61] Murakami Tetsuro *et al.* (2015) **Als/ftd mutation-induced phase transition of fus liquid droplets and reversible hydrogels into irreversible hydrogels impairs rnp granule function** *Neuron* **88**:678–690
- [62] Patel Avinash *et al.* (2015) **A liquid-to-solid phase transition of the als protein fus accelerated by disease mutation** *Cell* **162**:1066–1077
- [63] Murthy Anastasia C, Dignon Gregory L, Kan Yelena, Zerze Gül H, Parekh Sapun H, Mittal Jeetain, Fawzi Nicolas L (2019) **Molecular interactions underlying liquid-liquid phase separation of the fus low-complexity domain** *Nature structural & molecular biology* **26**:637–648
- [64] Zbinden Auré Lie, Rez-Berlanga Manuela Pé, De Rossi Pierre, Polymenidou Magdalini (2020) **Phase Separation and Neurodegenerative Diseases: A Disturbance in the Force** *Developmental Cell* **55**:45–68
- [65] Guenther Elizabeth L, Cao Qin, Trinh Hamilton, Lu Jiahui, Sawaya Michael R, Cascio Duilio, Boyer David R, Rodriguez Jose A, Hughes Michael P, Eisenberg David S (2018) **Atomic structures of tdp-43 lcd segments and insights into reversible or pathogenic aggregation** *Nature structural & molecular biology* **25**:463–471
- [66] Lotthammer Jeffrey M, Ginell Garrett M, Griffith Daniel, Emenecker Ryan, Holehouse Alex S (2024) **Direct prediction of intrinsically disordered protein conformational properties from sequence** *Biophysical Journal* **123**
- [67] Hennig Sven *et al.* (2015) **Prion-like domains in RNA binding proteins are essential for building subnuclear paraspeckles** *Journal of Cell Biology* **210**:529–539
- [68] Holehouse Alex S., Ginell Garrett M., Griffith Daniel, Böke Elvan (2021) **Clustering of aromatic residues in prion-like domains can tune the formation, state, and organization of biomolecular condensates** *Biochemistry* **60**:3566–3581
- [69] Lin Yuan, Protter David S.W., Rosen Michael K., Parker Roy (2015) **Formation and Maturation of Phase-Separated Liquid Droplets by RNA-Binding Proteins** *Molecular Cell* **60**:208–219
- [70] Ryan Veronica H, Dignon Gregory L, Zerze Gül H, Chabata Charlene V, Silva Rute, Conicella Alexander E, Amaya Joshua, Burke Kathleen A, Mittal Jeetain, Fawzi Nicolas L (2018) **Mechanistic view of hnnpa2 low-complexity domain structure, interactions, and phase separation altered by mutation and arginine methylation** *Molecular cell* **69**:465–479
- [71] Sanchez-Burgos Ignacio, Espinosa Jorge R, Joseph Jerelle A, Collepardo-Guevara Rosana (2021) **Valency and binding affinity variations can regulate the multilayered organization of protein condensates with many components** *Biomolecules* **11**
- [72] Choi Jeong-Mo, Pappu Rohit V (2020) **The stickers and spacers framework for describing phase behavior of multivalent intrinsically disordered proteins** *Biophysical Journal* **118**
- [73] Ginell Garrett M, Holehouse Alex S (2022) **An introduction to the stickers-and-spacers framework as applied to biomolecular condensates** *Phase-Separated Biomolecular Condensates: Methods and Protocols* Springer :95–116
- [74] Das Suman, Lin Yi Hsuan, Vernon Robert M., Forman-Kay Julie D., Chan Hue Sun (2020) **Comparative roles of charge, π , and hydrophobic interactions in sequence-dependent phase separation of intrinsically disordered proteins** *Proceedings of the National Academy of Sciences of the United States of America* **117**:28795–28805

- [75] Dyson H Jane, Wright Peter E, Scheraga Harold A (2006) **The role of hydrophobic interactions in initiation and propagation of protein folding** *Proceedings of the National Academy of Sciences* **103**:13057–13061
- [76] Andrew Charles D, Bhattacharjee Samita, Kokkoni Nicoleta, Hirst Jonathan D, Jones Gareth R, Doig Andrew J (2002) **Stabilizing interactions between aromatic and basic side chains in α -helical peptides and proteins. tyrosine effects on helix circular dichroism** *Journal of the American Chemical Society* **124**:12706–12714
- [77] Li Wei, Jiang Hao, Protein Interactions in Liquid-Liquid Phase Separation (2022) **Nuclear protein condensates and their properties in regulation of gene expression** *Journal of Molecular Biology* **434**
- [78] Tejedor Andrés R, Collepardo-Guevara Rosana, Ramirez Jorge, Espinosa Jorge R (2023) **Time-dependent material properties of aging biomolecular condensates from different viscoelasticity measurements in molecular dynamics simulations** *The Journal of Physical Chemistry B*
- [79] Alberti Simon, Gladfelter Amy, Mittag Tanja (2019) **Considerations and Challenges in Studying Liquid-Liquid Phase Separation and Biomolecular Condensates** *Cell* **176**:419–434
- [80] Ray Soumik, Buell Alexander K (2024) **Emerging experimental methods to study the thermodynamics of biomolecular condensate formation** *The Journal of Chemical Physics* **160**
- [81] Espinosa Jorge R., Joseph Jerelle A., Sanchez-Burgos Ignacio, Garaizar Adiran, Frenkel Daan, Collepardo-Guevara Rosana (2020) **Liquid network connectivity regulates the stability and composition of biomolecular condensates with many components** *Proceedings of the National Academy of Sciences* **117**:13238–13247
- [82] Hong Yuri, Naja? Saeed, Casey Thomas, Shea Joan-Emma, Han Song-I, Hwang Dong Soo (2022) **Hydrophobicity of arginine leads to reentrant liquid-liquid phase separation behaviors of arginine-rich proteins** *Nature Communications* **13**
- [83] Greig Jamie A, Nguyen Tu Anh, Lee Michelle, Holehouse Alex S, Posey Ammon E, Pappu Rohit V, Jedd Gregory (2020) **Arginine-enriched mixed-charge domains provide cohesion for nuclear speckle condensation** *Molecular cell* **77**:1237–1250
- [84] Strom Amy R, Emelyanov Alexander V, Mir Mustafa, Fyodorov Dmitry V, Darzacq Xavier, Karpen Gary H (2017) **Phase separation drives heterochromatin domain formation** *Nature* **547**:241–245
- [85] Mitchell Dennis J., Steinman Lawrence, Kim D T, Fathman C G, Rothbard Jonathan B. (2000) **Polyarginine enters cells more efficiently than other polycationic homopolymers** *Journal of Peptide Research* **56**:318–325
- [86] Das Rahul K., Pappu Rohit V. (2013) **Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues** *Proceedings of the National Academy of Sciences* **110**:13392–13397
- [87] Cohan Megan C., Shinn Min Kyung, Lalmansingh Jared M., Pappu Rohit V. (2022) **Uncovering non-random binary patterns within sequences of intrinsically disordered proteins** *Journal of Molecular Biology* **434**

- [88] Zeng Xiangze, Ruff Kiersten M., Pappu Rohit V. (2022) **Competing interactions give rise to two-state behavior and switch-like transitions in charge-rich intrinsically disordered proteins** *Proceedings of the National Academy of Sciences* **119**
- [89] De Sancho David (2022) **Phase separation in amino acid mixtures is governed by composition** *Biophysical Journal* **121**:4119–4127
- [90] Ahlers Jonas, Adams Ellen M., Bader Verian, Pezzotti Simone, Winklhofer Konstanze F., Tatzelt Jörg, Havenith Martina (2021) **The key role of solvent in condensation: Mapping water in liquid-liquid phase-separated fus** *Biophysical Journal* **120**:1266–1275
- [91] Nomoto Akira, Nishinami Suguru, Shiraki Kentaro (2021) **Solubility parameters of amino acids on liquid-liquid phase separation and aggregation of proteins** *Frontiers in Cell and Developmental Biology* **9**
- [92] Zheng Wenwei, Dignon Gregory L., Brown Matthew L., Kim Young C., Mittal Jeetain (2020) **Hydropathy patterning complements charge patterning to describe conformational preferences of disordered proteins** *Journal of Physical Chemistry Letters* **11**:3408–3415
- [93] Weiner Benjamin G, Pyo Andrew GT, Meir Yigal, Wingreen Ned S (2021) **Motif-pattern dependence of biomolecular phase separation driven by specific interactions** *PLoS computational biology* **17**
- [94] Huihui Jonathan, Ghosh Kingshuk (2021) **Intrachain interaction topology can identify functionally similar intrinsically disordered proteins** *Biophysical Journal* **120**:1860–1868
- [95] Ghosh Kingshuk, Huihui Jonathan, Phillips Michael, Haider Austin (2022) **Rules of physical mathematics govern intrinsically disordered proteins** *Annual Review of Biophysics* **51**:355–376
- [96] Rana Ushnish, Brangwynne Clifford P., Athanassios Z. (2021) **Panagiotopoulos. Phase separation vs aggregation behavior for model disordered proteins** *The Journal of Chemical Physics* **155**
- [97] Martin Erik W, Holehouse Alex S, Grace Christy R, Hughes Alex, Pappu Rohit V, Mittag Tanja (2016) **Sequence determinants of the conformational properties of an intrinsically disordered protein prior to and upon multisite phosphorylation** *Journal of the American Chemical Society* **138**:15323–15335
- [98] Holehouse Alex S, Ginell Garrett M, Griffith Daniel, Boke Elvan (2021) **Clustering of aromatic residues in prion-like domains can tune the formation, state, and organization of biomolecular condensates: published as part of the biochemistry virtual special issue “protein condensates”** *Biochemistry* **60**:3566–3581
- [99] Regy Roshan Mammen, Thompson Jacob, Kim Young C., Mittal Jeetain (2021) **Improved coarse-grained model for studying sequence dependent phase separation of disordered proteins** *Protein Science* **30**:1371–1379
- [100] Dignon Gregory L., Zheng Wenwei, Kim Young C., Best Robert B., Mittal Jeetain (2018) **Sequence determinants of protein phase behavior from a coarse-grained model** *PLoS Computational Biology* **14**

- [101] Tesei Giulio, Schulze Thea K., Crehuet Ramon, Lindorff-Larsen Kresten (2021) **Accurate model of liquid-liquid phase behavior of intrinsically disordered proteins from optimization of single-chain properties** *Proceedings of the National Academy of Sciences* **118**
- [102] Regy Roshan Mammen, Zheng Wenwei, Mittal Jeetain (2021) **Using a sequence-specific coarse-grained model for studying protein liquid-liquid phase separation** *Methods Enzymol.* **646**:1–17
- [103] Latham Andrew P., Zhang Bin (2019) **Maximum entropy optimized force field for intrinsically disordered proteins** *Journal of Chemical Theory and Computation* **16**:773–781
- [104] Park Sohee, Lee Yongjin, Jho Yongseok, Hwang Dong Soo (2022) **Molecular hydration tunes the cation- π interaction strength in aqueous solution** *Advanced Materials Interfaces* **10**
- [105] Gallivan Justin P., Dougherty Dennis A. (2000) **A computational study of cation- π interactions vs salt bridges in aqueous media: implications for protein engineering** *Journal of the American Chemical Society* **122**:870–874
- [106] Ukmar-Godec Tina, Hutten Saskia, Grieshop Matthew P., Rezaei-Ghaleh Nasrollah, Cima-Omori Maria-Sol, Biernat Jacek, Mandelkow Eckhard, Söding Johannes, Dormann Dorothee, Zweckstetter Markus (2019) **Lysine/RNA-interactions drive and regulate biomolecular condensation** *Nature Communications* **10**
- [107] Thompson A. P. *et al.* (2022) **LAMMPS - a flexible simulation tool for particle-based materials modeling at the atomic, meso, and continuum scales** *Comp. Phys. Comm* **271**
- [108] Grønbech-Jensen Niels, Farago Oded (2013) **A simple and effective verlet-type algorithm for simulating langevin dynamics** *Molecular Physics* **111**:983–991

Author information

M Julia Maristany

Department of Physics, University of Cambridge, Cambridge, UK
ORCID iD: [0009-0009-8875-9225](https://orcid.org/0009-0009-8875-9225)

Anne Aguirre Gonzalez

Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK

Jorge R Espinosa

Universidad Complutense de Madrid, Madrid, Spain
ORCID iD: [0000-0001-9530-2658](https://orcid.org/0000-0001-9530-2658)

Jan Huertas

Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK
ORCID iD: [0000-0003-2607-5214](https://orcid.org/0000-0003-2607-5214)

Rosana Colleparado-Guevara

Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK,
Department of Genetics, University of Cambridge, Cambridge, UK
ORCID iD: [0000-0003-1781-7351](https://orcid.org/0000-0003-1781-7351)

Jerelle A Joseph

Department of Chemical and Biological Engineering, Princeton University, Princeton, USA,
Omenn–Darling Bioengineering Institute, Princeton University, Princeton, USA
ORCID iD: [0000-0003-4525-180X](https://orcid.org/0000-0003-4525-180X)

For correspondence: jerellejoseph@princeton.edu

Editors

Reviewing Editor

Bin Zhang

Massachusetts Institute of Technology, Cambridge, United States of America

Senior Editor

Qiang Cui

Boston University, Boston, United States of America

Reviewer #1 (Public review):

Summary:

In this preprint, the authors systematically and rigorously investigate how specific classes of residue mutations alter the critical temperature as a proxy for the driving forces for phase separation. The work is well executed, the manuscript well-written, and the results reasonable and insightful.

Strengths:

The introductory material does an excellent job of being precise in language and ideas while summarizing the state of the art. The simulation design, execution, and analysis are exceptional and set the standard for large-scale simulation studies. The results, interpretations, and Discussion are largely nuanced, clear, and well-motivated, and the pedagogical nature with which sampling convergence is discussed is greatly appreciated. Finally, the underlying data are shared in a clear and accessible manner. Overall, the manuscript is a model

Weaknesses:

The simplicity of a one-bead-per-residue model parameterized to capture UCST-type phase behavior does perhaps impact some aspects of the generality of this work. That said, the authors carefully acknowledge these limitations, and overall, this is not seen as a major weakness of the conclusions drawn or the manuscript, given those conclusions are appropriately couched.

<https://doi.org/10.7554/eLife.99068.2.sa3>

Reviewer #2 (Public review):

This is an interesting manuscript where a CA-only CG model (Mpipi) was used to examine the critical temperature (T_c) of phase separation of a set of 140 variants of prion-like low complexity domains (PLDs). The key result is that T_c of these PLDs seems to have a linear dependence on substitutions of various sticker and space residues. This is potentially useful for estimating the T_c shift when making novel mutations of a PLD.

Comments on revisions: The authors have addressed concerns raised previously.

<https://doi.org/10.7554/eLife.99068.2.sa2>

Reviewer #3 (Public review):

Summary:

"Decoding Phase Separation of Prion-Like Domains through Data-Driven Scaling Laws" by Maristany et al. offers a significant contribution to the understanding of phase separation in prion-like domains (PLDs). The study investigates the phase separation behavior of PLDs, which are intrinsically disordered regions within proteins that have the propensity to undergo liquid-liquid phase separation (LLPS). This phenomenon is crucial in forming biomolecular condensates, which play essential roles in cellular organization and function. The authors employ a data-driven approach to establish predictive scaling laws that describe the phase behavior of these domains.

Strengths:

The study benefits from a robust dataset encompassing a wide range of PLDs, which enhances the generalizability of the findings. The authors' meticulous curation and analysis of this data add to the study's robustness. The scaling laws derived from the data provide predictive insights into the phase behavior of PLDs, which can be useful in the future for the design of synthetic biomolecular condensates.

<https://doi.org/10.7554/eLife.99068.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer # 1 (Public Review):

Summary:

In this preprint, the authors systematically and rigorously investigate how specific classes of residue mutations alter the critical temperature as a proxy for the driving forces for phase separation. The work is well executed, the manuscript well-written, and the results reasonable and insightful.

Strengths:

The introductory material does an excellent job of being precise in language and ideas while summarizing the state of the art. The simulation design, execution, and analysis are exceptional and set the standard for these types of large-scale simulation studies. The results, interpretations, and Discussion are largely nuanced, clear, and well-motivated.

We thank the reviewer for their assessment of our work and for highlighting the key strengths of the paper.

Weaknesses:

This is not exactly a weakness, but I think it would future-proof the authors' conclusions to clarify a few key caveats associated with this work. Most notably, given the underlying implementation of the Mpipi model, temperature dependencies for intermolecular interactions driven by solvent effects (e.g., hydrophobic effect and charge-mediated interactions facilitated by desolvation penalties) are not captured. This itself is not a "weakness" per se, but it means I would imagine CERTAIN types of features would not be wellcaptured; notably, my expectation is that at higher temperatures, proline-rich sequences drive intermolecular interactions, but at lower temperatures, they do not. This is likely also true for the aliphatic residues, although these are found less frequently in IDRs. As such, it may be worth the authors explicitly discussing.

We also thank the reviewer for pointing out that a more detailed discussion of the model limitations is needed. The original Mpipi model was designed to probe UCST-type transitions (that are associative in nature) of disordered sequences. The reviewer is correct, that in its current form, the model does not capture LCST-type transitions that depend on changes in solvation of hydrophobic residues with temperature. We have amended the discussion to highlight this fact.

Similarly, prior work has established the importance of an alpha-helical region in TDP-43, as well as the role of aliphatic residues in driving TDP-43's assembly (see Schmidt et al 2019). I recognize the authors have focussed here on a specific set of mutations, so it may be worth (in the Discussion) mentioning [1] what impact, if any, they expect transient or persistent secondary structure to have on their conclusions and [2] how they expect aliphatic residues to contribute. These can and probably should be speculative as opposed to definitive.

Again - these are not raised as weaknesses in terms of this work, but the fact they are not discussed is a minor weakness, and the preprint's use and impact would be improved on such a discussion.

We agree with the reviewer that the effects of structural changes/propensities on these scaling behaviors would be an interesting and important angle to probe. We also comment on this in the discussion.

Reviewer # 2 (Public Review):

This is an interesting manuscript where a CA-only CG model (Mpipi) was used to examine the critical temperature (T_c) of phase separation of a set of 140 variants of prion-like low complexity domains (PLDs). The key result is that T_c of these PLDs seems to have a linear dependence on substitutions of various sticker and space residues. This is potentially useful for estimating the T_c shift when making novel mutations of a PLD. However, I have strong reservations about the significance of this observation as well as some aspects of the technical detail and writing of the manuscript.

We thank the reviewer for their thoughtful and detailed feedback on the manuscript.

(1) Writing of the manuscript: The manuscript can be significantly shortened with more concise discussions. The current text reads as very wordy in places. It even appears that the authors may be trying a bit too hard to make a big deal out of the observed linear dependence.

The manuscript needs to be toned done to minimize self-promotion throughout the text. Some of the glaring examples include the wording "unprecedented", "our research marks a significant milestone in the field of computational studies of protein phase behavior ..",

“Our work explores a new framework to describe, quantitatively, the phase behavior ...”, and others.

We thank the reviewer for their suggestions on the writing of the manuscript. We understand the concern regarding the length and tone of the manuscript, and in response to their feedback, we have revised the language throughout the manuscript.

There is really little need to emphasize the need to manage a large number of simulations for all 140 variants. Yes, some thoughts need to go into designing and managing the jobs and organizing the data, but it is pretty standard in computational studies. For example, large-scale protein ligand-free energy calculations can require one to a few orders of magnitude larger number of runs, and it is pretty routine.

We fully agree with the reviewer that this aspect of the study is relatively standard in computational research and does not require special emphasis. In response, we have revised the manuscript to shorten the aforementioned section, focusing instead on the scientific insights gained from the simulations rather than the logistical challenges of managing them.

When discussing the agreement with experimental results on T_m , it should be noted that the values of $R > 0.93$ and $RMSD < 14$ K are based on only 16 data points. I am not sure that one should refer to this as “extended validation”. It is more like a limited validation given the small data size.

We thank the reviewer for their consideration of our validation set. Indeed, the agreement with experimental results is based on 16 data points, as this set represents the available published data at the time of writing of this manuscript. The term “extended validation” is used to signify that our current dataset builds upon previous validations (in Joseph, Reinhardt et al. Nat Comput. Sci. 2021), incorporating additional variants not previously examined. The metrics of an $r > 0.93$ and a low RMSD indicate a strong agreement between the model and experiments, and an improvement with respect to other reported models. We are committed to continue validating our methods.

Results of linear fitting shown in Eq 4-12 should be summarized in a single table instead of scattering across multiple pages.

We considered the reviewer’s suggestion to compile all the laws into a single table. However, we believe it would be more effective for readers to reference each relationship directly where it is first discussed in the text. That said, we do include Table 1 in the original manuscript, which provides a summary of all the laws.

The title may also be toned down a bit given the limited significance of the observed linear dependence.

We respectfully disagree with the reviewer and believe that the current title accurately captures the scope of the manuscript.

(2) Significance and reliability of T_c : Given the simplicity of Mpipi (a CA-only model that can only describe polymer chain dimension) and the low complexity nature of PLDs, the sequence composition itself is expected to be the key determinant of T_c . This is also reflected in various mean-field theories. It is well known that other factors will contribute, such as patterning (examined in this work as well), residual structures, and conformational preferences in dilute and dense phases. The observed roughly linear dependence is a nice confirmation but really unsurprising by itself. It appears how many

of the constructs deviate from the expected linear dependence (e.g., Figure 4A) may be more interesting to explore.

While linear dependencies in critical solution temperatures may appear expected for certain systems, for example, symmetric hard spheres, the heterogeneity of intrinsically disordered regions (IDRs), like prion-like domains (PLDs), make this finding notable. The simplicity of our linear scaling law belies the underlying complexity of multivalent interactions and sequence-dependent behaviors in a certain sequence regime, which has not been quantitatively characterized in this manner before. Likewise, although linear dependencies may be expected in simplified models, the real-world applicability and empirical validation of these laws in biologically relevant systems are not guaranteed. Our chemically based model provides the robustness needed to do that. The linear relationship observed is significant because it provides a predictive framework for understanding how specific mutations affect a diverse set of PLDs. The framework presented can be extended to other protein families upon the application of a validated model, which might or might not yield linear relationships depending on the cooperative effects of their collective behavior. This extends beyond confirming known theories—it offers a practical tool for predicting phase behavior based on sequence composition.

We agree with the reviewer that, while the overarching linear trend is clear, deviations from linearity observed in constructs like those in Figure 4A point to additional, and interesting, layers of complexity. These deviations offer interesting avenues for future research and suggest that while linearity might dominate PLD critical behavior, other factors may modulate this behavior under specific conditions.

This is an excellent suggestion from the reviewer that, while it falls outside the scope of the current study, we are interested in exploring in the future.

Finally, the relationships are all linear, they have been normalized in different ways—the strength of the study also lies in that. Instead of focusing solely on linearity, our study explores the physical mechanisms that underlie these relationships. This approach provides a more complete understanding of how sequence composition and the underlying chemistry of the mutated residues influence $T_{c/sub}$.

The assumption that all systems investigated here belong to the same universality class as a 3D Ising model and the use of Eqn 20 and 21 to derive T_c is poorly justified. Several papers have discussed this issue, e.g., see Pappu Chem Rev 2023 and others. Muthukumar and coworkers further showed that the scaling of the relevant order parameters, including the conserved order parameter, does not follow the 3D Ising model. More appropriate theoretical models including various mean field theories can be used to derive binodal from their data, such as using Rohit Pappu's FIREBALL toolset. Imposing the physics of the 3D Ising model as done in the current work creates challenges for equivalence relationships that are likely unjustified.

We thank the reviewer for raising this point and for highlighting the FIREBALL toolset. Based on our understanding, FIREBALL is designed to fit phase diagrams using mean-field theories, such as Flory–Huggins and Gaussian Cluster Theory. Our experience with this toolset suggests that it places a higher weight on the dilute arm of the binodal. However, in our slab simulations, we observe greater uncertainty in the density of the dilute arm. This leads to only a moderate fit of the data to the mean-field theories employed in the toolset. While we agree that there is no reason to assume the phase behavior of these systems is fully captured by the 3D Ising model, we expect that such a model will describe the behavior near the critical point better than mean-field theories. Testing our results further with different critical exponents would be valuable in assessing how these predictions compare to a broader set of experimental data. Additionally, we have made the raw data points for the

phase diagrams available on our GitHub, enabling practitioners to apply alternative fitting methods.

While it has been a common practice to extract T_c when fitting the coexistence densities, it is not a parameter that is directly relevant physiologically. Instead, C_{sat} would be much more relevant to think about if phase separation could occur in cells.

While it is true that C_{sat} is directly relevant to whether phase separation can occur in cells under physiological conditions, T_c should not be dismissed as irrelevant. T_c provides fundamental insights into the thermodynamics of phase separation, reflecting the overall stability and strength of interactions driving condensate formation. This stability is crucial for understanding how environmental factors, such as temperature or mutations, might affect phase behavior. In Figure 2C and D we compare experimental C_{sat} values with our predicted T_c from simulations. These quantities are roughly inversely proportional to each other and so we expect that, to a first approximation, the relationships recovered for T_c should hold when considering C_{sat} at a fixed temperature.

Reviewer # 3 (Public Review):

Summary:

“Decoding Phase Separation of Prion-Like Domains through Data-Driven Scaling Laws” by Maristany et al. offers a significant contribution to the understanding of phase separation in prion-like domains (PLDs). The study investigates the phase separation behavior of PLDs, which are intrinsically disordered regions within proteins that have a propensity to undergo liquid-liquid phase separation (LLPS). This phenomenon is crucial in forming biomolecular condensates, which play essential roles in cellular organization and function. The authors employ a data-driven approach to establish predictive scaling laws that describe the phase behavior of these domains.

Strengths:

The study benefits from a robust dataset encompassing a wide range of PLDs, which enhances the generalizability of the findings. The authors’ meticulous curation and analysis of this data add to the study’s robustness. The scaling laws derived from the data provide predictive insights into the phase behavior of PLDs, which can be useful in the future for the design of synthetic biomolecular condensates.

We thank the reviewer for highlighting the importance of our work and for their critical feedback.

Weaknesses:

While the data-driven approach is powerful, the study could benefit from more experimental validation. Experimental studies confirming the predictions of the scaling laws would strengthen the conclusions. For example, in Figure 1, the T_c of TDP-43 is below 300 K even though it can undergo LLPS under standard conditions. Figure 2 clearly highlights the quantitative accuracy of the model for hnRNPA1 PLD mutants, but its applicability to other systems such as TDP-43, FUS, TIA1, EWSR1, etc., may be questionable.

In the manuscript, we have leveraged existing experimental data for the A1-LCD variants, extracting critical temperatures and saturation concentrations to compare with our model and scaling law predictions. We acknowledge that a larger set of experiments would be beneficial. By selecting sequences that are related, we hypothesize that the scaling laws described herein should remain robust. In the case of TDP-43, to our knowledge this protein

does not phase separate on its own under standard conditions. In vitro experiments that report phase separation at/above 300 K involve either the use of crowding agents (such as dextran or PEG) or multicomponent mixtures that include RNA or other proteins. Therefore, our predictions for TDP-43 are consistent with experiments. In general, we hope that the scaling laws presented in our work will inspire other researchers to further test their validity.

The authors may wish to consider checking if the scaling behavior is only observed for T_c or if other experimentally relevant quantities such as C_{sat} also show similar behavior. Additionally, providing more intuitive explanations could make the findings more broadly accessible.

In Figure 2C and D we compare experimental C_{sat} values with our predicted T_c from simulations. These quantities are roughly inversely proportional to each other and so we expect that, to a first approximation, the relationships recovered for T_c should hold when considering C_{sat} at a fixed temperature.

The study focuses on a particular subset of intrinsically disordered regions. While this is necessary for depth, it may limit the applicability of the findings to other types of phase-separating biomolecules. The authors may wish to discuss why this is not a concern. Some statements in the paper may require careful evaluation for general applicability, and I encourage the authors to exercise caution while making general conclusions. For example, "Therefore, our results reveal that it is almost twice more destabilizing to mutate Arg to Lys than to replace Arg with any uncharged, non-aromatic amino acid..." This may not be true if the protein has a lot of negative charges.

A significant number of proteins, in addition to those mentioned in the manuscript, that contain prion-like low complexity domains have been reported to exhibit phase separation behaviors and/or are constituents of condensates inside cells. We therefore expect these laws to be applicable to such systems and have further revised the text to emphasize this point. As the reviewer suggests, we have also clarified that the reported scaling of various mutations applies to these systems.

I am surprised that a quarter of a million CPU hours are described as staggering in terms of computational requirements.

We have removed the note on CPU hours from the manuscript. However, we would like to clarify that the amount of CPU hours was incorrectly reported. The correct estimate is 1.25 million hours, but this value was unfortunately misrepresented during the editing process. We thank the reviewer for catching this mistake on our part.

Reviewer # 1 (Recommendations For The Authors):

Some minor points here:

"illustrating that IDPs indeed behave like a polymer in a good solvent [43]. " Whether or not an IDP depends as a polymer in a good solvent depends on the amino acid sequence - the referenced paper selected a set of sequences that do indeed appear on average to map to a good-solvent-like polymer, but lest we forget SAXS experiments require high protein concentrations and until the recent advent of SEC-SAXS, your protein essentially needed to be near infinitely soluble to be measured. As such, this paper's conclusions are, apparently, ignorant of the limitations associated with the data they are describing, drawing sweeping generalizations that are clearly not supported by a multitude of studies in which sequence-dependencies have led to ensembles with a scaling exponent far below 0.59 (See Riback et al 2017, Peng et al 2019, Martin et al 2020, etc).

We thank the reviewer for raising this point. To avoid making incorrect generalizations and potentially misleading readers, we have removed the quoted statement from our manuscript.

As of right now, the sequences are provided in a convenient multiple-sequence alignment figure. However, it would be important also to provide all sequences in an Excel table to make it easy for folks to compare.

In addition to the sequence alignment figure, we now provide all tested sequences in an Excel table format in the GitHub repository.

Maybe I'm missing it, but it would be extremely valuable if the coexistence points plot in all the figures were provided as so-called source data; this could just be on the GitHub repository, but I'm envisaging a scenario where for each sequence you have a 4 column file where Col1=concentration and Col2=temperature, col3=fit concentration and col4=fit temperature, such that someone could plot col1 vs. col2 and col3 vs. col4 and reproduce the binodals in the various figures. Given the tremendous amount of work done to achieve binodals:

The coexistence points used to plot the figures are now provided in the GitHub, in a format similar to that suggested by the reviewer.

It would be nice to visually show how finite size effects are considered/tested for (which they are very nicely) because I think this is something the simulation field should be thinking about more than they are.

Thank you for highlighting this point. In our previous work (supporting information of the original Mpipi paper), we demonstrated a thorough approach by varying both the cross-sectional area of the box and the long axis while keeping the overall density constant. In this work, we verified that the cross-sectional area was larger than the average R_g of the protein. We then maintained a fixed cross-sectional area to long-axis ratio, varying the number of proteins while keeping the overall density constant. We have updated Appendix 1–Figure 2 to clarify our procedure and revised the caption to better explain how we ensured the number of proteins was adequate.

When explaining the law of reticular diameters, it would be good to explain where the 3.06 exponent comes from.

Based on the reviewer's suggestion, we have added to the text: "The constant 3.06 in the equation is a dimensionless empirical factor that was derived from simulations of the 3D Ising model."

The NCPR scale in Figure 5 being viridis is not super intuitive and may benefit from being seismic or some other r-w-b colormap just to make it easier for a reader to map the color to meaning.

We thank the reviewer for this suggestion and have replaced the scale with a r-w-b colormap.

The "sticker and spacer" framework has received critiques recently given its perceived simplicity. However, this work seems to clearly illustrate that certain types of residues have a large effect on T_c when mutated, whereas others have a smaller effect. It may be worth re-phrasing the sticker-spacer introduction not as "everyone knows aromatic/arginine residues are stickers" but as "aromatic and arginine residues have been proposed to be stickers, yet other groups have argued all residues matter equally"

and then go on to make the point that while a black-and-white delineation is probably not appropriate, based on the data, certain residues ARE demonstrably more impactful on Tc than others, which is the definition of stickers. With this in mind, it may be useful to separate out a sticker and a spacer distribution in Figure 1D, because the different distribution between the two residues types is not particularly obvious from the overlapping points.

We have revised the introduction of the sticker–spacer model in the manuscript for clarity. As the reviewer suggests, we have also separated the sticker and spacer distribution, which is now summarized in new Appendix 0–figure 8.

Reviewer # 3 (Recommendations For The Authors):

Figure 2 clearly highlights the quantitative accuracy of the model for hnRNPA1 PLD mutants, but its applicability to other systems such as TDP-43, FUS, TIA1, EWSR1, etc., may be questionable. The following sentence may be revised to reflect this: “Our extended validation set confirms that the Mpipi potential can ...”

Based on the reviewer’s suggestion, we have revised the text: “Our validation set, which expands the range of proteins variants originally tested [32], highlights that the Mpipi potential can effectively capture the thermodynamic behavior of a wide range of hnRNPA1-PLD variants, and suggests that Mpipi is adequate for proteins with similar sequence compositions, as in the set of proteins analyzed in this study. In recent work by others [66], Mpipi was tested against experimental radius of gyration data for 137 disordered proteins and the model produced highly accurate results, which further suggests the applicability of the approach to a broad range of sequences.”

<https://doi.org/10.7554/eLife.99068.2.sa0>