

Stumpy forms are the predominant transmissible forms of *Trypanosoma brucei*

Reviewed Preprint

v3 • May 15, 2025

Revised by authors

Reviewed Preprint

v2 • January 11, 2024

Reviewed Preprint

v1 • October 9, 2023

Jean Marc Tsagmo Ngoune, Parul Sharma, Aline Crouzols, Nathalie Petiot, Brice Rotureau 

Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France • Sorbonne Université, ED515 Complexité du Vivant, Paris, France • Parasitology Unit, Institut Pasteur of Guinea, Conakry, Guinea

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

African (or Salivarian) trypanosomes are significant pathogens of humans and domestic animals. For many decades it was accepted that only the "stumpy" non-proliferative form was capable of infecting the Tsetse-fly vector, but recent work challenged this, suggesting that the proliferative "slender" form is also infective. The current paper provides **important** and **convincing** laboratory evidence that the original concept is probably correct for most infections: the slender form was not infective for adult Tsetse, and was only able to infect young, less immunocompetent flies if N-acetyl glucosamine was added to the feed.

<https://doi.org/10.7554/eLife.91602.3.sa3>

Abstract

Schuster *et al.* demonstrated that bloodstream slender forms of African trypanosomes are readily transmissible to young tsetse flies where they can complete their complex life cycle (1). In their experimental conditions, a single slender parasite was sufficient for productive infection. Here, we compared the infectivity of slender and stumpy bloodstream forms in adult flies with a mature immune system, and without using any chemical compounds that would alter the insect immune response and/or promote the infection. After ingestion of slender forms, infected flies were observed only in one out of 24 batches of non-immunocompetent teneral flies and with a high number of parasites. In contrast, infected flies were detected in 75% (18/24) of the batches infected with stumpy parasites, and as few as 10 stumpy parasites produced mature infections in immune adult flies. We discuss that, although Schuster *et al.* have demonstrated the intrinsic capacity of slender form trypanosomes to infect young and naive tsetse flies, highlighting the remarkable plasticity and adaptability of these protists, this phenomenon is unlikely to significantly contribute to the epidemiology of African trypanosomiasis. According to both experimental and field observations, stumpy forms appear to be the most adapted forms for African trypanosome transmission from the mammalian host to the tsetse fly vector in natural conditions.

Introduction

Protist parasites of the *Trypanosoma brucei* group cause Human African Trypanosomiasis (HAT), or sleeping sickness in humans, and nagana in cattle (2). They are transmitted by the blood feeding tsetse fly following a long (at least 3 weeks) and complex (at least 9 distinct stages) cyclical development (review in (3)). In the mammalian host's blood circulation, proliferating slender trypanosomes differentiate into cell cycle-arrested stumpy cells upon quorum sensing when they reach high parasite densities (4–7). This differentiation is thought not only to regulate the parasite load in the reservoir host (8), but also to provide transmissible parasites adapted to pursue the life cycle in the vector host (9). Indeed, stumpy forms express several transcripts and proteins necessary to the next developmental stage in the insect, the procyclic form, including the Protein Associated to Differentiation 1 or PAD1 (5). For decades, arrest of the cell cycle and differentiation to the stumpy stage were presumed essential for the developmental progression of bloodstream trypanosomes to the insect stages.

Recently, Schuster *et al.* demonstrated that slender trypanosomes can also present some intrinsic characteristic of transmissible forms (PAD1 mRNAs and proteins) and are readily transmissible to both young male and female tsetse flies, where they can complete their complex life cycle (1), yet with a lower efficiency than stumpy forms (10). In their experimental conditions, a single slender parasite was sufficient for productive infection. These laboratory conditions are however significantly different from what is encountered in the field. First, only young teneral flies (1–3 days post-eclosion) with an immature immune system were used. Second, in some experiments, chemical compounds altering the insect immune response (glutathione) and/or promoting the infection (N-acetyl-glucosamine) were added to the infective meal. To assess the importance of these parameters, we challenged the infectivity of slender bloodstream forms in adult tsetse flies, i.e. in conditions closer to the natural situation.

Material and methods

Strains, culture and *in vitro* differentiation

The AnTat 1.1E Paris pleomorphic strain of *Trypanosoma brucei brucei* was derived from a strain originally isolated from a bushbuck in Uganda in 1966 (11). Bloodstream form trypanosomes were cultivated in HMI-9 medium supplemented with 10% (v/v) FBS (12) at 37°C in 5% CO₂. Proliferative slender cells were maintained at densities lower than 5.10⁵ parasites/ml to prevent their natural quorum-sensing-dependent differentiation into stumpy forms. For *in vitro* slender to stumpy BSF differentiation, we used 8-pCPT-2'-O-Me-5'-AMP, a nucleotide analogue of 5'-AMP (BIOLOG Life Science Institute, Germany). Briefly, 2×10⁶ pleomorphic AnTat 1.1E slender forms were incubated with 8-pCPT-2'-O-Me-5'-AMP (5 μM) for 48 h (13). Freshly differentiated stumpy forms and slender cells were then centrifuged at 1,400 × g for 10 minutes and resuspended at the appropriate densities in SDM-79 medium supplemented with 10% FBS. Cells were resuspended at either 10³, 10⁴ or 10⁵ parasites / ml. Assuming individual bloodmeal volumes ranging between 10–100 μl, this would correspond to ingestions of 10–100, 100–1,000 or 1,000–10,000 parasites per condition.

Tsetse fly maintenance, infection and dissection

Glossina morsitans morsitans tsetse flies were maintained in Roubaud cages at 27°C and 70% hygrometry and fed through a silicone membrane with fresh mechanically defibrinated sheep blood (BCL, France). Adult (between 2 and 3 weeks after emergence) or teneral males (between 24h and 72h post-emergence) were allowed to ingest parasites through a silicone membrane. No

chemical supplement was used in the first set of experiments. For assessing the effect of immunomodulatory compounds in the second set of experiments, 60mM *N*-acetylglucosamine were added to the infective meal. A total of 3 to 5 independent biological replicates per condition were performed with batches of 50 flies per condition.

Flies were starved for at least 24 hours before being dissected blindly 28 to 31 days post-ingestion for isolation of all stages from the midgut and salivary glands. For recovery of all tsetse organs, after rapid isolation of the salivary glands in a first drop of phosphate buffer saline (PBS), whole tsetse alimentary tracts, from the distal part of the foregut to the Malpighian tubules, were dissected and arranged lengthways in another drop of PBS as previously described (14, 15). Isolated organs were then scrutinized under a microscope at 40x magnification by two independent readers and infection rates per organ were scored (16).

Immuno-fluorescence analysis (IFA)

Cultured parasites were washed in TDB and spread onto poly-L-lysine coated slides. For flash methanol fixation, slides were air-dried for 10 min, fixed in methanol at -20°C for 5 seconds and rehydrated for 20 min in PBS. For immunodetection of stumpy forms, slides were incubated for 1 h at 37°C with a rabbit polyclonal anti-PAD1 antibody (kindly provided by Keith Matthews, University of Edinburgh) (5) diluted at 1:300 in PBS containing 0.1% Bovine Serum Albumin (BSA). After 3 consecutive 5 min washes in PBS, a species and subclass-specific secondary antibody coupled to the Alexa 488 fluorochrome (Jackson ImmunoResearch) diluted at 1:1000 in PBS containing 0.1% BSA was applied for 1 h at 37°C . After washing in PBS, slides were finally stained with 4',6-diamidino-2-phenylindole (DAPI, 1 $\mu\text{g}/\text{ml}$) for visualization of kinetoplast and nuclear DNA content and mounted under coverslips with ProLong antifade reagent (Invitrogen), as previously described (14). Slides were observed under an epifluorescence DMI4000 microscope (Leica) with a 100x oil objective (NA 1.4) to assess the proportion of PAD1-positive cells in the infective meals ($n > 100$ cells / condition).

Statistical analysis

Infections rates were compared by a two-sided ANOVA at 95% confidence with Prism V10.0.3 (GraphPad). MG infection rate comparisons were statistically significant between teneral and adult flies infected with ST in each amount ($p < 0.02$ with 10 parasites; $p < 0.0001$ with 100 and 1,000 parasites) and with 1,000 SL ($p < 0.0001$). MG infection rate comparisons were statistically significant ($p < 0.0001$) between parasite stages (SL and ST) in each amount (10, 100 and 1,000) and for each fly group (teneral and adult), excepted in teneral flies infected with 1,000 parasites ($p = 0.2356$).

Results

Pleomorphic *T. b. brucei* bloodstream forms were either maintained in culture at a density lower than 5.10^5 parasites / ml to prevent quorum-sensing-induced differentiation and obtain only slender forms or induced for differentiation with a 5'- AMP nucleotide analogue to obtain mostly stumpy forms. The expression of PAD1 at the cell surface was assessed by immunofluorescence analysis prior to each experimental infection: no PAD1 expression was detected in the slender group, whereas an average of 63% (52% to 71%, $n = 12$ replicates) of the induced cells were expressing PAD1. Batches of 50 teneral ($< 72\text{h}$) or adult (2-3 weeks) male tsetse flies were fed in parallel with either slender or stumpy forms at densities corresponding to individual ingestion of about 10, 100 or 1,000 parasites per bloodmeal. In total, 1,384 flies from 12 distinct experimental infections were dissected about 4 weeks (28 to 31 days) after parasite ingestion. Infection rates in midguts and salivary glands were quantified and plotted for each condition (Figure 1 and Table S1).




Figure 1.

Stumpy forms are the predominant transmissible forms.

Comparison of (A) midgut and (B) salivary gland infection rates in teneral (T) Vs. adult (A) tsetse flies (batches of 50 flies) infected with 10 to 100 (circles, 3 independent experiments), 100 to 1,000 (squares, 4 independent experiments) or 1,000 to 10,000 (triangles, 5 independent experiments) parasites in the slender (SL in blue) or stumpy (ST in red) forms.

After ingestion of slender forms, infected flies were observed in only 1 batch out of 24. This occurred in not yet fully immunocompetent teneral flies and with the highest number of ingested parasites (1,000 to 10,000 parasites). In contrast, midgut and salivary glands infected flies were observed in 75% (18/24) and 62.5% (15/24) of the batches infected with stumpy parasites, respectively. As few as 10 stumpy parasites produced mature infections in immunocompetent adult flies and the infection rates were similar whatever the amounts of stumpy forms ingested. In more susceptible non-immune teneral flies, the infection rates were increasing with the number of stumpy forms ingested.

Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster *et al.* study (1). Nevertheless, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

Discussion

The findings of Schuster *et al.* (1) have opened a debate on the traditional view of the trypanosome life cycle where slender trypanosomes are considered as non-competent for cyclical development in the insect vector (18). The authors proposed that their observations could provide a solution to a long-lasting paradox, namely the successful transmission of parasites in chronic infections, despite low parasitemia. Nonetheless, Schuster *et al.* performed all their experimental infections in laboratory conditions that were optimized for maximum transmission efficiency, with the use of teneral flies, and for some experiments, the addition of N-acetylglucosamine or glutathione (1).

Teneral flies are young flies that remain unfed for up to 3-5 days after emergence from their puparium. They are known to be significantly more susceptible to trypanosome infection as multiple studies demonstrated they are immunologically immature (weak immune system and, leaky peritrophic matrix) (19–22). According to capture-recapture studies (23), teneral flies however represent a minority of individuals in wild tsetse populations. Hence, knowing that adults can live up to 9 months (24), the impact of teneral flies on trypanosome transmission may be limited, if not incidental.

In addition, Schuster *et al.* supplemented most infective meals with 60 mM N-acetylglucosamine, an inhibitor of tsetse midgut lectins (17) that was also confirmed to enhance trypanosome infectivity in teneral flies in the present study. For infections with monomorphic parasites, the addition of 12.5 mM glutathione, an antioxidant that reduces the midgut environment, protected trypanosomes from cell death induced by reactive oxygen species (25). In total, these two chemical compounds used in Schuster *et al.* (1) have inhibited the immune response in teneral flies and substantially enhanced the chances for slender trypanosomes to develop in the insect vector.

Conditions are far less favourable in the field; hence we investigated and compared the infection potential of slender and stumpy forms in adult and teneral flies without the addition of chemicals. We observed that slender forms were not infective to adult tsetse flies and only at densities higher than 10^5 parasites / ml. Nevertheless, in endemic areas, especially in Western Africa, parasitemia in confirmed cases are usually very low ($< 10^4$ parasites / ml in Guinea for instance) and it is necessary to concentrate parasites in blood prior to microscopic examination to increase

sensitivity of parasitological diagnosis (26). Hence, the possible transmission of a few slender trypanosomes from the blood of individuals with a chronic infection is unlikely to explain the maintenance of the parasite circulation in tsetse populations.

By contrast, we observed that as few as 10 stumpy parasites are enough to produce mature infections in both teneral and adult flies, already with a significant efficiency. In patients with low parasitemia, the quorum-sensing-triggered differentiation of slender to stumpy forms could be compatible with, or even more adapted to, extravascular forms present in some tissues and organs, with a limited dilution of the parasites and parasite factors remaining concentrated locally. Indeed, extravascular PAD1-positive trypanosomes were detected in high numbers at least in adipose tissues (26) and in the dermis (27) of experimentally infected mice. The presence of PAD1-positive extravascular trypanosomes was also assessed in the skin of confirmed gambiense HAT cases and unconfirmed seropositive individuals in endemic areas (28) (and unpublished data). This suggests that stumpy trypanosomes accessible to tsetse flies are likely more abundant than previously estimated in individuals with low parasitemia.

Schuster *et al.* have demonstrated the intrinsic capacity of slender form trypanosomes to infect young and naive tsetse flies, highlighting the remarkable plasticity and adaptability of these protists. The fine understanding of the underlying cellular mechanisms and / or transient adaptations involved in this process remains an exciting challenge. This event is however unlikely to contribute to the epidemiology of African trypanosomiasis in natural settings. According to both experimental and field observations, stumpy forms appear to be the most adapted forms for African trypanosome transmission from the mammalian host to the tsetse fly vector in natural conditions.

Acknowledgements

We thank K. Matthews for providing the anti-PAD1 antibody and P. Bastin for his critical reading of the manuscript.

Additional information

Funding

This work was supported by the Institut Pasteur, the Programme Investissement d'Avenir of the French Government, Laboratoire d'Excellence, ANR-10-LABX-62- IBEID and ANR-11-LABX-0024-PARAFRAP. This work was supported the French National Agency for Scientific Research projects ANR-18-CE15-0012 TrypaDerm and ANR-19-CE15-0004-02 AdipoTryp. None of these funding sources has a direct scientific or editorial role in the present study.

Author contributions

AC, NP, JMTN and PS performed the experiments. BR designed the study, analysed the data, and wrote the manuscript. JMTN, PS and BR discussed the manuscript.

Additional files

Sup Table 1

References

1. Schuster S, Lisack J, Subota I, Zimmermann H, Reuter C, Mueller T, et al. (2021) **Unexpected plasticity in the life cycle of *Trypanosoma brucei*** *eLife* **10**
2. Buscher P, Cecchi G, Jamonneau V, Priotto G. (2017) **Human African trypanosomiasis** *Lancet* **390**:2397–409
3. Rotureau B, Van Den Abbeele J. (2013) **Through the dark continent: African trypanosome development in the tsetse fly** *Frontiers in cellular and infection microbiology* **3**:53
4. Vassella E, Reuner B, Yutzy B, Boshart M. (1997) **Differentiation of African trypanosomes is controlled by a density sensing mechanism which signals cell cycle arrest via the cAMP pathway** *J Cell Sci* **110**:2661–71
5. Dean S, Marchetti R, Kirk K, Matthews KR (2009) **A surface transporter family conveys the trypanosome differentiation signal** *Nature* **459**:213–7
6. Mony BM, MacGregor P, Ivens A, Rojas F, Cowton A, Young J, et al. (2014) **Genome-wide dissection of the quorum sensing signalling pathway in *Trypanosoma brucei*** *Nature* **505**:681–5
7. Rojas F, Silvester E, Young J, Milne R, Tetley M, Houston DR, et al. (2019) **Oligopeptide Signaling through TbGPR89 Drives Trypanosome Quorum Sensing** *Cell* **176**:306–17
8. Turner CM, Aslam N, Dye C. (1995) **Replication, differentiation, growth and the virulence of *Trypanosoma brucei* infections** *Parasitology* **111**:289–300
9. Rico E, Rojas F, Mony BM, Szoor B, Macgregor P, Matthews KR (2013) **Bloodstream form pre-adaptation to the tsetse fly in *Trypanosoma brucei*** *Frontiers in cellular and infection microbiology* **3**:78
10. Matthews KR, Larcombe S. (2022) **Comment on ‘Unexpected plasticity in the life cycle of *Trypanosoma brucei*’** *eLife* **11**
11. Le Ray D, Barry JD, Easton C, Vickerman K. (1977) **First tsetse fly transmission of the “AnTat” serodeme of *Trypanosoma brucei*** *Ann Soc Belg Med Trop* **57**:369–81
12. Hirumi H, Hirumi K. (1989) **Continuous cultivation of *Trypanosoma brucei* blood stream forms in a medium containing a low concentration of serum protein without feeder cell layers** *J Parasitol* **75**:985–9
13. Barquilla A, Saldivia M, Diaz R, Bart JM, Vidal I, Calvo E, et al. (2012) **Third target of rapamycin complex negatively regulates development of quiescence in *Trypanosoma brucei*** *Proc Natl Acad Sci U S A* **109**:14399–404
14. Rotureau B, Subota I, Bastin P. (2011) **Molecular bases of cytoskeleton plasticity during the *Trypanosoma brucei* parasite cycle** *Cell Microbiol* **13**:705–16

15. Rotureau B, Subota I, Buisson J, Bastin P. (2012) **A new asymmetric division contributes to the continuous production of infective trypanosomes in the tsetse fly** *Development* **139**:1842–50
16. Rotureau B, Blisnick T, Subota I, Julkowska D, Cayet N, Perrot S, et al. (2014) **Flagellar adhesion in *Trypanosoma brucei* relies on interactions between different skeletal structures in the flagellum and cell body** *J Cell Sci* **127**:204–15
17. Peacock L, Ferris V, Bailey M, Gibson W. (2006) **Multiple effects of the lectin-inhibitory sugars D-glucosamine and N-acetyl-glucosamine on tsetse-trypanosome interactions** *Parasitology* **132**:651–8
18. Guegan F, Figueiredo L. (2021) **A two-stage solution** *eLife* **10**
19. Wijers DJ (1958) **Factors that may influence the infection rate of *Glossina palpalis* with *Trypanosoma gambiense*. I. The age of the fly at the time of the infected feed** *Ann Trop Med Parasitol* **52**:385–90
20. Walshe DP, Lehane MJ, Haines LR (2011) **Post eclosion age predicts the prevalence of midgut trypanosome infections in *Glossina*** *PLoS ONE* **6**:e26984
21. Aksoy S, Gibson WC, Lehane MJ (2003) **Interactions between tsetse and trypanosomes with implications for the control of trypanosomiasis** *Adv Parasitol* **53**:1–83
22. Weiss BL, Wang J, Maltz MA, Wu Y, Aksoy S. (2013) **Trypanosome infection establishment in the tsetse fly gut is influenced by microbiome-regulated host immune barriers** *PLoS Pathog* **9**:e1003318
23. Hargrove JW (1990) **Age-Dependent Changes in the Probabilities of Survival and Capture of the Tsetse, *Glossina-Morsitans-Morsitans* Westwood** *Insect Sci Appl* **11**:323–30
24. Challier A. (1982) **The Ecology of Tsetse (*Glossina* Spp) (Diptera, Glossinidae) - a Review (1970-1981)** *Insect Sci Appl* **3**:97–143
25. MacLeod ET, Maudlin I, Darby AC, Welburn SC (2007) **Antioxidants promote establishment of trypanosome infections in tsetse** *Parasitology* **134**:827–31
26. Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-Branco F, et al. (2016) ***Trypanosoma brucei* Parasites Occupy and Functionally Adapt to the Adipose Tissue in Mice** *Cell Host Microbe*
27. Capewell P, Cren-Travaille C, Marchesi F, Johnston P, Clucas C, Benson RA, et al. (2016) **The skin is a significant but overlooked anatomical reservoir for vector-borne African trypanosomes** *eLife* **5**
28. Camara M, Soumah AM, Ilboudo H, Travaille C, Clucas C, Cooper A, et al. (2021) **Extravascular Dermal Trypanosomes in Suspected and Confirmed Cases of gambiense Human African Trypanosomiasis** *Clin Infect Dis* **73**:12–20

Author information

Jean Marc Tsagmo Ngoune

Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France

Parul Sharma

Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France, Sorbonne Université, ED515 Complexité du Vivant, Paris, France

Aline Crouzols

Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France

Nathalie Petiot

Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France

Brice Rotureau

Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Institut Pasteur, Université Paris Cité, INSERM U1347, Paris, France, Parasitology Unit, Institut Pasteur of Guinea, Conakry, Guinea
ORCID iD: [0000-0003-0671-8999](https://orcid.org/0000-0003-0671-8999)

For correspondence: brice.rotureau@pasteur.fr

Editors

Reviewing Editor

Christine Clayton

Centre for Molecular Biology of Heidelberg University (ZMBH), Heidelberg, Germany

Senior Editor

Dominique Soldati-Favre

University of Geneva, Geneva, Switzerland

Reviewer #2 (Public review):

Summary:

In contrast to the recent findings reported by Schuster S et al., this brief paper presents evidence suggesting that the stumpy form of *T. brucei* is likely the most pre-adapted form to progress through the life cycle of this parasite in the tsetse vector.

Strengths:

One significant experimental point is that all fly infection experiments are conducted in the absence of "boosting" metabolites like GlcNAc or S-glutathione. As a result, flies infected with slender trypanosomes present very low or nonexistent infection rates. This provides important experimental evidence that the findings of Schuster S and colleagues may need to be revisited.

In the revised submission the authors also compared trypanosome midgut infection levels in tsetse flies when either young (teneral) or mature adult flies received infected bloodmeals,

with or without 60 mM GlcNAc. The data clearly show that, unlike in teneral flies, the addition of GlcNAc to the trypanosome-infected bloodmeal does not enhance midgut infection in mature adult flies. This is now convincingly demonstrated in Figure 2 and provides strong experimental support for the suggestion that the effect reported by Schuster S. et al. may have been influenced by both fly age and the inclusion of metabolic "boosters" in the bloodmeal.

<https://doi.org/10.7554/eLife.91602.3.sa2>

Reviewer #3 (Public review):

The dogma in the Trypanosome field is that transmission by Tsetse flies is ensured by stumpy forms. This has been recently challenged by the Engstler lab (Schuster et al.), who showed that slender forms can also be transmitted by teneral flies. In this work, the authors aimed to test whether transmission by slender forms is possible and frequent. The authors observed that most stumpy forms infections with teneral and adult flies were successful while only 1 out of 24 slender form infections were successful.

The comparison of midgut infection in adult vs teneral flies was significant in most of the conditions. However, the critical comparison is still missing: within each type of fly (adult or teneral), was the MG infection significantly different between slender and stumpy forms?

Figure 2 convincingly demonstrates the effect of the metabolite N-acetylglucosamine on Tsetse infection. This addition helps better integrate the study with previous work. I thank the authors for their effort in performing this experiment.

It still remains unknown why this work and Schuster et al. reached different conclusions. As a result it remains unclear in which conditions slender forms could be important for transmission. Several variables could explain differences between the two groups: the strain used, the presence or absence of glutathione, how Tsetse colonies were maintained, thorough molecular and cellular characterisation of slender and stumpy forms (to avoid using intermediate forms as slender forms), comparison to recent field parasite strains.

<https://doi.org/10.7554/eLife.91602.3.sa1>

Author response:

The following is the authors' response to the previous reviews

eLife Assessment

For decades it has been accepted that only the growth-arrested "stumpy" form of Trypanosoma brucei can infect the arthropod vector, the Tsetse fly, but this was recently challenged by a demonstration that - under artificial conditions that are known to enhance infectivity - the proliferative "slender" form can also establish Tsetse infections. The infectiousness of the two forms is a fundamental question in trypanosome biology and epidemiology, concerning both infection dynamics and parasite differentiation. The authors of the current study provide compelling evidence that without artificial enhancement, the "stumpy" form is indeed much more infective for Tsetse than the slender form; they suggest that this is probably also true in the wild.

Since the authors of this paper did not themselves test the effect of enhancing conditions, the precise reason for the discrepancy in results between the two laboratories has not been demonstrated conclusively.

This specific comment was addressed in the revision and illustrated with new data.

Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

Public Reviews:

Reviewer #1 (Public Review):

Summary:

Ngoune et al. present compelling evidence that Slender cells are challenged to infect tsetse flies. They explore the experimental context of a recent important paper in the field, Schuster et al., that presents evidence suggesting the proliferative Slender bloodstream T.brucei can infect juvenile tsetse flies. Schuster et al. was disruptive to the widely accepted paradigm that the Stumpy bloodstream form is solely responsible for tsetse infection and T.brucei transmission potential. Evidence presented here shows that in all cases, Stumpy form parasites are exponentially more capable of infecting tsetse flies. They further show that Slender cells do not infect mature flies.

However, they raise questions of immature tsetse immunological potential and field transmission potential that their experiments do not address. Specifically, they do not show that teneral tsetse flies are immunocompromised, that tsetse flies must be immunocompromised for Slender infection nor that younger teneral tsetse infection is not pertinent to field transmission.

All these specific comments were addressed in the revision and illustrated with new data and references.

- The limited immunocompetence of teneral flies has been extensively studied by the labs of S. Aksoy at Yale and M. Lehane at Liverpool. In the discussion, we provide key references from these two labs 19-22.
- Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).
- Our comment on the relevance to field transmission is simply based on field observations of the fly biology. For example, according to the capture-recapture experiments described in HARGROVE JW insect sci applic 1990 (new ref 23), wild female mortality was reported 6.8% shortly after emergence, <1% for ages 20-50 days and rose to 5% by 130 day (a pattern similar

to that for laboratory reared tsetse), while wild male daily mortality was 8.3% after emergence, fell to 5.5% by 9 days, then rose continuously to more than 10% by 30 days. This means that adult flies represent the majority of individuals in a wild tsetse population. Hence, knowing that both males and females are strictly hematophagous and that they can live up to nine months, the impact of teneral flies (up to 4 days after emergence) on trypanosome transmission appears limited, if not incidental.

Strengths:

Experimental Design is precise and elegant, outcomes are convincing. Discussion is compelling and important to the field. This is a timely piece that adds important data to a critical discussion of host:parasite interactions, of relevance to all parasite transmission.

Thank you

Weaknesses:

As above, the authors dispute the biological relevance of teneral tsetse infection in the wild, without offering evidence to the contrary. Statements need to be softened for claims regarding immunological competence or relevance to field transmission.

All these specific comments were addressed in the revision and illustrated with new data and references.

- The limited immunocompetence of teneral flies has been extensively studied by the labs of S. Aksoy at Yale and M. Lehane at Liverpool. In the discussion, we provide key references from these two labs 19-22.

- Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

- Our comment on the relevance to field transmission is simply based on field observations of the fly biology. For example, according to the capture-recapture experiments described in HARGROVE JW insect sci applic 1990 (new ref 23), wild female mortality was reported 6.8% shortly after emergence, <1% for ages 20-50 days and rose to 5% by 130 day (a pattern similar to that for laboratory reared tsetse), while wild male daily mortality was 8.3% after emergence, fell to 5.5% by 9 days, then rose continuously to more than 10% by 30 days. This means that adult flies represent the majority of individuals in a wild tsetse population. Hence, knowing that both males and females are strictly hematophagous and that they can live up to nine months, the impact of teneral flies (up to 4 days after emergence) on trypanosome transmission appears limited, if not incidental.

Reviewer #2 (Public Review):

Summary:

*In contrast to the recent findings reported by Schuster S et al., this brief paper presents evidence suggesting that the stumpy form of *T. brucei* is likely the most pre-adapted form to progress through the life cycle of this parasite in the tsetse vector.*

Strengths:

One significant experimental point is that all fly infection experiments are conducted in the absence of "boosting" metabolites like GlcNAc or S-glutathione. As a result, flies infected with slender trypanosomes present very low or nonexistent infection rates. This provides important experimental evidence that the findings of Schuster S and colleagues may need to be revisited.

Thank you

Weaknesses:

However, I believe the authors should have included their own set of experiments demonstrating that the presence of these metabolites in the infectious bloodmeal enhances infection rates in flies receiving blood meals containing slender trypanosomes. Considering the well-known physiological variabilities among flies from different facilities, including infection rates, this would have strengthened the experimental evidence presented by the authors.

This specific comment was addressed in the revision and illustrated with new data.

Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

Reviewer #3 (Public Review):

The dogma in the Trypanosome field is that transmission by Tsetse flies is ensured by stumpy forms. This has been recently challenged by the Engstler lab (Schuster et al.), who showed that slender forms can also be transmitted by teneral flies. In this work, the authors aimed to test whether transmission by slender forms is possible and frequent. The authors observed that most stumpy forms infections with teneral and adult flies were successful while only 1 out of 24 slender form infections were successful.

In this revised version of the manuscript, the authors made some text changes and included statistical testing as a new section of the Materials and Methods. It seems the comparison of midgut infection in adult vs teneral flies was significant in most of the conditions. However, the critical comparison is still missing: within each type of fly (adult or teneral), was the MG infection significantly different between slender and stumpy forms?

An ANOVA statistical analysis was performed and a dedicated section added to the revised version. MG infection rate comparisons were statistically significant between teneral and adult flies infected with ST in each amount ($p < 0.02$ with 10 parasites; $p < 0.0001$ with 100 and

1,000 parasites) and with 1,000 SL ($p < 0.0001$). MG infection rate comparisons were statistically significant ($p < 0.0001$) between parasite stages (SL and ST) in each amount (10, 100 and 1,000) and for each fly group (teneral and adult), excepted in teneral flies infected with 1,000 parasites ($p = 0.2356$).

Given no additional experiments were performed, it remains unknown why this work and Schuster et al. reached different conclusions. As a result it remains unclear in which conditions slender forms could be important for transmission. Several variables could explain differences between the two groups: the strain used, the presence or absence of N-acetylglucosamine and/or glutathione, how Tsetse colonies were maintained, thorough molecular and cellular characterisation of slender and stumpy forms (to avoid using intermediate forms as slender forms), comparison to recent field parasite strains.

This specific comment was addressed in the revision and illustrated with new data.

Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

The manuscript is improved, but the author has not addressed much of the constructive criticism offered that would benefit the manuscript.

To clarify, evidence from Schuster et al did not demonstrate, rather it suggested. That is a major point of this paper - that the previous evidence presented had caveats. Terms such as demonstrate or prove are inappropriate in most biological contexts, unless evidence is without caveat.

This specific comment was addressed in the revision and illustrated with new data.

Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

Statements regarding teneral flies in the field are softened. Yet the referenced papers pertain more to commensurate coinfections rather than reduced immunocapacity of immature teneral flies in the field. This should be clarified.

The limited immunocompetence of teneral flies has been extensively studied by the labs of S. Aksoy at Yale and M. Lehane at Liverpool. In the discussion, we provide key references from these two labs 19-22.

The text remains convoluted to read with grammatical errors in places. For example, it is incorrect to begin a sentence with However. There are far too many run-on sentences in the manuscript that confuse this straightforward story.

The revised text was improved as much as possible.

All text requires grammatical refinement and softer claims unless additional experiments are undertaken.

Reviewer #2 (Recommendations For The Authors):

I continue to endorse the publication of this manuscript; however, I am somewhat disappointed by the authors' justifications for not conducting additional experiments or exploring other factors that might influence the infection phenotypes in the fly.

This specific comment was addressed in the revision and illustrated with new data.

Differences between the strain clones, the cell culture conditions and/or the fly colony maintenance conditions could explain part of the differences in infection rates observed here as compared to the Schuster et al. study (1). However, the use of the lectin-inhibitory sugar N-acetyl-glucosamine to enhance infection rates in the latter study could be a more likely explanation. To assess this hypothesis, an additional experimental challenge was performed to compare infection rates in teneral versus adult flies, with or without N-acetyl-glucosamine supplement in an infective meal containing 10^5 slender parasites / ml (Figure 2). Whereas no infection was detected in adult flies, the N-acetyl-glucosamine supplementation of the infective meal led to an increase of the infection rates from 2,4% to 13,3% in teneral flies (Figure 2).

<https://doi.org/10.7554/eLife.91602.3.sa0>