

Molecular dynamics of the matrisome across sea anemone life history

Reviewed Preprint

v2 • September 26, 2025

Revised by authors

Reviewed Preprint

v1 • April 2, 2025

B Gideon Bergheim, Alison G Cole, Mandy Rettel, Frank Stein, Stefan Redl, Michael W Hess, Aissam Ikmi, Suat Özbek 

University of Heidelberg, Centre for Organismal Studies, Department of Evolutionary Neurobiology, Heidelberg, Germany • Department of Neurosciences and Developmental Biology, Faculty of Life Sciences, University of Vienna, Vienna, Austria • European Molecular Biology Laboratory, Heidelberg, Germany • Institute of Neuroanatomy, Medical University of Innsbruck, Innsbruck, Austria • Innsbruck Medical University, Innsbruck, Austria

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **valuable** study provides a comprehensive description of the *Nematostella vectensis* matrisome - the genes encoding the proteins of the extracellular matrix. The authors combine new mass spectrometry data with bioinformatic analyses of previously published genomic and single-cell RNAseq data. The analysis is thorough, and the discussion and conclusions are **convincing**. This work will be of interest to biologists working on the evolution of the matrisome, as well as more broadly those working with non-bilaterian animals.

<https://doi.org/10.7554/eLife.105319.2.sa2>

Abstract

The evolutionary expansion of extracellular matrix (ECM) molecules has been crucial for the establishment of cell adhesion and the transition from unicellular to multicellular life. Members of the early diverging metazoan phylum cnidaria offer an exceptionally rich perspective into the metazoan core adhesome and its original function in developmental and morphogenetic processes. Here, we present the ensemble of ECM proteins and associated factors for the starlet sea anemone *Nematostella vectensis* based on *in silico* prediction and quantitative proteomic analysis of decellularized mesoglea from different life stages. The integration of the matrisome with single cell transcriptome atlases shows that gastrodermal cells are the primary producers of *Nematostella*'s complex ECM, confirming the homology of the cnidarian inner cell layer with bilaterian mesoderm. The transition from larva to polyp is marked by an upregulation of metalloproteases and basement membrane components including all members of an unusually diversified SVEP1/Polydom family, suggesting massive epithelial remodeling. The enrichment of Wnt/PCP pathway factors during this process further indicates directed cell rearrangements as a key contributor to the polyp's morphogenesis. Mesoglea maturation in adult polyps involves wound response proteins indicating shared molecular patterns in growth and regeneration. Our study identifies conserved matrisomal networks that coordinate transitions in *Nematostella*'s life history.

Introduction

The evolution of extracellular matrix, a complex network of secreted, typically modular proteins, is closely linked to the emergence of metazoan life forms (Rokas 2008 [↗](#); Hynes 2009 [↗](#); Ozbek, et al. 2010 [↗](#); Hynes 2012 [↗](#); Naba 2024 [↗](#)). While some ECM components, such as integrins and cadherin receptors can be traced back to unicellular organisms (Sebe-Pedros, et al. 2010 [↗](#); Nichols, et al. 2012 [↗](#)), early diverging cnidarians (hydroids, jellyfish, corals, and sea anemones) are believed to possess one of the most complete adhesomes among non-bilaterian clades (Ozbek, et al. 2010 [↗](#); Tucker and Adams 2014 [↗](#)). Cnidarians, the sister group to bilaterians, are characterized by a simple body plan with a central body cavity and a mouth opening surrounded by tentacles. They are diploblastic organisms, consisting of an outer epithelium and an inner gastrodermis separated by a complex ECM called mesoglea (Sarras 2012; Bergheim and Ozbek 2019 [↗](#)). The mesoglea, which is best studied in the freshwater polyp *Hydra*, forms a flexible, tri-laminar structure. It is composed of a central, amorphous interstitial matrix (IM) interspersed with collagenous filaments, sandwiched between two thin layers of basement membrane (BM) (Sarras, et al. 1991; Shimizu, et al. 2008 [↗](#); Aufschnaiter, et al. 2011 [↗](#); Bergheim and Ozbek 2019 [↗](#)). Studies in *Hydra* have shown that the mesoglea can be separated intact from the epithelial cell sheets by a freeze-thaw technique (Day and Lenhoff 1981 [↗](#); Veschgini, et al. 2023 [↗](#)). Previously, we analyzed the proteome of decellularized *Hydra* mesoglea and identified 37 unique protein sequences (Lommel, et al. 2018 [↗](#)), including most of the described core matrisome components (Sarras 2012). Among medusozoans (jellyfish and hydroids), hydras stand out for having lost the free-swimming medusa form. They also lack a planula larva stage from which anthozoans (corals and sea anemones) typically produce sessile polyps. We therefore hypothesized that a cnidarian species with a complex life cycle could offer a more comprehensive picture of the non-bilaterian ECM repertoire. Here, we analyzed the matrisome of the anthozoan starlet sea anemone *Nematostella vectensis*, by employing *in silico* predictions of ECM proteins that were partially confirmed by a subsequent proteomic analysis of decellularized mesoglea. We detected a rich collection of matrisome proteins, comparable in its core matrisome complexity to vertebrate species (Naba, et al. 2012 [↗](#)). Furthermore, mapping of our matrisome data onto a previously established single-cell transcriptome dataset (Steger, et al. 2022 [↗](#); Cole, et al. 2024 [↗](#)) revealed a prominent role of the gastroderm in ECM production. Cnidocytes, which produce the cnidarian stinging organelle, are characterized by a distinct set of ECM proteins that significantly contribute to the complexity of the cnidarian matrisome. Quantitative proteomics of mesoglea samples from different life stages (larva, primary and adult polyp) showed that while the larval mesoglea contained only few exclusively enriched factors, the transition from the larval stage to primary polyp was marked by an upregulation of a large fraction of the matrisome. This set of proteins included many metalloproteases and basement membrane factors, indicating significant epithelial reorganization. Remarkably, all members of an unusually diverse SVEP1/Polydom family were upregulated during this morphogenetic process, implicating a conserved role of this protein family for epithelial morphogenesis. Additionally, a significant enrichment of Wnt/planar cell polarity (PCP) signaling components, such as ROR2 and protocadherin Fat4, supports that directed cell movements underlie the axial elongation and morphogenesis of the polyp (Stokkermans, et al. 2022 [↗](#)). The final transition to the adult animal involves an increased addition of elastic fiber components to the mesoglea and matricellular factors associated with wound healing, indicating common ECM-associated mechanisms in regeneration, growth and tissue differentiation.

Results

Molecular composition and structure of the *Nematostella* ECM

To investigate the components and dynamics of the ECM throughout various life stages of *Nematostella vectensis*, we employed a protocol for obtaining decellularized mesoglea, originally developed for *Hydra* (Day and Lenhoff 1981 [↗](#); Lommel, et al. 2018 [↗](#); Veschgini, et al. 2023 [↗](#)). We isolated mesogleas from larvae at three days post-fertilization (3 dpf), primary polyps at 10 dpf, and from small adult polyps that were at least 1 year old. Protein extraction was performed under strongly reducing conditions at high temperatures (90°C) to solubilize the cross-linked protein network of the ECM. The extracted proteins were digested with trypsin and analyzed using quantitative mass spectrometry (Fig. 1A [↗](#)). Extensive studies in both vertebrates and invertebrates have identified specific characteristics of ECM proteins, typically based on conserved domains and domain arrangements (Engel 1996 [↗](#); Hohenester and Engel 2002 [↗](#); King, et al. 2008 [↗](#); Ozbek, et al. 2010 [↗](#); Naba, et al. 2012 [↗](#); Tucker and Adams 2014 [↗](#)). Building on this knowledge and a *de novo* annotation of all predicted *Nematostella* protein models using InterProScan (Jones, et al. 2014 [↗](#)), we identified 1,812 potential ECM proteins (Fig. 1B [↗](#)). These were further classified into orthogroups based on similarity using OrthoFinder (Emms and Kelly 2019 [↗](#)). To refine this analysis, we also predicted *in silico* matrisomes from protein models of several early-branching metazoan species, including two choanoflagellates, two sponges, three ctenophores, 9 additional representative cnidarians, and two placozoan species. SignalP (Teufel, et al. 2022 [↗](#)) was used to confirm the presence of signal peptides, while DeepLoc (Thummuluri, et al. 2022 [↗](#)) was employed to predict the cellular localization of the matrisome draft. Additionally, we explored the closest BLAST hits in the NCBI and SwissProt databases. After manually excluding duplicates and sequencing artifacts, we narrowed down the number of high-confidence ECM genes defined by the possession of a *bona fide* ECM domain (Hynes and Naba 2012 [↗](#)) to 829 (Fig. 1B [↗](#)). Each protein sequence was manually reviewed, with annotations assigned based on domain predictions, UniProt identifiers, and comparisons to known ECM proteins. Our subsequent mass spectrometry analysis initially identified a total of 5,286 proteins. However, we suspected that a significant portion might be cellular contaminants originating from residual amoebocytes within the mesoglea (Tucker, et al. 2011 [↗](#)), which could not be completely removed during the isolation procedure. To address this, we used our curated *in silico* list of 829 candidate ECM genes as a filter and identified 287 ECM components in the isolated *Nematostella* mesogleas of which 210 belonged to the matrisome (Fig. 1B [↗](#), supplementary tables S1, S2). This number exceeds the previously reported 37 matrisomal proteins in *Hydra* (Lommel, et al. 2018 [↗](#)), suggesting a greater compositional diversity in *Nematostella*. A comparison of mesoglea samples of both species resolved by protein gel electrophoresis confirmed the higher complexity of the *Nematostella* mesoglea, particularly in the lower molecular weight fraction (supplementary figure S1). The discrepancy between the number of predicted ECM proteins and those confirmed by mass spectrometry can be attributed to the exclusion of nematocysts from the mesoglea isolates, despite their substantial contribution to the matrisome (as shown below). Furthermore, soluble factors and transmembrane receptors are likely underrepresented in the mesoglea isolates. We further organized our curated dataset following the classification proposed by Hynes and Naba, who introduced the concept of a “core” matrisome, characterized by 55 signature InterPro domains, including EGF, LamG, TSP1, vWFA, and collagen (Hynes and Naba 2012 [↗](#)). Within this core matrisome, we identified 241 proteins, including collagens, proteoglycans, and ECM glycoproteins (e.g., laminins (Fahey and Degan 2012 [↗](#)) and thrombospondins (Tucker, et al. 2013 [↗](#); Shoemark, et al. 2019 [↗](#))) (Fig. 1C [↗](#), supplementary table S1). Additionally, we identified a set of 310 “matrisome-associated” factors. This group includes molecules that (i) have structural or functional associations with the core matrisome, (ii) are involved in ECM remodeling (e.g., metalloproteases), or (iii) are secreted proteins, including growth factors. In summary, the *Nematostella* matrisome comprises 551 proteins, representing approximately 3% of its proteome (Artamonova and Mushegian 2013 [↗](#)) and roughly half the size of the human matrisome with 1,056 proteins (Naba, et al. 2012 [↗](#)). While the *Nematostella* core matrisome is comparable to that of bilaterians, vertebrate species exhibit a dramatic expansion in ECM-associated factors (Fig. 1D [↗](#)). The remaining non-matrisomal 278 proteins, identified through negative selection based

on exclusive domain lists for each ECM category (Naba, et al. 2012 [DOI](#)), were assigned to the “adhesome” category primarily comprising transmembrane adhesion receptors (e.g., cadherins and IgCAM-like molecules) or categorized as “other” ECM domain proteins including proteins with specialized functions, such as venoms or stress and injury response proteins (**Fig. 1B,C** [DOI](#), Table S1). Additionally, 17 proteins could not be confidently assigned to any category but were retained in the dataset as candidates for future functional characterization.

To validate the preservation of the isolated ECM, we generated polyclonal antibodies targeting unique peptide sequences in laminin gamma 1 (anti-Lam), type IV collagen NvCol4b (anti-Col4), a specific fibrillar collagen NvCol2c (anti-Col2), and a consensus motif for several *Nematostella* fibrillar collagens (anti-PanCol) (supplementary figure S2). We then performed immunofluorescence confocal microscopy on *Nematostella* whole mounts and decellularized mesoglea from all life stages (**Fig. 1E-K** [DOI](#)). Consistent with recent findings in *Hydra* (Veschgini, et al. 2023 [DOI](#)), the *in vivo* and *ex vivo* images exhibited similar patterns along the polyp’s body, indicating the structural integrity of the mesoglea after decellularization (**Fig. 1F-K** [DOI](#)). In cross-sections, the laminin antibody stained the thin double layer of the basement membrane (BM) lining the two epithelia, while the PanCol antibody detected the intervening fibrous layer of the interstitial matrix (IM) (**Fig. 1E** [DOI](#)). Both antibodies also showed diffuse staining at the apical surface of ectodermal cells, likely due to the sticky nature of the glycocalyx. Ultrastructural examination of the mesoglea revealed a mean thickness of approximately 0.5 μm in larvae and 1.5 μm in primary polyps (supplementary figure S3A-B, G-H, supplementary table S3). In the triangular areas at the base of the gastrodermal mesenteric folds, the mesoglea was expanded (supplementary figure S3C-F). The BM lining the epithelia of larvae was very delicate, measuring only 70 nm, and the IM appeared as a loose array of thin fibrils (supplementary figure S3B). Primary polyps possessed a distinct BM, with a thickness of approximately 130 nm, appearing as a dense meshwork of fibrils (supplementary figure S3H, supplementary table S3). Previous reports (Tucker, et al. 2011 [DOI](#)) indicated that the IM in older primary polyps was interspersed with thin fibrils of about 5 nm and extended thick fibrils of about 20-25 nm. Our samples from younger primary polyps (supplementary figure S3H) showed thin fibrils of approximately 6 nm and thick ones of about 13 nm, with occasional fibrils of up to 27 nm (supplementary table S3). Immunostainings with the laminin antibody revealed a thickened mesoglea at the aboral pole of the polyp, forming an unusual knot-like structure (supplementary figure S3J). Ultrastructural analysis showed that the fibrils in this region were densely packed and aligned along the oral-aboral axis (supplementary figure S3K-L), possibly providing a rigid attachment site for the mesentery retractor muscles. Immunoelectron microscopy confirmed the observations from immunofluorescence: anti-Lam immunogold labeling was primarily localized along the plasma membrane of ECM-lining cells, anti-Col4 labeling was found at the BM (supplementary figure S3M-N), and anti-PanCol labeling was predominantly distributed throughout the IM, with a similar pattern observed for the anti-Col2 antibody (supplementary figure S3O-P).

Cell type specificity of matrisome expression

Recently, single-cell RNA sequencing has been applied to identify the origin of neuroglandular cell lineages in *Nematostella* (Steger, et al. 2022 [DOI](#)), hypothesize on the origin of muscle cell types (Cole, et al. 2023 [DOI](#)), and catalog the distribution of cell states associated within all tissue types (Cole, et al. 2024 [DOI](#)). We made use of this developmental cell type atlas to determine the cell type specificity of matrisomal gene expression. Expression profiles for all ECM genes across the entire life cycle are available in the supplementary data (supplementary table S4). We calculated an average expression score for each of the ECM gene sets (core and associated matrisome, adhesome/other) and found above average scores for the core matrisome associated with the mature gastrodermis and developing cnidocytes, and to a lesser extent also for the other two categories (**Fig. 2** [DOI](#)). Core matrisome genes also showed additional high expression scores within an uncharacterized gland cell type (GD.1), matrisome-associated genes within the digestive gland set, and adhesome/other genes within the maturing cnidocytes (**Fig. 2A** [DOI](#)). We looked specifically at the distribution of core matrisome genes across all cell states and generated a list of




Figure 1.

Analysis of the *Nematostella* matrisome.

(A) Mesoglea from larvae, primary polyps and adults was decellularized and analyzed by mass spectrometry. In parallel, an *in silico* matrisome was predicted using a computational approach, and curated manually. (B-C) 1812 potential ECM proteins were predicted bioinformatically. The manually curated list of ECM factors consists of 829 proteins. The curated ECM proteins were sorted into core matrisome and matrisome-associated groups, which together constitute the *Nematostella* matrisome (551 proteins). The remaining non-matrisomal proteins are categorized as “adhesome” that include transmembrane receptors, and “other” ECM domain proteins, which include adhesive proteins, venoms, enzymes, ion channels, stress and injury response factors and diverse uncharacterized proteins (see supplementary table S1 for detailed annotations and sub-categories). In total 287 ECM proteins were confirmed by mass spectrometry analysis, 210 of which belong to the matrisome and 47 to the “adhesome”. (D) Comparison of the *Nematostella* matrisome size with published matrisomes of other species. While the complexity of the *Nematostella* core matrisome is comparable to that of vertebrates, the number of ECM-associated proteins is disproportionally lower. The *Drosophila* core matrisome is characterized by significant secondary reduction. (E) Laminin antibody stains the bilaminar structure of the BL (magenta) at the base of the epithelial cell layers, while the pan-Collagen antibody (yellow) detects the central IM. Scale bar, 10 µm. The three life stages of *Nematostella* before (F-H) and after (I-K) decellularization. The mesoglea is stained with Laminin antibody to demonstrate its structural preservation and by DAPI (cyan) to visualize residual nuclei and nematocysts. The decellularized mesoglea retains morphological structures such as tentacles (t) and mesenteries (m). Scale bars: F, G, I, J, 100 µm; H, K, 1 mm.

differentially up-regulated genes (supplementary table S5, supplementary figure S4). Of note is the absence of any differentially expressed core matrisome factors within the ectodermal tissues, and contrastingly a large set that are specific to either the mesoendodermal inner cell layer (gastrodermis) or cnidocytes (**Fig. 2B**). We also find sets of core matrisome genes that are specific to different secretory gland cell types, including mucin-producing, digestive-enzyme producing, and uncharacterized S2-class cell types. Interestingly, the gland cell-specific matrisome genes include several of the Polydom family members upregulated during larva-to-primary polyp transition (see below). Altogether these expression profiles suggest that core components of the mesoglea (collagens, laminins) are produced from the inner cell layer, and that a large set of ECM glycoproteins and all of the minicollagens (David, et al. 2008; Zenkert, et al. 2011) have been recruited into the formation of the cnidarian synapomorphy, the cnidocytes.

Collagens constitute the primary structural components of the animal ECM and, being a highly diverse family of triple helical proteins (Fidler, et al. 2018), form a significant part of the core matrisome. Vertebrate collagens consist of 28 types (I-XXVIII) categorized as fibril-forming, network or beaded filament-forming, and transmembrane collagens (Kadler, et al. 2007). Type IV collagen, a major constituent of basement membranes, is considered to be a primordial component of the animal ECM based on studies in ctenophores that lack fibrillar collagens but display a remarkable diversity of collagen IV genes (Fidler, et al. 2017; Draper, et al. 2019). Our analysis revealed 12 bilaterian-type collagens (**Fig. 2C**) with conserved C-terminal non-collagenous trimerization domains (NC1) and extended triple helical stretches of ~1000 residues, including two sequences for type IV collagen (NvCol4a/b) together with a peroxidase homolog (NV2.13306), indicating the presence of sulfilimine cross-links that stabilize the *Nematostella* BM (Fidler, et al. 2014). In *Hydra*, which lacks this specific post-translational modification, six collagen genes have been identified through cDNA cloning and proteomic analysis (Deutzmann, et al. 2000; Fowler, et al. 2000; Zhang, et al. 2007; Lommel, et al. 2018). We have classified the collagens according to the *Hydra* nomenclature (Zhang, et al. 2007) and identified three isoforms of NvCol1, each comprising an isolated minor triple helical domain at the N-terminus (**Fig. 2C**, upper panel). In addition, we detected four NvCol2 paralogs, which contain an additional whey acidic protein 4 disulfide core (WAP) domain at the N-terminus. However, our dataset lacks a Hcol3-like collagen with alternating N-terminal WAP and vWFA domains, although it includes a protein with extensive WAP/vWFA repeats but lacking a collagen triple helix or NC1 domain (NV2.11346). NvCol5 consists of a continuous collagen domain with two minor interruptions following the signal peptide and is otherwise similar to NvCol1. We have not found sequences resembling Hcol6, which is characterized by triple helical sequences interrupted by multiple vWFA domains. Collagen XVIII-like, which is predicted to contain an unrelated Mucin-like insertion, includes an N-terminal Laminin-G/TSPN motif followed by a discontinuous central collagenous domain, similar to the BM-associated vertebrate $\alpha 1$ (XVIII) collagen chains (Heljasvaara, et al. 2017) and *Drosophila* Multiplexin (Meyer and Moussian 2009). NvCol7 shares a similar domain organization but differs by having only a single interruption of the triple helix near the N-terminus. According to Exposito et al., the *Nematostella* fibrillar collagen sequences are phylogenetically related to A clade collagens that in mammals possess a vWFC module in their N-propeptide supposed to have evolved from the cnidarian WAP domain (Exposito, et al. 2008). NvCol7 is an exception and belongs to B clade collagens characterized by the possession of an N-terminal Laminin-G/TSPN domain.

Most of the *Nematostella* collagens show a broad expression in diverse gastrodermal cell populations throughout all life stages (supplementary figure S5). NvCol2c is an exception, as it is predominantly expressed in neuronal cells of larvae indicating a specialized function in neurogenesis. NvCol7 is distinguished by showing an expression both in gastrodermal cells and two small cnidocyte cell populations described by Steger et al. to be exclusive for planula larvae (Steger, et al. 2022). Unlike the IM collagens, NvCol4a is additionally expressed during embryogenesis, emphasizing the pivotal role of the BM in organizing the epithelial tissue architecture during early development. In addition to these bilaterian-type collagens, we have




Figure 2.

Single cell atlas of core matrisome genes.

(A) Dimensional reduction cell plot (UMAP) highlighting cell clusters showing over-abundant expression of the core matrisome, matrisome-associated, and adhesome/other gene sets. Expression values correspond to gene module scores for each set of genes. (B) Dotplot expression profiles of upregulated genes of the core matrisome across cell type partitions, separated across phases of the life cycle. Illustrated are the top 5 genes with expression in at least 20% of any cell state cluster, calculated to be upregulated with a p -value of ≤ 0.001 . See supplementary table S5 for a full list of differentially expressed core matrisome genes. Larva = 18hr:4day samples; Primary Polyp = 5:16 day samples; Adult = tissue catalog from juvenile and adult specimens. (C) *Nematostella* collagens. Domain organization of matrisome proteins containing a collagen triple helix as core element. The proteins are categorized into fibrillar and basement membrane collagens, short-chain collagens and nematocyte-specific minicollagens.

identified four spongin-like proteins, which are short-chain collagens with derived NC1 domains (Fig. 2C, middle panel). These molecules have been described as truncated variants of collagen IV with a wide distribution in several invertebrate phyla (Aouacheria, et al. 2006). Except for NvSpongin-like-2, their expression is not aligned with that of fibrillar collagens, but is mostly restricted to neuroglandular cells (supplementary figure S5). An additional set of “collagen-like” sequences that comprise short triple helical stretches without additional domain motifs (Fig. 2C, middle panel) is broadly expressed across cell types and developmental stages (supplementary figure S5, supplementary table S4). In contrast, the large set of minicollagens (Fig. 2C, lower panel) is strictly aligned with the cnidocyte lineage as detailed below.

A specialized cnidocyte matrisome

Cnidocytes, the stinging cells of cnidarians, are a key synapomorphy of this clade. Previous studies have demonstrated that, from a transcriptomic perspective, nematocyst capsule formation is distinct from the mature profile (Chari, et al. 2021; Steger, et al. 2022). We filtered the dataset of the 829 curated ECM protein-coding sequences for genes expressed within cnidocytes and found 298 genes that are detectable above average in at least 5% of any cnidocyte transcriptomic state (supplementary figure S6, supplementary table S6). We further separated this list of genes into those that are absent from non-cnidocyte states (101 genes: ‘exclusive’) or are also detected within more than 50% of the non-cnidocyte transcriptomic states (41 genes: ‘ubiquitous’) (Fig. 3A). We consider the remaining genes shared across cnidocyte and non-cnidocyte profiles (156: ‘shared’). Of the cnidocyte-specific genes, we examined the proportion of genes specific to either the capsule-building specification profiles (79: ‘specification’) or the maturation phase of cnidogenesis (24: ‘maturation’). Most of these genes are restricted to the specification phase, with a smaller subset associated with the mature transcriptomic profile (Fig. 3A). The latter group includes several members of a vastly expanded family of Fibrinogen-related proteins (FREPs) (supplementary table S1), which have been implicated in innate immunity across various phyla (Zhou, et al. 2024) and may function as venom components. In the ‘shared’ gene set, most genes are associated with the mature cnidocyte profile and overlap with various neuroglandular subtypes (supplementary figure S6). This observation supports the hypothesis that the cnidocytes arose from an ancestral neuronal population (Richards and Rentzsch 2014; Tournière, et al. 2020). We then calculated a gene module score for each gene set to estimate specificity across the dataset, summarizing the specificity of each gene set (Fig. 3B). Further examination of the gene lists reveals cnidocyte specificity of nematogalectins and minicollagens that serve as structural components for cnidocysts (Kurz, et al. 1991; Hwang, et al. 2010). Minicollagens, that have served as important phylum-specific genes (Holland, et al. 2011), comprise a short collagen domain (about 15 Gly-X-Y repeats) flanked by proline-rich regions and terminal cysteine-rich domains (CRDs) (David, et al. 2008). We identified all five previously described *Nematostella* minicollagen sequences (Zenkert, et al. 2011; Steger, et al. 2022) (NvCol-1, -3, -4, -5, -6) along with four additional proteins with incomplete minicollagen sequence features, which we termed “minicollagen-like” (Fig. 2C). Intriguingly, we also identified a protein that combines features of both minicollagens and extended ECM-type collagens, suggesting a possible evolutionary origin of minicollagens from this gene family. This protein, NvNCol-7, contains a minicollagen pro-peptide sequence, proline-rich regions, and canonical N- and C-terminal CRDs (Tursch, 2016 #1245). Unlike previously described minicollagens, it includes an extended discontinuous collagen sequence of ~1000 residues, comprising 25 alternating blocks of mostly 12 or 15 Gly-X-Y repeats. These blocks are interrupted by either a single alanine or MPP/SPPSPP sequences, resembling degenerated collagen triplets. The presence of a minicollagen pro-peptide in this collagen suggests its expression in the cnidocyte lineage and secretion into the nematocyst vesicle as a structural component of cnidocyst walls or tubules (Adamczyk, et al. 2010; Garg, et al. 2023). This is confirmed by the single cell expression data, which show prominent and exclusive expression in cnidocyte lineages (supplementary figures S5, S6). Interestingly, *Nematostella* minicollagens exhibit differential expression across different cnidocyte subtypes, nematocytes and spirocytes (Fig. 3C). Spirocytes express NvCol-5 and NvNcol-like-3 and 4, while NvCol-1, 4 and 6 and NvNCol-7 are expressed within the nematocytes and NvCol-3 is expressed in both. Other cnidocyte-

specific genes also show differential paralog expression between cnidocyte types, including the four NOWAs (Engel, et al. 2002 [↗](#); Garg, et al. 2023 [↗](#)), and the vWFA domain proteins (**Fig. 3C** [↗](#)). NvTSR-2 vs NvTSR-3 distinguish between the two nematocyte lineages, nem.1 and nem.2. These are postulated to be basitrichous haplonemas/isorhizas based on abundance and distribution across the adult tissue libraries, although these identities have not been validated. In summary, a significant fraction (32 %) of matrisomal genes (176 of 551, supplementary table S6) are expressed within the cnidocyte lineage. The proportion of cnidarian-specific factors in this subset is increased (85 of 176, 48 %) as compared to the full matrisome (234 of 551, 42%, supplementary table S1), supporting the notion that the cnidocyte proteome represents a specialized ECM adapted to the unique assembly process and biophysical requirements of the cnidarian stinging organelle (Ozbek 2011 [↗](#); Balasubramanian, et al. 2012 [↗](#)).

Larva-to-polyp transition is marked by factors of basement membrane remodeling and Wnt/PCP signaling

Larva-polyp morphogenesis in *Nematostella* involves significant changes in body shape including the elongation of the body axis and the development of oral tentacles and internal mesenteries (Stokkermans, et al. 2022 [↗](#)). We performed quantitative proteomics using tandem mass tag labeling (TMT) (see supplementary figure S7 for normalization steps) to examine whether this process is accompanied by stage-specific variations of matrisome components. As shown above, 38 % (210 of 551) of the matrisomal factors were detected by our mass spectrometry analysis (**Fig. 1B** [↗](#)). We identified stage-specific mesoglea components by assigning hits (>2-fold change and <0.05 false discovery rate (fdr)) using a modified t-test limma for each of the three life stage comparisons (supplementary table S7). Globally, collagens and basement membrane proteoglycans constitute the bulk of the ECM in the core matrisome across all life stages while secreted factors are the most abundant subcategory among matrisome-associated proteins (**Fig. 4A** [↗](#)). As illustrated in the heat map in **Fig. 4B** [↗](#), the transition from larvae to primary polyp is characterized by a general increase of mesoglea components. 94 proteins were differentially upregulated in primary polyps while only four, including vitellogenin and its receptor that are crucial for lipid transport from the ECM into the oocyte (Lebouvier, et al. 2022 [↗](#)), were differentially abundant in larvae. The proteins enriched in primary polyps as compared to larvae include, in addition to various factors involved in general cell adhesion (e.g., cadherin-1, coadhesin-like proteins), two major functional groups (**Fig. 4C** [↗](#), supplementary table S7): (1) factors involved in BM establishment and remodeling and (2) components of the Wnt/PCP signaling pathway. Both groups are indicative of a massive epithelial rearrangement, rather than cell proliferation, as a driver of larva-to-polyp transition (Stokkermans, et al. 2022 [↗](#)). Notably, in addition to laminins, the BM proteoglycan perlecan and numerous astacin, ADAMTS and MMP family metalloproteases (see supplementary figure S9 for an overview), the first group contained all members of an unusually expanded Polydom protein family (**Fig. 4E** [↗](#)). Polydom/SVEP1 is a secreted multidomain ECM protein initially discovered in a murine bone-marrow stromal cell line (Gilges, et al. 2000 [↗](#)). In humans, it is composed of eight different domains including an N-terminal vWFA domain, followed by a cysteine-rich ephrin receptor motif, Sushi/CCP and hyalin repeat (HYR) units, EGF-like domains, a central pentraxin domain and a long tail of 28 Sushi domains terminating in three EGF repeats (**Fig. 4E** [↗](#)). Polydom has recently been shown to be a ligand for the orphan receptor Tie1 and induce lymphatic vessel remodeling via PI3K/Akt signaling (Sato-Nishiuchi, et al. 2023 [↗](#)). Earlier studies have shown basement membrane deposition of Polydom and a role in epithelial cell-cell adhesion via integrin binding (Sato-Nishiuchi, et al. 2012 [↗](#); Samuelov, et al. 2017 [↗](#)). The *Hydractinia* homolog has been characterized as a factor specific to i-cells and to be potentially involved in innate immunity (Schwarz, et al. 2008 [↗](#)). In comparison to vertebrate Polydoms, *Hydractinia* Polydom contains additional Pan/Apple, FA58C, and CUB domains, but has a reduced number of terminal Sushi repeats (**Fig. 4E** [↗](#)). In our study, we identified a *Nematostella* Polydom homolog (NvPolydom1) that shares high similarity with the *Hydractinia* and *Hydra* proteins, suggesting a conserved arrangement of domains in cnidarians




Figure 3.

Cell-type specificity of cnidocyte-expressed ECM genes.

A) The distribution of cnidocyte-expressed genes categorized as 'ubiquitous' (blue: 41), 'shared' (red: 27), 'mature-specific' (green: 38), or 'specification-specific' (purple: 88). (B) Expression of the module scores of each gene-subset across the main tissue-type data partitions, illustrated on UMAP dimensional reduction. (C) Sequential gene expression activation illustrated on a dotplot of top 5 differentially expressed genes (p -value ≤ 0.001) for each cnidocyte cell state. Nematocyte specification shares many genes, while spirocyte specification uses a distinct gene set. Nep.8, nematocyst-expressed protein 8 categorized as venom protein.

(Fig. 4E [↗](#)). This includes a pentraxin-PAN/Apple-FA58C core structure, a tail consisting of six Sushi repeats, and 1-2 terminal CUB domains. Two additional paralogs, NvPolydom2 and NvPolydom3, exhibit differences in their central region, with NvPolydom2 lacking the PAN/Apple domain and NvPolydom3 lacking both PAN/Apple and FA58C domains. Furthermore, we discovered four shorter Polydom-like sequences that share a common EGF-Sushi-HYR-TKE structure but lack vWFA and Pentraxin domains, as well as the terminal Sushi repeats. These Polydom-like proteins, resembling a truncated N-terminal part of canonical Polydoms, possess additional domains at the N- or C-termini, such as thrombospondin type-1 repeat (TSR) or Ig-like domains. The shortest member, referred to as Polydom-related, lacks EGF-like modules and consists of the central Sushi, HYR, and TKE domains found in all Polydom-like sequences, suggesting that this core motif may be essential for biological function. All Polydom and Polydom-like paralogs are expressed within the putative digestive cell state GD.1, with two of these showing additional expression within the uncharacterized secretory cell type S2.tll.2&3 (supplementary figure S8). This, together with the exceptionally high isoform diversity, indicates a requirement for genetic robustness to account for perturbations of the developmental process regulated by the *Nematostella* Polydoms. Given that mouse Polydom is essential for endothelial cell migration in a Tie-dependent manner ([Sato-Nishiuchi, et al. 2023 \[↗\]\(#\)](#)), it is plausible that the *Nematostella* homologs could serve a similar function for rearrangement of epithelial cells along the BM during primary polyp morphogenesis. Interestingly, the top differentially abundant factor in the primary polyp mesoglea is a secreted integrin- α -related protein (sIntREP) containing three integrin- α N-terminal domains followed by a stretch of EGF repeats. It is an attractive hypothesis that sIntREP modulates Integrin-dependent cell adhesion by Polydoms and other factors to facilitate cell migration. The second group of proteins that contains several components of the Wnt/PCP pathway, including ROR2, protocadherin Fat4-like, and hedgling, further supports the assumption of epithelial cell migration as a main driver of primary polyp morphogenesis. Wnt/PCP signaling, originally described in *Drosophila melanogaster* ([Gubb and Garcia-Bellido 1982 \[↗\]\(#\)](#); [Adler 2002 \[↗\]\(#\)](#)), has a well-established role in convergent extension movements of cells during gastrulation to facilitate the elongation of the embryo along its oral-aboral axis ([Gao 2012 \[↗\]\(#\)](#)). Its “core” factors include the transmembrane proteins Frizzled, Van Gogh/Strabismus, Flamingo, and the intracellular components Dishevelled, Prickle, and Diego ([Simons and Mlodzik 2008 \[↗\]\(#\)](#)). An additional level of PCP within tissues is regulated by the unusually large protocadherins Fat and Dachous that interact in a heterophilic manner and display cellular asymmetries ([Matakatsu and Blair 2004 \[↗\]\(#\)](#)). Recently, the *Hydra* Fat-like homolog has been reported to be polarized along the oral-aboral axis and to organize epithelial cell alignment via organization of the actin cytoskeleton ([Brooun, et al. 2020 \[↗\]\(#\)](#)). Hedgling is an ancestral, non-bilaterian member of the cadherin superfamily with high similarity to FAT and Flamingo cadherins ([Adamska, et al. 2007 \[↗\]\(#\)](#)). It has a gastrodermal expression in the *Nematostella* primary polyp ([Adamska, et al. 2007 \[↗\]\(#\)](#)), which it shares with Wnt5a and ROR2 (supplementary table S4). ROR2, a highly conserved receptor tyrosine kinase, is the principal transducer of PCP signaling via Wnt5a in the *Xenopus* embryo ([Hikasa, et al. 2002 \[↗\]\(#\)](#); [Schambony and Wedlich 2007 \[↗\]\(#\)](#)). It has also been reported to induce directional cell movements in mammals ([He, et al. 2008 \[↗\]\(#\)](#)) and to induce filopodia formation as a prerequisite for directed cell migration ([Nishita, et al. 2006 \[↗\]\(#\)](#)). Taken together, the molecular dynamics of the mesoglea revealed by quantitative mass spectrometry suggests a massive epithelial rearrangement and directed cell migration as the main morphogenetic process underlying axial elongation during primary polyp morphogenesis. Interestingly, IM components such as fibrillar collagens do not appear to contribute largely to this process. In contrast, the adult mesoglea is significantly enriched in elastic fiber components, such as fibrillins and fibulin. This compositional shift likely adds to the visco-elastic properties ([Gosline 1971a \[↗\]\(#\)](#), [b \[↗\]\(#\)](#)) of the growing body column (Fig. 4B,D [↗](#), supplementary table S7). In addition, the adult mesoglea contains several matricellular factors associated with different aspects of wound healing in vertebrate organisms ([Cárdenas-León, et al. 2022 \[↗\]\(#\)](#)). These include SPARC-related follistatin domain proteins, uromodulin, and periostin. The *Nematostella* uromodulin gene was previously reported to be highly upregulated in the wound ectoderm, likely contributing to the innate immune response ([DuBuc, et al. 2014 \[↗\]\(#\)](#)). The same study showed a circular upregulation of the MMP inhibitor NvTIMP around the wound site.

Indeed, while metalloproteases are similarly upregulated in adults and primary polyps, we observed a noticeable increase of diverse classes of protease inhibitors in adults (7 vs 2 in primary polyps), including TIMP, Kunitz and Kazal-type protease inhibitors, as well as thyroglobulin repeat proteins (Novinec, et al. 2006 [DOI](#)) (supplementary table S7). This is indicative of a high degree of protease activity regulation in tissue morphogenesis during growth as also observed during *Nematostella* whole-body regeneration (Schaffer, et al. 2016 [DOI](#)). Taken together, the transition from the primary polyp to the adult is characterized by an increase of elastic fibrillar IM components contributing to the long-range elasticity and resilience of the mesoglea, and a recruitment of wound response factors that together with a complex network of proteases and protease inhibitors likely regulate growth, organogenesis and tissue differentiation.

Discussion

The evolution of the ECM, a complex proteinaceous network that connects cells and organizes their spatial arrangement in tissues, has been a key innovation driving the emergence of multicellular life forms (Rokas 2008 [DOI](#); Brunet and King 2017 [DOI](#)). Although ctenophores have recently been identified as the sister group to all other animals (Schultz, et al. 2023 [DOI](#)), the hitherto available genomic evidence suggests that they possess only a minimal repertoire of conserved ECM and ECM-affiliated proteins, limiting comparative studies with bilaterians (Draper, et al. 2019 [DOI](#)). In contrast, cnidarian genome data have offered broad evidence for conserved matrisomes anticipating the complexity of mammalian species (Tucker and Adams 2014 [DOI](#)). Here, we identified 551 ECM proteins that comprise the matrisome of the sea anemone *Nematostella vectensis* through *in silico* prediction using transcriptome databases and analyzed the dynamics of 287 ECM factors by TMT labeling and LC-MS/MS of decellularized mesoglea samples from different life stages. Utilizing cell-type specific atlases we showed that the inner gastroderm is the major source of the *Nematostella* ECM, including all 12 collagen-encoding genes. This finding supports the model of germ layer evolution proposed by Steinmetz et al. (Steinmetz, et al. 2017 [DOI](#)) where, based on both transcription factor profiles and structural gene sets, the cnidarian inner cell layer (endoderm) is homologous to the bilaterian mesoderm that gives rise to connective tissues. It also contrasts the situation in *Hydra* where both germ layers contribute to the synthesis of core matrisome proteins (Epp, et al. 1986 [DOI](#); Zhang, et al. 2007 [DOI](#)). The primacy of the gastrodermis in ECM synthesis might be related to the anthozoan-specific mesenteries, which represent extensions of the mesoglea into the body cavity sandwiched by two endodermal layers. Discrete endo- and ectodermal ECM transcript repertoires would result in a restricted composition of the mesoglea in these mesenteric folds. Whether a gastroderm-based matrisome represents an ancestral state of the cnidarian phylum can only be resolved by the inclusion of omics data from a larger diversity of cnidarian species. While anthozoans such as *Nematostella* have previously been considered a basal group among cnidarians (Bridge, et al. 1992 [DOI](#); Miller, et al. 2000 [DOI](#)), more recent phylogenomic studies consider them as a sister clade to medusozoans (DeBiasse, et al. 2024 [DOI](#)). To evaluate the complexity of *Nematostella*'s matrisome across cnidarians and other metazoan phyla, we plotted matrisome sizes from published databases and newly generated *in silico* matrisomes of representative species against orthogroup counts (Fig. 5 [DOI](#)). For the *in silico* matrisomes, only orthogroups shared with at least one found in published matrisomes were counted. Interestingly, anthozoan species generally exhibit a higher complexity than medusozoans and populate a transitory region between bilaterians and non-bilaterians in the evolutionary trajectory. This indicates that the acquisition of complex life cycles in medusozoa, that are distinguished by the pelagic medusa stage, was not accompanied by a diversification of the matrisome repertoire. This is in line with findings from genome data in *Aurelia*, a cnidarian with a medusa stage, that questioned the hypothesis of the medusozoan body-plan being derived. Rather, the authors found a redeployment of the existing genetic repertoire (Gold, et al. 2019 [DOI](#)). The reduced complexity of the medusozoan ECM might therefore represent a strategy to minimize the cost for ECM remodeling during metamorphosis and rely on a restricted set of conserved genes to form the expanded jellyfish mesoglea.




Figure 4.

Mesoglea dynamics across life stages assessed by quantitative proteomics of isolated mesoglea.

(A) Boxplot representation of normalized log₂ TMT reporter ion intensities for different protein subgroups of the matrisome. 'All' represents all proteins in each respective dataset. Horizontal line indicates median TMT intensity in the complete dataset. (B) 2-log transformed median abundances of proteins across different life stages. The curated ECM proteins were filtered for proteins with a 2-fold change in any of the life stages and a false discovery rate of 0.05 using a moderated t-test (limma). The heatmap shows the 2-log transformed median abundance of 4 samples per life stage. Most proteins are upregulated in only one of the life stages. Notably, BM factors including all polydoms are upregulated in the primary polyp. Most ECM protein categories can be clearly divided into adult specific and primary polyp specific proteins underscoring the differential composition of the mesoglea at different life stages. (C-D) Volcano plots showing the differential abundance of proteins in the mesoglea extracts of the three different life stages. (C) Proteins involved in BM organization including all polydoms and in Wnt/PCP signaling are upregulated during larva-to-primary polyp transition as highlighted. (D) The adult mesoglea compared to primary polyps is characterized by an enrichment of elastic fibril components and matricellular glycoproteins involved in wound response and regeneration. gray = non-matrimonial background, orange = insignificant, magenta = differentially abundant matrisome proteins. (E) Domain organization of bilaterian and cnidarian polydoms. The *Nematostella* matrisome contains an expanded group of polydoms and polydom-like proteins, including three cnidarian-type polydom paralog, four shorter polydom-like sequences and a polydom-related protein, which contains only the core Sushi-HYR-TKE motif. Domain symbols: vWFA (light blue), EGF-like (purple), Sushi/SCR/CCP (orange), Hyalin repeat (red), Pentraxin (yellow), CUB (light green), Tyrosine-protein kinase ephrin (dark green), PAN/Apple (olive green), Ricin B-like (pink), Thrombospondin type-1 repeat (brown), Coagulation factor 5/8 (dark purple), Ig-like (dark grey).




Figure 5.

Matrisome complexity across metazoan phyla.

Matrisome sizes of published and newly generated *in silico* matrisomes of representative cnidarians and other metazoan species were plotted against their respective orthogroup count. Only proteins from orthogroups shared with at least one published matrisome were counted. Anthozoans generally show a higher matrisome complexity than medusozoan species populating a transitory region between bilaterians and non-bilaterians in the evolutionary trajectory.

A significant fraction of *Nematostella*'s exceptionally rich matrisome is devoted to the formation of the cnidocyst, a unique cellular novelty of the cnidarian clade (Jékely, et al. 2015 [↗](#); Babonis, et al. 2023 [↗](#)). Cnidocyst-specific proteins follow an unusual secretion route into the lumen of the growing cnidocyst vesicle, which topologically represents extracellular space (Ozbek 2011 [↗](#)). It has therefore been speculated that they originated from neurosecretory vesicles used for predation in early metazoans (Balasubramanian, et al. 2012 [↗](#)). The recent finding that cnidocysts are instrumental for the predatory lifestyle of the *Aiptasia* larvae (Maegelé, et al. 2023 [↗](#)) supports the hypothesis of a deeply-rooted extrusive mechanism for prey capture in metazoan evolution. In this context, it is intriguing that the majority of *Nematostella* cnidocyte genes shared with other cell types is expressed within neuroglandular subtypes (supplementary figure S6). In addition, the cnidocyst-specific matrisome contains diverse proteins with repetitive ECM domains that likely have general bioadhesive or fibrous properties that might play a role in entangling and ingesting prey organisms. These include several fibropellin-like and other EGF repeat proteins (poly-EGF) as well as thrombospondin type-1 repeat (TSR) proteins, such as properdin-like.

The changes in the matrisome profiles across *Nematostella*'s major life stages suggest a highly dynamic epithelial rearrangement during primary polyp morphogenesis. The involvement of Wnt/PCP factors in this process indicates similar cell migration and re-orientation events as during convergent extension processes in gastrulation. Kumburegama et al. have shown that primary archenteron invagination and apical constriction of bottle cells in *Nematostella* is dependent on the PCP components *strabismus* (Kumburegama, et al. 2011 [↗](#)) and *Fzd10* (Wijesena, et al. 2022 [↗](#)). Mesentery formation in primary polyps, which involves sequential folding events of the endodermal epithelium (Berking 2007 [↗](#)), likely involves similar molecular pathways. As already observed by Appelöf (Appelöf 1900 [↗](#)), mesoglea synthesis follows invagination during this process. The molecular network composed of Wnt/PCP and basal membrane factors that our data revealed might therefore indicate an actomyosin-controlled invagination of the endodermal layer followed by BM production to re-align the cells in their apico-basal architecture. The upregulation of wound response factors in the adult animal might indicate a transient loss of tissue integrity during collective cell migration which could entail an actin-based purse-string mechanism as during wound healing (Begnaud, et al. 2016 [↗](#); Bischoff, et al. 2021 [↗](#)). Future work will further decipher the gene-regulatory network controlling polyp morphogenesis in this anthozoan model.

Materials and Methods

Nematostella culture

Adult *Nematostellas* were kept in plastic boxes at 18 °C in 1/3 artificial sea water (~11 ppm) (*Nematostella* medium) in the dark. They were fed with freshly hatched *Artemia* nauplii and cleaned once per week. To induce spawning, animals were transferred to 27 °C *Nematostella* medium in light for 8 h and then washed with 18 °C *Nematostella* medium. The egg patches were collected and dejellied in 5 % cysteine solution for 15 min. Unless otherwise stated, the embryos were left to develop in *Nematostella* medium at room temperature (RT) in normal day/night cycles.

Immunocytochemistry

Larvae on 3 dpf, primary polyps on 10 dpf, and small adult polyps were collected and left to relax at 27 °C in direct light for 30 min. A solution of 7 % MgCl₂ in seawater was slowly added, and the animals were anesthetized for 20 min. Fixation was performed with Lavdovsky's fixative (50 % ethanol, 36 % H₂O, 10 % formaldehyde, 4 % acetic acid) for 30 min at RT after which the samples were incubated in 150 mM Tris, pH 9.0, 0.05 % Tween-20, for 10 min. An incubation step at 70 °C for 10 min and a subsequent cooling to RT was followed by three 10 min washing steps in PBS, 0.1 % Tween-20, PBS, 0.1% Tween-20, 0.1% Triton-X100, and PBS, 0.1% Tween-20, respectively. Primary antibody incubation (rabbit anti-Laminin, rat anti-Pan-Collagen) was performed at 1:100 in 0.5 % milk powder overnight at 4 °C. The samples were washed 3 times for 10 min in PBS, 0.1%

Tween-20 and incubated with secondary antibodies at 1:400 for 2.5 h at RT. Prior to mounting on object slides with Mowiol, DAPI was added at 1:1000 for 30 min. Decellularized mesogleas were transferred onto a microscopy slide lined with liquid blocker. The mesogleas of larvae and primary polyps were carefully stuck to the slide using an eyelash. The staining protocol followed the whole mount immunocytochemistry protocol with 10 min fixation and only one washing step per wash to avoid washing off of the mesogleas. The antibodies were incubated on the slides in a Petri dish with a wet paper towel to prevent evaporation. Images were acquired with an A1R microscope at the Nikon Imaging Facility Heidelberg. Further image processing was performed with Fiji ImageJ v1.53t.

Mesoglea decellularization

About 500.000 larvae (3 dpf) and primary polyps (10 dpf), and four adult animals were collected. The adult polyps were cut open along the oral-aboral axis using a scalpel to ease the decellularization of the endoderm. All samples were incubated in 0.5 % N-lauryl-sarcosinate for 5 min and then frozen in liquid nitrogen. After thawing at RT, the samples were transferred to ddH₂O using a 70 µm sieve for larvae and primary polyps. The mesogleas were decellularized in ddH₂O by repeated pipetting with a flamed glass pipette and frequent water changes. The progress of decellularization was checked repeatedly by phase contrast microscopy at 60x magnification using a Nikon 80i microscope. As a final quality control, a few sample mesogleas were stained with DAPI for 10 min in PBS to visualize residual cells or nematocysts. Decellularized mesogleas were then picked individually for further analysis.

Sample preparation for SP3 and TMT labeling

Isolated mesogleas (larvae and primary polyps, N = 150, adults, N = 4) were dissolved in 1 M dithiothreitol (DTT) for 30 min at 90 °C and protein extraction was performed using trichloroacetic acid (TCA) precipitation. TCA pellets were resuspended in 50 µL 1% SDS, 50 mM HEPES pH 8.5. Reduction of disulfide bridges in cysteine containing proteins was performed with DTT (56 °C, 30 min, 10 mM in 50 mM HEPES, pH 8.5). Reduced cysteines were alkylated with 2-chloroacetamide (RT, in the dark, 30 min, 20 mM in 50 mM HEPES, pH 8.5). Samples were prepared using the SP3 protocol (Hughes, et al. 2014 [↗](#); Hughes, et al. 2019 [↗](#)) and trypsin (sequencing grade, Promega) was added in an enzyme to protein ratio of 1:50 for overnight digestion at 37°C. Then, peptide recovery was performed in HEPES buffer by collecting the supernatant on a magnet and combining it with the second elution wash of beads with HEPES buffer. Peptides were labelled with TMT10plex (Werner, et al. 2014 [↗](#)) Isobaric Label Reagent (ThermoFisher) according to the manufacturer's instructions. For further sample clean up an OASIS® HLB µElution Plate (Waters) was used for each sample separately. A control run was performed to be able to mix equal peptide amounts based on the MS signal in each run and samples were combined for TMT9plex accordingly. Offline high pH reverse phase fractionation was carried out on an Agilent 1200 Infinity high-performance liquid chromatography system, equipped with a Gemini C18 column (3 µm, 110 Å, 100 x 1.0 mm, Phenomenex) (Reichel, et al. 2016 [↗](#)).

Mass spectrometry and data analysis

LC-MS/MS, Liquid Chromatography (LC) was performed as previously described for *Hydra* mesoglea (Veschgini, et al. 2023 [↗](#)). IsobarQuant (Franken, et al. 2015 [↗](#)) and Mascot (v2.2.07) were used to process the acquired data, which was searched against the *Nematostella vectensis* NV2 (wein_nvec200_tcsv2) protein models (<https://simrbase.stowers.org/starletseanemone> [↗](#)) containing common contaminants and reversed sequences. The following modifications were included into the search parameters: Carbamidomethyl (C) and TMT10 (K) (fixed modification), Acetyl (Protein N-term), Oxidation (M) and TMT10 (N-term) (variable modifications). For the full scan (MS1) a mass error tolerance of 10 ppm and for MS/MS (MS2) spectra of 0.02 Da was set. Further parameters were: trypsin as protease with an allowance of maximum two missed cleavages, a minimum peptide length of seven amino acids, and at least two unique peptides were required for

a protein identification. The false discovery rate on peptide and protein level was set to 0.01. The raw output files of IsoBarQuant (protein.txt – files) were processed using R. Contaminants were filtered out and only proteins that were quantified with at least two unique peptides were considered for the analysis. 5056 proteins passed the quality control filters. Log2-transformed raw TMT reporter ion intensities ('signal_sum' columns) were first cleaned for batch effects using limma (Ritchie, et al. 2015 [↗](#)) and further normalized using variance stabilization normalization (Huber, et al. 2002 [↗](#)) (see supplementary figure S7 for an overview of these steps). Proteins were tested for differential expression using the limma package. The replicate information was added as a factor in the design matrix given as an argument to the 'lmFit' function of limma. A protein was annotated as a hit with a false discovery rate (fdr) smaller 5% and a fold-change of at least 2. For the heatmap shown in **Fig. 4B** [↗](#), the log2-transformed median abundance of all samples for each life stage was calculated.

In silico matrisome prediction

Domains for the protein models of the SIMRbase *Nematostella vectensis* NV2 (wein_nvec200_tcsv2) transcriptome (<https://simrbase.stowers.org/starletseanemone> [↗](#)) were *de novo* annotated using InterProScan (Jones, et al. 2014 [↗](#)). The proteins were then filtered positively using a list of known ECM protein domains and negatively by the presence of the respective exclusive domains according to Naba et al. (Naba, et al. 2012 [↗](#)). Signal peptides were predicted using SignalP-6.0 (Teufel, et al. 2022 [↗](#)) and DeepLoc-2 (Thummuluri, et al. 2022 [↗](#)). The latter was also used to predict the cellular localization of proteins. In addition, OrthoFinder vers. 2.5.4 was used with default settings to predict orthogroups from all predicted matrisomes and published matrisomes from *M. musculus*, *H. sapiens* (Naba, et al. 2012 [↗](#)), *B. taurus* (Listrat, et al. 2023 [↗](#)), *C. japonica* (Huss, et al. 2019 [↗](#)), *D. melanogaster* (Davis, et al. 2019 [↗](#)), *D. rerio* (Nauroy, et al. 2018 [↗](#)), *S. mediterranea* (Cote, et al. 2019 [↗](#)) and *C. elegans* (Teuscher, et al. 2019 [↗](#)). Finally, all *Nematostella* sequences were manually annotated by comparing their domain architecture to published protein groups. Domain and orthogroup analysis were performed using custom python scripts (See supplementary data). For the orthogroup analysis, the phylogenetically hierarchical orthogroups predicted by OrthoFinder were analyzed. To prevent domain redundancy, we restricted the analysis to SMART domains for the domain comparison, as domain comparisons for other domain databases showed similar results. To achieve a better orthogroup definition we predicted additional *in silico* matrisomes for a number of available protein model datasets in non-bilaterian species using the same bioinformatic pipeline as for *Nematostella*: Choanoflagellata: *Monosiga brevicollis*, *Salpingoeca rosetta*; Porifera: *Amphimedon queenslandica*, *Ephydatia muelleri*; Ctenophora: *Mnemiopsis leidyi*, *Pleurobrachia bachei*, *Beroe ovata* (<http://ryanlab.whitney.ufl.edu/bovadb> [↗](#)), *Pleurobrachia bachei* (Moroz, et al. 2014 [↗](#)); Placozoa: *Tricoplax adhaerens*, *Tricoplax spec*; Cnidaria: *Aurelia aurita* (Gold, et al. 2019 [↗](#)), *Clytia hemispherica* (<http://marimba.obs-vlfr.fr> [↗](#)), *Exaiptasia diaphana* (Oakley, et al. 2016 [↗](#)), *Hydra vulgaris*, *Acropora digitifera* (Shinzato, et al. 2021 [↗](#)), *Stylophora pistillata*, *Calvadosia cruxmelitensis* (Ohdera, et al. 2019 [↗](#)), *Morbakka virulenta* (Khalturin, et al. 2019 [↗](#)), *Thelohanellus kitauei*, *Acropora digitifera*, *Porites asteroides* (Kenkel, et al. 2013 [↗](#)), *Xenia spec.* (Hu, et al. 2020 [↗](#)), Chordata: *Branchiostoma belcheri*, *Branchiostoma floridae*, *Branchiostoma lanceolatum* (Uniprot Reference Proteomes). To identify orthogroups specific to cnidarians, we filtered the OrthoFinder-derived orthogroups and phylogenetic hierarchies, selecting only those that exhibited cnidarian exclusivity. To identify potential venoms in our matrisome dataset, we performed BLAST searches against the ToxProt database of known animal toxins (Jungo, et al. 2012 [↗](#)). We categorized as “putative venoms” (supplementary table S1) candidates that exhibited significant matches in this database ($E > 1e-03$).

Single cell RNA expression

The expression matrix corresponding to the predicted ECM genes was extracted from the updated single cell atlas (<https://doi.org/10.1186/s12983-024-00529-z> [↗](#)). To generate expression data, the dataset was first separated into three life-cycle stages: samples from 18 h gastrula until 4 d planula were classified as 'larva', samples from 5 d through 16 d primary polyps were classified as 'primary

polyp', and all samples derived from juvenile or adult tissues were classified as 'adult'. The full gene matrix was filtered down to only the 829 models corresponding to the curated list of ECM genes. For plotting expression values, the principal tissue-type annotations were further collapsed to cluster together early and late ectodermal clusters, specification and mature cnidocyte states, and to collapse the primary germ cells and putative stem cells into a single data partition. Differentially expressed genes were calculated across all annotated cell-type states using the Seurat vs.4 function `Seurat::FindAllMarkers`, requiring a return-threshold of 0.001, and a minimum detection in 20% of any cluster. Module expression scores for different gene sets (core, associated, and other for the full dataset, and 'ubiquitous', 'shared', 'specification-specific' and 'mature-specific' for the cnidocyte subset) were calculated using the Seurat function `Seurat::AddModuleScore`. The cnidocyte genes were binned as described above according to summarized expression data generated by the `Seurat::DotPlot` function, in the 'data' matrix of the resulting ggplot.

Electron microscopy

For morphology, *Nematostella* larvae and primary polyps were processed as previously described for *Hydra* (Bottger, et al. 2012 [DOI](#); Garg, et al. 2023 [DOI](#)). Briefly, animals were subjected to cryofixation (high-pressure freezing, freeze-substitution and epoxy resin embedding: HPF/FS: supplementary figure S3A, B, G, H) or to standard chemical fixation (glutaraldehyde, followed by OsO₄, resin embedding: CF: supplementary figure S3F, L).

Ultrathin sections were optionally stained with uranyl acetate and lead for general contrast enhancement or with periodic acid, thiocarbohydrazide and silver proteinate to highlight periodic acid-Schiff-positive constituents (supplementary figure S3H). For Tokuyasu-immunoelectron microscopy (Tokuyasu 1973 [DOI](#)) samples were either fixed for >3 days at RT with 4 % w/v formaldehyde solution in PHEM (5 mM HEPES, 60 mM PIPES, 10 mM EGTA, 2 mM MgCl₂), pH 7.0 (TOK: supplementary figure S3M, O) or by using a new modification of established HPF/FS-sample rehydration methods (Ripper, et al. 2008 [DOI](#); Schmiedinger, et al. 2013 [DOI](#)); this modification included freeze substitution with methanol containing 3.2 % w/v formaldehyde, 0.08 % w/v uranyl acetate and 8.8 % H₂O, removal of uranyl acetate at 4° C (on ice) and partial sample rehydration and postfixation through incubation in Lavdovski's fixative for 1 hour at RT (HPF/FS/RH-Lav: supplementary figure S3N, P). Fixed samples were rinsed with PHEM buffer and further processed for thawed cryosection immunogold labelling (Tokuyasu 1973 [DOI](#)) as previously described (Garg, et al. 2023 [DOI](#)). Anti-Lam and anti-PanCol label on standard TOK-sections were visualized with goat anti-rabbit or goat anti-rat secondary antibodies coupled to 10 nm colloidal gold (#EM.GAR10/1, #EM.GAT10/1, British Biocell). Anti-Col4 and anti-Col2c labelling was performed on HPF/FS/RH-Lav-samples by using Nanogold®-IgG Goat anti-Guinea Pig IgG or Nanogold®-IgG Goat anti-Rat IgG (H+L) (#2054, 2007, Nanoprobes), respectively, followed by silver enhancement with HQ silver™ (#2012, Nanoprobes).

Acknowledgements

This work was supported by the German Science Foundation (DFG) (Collaborative Research Center 1324 (B07)) and OE 416/8-1) to S.Ö. G.B. acknowledges the Schmeil Foundation Heidelberg for its support. We thank Josephine C. Adams for fruitful discussions.

Additional information

Data Availability Statement

The MS run results can be found in the PRIDE Database (PXD045345). An R script for generating all single cell RNA expression figures and the sequences and OrthoFinder results of the *in silico* matrisomes are available at <https://github.com/suatoezbek/Oezbek-matrisome>.

Funding

Deutsche Forschungsgemeinschaft (Collaborative Research Center 1324 (B07); OE 416/8-1)

Additional files

Supplementary Figures. [↗](#)

README File for Suppl. Tables. [↗](#)

Table S1. *In silico* matrisome prediction of the *Nematostella mesoglea* including results of differential abundance analysis using moderated t-test (limma) for each developmental comparison (Adult - Larvae, Larvae - Primary Polyp, Adult - Primary Polyp). Related to Figs. 1 and 4.. [↗](#)

Table S2. Table S2. Full mass spectrometry data of the *Nematostella mesoglea*. Related to Figs. 1 and 4. *qpm*: quantified unique peptide matches, *top3*: average log₁₀ MS1 intensity of the three most abundant peptides for a given protein. *norm_batchcl_raw_signal_sum*: normalized and batch-cleaned reporter ion intensities from the TMT experiment. [↗](#)

Table S3. Raw data of mesoglea and BM thickness as well as fibril thickness measurements in electron micrographs of larva and primary polyp cryosections. Related to Fig. S3. [↗](#)

Table S4. Gene expression profiles indicating the average expression (*avg.exp*), the percentage of the cells with gene expression (*pct.exp*), and the relative expression levels (*avg.exp.scaled*) for all matrisomal genes across all cell state identities (*cell.state.id*) in Cole et al. 2024, *Frontiers in Zoology*. *pSC*, putative germ cell, *PGC*, primary germ cell, *NPC*, neuroglandular progenitor cell. *GD*, digestive gland cell, *NGD*, digestive neuroglandular cell. Related to Fig. 2. [↗](#)

Table S5. Single cell expression profiles of core matrisomal genes. Related to Fig. 2. Output of the *FindAllMarkers* function of the Seurat package (vs.4.4.0) run with default parameters using all core matrisome genes across all clusters annotated in Cole et al. 2024 *Frontiers in Zoology* (*cell.state.id*). *gene*: core matrisome gene tested, *p_val*: calculated p-value of the expression difference; *avg_log2FC*: calculated average log₂ fold change between cluster 1 (*cell.state.id*) and the rest of the dataset; *pct.1*: percentage of cells in cluster 1 (*cell.state.id*) with gene expression; *pct.2*: percentage of cells within the remaining clusters with gene expression; *p_val_adj*: adjusted p-value. [↗](#)

Table S6. Matrix of average expression values for all cnidocyte-specific genes (rows) across all cell state identities in Cole et al. 2024 *Frontiers in Zoology* (columns). Related to Fig. 3. [↗](#)

Table S7. Quantitative mass spectrometry data of mesoglea samples from different life stages. Related to Fig. 4. [↗](#)

References

- Adamczyk P, Zenkert C, Balasubramanian PG, Yamada S, Murakoshi S, Sugahara K, Hwang JS, Gojobori T, Holstein TW, Ozbek S (2010) **A non-sulfated chondroitin stabilizes membrane tubulation in cnidarian organelles** *J Biol Chem* **285**:25613–25623 [Google Scholar](#)
- Adamska M, Matus DQ, Adamski M, Green K, Rokhsar DS, Martindale MQ, Degnan BM (2007) **The evolutionary origin of hedgehog proteins** *Curr Biol* **17**:R836–837 [Google Scholar](#)
- Adler PN (2002) **Planar signaling and morphogenesis in Drosophila** *Dev Cell* **2**:525–535 [Google Scholar](#)
- Aouacheria A, Geourjon C, Aghajari N, Navratil V, Deleage G, Lethias C, Exposito JY. (2006) **Insights into early extracellular matrix evolution: Spongin short chain collagen-related proteins are homologous to basement membrane type IV collagens and form a novel family widely distributed in invertebrates** *Molecular biology and evolution* **23**:2288–2302 [Google Scholar](#)
- Appelöf A (1900) **Studien über Actinien-Entwicklung: Bergens Museums Aarbog** [Google Scholar](#)
- Artamonova II, Mushegian AR. (2013) **Genome Sequence Analysis Indicates that the Model Eukaryote Nematostella vectensis Harbors Bacterial Consorts** *Applied and Environmental Microbiology* **79**:6868–6873 [Google Scholar](#)
- Aufschnaiter R, Zamir EA, Little CD, Ozbek S, Munder S, David CN, Li L, Sarras MP, Zhang X (2011) **In vivo imaging of basement membrane movement: ECM patterning shapes Hydra polyps** *J Cell Sci* **124**:4027–4038 [Google Scholar](#)
- Babonis LS, Enjolras C, Reft AJ, Foster BM, Hugosson F, Ryan JF, Daly M, Martindale MQ (2023) **Single-cell atavism reveals an ancient mechanism of cell type diversification in a sea anemone** *Nat Commun* **14**:885 [Google Scholar](#)
- Balasubramanian PG, Beckmann A, Warnken U, Schnolzer M, Schuler A, Bornberg-Bauer E, Holstein TW, Ozbek S (2012) **Proteome of Hydra nematocyst** *J Biol Chem* **287**:9672–9681 [Google Scholar](#)
- Begnaud S, Chen T, Delacour D, Mege RM, Ladoux B (2016) **Mechanics of epithelial tissues during gap closure** *Curr Opin Cell Biol* **42**:52–62 [Google Scholar](#)
- Bergheim BG, Ozbek S (2019) **Extracellular matrix and morphogenesis in cnidarians: a tightly knit relationship** *Extracellular Matrix* **63**:407–416 [Google Scholar](#)
- Berking S (2007) **Generation of bilateral symmetry in Anthozoa: a model** *J Theor Biol* **246**:477–490 [Google Scholar](#)
- Bischoff MC, Lieb S, Renkawitz-Pohl R, Bogdan S (2021) **Filopodia-based contact stimulation of cell migration drives tissue morphogenesis** *Nat Commun* **12**:791 [Google Scholar](#)

Bottger A, Doxey AC, Hess MW, Pfaller K, Salvenmoser W, Deutzmann R, Geissner A, Pauly B, Altstatter J, Munder S, et al. (2012) **Horizontal gene transfer contributed to the evolution of extracellular surface structures: the freshwater polyp Hydra is covered by a complex fibrous cuticle containing glycosaminoglycans and proteins of the PPOD and SWT (sweet tooth) families** *PLoS One* **7**:e52278 [Google Scholar](#)

Bridge D, Cunningham CW, Schierwater B, DeSalle R, Buss LW (1992) **Class-level relationships in the phylum Cnidaria: evidence from mitochondrial genome structure** *Proc Natl Acad Sci U S A* **89**:8750–8753 [Google Scholar](#)

Brooun M, Klimovich A, Bashkurov M, Pearson BJ, Steele RE, McNeill H (2020) **Ancestral roles of atypical cadherins in planar cell polarity** *Proc Natl Acad Sci U S A* **117**:19310–19320 [Google Scholar](#)

Brunet T, King N (2017) **The Origin of Animal Multicellularity and Cell Differentiation** *Dev Cell* **43**:124–140 [Google Scholar](#)

Cárdenas-León CG, Mäemets-Allas K, Klaas M, Lagus H, Kankuri E, Jaks V (2022) **Matricellular proteins in cutaneous wound healing** *Frontiers in Cell and Developmental Biology* **10** [Google Scholar](#)

Chari T, Weissbourd B, Gehring J, Ferraioli A, Leclerc L, Herl M, Gao F, Chevalier S, Copley RR, Houlston E, et al. (2021) **Whole-animal multiplexed single-cell RNA-seq reveals transcriptional shifts across Clytia medusa cell types** *Sci Adv* **7**:eabh1683 [Google Scholar](#)

Cole AG, Jahnel SM, Kaul S, Steger J, Hagauer J, Denner A, Murguía PF, Taudes E, Zimmermann B, Reischl R, et al. (2023) **Muscle cell-type diversification is driven by bHLH transcription factor expansion and extensive effector gene duplications** *Nature communications* **14** [Google Scholar](#)

Cole AG, Steger J, Hagauer J, Denner A, Ferrer Murguía P, Knabl P, Narayanaswamy S, Wick B, Montenegro JD, Technau U (2024) **Updated single cell reference atlas for the starlet anemone Nematostella vectensis** *Front Zool* **21**:8 [Google Scholar](#)

Cote LE, Simental E, Reddien PW (2019) **Muscle functions as a connective tissue and source of extracellular matrix in planarians** *Nature communications* **10** [Google Scholar](#)

David CN, Ozbek S, Adamczyk P, Meier S, Pauly B, Chapman J, Hwang JS, Gojobori T, Holstein TW (2008) **Evolution of complex structures: minicollagens shape the cnidarian nematocyst** *Trends Genet* **24**:431–438 [Google Scholar](#)

Davis MN, Horne-Badovinac S, Naba A (2019) **In-silico definition of the Drosophila melanogaster matrisome** *Matrix Biol Plus* **4**:100015 [Google Scholar](#)

Day RM, Lenhoff HM (1981) **Hydra mesoglea: a model for investigating epithelial cell - basement membrane interactions** *Science* **211**:291–294 [Google Scholar](#)

DeBiasse MB, Buckenmeyer A, Macrander J, Babonis LS, Bentlage B, Cartwright P, Prada C, Reitzel AM, Stampar SN, Collins A, et al. (2024) **A Cnidarian Phylogenomic Tree Fitted With Hundreds of 18S Leaves** *Bulletin of the Society of Systematic Biologists* **3** [Google Scholar](#)

Deutzmann R, Fowler S, Zhang X, Boone K, Dexter S, Boot-Handford RP, Rachel R, Sarras MP (2000) **Molecular, biochemical and functional analysis of a novel and developmentally important fibrillar collagen (Hcol-I) in hydra** *Development* **127**:4669–4680 [Google Scholar](#)

- Draper GW, Shoemark DK, Adams JC (2019) **Modelling the early evolution of extracellular matrix from modern Ctenophores and Sponges** *Extracellular Matrix* **63**:389–405 [Google Scholar](#)
- DuBuc TQ, Traylor-Knowles N, Martindale MQ (2014) **Initiating a regenerative response; cellular and molecular features of wound healing in the cnidarian Nematostella vectensis** *BMC Biol* **12**:24 [Google Scholar](#)
- Emms DM, Kelly S (2019) **OrthoFinder: phylogenetic orthology inference for comparative genomics** *Genome Biol* **20**:238 [Google Scholar](#)
- Engel J (1996) **Domain organizations of modular extracellular matrix proteins and their evolution** *Matrix Biology* **15**:295–299 [Google Scholar](#)
- Engel U, Ozbek S, Engel R, Petri B, Lottspeich F, Holstein TW (2002) **NOWA, a novel protein with minicollagen Cys-rich domains involved in nematocyst formation in Hydra** *J. Cell Sci* **115** [Google Scholar](#)
- Epp L, Smid I, Tardent P (1986) **Synthesis of the mesoglea by ectoderm and endoderm in reassembled hydra** *J Morphol* **189**:271–279 [Google Scholar](#)
- Exposito JY, Larroux C, Cluzel C, Valcourt U, Lethias C, Degnan BM (2008) **Demosponge and sea anemone fibrillar collagen diversity reveals the early emergence of A/C clades and the maintenance of the modular structure of type V/XI collagens from sponge to human** *J Biol Chem* **283**:28226–28235 [Google Scholar](#)
- Fahey B, Degnan BM (2012) **Origin and evolution of laminin gene family diversity** *Mol Biol Evol* **29**:1823–1836 [Google Scholar](#)
- Fidler AL, Boudko SP, Rokas A, Hudson BG (2018) **The triple helix of collagens - an ancient protein structure that enabled animal multicellularity and tissue evolution** *Journal of Cell Science* **131** [Google Scholar](#)
- Fidler AL, Darris CE, Chetyrkin SV, Pedchenko VK, Boudko SP, Brown KL, Jerome WG, Hudson JK, Rokas A, Hudson BG (2017) **Collagen IV and basement membrane at the evolutionary dawn of metazoan tissues** *eLife* **6** [Google Scholar](#)
- Fidler AL, Vanacore RM, Chetyrkin SV, Pedchenko VK, Bhawe G, Yin VP, Stothers CL, Rose KL, McDonald WH, Clark TA, et al. (2014) **A unique covalent bond in basement membrane is a primordial innovation for tissue evolution** *Proceedings of the National Academy of Sciences of the United States of America* **111**:331–336 [Google Scholar](#)
- Fowler SJ, Jose S, Zhang X, Deutzmann R, Sarras MP, Boot-Handford RP (2000) **Characterization of hydra type IV collagen. Type IV collagen is essential for head regeneration and its expression is up-regulated upon exposure to glucose** *J Biol Chem* **275**:39589–39599 [Google Scholar](#)
- Franken H, Mathieson T, Childs D, Sweetman GM, Werner T, Togel I, Doce C, Gade S, Bantscheff M, Drewes G, et al. (2015) **Thermal proteome profiling for unbiased identification of direct and indirect drug targets using multiplexed quantitative mass spectrometry** *Nat Protoc* **10**:1567–1593 [Google Scholar](#)
- Gao B (2012) **Wnt regulation of planar cell polarity (PCP)** *Curr Top Dev Biol* **101**:263–295 [Google Scholar](#)

Garg N, Stibler UK, Eismann B, Mercker M, Bergheim BG, Linn A, Tuchscherer P, Engel U, Redl S, Marciniak-Czochra A, et al. (2023) **Non-muscle myosin II drives critical steps of nematocyst morphogenesis** *iScience* **26**:106291 [Google Scholar](#)

Gilges D, Vinit MA, Callebaut I, Coulombel L, Cacheux V, Romeo PH, Vigon I (2000) **Polydom: a secreted protein with pentraxin, complement control protein, epidermal growth factor and von Willebrand factor A domains** *Biochem J* **352**:49–59 [Google Scholar](#)

Gold DA, Katsuki T, Li Y, Yan X, Regulski M, Ibberson D, Holstein T, Steele RE, Jacobs DK, Greenspan RJ (2019) **The genome of the jellyfish *Aurelia* and the evolution of animal complexity** *Nat Ecol Evol* **3**:96–104 [Google Scholar](#)

Gosline JM (1971a) **Connective Tissue Mechanics of *Metridium-Senile*.1. Structural and Compositional Aspects** *Journal of Experimental Biology* **55**:763–774 [Google Scholar](#)

Gosline JM (1971b) **Connective Tissue Mechanics of *Metridium-Senile*.2. Visco-Elastic Properties and Macromolecular Model** *Journal of Experimental Biology* **55**:775–795 [Google Scholar](#)

Gubb D, Garcia-Bellido A (1982) **A genetic analysis of the determination of cuticular polarity during development in *Drosophila melanogaster*** *J Embryol Exp Morphol* **68**:37–57 [Google Scholar](#)

He F, Xiong W, Yu X, Espinoza-Lewis R, Liu C, Gu S, Nishita M, Suzuki K, Yamada G, Minami Y, et al. (2008) **Wnt5a regulates directional cell migration and cell proliferation via Ror2-mediated noncanonical pathway in mammalian palate development** *Development* **135**:3871–3879 [Google Scholar](#)

Heljasvaara R, Aikio M, Ruotsalainen H, Pihlajaniemi T (2017) **Collagen XVIII in tissue homeostasis and dysregulation - Lessons learned from model organisms and human patients** *Matrix Biology* :57–58 [Google Scholar](#)

Hikasa H, Shibata M, Hiratani I, Taira M (2002) **The *Xenopus* receptor tyrosine kinase Xrnr2 modulates morphogenetic movements of the axial mesoderm and neuroectoderm via Wnt signaling** *Development* **129**:5227–5239 [Google Scholar](#)

Hohenester E, Engel J (2002) **Domain structure and organisation in extracellular matrix proteins** *Matrix Biology* **21**:115–128 [Google Scholar](#)

Holland JW, Okamura B, Hartikainen H, Secombes CJ (2011) **A novel minicollagen gene links cnidarians and myxozoans** *Proc Biol Sci* **278**:546–553 [Google Scholar](#)

Hu M, Zheng X, Fan CM, Zheng Y (2020) **Lineage dynamics of the endosymbiotic cell type in the soft coral *Xenia*** *Nature* **582**:534–538 [Google Scholar](#)

Huber W, von Heydebreck A, Sultmann H, Poustka A, Vingron M. (2002) **Variance stabilization applied to microarray data calibration and to the quantification of differential expression** *Bioinformatics* **18**:S96–104 [Google Scholar](#)

Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, Krijgsvelde J (2014) **Ultrasensitive proteome analysis using paramagnetic bead technology** *Molecular Systems Biology* **10**:757 [Google Scholar](#)

- Hughes CS, Moggridge S, Muller T, Sorensen PH, Morin GB, Krijgsveld J (2019) **Single-pot, solid-phase-enhanced sample preparation for proteomics experiments** *Nat Protoc* **14**:68–85 [Google Scholar](#)
- Huss DJ, Saias S, Hamamah S, Singh JM, Wang JH, Dave M, Kim J, Eberwine J, Lansford R. (2019) **Avian Primordial Germ Cells Contribute to and Interact With the Extracellular Matrix During Early Migration** *Frontiers in Cell and Developmental Biology* **7** [Google Scholar](#)
- Hwang JS, Takaku Y, Momose T, Adamczyk P, Ozbek S, Ikeo K, Khalturin K, Hemmrich G, Bosch TC, Holstein TW, et al. (2010) **Nematogalectin, a nematocyst protein with GlyXY and galectin domains, demonstrates nematocyte-specific alternative splicing in Hydra** *Proc Natl Acad Sci U S A* **107**:18539–18544 [Google Scholar](#)
- Hynes RO (2012) **The evolution of metazoan extracellular matrix** *Journal of Cell Biology* **196**:671–679 [Google Scholar](#)
- Hynes RO (2009) **The extracellular matrix: not just pretty fibrils** *Science* **326**:1216–1219 [Google Scholar](#)
- Hynes RO, Naba A (2012) **Overview of the matrisome--an inventory of extracellular matrix constituents and functions** *Cold Spring Harb Perspect Biol* **4**:a004903 [Google Scholar](#)
- Jekely G, Keijzer F, Godfrey-Smith P (2015) **An option space for early neural evolution** *Philos Trans R Soc Lond B Biol Sci* **370** [Google Scholar](#)
- Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, et al. (2014) **InterProScan 5: genome-scale protein function classification** *Bioinformatics* **30**:1236–1240 [Google Scholar](#)
- Jungo F, Bougueleret L, Xenarios I, Poux S (2012) **The UniProtKB/Swiss-Prot Tox-Prot program: A central hub of integrated venom protein data** *Toxicon* **60**:551–557 [Google Scholar](#)
- Kadler KE, Baldock C, Bella J, Boot-Handford RP (2007) **Collagens at a glance** *Journal of Cell Science* **120**:1955–1958 [Google Scholar](#)
- Kenkel CD, Meyer E, Matz MV (2013) **Gene expression under chronic heat stress in populations of the mustard hill coral (*Porites astreoides*) from different thermal environments** *Mol Ecol* **22**:4322–4334 [Google Scholar](#)
- Khalturin K, Shinzato C, Khalturina M, Hamada M, Fujie M, Koyanagi R, Kanda M, Goto H, Anton-Erxleben F, Toyokawa M, et al. (2019) **Medusozoan genomes inform the evolution of the jellyfish body plan** *Nat Ecol Evol* **3**:811–822 [Google Scholar](#)
- King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough S, Hellsten U, Isogai Y, Letunic I, et al. (2008) **The genome of the choanoflagellate *Monosiga brevicollis* and the origin of metazoans** *Nature* **451**:783–788 [Google Scholar](#)
- Kumburegama S, Wijesena N, Xu R, Wikramanayake AH (2011) **Strabismus-mediated primary archenteron invagination is uncoupled from Wnt/beta-catenin-dependent endoderm cell fate specification in *Nematostella vectensis* (Anthozoa, Cnidaria): Implications for the evolution of gastrulation** *Evodevo* **2**:2 [Google Scholar](#)

Kurz EM, Holstein TW, Petri BM, Engel J, David CN (1991) **Mini-collagens in hydra nematocytes** *J Cell Biol* **115**:1159–1169 [Google Scholar](#)

Lebouvier M, Miramon-Puertolas P, Steinmetz PRH (2022) **Evolutionarily conserved aspects of animal nutrient uptake and transport in sea anemone vitellogenesis** *Curr Biol* **32**:4620–4630 [Google Scholar](#)

Listrat A, Bobby C, Tournayre J, Jousse C (2023) **Bovine extracellular matrix proteins and potential role in meat quality: First in silico Bos taurus compendium** *J Proteomics* **279**:104891 [Google Scholar](#)

Lommel M, Strompen J, Hellewell AL, Balasubramanian GP, Christofidou ED, Thomson AR, Boyle AL, Woolfson DN, Puglisi K, Hartl M, et al. (2018) **Hydra Mesoglea Proteome Identifies Thrombospondin as a Conserved Component Active in Head Organizer Restriction** *Sci Rep* **8**:11753 [Google Scholar](#)

Maegele I, Rupp S, Ozbek S, Guse A, Hambleton EA, Holstein TW (2023) **A predatory gastrula leads to symbiosis-independent settlement in Aiptasia** *Proc Natl Acad Sci U S A* **120**:e2311872120 [Google Scholar](#)

Matakatsu H, Blair SS (2004) **Interactions between Fat and Dachous and the regulation of planar cell polarity in the Drosophila wing** *Development* **131**:3785–3794 [Google Scholar](#)

Meyer F, Moussian B (2009) **Drosophila multiplexin (Dmp) modulates motor axon pathfinding accuracy** *Dev Growth Differ* **51**:483–498 [Google Scholar](#)

Miller DJ, Hayward DC, Reece-Hoyes JS, Scholten I, Catmull J, Gehring WJ, Callaerts P, Larsen JE, Ball EE (2000) **Pax gene diversity in the basal cnidarian Acropora millepora (Cnidaria, Anthozoa): implications for the evolution of the Pax gene family** *Proc Natl Acad Sci U S A* **97**:4475–4480 [Google Scholar](#)

Moroz LL, Kocot KM, Citarella MR, Dosung S, Norekian TP, Povolotskaya IS, Grigorenko AP, Dailey C, Berezikov E, Buckley KM, et al. (2014) **The ctenophore genome and the evolutionary origins of neural systems** *Nature* **510**:109–114 [Google Scholar](#)

Naba A (2024) **Mechanisms of assembly and remodelling of the extracellular matrix** *Nat Rev Mol Cell Biol* [Google Scholar](#)

Naba A, Clauser KR, Hoersch S, Liu H, Carr SA, Hynes RO (2012) **The matrisome: in silico definition and in vivo characterization by proteomics of normal and tumor extracellular matrices** *Mol Cell Proteomics* **11**:014647 [Google Scholar](#)

Nauroy P, Hughes S, Naba A, Ruggiero F (2018) **The in-silico zebrafish matrisome: A new tool to study extracellular matrix gene and protein functions** *Matrix Biology* **65**:5–13 [Google Scholar](#)

Nichols SA, Roberts BW, Richter DJ, Fairclough SR, King N (2012) **Origin of metazoan cadherin diversity and the antiquity of the classical cadherin/beta-catenin complex** *Proc Natl Acad Sci U S A* **109**:13046–13051 [Google Scholar](#)

- Nishita M, Yoo SK, Nomachi A, Kani S, Sougawa N, Ohta Y, Takada S, Kikuchi A, Minami Y (2006) **Filopodia formation mediated by receptor tyrosine kinase Ror2 is required for Wnt5a-induced cell migration** *J Cell Biol* **175**:555–562 [Google Scholar](#)
- Novinec M, Kordis D, Turk V, Lenarcic B (2006) **Diversity and evolution of the thyroglobulin type-1 domain superfamily** *Molecular biology and evolution* **23**:744–755 [Google Scholar](#)
- Oakley CA, Ameismeier MF, Peng L, Weis VM, Grossman AR, Davy SK (2016) **Symbiosis induces widespread changes in the proteome of the model cnidarian Aiptasia** *Cell Microbiol* **18**:1009–1023 [Google Scholar](#)
- Ohdera A, Ames CL, Dikow RB, Kayal E, Chiodin M, Busby B, La S, Pirro S, Collins AG, Medina M, et al. (2019) **Box, stalked, and upside-down? Draft genomes from diverse jellyfish (Cnidaria, Acraspeda) lineages: Alatina alata (Cubozoa), Calvadosia cruxmelitensis (Staurozoa), and Cassiopea xamachana (Scyphozoa)** *Gigascience* **8** [Google Scholar](#)
- Ozbek S (2011) **The cnidarian nematocyst: a miniature extracellular matrix within a secretory vesicle** *Protoplasma* **248**:635–640 [Google Scholar](#)
- Ozbek S, Balasubramanian PG, Chiquet-Ehrismann R, Tucker RP, Adams JC (2010) **The evolution of extracellular matrix** *Mol Biol Cell* **21**:4300–4305 [Google Scholar](#)
- Reichel M, Liao Y, Rettel M, Ragan C, Evers M, Alleaume AM, Horos R, Hentze MW, Preiss T, Millar AA (2016) **In Planta Determination of the mRNA-Binding Proteome of Arabidopsis Etiolated Seedlings** *Plant Cell* **28**:2435–2452 [Google Scholar](#)
- Richards GS, Rentzsch F (2014) **Transgenic analysis of a SoxB gene reveals neural progenitor cells in the cnidarian Nematostella vectensis** *Development* **141**:4681–4689 [Google Scholar](#)
- Ripper D, Schwarz H, Stierhof YD (2008) **Cryo-section immunolabelling of difficult to preserve specimens: advantages of cryofixation, freeze-substitution and rehydration** *Biology of the Cell* **100**:109–123 [Google Scholar](#)
- Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) **. limma powers differential expression analyses for RNA-sequencing and microarray studies** *Nucleic Acids Res* **43**:e47 [Google Scholar](#)
- Rokas A (2008) **The origins of multicellularity and the early history of the genetic toolkit for animal development** *Annu Rev Genet* **42**:235–251 [Google Scholar](#)
- Samuelov L, Li Q, Bochner R, Najor NA, Albrecht L, Malchin N, Goldsmith T, Grafi-Cohen M, Vodo D, Fainberg G, et al. (2017) **SVEP1 plays a crucial role in epidermal differentiation** *Exp Dermatol* **26**:423–430 [Google Scholar](#)
- Sarras MP Jr (2012) **Components, structure, biogenesis and function of the Hydra extracellular matrix in regeneration, pattern formation and cell differentiation** *Int J Dev Biol* **56**:567–576 [Google Scholar](#)
- Sarras MP Jr, Madden ME, Zhang XM, Gunwar S, Huff JK, Hudson BG (1991) **Extracellular matrix (mesoglea) of Hydra vulgaris. I. Isolation and characterization** *Dev Biol* **148**:481–494 [Google Scholar](#)

- Sato-Nishiuchi R, Doiguchi M, Morooka N, Sekiguchi K (2023) **Polydom/SVEP1 binds to Tie1 and promotes migration of lymphatic endothelial cells** *J Cell Biol* **222** [Google Scholar](#)
- Sato-Nishiuchi R, Nakano I, Ozawa A, Sato Y, Takeichi M, Kiyozumi D, Yamazaki K, Yasunaga T, Futaki S, Sekiguchi K (2012) **Polydom/SVEP1 Is a Ligand for Integrin $\alpha 9 \beta 1$** *Journal of Biological Chemistry* **287**:25615–25630 [Google Scholar](#)
- Schaffer AA, Bazarsky M, Levy K, Chalifa-Caspi V, Gat U (2016) **A transcriptional time-course analysis of oral vs. aboral whole-body regeneration in the Sea anemone** *Bmc Genomics* **17** [Google Scholar](#)
- Schambony A, Wedlich D (2007) **Wnt-5A/Ror2 regulate expression of XPAPC through an alternative noncanonical signaling pathway** *Dev Cell* **12**:779–792 [Google Scholar](#)
- Schmiedinger T, Vogel GF, Eiter O, Pfaller K, Kaufmann WA, Flörl A, Gutleben K, Schönherr S, Witting B, Lechleitner TW, et al. (2013) **Cryo-Immuno-electron Microscopy of Adherent Cells Improved by the Use of Electrospun Cell Culture Substrates** *Traffic* **14**:886–894 [Google Scholar](#)
- Schultz DT, Haddock SHD, Bredeson JV, Green RE, Simakov O, Rokhsar DS (2023) **Ancient gene linkages support ctenophores as sister to other animals** *Nature* **618**:110–117 [Google Scholar](#)
- Schwarz RS, Bosch TC, Cadavid LF (2008) **Evolution of polydom-like molecules: identification and characterization of cnidarian polydom (Cnpolydom) in the basal metazoan Hydractinia** *Dev Comp Immunol* **32**:1192–1210 [Google Scholar](#)
- Sebe-Pedros A, Roger AJ, Lang FB, King N, Ruiz-Trillo I (2010) **Ancient origin of the integrin-mediated adhesion and signaling machinery** *Proc Natl Acad Sci U S A* **107**:10142–10147 [Google Scholar](#)
- Shimizu H, Aufschnaiter R, Li L, Sarras MP, Borza DB, Abrahamson DR, Sado Y, Zhang X. (2008) **The extracellular matrix of hydra is a porous sheet and contains type IV collagen** *Zoology (Jena)* **111**:410–418 [Google Scholar](#)
- Shinzato C, Khalturin K, Inoue J, Zayasu Y, Kanda M, Kawamitsu M, Yoshioka Y, Yamashita H, Suzuki G, Satoh N (2021) **Eighteen Coral Genomes Reveal the Evolutionary Origin of Acropora Strategies to Accommodate Environmental Changes** *Mol Biol Evol* **38**:16–30 [Google Scholar](#)
- Shoemark DK, Ziegler B, Watanabe H, Strompen J, Tucker RP, Ozbek S, Adams JC (2019) **Emergence of a Thrombospondin Superfamily at the Origin of Metazoans** *Mol Biol Evol* **36**:1220–1238 [Google Scholar](#)
- Simons M, Mlodzik M (2008) **Planar cell polarity signaling: from fly development to human disease** *Annu Rev Genet* **42**:517–540 [Google Scholar](#)
- Steger J, Cole AG, Denner A, Lebedeva T, Genikhovich G, Ries A, Reischl R, Taudes E, Lassnig M, Technau U (2022) **Single-cell transcriptomics identifies conserved regulators of neuroglandular lineages** *Cell Rep* **40**:111370 [Google Scholar](#)

- Steinmetz PRH, Aman A, Kraus JEM, Technau U (2017) **Gut-like ectodermal tissue in a sea anemone challenges germ layer homology** *Nat Ecol Evol* **1**:1535–1542 [Google Scholar](#)
- Stokkermans A, Chakrabarti A, Subramanian K, Wang L, Yin S, Moghe P, Steenbergen P, Monke G, Hiiragi T, Prevedel R, et al. (2022) **Muscular hydraulics drive larva-polyp morphogenesis** *Curr Biol* **32**:4707–4718 [Google Scholar](#)
- Teufel F, Armenteros JJA, Johansen AR, Gislason MH, Pihl SI, Tsirigos KD, Winther O, Brunak S, von Heijne G, Nielsen H. (2022) **SignalP 6.0 predicts all five types of signal peptides using protein language models** *Nature Biotechnology* **40**:1023–1025 [Google Scholar](#)
- Teuscher AC, Jongsma E, Davis MN, Statzer C, Gebauer JM, Naba A, Ewald CY (2019) **The in-silico characterization of the *Caenorhabditis elegans* matrisome and proposal of a novel collagen classification** *Matrix Biol Plus* **1**:100001 [Google Scholar](#)
- Thummuluri V, Almagro Armenteros JJ, Johansen AR, Nielsen H, Winther O (2022) **DeepLoc 2.0: multi-label subcellular localization prediction using protein language models** *Nucleic Acids Res* **50**:W228–W234 [Google Scholar](#)
- Tokuyasu KT (1973) **A technique for ultracryotomy of cell suspensions and tissues** *J Cell Biol* **57**:551–565 [Google Scholar](#)
- Tournière O, Dolan D, Richards GS, Sunagar K, Columbus-Shenkar YY, Moran Y, Rentzsch F (2020) **NvPOU4/Brain3 Functions as a Terminal Selector Gene in the Nervous System of the Cnidarian** *Cell Reports* **30**:4473–4489 [Google Scholar](#)
- Tucker RP, Adams JC (2014) **Adhesion Networks of Cnidarians: A Postgenomic View** *International Review of Cell and Molecular Biology* **308**:323–377 [Google Scholar](#)
- Tucker RP, Hess JF, Gong Q, Garvey K, Shibata B, Adams JC (2013) **A thrombospondin in the anthozoan *Nematostella vectensis* is associated with the nervous system and upregulated during regeneration** *Biol Open* **2**:217–226 [Google Scholar](#)
- Tucker RP, Shibata B, Blankenship TN (2011) **Ultrastructure of the mesoglea of the sea anemone *Nematostella vectensis* (Edwardsiidae)** *Invertebrate Biology* **130**:11–24 [Google Scholar](#)
- Veschgini M, Suzuki R, Kling S, Petersen HO, Bergheim BG, Abuillan W, Linke P, Kaufmann S, Burghammer M, Engel U, et al. (2023) **Wnt/beta-catenin signaling induces axial elasticity patterns of *Hydra* extracellular matrix** *iScience* **26**:106416 [Google Scholar](#)
- Werner T, Sweetman G, Savitski MF, Mathieson T, Bantscheff M, Savitski MM (2014) **Ion coalescence of neutron encoded TMT 10-plex reporter ions** *Anal Chem* **86**:3594–3601 [Google Scholar](#)
- Wijesena N, Sun HY, Kumburegama S, Wikramanayake AH (2022) **Distinct Frizzled receptors independently mediate endomesoderm specification and primary archenteron invagination during gastrulation in *Nematostella*** *Developmental biology* **481**:215–225 [Google Scholar](#)
- Zenkert C, Takahashi T, Diesner MO, Ozbek S (2011) **Morphological and molecular analysis of the *Nematostella vectensis* cnidom** *PLoS One* **6**:e22725 [Google Scholar](#)

Zhang X, Boot-Handford RP, Huxley-Jones J, Forse LN, Mould AP, Robertson DL, Lili Athiyal M, Sarras MP Jr (2007) **The collagens of hydra provide insight into the evolution of metazoan extracellular matrices** *J Biol Chem* **282**:6792–6802 [Google Scholar](#)

Zhou Y, Chen L, Hao ST, Cao XK, Ni SS (2024) **Zebrafish ANGPT4, member of fibrinogen-related proteins, is an LTA-, LPS and PGN-binding protein with a bacteriolytic activity** *Fish & Shellfish Immunology* **147** [Google Scholar](#)

Author information

B Gideon Bergheim

University of Heidelberg, Centre for Organismal Studies, Department of Evolutionary Neurobiology, Heidelberg, Germany
ORCID iD: [0000-0002-3922-8147](#)

Alison G Cole

Department of Neurosciences and Developmental Biology, Faculty of Life Sciences, University of Vienna, Vienna, Austria
ORCID iD: [0000-0002-7515-7489](#)

Mandy Rettel

European Molecular Biology Laboratory, Heidelberg, Germany
ORCID iD: [0000-0002-8304-3385](#)

Frank Stein

European Molecular Biology Laboratory, Heidelberg, Germany
ORCID iD: [0000-0001-9695-1692](#)

Stefan Redl

Institute of Neuroanatomy, Medical University of Innsbruck, Innsbruck, Austria

Michael W Hess

Innsbruck Medical University, Innsbruck, Austria

Aissam Ikmi

European Molecular Biology Laboratory, Heidelberg, Germany
ORCID iD: [0000-0001-6310-8404](#)

Suat Özbek

University of Heidelberg, Centre for Organismal Studies, Department of Evolutionary Neurobiology, Heidelberg, Germany
ORCID iD: [0000-0003-2569-3942](#)

For correspondence: suat.oezbek@cos.uni-heidelberg.de

Editors

Reviewing Editor

Ariel Chipman

The Hebrew University of Jerusalem, Jerusalem, Israel

Senior Editor

Kathryn Cheah

University of Hong Kong, Hong Kong, Hong Kong

Reviewer #1 (Public review):

Summary:

In this manuscript entitled "Molecular dynamics of the matrisome across sea anemone life history", Bergheim and colleagues report the prediction, using an established sequence analysis pipeline, of the "matrisome" - that is, the compendium of genes encoding constituents of the extracellular matrix - of the starlet sea anemone *Nematostella vectensis*. Re-analysis of an existing scRNA-Seq dataset allowed the authors to identify the cell types expressing matrisome components and different developmental stages. Last, the authors apply time-resolved proteomics to provide experimental evidence of the presence of the extracellular matrix proteins at three different stages of the life cycle of the sea anemone (larva, primary polyp, adult) and show that different subsets of matrisome components are present in the ECM at different life stages with, for example, basement membrane components accompanying the transition from larva to primary polyp and elastic fiber components and matricellular proteins accompanying the transition from primary polyp to the adult stage.

Strengths:

The ECM is a structure that has evolved to support the emergence of multicellularity and different transitions that have accompanied the complexification of multicellular organisms. Understanding the molecular makeup of structures that are conserved throughout evolution is thus of paramount importance.

The in-silico predicted matrisome of the sea anemone has the potential to become an essential resource for the scientific community to support big data annotation efforts and better understand the evolution of the matrisome and of ECM proteins, an important endeavor to better understand structure/function relationships. Toward this goal, the authors provide a comprehensive list with extensive annotations and cross-referencing of the 551 genes encoding matrisome proteins in the sea anemone genome.

This study is also an excellent example of how integrating datasets generated using different -omic modalities can shed light on various aspects of ECM metabolism, from identifying the cell types of origins of matrisome components using scRNA-Seq to studying ECM dynamics using proteomics.

Weakness:

- Prior proteomic studies on the ECM of vertebrate organisms have shown the importance of allowing certain post-translational modifications during database search to ensure maximizing peptide-to-spectrum matching and accurately evaluating protein quantification. Such PTMs include the hydroxylation of lysines and prolines that are collagen-specific PTMs. Multiple reports have shown that omitting these PTMs while analyzing LC-MS/MS data would lead to underestimating the abundance of collagens and the misidentification of certain

collagens. While the authors in their response state that the inclusion of these PTMs only led to a modest increase in protein identification, they do not comment on the impact of including these PTMs on PSMs or protein abundance (precursor ion intensity).

<https://doi.org/10.7554/eLife.105319.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

Summary:

*In this manuscript entitled "Molecular dynamics of the matrisome across sea anemone life history", Bergheim and colleagues report the prediction, using an established sequence analysis pipeline, of the "matrisome" - that is, the compendium of genes encoding constituents of the extracellular matrix - of the starlet sea anemone *Nematostella vectensis*. Re-analysis of an existing scRNA-Seq dataset allowed the authors to identify the cell types expressing matrisome components and different developmental stages. Last, the authors apply time-resolved proteomics to provide experimental evidence of the presence of the extracellular matrix proteins at three different stages of the life cycle of the sea anemone (larva, primary polyp, adult) and show that different subsets of matrisome components are present in the ECM at different life stages with, for example, basement membrane components accompanying the transition from larva to primary polyp and elastic fiber components and matricellular proteins accompanying the transition from primary polyp to the adult stage.*

Strengths:

The ECM is a structure that has evolved to support the emergence of multicellularity and different transitions that have accompanied the complexification of multicellular organisms. Understanding the molecular makeup of structures that are conserved throughout evolution is thus of paramount importance.

The in-silico predicted matrisome of the sea anemone has the potential to become an essential resource for the scientific community to support big data annotation efforts and understand better the evolution of the matrisome and of ECM proteins, an important endeavor to better understand structure/function relationships. This study is also an excellent example of how integrating datasets generated using different -omic modalities can shed light on various aspects of ECM metabolism, from identifying the cell types of origins of matrisome components using scRNA-Seq to studying ECM dynamics using proteomics.

We greatly appreciate the positive feedback regarding the design of our study and the evolutionary significance of our findings.

Weaknesses:

My concerns pertain to the three following areas of the manuscript:

(1) In-silico definition of the anemone matrisome using sequence analysis:

a) While a similar computational pipeline has been applied to predict the matrisome of several model organisms, the authors fail to provide a comprehensive definition of the anemone matrisome: In the text, the authors state the anemone matrisome is composed

of "551 proteins, constituting approximately 3% of its proteome (see page 6, line 14), but Figure 1 lists 829 entries as part of the "curated" matrisome, Supplementary Table S1 lists the same 829 entries and the authors state that "Here, we identified 829 ECM proteins that comprise the matrisome of the sea anemone *Nematostella vectensis*" (see page 17, line 10). Is the sea anemone matrisome composed of 551 or 829 genes? If we refer to the text, the additional 278 entries should not be considered as part of the matrisome, but what is confusing is that some are listed as glycoproteins and the "new_manual_annotation" proposed by the authors and that refer to the protein domains found in these additional proteins suggest that in fact, some could or should be classified as matrisome proteins. For example, shouldn't the two lectins encoded by NV2.3951 and NV2.3157 be classified as matrisome-affiliated proteins? Based on what has been done for other model organisms, receptors have typically been excluded from the "matrisome" but included as part of the "adhesome" for consistency with previously published matrisome; the reviewer is left wondering whether the components classified as "Other" / "Receptor" should not be excluded from the matrisome and moved to a separate "adhesome" list.

In addition to receptors, the authors identify nearly 70 glycoproteins classified as "Other". Here, does other mean "non-matrisome" or "another matrisome division" that is not core or associated? If the latter, could the authors try to propose a unifying term for these proteins? Unfortunately, since the authors do not provide the reasons for excluding these entries from the bona fide matrisome (list of excluding domains present, localization data), the reader is left wondering how to treat these entries.

Overall, the study would gain in strength if the authors could be more definitive and, if needed, even propose novel additional matrisome annotations to include the components for now listed as "Other" (as was done, for example, for the *Drosophila* or *C. elegans* matrisomes).

The reviewer is correct to point out the confusing terminology used throughout our manuscript, where both the total of 829 proteins constituting the curated list of ECM domain proteins and the actual matrisome (excluding "others") were referred to as "matrisomes". In general, we followed the example set by Naba & Hynes in their 2012 paper (Mol Cell Proteomics. 2012 Apr;11(4):M111.014647. doi: 10.1074/mcp.M111.014647), where they define the "matrisome" as encompassing all components of the extracellular matrix ("core matrisome") and those associated with it ("matrisome-associated" proteins). This corresponds to our group of 551 proteins, comprising both core matrisome and matrisome-associated proteins. The Naba & Hynes paper also contains the inclusive and exclusive domain lists for the matrisome that we applied for our dataset. In the revised manuscript, we have now labelled the group of 829 proteins as "curated ECM domain proteins/genes", which includes all proteins positively selected for containing a bona fide ECM domain. After excluding non-matrisomal proteins such as receptors, we arrive at the 551 proteins that constitute the "*Nematostella* matrisome". We have maintained this terminology throughout the revised manuscript and have revised Figures 1B and 4B accordingly.

Regarding the category of "other" proteins, which by definition are not part of the matrisome although containing ECM domains, we have taken the reviewer's advice and classified these in more detail. We categorized all receptors as "adhesome" (202 proteins). The remaining group of "other" secreted ECM domain proteins were then further subcategorized. Those exhibiting significant matches in the ToxProt database were subclassified as "putative venoms" (15 proteins). This group also includes the two lectins (NV2.3951 and NV2.3157), which had been originally shifted to the "other" category due to their classification as venoms. We categorized as "adhesive proteins" (28 proteins) factors such as coadhesins that due to their domain architecture resemble bioadhesive proteins described in proteomic studies of other invertebrate species, such as corals or sponges (see also <https://doi.org/10>

[.1016/j.jprot.2022.104506](https://doi.org/10.1016/j.jprot.2022.104506)). Further sub-categories are stress/injury response proteins (9 proteins) and ion channels (6 proteins). The remaining 17 proteins were categorized as “uncharacterized ECM domain proteins”. These include highly diverse proteins possessing either single ECM domains or novel domain combinations. We decided to retain those in our dataset as candidates for future functional characterization.

b) It is surprising that the authors are not providing the full currently accepted protein names to the entries listed in Supplementary Table S1 and have used instead "new_manual_annotation" that resembles formal protein names. This liberty is misleading. In fact, the "new_manual_annotation" seems biased toward describing the reason the proteins were positively screened for through sequence analysis, but many are misleading because there is, in fact, more known about them, including evidence that they are not ECM proteins. The authors should at least provide the current protein names in addition to their "new_manual_annotations".

c) To truly serve as a resource, the Table should provide links to each gene entry in the Stowers Institute for Medical Research genome database used and some sort of versioning (this could be added to columns A, B, or D). Such enhancements would facilitate the assessment of the rigor of the list beyond the manual QC of just a few entries.

d) Since UniProt is the reference protein knowledge database, providing the UniProt IDs associated with the predicted matrisome entries would also be helpful, giving easy access to information on protein domains, protein structures, orthology information, etc.

e) In conclusion, at present, the study only provides a preliminary draft that should be more rigorously curated and enriched with more comprehensive and authoritative annotations if the authors aspire the list to become the reference anemone matrisome and serve the community.

Table S1 has been updated to include links to the respective Stowers Institute IDs (first two columns), as well as SwissProt IDs and current descriptions from both the Stowers Institute (SI) and Swissprot.

In our manual annotations, we prioritized these over automated ones due to the considerable effort invested in examining each sequence individually. The cnidaria-specific minicollagens and NOWA proteins might serve as an example. According to the SI descriptions, the minicollagens are annotated as “keratin-associated protein, predicted or hypothetical protein, collagen-like protein and pericardin”. We classified these as minicollagens on the basis of overall domain architecture and of signature domains and sequence motifs, such as minicollagen cysteine-rich domains (CRDs) and polyproline stretches (doi: [10.1016/j.tig.2008.07.001](https://doi.org/10.1016/j.tig.2008.07.001)). NOWA is a CTLD/CRD-containing protein that is part of nematocyst tubules (doi: [10.1016/j.isci.2023.106291](https://doi.org/10.1016/j.isci.2023.106291)). The first two NOWA isoforms, according to SI descriptions, were annotated as aggrecan and brevican core proteins, which is very misleading. We therefore feel that our manual annotations better serve the cnidarian research community in classifying these proteins.

Automated annotations of ECM proteins often rely on similarities between individual domains, neglecting overall domain composition. For example, Swissprot descriptions annotate 31 TSP1 domain-containing proteins in our list as “Hemicentin-1”, but closer inspection reveals that only one sequence (NV2.24790) qualifies as Hemicentin-1 due to its characteristic vWFA, Ig-like, TSP1, G2 nidogen, and EGF-like domain architecture. Regarding novel protein annotations, NV2.650 might serve as an example. While SI descriptions annotate this protein as “epidermal growth factor” based on the presence of several EGF-like domains, our analysis reveals two integrin alpha N-terminal domains that classify this sequence as integrin-related. We have therefore assigned a description (Secreted integrin-N-

related protein) that references this defining domain and avoids misclassification within the EGF family.

In cases where the automated annotation (including those in Genbank) matched our own findings, we adopted the existing description, as seen with netrin-1 (NV2.7734). We acknowledge that our manual annotations are not flawless and will be refined by future research. Nonetheless, we offer them as an approximation to a more accurate definition of the identified protein list.

(2) Proteomic analysis of the composition of the mesoglea during the sea anemone life cycle:

a) The product of 287 of the 829 genes proposed to encode matrisome components was detected by proteomics. What about the other ~550 matrisome genes? When and where are they expressed? The wording employed by the authors (see line 11, page 13) implies that only these 287 components are "validated" matrisome components. Is that to say that the other ~550 predicted genes do not encode components of the ECM? This should be discussed.

Obviously, our wording was not sufficiently accurate here. In the revised Fig. 1B we indicated that 210 of the 551 matrisome (core and associated) proteins were confirmed by mass spectrometry. In total, 287 proteins were identified by mass spectrometry, meaning that 77 of those are non-matrisomal proteins belonging to the “adhesome” (47) and “other” (30) groups. The fact that the remaining 542 proteins of the matrisome predicted by our in silico analysis could not be identified has two major reasons: (1) Our study was focussed on the molecular dynamics of the mesoglea. Therefore, only mesogleas were isolated for the mass spectrometry analysis and nematocysts were mostly excluded by extensive washing steps. As nematocysts contribute significantly to the predicted matrisome, this group of proteins is underrepresented in the mass spectrometry analysis. (2) A significant fraction of the predicted ECM proteins constitutes soluble factors and transmembrane receptors. These might not be necessarily part of the mesoglea isolates. In addition, the isolation and solubilization method we applied might have technical limitations. Although we used harsh conditions for solubilizing the mesoglea samples (90°C and high DTT concentrations), we cannot exclude that we missed proteins which resisted solubilization and thus trypsinization. We confirmed that all genes predicted by the in silico analysis have transcriptomic profiles as demonstrated in supplementary table S4. We have clarified these points in the revised results part (p.6) and also revised the statement in line 16, page 13.

b) Can the authors comment on how they have treated zero TMT values or proteins for which a TMT ratio could not be calculated because unique to one life stage, for example?

We did not include these proteins in the analysis of the respective statistical comparison. This involved only very few proteins (about 10).

c) Could the authors provide a plot showing the distribution of protein abundances for each matrisome category in the main figure 4? In mammals, the bulk of the ECM is composed of collagens, followed by fibrillar ECM glycoproteins, the other matrisome components being more minor. Is a similar distribution observed in the sea anemone mesoglea?

We have included such a plot showing protein abundances across life stages and protein categories (Fig. 4A). Collagens and basement membrane proteoglycans (perlecan) are the most abundant protein categories in the core matrisome while secreted factors dominate in the matrisome-associated group.

d) Prior proteomic studies on the ECM of vertebrate organisms have shown the importance of allowing certain post-translational modifications during database search to ensure maximizing peptide-to-spectrum matching. Such PTMs include the hydroxylation of lysines and prolines that are collagen-specific PTMs. Multiple reports have shown that omitting these PTMs while analyzing LC-MS/MS data would lead to underestimating the abundance of collagens and the misidentification of certain collagens. The authors may want to reanalyze their dataset and include these PTMs as part of their search criteria to ensure capturing all collagen-derived peptides.

Thank you for this suggestion. We have re-analyzed our dataset including lysine and proline hydroxylation as PTM. While we obtained in total 70 more proteins using this approach, this additional group did not contain any large collagen or minicollagen we had not detected before. We only obtained two additional collagen-like proteins with very short triple helical domains (V2t013973001.1, NV2t024002001.1), one being a fragment. We don't feel this justifies implementing a re-analysis of the proteome in our study.

e) The authors should ensure that reviewers are provided with access to the private PRIDE repository so the data deposited can also be evaluated. They should also ensure that sufficient meta-data is provided using the SRDF format to allow the re-use of their LCMS/MS datasets.

We apologize for not providing the reviewer access in our initial submission and have asked the editorial office to forward the PRIDE repository link to all reviewers immediately after receiving the reviews. We did upload a metadata.csv file with the proteomics dataset. This file contains an annotation of all TMT labels to the samples and conditions and replicates used in the manuscript. It contains similar information as an SRDF format file. In addition, the search output files on protein and psm level have been provided. So, from our point of view, we provided all necessary information to reproduce the analysis.

(3) Supplementary tables:

The supplementary tables are very difficult to navigate. They would become more accessible to readers and non-specialists if they were accompanied by brief legends or "README" tabs and if the headers were more detailed (see, for example, Table S2, what does "ctrl.ratio_Larvae_rep2" exactly refer to? Or Table S6 whose column headers using extensive abbreviations are quite obscure). Similarly, what do columns K to BX in Supplementary Table S1 correspond to? Without more substantial explanations, readers have no way of assessing these data points.

We have revised the tables and removed any redundant data columns. We also included detailed explanations of the used abbreviations, both in the headers and in a separate README file. Some of the information was apparently lost during the conversion to pdf files. We will therefore upload the original .xls files when submitting the revised manuscript.

Reviewer #2 (Public review):

This work set out to identify all extracellular matrix proteins and associated factors present within the starlet sea anemone Nematostella vectensis at different life stages. Combining existing genomic and transcriptomic datasets, alongside new mass spectrometry data, the authors provide a comprehensive description of the Nematostella matrisome. In addition, immunohistochemistry and electron microscopy were used to image whole mount and decellularized mesoglea from all life stages. This served to validate the de-cellularization methods used for proteomic analyses, but also resulted in a very nice description of mesoglea structure at different life stages. A previously

published developmental cell type atlas was used to identify the cell type specificity of the matrisome, indicating that the core matrisome is predominantly expressed in the gastrodermis, as well as cnidocytes. The analyses performed were rigorous and the results were clear, supporting the conclusions made by the authors.

Thank you. We greatly appreciate the positive assessment of our study.

Reviewer #3 (Public review):

Summary:

*This manuscript by Bergheim et al investigates the molecular and developmental dynamics of the matrisome, a set of gene products that comprise the extracellular matrix, in the sea anemone *Nematostella vectensis* using transcriptomic and proteomic approaches. Previous work has examined the matrisome of the hydra, a medusozoan, but this is the first study to characterize the matrisome in an anthozoan. The major finding of this work is a description of the components of the matrisome in *Nematostella*, which turns out to be more complex than that previously observed in hydra. The authors also describe the remodeling of the extracellular matrix that occurs in the transition from larva to primary polyp, and from primary polyp to adult. The authors interpret these data to support previously proposed (Steinmetz et al. 2017) homology between the cnidarian endoderm with the bilaterian mesoderm.*

Strengths:

*The data described in this work are robust, combining both transcriptome and proteomic interrogation of key stages in the life history of *Nematostella*, and are of value to the community.*

Thank you for your positive assessment of our dataset.

Weaknesses:

The authors offer numerous evolutionary interpretations of their results that I believe are unfounded. The main problem with extending these results, together with previous results from hydra, into an evolutionary synthesis that aims to reconstruct the matrisome of the ancestral cnidarian is that we are considering data from only two species. I agree with the authors' depiction of hydra as "derived" relative to other medusozoans and see it as potentially misleading to consider the hydra matrisome as an exemplar for the medusozoan matrisome. Given the organismal and morphological diversity of the phylum, a more thorough comparative study that compares matrisome components across a selection of anthozoan and medusozoan species using formal comparative methods to examine hypotheses is required.

Specifically, I question the author's interpretation of the evolutionary events depicted in this statement:

*"The observation that in *Hydra* both germ layers contribute to the synthesis of core matrisome proteins (Epp et al. 1986; Zhang et al. 2007) might be related to a secondary loss of the anthozoan-specific mesenteries, which represent extensions of the mesoglea into the body cavity sandwiched by two endodermal layers."*

*Anthozoans and medusozoans are evolutionary sisters. Therefore, the secondary loss of "anthozoan-like mesenteries" in hydrozoans is at least as likely as the gain of this character state in anthozoans. By extension, there is no reason to prefer the hypothesis that the state observed in *Nematostella*, where gastroderm is responsible for the synthesis of the core matrisome components, is the ancestral state of the phylum.*

Moreover, the fossil evidence provided in support of this hypothesis (Ou et al. 2022) is not relevant here because the material described in that work is of a crown group anthozoan, which diversified well after the origin of Anthozoa. The phylogenetic structure of Cnidaria has been extensively studied using phylogenomic approaches and is generally well supported (Kayal et al. 2018; DeBiasse et al. 2024). Based on these analyses, anthozoans are not on a "basal" branch, as the authors suggest. The structure of cnidarian phylogeny bifurcates with Anthozoa forming one clade and Medusozoa forming the other. From the data reported by Bergheim and coworkers, it is not possible to infer the evolutionary events that gave rise to the different matrisome states observed in *Nematostella* (an anthozoan) and *hydra* (a medusozoan). Furthermore, I take the observation in Fig 5 that anthozoan matrisomes generally exhibit a higher complexity than other cnidarian species to be more supportive of a lineage-specific expansion of matrisome components in the Anthozoa, rather than those components being representative of an ancestral state for Cnidaria. Whatever the implication, I take strong issue with the statement that "the acquisition of complex life cycles in medusozoa, that are distinguished by the pelagic medusa stage, led to a secondary reduction in the matrisome repertoire." There is no causal link in any of the data or analyses reported by Bergheim and co-workers to support this statement and, as stated above, while we are dealing with limited data, insufficient to address this question, it seems more likely to me that the matrisome expanded in anthozoans, contrasting with the authors' conclusions. While the discussion raises many interesting evolutionary hypotheses related to the origin of the cnidarian matrisome, which is of vital interest if we are to understand the origin of the bilaterian matrisome, a more thorough comparative analysis, inclusive of a much greater cnidarian species diversity, is required if we are to evaluate these hypotheses.

DeBiasse MB, Buckenmeyer A, Macrander J, Babonis LS, Bentlage B, Cartwright P, Prada C, Reitzel AM, Stampar SN, Collins A, et al. 2024. A Cnidarian Phylogenomic Tree Fitted With Hundreds of 18S Leaves. *Bulletin of the Society of Systematic Biologists* [Internet] 3. Available from: <https://ssbulletin.org/index.php/bssb/article/view/9267>

Epp L, Smid I, Tardent P. 1986. Synthesis of the mesoglea by ectoderm and endoderm in reassembled hydra. *J Morphol* [Internet] 189:271-279. Available from: <https://pubmed.ncbi.nlm.nih.gov/29954165/>

Kayal E, Bentlage B, Sabrina Pankey M, Ohdera AH, Medina M, Plachetzki DC, Collins AG, Ryan JF. 2018. Phylogenomics provides a robust topology of the major cnidarian lineages and insights on the origins of key organismal traits. *BMC Evol Biol* [Internet] 18:1-18. Available from: <https://bmcecol-evol.biomedcentral.com/articles/10.1186/s12862-018-1142-0>

Ou Q, Shu D, Zhang Z, Han J, Van Iten H, Cheng M, Sun J, Yao X, Wang R, Mayer G. 2022. Dawn of complex animal food webs: A new predatory anthozoan (Cnidaria) from Cambrian. *The Innovation* 3:100195

Steinmetz PRH, Aman A, Kraus JEM, Technau U. 2017. Gut-like ectodermal tissue in a sea anemone challenges germ layer homology. *Nature Ecology & Evolution* 2017 1:10 [Internet] 1:1535-1542. Available from: <https://www.nature.com/articles/s41559-017-0285-5>

Zhang X, Boot-Handford RP, Huxley-Jones J, Forse LN, Mould AP, Robertson DL, Li L, Athiyal M, Sarras MP. 2007. The collagens of hydra provide insight into the evolution of metazoan extracellular matrices. *J Biol Chem* [Internet] 282:6792-6802. Available from: <https://pubmed.ncbi.nlm.nih.gov/17204477/>

We agree with the reviewer that only the analysis of several additional anthozoan and medusozoan representatives will yield a valid basis for a reconstruction of the ancestral cnidarian matrisome and allow statements about ancestral or novel features within the phylum. We have therefore revised our statements in the discussion part of the manuscript by implementing the cited literature and also findings from medusozoan genome analysis (e.g. Gold et al., 2018) demonstrating that changes in gene content are as common in the anthozoans as in medusozoans, which questioned the previously stated “basal” state of Nematostella or of anthozoans in general.

Reviewer #1 (Recommendations for the authors):

(1) In Figure 2A, an "o" is missing in the labeling of the "developing cnidocytes" population.

Thank you, we have corrected the typo.

(2) It would be helpful to have the different life stages indicated as headers of the heat maps presented in Figure 4.

We have included symbolic representations for the different life stages on top of the heat maps in addition to the respective labels at the bottom.

Reviewer #2 (Recommendations for the authors):

Important changes:

(1) Figure 2B The x-axis tissue names should be changed to something more easily readable/understandable - some are clear, but others are not. Perhaps abbreviations could be expanded in the legend.

We have expanded the legend in Fig. 2B to render it more easily readable. We have also rotated the maps in A to have them aligned with the ones in Fig.3B.

(2) Figure 3B This figure would be improved by the inclusion of cluster names, to understand better the mapping.

We have added relevant cluster names to Fig. 3B and as stated above aligned the orientation of the maps in Fig. 2B and Fig. 3B.

(3) Figure 3C As with 2B, I find the y-axis cnidocyte cell state names to be unclear at times. Perhaps abbreviations could be expanded in the legend.

All abbreviations were expanded in Fig.3C axis labels.

(4) Many of the supplementary tables are not well exported or easily readable as is (gene names are truncated, headers truncated, etc), which means that they may not be easily usable by researchers in the field interested in following up on this work in other contexts. Indeed, to be more usable, please consider sharing these supplementary data as .csv files, for example, instead of as .pdfs.

We are sorry for this inconvenience, which was obviously caused by the conversion to pdf files. We will upload the original csv files when submitting the revised manuscript.

Smaller nitpicky comments:

(5) Page 2 line 4 & page 3 line 7: Please consider a term other than "pre-bilaterian". The drawing/ordering of a phylogeny of extant species is not meaningful in terms of more or less ancestral. e.g. if the tips are flipped in the drawing of the tree, can we say that bilaterians are pre-cnidarians? What does that mean?

We have used that term on the basis that cnidarians existed before the appearance of bilaterians according to the fossil record and molecular phylogenies (McFadden et al., 2021; Adoutte et al., 2000; Cavalier-Smith et al., 1996; Collins, 1998; Kim et al., 1999; Medina et al., 2001; Wainright et al., 1993). To acknowledge remaining uncertainties in the timing of origin of animals, we will use the term "early-diverging metazoans" instead, which is widely accepted in the cnidarian community.

(6) Page 3 line 9 I was confused by the use of "gastrula-shaped body" to describe cnidarians, which are on the whole very morphologically diverse and don't all resemble gastrulae (that can also be quite diverse).

This term is sometimes used to refer to the diploblastic cnidarian body plan (outer ectoderm, inner endoderm) with a mouth that corresponds to the blastopore. To avoid misunderstandings, we changed it in the revised manuscript to "Cnidarians, the sister group to bilaterians, are characterized by a simple body plan with a central body cavity and a mouth opening surrounded by tentacles."

Reviewer #3 (Recommendations for the authors):

(1) In general, I felt there was a lot of discussion about protein structure and diversity that is difficult to follow without a figure. I think some of the information in Supplementary Figures S5, S9, and S11 should be in the main figures.

Following the reviewer's suggestion, we have integrated Fig. S5 (collagens) into the main Fig. 2 and Fig. S9 (polydoms) into Fig. 4. As metalloproteases are not extensively discussed in the manuscript (and also due to the large size of the figure) we have kept Fig. S11 as a supplementary figure.

(2) Page 3, Line 7: The use of the term "pre-bilaterian" is inappropriate. Cnidarians and bilaterians are evolutionary sisters. Therefore, each lineage derives from the same split and is the same age. The cnidarian lineage is not older than the bilaterian lineage.

Following a similar request by reviewer 2 we have replaced this term by "early diverging metazoans".

(3) Page 5, Line 10. How were *in silico* matrisomes from early-branching metazoan species predicted?

We applied the same bioinformatic pipeline as for the *Nematostella* matrisome. We clarified this in the respective methods part.

(4) Page 16, Line 8: This should be *Thus*.

Obviously, the wording of this sentence was ambiguous. We changed it to "In contrast, the adult mesoglea is significantly enriched in elastic fiber components, such as fibrillins and fibulin. This compositional shift likely adds to the visco-elastic properties (Gosline 1971a, b) of the growing body column (Fig. 4B,D, supplementary table S7)."

<https://doi.org/10.7554/eLife.105319.2.sa0>