

Reviewed Preprint

v1 • November 28, 2023

Not revised

Reviewed Preprint

v2 • March 10, 2026

Revised by authors

✉ For correspondence:

zhangluoying@hust.edu.cn

Competing interests:

No competing interests declared

Funding:

See [page 26](#)

Reviewing editor:

Krishna Melnattur, Ashoka University, India

© 2023, Chen et al. This article is distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Circadian photoreceptor CRYPTOCHROME promotes wakefulness under short winter-like days via a GABAergic circuitry

Lixia Chen^{1,2}, Danya Tian¹, Chang Su¹, Luoying Zhang^{1,3} ✉

¹Key Laboratory of Molecular Biophysics of Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, China • ²National "111" Center for Cellular Regulation and Molecular Pharmaceutics, Key Laboratory of Fermentation Engineering (Ministry of Education), Hubei University of Technology, Wuhan, China • ³Hubei Province Key Laboratory of Oral and Maxillofacial Development and Regeneration, Wuhan, China

eLife Assessment

Winter months with short days are commonly associated with seasonal depression and hypersomnolence; the mechanisms behind this hypersomnolence however, remain unclear. Chen and colleagues identify a genetic basis for this phenomenon in the fly *Drosophila* - mutations in the circadian photoreceptor cryptochrome resulted in increased sleep under short photoperiods. These findings are **valuable** insights into the genetic mechanisms regulating sleep under short days. The data supporting the precise neurobiological basis of these effects however, remains **incomplete**.

<https://doi.org/10.7554/eLife.92608.2.sa4>

Abstract

A cardinal symptom of seasonal affective disorder (SAD, also known as winter depression) is hypersomnolence, while the cause of this “winter sleepiness” is not known. Here we found that lack of the circadian photoreceptor *cryptochrome* (*cry*) leads to increased sleep under short winter-like days in fruit flies, reminiscent of the hypersomnolence in SAD. CRY functions in neurons that synthesize the major inhibitory neurotransmitter GABA, including the small ventral lateral neurons which are known to be circadian pacemakers, and down-regulates the GABAergic tone. This in turn leads to increased neural activity of the wake-promoting large ventral lateral neurons, a subset of circadian neurons that are inhibited by GABA-A receptor. CRY protein is known to be degraded by light, thus rendering CRY to be functional within this GABAergic circuitry to enhance wakefulness only under short day length. Taken together, we demonstrate a mechanism that specifically regulates wakefulness under short winter-like days, which may provide insights regarding the winter sleepiness in SAD.

Introduction

Seasonal affective disorder (SAD), also known as winter depression, is characterized by the onset of depression in fall/winter months and spontaneous remission in the spring/summer (Rosenthal et al., 1984 [\[1\]](#)). It is generally agreed that SAD is caused by lack of day light in the fall/winter months due to the short day length (or photoperiod), as bright light therapy is effective and most commonly used for treating the depression associated with SAD (Golden et al., 2005 [\[2\]](#)). The prevalence of SAD is ~1-10% world-wide, with symptoms lasting for approximately 40% of the year (Kurlansik and Ibay, 2012 [\[3\]](#)). About 64-80% of SAD patients report a winter increase in sleep, ranging from 30 min to 2 h longer in duration compared to controls, which is considered to be a

distinguishing symptom in the characterization and diagnosis of SAD (Wescott et al., 2020 [↗](#)). Currently almost nothing is known regarding the underlying mechanism of this winter hypersomnolence.

This phenomenon of winter hypersomnolence in SAD patients implicates distinct mechanisms that regulate sleep under short winter-like photoperiods vs. longer non-winter-like photoperiods, as the sleep of these individuals appear to be selectively perturbed under short photoperiods. However, the mechanisms by which sleep duration is determined under different photoperiods have not been characterized. Since the circadian clock is believed to be important for adaptations to seasonal changes in the environment and in particular, seasonal changes of photoperiod, we hypothesize that the circadian clock may also participate in regulating sleep duration under different photoperiods (Wood and Loudon, 2014 [↗](#)).

To address our hypothesis, we tested fruit flies mutant for different circadian clock genes under a range of photoperiods. We found that flies lacking the circadian photoreceptor CRYPTOCHROME (CRY) display increased sleep duration specifically under short photoperiods, similar to the winter hypersomnolence in SAD. Genetic and pharmacological analysis identified that CRY is functioning in GABAergic neurons and acts upon GABA-A receptor to promote wakefulness. We further narrowed down the neural circuitry mediating the influences of CRY on sleep by demonstrating that *cry* deficiency increases the GABAergic tone and reduces calcium concentration in the wake-promoting large ventral lateral neurons (l-LNvs) which are known to be GABA-A+ (Hamasaka et al., 2005 [↗](#); Parisky et al., 2008 [↗](#)). Consistently, inhibiting these neurons increases sleep while activating them blocks the effects of *cry* deficiency on sleep. CRY likely functions in part in the GABAergic small ventral lateral neurons (s-LNvs), and lack of *cry* increases GABA level and the activity of these cells while impairing their GABA transmission suppresses the sleep phenotype of *cry* mutants (Allada and Chung, 2010 [↗](#)). In summary, here we identify a potential role for CRY in down-regulating the GABAergic signaling from the s-LNvs to the l-LNvs specifically under short photoperiod. This in turn enhances the neural activity of the wake-promoting l-LNvs, resulting in increased wakefulness during short winter-like days. These findings reveal a mechanism underlying how sleep duration is determined under winter-like photoperiod, while disruptions of this regulatory system may be related to the winter hypersomnolence associated with SAD.

Results

***cry* mutation increases sleep duration specifically under short photoperiods**

We assessed the sleep of flies mutant for circadian clock gene *period* (*per*⁰), *timeless* (*tim*⁰), *clock* (*clk*^{*irr*k}), *cycle* (*cyc*⁰) and *cry* (*cry*^{*b*}) under a range of photoperiods (Figure 1A and B [↗](#)) (Allada et al., 1998 [↗](#); Konopka and Benzer, 1971 [↗](#); Sehgal et al., 1994 [↗](#); Stanewsky et al., 1998 [↗](#)). We found that *cry*^{*b*} mutation, which is known to be a loss of function or severe hypomorphic allele, leads to increased sleep duration under 4 h light: 20 h dark condition (4L20D) and 8L16D but not under longer photoperiods (Stanewsky et al., 1998 [↗](#)). Since this phenotype recapitulates the winter hypersomnolence of SAD patients, we further characterized the effects of *cry* deficiency on sleep. We focused on sleep under 4L20D, as the extent of sleep increase is slightly larger than that of 8L16D. We also demonstrated that *cry* mutation lengthens sleep duration in both male and female flies, while waking activity is not significantly reduced in the mutants (Figure 1C and D [↗](#)). This means the increased sleep in the mutants is not caused by defects of locomotor ability. We next examined the sleep architecture of these flies and found that *cry* mutation enhances sleep by extending the duration of average sleep bout rather than increasing sleep bout number, indicating that *cry* deficiency promotes sleep consolidation under short photoperiod (Figure 1E and F [↗](#)). Because CRY exerts influences on sleep/wakefulness in a gender-independent manner, we used male flies for the remainder of the study. In addition, we tested the effects of a *cry* knock-out allele (*cry*⁰) on sleep under 4L20D and also observed significantly prolonged sleep duration, similar to *cry*^{*b*} mutation (Figure 1—figure supplement 1 [↗](#)) (Dolezelova et al., 2007 [↗](#)).




Figure 1. *cry* mutation increases sleep duration selectively under short photoperiods.

(A) Daily sleep duration in male circadian clock gene mutants *per⁰*, *tim⁰*, *clk^{rk}*, *cyc⁰*, and *cry^b* compared to wild-type (WT) males under 4L20D, 8L16D, 12L12D, 16L8D, 20L4D (12L12D, n = 86, 63, 91, 14, 91, 53 flies; 8L16D, n = 21, 31, 31, 10, 32, 31 flies; 12L12D, n = 31, 14, 32, 14, 27, 32 flies; 16L8D, n = 36, 30, 33, 17, 36, 30 flies; 20L4D, n = 42, 55, 83, 26, 91, 89 flies). (B) Average sleep traces of *cry^b* and WT male flies under 4L20D. (C-F) The daily sleep duration (C), waking activity (D), sleep bout number (E), average sleep bout length (F) for *cry^b* and WT female and male flies under 4L20D (n = 56, 88, 58, 91 flies). For (B), statistical differences between *cry^b* and WT is determined by paired two-tailed *t*-test, $*P < 0.001$. For A and C-E, statistical differences is determined by two-tailed Student's *t*-test, $***P < 0.001$. Error bars represent standard error of the mean (SEM). ZT, Zeitgeber Time (ZT0 is the time of lights on).

CRY functions in GABAergic neurons and promotes wakefulness via GABA/GABA-A

Upon light activation, CRY binds to the clock protein TIM and results in its degradation, which is followed subsequently by degradation of CRY itself (Busza et al., 2004 [DOI](#); Ceriani et al., 1999 [DOI](#); Lin et al., 2001 [DOI](#)). Therefore, we first tested whether the influences of CRY on sleep/wakefulness requires TIM. We monitored sleep in flies mutant for both *cry* and *tim*, and found that the sleep duration of these double mutants are comparable to that of *cry* mutants (Figure 2—figure supplement 1 [DOI](#)). This indicates that CRY promotes wakefulness in a TIM-independent manner.

To identify anatomical substrates that mediate the effects of CRY on wakefulness, we employed the UAS/GAL4 system to knock down *cry* in different brain structures and cell types and verified that *cry* is indeed knocked down by assessing its mRNA level (Figure 2—figure supplement 2 [DOI](#)). We found that knocking down *cry* in GABAergic neurons using *vesicular GABA transporter* (*VGAT*)GAL4 or two independent *glutamic acid decarboxylase 1* (*Gad1*)GAL4 lines results in increased sleep duration under 4L20D but not 12L12D, similar to the *cry* mutant phenotype (Figure 2A-D [DOI](#); Figure 2—figure supplement 3A-3L [DOI](#)) (Deng et al., 2019 [DOI](#)). Given that *Gad1*GAL4-2 generated a more prominent sleep phenotype, we used this driver for the remainder of this study.

Since CRY appears to act in GABAergic neurons to promote wakefulness, we tested whether GABA is involved in this regulatory process. We first took a pharmacological approach and fed flies with drugs to inhibit GABA transaminase (ethanolamine-O-sulphate, EOS) or GABA transporter (nipecotic acid, NipA). Both of these treatments are known to increase GABA level, and indeed lengthens the sleep duration in WT flies (Ki and Lim, 2019 [DOI](#); Leal and Neckameyer, 2002 [DOI](#)). *cry* mutation fails to enhance sleep when treated with NipA, while the extent of sleep increase caused by *cry* mutation is smaller when treated with EOS compared to the control (Figure 2E [DOI](#); Figure 2—figure supplement 4A and B [DOI](#)). Similarly, knocking down *cry* in GABAergic neurons no longer lengthens sleep duration after NipA treatment (Figure 2F [DOI](#); Figure 2—figure supplement 4C [DOI](#)). For further validation, we adopted a genetic approach. We found that knocking down *Gad1*, the synthetic enzyme for GABA, suppresses the long sleep phenotype of *cry* RNAi flies (Figure 2G [DOI](#); Figure 2—figure supplement 4D [DOI](#)) (Jackson et al., 1990 [DOI](#)). In addition, we knocked down *VGAT* which encodes an essential transporter responsible for packing GABA into synaptic vesicles (Enell et al., 2007 [DOI](#)). *cry* deficiency fails to lengthen sleep duration when *VGAT* is knocked down (Figure 2H and I [DOI](#); Figure 2—figure supplement 4E and F [DOI](#)). We verified that *Gad1* and *VGAT* are indeed knocked down by measuring their mRNA levels (Figure 2—figure supplement 4G and H [DOI](#)). As a control for the genetic interaction experiments, we co-expressed a GFP RNAi and found this does not significantly alter the sleep duration of *cry* RNAi flies (Figure 2—figure supplement 4I [DOI](#)). These series of results indicate that CRY acts via GABA to promote wakefulness.

We next sought to identify GABA receptor that mediates the effects of CRY on sleep/wakefulness. We fed flies with agonists of GABA-A (THIP) and GABA-B receptor (SKF-97541) (Ki and Lim, 2019 [DOI](#); Matsuda et al., 1996 [DOI](#); Mezler et al., 2001 [DOI](#)). Both drugs enhance sleep in WT, while *cry* mutation can increase sleep in flies fed with SKF-97541 but not THIP, implicating that CRY acts through GABA-A to promote wake (Figure 3A-C [DOI](#)). Consistently, GABA-A receptor antagonist carbamazepine (CBZ) reduces sleep in WT flies while *cry* mutation fails to lengthen sleep duration after CBZ treatment (Figure 3D and E [DOI](#)) (Agosto et al., 2008 [DOI](#)). We also treated *cry* RNAi flies with THIP or CBZ and found this treatment abolished the long sleep phenotype as well (Figure 3F-I [DOI](#)). To validate that CRY modulates sleep/wakefulness by acting upon GABA-A receptor, we tested for genetic interaction between *cry* and *Resistant to dieldrin* (*Rdl*), a gene that encodes GABA-A receptor in flies and has previously been shown to be involved in sleep regulation (Chung et al., 2009 [DOI](#); French-Constant et al., 1993 [DOI](#); Parisky et al., 2008 [DOI](#)). We found that *Rdl* mutation (*Rdl*^{MDRR}) blocks the sleep-enhancing effect of *cry* RNAi (Figure 3J-L [DOI](#)) (Agosto et al., 2008 [DOI](#)). These findings demonstrate that GABA-A receptor mediates the wake-promoting function of CRY.

In short, pharmacological and genetic approaches reveal that CRY acts in GABAergic neurons and impinges on GABA/GABA-A signaling to promote wakefulness.




Figure 2. CRY functions in GABAergic neurons and promotes wakefulness via GABA signaling.

(A, B, C) Daily sleep duration of male flies with *cry* knocked down in GABAergic neurons by *VGATGAL4* (A) (*n* = 72, 74, 96, 49, 54 flies), *Gad1GAL4-1* (B) (*n* = 29, 26, 29, 23, 32 flies) or *Gad1GAL4-2* (C) (*n* = 34, 31, 31, 25, 23 flies) monitored under 4L20D. (D) Daily sleep duration of *cry* knocked down in GABAergic neurons by *Gad1GAL4* monitored under 12L12D (*n* = 20, 19, 30, 26, 32, 20 flies). For A-D, One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, ****P* < 0.001; compared to UAS control, #*P* < 0.05, ###*P* < 0.001. (E) Daily sleep duration of male WT and *cry*^b flies fed with EOS or NipA under 4L20D (*n* = 31, 32, 30, 30, 25, 32 flies). Two-tailed Student's *t*-test: compared to WT, ###*P* < 0.001; compared to vehicle control, **P* < 0.05, ****P* < 0.001. (F) Daily sleep duration of *cry* RNAi and control flies fed with NipA under 4L20D (*n* = 72, 74, 95, 49, 54, 23, 24, 26, 15, 24 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ****P* < 0.001; compared to UAS control, ###*P* < 0.001. For comparing to vehicle control, two-tailed Student's *t*-test was used, \$\$\$*P* < 0.001. (G) Daily sleep duration of male flies with *cry* and *gad1* knocked down in GABAergic neurons monitored under 4L20D (*n* = 72, 73, 24, 33, 45, 33, 27 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ***P* < 0.01, ****P* < 0.001; compared to UAS control, ###*P* < 0.001. For comparing to RNAi control, two-tailed Student's *t*-test was used: compared to UAS*cry*RNAi/GAL4 control, \$\$\$*P* < 0.001; compared to UAS*Gad1*RNAi/GAL4 control, &&&*P* < 0.001. (H) Daily sleep duration of male flies with *cry* and *VGAT* knocked down in GABAergic neurons monitored under 4L20D (*n* = 72, 73, 46, 41, 45, 66, 34 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ****P* < 0.001; compared to UAS control, #*P* < 0.05, ###*P* < 0.001. For comparing to RNAi control, two-tailed Student's *t*-test was used: compared to UAS*cry*RNAi/GAL4 control, \$\$\$*P* < 0.001; compared to UAS*VGAT*RNAi/GAL4 background, not significant. (I) Daily sleep duration of *cry* mutant flies with *VGAT* knocked down in GABAergic neurons monitored under 4L20D (*n* = 72, 59, 20, 23, 69, 22 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ***P* < 0.01, ****P* < 0.001; compared to UAS control, #*P* < 0.05, ###*P* < 0.001. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: compared to WT background, &&&*P* < 0.001; compared to *UVGAT*RNAi/GAL4, not significant. Error bars represent SEM; G4, GAL4; U, UAS; NS, not significant.




Figure 3. GABA-A receptor mediates the effects of CRY on sleep/wakefulness.

(A, C, D) Daily sleep duration of male WT and *cry* mutant flies fed with THIP (A) ($n = 28, 29, 16, 19$ flies), SKF-97541 (C) ($n = 25, 21, 23, 18$ flies) or CBZ (D) ($n = 28, 25, 31, 29$ flies) under 4L20D. Two-tailed Student's *t*-test: compared to WT, $###P < 0.001$; compared to vehicle control, $***P < 0.001$. (B, E) Sleep profile of male WT and *cry* mutant flies fed with THIP (B) or CBZ (E) under 4L20D in A and D, respectively. White box indicates light period while black box indicates dark period. (F, H) Daily sleep duration of male *cry* RNAi and control flies fed with THIP (F) ($n = 72, 74, 95, 49, 54, 23, 30, 27, 27, 20$ flies) or CBZ (H) ($n = 72, 74, 95, 49, 54, 18, 22, 26, 30, 24$ flies) under 4L20D. For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, $***P < 0.001$; compared to UAS control, $##P < 0.01$, $###P < 0.001$. For comparing to vehicle control, two-tailed Student's *t*-test was used, $$$$P < 0.001$. (G, I) Sleep profile of male *cry* RNAi and control flies fed with THIP (G) or CBZ (I) under 4L20D in F and H, respectively. White box indicates light period while black box indicates dark period. (J) Daily sleep duration of male *rd^{MD-RR}* flies with *cry* knocked down in GABAergic neurons under 4L20D, along with relevant controls ($n = 72, 73, 66, 44, 45, 33, 45, 54, 38, 33$ flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, $***P < 0.001$; compared to UAS control, $###P < 0.001$. For comparing to WT or UAS/cryRNAi/GAL4 control, two-tailed Student's *t*-test was used: compared to WT, $&P < 0.05$; compared to UAS/cryRNAi/GAL4 control, $$$$P < 0.001$. (K, L) Sleep profile of male *rd^{MD-RR}* flies with *cry* knocked down in GABAergic neurons using *cry*RNAi-1 (K) or *cry*RNAi-2 (L) under 4L20D in J. Error bars represent SEM; G4, GAL4; U, UAS.

CRY acts upon l-LNvs to promote wakefulness

Previous studies reported that GABA enhances sleep at least in part by activating RDL on the l-LNvs, thus inhibiting the activities of these cells and reducing the release of the arousal-promoting neuropeptide pigment dispersing factor (PDF) (Chung et al., 2009 [↗](#); Parisky et al., 2008 [↗](#)). Therefore, we tested whether the l-LNvs mediate the effects of CRY on sleep/wakefulness. We first tested whether the l-LNvs indeed receive projections from GABAergic neurons. We expressed a GFP-tagged synaptotagmin (*synt-GFP*) which labels axon terminals using *Gad1GAL4*, and GFP signal was observed at the l-LNv soma (Figure 4—figure supplement 1A [↗](#)) (Zhang et al., 2002 [↗](#)). We employed the GFP Across Synaptic Partners (GRASP) method with enhanced specificity (t-GRASP) to check for synaptic connections between GABAergic neurons and l-LNvs (Shearin et al., 2018 [↗](#)). Pre-mGRASP was expressed in the axon terminal of GABAergic neurons (using *Gad1GAL4*), whereas its t-GRASP partner post-mGRASP was expressed in the l-LNvs (using the driver *PdfLexA*) (Shang et al., 2008 [↗](#)). GFP signal can be detected at the l-LNv soma, which means GABAergic neurons may send synaptic projections to the l-LNvs (Figure 4—figure supplement 1B [↗](#)). We further validated this connection using the *trans*-Tango technique which labels down-stream synaptic targets of GABAergic neurons with the HA tag, and we were able to observe HA expression in the l-LNvs (Figure 4—figure supplement 1C [↗](#)) (Talay et al., 2017 [↗](#)). All in all, these results strongly suggest that GABAergic neurons project to the l-LNvs and form synaptic connections. Consistently, knocking down *Rdl* in the l-LNvs rescues the long sleep phenotype of *cry* mutants (Figure 4—figure supplement 1D [↗](#)).

We next examined the effects of *cry* mutation on GABA level at the l-LNv soma by immunostaining. As expected, GABA signal is significantly increased in *cry* mutants under 4L20D but not 12L12D (Figure 4A-D [↗](#)). Consistent with elevated GABA, GCaMP6m signal, an indicator of intracellular calcium concentration, is significantly reduced in the l-LNvs under 4L20D (Figure 4E and F [↗](#)) (Chen et al., 2013 [↗](#)). This implies that the neural activity of these cells is down-regulated, possibly due to increased GABA signaling.

To test whether the long sleep phenotype in *cry* mutants is due to reduced activity of the l-LNvs, we electrically silenced these neurons by expressing an inward rectifying potassium channel Kir2.1 (Baines et al., 2001 [↗](#)). This results in lengthened sleep duration under 4L20D, mimicking the sleep phenotype of *cry* mutants (Figure 4G [↗](#); Figure 4—figure supplement 2A [↗](#)). On the other hand, activating the l-LNvs by expressing the temperature-gated depolarizing cation channel TrpA1 or the bacterial depolarization-activated sodium channel NachBac blocks the sleep enhancing effect of *cry* mutation (Figure 4H-K [↗](#); Figure 4—figure supplement 2B-E [↗](#)) (Hamada et al., 2008 [↗](#); Nitabach et al., 2006 [↗](#)). These findings support the notion that *cry* mutation lengthens sleep duration by down-regulating the neural activity of the l-LNvs. Since it has been shown that the l-LNvs promote arousal by releasing PDF, we assessed whether PDF or its receptor PDFR is required by CRY to exert influence on sleep/wakefulness (Chung et al., 2009 [↗](#); Parisky et al., 2008 [↗](#)). *cry* mutation fails to lengthen sleep duration on *Pdf* or *Pdfr* mutant background, indicating the necessity of PDF signaling in mediating the arousal-promoting function of CRY (Figure 4L and M [↗](#); Figure 4—figure supplement 2F and G [↗](#)) (Hyun et al., 2005 [↗](#); Lear et al., 2005 [↗](#); Mertens et al., 2005 [↗](#); Renn et al., 1999 [↗](#)).

To summarize, these results strongly suggest that CRY promotes wakefulness by reducing GABA signaling and thus increasing the activity of the l-LNvs, which may in turn lead to increased release of PDF.

CRY acts in the s-LNvs to promote wakefulness

To identify the subset of GABAergic neurons in which CRY functions to regulate wakefulness, we first examined the expression pattern of *Gad1GAL4* by labeling the GAL4⁺ cells with a nuclear GFP (nls-GFP) (Shiga et al., 1996 [↗](#)). While the l-LNvs are not GFP⁺, we noticed that the s-LNvs which are close to the l-LNvs and also express PDF appear to be GFP⁺ (Figure 5A and B [↗](#); Figure 5-figure supplement 1A [↗](#)). In addition, nls-GFP was expressed in GABAergic neurons (using *Gad1GAL4*), whereas RFP was expressed in the l- and s-LNvs (using the driver *PdfLexA*). GFP signal can be




Figure 4. CRY increases the neural activity of the l-LNVs to promote wakefulness.

(A-D) Brains from male WT and *cry* mutant flies dissected at ZT1, 7, 13, 19 under 4L20D (A) or 12L12D (C) are immunostained with GABA (cyan) and PDF (magenta) antisera, and representative I-LNVs are displayed. Merged signal is shown as yellow. Bar graphs represent normalized GABA intensity in the I-LNVs under 4L20D (B) (n = 26-56 cells) and 12L12D (D) (n = 32-128 cells). The average value of the control group at ZT1 is set to 1. (E) Representative live image of the I-LNVs expressing GCaMP6m and tdTomato using *Pdf*GAL4. Brain samples are dissected at the indicated time points under 4L20D. (F) Quantification of GCaMP6m signal intensity normalized to that of tdTomato (n = 25-54 cells). Student's *t*-test: **P* < 0.05, ***P* < 0.01, ****P* < 0.001. (G) Daily sleep duration of male flies expressing Kir2.1 in PDF neuron using *Pdf*GAL4 and controls, monitored under 4L20D (n = 32, 42, 30 flies). One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, **P* < 0.05; compared to UAS control, *****P* < 0.001. (H, I) Daily sleep duration of male *cry* mutant flies expressing TrpA1 in the I-LNVs using c929GAL4 (H) (n = 42, 42, 38, 35, 45, 37 flies) or R10H10GAL4 (I) (n = 62, 62, 17, 30, 29, 30 flies) and relevant controls, monitored under 4L20D and 29°C to activate TrpA1. For comparison with UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, ****P* < 0.001; compared to UAS control, *****P* < 0.001. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: &*P* < 0.05, &&*P* < 0.001. (J, K) Daily sleep duration of *cry* mutant flies expressing NachBac in the I-LNVs using c929GAL4 (J) (n = 45, 30, 32, 48, 55, 48 flies) or R10H10GAL4 (K) (n = 25, 59, 29, 74, 25, 32 flies) and relevant controls, monitored under 4L20D. For comparison with UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, **P* < 0.05, ****P* < 0.001; compared to UAS control, *****P* < 0.001. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: &*P* < 0.05, &&*P* < 0.001. (L, M) Daily sleep duration of male *Pdf⁰¹-cry^b* (L) (n = 58, 91, 89, 31 flies) and *Pdf^rhan5304;cry^b* (M) (n = 58, 91, 31, 31 flies) mutants along with relevant controls, monitored under 4L20D. Two-tailed Student's *t*-test: compared to WT background, ****P* < 0.001; compared to *Pdf⁰¹* or *Pdf^rhan5304*, *****P* < 0.001; compared to *cry^b*, &&*P* < 0.01, &&&*P* < 0.001. The scale bar represents 15 μm. Error bars represent SEM. G4, GAL4; U, UAS.

detected in the s-LNvs but not the l-LNvs (Figure 5-figure supplement 1B [↗](#)). Because the s-LNvs are known to express CRY, we suspected that CRY may be acting in these s-LNvs to regulate the activity of the l-LNvs via GABA signaling (Benito et al., 2008 [↗](#); Yoshii et al., 2008 [↗](#)). To test this idea, we knocked down *cry* in the s-LNvs using R6GAL4 while over-expressing *dicer2* (*dcr2*) to enhance RNAi efficiency and observed a modest but significant lengthening of sleep duration (Figure 5C [↗](#); Figure 5—figure supplement 2A [↗](#)) (Helfrich-Forster et al., 2007 [↗](#)). However, when we adopted a *Pdf*GAL80 to block the actions of GAL4 in the PDF neurons in *Gad1*GAL4/UAS*cry*RNAi, this does not alter the long-sleep phenotype (Figure 5—figure supplement 2B [↗](#)) (Stoleru et al., 2004 [↗](#)). These series of results suggest that *cry* expression in the s-LNvs are necessary but not sufficient to maintain normal sleep/wakefulness. We were indeed able to detect GABA signal at the s-LNvs, while GABA intensity is enhanced in *cry* mutants under 4L20D but not 12L12D (Figure 5D-G [↗](#)). To further validate that the s-LNvs are GABAergic, we knocked down VGAT in these cells and observed a decrease of GABA intensity (Figure 5—figure supplement 3A and B [↗](#)).

Next, we examined the effects of *cry* deficiency on the activity level of the s-LNvs. We found that *cry* mutation increases calcium concentration in these cells under 4L20D, in stark contrast to that of the l-LNvs (Figure 5H and I [↗](#)). This strongly suggests that *cry* deficiency leads to elevated neural activity in the s-LNvs. Consistently, we observed increased sleep when we activated these cells using TrpA1, similar to the long-sleep phenotype of *cry* mutants (Figure 5J [↗](#); Figure 5—figure supplement 4A [↗](#)). Because R6GAL4 is also expressed in several other cells in the brain, we combined *Pdf*GAL80 to verify that the sleep-enhancing effect is caused by over-activation of the s-LNvs (Helfrich-Forster et al., 2007 [↗](#)). Indeed, we no longer observed the long-sleep phenotype when GAL4 expression is inhibited in the PDF neurons by GAL80 (Figure 5J [↗](#); Figure 5—figure supplement 4A [↗](#)). Moreover, we found that *cry* mutation can no longer exert effects on sleep when the s-LNvs are activated, further validating that the s-LNvs mediate the influences of CRY on sleep (Figure 5K [↗](#); Figure 5—figure supplement 4B [↗](#)). Because CRY has been shown to regulate membrane depolarization and input resistance together with the redox sensor of the voltage-gated potassium channel β -subunit HYPERKINETIC (HK), we tested whether CRY and HK also function together in the s-LNvs to regulate arousal (Agrawal et al., 2017 [↗](#); Fogle et al., 2015 [↗](#)). We found that while over-expressing *hk* in the s-LNvs of WT flies does not alter sleep duration under 4L20D, over-expressing it in *cry* mutants significantly increases sleep (Figure 5L [↗](#); Figure 5—figure supplement 4C [↗](#)). This genetic interaction indicates that CRY and HK cooperate to regulate wakefulness, possibly by modulating the electric activity of the s-LNvs.

Next, we expressed syt-GFP in the s-LNvs and observed GFP signal at the soma of the l-LNvs, indicating that the s-LNvs send axonal terminals to the l-LNvs (Figure 6A [↗](#)). *trans*-Tango technique further demonstrated that the s-LNvs project to form synaptic connections with the l-LNvs (Figure 6B [↗](#)). Moreover, when we disrupted GABA transmission in the s-LNvs by knocking down VGAT or *Gad1*, this reduced GABA intensity in the l-LNvs, implicating that the s-LNvs release GABA onto the l-LNvs (Figure 6C-E [↗](#)). At the behavioral level, these manipulations suppress the long-sleep phenotype of *cry* mutation (Figure 6F-I [↗](#)).

Taken together, these results suggest that *cry* deficiency in the s-LNvs results in increased neuronal activity and thus enhanced GABA release at the l-LNvs via direct synaptic projections, ultimately leading to decreased wakefulness and increased sleep.

Short photoperiod reduces GABA level at the l-LNvs

Our findings thus far point to an inhibitory role of CRY on the GABAergic tone, and this in turn removes the inhibition on the neural activities of the l-LNvs. Since CRY is degraded by light, we hypothesized that CRY should exert a stronger effect on GABA under short photoperiod (Emery et al., 1998 [↗](#)). Consistently, GABA intensity does appear to be down-regulated at the l-LNvs under 4L20D vs. 12L12D (Figure 7A and B [↗](#)). On the other hand, GABA intensity does not exhibit photoperiod-dependent alteration in the s-LNvs (Figure 7C and D [↗](#)). Presumably, this reduced GABA level at the l-LNvs under 4L20D can result in elevated activation of these cells and increased wakefulness. In line with this, WT flies display shortened sleep duration under 4L20D compared to




Figure 5. CRY acts in the s-LNvs to inhibit their neural activity and promote wakefulness.

(A) Brains of male flies expressing nls-GFP driven by *Gad1*GAL4 maintained under 4L20D and immunostained with PDF antisera. Representative I-LNv and s-LNv are displayed. (B) Quantification of GFP signal intensity of PDF neuron ($n = 20, 49, 43$ cells). Two-tailed Student's t test: $***P < 0.001$. (C) Daily sleep duration of flies with *cry* knocked down in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D ($n = 45, 60, 63, 34, 39$ flies). (D-G) Brains from male WT and *cry* mutant flies dissected at ZT1, 7, 13, 19 under 4L20D (D) or 12L12D (F) are immunostained with GABA (cyan) and PDF (magenta) antisera, and representative s-LNvs are displayed. Merged signal is shown as yellow. Bar graphs represent normalized GABA intensity in the s-LNvs under 4L20D (E) ($n = 20-53$ cells) and 12L12D (G) ($n = 29-67$ cells). The average value of the control group at ZT1 is set to 1. (H) Representative live image of the s-LNvs expressing GCaMP6m and tdTomato using *Pdf*GAL4. Brain samples are dissected at the indicated time points under 4L20D. (I) Quantification of GCaMP6m signal intensity normalized to that of tdTomato ($n = 29-47$ cells). Two-tailed Student's t -test: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. (J) Daily sleep duration of male flies expressing TrpA1 in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D and 29°C to activate TrpA1 ($n = 45, 54, 29, 22, 56, 22$ flies). (K) Daily sleep duration of male *cry* mutant flies expressing TrpA1 in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D and 29°C to activate TrpA1 ($n = 33, 42, 23, 35, 36, 20$ flies). (L) Daily sleep duration of male *cry* mutant flies over-expressing HK in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D ($n = 24, 29, 23, 30, 37, 22$ flies). For comparison with UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, $*P < 0.05$, $***P < 0.001$; compared to UAS control, $##P < 0.01$, $###P < 0.001$. For comparison between mutant vs. control, two-tailed Student's t -test was used: compared to WT background, $&&P < 0.01$, $&&&P < 0.001$. The scale bar represents 15 μ m. Error bars represent SEM. G4, GAL4; U, UAS; NS, not significant.




Figure 6. s-LNvs release GABA onto the I-LNvs.

(A) I-LNvs of male flies expressing syt-GFP in the s-LNvs using R6GAL4 maintained under 4L20D and immunostained with PDF antisera. (B) I-LNvs of male flies expressing trans-Tango in the s-LNvs using R6GAL4 maintained under 4L20D and immunostained with HA and PDF antisera. (C) Brains from male flies with VGAT (top) or *Gad1* (bottom) knocked down in the s-LNvs using R6GAL4 and controls maintained under 4L20D and immunostained with GABA (cyan) and PDF (magenta) antisera. Representative I-LNvs are displayed. Merged signal is shown as yellow. (D, E) Bar graphs represent normalized GABA intensity in the I-LNvs of flies with VGAT (D) ($n = 29, 37$ cells) or *Gad1* (E) ($n = 37, 51$ cells) knocked down in the s-LNvs using R6GAL4, monitored under 4L20D. The average value of the control group is set to 1. Two-tailed Student's *t*-test: $**P < 0.01$, $***P < 0.001$. (F) Daily sleep duration of male *cry* mutant flies with VGAT knocked down in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D ($n = 31, 31, 32, 29, 43, 19$ flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, $*P < 0.05$, $**P < 0.01$; compared to UAS control, $###P < 0.01$. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: compared to WT background, $&&&P < 0.001$. (G) Sleep profile of male *cry* mutant flies with VGAT knocked down in the s-LNvs using R6GAL4 and relevant controls in F, monitored under 4L20D. (H) Daily sleep duration of male *cry* mutant flies with *Gad1* knocked down in the s-LNvs using R6GAL4 and relevant controls, monitored under 4L20D ($n = 59, 59, 53, 60, 62, 45$ flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$; compared to UAS control, $\#P < 0.05$, $\#\#P < 0.01$. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: compared to WT background, $&&&P < 0.001$. (I) Sleep profile of male *cry* mutant flies with *Gad1* knocked down in the s-LNvs using R6GAL4 and relevant controls in H, monitored under 4L20D. The scale bar represents 15 μm . Error bars represent SEM. G4, GAL4; U, UAS; NS, not significant.

12L12D (Figure 7E and F). This sleep reduction is a result of decreased sleep bout length but not bout number, while wake activity is not altered by photoperiod (Figure 7G-I). These observations indicate that short photoperiod hampers sleep maintenance, similar to the effects of CRY.

Discussion

Previous studies have shown that CRY mediates light-induced electrical activity of the l-LNvs and acute arousal, which indicates that CRY can immediately act to promote wake and terminate sleep in response to light pulse (Fogle et al., 2015; Sheeba et al., 2008). At the molecular level, this is believed to be accomplished by a direct coupling of light-activated CRY with HK in the l-LNvs (Fogle et al., 2015). Here we found that CRY promotes extended wakefulness under short photoperiod by functioning as an inhibitor of GABAergic tone. In contrast to the previously characterized roles of CRY which are activated by light, we believe this novel function we identify here reflects a role for CRY in the dark. Under short photoperiods, CRY acts in the dark to inhibit GABA signaling and thus promote wakefulness, leading to reduced sleep during the dark phase compared to longer photoperiods. Indeed, CRY is an ideal signal for conveying photoperiodic information, for it is degraded by light and only accumulates during darkness (Emery et al., 1998). It can measure the length of the day/night and thus instruct downstream signaling components to modulate photoperiod-dependent processes. Consistent with this idea, the largest increases of sleep in *cry* mutants occur immediately after lights-off and prior to lights-on (Fig 1B), indicating that CRY functions to promote wakefulness during these time windows which are exactly the time windows that would be affected by photoperiod changes (i.e. there will be light during these time windows under longer photoperiods). In other words, CRY appears to act at the time of the day that is most sensitive to photoperiodic changes, which fits perfectly with the idea that it serves as an instructive signal of day/night length.

While EOS and NipA treatment lengthen sleep duration as previously reported, we are somewhat surprised by the observation that knocking down *Gad1* or *VGAT* in GABAergic neurons also extends sleep duration (Ki and Lim, 2019; Leal and Neckameyer, 2002). We reason this may be due to some sort of over-compensation induced by chronic GABA deficiency to maintain excitation/inhibition balance, as previous studies have reported that the amplitude of glutamatergic current is substantially down-regulated in *Gad1* mutant flies and increased in *Gad1* over-expression flies (Featherstone et al., 2000), (Featherstone et al., 2002). Consistent with this idea, we noticed when *cry* is knocked down in *Gad1/VGAT* RNAi flies, sleep duration is shortened and comparable to that of the controls. This is probably because *cry* depletion enhances the GABAergic tone, thus normalizing GABA signaling in these flies which results in normal sleep duration.

We acknowledge that THIP treatment leads to prominent lengthening of sleep duration, and thus it is possible in this case *cry* deficiency no longer increases sleep duration due to a ceiling effect rather than epistatic interaction. Nonetheless, considering that CBZ and *rdl* mutation also block the effect of *cry* deficiency on sleep duration, it is highly likely that GABA-A receptor mediates the influences of CRY on sleep. One caveat is that the *rdl*^{MD-RR} mutation has been shown to diminish the desensitization of GABA-A receptor and is thus believed to be a gain-of-function allele, but here we found that it eliminates the effect of *cry* RNAi on sleep (Zhang et al., 1994). Since treatment with GABA-A agonist THIP leads to substantially increased sleep while under the same condition *rdl*^{MD-RR} mutation does not appear to significantly alter sleep duration, we suspect that similar to *Gad1/VGAT* RNAi, chronic enhancement of GABA-A function associated with *rdl*^{MD-RR} mutation may also trigger some kind of compensatory mechanism that counteracts this increased GABA-A activity. Consequently, the influences of *cry* RNAi on sleep are blocked. Of course, a lot more in depth characterizations are required to elucidate these issues.

The s-LNvs have been reported to receive GABAergic inputs possibly via GABA-B receptors, but have not been shown to be able to synthesize GABA (Dahdal et al., 2010; Hamasaka et al., 2005). One study using cell-type specific gene expression profiling demonstrates *Gad1* and *VGAT* expression in both s-LNvs and l-LNvs, although with relatively low signal (Nagoshi et al., 2010).




Figure 7. Short photoperiod reduces GABA level at the I-LNvs and increases sleep duration.

(A-D) Brains from male WT flies dissected at ZT1, 7, 13, 19 under 4L20D or 12L12D are immunostained with GABA (cyan) and PDF (magenta) antisera. Representative I-LNvs (A) and s-LNvs (C) are displayed. Merged signal is shown as yellow. Bar graphs represent normalized GABA intensity in the I-LNvs (B) ($n = 28-55$ cells) and s-LNvs (D) ($n = 21-59$ cells). The average value of the control group at ZT1 is set to 1. Student's t -test: *** $P < 0.001$. (E-I) The daily sleep duration (E), sleep bout number (F), average sleep bout length (G), waking activity (H) and sleep profile (I) of male WT flies under 4L20D and 12L12D ($n = 91, 32$ flies). Two-tailed Student's t -test: *** $P < 0.001$. The scale bar represents 15 μ m. Error bars represent SEM. (J) A model demonstrating how CRY regulates the GABAergic s-LNv/I-LNv circuitry under short vs. longer photoperiods to promote arousal.

Here we observed that *Gad1*GAL4 is expressed in the s-LNvs, and their GABA intensity is reduced when we use R6GAL4 to knock down *VGAT* in these cells. R6GAL4 drives prominent expression in the s-LNvs with very little if any expression in the l-LNvs, and weaker (and not very consistent) expression in several other neurons in the protocerebrum, pars intercerebralis and subesophageal area which are all believed to lie outside of the circadian neuron network (Helfrich-Forster et al., 2007). Therefore, we reason that the alteration of GABA signal associated with knocking down *VGAT* should arise from *VGAT* deficiency within the s-LNvs. Taken together, these results suggest that the s-LNvs are GABAergic. Although knocking down *VGAT* or *Gad1* in these cells can suppress the long sleep phenotype of *cry* mutants, knocking down *cry* in these cells only lead to a modest lengthening of sleep duration. Moreover, inhibiting *cry* RNAi expression in PDF neurons does not eliminate the long-sleep phenotype of *Gad1*GAL4/UAS*cry*RNAi flies. Therefore, we suspect that *cry* deficiency in other GABAergic neurons are also required for the long-sleep phenotype. Given that the s-LNvs are known to express CRY and appear to be GABAergic based on our findings here, we believe that CRY acts at least in part in the s-LNvs to promote wakefulness under short photoperiod.

The molecular mechanism by which CRY down-regulates the GABAergic tone remains unclear. Besides conducting light-driven depolarization via HK, CRY has also been shown to act in synergy with HK to prevent membrane input resistance from falling to a low level in larval salivary glands (Agrawal et al., 2017; Fogle et al., 2015). In contrast to previous studies, here we found that lack of CRY increases the activity of the s-LNvs. Instead of functioning in synergy with HK, CRY appears to act in the opposite direction of HK as over-expressing *hk* enhances the long-sleep phenotype caused by *cry* mutation. We reason that the coupling between CRY and HK as well as their influences on the electric activity in the s-LNvs may be different from that of the l-LNvs and larval salivary glands. Nonetheless, our results also support an interaction between CRY and HK to promote arousal. We suspect that CRY acts via HK to inhibit the activity of the s-LNvs, which results in decreased GABA release and dis-inhibition of the l-LNvs. Extensive further investigations will be needed to elucidate the mechanism by which CRY regulates the activity of the s-LNvs.

In conclusion, here we describe a CRY-controlled GABAergic circuitry involving the l-LNvs and the s-LNvs that adjusts sleep duration in adaptation to changes in day length and propose a mechanistic explanation regarding how this circuitry functions (Fig 7). Under short photoperiods, more CRY accumulates and inhibits the activity of the GABAergic s-LNvs, leading to a dis-inhibition of the l-LNvs which can release more PDF and promote arousal. Under longer photoperiods, on the other hand, less CRY can accumulate and thus the s-LNvs will exert more inhibitory influences on the l-LNvs, leading to decreased release of PDF and wakefulness. Notably, almost all neurons in the mammalian pacemaker, the suprachiasmatic nucleus, are GABAergic, and GABA/GABA-A signaling have been shown to mediate neuronal coupling in response to photoperiod changes (Ono et al., 2021). We believe a similar GABAergic circuitry may exist in the mammalian system that adjusts sleep/wakefulness to photoperiodic changes.

Materials and Methods

Fly strains

All strains were obtained from Bloomington *Drosophila* Stock Center, Vienna *Drosophila* Resource Center and TsingHua Fly Center or as gifts from colleagues. Except for *Gad1*GAL4-2, all neurotransmitter related GAL4 lines were generated in Dr. Yi Rao's laboratory (Deng et al., 2019). The following fly strains were used in this study: *cry*^b (BDSC:80921), *per*⁰ (BDSC:80917), *tim*⁰ (BDSC:80922), *clk*^{rk} (BDSC:80927), *cyc*⁰ (BDSC:80929), *Pdf*⁰¹ (BDSC:26654), *Pdf*^{han5304} (BDSC:33068), UAS-*cry*RNAi-1 (THU4761), UAS-*cry*RNAi-2 (THU5148), *cry*⁰³ (Dolezelova et al., 2007), UAS-*gad1*RNAi-1 (THU2920), *Rdl*^{MD-RR} (BDSC:1675), UAS-*VGAT*RNAi-1 (THU4303), UAS-*VGAT*RNAi-2 (V45916), UAS-*hk* (BDSC:86270), isogenic *w*¹¹¹⁸ (BDSC:5905), UAS-*NachBac* (BDSC:9469), *Pdf*GAL4 (BDSC:6899), *VGAT*GAL4 (BDSC:58980), UAS-*TrpA1* (BDSC:26263), UAS-*Kir2.1* (BDSC:6596), UAS-*dcr2* (V6009), UAS-*sytGFP* (BDSC:6925), UAS-*nlsGFP* (BDSC:4775),

UAS-*myrGFP*, UAS-*mtdtomato*(3XHA); *Trans-tango* (BDSC:77480) 13XLexAop2-*post-t-GRASP*, 20XUAS-*pre-mGRASP* (BDSC:79040), LexAop-rCD2-RFP (Perisse et al., 2013 [DOI](#)), *Gad1*GAL4-1, *Gad1*GAL4-2 (BDSC:51630), *Dop2*RGAL4, *TH*GAL4, *Trh*GAL4, *Ado*RGAL4, *DAT*GAL4, *GABAR1*GAL4, *GABAR3*GAL4, *SerT*GAL4, *GluRI*A GAL4, *R6*GAL4 (Helfrich-Forster et al., 2007 [DOI](#)), *c929*GAL4 (Park et al., 2008 [DOI](#)), *R10H10*GAL4 (Sekiguchi et al., 2020 [DOI](#)), *R78G01*GAL4 (Yoshii et al., 2015 [DOI](#)), *Pdf*GAL80 (Stoleru et al., 2004 [DOI](#)), UAS-GCaMP6m-Tdtomato (Chen et al., 2013 [DOI](#)) and *Pdf*-LexA (Shang et al., 2008 [DOI](#)). All flies used for sleep monitoring were backcrossed with the isogenic *w*¹¹¹⁸ strain for at least 5 times except for *cry*⁰³ which was backcrossed three times. All experiments were conducted in male flies unless otherwise specified.

Fly sleep monitoring and analysis

Flies were raised on standard cornmeal-yeast-sucrose medium and kept in 12L12D at 25°C until behavior monitoring. ~3 to 4-day-old flies were entrained under different photoperiods at 25°C for 4 days, and then their activities in the next 3 days were analyzed. Sleep is defined as 5 min consecutive inactivity. Sleep was analyzed with Counting Macro written in Excel (Microsoft) following previously published protocol (Pfeiffenberger et al., 2010 [DOI](#)). Flies were fed with agar-sucrose food (2% agar, 5% sucrose) during entire sleep monitoring. *TrpA1* flies were raised at 21°C and baseline sleep was monitored at 21°C. Temperature was then raised at lights on to 29°C for further sleep monitoring.

Drug treatment

For pharmacological experiments, drugs were mixed in the fly food at the following concentrations. For nipecotic acid (10 mg/ml, Sigma) and EOS (10 mM, Sigma), drugs were fed during the entire sleep monitoring. For THIP (10 µg/ml, Sigma), SKF-97541 (10 µg/ml, Tocris) and CBZ (0.15 mg/ml, Sinopharm Chemical Reagent), drugs were fed for 1 day after baseline sleep monitoring. The same amount of solvent was added into the fly food as vehicle control.

RNA extraction and quantitative real-time PCR (qRT-PCR)

Approximately 50 5-day-old flies were collected and frozen immediately on dry ice. Fly heads were isolated and homogenized in Trizol reagent (Life Technologies). Total RNA was extracted and qRT-PCR conducted following our previously published procedures (Bu et al., 2019 [DOI](#)). The following primers were used to quantify mRNA level, *cry* F: TGCAGGTACCAAGAATGTGG, R: GTCCACGTCCATCAGTTGC. *Gad1* F: TGCCACCACATTGAAGTACC, R: GGTGAACATGTTGGTGTTCG. *VGAT* F: ACGGCTTTAGCAAGGTAGC, R: TTGGAATTCGTCGATTTTGC.

Immunostaining

Male flies were entrained for 3 days under indicated photoperiod and collected on Day 4. flies were anesthetized with CO₂ and dissected and fixed with 4% paraformaldehyde diluted in PBS at the indicated time points. The brains were then fixed with 4% paraformaldehyde for 30 minutes. Samples were washed with PBT (PBS with 0.3% TritonX-100) for 3 × 10min. Then samples were incubated in PBS with 1% TritonX-100 for 20 min at room temperature. Brain samples were then blocked in PBT with 5% fetal bovine serum (Hyclone) for 30 minutes and subsequently incubated with mouse anti-PDF (1:100, DSHB), rabbit anti-GABA (1:200, Sigma), and anti-HA (1:100, DSHB) for 2-4 days at 4°C (4 days for anti-GABA and 2 days for the other antibodies). After PBT rinses for 3 times, the brains were incubated with donkey anti-mouse Alexa Fluor-594 (1:1000, Life Technologies), donkey anti-rabbit Alexa Fluor-488 (1:1000, Abcam) and donkey anti-rabbit Alexa Fluor-647 (1:1000, Abcam) overnight at 4°C. Then the brains were rinsed 3 times in PBS and mounted and imaged using Olympus FV3000 confocal microscope with a 60× objective lens. The intensity of GABA signal was quantified by ImageJ software. For each cell, the image slice with the strongest signal in the Z stack was selected and average intensity was quantified. A region on the same image slice was then selected as background and its intensity value was subtracted from the average intensity value of the cell. This subtracted value was subsequently normalized to the

average intensity of the control group as indicated in figure legends. Sample sizes in the legends indicate number of cells examined per genotype. For *trans*-Tango experiment, offsprings were raised at 18°C for 3-4 weeks before HA and PDF immunostaining.

Calcium imaging

For live imaging experiments (GCaMP and tdTomato), flies 2-3 days old were collected into tubes with standard food and entrained under LD for 3 days. Flies were anesthetized with CO₂ and brains were dissected at the indicated time points in *Drosophila* adult hemolymph-like saline solution. The dissected brain samples were put on glass slide and sealed with cover slide. The duration for dissection and microscopy should be completed within 0.5 hours for each time point. Images were captured with Olympus FV3000 confocal microscopy with a 20× objective lens. The intensity of GCaMP and tdTomato signals were quantified by ImageJ software. Sample sizes in the legends indicate number of cells examined per genotype.

Statistical Analysis

For data that fit normal distribution, two-tailed Student's *t*-test (Microsoft Excel) was used to compare the difference between two genotypes. For data that do not fit normal distribution, Mann-Whitney test (Prism Graphpad) was used to compare the difference between two genotypes. For multiple comparisons, one-way ANOVA with Bonferroni multiple comparison test (Prism Graphpad) was used. Sample size, statistical test, and significance values are indicated in figure legends. Sample size is determined based on previous studies with similar experimental assays.




Figure 1—figure supplement 1. *cry* knock-out mutation lengthens sleep duration under 4L20D. Daily sleep duration of male WT flies and *cry* knock-out flies, monitored under 4L20D (n = 53, 35, 47). Student's *t*-test: **P* < 0.05. Error bars represent SEM.




Figure 2—figure supplement 1. CRY does not regulate sleep duration via TIM.

Daily sleep duration of male WT flies and flies mutant for *cry* and/or *tim*, monitored under 4L20D (n = 58, 63, 91, 33 flies). Two-tailed Student's *t*-test: compared to WT background, ****P* < 0.001; compared to *tim*⁰, ###*P* < 0.001. Error bars represent SEM.




Figure 2—figure supplement 2. Screening for anatomical substrates that mediate the effects of CRY on sleep/wakefulness.

(A) Difference in sleep duration between male *cry* RNAi flies compared to GAL4 and UAS controls under 4L20D (n = 20-92). (B) Plots of relative mRNA abundance for *cry* determined by qRT-PCR in whole-head extracts of *cry*RNAi and control flies under 4L20D (n = 4). The value of the GAL4 control group was set to 1. Mann-Whitney test: compared to GAL4 control, **P* < 0.05; compared to UAS control, #*P* < 0.05. G4, GAL4; U, UAS.




Figure 2—figure supplement 3. CRY promotes wakefulness in GABAergic neurons.

(A-D) Sleep profile (A), daily waking activity (B), sleep bout number (C), average sleep bout length (D) of male flies with *cry* knocked down in GABAergic neurons by VGATG4L4 monitored under 4L20D (n = 72, 74, 96, 49, 54 flies). (E-H) Sleep profile (E), daily waking activity (F), sleep bout number (G), average sleep bout length (H) of male flies with *cry* knocked down in GABAergic neurons by *Gad1G4L4-1* monitored under 4L20D (n = 29, 26, 29, 23, 32 flies). (I-L) Sleep profile (I), daily waking activity (J), sleep bout number (K), average sleep bout length (L) of male flies with *cry* knocked down in GABAergic neurons by *Gad1G4L4-2* monitored under 4L20D (n = 34, 31, 31, 25, 23 flies). One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, ****P* < 0.001; compared to UAS control, #*P* < 0.05, ###*P* < 0.001.




Figure 2—figure supplement 4. CRY promotes wakefulness via GABA signaling.

(A, B) Sleep profile of male WT and *cry* mutant flies fed with NipA (A) or EOS (B) under 4L20D in Fig 2E. White box indicates light period while black box indicates dark period. (C) Sleep profile of *cry* RNAi and control flies fed with NipA under 4L20D in Fig 2F. (D) Sleep profile of male flies with *cry* and *gad1* knocked down in GABAergic neurons monitored under 4L20D in Fig 2G. (E) Sleep profile of male flies with *cry* and *VGAT* knocked down in GABAergic neurons monitored under 4L20D in Fig 2H. (F) Sleep profile of *cry* mutant flies with *VGAT* knocked down in GABAergic neurons monitored under 4L20D in Fig 2I. (G, H) Plots of relative mRNA abundance for *Gad1* (G) and *VGAT* (H) determined by qRT-PCR in whole-head extracts of RNAi and control flies under 4L20D (n = 6). The value of the GAL4 control group was set to 1. Mann-Whitney test: compared to GAL4 control, **P < 0.01; compared to UAS control, ###P < 0.01. (I) Daily sleep duration of male flies with *cry* and *gfp* knocked down in GABAergic neurons by *Gad1*GAL4 monitored under 4L20D (n = 72, 26, 29, 20, 30, 31, 68, 32, 31, 29, 29). One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, ***P < 0.001; compared to UAS control, #P < 0.05, ###P < 0.001. Error bars represent SEM; G4, GAL4; U, UAS; NS, not significant.




Figure 4—figure supplement 1. GABAergic neurons project to and form synapses with the I-LNvs.

(A) I-LNvs of flies expressing syt-GFP in GABAergic neurons using *Gad1Gal4* maintained under 4L20D and immunostained with PDF antisera. (B) I-LNvs of flies expressing Pre-mGRASP in GABAergic neurons using *Gad1Gal4* and t-GRASP in the I-LNvs using *PdfLexA* maintained under 4L20D. Brains are immunostained with PDF antisera. (C) I-LNvs of flies expressing trans-Tango in GABAergic neurons using *Gad1Gal4* maintained under 4L20D and immunostained with HA and PDF antisera. (D) Daily sleep duration of *cry* mutant flies with *rdl* knocked down in the I-LNvs using R78G01GAL4 and relevant controls, monitored under 4L20D (n = 31, 31, 64, 32, 31, 32 flies). For comparison between RNAi flies vs. UAS/GAL4 controls, one-way ANOVA with Bonferroni multiple comparison test was used: compared to GAL4 control, **P* < 0.05; compared to UAS control, ###*P* < 0.001. For comparison between mutant vs. control, two-tailed Student's *t*-test was used: compared to WT background, &*P* < 0.05, &&*P* < 0.001; compared to *Urd/RNAi/GAL4*, not significant. The scale bar represents 15 μ m. Error bars represent SEM. G4, GAL4; U, UAS; LexAop, LexA operator.

Figure 4—figure supplement 2. CRY promote wakefulness by impinging on the excitability of I-LNV and PDF signaling.

(A) Sleep profile of male flies expressing Kir2.1 in PDF neuron using *PdfGAL4* and controls, monitored under 4L20D in Fig 4G. White box indicates light period while black box indicates dark period. (B, C) Sleep profile of male *cry* mutant flies expressing TrpA1 in the I-LNVs using *c929GAL4* (B) or *R10H10GAL4* (C) and relevant controls in Fig 4H and I, monitored under 4L20D and 29°C to activate TrpA1. (D, E) Sleep profile of *cry* mutant flies expressing NachBac in the I-LNVs using *c929GAL4* (D) or *R10H10GAL4* (E) and relevant controls in Fig 4J and K, monitored under 4L20D. (F, G) Daily sleep duration of male *Pdf⁰¹-cry^b* (D) and *Pdfr^{han5304};cry^b* (E) mutants along with relevant controls in Fig 4L and M, respectively, monitored under 4L20D.





Figure 5—figure supplement 1. *Gad1GAL4* drives expression in the s-LNVs but not I-LNVs.

(A) Brains of male flies expressing nls-GFP driven by *Gad1GAL4* maintained under 4L20D and immunostained with PDF antisera. (B) Brains of male flies expressing nls-GFP in GABAergic neurons using *Gad1GAL4* and rCD2-RFP in the PDF neurons using *PdflLexA* maintained under 4L20D. The scale bar represents 15 μ m.






Figure 5—figure supplement 2. *cry* expression in the s-LNvs is necessary but not sufficient for maintaining normal sleep/wakefulness.

(A) Sleep profile of *cry* knocked down in s-LNvs by R6GAL4 monitored under 4L20D in Fig 5C. White box indicates light period while black box indicates dark period. (B) Daily sleep duration of *cry* knocked down in PDF- GABAergic neurons using *Gad1GAL4* and *PdfGAL80*, monitored under 4L20D (n = 101, 26, 29, 27, 31, 23, 32, 73, 63). One-way ANOVA with Bonferroni multiple comparison test: compared to GAL4 control, ****P* < 0.001; compared to UAS control, #*P* < 0.05, ###*P* < 0.001. Error bars represent SEM. G4, GAL4; G80, GAL80; U, UAS.

Figure 5—figure supplement 3. s-LNvs appear to be GABAergic neurons.

(A) Brains from male flies with *VGAT* knocked down in the s-LNvs using *R6GAL4* and control maintained under 4L20D and immunostained with GABA (cyan) and PDF (magenta) antisera. Representative s-LNvs are displayed. Merged signal is shown as yellow. (B) Bar graphs represent normalized GABA intensity in the s-LNvs with *VGAT* knocked down using *R6GAL4* ($n = 25, 32$ cells). The average value of the control group is set to 1. Two-tailed Student's *t*-test, $***P < 0.001$. The scale bar represents 15 μm . Error bars represent SEM. G4, *GAL4*; U, *UAS*.





Figure 5—figure supplement 4. CRY promotes wakefulness by impinging on the excitability of the s-LNvs.

(A) Sleep profile of male flies expressing *TrpA1* in the s-LNvs using *R6GAL4* and relevant controls in Fig 5J, monitored under 4L20D and 29°C to activate *TrpA1*. White box indicates light period while black box indicates dark period. (B) Sleep profile of male *cry* mutant flies expressing *TrpA1* in the s-LNvs using *R6GAL4* and relevant controls in Fig 5K, monitored under 4L20D and 29°C to activate *TrpA1*. (C) Sleep profile of male *cry* mutant flies over-expressing HK in the s-LNvs using *R6GAL4* and relevant controls in Fig 5L, monitored under 4L20D. Error bars represent SEM. G4, *GAL4*; G80, *GAL80*; U, *UAS*.



Data availability

All data generated or analyzed during this study are included in the manuscript and supporting files.

Acknowledgements

We would like to thank Drs. Chang Liu, Yi Rao, Liming Wang, Junhai Han, Zhihua Liu and Yi Zhong for kindly providing flies used in this study. We would also like to thank Dr. Chang Liu for helpful advice on GABA immunostaining. This work was supported by grants from the Ministry of Science and Technology of China STI 2030-Major Projects (2021ZD0203202 and 2024YFA1803201), Natural Science Foundation of China (82530044 and 32341021) and Science and Technology Department of Hubei Province (2022CFA049) to Luoying Zhang and grant from the Natural Science Foundation of China (32300984) to Lixia Chen.

Additional information

Author contributions

L.C. and L.Z. designed the experiments. L.C. and D.T. conducted the experiments. L.C., C.S. and L.Z. wrote the paper.

Funding

Funder	Grant reference number	Author
Ministry of Science and Technology of the People's Republic of China (MOST)	2021ZD0203202	Luoying Zhang
MOST National Natural Science Foundation of China (NSFC)	82530044	Luoying Zhang
National Natural Science Foundation of China	32341021	Luoying Zhang
Ministry of Science and Technology of the People's Republic of China (MOST)	2024YFA1803201	Luoying Zhang
MOST National Natural Science Foundation of China (NSFC)	32300984	Lixia Chen
Science and Technology Department of Hubei Province	2022CFA049	Luoying Zhang

Author ORCID iDs

Luoying Zhang: <https://orcid.org/0000-0001-5996-6975>

Additional files

Figure 1 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 1 [↗](#).

Figure 2 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 2 [↗](#).

Figure 3 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 3 [↗](#).

Figure 4 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 4 [↗](#).

Figure 5 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 5 [↗](#).

Figure 6 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 6 [↗](#).

Figure 7 source data [↗](#) Numerical data with associated statistical analyses underlying Figure 7 [↗](#).

References

1. **Agosto J.**, Choi J.C., Parisky K.M., Stilwell G., Rosbash M., Griffith L.C (2008) Modulation of GABA(A) receptor desensitization uncouples sleep onset and maintenance in *Drosophila*. *Nat Neurosci* **11**:354-359
2. **Agrawal P.**, Houl J.H., Gunawardhana K.L., Liu T., Zhou J., Zoran M.J., Hardin P.E (2017) *Drosophila* CRY Entrains Clocks in Body Tissues to Light and Maintains Passive Membrane Properties in a Non-clock Body Tissue Independent of Light. *Curr Biol* **27**:2431-2441.
3. **Allada R.**, Chung B.Y (2010) Circadian organization of behavior and physiology in *Drosophila*. *Annu Rev Physiol* **72**:605-624
4. **Allada R.**, White N.E., So W.V., Hall J.C., Rosbash M (1998) A mutant *Drosophila* homolog of mammalian Clock disrupts circadian rhythms and transcription of *period* and *timeless*. *Cell* **93**:791-804
5. **Baines R.A.**, Uhler J.P., Thompson A., Sweeney S.T., Bate M (2001) Altered electrical properties in *Drosophila* neurons developing without synaptic transmission. *J Neurosci* **21**:1523-1531
6. **Benito J.**, Houl J.H., Roman G.W., Hardin P.E. (2008) The blue-light photoreceptor CRYPTOCHROME is expressed in a subset of circadian oscillator neurons in the *Drosophila* CNS. *J Biol Rhythms* **23**:296-307
7. **Bu B.**, He W., Song L., Zhang L (2019) Nuclear Envelope Protein MAN1 Regulates the *Drosophila* Circadian Clock via *Period*. *Neurosci Bull* **35**:969-978
8. **Busza A.**, Emery-Le M., Rosbash M., Emery P (2004) Roles of the two *Drosophila* CRYPTOCHROME structural domains in circadian photoreception. *Science* **304**:1503-1506
9. **Ceriani M.F.**, Darlington T.K., Staknis D., Mas P., Petti A.A., Weitz C.J., Kay S.A (1999) Light-dependent sequestration of TIMELESS by CRYPTOCHROME. *Science* **285**:553-556
10. **Chen T.W.**, Wardill T.J., Sun Y., Pulver S.R., Renninger S.L., Baohan A., Schreiter E.R., Kerr R.A., Orger M.B., Jayaraman V., *et al.* (2013) Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**:295-300
11. **Chung B.Y.**, Kilman V.L., Keath J.R., Pitman J.L., Allada R (2009) The GABA(A) receptor RDL acts in peptidergic PDF neurons to promote sleep in *Drosophila*. *Curr Biol* **19**:386-390
12. **Dahdal D.**, Reeves D.C., Ruben M., Akabas M.H., Blau J (2010) *Drosophila* pacemaker neurons require G protein signaling and GABAergic inputs to generate twenty-four hour behavioral rhythms. *Neuron* **68**:964-977
13. **Deng B.**, Li Q., Liu X., Cao Y., Li B., Qian Y., Xu R., Mao R., Zhou E., Zhang W., *et al.* (2019) Chemoconnectomics: Mapping Chemical Transmission in *Drosophila*. *Neuron* **101**:876-893.
14. **Dolezelova E.**, Dolezel D., Hall J.C (2007) Rhythm defects caused by newly engineered null mutations in *Drosophila's cryptochrome* gene. *Genetics* **177**:329-345
15. **Emery P.**, So W.V., Kaneko M., Hall J.C., Rosbash M (1998) CRY, a *Drosophila* clock and light-regulated cryptochrome, is a major contributor to circadian rhythm resetting and photosensitivity. *Cell* **95**:669-679
16. **Enell L.**, Hamasaka Y., Kolodziejczyk A., Nassel D.R (2007) gamma-Aminobutyric acid (GABA) signaling components in *Drosophila*: immunocytochemical localization of GABA(B) receptors in relation to the GABA(A) receptor subunit RDL and a vesicular GABA transporter. *J Comp Neurol* **505**:18-31
17. **Featherstone D.E.**, Rushton E., Broadie K (2002) Developmental regulation of glutamate receptor field size by nonvesicular glutamate release. *Nat Neurosci* **5**:141-146
18. **Featherstone D.E.**, Rushton E.M., Hilderbrand-Chae M., Phillips A.M., Jackson F.R., Broadie K (2000) Presynaptic glutamic acid decarboxylase is required for induction of the postsynaptic receptor field at a glutamatergic synapse. *Neuron* **27**:71-84
19. **ffrench-Constant R.H.**, Steichen J.C., Rocheleau T.A., Aronstein K., Roush R.T. (1993) A single-amino acid substitution in a gamma-aminobutyric acid subtype A receptor locus is associated with cyclodiene insecticide resistance in *Drosophila* populations. *Proc Natl Acad Sci U S A* **90**:1957-1961

20. Fogle K.J., Baik L.S., Houl J.H., Tran T.T., Roberts L., Dahm N.A., Cao Y., Zhou M., Holmes T.C (2015) CRYPTOCHROME-mediated phototransduction by modulation of the potassium ion channel beta-subunit redox sensor. *Proc Natl Acad Sci U S A* **112**:2245-2250
21. Golden R.N., Gaynes B.N., Ekstrom R.D., Hamer R.M., Jacobsen F.M., Suppes T., Wisner K.L., Nemeroff C.B (2005) The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. *Am J Psychiatry* **162**:656-662
22. Hamada F.N., Rosenzweig M., Kang K., Pulver S.R., Ghezzi A., Jegla T.J., Garrity P.A (2008) An internal thermal sensor controlling temperature preference in *Drosophila*. *Nature* **454**:217-220
23. Hamasaka Y., Wegener C., Nassel D.R (2005) GABA modulates *Drosophila* circadian clock neurons via GABAB receptors and decreases in calcium. *J Neurobiol* **65**:225-240
24. Helfrich-Forster C., Shafer O.T., Wulbeck C., Grieshaber E., Rieger D., Taghert P (2007) Development and morphology of the clock-gene-expressing lateral neurons of *Drosophila melanogaster*. *J Comp Neurol* **500**:47-70
25. Hyun S., Lee Y., Hong S.T., Bang S., Paik D., Kang J., Shin J., Lee J., Jeon K., Hwang S., *et al.* (2005) *Drosophila* GPCR *Han* is a receptor for the circadian clock neuropeptide PDF. *Neuron* **48**:267-278
26. Jackson F.R., Newby L.M., Kulkarni S.J (1990) *Drosophila* GABAergic systems: sequence and expression of glutamic acid decarboxylase. *J Neurochem* **54**:1068-1078
27. Ki Y., Lim C (2019) Sleep-promoting effects of threonine link amino acid metabolism in *Drosophila* neuron to GABAergic control of sleep drive. *eLife* **8**
28. Konopka R.J., Benzer S (1971) Clock mutants of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* **68**:2112-2116
29. Kurlansik S.L., Ibay A.D (2012) Seasonal affective disorder. *Am Fam Physician* **86**:1037-1041
30. Leal S.M., Neckameyer W.S. (2002) Pharmacological evidence for GABAergic regulation of specific behaviors in *Drosophila melanogaster*. *J Neurobiol* **50**:245-261
31. Lear B.C., Merrill C.E., Lin J.M., Schroeder A., Zhang L., Allada R (2005) A G protein-coupled receptor, groom-of-PDF, is required for PDF neuron action in circadian behavior. *Neuron* **48**:221-227
32. Lin F.J., Song W., Meyer-Bernstein E., Naidoo N., Sehgal A (2001) Photoc signaling by *cryptochrome* in the *Drosophila* circadian system. *Mol Cell Biol* **21**:7287-7294
33. Matsuda K., Hosie A.M., Buckingham S.D., Squire M.D., Baylis H.A., Satella D.B (1996) pH-Dependent actions of THIP and ZAPA on an ionotropic *Drosophila melanogaster* GABA receptor. *Brain Res* **739**:335-338
34. Mertens I., Vandingenen A., Johnson E.C., Shafer O.T., Li W., Trigg J.S., De Loof A., Schoofs L., Taghert P.H. (2005) PDF receptor signaling in *Drosophila* contributes to both circadian and geotactic behaviors. *Neuron* **48**:213-219
35. Mezler M., Müller T., Raming K (2001) Cloning and functional expression of GABA receptors from *Drosophila*. *European Journal of Neuroscience* **13**:477-486
36. Nagoshi E., Sugino K., Kula E., Okazaki E., Tachibana T., Nelson S., Rosbash M (2010) Dissecting differential gene expression within the circadian neuronal circuit of. *Nat Neurosci* **13**:60-U220
37. Nitabach M.N., Wu Y., Sheeba V., Lemon W.C., Strumbos J., Zelensky P.K., White B.H., Holmes T.C (2006) Electrical hyperexcitation of lateral ventral pacemaker neurons desynchronizes downstream circadian oscillators in the fly circadian circuit and induces multiple behavioral periods. *J Neurosci* **26**:479-489
38. Ono D., Honma K.I., Honma S (2021) GABAergic mechanisms in the suprachiasmatic nucleus that influence circadian rhythm. *J Neurochem* **157**:31-41
39. Parisky K.M., Agosto J., Pulver S.R., Shang Y., Kuklin E., Hodge J.J., Kang K., Liu X., Garrity P.A., Rosbash M., *et al.* (2008) PDF cells are a GABA-responsive wake-promoting component of the *Drosophila* sleep circuit. *Neuron* **60**:672-682

40. Park D., Veenstra J.A., Park J.H., Taghert P.H (2008) Mapping peptidergic cells in *Drosophila*: where DIMM fits in. *PLoS One* **3**:e1896
41. Perisse E., Yin Y., Lin A.C., Lin S.W., Huetteroth W., Waddell S (2013) Different Kenyon Cell Populations Drive Learned Approach and Avoidance in. *Neuron* **79**:945-956
42. Pfeifferberger C., Lear B.C., Keegan K.P., Allada R (2010) Processing sleep data created with the *Drosophila* Activity Monitoring (DAM) System. *Cold Spring Harb Protoc* **2010**
43. Renn S.C., Park J.H., Rosbash M., Hall J.C., Taghert P.H (1999) A *pdf* neuropeptide gene mutation and ablation of PDF neurons each cause severe abnormalities of behavioral circadian rhythms in *Drosophila*. *Cell* **99**:791-802
44. Rosenthal N.E., Sack D.A., Gillin J.C., Lewy A.J., Goodwin F.K., Davenport Y., Mueller P.S., Newsome D.A., Wehr T.A (1984) Seasonal affective disorder. A description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry* **41**:72-80
45. Sehgal A., Price J.L., Man B., Young M.W (1994) Loss of circadian behavioral rhythms and *per* RNA oscillations in the *Drosophila* mutant *timeless*. *Science* **263**:1603-1606
46. Sekiguchi M., Inoue K., Yang T., Luo D.G., Yoshii T (2020) A Catalog of GAL4 Drivers for Labeling and Manipulating Circadian Clock Neurons in. *J Biol Rhythm* **35**:207-213
47. Shang Y., Griffith L.C., Rosbash M (2008) From the Cover: Feature Article: Light-arousal and circadian photoreception circuits intersect at the large PDF cells of the *Drosophila* brain. *Proc Natl Acad Sci U S A* **105**:19587-19594
48. Shearin H.K., Quinn C.D., Mackin R.D., Macdonald I.S., Stowers R.S (2018) t-GRASP, a targeted GRASP for assessing neuronal connectivity. *J Neurosci Methods* **306**:94-102
49. Sheeba V., Gu H., Sharma V.K., O'Dowd D.K., Holmes T.C (2008) Circadian- and light-dependent regulation of resting membrane potential and spontaneous action potential firing of *Drosophila* circadian pacemaker neurons. *J Neurophysiol* **99**:976-988
50. Shiga Y., Tanaka-Matakatsu M., Hayashi S (1996) A nuclear GFP/ beta-galactosidase fusion protein as a marker for morphogenesis in living *Drosophila*. *Dev Growth Differ* **38**:99-106
51. Stanewsky R., Kaneko M., Emery P., Beretta B., Wagner-Smith K., Kay S.A., Rosbash M., Hall J.C (1998) The *cry^b* mutation identifies cryptochrome as a circadian photoreceptor in *Drosophila*. *Cell* **95**:681-692
52. Stoleru D., Peng Y., Agosto J., Rosbash M (2004) Coupled oscillators control morning and evening locomotor behaviour of *Drosophila*. *Nature* **431**:862-868
53. Talay M., Richman E.B., Snell N.J., Hartmann G.G., Fisher J.D., Sorkac A., Santoyo J.F., Chou-Freed C., Nair N., Johnson M., et al. (2017) Transsynaptic Mapping of Second-Order Taste Neurons in Flies by trans-Tango. *Neuron* **96**:783-795.
54. Wescott D.L., Soehner A.M., Roeklein K.A (2020) Sleep in seasonal affective disorder. *Curr Opin Psychol* **34**:7-11
55. Wood S., Loudon A (2014) Clocks for all seasons: unwinding the roles and mechanisms of circadian and interval timers in the hypothalamus and pituitary. *J Endocrinol* **222**:R39-59
56. Yoshii T., Hermann-Luibl C., Kistenpennig C., Schmid B., Tomioka K., Helfrich-Förster C (2015) Cryptochrome-Dependent and -Independent Circadian Entrainment Circuits in. *Journal of Neuroscience* **35**:6131-6141
57. Yoshii T., Todo T., Wulbeck C., Stanewsky R., Helfrich-Forster C (2008) Cryptochrome is present in the compound eyes and a subset of *Drosophila*'s clock neurons. *J Comp Neurol* **508**:952-966
58. Zhang H.G., French-Constant R.H., Jackson M.B. (1994) A unique amino acid of the *Drosophila* GABA receptor with influence on drug sensitivity by two mechanisms. *J Physiol* **479**:65-75
59. Zhang Y.Q., Rodesch C.K., Broadie K (2002) Living synaptic vesicle marker: synaptotagmin-GFP. *Genesis* **34**:142-145

Peer reviews

Reviewer #1 (Public review):

Summary:

In this paper, Chen et al. identified a role for the circadian photoreceptor CRYPTOCHROME (CRY) in promoting wakefulness under short photoperiods. This research is potentially important as hypersomnolence is often seen in patients suffering from SAD during winter times. The mechanisms underlying these sleep effects are poorly known.

Strengths:

The authors clearly demonstrated that mutations in *cry* lead to elevated sleep under 4:20 Light-Dark (LD) cycles. Furthermore, using RNAi, they identified GABAergic neurons as a primary site of CRY action to promote wakefulness under short photoperiods. They then provide genetic and pharmacological evidence demonstrating that CRY acts on GABAergic transmission to modulate sleep under such conditions.

Weaknesses:

The authors then went on to identify the neuronal location of this CRY action on sleep. This is where this reviewer is much more circumspect about the data provided. The authors hypothesize that the l-LNvs which are known to be arousal promoting may be involved in the phenotypes they are observing. To investigate this, they undertook several imaging and genetic experiments.

While the authors have made improvements in this resubmitted manuscript, there are still multiple concerns about the paper. I think the authors provide enough evidence suggesting that CRY plays a role in sleep under short photoperiod. The data also supports that CRY acts in GABAergic neurons. However, there are still major issues with the quality of the confocal images presented throughout the paper. In many cases it appears that the images are oversaturated with poor resolution, making it hard to understand what is going on. In addition, none of the drivers used in this study are specific to the neurons the authors aim to manipulate. Therefore, the identity of the GABAergic neurons involved in this CRY dependent sleep mechanism remains unclear. Similarly, whether l-LNvs are the target of this GABA mediated sleep regulation under short photoperiod is not fully demonstrated. The data presented suggests that but does not prove it.

Major concerns:

(1) While the authors provided sleep parameters like consolidation or waking activity for some experiments. These measurements are still not shown for several experiments (for example Figures 2E, 3, 4, 5, and 6). These data are essential, these metrics must be reported for all sleep experiments.

(2) Line 144 "We fed flies with agonists of GABA-A (THIP) and GABA-B receptor (SKF-97541) (Ki and Lim, 2019; Matsuda et al., 1996; Mezler et al., 2001). Both drugs enhance sleep in WT," The proper citation is needed here, Dissel et al., 2015 PMID:25913403. Both THIP and SKF-97541 were used in that paper.

(3) Figure 2C and 2F: it appears that the control data is the same in both panels. That is not acceptable.

(4) Figure 4A: With the quality of the images, it is impossible to assess whether GABA levels are increased at the l-LNvs soma.

(5) Fig 4 S1A shows colabeling of l-LNvs and Gad1-Gal4 expressing neurons. They are almost 100% overlapping signals. This would indicate that the l-LNvs are GABAergic themselves, or that there is a problem with this experiment.

(6) Fig 4 S1B: Again, I can see colabelling of the GFP and PDF staining, suggesting that Gad1-Gal4 expresses in l-LNvs.

(7) Line 184: "Consistently, knocking down Rdl in the l-LNvs rescues the long sleep phenotype of cry mutants (Figure 4-figure supplement 1D)." This statement is incorrect as the driver used for this experiment, 78G01-GAL4 is not specific to the l-LNvs, so it is possible that the phenotypes observed are not coming from these neurons.

(8) Figure 4G-K: None of these manipulations are specific to the l-LNvs. The authors describe 10H10-GAL4 and 78G01-GAL4 as l-LNvs specific tools, but this is not the case. Why not use the SS00681 Split-GAL4 line described in Liang et al., 2017 PMID: 28552314? It is possible that some of the effects reported in this manuscript are not caused by manipulating the l-LNvs.

(9) Similarly for the manipulation of s-LNvs, the authors cannot rule out effect that are coming from other cells as R6-GAL4 is not specific to s-LNvs.

(10) The staining presented in Fig 5 S1 is not very convincing. Difficult to see whether Gad1-GAL4 only expresses in the s-LNvs.

<https://doi.org/10.7554/eLife.92608.2.sa3>

Reviewer #3 (Public review):

Summary:

In humans, short photoperiods are associated with hypersomnolence. The mechanisms underlying these effects is however, unknown. Chen et al. use the fly *Drosophila* to determine the mechanisms regulating sleep under short photoperiods. They find that mutations in the circadian photoreceptor cryptochrome (cry) increase sleep specifically under short photoperiods (e.g. 4h light : 20 h dark). They go on to show that cry is required in GABAergic neurons and that the effects of the cry mutation on sleep are mediated by alterations in GABA signalling. Further, they suggest that the relevant subset of GABAergic neurons are the well-studied small ventral lateral neurons that they suggest inhibit the arousal promoting large ventral neurons via GABA signalling

Strengths:

Genetic analysis to show that cryptochrome (but not other core clock genes) mediates the increase in sleep in short photoperiods, and circuit analysis to localise cry function to GABAergic neurons.

Weaknesses:

The authors' have substantially revised their manuscript, and the manuscript is better for the revisions. However, the conclusion that the sLNvs are GABAergic is unfortunately still not well supported by the data. A key sticking point remains the anti GABA immunostaining, and specific driver lines for sLNvs and lLNvs.

The authors should tone down their conclusions to reflect the fact that their data, as presented, does not support the model that cry acts in sLNvs to modulate GABA signalling onto lLNvs and thus modulate sleep.

<https://doi.org/10.7554/eLife.92608.2.sa2>

Reviewer #4 (Public review):

Summary:

Short photoperiod is an important experimental manipulation in neurobiology, endocrinology, and metabolism studies. However, the molecular mechanisms by which short photoperiod gives rise to behavioral phenotypes that are seen in seasonal affective disorders remain unknown. Using the classic circadian model organism *Drosophila*, this study examines short photoperiod-induced hypersomnolence and identifies the circadian photoreceptor cryptochrome as a regulator of GABAergic tone within the clock neural circuit to promote wakefulness under short photoperiod conditions. The discovery has broad implications for understanding how short photoperiod modulates neural inhibition in circadian circuits in regulating sleep.

Strengths:

The *Drosophila* model provided a powerful platform to dissect the molecular mechanisms underlying short photoperiod-induced hypersomnolence. A battery of behavioral, imaging, circuit-manipulation approaches was employed to test the novel hypothesis that the circadian photoreceptor cryptochrome modulates GABAergic tone within the clock neural circuit to promote wakefulness under short photoperiod conditions.

Weaknesses:

The current model proposed by the authors suggests that the small ventral lateral neurons of the *Drosophila* clock circuit are GABAergic; however, this remains unclear. At present, the field lacks sufficient data and validated reagents to definitively establish the GABAergic identity of these neuropeptidergic neurons.

<https://doi.org/10.7554/eLife.92608.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public Review):

Summary:

In this paper, Chen et al. identified a role for the circadian photoreceptor CRYPTOCHROME (cry) in promoting wakefulness under short photoperiods. This research is potentially important as hypersomnolence is often seen in patients suffering from SAD during winter times. The mechanisms underlying these sleep effects are poorly known.

Strengths:

The authors clearly demonstrated that mutations in cry lead to elevated sleep under 4:20 Light-Dark (LD) cycles. Furthermore, using RNAi, they identified GABAergic neurons as a primary site of cry action to promote wakefulness under short photoperiods. They then provide genetic and pharmacological evidence demonstrating that cry acts on GABAergic transmission to modulate sleep under such conditions.

Weaknesses:

The authors then went on to identify the neuronal location of this cry action on sleep. This is where this reviewer is much more circumspect about the data provided. The

authors hypothesize that the l-LNvs which are known to be arousal-promoting may be involved in the phenotypes they are observing. To investigate this, they undertook several imaging and genetic experiments.

Major concerns:

(1) Figure 2 A-B: The authors show that knocking down cry expression in GABAergic neurons mimics the sleep increase seen in cryb mutants under short photoperiod. However, they do not provide any other sleep parameters such as sleep bout numbers, sleep bout duration, and more importantly waking activity measurements. This is an essential parameter that is needed to rule out paralysis and/or motor defects as the cause of increased "sleep". Any experiments looking at sleep need to include these parameters.

Thank you for bringing up these points. We have now included these sleep parameters in Figure 2—figure supplement 3.

(2) For all Figures displaying immunostaining and imaging data the resolution of the images is quite poor. This makes it difficult to assess whether the authors' conclusions are supported by the data or not.

We apologize for the poor resolution. This is probably due to the compression of the figures in the merged PDF file. We are now uploading the figures individually and hopefully this can resolve the resolution issue.

(3) In Figure 4-S1A it appears that the syt-GFP signal driven by Gad1-GAL4 is colabeling the l-LNvs. This would imply that the l-LNvs are GABAergic. The authors suggest that this experiment suggests that l-LNvs receive input from GABAergic neurons. I am not sure the data presented support this.

We agree that this piece of data alone is not sufficient to demonstrate that the l-LNvs receive GABAergic inputs rather than the l-LNvs are GABAergic. However, when nlsGFP signal is driven by two independent Gad1-GAL4 lines (one generated by P element insertion while the other generated by GAL4 inserted into the Gad1 locus), we do not observe any prominent signal in the l-LNvs (Figure 5A and B; Figure 5-figure supplement 1A). We have also co-labeled using Gad1GAL4 and PdfLexA (Figure 5-figure supplement 1B). As can be seen, Gad1GAL4-driven GFP signal is present only in the s-LNvs but not the l-LNvs. This further supports the idea that the l-LNvs are not GABAergic, and that the syt-GFP signal likely arises from GABAergic neurons projecting to the l-LNvs.

(4) In Figure 4-S1B. The GRASP experiment is not very convincing. The resolution of the image is quite poor. In addition, the authors used Pdf-LexA to express the post t-GRASP construct in l-LNvs, but Pdf-LexA also labels the s-LNvs, so it is possible that the GRASP signal the authors observe is coming from the s-LNvs and not the l-LNvs. The authors could use a l-LNvs specific tool to do this experiment and remove any doubts. Altogether this reviewer is not convinced that the data presented supports the conclusion "All in all, these results demonstrate that GABAergic neurons project to the l-LNvs and form synaptic connections." (Line 176). In addition, the authors could have downregulated the expression of Rdl specifically in l-LNvs to support their conclusions. The data they are providing supports a role for RDL but does not prove that RDL is involved in l-LNvs.

Thank you for these wonderful suggestions. Again we apologize for the poor resolution and hopefully by uploading the images separately we can resolve this issue. We agree that the GRASP signal could be coming from the s-LNvs and not the l-LNvs but unfortunately we are not able to find a LexA that is specifically expressed in the l-LNvs. We believe the trans-Tango data further support the idea that GABAergic neurons project to and form synaptic connections with the l-LNvs. Nonetheless, we have changed our conclusion to "All in all, these


results strongly suggest that GABAergic neurons project to the l-LNvs and form synaptic connections” to be more rigorous. In addition, we have obtained R78G01GAL4 which is specifically expressed in the l-LNvs, and using this GAL4 to knock down Rdl rescues the long-sleep phenotype of cry mutants (Figure 4—figure supplement 1D).

(5) In Figures 4 A and C: it appears that GABA is expressed in the l-LNvs. Is this correct? Can the authors clarify this? Maybe the authors could do an experiment where they co-label using Gad1-GAL4 and Pdf-LexA to clearly demonstrate that l-LNvs are not GABAergic. Also, the choice of colors could be better. It is very difficult to see what GABA is and what is PDF.

Thank you for this wonderful suggestion. We have now co-labeled using Gad1GAL4 and PdfLexA (Figure 5-figure supplement 1B). As can be seen, Gad1GAL4-driven GFP signal is present only in the s-LNvs but not the l-LNvs. We suspect the GABA signal at the l-LNvs may arise from the GABAergic projections received by these cells. We have now changed the color of the GABA/PDF signals in these images and have reduced the intensity of the PDF signal. Hopefully, it would be easier to visualize in this revised version.

(6) Figure 4G: Pdf-GAL4 expresses in both s-LNvs and l-LNvs. So, in this experiment, the authors are silencing both groups, not only the l-LNvs. Why not use a l-LNvs specific tool?

Thank you for bring up this important point. We have previously used c929GAL4 to express Kir2.1 and this led to lethality. We have now used two l-LNv-specific GAL4 drivers (R78G01GAL4 and R10H10GAL4) that we newly obtained to express Kir2.1 but did not observe significant effect on sleep. Please see Author response image 1 for the results.



Author response image 1. Daily sleep duration of male flies expressing Kir2.1 in l-LNvs using R78G01GAL4 (A) (n = 40, 41, 30 flies) and R10H10GAL4 (B) (n = 40, 41, 32 flies) and controls, monitored under 4L20D. One-way ANOVA with Bonferroni multiple comparison test was used to calculate the difference between experimental group and control group.

(7) Figure 4H-I: The C929-GAL4 driver expresses in many peptidergic neurons. This makes the interpretation of these data difficult. The effects could be due to peptidergic cells being different than the l-LNvs. Why not use a more specific l-LNvs specific tool? I am also confused as to why some experiments used Pdf-GAL4 and some others used C929-GAL4

in a view to specifically manipulate l-LNvs? This is confusing since both drivers are not specific to the l-LNvs.

Thank you for bring up these important points. We have now used the l-LNv-specific R10H10GAL4 and the results are more or less comparable with that of c929GAL4 (Figure 4I and K), i.e. activating the l-LNvs blocks the long-sleep phenotype of cry mutants. The reason PdfGAL4 is used in 4G is because c929GAL4 leads to lethality while the l-LNv-specific GAL4 lines do not alter sleep.

(8) Figure 5-S1B: Why does the pdf-GAL80 construct not block the sleep increase seen when reducing expression of cry in Gad1-GAL4 neurons? This suggests that there are GABAergic neurons that are not PDF expressing involved in the cry-mediated effect on sleep under short photoperiods.

Yes, this is indeed the conclusion we draw from this result, and we commented on this in the Discussion: “Moreover, inhibiting cry RNAi expression in PDF neurons does not eliminate the long-sleep phenotype of Gad1GAL4/UAScryRNAi flies. Therefore, we suspect that cry deficiency in other GABAergic neurons is also required for the long-sleep phenotype. Given that the s-LNvs are known to express CRY and appear to be GABAergic based on our findings here, we believe that CRY acts at least in part in the s-LNvs to promote wakefulness under short photoperiod.”

In conclusion, it is not clear that the authors demonstrated that they are looking at a cry-mediated effect on GABA in s-LNvs resulting in a modulation of the activity of the l-LNvs. Better images and more-suited genetic experiments could be used to address this.

Thank you very much for all the comments. They are indeed quite helpful for improving our manuscript. Hopefully, with images of higher quality and the additional experiments described above, we have now provided more evidence supporting our major conclusion.

Reviewer #2 (Public Review):

Summary:

The sleep patterns of animals are adaptable, with shorter sleep durations in the winter and longer sleep durations in the summer. Chen and colleagues conducted a study using Drosophila (fruit flies) and discovered that a circadian photoreceptor called cryptochrome (cry) plays a role in reducing sleep duration during day/night cycles resembling winter conditions. They also found that cry functions in specific GABAergic circadian pacemaker cells known as s-LNvs inhibit these neurons, thereby promoting wakefulness in the animals in the winter. They also identified l-LNvs, known as arousal-promoting cells, as the downstream neurons.

Strengths:

Detailed mapping of the neural circuits cry acts to mediate the shortened sleep in winter-like day/night cycles.


Weaknesses:

The supporting evidence for s-LNvs being GABAergic neurons is not particularly strong. Additionally, there is a lack of direct evidence regarding changes in neural activity for s-LNvs and l-LNvs under varying day/night cycles, as well as in cry mutant flies.

Thank you very much for all the comments. We have now expressed nlsGFP by two independent Gad1-GAL4 lines (one generated by P element insertion while the other generated by GAL4 inserted into the Gad1 locus), and positive signals in the s-LNvs can be

observed (Figure 5A and B; Figure 5-figure supplement 1A). Hopefully, this can provide some further support regarding the s-LNvs being GABAergic neurons.

We have now examined GCaMP signals in the l- and s-LNvs of WT and cry mutants under 4L20D/12L12D. Please see Author response image 2 for the results. As can be seen, both WT and cry mutants show photoperiod-dependent changes. Interestingly, cry mutants show more prominent reduction of GCaMP signal in the l-LNvs compared to WT under 12L12D vs. 4L20D, but the sleep duration phenotype is observed only under 4L20D. Moreover, GCaMP signal is elevated in the s-LNvs of cry mutants relative to WT under 4L20D but decreased under 12L12D. These results indicate that there are distinct mechanisms regulating sleep under short vs. normal photoperiod (with CRY being dispensable under 12L12D), and the role of CRY in modulating the activity of these neurons are also photoperiod-dependent. Further in-depth characterizations are needed to delineate these complex issues.



Author response image 2. Quantification of GCaMP6m signal intensity normalized to that of tdTomato under 12L12D and 4L20D (n = 25-45 cells). Student's t-test: compared to WT, #P < 0.05, ##P < 0.01; 12L12D vs. 4L20D, *P < 0.05, ***P < 0.001.

Reviewer #3 (Public Review):

Summary:

In humans, short photoperiods are associated with hypersomnolence. The mechanisms underlying these effects are, however, unknown. Chen et al. use the fly Drosophila to determine the mechanisms regulating sleep under short photoperiods. They find that mutations in the circadian photoreceptor cryptochrome (cry) increase sleep specifically under short photoperiods (e.g. 4h light: 20 h dark). They go on to show that cry is required in GABAergic neurons. Further, they suggest that the relevant subset of GABAergic neurons are the well-studied small ventral lateral neurons that they suggest inhibit the arousal-promoting large ventral neurons via GABA signalling.

Strengths:

Genetic analysis to show that cryptochrome (but not other core clock genes) mediates the increase in sleep in short photoperiods, and circuit analysis to localise cry function to GABAergic neurons.

Weaknesses:

The authors' conclusion that the sLNvs are GABAergic is not well supported by the data. Better immunostaining experiments and perhaps more specific genetic driver lines would help with this point (details below).

(1) The sLNvs are well known as a key component of the circadian network. The finding that they are GABAergic would if true, be of great interest to the community. However, the data presented in support of this conclusion are not convincing. Much of the confocal images are of insufficient resolution to evaluate the paper's claims. The Anti-GABA immunostaining in Fig 4 and 5 seem to have a high background, and the GRASP experiments in Fig 4 supplement 1 low signal.

We apologize for the poor resolution. This is probably due to the compression of the figures in the merged PDF file. We are now uploading the figures individually and hopefully this can resolve the resolution issue. Unfortunately, the GABA immunostaining does not work very well in our hands and thus the background is high. We have now adjusted the images by changing the minimum lookup table (LUT) value in the green channel to 213, which removes all pixels below 213. This can remove background without changing the gray values, so the analysis is not affected. We have modified all images the exact same way and hopefully this can improve the contrast. Furthermore, we have now expressed nlsGFP by two independent Gad1-GAL4 lines (one generated by P element insertion while the other generated by GAL4 inserted into the Gad1 locus), and positive signals in the s-LNvs can be observed (Figure 5A and B; Figure 5-figure supplement 1A). Hopefully, this can provide some further support regarding the s-LNvs being GABAergic neurons.

Transcriptomic datasets are available for the components of the circadian network (e.g. PMID 33438579, and PMID 19966839). It would be of interest to determine if transcripts for GAD or other GABA synthesis/transport components were detected in sLNvs. Further, there are also more specific driver lines for GAD, and the lLNvs, sLNvs that could be used.

Thank you for these wonderful suggestions. Based on PMID 19966839, both the s-LNvs and l-LNvs express Gad1 and VGAT at a relatively low level, although here in our study Gad1GAL4 expression is observed only in the s-LNvs and not l-LNvs. We have commented on this in the 4th paragraph of Discussion: "One study using cell-type specific gene expression profiling demonstrates Gad1 and VGAT expression in both s-LNvs and l-LNvs, although with relatively low signal (Nagoshi et al., 2010). Here we observed that Gad1GAL4 is expressed in the s-LNvs, and their GABA intensity is reduced when we use R6GAL4 to knock down VGAT in these cells." PMID 33438579 does not report expression of these genes in either s-LNvs or l-LNvs, likely due to insufficient sequencing depth. Furthermore, we have now used two l-LNv-specific GAL4 lines (R78G01GAL4 and R10H10GAL4) to conduct some of the experiments that we previously used c929GAL4 for, and obtained comparable results (Figure 4I and K).

(2) The authors' model posits that in short photoperiods, cry functions to suppress GABA secretion from sLNvs thereby disinhibiting the lLNvs. In Fig 4I they find that activating the lLNvs (and other peptidergic cells) by c929>NaChBac in a cryb background reduces sleep compared to activating lLNvs in a wild-type background. It's not clear how this follows from the model. A similar trend is observable in Fig 4H with TRP-mediated activation of lLNvs, although it is not clear from the figure if the difference b/w cryb vs wild-type background is significant.


Thank you for bring up this important point. This does appear to be counterintuitive. We suspect that in cry mutants, there is more inhibition occurring at the l-LNvs and thus the system may be particularly sensitive to their activation. Therefore, activating these neurons on the mutant background can result in a more prominent wake-promoting effect compared to that of WT.

Recommendations for the authors:

Our major concern centers around the claim that the sLNvs are GABAergic and secrete GABA onto the lLNvs. As it stands, this is not well supported by the data.

The authors could substantiate these findings by using more specific driver lines for GAD / vGAT (MiMic based lines are available that should better recapitulate endogenous expression). Transcriptomic data for circadian neurons are available, the FlyWire consortium also predicts neurotransmitter identities for specific neural circuits. These datasets could be mined for evidence to support the claim of sLNvs being GABAergic

Thank you for these wonderful suggestions. We have now used MiMic-based lines for Gad1 (BS52090, Mi{MIC}Gad1MI09277) and VGAT (BS23022, Mi{ET1}VGATMB01219) to knock down cry but unfortunately were not able to observe changes in sleep. Please see Author response image 3 for the results.



Author response image 3. Daily sleep duration of male flies with cry knocked down in GABAergic neurons by Gad1GAL4 (A) (n = 30, 38, 50, 18, 31 flies) or VGATGAL4 (B) (n = 28, 38, 50, 18, 30 flies) monitored under 4L20D. One-way ANOVA with Bonferroni multiple comparison test: compared to UAS control, ###P < 0.001.

Furthermore, we have now included another Gad1GAL4 line which is generated by knocking GAL4 transgene into the Gad1 locus. We are also able to observe increased sleep when using this GAL4 to knock down cry, and positive signals in the s-LNvs can be observed when using this GAL4 to drive nlsGFP (Figure 2B; Figure 5-figure supplement 1A).

Based on PMID 19966839, both the s-LNvs and l-LNvs express Gad1 and VGAT at a relatively low level, although here in our study Gad1GAL4 expression is observed only in the s-LNvs and not l-LNvs. We have commented on this in the 4th paragraph of Discussion: “One study using cell-type specific gene expression profiling demonstrates Gad1 and VGAT expression in both s-LNvs and l-LNvs, although with relatively low signal (Nagoshi et al., 2010). Here we observed that Gad1GAL4 is expressed in the s-LNvs, and their GABA intensity is reduced when we use R6GAL4 to knock down VGAT in these cells.” The FlyWire does not have prediction for this particular circuit that we are interested in.

Further, many of the immunostaining images have high background / low signal - so better confocal images would help, as would the use of more specific driver lines for the LNvs as it is sometimes hard to distinguish the lLNvs from sLNvs.

We have now adjusted all images by changing the minimum lookup table (LUT) value in the green channel to 213 and that of the red channel to 279, which removes all pixels below 213

and 279, respectively. This can remove background without changing the gray values, so the analysis is not affected. We have modified all images the exact same way and hopefully this can improve the signal to noise ratio. We were not able to find a LexA line that is specifically expressed in the l-LNVs but we have found two l-LNV-specific GAL4 lines (R78G01GAL4 and R10H10GAL4). We used these lines to conduct some of the experiments that we previously used c929GAL4 for, and obtained comparable results (Figure 4I and 4K).

Additional specific comments are in the reviews above.

Minor points:

(1) Line 55: CRYPTOCHROME is misspelled.

This has been fixed.

(2) Line 140: The authors need to provide the appropriate references for the use of THIP and SKF-97541.

This has been added.

(3) Line 149: there are multiple GABA-A receptors in flies, the authors should acknowledge that. What about Lcch3 or Grd?

Thank you for bring up this important point. Here we focused only on Rdl because it is the only GABA-A receptor known to be involved in sleep regulation. We have modified our description regarding this issue: “We tested for genetic interaction between cry and Resistant to dieldrin (Rdl), a gene that encodes GABA-A receptor in flies and has previously been shown to be involved in sleep regulation.”

<https://doi.org/10.7554/eLife.92608.2.sa0>