

Recurrent Mutations Drive the Rapid Evolution of Pesticide Resistance in the Two-spotted Spider Mite *Tetranychus urticae*

Reviewed Preprint

v2 • July 3, 2025

Revised by authors

Reviewed Preprint

v1 • May 7, 2025

Li-Jun Cao, Jin-Cui Chen, Joshua A Thia, Thomas L Schmidt, Richard Ffrench-Constant, Lin-Xi Zhang, Yu Yang, Meng-Chu Yuan, Jia-Yue Zhang, Xiao-Yang Zhang, Qiong Yang, Ya-Jun Gong, Hu Li, Xue-Xin Chen, Ary A Hoffmann , Shu-Jun Wei 

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China • Bio21 Institute, School of BioSciences, University of Melbourne, Parkville, Australia • Biosciences, University of Exeter, Exeter, United Kingdom • Department of Entomology, College of Plant Protection, China Agricultural University, Beijing, China • College of Plant Science and Technology, Beijing University of Agriculture, Beijing, China • Hebei North University, College of Agriculture and Forestry Technology, Zhangjiakou, China • Institute of Insect Science, Zhejiang University, Hangzhou, China

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This study provides **important** insights into the evolution of pesticide resistance, demonstrating that resistance can arise rapidly and repeatedly, which complements prior work on parallel evolution across species. The combination of extensive temporal sampling in the field, experimental evolution, and genomics makes for **compelling** findings. The authors are to be commended for acknowledging the main limitations of their study in the Discussion. Framing the work in a broader context of resistance beyond arthropod pests would further increase the appeal of the study, which is of relevance for both agronomic practitioners and evolutionary biologists.

<https://doi.org/10.7554/eLife.106288.2.sa3>

Abstract

The genetic basis of pesticide resistance has been widely studied, but the exact nature of this evolutionary process in the field is often unclear, particularly when a limited number of populations is considered and when there is a lag between the evolutionary event and its investigation. We showed that an unprecedented number of recurrently evolved mutations in an arthropod pest, the two-spotted spider mite *Tetranychus urticae*, drive the rapid evolution of resistance to a recently commercialized acaricide, cyetpyrafen. We first observed high levels of resistance that appeared and became widespread within three years. Genome scans revealed genetic heterogeneity of resistance among populations and identified 15 target mutations, including six mutations on five amino acid residues of subunit *sdhB*, and nine mutations on three amino acid residues of subunit *sdhD* of the pesticide target succinate dehydrogenase, with as many as five substitutions on one residue. No mutations were

present in 2666 screened historical specimens, suggesting that mutations arose rapidly through *de novo* substitutions or from very rare segregating mutations. Identical mutations recurrently appeared in different genetic backgrounds, increasing the likelihood of resistance evolution. The high number of mutational options available for the evolution of target site resistance in this pest challenges resistance management practices.

1. Introduction

The development of pesticide resistance represents rapid contemporary evolution of organisms in response to human-induced stressors (Ffrench-Constant et al., 2004 [↗](#), McKenzie and Batterham, 1994 [↗](#), Tabashnik et al., 2013 [↗](#), Gould et al., 2018 [↗](#)). Since the first instance of pesticide resistance in a scale insect was documented in 1914, cases have exponentially increased in arthropods, weeds and pathogens, posing threats to agricultural production, human health, and the environment (Sparks et al., 2021 [↗](#), Melander, 1914 [↗](#), Gould et al., 2018 [↗](#), Fritz, 2022 [↗](#), Peterson et al., 2018 [↗](#)). This has led to attempts to stop the evolution of resistance, or at least slow its appearance within populations (Gould et al., 2018 [↗](#)). A key task in resistance management is to understand the evolution of resistance in field populations, where a given mutation may have originated once and spread via gene flow or evolved through multiple independent substitutions, and where resistance-associated mutations can pre-date (standing genetic variation) or arise subsequent to (*de novo* mutation) the introduction of pesticide selection pressures (Ffrench-Constant, 2007 [↗](#), Daborn et al., 2002 [↗](#), Kersten et al., 2023 [↗](#)).

A challenge in understanding the evolution of resistance is that the genetic basis of pesticide resistance observed in laboratory-generated resistant strains often does not reflect that in field populations, especially for arthropod pests (Chen et al., 2023 [↗](#), Walsh et al., 2022 [↗](#), Ffrench-Constant, 2013 [↗](#), Legan et al., 2024 [↗](#)). Field populations are typically much larger and more genetically heterogeneous than laboratory populations. Because of the greater potential availability of rare mutations with large effect sizes in field populations, monogenic resistance to pesticides involving these rare mutations may occur relatively more frequently than in laboratory-generated strains (McKenzie and Batterham, 1994 [↗](#)), though small effect loci can also remain important in field resistance (Schmidt et al., 2024 [↗](#), Lucas et al., 2023 [↗](#)). In contrast, artificial selection in the laboratory usually takes place in populations with much smaller effective population sizes, as well as a constant intensity and pattern of selection expected to favor the accumulation of multiple changes of small effect (Ffrench-Constant, 2013 [↗](#)). Where alleles with large effects are selected in the laboratory, these may not necessarily reflect the full set of resistant mutations in the field (Legan et al., 2024 [↗](#)). Such mismatches between laboratory-selected and field-evolved resistance may lead to an inaccurate understanding of resistance evolution (Ffrench-Constant, 2013 [↗](#), Bras et al., 2022 [↗](#), Legan et al., 2024 [↗](#)).

A further challenge in understanding resistance evolution in the field is that the genetic basis of resistance is often examined retrospectively well after the emergence of resistance in the field. Retrospective studies can fail to capture processes that have occurred in the intervening period (Andreev et al., 1999 [↗](#), Raymond et al., 1991 [↗](#), Ffrench-Constant et al., 1993b [↗](#), Adesanya et al., 2021 [↗](#)), such as the introduction of new alleles through gene flow or *de novo* mutations, and fitness costs as well as changing selection regimes through pesticide rotations that can lead to resistance alleles being transient (Schmidt et al., 2010 [↗](#), Ffrench-Constant, 2013 [↗](#), Kreiner et al., 2022b [↗](#), Kreiner et al., 2019 [↗](#), Groeters and Tabashnik, 2000 [↗](#)). Preserved specimens before and following the introduction of pesticides can allow for resistance processes to be investigated retrospectively, but samples are often unavailable (Hartley et al., 2006 [↗](#), D'Costa et al., 2011 [↗](#)). Ongoing genetic monitoring of populations can help in overcoming some of these issues, particularly if population processes affecting a species are understood and if the genetic basis and new resistant alleles are identified as they evolve in the field (Van Leeuwen et al., 2012 [↗](#), Pélissié et al., 2022 [↗](#), Pezzini et al., 2024 [↗](#), Taylor et al., 2021 [↗](#)).

Here, we take advantage of the recent evolution of resistance in the two-spotted spider mite (TSSM), *Tetranychus urticae* (Acari: Tetranychidae), to a novel acaricide, cyetpyrafen, to provide a detailed understanding of contemporary resistance evolution. TSSM is a global super pest that has evolved resistance to at least 11 groups of acaricides based on their modes of action and is the most resistant invertebrate pest worldwide (De Rouck et al., 2023 [↗](#), Van Leeuwen et al., 2010 [↗](#)). Succinate dehydrogenase inhibitor (SDHi) acaricides (IRAC group 25) were first launched in 2007 for mite control. Four chemicals consisting of cyetpyrafen (first launched in 2017 in China), cyenopyrafen (2009 in Japan), cyflumetofen (2007 in Japan), and pyflubumide (2015 in Japan) fall within this group of acaricides globally (Hayashi et al., 2013 [↗](#), Nakahira, 2011 [↗](#)). Laboratory-selected resistance has already been identified to these new acaricides, and is conferred by target-site mutations of the SDH coding *sdhB* and *sdhD* genes, with mutations identified at three positions (Njiru et al., 2022 [↗](#), Sugimoto et al., 2020 [↗](#)). In China, three acaricidal SDH inhibitors have been registered over the last decade: cyflumetofen (2013), cyetpyrafen (2017) and cyenopyrafen (2019). Since its release in 2017, cyetpyrafen has become the most popular pesticide for controlling spider mites, though low-level resistance was detected in the field only one year after its release (Chen et al., 2019 [↗](#)).

We have now tracked resistance through repeatedly monitoring field-evolved resistance of TSSM to cyetpyrafen at a time when resistance has developed rapidly to high levels across geographically distant populations. We investigated resistance using a combination of population genomics, laboratory selection, bioassays, and studies on historically preserved specimens before the introduction of cyetpyrafen. We find that multiple and recurrently evolved mutations drive the rapid development of high-level resistance to cyenopyrafen in field populations across a widespread geographical range. We also demonstrate that these mutations likely evolved *de novo* after the introduction of cyetpyrafen, though they may also have originated from extremely rare mutations segregating in populations before cyetpyrafen selection. We emphasize the benefits of integrating genomics with other approaches in longitudinal studies to capture the dynamics of rapid evolutionary changes.

2. Results

2.1 Strong resistance evolves rapidly in field and lab-selected populations

To monitor the evolution and spread of cyetpyrafen resistance in the field, we conducted bioassays on 46 mite population samples collected from various regions in China between 2020 and 2024 (Figure 1 [↗](#), Tables S1-S2). The susceptibility of these population samples to cyetpyrafen was compared with previously reported data for populations collected between 2017 and 2018 during the early release stage of cyetpyrafen in China (Chen et al., 2019 [↗](#), Gong et al., 2017 [↗](#)). In 2017-2018, all seven tested populations remained susceptible, with LC_{50} values ranging from 1.4 to 21.6 mg/L (Figure 1a [↗](#)). In 2020, one population from Shouguang City, Shandong Province (SDSG2), showed resistance with an $LC_{50} > 32,000$ mg/L, while the other 13 populations remained susceptible (Figure 1b [↗](#)). By 2021, high levels of resistance to cyetpyrafen quickly became widespread in field populations: eight of 10 tested populations had an LC_{50} higher than 1606.2 mg/L (Figure 1c [↗](#)). In 2024, 11 of the 13 tested populations evolved resistance with an LC_{50} higher than 1498.6 mg/L (Figure 1d [↗](#)). Overall, the susceptibility of all populations was bimodal, with populations being highly susceptible (LC_{50} ranges from 0.61 to 30.1 mg/L, $\bar{x} = 6.72$ mg/L) or highly resistant ($LC_{50} > 1498.6$ mg/L, $\bar{x} > 5892.5$ mg/L) and an 877-fold difference between mean LC_{50} s of the susceptible and resistant groups. While only a single sample was resistant in 2020, by 2021 resistance was observed across the entire range of China (Figure 1c [↗](#)). This pattern indicates that field populations of TSSMs have rapidly developed high levels of resistance to cyetpyrafen across a wide geographical area.




Figure 1

Susceptibility of the two-spotted spider mite *Tetranychus urticae* to cyetpyrafen.

(a-d) Collection maps of populations used for bioassay in different years. Red and green points represent resistant and susceptible populations, respectively. (e) LC_{50} values of each population collected in 2018 (7 populations) (Chen et al., 2019), 2020 (14), 2021 (10), 2024 (13), and a laboratory-selected resistant population (LabR) and a parallel reared susceptible population (LabS) to cyetpyrafen. The green bars show the susceptible populations, while the red bars show the resistant populations.

We conducted artificial selection to further investigate resistance evolution, using two laboratory strains derived from the same field population and reared in parallel. The field population for selection was collected from Hangzhou, Zhejiang province (ZJXS1), with an LC_{50} of 21.6 mg/L. One laboratory strain, 'LabR', was exposed repeatedly to cyetpyrafen, while the other, 'LabS', was not. LabR underwent a gradual increase in cyetpyrafen resistance for the first 15 selection rounds (31 generations), but a much sharper increase over the remaining 18 selection rounds (35 generations) (Table S3). By the end of the experiment, LabR possessed an LC_{50} of 54,336 mg/L, a 2,521-fold increase in resistance relative to the ZJXS1 field population. In contrast, the LC_{50} of the LabS strain decreased to 1.21 mg/L over the same period (**Figure 1e**, Table S3), due to unknown reasons, perhaps linked to adaptation to the laboratory environment. However, this is a very minor change in resistance compared to the large increase seen in the selected strain.

Additional bioassays were used to test whether the selected LabR strain showed cross-resistance to other pesticides that target proteins in the mitochondrial electron transport chain. For the other two mitochondrial complex II inhibitors (IRAC group 25), the LC_{50} of cyenopyrafen increased 21,254-fold (3.76 to 79918 mg/L), and the LC_{50} of cyflumetofen increased 69,861-fold (6.47 to 452,003 mg/L). For the mitochondrial complex III inhibitors, the LC_{50} of bifenazate (group 20D) increased 2-fold (9.87 to 18 mg/L), and the LC_{50} of pyridaben (group 21A) increased 3-fold (3,763.4 to 10,352 mg/L). The results indicate that cyetpyrafen selection led to high cross-resistance to other mitochondrial complex II inhibitors but weak cross-resistance to the mitochondrial complex III inhibitors (Table S3).

2.2 Selective sweep signals are clear but heterogeneous

To identify genes potentially associated with cyetpyrafen resistance, we conducted pooled whole-genome resequencing on 22 TSSM population samples, including historic samples of eight geographical populations collected in 2017 before the first field releases of cyetpyrafen, 12 samples (10 resistant and two susceptible) collected afterwards, and two laboratory populations (LabS and LabR) which originated from the same field population (Tables S1, S4). To allow for variation in the genetic basis of resistance and genetic backgrounds among populations, we chose the Population Branch Excess (PBE) analysis (Yassin et al., 2016) method to scan selection signals across the genome; we used two susceptible populations (NMHS2 and LabS) as non-focal populations to correct for the varied genetic backgrounds, and one resistant population as the focal population to identify population-specific signals of pesticide selection. No clear and coincident selection signal was identified among historical population samples (**Figure 2a**). The laboratory-selected resistant LabR population had clear selection signals at two genes, *sdhB* and *sdhD*, that encode two subunits of succinate dehydrogenase, the target of SDHi acaricides (**Figure 2b**). None of the 10 field-collected resistant populations had selection signals at both of these genes, but four (BJDX6, ZJHZ1, ZJHZ2, and LNDD) had selection signals at *sdhB*, and one (BJCP4) had a selection signal at *sdhD*. Five of the resistant populations (QHHD, GXNN, SDSG3, SDQZ and YNYX) showed no selection signal at either gene; this same pattern was observed in the one population sample (HNCS2) that remained susceptible after cyetpyrafen was introduced. Additional PBE peaks were observed at other genomic regions in historical, resistant, and susceptible populations (**Figure 2a**, 2b); these could reflect demographic processes or other selective forces not considered in this study.

Selection for cyetpyrafen resistance could impact patterns of genetic diversity in the regions of interest or more widely across the genome if selection decreases overall population size. We investigated these impacts focusing on genome-wide patterns of nucleotide diversity (π) and on Tajima's D . Five resistant populations with selection signals identified by PBE analysis (**Figure 2b**) showed corresponding decreases in π and Tajima's D near the two SDH genes (Figure S1). The overall genetic diversity and Tajima's D across the genome were not significantly associated with resistance level, genetic structure or geographic coordinates ($P > 0.05$) (Figure S2, Table S4), suggesting that resistant populations had not experienced strong bottlenecks. Five resistant




Figure 2

Signals of selective sweep across genomes of the two-spotted spider mite *Tetranychus urticae*.

(a) PBE values across the genome for eight historic populations collected in 2017 before the commercial release of cyetpyrafen. No clear and consistent selection signal was identified among these samples. Chromosomes are colored alternately dark grey and light grey. (b) PBE values for one lab-selected resistant population (LabR), 10 resistant field populations and one susceptible population (HNCS) collected in 2021 and 2024 after the release of cyetpyrafen. The laboratory-selected resistant LabR population had clear selection signals at two genes, *sdhB* and *sdhD*. Four (BJDX6, ZJHZ1, ZJHZ2, and LNDD) field-collected resistant populations had selection signals at *sdhB*, and one (BJCP4) had a selection signal at *sdhD*. Chromosomes are colored alternately dark red and light red for resistant populations, dark green and light green for susceptible population HNCS. Allele frequency (pool-seq data) and haplotypes of mutations are indicated near the corresponding *sdhB* and *sdhD* genes of each population in (a) and (b). (c) PBE values for genomic regions around the *sdhB* gene, comparing values between the historic (left) and recently collected (right) populations. (d) As above but for the *sdhD* gene. Each point represents a genomic window of 5-kp wide when compared to reference genomes involving the susceptible laboratory line (LabS) and a susceptible field population (NMHH2). The vertical lines indicate the position of *sdh* genes, while the horizontal lines represent the 1% threshold of PBE values.

populations without selection signals did not show obvious reduced genetic diversity or altered Tajima's *D* values around the SDH genes (Figure S1) suggesting that these had not experienced strong selection or selective sweeps.

2.3 Multiple mutations have arisen in the field

To identify specific mutations conferring resistance, we examined the two SDH genes, *sdhB* and *sdhD*, where selective sweep signals were detected in some populations. We used whole-genome resequencing data from pooled individuals; we also conducted Sanger sequencing of partial *sdhB* (216 bp of 1001 bp) and *sdhD* (155 bp of 1,008 bp) genes to validate the pool-seq data and examined mutation frequencies in additional populations; this involved 1268 individuals from 52 populations, including samples collected before and after the release of cyetpyrafen (Table S1). Allele frequencies identified by pool-seq and Sanger sequencing were correlated (Figure S4). A total of 15 amino acid substitutions were identified, including six mutations on five amino acid residues of *sdhB*: H146Q (CAT to CAA), S212I (AGC to ATC), H258Y (CAT to TAT), I260T (ATC to ACC), I260V (ATC to GTC), and A285S (GCA to TCA), and nine mutations on three amino acid residues of *sdhD*: D116G (GAC to GGC), D116E (GAC/T to GAA), D116N (GAC to AAC), R119C (CGT to TGT), R119L (CGT to CTT), R119G (CGT to GGT), R119P (CGT to CCT), R119H (CGT to CAT), and P120L (CCC to CTC) (Figure 3b, Table S2). The residue R119 on *sdhD* was a notable extreme case, involving five amino acid changes. The second codon position of R119 (G) was substituted by the other three nucleotides, while the first codon position (C) was substituted by two other nucleotides (T and G), indicating that the first two codon positions of R119 are almost saturated in terms of possible nucleotide substitutions of these codon positions.

Multiple mutations were commonly found in a single population, while none of the mutations were fixed in any field population (Figure 2b, Table S2). Populations sampled in 2020 had at most two mutations, while those sampled in 2021 and 2024 had at most five. Populations sampled in 2020, 2021 and 2024 had on average 1.17, 2.46 and 2.55 mutations, respectively. We also examined the other two SDH subunit genes, *sdhA* and *sdhC*, using pool-seq data, but no amino acid replacements were identified.

We checked the evolutionary conservation of these mutations across animals by aligning homologous sequences from ~700 animal species, encompassing nematodes, insects, fishes, and mammals (Figs. 3a and 3b). The protein sequences of *sdhB* and *sdhD* were highly conserved among various animal species, particularly around the mutant residues H146, S212, I260, D116, and R119 observed in this study. In a structural model of human SDH, eight amino acid residues were predicted at the ubiquinone binding site (Q-site), corresponding to 211, 215 and 260 on mite subunit *sdhB*, 58, 63, 67 and 71 on *sdhC*, and 117 on *sdhD* (Njiru et al., 2022, Du et al., 2023). All detected mutations in the mite are located around or in the Q-site, although differences exist between human and mite sequences (Figure 3d).

Although cyetpyrafen only started to be used in 2017, it is possible that these potential resistance mutations were present as standing genetic variation before this time. To investigate this, we obtained Sanger sequencing and pool-seq data from 2317 female mites collected before the cyetpyrafen release and 4631 after its release (Figure 3e, Table S2). Of the 15 putative resistance mutations, only one (A285S) was observed in the pre-release specimens, which was present in population SDRZ (unclear resistance status), but was not detected in specimens collected after the cyetpyrafen release. The historical trajectory of resistance alleles revealed extreme selection after release of cyetpyrafen (Figure 3e). These 14 resistance alleles have collectively experienced a selective strength of $\bar{S} = 0.707$ ($Z = 3.372$, $p = 0.0007$) per year.

However, based on logistic regression, the estimate of the strength of selection on each resistance allele was only significant for I260V ($p = 0.011$). Seven mutations were first detected in 2020 (I260T, D116G, D116N, R119C, R119L, R119P, and R119H), three mutations were first detected in 2021 (H146Q, S212I, and I260V), and the remaining four mutations were first detected in 2024 (H258Y,




Figure 3

Amino acid mutations on *sdhB* and *sdhD* genes in populations of the two-spotted spider mite *Tetranychus urticae*.

(a, b) Sequence logos at mutations related to cyetpyrafen resistance in *sdhB* (a, 757 sequences) and *sdhD* (b, 158 sequences). The numbers under logos indicate the position of amino acid residues on protein chains. Excluding *T. urticae*, conserved amino acids without any alternative are labeled with asterisks. Amino acids of *T. urticae* are shown below. Amino acids in red indicate mutations. (c) 3D structure of the wild-type succinate dehydrogenase of *T. urticae*. Solid 3D shapes show eight amino acid residues at the Q-site, which is the binding site of cyetpyrafen. The *sdhA*, *sdhB*, *sdhC* and *sdhD* chains are shown in yellow, blue, green, and red, respectively. (d) Amino acid mutation sites on a 3D structure of the wild-type succinate dehydrogenase of *T. urticae*. Solid 3D shapes indicate mutations on five residues identified in this study (His258, Ile260, Asp116, Arg119, and Pro120) and one residue (Ser56) reported by previous studies (Njiru et al., 2022, Sugimoto et al., 2020) that confer resistance to SDH inhibitors. (e) The trajectory of allele frequencies of resistance mutations through time fitted by logistic regression. Each dot represents one population. A258S was omitted as it was only present in one population (SDRZ) sampled in 2017.

D116E, R119G, and P120L) (**Figure 3e** [↗](#), Table S2). Among the target-site mutations, I260V and R119C were the most common, found in 65.9% and 46.3% (respectively) of the populations collected after the release of cyetpyrafen, with their frequency increasing rapidly (**Figure 3e** [↗](#)). In the laboratory-selected population LabR, two mutations (I260V and R119C) were fixed, but neither was detected in either the field population from which it was sourced (ZJXS1) or the unselected susceptible laboratory population derived from ZJXS1 (LabS). These results indicate that the putative resistance mutations were not likely to be present as standing genetic variation unless they were segregating at very low frequencies, and they may have evolved recently through *de novo* substitutions although this cannot be demonstrated definitively.

2.4 Recurrent evolution of identical mutations

When a specific resistance mutation is found in multiple populations, this can be the result of migration between populations or to multiple independent origins of the same mutation. We evaluated these evolutionary hypotheses using two methods: TreeMix (Pickrell and Pritchard, 2012 [↗](#)) and haplotype network analysis.

We used TreeMix to investigate whether populations with identical alleles were more genetically similar or had recent histories of migration. TreeMix builds maximum likelihood trees from population allele frequencies and then adds optional migration edges between populations. We set the SCCD population from southwest China as the root population as it had the highest genetic diversity, and we built trees that followed a strict drift model (i.e. no migration) and trees with up to 10 migration edges. The strict drift model explained most of the variance in relatedness among populations (mean over 1000 replicates of 0.930). The best estimate of the number of historical migration events was one. The inclusion of a single gene flow event resulted in only a very small increase in explained genetic variance compared to 0.930 for the drift model, suggesting that migration among populations is low. The consensus tree over the 1000 replicates assigned most of the populations into northern and southern groups by geography, but not by resistance status (**Figure 4a** [↗](#)). We also observed several instances where geographically distant populations grouped together. Some southern populations were placed with the northern group (YNYX), and vice versa (BJDX6); these populations may have been recently established via long-distance dispersal, possibly through the transport of seedlings.

The above-mentioned mutations in *sdhB* and *sdhD* genes were detected in six populations from the southern group and seven populations from the northern group. No significant isolation-by-distance pattern was detected in resistance allele frequencies across survey years (Mantel test: 2020, $P = 0.73$; 2021, $P = 0.52$; 2023, $P = 0.16$), indicating that the spread of resistant alleles was not linked to geographical factors. The mutation I260V was detected in every field population sampled after the release of cyetpyrafen (**Figure 4a** [↗](#), Table S2), based on pool-seq data. The mutation I260T was only detected in the northern group and R119L was mostly detected in the southern group. The mutation H146Q was found in four adjacent populations (ZJHZ1, ZJHZ2, GXNN and BJCP4). This geographical heterogeneity in the distribution of resistance mutations suggests multiple mechanisms of resistance evolution underlying these patterns, involving some spread via migration between populations (e.g. H146Q) as well as possible independent origins of the same mutation (e.g. I260V).

We explored these ideas further using haplotype network analysis. When resistance spreads via migration, these mutations should have identical haplotypes, unless recombination has taken place which will introduce a second haplotype that segregates locally alongside the initial haplotype. We constructed the haplotype network for the partial *sdhB* (216 of 1001 bp) (**Figure 4b** [↗](#)) and *sdhD* (155 of 1,008 bp) (**Figure 4c** [↗](#)) genes determined from Sanger sequencing. Since haplotypes could not be determined when two loci were heterozygous, we detected haplotypes from sequencing data with at most one heterozygous locus. In total 844 and 696 individuals were used to detect haplotypes of *sdhB* and *sdhD*, respectively. The genes showed high genetic diversity, with 8 SNPs (*sdhB*) and 11 SNPs (*sdhD*) detected, which generated 11 (*sdhB*) and 24 (*sdhD*)




Figure 4
Genetic structure and haplotype networks of *sdhB* and *sdhD* genes of the two-spotted spider mite *Tetranychus urticae*.

(a) Treemix result showing genetic structure and gene flow among populations. The mutations detected by pool-seq data were labeled near the corresponding population. (b) Haplotype network of *sdhB* gene inferred from homozygous individuals. In figures (a) and (d), the red population codes show resistant populations, the green ones show susceptible populations, while the black ones show populations of unknown resistance status. (c) Haplotype network of *sdhD* gene inferred from homozygous individuals. The colored haplotypes are those carrying the resistant mutation. (d) Presence of each haplotype carrying the target mutations in two periods. In the early resistance development stage of 2020-2021, each haplotype was mainly distributed across a limited range, while in 2024, the same haplotype was distributed across a wide geographical range. The filled rectangle indicates that the entire haplotype is present in the population, while a colored border suggests that only the mutations were detected, but the haplotype was not assigned. The number of populations carrying the corresponding mutation are listed in the right column for the two time periods.

haplotypes, respectively. We assumed that common haplotypes located at the center of the network are the ancestral haplotypes (H1 and H4 in the case of *sdhB*; H1, H2, H3, and H11 in the case of *sdhD*). We assumed all other haplotypes were derived. Each derived haplotype was only one nucleotide different from its nearest ancestral haplotype. Five of the nine derived *sdhB* haplotypes carried a putative resistance mutation, as did 13 of the 22 derived *sdhD* haplotypes. Combinations of these mutations between the two subunits were present. However, there was no instance of two or more resistance mutations occurring within the same haplotype.

Of the 10 putative resistance mutations detected through Sanger sequencing, all but four (H258Y, D116N, R119G, and P120L) were found as multiple haplotypes (Figs. 4b and 4c). To test whether these haplotypes were produced by multiple origins or by recombination, we inferred the possible means by which recombination could produce each haplotype, and investigated the geographical distributions of the requisite ancestral and derived haplotypes. For mutations in *sdhB*, we inferred that recombination would have to involve the H4 ancestral haplotype as donor, while the H11 ancestral haplotype was required for *sdhD* (Figure S3). Thus, recombination could have produced the *sdhB* haplotypes H5 (H4 x H6) and H8 (H4 x H11), and the *sdhD* haplotypes H17 (H11 x H10), H16 (H11 x H13) and H21 (H11 x H20). However, only three populations (GXNN, YNKM3, and YNYX) had both of the two recombining haplotypes required and one derived haplotype (Table S5). Additionally, the proximity of the mutations was such that multiple recombination events would be required within the same 5 – 60 bp regions, which seems unlikely even with a high recombination rate of (say) 10 cM/Mb where two mutations might only be expected to occur once in 2.5×10^8 effective individuals across 60 bp; this assumes a population recombination rate $\rho = 2cN_e r$, where r refers to recombination rate per base-pair per generation, N_e refers to effective population size, and c refers to ploidy of the genome (Li and Stephens, 2003). These results do not indicate an obvious role for recombination in generating the derived haplotypes.

All haplotypes carrying putative resistance mutations in the *sdhB* gene arose in 2020–2021, along with all haplotypes of the *sdhD* gene apart from H6, H7, H16 and H23 (Figure 4d, Table S5). By 2024, the *sdhB* H6 haplotype became the most prevalent haplotype, while the non-resistant *sdhB* H3 haplotype was almost lost despite it being prevalent between 2013 and 2021 (Figure 4d, Table S5). Some individual populations had multiple haplotypes of the same putative resistance mutation (Figure 4d), and different haplotypes of the same mutation were often found in adjacent populations (Figure 4a, Figure 4d, Table S5). In the ZJXS1 population used to produce the LabR and LabS strains, four haplotypes of the *sdhB* gene (H1, H3, H4 and H7) and six haplotypes of the *sdhD* gene (H1–H4, H11 and H24) were found, none of which had putative resistance mutations. After 66 generations of cyetpyrafen selection, two resistant haplotypes were fixed in the LabR population: H6 of *sdhB* and H10 of *sdhD*, which were likely derived from H1 of *sdhB* and H2 of *sdhD*, respectively (Table S5). These results illustrate the rapid and geographically widespread emergence of resistant alleles in these populations, linked to several mutations found in a range of different haplotypes.

2.5 Single mutations have a large effect on resistance

We next tested resistance of homozygous individuals carrying mutations singly or in combination with other mutations. We first tested the population-level association between allele frequencies of the mutations and levels of resistance, using the five mutations present in at least ten populations (I260T, I260V, D116G, R119C, and R119L) and resistance indicated by survival at 1000 mg/L cyetpyrafen. When each mutation was analyzed separately, a weak association was found between survival and the frequency of the mutant allele (Table S6, linear regression $R^2 = 0.091$ – 0.303). The multivariate regression analysis revealed that frequency of three mutations, I260T ($P = 0.00128$), I260V ($P = 0.00423$) and D116G ($P = 0.00058$), were significantly correlated with resistance of the field populations. When we considered the frequency of individuals having any of the five mutations, we found a strong positive association between survival and the frequency of the predominant allele ($R^2 = 0.707$), the frequency of individuals carrying at least one resistant allele

(heterozygous and homozygous mutant genotypes) ($R^2 = 0.747$), and the frequency of individuals with homozygous mutant genotypes ($R^2 = 0.606$) (**Figure 5a–5c**). These results showed that single mutations explained a low proportion of resistance variation across populations. But when considering all five mutations together, a substantial proportion of resistance variation was explained.

Homozygous lines at the mutation site(s) were created and tested for susceptibility to cyetpyrafen (**Figure 5d**). These lines carried homozygous I260V, D116G, R119E, R119L, and I260V+R119C mutations. When treated with 1000 mg/L cyetpyrafen, mortality of the susceptible line was 100%, but was lower than 25.2% in all other lines, indicating that each mutation and a combination of two mutations conferred high resistance to cyetpyrafen. These results demonstrate that cyetpyrafen resistance is conferred by multiple large-effect mutations in the *sdhB* and *sdhD* genes, although further functional validations are needed.

3. Discussion

Pesticide resistance can evolve rapidly, in contrast to traditional views of adaptation which often emphasize a gradual process (Fisher, 1930). Due to the time taken for resistance to be recognized, evolutionary studies of pesticide resistance are often *post hoc*. Here, we conducted field monitoring of resistance to a novel acaricide, cyetpyrafen, following its 2017 release, with the opportunity to track resistance as it develops. We provided a longitudinal picture of resistance development in the field involving multiple and recurrently evolved large-effect mutations. Cross-resistance was found between cyetpyrafen and two other SDHi acaricides—cyflumetofen and cyenopyrafen—which were released in China before and after cyetpyrafen, respectively. Although we could not clearly identify the contributions of these acaricides to resistance evolution, they have only held a very small market share compared to cyetpyrafen which is expected to be the main contributor to rapid resistance evolution.

Currently available pesticides mainly act by direct binding to receptors and enzymes, resulting in the inhibition of physiological functions associated with the nervous system, energy production system, growth and development, and midgut absorption (Sparks et al., 2021). Mutations that alter the binding kinetics of pesticides on these proteins and thus potentially confer pesticide resistance have been widely identified in agricultural pests and disease vectors (De Rouck et al., 2023; Acford-Palmer et al., 2023). Resistant mutations of some target genes are highly conserved both among populations and among species, perhaps due to the functional constraints of these targets (Ffrench-Constant et al., 1993a; Hawkins et al., 2019).

However, multiple mutations on single pesticide targets have also been identified in the same pest species and among different species. These cases often involve the voltage gated sodium channel (VGSC) gene, where at least 5 and 10 mutations have been cumulatively reported across studies on TSSM and the ectoparasite *Dermanyssus gallinae* respectively (De Rouck et al., 2023). Similarly, the evolution of resistance to acetolactate synthase (ALS)-inhibiting herbicides exhibits high repeatability across weed species, primarily driven by numerous SNPs within the ALS target gene (Gaines et al., 2020). To date, 29 distinct target-site mutations have been identified on eight amino acid residues across 67 weed species (Tranel et al., 2025). Notably, even within a single weed species, up to eight different mutations can occur on three amino acid residues (Tranel et al., 2025).

Here we found 15 mutations on eight amino acid residues in TSSM that all arose within a few years following the introduction of cyetpyrafen. Notably, five of the substitutions involve R119 of the *sdhD* gene. As far as we know, this is the highest number of both mutations and altered residues ever found in a single pesticide target arthropod pests, all emerging in a short period following initial detection of resistance. These mutations were located at the Q-site of SDH targeted




Figure 5

Function of target-site mutations in the *sdhB* and *sdhD* genes and survival of two-spotted spider mite *Tetranychus urticae* in cyetpyrafen resistance.

(a-c) Association between frequency of target-site mutations in the *sdhB* and *sdhD* genes and survival of two-spotted spider mite *T. urticae* exposed to 1000 mg/L of cyetpyrafen. (a) Allele frequencies are measured as the predominant mutation in the population at either the *sdhB* or *sdhD* genes. (b) Allele frequency is the proportion of individuals carrying a resistant allele at either the *sdhB* or *sdhD* genes. (c) Allele frequency is the proportion of individuals carrying a homozygous genotype for either the *sdhB* or *sdhD* genes. Red and dark green points represent resistant and susceptible populations, respectively. Blue dashed lines indicate the expected association between two variables, i.e. $y = x$. (d) Mortality of mite lines carrying one of the homozygous mutations or a combination of two mutations exposed to 1000 mg/L of cyetpyrafen. V, I260V; G, D116G; E, R119E; L, R119L; V and C, I260V and R119C; S, susceptible populations LabS. Populations with the same homozygous mutations are shown in the same color. Error bars show standard error of the mean.

by acaricidal inhibitors of SDH. Among these mutations, I260T was previously found in a cyflumetofen-resistant population (Sugimoto et al., 2020 [↗](#)), while I260V (with S56L in subunit C) was found in a pyflubumide and cyenopyrafen-resistant population (Maeoka and Osakabe, 2021 [↗](#)). R119L has previously been reported in cyflumetofen resistant and susceptible strains of TSSM but had not yet been validated as having a role in resistance (Sugimoto et al., 2020 [↗](#)). These results provide information on the nature of novel mutations in a pesticide target and the high level of resistance to cyetpyrafen that they can confer. It would be interesting to further validate the impacts of the mutations across chemicals to establish patterns of cross resistance given that the same mutation may confer a different level of resistance to different chemicals (Maeoka and Osakabe, 2021 [↗](#)).

Our temporal collections of TSSM preceding cyetpyrafen release in the field indicated that the resistant mutations were not at an appreciable frequency prior to pesticide selection. Sequencing of 13 populations (involving a total of 2666 TSSMs) from 2013 to 2018 did not recover any of the identified mutations. This points to a possible role of recurrent *de novo* mutation, although resistant alleles could have been present at very low frequencies before pesticide selection which can be difficult to detect (Hawkins et al., 2019 [↗](#)). In laboratory populations of TSSM, the H258Y mutation arose *de novo* and was subsequently favored by continuous selection, resulting in resistance to pyflubumide and cyenopyrafen (Njiru et al., 2023 [↗](#)). Our selection experiments also led to new mutations (I260V, R119C) arising over 66 generations (2 years) of cyetpyrafen selection in the LabR population. Although it is hard to completely exclude the possibility of contamination during laboratory selection, because TSSM is tiny and can easily be dispersed passively, the haplotype analysis showed that the two mutations in the lab-selected population likely arose *de novo*. In the same way, multiple *de novo* mutations could have contributed to the rapid evolution of resistance in different populations and regions as documented in our field surveys.

Although few studies have traced resistance evolution in the field, they suggest that both *de novo* mutations and standing genetic variation can contribute to resistance, with their relative importance depending on the specific pesticides and pests involved (Hawkins et al., 2019 [↗](#)). Resistant alleles can originate from a single event and spread to distant populations by migration (Raymond et al., 1991 [↗](#), Daborn et al., 2002 [↗](#)), but resistant mutations can also have multiple origins (Guan et al., 2021 [↗](#), Ilias et al., 2014 [↗](#), Troczka et al., 2012 [↗](#), Kreiner et al., 2022b [↗](#), Schmidt et al., 2024 [↗](#)). Direct evidence for multiple origins comes from the presence of the same mutation on multiple haplotypes, assuming that recombination can be excluded as an explanation because insufficient time has elapsed after a resistant mutation is introduced into a new background (Semenov et al., 2019 [↗](#)). In our study, the requisite recombinant haplotypes were rarely found in the same population, and we therefore suspect that recombination is unlikely to be the source of patterns of haplotype variation over short DNA sequences. We suspect that populations of mites like TSSM have a strong evolutionary potential to evolve *in situ* rather than relying on gene flow, and that there is a repeatability of evolutionary changes when an organism is exposed to the same selection pressure, a long-standing issue in evolutionary biology (Konecna et al., 2021 [↗](#)). The recurrent appearance of the same mutation in different genetic backgrounds suggests that the likelihood of resistance evolution is high in TSSM and not dependent on gene flow, and the likelihood of this scenario is further increased by the fact that resistance can develop through several mutational options. We observed 2.7 resistant mutations per population in samples collected in 2024, seven years after the release of cyetpyrafen. The estimated mutation rate (θ) is 0.0193, assuming 20 generations per year for TSSM. An effective population size (N) of 2.29×10^6 would be necessary to reach the number of *de novo* mutation observed in this study, given $\theta = 4N\mu$ (haplodiploid sex determination of TSSM) and a mutation rate of $\mu = 2.8 \times 10^{-9}$ per base pair per generation as estimated for *Drosophila melanogaster* (Keightley et al., 2014 [↗](#)). The high reproductive capacity (> 100 eggs per female) and short generation time of TSSM make it possible to reach such a large population size in the field.

Because field populations of TSSM can use multiple mutations to reach the same phenotypic endpoint (resistance), it is unlikely that all resistant mutations will be identified from laboratory selection studies particularly if these involve a single laboratory population. Even in the case where multiple populations of TSSM have been selected for resistance (to cyenopyrafen and the related chemicals cyflumetofen and pyflubumide), QTL mapping still only identified a few of the relevant mutations providing resistance to each chemical (Sugimoto et al., 2020 [↗](#)). Additionally, mutations identified in laboratory selection may not be involved in resistance in the field because of fitness costs. For instance, H258Y conferred resistance to cyenopyrafen and pyflubumide in laboratory-selected populations is associated with a high fitness cost in TSSM (Njiru et al., 2023 [↗](#)), and also likely reinforces cyflumetofen binding and toxicity (Njiru et al., 2022 [↗](#)). Relying on a limited set of resistance alleles identified in the laboratory for monitoring may bias inferences of resistance evolution in the field where many alleles can produce resistance. Genetic and evolutionary studies on field populations are then essential to develop a full picture of resistance evolution (Konecna et al., 2021 [↗](#), Lucas et al., 2023 [↗](#), Schmidt et al., 2024 [↗](#), Kersten et al., 2023 [↗](#), Pezzini et al., 2024 [↗](#)).

Theoretical and empirical studies have showed that field-evolved resistance often involves large-effect alleles with a simple genetic basis, while laboratory selection may result in polygenic genetic architectures, due to differences in the occurrence of rare alleles (more likely in large field populations) and different selection pressures (more variable with intense periods in the field on occasions) (Ffrench-Constant, 2013 [↗](#), Roush and McKenzie, 1987 [↗](#), Ffrench-Constant et al., 1990 [↗](#), McKenzie and Batterham, 1994 [↗](#), Crow, 1954 [↗](#)). This could potentially be mitigated by establishing laboratory selection lines using large starting populations from the field with pre-existing resistance alleles (Gould et al. 1997). Until recently, few studies have directly compared the genetic basis of resistance in laboratory populations with field-evolved resistance across a large geographical scale (McKenzie et al., 1992 [↗](#), Ffrench-Constant et al., 1990 [↗](#), Crow, 1954 [↗](#), Legan et al., 2024 [↗](#)). Here we identified large-effect alleles in both field- and laboratory-selected resistant populations. High-level resistance was generated after lab selection with cyetpyrafen for 33 of 65 generations, almost at the same rate as the field populations where high-level resistance was observed three years after the commercial release of cyetpyrafen. In field populations, it seems that multiple allelic options can be favored by selection due to parallel adaptation, and that the same allele may also end up residing in different haplotypes in different areas. Through migration these alleles/backgrounds may eventually end up in the same population to produce a polyallelic basis of adaptation where particular alleles may then not reach high frequencies unless one haplotype is favored (Ralph and Coop, 2010 [↗](#)).

Pesticide resistance provides many fascinating examples of rapid and recent evolution under high selection pressure, which can be leveraged to address fundamental issues about the evolutionary basis of adaptations to new environments (Hawkins et al., 2019 [↗](#)), and also to examine the reliability of commonly used research methods. The development of high-throughput sequencing and population genomics methodologies provides an opportunity to detect selective signatures across the genome (Van Etten et al., 2020 [↗](#), Cohen et al., 2022 [↗](#), Pélissié et al., 2022 [↗](#)). Although pesticide resistance is a rapid evolutionary process under strong selective pressures, identifying candidate genes directly from genomic signals remains difficult when adaptation occurs via soft sweeps (Pennings and Hermisson, 2006 [↗](#), Schrider et al., 2015 [↗](#)). Soft sweeps may be common in adaptive evolution if the effective mutation rate ($4N_e\mu$) $\gg$ 0.01, e.g., through large population sizes (N_e), and/or high allelic mutation rates (Hermisson and Pennings, 2005 [↗](#), Pennings and Hermisson, 2006 [↗](#)). The multiple haplotypes and distinct evolutionary history of SDH genes among resistant field populations suggest a prevalence of soft selective sweeps through recurrent mutation as implicated in pesticide resistance in other pest systems, such as *Drosophila* (Garud and Petrov, 2016 [↗](#)), and the oomycete *Plasmopara viticola* (Delmas et al., 2017 [↗](#)). This is congruent with the 'messier' signals found in field populations relative to the artificially-selected resistant laboratory population. Sweeps around the SDH genes are incomplete, or other contributing loci are undergoing selective sweeps, which may further reduce signal to detect

adaptation (Pennings and Hermisson, 2006 [↗](#), Schrider et al., 2015 [↗](#)). Fewer mutation types and haplotypes along with higher mutation frequency are more likely to be detected. Clearer signals of selective sweeps could emerge from haplotype-based linkage-disequilibrium methods using individual-level genomic data (Hermisson and Pennings, 2017 [↗](#)).

In conclusion, we have captured a detailed picture of contemporary evolutionary processes under pesticide selection in field populations of an invertebrate pest. We demonstrate that the rapid evolution of pesticide resistance is driven by an unprecedented number of population-specific mutations in target genes that arose recurrently. These findings reveal complex origins and diverse genetic bases of pesticide resistance in the field. These findings represent a challenge for SNP-specific molecular monitoring of resistance and for the implementation of effective resistance management strategies. They also highlight the value of amplicon and whole-genome sequencing to detect multiple target-site mutations. Finally, we call for future studies on field-evolved resistance to provide theoretical and practical insights into effective pest control, leveraging the genomics era.

4. Materials and Methods

4.1 Samples

We used 72 field-collected and laboratory-selected populations of TSSM across China for bioassays, whole-genome resequencing, Sanger sequencing of target genes, and establishment of mutant strains (Table S1), including 15 populations collected before (2013 - 2017) and 55 populations collected after (2018 - 2024) the release of cyetpyrafen, as well as one laboratory-selected resistant population, and one laboratory-maintained susceptible population. More than 500 adults were collected from at least three points separated by 100 meters for each population. Samples were kept in 100% alcohol or transported live to the laboratory for chemical bioassays. Because TSSM has a haplodiploid sex-determination system, only female adults were randomly selected for bioassays and genotyping. A initial colony of about 4,000 individuals was established from the field population ZJXS1 in 2018, which was subsequently divided into two parallel cohorts under controlled laboratory conditions. The selection group (LabR) was subjected to continuous selection pressure using cyetpyrafen, while the control group (LabS) was maintained under identical laboratory conditions without exposure to acaricidal agents. The TSSM strain was reared on French bean, *Pemphigus vulgaris* L., under 25±1 °C, 60±5% relative humidity and a photoperiod of L16h: D8h. We continuously selected this population in the laboratory by killing about 60% to 80% of randomly selected individuals every two generations, to obtain a laboratory-selected resistant strain. When the mortality was lower than 60%, another round of spraying was applied by increasing the dosage of cyetpyrafen. The LC₅₀ values were tested at F₁, F₃₂, F₅₄, F₆₀, F₆₂, and F₆₆ generations to quantify resistance development trajectories.

4.2 Bioassay

Bioassays were conducted to test the susceptibility of TSSM to acaricides using a Potter Spray Tower (Burkard Scientific, London, UK). We used transparent plastic cups (6 cm in diameter and 4 cm in height) to keep the female adult mites. A layer of 0.2% agar was placed on the bottom of the cups to avoid leaves drying out. Fresh leaves of *P. vulgaris* were cut into a circle with a diameter of 6 cm to fit the cup and put onto the agar. The borders of the leaves were sealed using 0.2% agar. Thirty female adults were carefully moved into a plastic cup by suction trap.

Two ml acaricide solution was applied through a Potter spray tower (Burkard Manufacturing Co. Ltd., UK) at 68.9 kPa and a sedimentation time of 30 s. Each population was exposed to between five and eight pre-estimated concentrations of acaricides, diluted with water containing 0.1% Triton X-100 (Beijing Solar BioScience and Technology Limited Company, China). The 0.1% Triton X-100 solution was used as a control treatment to correct the potential influence of adjuvants in

the acaricides. The treated mites were kept under 25 °C, 60-70% relative humidity, and a photoperiod of 16 h:8 h (L:D). Mortality was assessed under a Stemi 305 stereomicroscope (Zeiss, Germany), 48 hours after spraying. The tested TSSMs were considered dead if no movement of appendages was observed when they were prodded with a fine brush at 24 hours and 48 hours after spraying. The 50% lethal concentration (LC_{50}) and its 95% confidence intervals were calculated using probit regressions implemented in the statistical software DPS v12.01 (Tang and Zhang, 2013 [↗](#)).

4.3 Pooled whole-genome resequencing and SNP calling

Between 50 and 400 individuals per population were used for pooled genome resequencing to obtain population genomic data for diversity estimates and demographic analyses. DNA libraries with insert sizes of 400 bp were constructed and sequenced on the Illumina NovaSeq 6000 platform to obtain 150 bp paired-end reads. Raw reads were filtered using fastp v0.23.0 (Chen et al., 2018 [↗](#)). The filtered reads were mapped to our recently assembled chromosome-level genome (Cao et al., 2024 [↗](#)) using bwa-mem2 2.2.1 with default parameters (Vasimuddin et al., 2019 [↗](#)). SAMtools v1.14 (Li et al., 2009 [↗](#)) was used to sort the sequence data, removing duplicated reads, removing reads with mapping quality < 20, and converting bam files to mpileup files. Indels in mpileup files were filtered out (with a 5-bp margin on each side) to avoid spurious SNP calling using scripts from PoPoolation (Kofler et al., 2011a [↗](#)). Then mpileup files were converted using scripts from PoPoolation2 (Kofler et al., 2011b [↗](#)).

4.4 Population genetic diversity and demography analyses

Genetic diversity in each population was quantified by estimating nucleotide diversity (π) and Tajima's D across 5 kb non-overlapping windows, using PoPoolation (Kofler et al., 2011a [↗](#)). To reduce the impact of read depth on estimates of nucleotide diversity, each alignment was converted into a population-specific pileup file using SAMtools, and reads were subsampled without replacement up to a depth of 50 (Hoban et al., 2016 [↗](#)). The mean F_{ST} across all windows for each pairwise population comparison was calculated using PoPoolation2 (Kofler et al., 2011a [↗](#)) across 5 kb non-overlapping windows. To infer population relationships and historical migration, a rooted maximum likelihood tree was constructed with a strict drift model (no migration) using Treemix v1.13 (Pickrell and Pritchard, 2012 [↗](#)). The SCCD population was set as root and we tested migration events from 0 to 10 with at least 100 runs for each case. The best number of migration events was determined by the second-order rate of change in likelihood (Δm) across incremental migration events (m) using the Evanno method (Evanno et al., 2005 [↗](#)) embedded in R package 'optM' (Fitak, 2021 [↗](#)). For the best m , we performed 1000 runs to generate 1000 maximum likelihood trees and obtained a consensus tree using the program SumTrees version 4.10 (Sukumaran and Holder, 2010 [↗](#)).

4.5 Genome scan of outlier SNPs

We investigated the genomic signals of selection around the four SDH genes (± 1 Mb) and across the whole genome. The main statistic used to identify outlier regions was the population branch excess (PBE) statistic (Yassin et al., 2016 [↗](#)). The PBE statistic is based on comparisons of F_{ST} values among three populations, comprising a focal population and two nonfocal populations (Pool et al., 2017 [↗](#)). In the analysis, each resistant population was set as the focal population, and two susceptible populations (LabS and NMHH2) were set as the nonfocal populations. PBE values were calculated for 5 kb non-overlapping sliding windows. Windows with top 5% PBE value were considered outliers. Patterns of nucleotide diversity (π) estimates and Tajima's D values around the SDH genes were also examined. We expected resistant populations to have lower nucleotide diversity and Tajima's D values near the SDH genes due to directional selection favoring resistant haplotypes. The coding effects of SNPs within the SDH genes were obtained using SnpEff v4.3 (Cingolani et al., 2012 [↗](#)).

4.6 SDH homology modeling

The three-dimensional (3D) structures of the wild and mutational SDH from TSSM were modeled with SWISS-MODEL (Waterhouse et al., 2018) using 8GS8.1 (Du et al., 2023) as a template. To examine the conservation of *SdhB* and *SdhD* genes, we collected the protein sequences of two genes of different animals in the protein database of NCBI (<https://www.ncbi.nlm.nih.gov/protein>). Then, MUSCLE in MEGA v11 (Tamura et al., 2021) were used to perform sequence alignment and trim protein sequences around mutation points. After removing identical sequences of the same species, the multiple sequence alignment sequences were used to determine conservation and visualized with WebLogo 3 (Crooks et al., 2004).

4.7 PCR amplification and sequencing of target genes

The target genes sequences of 16 - 48 individuals from each population were used to characterize the spatiotemporal variation of these target-site mutations in these populations before and after the release of cyetpyrafen (Table S1). PCR primers were designed for sequences of the *sdhB* gene (*sdhb_F*: GAGTCTAGCCGAGTCTGAT, *sdhb_R*: ACTTGGGACCTGCTGTACTC) and the *sdhD* gene (*sdhd_F*: GCGATGGATAAGGCATAGACG, *sdhd_R*: TCTGGTCAGTGTGGTTCCTC) to capture the candidate mutation sites. Polymerase chain reaction (PCR) was conducted in the Mastercycler pro system (Eppendorf, Germany) under an annealing temperature of 54 °C for 1 min using Promega Go Taq™ Master Mixes (Madison, Wisconsin, USA). Amplified products were purified and sequenced directly from one strand using an ABI 3730xl DNA Analyzer by Sangon Biotech Co. Ltd. (Shanghai, China).

4.8 Haplotype analysis

Haplotypes of each gene were identified from Sanger sequencing. We constructed haplotype networks for *sdhB* and *sdhD* from individuals that contained homozygous resistant genotypes. Haplotype networks were constructed using the R package *pegas* (Paradis, 2010), and were visualized with the R package *ggplot2*.

4.9 Association analysis between mutation frequency and resistance level

Regressions were run to assess the strength of association between target-site mutation frequencies and mean population-level resistance phenotypes. Since LC_{50} of some populations could not be calculated or were inaccurate due to high resistance levels, the survival rate after spraying with 1000 mg/L cyetpyrafen, which is about 6.7 to 10-fold of field-recommended dose, was used to quantify the resistance level of TSSM to cyetpyrafen. Because multiple target-site mutations existed, we evaluated the associations using different metrics of target-site mutation frequencies: (i) the frequency of the predominant mutation in each population, (ii) the frequency of individuals carrying any mutation in each population, and (iii) the frequency of individuals with homozygous mutations in each population. Regressions were fitted as linear models using the *lm* function from the R package *stats* and visualized with *ggplot2*. We did not adjust associations for relatedness among populations because the evolutionary events leading to resistance appear to be largely independent. A logistic regression model was used to track trajectories of allele frequencies. The selection coefficient of each allele (s) and their joint effect ($\bar{S}$) were estimated (Kreiner et al., 2022a). We calculated F_{ST} using the frequency of resistance alleles following Hartl et al. (1997) and explored isolation by distance (IBD) with Mantel tests in the R package *vegan*.

4.10 Establishment of homozygous lines

Single immature TSSM females were picked from resistant populations and reared individually in a petri dish. When unmated, females will only produce haploid male eggs. To obtain the genotype of males for the *sdh* genes, the DNA of each female was extracted and genotyped as mentioned

above after it had laid more than 10 eggs, while the eggs (> 10) were allowed to develop into male adults (F_1 male adults). To obtain the genotype of virgin females, the first six eggs were genotyped while the females were kept live (F_0 females). Then, each pair of F_0 female and F_1 male adults with the same homozygous genotypes for the six mutation sites was placed into the same petri dish and left to mate and produce male and female progeny, with females from the line generated defined as a line homozygous for the target resistance mutation(s). The mated male and female pairs were from the same population to reduce differences in their genetic backgrounds. The homozygous lines at the mutation sites were tested for susceptibility to cyetpyrafen with the bioassay method described above.

Acknowledgements

We thank Ming-Liang Li, Hong-Ping Tian, Liang-Bin Zeng, for their assistance with sample collections. This study is supported by National Key R&D Program of China (2023YFD1401200) and Program of Beijing Academy of Agriculture and Forestry Sciences (QNJJ202006, JKZX202208). Ary Hoffmann and Joshua Thia are supported by the Grains Research and Development Corporation and Horticulture Innovation (Australia).

Additional information

Author contributions

S.J.W. designed research; J.C.C., L.J.C., L.X.Z., Y.Y., M.C.Y., J.Y.Z., X.Y.Z., and Y.J.G., performed research; L.J.C., S.J.W., J.A.T., T.L.S., Q.Y., and A.A.H. analyzed data; S.J.W., A.A.H., L.J.C., J.A.T., T.L.S., H.L., R.F.C., and X.X.C. wrote the paper.

Data availability statement

Raw reads of whole genome resequencing data were submitted to NCBI Sequence Read Archive, with accession numbers SRR30666541 - SRR30666561 and SRR28000465. All scripts and data used in the study were deposited to GitHub: https://github.com/caolijun/TU_pool-seq and Figshare: 10.6084/m9.figshare.27231003 (Wei and Cao, 2025).

Funding

National Key R&D Program of China (2023YFD1401200)

Program of Beijing Academy of Agriculture and Forestry Sciences (QNJJ202006)

Program of Beijing Academy of Agriculture and Forestry Sciences (JKZX202208)

Grains Research and Development Corporation and Horticulture Innovation (Australia)

Additional files

Supplemental materials [↗](#)

References

- Acford-Palmer H., Campos M., Bandibabone J., N'do S., Bantuzeko C., Zawadi B., Walker T., Phelan J.E., Messenger L.A., Clark T.G., Campino S (2023) **Detection of insecticide resistance markers in *Anopheles funestus* from the Democratic Republic of the Congo using a targeted amplicon sequencing panel** *Scientific Reports* **13**:17363 [Google Scholar](#)
- Adesanya A. W., Lavine M. D., Moural T. W., Lavine L. C., Zhu F., Walsh D. B (2021) **Mechanisms and management of acaricide resistance for *Tetranychus urticae* in agroecosystems** *Journal of Pest Science* **94**:639–663 [Google Scholar](#)
- Andreev D., Kreitman M., Phillips T. W., Beeman R. W (1999) **Multiple origins of cyclodiene insecticide resistance in *Tribolium castaneum* (Coleoptera: Tenebrionidae)** *Journal of Molecular Evolution* **48**:615–624 [Google Scholar](#)
- Bras A., Roy A., Heckel D. G., Anderson P., Karlsson Green K. (2022) **Pesticide resistance in arthropods: Ecology matters too** *Ecology Letters* **25**:1746–1759 [Google Scholar](#)
- Cao L. J., Guan T. B., Chen J. C., Yang F. Y., Liu J. X., Jin F. L., Wei S. J (2024) **Chromosome-level genome assembly of the two-spotted spider mite *Tetranychus urticae*** *Scientific Data* **11**:798 [Google Scholar](#)
- Chen J. C., Gong Y. J., Shi P., Wang Z. H., Cao L.-J., Wang P., Wei S. J (2019) **Field-evolved resistance and cross-resistance of the two-spotted spider mite, *Tetranychus urticae*, to bifenthrin, cyenopyrafen and SYP-9625** *Experimental and Applied Acarology* **77**:545–554 [Google Scholar](#)
- Chen S., Zhou Y., Chen Y., Gu J (2018) . **fastp: an ultra-fast all-in-one FASTQ preprocessor** *Bioinformatics* **34**:i884–i890 [Google Scholar](#)
- Chen Y. H., Cohen Z. P., Bueno E. M., Christensen B. M., Schoville S. D. (2023) **Rapid evolution of insecticide resistance in the Colorado potato beetle, *Leptinotarsa decemlineata*** *Current Opinion in Insect Science* **55**:101000 [Google Scholar](#)
- Cingolani P., Platts A., Wang L.L., Coon M., Nguyen T., Wang L., Land S.J., Lu X., Ruden D.M (2012) **A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3** *Fly (Austin)* **6**:80–92 [Google Scholar](#)
- Cohen Z. P., Chen Y. H., Groves R., Schoville S. D (2022) **Evidence of hard-selective sweeps suggests independent adaptation to insecticides in Colorado potato beetle (Coleoptera: Chrysomelidae) populations** *Evolutionary Applications* **15**:1691–1705 [Google Scholar](#)
- Crooks G. E., Hon G., Chandonia J.-M., Brenner S. E (2004) **WebLogo: A sequence logo generator** *Genome Research* **14**:1188–1190 [Google Scholar](#)
- Crow J. F (1954) **Analysis of a DDT-resistant strain of *Drosophila*** *Journal of Economic Entomology* **47**:393–398 [Google Scholar](#)

- D'costa V. M., King C. E., Kalan L., Morar M., Sung W. W., Schwarz C., Froese D., Calmels F., Debruyne R. J. N (2011) **Antibiotic resistance is ancient** *Nature* **477**:457–461 [Google Scholar](#)
- Daborn P. J., Yen J. L., Bogwitz M. R., Le Goff G., Feil E, Jeffers S, Tijet N, Perry T, Heckel D, Batterham P, Feyereisen R, Wilson T G, Ffrench-Constant R H (2002) **A single p450 allele associated with insecticide resistance in Drosophila** *Science* **297**:2253–2256 [Google Scholar](#)
- De Rouck S., Inak E., Dermauw W., Van Leeuwen T. (2023) **A review of the molecular mechanisms of acaricide resistance in mites and ticks** *Insect Biochemistry and Molecular Biology* **159**:103981 [Google Scholar](#)
- Delmas C. E., Dussert Y., Delière L., Couture C., Mazet I. D., Richart Cervera S., Delmotte F. (2017) **Soft selective sweeps in fungicide resistance evolution: recurrent mutations without fitness costs in grapevine downy mildew** *Molecular Ecology* **26**:1936–1951 [Google Scholar](#)
- Du Z., Zhou X., Lai Y., Xu J., Zhang Y., Zhou S., Feng Z., Yu L., Tang Y., Wang W., Yu L., Tian C., Ran T., Chen H., Guddat L. W., Liu F., Gao Y., Rao Z., Gong H (2023) **Structure of the human respiratory complex II** *Proceedings of the National Academy of Sciences of the United States of America* **120**:e2216713120 [Google Scholar](#)
- Evanno G., Regnaut S., Goudet J (2005) **Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study** *Molecular Ecology* **14**:2611–20 [Google Scholar](#)
- Ffrench-Constant R., Roush R. T., Mortlock D., Dively G. (1990) **Isolation of dieldrin resistance from field populations of Drosophila melanogaster (Diptera: Drosophilidae)** *Journal of Economic Entomology* **83**:1733–1737 [Google Scholar](#)
- Ffrench-Constant R., Steichen J., Rocheleau T., Aronstein K., Roush R. (1993a) **A single-amino acid substitution in a gamma-aminobutyric acid subtype A receptor locus is associated with cyclodiene insecticide resistance in Drosophila populations** *Proceedings of the National Academy of Sciences of the United States of America* **90**:1957–1961 [Google Scholar](#)
- Ffrench-Constant R. H. (2007) **Which came first: insecticides or resistance?** *Trends in Genetics* **23**:1–4 [Google Scholar](#)
- Ffrench-Constant R. H. (2013) **The molecular genetics of insecticide resistance** *Genetics* **194**:807–815 [Google Scholar](#)
- Ffrench-Constant R. H., Daborn P. J., Le Goff G (2004) **The genetics and genomics of insecticide resistance** *Trends in Genetics* **20**:163–170 [Google Scholar](#)
- Ffrench-Constant R. H., Rocheleau T. A., Steichen J. C., Chalmers A. E. (1993b) **A point mutation in a Drosophila GABA receptor confers insecticide resistance** *Nature* **363**:449–451 [Google Scholar](#)
- Fisher R. A (1930) **The genetical theory of natural selection** Oxford University Press [Google Scholar](#)
- Fitak R. R (2021) **OptM: estimating the optimal number of migration edges on population trees using Treemix** *Biology Methods and Protocols* **6**:bpab017 [Google Scholar](#)

- Fritz M. L. (2022) **Utility and challenges of using whole-genome resequencing to detect emerging insect and mite resistance in agroecosystems** *Evolutionary Applications* **15**:1505–1520 [Google Scholar](#)
- Gaines T. A., Duke S. O., Morran S., Rigon C. A. G., Tranel P. J., Küpper A., Dayan F. E. (2020) **Mechanisms of evolved herbicide resistance** *Journal of Biological Chemistry* **295**:10307–10330 [Google Scholar](#)
- Garud N. R., Petrov D. A. (2016) **Elevated linkage disequilibrium and signatures of soft sweeps are common in *Drosophila melanogaster*** *Genetics* **203**:863–880 [Google Scholar](#)
- Gong Y.-J., Chen J.-C., Jiang J.-Y., Wang Z.-H., Cao L.-J., Wei S.-J. (2017) **Toxicity and field control efficacy of the new acaricide SYP-9625 to the two-spotted spider mite (*Tetranychus urticae* koch)** *Agrochemicals* **56**:561–563 [Google Scholar](#)
- Gould F., Brown Z. S., Kuzma J. (2018) **Wicked evolution: Can we address the sociobiological dilemma of pesticide resistance?** *Science* **360**:728–732 [Google Scholar](#)
- Groeters F. R., Tabashnik B. E. (2000) **Roles of selection intensity, major genes, and minor genes in evolution of insecticide resistance** *Journal of Economic Entomology* **93**:1580–1587 [Google Scholar](#)
- Guan F., Hou B., Dai X., Liu S., Liu J., Gu Y., Jin L., Yang Y., Fabrick J. A., Wu Y. (2021) **Multiple origins of a single point mutation in the cotton bollworm tetraspanin gene confers dominant resistance to Bt cotton** *Pest Management Science* **77**:1169–1177 [Google Scholar](#)
- Hartl D. L., Clark A. G., Clark A. G. (1997) **Principles of population genetics** Sinauer associates Sunderland, MA [Google Scholar](#)
- Hartley C., Newcomb R., Russell R., Yong C., Stevens J., Yeates D., La Salle J., Oakeshott J. (2006) **Amplification of DNA from preserved specimens shows blowflies were preadapted for the rapid evolution of insecticide resistance** *Proceedings of the National Academy of Sciences of the United States of America* **103**:8757–8762 [Google Scholar](#)
- Hawkins N. J., Bass C., Dixon A., Neve P. (2019) **The evolutionary origins of pesticide resistance** *Biological Reviews* **94**:135–155 [Google Scholar](#)
- Hayashi N., Sasama Y., Takahashi N., Ikemi N. (2013) **Cyflumetofen, a novel acaricide—its mode of action and selectivity** *Pest Management Science* **69**:1080–1084 [Google Scholar](#)
- Hermisson J., Pennings P. S. (2005) **Soft sweeps: molecular population genetics of adaptation from standing genetic variation** *Genetics* **169**:2335–52 [Google Scholar](#)
- Hermisson J., Pennings P. S. (2017) **Soft sweeps and beyond: understanding the patterns and probabilities of selection footprints under rapid adaptation** *Methods in Ecology and Evolution* **8**:700–716 [Google Scholar](#)
- Hoban S., Kelley J. L., Lotterhos K. E., Antolin M. F., Bradburd G., Lowry D. B., Poss M. L., Reed L. K., Storfer A., Whitlock M. C. (2016) **Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future directions** *American Naturalist* **188**:379–397 [Google Scholar](#)
- Ilias A., Vontas J., Tsagkarakou A. (2014) **Global distribution and origin of target site insecticide resistance mutations in *Tetranychus urticae*** *Insect Biochemistry and Molecular Biology* **48**:17–28 [Google Scholar](#)

Keightley P. D., Ness R. W., Halligan D. L., Haddrill P. R (2014) **Estimation of the spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family** *Genetics* **196**:313–20 [Google Scholar](#)

Kersten S., Chang J., Huber C. D., Voichek Y., Lanz C., Hagmaier T., Lang P., Lutz U., Hirschberg I., Lerchl J., Porri A., Peer Y. V. D., Schmid K., Weigel D., Rabanal F. A (2023) **Standing genetic variation fuels rapid evolution of herbicide resistance in blackgrass** *Proceedings of the National Academy of Sciences of the United States of America* **120**:e2206808120 [Google Scholar](#)

Kofler R., Orozco-Terwengel P., De Maio N., Pandey R.V., Nolte V., Futschik A., Kosiol C., Schlötterer C. (2011a) **PoPoolation: a toolbox for population genetic analysis of next generation sequencing data from pooled individuals** *PLoS One* **6**:e15925 [Google Scholar](#)

Kofler R., Pandey R. V., Schlotterer C (2011b) **PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq)** *Bioinformatics* **27**:3435–3436 [Google Scholar](#)

Konecna V., Bray S., Vlcek J., Bohutinska M., Pozarova D., Choudhury R. R., A. Bollmann-Giolai, Flis P., Salt D.E., Parisod C., Yant L., Kolář F. (2021) **Parallel adaptation in autopolyploid *Arabidopsis arenosa* is dominated by repeated recruitment of shared alleles** *Nature Communications* **12**:4979 [Google Scholar](#)

Kreiner J. M., Giacomini D. A., Bemm F., Waithaka B., Regalado J., Lanz C., Hildebrandt J., Sikkema P. H., Tranel P. J., Weigel D., Stinchcombe J. R., Wright S. I (2019) **Multiple modes of convergent adaptation in the spread of glyphosate-resistant *Amaranthus tuberculatus*** *Proceedings of the National Academy of Sciences of the United States of America* **116**:21076–21084 [Google Scholar](#)

Kreiner J. M., Latorre S. M., Burbano H. A., Stinchcombe J. R., Otto S. P., Weigel D., Wright S. I (2022a) **Rapid weed adaptation and range expansion in response to agriculture over the past two centuries** *Science* **378**:1079–1085 [Google Scholar](#)

Kreiner J. M., Sandler G., Stern A. J., Tranel P. J., Weigel D., Stinchcombe J. R., Wright S. I (2022b) . **Repeated origins, widespread gene flow, and allelic interactions of target-site herbicide resistance mutations** *eLife* **11**:70242 [Google Scholar](#)

Legan A. W., Allan C. W., Jensen Z. N., Degain B. A., Yang F., Kerns D. L., Benowitz K. M., Fabrick J. A., Li X., Carrière Y., Matzkin L. M., Tabashnik B. E. (2024) **Mismatch between lab-generated and field-evolved resistance to transgenic Bt crops in *Helicoverpa zea*** *Proceedings of the National Academy of Sciences of the United States of America* **121**:e2416091121 [Google Scholar](#)

Li H., Handsaker B., Wysoker A., Fennell T., Ruan J., Homer N., Marth G., Abecasis G., Durbin R., 1000 Genome Project Data Processing Subgroup (2009) **The sequence alignment/map format and SAMtools** *Bioinformatics* **25**:2078–2079 [Google Scholar](#)

Li N., Stephens M (2003) **Modeling linkage disequilibrium and identifying recombination hotspots using single-nucleotide polymorphism data** *Genetics* **165**:2213–2233 [Google Scholar](#)

Lucas E. R., Nagi S. C., Egyir-Yawson A., Essandoh J., Dadzie S., Chabi J., Djogbenou L. S., Medjigbodo A. A., Edi C. V., Ketoh G. K., Koudou B. G., Van't Hof A. E., Rippon E. J., Pipini D., Harding N. J., Dyer N. A., Cerdeira L. T., Clarkson C. S., Kwiatkowski D. P., Miles A., Donnelly M. J., Weetman D. (2023) **Genome-wide association studies reveal novel loci associated with**

pyrethroid and organophosphate resistance in *Anopheles gambiae* and *Anopheles coluzzii* *Nature Communications* **14**:4946 [Google Scholar](#)

Maeoka A., Osakabe M (2021) **Co-occurrence of subunit B and C mutations in respiratory complex II confers high resistance levels to pyflubumide and cyenopyrafen in the two-spotted spider mite *Tetranychus urticae* (Acari: Tetranychidae)** *Pest Management Science* **77**:5149–5157 [Google Scholar](#)

Mckenzie J. A., Batterham P (1994) **The genetic, molecular and phenotypic consequences of selection for insecticide resistance** *Trends in Ecology and Evolution* **9**:166–169 [Google Scholar](#)

Mckenzie J. A., Parker A., Yen J (1992) **Polygenic and single gene responses to selection for resistance to diazinon in *Lucilia cuprina*** *Genetics* **130**:613–620 [Google Scholar](#)

Melander A (1914) **Can insects become resistant to sprays?** *Journal of Economic Entomology* **7**:167 [Google Scholar](#)

Nakahira K (2011) **Strategy for discovery of a novel miticide Cyenopyrafen which is one of electron transport chain inhibitors** *Journal of Pesticide Science* **36**:511–515 [Google Scholar](#)

Njiru C., Saalwaechter C., Gutbrod O., Geibel S., Wybouw N., Van Leeuwen T. (2022) **A H258Y mutation in subunit B of the succinate dehydrogenase complex of the spider mite *Tetranychus urticae* confers resistance to cyenopyrafen and pyflubumide, but likely reinforces cyflumetofen binding and toxicity** *Insect Biochemistry and Molecular Biology* **144**:103761 [Google Scholar](#)

Njiru C., Saalwaechter C., Mavridis K., Vontas J., Geibel S., Wybouw N., Van Leeuwen T. (2023) **The complex II resistance mutation H258Y in succinate dehydrogenase subunit B causes fitness penalties associated with mitochondrial respiratory deficiency** *Pest Management Science* **79**:4403–4413 [Google Scholar](#)

Paradis E (2010) . **pegas: an R package for population genetics with an integrated-modular approach** *Bioinformatics* **26**:419–20 [Google Scholar](#)

PÉLISSIÉ B., Chen Y. H., Cohen Z. P., Crossley M. S., Hawthorne D. J., Izzo V., Schoville S. D (2022) **Genome resequencing reveals rapid, repeated evolution in the Colorado potato beetle** *Molecular Biology and Evolution* **39**:msac016 [Google Scholar](#)

Pennings P. S., Hermisson J (2006) **Soft sweeps II—molecular population genetics of adaptation from recurrent mutation or migration** *Molecular Biology and Evolution* **23**:1076–1084 [Google Scholar](#)

Peterson M. A., Collavo A., Ovejero R., Shivrain V., Walsh M. J (2018) **The challenge of herbicide resistance around the world: a current summary** *Pest Management Science* **74**:2246–2259 [Google Scholar](#)

Pezzini D., Taylor K. L., Reisig D. D., Fritz M. L (2024) **Cross-pollination in seed-blended refuge and selection for Vip3A resistance in a lepidopteran pest as detected by genomic monitoring** *Proceedings of the National Academy of Sciences of the United States of America* **121**:e2319838121 [Google Scholar](#)

- Pickrell J. K., Pritchard J. K (2012) **Inference of population splits and mixtures from genome-wide allele frequency data** *PLoS Genetics* **8**:e1002967 [Google Scholar](#)
- Pool J. E., Braun D. T., Lack J. B (2017) **Parallel evolution of cold tolerance within *Drosophila melanogaster*** *Molecular Biology and Evolution* **34**:349–360 [Google Scholar](#)
- Ralph P., Coop G (2010) **Parallel adaptation: one or many waves of advance of an advantageous allele?** *Genetics* **186**:647–668 [Google Scholar](#)
- Raymond M., Callaghan A., Fort P., Pasteur N (1991) **Worldwide migration of amplified insecticide resistance genes in mosquitoes** *Nature* **350**:151 [Google Scholar](#)
- Roush R. T., McKenzie J. A (1987) **Ecological genetics of insecticide and acaricide resistance** *Annual Review of Entomology* **32**:361–380 [Google Scholar](#)
- Schmidt J. M., Good R. T., Appleton B., Sherrard J., Raymant G. C., Bogwitz M. R., Martin J., Daborn P. J., Goddard M. E., Batterham P (2010) **Copy number variation and transposable elements feature in recent, ongoing adaptation at the Cyp6g1 locus** *PLoS Genetics* **6**:e1000998 [Google Scholar](#)
- Schmidt T. L., Endersby-Harshman N. M., van Rooyen A.R.J., Katusele M., Vinit R., Robinson L.J., Laman M., Karl S., Hoffmann A.A (2024) **Global, asynchronous partial sweeps at multiple insecticide resistance genes in *Aedes* mosquitoes** *Nature Communications* **15**:6251 [Google Scholar](#)
- Schrider D. R., Mendes F. K., Hahn M. W., Kern A. D (2015) **Soft shoulders ahead: spurious signatures of soft and partial selective sweeps result from linked hard sweeps** *Genetics* **200**:267–284 [Google Scholar](#)
- Semenov G. A., Safran R. J., Smith C. C. R., Turbek S. P., Mullen S. P., Flaxman S. M (2019) **Unifying theoretical and empirical perspectives on genomic differentiation** *Trends in Ecology and Evolution* **34**:987–995 [Google Scholar](#)
- Sparks T. C., Storer N., Porter A., Slater R., Nauen R (2021) **Insecticide resistance management and industry: the origins and evolution of the Insecticide Resistance Action Committee (IRAC) and the mode of action classification scheme** *Pest Management Science* **77**:2609–2619 [Google Scholar](#)
- Sugimoto N., Takahashi A., Ihara R., Itoh Y., Jouraku A., van Leeuwen T., & Osakabe M. (2020) **QTL mapping using microsatellite linkage reveals target-site mutations associated with high levels of resistance against three mitochondrial complex II inhibitors in *Tetranychus urticae*** *Insect Biochemistry and Molecular Biology* **123**:103410 [Google Scholar](#)
- Sukumaran J., Holder M. T (2010) **DendroPy: a Python library for phylogenetic computing** *Bioinformatics* **26**:1569–1571 [Google Scholar](#)
- Tabashnik B. E., Brévault T., Carrière Y. (2013) **Insect resistance to Bt crops: lessons from the first billion acres** *Nature Biotechnology* **31**:510–521 [Google Scholar](#)
- Tamura K., Stecher G., Kumar S (2021) **MEGA11: Molecular Evolutionary Genetics Analysis Version 11** *Molecular Biology and Evolution* **38**:3022–3027 [Google Scholar](#)
- Tang Q. Y., Zhang C. X (2013) **Data Processing System (DPS) software with experimental design, statistical analysis and data mining developed for use in entomological research**

Insect science **20**:254–260 [Google Scholar](#)

Taylor K. L., Hamby K. A., Deyonke A. M., Gould F., Fritz M. L. (2021) **Genome evolution in an agricultural pest following adoption of transgenic crops** *Proceedings of the National Academy of Sciences of the United States of America* **118**:e2020853118 [Google Scholar](#)

Tranel P. J., Wright T. R., Heap I. M. (2025) **Mutations in herbicide-resistant weeds to Inhibition of Acetolactate Synthase [Online]** <http://www.weedscience.org>

Trocza B., Zimmer C. T., Elias J., Schorn C., Bass C., Davies T. E., Field L. M., Williamson M. S., Slater R., Nauen R. (2012) **Resistance to diamide insecticides in diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae) is associated with a mutation in the membrane-spanning domain of the ryanodine receptor** *Insect Biochemistry and Molecular Biology* **42**:873–880 [Google Scholar](#)

Van Etten M., Lee K. M., Chang S.-M., Baucom R. S. (2020) **Parallel and nonparallel genomic responses contribute to herbicide resistance in *Ipomoea purpurea*, a common agricultural weed** *PLoS Genetics* **16**:e1008593 [Google Scholar](#)

Van Leeuwen T., Demaeght P., Osborne E. J., Dermauw W., Gohlke S., Nauen R., Grbic M., Tirry L., Merzendorfer H., Clark R. M. (2012) **Population bulk segregant mapping uncovers resistance mutations and the mode of action of a chitin synthesis inhibitor in arthropods** *Proceedings of the National Academy of Sciences of the United States of America* **109**:4407–4412 [Google Scholar](#)

Van Leeuwen T., Vontas J., Tsagkarakou A., Dermauw W., Tirry L. (2010) **Acaricide resistance mechanisms in the two-spotted spider mite *Tetranychus urticae* and other important Acari: a review** *Insect Biochemistry and Molecular Biology* **40**:563–572 [Google Scholar](#)

Vasimuddin M., Misra S., Li H., Aluru S. (2019) **Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems** In: *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)* pp. 314–324 [Google Scholar](#)

Walsh T., Heckel D. G., Wu Y., Downes S., Gordon K., Oakeshott J. (2022) **Determinants of insecticide resistance evolution: comparative analysis among Heliothines** *Annual Review of Entomology* **67**:387–406 [Google Scholar](#)

Waterhouse A., Bertoni M., Bienert S., Studer G., Tauriello G., Gumienny R., Heer F. T., de Beer T.A.P., Rempfer C., Bordoli L., Lepore R., Schwede T. (2018) **SWISS-MODEL: homology modelling of protein structures and complexes** *Nucleic Acids Research* **46**:W296–W303 [Google Scholar](#)

Wei S.-J., Cao L.-J. (2025) **Unprecedented number of recurrent mutations drive the rapid evolution of pesticide resistance in a notorious pest** *eLife* [Google Scholar](#)

Yassin A., Debat V., Bastide H., Gidaszewski N., David J. R., Pool J. E. (2016) **Recurrent specialization on a toxic fruit in an island *Drosophila* population** *Proceedings of the National Academy of Sciences of the United States of America* **113**:4771–4776 [Google Scholar](#)

Author information

Li-Jun Cao[#]

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China

[#]equal contributions

Jin-Cui Chen[#]

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China

[#]equal contributions

Joshua A Thia[#]

Bio21 Institute, School of BioSciences, University of Melbourne, Parkville, Australia

[#]equal contributions

Thomas L Schmidt

Bio21 Institute, School of BioSciences, University of Melbourne, Parkville, Australia

Richard Ffrench-Constant

Biosciences, University of Exeter, Exeter, United Kingdom

ORCID iD: [0000-0001-5385-9888](https://orcid.org/0000-0001-5385-9888)

Lin-Xi Zhang

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China, Department of Entomology, College of Plant Protection, China Agricultural University, Beijing, China

Yu Yang

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China, College of Plant Science and Technology, Beijing University of Agriculture, Beijing, China

Meng-Chu Yuan

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China, College of Plant Science and Technology, Beijing University of Agriculture, Beijing, China

Jia-Yue Zhang

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China, Hebei North University, College of Agriculture and Forestry Technology, Zhangjiakou, China

Xiao-Yang Zhang

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China, College of Plant Science and Technology, Beijing University of Agriculture, Beijing, China

Qiong Yang

Bio21 Institute, School of BioSciences, University of Melbourne, Parkville, Australia
ORCID iD: [0000-0003-3336-5703](https://orcid.org/0000-0003-3336-5703)

Ya-Jun Gong

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China

Hu Li

Department of Entomology, College of Plant Protection, China Agricultural University, Beijing, China

Xue-Xin Chen

Institute of Insect Science, Zhejiang University, Hangzhou, China

Ary A Hoffmann

Bio21 Institute, School of BioSciences, University of Melbourne, Parkville, Australia
ORCID iD: [0000-0001-9497-7645](https://orcid.org/0000-0001-9497-7645)

For correspondence: ary@unimelb.edu.au

Shu-Jun Wei

Institute of Plant Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China
ORCID iD: [0000-0001-7398-0968](https://orcid.org/0000-0001-7398-0968)

For correspondence: shujun268@163.com

Editors

Reviewing Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #1 (Public review):

Summary:

The study by Cao et al. provides a compelling investigation into the role of mutational input in the rapid evolution of pesticide resistance, focusing on the two-spotted spider mite's response to the recent introduction of the acaricide cyetpyrafen. This well-documented introduction of the pesticide-and thus a clearly defined history of selection-offers a powerful framework for studying the temporal dynamics of rapid adaptation. The authors combine resistance phenotyping across multiple populations, extensive resequencing to track the frequency of resistance alleles, and genomic analyses of selection in both contemporary and historical samples. These approaches are further complemented by laboratory-based experimental evolution, which serves as a baseline for understanding the genetic architecture of resistance across mite populations in China. Their analyses identify two key

resistance-associated genes, *sdhB* and *sdhD*, within which they detect 15 mutations in wild-collected samples. Protein modeling reveals that these mutations cluster around the pesticide's binding site, suggesting a direct functional role in resistance. The authors further examine signatures of selective sweeps and their distribution across populations to infer the mechanisms-such as de novo mutation or gene flow-driving the spread of resistance, a crucial consideration for predicting evolutionary responses to extreme selection pressure. Overall, this is a well-rounded, thoughtfully designed and well-written manuscript. It shows significant novelty, as it is relatively rare to integrate broad-scale evolutionary inference from natural populations with experimentally informed bioassays, however, follow up work will be needed to fully resolve haplotype structure and the functional effects of resistance mutations in the system.

Strengths:

One of the most compelling aspects of this study is its integration of genomic time-series data in natural populations with controlled experimental evolution. By coupling genome sequencing of resistant field populations with laboratory selection experiments, the authors tease apart the individual effects of resistance alleles along with regions of the genome where selection is expected to occur, and compare that to the observed frequency in the wild populations over space and time. Their temporal data clearly demonstrates the pace at which evolution can occur in response to extreme selection. This type of approach is a powerful roadmap for the rest of the field of rapid adaptation.

The study effectively links specific genetic changes to resistance phenotypes. The identification of *sdhB* and *sdhD* mutations as major drivers of cyetpyrafen resistance is well supported by allele frequency shifts in both field and experimental populations. The scope of their sampling clearly facilitated the remarkable number of observed mutations within these target genes, and the authors provide a careful discussion of the likelihood of these mutations from de novo or standing variation. Furthermore, the discovered cross-resistance that these mutations confer to other mitochondrial complex II inhibitors highlights the potential for broader resistance management and evolution.

Weaknesses:

- (1) Pleiotropy without pesticide modes of action (cyflumetofen and cyetpyrafen) may also play a role in the rapid response to the focal pesticide in this study
- (2) Other aspects of the environment that might influence selection were not considered in the structure of resistance alleles (i.e. climate, elevation)
- (3) Very little data were used for haplotype reconstruction, only 8 SNPs, and this excluded all heterozygous alleles, which could dramatically influence the complexity of these inferred haplotype networks.
- (4) Single Mutations and Their Effects:
 - Allelic effects were not estimated in isogenic lines, so the effects presented also include heterogeneity from allelic interactions with the genomic background
 - The authors see populations that segregate for resistance mutations but that have no survival to pesticides. This suggests either not all of the resistance mutations studied here actually have functional effects or that dominance is playing a role in masking their effects in the heterozygous state.

<https://doi.org/10.7554/eLife.106288.2.sa2>

Reviewer #2 (Public review):

Summary:

This paper investigates the evolution of pesticide resistance in the two-spotted spider mite following the introduction of an SDHI acaricide, cyatpyrafen, in China. The authors make use of cyatpyrafen-naive populations collected before that pesticide was first used, as well as more recent populations (both sensitive and resistant) to conduct comparative population genomics. They report 15 different mutations in the insecticide target site from resistant populations, many reported here for the first time, and look at the mutation and selection processes underlying the evolution of resistance, through GWAS, haplotype mapping, and testing for loss of diversity indicating selective sweeps. None of the target site mutations found in resistant populations was found in pre-exposure populations, suggesting that the mutations may have arisen *de novo* rather than being present as standing variation, unless initially present at very low frequencies; a *de novo* origin is also supported by evidence of selective sweeps in some resistant populations. Furthermore, there is no significant evidence of migration of resistant genotypes between the sampled field populations indicating multiple origins of common mutations. Overall, this indicates a very high mutation rate and a wide range of mutational pathways to resistance for this target site in this pest species. The series of population genomic analyses carried out here, in addition to the evolutionary processes that appear to underly resistance development in this case, could have implications for the study of resistance evolution more widely.

Strengths:

This paper combines phenotypic characterisation with extensive comparative population genomics, made possible by the availability of multiple population samples (each with hundreds of individuals) collected before as well as after then introduction of the pesticide cyatpyrafen, as well as lab-evolved lines. This results in findings of mutation and selection processes that can be related back to the pesticide resistance trait of concern. Large numbers of mites were tested phenotypically to show the levels of resistance present, and the authors also made near-isogenic lines to confirm the phenotypic effects of key mutations. The population genomic analyses consider a range of alternative hypotheses, including mutations arising by *de novo* mutation or selection from standing genetic variation; and mutations in different populations arising independently or arriving by migration. The claim that mutations most likely arose by multiple repeated *de novo* mutations is therefore supported by multiple lines of evidence: the direct evidence of none of the mutations being found in over 2000 individuals from naive populations, and the indirect evidence from population genomics showing evidence of selective sweeps but not of significant migration between the sampled populations.

Weaknesses:

As acknowledged within the discussion, whilst evidence supports a *de novo* origin of the resistance associated mutations, this cannot be proven definitively as mutations may have been present at a very low frequency and therefore not found within the tested pesticide-naive population samples.

Near-isofemale lines were made to confirm the resistance levels associated with five of the 15 mutations, but otherwise the genotype-phenotype associations are correlative as confirmation by functional genetics was beyond the scope of this study.

Comments on revisions:

My recommendations have all been addressed in the revised version.

<https://doi.org/10.7554/eLife.106288.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

Summary:

*The study by Cao et al. provides a compelling investigation into the role of mutational input in the rapid evolution of pesticide resistance, focusing on the two-spotted spider mite's response to the recent introduction of the acaricide cyetpyrafen. This well-documented introduction of the pesticide - and thus a clearly defined history of selection - offers a powerful framework for studying the temporal dynamics of rapid adaptation. The authors combine resistance phenotyping across multiple populations, extensive resequencing to track the frequency of resistance alleles, and genomic analyses of selection in both contemporary and historical samples. These approaches are further complemented by laboratory-based experimental evolution, which serves as a baseline for understanding the genetic architecture of resistance across mite populations in China. Their analyses identify two key resistance-associated genes, *sdhB* and *sdhD*, within which they detect 15 mutations in wild-collected samples. Protein modeling reveals that these mutations cluster around the pesticide's binding site, suggesting a direct functional role in resistance. The authors further examine signatures of selective sweeps and their distribution across populations to infer the mechanisms - such as de novo mutation or gene flow-driving the spread of resistance, a crucial consideration for predicting evolutionary responses to extreme selection pressure. Overall, this is a well-rounded, thoughtfully designed, and well-written manuscript. It shows significant novelty, as it is relatively rare to integrate broad-scale evolutionary inference from natural populations with experimentally informed bioassays, however, some aspects of the methods and discussion have an opportunity to be clarified and strengthened.*

Strengths:

One of the most compelling aspects of this study is its integration of genomic time-series data in natural populations with controlled experimental evolution. By coupling genome sequencing of resistant field populations with laboratory selection experiments, the authors tease apart the individual effects of resistance alleles along with regions of the genome where selection is expected to occur, and compare that to the observed frequency in the wild populations over space and time. Their temporal data clearly demonstrates the pace at which evolution can occur in response to extreme selection. This type of approach is a powerful roadmap for the rest of the field of rapid adaptation.

*The study effectively links specific genetic changes to resistance phenotypes. The identification of *sdhB* and *sdhD* mutations as major drivers of cyetpyrafen resistance is well-supported by allele frequency shifts in both field and experimental populations. The scope of their sampling clearly facilitated the remarkable number of observed mutations within these target genes, and the authors provide a careful discussion of the likelihood of these mutations from de novo or standing variation. Furthermore, the discovered cross-resistance that these mutations confer to other mitochondrial complex II inhibitors highlights the potential for broader resistance management and evolution.*

Weaknesses:

(1) Experimental Evolution:

- Additional information about the lab experimental evolution would be useful in the main text. Specifically, the dose of cyetpyrafen used should be clarified, especially with respect to the LD50 values. How does it compare to recommended field doses? This is expected to influence the architecture of resistance evolution. What was the sample size? This will help readers contextualize how the experimental design could influence the role of standing variation.

The experimental design involved sampling approximately 6,000 individuals from the wild population ZJSX1, which were subsequently divided into two parallel cohorts under controlled laboratory conditions. The selection group (LabR) was subjected to continuous selection pressure using cyetpyrafen, while the control group (LabS) was maintained under identical laboratory conditions without exposure to acyetypyrafen. A dynamic selection regime was implemented wherein the acaricide dosage was systematically adjusted every two generations to maintain a consistent selection intensity, achieving a mortality rate of 60% $\pm$ 10% in the LabR population. This adaptive dosage strategy ensured sustained evolutionary pressure while preventing population collapse. The LC₅₀ values were tested at F1, F32, F54, F60, F62, and F66 generations using standardized bioassay protocols to quantify resistance development trajectories and optimize dosage for subsequent selection cycles. We provided the additional information in subsection 4.1 of the materials and methods section.

- The finding that lab-evolved strains show cross-resistance is interesting, but potentially complicates the story. It would help to know more about the other mitochondrial complex II inhibitors used across China and their impact on adaptive dynamics at these loci, particularly regarding pre-existing resistance alleles. For example, a comparison of usage data from 2013, 2017, and 2019 could help explain whether cyetpyrafen was the main driver of resistance or if previous pesticides played a role. What happened in 2020 that caused such rapid evolution 3 years after launch?

Although the introduction of the other two SDHI acaricides complicates the story, we would like to provide a complete background on the usage of acaricides with this mode of action in China. Although cyflumetofen was released in 2013 before cyetpyrafen, and cyenopyrafen was released in 2019 after cyetpyrafen, their market share is minor (about 3.2%) compared to cyetpyrafen (about 96.8%, personal communication). Since cross-resistance is reported among SDHIs, we could not exclude the contribution of cyflumetofen to the initial accumulation of resistance alleles, but the effect should be minor, both because of their minimal market share and because of the independent evolution of resistance in the field as found in our study. Although the contribution of cyflumetofen and cyenopyrafen cannot be entirely excluded, the rapid evolution of resistance seems likely to be mainly explained by the intensive application of cyetpyrafen. To clarify this issue, we added relevant information in the first paragraph of the discussion section.

(2) Evolutionary history of resistance alleles:

- It would be beneficial to examine the population structure of the sampled populations, especially regarding the role of migration. Though resistance evolution appears to have had minimal impact on genome-wide diversity (as shown in Supplementary Figure 2), could admixture be influencing the results? An explicit multivariate regression framework could help to understand factors influencing diversity across populations, as right now much is left to the readers' visual acuity.

The genetic structure of the populations was examined by Treemix analysis. We detected only one migration event from JXNC to SHPD (no resistance data available for these two populations), suggesting a limited role for migration to resistance evolution. The multiple regression analysis revealed that overall genetic diversity and Tajima's D across the genome

were not significantly associated with resistance levels, genetic structure or geographic coordinates ($P > 0.05$), which all support a limited role of migration in resistance development.

- It is unclear why lab populations were included in the migration/treemix analysis. We might suggest redoing the analysis without including the laboratory populations to reveal biologically plausible patterns of resistance evolution.

Thank you for the constructive suggestion. The Treemix analysis was redone by removing laboratory populations and is now reported.

- Can the authors explore isolation by distance (IBD) in the frequency of resistance alleles?

Thank you for the constructive suggestion. No significant isolation-by-distance pattern was detected for resistance allele frequencies across all surveyed years (2020: $P=0.73$; 2021: $P=0.52$; 2023: $P=0.16$; Mantel test). We added these results to the text.

- Given the claim regarding the novelty of the number of pesticide resistance mutations, it is important to acknowledge the evolution of resistance to all pesticides (antibiotics, herbicides, etc.). ALS-inhibiting herbicides have driven remarkable repeatability across species based on numerous SNPs within the target gene.

We appreciate this comment, which highlights the need to place our findings within the broader evolutionary context of pesticide resistance. We have investigated references relevant to the evolution of resistance to diverse pesticides. As far as we can tell, the 15 target mutations in eight amino acid residues are among the highest number of pesticide resistance mutations detected, especially within the context of animal studies. We have added relevant text to the second paragraph of the discussion.

- Figure 5 A-B. Why not run a multivariate regression with status at each resistance mutation encoded as a separate predictor? It is interesting that focusing on the predominant mutation gives the strongest r^2 , but it is somewhat unintuitive and masks some interesting variation among populations.

We conducted a multiple regression analysis to explore the influence of multiple mutations on resistance levels of field populations. However of 15 putative resistant mutations, only five were detected in more than three populations where bioassay data are available, i.e. I260T, I260V, D116G, R119C, R119L. The frequency of three of these mutations, I260T ($P = 0.00128$), I260V ($P = 0.00423$) and D116G ($P = 0.00058$), are significantly correlated with the resistance level of field populations. This has been added.

(3) Haplotype Reconstruction (Line 271-):

- We are a bit sceptical of the methods taken to reconstruct these haplotypes. It seems as though the authors did so with Sanger sequencing (this should be mentioned in the text), focusing only on homozygous SNPs. How many such SNPs were used to reconstruct haplotypes, along what length of sequence? For how many individuals were haplotypes reconstructed? Nonetheless, I appreciated that the authors looked into the extent to which the reconstructed haplotypes could be driven by recombination. Can the authors elaborate on the calculations in line 296? Is that the census population size estimate or effective?

Because haplotypes could not be determined when more than two loci were heterozygous, we detected haplotypes from sequencing data with at most one heterozygous locus. In total 844 individuals and 696 individuals were used to detect haplotypes of *sdhB* and *sdhD*. We detected 11 haplotypes (with 8 SNPs) and 24 haplotypes (with 11 SNPs) along 216 bp of the *sdhB* and 155 bp of the *sdhD* genes, respectively. Please see the fifth paragraph of subsection 2.4. We used $p = 4 \times N_e \times d$ (genetic distance) (Li and Stephens, 2003) to calculate the number of effective individuals for one recombination event.

(4) Single Mutations and Their Effect (line 312-):

- It's not entirely clear how the breeding scheme resulted in near-isogenic lines. Could the authors provide a clearer explanation of the process and its biological implications?

To investigate the effect of single mutations or their combination on resistance levels, we isolated the females and males with the same homozygous/ hemizygous genotypes for creating homozygous lines. Females from these lines were not near-isogenic, but homozygous for the critical mutations. We revised the description in the methods section to clearly define these lines.

- If they are indeed isogenic, it's interesting that individual resistance mutations have effects on resistance that vary considerably among lines. Could the authors run a multivariate analysis including all potential resistance SNPs to account for interactions between them? Given the variable effects of the D116G substitution (ranging from 4-25%), could polygenic or epistatic factors be influencing the evolution of resistance?

We couldn't conduct multivariate analysis because most lines have only one resistant SNP. The four lines homozygous for 116G were from the same population. The variable mortality may reflect other unknown mechanisms but these are beyond the scope of this study.

- Why are there some populations that segregate for resistance mutations but have no survival to pesticides (i.e., the green points in Figure 5)? Some discussion of this heterogeneity seems required in the absence of validation of the effects of these particular mutations. Could it be dominance playing a role, or do the authors have some other explanation?

We didn't investigate the degree of dominance of each mutation. The mutation I260V shows incompletely dominant inheritance (Sun, et al. 2022). To investigate survival rate of different populations, the two-spotted spider mite *T. urticae* was exposed to 1000 mg/L of cyetpyrafen, higher than the recommended field dose of 100 mg/L. Such a high concentration may lead to death of an individual heterozygous for certain mutations, such as I260V.

- The authors mention that all resistance mutations co-localized to the Q-site. Is this where the pesticide binds? This seems like an important point to follow their argument for these being resistance-related.

Yes. We revised Fig. 3c to show the Q-site.

(5) Statistical Considerations for Allele Frequency Changes (Figure 3):

- It might be helpful to use a logistic regression model to assess the rate of allele frequency changes and determine the strength of selection acting on these alleles (e.g., Kreiner et al. 2022; Patel et al. 2024). This approach could refine the interpretation of selection dynamics over time.

Thank you for this suggestion. A logistic regression model was used to track allele frequencies trajectories. The selection coefficient of each allele and their joint effects were estimated.

Reviewer #2 (Public review):

Summary:

This paper investigates the evolution of pesticide resistance in the two-spotted spider mite following the introduction of an SDHI acaricide, cyatpyrafen, in China. The authors make use of cyatpyrafen-naïve populations collected before that pesticide was first used, as well as more recent populations (both sensitive and resistant) to conduct comparative population genomics. They report 15 different mutations in the insecticide target site from resistant populations, many reported here for the first time, and look at the mutation and selection processes underlying the evolution of resistance, through GWAS, haplotype mapping, and testing for loss of diversity indicating selective sweeps. None of the target site mutations found in resistant populations was found in pre-exposure populations, suggesting that the mutations may have arisen de novo rather than being present as standing variation, unless initially present at very low frequencies; a de novo origin is also supported by evidence of selective sweeps in some resistant populations. Furthermore, there is no significant evidence of migration of resistant genotypes between the sampled field populations, indicating multiple origins of common mutations. Overall, this indicates a very high mutation rate and a wide range of mutational pathways to resistance for this target site in this pest species. The series of population genomic analyses carried out here, in addition to the evolutionary processes that appear to underlie resistance development in this case, could have implications for the study of resistance evolution more widely.

Strengths:

This paper combines phenotypic characterisation with extensive comparative population genomics, made possible by the availability of multiple population samples (each with hundreds of individuals) collected before as well as after the introduction of the pesticide cyatpyrafen, as well as lab-evolved lines. This results in findings of mutation and selection processes that can be related back to the pesticide resistance trait of concern. Large numbers of mites were tested phenotypically to show the levels of resistance present, and the authors also made near-isogenic lines to confirm the phenotypic effects of key mutations. The population genomic analyses consider a range of alternative hypotheses, including mutations arising by de novo mutation or selection from standing genetic variation, and mutations in different populations arising independently or arriving by migration. The claim that mutations most likely arose by multiple repeated de novo mutations is therefore supported by multiple lines of evidence: the direct evidence of none of the mutations being found in over 2000 individuals from naïve populations, and the indirect evidence from population genomics showing evidence of selective sweeps but not of significant migration between the sampled populations.

Weaknesses:

As acknowledged within the discussion, whilst evidence supports a de novo origin of the resistance-associated mutations, this cannot be proven definitively as mutations may have been present at a very low frequency and therefore not found within the tested pesticide-naïve population samples.

We agree that we could not definitively exclude the presence of a very low incidence of favoured mutations before the introduction of this novel acaricide.

Near-isofemale lines were made to confirm the resistance levels associated with five of the 15 mutations, but otherwise, the genotype-phenotype associations are correlative, as confirmation by functional genetics was beyond the scope of this study.

We hope that future functional studies will validate the effects of these mutations on resistance in both the two-spotted spider mite *T. urticae* and other spider mite species. This could be done by creating near-isogenic female lines or using CRISPR-Cas9 technology, as gene knockouts have recently been established for *T. urticae*.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

(1) Could the authors elaborate on the environmental context (e.g., climate, geography) of the sampled populations to give more nuance to the analysis of genetic differentiation and resistance evolution?

We have explored the influence of geographic isolation on the frequency of resistance alleles by Mantel tests (isolation by distance). We didn't investigate the influence of climate, because most of the samples were from greenhouses, where the climate to which the pest is exposed is unclear.

(2) Line 161: is this supposed to be one R and one S?

Yes, we added this information (LabR and LabS).

(3) Line 207: variation is not saturated at the first two sites because the different combinations are not seen. This is a bit misleading.

What we wanted to indicate was that the two codon positions are saturated, rather than their combinations. We revised this sentence by adding "of each codon position".

(4) Line 376: continuous selection did not "result in a new mutation arising". Rather, the mutation arose and was subsequently selected on.

We revised the expression of this *de novo* mutation and selection process.

(5) Line 402: can the authors explore what N_e would be necessary to drive the number of mutational origins they observe, as in (Karasov et al. 2010)?

It is challenged to estimate N_e , especially when mutation rate data from the two-spotted spider mite *T. urticae* is unavailable. We observed 2.7 resistant mutations per population in samples collected in 2024, seven years after the release of cyetpyrafen. The estimated mutation rate (Θ) is 0.0193, given 20 generations per year for *T. urticae*. An effective population size (N_e) of 2.29×10^6 would be necessary to reach the number of *de novo* mutations observed in this study, given $\Theta = 3N_e\mu$ (haplodiploid sex determination of *T. urticae*) and a mutation rate of $\mu = 2.8 \times 10^{-9}$ per base pair per generation as estimated for *Drosophila melanogaster* (Keightley et al., 2014). The high reproductive capacity of *T. urticae* (> 100 eggs per female) and short generation time makes it easier to reach such a population size in the field as we now note.

(6) Line 482: how did the authors precisely kill 60% of samples with their selection? What was the applied rate? In general, listing the rates of insecticide used in dose response would be useful to decipher if LD50s are projected outside of the doses used (seems like

they are). In this case, authors should limit their estimates to those > the highest rate used in the dose response.

It is difficult to control mortality precisely. We applied cyetpyrafen every two generations but did not determine the LC_{50} every two generations. When mortality was lower than 60%, another round of spraying was applied by increasing the dosage of the pesticide. The LC_{50} values were tested at F_1 , F_{32} , F_{54} , F_{60} , F_{62} , and F_{66} generations to establish the trajectories around resistance.

(7) The light pink genomic region in Figure 2 was distracting. Why is it included if there is no discussion of genomic regions outside the *sdh* genes? Generally, there was a lot going on in this figure, and some guiding categories (i.e., lab selected vs wild population) on the figure itself could help orient the reader.

We included chromosome 2 colored in light pink/ red to show the selection signal across a wider genomic region. In the figure legend, we added a description of the lab selected, field resistant and field susceptible populations. Very little common selection signal was detected among resistant populations on chromosome 2, indicating this region was less likely to be involved in resistance evolution of *T. urticae* to cyetpyrafen. We also described the result briefly in the figure legend.

Reviewer #2 (Recommendations for the authors):

(1) The most significant aspect of this study is the use of multiple pest population samples taken before as well as after the introduction of a class of pesticides, allowing a thorough comparative population genomics study in a species where a range of resistance mutations have appeared within a few years. I would prefer to see a title conveying this significance, rather than the current study, which focuses on the total number of mutations and claimed notoriety of the (at that point unnamed) study species. Similarly, I would prefer an abstract that relies less on superlative claims and includes more details: the scientific name of the study species; the number of years in which resistance evolved; the number of historical specimens; how the resistance levels for single mutations were shown.

(1) The title was changed by adding “the two-spotted spider mite *Tetranychus urticae*” and removing the “unprecedented number” to emphasize that “recurrent mutations drive rapid evolution”, i.e., “Recurrent Mutations Drive the Rapid Evolution of Pesticide Resistance in the Two-spotted Spider Mite *Tetranychus urticae*.”

(2) The scientific name of the study species was added.

(3) The number of years in which resistance evolved was added.

(4) The number of historical specimens was added (2666).

(5) Because we used homozygous lines but not iso-genic lines or gene-edited lines, our bioassay data could not provide direct evidence on the level of resistance conferred by each mutation. We revised our description of the results and removed this content from the abstract.

Line 29: if you want to claim the number is unprecedented, please specify the context: unprecedented for a pesticide target in an arthropod pest? (more resistance mutations may have been found in bacteria/fungi...).

We revised the sentence by adding “in an arthropod pest”.

Line 30: rather than a claim of notoriety, it may be better to specify what damage this pest causes.

Revised by describing it as an arthropod pest.

Line 34: please clarify, was this all in different haplotypes, or were some mutations found in combination?

Done: We identified 15 target mutations, including six mutations on five amino acid residues of subunit sdhB, and nine mutations on three amino acid residues of subunit sdhD, with as many as five substitutions on one residue.

(2) The introduction begins by framing the context as resistance evolution in invertebrate pests. However, the evolutionary processes examined in the study are applicable to resistance in other systems, and potentially to other cases of rapid contemporary evolution. The authors could show wider significance for their work beyond the subfield of invertebrate pests by including more of this wider context in their introduction and discussion: even if this means they can no longer claim novelty based on the number of mutations alone, the study is a strong example of the use of population genomics combined with functional and phenotypic characterisation to investigate the evolutionary processes underlying the emergence of resistance, so could have wider importance than within its current framing.

The background was revised as mentioned above to take this into account.

For example, in lines 48-50, please clarify what is meant by pesticides here (insects/arthropods? weeds and pathogens too?) In lines 69-73, the opposite is sometimes seen in fungal pathogens, with large numbers of mutations generated in lab-evolved strains.

We extended pesticides to those targeting arthropods, weeds and pathogens. We still emphasize the situation mainly with respect to arthropod pests.

(3) Lines 91-93: how many modes of action? How recently were SDHI acaricides introduced?

Added: at least 11 groups of acaricides based on their modes of action. SDHI was launched in 2007.

(4) Line 98-102: Use in China is a useful background for the study populations, but the global context should be included too.

Yes, four SDHI acaricides developed around the globe were introduced.

(5) Line 113: They show diverse mutations, but all within the mechanism of target-site point mutations.

We agree to your suggestion. This sentence has been removed as it repeats information stated above it.

(6) Line 115-116: Yes, agreed; I think this is the main strength of the current study and should be emphasised sooner.

Thanks.

(7) Line 158: Selective sweep signals were clear in half of the resistant populations but not in the others. The suggestion that the others had undergone soft sweeps, with multiple mutations increasing in frequency simultaneously but no one reaching fixation, seems reasonable; but the authors could compare the populations that did show a sweep with those that did not (for example, was there greater diversity or evenness of genotypes in those that did not?).

Five resistant populations with selection signals identified by PBE analysis (Figure 2b) showed corresponding decreases in π and Tajima's D near the two SDH genes but not across the genome (Figure S1).

(8) Line 313: please clarify "in combination with other mutations" within a mixed population or combined in one individual/haplotype? Also, the phrase "characterised the function" may be a little misleading, as this is a correlative analysis, not functional confirmation.

None of the combinations of different resistant mutations was observed in a single haplotype. Here, we examine resistance levels associated with a single mutation or two mutations on *sdhB* and *sdhD* in one individual, i.e. *sdhB*_I260V and *sdhD*_R119C. We revised the sentences to avoid any implication of functional confirmation.

(9) Line 358: again, please clarify the context: among arthropod pests?

Done.

(10) Line 360-363: please give some background on when and where these related compounds were introduced.

Added.

(11) Line 410: yes fitness costs may be a factor, but you could also give an example of a cost expressed in the absence of any pesticides, as well as the given example of negative cross-resistance.

We added the example of the H258Y mutation which causes both fitness costs and negative cross-resistance.

(12) Lines 419-438: this is one aspect where the situation for insecticides is in contrast with some other resistance areas.

Yes, we restricted these statements to arthropod pests.

(13) Line 466: some more detail could be given here: for example, SNP-specific monitoring would be less effective, but amplicon sequencing would be more suitable.

Yes, revised.

(14) Lines 472-475: Please list the numbers of field/lab, pre/post exposure, and sensitive/resistant populations within the main text.

Done. The number of sensitive/resistant populations was reported in the result section.

| (15) Line 483: randomly selected individuals?

Yes, added randomly selected individuals.

| (16) Line 556: Sanger sequencing to characterise populations? Or a number of individuals from each population?

Revised.

| (17) References: there are some duplicate entries, please check this.

Checked.

| (18) Figure 1e: consider a log(10) scale to better show large fold changes and avoid multiple axis breaks.

Thanks for your suggestions. However we didn't scale the LC_{50} value, because we wanted to show the specific impact of 1,000 mg/L. The breaks in the Y axis around 30 mg/L -1,000 mg/L reveal that the LC_{50} s of the resistant populations were all greater than 1000 mg/L, while those of the susceptible populations were all below 30 mg/L. This justified the use 1000 mg/L as a discriminating dose to investigate resistance status and level in subsequent work.

<https://doi.org/10.7554/eLife.106288.2.sa0>