

Mechanistic insights into transcriptional regulation of ARHGAP36 expression identify a factor predictive of neuroblastoma survival

Serhiy Havrylov, Armin M Gamper, Ordan J Lehmann 

Department of Medical Genetics, University of Alberta, Edmonton, Canada • Department of Ophthalmology, University of Alberta, Edmonton, Canada • Department of Oncology, Cross Cancer Institute, University of Alberta, Edmonton, Canada

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study examines the potential role of ARHGAP36 transcriptional regulation by FOXC1 in controlling sonic hedgehog signaling in human neuroblastoma. While there are many **solid** findings that strongly support this signaling pathway, there are some aspects of the study that are underdeveloped.

<https://doi.org/10.7554/eLife.108827.1.sa3>

Abstract

Cancer repeatedly exploits attributes fundamental for morphogenesis to advance malignancy and metastasis. This is illustrated by lineage specific transcription factors that regulate neural crest migration representing frequent drivers of malignancy. One such example is the *forkhead* transcription factor FOXC1 where gain of function is a feature of diverse cancers that is associated with an unfavourable prognosis. Using RNA-, ChIP-sequencing and CRISPR interference, we show that Foxc1 binds a locus in a region of closed chromatin to induce expression of Arhgap36, a tissue-specific inhibitor of Protein Kinase A. Because PKA is a core Hedgehog (Hh) pathway inhibitor, Foxc1's induction of Arhgap36 expression increases Hh activity. The function of Sufu, a PKA substrate and a second essential Hh pathway inhibitor, is likewise impaired. The resulting increased Hh pathway output is resistant to pharmacological inhibition of *Smoothened*, a phenotype of more aggressive cancers. The Foxc1-Arhgap36 relationship identified in murine cells was further evaluated in neuroblastoma, a neural crest derived pediatric malignancy. This demonstrated in a cohort of 1348 patients that high levels of ARHGAP36 are predictive of improved five-year survival. Accordingly, this study has identified as a novel transcription factor which enhances ARHGAP36 expression, one that induces Hh activity in multiple tissues during development. It also establishes a model by which increased levels of FOXC1 via ARHGAP36 and PKA inhibition dysregulate multiple facets of Hh signaling, and provides evidence demonstrating relevance to a common neural-crest derived malignancy.

Introduction

The neural crest comprises a multi-potent stem cell population that contributes to formation of diverse tissues. Through an epithelial to mesenchymal transition (EMT) that permits delamination from the neural plate, the acquisition of mesenchymal characteristics enables neural crest cells to transition towards a migratory phenotype. During a frequently lengthy path, neural crest cells proliferate, generating precursor cells that populate targets in multiple tissues, and so contribute to the formation of diverse organs. Such attributes essential to embryonic morphogenesis, are reminiscent of the invasive steps efficiently co-opted by cancer cells to drive malignancy and metastasis.

The lineage-specific transcription factors that regulate neural crest development represent frequent drivers of malignancy. An example of this paradigm is the *forkhead* box transcription factor *FOXC1*, that influences the migration and differentiation of neural crest cells. In addition to key roles in brain, ocular, cardiac, renal and skeletal development [1], *FOXC1* is aberrantly expressed in diverse cancers. This overexpression was first identified in basal like breast cancer [2, 3], the subtype with the worst prognosis, and subsequently in at least fifteen additional tumor types, encompassing both solid tumors and hematological malignancies, such as AML [4, 5]. *FOXC1* is also a prognostic factor for metastasis and survival, with its expression increasing tumour proliferation, invasion and dissemination [6]. Roles that include EMT induction, cell cycle control, angiogenesis, and regulation of stem cell populations are thought to promote *FOXC1*'s adverse oncological effects. However, despite such insights, the precise mechanisms behind *FOXC1*'s contributions to diverse malignancies remain incompletely defined.

Dysregulation of Hedgehog (Hh) signaling occurs frequently in sporadic and inherited cancers [7–10] and is estimated to underlie some 30% of malignancies. Hedgehog's critical role in malignancy is linked to the pathway's regulation of the Glioma (Gli) transcription factors and the requirement of Hh activity to maintain normal and cancer cell stemness. Notably, *FOXC1* and several of its paralogs are able to induce Hh expression and Hh pathway activity [11–15]. The output of the Hh pathway is tightly regulated by three equipotent inhibitors: the Patched transmembrane receptors, Suppressor of Fused (Sufu), and Protein Kinase A (PKA). Of these, PKA is positioned comparatively late in the signal transduction cascade and, by phosphorylating the GLI oncoproteins, directs their conversion from full length active to truncated repressor forms [8, 16, 17]. As a result, PKA controls the transcriptional outcome of Hh signaling. In recent years, a tissue-specific PKA antagonist has been identified. This Rho GTPase activating protein, Arhgap36, alters the balance between activator and repressor categories of GLI by inhibiting PKA, thereby inducing strong Hh pathway activation [18–20].

In this study, RNA- and ChIP-sequencing coupled with CRISPR interference were used to demonstrate that *Foxc1* binds a locus located in a region of closed chromatin to induce *Arhgap36* expression. In turn, *Arhgap36* depletes the levels of PKA and its catalytic subunit PKAC, both strongly activating Hh signaling and making signal transduction less dependent on regulation via *Smoothed*. The oncological relevance of the *Foxc1*-*Arhgap36* relationship is demonstrated in neuroblastoma, a neural crest derived pediatric malignancy where *ARHGAP36* levels predict five-year mortality. Consequently, our study identifies a mechanism through which overexpression of *FOXC1* dysregulates the Hedgehog pathway and contributes to malignancy.

Results

To discern pathways by which *Foxc1* may contribute to tumorigenesis, *Foxc1* was stably overexpressed in NIH3T3 fibroblasts. A retroviral construct was used that recapitulated the high levels of FOXC1 expression observed in a range of malignancies [15]. After RNA-sequencing, analysis using a stringent false discovery rate ($q \leq 0.01$; $\log_2$ fold change ≥ 1.0) defined a set of differentially expressed genes that were each distinguished by two independent bioinformatic workflows (DEseq2, and Cuffdiff; Table S1). The majority were up-regulated by *Foxc1* expression ($n=192$ genes; total 285; Fig. 1A). Gene-set enrichment analysis demonstrated significant enrichment for cancer invasiveness phenotypes, and malignancies, that importantly included neural crest associated malignancies such as melanoma and neuroblastoma (EMT $P = 4.8 \times 10^{-11}$; Malignancies $P = 10^{-3} - 10^{-4}$; Table S2). Also enriched were phenotypes either induced by *FOXC1* mutation, (glaucoma and abnormal intraocular pressure, $P \leq 4.7 \times 10^{-4}$) [21, 22] or present in *Foxc1*^{-/-} murine mutants (abnormal skeletal morphology and renal anomalies, $P = 0.05 - 10^{-3}$) [1, 23, 24]. Finally, genes relevant to *Foxc1*'s biological functions (angiogenesis, endochondral ossification, and hair follicle development) [13, 14, 25–30] were also enriched in relevant cell types (vasculature $P = 2.1 \times 10^{-5}$; neural crest $P = 3.0 \times 10^{-4}$; Table S2). Together, these results support the differentially expressed genes being representative of *Foxc1* functions in development and disease.

20% of the significantly dysregulated genes are implicated in Hh signalling (62 of 285) either as regulators or targets of the pathway (Table S1). Besides a cytokine that regulates inflammation and immunity (Interleukin Receptor 1 antagonist), *Arhgap36* mRNA was the most upregulated in the RNA-sequencing dataset (Fig. 1A). To test whether *Foxc1* consistently induces *Arhgap36* expression in mesenchymal-derived cells, qPCR measurements were performed and strong induction of *Arhgap36* mRNA observed after *Foxc1* overexpression in several relevant cell lines [C2C12 (murine myoblast) 10^5 -fold, ATDC5 (chondrogenic) 10^2 -fold, NIH3T3 10^3 -fold increase; Fig. 1B and Fig. S1]. Moreover, in a Hh signalling reporter cell line (NIH3T3-Gli2-mGFP) [15, 31], *Foxc1* overexpression induced *Arhgap36* protein expression at levels comparable to ectopically expressed Myc-FLAG tagged *Arhgap36* (Fig. 1C). Both *Foxc1* overexpression and ectopically expressed *Arhgap36* elevated protein expression of Gli1, a terminal effector and major readout of Hh pathway activity (Fig. 1C). Notably, the cellular localisation patterns of *Foxc1*-induced endogenous and ectopically expressed *Arhgap36* are similar (Fig. 1D, Fig. S2A). Collectively, these data demonstrate *Foxc1*'s ability to induce *Arhgap36* expression and activate Hedgehog signalling.

To characterize the functional interaction between the transcription factor *Foxc1* and *Arhgap36*, whole genome ChIP-sequencing (ChIP-seq) was performed on Gli2-mGFP NIH3T3 cells stably expressing *Foxc1*. Two independent *Foxc1* antibodies were used, together with an isotype-matched control immunoglobulin as control for *Foxc1* peak specificity. The number of uniquely mapped 60 bp reads (Input $> 5.3 \times 10^8$, *Foxc1*-ChIP 4.8×10^8) corresponds with deep sequencing coverage. Examination of chromatin accessibility demonstrates that, with the exception of embryonic stem cells, the *Arhgap36* locus is inactive (Fig. S3), an observation consistent with *Arhgap36*'s highly restricted tissue expression. Within a 200 kb interval encompassing the *Arhgap36* locus, five regions displayed strongly overlapping ChIP-seq peaks with both anti-*Foxc1* antibodies (Fig. 2A). Three (Prox-1 to Prox-3) were 0.2 - 2.5 kb from the transcription start sites, while two (Dist-1, Dist-2) were 55 and 71 kb upstream (Fig. 2A). Two amplicons were selected within each peak, and ChIP-qPCR demonstrated strongly increased *Foxc1* signal at each peak, relative to DNA precipitated using a non-specific normal IgG control (Fig. S4). Two groups of position weight matrices were significantly enriched in the bulk ChIP-seq peak data (STREME sequence motif discovery algorithm) (Fig. 2B, Fig. S4). The first comprised consensus sequences closely resembling known *Foxc1*-binding motifs ($P = 10^{-41} - 10^{-65}$), while the second corresponded closely




Figure 1.

Foxc1 induces expression of Arhgap36 and activates Hedgehog signalling.

(A) Volcano plot depicting in green differentially expressed genes with Foxc1 expression in NIH3T3 cells. (B) Confirmation of the robust Foxc1-induced increase in Arhgap36 mRNA in NIH3T3 cells by qPCR. (C) Foxc1 drives endogenous Arhgap36 protein expression, and that of Gli1, in NIH3T3-Gli2-mGFP cells to comparable levels of Myc-FLAG Arhgap36 ectopically expressed in parental NIH3T3 cells. (D) Immunofluorescence imaging demonstrates strong endogenous Arhgap36 expression, with the membrane staining in Foxc1-expressing cells recapitulating that of ectopically expressed Arhgap36-MF protein. [RNA-seq: n=3; quantitative Western blots: n=4 replicates; MF denotes Myc-FLAG tagged Arhgap36]

with that of Fos-Jun transcription factor dimers ($P = 10^{-13} - 10^{-20}$, **Fig. 2B**, **Fig. S5**). Each of the five ChIP-seq peaks contains the identified Foxc1 motifs, supporting Arhgap36 being a direct transcriptional target of Foxc1.

We next investigated in a Hh reporter line [15] if Foxc1 overexpression recapitulated Arhgap36's augmentation of Hh signaling, via down-regulation of PKA. As expected, ectopic expression of Foxc1 depleted the level of PKAC, PKA's catalytic subunit (63% reduction, $P = 1.4 \times 10^{-7}$; **Fig. 3A**). Foxc1 overexpression also reduced phosphorylation of threonine 197 in PKAC's activation loop, a residue essential to PKA enzymatic function and protein stability [32, 33]. This ~70% depletion of pT197 PKAC ($P = 1.3 \times 10^{-8}$) is analogous to the effect of ectopic Arhgap36 expression in parental NIH3T3 cells (**Fig. 3A**). Immunofluorescent staining demonstrated in cells with ectopic Foxc1 or Arhgap36 expression the near complete absence of PKAC and pT197 PKAC in the cytoplasm, especially at the basal body, where compartmentalized PKA-dependent control of Hh pathway occurs (**Fig. 3B**). Notably, in a transduced cell population with heterogeneous Foxc1 expression, PKAC signal is almost entirely depleted from cells with a high Foxc1 nuclear signal, demonstrating an inverse correlation between levels of PKAC and nuclear Foxc1 immunostaining (**Fig. S2B**). Similar results were observed in a second cell model, pre-adipocyte cells (3T3-L1) where expression of Foxc1 induced expression of Arhgap36 (albeit at lower levels) and a 40-50% reduction in protein levels of PKAC and pT197 PKAC (**Fig. S2C,D**). Direct overexpression of Arhgap36 led to strong downregulation of PKAC. These data, from two cell-based systems, demonstrate that Foxc1 expression depletes PKAC, the enzyme essential for promoting repressor forms of Gli, which regulate Hh target gene expression.

We next examined using CRISPR interference whether steric repression of the five Foxc1 binding loci affected Arhgap36 transcription. Two clones of NIH3T3-Gli2-mGFP cells that overexpress Foxc1 were created that each stably expressed an inactive form of Cas9 fused with the Krüppel-associated box (KRAB) repressor domain of ZIM3 protein. Recruitment of ZIM3 KRAB-dCas9 to each of the five Foxc1-binding regions by retroviral delivery of five separate pools each comprising 3-4 single guide RNAs, altered Arhgap36 expression, although to varying degrees (**Fig. 4A**, **Fig. S6**). The sgRNA pools targeting Prox-3 and Dist-2 reduced Arhgap36 mRNA expression by 98-99% and 67-76% respectively (**Figure 4A**). In both cases, reduced expression of Arhgap36 was associated with lower levels of Gli1 mRNA, and the degree of Arhgap36 mRNA reduction correlated with the decrease in Gli1 mRNA levels. Western immunoblot analysis confirmed that silencing either the Prox-3 or Dist-2 regions decreased Arhgap36 protein expression. This in turn correlated with an increase in PKAC and pT197 PKAC levels and a reduction in Gli1 protein levels (**Fig. 4B**). Since these data support Prox-3 and Dist-2 functioning, respectively, as promoter and distal enhancer for murine *Arhgap36*, evolutionary conservation of the Foxc1-binding loci was assessed. Multiple alignment across 60 vertebrates (Multiz) and measurements of evolutionary conservation (PhastCons) demonstrated that Prox-3 and Dist-2 are both conserved in placental, but not in marsupial mammals, nor in other vertebrates (**Fig. S7**). Multiple potential binding motifs for Foxc1 and Fos-Jun family transcription factors were identified in both regions; as illustrated by 16 Foxc1-binding motifs in both Prox-3 (8 motifs) and Dist-2 (8 motifs) of the murine locus, that were conserved or partially conserved in human (**Fig. S7**; Table S3). Taken together, the data presented support the Prox-3 and Dist-2 regions functioning as cis-regulatory elements for murine *Arhgap36*, and other placental mammals, including human.

Since PKA depletion de-represses Hh signaling, the pathway activation is predicted to become less dependent on Hh ligand, which pharmacologically would manifest as tumour resistance to inhibition of *Smoothened* (*Smo*). To test whether Foxc1 expression induces such ligand-independent Hh signal transduction, Gli1 levels were measured with and without two *Smoothened*-specific antagonists (Sonidegib and Cyclopamine). The results revealed that the Foxc1-induced increase of Gli1 mRNA and protein levels is resistant to inhibition by either antagonist (**Fig. 5A,B**). Furthermore, the Foxc1-induced resistance to Sonidegib is phenocopied by the ectopic expression of Arhgap36 (**Fig. S8A**). In both cases Gli1 upregulation, whether by Foxc1 or




Figure 2.

ChIP-seq identification of Foxc1-binding sites at the Arhgap36 locus.

(A) ChIP sequencing with independent anti-Foxc1 antibodies revealed substantial peak overlap for both ChIP samples, consistent with high antibody specificity. Peak calling identified five significant ChIP signal regions within ± 100 kb of *Arhgap36* [2 distal, 3 proximal; q -value ≤ 0.05]. (B) Within these Foxc1 ChIP peaks, the discovery algorithm STREME identified two major groups of significantly enriched motifs [$p = 2.8 \times 10^{-20}$, 2.5×10^{-65}]. The group 1 position weight matrices are highly similar to known Foxc1 motifs, while the group 2 PWMs very closely resemble the heptanucleotide recognition sequence bound by Fos-Jun transcription factor dimers [most prominently Fosl2]. The distribution of both PWM groups in the vicinity of Arhgap36 is shown on plot A.




Figure 3.

Foxc1-induced Arhgap36 reduces levels of protein kinase A catalytic subunit (PKAC).

(A) Foxc1 expression in Gli2-mGFP NIH3T3 cells strongly reduces PKAC, and catalytically active pT197 PKAC, to comparable levels to those observed with ectopic expression of Arhgap36-MF. Quantification shows > 2-fold reduction of PKAC/pT197 PKAC protein levels [Western blots: n=4 replicates]. (B) Immunofluorescent staining demonstrates equivalent reductions in PKAC/pT197 PKAC signal in the cytoplasm and at the basal bodies of cells expressing either Foxc1 or ectopic Arhgap36-MF. [Dashed box: 3x insets, basal body; white arrows].




Figure 4.

CRISPRi at potential Foxc1-binding sites diminishes Arhgap36 expression and Hedgehog activity.

(A) In independent CRISPRi-competent clonal cell lines, pools of guide RNAs targeting the Prox-3 and Dist-2 regions, robustly reduced Arhgap36 and Gli1 mRNA expression. (B) The reduced Arhgap36 and Gli1 protein expression with CRISPRi sgRNA targeting, implicates the same two ChIP-seq identified peaks in Foxc1's transcriptional control of the Arhgap36 locus.

by Arhgap36 overexpression, is not significantly reduced by co-treatment with Sonidegib. Indeed, Foxc1 overexpression leads to a similar level of Gli1 upregulation in NIH3T3 cells as treatment with a *Smoothed* agonist (SAG), and co-treatment of cells with SAG does not significantly enhance Gli1 levels beyond Foxc1-induced upregulation alone (**Fig. 5B**). Consequently, these data demonstrate that Foxc1 induces strong non-canonical activation of Hh signaling, a characteristic of multiple malignancies, including advanced and metastatic tumors.

PKA's roles regulating the Hh pathway extend beyond repressing Gli activity. PKA also phosphorylates key serine and threonine residues within Sufu. This inhibits ciliary trafficking of Sufu-Gli complexes, resulting in retention of Sufu at the tip of the primary cilium [34, 35]. We found that in addition to substantially increased axonemal tip accumulation of Gli2-mGFP (**Fig. 6A**, **Fig. S8B,C**), expression of Foxc1 also increases Sufu levels at the tip of the axoneme 3-fold ($P \leq 2 \times 10^{-10}$) while a similar 2-fold increase is induced by ectopic expression of Arhgap36 (**Fig. 6A-C**). The status of Sufu's PKA-dependent phosphorylation sites is known to influence Sufu activity. Two of these adjacent serine residues comprise a classical dual phosphorylation site for PKA (pS346) and GSK3 β (pS342) [34]. Consequently, the ~50% reduction in levels of Sufu pS342 after Foxc1 ($P = 6.4 \times 10^{-5}$) or Arhgap36 expression (**Fig. 6D**), provides further support for Hh pathway regulation by Foxc1. Together these data demonstrate that the effects of Foxc1 expression on Hh signal transduction involve several core regulators of Hh pathway activity.

The presented data support a model by which Foxc1-dependent Arhgap36 expression attenuates PKA activity and Sufu function. To verify that the observed changes in Sufu and Gli2 ciliary accumulation induced by Foxc1 overexpression are mediated by Arhgap36, shRNA silencing of Arhgap36 was performed in Foxc1-expressing Gli2-mGFP cells (**Fig. 7**). Two different Arhgap36-targeting shRNAs each substantially diminished ciliary Gli2-mGFP accumulation (shRNA1: 35%, shRNA2: 64% reduction; $P = 4.9 \times 10^{-7}$ and 4.5×10^{-10} ; **Fig. 7A-C**) while Gli2-mGFP levels in control cells that express very low levels of Arhgap36 were unchanged. The decrease in axonemal Gli2-mGFP correlated with the reduction in Gli1 mRNA (**Fig. 7D**), and the shRNA that more efficiently inhibited Arhgap36 induced a greater decrease in both Gli2-mGFP accumulation and Gli1 expression. The specificity of Arhgap36 inhibition is apparent from marked reductions in Gli1 and Arhgap36 mRNA expression, while levels of Foxc1 were unaltered (**Fig. 7D**). Analysis of Gli1 expression in parental Foxc1-expressing NIH3T3 cells treated with the same Arhgap36-targeting shRNA recapitulated the prior findings (**Fig. S9**). Taken together, these data demonstrate that the effects of Foxc1 expression on Hh pathway activity are Arhgap36-mediated.

To test the oncologic relevance of the Foxc1 - Arhgap36 relationship, *ARHGAP36* mRNA levels were first surveyed in cancer expression datasets. This revealed high expression in a common pediatric malignancy, neuroblastoma, as well as specific CNS, breast, lung, and neuroendocrine tumors. Due to neuroblastoma's neural crest origin and Foxc1's roles in this stem cell population, we focused on neuroblastoma data. Survival was evaluated in three independent neuroblastoma patient cohorts using Kaplan-Meier analyses after stratification into terciles of high, medium and low *ARHGAP36* mRNA expression. The highest tercile of *ARHGAP36* expression correlated with a favorable overall survival compared to the lowest: GSE49711, 91 vs 63% 5-year survival; E-MTAB-178191, 91 vs 61%; TARGET study, 60 vs 31%, (**Fig. 8A-C**) [36–38]. The effect was consistent across all three cohorts, despite differences in cancer staging - the first two cohorts primarily comprise patients with early disease, while TARGET mainly consists of advanced or metastatic neuroblastoma cases. Merging the data into a single dataset of 1348 patients narrowed the confidence intervals to provide more precise estimates. The highest tercile of *ARHGAP36* expression was associated with 87% five-year survival (range 86-92%) in contrast to 58% (53-63%) for the lowest tercile ($P = 1.7 \times 10^{-19}$). Illustrating these findings in the context of *MYCN*, whose amplification represents the primary genetic marker for high-risk and prognostically poor neuroblastoma, high *ARHGAP36* expression was associated with an 89% five-year survival compared to 37% with *MYCN* amplification (**Fig. S10**).




Figure 5.

Foxc1-induced Hh signalling has reduced dependence on *Smoothed*.

(A) Elevated levels of *Gli1* mRNA in Foxc1-expressing NIH3T3 cells are resistant to inhibition by the *Smoothed* antagonists sonidegib and cyclopamine. Wild-type NIH3T3 cells stimulated with *Smoothed* agonist (SAG) and treated with sonidegib, provide a control for inhibitor efficiency. (B) Resistance to *Smoothed* inhibition is supported by the elevated *Gli1* protein levels in Foxc1-expressing Gli2-mGFP NIH3T3 cells treated with sonidegib. Note that expression of Foxc1 induces comparable *Gli1* protein levels to vector control cells [pLXSH] treated with SAG; and that levels of Arhgap36 protein itself are unaffected by either sonidegib, or SAG treatment. [qPCRs, quantitative Western blots: n=4 replicates].




Figure 6.

Foxc1 promotes ciliary accumulation and decreases phosphorylation of SuFu.

(A) Representative immunofluorescence images demonstrate increased SuFu accumulation at axonemal tips of Foxc1-expressing cells, and NIH3T3 cells that express Myc-FLAG Arhgap36. Note, SAG stimulation *per se* does not substantially affect ciliary accumulation of SuFu. (B) Distribution of SuFu intensity in individual cilia (C) Mean ciliary SuFu intensity values. These demonstrate that the ciliary SuFu signal in cells expressing Foxc1, and separately Arhgap36, is substantially increased relative to empty vector controls [pLXSH and pLXSN; n = 9 combined experiments]. (D) Decreased phosphorylation of SuFu at the S342 residue in Gli2-mGFP NIH3T3s expressing Foxc1, and NIH3T3 cells expressing Myc-FLAG Arhgap36, relative to vector controls. Expression of Foxc1 also significantly impacts the total protein levels of SuFu, in contrast to the non-significant effect of Arhgap36 [quantitative Western blots: n=4 replicates].




Figure 7.

Arhgap36 is required for Foxc1-induced activation of Hh signalling.

(A) Immunofluorescence images demonstrate that two Arhgap36-targeting shRNAs each substantially reduce axonemal tip accumulation of Gli2 in cells that express Foxc1. (B, C) Individual and mean ciliary Gli2 intensity values demonstrate the decrease of ciliary Gli2 signal with Arhgap36 shRNA inhibition [$n = 3$ combined experiments]. (D) qPCR analyses demonstrate a strong decrease in Gli1 mRNA levels in Foxc1-expressing cells treated with Arhgap36 shRNAs, relative to pLKO.5 control. Note, concordant Gli1 and Arhgap36 expression levels across all conditions.




Figure 8.

Overall survival of neuroblastoma patients stratified by Arhgap36 expression levels.

(A-C) Kaplan-Meier plots show poor overall survival with low Arhgap36 expression across three independent neuroblastoma datasets. The significant reduced 5-year survival for cases with low levels of Arhgap36 expression is evident from the range of Hazard Ratios (2.7 – 8.0). Panel (D) shows the same analysis once the data are merged into a single dataset comprising n=1348 individuals (HR 2.8 – 4.8). [Cases stratified into terciles of Arhgap36 expression as shown in boxplots: first tercile (T1) “low”; second (T2) “medium”; third (T3) “high”].

Discussion

Here we show that the PKA antagonist *Arhgap36* is a central gene dysregulated by *Foxc1* overexpression, and that this linkage provides new perspectives in *Foxc1*'s contributions to malignancy. *Foxc1*'s transcriptional regulation of *Arhgap36*, and the consequent inhibition of PKA, alters the equilibrium between full length transcriptional GLI activators and truncated transcriptional repressors, weakening regulation and increasing activity of a pathway fundamental to development. It is therefore logical that a transcription factor identified to enhance *Arhgap36* expression is one that induces Hh activity in multiple tissues during development [13–15]. In some contexts, Hedgehog functions as a morphogen that confers positional information, with a gradient of activity specifying individual cell types and behaviors required for establishment of complex tissue patterns [39–41]. Increased gene dosage from *FOXC1* copy number variation causes phenotypes indicative of impaired Hedgehog signaling, including cerebellar hypoplasia and anomalous facial skeletal development [42–45]. Consequently, *Foxc1*'s ability to induce *Arhgap36* expression establishes a new mechanism by which the transcription factor may modulate Hh pathway activity. This has particular relevance for oncology, where dysregulated Hh expression makes major contributions and larger fold increases in *FOXC1* expression occur.

A second finding concerns the breadth of alteration to Hh signaling. *Foxc1*'s induction of *Arhgap36* depletes an evolutionary conserved negative regulator of Hh signaling (PKA) from the cytoplasm and base of the primary cilium. Such wide loss of PKA activity should release Gli transcription factors from phosphorylation-induced direct inhibition (in case of Gli2) and proteolytic processing (in case of Gli2 and Gli3). In parallel, *Foxc1*-induced depletion of PKAC predictably dysregulates the PKA substrate Sufu: a second major negative regulator of the Hh pathway, whose roles encompass development, disease, and notably tumour suppression. Sufu acts as a scaffold between PKA and Gli proteins, that facilitates their phosphorylation and proteolytic processing, while also directly inhibiting transcriptional activity of active Gli forms [35]. The reduced phosphorylation of Sufu and increased translocation of Sufu and Gli2 to the tip of the cilium represent alterations to fundamental features of Hh signal transduction. As *Arhgap36* acts independently of Smo [18], another consequence of *Foxc1*-driven *Arhgap36* expression observed in our experiments is acquired resistance of Hh pathway activity to Smo antagonists. Consequently, by altering activity of PKA, *Foxc1* overexpression impairs central elements of the Hh pathway resulting in increased transcription of Hh-target genes and, via augmented Smo-independent activity, tumor resistance to *Smoothed* antagonists. Collectively, such properties are consistent with the overexpression of *FOXC1* in multiple tumors being associated with more aggressive cancer phenotypes, drug resistance and metastasis.

Forkhead box transcription factors' ability to access areas of closed chromatin is mediated by a conserved winged-helix DNA binding domain that is structurally similar to histones H1 and H5 [46]. This capacity to directly bind DNA targets in condensed chromatin enables pioneer factors to activate enhancers and, by opening condensed chromatin, provide access to transcription factors that lack pioneering ability. The closely correlated ChIP-seq peaks identified in this study, which contain consensus sequences for *Foxc1* and Fos-Jun dimers, support coordinated transcription factor binding at the *Arhgap36* locus. The consistent co-localisation of these transcription factors' binding motifs, including in other placental mammals, accords with enrichment of FOSL2 and JUNB motifs at the *FOXC1* ChIP-seq peaks noted in a previous study [3]. Together, such data support a model in which the overexpression of *FOXC1*, by initiating local opening of chromatin [47], makes the *ARHAGP36* locus accessible to other transcription factors to induce gene expression. Such combinatorial control of *Arhgap36*'s transcriptional regulation is predicted to have heterogeneous effects on Hh pathway activity, varying in tissue and cell-specific manners and being determined by levels of *FOXC1* and FOS-JUN dimers.

FOXC1's induction of Hh signaling via ARHGAP36 and PKA inhibition is expected to modify processes in both development and disease. Since our data were derived with retroviral constructs, where 10-20 fold increases in Foxc1 protein levels reiterate overexpression in FOXC1-associated tumours [2], we evaluated the relevance of the Foxc1-Arhgap36 interaction in human malignancy. Analyses in neuroblastoma, a heterogeneous, life-threatening and common pediatric malignancy that originates from sympathoadrenal neural crest progenitors, revealed higher levels of ARHGAP36 expression were comparatively protective. The 2.8 to 8-fold greater five-year survival identified in three independent patient cohorts suggests ARHAGP36 may represent an informative prognostic factor. Because neuroblastoma treatment regimens frequently impact the development of surviving children [48], the ability to stratify by prognosis may facilitate treatment choice and minimize morbidity for a portion of patients. Although the mechanism(s) mediating this effect remain to be explored, this observation is consistent with the requirement of Hh activity for many tumours to proliferate [49, 50] and for maintenance of stem cell self-renewal, including that of cancer stem cells [51].

In summary, dysregulated expression of *FOXC1* is well-implicated in malignancy, with expanding evidence that elevated *FOXC1* levels are associated with increased mortality in breast and hepatic cancer and malignancies as diverse as AML and low grade glioma [4, 52–54]. In contrast with such progress, determining how FOXC1 promotes malignancy has proved more elusive. This work establishes that overexpression of FOXC1, at levels consistent with FOXC1-associated malignancies, transcriptionally activates Arhgap36 expression which in turn dysregulates multiple fundamental aspects of Hh pathway activity. The presented data further demonstrate that ARHGAP36 levels are predictive of five-year survival in a common pediatric malignancy and identify a biological characteristic that merits further investigation. Overall, this study provides additional mechanistic insight into FOXC1's capacity for stimulating malignancy.

Supplementary figures




Figure S1.

Foxc1 induces expression of *Arhgap36* and activates Hedgehog signalling.

(A,B) Stable overexpression of *Foxc1* in C2C12 and ATDC5 cells induces strong increases in *Arhgap36*, and *Gli1*, mRNA levels.




Figure S2.

Foxc1-driven Arhgap36 reduces levels of protein kinase A catalytic subunit (PKAC) in Gli2-mGFP NIH3T3 cells and 3T3-L1 preadipocytes.

(A) Immunofluorescent staining confirms substantial reduction of PKAC signal in Foxc1-expressing Gli2-mGFP NIH3T3 cells. Note correlation between positive Arhgap36 staining and PKAC loss in individual cells. (B) Immunofluorescent staining of Foxc1 and PKAC under the same conditions. Note nearly complete loss of PKAC signal in cells with high Foxc1 nuclear signal. (C) Foxc1 induced expression of endogenous Arhgap36 in 3T3-L1 cells is accompanied by 40-50% reduction in PKAC/pT197 PKAC protein levels. (D) Arhgap36-MycFLAG expression in 3T3-L1 cells strongly reduces PKAC, and catalytically active pT197 PKAC.




Figure S3.

The *ARHGAP36* locus is located in a region of predominantly closed chromatin.

(A) DNase I hypersensitivity data from the UCSC genome browser for the 100 kb regions surrounding human *ARHGAP36*, and *GLI1* and *PRKACA* (for comparison), from diverse human cell lines (ENCODE). DNase I HS is a marker for open chromatin that is commonly found at active cis-regulatory sequences including promoters and enhancers. At the *GLI1* and *PRKACA* loci, note the multiple, relatively uniform density signal peaks (appearing as dark vertical lines) for the majority of cell lines particularly immediately adjacent to transcription start sites (TSSs), illustrating “open chromatin” that is consistent with expression of both genes in a wide range of tissues. In contrast, strong DNase I HS signal at the *ARHGAP36* locus is observed in the subset of the 75 cell lines of developmental origin marked with a purple vertical bar [Embryonic stem cells (H1hESC, H7hESC) and induced pluripotent stem (iPS CWRU1, iPS NIH11, iPS NIH17) cell lines]. The signal density for these is highest adjacent to several *ARHGAP36* TSSs, corresponding to the multiple known transcripts, with weak signal levels observed for most of the other cell lines.




Figure S4.

ChIP-qPCR validation of Foxc1 ChIP-Seq peaks in the vicinity of Arhgap36 gene.

(A,B) Plots demonstrate read distribution for Foxc1 ChIP-Seq data from [Figure 2](#) [A, entire locus (within ± 100 kb of Arhgap36 ORFs); B, individual peaks (5 kb regions); location of amplicons (a-j) that were analysed by ChIP-qPCR is indicated in red]. (C) ChIP-qPCR data confirm substantial enrichment of signal in all proximal and distal peaks identified by ChIP-Seq ($N = 1$).




Figure S5.

***De novo* motifs detected in Foxc1 ChIP-Seq peaks using STREME closely resemble known binding motifs of Foxc1 and Fosl2.**

(A) *De novo* motif analysis conducted using STREME tool identified two major groups of significantly enriched motifs (PWMs groups 1 and 2). Group 1 PWMs (ranked top #1 for intersected peaks) share high degree of similarity with known Foxc1 motifs, while Group 2 PWMs (ranked top #2 for intersected peaks) are highly similar to motifs of Fos-Jun family TFs, most prominently Fosl2. Notably, when analysed separately, ChIP-Seq data obtained using each of two anti-Foxc1 antibodies produce highly similar *de novo* PWMs. Known motifs of Foxc1 and Fosl2 transcription factors were retrieved from JASPAR and HOCOMOCO databases for comparison.




Figure S6.

CRISPRi analysis of potential Foxc1-binding sites in the vicinity of Arhgap36 gene.

(A) Plot shows read distribution for the ChIP samples and input chromatin control with five identified peaks (Prox-1 to Prox-3, Dist-1 and Dist-2). Dashed line demonstrates potential relationship between Prox-3 and Dist-2, as promoter and enhancer regions for murine *Arhgap36*. Panel (B) shows 1) ChIP-Seq coverage at both affected peaks; 2) nearby location of TSS for a major *Arhgap36* isoform within the wide Prox-3 peak (approximately 80 – 110 bp downstream of narrow peak Prox-3a, and 500 bp upstream narrow peak Prox-3b); 3) location of individual sgRNA sites used for CRISPRi targeting.




Figure S7.

Conservation in Prox 3 and Dist 2 regions of Arhgap36 locus

(A,B) Representation of portions of the Arhgap36 locus corresponding to ChIP-seq peaks Prox-3 and Dist-2. Depicted in order are: ChIP-seq tracks with read coverage for two Foxc1 antibodies, UCSC-derived placental and non-placental mammal conservation data [Multiz alignment for representative vertebrate species, black lines], and distribution of predicted Fox and Fos-Jun transcription factor binding sites. Note, the heavy conservation in the Prox-3 region across all placental mammals, and in the Dist-2 region, that this is confined to the short interval detected by both Foxc1 antibodies. In addition, Prox-3 contains a cluster of 3 predicted binding sites for Foxc1, while Dist-2 includes a cluster of Foxc1 and Fos-Jun predicted binding motifs. [Placental mammal conservation by PhastCons (dark green “Placental Cons” track); placental mammal conserved elements (cherry “Placental EI” track)].




Figure S8.

Foxc1 expression phenocopies Arhgap36-induced resistance to *Smoothened* inhibition and facilitates ciliary accumulation of Gli2.

(A) Quantitative western blotting demonstrates that resistance to sonidegib inhibition observed in Foxc1-expressing Gli2-mGFP NIH3T3 cells recapitulates observations in NIH3T3 cells with ectopic expression of Myc-FLAG Arhgap36 [Arhgap36-MF]. (B,C) Accumulation of Gli2-mGFP at the tips of primary cilia in Gli2-mGFP NIH3T3 cells that express Foxc1, relative to relevant empty vector control [pLXSH]. Cells serum-starved for 20 h in media containing 0% FBS, 0.5% BSA, and next to stimulated with either vehicle or *Smoothened* agonist (SAG). Beeswarm/box plot **B** shows distribution of Gli2-mGFP intensity in individual cilia across all conditions tested [boxplot: box/whiskers – quartiles; black bar – median; black dot – mean; N = 9 combined experiments]. Mean ciliary Gli2-mGFP intensity values are provided in plot **C**.

Figure S9.

Foxc1-induced Gli1 expression is attributable to Arhgap36.

(A) qPCR analyses demonstrate substantial increase in Gli1 mRNA levels in NIH3T3 cells expressing Foxc1. This effect is reversed by shRNA-mediated knock-down of Arhgap36 expression.





Figure S10.

Overall survival of neuroblastoma patients stratified by Arhgap36 expression levels and MYCN amplification status.

(A) Analysis was performed on the GSE49711, E-MTAB-1781 and TARGET 2018 data merged into a single dataset comprising n=1348 individuals (Figure 8D). MYCN-amplified cases were segregated into a separate group. Kaplan-Meier plots show: a) best overall survival for patients with high Arhgap36 expression and without MYCN amplification, b) preserved lower overall survival rate among patients with low Arhgap36 expression and without MYCN amplification (HR 1.7 - 3.4), c) predictably poor survival among high-risk MYCN-amplified cases (HR 6.1 - 11.3 compared to Arhgap36 high reference).



Acknowledgements

The authors are grateful to the patients who have contributed to this study. The authors thank Dr. Fred Berry for numerous fruitful discussions that facilitated and helped guide this research. We also thank multiple colleagues for critically reviewing earlier versions of the manuscript. Funding was provided by the Canadian Institutes of Health Research (CIHR) (MOP-133658). Experiments were performed at the Cell Imaging Core (*RRID:SCR_019200* [↗](#)), which is supported by the University of Alberta Faculty of Medicine & Dentistry, University Hospital Foundation, Striving for Pandemic Preparedness – The Alberta Research Consortium, and Canada Foundation for Innovation.

Additional Information

Data Availability Statement

All data generated and/or analysed in this study are available from the corresponding author upon reasonable request. The RNA-sequencing and ChIP-sequencing data have been deposited to the NCBI GEO database with the following accession numbers: GSE297719 (RNA-seq), GSE297865 (ChIP-seq).

Funding

Canadian Institutes of Health Research (MOP-133658)

Additional files

Additional experimental methods. [↗](#)

File S1. List of differentially expressed genes in NIH3T3 cells expressing Foxc1 or pLXSH empty vector control. Complete list of differentially expressed genes identified in NIH3T3 cells expressing Foxc1 or empty vector control [pLXSH] by RNA-sequencing ($n = 3$) and used for volcano plot (Figure 1A). $n = 292$ differentially expressed genes; 195 upregulated, 97 downregulated. Selection criteria: q -value (FDR-adjusted p -value) ≤ 0.01 , absolute $\log_2$ ratio ≥ 1 , signal cut-off ≥ 1 FPKM in either condition]. [↗](#)

File S2. Functional profiling of the list of differentially expressed genes from File S1. Functional profiling of the list of genes differentially expressed in Foxc1-expressing NIH3T3 cells vs empty vector control [pLXSH] was performed using gene set enrichment analysis (GSEA) platform Enrichr [<https://maayanlab.cloud/Enrichr/>]. The list was analysed against multiple gene set libraries and databases. Results demonstrate significant associations between differentially expressed genes and molecular pathways / biological processes / pathological conditions with previous implication of Foxc1, as well as Foxc1-specific cell and tissue expression. Overall, these results indicate that ectopic expression of Foxc1 in NIH3T3 affects functionally relevant targets. Examples include association with ocular and vascular disease [glaucoma $p = 5.8 \times 10^{-4}$; intraocular pressure $p = 5.9 \times 10^{-3}$; hypertension $p = 5.7 \times 10^{-4}$; myocardial ischemia $p = 2.4 \times 10^{-4}$; coronary artery disease $p = 0.01$], vascular development [regulation of angiogenesis $p = 9.4 \times 10^{-5}$], bone development [endochondral ossification 2.8×10^{-4}], a number of mouse skeletal and renal phenotypes [MGI phenotypes]; multiple cancers and cancer invasiveness phenotypes [epithelial mesenchymal transition $p = 7.7 \times 10^{-11}$]. Relevant cell and tissue

expression signatures [mesenchyme $p = 3.4 \times 10^{-8}$; mesenchymal stem cells $p = 5.8 \times 10^{-4}$; (mesenchymal) stromal cells $p = 1.8 \times 10^{-6}$; vasculature $p = 2.7 \times 10^{-5}$; neural crest $p = 3.8 \times 10^{-4}$; several skeletal tissues; differentiating osteoblasts $p \leq 1.4 \times 10^{-6}$]. p -values are FDR-adjusted. [↗](#)

File S3. List of predicted Fox and Fos-Jun transcription factor binding sites in Prox 3 and Dist 2 regions of mouse *Arhgap36*. [↗](#)

References

1. Kume T., et al. (1998) **The forkhead/winged helix gene Mf1 is disrupted in the pleiotropic mouse mutation congenital hydrocephalus** *Cell* **93**:985–96 [Google Scholar](#)
2. Ray P.S., et al. (2010) **FOXC1 is a potential prognostic biomarker with functional significance in basal-like breast cancer** *Cancer Res* **70**:3870–6 [Google Scholar](#)
3. Ramachandran R., et al. (2024) **Conserved role of FOXC1 in TNBC is parallel to FOXA1 in ER+ breast cancer** *iScience* **27**:110500 [Google Scholar](#)
4. Somerville T.D., et al. (2015) **Frequent Derepression of the Mesenchymal Transcription Factor Gene FOXC1 in Acute Myeloid Leukemia** *Cancer Cell* **28**:329–42 [Google Scholar](#)
5. Simeoni F., et al. (2021) **Enhancer recruitment of transcription repressors RUNX1 and TLE3 by mis-expressed FOXC1 blocks differentiation in acute myeloid leukemia** *Cell Rep* **36**:109725 [Google Scholar](#)
6. Han B., et al. (2017) **FOXC1: an emerging marker and therapeutic target for cancer** *Oncogene* **36**:3957–3963 [Google Scholar](#)
7. Raleigh D.R., Reiter J.F. (2019) **Misactivation of Hedgehog signaling causes inherited and sporadic cancers** *J Clin Invest* **129**:465–475 [Google Scholar](#)
8. Zhang Y., Beachy P.A. (2023) **Cellular and molecular mechanisms of Hedgehog signalling** *Nat Rev Mol Cell Biol* **24**:668–687 [Google Scholar](#)
9. Cong G., et al. (2025) **Mechanisms and therapeutic potential of the hedgehog signaling pathway in cancer** *Cell Death Discov* **11**:40 [Google Scholar](#)
10. Johnson R.L., et al. (1996) **Human homolog of patched, a candidate gene for the basal cell nevus syndrome** *Science* **272**:1668–71 [Google Scholar](#)
11. Chang B.E., et al. (1997) **Axial (HNF3beta) and retinoic acid receptors are regulators of the zebrafish sonic hedgehog promoter** *EMBO J* **16**:3955–64 [Google Scholar](#)
12. Han B., et al. (2015) **FOXC1 Activates Smoothed-Independent Hedgehog Signaling in Basal-like Breast Cancer** *Cell Rep* **13**:1046–58 [Google Scholar](#)
13. Yoshida M., et al. (2015) **The transcription factor Foxc1 is necessary for Ihh-Gli2-regulated endochondral ossification** *Nat Commun* **6**:6653 [Google Scholar](#)
14. Almubarak A., et al. (2024) **FOXC1 and FOXC2 regulate growth plate chondrocyte maturation towards hypertrophy in the embryonic mouse limb skeleton** *Development* **151** [Google Scholar](#)

15. Havrylov S., et al. (2024) **Pleiotropy in FOXC1-attributable phenotypes involves altered ciliation and cilia-dependent signaling** *Sci Rep* **14**:20278 [Google Scholar](#)
16. Wang B., Fallon J.F., Beachy P.A. (2000) **Hedgehog-regulated processing of Gli3 produces an anterior/posterior repressor gradient in the developing vertebrate limb** *Cell* **100**:423–34 [Google Scholar](#)
17. Niewiadowski P., et al. (2014) **Gli protein activity is controlled by multisite phosphorylation in vertebrate Hedgehog signaling** *Cell Rep* **6**:168–181 [Google Scholar](#)
18. Rack P.G., et al. (2014) **Arhgap36-dependent activation of Gli transcription factors** *Proc Natl Acad Sci U S A* **111**:11061–6 [Google Scholar](#)
19. Eccles R.L., et al. (2016) **Bimodal antagonism of PKA signalling by ARHGAP36** *Nat Commun* **7**:12963 [Google Scholar](#)
20. Zhang B., et al. (2019) **Patched1-ArhGAP36-PKA-Inversin axis determines the ciliary translocation of Smoothened for Sonic Hedgehog pathway activation** *Proc Natl Acad Sci U S A* **116**:874–879 [Google Scholar](#)
21. Nishimura D.Y., et al. (1998) **The forkhead transcription factor gene FKHL7 is responsible for glaucoma phenotypes which map to 6p25** *Nat Genet* **19**:140–7 [Google Scholar](#)
22. Mears A.J., et al. (1998) **Mutations of the forkhead/winged-helix gene, FKHL7, in patients with Axenfeld-Rieger anomaly** *Am J Hum Genet* **63**:1316–28 [Google Scholar](#)
23. Hong H.K., Lass J.H., Chakravarti A. (1999) **Pleiotropic skeletal and ocular phenotypes of the mouse mutation congenital hydrocephalus (ch/Mf1) arise from a winged helix/forkhead transcriptionfactor gene** *Hum Mol Genet* **8**:625–37 [Google Scholar](#)
24. Kume T., Deng K., Hogan B.L. (2000) **Murine forkhead/winged helix genes Foxc1 (Mf1) and Foxc2 (Mfh1) are required for the early organogenesis of the kidney and urinary tract** *Development* **127**:1387–95 [Google Scholar](#)
25. Kume T., et al. (2001) **The murine winged helix transcription factors, Foxc1 and Foxc2, are both required for cardiovascular development and somitogenesis** *Genes Dev* **15**:2470–82 [Google Scholar](#)
26. Seo S., et al. (2012) **Forkhead box transcription factor FoxC1 preserves corneal transparency by regulating vascular growth** *Proc Natl Acad Sci U S A* **109**:2015–20 [Google Scholar](#)
27. Siegenthaler J.A., et al. (2013) **Foxc1 is required by pericytes during fetal brain angiogenesis** *Biol Open* **2**:647–59 [Google Scholar](#)
28. Lay K., Kume T., Fuchs E. (2016) **FOXC1 maintains the hair follicle stem cell niche and governs stem cell quiescence to preserve long-term tissue-regenerating potential** *Proc Natl Acad Sci U S A* **113**:E1506–15 [Google Scholar](#)
29. Bhakuni T., et al. (2024) **FOXC1 regulates endothelial CD98 (LAT1/4F2hc) expression in retinal angiogenesis and blood-retina barrier formation** *Nat Commun* **15**:4097 [Google Scholar](#)

30. Wang L., et al. (2016) **Foxc1 reinforces quiescence in self-renewing hair follicle stem cells** *Science* **351**:613–7 [Google Scholar](#)
31. Kim J., Kato M., Beachy P.A. (2009) **Gli2 trafficking links Hedgehog-dependent activation of Smoothened in the primary cilium to transcriptional activation in the nucleus** *Proc Natl Acad Sci U S A* **106**:21666–71 [Google Scholar](#)
32. Steinberg R.A., et al. (1993) **Autoactivation of catalytic (C alpha) subunit of cyclic AMP-dependent protein kinase by phosphorylation of threonine 197** *Mol Cell Biol* **13**:2332–41 [Google Scholar](#)
33. Cheng X., et al. (1998) **Phosphorylation and activation of cAMP-dependent protein kinase by phosphoinositide-dependent protein kinase** *Proc Natl Acad Sci U S A* **95**:9849–54 [Google Scholar](#)
34. Chen Y., et al. (2011) **Dual Phosphorylation of suppressor of fused (Sufu) by PKA and GSK3beta regulates its stability and localization in the primary cilium** *J Biol Chem* **286**:13502–11 [Google Scholar](#)
35. Tukachinsky H., Lopez L.V., Salic A. (2010) **A mechanism for vertebrate Hedgehog signaling: recruitment to cilia and dissociation of SuFu-Gli protein complexes** *J Cell Biol* **191**:415–28 [Google Scholar](#)
36. Consortium S.M.-I (2014) **A comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the Sequencing Quality Control Consortium** *Nat Biotechnol* **32**:903–14 [Google Scholar](#)
37. Oberthuer A., et al. (2015) **Revised risk estimation and treatment stratification of low- and intermediate-risk neuroblastoma patients by integrating clinical and molecular prognostic markers** *Clin Cancer Res* **21**:1904–15 [Google Scholar](#)
38. Ma X., et al. (2018) **Pan-cancer genome and transcriptome analyses of 1,699 paediatric leukaemias and solid tumours** *Nature* **555**:371–376 [Google Scholar](#)
39. Heemskerk J., DiNardo S. (1994) **Drosophila hedgehog acts as a morphogen in cellular patterning** *Cell* **76**:449–60 [Google Scholar](#)
40. Helms J., Thaller C., Eichele G. (1994) **Relationship between retinoic acid and sonic hedgehog, two polarizing signals in the chick wing bud** *Development* **120**:3267–74 [Google Scholar](#)
41. Marti E., et al. (1995) **Requirement of 19K form of Sonic hedgehog for induction of distinct ventral cell types in CNS explants** *Nature* **375**:322–5 [Google Scholar](#)
42. Aldinger K.A., et al. (2009) **FOXC1 is required for normal cerebellar development and is a major contributor to chromosome 6p25.3 Dandy-Walker malformation** *Nat Genet* **41**:1037–42 [Google Scholar](#)
43. French C.R., et al. (2014) **Mutation of FOXC1 and PITX2 induces cerebral small-vessel disease** *J Clin Invest* **124**:4877–81 [Google Scholar](#)
44. Haldipur P., et al. (2014) **Foxc1 dependent mesenchymal signalling drives embryonic cerebellar growth** *eLife* **3** [Google Scholar](#)

45. Haldipur P., et al. (2017) **Phenotypic outcomes in Mouse and Human Foxc1 dependent Dandy-Walker cerebellar malformation suggest shared mechanisms** *eLife* **6** [Google Scholar](#)
46. Cirillo L.A., et al. (1998) **Binding of the winged-helix transcription factor HNF3 to a linker histone site on the nucleosome** *EMBO J* **17**:244–54 [Google Scholar](#)
47. Xu P., et al. (2021) **Foxc1 establishes enhancer accessibility for craniofacial cartilage differentiation** *eLife* **10** [Google Scholar](#)
48. Sainero-Alcolado L., et al. (2024) **Defining neuroblastoma: From origin to precision medicine** *Neuro Oncol* **26**:2174–2192 [Google Scholar](#)
49. Dahmane N., et al. (1997) **Activation of the transcription factor Gli1 and the Sonic hedgehog signalling pathway in skin tumours** *Nature* **389**:876–81 [Google Scholar](#)
50. Duman-Scheel M., et al. (2002) **Hedgehog regulates cell growth and proliferation by inducing Cyclin D and Cyclin E** *Nature* **417**:299–304 [Google Scholar](#)
51. Beachy P.A., Karhadkar S.S., Berman D.M. (2004) **Tissue repair and stem cell renewal in carcinogenesis** *Nature* **432**:324–31 [Google Scholar](#)
52. Ray P.S., et al. (2011) **Basal-like breast cancer defined by FOXC1 expression offers superior prognostic value: a retrospective immunohistochemical study** *Ann Surg Oncol* **18**:3839–47 [Google Scholar](#)
53. Xia L., et al. (2013) **Overexpression of forkhead box C1 promotes tumor metastasis and indicates poor prognosis in hepatocellular carcinoma** *Hepatology* **57**:610–24 [Google Scholar](#)
54. Cao Q., et al. (2019) **FOXC1 silencing inhibits the epithelial-to-mesenchymal transition of glioma cells: Involvement of beta-catenin signaling** *Mol Med Rep* **19**:251–261 [Google Scholar](#)

Author information

Serhiy Havrylov

Department of Medical Genetics, University of Alberta, Edmonton, Canada, Department of Ophthalmology, University of Alberta, Edmonton, Canada

Armin M Gamper

Department of Oncology, Cross Cancer Institute, University of Alberta, Edmonton, Canada

Ordan J Lehmann

Department of Medical Genetics, University of Alberta, Edmonton, Canada, Department of Ophthalmology, University of Alberta, Edmonton, Canada

ORCID iD: [0000-0003-4141-7411](https://orcid.org/0000-0003-4141-7411)

For correspondence: olehmann@ualberta.ca

Editors

Reviewing Editor

Samuel Pleasure

University of California, San Francisco, San Francisco, United States of America

Senior Editor

Richard White

University of Oxford, Oxford, United Kingdom

Reviewer #1 (Public review):

This thoughtful and thorough mechanistic and functional study reports ARHGAP36 as a direct transcriptional target of FOXC1, which regulates Hedgehog signaling (SUFU, SMO, and GLI family transcription factors) through modulation of PKAC. Clinical outcome data from patients with neuroblastoma, one of the most common extracranial solid malignancies in children, demonstrate that ARHGAP36 expression is associated with improved survival. Although this study largely represents a robust and near-comprehensive set of focused investigations on a novel target of FOXC1 activity, several significant omissions undercut the generalizability of the findings reported.

(1) It is notable that the volcano plot in Figure 1a does not show evidence of canonical Hedgehog gene regulation, even though the subsequent studies in this paper clearly demonstrate that ARHGAP36 regulates Hedgehog signal transduction. Is this because canonical Hedgehog target genes (GLI1, PTCH1, SUFU) simply weren't labeled? Or is there a technical limitation that needs to be clarified? A note about Hedgehog target genes is made in conjunction with Table S1, but the justification or basis of defining these genes as Hedgehog targets is unclear. More broadly, it would be useful to see ontology analyses from these gene expression data to understand FOXC1 target genes more broadly. Ontology analyses are included in a supplementary table, but network visualizations would be much preferred.

(2) Likewise, the ChIP-seq data in Figure 2 are under-analyzed, focusing only on the ARHGAP36 locus and not more broadly on the FOXC1 gene expression program. This is a missed opportunity that should be remedied with unbiased analyses intersecting differentially expressed FOXC1 peaks with differentially expressed genes from RNA-sequencing data displayed in Figure 1.

(3) RNA-seq and ChIP-seq data strongly suggest that FOXC1 regulates ARHGAP36 expression, and the authors convincingly identify genomic segments at the ARHGAP36 locus where FOXC1 binds, but they do not test if FOXC1 specifically activates this locus through the creation of a luciferase or similar promoter reporter. Such a reagent and associated experiments would not only strengthen the primary argument of this investigation but could serve as a valuable resource for the community of scientists investigating FOXC1, ARHGAP36, the Hedgehog pathway, and related biological processes. CRISPRi targeting of the identified regions of the ARHGAP locus is a useful step in the right direction, but these experiments are not done in a way to demonstrate FOXC1 dependency.

(4) It would be useful to see individual fluorescence channels in association with images in Figure 3b.

(5) Perhaps the most significant limitation of this study is the omission of in vivo data, a shortcoming the authors partly mitigate through the incorporation of clinical outcome data from pediatric neuroblastoma patients in the context of ARHGAP36 expression. The authors also mention that high levels of ARHGAP36 expression were also detected in "specific CNS, breast, lung, and neuroendocrine tumors," but do not provide clinical outcome data for these

cohorts. Such analyses would be useful to understand the generalizability of their findings across different cancer types. More broadly, how were high, medium, and low levels of ARHGAP36 expression identified? "Terciles" are mentioned, but such an approach is not experimentally rigorous, and RPA or related approaches (nested rank statistics, etc) are recommended to find optimal cutpoints for ARHGAP36 expression in the context of neuroblastoma, "specific CNS, breast, lung, and neuroendocrine" tumor outcomes.

<https://doi.org/10.7554/eLife.108827.1.sa2>

Reviewer #2 (Public review):

FOXC1 is a transcription factor essential for the development of neural crest-derived tissues and has been identified as a key biomarker in various cancers. However, the molecular mechanisms underlying its function remain poorly understood. In this study, the authors used RNA-seq, ChIP-seq, and FOXC1-overexpressing cell models to show that FOXC1 directly activates transcription of ARHGAP36 by binding to specific cis-regulatory elements. Elevated expression of FOXC1 or ARHGAP36 was found to enhance Hedgehog (Hh) signaling and suppress PKA activity. Notably, overexpression of either gene also conferred resistance to Smoothened (SMO) inhibitors, indicating ligand-independent activation of Hh signaling. Analysis of public gene expression datasets further revealed that ARHGAP36 expression correlates with improved 5-year overall survival in neuroblastoma patients. Together, these findings uncover a novel FOXC1-ARHGAP36 regulatory axis that modulates Hh and PKA signaling, offering new insights into both normal development and cancer progression.

The main strengths of the study are:

- (1) Identification of a novel signaling pathway involving FOXC1 and ARHGAP36, which may play a critical role in both normal development and cancer biology.
- (2) Mechanistic investigation using RNA-seq, ChIP-seq, and functional assays to elucidate how FOXC1 regulates ARHGAP36 and how this axis modulates Hh signaling.
- (3) Clinical relevance demonstrated through analysis of neuroblastoma patient datasets, linking ARHGAP36 expression to improved 5-year overall survival.

The main weaknesses of the study are:

- (1) Lack of validation in neuroblastoma models - the study does not directly test its findings in neuroblastoma cell models, limiting translational relevance.
- (2) Incomplete mechanistic insight into PKA regulation - the study does not fully elucidate how FOXC1-ARHGAP36 regulates PKAC activity at the molecular level.
- (3) Insufficient discussion of clinical outcome data - while ARHGAP36 expression correlates with improved survival in neuroblastoma, the manuscript lacks a clear interpretation of this unexpected finding, especially given the known oncogenic roles of FOXC1, ARHGAP36, and Hh signaling.

<https://doi.org/10.7554/eLife.108827.1.sa1>

Reviewer #3 (Public review):

Summary:

The focus of the research is to understand how transcription factors with high expression in neural crest cell-derived cancers (e.g., neuroblastoma) and roles in neural crest cell

development function to promote malignancy. The focus is on the transcription factor FOXC1 and using murine cell culture, gain- and loss-of-function approaches, and ChIP profiling, among other techniques, to place PKC inhibitor ARHGAP36 mechanistically between FOXC1 and another pathway associated with malignancy, Sonic Hedgehog (SHH).

Strengths:

Major strengths are the mechanistic approaches to identify FOXC1 direct targets, definitively showing that FOXC1 transcriptional regulation of ARHGAP36 leads to dysregulation of SHH signaling downstream of ARHGAP36 inhibition of PKC. Starting from a screen of Foxc1 OE to get to ARHGAP36 and then using genetic and pharmacological manipulation to work through the mechanism is very well done. There is data that will be of use to others studying FOXC1 in mesenchymal cell types, in particular, the FOXC1 ChIP-seq.

Weaknesses:

Work is almost all performed in NIH3T3 or similar cells (mouse cells, not patient or mouse-derived cancer cells), so the link to neuroblastoma that forms the major motivation of the work is not clear. The authors look at ARHGAP36 levels in association with the neuroblastoma patient survival; however, the finding, though interesting and quite compelling, is misaligned with what the literature shows about FOXC1 and SHH, their high expression is associated with increased malignancy (also maybe worse outcomes?). Therefore, ARHGAP36 expression may be more complicated in a tumor cell or may be unrelated to FOXC1 or SHH, leaving one to wonder what the work in NIH3T3 cells, though well done, is telling us about the mechanisms of FOXC1 as an oncogene in neuroblastoma cells or in any type of cancer cell. Does it really function as an SHH activator to drive tumor growth? The 'oncogenic relevance' and 'contribution to malignancy' claimed in the last paragraph of the introduction are currently weakly supported by the data as presented. This could be improved by studying some of these mechanisms in patient-derived neuroblastoma cells with high FOXC1 expression. Does inhibiting FOXC1 change SHH and ARHGAP36 and have any effect on cell proliferation or migration? Alternatively, does OE of FOXC1 in NIH3T3 cells increase their migration or stimulate proliferation in some way, and is this dependent on ARHGAP36 or SHH? Application of their mechanistic approaches in cancer cells or looking for hallmarks of cancer phenotypes with FOXC1 OE (and dependent on SHH or ARHGAP36) could help to make a link with cellular phenotypes of malignant cells.

<https://doi.org/10.7554/eLife.108827.1.sa0>

Author Response:

Thank you for forwarding these helpful and thoughtful reviews - at a time when the review process in some journals can be a bit of a 'bloodsport', it is refreshing to receive such constructive and excellent comments. We essentially agree with the key points the reviewers have made, and as an interim response provide clarification of two areas:

1. As the reviewers highlighted, genome-wide analysis of ChIP-seq data from Foxc1 over-expression is indeed very worthwhile, and may offer insights for diverse malignancies where FOXC1 is over-expressed. We have a manuscript in preparation integrating this data set with ATAC-and RNA-seq data to identify genes transcriptionally regulated by elevated levels of Foxc1. In the interim, our full ChIP-seq data are available via the GEO accession number listed in the manuscript.

2. Analysis in neuroblastoma cell lines and then xenografts is equally important. Experiments manipulating ARHGAP36 levels in human neuroblastoma cell lines are underway, however a detailed mechanistic understanding of how ARHGAP36 influences neuroblastoma prognosis will take time, and lies beyond the scope of the current manuscript.

<https://doi.org/10.7554/eLife.108827.1.sa4>