

Lipopolysaccharide stimulates dynamic changes in B cell metabolism to promote proliferation

Reviewed Preprint

v1 • October 17, 2025

Not revised

Dana MS Cheung, Momchil Razsolkov, Fabrizia Bonacina, Stephen Andrews, Megan Sumoreeah, Linda V Sinclair, Andrew JM Howden, J Simon C Arthur 

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom • Department of Pharmacological and Biomolecular sciences “Rodolfo Paoletti”, University of Milan, Milan, Italy

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This work reveals metabolic pathways and molecular events mechanistically linked to B cell activation. Using an unbiased, comprehensive proteome profiling method and various functional validation approaches, this study generated **convincing** evidence suggesting a role for amino acid uptake, cholesterol accumulation, and protein prenylation in the proliferation, survival, and biogenesis of B cells stimulated with LPS and other activating stimuli. The significance of the findings is considered to be **fundamental**, in that they will advance our understanding of cell metabolism during B cell activation.

<https://doi.org/10.7554/eLife.109093.1.sa2>

Abstract

Naïve B cells exit quiescence and enter a proliferative state upon activation, ultimately differentiating into antibody-secreting or memory B cells. Toll-like receptor (TLR) ligands, such as lipopolysaccharide (LPS), can serve as physiological stimuli to initiate this transition. Using quantitative proteomics, we show that TLR4 engagement induces metabolic reprogramming in murine B cells, increasing the expression of amino acid transporters and cholesterol biosynthetic enzymes. The amino acid transporter SLC7A5 is markedly upregulated following LPS stimulation, and conditional deletion of *Slc7a5* impairs B cell proliferation, underscoring its essential role in B cell activation. LPS also elevates intracellular cholesterol levels, and inhibition of the rate-limiting enzyme HMG-CoA reductase blocks proliferation. This effect was mediated by a dual requirement for cholesterol metabolism and protein prenylation downstream of HMG-CoA reductase. Notably, this was not unique to TLR4 signalling but is also observed in B cells activated via TLR7, TLR9, CD40, or the B cell receptor. Together, these findings reveal that metabolic rewiring, including amino acid uptake and cholesterol metabolism, is an essential feature of B cell activation and proliferation.

Introduction

B cell activation can be initiated through either T cell-dependent (TD) or T cell-independent (TI) pathways, with TI activation further subdivided into a type I or type II response depending on the stimuli. During TD activation, following recognition of a protein antigen via the B cell receptor (BCR), the B cell can present peptides from the antigen to CD4 T cells, resulting in T cell activation. Additionally, the B cell receives co-stimulation through the binding of its CD40 receptor to the CD40 ligand found on T cells, which promotes differentiation, proliferation, somatic hypermutation, and class switch recombination (CSR) (Hoffman et al, 2016 [DOI](#)). Type I TI responses are activated by agonists such as lipopolysaccharide (LPS), which bind to toll-like receptors (TLRs) on B cells to induce a strong proliferative signal and increase antibody production. Longer-term exposure to LPS can also promote the differentiation of plasma cells (Genestier et al, 2007 [DOI](#)). Conversely, type II TI antigens consist of long repetitive structures, such as polysaccharides, which at high concentrations crosslink multiple BCRs to stimulate proliferation and antibody production (Obukhanych & Nussenzweig, 2006 [DOI](#)).

A well-established model to study TI B cell activation is to use LPS, a component of the cell wall of gram-negative bacteria, which stimulates TLR4 (Hoshino et al, 1999 [DOI](#); Poltorak et al, 1998 [DOI](#)) to induce robust B cell proliferation and antibody secretion, both *ex vivo* (Coutinho et al, 1974 [DOI](#); Dziarski, 1982 [DOI](#)) and *in vivo* (Johnson et al, 1956 [DOI](#)). Upon activation, TLR4 propagates downstream signalling through the adaptor proteins MyD88 and TRIF. This in turn activates pathways, including MAPK and NF- κ B signalling, to elicit the required physiological response (Arthur & Ley, 2013 [DOI](#); Kawai et al, 2024 [DOI](#)). The essential role of TLR4 in LPS-induced activation of B cells has been demonstrated by studies using TLR4-deficient mice. B cells from TLR4 or Myd88 knockout mice did not proliferate in response to LPS or lipid A stimulation *ex vivo* (Hoshino et al, 1999 [DOI](#); Kawai et al, 1999 [DOI](#)). Similarly, in studies where B cells from MyD88- and TLR4-knockout mice were transferred into μ MT mice (that lack mature B cells), a significant reduction in IgG1 production was observed (Pasare & Medzhitov, 2005 [DOI](#)).

Naïve B cells have been shown to be in a quiescent state with low metabolic demand (Waters et al, 2018 [DOI](#)). Conversely, B cell activation must require a significant increase in biomolecules such as proteins, lipids, and nucleic acids to support their rapid expansion and proliferation. The existing literature has focused on the role of glucose and glucose transporters in B cell function. Previous studies have demonstrated that activation using LPS or anti-IgM to stimulate the BCR increases glucose uptake and leads to an increased oxygen consumption rate (Dufort et al, 2007 [DOI](#); Jellusova et al, 2017 [DOI](#)). Whether this corresponds to changes in metabolic enzymes at a proteomic level, and whether TI B cell activation utilises other metabolic pathways, remains unknown. This increased glucose uptake has been shown to provide precursors for ribonucleotide and lipid synthesis (Waters et al, 2018 [DOI](#)). Similarly, the deletion of *Slc2a1* (which encodes for glucose transporter GLUT1) impaired germinal centre formation, plasma cell differentiation, and the production of high-affinity antibodies (Bierling et al, 2024 [DOI](#); Brookens et al, 2023 [DOI](#)). Recent reviews have expressed the need for studies investigating metabolic changes in activated B cells, especially the role of lipid metabolism, which is poorly understood (Johnstone et al, 2024 [DOI](#); Peeters & Jellusova, 2023 [DOI](#)).

T cell activation has been shown to stimulate global changes in metabolism at a proteomic level (Procaccini et al, 2016 [DOI](#); Howden et al, 2019 [DOI](#); Tan et al, 2017 [DOI](#)). However, there are no comparable proteomic studies that demonstrate how B cells support the change in metabolic requirements and energy demand from quiescence to activation upon LPS stimulation. Therefore, we used Data Independent Acquisition (DIA) based mass spectrometry to profile the B cell response to a combination of LPS + Interleukin 4 (IL-4) stimulation, a well-established *ex vivo* model of TI B cell activation. From this, we identified a substantial upregulation of proteins

involved in multiple metabolic pathways, including amino acid uptake and cholesterol biosynthesis. We demonstrate the requirement of L-amino acid transporter SLC7A5 for B cell survival and proliferation *ex vivo*. Similarly, we highlight the necessity of both cholesterol metabolism and protein prenylation downstream of the rate-limiting enzyme HMG-CoA reductase for B cell growth, survival, and proliferation. These results highlight the pronounced metabolic rewiring induced downstream of LPS + IL-4 stimulation.

Results

LPS + IL-4 stimulation of murine B cells promotes proliferation and changes the expression of enzymes involved in cellular metabolism

Previous studies have shown that *ex vivo* stimulation of murine B cells with LPS and IL-4 can drive B cell proliferation and class switching to IgG1 (Snapper et al, 1988 [\[1\]](#)). Consistent with this, we found that lymphocytes isolated from lymph nodes and subsequently stimulated with LPS + IL-4 resulted in the proliferation of CD19⁺ B cells, as judged by dilution of Cell Trace Violet (CTV) labelling. LPS + IL-4 stimulation led to extensive proliferation, with up to 6 generations of B cells present after 72 hours (**Figure 1A** [\[2\]](#)). Proliferation itself is thought to be a prerequisite for cells to undergo class-switching (Hodgkin et al, 1996 [\[3\]](#)). In line with this, only LPS + IL-4 stimulated B cells found in later generations were positive for IgG1 (**Figure 1A** [\[2\]](#)). On a similar note, cell size is considered a key regulator of the G1/S transition, as progression through the cell cycle can only continue once a certain threshold in size has been reached (Barberis et al, 2007 [\[4\]](#)). Accordingly, we found that 72 hours of LPS + IL-4 stimulation led to a marked increase in cell size and granularity as indicated by a higher forward/side scatter compared to naïve B cells (**Figure 1B-D** [\[2\]](#)). Stimulation with IL-4 alone resulted in a smaller increase in cell size than a combination of LPS + IL-4, but did not give rise to a strong proliferative response or class switching (**Figure 1A-D** [\[2\]](#)). Naïve B cells were used for comparison in ongoing experiments, as surviving unstimulated B cells exhibited a similar phenotype in the flow cytometry analysis; however, the lack of stimulation led to significant cell death from 24 hours onwards (**Figure 1A-D** [\[2\]](#), S1A-B).

These functional differences following activation suggest a significant change may be occurring at the protein level. To investigate this further, we analysed changes in the proteome after 24 hours of stimulation, as this time point was sufficient for B cells to increase in size but retain their synchrony in terms of the number of divisions they had undergone *ex vivo*. DIA-based mass spectrometry was used to determine proteomes of naïve B cells to activated B cells after 24 hours of LPS + IL-4 stimulation. The proteomic ruler method (Wiśniewski et al, 2014 [\[5\]](#)) was then used to estimate protein copy numbers and concentration. Using this approach, a total of 8030 protein species were identified, of which 7187 were quantified in both naïve and LPS + IL-4 stimulated B cells. Based on the mass spectrometry data, the predicted total protein mass in the B cells increased by approximately 2-fold following stimulation with a combination of LPS + IL-4 (**Figure 1E** [\[2\]](#)). This change in protein mass was due to an increased copy number of most proteins in the cell (**Figure 1F** [\[2\]](#)), with a median log₂ fold increase of 1.04. To adjust for the increase in cell size, the fold change in protein concentration was analysed, as this provides an indication of what proteins are selectively upregulated or downregulated as opposed to those that scale with cell size.

For the data using estimated cellular concentrations, using cutoffs of $q < 0.05$ and a log₂ fold change more than one standard deviation away from the median, 863 proteins were found to be upregulated following activation, while a further 301 were detected in all of the stimulated B cells but none of the naïve ones. 409 proteins were downregulated in the LPS + IL-4 stimulated B cells in comparison to naïve B cells, while a further 57 were detected in the naïve B cells, but none of the activated B cells (**Figure 1G** [\[2\]](#)).




Figure 1.

Stimulation with LPS + IL-4 promotes B cell proliferation and class switch recombination

(A-D) Cells from the lymph nodes of C57BL/6J mice were stained with Cell Trace Violet (CTV) and stimulated with IL-4 (10ng/ml) +/- LPS (20μg/ml) for 72 hours and analysed by flow cytometry. Data for naive B cells (stained on day 0) is also shown. Live CD19+ B cells were identified using the gating strategy described in Figure S7A. **(A)** Representative flow cytometry plots comparing IgG1 expression and CTV staining after 72 hours of IL-4 or LPS + IL-4 stimulation. **(B)** Representative plots for FSC and SSC at 72 hours of IL-4 or LPS + IL-4 stimulation with geometric means for the **(C)** forward scatter and **(D)** side scatter. Data shows the results of three biological replicates. Data was analysed by one-way ANOVA followed by multiple comparison testing via Sidak's analysis, for comparisons to naive B cells, where ****p<0.0001. Full ANOVA results are given in [Supplementary Table S4](#). **(E-H)** Cells from the lymph nodes of C57BL/6J mice, stimulated with LPS (20μg/ml) and IL-4 (10ng/ml) for 24 hours, and CD19+ B cells were isolated by FACS. Alternatively, naive CD19+ B cells were sorted directly from ex vivo lymph node cells. Cells were lysed and analysed by proteomics as described in the methods. Samples from four mice for LPS + IL-4 and three for naive were generated. **(E)** Total protein content (pg/cell) was estimated from the proteomic data. Statistical power was determined using an unpaired two-tailed t-test, where **p<0.01. **(F)** Volcano plot depicting changes in estimated protein copy number in naive vs LPS + IL-4 stimulated B cells. **(G)** Volcano plot showing the estimated cellular protein concentration (μM) in naive vs LPS + IL-4 stimulated B cells. Horizontal dashed lines indicate q < 0.05. Vertical dashed lines indicate log2 fold change of one standard deviation away from the median. **(H)** Enrichment analysis of the upregulated proteins in **(G)** against GO-term and KEGG databases.

To identify the biological pathways that were affected by LPS + IL-4 stimulation, proteins that were classified as regulated using the criteria above were searched against Gene Ontology (GO) Biological Processes terms and *Kyoto Encyclopaedia of Genes and Genomes* (KEGG) databases. Enrichment analysis on the downregulated proteins did not highlight any specific processes. However, upregulated proteins were associated with terms related to the cell cycle, ribosome biogenesis, protein translation, and steroid biosynthesis (**Figure 1H** [↗](#), **Table S1** [↗](#)).

While changes in the above processes would make sense in terms of cell proliferation, it was notable that changes in the glycolytic and oxidative phosphorylation pathways were not highlighted in this analysis. Previous studies have indicated that there is an increased demand for glucose in LPS-activated B cells (Dufort et al, 2007 [↗](#); Johnstone et al, 2024 [↗](#)). Isotope tracing experiments following CD40L and IL-4 activation to stimulate proliferation have suggested that this increased glucose uptake is used in the pentose phosphate pathway, potentially to provide NADPH and precursors for nucleotide biosynthesis, while alternative carbon sources may be used to drive an increase in the TCA cycle and oxidative phosphorylation (Waters et al, 2018 [↗](#)). The levels of enzymes involved in glucose metabolism were therefore examined in the proteomic data. In general, changes in the estimated concentrations of proteins involved in glycolysis, pentose phosphate pathway, TCA cycle, or oxidative phosphorylation were not observed (Figure S2A, B), although their copy numbers did increase in proportion to the increase in cell size (Figure S2C, D). B cells express the SLC2A1 (Glut1) glucose transporter, and deletion of this gene has been shown to reduce B cell activation *in vivo* (Bierling et al, 2024 [↗](#); Brookens et al, 2023 [↗](#)). There was an increase in the concentration of SLC2A1, which would be consistent with an elevated ability to take up glucose (Figure S2E). In addition, there was a 13-fold increase in hexokinase 2 (HK2), one of the enzymes involved in catalysing the 1st committed step for glucose to enter the glycolytic or pentose phosphate pathways (Tanner et al, 2018 [↗](#)) (Figure S2F) and has recently been shown to be required for maximal B cell responses to *ex vivo* stimulation with LPS (Paradoski et al, 2024 [↗](#)). To investigate if the increased copy number of Electron Transport Chain proteins correlated with changes in mitochondrial volume, B cells were stained with MitoTracker Red, a fluorescent dye that binds to mitochondria. We found that MitoTracker staining was significantly increased following 24 hours of LPS + IL-4 stimulation (Figure S2G, H).

LPS + IL-4 stimulation upregulates proteins involved in the cell cycle

Stimulation with LPS + IL-4 causes B cells to exit from a quiescent state and start to proliferate (**Figure 1** [↗](#)). This was confirmed using DAPI staining, which demonstrated that while naïve B cells were predominantly in a G0/1 state, after 24 hours of LPS + IL-4 stimulation, the proportion of B cells in the S and G2 stages of the cell cycle were increased (**Figure 2A** [↗](#), B). Progression through G1 is associated with an increase in Cyclin D expression, and activation of CDK4 and 6 (Rubin et al, 2020 [↗](#)). At the same time, inhibitory cell cycle proteins such as p27Kip1 (Cdkn1b) are downregulated (Wagner et al, 1998 [↗](#)). Therefore, we analysed the proteomic data for evidence of these changes. In naïve B cells, Cyclin D was not detected, while levels of CDK4 and 6 were low. The levels of Cyclin D, CDK4, and CDK6 were all increased following stimulation (**Figure 2C-E** [↗](#)). Naïve B cells also expressed high levels of p27Kip1, and this was reduced by LPS + IL-4 stimulation (**Figure 2F** [↗](#)).

Unphosphorylated RB1 binds to the E2F1/2/3 transcription factor, preventing the transcription of genes required for cell cycle progression. Increased CDK4/6 activity initiates RB1 phosphorylation, reducing its interaction with E2F, and resulting in the induction of E2F target genes, allowing progression through G1 to the S phase of the cell cycle (Engeland, 2022 [↗](#); Rubin et al, 2020 [↗](#)). Increased phosphorylation of RB1 was observed in B cells following LPS + IL-4 stimulation (**Figure 2G** [↗](#), H). In line with the increased RB1 phosphorylation, the levels of protein expressed from known E2F target genes, such as *Mcm2-7*, *Pcna*, *Dhfr*, *Rrm1/2*, and *Pola3* (Chicas et al, 2010 [↗](#); Engeland, 2022 [↗](#); Leone et al, 1998 [↗](#); Wagner et al, 1998 [↗](#)) were increased following LPS + IL-4




Figure 2.

Proteins involved in cell cycle progression are upregulated in LPS + IL-4 activated B cells

(A-B) B cells were purified from the spleens of C57BL/6J mice and either fixed on isolation (naive) or stimulated with LPS (20 μ g/ml) and IL-4 (10ng/ml) for 24 hours before fixation. Cells were then stained with DAPI and CD19. The cell cycle stage were analysed using the gating strategy shown in Figure S7B. **(A)** Representative histograms showing the proportion of B cells in different phases of the cell cycle. **(B)** Quantification shows three technical replicates from cells isolated from one mouse and is representative of three independent experiments. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Sidak's analysis, where $p < 0.01$ is indicated by **, $p < 0.001$ by *** and $p < 0.0001$ by **** for comparisons between the naive and LPS + IL-4 conditions. **(C-F)** Graphs depicting changes in the estimated cellular concentration (nM) of proteins implicated in entry into the cell cycle, determined from the proteomic dataset described in **Figure 1**. **(C)** Cyclin D, **(D)** CDK4, **(E)** CDK6, **(F)** p27Kip1. **(G-H)** B cells were purified from the spleens of C57BL/6J mice and stimulated with LPS (20 μ g/ml) and IL-4 (10ng/ml) for 24 hours before fixing and staining for phospho-retinoblastoma (p-Rb). Gating strategy for p-Rb staining is shown in Figure S7B. **(G)** Representative histogram comparing p-Rb staining in naive and LPS + IL-4 stimulated B cells. **(H)** Quantification shows three technical replicates from cells isolated from one mouse and is representative of three independent experiments. **(I)** Heat map showing the expression of proteins encoded by E2F target genes, derived from the proteomic data. **(J-M)** Graphs depicting changes in the estimated cellular concentration (nM) of proteins implicated in cell cycle progression, determined from the proteomic dataset described in **Figure 1**. **(J)** Cyclin A, **(K)** Cyclin B, **(L)** CDK2, **(M)** CDK1. p values were determined using an unpaired two-tailed Student's t-test for (H) or represent adjusted p values from the FDR calculations applied to the proteomic dataset (**D, E, F, L**), where $p < 0.01$ is indicated by **, $p < 0.001$ by *** and $p < 0.0001$ by ****.

stimulation (**Figure 2I**). Further progression through the cell cycle requires Cyclins E, A, and B, as well as CDK2 and CDK1. While Cyclin E was not detected in the proteomic data, the levels of Cyclin A, Cyclin B, CDK2, and CDK1 were all increased by LPS + IL-4 stimulation (**Figure 2J-M**).

Protein synthesis in LPS + IL-4 stimulated B cells is dependent on the uptake of amino acids

The cell growth and proliferation promoted by LPS + IL-4 stimulation would be expected to require an increased rate of protein synthesis. This is consistent with the proteomic analysis, which demonstrated an increase in total protein content within activated B cells (**Figure 1E**) and an enrichment of proteins associated with protein translation, including proteins that regulate ribosome biogenesis and the ribosomes themselves (**Figure 1H**). The ribosome is composed of 40S and 60S subunits, each of which is composed of multiple proteins (Doudna & Rath, 2002). LPS + IL-4 stimulation increased the total concentration of ribosomal proteins that make up both the 40S and 60S subunits (**Figure 3A, B**). This overall increase was due to higher expression of most of the individual proteins that make up the subunits, rather than a large increase in one or two of the proteins (**Figure 3C, D**). In line with the increase in ribosomal proteins, there was also an upregulation of proteins associated with ribosome biogenesis (**Figure 3E-G**). To determine if this correlated with an increased rate of protein synthesis, we measured the incorporation of the puromycin analogue o-propargyl-puromycin (OPP) into newly synthesised proteins. Consistent with the upregulation of ribosomal proteins, OPP incorporation was significantly higher in B cells stimulated with LPS + IL-4 for 24 hours compared to naive B cells (**Figure 3H-I**).

Elevated protein synthesis during rapid proliferation would require an increased supply of amino acids. B cells have also been reported to increase amino acid biosynthesis upon stimulation. ssRNAseq has shown that cycling B cells express higher levels of mRNA for enzymes in the serine biosynthetic pathway (D'Avola et al, 2022). Furthermore, the level of these proteins, as well as rates of serine biosynthesis, were found to be increased by stimuli such as anti-IgM, CD40L, and CpG, which are known to induce B cell proliferation (D'Avola et al, 2022). While this was not further addressed in the current study, the proteomic data reported here did show that serine biosynthetic enzymes, including the rate-limiting enzyme PHDGH, were increased in response to LPS + IL-4 stimulation (Figure S3).

Increased synthesis alone cannot satisfy the increased need for amino acids in proliferating cells, as the cell is unable to make essential amino acids. Several plasma membrane amino acid transporters were detected in the proteomic analysis, and these were all found to be upregulated in response to stimulation with LPS + IL-4 (**Figure 4A**). To look at the requirement for increased amino acid uptake during B cell proliferation more closely, we focused on Lat1, a member of the System L transporters. Lat1 is a heterodimer composed of an amino acid transporting light chain subunit, SLC7A5, and a chaperone heavy chain subunit, SLC3A2 (CD98) (Napolitano et al, 2015). Both of these subunits were upregulated in the proteomic data following LPS + IL-4 stimulation (**Figure 4B, C**). Additionally, the upregulation of SLC3A2 was confirmed by flow cytometry (**Figure 4D, E**).

Slc7a5^{fl/fl}/Vav-iCre^{+/-} mice, which have a deletion of SLC7A5 in all hematopoietic cells, have a comparable number of naive B cells to wild-type (WT) littermate controls, indicating that SLC7A5 is not critical for initial B cell development (Sinclair et al, 2013). To examine how the loss of SLC7A5 affects B cell proliferation in response to LPS + IL-4 stimulation, B cells were purified from WT and *Slc7a5*^{fl/fl}/Vav-iCre^{+/-} mice, labelled with CTV, and stimulated for 72 hours with LPS + IL-4 *ex vivo*. The percentage of viable B cells after 72 hours was much lower in *Slc7a5*^{fl/fl}/Vav-iCre^{+/-} mice compared to WT B cells (**Figure 4F**). Similarly, while WT B cells proliferated, the majority of the *Slc7a5*^{fl/fl}/Vav-iCre^{+/-} B cells failed to divide, and the number of live B cells after 72 hours was much lower compared to WT B cells (**Figure 4G, H**). In line with the failure to proliferate,




Figure 3.

LPS + IL-4 stimulation promotes protein synthesis

(A, B) Graphs show changes in cellular concentration (nM) of the sum of proteins that make up the large (A) and small (B) ribosomal subunits with heat maps showing the expression of the individual proteins making up the large (C) and small (D) subunits. (E-G) The proteomic dataset was mined for proteins involved in the biogenesis of the large and small ribosomal subunits based on the GO terms: GO:0000027, GO:0042273, GO:0000028, GO:0042274. (E) shows the sum of the proteins involved in the biogenesis of the large subunit and (F) shows the small subunit, with individual proteins represented on the volcano plot (G). Horizontal dashed lines on (G) indicate $q < 0.05$ while vertical dashed lines indicate $\log_2$ fold change more than one standard deviation away from the median. (H-I) Splenocytes from C57BL/6 mice were stimulated with LPS (20 $\mu\text{g}/\text{ml}$) and IL-4 (10 ng/ml) for 24 hours before fixing and staining for the uptake of puromycin analog O-propargyl-puromycin (OPP) to measure protein synthesis. Gating strategy for OPP staining is shown in Figure S7C. (H) Representative histogram comparing OPP uptake between naïve and LPS + IL-4 stimulated B cells. (I) Quantification shows three technical replicates from cells isolated from one mouse. Statistical power was determined using an unpaired two-tailed Student's t-test, where $p < 0.001$ is indicated by *** and $p < 0.0001$ by ****.




Figure 4.

Amino acid transporter SLC7A5 is required for key B cell functions

(A) Heat map showing the expression of genes encoding for proteins involved in plasma membrane amino acid transport determined from the proteomic dataset described in Figure 1. (B, C) Graphs depicting changes in cellular concentration (nM) of (B) SLC7A5 and (C) SLC3A2 derived from the proteomic data. p(adj)<0.0001, based on the proteomic FDR calculations, is indicated by ****. (D-E) Lymph node cells from C57BL6/J mice were stimulated with LPS (20µg/ml) and IL-4 (10ng/ml) for 24 hours before staining for CD98. (D) Representative FACS plot comparing CD98 expression. (E) Quantification shows three technical replicates from cells isolated from one mouse and is representative of two independent experiments. Statistical power was determined using one-way ANOVA followed by multiple comparison testing via Dunnett's analysis, for comparison to naïve B cells, where ***p<0.001 and ****p<0.0001. (F-I) B cells were purified from the spleens of wild type (WT) and SLC7A5 KO mice and stained with CTV before stimulation with LPS (20µg/ml) and IL-4 (10ng/ml) for 72 hours. (F) Percentage of live B cells (7AAD-ve) in WT and SLC7A5 KO mice. (G) Histogram representing CTV staining of WT and SLC7A5 KO B cells. (H) Live B cell number of WT and SLC7A5 KO mice. (I) Percentage of B cells that are IgG1+ve in WT and SLC7A5 KO mice. Data shows the results of four biological replicates per genotype. p<0.0001 is indicated by **** (unpaired two-tailed Student's t-test). (J-K) B cells were purified from the spleens of WT and SLC7A5 KO mice and stimulated with LPS (20µg/ml) +/- IL-4 (10ng/ml) for 24 hours before fixing and staining for the uptake of kynurenine to measure amino acid uptake. Gating strategy for kynurenine uptake in Figure S7D. (J) Quantification of kynurenine MFI between B cells from WT and SLC7A5 KO mice with (+) or without (-) LPS + IL-4, kynurenine or aminobicyclo-(2,2,1)-heptane-2-carboxylic acid (BCH). (K) Quantification of kynurenine MFI between B cells from WT and SLC7A5 KO mice with (+) or without (-) LPS, kynurenine or BCH. Data shows the results of four biological replicates per genotype. Statistical power was determined for using two-way ANOVA followed by multiple comparison testing via Sidak's analysis, where **** indicates p<0.0001 for comparisons between genotypes.

there was no evidence of class switching in the remaining *Slc7a5* fl/fl/Vav-iCre^{+/−} B cells (**Figure 4I** [↗](#)). SLC7A5 transports large neutral amino acids, including phenylalanine, histidine, methionine, tryptophan, and isoleucine, as well as the tryptophan metabolite kynurenine (Fotiadis et al, 2013 [↗](#)). As kynurenine is fluorescent and its uptake can be measured by flow cytometry, the effect of LPS + IL-4 treatment on kynurenine uptake in B cells was determined. In line with the increased expression of SLC7A5 in LPS + IL-4 treated B cells, kynurenine uptake was also higher in B cells stimulated with LPS + IL-4 for 24 hours compared to naive B cells (**Figure 4J** [↗](#)). Kynurenine uptake in the LPS + IL-4 stimulated B cells was reduced by the competitive substrate 2-aminobicyclo-(2,2,1)-heptane-2-carboxylic acid (BCH), which inhibits uptake through System L transporters. Kynurenine uptake in *Slc7a5* fl/fl/Vav-iCre^{+/−} B cells was similar to the uptake in BCH-treated WT cells, suggesting that SLC7A5 was the major system L transporter in LPS + IL-4 stimulated B cells (**Figure 4J** [↗](#)). BCH or SLC7A5 knockout did not reduce kynurenine uptake to baseline levels, indicating that other transport mechanisms for kynurenine uptake exist in B cells. Similar results were obtained for B cells stimulated with LPS alone, suggesting that LPS rather than IL-4 is the main driver of kynurenine uptake (**Figure 4K** [↗](#)).

Cholesterol metabolism is upregulated in activated B cells

The enrichment analysis (**Figure 1H** [↗](#)) suggested that sterol biosynthesis may be upregulated by LPS + IL-4 stimulation. Sterol biosynthesis allows the generation of cholesterol from acetyl-CoA within the cell via a complex multistep process (Figure S4). The 1st half of this pathway, often referred to as the mevalonate pathway, results in the production of farnesyl pyrophosphate (Buhaescu & Izzedine, 2007 [↗](#)). This can be used for several purposes in the cell, including the synthesis of cholesterol via the Bloch or Kandutsch-Russell pathways (Mitsche et al, 2015 [↗](#)), prompting us to investigate this pathway further. Analysis of proteomic data showed that the majority of enzymes involved in cholesterol biosynthesis were increased in LPS + IL-4 stimulated B cells relative to naive B cells (**Figure 5A** [↗](#)). Cholesterol biosynthesis is controlled by 2 rate-limiting steps: the conversion of HMG-CoA to mevalonic acid, which is catalysed by HMG-CoA-reductase (HMGCR), and the conversion of squalene to 2,3(S)-oxidosqualene, catalysed by squalene monooxygenase (SQLE) (Buhaescu & Izzedine, 2007 [↗](#)). While neither of these enzymes were detected in the proteomic dataset for naive B cells, they were both consistently expressed after LPS + IL-4 stimulation (**Figure 5B** [↗](#), C). Cells can also obtain cholesterol via the uptake of low-density lipoprotein (LDL) through the low-density lipoprotein receptor (LDLR) (Luo et al, 2020 [↗](#)). Similar to HMGCR and SQLE, the LDLR was not found in naive B cells but was detected in B cells following LPS + IL-4 stimulation (**Figure 5D** [↗](#)).

Cholesterol homeostasis is regulated by the transcription factor SREBP2, which is found as an immature precursor protein embedded in the endoplasmic reticulum (ER) with chaperones SCAP and INSIG (Luo et al, 2020 [↗](#); Brown et al, 2018 [↗](#)). Low sterol levels are detected by the sterol-sensing domain of SCAP, leading to the dissociation and degradation of INSIG. Subsequently, the SCAP-SREBP2 complex is transported from the ER to the Golgi apparatus, where it interacts with site 1 and site 2 proteases, allowing cleavage of SREBP2. The mature form of SREBP2 translocates to the nucleus, resulting in the upregulation of genes involved in cholesterol metabolism, including *hmgcr*, *sqle*, and *ldlr* (Luo et al, 2020 [↗](#); Brown et al, 2018 [↗](#)). Low levels of SREBP2 expression were detectable in naive B cells and became upregulated following LPS + IL-4 stimulation (**Figure 5E** [↗](#)). Together, the above results suggest that LPS + IL-4 stimulation would drive an increase in cellular cholesterol.

To test this, B cells were stained with the fluorescent dye filipin, which binds to cellular cholesterol. LPS + IL-4 stimulation resulted in a gradual increase in cholesterol levels from 8 hours to 24 hours of stimulation (**Figure 5F** [↗](#)). As cells can obtain cholesterol via *de novo* synthesis or uptake of LDL from serum, we compared the effects of culturing B cells in normal serum or serum treated with fumed silica to remove LDL (Brovkovich et al, 2019 [↗](#)). Analysis of the serum indicated that this treatment successfully removed LDL and HDL but did not affect the levels of triglycerides (Table S2 [↗](#)). Culture in media free of LDL (CF media) did not prevent the increase in




Figure 5.

LPS + IL-4 stimulation upregulates cholesterol metabolism in B cells

(A-E) Analysis of the genes involved in cholesterol metabolism in the proteomic dataset. (A) Heat map of all the enzymes involved in cholesterol biosynthesis. Cellular concentration (nM) of (B) HMGCR, (C) SQLE, (D) LDLR, and (E) SREBP2. A $p(\text{adj})$ of <0.01 is indicated by **. (F) Splenocytes from C57BL6/J mice were plated in cholesterol-free media and stimulated with LPS (20 μ g/ml) and IL-4 (10ng/ml). The cells were fixed at the stated time points and stained with filipin. The gating strategy for filipin staining in Figure S8A. Data shows three technical replicates from cells isolated from one mouse. Statistical power was determined by one-way ANOVA followed by multiple comparison testing via Dunnett's analysis, where * indicates $p<0.05$ and **** indicates $p<0.0001$ for comparisons to naïve B cells. (G-H) Splenocytes from C57BL6/J mice were plated in cholesterol-free media and pre-treated with DMSO as a vehicle control or varying concentrations of Fluvastatin or NB-598 for 45 minutes before stimulation with LPS (20 μ g/ml) and IL-4 (10ng/ml). The cells were fixed after 24 hours and stained with filipin before acquisition. (G) Filipin staining of Fluvastatin titration. (H) Filipin staining of NB-598 titration. Data shows three technical replicates from cells isolated from one mouse and is representative of two independent experiments, each with one biological replicate. Statistical power was determined using a one-way ANOVA followed by multiple comparison testing via Dunnett's analysis, where *** $p<0.001$ and **** $p<0.0001$ for comparison to LPS + IL-4 stimulated B cells. (I) Splenocytes from C57BL6/J mice were plated in normal or cholesterol-free (CF) media and pre-treated with Fluvastatin (10 μ M) or NB-598 (10 μ M) before stimulation with LPS + IL-4. The cells were fixed after 24 hours and stained with filipin before acquisition. Data shows the results of three biological replicates. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Tukey's analysis, where **** indicates $p<0.0001$ for comparisons to the LPS + IL-4 condition.

filipin staining stimulated by LPS + IL-4 (**Figure 5G-I**). Inhibitors of cholesterol biosynthesis were titrated to determine what concentration was required to reduce filipin staining in B cells stimulated in CF media. The increase in cholesterol following LPS + IL-4 stimulation could be reduced by the HMGCR inhibitor Fluvastatin or by NB-598, an inhibitor of SQLE (**Figure 5G**, H). Cells remained viable after 24 hours of inhibitor treatment with up to 10 μ M of either inhibitor (Figure S5A-F). Furthermore, the ability of HMGCR inhibition to reduce cholesterol levels did not correlate with effects on cell size, as judged by FSC measurements by flow cytometry after 24 hours (Figure S5B, E). From this, a concentration of 10 μ M was chosen for both NB-598 and Fluvastatin as this was sufficient to reduce cholesterol as close to naïve B cell levels as possible, but did not cause significant cell death after 24 hours. Different classes of statins have been described to exert different pleiotropic effects *in vivo* (Greenwood et al, 2006; Oesterle et al, 2017). Depending on their solubility, statins can be categorised as hydrophilic or lipophilic. Lipophilic statins, such as Fluvastatin, passively diffuse into cells, whereas hydrophilic statins may enter the cell through active transport. In the case of B cells, the use of a more hydrophilic statin, Rosuvastatin, gave similar results to Fluvastatin (Supplementary Figure S5G-J).

The increase in filipin staining following LPS + IL-4 stimulation for 24 hours in normal media was reduced, but not abolished, by either NB-598 or Fluvastatin treatment (**Figure 5J**). The ability of NB-598 and Fluvastatin treatment to reduce LPS + IL-4 induced filipin staining was greater in CF media relative to normal media that contains cholesterol (**Figure 5J**). Together, these results suggest that B cells may obtain cholesterol via a combination of biosynthesis and uptake.

The mevalonate pathway is critical for the survival and proliferation of LPS + IL-4 stimulated B cells

As the above data indicated that B cells increased their cholesterol levels following LPS + IL-4 stimulation, we assessed the importance of this process for B cell proliferation *ex vivo*.

Stimulation with LPS + IL-4 for 48 hours in normal media promoted proliferation, and this was reduced by treatment with either Fluvastatin or NB-598 (as shown by CTV staining, **Figure 6A**). In line with this decreased proliferation, both Fluvastatin and NB-598 greatly reduced the number of B cells present at 48 hours compared to cells treated with LPS + IL-4 alone (**Figure 6B**). The remaining cells in the Fluvastatin or NB-598 treated conditions also showed significantly lower levels of survival (based on 7-AAD staining, **Figure 6C**) and failed to increase in size (as judged by forward scatter, **Figure 6C**). Similar results were obtained for B cells grown in cholesterol-free media (**Figure 6A-D**).

In addition to cholesterol biosynthesis, the mevalonate pathway is also required for protein prenylation, a process that involves the addition of a farnesyl or geranylgeranyl motif to cysteine residues near the C-terminus of target proteins (Figure S4A) (Wang & Casey, 2016). As SQLE acts downstream of this branch point, the effects of NB-598 would argue for an important role of cholesterol biosynthesis in LPS + IL-4 induced proliferation; however, they would not exclude a parallel role for protein prenylation. To determine if the inhibitory effects of Fluvastatin in stimulated B cells may be in part due to changes in prenylation, B cells were treated with a prenylation inhibitor, FGTI-2734 (Kazi et al, 2019), prior to stimulation with LPS + IL-4 for 48 hours. Treatment of cells with FGTI-2734 decreased B cell proliferation in response to LPS + IL-4 stimulation in normal media (**Figure 6E**), and in line with this, the number of live B cells also decreased (**Figure 6F**). Cell survival was slightly decreased, although not to the same level as seen with Fluvastatin treatment (compare **Figure 6G** to **6C**). FGTI-2734 did not have a major impact on cell size in normal media (**Figure 6H**). Similar effects were observed in CF media, with the exception that FGTI-2734 treated cells were smaller than those stimulated with LPS + IL-4 alone (**Figure 6H**).




Figure 6.

Blocking rate-limiting enzymes in the cholesterol biosynthesis pathway reduces B cell growth, survival and proliferation

(A-D) B cells were purified from the spleens of C57BL6/J mice and cultured in normal or cholesterol-free (CF) media. The cells were stained with CTV, then pre-treated with DMSO as a vehicle control or Fluvastatin (10 μ M) or NB-598 (10 μ M), where indicated, for 45 minutes prior to stimulation with LPS (20 μ g/ml) and IL-4 (10 μ g/ml) for 48 hours. Gating strategy for CTV staining is shown in Figure S8B. (A) shows representative CTV staining, (B) live B cell number, (C) percentage of live B cells (7AAD-ve) (D) forward scatter. Data shows the results of three biological replicates. (E-H) Same as (A-D) but with pretreatment using FGTI-2734 (10 μ M). (E) shows representative CTV staining, (F) live B cell number, (G) percentage of live B cells (7AAD-ve) after FGTI-2734 treatment and (H) forward scatter of B cells. Data shows the results of three biological replicates. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Tukey's analysis, $p < 0.01$ is indicated by **, $p < 0.001$ by *** and $p < 0.0001$ by **** for comparisons to the LPS + IL-4 condition. ns indicates $p > 0.05$. For all panels, cells in the absence of LPS + IL-4 were naïve B cells analysed on the day of isolation.

In theory, the effects of HMGCR inhibitors should be reduced if cells were supplemented with a metabolite downstream of HMGCR in the mevalonate pathway. To test this, the effects of supplementing growth medium with mevalonate, the product generated in the reaction catalysed by HMGCR, were tested. Initially, the ability of mevalonate to rescue the effects of Fluvastatin on filipin staining was examined. For this, a concentration of 2mM mevalonate was used, as initial experiments showed that higher concentrations resulted in cell death and reduced filipin staining (Figure S6). In cells treated for 48 hours with LPS + IL-4, the reduction in filipin caused by Fluvastatin treatment in normal media was reversible by the addition of mevalonate (**Figure 7A** [↗](#)). In contrast, mevalonate had less effect in cells stimulated for 24 hours with LPS + IL-4, although at this time point, the ability of Fluvastatin to reduce filipin staining was also less pronounced than at 48 hours (**Figure 7A** [↗](#)). Surprisingly, in CF media, mevalonate was unable to rescue the effects of Fluvastatin at either time point (**Figure 7B** [↗](#)).

The effect of Fluvastatin on cell survival was partially reversible by the addition of mevalonate in normal media, which restored cell survival to 80% (**Figure 7C** [↗](#)) and fully restored cell size relative to cells stimulated with LPS and IL-4 in the absence of Fluvastatin (**Figure 7D** [↗](#)). However, mevalonate supplementation was not sufficient to completely rescue proliferation, as CTV staining demonstrated that 2 generations of B cells were present rather than 4 generations after LPS + IL-4 stimulation alone (**Figure 7E** [↗](#)). The number of live B cells following mevalonate addition was significantly higher compared to the Fluvastatin only condition, but did not reach the same level as LPS + IL-4 stimulation alone (**Figure 7F** [↗](#)). Similar to what was seen for filipin staining, mevalonate was unable to rescue any of the inhibitory effects of Fluvastatin treatment in CF media (**Figure 7C-F** [↗](#)).

The role of prenylation was also tested by supplementing cells with geranylgeranyl pyrophosphate (GGPP), which can be used for prenylation but not cholesterol biosynthesis. Surprisingly, GGPP was more effective than mevalonate in rescuing the effect of Fluvastatin on LPS + IL-4 induced proliferation (**Figure 8A** [↗](#)), and in line with this, there was a greater number of live B cells in the GGPP treated condition relative to the mevalonate treated condition (**Figure 8B** [↗](#)). A combination of GGPP and mevalonate did not have a synergistic effect compared to GGPP alone (**Figure 8A** [↗](#), B). GGPP was also able to rescue the effects of Fluvastatin on both cell survival and cell size (**Figure 8C** [↗](#), D). Conversely, in CF media, the addition of GGPP was unable to rescue any of the inhibitory effects of Fluvastatin (**Figure 8E-H** [↗](#)).

Given that GGPP should rescue prenylation but not cholesterol metabolism, its effects on proliferation are unexpected if B cells need cholesterol to proliferate. We therefore tested if GGPP affected cholesterol levels in Fluvastatin treated B cells. As seen in previous experiments, Fluvastatin reduced the ability of LPS + IL-4 stimulation to increase cholesterol levels at both 24 hours and 48 hours of treatment. GGPP was able to partially rescue this at both time points (**Figure 8I** [↗](#)). This was in contrast to mevalonate, which could only be rescued at 48 hours (**Figure 7A** [↗](#)), an observation that may, in part, explain why GGPP was more effective at restoring proliferation in the presence of Fluvastatin than mevalonate. The effects of GGPP on cholesterol levels were, however, limited to normal media, as similar to mevalonate, GGPP did not rescue to effect of Fluvastatin on filipin staining in CF media (**Figure 8J** [↗](#)).

The mevalonate pathway is required for B cell proliferation in response to multiple agonists

While the above results indicate that cholesterol is required for B cell proliferation in response to LPS + IL-4 stimulation, they do not demonstrate whether the primary signal for this is LPS or IL-4. To test this, B cells were stimulated with either IL-4, LPS, or a combination of both, and changes in cholesterol were measured by filipin staining. Stimulation with either LPS or IL-4 for 24 hours was able to increase cholesterol levels, although LPS had a greater effect (**Figure 9A** [↗](#)).




Figure 7.

Mevalonate supplementation rescues the effect of Fluvastatin treatment in B cells

(A-B) Splenocytes from C57BL6/J mice were plated in normal or cholesterol-free (CF) media and pre-treated with HEPES as a vehicle control or mevalonate (2mM) for 1 hour prior to treatment with DMSO or Fluvastatin (10 μ M), where indicated, for 45 minutes before stimulation with LPS (20 μ g/ml) and IL-4 (10ng/ml). The cells were fixed after 24 hours or 48 hours of LPS + IL-4 stimulation and stained with filipin. **(A)** Filipin staining comparing B cells +/- Fluvastatin or mevalonate after 24 or 48 hours of LPS + IL-4 stimulation in normal media. **(B)** Filipin staining comparing B cells +/- Fluvastatin or mevalonate after 24 or 48 hours of LPS + IL-4 stimulation in cholesterol-free media. Data shows the results of three biological replicates. **(C-F)** B cells were purified from the spleens of C57BL6/J mice and cultured in normal or cholesterol-free media. The cells were stained with CTV, then pre-treated with HEPES as a vehicle control or mevalonate (2mM) for 1 hour. The cells were treated with DMSO or Fluvastatin (10 μ M), where indicated, for 45 minutes prior to stimulation with LPS (20 μ g/ml) and IL-4 (10 μ g/ml) for 48 hours. **(C)** Shows percentage of live B cells (7AAD^{-ve}), **(D)** forward scatter of B cells, **(E)** representative CTV staining, **(F)** live B cell number. Data shows the results of three biological replicates. Where shown, statistical power was determined using two-way ANOVA followed by multiple comparison testing via Sidak's analysis. For comparison to Fluvastatin treated B cells, $p < 0.05$ is indicated by *, < 0.01 by **, < 0.001 by ***, < 0.0001 by **** and > 0.05 by ns. For all panels, cells in the absence of LPS + IL-4 were naïve B cells analysed on the day of isolation.




Figure 8.

GGPP supplementation rescues the effect of Fluvastatin treatment in B cells

(A-H) B cells were purified from the spleens of C57BL6/J mice and cultured in normal (A-D) or cholesterol free (CF, E-H) media. The cells were stained with CTV, pre-treated with methanol:ammonium hydroxide solution (CH₃OH:NH₄OH) as a vehicle control or geranylgeranyl pyrophosphate (GGPP) (10μM) for 1 hour before treatment with Fluvastatin (10μM), where indicated. The cells were stimulated with LPS (20μg/ml) and IL-4 (10μg/ml) and cultured for 48 hours. (A) shows representative CTVstaining for B cells cultured in normal media, (B) live B cell number, (C) percentage of live B cells and (D) forward scatter of B cells. (E) shows representative CTV staining for B cells cultured in CF media, (F) live B cell number, (G) percentage of live B cells and (H) forward scatter of B cells. Data shows the results of three biological replicates. Statistical power was determined using one-way ANOVA followed by multiple comparison testing via Dunnett's analysis. For comparison to Fluvastatin treated B cells, p<0.001 is indicated by *** and p<0.0001 by ****. ns indicated p<0.05. For (B) and (F) this data was log-transformed then statistically analysed due to unequal variance. (I-J) Splenocytes were plated in normal or cholesterol-free media and pre-treated with methanol:ammonium hydroxide solution (CH₃OH:NH₄OH) as a vehicle control or geranylgeranyl pyrophosphate (GGPP) (10μM), where indicated for 1 hour before treatment with DMSO or Fluvastatin (10μM). The cells were then stimulated with LPS (20μg/ml) and IL-4 (10ng/ml). The cells were fixed after 24 hours and stained with filipin before acquisition. (I) Filipin staining comparing B cells +/- Fluvastatin or GGPP after 24 or 48 hours of LPS + IL-4 stimulation in normal media. (J) Filipin staining comparing B cells +/- Fluvastatin or GGPP after 24 or 48 hours of LPS + IL-4 stimulation in cholesterol-free media. Data shows the results of three biological replicates. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Sidak's analysis. For comparison to Fluvastatin treated B cells p<0.01 is indicated by **, <0.001 by *** and p<0.0001 by ****. ns indicated p<0.05. For all panels, cells in the absence of LPS + IL-4 were naïve B cells analysed on the day of isolation.




Figure 9.

Cholesterol is required for the growth and proliferation of B cells regardless of stimuli

(A) Splenocytes from C57BL6/J mice were plated in normal or cholesterol-free (CF) media, before stimulation with LPS (20 μ g/ml), IL-4 (10 μ g/ml) or a combination of LPS and IL-4. The cells were fixed after 24 hours and stained with filipin. Data shows the results of three biological replicates. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Tukey's analysis. For comparisons to naïve B cells, $p < 0.0001$ is indicated by ****. (B) Splenocytes from C57BL6/J mice were plated in normal media before stimulation with LPS (20 μ g/ml), Resiquimod (1 μ g/ml), ODN 1826 (1 μ g/ml), Anti-IgM (10 μ g/ml), or CD40L (500ng/ml). The cells were fixed after 24 hours and stained with filipin. (B) Filipin staining comparing cholesterol content between different stimuli. Data shows the results of three biological replicates. Statistical power was determined using one-way ANOVA followed by multiple comparison testing via Dunnett's analysis. For comparisons to naïve B cells, $p < 0.0001$ is indicated by ****. (C-E) B cells were purified from the spleens of C57BL6/J mice and cultured in normal media. The cells were stained with CTV, then pre-treated with DMSO as a vehicle control or Fluvastatin (10 μ M) for 45 minutes before stimulation with LPS (20 μ g/ml), Resiquimod (1 μ g/ml), ODN 1826 (1 μ g/ml), Anti-IgM (10 μ g/ml), or CD40L (500ng/ml) as indicated for 72 hours. (C) shows representative histogram for CTV staining, (D) the percentage of live B cells (7AAD-ve), (E) live B cell number. Data shows the results of three biological replicates. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Tukey's analysis, where $p < 0.0001$ is indicated. For (E) this data was log-transformed and then statistically analysed due to unequal variance. Statistical power was determined using two-way ANOVA followed by multiple comparison testing via Sidak's analysis, where **** $p < 0.0001$.

Given the effects of HMGCR inhibition on cholesterol levels following LPS stimulation, we also tested whether other stimuli known to activate B cell proliferation could increase cholesterol levels. Murine B cells also respond to TLR7 (Weeratna et al, 2005) and TLR9 (Arunkumar et al, 2013; Yi et al, 1998) agonists. Both the TLR7 agonist Resiquimod and the TLR9 agonist CpG increased cholesterol levels in B cells (**Figure 8B**). Naive B cells can also be activated *ex vivo* by ligation of their BCR with anti-IgM antibodies or by stimulation with CD40 ligand (CD40L). Similar to TLR agonists, both stimuli also increased cholesterol levels in B cells (**Figure 9B**). Resiquimod, CpG, anti-IgM, and CD40L were all able to stimulate proliferation, as evidenced by CTV labelling and live B cell number after 72 hours of treatment (**Figure 9D**, E). While none of these stimuli were as effective at inducing proliferation as LPS, in each case, treatment with Fluvastatin reduced survival and blocked proliferation of the B cells (**Figure 9C-E**).

Discussion

In this study, we have used proteomic analysis to highlight the switch in metabolic demand from naïve B cells in a resting state to LPS + IL-4 activated B cells, which significantly increase their energy and biosynthetic requirements to sustain growth, proliferation, and class switching.

Upon activation, B cells upregulated several amino acid transporters, including the system L transporters SLC7A5 and 6 (**Figure 4A**), which can import large neutral amino acids, including the essential amino acids histidine, isoleucine, leucine, methionine, phenylalanine, threonine, tryptophan, and valine (Fotiadis et al, 2013). Of these 2 transporters, SLC7A5 was expressed at a higher level and was responsible for the majority of amino acid uptake via system 1 transporters (**Figure 4I**). SLC7A5 forms a dimer with CD98 (SLC3A2) to form an active transporter. SLC7A5 is not, however, the only binding partner of CD98, as it can also bind to the SLC7A11 and SLC7A6 transporters or interact with integrins (Kahya et al, 2021). Previous work has shown that mice with B cell-specific deletions of CD98 presented normal development of B cells in the bone marrow, but significantly lower levels of class-switched IgG due to the inhibition of B cell proliferation and plasma cell formation following the activation of naive B cells in the periphery (Cantor et al, 2009). The defect in proliferation was attributed to defective integrin signalling rather than an impairment in amino acid transport, based on the deletion of the integrin or amino acid transporter regions in CD98. Knockout of SLC7A5 in either hematopoietic cells or B cells has been previously shown not to affect the number of splenic B cells in mice (Sinclair et al, 2013), although a separate study found a decrease in the number of peritoneal B1 cells (Sun et al, 2023). While not critical for initial B cell development, SLC7A5 was critical for the survival and expansion of B cells in response to *ex vivo* stimulation with LPS + IL-4 (**Figure 4F-I**). Of note, recent research in human B cells has shown that stimulation with CpG increases SLC7A5 expression, which in turn transports L-leucine into the cell and facilitates mTORC1 activation, which was required for IgG production and inflammatory cytokine secretion (Torigoe et al, 2019).

Similar observations have been made in T cells, where SLC7A5 is not required for initial T cell development but is required for the clonal expansion and effector functions of T cells following activation via the T cell receptor in the periphery (Sinclair et al, 2013). This suggests that B cells, similar to T cells, may rely on different amino acid transporters at different points in their development. The expression of SLC7A5 following T cell activation has been shown to require the transcription factor c-Myc (Marchingo et al, 2020). While the role of c-Myc was not directly addressed in this study, analysis of the proteomic data presented here does show that c-Myc was increased by LPS stimulation (**Table S3**). This is consistent with previous reports showing an upregulation of c-Myc in response to several B cell activating stimuli, including LPS, CD40, and the BCR (Luo et al, 2018; Dominguez-Sola et al, 2012; Calado et al, 2012; Tesi et al, 2019). While in T cells, c-Myc regulates metabolic changes, including the expression of SLC7A5 (Sinclair & Cantrell, 2025), this has not been studied to the same degree in B cells. Of note, transcriptional

profiling following LPS stimulation of B cells has suggested that SLCA5 mRNA induction in response to LPS can occur in the absence of c-Myc (Tesi et al, 2019 [DOI](#)), suggesting the regulation of SLC7A5 may not be identical between T and B cells.

As B cells expand and divide, they will require an increase in lipids to generate new membrane. Cholesterol is an important lipid component of the plasma membrane, which affects membrane fluidity (Luo et al, 2020 [DOI](#)) and has also been implicated in the regulation of several receptors, including the BCR (Bléry et al, 2006 [DOI](#)). The regulation of cholesterol levels in cells is therefore closely controlled at the level of both cholesterol biosynthesis and its uptake via the endocytosis of LDL particles. Low levels of cholesterol in the endoplasmic reticulum are detected by the transcription factor SREBP2 (Luo et al, 2020 [DOI](#); Brown et al, 2018 [DOI](#)). The importance of this system in B cells was recently demonstrated by the conditional knockout of SCAP in B cells, a key regulator of SREBP1 and 2. While this did not prevent the development of B cells or their levels at steady state in mice, it did prevent the activation and proliferation of B cells in response to antigens and thus compromised the humoral immune response. SCAP deletion will, however, affect both SREBP1 and 2 function and therefore have wider effects on lipid metabolism beyond its effects on cholesterol. We show here that in response to several activating stimuli *ex vivo*, including LPS, B cells upregulate cholesterol, and that blocking HMGCR, a key rate-limiting enzyme in the cholesterol biosynthesis pathway, inhibits B cell proliferation. In the case of LPS stimulation, B cells seem to be able to obtain cholesterol by both uptake and biosynthesis. For example, B cells can still increase their cholesterol levels in the absence of LDL in the cell culture medium. In addition, NB-598, an inhibitor of the 2nd rate-limiting enzyme in the cholesterol biosynthesis pathway, can reduce the ability of LPS to increase cholesterol levels. In contrast, the greater effect of NB-598 on cholesterol levels of B cells grown in CF media relative to normal media, and the inability of mevalonate to rescue the effects of statin treatment in CF media, would indicate that B cells also use LDL uptake as a source of cholesterol. In line with this, B cells upregulated the LDLR following activation (**Figure 5D** [DOI](#)). The precise balance between these two mechanisms may depend on the duration of stimulation and the availability of LDL in the extracellular environment.

In line with our results, following CD40 mediated activation of human B cells, Atorvastatin was reported to inhibit human B cell proliferation and the expression of the activation markers CD80 and CD86 (Shimabukuro-Vornhagen et al, 2014 [DOI](#)). Interpretation of the effects of statins on B cells is, however, complex. Statins inhibit HMGCR, the rate-limiting enzyme in the mevalonate pathway that is required for the initial steps in cholesterol biosynthesis. This pathway generates farnesyl pyrophosphate, an intermediate that can feed into not only cholesterol biosynthesis, but also protein prenylation and ubiquinone pathways (Mullen et al, 2016 [DOI](#); Wang & Casey, 2016 [DOI](#)). Thus, the effects of statins reported here could be due not only to their effects on cholesterol, but also via these other pathways. Prenylation has been linked to proliferation in several cell types (Su et al, 2020 [DOI](#)), including T cells, where Simvastatin inhibits activation via two mechanisms, one of which involves blocking the prenylation of Ras and Rac (Ghittoni et al, 2005 [DOI](#)). We therefore investigated whether statin-mediated effects on B cell survival and proliferation were in part due to prenylation. We found that FGTI-2734, an inhibitor of FTase and GGTase, blocked LPS + IL-4 induced B cell proliferation, suggesting an important role for prenylation in B cells. In agreement with this, prenylation has also been reported to be important in B cells for the expression of CD80 and CD86 following stimulation via CD40, although its role in proliferation was not directly addressed (Shimabukuro-Vornhagen et al, 2014 [DOI](#)). Given that prenylation can target a large number of proteins, including many members of the Ras GTPase superfamily, to regulate their cellular localisation, the effects of blocking prenylation in B cells are likely to be complex. Interestingly, GGPP, which is needed for protein geranylgeranylation (a form of protein prenylation), was able to rescue the effects of Fluvastatin on LPS + IL-4 stimulated B cells. Notably, it was more efficient than mevalonate at restoring proliferation in the Fluvastatin treated cells. This ability to rescue proliferation was unexpected, as GGPP would not directly feed into the cholesterol biosynthesis pathway, and other experiments described here using CF media and

squalene monooxygenase inhibitors argue for a requirement for cholesterol for B cell proliferation. We therefore looked at its effect on cholesterol levels and found that it was able to rescue the effects of Fluvastatin in normal but not CF media. This could be explained by prenylation being required for cholesterol uptake but not biosynthesis. Cholesterol uptake via the LDLR requires the clathrin-mediated endocytosis of the LDLR and its trafficking to endosomes, where cholesterol is released from LDL (Nguyen et al, 2025 [DOI](#)). Vesicle trafficking is regulated via the action of a number of Rab GTPases, including Rab5, which is recruited to early endosomes and regulates their maturation (Zhao et al, 2025 [DOI](#); Zeigerer et al, 2012 [DOI](#)). More significantly, Rab5 prenylation has been shown to be important for its correct targeting to endosomes and therefore its biological function (Gomes et al, 2003 [DOI](#)). Treatment of cells with statins to inhibit prenylation has previously been shown to affect Rab subcellular localisation and vesicle trafficking (Ali et al, 2010 [DOI](#); Ronzier et al, 2019 [DOI](#); Xia et al, 2018 [DOI](#)). Loss of the prenylation of the Rab proteins involved in LDLR endocytosis could therefore negatively impact this process, leading to a failure of LDL uptake into the B cell. Prenylation is also likely to impact other processes in B cells, for example, TLR9-induced IL-10 production by regulatory B cells has been found to require geranylgeranylation and not farnesylation (Bibby et al, 2020 [DOI](#)). Further work will be needed to uncover the proteins prenylated upon B cell activation and determine the effects of prenylation on their function.

Overall, the proteomic analysis described here supports that B cells undergo a major reorganisation of metabolic pathways following TLR activation. Naïve B cells exist in a quiescent state where they do not grow or proliferate and express low levels of metabolic enzymes. Conversely, B cells stimulated with LPS + IL-4 massively upregulate metabolic enzymes involved in processes including protein synthesis, amino acid uptake, and cholesterol metabolism, which is required to support an increase in cell size, proliferation, and antibody class switching. Most significantly, we found that LPS + IL-4 stimulated B cells depended on both cholesterol biosynthesis, cholesterol uptake, and prenylation for expansion and proliferation.

Methods

Mice

Wildtype C57BL/6 mice (Charles Rivers) aged between 8 and 12 weeks were used for the majority of experiments. Conditional *SLC7A5* knockout mice bred to Vav-iCre mice have been described previously and were maintained on a C57BL6/J background (de Boer et al, 2003 [DOI](#); Poncet et al, 2014 [DOI](#); Sinclair et al, 2013 [DOI](#)). The mice were kept in individually ventilated cages, under specific pathogen-free conditions. Mice were given free access to food (RM3 irradiated pelleted diet, Special Diet Services) and water. Animal rooms were maintained at 21°C and a humidity of 45 to 55% with a 12/12 h light cycle. Mice were sacrificed using a rising concentration of CO₂ and death was confirmed by either cervical dislocation or exsanguination. All animal work was approved by the University of Dundee's Welfare and Ethical Use of Animals Committee and in accordance with UK and EU law.

Cell isolation

Lymph nodes were homogenised through a 40µm cell strainer in DPBS buffer (Gibco), and after centrifugation at 450g for 4 minutes, the pellet was resuspended in DPBS. Lymphocytes were counted on a Novocyte (Agilent Technologies) and seeded at 1×10⁶ cells/ml in B cell medium (RPMI-1640 supplemented with 10% FBS, 50 U/ml penicillin-streptomycin, 5mM L-glutamine, 10mM HEPES buffer, 1mM sodium pyruvate, 50 µM 2-mercaptoethanol and non-essential amino acids) for stimulation.

Spleens were dissociated through a 40µM filter in DPBS buffer and after centrifugation at 450g for 4 minutes, the pellet was resuspended in 1ml of Red Blood Cell Lysing Buffer (Hybri-Max™, Sigma) for 4 minutes at room temperature to lyse red blood cells. The reaction was quenched by the addition of DPBS, and the cells were pelleted by centrifugation at 450g for 4 minutes. The pellet was resuspended in 10ml DPBS before counting the splenocytes on a Novocyte. Splenocytes were seeded at 1×10^6 cells/ml in B cell medium for stimulation or further purification of B cells.

B cell isolation

To obtain an enriched population of B cells, magnetic sorting was used to remove non-B cells. Briefly, splenocytes were incubated with Fc receptor-blocking anti-CD16/CD32 antibody (BD Pharmingen) for 5 minutes on ice. The cells were incubated with a mixture of biotin-labelled anti-mouse antibodies (10µg/ml) ([Table 1](#)) for 20 minutes on ice. Unbound antibody was removed by washing with 10ml Magnetic Activated Cell Sorting (MACS) buffer (DPBS + 0.5% BSA + 2mM EDTA) followed by centrifugation. The cells were then resuspended in MACS buffer with streptavidin microbeads (1:10) (Miltenyi Biotec) and incubated at 4 C for 15 minutes. Miltenyi LD columns were calibrated by passing through 3ml of MACS buffer. Cells were washed with 10ml MACS buffer and centrifuged at 450g for 4 minutes to remove excess beads. The cell pellet was then resuspended in MACS buffer and separated using magnetic-activated cell sorting columns (LD) according to the manufacturer's instructions (Miltenyi Biotec). Single-cell suspensions were collected, and B cell numbers and purity (CD19+) were measured on a NovoCyt. Typical purity levels were 97% CD19+ B cells. B cells were seeded at 2×10^5 cells/ml of B cell medium for stimulation for flow cytometry experiments. When stated, purified B cells were stained with 2.5µM CellTrace™ Violet (ThermoFisher) at 37 C for 20 minutes, then quenched in 10ml B cell medium before seeding at the required density.

Stimulations and inhibitors

Cells were rested at 37 C for 1 hour. When required, the cells were pre-treated with the appropriate inhibitor for 45 minutes before stimulation, at the concentrations indicated in the figure legend: NB-598 (MedChemExpress), Fluvastatin sodium (TOCRIS), Rosuvastatin calcium (TOCRIS), FGTL-2734 (MedChemExpress). Inhibitors were dissolved in DMSO (Thermo Fisher) and used at a dilution of 1µl per 1000µl of culture medium. As indicated in the figure legends, cells were treated with 1µl DMSO as a vehicle control.

Cells were pre-incubated with the following metabolites at 37 C for 1 hour before inhibitor treatment/stimulation. Metabolites were used at the concentrations indicated in the figure legend: Mevalonic acid 5-phosphate lithium salt hydrate (Sigma Aldrich) which was dissolved in HEPES (Thermo Fisher). As indicated in the figure legends, cells were treated with 20µl HEPES as a vehicle control. Geranylgeranyl pyrophosphate ammonium salt (Sigma Aldrich) which comes in a 7:3 methanol: ammonium hydroxide solution. As indicated in the figure legends, cells were treated with 4.55µl 7:3 methanol: ammonium hydroxide solution as a vehicle control.

A combination of 20µg/ml LPS (L2654, Sigma) and 10ng/ml IL-4 (214-14, Peprotech) was used to stimulate the cells unless otherwise stated in the figure legend. For some experiments, cells were treated with LPS or IL-4 alone, or with 1µg/ml Resiquimod/R848 (tlrl-r848, Invivogen), 1µg/ml ODN 2006 (tlrl-2006, Invivogen), 10µg/ml Anti-Mouse IgM µ chain (115-006-075, Stratech) or 500ng/ml CD40 ligand (8230-CL-050/CF, R&D).

Cholesterol free serum

Silica-depleted FBS was purchased from Biowest and used in place of normal FBS when making B cell culture medium. Samples of normal and cholesterol-free FBS were sent to London Health Company, who carried out tests to measure changes in triglycerides, cholesterol, high-density lipoprotein (HDL) and low-density lipoprotein (LDL) ([Table S2](#)).

Antibody	Clone	Concentration	Catalogue number	Supplier
Anti-CD11b	M1/70	10µg/ml	101204	BioLegend
Anti-CD11c	N418	10µg/ml	117304	BioLegend
Anti-CD3ε	145-2C11	10µg/ml	100304	BioLegend
Anti-TER119	TER-119	10µg/ml	116204	BioLegend

Table 1

MACS antibodies.

Cell survival, proliferation and class switching assays

Following stimulation, cells were washed in Fluorescence Activated Cell Sorting (FACS) buffer (1% BSA, DPBS) and then incubated with Fc receptor-blocking anti-CD16/CD32 antibody in FACS buffer for 15 minutes at 4 C before staining with the appropriate fluorochrome-conjugated antibodies (Table 2) in FACS buffer for 20 minutes at 4 C. The samples were washed with FACS buffer prior to resuspension in 0.25µg of 7AAD. Flow cytometry data was collected on a Novocyte and analysed using FlowJo V.10 software (BD Biosciences). A representative gating strategy is shown in Figure S7A for lymph node/splenocyte isolation, and Figure S8B for B cell isolation.

Cell cycle analysis and p-Rb staining

Cells were washed in FACS buffer and then fixed using 1:1 IC fixation buffer (catalogue number: 00-8222-49, eBiosciences) and FACS buffer and incubated at 4 C for 20 minutes. Following another wash in FACS buffer, cells were resuspended in 100µl of cold ethanol and stored at -20°C for 30 minutes. The samples were washed with FACS buffer twice before incubation with Fc receptor-blocking anti-CD16/CD32 antibody in FACS buffer for 15 minutes at 4 C. Surface stains (Table 2) were incubated for 1 hour at room temperature before washing with FACS buffer and resuspension in 0.5µg/ml of DAPI. Flow cytometry data was collected on a Novocyte and analysed using FlowJo V.10 software. A representative gating strategy is shown in Figure S7B.

O-propargyl-puromycin uptake assay

Uptake of the puromycin analogue O-propargyl-puromycin was used to estimate protein synthesis rates. Splenocytes were stimulated with 20µg/ml LPS and 10ng/ml IL-4. After 24 hours the cells were treated with 20µM of O-propargyl-puromycin for 30 minutes at 37°C. 100 µL of 2% paraformaldehyde in B cell medium was added to 100µl of cells before incubation in complete darkness at room temperature for 30 minutes. The samples were pelleted by centrifugation at 450g for 4 minutes before resuspension in 100µl of DPBS. Following this, the samples were washed with FACS buffer and then resuspended in 100µl of 0.01% Saponin (Sigma) in DPBS, before incubation at room temperature for 20 minutes. The samples were pelleted by centrifugation at 450g for 4 minutes before resuspension in 30µl of Click mixture: 1mM Copper sulphate (209198, Sigma), 10mM Sodium ascorbate (A7631, Sigma), 1mM BTTAA (906328, Sigma), 10mM Aminoguanidine (81530, Cayman Chemicals), DPBS (14190144, Gibco), 5µM Alexafluor 647 azide (A10277, Invitrogen) and incubation in complete darkness at room temperature for 1 hour. The samples were pelleted by centrifugation at 450g for 4 minutes before resuspension in 200µl of FACS buffer and incubation in complete darkness at room temperature for 30 minutes. Again, the samples were pelleted by centrifugation at 450g for 4 minutes before washing again in FACS buffer. The cells were incubated with Fc receptor-blocking anti-CD16/CD32 antibody in FACS buffer for 15 minutes at 4 C. Surface stains (Table 1) were incubated for 1 hour at room temperature. The samples were washed with FACS buffer prior to resuspension in 300µl FACS buffer. Flow cytometry data was collected on a Novocyte and analysed using FlowJo V.10 software. A representative gating strategy is shown in Figure S7C.

Kynurenine uptake assay

Uptake of the tryptophan metabolite kynurenine was used to estimate the uptake rates via neutral amino acid transporters (Sinclair et al, 2018). Purified B cells were stimulated with 20µg/ml LPS and 10ng/ml IL-4. After 24 hours, the cells were washed in FACS buffer and then incubated with Fc receptor-blocking anti-CD16/CD32 antibody in FACS buffer for 15 minutes at 37 C before staining with CD19 APC (Table 2) in FACS buffer for 20 minutes at 37 C. The samples were washed with FACS buffer and then resuspended in 0.2ml HBSS. Either 0.1ml of HBSS or 0.1ml of 40mM BCH was then added to each sample. After this, 100µl of 800µM Kynurenine was added and the samples were incubated for 5 minutes at 37°C. After 5 minutes, the samples were fixed by adding 125µl of

Antibody	Fluorophore	Clone	Dilution	Catalogue number	Supplier
Anti-CD19	APC	6D5	1:300	115512	Biolegend
Anti-CD19	PE	1D3	1:200	557399	BD Biosciences
Anti-CD19	FITC	1D3	1:200	152403	BioLegend
Anti-CD93	APC	AA4.1	1:200	136510	BioLegend
Anti-CD98	PE	RL388	1:200	128208	BioLegend
dsDNA	eFluor 660	-	1:1000	65-0864-14	eBioscience
Anti-IgG1	FITC	A85-1	1:100	553443	BD Biosciences
Anti-pRb	FITC	D20B12	1:300	4277S	Cell Signaling Technology

Table 2

Flow cytometry antibodies

4% paraformaldehyde made in DPBS for 18 minutes at 4°C. The samples were then washed 2 times with FACS buffer and re-suspended in 400 µl of FACS buffer. Flow cytometry data was collected on an LSR Fortessa (BD biosciences) and analysed using FlowJo V.10 software. A representative gating strategy is shown in Figure S7D.

Cholesterol staining assay

Changes in cellular cholesterol content were determined using the cholesterol-binding agent filipin (Miller, 1984 [\[1\]](#)). Splenocytes were stimulated with 20 µg/ml LPS and 10 ng/ml IL-4. After 24 hours, the cells were washed twice in DPBS before staining for viability using eFluor 660 fixable viability dye (Table 2 [\[2\]](#)) for 30 minutes at 4°C. After two washes in FACS buffer, the cells were fixed using 1:1 IC fixation buffer (catalogue number: 00-8222-49, eBiosciences) and FACS buffer, before incubation at 4°C for 20 minutes. Following another wash in FACS buffer, cells were incubated with Fc receptor-blocking anti-CD16/CD32 antibody in FACS buffer for 15 minutes at 4°C. Surface stains (Table 2 [\[2\]](#)) were incubated for 20 minutes at 4°C. The samples were washed with FACS buffer prior to resuspension in 270 µl FACS buffer and 30 µl of 1 µg/µl filipin (Merck) before incubation in complete darkness at room temperature for 30 minutes. Flow cytometry data was collected on an LSR Fortessa and analysed using FlowJo V.10 software. A representative gating strategy is shown in Figure S8A.

Mitochondria staining assay

Splenocytes were stimulated with 20 µg/ml LPS and 10 ng/ml IL-4. After 24 hours, the cells were washed twice in FACS buffer. 50 µl of prewarmed (37°C) staining solution containing 100 nM MitoTracker Red probe (Invitrogen) in FACS buffer was added into the cells before incubation at 37°C for 20 minutes. The samples were washed with FACS buffer before incubation with Fc receptor-blocking anti-CD16/CD32 antibody (BD Pharmingen) in FACS buffer for 15 minutes at 4°C before staining with the appropriate fluorochrome-conjugated antibodies (Table 2 [\[2\]](#)) in FACS buffer for 20 minutes at 4°C. The samples were washed with FACS buffer prior to resuspension in 0.25 µg of 7AAD. Flow cytometry data was collected on a Novocyte and analysed using FlowJo V.10 software. A representative gating strategy is shown in Figure S8C.

Proteomics

Wildtype C57BL/6 mice (Charles Rivers) aged between 8 and 12 weeks were used for all proteomic experiments. Lymph nodes were extracted from mice, mashed in RPMI media before filtering through a 70 µm cell strainer. Naïve B cells were purified by fluorescence-activated cell sorting. Lymphocytes were incubated with Fc receptor-blocking anti-CD16/CD32 antibody (BD Pharmingen) in FACS buffer for 15 minutes at 4°C before staining with fluorochrome-conjugated antibodies (Table 2 [\[2\]](#)) in FACS buffer for 20 minutes at 4°C. The samples were washed with FACS buffer prior to resuspension in 0.5 µg/ml of DAPI. CD19⁺, CD93⁻, DAPI-cells were sorted using a Sony SH800 cell sorter. Sorted cells were washed twice with HBSS and snap frozen in liquid nitrogen to be stored at -80 °C until processing for mass spectrometry. For B cell activation, lymphocytes were suspended at a final density of 1.5 million cells/ml in medium (RPMI-1640 containing glutamine supplemented with 10% FBS, 50 U/ml penicillin-streptomycin, 50 µM 2-mercaptoethanol) and activated for 24 hours in the presence of 20 µg/ml LPS and 10 ng/ml IL-4. After 24 hours, the cells were incubated with Fc receptor-blocking anti-CD16/CD32 antibody (BD Pharmingen) in FACS buffer for 15 minutes at 4°C before staining with fluorochrome-conjugated antibodies (Table 2 [\[2\]](#)) in FACS buffer for 20 minutes at 4°C. The samples were washed with FACS buffer prior to resuspension in 0.5 µg/ml of DAPI. CD19⁺ and DAPI-cells were sorted and collected as described above. Cells were then washed 3 times in PBS and cell pellets frozen at -80°C.

Cell pellets were lysed and proteins were digested following the protocol described by (Baker et al, 2022 [\[3\]](#)). In brief, 400 µl of lysis buffer (5% sodium dodecyl sulfate, 50 mM triethylammonium bicarbonate (pH 8.5) and 10 mM tris(2-carboxyethyl)phosphine-hydrochloride) was added to each

sample and lysates shaken at room temperature at 1000 rpm for 5 minutes before being incubated at 95 °C at 500 rpm for 5 minutes. Samples were allowed to cool and were then sonicated using a BioRuptor (15 cycles: 30 sec on and 30 sec off) and alkylated with 20mM iodoacetamide for 1 h at 22 °C in the dark. To determine protein concentration the EZQ protein quantitation kit (Thermo) was used and protein cleanup and digestion was performed using S-TRAP mini columns (Protifi). Proteins were digested with trypsin at 1:20 ratio (enzyme:protein) for 2 hours at 47°C. Digested peptides were eluted from S-TRAP columns using 50mM ammonium bicarbonate, followed by 0.2% aqueous formic acid and 50% aqueous acetonitrile containing 0.2% formic acid. Peptides were dried by speedvac before resuspending in 1% formic acid. Peptide quantity was measured using the CBQCA kit (Thermo).

LC-MS/MS analysis

Peptides were analysed by single shot Data Independent Acquisition (DIA) mass spectrometry as previously described (Molina-Gonzalez et al, 2023 [DOI](#); Sollberger et al, 2024 [DOI](#)). 1.5 µg of peptide from each sample was injected onto a nanoscale C18 reverse-phase chromatography system (UltiMate 3000 RSLC nano, Thermo Scientific) and electrosprayed into an Orbitrap Exploris 480 Mass Spectrometer (Thermo Fisher). The following buffers were used for liquid chromatography: buffer A (0.1% formic acid in Milli-Q water (v/v)) and buffer B (80% acetonitrile and 0.1% formic acid in Milli-Q water (v/v)). Samples were loaded at 10 µl/min onto a trap column (100 µm × 2 cm, PepMap nanoViper C18 column, 5 µm, 100 Å, Thermo Scientific) equilibrated in 0.1% trifluoroacetic acid (TFA). The trap column was washed for 3 min at the same flow rate with 0.1% TFA then switched in-line with a Thermo Scientific, resolving C18 column (75 µm × 50 cm, PepMap RSLC C18 column, 2 µm, 100 Å). Peptides were eluted from the column at a constant flow rate of 300 nl/min with a linear gradient from 3% buffer B to 6% buffer B in 5 min, then from 6% buffer B to 35% buffer B in 115 min, and finally to 80% buffer B within 7 min. The column was then washed with 80% buffer B for 4 min and re-equilibrated in 3% buffer B for 15 min. Two blanks were run between each sample to reduce carry-over. The column was kept at a constant temperature of 50°C.

The data was acquired using an easy spray source operated in positive mode with spray voltage at 2.445 kV, and the ion transfer tube temperature at 250°C. The MS was operated in DIA mode. A scan cycle comprised a full MS scan (m/z range from 350 to 1650), with RF lens at 40%, AGC target set to custom, normalised AGC target at 300%, maximum injection time mode set to custom, maximum injection time at 20 ms, microscan set to 1 and source fragmentation disabled. MS survey scan was followed by MS/MS DIA scan events using the following parameters: multiplex ions set to false, collision energy mode set to stepped, collision energy type set to normalized, HCD collision energies set to 25.5, 27 and 30%, orbitrap resolution 30,000, first mass 200, RF lens 40%, AGC target set to custom, normalized AGC target 3000%, microscan set to 1 and maximum injection time 55 ms. Data for both MS scan and MS/MS DIA scan events were acquired in profile mode.

Proteomic data analysis

For comparing naïve and LPS stimulated B cells, raw mass spec data files were searched using Spectronaut version 19 (Biognosys). Data was analysed by DirectDIA. Raw mass spec data files were searched against a mouse database (Swissprot Trembl November 2023) with the following parameters: directDIA, false discovery rate set to 1%, protein N-terminal acetylation and methionine oxidation were set as variable modifications and carbamidomethylation of cysteine residues was selected as a fixed modification.

The following Spectronaut settings were used (Baker et al, 2024 [DOI](#)): all identification settings were set to 0.01 (precursor q-value cut-off, precursor PEP cut-off, protein FDR strategy (accurate), protein q-value cut-off (experiment), protein q-value cut-off (run) and protein PEP cut-off). The

following quantification settings were used: ‘Quant 2.0’, the MS-Level quantity was set to ‘MS2’, imputation was disabled, major group Top N and minor group Top N were set as ‘False’ and cross run normalisation was set as ‘False’.

Intensity vales from Spectronaut were further analysed in Perseus (Tyanova et al, 2016 [↗](#)) and the histone ruler method (Wiśniewski et al, 2014 [↗](#)) used to estimate copy number and concentrations of the identified proteins. Fold change was calculated relative to levels in naïve B cells and significance determined using unpaired Students t tests on $\log_{10}$ transformed data (**Supplementary table S3** [↗](#)). The False Discovery Rate approach using the 2-stage step up method Benjamini, Krieger, and Yekutieli (Benjamini et al, 2006 [↗](#)) with a Q value of 0.01. Enrichment against the Gene Ontology (GO) Biological Processes, Uniprot Keyword Biological Process and *Kyoto Encyclopaedia of Genes and Genomes* (KEGG) databases. Biological Process GO terms (Gene Ontology Consortium et al, 2023 [↗](#)) and KEGG pathways (Kanehisa et al, 2016 [↗](#)) was carried out using the DAVID gene enrichment site (Sherman et al, 2022 [↗](#)) (**Supplementary Table S1** [↗](#)).

Statistical analysis

Unless otherwise stated, graphs show mean values + SD with individual replicates shown by symbols. Unpaired two-tailed Student’s *t* test, one-way ANOVA, two-way ANOVA with post-hoc testing were performed in Prism, and the tests used are indicated in the figure legends. Full results of the ANOVA calculations are given in **Supplemental Table S4** [↗](#).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium (<http://proteomecentral.proteomexchange.org> [↗](#)) via the PRIDE partner repository (Perez-Riverol et al, 2024 [↗](#)) with the dataset identifier PXD057739. Flow cytometry data that support the findings of this study are available from the corresponding author upon request.

Any additional information required to reanalyse the data reported in this paper is available from the lead contact upon request.

Acknowledgements

The authors thank A. Rennie and R. Clarke from the Flow Cytometry Facility for cell sorting and advice on flow cytometry. We also thank the FingerPrints Proteomics facility for running our samples for mass spectrometry and for advice on proteomics. This research was supported by a Wellcome Trust (DMSC, UNS148487) and MRC (MR/N013735/1, MR) PhD studentship awards. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Additional information

Author contributions

D.M.S.C., M.R., S.A., performed the experiments. A.J.M. performed the liquid chromatography–mass spectrometry. D.M.S.C and J.S.C.A analysed the proteomics data. L.V.S designed the kynurenine uptake and protein synthesis experiments. M.S optimised the silica depletion and filipin staining experiments. D.M.S.C and J.S.C.A wrote the manuscript.

Resource availability Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Simon Arthur (j.s.c.arthur@dundee.ac.uk).

Materials availability

This study did not generate new unique reagents.

Additional files

Document S1. *Figures S1–S8 and Table S2 - Serum tests related to Figure 5 and Table S4 - Statistics table related to all figures.* [↗](#)

Supplemental Table S1. *Excel file containing terms for protein enrichment too large to fit in a pdf. Related to Figure 1H.* [↗](#)

Supplemental Table S3. *Figures S1–S8 and Table S2 - Serum tests related to Figure 5 and Table S4 - Statistics table related to all figures.* [↗](#)

References

1. Ali BR, Nouvel I, Leung KF, Hume AN, Seabra MC (2010) **A novel statin-mediated “prenylation block-and-release” assay provides insight into the membrane targeting mechanisms of small GTPases** *Biochemical and Biophysical Research Communications* **397**:34–41 [Google Scholar](#)
2. Arthur JSC, Ley SC (2013) **Mitogen-activated protein kinases in innate immunity** *Nat Rev Immunol* **13**:679–692 [Google Scholar](#)
3. Arunkumar N, Liu C, Hang H, Song W (2013) **Toll-like receptor agonists induce apoptosis in mouse B-cell lymphoma cells by altering NF- κ B activation** *Cell Mol Immunol* **10**:360–372 [Google Scholar](#)
4. Baker CP, Bruderer R, Abbott J, Arthur JSC, Brenes AJ (2024) **Optimizing Spectronaut Search Parameters to Improve Data Quality with Minimal Proteome Coverage Reductions in DIA Analyses of Heterogeneous Samples** *J Proteome Res* **23**:1926–1936 [Google Scholar](#)
5. Baker CP, Phair IR, Brenes AJ, Atrih A, Ryan DG, Bruderer R, Dinkova-Kostova AT, Lamont DJ, Arthur JSC, Howden AJM (2022) **DIA label-free proteomic analysis of murine bone-marrow-derived macrophages** *STAR Protocols* **3**:101725 [Google Scholar](#)
6. Barberis M, Klipp E, Vanoni M, Alberghina L (2007) **Cell Size at S Phase Initiation: An Emergent Property of the G1/S Network** *PLOS Computational Biology* **3**:e64 [Google Scholar](#)
7. Benjamini Y, Krieger AM, Yekutieli D (2006) **Adaptive linear step-up procedures that control the false discovery rate** *Biometrika* **93**:491–507 [Google Scholar](#)
8. Bibby JA, Purvis HA, Hayday T, Chandra A, Okkenhaug K, Rosenzweig S, Aksentijevich I, Wood M, Lachmann HJ, Kemper C, et al (2020) **Cholesterol metabolism drives regulatory B cell IL-10 through provision of geranylgeranyl pyrophosphate** *Nat Commun* **11**:3412 [Google Scholar](#)
9. Bierling TEH, Gumann A, Ottmann SR, Schulz SR, Weckwerth L, Thomas J, Gessner A, Wichert M, Kuwert F, Rost F, et al (2024) **GLUT1-mediated glucose import in B cells is critical for anaplerotic balance and humoral immunity** *Cell Reports* **43**:113739 [Google Scholar](#)
10. Bléry M, Tze L, Miosge LA, Jun JE, Goodnow CC (2006) **Essential role of membrane cholesterol in accelerated BCR internalization and uncoupling from NF- κ B in B cell clonal anergy** *J Exp Med* **203**:1773–1783 [Google Scholar](#)
11. de Boer J, Williams A, Skavdis G, Harker N, Coles M, Tolaini M, Norton T, Williams K, Roderick K, Potocnik AJ, et al (2003) **Transgenic mice with hematopoietic and lymphoid specific expression of Cre** *Eur J Immunol* **33**:314–325 [Google Scholar](#)
12. Brookens SK, Cho SH, Paik Y, Meyer K, Raybuck AL, Park C, Greenwood DL, Rathmell JC, Boothby MR (2023) **Plasma Cell Differentiation, Antibody Quality, and Initial Germinal Center B Cell Population Depend on Glucose Influx Rate** *The Journal of Immunology* **212**:43–56 [Google Scholar](#)

13. Brovkovich V, Aldrich A, Li N, Atilla-Gokcumen GE, Frasier J (2019) **Removal of Serum Lipids and Lipid-Derived Metabolites to Investigate Breast Cancer Cell Biology** *Proteomics* **19**:1800370 [Google Scholar](#)
14. Brown MS, Radhakrishnan A, Goldstein JL (2018) **Retrospective on Cholesterol Homeostasis: The Central Role of Scap** *Annu Rev Biochem* **87**:783–807 [Google Scholar](#)
15. Buhaescu I, Izzedine H (2007) **Mevalonate pathway: A review of clinical and therapeutical implications** *Clinical Biochemistry* **40**:575–584 [Google Scholar](#)
16. Calado DP, Sasaki Y, Godinho SA, Pellerin A, Köchert K, Sleckman BP, de Alborán IM, Janz M, Rodig S, Rajewsky K (2012) **The cell-cycle regulator c-Myc is essential for the formation and maintenance of germinal centers** *Nat Immunol* **13**:1092–1100 [Google Scholar](#)
17. Cantor J, Browne CD, Ruppert R, Féral CC, Fässler R, Rickert RC, Ginsberg MH (2009) **CD98hc facilitates B cell proliferation and adaptive humoral immunity** *Nat Immunol* **10**:412–419 [Google Scholar](#)
18. Chicas A, Wang X, Zhang C, McCurrach M, Zhao Z, Mert O, Dickins RA, Narita M, Zhang M, Lowe SW (2010) **Dissecting the unique role of the retinoblastoma tumor suppressor during cellular senescence** *Cancer Cell* **17**:376–387 [Google Scholar](#)
19. Coutinho A, Gronowicz E, Bullock WW, Möller G (1974) **Mechanism of thymus-independent immunocyte triggering** *The Journal of Experimental Medicine* **139**:74–92 [Google Scholar](#)
20. D’Avola A, Legrave N, Tajan M, Chakravarty P, Shearer RL, King HW, Kluckova K, Cheung EC, Clear AJ, Gunawan AS, et al (2022) **PHGDH is required for germinal center formation and is a therapeutic target in MYC-driven lymphoma** *J Clin Invest* **132**:e153436 [Google Scholar](#)
21. Dominguez-Sola D, Vitorica GD, Ying CY, Phan RT, Saito M, Nussenzweig MC, Dalla-Favera R (2012) **The proto-oncogene MYC is required for selection in the germinal center and cyclic reentry** *Nat Immunol* **13**:1083–1091 [Google Scholar](#)
22. Doudna JA, Rath VL (2002) **Structure and Function of the Eukaryotic Ribosome: The Next Frontier** *Cell* **109**:153–156 [Google Scholar](#)
23. Dufort FJ, Bleiman BF, Gumina MR, Blair D, Wagner DJ, Roberts MF, Abu-Amer Y, Chiles TC (2007) **Cutting Edge: IL-4-Mediated Protection of Primary B Lymphocytes from Apoptosis via Stat6-Dependent Regulation of Glycolytic Metabolism** *The Journal of Immunology* **179**:4953–4957 [Google Scholar](#)
24. Dziarski R (1982) **Preferential induction of autoantibody secretion in polyclonal activation by peptidoglycan and lipopolysaccharide. I. In vitro studies** *The Journal of Immunology* **128**:1018–1025 [Google Scholar](#)
25. Engeland K (2022) **Cell cycle regulation: p53-p21-RB signaling** *Cell Death Differ* **29**:946–960 [Google Scholar](#)
26. Fotiadis D, Kanai Y, Palacín M (2013) **The SLC3 and SLC7 families of amino acid transporters** *Molecular Aspects of Medicine* **34**:139–158 [Google Scholar](#)
27. Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, Ebert D, Feuermann M, Gaudet P, Harris NL, et al (2023) **The Gene Ontology knowledgebase in 2023** *Genetics* **224**:iyad031 [Google Scholar](#)

28. Genestier L, Taillardet M, Mondiere P, Gheit H, Bella C, Defrance T (2007) **TLR Agonists Selectively Promote Terminal Plasma Cell Differentiation of B Cell Subsets Specialized in Thymus-Independent Responses¹** *The Journal of Immunology* **178**:7779–7786 [Google Scholar](#)
29. Ghittoni R, Patrussi L, Pirozzi K, Pellegrini M, Lazzerini PE, Capecchi PL, Pasini FL, Baldari CT (2005) **Simvastatin inhibits T-cell activation by selectively impairing the function of Ras superfamily GTPases** *The FASEB Journal* **19**:1–24 [Google Scholar](#)
30. Gomes AQ, Ali BR, Ramalho JS, Godfrey RF, Barral DC, Hume AN, Seabra MC (2003) **Membrane Targeting of Rab GTPases Is Influenced by the Prenylation Motif** *Mol Biol Cell* **14**:1882–1899 [Google Scholar](#)
31. Greenwood J, Steinman L, Zamvil SS (2006) **Statin therapy and autoimmune disease: from protein prenylation to immunomodulation** *Nat Rev Immunol* **6**:358–370 [Google Scholar](#)
32. Hodgkin PD, Lee JH, Lyons AB (1996) **B cell differentiation and isotype switching is related to division cycle number** *Journal of Experimental Medicine* **184**:277–281 [Google Scholar](#)
33. Hoffman W, Lakkis FG, Chalasani G (2016) **B Cells, Antibodies, and More** *Clinical Journal of the American Society of Nephrology* **11**:137 [Google Scholar](#)
34. Hoshino K, Takeuchi O, Kawai T, Sanjo H, Ogawa T, Takeda Y, Takeda K, Akira S (1999) **Cutting Edge: Toll-Like Receptor 4 (TLR4)-Deficient Mice Are Hyporesponsive to Lipopolysaccharide: Evidence for TLR4 as the Lps Gene Product¹** *The Journal of Immunology* **162**:3749–3752 [Google Scholar](#)
35. Howden AJM, Hukelmann JL, Brenes A, Spinelli L, Sinclair LV, Lamond AI, Cantrell DA (2019) **Quantitative analysis of T cell proteomes and environmental sensors during T cell differentiation** *Nat Immunol* **20**:1542–1554 [Google Scholar](#)
36. Jellusova J, Cato MH, Apgar JR, Ramezani-Rad P, Leung CR, Chen C, Richardson AD, Conner EM, Benschop RJ, Woodgett JR, et al (2017) **Gsk3 is a metabolic checkpoint regulator in B cells** *Nat Immunol* **18**:303–312 [Google Scholar](#)
37. Johnson AG, Gaines S, Landy M (1956) **Studies on the O antigen of Salmonella typhosa. V. Enhancement of antibody response to protein antigens by the purified lipopolysaccharide** *The Journal of experimental medicine* **103**:225–246 [Google Scholar](#)
38. Johnstone JC, Yazicioglu YF, Clarke AJ (2024) **Fuelling B cells: dynamic regulation of B cell metabolism** *Current Opinion in Immunology* **91**:102484 [Google Scholar](#)
39. Kahya U, Köseer AS, Dubrovskaya A (2021) **Amino Acid Transporters on the Guard of Cell Genome and Epigenome** *Cancers* **13**:125 [Google Scholar](#)
40. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M (2016) **KEGG as a reference resource for gene and protein annotation** *Nucleic Acids Res* **44**:D457–462 [Google Scholar](#)
41. Kawai T, Adachi O, Ogawa T, Takeda K, Akira S (1999) **Unresponsiveness of MyD88-Deficient Mice to Endotoxin** *Immunity* **11**:115–122 [Google Scholar](#)
42. Kawai T, Ikegawa M, Ori D, Akira S (2024) **Decoding Toll-like receptors: Recent insights and perspectives in innate immunity** *Immunity* **57**:649–673 [Google Scholar](#)

43. Kazi A, Xiang S, Yang H, Chen L, Kennedy P, Ayaz M, Fletcher S, Cummings C, Lawrence HR, Beato F, et al (2019) **Dual Farnesyl and Geranylgeranyl Transferase Inhibitor Thwarts Mutant KRAS-Driven Patient-Derived Pancreatic Tumors** *Clinical Cancer Research* **25**:5984–5996 [Google Scholar](#)
44. Leone G, DeGregori J, Yan Z, Jakoi L, Ishida S, Williams RS, Nevins JR (1998) **E2F3 activity is regulated during the cell cycle and is required for the induction of S phase** *Genes Dev* **12**:2120–2130 [Google Scholar](#)
45. Luo J, Yang H, Song B-L (2020) **Mechanisms and regulation of cholesterol homeostasis** *Nat Rev Mol Cell Biol* **21**:225–245 [Google Scholar](#)
46. Luo W, Weisel F, Shlomchik MJ (2018) **B Cell Receptor and CD40 Signaling Are Rewired for Synergistic Induction of the c-Myc Transcription Factor in Germinal Center B Cells** *Immunity* **48**:313–326 [Google Scholar](#)
47. Marchingo JM, Sinclair LV, Howden AJ, Cantrell DA (2020) **Quantitative analysis of how Myc controls T cell proteomes and metabolic pathways during T cell activation** *eLife* **9**:e53725 <https://doi.org/10.7554/eLife.53725> | [Google Scholar](#)
48. Miller RG (1984) **The use and abuse of filipin to localize cholesterol in membranes** *Cell Biol Int Rep* **8**:519–535 [Google Scholar](#)
49. Mitsche MA, McDonald JG, Hobbs HH, Cohen JC (2015) **Flux analysis of cholesterol biosynthesis in vivo reveals multiple tissue and cell-type specific pathways** *eLife* **4**:e07999 <https://doi.org/10.7554/eLife.07999> | [Google Scholar](#)
50. Molina-Gonzalez I, Holloway RK, Jiwaji Z, Dando O, Kent SA, Emelianova K, Lloyd AF, Forbes LH, Mahmood A, Skripuletz T, et al (2023) **Astrocyte-oligodendrocyte interaction regulates central nervous system regeneration** *Nat Commun* **14**:3372 [Google Scholar](#)
51. Mullen PJ, Yu R, Longo J, Archer MC, Penn LZ (2016) **The interplay between cell signalling and the mevalonate pathway in cancer** *Nat Rev Cancer* **16**:718–731 [Google Scholar](#)
52. Napolitano L, Scalise M, Galluccio M, Pochini L, Albanese LM, Indiveri C (2015) **LAT1 is the transport competent unit of the LAT1/CD98 heterodimeric amino acid transporter** *Int J Biochem Cell Biol* **67**:25–33 [Google Scholar](#)
53. Nguyen MKL, Pinkenburg C, Du JJ, Bernaus-Esqu   M, Enrich C, Rentero C, Grewal T (2025) **The multiple facets of Rab proteins modulating the cellular distribution of cholesterol from the late endosomal compartment** *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1872**:119896 [Google Scholar](#)
54. Obukhanych TV, Nussenzweig MC (2006) **T-independent type II immune responses generate memory B cells** *Journal of Experimental Medicine* **203**:305–310 [Google Scholar](#)
55. Oesterle A, Laufs U, Liao JK (2017) **Pleiotropic Effects of Statins on the Cardiovascular System** *Circulation Research* **120**:229–243 [Google Scholar](#)
56. Paradoski BT, Hou S, Mejia EM, Olayinka-Adefemi F, Fowke D, Hatch GM, Saleem A, Banerji V, Hay N, Zeng H, et al (2024) **PI3K-dependent reprogramming of hexokinase isoforms**

controls glucose metabolism and functional responses of B lymphocytes *iScience*
27 [Google Scholar](#)

57. Pasare C, Medzhitov R (2005) **Control of B-cell responses by Toll-like receptors** *Nature* **438**:364–368 [Google Scholar](#)
58. Peeters R, Jellusova J (2023) **Lipid metabolism in B cell biology** *Molecular Oncology* **18**:1795–1813 [Google Scholar](#)
59. Perez-Riverol Y, Bandla C, Kundu DJ, Kamatchinathan S, Bai J, Hewapathirana S, John NS, Prakash A, Walzer M, Wang S, et al (2024) **The PRIDE database at 20 years: 2025 update** *Nucleic Acids Res* :gkae1011 [Google Scholar](#)
60. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, et al (1998) **Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene** *Science* **282**:2085–2088 [Google Scholar](#)
61. Poncet N, Mitchell FE, Ibrahim AFM, McGuire VA, English G, Arthur JSC, Shi Y-B, Taylor PM (2014) **The catalytic subunit of the system L1 amino acid transporter (slc7a5) facilitates nutrient signalling in mouse skeletal muscle** *PLoS One* **9**:e89547 [Google Scholar](#)
62. Procaccini C, Carbone F, Di Silvestre D, Brambilla F, De Rosa V, Galgani M, Faicchia D, Marone G, Tramontano D, Corona M, et al (2016) **The Proteomic Landscape of Human Ex Vivo Regulatory and Conventional T Cells Reveals Specific Metabolic Requirements** *Immunity* **44**:406–421 [Google Scholar](#)
63. Ronzier E, Parks XX, Qudsi H, Lopes CM (2019) **Statin-specific inhibition of Rab-GTPase regulates cPKC-mediated IKs internalization** *Sci Rep* **9**:17747 [Google Scholar](#)
64. Rubin SM, Sage J, Skotheim JM (2020) **Integrating Old and New Paradigms of G1/S Control** *Mol Cell* **80**:183–192 [Google Scholar](#)
65. Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, Imamichi T, Chang W (2022) **DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update)** *Nucleic Acids Res* **50**:W216–W221 [Google Scholar](#)
66. Shimabukuro-Vornhagen A, Zoghi S, Liebig TM, Wennhold K, Chemitz J, Draube A, Kochanek M, Blaschke F, Pallasch C, Holtick U, et al (2014) **Inhibition of Protein Geranylgeranylation Specifically Interferes with CD40-Dependent B Cell Activation, Resulting in a Reduced Capacity To Induce T Cell Immunity** *The Journal of Immunology* **193**:5294–5305 [Google Scholar](#)
67. Sinclair LV, Cantrell DA (2025) **Protein Synthesis and Metabolism in T Cells** *Annu Rev Immunol* **43**:343–366 [Google Scholar](#)
68. Sinclair LV, Neyens D, Ramsay G, Taylor PM, Cantrell DA (2018) **Single cell analysis of kynurenine and System L amino acid transport in T cells** *Nat Commun* **9**:1981 [Google Scholar](#)
69. Sinclair LV, Rolf J, Emslie E, Shi Y-B, Taylor PM, Cantrell DA (2013) **Control of amino-acid transport by antigen receptors coordinates the metabolic reprogramming essential for T cell differentiation** *Nat Immunol* **14**:500–508 [Google Scholar](#)

70. Snapper CM, Finkelman FD, Stefany D, Conrad DH, Paul WE (1988) **IL-4 induces co-expression of intrinsic membrane IgG1 and IgE by murine B cells stimulated with lipopolysaccharide** *The Journal of Immunology* **141**:489–498 [Google Scholar](#)
71. Sollberger G, Brenes AJ, Warner J, Arthur JSC, Howden AJM (2024) **Quantitative proteomics reveals tissue-specific, infection-induced and species-specific neutrophil protein signatures** *Sci Rep* **14**:5966 [Google Scholar](#)
72. Su W, Chapman NM, Wei J, Zeng H, Dhungana Y, Shi H, Saravia J, Zhou P, Long L, Rankin S, et al (2020) **Protein Prenylation Drives Discrete Signaling Programs for the Differentiation and Maintenance of Effector Treg Cells** *Cell Metabolism* **32**:996–1011 [Google Scholar](#)
73. Sun Y, Wen J, Xu T, Meng L (2023) **Reduction of peritoneal cavity B1a cells in adult Slc7a5 knockdown mice via dysregulating the mTOR pathway** *Int Immunopharmacol* **117**:109939 [Google Scholar](#)
74. Tan H, Yang K, Li Y, Shaw TI, Wang Y, Blanco DB, Wang X, Cho J-H, Wang H, Rankin S, et al (2017) **Integrative Proteomics and Phosphoproteomics Profiling Reveals Dynamic Signaling Networks and Bioenergetics Pathways Underlying T Cell Activation** *Immunity* **46**:488–503 [Google Scholar](#)
75. Tanner LB, Goglia AG, Wei MH, Sehgal T, Parsons LR, Park JO, White E, Toettcher JE, Rabinowitz JD (2018) **Four Key Steps Control Glycolytic Flux in Mammalian Cells** *Cell Syst* **7**:49–62 [Google Scholar](#)
76. Tesi A, de Pretis S, Furlan M, Filipuzzi M, Morelli MJ, Andronache A, Doni M, Verrecchia A, Pelizzola M, Amati B, et al (2019) **An early Myc-dependent transcriptional program orchestrates cell growth during B-cell activation** *EMBO reports* **20**:e47987 [Google Scholar](#)
77. Torigoe M, Maeshima K, Ozaki T, Omura Y, Gotoh K, Tanaka Y, Ishii K, Shibata H (2019) **I-Leucine influx through Slc7a5 regulates inflammatory responses of human B cells via mammalian target of rapamycin complex 1 signaling** *Modern Rheumatology* **29**:885–891 [Google Scholar](#)
78. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J (2016) **The Perseus computational platform for comprehensive analysis of (prote)omics data** *Nat Methods* **13**:731–740 [Google Scholar](#)
79. Wagner EF, Hleb M, Hanna N, Sharma S (1998) **A pivotal role of cyclin D3 and cyclin-dependent kinase inhibitor p27 in the regulation of IL-2-, IL-4-, or IL-10-mediated human B cell proliferation** *J Immunol* **161**:1123–1131 [Google Scholar](#)
80. Wang M, Casey PJ (2016) **Protein prenylation: unique fats make their mark on biology** *Nat Rev Mol Cell Biol* **17**:110–122 [Google Scholar](#)
81. Waters LR, Ahsan FM, Wolf DM, Shirihi O, Teitell MA (2018) **Initial B Cell Activation Induces Metabolic Reprogramming and Mitochondrial Remodeling** *iScience* **5**:99–109 [Google Scholar](#)
82. Weeratna RD, Makinen SR, McCluskie MJ, Davis HL (2005) **TLR agonists as vaccine adjuvants: comparison of CpG ODN and Resiquimod (R-848)** *Vaccine* **23**:5263–5270 [Google Scholar](#)
83. Wiśniewski JR, Hein MY, Cox J, Mann M (2014) **A “Proteomic Ruler” for Protein Copy Number and Concentration Estimation without Spike-in Standards** *Molecular & Cellular Proteomics*

13:3497–3506 [Google Scholar](#)

84. Xia Y, Xie Y, Yu Z, Xiao H, Jiang G, Zhou X, Yang Y, Li X, Zhao M, Li L, et al (2018) **The Mevalonate Pathway Is a Druggable Target for Vaccine Adjuvant Discovery** *Cell* **175**:1059–1073 [Google Scholar](#)
85. Yi A-K, Chang M, Peckham DW, Krieg AM, Ashman RF (1998) **CpG Oligodeoxyribonucleotides Rescue Mature Spleen B Cells from Spontaneous Apoptosis and Promote Cell Cycle Entry** *The Journal of Immunology* **160**:5898–5906 [Google Scholar](#)
86. Zeigerer A, Gilleron J, Bogorad RL, Marsico G, Nonaka H, Seifert S, Epstein-Barash H, Kuchimanchi S, Peng CG, Ruda VM, et al (2012) **Rab5 is necessary for the biogenesis of the endolysosomal system in vivo** *Nature* **485**:465–470 [Google Scholar](#)
87. Zhao X, Long S, Zhu M, Hao H, Liao Y, Zhang C, Liu Z (2025) **Potential Roles of SNX17, Rab11, and Rab5 in LDLR Recycling Arteriosclerosis, Thrombosis, and Vascular Biology** **45**:e338–e354 [Google Scholar](#)

Author information

Dana MS Cheung

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom
ORCID iD: [0000-0003-4920-8876](#)

Momchil Razsolkov

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom

Fabrizia Bonacina

Department of Pharmacological and Biomolecular sciences “Rodolfo Paoletti”, University of Milan, Milan, Italy

Stephen Andrews

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom

Megan Sumoreeah

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom

Linda V Sinclair

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom

Andrew JM Howden

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee, Dundee, United Kingdom

J Simon C Arthur

Division of Cell Signalling and Immunology, School of Life Sciences, University of Dundee,
Dundee, United Kingdom

For correspondence: j.s.c.arthur@dundee.ac.uk

Editors

Reviewing Editor

Mia Smith

University of Colorado Anschutz Medical Campus, Aurora, United States of America

Senior Editor

Tadatsugu Taniguchi

The University of Tokyo, Tokyo, Japan

Reviewer #1 (Public review):

The work presented by Cheung et al. used a quantitative proteomics method to capture molecular changes in B cells exposed to LPS and IL-4, a combination of stimuli activating naive B cells. Amino acid transporters, cholesterol biosynthetic enzymes, ribosomal components, and other proteins involved in cell proliferation were found to increase in stimulated B cells. Experiments involving genetic loss-of-function (SLC7A5), pharmacological inhibition (HMGCR, SQLE, prenylation), and functional rescue by metabolites (mevalonate, GGPP) validated the proteomics data and revealed that amino acid uptake, cholesterol/mevalonate biosynthesis, and cholesterol uptake played a crucial role in B cell proliferation, survival, biogenesis, and immunoglobulin class switching. Experiments involving cholesterol-free medium showed that both biosynthesis and LDLR-mediated uptake catered to the cholesterol demand of LPS/IL-4-stimulated B cells. A role for protein prenylation in LDLR-mediated cholesterol uptake was postulated and backed by divergent effects of GGPP rescue in the presence and absence of cholesterol in culture medium.

Strengths:

The discovery was made by proteome-wide profiling and unbiased computational analysis. The discovered proteins were functionally validated using appropriate tools and approaches. The metabolic processes identified and prioritized from this comprehensive survey and systematic validation are highly likely to represent mechanisms of high importance and influence. Analysis of immune cell metabolism at the protein level is relatively compared to transcriptomic and metabolomic analysis.

The conclusions from functional validation experiments were supported by clear data and based on rational interpretations. This was enabled by well-established readouts/analytical methods used to analyze cell proliferation, viability, size, cholesterol content, and transporter/enzyme function. The data generated from these experiments strongly support the conclusions.

This work reveals a complex, yet intriguing, relationship between cholesterol metabolism and protein prenylation as they serve to promote B cell activation. The effects of pharmacological inhibition and metabolite replenishment on the cholesterol content and activation of B cells were precisely determined and logically interpreted.

Weaknesses:

The findings of this study were obtained almost exclusively from ex vivo B cell stimulation experiments. Their contribution to B cell state and B-cell-mediated immune responses in vivo was not explored. Without in vivo data, the study still provides valuable mechanistic information and insights, but it remains unknown, and there is no discussion about how the identified mechanisms may play out in B cell immunity.

The role of HMGCR, SQLE, and prenylation in B cell activation was assessed using pharmacological inhibitors. Evidence from other loss-of-function approaches, which could strengthen the conclusions, does not exist. This is a moderate weakness.

<https://doi.org/10.7554/eLife.109093.1.sa1>

Reviewer #2 (Public review):

This study uses mass spectrometry to quantify how LPS and IL-4 modify the mouse B cell proteome as naïve cells undergo blastogenesis and enter the cell cycle. This analysis revealed changes in key proteins involved in amino acid transport and cholesterol biosynthesis. Genetic and pharmacological experiments indicated important roles for these metabolic processes in B cell proliferation.

This work provides new information about the regulation of TI B cell responses by changes in cell metabolism and also a comprehensive mass spectrometry dataset, which will be an important general resource for future studies. The experiments are thorough and carefully carried out. The majority of conclusions are backed up by data that is shown to be highly significant statistically.

The study would be strengthened by additional experiments to determine whether the detected changes are unique to stimulation with LPS + IL-4 or more generic responses of resting B cells to mitogenic agonists.

<https://doi.org/10.7554/eLife.109093.1.sa0>

Author response:

Reviewer #1:

We agree with the reviewer that a limitation of our study is its focus on cell-based assays rather than *in vivo* experiments. We did consider evaluating the effects of statins on B cell responses *in vivo*; however, this approach is complicated by findings that statins can influence antigen presentation by dendritic cells, thereby impacting antibody responses (Xia *et al*, 2018). One possible solution would be to use B cell-specific conditional knockout models to study the roles of the identified proteins in an *in vivo* context. However, we currently do not have access to these models and were therefore unable to include such experiments within a feasible timeframe. We will revise the discussion section to acknowledge these points.

The reviewer also noted that our study assessed the roles of HMGCR, SQLE, and prenylation in B cell activation using pharmacological inhibitors and genetic knockdown/out approaches. Loss-of-function techniques such as RNAi, siRNA, and CRISPR can be challenging to apply to primary B cells, but we are exploring their feasibility for future revisions. While we acknowledge the limitations of using pharmacological inhibitors, we have taken several steps to mitigate these, including targeting multiple steps in the cholesterol biosynthetic pathway using structurally distinct inhibitors and conducting rescue experiments by supplementing downstream metabolites. To further investigate potential off-target effects of statins, we have

recently performed proteomic analysis of B cells treated with and without fluvastatin. The data suggest that fluvastatin primarily affects cholesterol metabolism and does not cause widespread off-target effects. We will include this new data in the revised manuscript.

Reviewer #2:

The reviewer suggested that the study would be strengthened by determining whether the observed changes are specific to LPS + IL-4 stimulation or represent a more general B cell response to mitogenic signals.

A complementary study by James et al. (James *et al*, 2024) investigated murine B cells stimulated via the B cell receptor (BCR) and CD40, using anti-IgM and anti-CD40 antibodies alongside IL-4. Their proteomic analysis showed that such co-stimulation induces a fivefold increase in total cellular protein mass within 24 hours, mirroring our findings with LPS + IL-4. They also reported upregulation of proteins associated with cell cycle progression, ribosome biogenesis, and amino acid transport. Furthermore, by using SLC7A5 knockout mice, they demonstrated that this transporter is required for B cell activation. We will expand our discussion to include and these findings. We will also expand on the final figure in our paper showing that the effects of statins are not limited to LPS.

References:

James O, Sinclair LV, Lefter N, Salerno F, Brenes A & Howden AJM (2024) A proteomic map of B cell activation and its shaping by mTORC1, MYC and iron. *bioRxiv* 2024.12.19.629506 doi:10.1101/2024.12.19.629506 [PREPRINT]

Xia Y, Xie Y, Yu Z, Xiao H, Jiang G, Zhou X, Yang Y, Li X, Zhao M, Li L, *et al* (2018) The Mevalonate Pathway Is a Druggable Target for Vaccine Adjuvant Discovery. *Cell* 175: 1059-1073.e21

<https://doi.org/10.7554/eLife.109093.1.sa3>