

Conservation of the cooling agent binding pocket within the TRPM subfamily

Reviewed Preprint

v2 • October 2, 2024

Revised by authors

Reviewed Preprint

v1 • July 10, 2024

Kate Huffer, Matthew CS Denley, Elisabeth V Oskoui, Kenton J Swartz 

Molecular Physiology and Biophysics Section, Porter Neuroscience Research Center, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, United States

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

Abstract

Transient Receptor Potential (TRP) channels are a large and diverse family of tetrameric cation selective channels that are activated by many different types of stimuli, including noxious heat or cold, organic ligands such as vanilloids or cooling agents, or intracellular Ca^{2+} . Structures available for all subtypes of TRP channels reveal that the transmembrane domains are closely related despite their unique sensitivity to activating stimuli. Here we use computational and electrophysiological approaches to explore the conservation of the cooling agent binding pocket identified within the S1-S4 domain of the Melastatin subfamily member TRPM8, the mammalian sensor of noxious cold, with other TRPM channel subtypes. We find that a subset of TRPM channels, including TRPM2, TRPM4 and TRPM5, contain pockets very similar to the cooling agent binding pocket in TRPM8. We then show how the cooling agent icilin modulates activation of TRPM4 to intracellular Ca^{2+} , enhancing the sensitivity of the channel to Ca^{2+} and diminishing outward-rectification to promote opening at negative voltages. Mutations known to promote or diminish activation of TRPM8 by cooling agents similarly alter activation of TRPM4 by icilin, suggesting that icilin binds to the cooling agent binding pocket to promote opening of the channel. These findings demonstrate that TRPM4 and TRPM8 channels share related ligand binding pockets that are allosterically coupled to opening of the pore.

eLife Assessment

In this **valuable** study, Huffer et al posit that non-cold sensing members of the TRPM subfamily of ion channels (e.g., TRPM2, TRPM4, TRPM5) contain a binding pocket for icilin that overlaps with the one found in the cold-activated TRPM8 channel. After examining a body of TRP channel cryo-EM structures to identify the conserved site, this study presents **convincing** electrophysiological evidence supporting the presence of an icilin binding pocket within TRPM4. This study shows that icilin has modulatory effects on the TRPM4 channel and will be of direct interest to those working in the TRP-channel field, but it also has implications for studies of somatosensation, taste, as well as pharmacological targeting of the TRPM subfamily.

<https://doi.org/10.7554/eLife.99643.2.sa2>

Introduction

The founding member of the Transient Receptor Potential (TRP) channel family was cloned two decades after a *Drosophila trp* mutant was discovered to exhibit a transient receptor potential in electroretinogram recordings (Cosens and Manning 1969 [↗](#), Montell and Rubin 1989 [↗](#), Wong, Schaefer et al. 1989 [↗](#)). Many other TRP channels have been subsequently identified in eukaryotes and classified into ten subfamilies based on sequence similarity (Himmel and Cox 2020 [↗](#), Cabezas-Bratesco, McGee et al. 2022 [↗](#)). All TRP channels are tetrameric cation channels that contain six transmembrane (TM) segments (S1 to S6) per subunit (**Figure 1** [↗](#)), similar to voltage-activated K⁺ channels (Cao 2020 [↗](#), Huffer, Aleksandrova et al. 2020 [↗](#)). However, individual TRP channel family members respond to multiple different types of stimuli, including chemical ligands, temperature, voltage, membrane lipids and mechanical deformation, and they often function as coincidence detectors that integrate multiple stimuli (Chuang, Neuhausser et al. 2004 [↗](#)). TRP channels can be regulated by G-protein pathways, and in many cases may function as receptor-operated channels, including the founding member in *Drosophila* photoreceptors (Plant and Schaefer 2003 [↗](#), Vazquez, Wedel et al. 2004 [↗](#), Kukkonen 2011 [↗](#), Shalygin, Kolesnikov et al. 2021 [↗](#), Rohacs 2024 [↗](#)). Two of the most extensively studied TRP channels are TRPV1 and TRPM8, which in mammals function as detectors of noxious heat (Caterina, Schumacher et al. 1997 [↗](#), Tominaga, Caterina et al. 1998 [↗](#), Caterina, Leffler et al. 2000 [↗](#)) and noxious cold (McKemy, Neuhausser et al. 2002 [↗](#), Peier, Moqrich et al. 2002 [↗](#), Bautista, Siemens et al. 2007 [↗](#), Dhaka, Murray et al. 2007 [↗](#)), respectively. Although the mechanisms of temperature sensing in TRPV1 and TRPM8 remain incompletely understood (Bandell, Dubin et al. 2006 [↗](#), Grandl, Hu et al. 2008 [↗](#), Grandl, Kim et al. 2010 [↗](#), Clapham and Miller 2011 [↗](#), Qin 2014 [↗](#), Jara-Oseguera, Bae et al. 2016 [↗](#), Liu and Qin 2017 [↗](#), Zhang, Jara-Oseguera et al. 2018 [↗](#), Garcia-Avila and Islas 2019 [↗](#), Diaz-Franulic, Verdugo et al. 2021 [↗](#), Lu, Yao et al. 2022 [↗](#), Yeh, Jara-Oseguera et al. 2023 [↗](#)), there is consensus about where vanilloids like capsaicin bind to activate TRPV1 and cooling agents such as icilin bind to activate TRPM8 (Cao, Liao et al. 2013 [↗](#), Liao, Cao et al. 2013 [↗](#), Gao, Cao et al. 2016 [↗](#), Yin, Wu et al. 2018 [↗](#), Yin, Le et al. 2019 [↗](#), Yin and Lee 2020 [↗](#), Zhang, Julius et al. 2021 [↗](#), Zhao, Xie et al. 2022 [↗](#)).

Vanilloids and cooling agents were originally proposed to bind to similar regions of TRPV1 and TRPM8 because mutations affecting channel activation by these compounds could be found in similar TM regions (Chuang, Neuhausser et al. 2004 [↗](#)). However, structures solved using cryogenic electron microscopy (cryo-EM) have since revealed that vanilloids bind to a membrane-facing pocket positioned at the interface between the pore domain formed by S5-S6 and the peripheral S1-S4 domains (Cao, Liao et al. 2013 [↗](#), Liao, Cao et al. 2013 [↗](#), Gao, Cao et al. 2016 [↗](#), Zhang, Julius et al. 2021 [↗](#)), whereas cooling agents bind within a pocket formed entirely by the S1-S4 helices that opens to the cytoplasm (Yin, Wu et al. 2018 [↗](#), Yin, Le et al. 2019 [↗](#), Yin and Lee 2020 [↗](#)) (**Figure 1** [↗](#)). The vanilloid binding pocket in TRPV1 can be occupied by vanilloid agonists and antagonists (Cao, Liao et al. 2013 [↗](#), Liao, Cao et al. 2013 [↗](#), Gao, Cao et al. 2016 [↗](#), Kwon, Zhang et al. 2021 [↗](#), Zhang, Julius et al. 2021 [↗](#), Kwon, Zhang et al. 2022 [↗](#), Neuberger, Oda et al. 2023 [↗](#)), in addition to membrane lipids that have been proposed to stabilize a closed state of the channel (Cao, Cordero-Morales et al. 2013 [↗](#), Gao, Cao et al. 2016 [↗](#), Zhang, Julius et al. 2021 [↗](#), Arnold, Mancino et al. 2024 [↗](#)). Cryo-EM maps for all five available structures with the cooling agent icilin bound (6nr3, 7wrc, 7wrd, 7wre, and 7wrf) show a large non-protein density consistent with icilin nestled within the S1-S4 domain (Yin, Wu et al. 2018 [↗](#), Yin, Le et al. 2019 [↗](#), Yin and Lee 2020 [↗](#), Zhao, Xie et al. 2022 [↗](#)) (**Figure 1** [↗](#)). This cooling agent binding pocket is lined by residues where mutations are known to modulate sensitivity to icilin and/or the cooling agent menthol (Chuang, Neuhausser et al. 2004 [↗](#), Bandell, Dubin et al. 2006 [↗](#), Yin, Le et al. 2019 [↗](#), Yin and Lee 2020 [↗](#), Zhao, Xie et al. 2022 [↗](#)), supporting the assignment of this density to icilin during model building (**Figure 2** [↗](#)). Other TRPM8 ligands have also been observed in this pocket, including cooling agents WS-12 (6nr2) and C3 (8e4l, 8e4m) and the antagonists AMTB (6o6r) and




Figure 1

Structures of vanilloid bound to TRPV1 and cooling agent bound to TRPM8

Side views of **A**) TRPM8 bound to icilin (CPK) and Ca²⁺ (green sphere) and **B**) TRPV1 bound to RTX (CPK). Intracellular views of **C**) TRPM8 bound to icilin and Ca²⁺ and **D**) TRPV1 bound to RTX.

TC-I 2014 (6o72) (Diver, Cheng et al. 2019 [DOI](#), Yin, Le et al. 2019 [DOI](#), Yin and Lee 2020 [DOI](#), Yin, Zhang et al. 2022 [DOI](#), Zhao, Xie et al. 2022 [DOI](#)). Interestingly, other TRP channel structures from the Canonical and Vanilloid subfamilies also show small molecule ligands bound to a similar location within the S1-S4 domain (7b0s, 7b05, 7wdb, 7d4p, 6uza, 7dxg, 6dvy, 6dvz, 7ras, 7rau, 6pbe, 6d7o, 6d7q, 6d7t, 6d7v, 6d7x) (Singh, McGoldrick et al. 2018 [DOI](#), Singh, Saotome et al. 2018 [DOI](#), Hughes, Del Rosario et al. 2019 [DOI](#), Bai, Yu et al. 2020 [DOI](#), Vinayagam, Quentin et al. 2020 [DOI](#), Neuberger, Nadezhdin et al. 2021 [DOI](#), Song, Wei et al. 2021 [DOI](#), Guo, Tang et al. 2022 [DOI](#)), raising the possibility that this pocket may be a hot spot for ligand modulation across the TRP channel family.

The cytoplasmic entrance to the cooling agent binding pocket in TRPM8 also contains a binding site for a Ca^{2+} ion (6nr3, 7wrb, 7wrc, 7wrd, 7wre, 7wrf, 8e4l, 8e4m, 6o77) (**Figure 1** [DOI](#)) (Diver, Cheng et al. 2019 [DOI](#), Yin, Le et al. 2019 [DOI](#), Yin, Zhang et al. 2022 [DOI](#), Zhao, Xie et al. 2022 [DOI](#)). While Ca^{2+} alone has not been described to modulate TRPM8 activity, it is an essential cofactor for activation of TRPM8 by icilin (Chuang, Neuhausser et al. 2004 [DOI](#)), and several residues, including E782 in S2 and D802 in S3 of TRPM8 from *Ficedula albicollis* (faTRPM8) and mouse (mTRPM8), are located within 4 Å of both the Ca^{2+} ion and the icilin molecule (Yin, Le et al. 2019 [DOI](#), Zhao, Xie et al. 2022 [DOI](#)). Mutations in N799 and D802 in S3 of TRPM8 from rats (rTRPM8) and humans (hTRPM8) have been implicated in selective loss of icilin sensitivity (Chuang, Neuhausser et al. 2004 [DOI](#), Winking, Hoffmann et al. 2012 [DOI](#), Kuhn, Winking et al. 2013 [DOI](#), Beccari, Gemei et al. 2017 [DOI](#)). Because N799 does not contact the icilin molecule directly, it presumably perturbs icilin sensitivity by disrupting binding of the obligate cofactor, Ca^{2+} . Interestingly, the equivalent Ca^{2+} binding site has been observed in several other TRPM channel structures (6bqv, 6co7, 6d73, 6drj, 6nr3, 6o77, 6pkx, 6pus, 6puu, 7mbq, 7mbs, 7mbu, 7mbv, 7wrb, 7wrc, 7wrd, 7wre, 7wrf) (Autzen, Myasnikov et al. 2018 [DOI](#), Huang, Winkler et al. 2018 [DOI](#), Zhang, Toth et al. 2018 [DOI](#), Diver, Cheng et al. 2019 [DOI](#), Huang, Roth et al. 2019 [DOI](#), Yin, Le et al. 2019 [DOI](#), Yin, Wu et al. 2019 [DOI](#), Ruan, Haley et al. 2021 [DOI](#), Zhao, Xie et al. 2022 [DOI](#)), including TRPM2, TRPM4, and TRPM5, for which intracellular Ca^{2+} is the key physiological agonist, and where mutations to the Ca^{2+} -coordinating residues have been shown to alter channel activation by intracellular Ca^{2+} (Guo, She et al. 2017 [DOI](#), Autzen, Myasnikov et al. 2018 [DOI](#), Zhang, Toth et al. 2018 [DOI](#), Yamaguchi, Tanimoto et al. 2019 [DOI](#)). Structures from other TRP channel subfamilies—including Canonical (5z96, 6aei, 6jzo, 7b05, 7b0j, 7b0s, 7b16, 7b1g, 7d4p, 7d4q, 7dxb, 7dxf, 7e4t, 7wdb) (Duan, Li et al. 2018 [DOI](#), Vinayagam, Mager et al. 2018 [DOI](#), Duan, Li et al. 2019 [DOI](#), Song, Wei et al. 2021 [DOI](#), Guo, Tang et al. 2022 [DOI](#), Yang, Wei et al. 2022 [DOI](#)), Ankyrin (6v9v, 6v9w, 7or0, 7or1) (Zhao, Lin King et al. 2020 [DOI](#)), and Polycystin (7d7e, 7d7f) (Su, Chen et al. 2021 [DOI](#)) families—also contain cryo-EM density attributed to ions in this location. As in the cooling agent binding pocket, the presence of this ion binding pocket within the S1-S4 domains across the TRP channel family indicates that it is another key regulatory location.

The structural observations of related binding pockets for different activators across the TRP channel family may be connected to the conservation of mechanisms of channel activation between channels that exhibit different sensitivity to activating stimuli. For example, TRPV2 and TRPV3 are insensitive to the TRPV1-specific high-affinity vanilloid agonist Resiniferatoxin (RTx), but sensitivity to this vanilloid can be readily engineered into both TRPV channels by mutating only a few key residues within the vanilloid binding site (Yang, Vu et al. 2016 [DOI](#), Zhang, Hanson et al. 2016 [DOI](#), Zubcevic, Le et al. 2018 [DOI](#), Zhang, Swartz et al. 2019 [DOI](#)), revealing that the mechanisms responsible for coupling occupancy of the vanilloid site to channel opening are conserved in these three TRPV channels. In the present study, our aim was to explore the extent to which the cooling agent binding pocket described in TRPM8 is conserved in other TRPM channels and to then determine whether cooling agents can modulate channel opening in channels other than TRPM8. Our results suggest that the cooling agent binding pocket is well-conserved in several TRPM channels that were not previously reported to be sensitive to cooling agents and we find that icilin can bind to this well-conserved site in TRPM4 and promote opening of the pore by intracellular Ca^{2+} and alter the outwardly-rectifying properties of the channel.




Figure 2

Sequence and structure conservation of the icilin binding pocket in TRPM and TRPA channels

A) Structure-based sequence alignment of S1-S4 peripheral domains and TRP helix of selected TRP channel structures, with residues contributing to the icilin binding pocket in TRPM8 structures (7wre and 6nr3) highlighted in blue. The equivalent residues in other channels are colored according to the alignment quality score calculated from multiple sequence alignments, where highly conserved residues are color blue and poorly conserved residues are colored in white. Alignment quality score calculated in Jalview based on BLOSUM 62 scores (Henikoff and Henikoff 1992). Teal asterisks indicate Ca^{2+} coordinating residues in structures of TRPM channels. Black asterisks indicated Ca^{2+} coordinating residues in TRPA1. Red asterisks indicated residues where mutation influence cooling agent sensitivity in TRPM8. Gold asterisks indicate residues mutated in the present study. **B)** Chemical structure of icilin. **C)** S1-S4 residues contributing to the icilin binding pocket in TRPM8 structures (7wre and 6nr3) are shown as blue licorice, viewed from the intracellular side of the membrane as in Figure 1C, with the TRP box omitted for clarity. Cooling agent binding pocket mutations used in the present study are shown with carbon atoms colored gold and labeled in TRPM8 and TRPM4, and the equivalent residues in other channels are colored based on the alignment quality score, as in panel A. 7wre is mTRPM8, 6nr3 is faTRPM8 containing the A805G mutation, 6co7 is *Nematostella vectensis* TRPM2, 8ddr is mTRPM3, 6bqv is hTRPM4 and 7mbq is zebra fish TRPM5. Sequence identity between residues within the icilin binding pocket of TRPM8 and corresponding residues in the other TRP channel is as follows: TRPM5 (94%), TRPM4 (89%), TRPM2 (78%), TRPM3 and TRPM7 (44%), TRPA1 (22%), TRPV3 (11%).




Figure 2 - Figure Supplement 1

S1-S4 residues contributing to the icilin binding pocket in TRPM8 structures 7wre and 6nr3 are shown as blue licorice, viewed from the intracellular side of the membrane, with the TRP box omitted for clarity. Cooling agent binding pocket mutations used in the present study are shown with carbon atoms colored gold and labeled in TRPM8 and TRPM4, and the equivalent residues in other channels are colored based on the alignment quality score, as in [Figure 2A](#).

Results

Identification of residues lining the icilin binding pocket

We first investigated the icilin binding pocket using available structures of TRPM8 in complex with icilin and its obligate cofactor, Ca^{2+} (Yin, Wu et al. 2018 [\[1\]](#), Yin, Le et al. 2019 [\[2\]](#), Yin and Lee 2020 [\[3\]](#), Zhao, Xie et al. 2022 [\[4\]](#)). We defined the icilin binding pocket to include residues located near the icilin molecules in the available icilin-bound TRPM8 structures. With sufficient structural resolution, icilin's asymmetric arrangement of a hydroxyl group on one terminal benzene ring and a nitro group on the other benzene ring (**Figure 2** [\[5\]](#)) should facilitate identification of the ligand's physiological binding pose. Unfortunately, the electron densities attributed to icilin lack sufficient definition to fit these asymmetric functional groups in all but one of the five icilin-bound structures, even though the S1-S4 helices have relatively high local resolution in these structures. Interestingly, the structure with the most asymmetric icilin density (7wrd, with 2.98 Å overall nominal resolution) is not the highest resolution structure (7wre, with 2.52 Å overall nominal resolution). Previous comparison of these two binding poses has indicated that both poses are similarly plausible (Palchevskiy, Czarnocki-Cieciura et al. 2023 [\[6\]](#)), although the energy minimization performed in this study was conducted in the absence of the obligate cofactor, Ca^{2+} , which may be important for stabilizing icilin binding. We therefore decided to consider both icilin binding poses as possibly valid in defining the icilin binding pocket.

We identified the residues lining the icilin binding pocket based on residue proximity to icilin and on the ligand-protein interaction fingerprint generated by PoseFilter (Williams and Kalyanamoorthy 2021 [\[7\]](#)), which classifies interactions between the protein and ligand based on their chemistry in addition to proximity. We chose to use the most inclusive definition of binding pocket lining residues, counting any residue that was within 4 Å of the icilin molecule in either pose (6nr3 or 7wre), which also included all interacting residues identified by PoseFilter (**Figure 2** [\[5\]](#)). In addition, we included the Ca^{2+} coordinating residues in the binding pocket because Ca^{2+} is an essential cofactor and mutations in Ca^{2+} coordinating residues have been described to specifically disrupt icilin sensitivity in TRPM8 (Chuang, Neuhausser et al. 2004 [\[8\]](#), Winking, Hoffmann et al. 2012 [\[9\]](#), Kuhn, Winking et al. 2013 [\[10\]](#), Zhao, Xie et al. 2022 [\[4\]](#)).

Conservation of the icilin binding site in the Melastatin subfamily

To compare identified icilin binding pocket residues to other TRP channels, we utilized a structural alignment approach as described previously (Huffer, Aleksandrova et al. 2020 [\[11\]](#)), but expanding the scope of analysis to include a total of 264 structures of TRP channels determined to date. Comparing other Melastatin subfamily channels to TRPM8 revealed unexpectedly high conservation in the residues lining the cooling agent binding pocket in TRPM2 (78% identical), TRPM4 (89% identical) and TRPM5 (94% identical) (**Figure 2** [\[5\]](#)), channels that have not previously been reported to be sensitive to icilin. Mutations of residues in the structurally identified cooling agent binding pocket located within the S1-S4 domain of rTRPM8, including those corresponding to Y745 in S1, Q785 in S2, N799 and D802 in S3, R842 and H845 in S4, and Y1005 in the TRP box, are known to functionally influence cooling agent sensitivity in mTRPM8, rTRPM8, hTRPM8, *Parus major* TRPM8 (pmTRPM8) or faTRPM8 (Chuang, Neuhausser et al. 2004 [\[8\]](#), Bandell, Dubin et al. 2006 [\[12\]](#), Voets, Owsianik et al. 2007 [\[13\]](#), Malkia, Pertusa et al. 2009 [\[14\]](#), Winking, Hoffmann et al. 2012 [\[9\]](#), Kuhn, Winking et al. 2013 [\[10\]](#), Beccari, Gemei et al. 2017 [\[15\]](#), Yin, Wu et al. 2018 [\[1\]](#), Diver, Cheng et al. 2019 [\[16\]](#), Yin, Le et al. 2019 [\[2\]](#), Yin and Lee 2020 [\[3\]](#), Plaza-Cayon, Gonzalez-Muniz et al. 2022 [\[17\]](#), Zhao, Xie et al. 2022 [\[4\]](#)). These important residues (**Figure 2A; red asterisks** [\[5\]](#)), along with other residues located within 4 Å of icilin or Ca^{2+} in TRPM8 (**Figure 2A; blue highlighting** [\[5\]](#)), are highly conserved between TRPM8 and TRPM2, TRPM4, and TRPM5. These channels also show structural similarity in the shape of the pocket and orientation of the equivalent residues (**Figure 2C** [\[5\]](#)). Although a recently reported structure of TRPM4 prepared at physiological temperatures

reveals interesting structural changes within the intracellular melastatin domains compared to earlier structures, the structure of the cooling agent binding pocket is very similar (Hu, Park et al. 2024). The one notable and important difference near the cooling agent binding pocket between TRPM8, TRPM2, TRPM4 and TRPM5 is at the position corresponding to G805 in rTRPM8, a position that is conserved in mammalian TRPM8 channels that are sensitive to icilin, but is substituted by an Ala in avian TRPM8 channels that are insensitive to icilin (Chuang, Neuhausser et al. 2004). Because this critical position in TRPM8 is an Ala in most other TRPM channels (Figure 2), we predicted that TRPM2, TRPM4, and TRPM5 might be insensitive to icilin but could be rendered sensitive to icilin by substituting a Gly, similar to the icilin-sensitizing mutants engineered into avian TRPM8 channels (Chuang, Neuhausser et al. 2004, Yin, Le et al. 2019). In the case of TRPM3 and TRPM7, in addition to lacking the critical Gly residue, there is less conservation in the icilin- and Ca^{2+} - adjacent residues (44% identity in both cases) (Figure 2), suggesting that these channels may not be sensitive to icilin. TRPM1 and TRPM6, which were not included in the structural alignment because structures are not available, were predicted to behave similarly to TRPM3 and TRPM7 based on conventional sequence alignments in the S1-S4 region (not shown). Interestingly, TRPA1 and TRPV3, which have both been previously described to be sensitive to icilin (Story, Peier et al. 2003, Doerner, Gisselmann et al. 2007, Sherkheli, Vogt-Eisele et al. 2010, Sherkheli, Gisselmann et al. 2012, Billen, Brams et al. 2015), do not show homology in the equivalent binding pocket (Figure 2; Figure 2 – Figure Supplement 1; see Discussion).

Characterization of TRPM4 sensitivity to icilin

Based on the observed structural conservation, we considered attempting to engineer icilin sensitivity into TRPM2, TRPM4, or TRPM5. TRPM2 requires co-activation by intracellular Ca^{2+} and ADP ribose (ADPR) and its permeability to Ca^{2+} alters the local intracellular Ca^{2+} concentration (Perraud, Fleig et al. 2001, Sano, Inamura et al. 2001), making TRPM2 more challenging to study. In contrast, TRPM4 and TRPM5 are monovalent-selective and are activated by intracellular Ca^{2+} binding to the conserved Ca^{2+} binding site near the cooling agent binding pocket (Launay, Fleig et al. 2002, McKemy, Neuhausser et al. 2002, Hofmann, Chubanov et al. 2003, Liu and Liman 2003, Prawitt, Montellh-Zoller et al. 2003, Story, Peier et al. 2003, Andersson, Chase et al. 2004, Chuang, Neuhausser et al. 2004, Yamaguchi, Tanimoto et al. 2019). In both TRPM4 and TRPM5, Ca^{2+} dependence is also conferred by a second, intracellular Ca^{2+} binding site unrelated to the Ca^{2+} binding site near the cytoplasmic entrance of the cooling agent binding pocket in TRPM8 (Ruan, Haley et al. 2021, Hu, Park et al. 2024). At the time this study was initiated, the presence of this second Ca^{2+} site in TRPM4 was not appreciated, so we chose to focus on TRPM4 as a simpler system, based on the fact that it is relatively impermeable to Ca^{2+} (Launay, Fleig et al. 2002) and the assumption that activation is controlled by a single stimulus binding to a single binding site in each subunit. TRPM4 is also widely expressed in the body and plays important physiological roles in the cardiovascular, immune, endocrine, and nervous systems (Hasan and Zhang 2018, Wang, Xu et al. 2019). Pharmacological modulators of TRPM4 have been explored for a variety of conditions, including cancer, stroke, multiple sclerosis and heart disease (Bianchi, Smith et al. 2018, Dienes, Kovacs et al. 2021, Kovacs, Dienes et al. 2022).

We began by characterizing WT mouse TRPM4 (mTRPM4) using the inside-out configuration of the patch-clamp recording technique (Hamill, Marty et al. 1981) with Na^+ as the primary charge carrier in both intracellular and extracellular solutions. As described previously (Launay, Fleig et al. 2002, Nilius, Prenen et al. 2003, Ullrich, Voets et al. 2005, Zhang, Okawa et al. 2005), we observed that mTRPM4 is activated by intracellular Ca^{2+} in a concentration-dependent fashion and exhibits an outwardly-rectifying current-voltage (I-V) relation in the presence of Ca^{2+} (Figure 3). Partial rundown during the first Ca^{2+} application in each patch was observed, consistent with activity-dependent PIP2 depletion previously reported (Zhang, Okawa et al. 2005, Nilius, Mahieu et al. 2006, Guo, She et al. 2017), and the Ca^{2+} sensitivity we measured after the first Ca^{2+} application is similar to previously reported EC_{50} values for TRPM4 in inside-out patches that have not been supplemented with PIP2 (Zhang, Okawa et al. 2005, Nilius, Mahieu et al. 2006, Guo, She et al. 2017).




Figure 3

WT TRPM4 is sensitive to intracellular Ca^{2+} , voltage and icilin

A) Sample current families obtained using a holding voltage of -60 mV with 200 msec steps to voltages between -100 mV and +160 mV (Δ 20 mV) before returning to -60 mV. Control traces in the left column were obtained with TRPM4 in the absence of icilin and the presence of the labeled Ca^{2+} concentrations, and traces in the right column were obtained in the presence of 25 μM icilin and the labeled Ca^{2+} concentrations. **B)** Normalized I-V and **C)** normalized G-V plots for populations of cells in the absence (left, triangles) or presence (right, circles) of 25 μM icilin. Conductance values were obtained from tail current measurements. For each cell, values are normalized to the steady-state current or conductance at +160 mV in the presence of 500 μM Ca^{2+} . Error bars indicate standard error of the mean.

We first tested the sensitivity of mTRPM4 to icilin and found that the channel was not activated by 25 μM icilin applied alone over a wide range of voltages (**Figure 3A-C**). Icilin sensitivity in TRPM8 is known to require Ca^{2+} as a co-agonist, so we next tested whether icilin affects the activation of mTRPM4 by different concentrations of Ca^{2+} . In contrast to our prediction that TRPM4 would be insensitive to icilin based on G805 in TRPM8 not being conserved in TRPM4 (A867), we observed that icilin potentiates the activation mTRPM4 by intracellular Ca^{2+} (**Figure 3A-C**). This potentiation occurs at both positive and negative voltages at subsaturating Ca^{2+} concentrations (500 μM). In contrast, at saturating concentrations of Ca^{2+} (3 mM) (Nilius, Mahieu et al. 2006, Guo, She et al. 2017), minimal potentiation is observed at positive voltages, but notable potentiation is observed at negative voltages in both I-V and conductance-voltage (G-V) relations, indicating that icilin also diminishes the extent of outward-rectification (**Figure 3A-C**). Icilin also changes the kinetics of TRPM4 activation by voltage steps by enhancing the fraction of current elicited instantaneously after voltage steps and diminishing the fraction that activates more slowly on the timescale of 100-200 ms (**Figure 3A**). To quantify this, we defined steady-state currents (I_{ss}) observed at the end of 200 ms voltage steps to a maximally-activating voltage of +160 mV as the sum of current that activates instantaneously upon depolarization (I_{inst}) and the relaxing current that slowly activates during the voltage step (**Figure 4A**). The fraction of instantaneous current ($I_{\text{inst}}/I_{\text{ss}}$) increased with both Ca^{2+} and icilin (**Figure 4B**), indicating that both stimuli diminish outward-rectification. We also noticed that closure of TRPM4 channels following removal of both Ca^{2+} and icilin appeared to be slower compared to when the channel was only activated by Ca^{2+} alone (**Figure 4C,D**). From these results, we concluded that icilin interacts with TRPM4 and modulates activation of the channel by intracellular Ca^{2+} .

We also tested whether icilin modulates TRPM3 as several of the residues in TRPM8 that are critical for activation by cooling agents are not conserved in TRPM3. TRPM3 is activated by pregnenolone sulfate (PS) and exhibits an outward-rectifying I-V relationship with that agonist (Oberwinkler, Lis et al. 2005, Vriens, Owsianik et al. 2011, Held, Kichko et al. 2015), a property that is also common to both TRPM8 and TRPM4 in response to their activators. Because TRPM3 is inhibited by external Na^+ ions (Oberwinkler, Lis et al. 2005), similar to TRPV1 (Jara-Oseguera, Bae et al. 2016), we used Cs^+ as the primary charge carrier and recorded the activity of TRPM3 over a wide range of voltages before and after application of PS. Application of 25 μM icilin did not appear to activate TRPM3 when applied alone and we observed outwardly-rectifying currents in response to PS application, but these were not detectably altered by the prior application of icilin (**Figure 4 – Figure Supp. 1**).

TRPM4 mutations altering the effects of icilin

We next tested whether the icilin sensitivity observed in TRPM4 is mediated by binding of the cooling agent to the site identified in TRPM8. We first identified cooling agent binding pocket residues where mutations have been shown to specifically affect icilin sensitivity in TRPM8 without disrupting Ca^{2+} activation of TRPM4. Although some of the previously identified mutations like N799 and D802 in rTRPM8 and hTRPM8 disrupt icilin sensitivity (Chuang, Neuhausser et al. 2004, Winking, Hoffmann et al. 2012, Kuhn, Winking et al. 2013, Beccari, Gemei et al. 2017), they likely do so by disrupting binding of the obligate cofactor Ca^{2+} (Yin, Le et al. 2019, Zhao, Xie et al. 2022) and mutations in the equivalent Ca^{2+} -coordinating residues in rTRPM4 (N859, D862) have been shown to decrease Ca^{2+} affinity (Yamaguchi, Tanimoto et al. 2019). In contrast, one of the key determinants of icilin sensitivity in rTRPM8 is G805 (Chuang, Neuhausser et al. 2004), which is directly adjacent to L806 within the icilin binding pocket even though G805 itself is not within 4 Å of icilin in the available structures (**Figure 2**). G805 in rTRPM8 corresponds to an Ala in icilin-insensitive chicken TRPM8 (cTRPM8), and the G805A mutation decreases the sensitivity of rTRPM8 to icilin while the inverse Ala to Gly mutation in cTRPM8 or faTRPM8 introduces sensitivity to icilin (Chuang, Neuhausser et al. 2004, Yin, Le et al. 2019). We hypothesized that because mTRPM4 is already sensitive to icilin, making the equivalent A867G mutation might further enhance the sensitivity of the channel to icilin.




Figure 4

Icilin modulates voltage-dependent activation and closure kinetics of TRPM4

A) Sample current traces illustrating the fraction of current that activates rapidly (I_{inst}) compared to the steady-state current at the end of the pulse (I_{SS}). The pulse protocols used a holding voltage of -60 mV with 200 msec steps to +160 mV in the presence of varying concentrations of intracellular Ca^{2+} . Traces were obtained in the absence (left) or presence (right) of 25 μ M icilin. **B)** Instantaneous fraction of current (I_{inst}/I_{SS}) calculated using voltage steps to +160 mV at various concentrations of intracellular Ca^{2+} for individual cells in the absence (left, triangles) or presence (right, circles) of 25 μ M icilin. Error bars indicate standard error of the mean. **C)** Fraction of current remaining after application of 3 mM Ca^{2+} alone (triangles) or both 3 mM Ca^{2+} and 25 μ M icilin (squares) for WT TRPM4. Currents were elicited by voltage steps from -100 to +100 mV. **D)** Fraction of current remaining 14 seconds after removal of 3 mM Ca^{2+} alone (triangles) or both 3 mM Ca^{2+} and 25 μ M icilin (squares) for WT TRPM4. Currents were elicited by voltage steps from -100 mV to +160 mV.




Figure 4 – Figure Supplement 1

Icilin does not modulate voltage-dependent activation of TRPM3 α 2.

A) Sample current families obtained using a holding voltage of -60 mV with 50 msec prepulses to -150 mV before 100 msec steps to voltages between -160 mV and +160 mV (Δ 20 mV), and then returning to -60 mV. Control traces in the left column were obtained with TRPM3 α 2 in the absence of icilin (left, triangles), and traces in the right column were obtained in the presence of 25 μM icilin (right, circles). Activation of TRPM3 current is achieved with 40 μM pregnenolone sulphate (PregS, navy). **B**) Corresponding normalized I-V relations and **C**) plots of normalized current evoked at +160 mV in populations of cells in the absence (left, triangles) or presence (right, circles) of 25 μM icilin. Absolute values were obtained from steady-state current measurements. For each cell, values are normalized to the absolute current at +160 mV in the presence of 40 μM PregS. Error bars indicate standard error of the mean.

We first tested whether the A867G mutation in mTRPM4 alters channel activation and observed Ca^{2+} sensitivity and outward-rectification (**Figure 5**) that was similar to WT (**Figure 3**). We then applied icilin and observed that, as observed in TRPM4 and TRPM8, icilin applied in the absence of Ca^{2+} was not sufficient to robustly activate the A867G mutant (**Figure 5A-C**). When applied in the presence of intracellular Ca^{2+} , icilin potentiated Ca^{2+} -evoked currents even more dramatically in the A867G mutant than in the WT channel (**Figure 5**). Notably, maximal currents were observed by application of 500 μM Ca^{2+} in the presence of icilin, which is a subsaturating concentration in the WT channel, even in the presence of icilin (**Figure 3**). Icilin also diminished the extent of outward-rectification dramatically in the A867G mutant channel, which was observed as particularly dramatic potentiation of currents elicited at negative voltages compared to positive voltages and a nearly linear I-V relation at a saturating concentration of Ca^{2+} (**Figure 5**). This loss of outward-rectification could also be observed by quantifying the fraction of current activated instantaneously, where A867G mutant channels showed Ca^{2+} -dependent increases in $I_{\text{inst}}/I_{\text{ss}}$ similar to WT in the absence of icilin, but $I_{\text{inst}}/I_{\text{ss}}$ values near 1 in the presence of 25 μM icilin (**Figure 6**). The A867G mutation does not modulate TRPM4 sensitivity to Ca^{2+} applied alone, indicating that this mutation specifically altered modulation by icilin. The selective influence of the A867G mutation on icilin may be related to the observation in TRPM8 where the G805A mutation affects icilin sensitivity without altering sensitivity to menthol (Chuang, Neuhauser et al. 2004). We also noticed that closure of TRPM4 following removal of both Ca^{2+} and icilin was considerably slower in the A867G mutant compared to the WT channel (**Figure 6C,D**), suggesting that the mutant binds icilin with higher affinity than the WT channel and that the slowing of channel closure is due to slower icilin unbinding (**Figure 4C,D**). Taken together, the effects of the A867G mutation are consistent with the icilin sensitivity of TRPM4 being mediated by the equivalent binding pocket as in TRPM8.

Another critical determinant of icilin sensitivity in hTRPM8 is R842 (**Figure 2**), a residue where mutations to His dramatically diminish activation by the cooling agents icilin and menthol and that is within 4 Å of key substituent groups in icilin in the two possible docking orientations of the cooling agent (Voets, Owsianik et al. 2007, Palchevskiy, Czarnocki-Cieciora et al. 2023). If icilin binds to the equivalent pocket in TRPM4 as in TRPM8, we hypothesized that the equivalent mutation in TRPM4 (R901H) would disrupt icilin sensitivity. As with the A867G mutant, we began by testing whether R901H in TRPM4 exhibits similar behavior to the WT channel and observed Ca^{2+} sensitivity similar to WT TRPM4, though it notably enhanced outward-rectification (**Figure 7**). The R901H mutant was insensitive to icilin applied alone, as observed for WT and the A867G mutant in mTRPM4, however, unlike WT and the A867G mutant in mTRPM4, activation of the R901H mutant is not enhanced by icilin, regardless of whether subsaturating or saturating concentrations of intracellular Ca^{2+} are tested (**Figure 7**), and icilin does not alter the instantaneously-activating fraction of current or prolong channel closure (**Figure 8**). These results in the R901H mutant of TRPM4 are consistent with the loss of icilin sensitivity in R842H mutant of rTRPM8, providing further evidence that icilin binds to the conserved binding pocket in hTRPM4 to modulate channel activation by intracellular Ca^{2+} .

Discussion

The objective of the present study was to explore the conservation of the cooling-agent binding pocket within TRPM channels. This site has been carefully interrogated in TRPM8, where structures are available with different cooling agents bound and where extensive mutagenesis has determined the influence of key residues in activation of the channel by cooling agents (Chuang, Neuhauser et al. 2004, Bandell, Dubin et al. 2006, Voets, Owsianik et al. 2007, Malkia, Pertusa et al. 2009, Winking, Hoffmann et al. 2012, Kuhn, Winking et al. 2013, Beccari, Gemei et al. 2017, Yin, Wu et al. 2018, Diver, Cheng et al. 2019, Yin, Le et al. 2019, Yin and Lee 2020, Plaza-Cayon, Gonzalez-Muniz et al. 2022, Zhao, Xie et al. 2022). In TRPM8, the cooling agent binding pocket is contained within the S1-S4 domain and opens into the intracellular side of




Figure 5

A867G mutant TRPM4 retains sensitivity to Ca^{2+} and voltage, but has enhanced sensitivity to icilin

A) Sample current families obtained using a holding voltage of -60 mV with 200 msec steps to voltages between -100 mV and +160 mV (Δ 20 mV) before returning to -60 mV. Control traces in the left column were obtained with A867G mTRPM4 in the absence of icilin and the presence of the labeled Ca^{2+} concentrations, and traces in the right column were obtained in the presence of 25 μM icilin and the labeled Ca^{2+} concentrations. **B)** Normalized I-V and **C)** normalized G-V plots for populations of cells in the absence (left, triangles) or presence (right, circles) of 25 μM icilin. Conductance values were calculated from steady-state currents. For each cell, values are normalized to the steady-state current or conductance at +160 mV in the presence of 500 μM Ca^{2+} . Error bars indicate standard error of the mean.




Figure 6

Icilin modulation of TRPM4 is enhanced in the A867G mutant.

A) Sample current traces illustrating the fraction of current that activates rapidly (I_{inst}) compared to the steady-state current at the end of the pulse (I_{SS}). The pulse protocols used a holding voltage of -60 mV with 200 msec steps to +160 mV in the presence of varying concentrations of intracellular Ca^{2+} . Traces were obtained in the absence (left) or presence (right) of 25 μM icilin. **B**) Instantaneous fraction of current ($I_{\text{inst}}/I_{\text{SS}}$) calculated using +160 mV voltage steps at various concentrations of intracellular Ca^{2+} for individual cells in the absence (left, triangles) or presence (right, circles) of 25 μM icilin. Error bars indicate standard error of the mean. **C**) Fraction of current remaining after application of 3 mM Ca^{2+} alone (triangles) or both 3 mM Ca^{2+} and 25 μM icilin (squares) for WT TRPM4 (gray) or A867G TRPM4 (purple). Currents were elicited by voltage steps from -100 to +100 mV. **D**) Fraction of current remaining 14 seconds after removal of 3 mM Ca^{2+} alone (triangles) or both 3 mM Ca^{2+} and 25 μM icilin (squares) for WT (left) or A867G TRPM4 (right). Currents were elicited by voltage steps from -100 mV to +160 mV.




Figure 7

R901H mutant TRPM4 is sensitive to intracellular Ca^{2+} and voltage, but icilin does not promote opening

A) Sample current families obtained using a holding voltage of -60 mV with 200 msec steps to voltages between -100 mV and +160 mV (Δ 20 mV) before returning to -60 mV. Control traces in the left column were obtained with R901H mTRPM4 in the absence of icilin and the presence of the labeled Ca^{2+} concentrations, and traces in the right column were obtained in the presence of 25 μM icilin and the labeled Ca^{2+} concentrations. For the cell shown, current families were not obtained in the presence of icilin and the absence of Ca^{2+} . **B)** Normalized I-V and **C)** normalized G-V plots for populations of cells in the absence (left, triangles) or presence (right, circles) of 25 μM icilin. Conductance values were obtained from tail current measurements. For each cell, values are normalized to the steady-state current or conductance at +160 mV in the presence of 500 μM Ca^{2+} . Error bars indicate standard error of the mean.




Figure 8

Icilin modulation of the voltage-dependent activation of TRPM4 is disrupted in the R901H mutant

A) Sample current traces illustrating the fraction of current that activates rapidly (I_{inst}) compared to the steady-state current at the end of the pulse (I_{ss}). The pulse protocols used a holding voltage of -60 mV with 200 msec steps to +160 mV in the presence of varying concentrations of intracellular Ca^{2+} . Traces were obtained in the absence (left) or presence (right) of 25 μ M icilin. For the cell shown, current families were not obtained in the presence of icilin and absence of Ca^{2+} . **B)** Instantaneous fraction of current (I_{inst}/I_{ss}) calculated using +160 mV voltage steps at various concentrations of intracellular Ca^{2+} for individual cells in the absence (left, triangles) or presence (right, circles) of 25 μ M icilin. Error bars indicate standard error of the mean. **C)** Fraction of current remaining after 14 seconds of 0 mM Ca^{2+} wash, following removal of 3 mM Ca^{2+} alone (triangles) or both 3 mM Ca^{2+} and 25 μ M icilin (squares) for WT (left) or R901H TRPM4 (right). Currents were measured between -100 mV and +160 mV (Δ 20 mV), but only +160 mV current fractions are shown.

the membrane where a Ca^{2+} ion binding site is located. Our analysis across the available TRPM structures shows that the cooling agent binding pocket in TRPM8 is well-conserved in TRPM2, TRPM4 and TRPM5, but not in TRPM3 or TRPM7 (**Figure 2**). Using functional approaches, we explored the interaction of the icilin with TRPM4 and found that the channel can be modulated by icilin (**Figure 3**; **Figure 4**). Two hallmark features of TRPM4 are that it is activated by intracellular Ca^{2+} and that its I-V relation is outwardly-rectifying, permeating cations out of the cell considerably more favorably than into the cell (**Figure 3**, **5**, **7**). We found that icilin enhanced the sensitivity of the channel to intracellular Ca^{2+} and reduced the extent of outward-rectification such that the channel conducts cations into the cell considerably more favorably at negative membrane voltages (**Figure 3**; **Figure 4**). Mutations established to promote or disrupt the actions of icilin in TRPM8 had similar effects in TRPM4 (**Figure 5**–**8**), leading us to conclude that icilin is likely binding to a conserved cooling agent binding pocket in both channels. Our analysis of TRPM3 channels revealed that key residues within the cooling agent binding pocket of TRPM8 are not conserved in TRPM3 (**Figure 2**) and our results also show that icilin does not detectably alter activation by PS (**Figure 4** – **Figure Supp. 1**), providing a negative control for our findings with TRPM4. Collectively, these findings demonstrate that not only is the cooling agent binding pocket conserved between TRPM8 and TRPM4, but that coupling between occupancy of that pocket and opening of the channel are also conserved in both TRPM channels. The conserved nature of the coupling mechanism is particularly interesting when one considers that although the cooling agent binding pocket is highly conserved (89% identity), the S1-S6 regions exhibit much lower conservation (34% identity).

The identification of a conserved S1-S4 binding pocket in TRPM4 and TRPM8 raises many questions that would be interesting to interrogate in future studies. For example, there are many structurally distinct cooling agents that have been developed as agonists of TRPM8, some of which require Ca^{2+} as a co-agonist and some that do not, and there are many antagonists thought to bind within the cooling agent binding pocket in TRPM8 (Sherkheli, Vogt-Eisele et al. 2010, De Petrocellis, Ortar et al. 2015, Diver, Cheng et al. 2019, Gonzalez-Muniz, Bonache et al. 2019, Yin, Le et al. 2019, Yin and Lee 2020, Plaza-Cayon, Gonzalez-Muniz et al. 2022, Yin, Zhang et al. 2022, Zhao, Xie et al. 2022). It would be interesting to further explore the extent to which these ligands discriminate between the two channels and to try and solve structures of both TRPM4 with some of these ligands bound to determine the extent to which binding poses are similar. Our analysis of the cooling agent binding pocket in TRPM channels suggests that both TRPM2 and TRPM5 might also be sensitive to icilin or related compounds because most of the key residues are conserved, and these would be interesting directions to explore. Also, our experiments were done after depletion of PIP2 with the first intracellular Ca^{2+} application, which leads to partial rundown of the channel and a pronounced lowering of Ca^{2+} affinity (Zhang, Okawa et al. 2005, Nilius, Mahieu et al. 2006, Guo, She et al. 2017). It would be fascinating to investigate the extent to which PIP2 modulates the action of icilin on TRPM4. For example, the dramatic influence of PIP2 on Ca^{2+} sensitivity might enhance the affinity of icilin, which we only studied at a relatively high concentration that is saturating for TRPM8 (Andersson, Chase et al. 2004, Chuang, Neuhauser et al. 2004).

The conservation of cooling agent binding pockets in TRPM2, TRPM4 and TRPM5, and the finding that icilin can modulate TRPM4 is interesting because two other TRP channels have been reported to be sensitive to icilin even though our analysis shows that the cooling agent binding pocket in TRPM8 is not conserved in those channels. The first example is TRPA1, which can be activated by icilin even though the residues in TRPA1 correspond to those that form the cooling agent binding pocket in TRPM8 are poorly conserved (**Figure 2**). As observed for TRPM8, TRPA1 icilin sensitivity has been reported to require coactivation with Ca^{2+} (Doerner, Gisselmann et al. 2007), and the S1-S4 Ca^{2+} binding site observed in TRPMs is indeed conserved in TRPA1, though the S2 Ca^{2+} -coordinating residues are located one helical turn lower in TRPA1 when its structures are aligned with TRPM8 (**Figure 2**). Ca^{2+} acts directly to activate and inactivate TRPA1 at low and high concentrations, respectively (Wang, Chang et al. 2008, Hasan and Zhang 2018), and

can also act indirectly on TRPA1 through Ca^{2+} -binding proteins such as calmodulin (Hasan, Leeson-Payne et al. 2017 [DOI](#)). Mutations in this TRPA1 transmembrane Ca^{2+} binding site do affect Ca^{2+} potentiation and desensitization (Zimova, Sinica et al. 2018 [DOI](#), Zhao, Lin King et al. 2020 [DOI](#)). However, it is unclear whether this conserved transmembrane Ca^{2+} binding site directly mediates icilin sensitivity in TRPA1 as it does in TRPM8 because multiple other Ca^{2+} binding sites have been proposed in TRPA1, including EF hand motifs near the N-terminus (Doerner, Gisselmann et al. 2007 [DOI](#), Zurborg, Yurgionas et al. 2007 [DOI](#)), residues within the transmembrane helices S2 and S3 (Zhao, Lin King et al. 2020 [DOI](#)), and acidic residues at the C-terminus (Sura, Zima et al. 2012 [DOI](#)). Overall, the lack of conservation of the TRPM8 cooling agent binding pocket in TRPA1 raises the possibility that icilin binds to a different location to regulate the activity of that TRP channel.

A second example is TRPV3, which can be inhibited by icilin in a calcium-dependent manner (Sherkheli, Gisselmann et al. 2012 [DOI](#), Billen, Brams et al. 2015 [DOI](#)). It is unclear where icilin binds to TRPV3, though tryptophan fluorescence quenching experiments on mutants have revealed that residues W481 at the top of S2, W559 in the middle of S4, and W710 after the TRP box in TRPV3 are not directly involved in icilin binding (Billen, Brams et al. 2015 [DOI](#)). The residues equivalent to the icilin binding pocket in TRPM8 are not conserved in TRPV3 and the transmembrane Ca^{2+} ion binding site observed in TRPM channels and TRPA1 is also not conserved in TRPV3 (**Figure 2C**), suggesting that the observed Ca^{2+} dependence of the icilin sensitivity in TRPV3 is mediated by a different part of the channel, possibly through direct binding of Ca^{2+} ions or indirectly through calmodulin binding (Xiao, Tang et al. 2008 [DOI](#), Phelps, Wang et al. 2010 [DOI](#)). An arginine in the TRP box and an aspartate in the pore loop have been implicated in the Ca^{2+} dependence of TRPV3 and other TRPV channels (Garcia-Martinez, Morenilla-Palao et al. 2000 [DOI](#), Voets, Prenen et al. 2002 [DOI](#), Chung, Lee et al. 2004 [DOI](#), Xiao, Tang et al. 2008 [DOI](#), Hu, Grandl et al. 2009 [DOI](#), Xiao et al., 2008 [DOI](#)), so these would be potential sites to explore. As in TRPA1, the lack of conservation of the cooling agent binding pocket found in TRPM8 with the corresponding region of TRPV3 raises the possibility that icilin and Ca^{2+} bind to different regions in TRPV3 compared to TRPM channels.

The discovery that the cooling agent icilin can interact with TRPM4 and modulate its activation by Ca^{2+} and voltage has several important implications for understanding key mechanisms in TRP channels. First, it provides another clear example of two TRP channels that are activated by very different stimuli, cooling agents and noxious cold in the case of TRPM8 and intracellular Ca^{2+} in the case of TRPM4, but that share common underlying mechanisms wherein binding sites for ligands within the S1-S4 domain are coupled to channel opening. Conceptually, our findings with TRPM4 relate to those in TRPV channels, where sensitivity to activation by vanilloids can be engineered into the vanilloid-insensitive TRPV2 and TRPV3 channels with relatively few mutations within the pocket corresponding to where vanilloids bind in TRPV1 (Yang, Vu et al. 2016 [DOI](#), Zhang, Hanson et al. 2016 [DOI](#), Zubcevic, Le et al. 2018 [DOI](#), Zhang, Swartz et al. 2019 [DOI](#)). In the case of TRPV2 and TRPV3, the ability to open in response to occupancy of the vanilloid binding site is intrinsic to those channels and requires only sculpting of the vanilloid site to enable ligand binding. Although we imagined that we might need to undertake similar engineering of TRPM4 to make it sensitive to cooling agents given the presence of an Ala at 867, we instead discovered intrinsic sensitivity to icilin that could be further tuned by the A867G mutation. It is fascinating that coupling between two distinct types of ligand binding pockets and channel opening are conserved in these examples of TRPV and TRPM channels, raising the possibility that key mechanisms of channel gating will turn out to be shared features of other TRP channels.

Second, uncovering a relationship between TRPM4 and TRPM8 provides motivation to explore the mechanistic relationships between TRPM channels further because their functional properties can vary quite dramatically, from being heat activated in TRPM3 (Vriens, Owsianik et al. 2011 [DOI](#), Vandewauw, De Clercq et al. 2018 [DOI](#)) to cold activated in TRPM8 (McKemy, Neuhauser et al. 2002 [DOI](#), Peier, Moqrich et al. 2002 [DOI](#), Bautista, Siemens et al. 2007 [DOI](#), Dhaka, Murray et al. 2007 [DOI](#)). TRPM channels also exhibit varying rectification properties, for example, with TRPM4 exhibiting pronounced outward-rectification at all Ca^{2+} concentrations (Launay, Fleig et al. 2002 [DOI](#), Nilius,

Prenen et al. 2003 [↗](#), Ullrich, Voets et al. 2005 [↗](#), Zhang, Okawa et al. 2005 [↗](#)) and TRPM5 exhibiting outward-rectification at low Ca^{2+} concentrations and a nearly linear I-V relationship at high concentrations (Liu and Liman 2003 [↗](#), Yamaguchi, Tanimoto et al. 2019 [↗](#), Ruan, Haley et al. 2021 [↗](#)). Here we show that TRPM4 can exhibit these two extremes of rectification, from outwardly-rectifying in response to Ca^{2+} alone to a linear I-V in the presence of Ca^{2+} and icilin (Figure 3 [↗](#), 5 [↗](#)). Related differences in rectification in TRPM3 when activated by different agonists have factored into proposals for distinct ion permeation pathways; a conventional pore at the central axis and second within the S1-S4 domains (Vriens, Held et al. 2014 [↗](#)). Our observation that cooling agents can modulate the rectifying properties of TRPM4 quite dramatically may be mechanistically related to observations in TRPM3, but at least in TRPM4 it's hard to imagine how ions could conceivably permeate through the S1-S4 domain with icilin bound (Figure 1 [↗](#), 2 [↗](#)). A more likely explanation is that cooling agent binding to TRPM4 not only couples to channel opening, but also to a voltage-dependent mechanism that underlies rectification. It will be fascinating to explore how Ca^{2+} and cooling agent binding in such close proximity within the S1-S4 domain of TRPM4 can have such divergent effects on the mechanism of rectification.

Third, the discovery that cooling agents can bind to and modulate TRPM4 raises intriguing questions about why this site is conserved in so many TRPM channels. One possible answer is that there are endogenous, yet to be identified molecules that can bind to the cooling agent binding site to regulate the activity of TRPM channels. Conceptually, a role of endogenous modulators binding to the cooling agent binding pocket is similar to lipids occupying the vanilloid binding site in TRPV channels to stabilize the closed state of the channel, which are displaced when activators bind to activate the channel (Cao, Cordero-Morales et al. 2013 [↗](#), Gao, Cao et al. 2016 [↗](#), Nadezhdin, Neuberger et al. 2021 [↗](#), Zhang, Julius et al. 2021 [↗](#), Arnold, Mancino et al. 2024 [↗](#), Nadezhdin, Neuberger et al. 2024 [↗](#)).

Finally, our findings here with TRPM4 may also be relevant for understanding the neuroprotective actions of icilin in mouse models of multiple sclerosis and seizures (Pezzoli, Elhamdani et al. 2014 [↗](#), Ewanchuk, Allan et al. 2018 [↗](#), Moriyama, Nomura et al. 2019 [↗](#)). Some of these neuroprotective effects remain in TRPM8 or TRPA1 knockout animals, raising the intriguing possibility that TRPM4 may underlie the actions of icilin in these disease models. Indeed, TRPM4 inhibitors and TRPM4 knockouts have been previously described to play a role in these same mouse models of multiple sclerosis, inviting further investigation of TRPM4 modulators in this disease (Bianchi, Smith et al. 2018 [↗](#)).

Materials and methods

Structure-based sequence alignment

Structural alignment of TRPM and TRPA1 structure TM domains (pre-S1-TRP box) was performed using Fr-TM-Align as described previously (Huffer, Aleksandrova et al. 2020 [↗](#)). Representative structures were selected based on nominal resolution. Sequence alignments were made in Jalview. Structures were visualized with PyMOL.

Channel Constructs

The mTRPM4b DNA plasmid tagged with EGFP at the C-terminus was a generous gift from Dr. Youxing Jiang (UT Southwestern) and was transfected into HEK293 cells using FuGENE6 transfection reagent. Mouse TRPM3a2 DNA in the bicistronic pCAGGS/IRES-GFP vector (Vriens, Held et al. 2014 [↗](#)) was provided by Dr. Thomas Voets (Catholic University, Leuven, Belgium) and transfected as for mTRPM4. All mutations in mTRPM4 were made using the QuikChange Lightning technique (Agilent Technologies) and confirmed by DNA sequencing (Macrogen).

Cell Culture

Human Embryonic Kidney (HEK293) cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 10 mg L⁻¹ of gentamicin. HEK293 cells between passage numbers 5-25 were used and passaged when cells were between 40-80% confluent. The cells were treated with trypsin and then seeded on glass coverslips at about 15% of the original confluency in 35 mm petri dishes. Transfections were done using the FuGENE6 Transfection Reagent (Promega). Transfected cells were incubated at 37°C with 95% air and 5% CO₂ overnight for use in patch-clamp recordings; 16-48h depending on the construct.

Electrophysiology

For recording the activity of TRPM4 in inside-out patches, the pipette (extracellular) solution contained 130 mM NaCl, 2 mM MgCl₂, 0.5 mM CaCl₂, 10 mM HEPES, with pH adjusted to 7.4 using NaOH. The bath (intracellular) solutions all contained 125 mM CsCl, 5 mM NaCl, 2 mM MgCl₂, 10 mM HEPES, with pH adjusted to 7.4 using CsOH. CaCl₂ was varied, and 0 mM CaCl₂ solutions contained 1 mM EGTA. For EGTA-containing solutions, MgCl₂ concentrations were adjusted using MaxChelator such that there was 2 mM free MgCl₂ (Bers, Patton et al. 2010 [\[1\]](#)). For recording the activity of TRPM3 in whole-cell recordings, the pipette (intracellular) solution contained 130 mM CsCl, 1 mM (free) MgCl₂, 10 mM HEPES, 10 mM EGTA, with pH adjusted to 7.2 using CsOH. The external (extracellular) solution contained 130 mM CsCl, 1 mM MgCl₂, 10 mM HEPES, with pH adjusted to 7.4 using CsOH. The bathing solution in which seals were obtained contained 130 mM NaCl, 1 mM MgCl₂, 10 mM HEPES, with pH adjusted to 7.4 with NaOH. Bath and ground chambers were connected by an agar bridge containing 3M KCl. Icilin powder (Sigma) was dissolved in DMSO to a stock concentration of 40 mM, aliquoted, and stored at -80°C. As icilin is known to degrade (Kuhn, Kuhn et al. 2009 [\[2\]](#)), the stock was not subjected to repeated freeze-thaw cycles. Cs⁺-based solutions increased successful Giga seal formation over Na-based solutions. K⁺ was omitted to prevent contamination of Kv channel currents. Mg²⁺ was added to all solutions to inhibit endogenous TRPM7 currents that were observed in some passages (Nadler, Hermosura et al. 2001 [\[3\]](#), Hermosura, Monteilh-Zoller et al. 2002 [\[4\]](#), Kozak and Cahalan 2003 [\[5\]](#)). Patch pipette resistance ranged from 1 to 5 MOhms, with a typical value around 3 MOhms. Inside-out patch clamp recordings were performed using a holding voltage of -60 mV with 200 msec steps to voltages between -100 and +160 mV (Δ20 mV). Electrophysiology data was acquired with an Axopatch 200B amplifier at a sampling frequency of 10 kHz and filtered to 5 kHz with a low-pass filter. Variable PIP₂-depletion-induced current rundown was observed (Nilius, Mahieu et al. 2006 [\[6\]](#)), so all measurements were taken after current had reached a steady state. Patches that did not exhibit response to Ca²⁺ were excluded because this either indicated that the pulled patch had formed a vesicle, precluding access to the intracellular face of the membrane, or that TRPM4 expression was too low for our experiments. For long timecourses with multiple exposures, only currents that returned to baseline upon removal of Ca²⁺ were considered. Because icilin is thought to partition into the membrane, all non-icilin traces included in population data for I-V and G-V relations were collected prior to the application of icilin, even if icilin appeared to wash during the experiment. Patches that exhibited very large currents (>5 nA) were also excluded because they would result in substantial voltage errors and/or changes in the concentrations of ions. Currents were normalized to steady-state currents obtained in the presence of 500 μM Ca²⁺ at +160 mV for each cell. Baseline current subtraction was performed for each cell by subtracting leak currents obtained in the absence of Ca²⁺.

Statistical methods were not used to determine the sample size. Sample size for electrophysiological studies was determined empirically by comparing individual measurements with population data obtained under differing conditions until convincing differences or lack thereof were evident. Additional exploratory experiments performed to determine ideal recording conditions are consistent with the data reported here, but these pilot data are not included in our analysis due to changes in experimental conditions, including varying voltage pulse protocols and

different solution compositions. For all electrophysiological experiments, n values represent the number of patches studied from between 9 and 31 different batches of cells. Electrophysiology data analysis and visualization was performed with IgorPro and Python (matplotlib).

Acknowledgements

We thank Andres Jara-Oseguera, Surbhi Dhingra, Ana I. Fernández-Mariño, Xiao-Feng Tan and members of the Swartz laboratory for helpful discussions. This research was supported by the Intramural Research Program of the National Institute of Neurological Disorders and Stroke, NIH, Bethesda, MD to KJS (NS003018).

References

- Andersson D. A., Chase H. W., Bevan S. (2004) **TRPM8 activation by menthol, icilin, and cold is differentially modulated by intracellular pH** *J Neurosci* **24**:5364–5369
- Arnold W. R., Mancino A., Moss F. R., Frost A., Julius D., Cheng Y. (2024) **Structural basis of TRPV1 modulation by endogenous bioactive lipids** *Nat Struct Mol Biol*
- Autzen H. E., Myasnikov A. G., Campbell M. G., Asarnow D., Julius D., Cheng Y. (2018) **Structure of the human TRPM4 ion channel in a lipid nanodisc** *Science* **359**:228–232
- Bai Y. *et al.* (2020) **Structural basis for pharmacological modulation of the TRPC6 channel** *eLife* **9**
- Bandell M., Dubin A. E., Petrus M. J., Orth A., Mathur J., Hwang S. W., Patapoutian A. (2006) **High-throughput random mutagenesis screen reveals TRPM8 residues specifically required for activation by menthol** *Nat Neurosci* **9**:493–500
- Bautista D. M., Siemens J., Glazer J. M., Tsuruda P. R., Basbaum A. I., Stucky C. L., Jordt S. E., Julius D. (2007) **The menthol receptor TRPM8 is the principal detector of environmental cold** *Nature* **448**:204–208
- Beccari A. R. *et al.* (2017) **Novel selective, potent naphthyl TRPM8 antagonists identified through a combined ligand- and structure-based virtual screening approach** *Sci Rep* **7**
- Bers D. M., Patton C. W., Nuccitelli R. (2010) **A practical guide to the preparation of Ca(2+) buffers** *Methods Cell Biol* **99**:1–26
- Bianchi B., Smith P. A., Abriel H. (2018) **The ion channel TRPM4 in murine experimental autoimmune encephalomyelitis and in a model of glutamate-induced neuronal degeneration** *Mol Brain* **11**
- Billen B. *et al.* (2015) **Different ligands of the TRPV3 cation channel cause distinct conformational changes as revealed by intrinsic tryptophan fluorescence quenching** *J Biol Chem* **290**:12964–12974
- Cabezas-Bratesco D., McGee F. A., Colenso C. K., Zavala K., Granata D., Carnevale V., Opazo J. C., Brauchi S. E. (2022) **Sequence and structural conservation reveal fingerprint residues in TRP channels** *eLife* **11**
- Cao E (2020) **Structural mechanisms of transient receptor potential ion channels** *J Gen Physiol* **152**
- Cao E., Cordero-Morales J. F., Liu B., Qin F., Julius D. (2013) **TRPV1 channels are intrinsically heat sensitive and negatively regulated by phosphoinositide lipids** *Neuron* **77**:667–679
- Cao E., Liao M., Cheng Y., Julius D. (2013) **TRPV1 structures in distinct conformations reveal activation mechanisms** *Nature* **504**:113–118

- Caterina M. J., Leffler A., Malmberg A. B., Martin W. J., Trafton J., Petersen-Zeitz K. R., Koltzenburg M., Basbaum A. I., Julius D. (2000) **Impaired nociception and pain sensation in mice lacking the capsaicin receptor** *Science* **288**:306–313
- Caterina M. J., Schumacher M. A., Tominaga M., Rosen T. A., Levine J. D., Julius D. (1997) **The capsaicin receptor: a heat-activated ion channel in the pain pathway** *Nature* **389**:816–824
- Chuang H. H., Neuhausser W. M., Julius D. (2004) **The super-cooling agent icilin reveals a mechanism of coincidence detection by a temperature-sensitive TRP channel** *Neuron* **43**:859–869
- Chung M. K., Lee H., Mizuno A., Suzuki M., Caterina M. J. (2004) **2-aminoethoxydiphenyl borate activates and sensitizes the heat-gated ion channel TRPV3** *J Neurosci* **24**:5177–5182
- Clapham D. E., Miller C. (2011) **A thermodynamic framework for understanding temperature sensing by transient receptor potential (TRP) channels** *Proc Natl Acad Sci U S A* **108**:19492–19497
- Cosens D. J., Manning A. (1969) **Abnormal electroretinogram from a Drosophila mutant** *Nature* **224**:285–287
- De Petrocellis L., Ortar G., Schiano Moriello A., Serum E. M., Rusterholz D. B. (2015) **Structure-activity relationships of the prototypical TRPM8 agonist icilin** *Bioorg Med Chem Lett* **25**:2285–2290
- Dhaka A., Murray A. N., Mathur J., Earley T. J., Petrus M. J., Patapoutian A. (2007) **TRPM8 is required for cold sensation in mice** *Neuron* **54**:371–378
- Diaz-Franulic I., Verdugo C., Gonzalez F., Gonzalez-Nilo F., Latorre R. (2021) **Thermodynamic and structural basis of temperature-dependent gating in TRP channels** *Biochem Soc Trans* **49**:2211–2219
- Dienes C., Kovacs Z. M., Hezso T., Almasy J., Magyar J., Banyasz T., Nanasi P. P., Horvath B., Szentandrassy N. (2021) **Pharmacological Modulation and (Patho)Physiological Roles of TRPM4 Channel-Part 2: TRPM4 in Health and Disease** *Pharmaceuticals (Basel)* **15**
- Diver M. M., Cheng Y., Julius D. (2019) **Structural insights into TRPM8 inhibition and desensitization** *Science* **365**:1434–1440
- Doerner J. F., Gisselmann G., Hatt H., Wetzel C. H. (2007) **Transient receptor potential channel A1 is directly gated by calcium ions** *J Biol Chem* **282**:13180–13189
- Duan J. *et al.* (2019) **Cryo-EM structure of TRPC5 at 2.8-Å resolution reveals unique and conserved structural elements essential for channel function** *Sci Adv* **5**
- Duan J., Li J., Zeng B., Chen G. L., Peng X., Zhang Y., Wang J., Clapham D. E., Li Z., Zhang J. (2018) **Structure of the mouse TRPC4 ion channel** *Nat Commun* **9**
- Ewanchuk B. W., Allan E. R. O., Warren A. L., Ramachandran R., Yates R. M. (2018) **The cooling compound icilin attenuates autoimmune neuroinflammation through modulation of the T-cell response** *FASEB J* **32**:1236–1249
- Gao Y., Cao E., Julius D., Cheng Y. (2016) **TRPV1 structures in nanodiscs reveal mechanisms of ligand and lipid action** *Nature* **534**:347–351

- Garcia-Avila M., Islas L. D. (2019) **What is new about mild temperature sensing? A review of recent findings** *Temperature (Austin)* **6**:132–141
- Garcia-Martinez C., Morenilla-Palao C., Planells-Cases R., Merino J. M., Ferrer-Montiel A. (2000) **Identification of an aspartic residue in the P-loop of the vanilloid receptor that modulates pore properties** *J Biol Chem* **275**:32552–32558
- Gonzalez-Muniz R., Bonache M. A., Martin-Escura C., Gomez-Monterrey I. (2019) **Recent Progress in TRPM8 Modulation: An Update** *Int J Mol Sci* **20**
- Grandl J., Hu H., Bandell M., Bursulaya B., Schmidt M., Petrus M., Patapoutian A. (2008) **Pore region of TRPV3 ion channel is specifically required for heat activation** *Nat Neurosci* **11**:1007–1013
- Grandl J., Kim S. E., Uzzell V., Bursulaya B., Petrus M., Bandell M., Patapoutian A. (2010) **Temperature-induced opening of TRPV1 ion channel is stabilized by the pore domain** *Nat Neurosci* **13**:708–714
- Guo J., She J., Zeng W., Chen Q., Bai X. C., Jiang Y. (2017) **Structures of the calcium-activated, non-selective cation channel TRPM4** *Nature* **552**:205–209
- Guo W., Tang Q., Wei M., Kang Y., Wu J. X., Chen L. (2022) **Structural mechanism of human TRPC3 and TRPC6 channel regulation by their intracellular calcium-binding sites** *Neuron* **110**:1023–1035
- Hamill O. P., Marty A., Neher E., Sakmann B., Sigworth F. J. (1981) **Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches** *Pflugers Arch* **391**:85–100
- Hasan R., Leeson-Payne A. T., Jaggar J. H., Zhang X. (2017) **Calmodulin is responsible for Ca(2+)-dependent regulation of TRPA1 Channels** *Sci Rep* **7**
- Hasan R., Zhang X. (2018) **Ca(2+) Regulation of TRP Ion Channels** *Int J Mol Sci* **19**
- Held K. *et al.* (2015) **Activation of TRPM3 by a potent synthetic ligand reveals a role in peptide release** *Proc Natl Acad Sci U S A* **112**:E1363–1372
- Henikoff S., Henikoff J. G. (1992) **Amino acid substitution matrices from protein blocks** *Proc Natl Acad Sci U S A* **89**:10915–10919
- Hermosura M. C., Monteilh-Zoller M. K., Scharenberg A. M., Penner R., Fleig A. (2002) **Dissociation of the store-operated calcium current I(CRAC) and the Mg-nucleotide-regulated metal ion current MagNum** *J Physiol* **539**:445–458
- Himmel N. J., Cox D. N. (2020) **Transient receptor potential channels: current perspectives on evolution, structure, function and nomenclature** *Proc Biol Sci* **287**
- Hofmann T., Chubakov V., Gudermann T., Montell C. (2003) **TRPM5 is a voltage-modulated and Ca(2+)-activated monovalent selective cation channel** *Curr Biol* **13**:1153–1158
- Hu H., Grandl J., Bandell M., Petrus M., Patapoutian A. (2009) **Two amino acid residues determine 2-APB sensitivity of the ion channels TRPV3 and TRPV4** *Proc Natl Acad Sci U S A* **106**:1626–1631

- Hu J., Park S. J., Walter T., Orozco I. J., O'Dea G., Ye X., Du J., Lu W. (2024) **Physiological temperature drives TRPM4 ligand recognition and gating** *Nature*
- Huang Y., Roth B., Lu W., Du J. (2019) **Ligand recognition and gating mechanism through three ligand-binding sites of human TRPM2 channel** *eLife* 8
- Huang Y., Winkler P. A., Sun W., Lu W., Du J. (2018) **Architecture of the TRPM2 channel and its activation mechanism by ADP-ribose and calcium** *Nature* 562:145–149
- Huffer K. E., Aleksandrova A. A., Jara-Oseguera A., Forrest L. R., Swartz K. J. (2020) **Global alignment and assessment of TRP channel transmembrane domain structures to explore functional mechanisms** *eLife* 9
- Hughes T. E., Del Rosario J. S., Kapoor A., Yazici A. T., Yudin Y., Fluck E. C., Filizola M., Rohacs T., Moiseenkova-Bell V. Y. (2019) **Structure-based characterization of novel TRPV5 inhibitors** *eLife* 8
- Jara-Oseguera A., Bae C., Swartz K. J. (2016) **An external sodium ion binding site controls allosteric gating in TRPV1 channels** *eLife* 5
- Kovacs Z. M., Dienes C., Hezso T., Almasy J., Magyar J., Banyasz T., Nanasi P. P., Horvath B., Szentandrassy N. (2022) **Pharmacological Modulation and (Patho)Physiological Roles of TRPM4 Channel-Part 1: Modulation of TRPM4** *Pharmaceuticals (Basel)* 15
- Kozak J. A., Cahalan M. D. (2003) **MIC channels are inhibited by internal divalent cations but not ATP** *Biophys J* 84:922–927
- Kuhn F. J., Kuhn C., Luckhoff A. (2009) **Inhibition of TRPM8 by icilin distinct from desensitization induced by menthol and menthol derivatives** *J Biol Chem* 284:4102–4111
- Kuhn F. J., Winking M., Kuhn C., Hoffmann D. C., Luckhoff A. (2013) **Surface expression and channel function of TRPM8 are cooperatively controlled by transmembrane segments S3 and S4** *Pflugers Arch* 465:1599–1610
- Kukkonen J. P. (2011) **A menage a trois made in heaven: G-protein-coupled receptors, lipids and TRP channels** *Cell Calcium* 50:9–26
- Kwon D. H., Zhang F., Fedor J. G., Suo Y., Lee S. Y. (2022) **Vanilloid-dependent TRPV1 opening trajectory from cryoEM ensemble analysis** *Nat Commun* 13
- Kwon D. H., Zhang F., Suo Y., Bouvette J., Borgnia M. J., Lee S. Y. (2021) **Heat-dependent opening of TRPV1 in the presence of capsaicin** *Nat Struct Mol Biol* 28:554–563
- Launay P., Fleig A., Perraud A. L., Scharenberg A. M., Penner R., Kinet J. P. (2002) **TRPM4 is a Ca²⁺-activated nonselective cation channel mediating cell membrane depolarization** *Cell* 109:397–407
- Liao M., Cao E., Julius D., Cheng Y. (2013) **Structure of the TRPV1 ion channel determined by electron cryo-microscopy** *Nature* 504:107–112
- Liu B., Qin F. (2017) **Single-residue molecular switch for high-temperature dependence of vanilloid receptor TRPV3** *Proc Natl Acad Sci U S A* 114:1589–1594

- Liu D., Liman E. R. (2003) **Intracellular Ca²⁺ and the phospholipid PIP₂ regulate the taste transduction ion channel TRPM5** *Proc Natl Acad Sci U S A* **100**:15160–15165
- Lu X., Yao Z., Wang Y., Yin C., Li J., Chai L., Dong W., Yuan L., Lai R., Yang S. (2022) **The acquisition of cold sensitivity during TRPM8 ion channel evolution** *Proc Natl Acad Sci U S A* **119**
- Malkia A., Pertusa M., Fernandez-Ballester G., Ferrer-Montiel A., Viana F. (2009) **Differential role of the menthol-binding residue Y745 in the antagonism of thermally gated TRPM8 channels** *Mol Pain* **5**
- McKemy D. D., Neuhausser W. M., Julius D. (2002) **Identification of a cold receptor reveals a general role for TRP channels in thermosensation** *Nature* **416**:52–58
- Montell C., Rubin G. M. (1989) **Molecular characterization of the Drosophila trp locus: a putative integral membrane protein required for phototransduction** *Neuron* **2**:1313–1323
- Moriyama H., Nomura S., Kida H., Inoue T., Imoto H., Maruta Y., Fujiyama Y., Mitsushima D., Suzuki M. (2019) **Suppressive Effects of Cooling Compounds Icilin on Penicillin G-Induced Epileptiform Discharges in Anesthetized Rats** *Front Pharmacol* **10**
- Nadezhdin K. D., Neuberger A., Khosrof L. S., Talyzina I. A., Khau J., Yelshanskaya M. V., Sobolevsky A. I. (2024) **TRPV3 activation by different agonists accompanied by lipid dissociation from the vanilloid site** *Sci Adv* **10**
- Nadezhdin K. D., Neuberger A., Trofimov Y. A., Krylov N. A., Sinica V., Kupko N., Vlachova V., Zakharian E., Efremov R. G., Sobolevsky A. I. (2021) **Structural mechanism of heat-induced opening of a temperature-sensitive TRP channel** *Nat Struct Mol Biol* **28**:564–572
- Nadler M. J. *et al.* (2001) **LTRPC7 is a Mg.ATP-regulated divalent cation channel required for cell viability** *Nature* **411**:590–595
- Neuberger A., Nadezhdin K. D., Zakharian E., Sobolevsky A. I. (2021) **Structural mechanism of TRPV3 channel inhibition by the plant-derived coumarin osthole** *EMBO Rep* **22**
- Neuberger A., Oda M., Nikolaev Y. A., Nadezhdin K. D., Gracheva E. O., Bagriantsev S. N., Sobolevsky A. I. (2023) **Human TRPV1 structure and inhibition by the analgesic SB-366791** *Nat Commun* **14**
- Nilius B., Mahieu F., Prenen J., Janssens A., Owsianik G., Vennekens R., Voets T. (2006) **The Ca²⁺-activated cation channel TRPM4 is regulated by phosphatidylinositol 4,5-bisphosphate** *EMBO J* **25**:467–478
- Nilius B., Prenen J., Droogmans G., Voets T., Vennekens R., Freichel M., Wissenbach U., Flockerzi V. (2003) **Voltage dependence of the Ca²⁺-activated cation channel TRPM4** *J Biol Chem* **278**:30813–30820
- Oberwinkler J., Lis A., Giehl K. M., Flockerzi V., Philipp S. E. (2005) **Alternative splicing switches the divalent cation selectivity of TRPM3 channels** *J Biol Chem* **280**:22540–22548
- Palchevskyi S., Czarnocki-Cieciura M., Vistoli G., Gervasoni S., Nowak E., Beccari A. R., Nowotny M., Talarico C. (2023) **Structure of human TRPM8 channel** *Commun Biol* **6**
- Peier A. M. *et al.* (2002) **A TRP channel that senses cold stimuli and menthol** *Cell* **108**:705–715

- Perraud A. L. *et al.* (2001) **ADP-ribose gating of the calcium-permeable LTRPC2 channel revealed by Nudix motif homology** *Nature* **411**:595–599
- Pezzoli M., Elhamdani A., Camacho S., Meystre J., Gonzalez S. M., le Coutre J., Markram H. (2014) **Dampened neural activity and abolition of epileptic-like activity in cortical slices by active ingredients of spices** *Sci Rep* **4**
- Phelps C. B., Wang R. R., Choo S. S., Gaudet R. (2010) **Differential regulation of TRPV1, TRPV3, and TRPV4 sensitivity through a conserved binding site on the ankyrin repeat domain** *J Biol Chem* **285**:731–740
- Plant T. D., Schaefer M. (2003) **TRPC4 and TRPC5: receptor-operated Ca²⁺-permeable nonselective cation channels** *Cell Calcium* **33**:441–450
- Plaza-Cayon A., Gonzalez-Muniz R., Martin-Martinez M. (2022) **Mutations of TRPM8 channels: Unraveling the molecular basis of activation by cold and ligands** *Med Res Rev* **42**:2168–2203
- Prawitt D., Monteilh-Zoller M. K., Brixel L., Spangenberg C., Zabel B., Fleig A., Penner R. (2003) **TRPM5 is a transient Ca²⁺-activated cation channel responding to rapid changes in [Ca²⁺]_i** *Proc Natl Acad Sci U S A* **100**:15166–15171
- Qin F (2014) **Temperature Sensing by Thermal TRP Channels: Thermodynamic Basis and Molecular Insights** *Curr Top Membr* **74**:19–50
- Rohacs T (2024) **Phosphoinositide Regulation of TRP Channels: A Functional Overview in the Structural Era** *Annu Rev Physiol* **86**:329–355
- Ruan Z., Haley E., Orozco I. J., Sabat M., Myers R., Roth R., Du J., Lu W. (2021) **Structures of the TRPM5 channel elucidate mechanisms of activation and inhibition** *Nat Struct Mol Biol* **28**:604–613
- Sano Y., Inamura K., Miyake A., Mochizuki S., Yokoi H., Matsushime H., Furuichi K. (2001) **Immunocyte Ca²⁺ influx system mediated by LTRPC2** *Science* **293**:1327–1330
- Shalygin A., Kolesnikov D., Glushankova L., Gusev K., Skopin A., Skobeleva K., Kaznacheyeva E. V. (2021) **Role of STIM2 and Orai proteins in regulating TRPC1 channel activity upon calcium store depletion** *Cell Calcium* **97**
- Sherkheli M. A., Gisselmann G., Hatt H. (2012) **Supercooling agent icilin blocks a warmth-sensing ion channel TRPV3** *ScientificWorldJournal* **2012**
- Sherkheli M. A., Vogt-Eisele A. K., Bura D., Beltran Marques L. R., Gisselmann G., Hatt H. (2010) **Characterization of selective TRPM8 ligands and their structure activity response (S.A.R) relationship** *J Pharm Pharm Sci* **13**:242–253
- Singh A. K., McGoldrick L. L., Sobolevsky A. I. (2018) **Structure and gating mechanism of the transient receptor potential channel TRPV3** *Nat Struct Mol Biol* **25**:805–813
- Singh A. K., Saotome K., McGoldrick L. L., Sobolevsky A. I. (2018) **Structural bases of TRP channel TRPV6 allosteric modulation by 2-APB** *Nat Commun* **9**
- Song K., Wei M., Guo W., Quan L., Kang Y., Wu J. X., Chen L. (2021) **Structural basis for human TRPC5 channel inhibition by two distinct inhibitors** *eLife* **10**

- Story G. M. *et al.* (2003) **ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures** *Cell* **112**:819–829
- Su Q., Chen M., Wang Y., Li B., Jing D., Zhan X., Yu Y., Shi Y. (2021) **Structural basis for Ca(2+) activation of the heteromeric PKD1L3/PKD2L1 channel** *Nat Commun* **12**
- Sura L., Zima V., Marsakova L., Hynkova A., Barvik I., Vlachova V. (2012) **C-terminal acidic cluster is involved in Ca²⁺-induced regulation of human transient receptor potential ankyrin 1 channel** *J Biol Chem* **287**:18067–18077
- Tominaga M., Caterina M. J., Malmberg A. B., Rosen T. A., Gilbert H., Skinner K., Raumann B. E., Basbaum A. I., Julius D. (1998) **The cloned capsaicin receptor integrates multiple pain-producing stimuli** *Neuron* **21**:531–543
- Ullrich N. D., Voets T., Prenen J., Vennekens R., Talavera K., Droogmans G., Nilius B. (2005) **Comparison of functional properties of the Ca²⁺-activated cation channels TRPM4 and TRPM5 from mice** *Cell Calcium* **37**:267–278
- Vandewauw I. *et al.* (2018) **A TRP channel trio mediates acute noxious heat sensing** *Nature* **555**:662–666
- Vazquez G., Wedel B. J., Aziz O., Trebak M., Putney J. W. (2004) **The mammalian TRPC cation channels** *Biochim Biophys Acta* **1742**:21–36
- Vinayagam D., Mager T., Apelbaum A., Bothe A., Merino F., Hofnagel O., Gatsogiannis C., Raunser S. (2018) **Electron cryo-microscopy structure of the canonical TRPC4 ion channel** *eLife* **7**
- Vinayagam D. *et al.* (2020) **Structural basis of TRPC4 regulation by calmodulin and pharmacological agents** *eLife* **9**
- Voets T., Owsianik G., Janssens A., Talavera K., Nilius B. (2007) **TRPM8 voltage sensor mutants reveal a mechanism for integrating thermal and chemical stimuli** *Nat Chem Biol* **3**:174–182
- Voets T., Prenen J., Vriens J., Watanabe H., Janssens A., Wissenbach U., Bodding M., Droogmans G., Nilius B. (2002) **Molecular determinants of permeation through the cation channel TRPV4** *J Biol Chem* **277**:33704–33710
- Vriens J., Held K., Janssens A., Toth B. I., Kerselaers S., Nilius B., Vennekens R., Voets T. (2014) **Opening of an alternative ion permeation pathway in a nociceptor TRP channel** *Nat Chem Biol* **10**:188–195
- Vriens J. *et al.* (2011) **TRPM3 is a nociceptor channel involved in the detection of noxious heat** *Neuron* **70**:482–494
- Wang H. *et al.* (2019) **Gain-of-Function Mutations in TRPM4 Activation Gate Cause Progressive Symmetric Erythrokeratoderma** *J Invest Dermatol* **139**:1089–1097
- Wang Y. Y., Chang R. B., Waters H. N., McKemy D. D., Liman E. R. (2008) **The nociceptor ion channel TRPA1 is potentiated and inactivated by permeating calcium ions** *J Biol Chem* **283**:32691–32703
- Williams J. C., Kalyanamoorthy S. (2021) **PoseFilter: a PyMOL plugin for filtering and analyzing small molecule docking in symmetric binding sites** *Bioinformatics* **37**:3367–3368

- Winking M., Hoffmann D. C., Kuhn C., Hilgers R. D., Luckhoff A., Kuhn F. J. (2012) **Importance of a conserved sequence motif in transmembrane segment S3 for the gating of human TRPM8 and TRPM2** *PLoS One* 7
- Wong F., Schaefer E. L., Roop B. C., LaMendola J. N., Johnson-Seaton D., Shao D. (1989) **Proper function of the *Drosophila* trp gene product during pupal development is important for normal visual transduction in the adult** *Neuron* 3:81–94
- Xiao R., Tang J., Wang C., Colton C. K., Tian J., Zhu M. X. (2008) **Calcium plays a central role in the sensitization of TRPV3 channel to repetitive stimulations** *J Biol Chem* 283:6162–6174
- Yamaguchi S., Tanimoto A., Iwasa S., Otsuguro K. (2019) **TRPM4 and TRPM5 Channels Share Crucial Amino Acid Residues for Ca(2+) Sensitivity but Not Significance of PI(4,5)P(2)** *Int J Mol Sci* 20
- Yang F., Vu S., Yarov-Yarovoy V., Zheng J. (2016) **Rational design and validation of a vanilloid-sensitive TRPV2 ion channel** *Proc Natl Acad Sci U S A* 113:E3657–3666
- Yang Y., Wei M., Chen L. (2022) **Structural identification of riluzole-binding site on human TRPC5** *Cell Discov* 8
- Yeh F., Jara-Oseguera A., Aldrich R. W. (2023) **Implications of a temperature-dependent heat capacity for temperature-gated ion channels** *Proc Natl Acad Sci U S A* 120
- Yin Y., Le S. C., Hsu A. L., Borgnia M. J., Yang H., Lee S. Y. (2019) **Structural basis of cooling agent and lipid sensing by the cold-activated TRPM8 channel** *Science* 363
- Yin Y., Lee S. Y. (2020) **Current View of Ligand and Lipid Recognition by the Menthol Receptor TRPM8** *Trends Biochem Sci* 45:806–819
- Yin Y., Wu M., Hsu A. L., Borschel W. F., Borgnia M. J., Lander G. C., Lee S. Y. (2019) **Visualizing structural transitions of ligand-dependent gating of the TRPM2 channel** *Nat Commun* 10
- Yin Y., Wu M., Zubcevic L., Borschel W. F., Lander G. C., Lee S. Y. (2018) **Structure of the cold- and menthol-sensing ion channel TRPM8** *Science* 359:237–241
- Yin Y., Zhang F., Feng S., Butay K. J., Borgnia M. J., Im W., Lee S. Y. (2022) **Activation mechanism of the mouse cold-sensing TRPM8 channel by cooling agonist and PIP(2)** *Science* 378
- Zhang F., Hanson S. M., Jara-Oseguera A., Krepiy D., Bae C., Pearce L. V., Blumberg P. M., Newstead S., Swartz K. J. (2016) **Engineering vanilloid-sensitivity into the rat TRPV2 channel** *eLife* 5
- Zhang F., Jara-Oseguera A., Chang T. H., Bae C., Hanson S. M., Swartz K. J. (2018) **Heat activation is intrinsic to the pore domain of TRPV1** *Proc Natl Acad Sci U S A* 115:E317–E324
- Zhang F., Swartz K. J., Jara-Oseguera A. (2019) **Conserved allosteric pathways for activation of TRPV3 revealed through engineering vanilloid-sensitivity** *eLife* 8
- Zhang K., Julius D., Cheng Y. (2021) **Structural snapshots of TRPV1 reveal mechanism of polymodal functionality** *Cell* 184:5138–5150

- Zhang Z., Okawa H., Wang Y., Liman E. R. (2005) **Phosphatidylinositol 4,5-bisphosphate rescues TRPM4 channels from desensitization** *J Biol Chem* **280**:39185–39192
- Zhang Z., Toth B., Szollosi A., Chen J., Csanady L. (2018) **Structure of a TRPM2 channel in complex with Ca(2+) explains unique gating regulation** *eLife* **7**
- Zhao C., Xie Y., Xu L., Ye F., Xu X., Yang W., Yang F., Guo J. (2022) **Structures of a mammalian TRPM8 in closed state** *Nat Commun* **13**
- Zhao J., Lin King J. V., Paulsen C. E., Cheng Y., Julius D. (2020) **Irritant-evoked activation and calcium modulation of the TRPA1 receptor** *Nature* **585**:141–145
- Zimova L., Sinica V., Kadkova A., Vyklicka L., Zima V., Barvik I., Vlachova V. (2018) **Intracellular cavity of sensor domain controls allosteric gating of TRPA1 channel** *Sci Signal* **11**
- Zubcevic L., Le S., Yang H., Lee S. Y. (2018) **Conformational plasticity in the selectivity filter of the TRPV2 ion channel** *Nat Struct Mol Biol* **25**:405–415
- Zurborg S., Yurgionas B., Jira J. A., Caspani O., Heppenstall P. A. (2007) **Direct activation of the ion channel TRPA1 by Ca2+** *Nat Neurosci* **10**:277–279

Editors

Reviewing Editor

Lejla Zubcevic

University of Kansas Medical Center, Kansas City, United States of America

Senior Editor

Merritt Maduke

Stanford University, Stanford, United States of America

Reviewer #2 (Public review):

Summary:

The authors set out to study whether the cooling agent binding site in TRPM8, which is located between the S1-S4 and the TRP domain, is conserved within the TRPM family of ion channels. They specifically chose the TRPM4 channel as the model system, which is directly activated by intracellular Ca²⁺. Using electrophysiology, the authors characterized and compared the Ca²⁺ sensitivity and the voltage-dependence of TRPM4 channels in the absence and presence of synthetic cooling agonist icilin. They also analyzed the mutational effects of residues (A867G and R901H; equivalent mutations in TRPM8 were shown involved in icilin sensitivity) on Ca²⁺ sensitivity and voltage-dependence of TRPM4 in the absence and presence of Ca²⁺. Based on the results as well as structure/sequence alignment, the authors concluded that icilin likely binds to the same pocket in TRPM4 and suggested that this cooling agonist binding pocket is conserved in TRPM channels.

Strengths:

The authors gave a very thorough introduction of the TRPM channels. They have nicely characterized the Ca²⁺ sensitivity and the voltage-dependence of TRPM4 channels and demonstrated icilin potentiates the Ca²⁺ sensitivity and diminishes the outward rectification of TRPM4. These results indicate icilin modulates TRPM4 activation by Ca²⁺.

The authors have incorporated additional data analysis and control experiments in the revised manuscript to strengthen their findings. They have well addressed the concerns raised by reviewers in the responses.

Weaknesses:

The study is conducted based on an assumption that TRPM4 activation is controlled by Ca²⁺ binding to a single site in the S1-S4 pocket in each subunit, and the second Ca²⁺ site in the cytoplasmic MHRs is simplified.

Despite the technical reasons presented by the authors in the rebuttal, the conclusion of this study can be strengthened if more cooling compounds- the most well-studied natural cooling agonist menthol, and/or other cooling agonists such as WS-12 and/or C3-are tested for their effects on TRPM4 and several other TRPM channels.

<https://doi.org/10.7554/eLife.99643.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public Review):

In this important study, Huffer et al posit that non-cold sensing members of the TRPM subfamily of ion channels (e.g., TRPM2, TRPM4, TRPM5) contain a binding pocket for icilin which overlaps with the one found in the cold-activated TRPM8 channel.

The authors identify the residues involved in icilin binding by analyzing the existing TRPM8-icilin complex structures and then use their previously published approach of structure-based sequence comparison to compare the icilin binding residues in TRPM8 to other TRPM channels. This approach uncovered that the residues are conserved in a number of TRPM members: TRPM2, TRPM4, and TRPM5. The authors focus on TRPM4, with the rationale that it has the simplest activation properties (a single Ca²⁺-binding site). Electrophysiological studies show that icilin by itself does not activate TRPM4, but it strongly potentiates the Ca²⁺ activation of TRPM4, and introducing the A867G mutation (the mutation that renders avian TRPM8 sensitive to icilin) further increases the potentiating effects of the compound. Conversely, the mutation of a residue that likely directly interacts with icilin in the binding pocket, R901H, results in channels whose Ca²⁺ sensitivity is not potentiated by icilin.

The data indicate that, just like in TRPV channels, the binding pockets and allosteric networks might be conserved in the TRPM subfamily.

The data are convincing, and the authors employ good experimental controls.

We appreciate the supportive feedback of this reviewer.

Reviewer #2 (Public Review):

Summary:

The authors set out to study whether the cooling agent binding site in TRPM8, which is located between the S1-S4 and the TRP domain, is conserved within the TRPM family of ion channels. They specifically chose the TRPM4 channel as the model system, which is

directly activated by intracellular Ca²⁺. Using electrophysiology, the authors characterized and compared the Ca²⁺ sensitivity and the voltage dependence of TRPM4 channels in the absence and presence of synthetic cooling agonist icilin. They also analyzed the mutational effects of residues (A867G and R901H; equivalent mutations in TRPM8 were shown involved in icilin sensitivity) on Ca²⁺ sensitivity and voltage-dependence of TRPM4 in the absence and presence of Ca²⁺. Based on the results as well as structure/sequence alignment, the authors concluded that icilin likely binds to the same pocket in TRPM4 and suggested that this cooling agonist binding pocket is conserved in TRPM channels.

Strengths:

The authors gave a very thorough introduction to the TRPM channels. They have nicely characterized the Ca²⁺ sensitivity and the voltage-dependence of TRPM4 channels and demonstrated icilin potentiates the Ca²⁺ sensitivity and diminishes the outward rectification of TRPM4. These results indicate icilin modulates TRPM4 activation by Ca²⁺.

We appreciate the supportive feedback of this reviewer.

Weaknesses:

The reviewer has a few concerns. First, icilin alone (at 25 μM) and in the absence of Ca²⁺ does not activate the TRPM4 channel. Have the authors titrated a wide range of icilin concentrations (without Ca²⁺ present) for TRPM4 activation? It raises the question that whether icilin is indeed an agonist for TRPM4 channel. This has not been tested so it is unclear. One may argue that icilin needs Ca²⁺ as a co-factor for channel activation just like in TRPM8 channel. This leads to the second concern, which is a complication in the experimental design and data interpretation. TRPM4 itself requires Ca²⁺ for activation to begin with, thus it is hard to dissect whether the current observed here for TRPM4 is activated by Ca²⁺ or by icilin plus its cofactor Ca²⁺. This is the difference between TRPM8 and TRPM4, as TRPM8 itself is not activated by Ca²⁺, thus TRPM8 activation is through icilin and Ca²⁺ acts as a prerequisite for icilin activation.

We agree that the comparison between TRPM8 and TRPM4 is not perfect because TRPM4 requires Ca²⁺ for activation, but it is clear that the current activated by Ca²⁺ in the presence of icilin also involves icilin because it activates at lower Ca²⁺ concentrations and lower voltages. We have tested icilin at concentrations between 12.5 and 25 μM and at these concentrations icilin does not activate TRPM4 when applied alone, so we have no evidence that it is an agonist. Both of these concentrations are higher than those reported by Chuang et al. to be saturating for TRPM8 in the presence of Ca²⁺. We haven't tested icilin at higher concentrations because we wanted to keep the final concentration of DMSO low enough to avoid any effects of the vehicle. We now emphasize this even more clearly in the revised manuscript.

The results presented in this study are only sufficient to show that icilin modulates the Ca²⁺-dependent activation of TRPM4 and icilin at best may act as an allosteric modulator for TRPM4 function. One cannot conclude from the current work that icilin is an agonist or even specifically a cooling agonist for TRPM4. Icilin is a cooling agonist for TRPM8, but it does not mean that if icilin modulates TRPM4 activity then it serves as a cooling agonist for TRPM4.

We agree with these comments, and we believe that the intent of our statements in the manuscript are completely in line with this perspective. We never refer to icilin as a cooling agent for TRPM4 but rather refer to the cooling agent binding pocket in TRPM8 and how that appears to be conserved and functions in TRPM4 to modulate opening of the channel. We

have carefully gone through the manuscript to refer directly to icilin by name (rather than as a cooling agent) when referring to its actions on TRPM4 to make sure there is no confusion.

For the mutation data on A867G, Figure 4A-B, left panels, it looks like A867G has stronger Ca²⁺ sensitivity compared to the WT in the absence of icilin and the onset of current activation is faster than the WT, or this is simply due to the scale of the data figure are different between A867G and the WT. Overall the mutagenesis data are weak to support the conclusion that icilin binds to the S1-S4 pocket. The authors need to mutate more residues that are involved in direct interaction with icilin based on the available structural information, including but limited to residues equivalent to Y745 and H845 in human TRPM8.

The A867G mutant does seem to promote opening by Ca²⁺ in the absence of icilin, and we now comment on this in the manuscript. Having said that, we have not carefully studied the concentration-dependence for activation by Ca²⁺ because at higher concentrations we see evidence of desensitization. We think Ca²⁺, icilin and depolarized voltages promote an open state of TRPM4 and the A867G does so as well.

We respectfully disagree about the strength of mutagenesis results present in our manuscript. We present clear gain and loss of function for two mutants corresponding to influential residues within the cooling agent binding pocket of TRPM8. We agree that Y786 mutations would have been a valuable addition, and our plan was to include mutations of this residue. Unfortunately, both the Y786A and Y786H mutants exhibited rundown to repeated stimulation by Ca²⁺, making them challenging to obtain reliable results on their effects on modulation by icilin.

The authors set out to study the conservation of the cooling agonist binding site in TRPM family, but only tested a synthetic cooling agonist icilin on TRPM4. In order to draw a broad conclusion as the title and the discussion have claimed, the authors need to more cooling compounds, including the most well-known natural cooling agonist menthol, and other cooling agonists such as WS-12 and/or C3, and test their effects on several TRPM channels, not just TRPM4. With the current data, the authors need to significantly tone down the claim of a conserved cooling agonist binding pocket in the TRPM family.

We would have liked to broaden the scope to other ligands that modulate TRPM8 and we agree that including those data would certainly reinforce our conclusions. However, the first author recently moved on to a new faculty position and extending our findings would require enlisting another member of the lab and take away from their independent projects. We also do not agree that this is essential to support any of our conclusions. It is also important to keep in mind that icilin is a high-affinity ligand for TRPM8, such that weaker interactions with TRPM4 can still be readily observed. We think it is likely that lower affinity agonists like menthol might not have sufficient affinity to see activity in TRPM4. This scenario is not unlike our earlier experience with TRPV channels where we succeeded in engineering vanilloid sensitivity into TRPV2 and TRPV3 using the high affinity agonist resiniferatoxin (Zhang et al., 2016, eLife). In the case of TRPV2, another group had made the same quadruple mutant and failed to see activation by capsaicin even though resiniferatoxin also worked in their hands (see Fig. 2 in Yang et al., 2016, PNAS).

On page 11, the authors suggest based on the current data, that TRPM2 and TRPM5 may also be sensitive to cooling agonists because the key residues are conserved. TRPM2 is the closest homolog to TRPM8 but is menthol-insensitive. There are studies that attempted to convert menthol sensitivity to TRPM2, for example, Bandell 2006 attempted to introduce S2 and TRP domains from TRPM8 into TRPM2 but failed to make TRPM2 a menthol-sensitive channel. The sequence conservation or structural similarity is not

sufficient for the authors to suggest a shared cooling agonist sensitivity or even a common binding site in the TRPM2 and TRPM5 channels. Again, as pointed out above, the authors need to establish the actual activation of other TRPM channels by these agonists first, before proceeding to functionally probe whether other TRPM channels adopt a conserved agonist binding site.

We are somewhat confused by these comments because we do not comment about whether cooling agents can activate TRPM2 or TRPM5. We simply analyzed the structures to make the point that the key residues in the cooling agent binding pocket of TRPM8 are conserved in these other TRPM channels. The Bandell paper is relevant, but it is also possible that they failed to uncover a relationship because they only used an agonist that has relatively low affinity for TRPM8. It would have been interesting to see what they might have found if they had used a high-affinity ligand like icilin instead of a low affinity ligand like menthol.

Taken together, this current work presents data to show the modulatory effects of icilin on the Ca²⁺ dependent activation and voltage dependence of the TRPM4 channel.

We agree.

Reviewer #3 (Public Review):

Summary:

The family of transient receptor potential (TRP) channels are tetrameric cation selective channels that are modulated by a variety of stimuli, most notably temperature. In particular, the Transient receptor potential Melastatin subfamily member 8 (TRPM8) is activated by noxious cold and other cooling agents such as menthol and icilin and participates in cold somatosensation in humans. The abundance of TRP channel structural data that has been published in the past decade demonstrates clear architectural conservation within the ion channel family. This suggests the potential for unifying mechanisms of gating despite their varied modes of regulation, which are not yet understood. To address this question, the authors examine the 264 structures of TRP channels determined to date and observe a potential binding pocket for icilin in multiple members of the Melastatin subfamily, TRPM2, TRPM4, and TRPM5. Interestingly, none of the other Melastatin subfamily members had been shown to be sensitive to icilin apart from TRPM8. Each of these channels is activated by intracellular calcium (Ca²⁺) and a Ca²⁺ binding site neighbors the predicted pocket for icilin binding in all cryo-EM structures. The authors examined whether icilin could modulate the activation of TRPM4 in the presence of intracellular Ca²⁺. The addition of icilin enhances Ca²⁺-dependent activation of TRPM4, promotes channel opening at negative membrane potentials, and improves the kinetics of opening. Furthermore, mutagenesis of TRPM4 residues within the putative icilin binding pocket predicted to enhance or diminish TRPM4 activity elicit these behaviors. Overall, this study furthers our understanding of the Melastatin subfamily of TRP channel gating and demonstrates that a conserved binding pocket observed between TRPM4 and TRPM8 channel structures can function similarly to regulate channel gating.

Strengths:

This is a simple and elegant study capitalizing on a vast amount of high-resolution structural information from the TRP channel of ion channels to identify a conserved binding pocket that was previously unknown in the Melastatin subfamily, which is interrogated by the authors through careful electrophysiology and mutagenesis studies.

Weaknesses:

No weaknesses were identified by this reviewer.

We appreciate the supportive comments of the review.

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

I don't have any major asks, but a few questions did arise while reading your work.

(1) You refer multiple times to the VSLD pocket as being "open to the cytoplasm". It is not clear if you are implying that compounds such as icilin access the pocket via the cytoplasm (e.g., permeate the membrane to the cytosol, and then enter the binding site?) Is there data to support this? Some clarification here would be helpful, and perhaps explain if there is any distinction between how calcium might enter the VSLD binding site vs hydrophobic compounds like icilin.

This is an excellent point. Our reference to “open to the cytoplasm” was for Ca²⁺ ions and we have no evidence for how icilin enters the cooling agent binding pocket. We had tried to look for evidence that Ca²⁺ might trap icilin in TRPM4 but at the end of the day the results were not convincing enough to include in the manuscript. We have added data showing that icilin slows deactivation of TRPM4 after removing Ca²⁺, which is particularly evident in the A867G mutant, but this doesn’t inform on whether Ca²⁺ can trap icilin. We have added a statement about not knowing how icilin enters or leaves the cooling agent binding pocket in TRPM channels.

(2) Icilin is referred to as a "cooling compound", but its cooling effects are dependent on its interactions with TRPM8. This might be something to clarify, as it might otherwise be understood that other TRPM channels that interact with icilin also mediate the sensing of cool temperatures.

This is another excellent point and we have no reason to believe that icilin interacting with any TRPM channel other than TRPM8 mediates cooling sensations. We have added a statement to this effect in the discussion when considering actions of icilin that might be mediated by TRPM4 channels.

Reviewer #2 (Recommendations For The Authors):

(1) The title and statements in the results/discussion refer to icilin as a cooling agonist of TRPM4 and binds to a conserved "cooling agonist binding pocket", and the authors suggested a similar role and binding site for icilin in TRPM2 and TRPM5 channel. It is a too broad conclusion that is not fully supported by the current experimental data, which only shows icilin works as a modulator, not an agonist for TRPM4 channel. The authors should change the usage of cooling agonist or conserved cooling agonist binding pocket plus significantly tone down the conclusion of a conserved cooling agonist binding pocket, which is potentially misleading. Alternatively, if the authors insist on using cooling agonist in this context, they should establish the activation of TRPM4, TRPM2, and TRPM5 by icilin as the first step, because the current data only support icilin as a TRPM4 modulator but not an agonist.

We respectfully don’t agree with this opinion. We show broad conservation of the cooling agent binding pocket in structures of many TRPM channels, and we chose one of them to test for a functional relationship. We think that the title accurately reflects the topic of the paper and does not specify the extent to which functional conservation has been demonstrated and

we would like to keep it. The distinction between agonist and modulator is not even germane because icilin is not an agonist of TRPM8 either.

(2) The manuscript will be strengthened if the authors test additional cooling compounds of TRPM8, including menthol, the menthol analog WS-12, and C3. More importantly, distinct from icilin, these three compounds do not depend on Ca²⁺ to activate the TRPM8 channel. Thus when testing these compounds on TRPM4, it may reduce the complication of the role of Ca²⁺, as TRPM4 channel itself requires Ca²⁺ for activation.

We restate our response to this point in the public review...

We would have liked to broaden the scope to other ligands that modulate TRPM8 and we agree that including those data would certainly reinforce our conclusions. However, the first author recently moved on to a new faculty position and extending our findings would require enlisting another member of the lab and taking away from their independent projects. We also do not agree that this is essential to support any of our conclusions. It is also important to keep in mind that icilin is a high-affinity ligand for TRPM8, such that weaker interactions with TRPM4 can still be readily observed. We think it is likely that lower affinity agonists like menthol might not have sufficient affinity to see activity in TRPM4. This scenario is not unlike our earlier experience with TRPV channels where we succeeded in engineering vanilloid sensitivity into TRPV2 and TRPV3 using the high affinity agonist resiniferatoxin (Zhang et al., 2016, eLife). In the case of TRPV2, another group had made the same quadruple mutant and failed to see activation by capsaicin even though resiniferatoxin also worked in their hands (see Fig. 2 in Yang et al., 2016, PNAS).

(3) The manuscript will be strengthened if the authors test additional residues in the S1-S4 pocket that form direct interactions or are within interacting distances with icilin based on the cryo-EM structures.

We restate our response to this point in the public review...

We present clear gain and loss of function for two mutants corresponding to influential residues within the cooling agent binding pocket of TRPM8. We agree that Y786 mutations would have been a valuable addition and our plan was to include mutations of this residue. Unfortunately, both the Y786A and Y786H mutants exhibited rundown, making them challenging to obtain reliable results on their effects on modulation by icilin.

Furthermore, the ambiguity in the icilin binding pose based on available TRPM8 structures complicates structure-based identification of the most important interacting residues in TRPM8, and we would have needed to functionally validate the effects of any novel mutations we identified in TRPM8 prior to testing them in TRPM4. Instead, we have based our mutagenesis on constructs that have been previously characterized to affect the sensitivity of TRPM8 to cooling agents. A systematic mutagenesis scan of TRPM8 residues predicted to interact differentially with icilin in the two different available binding poses would likely help clarify the true binding pose of icilin and would be an interesting future study.

Reviewer #3 (Recommendations For The Authors):

I enjoyed reading this manuscript. It was well-executed and written. It will be interesting to corroborate these findings with a cryo-EM structure of TRPM2, TRPM4, or TRPM5 in the presence of icilin.

We agree and may pursue these in future studies. This would be particularly interesting given ambiguities in how icilin docks into TRPM8 in previously published structures.

Have the authors considered icilin accessibility to its binding pocket? In other words, could the presence of intracellular Ca²⁺ inhibit the accessibility of icilin to its binding pocket in TRPM4? It should be a straightforward experiment, I think it would be informative, and could further support the authors' conclusion of the location of the TRPM4 icilin binding pocket.

We completely agree and we had tried to look for evidence that Ca²⁺ might trap icilin in TRPM4 but at the end of the day the results were not convincing enough to include in the manuscript. We have added data showing that icilin slows deactivation of TRPM4 after removing Ca²⁺, which is particularly evident in the A867G mutant, but this doesn't inform on whether Ca²⁺ can trap icilin. We have added a statement about not knowing how icilin enters or leaves the cooling agent binding pocket in TRPM channels.

Figures 7 and 8 are missing the 0 μ M Ca²⁺ control trace in the presence of 25 μ M icilin.

All sample traces from Figures 7 and 8 are shown from a single cell for the sake of comparison (Likewise, the sample traces from Figures 3 and 4 come from a single cell, and the sample traces from Figures 5 and 6 come from a single cell). Unfortunately, we were unable to obtain data from an R901H mutant cell that contained all six conditions we wished to show, and there is no representative trace for 0 μ M Ca²⁺ in the presence of 25 μ M icilin for that cell.

This is up to the discretion of the authors, but perhaps a better way to arrange the paper Figures would be to combine Figures 5-6 and Figures 7-8 and rearrange the data to place some in a supplementary figure (e.g. Figure 5-6 = Figure 5 and Figure 5 - Figure Supplement 1, Figure 7-8 = Figure 6 and Figure 6 - Figure Supplement 1).

We carefully considered these suggestions and we appreciate the reviewers' flexibility but would prefer to retain the original arrangement of data in the figures.

Are there any mutations in the icilin binding pocket in TRPM4, and presumably TRPM2 and TRPM5, that are associated with human disease? This is a question that came to my mind and not one that needs to be addressed in the manuscript.

This is an interesting point. There are quite a few disease-associated mutants within TRPM4 at positions corresponding to the cooling agent binding pocket in TRPM8. We could not see an appropriate place in the discussion where we could concisely bring this information in so we decided against commenting.

<https://doi.org/10.7554/eLife.99643.2.sa0>