

Uev1A counteracts oncogenic *Ras* stimuli in both polyploid and diploid cells

Reviewed Preprint

v1 • May 30, 2025

Not revised

Qi Zhang, Yunfeng Wang, Xueli Fu, Ziguang Wang, Yang Zhang, Lizhong Yan, Yuejia Wang, Muhan Yang, Dongze Song, Ruixing Zhang, Hongru Zhang ✉, Shian Wu ✉, Shaowei Zhao ✉

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China • Department of Biochemistry and Molecular Biology, College of Life Sciences, Nankai University, Tianjin, China • Nankai International Advanced Research Institute (SHENZHEN FUTIAN), Shenzhen, China

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife assessment

This **valuable** study examines the role of E2 ubiquitin enzyme, Uev1a in tissue resistance to oncogenic RasV12 in *Drosophila melanogaster* polyploid germline cells and human cancer cell lines. The **incomplete** evidence suggests that Uev1a works with the E3 ligase APC/C to degrade Cyclin A, and the strength of evidence could be increased by addressing the expression of CycA in the ovaries and the uev1a loss of function in human cancer cells. This work would be of interest to researchers in germline biology and cancer.

<https://doi.org/10.7554/eLife.107104.1.sa3>

Abstract

Oncogenic *Ras* is known to induce DNA replication stress, leading to cellular senescence or death. In contrast, we found that it can also trigger polyploid *Drosophila* ovarian nurse cells to die by inducing aberrant division stress. To explore intrinsic protective mechanisms against this specific form of cellular stress, here we conducted a genome-wide genetic screen and identified the E2 enzyme Uev1A as a key protector. Reducing its expression levels exacerbates the nurse cell death induced by oncogenic *Ras*, while overexpressing it or its human homologs, UBE2V1 and UBE2V2, mitigates this effect. Although Uev1A is primarily known for its non-proteolytic functions, our studies demonstrate that it collaborates with the E3 APC/C complex to mediate the proteasomal degradation of Cyclin A, a key cyclin that drives cell division. Furthermore, Uev1A and UBE2V1/UBE2V2 also counteract oncogenic *Ras*-driven tumorigenesis in diploid cells, suppressing the overgrowth of germline tumors in *Drosophila* and human colorectal tumor xenografts in nude mice, respectively. Remarkably, elevated expression levels of UBE2V1/2 correlate with improved survival rates in human colorectal cancer patients harboring oncogenic *KRAS* mutations, indicating that their upregulation could represent a promising therapeutic strategy.

Introduction

Although the human body contains trillions of cells that could potentially be targeted by oncogenic mutations, cancers arise infrequently throughout a human lifetime, suggesting the presence of effective resistant mechanisms (Lowe et al., 2004). One key mechanism involves the induction of DNA replication stress by these oncogenic mutations, leading to cellular senescence or death (Hills and Diffley, 2014; Kotsantis et al., 2018). This mechanism is mediated through the activation of the DNA damage response (DDR) pathway and the tumor suppressor protein p53, which are triggered by DNA double-strand breaks. Only the cells that escape senescence and death, such as those harboring p53 mutations, can undergo transformation by oncogenic mutations, thereby initiating tumorigenesis (Gaillard et al., 2015; Igarashi et al., 2024; Macheret and Halazonetis, 2015). Among the most frequently mutated oncogenes in human cancers are the RAS genes (*KRAS*, *NRAS*, and *HRAS*), which mutations are present in 20-30% of all cancer cases (Gimple and Wang, 2019; Sanchez-Vega et al., 2018). These mutations often occur at codons 12, 13, and 61, resulting in RAS small GTPases being locked in a constitutively active, GTP-bound state (Moore et al., 2020). Notably, previous studies have shown that oncogenic *HRAS*^{G12V} can induce DNA replication stress via the DDR pathway, ultimately leading to cellular senescence (Di Micco et al., 2006; Serrano et al., 1997).

The *Ras oncogene at 85D* gene (*Ras85D*, hereafter referred to as *Ras*) in *Drosophila* exhibits high homology to human RAS genes (Neuman-Silberberg et al., 1984). Our previous research demonstrated that oncogenic *Ras*^{G12V} triggers cell death in *Drosophila* ovarian nurse cells (Zhang et al., 2024), a type of post-mitotic germ cells that undergo G/S endoreplication to become polyploid (Hammond and Laird, 1985) (Figure 1A). Notably, oncogenic *Ras*^{G12V} promotes the division of mitotic germ cells, the precursors to nurse cells. Furthermore, monoallelic deletion of the *cyclin A* (*cycA*) or *cyclin-dependent kinase 1* (*cdk1*) gene (*cycA*^{+/-} or *cdk1*^{+/-}) suppresses the nurse cell death induced by oncogenic *Ras*^{G12V} (Zhang et al., 2024). While *CycA*, a key cyclin promoting cell division, is not expressed in normal nurse cells (Lilly et al., 2000; Lilly and Spradling, 1996), its ectopic expression in nurse cells can trigger their death (as observed in this study). These findings suggest that the nurse cell death induced by oncogenic *Ras*^{G12V} is primarily due to aberrant promotion of cell division, rather than DNA hyper-replication. *Drosophila* ovarian nurse cells thus offer a valuable model for studying this specific form of cellular stress.

Ubiquitination not only mediates protein degradation through the proteasome or lysosome but also plays crucial regulatory roles in various biological processes (Dikic, 2017; Kwon and Ciechanover, 2017). These functions are largely determined by the topological structures of the polyubiquitin chains, where ubiquitin can be linked to another ubiquitin via any of its seven lysine (K) residues or its first methionine residue. Among these, K11- and K48-linked polyubiquitin primarily mediates the proteasomal degradation. In contrast, K63 linkage facilitates the lysosomal degradation of protein aggregates and damaged organelles through autophagy. Additionally, K63 linkage is involved in non-degradative processes such as DNA repair, kinase activation, and protein transport (Kwon and Ciechanover, 2017). Ubc13, a ubiquitin-conjugating (E2) enzyme (Ubc), is responsible for the K63 linkage, with its activity requiring a Ubc variant (Uev, E2) protein as a cofactor (Hofmann and Pickart, 1999; McKenna et al., 2001). These E2 enzymes have been shown to play regulatory roles in various biological processes, including cell death (Ma et al., 2013), DNA repair (Broomfield et al., 1998), DDR, and innate immunity (Andersen et al., 2005; Bai et al., 2020; Zhou et al., 2005).

However, it still remains unknown whether they also possess proteasomal proteolytic functions and whether such functions hold significant biological roles.




Figure 1.

Uev1A protects against the nurse cell death induced by oncogenic Ras^{G12V}.

(A) Schematic cartoon for *Drosophila* ovariole. During oogenesis, GSC undergoes asymmetric division to generate two daughter cells: one that self-renews and maintains GSC identity, and the other, called a cystoblast, that differentiates to support oogenesis. As differentiation progresses, each cystoblast performs four rounds of division with incomplete cytokinesis to produce 16 interconnected cystocytes, establishing a germline cyst. This germline cyst is then surrounded by epithelial follicle cells to form an egg chamber. Within each egg chamber, one of the 16 germ cells becomes the oocyte, while the remaining 15 differentiate into nurse cells. These nurse cells undergo G/S endocycling, becoming polyploid to aid in oocyte development. (B) Representative germarium and early-stage egg chamber with Flag-Ras^{G12V} overexpression driven by *bam*-GAL4-VP16. The red arrow denotes an early-stage egg chamber. (C) Genetic screening strategy. Genotype of "*bam>Ras^{G12V}*": *bam*-GAL4-VP16/FM7;; *UASp-Ras^{G12V}/TM6B*. (D, F, H, and K) Representative ovaries and egg chambers. The red arrows in (D) denote degrading egg chambers. Scale bars: 200 μ m. (E, I, and L) Quantification data. 30 ovaries from 7-day-old (E) or 3-day-old (I and L) flies were quantified for each genotype. Statistical significance was determined using t test (groups = 2) or one-way ANOVA (groups > 2): ** ($P < 0.01$), *** ($P < 0.001$), and **** ($P < 0.0001$). (G) Molecular information of the *uev1a*^{d1} and *uev1a*^{d2} mutations. The red dashed lines represent nucleotide deletions. (J) Protein sequence alignment of Uev1A, UBE2V1, and UBE2V2. It was performed using CLUSTALW and ESPrpt 3.0 software. The online version of this article includes the following source data and figure supplement for **figure 1**:

Source data 1. Screen results.

Source data 2. All genotypes.

Source data 3. Raw quantification data.

Figure supplement 1. Uev1A protects against the nurse cell death induced by oncogenic Yki^{3SA}.

In this study, to investigate the intrinsic protective mechanisms against the nurse cell death induced by oncogenic *Ras*^{G12V} in *Drosophila*, we conducted a genome-wide genetic screen and identified Uev1A as a crucial protector. Mechanistically, Uev1A works in conjunction with the anaphase-promoting complex or cyclosome (APC/C) to degrade the essential cyclin Cyclin A (CycA) via the proteasome, highlighting the critical role of its proteolytic function. Additionally, Uev1A and its human homologs, UBE2V1 and UBE2V2, also counteract oncogenic *Ras*-driven tumorigenesis in diploid cells, inhibiting the overgrowth of *Drosophila* germline tumors and human colorectal tumor xenografts in nude mice, respectively. These findings suggest that upregulation of UBE2V1/2 could represent a promising therapeutic approach for human colorectal cancers with *RAS* mutations.

Results

Genetic screen identifies Uev1A as a crucial protector

against the nurse cell death induced by oncogenic *Ras*^{G12V}

While oncogenic *Ras*^{G12V} triggers cell death in nurse cells (Zhang et al., 2024 [\[1\]](#)), it does not have the same effect in cystocytes, the precursor cells before differentiating into nurse cells (**Figure 1A, D, E** [\[1\]](#)). To confirm *Ras*^{G12V} expression in cystocytes, we generated a *UASz-flag-Ras*^{G12V} transgenic fly strain, which phenocopied the effects observed with the previously used *UASp-Ras*^{G12V}. *RAS* small GTPases need to be anchored to the inner cell membrane for their proper function (Willingham et al., 1980 [\[2\]](#)). Using *bam-GAL4-VP16* (Chen and McKearin, 2003 [\[3\]](#)), we overexpressed *Ras*^{G12V} specifically in cystocytes (*bam>flag-Ras*^{G12V}) at 29°C and observed membrane-localized Flag signals, validating the normal expression of *Ras*^{G12V} (**Figure 1B** [\[1\]](#)). Also, Flag signals were detected in some early-stage nurse cells, although these signals gradually diminished over time (**Figure 1B** [\[1\]](#)). However, nurse cell death remained very low in this context (**Figure 1D, E** [\[1\]](#)). This may be attributed to either insufficient levels of the oncoproteins or the presence of a protective mechanism.

To explore this potential protective mechanism, we conducted a genetic modifier screen by introducing individual genome-wide *Deficiencies* (361 lines) into *bam>Ras*^{G12V} flies (*bam>Ras*^{G12V}; *Deficiency*/+, **Figure 1C** [\[1\]](#)). Of note, ovaries from *bam>Ras*^{G12V}; *Df#7584*/+ flies exhibited the highest incidence of degrading egg chambers with dying nurse cells per ovary (see **Source data 1**). One gene deleted in this deficiency line is *uev1a*, and RNAi targeting *uev1a* reproduced the phenotype seen with *Df#7584* (**Figure 1D, E** [\[1\]](#)). In this and all subsequent experiments, we quantified the nurse-cell-death phenotype using the percentage of degrading to total egg chambers per ovary (**Figure 1E** [\[1\]](#)), a method that is more precise than our approach in the genetic screen. To further verify the role of *uev1a*, we generated a *UASz-uev1a* transgenic fly strain. Overexpression of Uev1A successfully rescued the nurse-cell-death phenotype in *bam>Ras*^{G12V}; *Df#7584*/+ fly ovaries (**Figure 1D, E** [\[1\]](#)), confirming that Uev1A is a crucial protector against the nurse cell death induced by oncogenic *Ras*^{G12V}.

Protective role of Uev1A against the nurse cell death induced by direct overexpression of *Ras*^{G12V}

In contrast to the scenario above (*bam>Ras*^{G12V}), direct overexpression of *Ras*^{G12V} in nurse cells using *nos-GAL4-VP16* (Rørth, 1998 [\[4\]](#); Van Doren et al., 1998 [\[5\]](#)) (*nos>Ras*^{G12V}) resulted in substantial cell death even at 25°C (**Figure 1F, I** [\[1\]](#)) (Zhang et al., 2024 [\[1\]](#)). Such observation prompted us to investigate the role of Uev1A in this context. Notably, the incidence of dying nurse cells was markedly elevated in *nos>Ras*^{G12V}+*uev1a-RNAi* ovaries compared to the

nos>Ras^{G12V}+GFP-RNAi control ovaries (**Figure 1F, I**). To further validate this, we generated two *uev1a* mutants using the CRISPR/Cas9 technique: *uev1a^{Δ1}* and *uev1a^{Δ2}*. Both mutations consist of small deletions that cause frame shifts within the second coding exon of the *uev1a* gene (**Figure 1G**). Of note, flies with the *uev1a^{Δ1/Δ1}*, *uev1a^{Δ1/Δ2}*, *uev1a^{Δ1}/Df#7584*, *uev1a^{Δ2/Δ2}*, or *uev1a^{Δ2}/Df#7584* genotype could not survive into adults, suggesting that both mutations are strong loss-of-function alleles. Consistent with the effects observed with *uev1a-RNAi*, either mutation markedly increased the incidence of dying nurse cells in *nos>Ras^{G12V}* ovaries (**Figure 1H, I**).

We next investigated whether upregulating Uev1A could mitigate the nurse cell death induced by direct overexpression of *Ras^{G12V}*. Remarkably, compared to the *nos>Ras^{G12V}+lacZ* control ovaries, the incidence of dying nurse cells was significantly reduced in *nos>Ras^{G12V}+Uev1A* ovaries (**Figure 1K, L**). Uev1A has two human homologs, UBE2V1 and UBE2V2, which share 67% (85%) and 67% (86%) sequence identities (similarities) with Uev1A, respectively (**Figure 1J**). We generated transgenic fly strains for both *UASz-UBE2V1* and *UASz-UBE2V2*. Notably, overexpression of either UBE2V1 or UBE2V2 significantly mitigated the nurse cell death induced by oncogenic *Ras^{G12V}*, similar to the effects observed with Uev1A overexpression (**Figure 1K, L**). Taken together, these results further confirm the protective role of Uev1A against the nurse cell death induced by oncogenic *Ras^{G12V}*.

Then, we were intrigued by the potential of Uev1A to protect against the nurse cell death induced by other oncogenic mutations. Yorkie (Yki), a key oncoprotein in the Hippo pathway (Huang et al., 2005), has a hyperactive form known as Yki^{3SA} (Oh and Irvine, 2009). Its overexpression using *nos-GAL4-VP16 (nos>Yki^{3SA})* at 29°C, but not at 25°C, could induce nurse cell death (**Figure 1**—**figure supplement 1**), albeit much less severe than that induced by *nos>Ras^{G12V}* at 25°C (compared with **Figure 1F, I**). Of note, the nurse cell death induced by oncogenic Yki^{3SA} was significantly alleviated by Uev1A overexpression (**Figure 1**—**figure supplement 1**), implying a broad role of Uev1A in this process. However, due to the mild effect of Yki^{3SA} on triggering nurse cell death, we focused on *Ras^{G12V}* in our subsequent studies.

Oncogenic *Ras^{G12V}* triggers nurse cells to die through activating the MAPK pathway

To investigate how Uev1A protects against the nurse cell death induced by oncogenic *Ras^{G12V}*, we first sought to further explore the mechanisms driving this cell death. Since egg chambers contain both nurse cells and oocytes (**Figure 1A**), we considered the possibility that the death signal could originate from the oocytes. The *Bicaudal D (BicD)* gene is essential for oocyte determination, and egg chambers deficient in it fail to specify oocytes (Suter and Steward, 1991). We confirmed this by using *nos>BicD-RNAi* (**Figure 2**—**figure supplement 1A**). Importantly, nurse cell death was much more pronounced in *nos>BicD-RNAi+Ras^{G12V}* egg chambers than in *nos>BicD-RNAi+GFP* control ones (**Figure 2**—**figure supplement 1A, B**), indicating that the death signal is intrinsic to the nurse cells rather than originating from the oocytes.

Our previous research demonstrated that oncogenic *Ras^{G12V}* promotes excessive proliferation in germline stem cells (GSCs) by activating the mitogen-activated protein kinase (MAPK) pathway (Zhang et al., 2024). This finding led us to investigate whether the same pathway influences the nurse cell death induced by oncogenic *Ras^{G12V}*. Notably, knockdown of *downstream of raf1 (dsor1*, encoding *Drosophila* MAPKK) (Tsuda et al., 1993) or *rolled (rl*, encoding *Drosophila* MAPK) (Biggs and Zipursky, 1992) significantly mitigated the nurse cell death induced by oncogenic *Ras^{G12V}* (**Figure 2A, B**), highlighting the pivotal role of the MAPK pathway in this process.




Figure 2.

Regulators that orchestrate the nurse cell death induced by oncogenic *Ras*^{G12V}.

(A, C, E, and G) Representative ovaries. The red arrows in (E) denote degrading egg chambers. Scale bars: 200 μ m. (B, D, F, and H) Quantification data. 30 ovaries from 3-day-old (B, D, and H) or 7-day-old (F) flies were quantified for each genotype. Statistical significance was determined using t test (groups = 2) or one-way ANOVA (groups > 2); **** ($P < 0.0001$).

The online version of this article includes the following source data and figure supplements for **figure 2**:

Source data 2. All genotypes.

Source data 3. Raw quantification data.

Figure supplement 1. Oncogenic *Ras*^{G12V} intrinsically triggers nurse cell death.

Figure supplement 2. Roles of the DDR pathway and p53 in the nurse cell death induced by oncogenic *Ras*^{G12V}.

Figure supplement 3. Uev1A does not directly degrade the *Ras*^{G12V} oncoproteins.

Figure supplement 4. Uev1A, Ben, and the APC/C complex work together to protect nurse cells from death during normal oogenesis.

CycA and the DDR pathway orchestrate the nurse cell death induced by oncogenic *Ras*^{G12V}

The Ras/MAPK pathway is well known to promote cell cycle progression (Gimple and Wang, 2019). Our previous work demonstrated that reducing the gene dosage of *CycA* or *Cdk1* suppresses the nurse cell death induced by oncogenic *Ras*^{G12V} (Zhang et al., 2024). In addition to *CycA* and *Cdk1*, Cyclin B (*CycB*) and String (*Stg*) are two other essential regulators that drive cell division (Figure 2C) (Edgar and Lehner, 1996). To investigate whether these regulators could individually induce nurse cell death upon overexpression, we tested each and found that only *CycA* triggered this phenotype (Figure 2C, D). Notably, co-overexpression of *CycA* with *Uev1A* significantly suppressed this effect (Figure 2C, D), suggesting a genetic interaction between the two.

Oncogenic *RAS* can also induce DNA replication stress in diploid cells through forced cell cycle progression, thereby activating the DDR pathway and p53 to trigger cellular senescence or death (Di Micco et al., 2006; Serrano et al., 1997). To investigate whether DDR plays a similar role in *Ras*-induced death of polyploid nurse cells, we targeted two key DDR genes: *telomere fusion* (*tefu*, encoding *Drosophila* ATM) (Oikemus et al., 2004; Silva et al., 2004; Song et al., 2004) and *loki* (*lok*, encoding *Drosophila* Chk2) (Masrouha et al., 2003; Xu et al., 2001). Strikingly, knockdown of either gene markedly enhanced this cell death (Figure 2—figure supplement 2A, B), revealing a protective role for the DDR pathway. By contrast, knockdown of p53 suppressed this cell death (Figure 2—figure supplement 2A, B), demonstrating opposing functions for DDR and p53 in this system.

Given that RNAi targeting *tefu* or *lok* phenocopied that of *uev1a*, we hypothesized that *Uev1A* may protect against the nurse cell death induced by oncogenic *Ras*^{G12V} through the DDR pathway. Since DDR is triggered by oncogenic *Ras*^{G12V}-induced stress in this context, we speculated that *Uev1A* should be upregulated as a response. To test this, we initially attempted to generate an antibody against *Uev1A* using its full protein sequence as the antigen; however, this approach was unsuccessful. As an alternative, we created a *uev1a-flag* knock-in fly strain by inserting the coding sequence of a 3xFlag tag immediately before the stop codon ("TAG") of the *uev1a* gene (Figure 2—figure supplement 2C). These knock-in flies were homozygous viable and fertile, indicating that the Flag insertion did not disrupt *Uev1A*'s normal function. Previous studies have shown that *Uev1A* can activate JNK signaling (Ma et al., 2013), which acts in the surrounding follicle cells to promote nurse cell removal during late oogenesis (Timmons et al., 2016). Indeed, we detected *Uev1A*-Flag signals in such follicle cells (Figure 2—figure supplement 2D), confirming that these signals reflect endogenous *Uev1a* expression in *Drosophila* ovaries. Surprisingly, *Uev1A*-Flag signals were low in both *nos>Ras*^{G12V}; *uev1a-flag*/+ and *uev1a-flag*/+ control germ cells, including nurse cells, with no significant differences between the two groups (Figure 2—figure supplement 2E). This suggests that *Uev1A* expression remains at basal levels, rather than being upregulated by oncogenic *Ras*^{G12V}-induced stress in this context. Although the possibility that *Uev1A* may play a role in the DDR pathway at these basal levels cannot be ruled out, we were then prompted to explore its potential proteolytic function as an E2 enzyme.

Uev1A collaborates with the APC/C complex to mitigate the nurse cell death induced by oncogenic *Ras*^{G12V}

We first investigated whether *Uev1A* is directly involved in degradation of the *Ras*^{G12V} oncoproteins. To address the challenges of analyzing membrane-localized signals in *nos>flag-Ras*^{G12V} nurse cells due to severe cell death, we performed this assay in *bam>flag-Ras*^{G12V} ovaries at 29°C. Notably, similar membrane-localized Flag signals were detected in both *bam>flag-*

Ras^{G12V}+*Uev1A* and *bam>flag-Ras*^{G12V}+*GFP* control germ cells, including nurse cells (**Figure 2—figure supplement 3** [↗](#)). This result suggests that Uev1A does not directly degrade the *Ras*^{G12V} oncoproteins.

APC/C complex primarily functions as an E3 ligase to facilitate the degradation of CycA during cell cycle progression (Sudakin et al., 1995 [↗](#)). A compelling alternative model to explain our findings is that Uev1A collaborates with the APC/C complex to degrade CycA. Cell division cycle 27 (*Cdc27*, *Drosophila* APC3) is an essential part of the substrate recognition TPR lobe within the APC/C complex (Yamano, 2019 [↗](#)). In *bam>Ras*^{G12V}+*cdc27-RNAi* ovaries, we observed dying nurse cells, a phenotype that was exacerbated with mutations in either *uev1a*^{Δ1} or *uev1a*^{Δ2} (**Figure 2E, F** [↗](#)). Furthermore, knocking down *cdc27* could increase the incidence of dying nurse cells in *bam>Ras*^{G12V}+*uev1a-RNAi* ovaries (**Figure 2E, F** [↗](#)). Additionally, we investigated Fizzy- related (*Fzr*), the *Drosophila* homolog of Cdh1, which plays a critical role as an activator of APC/C-dependent proteolysis and functions as a tumor suppressor in tumorigenesis (Greil et al., 2022 [↗](#); Sigrist and Lehner, 1997 [↗](#)). Remarkably, nearly all egg chambers in *bam>Ras*^{G12V}+*uev1a-RNAi+fzr-RNAi* ovaries exhibited degradation (**Figure 2E, F** [↗](#)).

We also knocked down three additional APC/C complex genes in *bam>Ras*^{G12V}+*uev1a-RNAi* ovaries: *shattered* (*shtd*, encoding *Drosophila* APC1) (Tanaka-Matakatsu et al., 2007 [↗](#)), *morula* (*mr*, encoding *Drosophila* APC2) (Kashevsky et al., 2002 [↗](#)), and *lemming A* (*lmgA*, encoding *Drosophila* APC11) (Nagy et al., 2012 [↗](#)). Among these factors, *Shtd* is a critical component of the scaffolding platform, while *Mr* and *LmgA* are essential components of the catalytic modules within the APC/C complex (Yamano, 2019 [↗](#)). However, no significant enhancement in nurse cell death was observed (data not shown). This may be due to the relatively mild phenotype of nurse cell death in *bam>Ras*^{G12V}+*uev1a-RNAi* ovaries. To address this, we switched to knocking down these genes in *nos>Ras*^{G12V} ovaries. Notably, knockdown of any of them could significantly exacerbated the nurse cell death induced by oncogenic *Ras*^{G12V} (**Figure 2G, H** [↗](#)), similar to the effect observed with Uev1A downregulation. Collectively, these findings provide genetic evidence that Uev1A collaborates with the APC/C complex to mitigate the nurse cell death induced by oncogenic *Ras*^{G12V}.

Then, we investigated whether Uev1A and the APC/C complex protect nurse cells from death during normal oogenesis. Given that the G2/M-promoting CycA is absent in the G/S-endocycling nurse cells (Lilly et al., 2000 [↗](#); Lilly and Spradling, 1996 [↗](#)), we used the cystocyte-specific driver, *bam-GAL4-VP16*, to conduct the assays at 29°C. Similar to nurse cells, Uev1A expression was at basal levels in cystocytes, with some expression detected in inner sheath cells and epithelial follicle cells (**Figure 2—figure supplement 2E** [↗](#)). Remarkably, nearly all egg chambers in *bam>fzr-RNAi* ovaries exhibited degradation (**Figure 2—figure supplement 4** [↗](#)), underscoring *Fzr*'s critical regulatory role in this context. Uev1A, lacking the conserved cysteine residue in its presumed active site for ubiquitin binding (Sancho et al., 1998 [↗](#); Zhou et al., 2005 [↗](#)), typically partners with Bendless (*Ben*), the *Drosophila* homolog of mammalian Ubc13 (Muralidhar and Thomas, 1993 [↗](#); Oh et al., 1994 [↗](#)), to perform the E2 enzyme function. Ovaries with the *bam>cdc27-RNAi*, *bam>cdc27-RNAi+uev1a-RNAi*, or *bam>cdc27-RNAi+ben-RNAi* genotype showed minimal degradation of egg chambers (**Figure 2—figure supplement 4** [↗](#)). However, degraded egg chambers were prevalent in *bam>cdc27-RNAi+ben-RNAi; uev1a*^{Δ1/+} and *bam>cdc27-RNAi+ben-RNAi; uev1a*^{Δ2/+} ovaries, where Uev1A, *Ben*, and *Cdc27* are all downregulated (**Figure 2—figure supplement 4** [↗](#)). These results suggest that Uev1A, *Ben*, and the APC/C complex also work together to protect nurse cells from death during normal oogenesis.

Uev1A and the APC/C complex work together to degrade CycA via the proteasome

The APC/C complex is known to collaborate with the E2 enzymes UBE2C (*Drosophila* homolog: Vihar) and UBE2S (*Drosophila* homolog: Ube2S) to facilitate the proteasomal degradation of CycA during cell cycle progression (Greil et al., 2022 [↗](#); Yamano, 2019 [↗](#)). This degradation involves the assembly of branched ubiquitin chains, incorporating K11, K48, and K63 linkages through a two-step mechanism (Meyer and Rape, 2014 [↗](#)). However, the involvement of Uev1A in this process remains unexplored. To explore it, we performed biochemical assays in cultured *Drosophila* Schneider 2 (S2) cells, overexpressing tag-fused proteins using *act-GAL4* to enhance expression efficiency. Co-immunoprecipitation (co-IP) assays revealed that Uev1A interacts with several components of the APC/C complex, including Mr, Cdc16 (*Drosophila* APC6), and Cdc23 (*Drosophila* APC8). In addition, Cdc27 was found to interact with CycA (**Figure 3A, B** [↗](#) and **Figure 3—figure supplement 1** [↗](#)). Remarkably, RNAi targeting *uev1a*, *ben*, or *cdc27* significantly stabilized CycA proteins after the treatment with cycloheximide (CHX), an inhibitor of protein synthesis (**Figure 3C-E** [↗](#)). Since the K63 linkage primarily mediates lysosomal degradation via autophagy (Kwon and Ciechanover, 2017 [↗](#)), we tested CycA stabilization with a chloroquine (CQ, a lysosome inhibitor) or MG132 (a proteasome inhibitor) treatment. Notably, MG132, but not CQ, treatment significantly stabilized CycA proteins (**Figure 3F** [↗](#)). These results suggest that Uev1A, Ben, and Cdc27 cooperate to promote CycA degradation through the proteasome rather than the lysosome.

To directly validate this, we performed ubiquitination assays for CycA in S2 cells. Cdc27 significantly enhanced CycA ubiquitination, while this effect was markedly reduced upon the knockdown of *uev1a*, *ben*, or both (**Figure 3G** [↗](#)). This result indicates that both Uev1A and Ben are essential for APC/C-mediated proteasomal degradation of CycA. Furthermore, we explored the roles of the seven lysine (K) residues in ubiquitin for polyubiquitin chain formation. The 7KR (lysine to arginine) mutation completely abolished Cdc27-promoted polyubiquitination of CycA. Intriguingly, the 6KR+K48 mutation, in which all K residues except K48 were mutated to R residues, failed to restore polyubiquitination. In contrast, either 6KR+K11 or 6KR+K63 mutation significantly restored polyubiquitination (**Figure 3H** [↗](#)). These findings suggest that the K11 and K63 linkages are primarily responsible for CycA polyubiquitination in *Drosophila* cells. Together with previous studies (Hofmann and Pickart, 1999 [↗](#); McKenna et al., 2001 [↗](#)), we propose that Uev1A and Ben mediate the K63 linkage in this process.

Uev1A inhibits the overgrowth of *Drosophila* germline tumors driven by oncogenic *Ras*^{G12V}

Given the absence of cell division in normal polyploid nurse cells (Hammond and Laird, 1985 [↗](#)), their death induced by division-promoting *Ras*^{G12V} represents an artificial stress. Notably, Uev1A was not upregulated in response to this stress (**Figure 2—figure supplement 2E** [↗](#)), and it executed the function through degrading CycA (**Figure 2** [↗](#) and **3** [↗](#)). These findings prompted us to investigate whether Uev1A also counteracts oncogenic *Ras*-driven tumorigenesis in diploid cells, which undergo normal cell division. Our prior research demonstrated that oncogenic *Ras*^{G12V} markedly promotes the overgrowth of diploid *bam*-deficient germline tumors (Zhang et al., 2024 [↗](#)), which are highly resistant to cell death (Zhang et al., 2023 [↗](#); Zhao et al., 2018 [↗](#)). Intriguingly, knocking down *uev1a* significantly enhanced the overgrowth of these tumors, while overexpressing Uev1A suppressed it (**Figure 4A, B** [↗](#)). These results indicate that Uev1A also plays a role in counteracting oncogenic *Ras*-driven tumorigenesis in diploid cells.

Considering the therapeutic potential of gene upregulation in inhibiting tumor growth, a critical concern is its impact on normal physiological processes. In this study, we examined the impact of Uev1A overexpression on *Drosophila* oogenesis and GSC maintenance. Notably, the *nos>Uev1A* flies remained fertile, and their ovaries appeared morphologically similar to those of the *nos>lacZ*




Figure 3.

Uev1A, Ben, and Cdc27 work together to degrade CycA through the proteasome.

(A and B) co-IP results. (C) RNAi efficiency experiments. The #1 dsRNAs were employed in RNAi assays targeting *uev1a*, *ben*, and *cdc27* in (D, E, and G). (D-F) CycA stability assays. In (E), the relative levels of CycA proteins were quantified using the following formula: (Mean gray value of the CycA/ β -Actin band at n hours post-treatment) $\div$ (Mean gray value of the CycA/ β -Actin band at 0 hour). Three independent replicates were conducted at each timepoint, and statistical significance was determined using two-way ANOVA with multiple comparisons: **** ($P < 0.0001$). In (F), cells harvested six hours post-treatment were analyzed. (G and H) Ubiquitination assays for CycA in S2 cells.

The online version of this article includes the following source data and figure supplement for **figure 3**:

Source data 3. Raw quantification data.

Figure supplement 1. Co-IP results.




Figure 4.

Uev1A inhibits the overgrowth of germline tumors induced by oncogenic *Ras*^{G12V}.

(A and C) Representative ovaries. All images in (A) are of the same magnification. Scar bars in (C): 200 μm . (B) Quantification data for ovarian size. The largest 2D area of each ovary in a single confocal focal plane was scanned, and its size was measured using ImageJ (NIH, Bethesda, MD, USA). 30 ovaries from 3-day-old flies were analyzed for each genotype. (D) Representative samples. The GSCs within stem cell niches are outlined by yellow dashed lines. Both images are of the same magnification. (E) Quantification data for GSC numbers per germarium. Germ cells that directly contact cap cells and contain dot-like spectrosomes were counted as GSCs. 100 germaria from 14-day-old flies were quantified for each genotype. In (B and E), statistical significance was determined using t test: *n.s.* ($P > 0.05$) and * ($P < 0.05$). (F) Working model. The E2 Uev1A/Ben complex, ubiquitin, and the E3 APC/C complex work together to degrade CycA, thereby preventing apoptosis or senescence, while also inhibiting cell overgrowth induced by oncogenic *Ras* through the MAPK signaling.

The online version of this article includes the following source data for **figure 4**:

Source data 2. All genotypes.

Source data 3. Raw quantification data.

control flies (**Figure 4C**), indicating normal oogenesis. Furthermore, each germarium in the ovaries of 14-day-old *nos>Uev1A* flies contained a similar number of GSCs as those in the *nos>lacZ* control flies (**Figure 4D, E**). These results suggest that Uev1A overexpression does not disrupt normal oogenesis and GSC maintenance.

UBE2V1 and UBE2V2 inhibit the overgrowth of human colorectal tumors driven by oncogenic *KRAS*

Our findings in *Drosophila* prompted us to explore the tumor-suppressive effects of UBE2V1 and UBE2V2 on human tumor growth. Using the Kaplan-Meier plotter (<https://kmplot.com/analysis>), we first evaluated the correlation between UBE2V1/2 expression and prognosis in several types of *RAS*-mutant cancer patients, including melanoma, myeloma, lung cancer, and colorectal cancer. Among them, higher expression levels of UBE2V1/2 were significantly associated with improved overall survival and relapse-free survival in *KRAS*-mutant colorectal cancer patients (**Figure 5A**). RNA-seq data from The Cancer Genome Atlas (TCGA) showed that *UBE2V1* and *UBE2V2* are transcribed at similar levels in colorectal cancer patients with oncogenic *RAS* mutations (including *KRAS*, *HRAS*, and *NRAS*) as in those without such mutations (**Figure 5B**). These findings suggest that UBE2V1 and UBE2V2 are not upregulated in response to oncogenic *RAS* mutations, paralleling the behavior of Uev1A in *Drosophila* ovarian nurse cells (**Figure 2—figure supplement 2E**).

To investigate the potential tumor-suppressive effects of UBE2V1 and UBE2V2 on colorectal cancer cells, we overexpressed them in human SW480 cells (carrying the *KRAS*^{G12V} mutation) and HCT116 cells (carrying the *KRAS*^{G13D} mutation), respectively. Overexpression efficiency was confirmed by Western blot analysis (**Figure 5—figure supplement 1A**). Proliferation assays using 5-ethynyl-2'-deoxyuridine (EdU) demonstrated that overexpression of either UBE2V1 or UBE2V2 significantly inhibited the proliferation of *KRAS*-mutant colorectal cancer cells *in vitro* (**Figure 5C, D**). This inhibitory effect was further corroborated by clonogenic and cell viability (CCK8) assays, which showed that UBE2V1 or UBE2V2 overexpression markedly impaired cell colony formation and survival (**Figure 5E–G**).

To validate the tumor-suppressive effects of UBE2V1 and UBE2V2 *in vivo*, we generated stable SW480 cell lines overexpressing either of them, with overexpression efficiency confirmed by Western blot analysis (**Figure 5—figure supplement 1B**). Subcutaneous tumorigenesis assays in *Balb/c* nude mice showed that UBE2V1 or UBE2V2 overexpression significantly suppressed tumor formation (**Figure 5H, I**). Immunohistochemical staining further revealed a marked reduction in Cyclin A expression and Ki-67 positivity rates in tumors overexpressing either UBE2V1 or UBE2V2 (**Figure 5J, K**). Notably, the tumor-suppressive effect of UBE2V1 was more pronounced than that of UBE2V2, aligning with their ability to protect against the nurse cell death induced by oncogenic *Ras*^{G12V} in *Drosophila* (**Figure 1K, L**). These results provide strong evidence for the tumor-suppressive effects of UBE2V1 and UBE2V2 on *KRAS*-mutant colorectal cancer.

Discussion

It is well established that oncogenic *Ras* can induce DNA replication stress, leading to cellular senescence or death (Hills and Diffley, 2014; Kotsantis et al., 2018). However, our previous (Zhang et al., 2024) and current findings demonstrated that oncogenic *Ras*^{G12V} can also trigger cell death in polyploid *Drosophila* ovarian nurse cells through aberrantly promoting their division. In this study, we performed a genome-wide genetic screen and identified the E2 enzyme Uev1A as a crucial protector against this specific form of cellular stress. Mechanistically, Uev1A collaborates with the APC/C complex (E3) to facilitate the proteasomal degradation of CycA, which overexpression alone can also trigger nurse cell death. Furthermore, Uev1A and its human homologs, UBE2V1 and UBE2V2, counteract oncogenic *Ras*-driven tumorigenesis in diploid




Figure 5.

Prognostic significance and tumor-suppressive effects of UBE2V1 and UBE2V2 on *KRAS*-mutant colorectal cancer.

(A) Kaplan-Meier analysis of overall survival and relapse-free survival in *KRAS*-mutant colorectal cancer patients with high or low expression levels of UBE2V1 and UBE2V2. (B) TCGA analysis comparing UBE2V1 and UBE2V2 expression levels in colorectal cancer patients with and without *RAS* mutations. The TCGA patient data were downloaded from the UCSC Xena website: the RNA-seq data (Version: 05-09-2024) and the somatic mutation data (Version: 08-05-2024). Statistical significance was determined using t test: *n.s.* ($P > 0.05$). (C and D) EdU incorporation assays to assess the effects of UBE2V1 and UBE2V2 overexpression (OE) on cell proliferation in SW480 and HCT116 cells. Empty OE vector was used as the control. All images in (C) are of the same magnification. (E-G) Assays to evaluate the effects of UBE2V1- and UBE2V2-OE on colony formation and cell viability in SW480 and HCT116 cells. (H and I) Subcutaneous tumorigenesis assays in nude mice, where tumors were excised, photographed, and weighed 28 days after tumor cell injection. (J and K) Immunohistochemical staining to assess CycA expression and Ki-67 positivity in tumor tissues. All images in (J) are of the same magnification. In (D, F, I-2, and K), statistical significance was determined using one-way ANOVA. In (G and I-1), statistical significance was determined using two-way ANOVA with multiple comparisons: * ($P < 0.05$), ** ($P < 0.01$), *** ($P < 0.001$), and **** ($P < 0.0001$).

The online version of this article includes the following source data and figure supplement for **figure 5**:

Source data 3. Raw quantification data.

Figure supplement 1. Validation of UBE2V1 and UBE2V2 overexpression in colorectal cancer cell lines.

Drosophila germline and human colorectal tumor cells, respectively (**Figure 4F**). These findings highlight the critical role of Uev1A in counteracting oncogenic *Ras* stimuli in both polyploid and diploid cells.

Our studies reveal that the DDR pathway protects against the nurse cell death induced by oncogenic *Ras*^{G12V} (**Figure 2—figure supplement 2A, B**), suggesting that this cellular stress both activates and is subsequently suppressed by the DDR. In contrast, p53 promotes nurse cell death under the same conditions (**Figure 2—figure supplement 2A, B**). Interestingly, previous research has also demonstrated distinct roles of p53 and Lok, a crucial DDR regulator, in regulating nurse cell death during mid-oogenesis. While Lok overexpression triggers nurse cell death, p53 overexpression does not. Furthermore, Lok-induced nurse cell death remains unaffected by p53 mutation, indicating that its mechanism operates independently of p53 (Bakhrat et al., 2010). The underlying mechanisms driving these differences warrant further investigation.

Previous studies have shown that Uev1A promotes cell death through activating JNK signaling (Ma et al., 2013). If this mechanism also applied to the nurse cell death induced by oncogenic *Ras*^{G12V}, we would expect a reduction in Uev1A levels to suppress this cell death. However, our observations revealed the opposite effect (**Figure 1F-I**). Additionally, we did not detect the expression of *puc-lacZ*, a reporter for JNK signaling (Glise et al., 1995), in *nos>Ras*^{G12V} nurse cells (data not shown). These findings suggest that Uev1A's protective role in this context is independent of JNK signaling.

Of note, the suppressive effects of Uev1A on *Ras*^{G12V}; *bam*^{RNAi} germline tumors in *Drosophila* were less pronounced than that of UBE2V1/2 on oncogenic *KRAS*^{G12V}-driven human colorectal tumor xenografts in nude mice (**compare Figure 4A, B** with **Figure 5H, I**). Our previous research has shown that *Ras*^{G12V}; *bam*^{-/-} germline tumor cells divide infrequently (Zhang et al., 2024). This suggests that the relatively weak suppression observed in these tumors may be due to the fact that Uev1A's mechanism of action is cell cycle-dependent: the more frequently tumor cells divide, the more effectively Uev1A can suppress tumor overgrowth. We did not observe significant negative effects of Uev1A overexpression on GSC maintenance in *Drosophila* ovaries (**Figure 4D, E**). This is likely because GSCs also divide infrequently (Morris and Spradling, 2011). Therefore, these findings underscore the importance of considering the potential negative effects of Uev1A/UBE2V1/2 overexpression in other contexts, particularly when cells divide rapidly.

RAS oncoproteins have long been considered "undruggable", due to the lack of deep-binding, targetable pockets (Moore et al., 2020). To date, targeted therapies have been approved only for *KRAS*^{G12C} (Canon et al., 2019; Ou et al., 2022; Skoulidis et al., 2021). Recently, small-molecule pan-KRAS degraders, developed using the proteolysis-targeting chimeras (PROTACs) strategy, have shown promise (Popow et al., 2024), though their clinical effectiveness and safety remain to be fully validated. Alternatively, targeting key regulators in the RAS signaling pathway—such as SOS, SHP2, Farnesyltransferase, Raf, MEK, ERK, and PI3K—has emerged as an attractive therapeutic approach (Moore et al., 2020). Recent research has also explored the strategy of hyperactivating oncogenic RAS to trigger cell death as a potential means to suppress the overgrowth of human colon tumor xenografts in nude mice (Dias et al., 2024). Additionally, targeting the cellular dependencies driven by oncogenic RAS may trigger RAS-specific synthetic lethality, presenting a promising therapeutic approach (Moore et al., 2020). Our findings highlight the tumor-suppressive effects of UBE2V1 and UBE2V2 on human colorectal tumors driven by oncogenic *KRAS* (**Figure 5**). Notably, both of these two E2 enzymes are relatively small, with UBE2V1 consisting of 147 amino acid residues and UBE2V2 145 residues (**Figure 1J**). This raises the exciting possibility that their upregulation, through mRNA delivery or small-molecule agonists, could offer a promising therapeutic approach for human cancer.

Materials and Methods

Key resources table

Reagent or resource	Source	Identifier
Antibodies		
mouse anti- α -Spectrin	Developmental Studies Hybridoma Bank, DSHB	3A9
mouse anti- β -Actin	Abmart	M20011
mouse anti- β -Actin	Zenbio	200068-8F10
mouse anti-Flag	Sigma	F1804
mouse anti-Flag	Utibody	UM3009
mouse anti-HA	Utibody	UM3004
mouse anti-Myc	Utibody	UM3011
mouse anti-UBE2V2	Santa Cruz Biotechnology	sc-377254
rabbit anti- α -Tubulin	Proteintech	14555-1-AP
rabbit anti- β -Tubulin	Zenbio	380628
rabbit anti-CycA	Immunoway	YT1167
rabbit anti-Flag	Zenbio	R24091
rabbit anti-HA	Zenbio	301113
rabbit anti-Ki67	Proteintech	27309-1-AP
rabbit anti-Myc	ABclonal	AE009
rabbit anti-UBE2V1	Wanleibio	WL04482
Alexa Fluor 546 goat anti-mouse	Invitrogen	A-11030
HRP Goat anti-Mouse IgG(H+L)	SIMUBIOTECH	S2002
HRP Goat anti-Rabbit IgG(H+L)	SIMUBIOTECH	S2001

Cell lines		
<i>Drosophila</i> Schneider 2 (S2) cells	Duojia Pan	N/A
human 293T cells	The American Type Culture Collection, ATCC	N/A
human HCT116 cells	ATCC	N/A
human SW480 cells	ATCC	N/A
Chemicals, solution, and kits		
BCA Protein Assay Kit	Beyotime Biotechnology	P0012
CCK8 Kit	APExBIO	K1018
Chloroquine (CQ)	Selleck	S6999
Cycloheximide (CHX)	MCE	HY-12320
DMSO	Macklin	D6258
Dulbecco's Modified Eagle Medium	Gibco	C11995500BT
EdU Incorporation Assay Kit	Beyotime Biotechnology	C0075
Fetal Bovine Serum (FBS)	Lonsera	S711-001S
Insect Culture Medium	Union	UK1000
Lipofectamine 2000	ThermoFisher	11668027
MG132	Selleck	S2619
Protein G Sepharose	Cytiva	17061801
Puromycin	Solarbio	P8230
T7 RiboMAX™ Express RNAi System	Promega	P1700
X-tremeGENE HP DNA Transfection Reagent	Roche	6366546001
<i>Drosophila</i> strains		
<i>bam-GAL4-VP16</i>	Erika Matunis	N/A
<i>nos-GAL4-VP16</i>	Ruth Lehmann	N/A

<i>nos-cas9; FRT2A</i>	this study	N/A
<i>nos-cas9 (on X)</i>	Zheng Guo	N/A
<i>puc-lacZ</i>	Bloomington <i>Drosophila</i> Stock Center, BDSC	98329
<i>UAS-bam-RNAi</i>	BDSC	33631
<i>UASp-BicD-RNAi</i>	TsingHua Fly Center, THFC	THU4454
<i>UASp-cdc27-RNAi</i>	THFC	TH201500102.S
<i>UASp-Cdk1</i>	BDSC	65396
<i>UASp-CycA</i>	BDSC	85308
<i>UASp-CycB</i>	BDSC	85312
<i>UASp-dsor1-RNAi</i>	THFC	THU0677
<i>UASp-fzr-RNAi</i>	THFC	TH201500745.S
<i>UASp-GFP</i>	Michael Buszczak	N/A
<i>UASp-GFP-RNAi</i>	BDSC	44412, 44415
<i>UASz-lacZ</i>	Shaowei Zhao	N/A
<i>UASp-ImgA-RNAi</i>	THFC	THU4085
<i>UASp-lok-RNAi</i>	THFC	TH01867.N
<i>UASp-mr-RNAi</i>	THFC	THU5250
<i>UASp-p53-RNAi</i>	THFC	THU5318
<i>UASp-Ras^{G12V}</i>	Shaowei Zhao	N/A
<i>UASp-rl-RNAi</i>	THFC	THU3530
<i>UASp-shtd-RNAi</i>	THFC	TH201500835.S
<i>UASp-Stg</i>	BDSC	58439
<i>UASp-tefu-RNAi</i>	THFC	THU5591

<i>UASp-uev1a-RNAi</i>	BDSC	66947
<i>UASz-flag-Ras^{G12V}</i>	this study	N/A
<i>UASz-UBE2V1</i>	this study	N/A
<i>UASz-UBE2V2</i>	this study	N/A
<i>UASz-uev1a</i>	this study	N/A
<i>UASz-Yki^{3SA}</i>	Shaowei Zhao	N/A
<i>y v nos-int; attP40</i>	this study	N/A
<i>y w nos-int; attP40</i>	BDSC	79604
<i>uev1a^{Δ1}</i>	this study	N/A
<i>uev1a^{Δ2}</i>	this study	N/A
<i>uev1a-flag</i>	this study	N/A
All Deficiency <i>Drosophila</i> strains (including 7584)	The Core Facility of <i>Drosophila</i> Resource and Technology (CEMCS), Chinese Academy of Sciences (CAS), China	
Mouse strain		
<i>Balb/c</i> (CAnN.Cg-Foxn1 ^{nu} /CrI) nude mice	Beijing Vital River Laboratory Animal Technology Co., Ltd.	401
Plasmids		
<i>pCDH-CMV</i>	Addgene	72265
<i>pCDH-CMV-UBE2V1</i>	this study	N/A
<i>pCDH-CMV-UBE2V2</i>	this study	N/A
<i>pCFD3</i>	Addgene	49410

<i>pCFD3-uev1a-gRNA-1</i>	this study	N/A
<i>pMD2.G</i>	Addgene	12259
<i>psPAX2</i>	Addgene	12260
<i>pU6-BbsI-chiRNA</i>	Eric H. Baehrecke	N/A
<i>pU6-uev1a-gRNA-2</i>	this study	N/A
<i>pUAS_t-attB</i>	GenBank	EF362409
<i>pUAS_t-APC7-Myc</i>	this study	N/A
<i>pUAS_t-Flag-ben</i>	this study	N/A
<i>pUAS_t-Flag-cycA</i>	this study	N/A
<i>pUAS_t-HA-Ub</i>	this study	N/A
<i>pUAS_t-HA-Ub^{7KR}</i>	this study	N/A
<i>pUAS_t-HA-Ub^{6KR+K11}</i>	this study	N/A
<i>pUAS_t-HA-Ub^{6KR+K48}</i>	this study	N/A
<i>pUAS_t-HA-Ub^{6KR+K63}</i>	this study	N/A
<i>pUAS_t-HA-uev1a</i>	this study	N/A
<i>pUAS_t-Myc-APC4</i>	this study	N/A
<i>pUAS_t-Myc-Cdc16</i>	this study	N/A
<i>pUAS_t-Myc-Cdc23</i>	this study	N/A
<i>pUAS_t-Myc-Cdc27</i>	this study	N/A
<i>pUAS_t-Myc-Fzr</i>	this study	N/A
<i>pUAS_t-Myc-Fzy</i>	this study	N/A
<i>pUAS_t-Myc-Ida</i>	this study	N/A
<i>pUAS_t-Myc-Mr</i>	this study	N/A
<i>pUASz1.0</i>	<i>Drosophila</i> Genomics Resource Center, DGRC	1431

<i>pUASz-flag-Ras^{G12V}</i>	this study	N/A
<i>pUASz-UBE2V1</i>	this study	N/A
<i>pUASz-UBE2V2</i>	this study	N/A
<i>pUASz-uev1a</i>	this study	N/A
<i>pUASz-uev1a-donor</i>	this study	N/A
<i>pUASz-Yki^{3SA}</i>	this study	N/A
Software		
Adobe Photoshop 2022	San Jose, CA, USA	
ImageJ	NIH	
Prism	Graphpad Software, Inc.	
ZEN 3.0 SR imaging software	Carl Zeiss	

Data availability

The results of the deficiency screening are provided in **Source data 1**. All genotypes are listed in **Source data 2**, and the raw quantification data can be found in **Source data 3**.

Fly husbandry

Cross experiments were conducted at 25°C, except for some performed at 29°C as noted.

Transgenic flies

UASz-flag-Ras^{G12V}, UASz-uev1a, UASz-UBE2V1, and UASz-UBE2V2

The coding sequences (CDSs) of *flag-Ras^{G12V}*, *uev1a-RA*, *UBE2V1*, and *UBE2V2* were cloned into the *pUASz1.0* vector (DeLuca and Spradling, 2018 [DOI](#)). These plasmids were then microinjected into fertilized fly embryos with the genotype “*y w nos-int; attP40*”, resulting in the generation of transgenic fly strains. The DNA sequence encoding 3xFlag was as follows:
ATGGACTACAAAGACCATGACGGTGATTATAAAGATCATGACATCGATTACAAGGAT
GACGATGACAAGCTT.

uev1a^{Δ1} and uev1a^{Δ2} mutants

The gRNA targeting the second coding exon of the *uev1a* gene was designed with the DNA sequence “GATCCAGCTCCTCCAGTAAGCGG” and cloned into the *pCFD3* vector (Port et al., 2014 [DOI](#)). The plasmids were then microinjected into fertilized fly embryos with the genotype “*y w nos-int; attP40*” to generate transgenic fly strain. These transgenic flies were subsequently crossed with “*nos-cas9; FRT2A*” flies to generate *uev1a* knockout mutants, which molecular information was identified by Sanger sequencing.

uev1a-flag knock in

The gRNA targeting the second-to-last coding exon of the *uev1a* gene was designed with the DNA sequence “GTTATAAACCGGAGCGTTGGTGG” and cloned into the *pU6-BbsI-chiRNA* vector. The homology arms consisted of 991 bp upstream of the stop codon (“TAG”), followed by a 3xFlag tag

sequence, the “TAG” stop codon, and 979 bp downstream of “TAG”. To prevent cleavage by the gRNA, the targeting sequence within the upstream homology arm was mutated to “CCTACCCTGCGCTTCATTA”, ensuring that the amino acid sequence remained unaltered. These DNA sequences were synthesized into the *pUASz1.0* vector between the *Bam*HI and *Kpn*I sites and used as a double-stranded DNA (dsDNA) donor for homologous recombination-mediated repair. The *pU6-uev1a-gRNA-2* (100 ng/μL) and *pUASz-uev1a-donor* (100 ng/μL) plasmids were then co-injected into fertilized fly embryos with the genotype “*nos-cas9 (on X)*”. The resulting *uev1a-flag* knock-in fly strain was identified by PCR and confirmed by Sanger sequencing.

Immuno-fluorescent staining

Fly ovaries were dissected in the PBS solution, fixed by the 4% paraformaldehyde solution (diluted in PBS) for 3 hours, washed by the PBST solution (0.3% Triton X-100 diluted in PBS) for 1 hour at room temperature (RT), then incubated with the primary antibodies (diluted in PBST) overnight at 4°C, washed by the PBST solution for 1 hour at RT, incubated with the secondary antibodies and 0.1 μg/mL DAPI (diluted in PBST) overnight at 4°C, washed by the PBST solution for 1 hour at RT, and subsequently mounted using 70% glycerol (autoclaved). The primary antibodies were used at the concentrations as follows: mouse anti-Flag (Sigma, #F1804) at 1:500 and mouse anti-α-Spectrin at 1:100. The Alexa Fluor conjugated secondary antibodies were used at 1:2,000.

Construction of plasmids used in S2 cells

The CDSs of the genes were amplified from the cDNAs derived from either S2 cells or adult wild-type (*w¹¹¹⁸*) flies and subsequently cloned into the *pUAS-attB* vector. N- or C-terminal tags were incorporated based on prior studies or structural predictions from AlphaFold. The following proteins were tagged at the N-terminus: APC4, Ben, CDC16, CDC23, Cdc27, CycA, Fzr, Fzy, Ida, Mr, Ub, Ub^{7KR}, Ub^{6KR+K11}, Ub^{6KR+K48}, Ub^{6KR+K63}, and Uev1A. In contrast, APC7 was tagged at the C-terminus.

Culture and transfection of S2 cells

S2 cells were maintained in insect culture medium supplemented with 10% FBS and incubated at 27°C without CO₂. For transient transfection, the X-tremeGENE HP DNA transfection reagent was used. Cells were harvested 36 hours post-transfection for further analysis.

RNAi assay in S2 cells

Two double-stranded RNAs (dsRNAs) targeting distinct regions of the relevant genes were synthesized using the T7 RiboMAX™ Express RNAi System. S2 cells were seeded in 6-well tissue culture plates and treated with 15 μg of dsRNA in the culture medium, followed by incubation for 24 hours. Expression plasmids were then transiently transfected into the cells, after which an additional 10 μg of dsRNA was added. The cells were cultured for another 36 hours before harvesting. As a negative control, dsRNA targeting the *AcGFP* gene was used. The templates for dsRNA synthesis were generated by PCR amplification from S2 cell genomic DNA using the following primers:

uev1a#1-F: GAATTAATACGACTCACTATAGGGAGAACGGAATTTCCGCTTACTG

uev1a#1-R: GAATTAATACGACTCACTATAGGGAGACGGACCGATGATCATGCC

uev1a#2-F: GAATTAATACGACTCACTATAGGGAGACACTAAAGATCGAGTGCG

uev1a#2-R: GAATTAATACGACTCACTATAGGGAGATGCCAGCTTCAGGTTCTC

ben#1-F: GAATTAATACGACTCACTATAGGGAGACCACGTCGCATCATCAAG

ben#1-R: GAATTAATACGACTCACTATAGGGAGAAGTCTTCGACGGCATATTTTC

ben#2-F: GAATTAATACGACTCACTATAGGGAGACAGATCCGGACCATATTG

ben#2-R: GAATTAATACGACTCACTATAGGGAGATCAGTCTTCGACGGCATATTTTC

cdc27#1-F: GAATTAATACGACTCACTATAGGGAGATCGCCAGGATCTGATTAAC

cdc27#1-R: GAATTAATACGACTCACTATAGGGAGAGCAGCGACAGATCCTTCTTC

cdc27#2-F: GAATTAATACGACTCACTATAGGGAGAGATGATGGGCAAAAAGCTAAAG

cdc27#2-R: GAATTAATACGACTCACTATAGGGAGACCATCGGCCGATTGTTTC

AcGFP-F: GAATTAATACGACTCACTATAGGGAGATGCACCACCGCAAGCTGCCTG

AcGFP-R: GAATTAATACGACTCACTATAGGGAGAGGCCAGCTGCACGCTGCCATC

Immunoprecipitation (IP), immunoblotting (ib), *in vivo* ubiquitination and protein stability assays in S2 cells

IP assay

The transfected S2 cells were lysed using NP-40 lysis buffer supplemented with protease inhibitors at 4°C for 30 minutes. The supernatants were incubated with primary antibodies at 4°C for 3 hours, followed by incubation with Protein G Sepharose for 2 hours. The beads were washed five times with NP-40 lysis buffer and subsequently analyzed by Western blotting. The following antibodies were used: mouse anti-Flag (Utibody, #UM3009) at 1:200, mouse anti-Myc at 1:200, and mouse anti-HA at 1:200.

IB assay

After 36 hours of transfection, the S2 cells were harvested and lysed in 2×SDS protein loading buffer [0.25 M Tris-HCl (pH 6.8), 78 mg/mL DTT, 100 mg/mL SDS, 50% glycerol, and 5 mg/mL bromophenol blue]. The lysates were then boiled for 5 minutes and subjected to western blot analysis. The following antibodies were used: mouse anti-Flag (Utibody, #UM3009) at 1:5000, rabbit anti-Flag at 1:5000, rabbit anti-Myc at 1:3000, rabbit anti-HA at 1:3000, mouse anti-β-Actin (Zenbio, #200068-8F10) at 1:5000, and rabbit anti-β-Tubulin at 1:5000.

***in vivo* ubiquitination assay**

30 hours post-transfection, S2 cells were treated with MG132 at a final concentration of 50 μM for 6 hours. Proteins were subsequently enriched through immunoprecipitation and analyzed by Western blot analysis.

Protein stability assay

S2 cells were treated with 20 μg/mL of the protein synthesis inhibitor CHX for specified time intervals prior to harvesting.

Human cell culture

The human colon cancer cell lines SW480 and HCT116, as well as the human embryonic kidney cell line 293T, were purchased from the American Type Culture Collection (ATCC) and cultured in Dulbecco's Modified Eagle Medium supplemented with 10% FBS. The cultures were maintained at

37°C with 5% CO₂ in a humidified incubator. The lentiviral vector *pCDH-CMV* was used for gene overexpression, while *pMD2.G* and *psPAX2* served as lentiviral packaging plasmids. Target plasmids, *pCDH-CMV-UBE2V1* and *pCDH-CMV-UBE2V2*, were synthesized by the GENEWIZ company (Suzhou, China). Plasmids were transfected into 293T cells using Lipofectamine 2000 to generate lentiviral particles, according to the manufacturer's protocol. SW480 and HCT116 cells were then transduced with these lentiviruses to establish stable cell lines expressing UBE2V1 and UBE2V2, respectively. To select for stably transduced cells, the cultures were maintained in medium containing 2 µg/mL puromycin.

EdU incorporation assay

Cells were seeded in 24-well plates at a density of 100,000 cells per well. Cell proliferation was assessed using the EdU incorporation assay kit, following the manufacturer's instructions. Briefly, EdU reagent was added to the culture medium, and cells were incubated for 2 hours to label DNA-synthesizing cells. After incubation, the medium containing EdU was removed, and cells were washed with PBS, fixed with fixative solution for 10 minutes, and then subjected to a click reaction for fluorescence labeling. Fluorescence microscopy was used to capture images and calculate the percentage of EdU⁺ cells.

Colony formation assay

Cells were seeded in 6-well plates at a density of 500 cells per well and cultured at 37°C in a 5% CO₂ incubator for 10 days without medium change. After 10 days, cells were gently washed with PBS to remove non-adherent cells, then stained with 0.5% crystal violet solution (containing 10% methanol and 1% acetic acid) for 15 minutes. The plates were subsequently rinsed with water to remove excess stain and air-dried. Colonies were photographed and counted using image analysis software.

CCK8 cell viability assay

Cells were seeded in 96-well plates at a density of 2,000 cells per well. Cell viability was assessed over five days using a CCK8 kit following the manufacturer's instructions. Measurements were taken on Day 0 (baseline) and daily from Day 1 to Day 4. At each time point, 10 µL of CCK8 solution was added to each well, and the plates were incubated for 2 hours at 37°C in a humidified incubator. Absorbance at 450 nm was measured using a microplate reader to assess cell viability and proliferation dynamics throughout the experimental period.

Western blot analysis

Cells were lysed with RIPA buffer containing protease inhibitors, and protein concentrations were determined using a BCA Protein Assay Kit. Protein samples (20 µg per lane) were separated by SDS-PAGE and transferred to PVDF membranes. Membranes were blocked with 5% non-fat milk and incubated overnight at 4°C with primary antibodies: mouse anti-β-actin (Abmart, #M20011) at 1:5,000, rabbit anti-α-Tubulin at 1:5,000, rabbit anti-UBE2V1 at 1:2,000, and mouse anti-UBE2V2 at 1:2,000. Following washes, membranes were incubated with HRP-conjugated secondary antibodies and developed using ECL. Band intensities were quantified by densitometry.

Immunohistochemistry assay

Tissue sections were deparaffinized with xylene, rehydrated through a graded ethanol series, and subjected to antigen retrieval in citrate buffer (pH 6.0) at 98°C for 10 minutes. Endogenous peroxidase activity was blocked using 3% hydrogen peroxide, and nonspecific binding was blocked with 5% BSA. Sections were incubated overnight at 4°C with primary antibodies: rabbit anti-CycA at 1:200 and rabbit anti-Ki67 at 1:2,000. After incubation with HRP-conjugated secondary antibodies, sections were developed using DAB, counterstained with hematoxylin, and mounted. Images were captured using a light microscope.

Animal experiment

Male *Balb/c* nude mice (7-week-old) were used to assess the tumorigenic potential of SW480 colon cancer cells. The mice were housed under specific pathogen-free conditions with a 12-hour light/dark cycle and had ad libitum access to food and water. SW480 cells were stably transfected with either the empty vector *pCDH-CMV* (control), *pCDH-CMV-UBE2V1* (UBE2V1-OE), or *pCDH-CMV-UBE2V2* (UBE2V2-OE), and 5×10^6 cells were subcutaneously injected into the right flank of each mouse. Each group consisted of six mice. Tumor growth was monitored approximately one week after injection. Tumor volumes were measured every four days using calipers and calculated using the formula: volume = length $\times$ width²/2. The experiment was terminated when tumors reached $\sim 1,000$ mm³, and mice were humanely euthanized according to institutional and national ethical guidelines.

Image collection and processing

Fluorescent images were captured using a Zeiss LSM 710 confocal microscope (Carl Zeiss AG, BaWü, GER) and processed with ZEN 3.0 SR imaging software (Carl Zeiss) and Adobe Photoshop 2022 (San Jose, CA, USA).

Acknowledgements

We gratefully thank Eric H. Baehrecke, Michael Buszczak, Zheng Guo, Ruth Lehmann, Erika Matunis, Duoqia Pan, Zhaohui Wang, Wei Xiao, Addgene, ATCC, BDSC, CEMCS, DGRC, GenBank, and THFC for providing plasmids, cell lines, and fly strains. This study was supported by grants 32270841 and 32070871 from the National Natural Science Foundation of China (NSFC) to Shaowei Zhao.

Additional information

Author contributions

S.Z. conceived the project. S.Z., S.W., and H.Z. supervised the studies. Q.Z., Yun.W., X.F., Z.W., Y.Z., L.Y., Yue.W., M.Y., D.S., and R.Z. performed the experiments. S.Z. wrote the manuscript and all authors commented on it.

Figure 1-figure supplement 1.

Uev1A protects against the nurse cell death induced by oncogenic Yki^{3SA} .

(A) Representative ovaries. The red arrows denote degrading egg chambers. Scale bars: 200 μ m. (B) Quantification data. 30 ovaries from 7-day-old flies were quantified for each genotype. Statistical significance was determined using t test: * ($P < 0.05$) and **** ($P < 0.0001$). See also **Source data 2 and 3**.





Figure 2-figure supplement 1.

Oncogenic Ras^{G12V} intrinsically triggers nurse cell death.

(A) Representative ovaries. Scale bars: 200 μ m. (B) Quantification data. 30 ovaries from 3-day-old flies were quantified for each genotype. Statistical significance was determined using t test: **** ($P < 0.0001$). See also **Source data 2 and 3**.






Figure 2-figure supplement 2.

Roles of the DDR pathway and p53 in the nurse cell death induced by oncogenic *Ras^{G12V}*.

(A) Representative ovaries. Scale bars: 200 μm. (B) Quantification data. 30 ovaries from 3-day-old flies were quantified for each genotype. Statistical significance was determined using one-way ANOVA: **** ($P < 0.0001$). (C) Schematic cartoon for *uev1a-flag* knock-in. (D and E) Representative samples. See also **Source data 2 and 3**.

Figure 2-figure supplement 3.

Uev1A does not directly degrade the Ras^{G12V} oncoproteins.

These experiments were performed at 29°C. All images are of the same magnification. See also **Source data 2**.





Figure 2-figure supplement 4.

Uev1A, Ben, and the APC/C complex work together to protect nurse cells from death during normal oogenesis.

These experiments were performed at 29°C. (A) Representative ovaries. The red arrows denote degrading egg chambers. Scale bars: 200 μm. (B) Quantification data. 30 ovaries from 7-day-old flies were quantified for each genotype. Statistical significance was determined using one-way ANOVA: **** ($P < 0.0001$). See also **Source data 2 and 3**.






Figure 3-figure supplement 1.

Co-IP results.

(A) Uev1A interacts with Ben. (B) Uev1A has no interaction with Cdc27. (C) Ben has no interaction with Cdc27. (D) Uev1A has no interaction with Fzr. (E) Uev1A has no interaction with Fzy.




Figure 5-figure supplement 1.

Validation of UBE2V1 and UBE2V2 overexpression in colorectal cancer cell lines.

(A) Western blot analysis confirming the transient overexpression of UBE2V1 and UBE2V2 in SW480 and HCT116 colorectal cancer cell lines. β -Actin was used as the loading control. (B) Western blot analysis confirming the stable overexpression of UBE2V1 and UBE2V2 in SW480 cells, with UBE2V1-OE #1 and UBE2V2 #3 cell lines utilized in subcutaneous tumorigenesis assays. α -Tubulin was used as the loading control. In both (A) and (B), cells transfected with an empty overexpression vector served as the control.

References

- Andersen P.L., Zhou H., Pastushok L., Moraes T., McKenna S., Ziola B., Ellison M.J., Dixit V.M., Xiao W (2005) **Distinct regulation of Ubc13 functions by the two ubiquitin-conjugating enzyme variants Mms2 and Uev1A** *J Cell Biol* **170**:745–755 [Google Scholar](#)
- Bai Z., Wei M., Li Z., Xiao W (2020) **Drosophila Uev1a is dually required for Ben-dependent DNA-damage response and fly mobility** *Cell Signal* **74**:109719 [Google Scholar](#)
- Bakhrat A., Pritchett T., Peretz G., McCall K., Abdu U (2010) **Drosophila Chk2 and p53 proteins induce stage-specific cell death independently during oogenesis** *Apoptosis* **15**:1425–1434 [Google Scholar](#)
- Biggs W.H., Zipursky S.L. (1992) **Primary structure, expression, and signal-dependent tyrosine phosphorylation of a Drosophila homolog of extracellular signal-regulated kinase** *Proc Natl Acad Sci U S A* [Google Scholar](#)
- Broomfield S., Chow B.L., Xiao W (1998) **MMS2, encoding a ubiquitin-conjugating-enzyme-like protein, is a member of the yeast error-free postreplication repair pathway** *Proc Natl Acad Sci U S A* **95**:5678–5683 [Google Scholar](#)
- Canon J., Rex K., Saiki A.Y., Mohr C., Cooke K., Bagal D., Gaida K., Holt T., Knutson C.G., Koppada N., et al. (2019) **The clinical KRAS(G12C) inhibitor AMG 510 drives anti-tumour immunity** *Nature* **575**:217–223 [Google Scholar](#)
- Chen D., McKearin D.M (2003) **A discrete transcriptional silencer in the bam gene determines asymmetric division of the Drosophila germline stem cell** *Development* **130**:1159–1170 [Google Scholar](#)
- DeLuca S.Z., Spradling A.C (2018) **Efficient Expression of Genes in the Drosophila Germline Using a UAS Promoter Free of Interference by Hsp70 piRNAs** *Genetics* **209**:381–387 [Google Scholar](#)
- Di Micco R., Fumagalli M., Cicalese A., Piccinin S., Gasparini P., Luise C., Schurra C., Garre M., Nuciforo P.G., Bensimon A., et al. (2006) **Oncogene-induced senescence is a DNA damage response triggered by DNA hyper-replication** *Nature* **444**:638–642 [Google Scholar](#)
- Dias M.H., Friskes A., Wang S., Fernandes Neto J.M., van Gemert F., Mourragui S., Papagianni C., Kuiken H.J., Mainardi S., Alvarez-Villanueva D., et al. (2024) **Paradoxical Activation of Oncogenic Signaling as a Cancer Treatment Strategy** *Cancer Discov* **14**:1276–1301 [Google Scholar](#)
- Dikic I (2017) **Proteasomal and Autophagic Degradation Systems** *Annu Rev Biochem* **86**:193–224 [Google Scholar](#)
- Edgar B.A., Lehner C.F (1996) **Developmental control of cell cycle regulators: a fly's perspective** *Science* **274**:1646–1652 [Google Scholar](#)
- Gaillard H., García-Muse T., Aguilera A (2015) **Replication stress and cancer** *Nat Rev Cancer* **15**:276–289 [Google Scholar](#)

- Gimple R.C., Wang X (2019) **RAS: Striking at the Core of the Oncogenic Circuitry** *Front Oncol* **9**:965 [Google Scholar](#)
- Glise B., Bourbon H., Noselli S (1995) **hemipterous encodes a novel Drosophila MAP kinase kinase, required for epithelial cell sheet movement** *Cell* **83**:451–461 [Google Scholar](#)
- Greil C., Engelhardt M., Wäsch R (2022) **The Role of the APC/C and Its Coactivators Cdh1 and Cdc20 in Cancer Development and Therapy** *Front Genet* **13**:941565 [Google Scholar](#)
- Hammond M.P., Laird C.D (1985) **Chromosome structure and DNA replication in nurse and follicle cells of Drosophila melanogaster** *Chromosoma* **91**:267–278 [Google Scholar](#)
- Hills S.A., Diffley J.F (2014) **DNA replication and oncogene-induced replicative stress** *Curr Biol* **24**:R435–444 [Google Scholar](#)
- Hofmann R.M., Pickart C.M (1999) **Noncanonical MMS2-encoded ubiquitin-conjugating enzyme functions in assembly of novel polyubiquitin chains for DNA repair** *Cell* **96**:645–653 [Google Scholar](#)
- Huang J., Wu S., Barrera J., Matthews K., Pan D (2005) **The Hippo signaling pathway coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the Drosophila Homolog of YAP** *Cell* **122**:421–434 [Google Scholar](#)
- Igarashi T., Yano K., Endo S., Shiotani B (2024) **Tolerance of Oncogene-Induced Replication Stress: A Fuel for Genomic Instability** *Cancers* **16** [Google Scholar](#)
- Kashevsky H., Wallace J.A., Reed B.H., Lai C., Hayashi-Hagihara A., Orr-Weaver T.L (2002) **The anaphase promoting complex/cyclosome is required during development for modified cell cycles** *Proc Natl Acad Sci U S A* **99**:11217–11222 [Google Scholar](#)
- Kotsantis P., Petermann E., Boulton S.J (2018) **Mechanisms of Oncogene-Induced Replication Stress: Jigsaw Falling into Place** *Cancer Discov* **8**:537–555 [Google Scholar](#)
- Kwon Y.T., Ciechanover A (2017) **The Ubiquitin Code in the Ubiquitin-Proteasome System and Autophagy** *Trends Biochem Sci* **42**:873–886 [Google Scholar](#)
- Lilly M.A., de Cuevas M., Spradling A.C. (2000) **Cyclin A associates with the fusome during germline cyst formation in the Drosophila ovary** *Dev Biol* **218**:53–63 [Google Scholar](#)
- Lilly M.A., Spradling A.C (1996) **The Drosophila endocycle is controlled by Cyclin E and lacks a checkpoint ensuring S-phase completion** *Genes Dev* **10**:2514–2526 [Google Scholar](#)
- Lowe S.W., Cepero E., Evan G (2004) **Intrinsic tumour suppression** *Nature* **432**:307–315 [Google Scholar](#)
- Ma X., Yang L., Yang Y., Li M., Li W., Xue L (2013) **dUev1a modulates TNF-JNK mediated tumor progression and cell death in Drosophila** *Dev Biol* **380**:211–221 [Google Scholar](#)
- Macheret M., Halazonetis T.D (2015) **DNA replication stress as a hallmark of cancer** *Annu Rev Pathol* **10**:425–448 [Google Scholar](#)
- Masrouha N., Yang L., Hijal S., Larochelle S., Suter B (2003) **The Drosophila chk2 gene loki is essential for embryonic DNA double-strand-break checkpoints induced in S phase or G2** *Genetics* **163**:973–982 [Google Scholar](#)

- McKenna S., Spyropoulos L., Moraes T., Pastushok L., Ptak C., Xiao W., Ellison M.J (2001) **Noncovalent interaction between ubiquitin and the human DNA repair protein Mms2 is required for Ubc13-mediated polyubiquitination** *J Biol Chem* **276**:40120–40126 [Google Scholar](#)
- Meyer H.J., Rape M (2014) **Enhanced protein degradation by branched ubiquitin chains** *Cell* **157**:910–921 [Google Scholar](#)
- Moore A.R., Rosenberg S.C., McCormick F., Malek S (2020) **RAS-targeted therapies: is the undruggable drugged?** *Nat Rev Drug Discov* **19**:533–552 [Google Scholar](#)
- Morris L.X., Spradling A.C (2011) **Long-term live imaging provides new insight into stem cell regulation and germline-soma coordination in the Drosophila ovary** *Development* **138**:2207–2215 [Google Scholar](#)
- Muralidhar M.G., Thomas J.B (1993) **The Drosophila bendless gene encodes a neural protein related to ubiquitin-conjugating enzymes** *Neuron* **11**:253–266 [Google Scholar](#)
- Nagy O., Pál M., Udvardy A., Shirras C.A., Boros I., Shirras A.D., Deák P (2012) **lemmingA encodes the Apc11 subunit of the APC/C in Drosophila melanogaster that forms a ternary complex with the E2-C type ubiquitin conjugating enzyme, Vihar and Morula/Apc** *Cell Div* **2** [Google Scholar](#)
- Neuman-Silberberg F.S., Schejter E., Hoffmann F.M., Shilo B.Z (1984) **The Drosophila ras oncogenes: structure and nucleotide sequence** *Cell* **37**:1027–1033 [Google Scholar](#)
- Oh C.E., McMahon R., Benzer S., Tanouye M.A (1994) **bendless, a Drosophila gene affecting neuronal connectivity, encodes a ubiquitin-conjugating enzyme homolog** *J Neurosci* **14**:3166–3179 [Google Scholar](#)
- Oh H., Irvine K.D (2009) **In vivo analysis of Yorkie phosphorylation sites** *Oncogene* **28**:1916–1927 [Google Scholar](#)
- Oikemus S.R., McGinnis N., Queiroz-Machado J., Tukachinsky H., Takada S., Sunkel C.E., Brodsky M.H (2004) **Drosophila atm/telomere fusion is required for telomeric localization of HP1 and telomere position effect** *Genes Dev* **18**:1850–1861 [Google Scholar](#)
- Ou S.I., Jänne P.A., Leal T.A., Rybkin II, Sabari J.K., Barve M.A., Bazhenova L., Johnson M.L., Velastegui K.L., Cilliers C., et al. (2022) **First-in-Human Phase I/IB Dose-Finding Study of Adagrasib (MRTX849) in Patients With Advanced KRAS(G12C) Solid Tumors (KRYSTAL-1)** *J Clin Oncol* **40**:2530–2538 [Google Scholar](#)
- Popow J., Farnaby W., Gollner A., Kofink C., Fischer G., Wurm M., Zollman D., Wijaya A., Mischerikow N., Hasenoehrl C., et al. (2024) **Targeting cancer with small-molecule pan-KRAS degraders** *Science* **385**:1338–1347 [Google Scholar](#)
- Port F., Chen H.M., Lee T., Bullock S.L (2014) **Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila** *Proc Natl Acad Sci U S A* **111**:E2967–2976 [Google Scholar](#)
- Rørth P (1998) **Gal4 in the Drosophila female germline** *Mech Dev* **78**:113–118 [Google Scholar](#)
- Sanchez-Vega F., Mina M., Armenia J., Chatila W.K., Luna A., La K.C., Dimitriadou S., Liu D.L., Kantheti H.S., Saghafein S., et al. (2018) **Oncogenic Signaling Pathways in The Cancer**

Genome Atlas *Cell* :321–337 [Google Scholar](#)

Sancho E., Vilá M.R., Sánchez-Pulido L., Lozano J.J., Paciucci R., Nadal M., Fox M., Harvey C., Bercovich B., Loukili N., et al. (1998) **Role of UEV-1, an inactive variant of the E2 ubiquitin-conjugating enzymes, in in vitro differentiation and cell cycle behavior of HT-29-M6 intestinal mucosecretory cells** *Mol Cell Biol* **18**:576–589 [Google Scholar](#)

Serrano M., Lin A.W., McCurrach M.E., Beach D., Lowe S.W (1997) **Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16INK4a** *Cell* **88**:593–602 [Google Scholar](#)

Sigrist S.J., Lehner C.F (1997) **Drosophila fizzy-related down-regulates mitotic cyclins and is required for cell proliferation arrest and entry into endocycles** *Cell* **90**:671–681 [Google Scholar](#)

Silva E., Tiong S., Pedersen M., Homola E., Royou A., Fasulo B., Siriaco G., Campbell S.D (2004) **ATM is required for telomere maintenance and chromosome stability during Drosophila development** *Curr Biol* **14**:1341–1347 [Google Scholar](#)

Skoulidis F., Li B.T., Dy G.K., Price T.J., Falchook G.S., Wolf J., Italiano A., Schuler M., Borghaei H., Barlesi F., et al. (2021) **Sotorasib for Lung Cancers with KRAS p.G12C Mutation** *N Engl J Med* [Google Scholar](#)

Song Y.H., Mirey G., Betson M., Haber D.A., Settleman J (2004) **The Drosophila ATM ortholog, dATM, mediates the response to ionizing radiation and to spontaneous DNA damage during development** *Curr Biol* **14**:1354–1359 [Google Scholar](#)

Sudakin V., Ganoth D., Dahan A., Heller H., Hershko J., Luca F.C., Ruderman J.V., Hershko A (1995) **The cyclosome, a large complex containing cyclin-selective ubiquitin ligase activity, targets cyclins for destruction at the end of mitosis** *Mol Biol Cell* **6**:185–197 [Google Scholar](#)

Suter B., Steward R (1991) **Requirement for phosphorylation and localization of the Bicaudal-D protein in Drosophila oocyte differentiation** *Cell* **67**:917–926 [Google Scholar](#)

Tanaka-Matakatsu M., Thomas B.J., Du W (2007) **Mutation of the Apc1 homologue shattered disrupts normal eye development by disrupting G1 cell cycle arrest and progression through mitosis** *Dev Biol* **309**:222–235 [Google Scholar](#)

Timmons A.K., Mondragon A.A., Schenkel C.E., Yalonskaya A., Taylor J.D., Moynihan K.E., Etchegaray J.I., Meehan T.L., McCall K (2016) **Phagocytosis genes nonautonomously promote developmental cell death in the Drosophila ovary** *Proc Natl Acad Sci U S A* **113**:E1246–1255 [Google Scholar](#)

Tsuda L., Inoue Y.H., Yoo M.A., Mizuno M., Hata M., Lim Y.M., Adachi-Yamada T., Ryo H., Masamune Y., Nishida Y (1993) **A protein kinase similar to MAP kinase activator acts downstream of the raf kinase in Drosophila** *Cell* **72**:407–414 [Google Scholar](#)

Van Doren M., Williamson A.L., Lehmann R. (1998) **Regulation of zygotic gene expression in Drosophila primordial germ cells** *Curr Biol* **8**:243–246 [Google Scholar](#)

Willingham M.C., Pastan I., Shih T.Y., Scolnick E.M (1980) **Localization of the src gene product of the Harvey strain of MSV to plasma membrane of transformed cells by electron microscopic immunocytochemistry** *Cell* **19**:1005–1014 [Google Scholar](#)

Xu J., Xin S., Du W (2001) **Drosophila Chk2 is required for DNA damage-mediated cell cycle arrest and apoptosis** *FEBS Lett* **508**:394–398 [Google Scholar](#)

Yamano H (2019) **APC/C: current understanding and future perspectives** *F1000Res* [Google Scholar](#)

Zhang Q., Wang Y., Bu Z., Zhang Y., Zhang Q., Li L., Yan L., Wang Y., Zhao S (2024) **Ras promotes germline stem cell division in Drosophila ovaries** *Stem Cell Reports* **19**:1205–1216 [Google Scholar](#)

Zhang Q., Zhang Y., Zhang Q., Li L., Zhao S (2023) **Division promotes adult stem cells to perform active niche competition** *Genetics* **224** [Google Scholar](#)

Zhao S., Fortier T.M., Baehrecke E.H (2018) **Autophagy Promotes Tumor-like Stem Cell Niche Occupancy** *Curr Biol* **28**:3056–3064 [Google Scholar](#)

Zhou R., Silverman N., Hong M., Liao D.S., Chung Y., Chen Z.J., Maniatis T (2005) **The role of ubiquitination in Drosophila innate immunity** *J Biol Chem* **280**:34048–34055 [Google Scholar](#)

Author information

Qi Zhang[†]

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

[†]These authors contributed equally

Yunfeng Wang[†]

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

[†]These authors contributed equally

Xueli Fu[†]

Department of Biochemistry and Molecular Biology, College of Life Sciences, Nankai University, Tianjin, China

[†]These authors contributed equally

Ziguang Wang

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

Yang Zhang

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

Lizhong Yan

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

Yuejia Wang

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

Muhan Yang

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

Dongze Song

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

Ruixing Zhang

Department of Biochemistry and Molecular Biology, College of Life Sciences, Nankai University, Tianjin, China

Hongru Zhang

Department of Biochemistry and Molecular Biology, College of Life Sciences, Nankai University, Tianjin, China, Nankai International Advanced Research Institute (SHENZHEN FUTIAN), Shenzhen, China

For correspondence: hrzhang@nankai.edu.cn

Shian Wu

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

For correspondence: wusa@nankai.edu.cn

Shaowei Zhao

Department of Genetics and Cell Biology, College of Life Sciences, Nankai University, Tianjin, China

ORCID iD: [0000-0002-4544-7215](https://orcid.org/0000-0002-4544-7215)

For correspondence: swzhao@nankai.edu.cn

Editors

Reviewing Editor

Erika Bach

NYU Grossman School of Medicine, New York, United States of America

Senior Editor

Lynne-Marie Postovit

Queens University, Kingston, Canada

Reviewer #1 (Public review):**Summary:**

This study uncovers a protective role of the ubiquitin-conjugating enzyme variant Uev1A in mitigating cell death caused by over-expressed oncogenic Ras in polyploid *Drosophila* nurse cells and by RasK12 in diploid human tumor cell lines. The authors previously showed that overexpression of oncogenic Ras induces death in nurse cells, and now they perform a deficiency screen for modifiers. They identified Uev1A as a suppressor of this Ras-induced cell death. Using genetics and biochemistry, the authors found that Uev1A collaborates with the APC/C E3 ubiquitin ligase complex to promote proteasomal degradation of Cyclin A. This function of Uev1A appears to extend to diploid cells, where its human homologs UBE2V1 and UBE2V2 suppress oncogenic Ras-dependent phenotypes in human colorectal cancer cells in vitro and in xenografts in mice.

Strengths:

- (1) Most of the data is supported by a sufficient sample size and appropriate statistics.
- (2) Good mix of genetics and biochemistry.
- (3) Generation of new transgenes and *Drosophila* alleles that will be beneficial for the community.

Weaknesses:

- (1) Phenotypes are based on artificial overexpression. It is not clear whether these results are relevant to normal physiology.
- (2) The phenotype of "degenerating ovaries" is very broad, and the study is not focused on phenotypes at the cellular level. Furthermore, no information is provided in the Materials and Methods on how degenerating ovaries are scored, despite this being the most important assay in the study.
- (3) In Figure 5, the authors want to conclude that *uev1a* is a tumor-suppressor, and so they over-express *ubev1/2* in human cancer cell lines that have RasK12 and find reduced proliferation, colony formation, and xenograft size. However, genes that act as tumor suppressors have loss-of-function phenotypes that allow for increased cell division. The *Drosophila uev1a* mutant is viable and fertile, suggesting that it is not a tumor suppressor in flies. Additionally, they do not deplete human *ubev1/2* from human cancer cell lines and assess whether this increases cell division, colony formation, and xenograph growth.
- (4) A critical part of the model does not make sense. *CycA* is a key part of their model, but they do not show *CycA* protein expression in WT egg chambers or in their over-expression models (*nos.RasV12* or *bam>RasV12*). Based on Lilly and Spradling 1996, Cyclin A is not expressed in germ cells in region 2-3 of the germarium; whether *CycA* is expressed in nurse cells in later egg chambers is not shown but is critical to document comprehensively.
- (5) The authors should provide more information about the knowledge base of *uev1a* and its homologs in the introduction.

<https://doi.org/10.7554/eLife.107104.1.sa2>

Reviewer #2 (Public review):**Summary:**

The authors performed a genetic screen using deficiency lines and identified Uev1a as a factor that protects nurse cells from RasG12V-induced cell death. According to a previous study from the same lab, this cell death is caused by aberrant mitotic stress due to CycA upregulation (Zhang et al.). This paper further reveals that Uev1a forms a complex with APC/C to promote proteasome-mediated degradation of CycA.

In addition to polyploid nurse cells, the authors also examined the effect of RasG12V-overexpression in diploid germline cells, where RasG12V-overexpression triggers active proliferation, not cell death. Uev1a was found to suppress its overgrowth as well.

Finally, the authors show that the overexpression of the human homologs, UBE2V1 and UBE2V2, suppresses tumor growth in human colorectal cancer xenografts and cell lines. Notably, the expression of these genes correlates with the survival of colorectal cancer patients carrying the Ras mutation.

Strength:

This paper presents a significant finding that UBE2V1/2 may serve as a potential therapy for cancers harboring Ras mutations. The authors propose a fascinating mechanism in which Uev1a forms a complex with APC/C to inhibit aberrant cell cycle progression.

Weakness:

The quantification of some crucial experiments lacks sufficient clarity.

<https://doi.org/10.7554/eLife.107104.1.sa1>

Author response:

Public Reviews:

Reviewer #1 (Public review):

Summary:

This study uncovers a protective role of the ubiquitin-conjugating enzyme variant Uev1A in mitigating cell death caused by over-expressed oncogenic Ras in polyploid Drosophila nurse cells and by RasK12 in diploid human tumor cell lines. The authors previously showed that overexpression of oncogenic Ras induces death in nurse cells, and now they perform a deficiency screen for modifiers. They identified Uev1A as a suppressor of this Ras-induced cell death. Using genetics and biochemistry, the authors found that Uev1A collaborates with the APC/C E3 ubiquitin ligase complex to promote proteasomal degradation of Cyclin A. This function of Uev1A appears to extend to diploid cells, where its human homologs UBE2V1 and UBE2V2 suppress oncogenic Ras-dependent phenotypes in human colorectal cancer cells in vitro and in xenografts in mice.

Strengths:

- (1) Most of the data is supported by a sufficient sample size and appropriate statistics.*
- (2) Good mix of genetics and biochemistry.*
- (3) Generation of new transgenes and Drosophila alleles that will be beneficial for the community.*

We greatly appreciate these comments.

Weaknesses:

(1) Phenotypes are based on artificial overexpression. It is not clear whether these results are relevant to normal physiology.

Downregulation of Uev1A, Ben, and Cdc27 together significantly increased the incidence of dying nurse cells in normal ovaries (Figure 2-figure supplement 4), indicating that the mechanism we uncovered also protects nurse cells from death during normal oogenesis.

(2) The phenotype of "degenerating ovaries" is very broad, and the study is not focused on phenotypes at the cellular level. Furthermore, no information is provided in the Materials and Methods on how degenerating ovaries are scored, despite this being the most important assay in the study.

Thanks for pointing out this issue. We quantified the phenotype of nurse cell death using "degrading/total egg chambers per ovary", not "degenerating ovaries" (see all quantification data in our manuscript). Notably, this phenotype ranges from mild to severe. In normal nurse cells, nuclei exhibit a large, round morphology in DAPI staining (see the first panel in Figure 1D). During early death, nurse cell nuclei become disorganized and begin to condense and fragment (see the third panel in Figure 2-figure supplement 2E). In late-stage death, the nuclei are completely fragmented into small, condensed spherical structures (see the second panel in Figure 1D), making cellular-level phenotypic quantification impossible. Since all nurse cells within the same egg chamber are interconnected, their death process is synchronous. Thus, quantifying the phenotype at the egg-chamber level is more practical than at the cellular level. To improve clarity, we will provide a detailed description of the phenotype and integrate this explanation into the main text of the revised manuscript.

(3) In Figure 5, the authors want to conclude that *uev1a* is a tumor-suppressor, and so they over-express *ubev1/2* in human cancer cell lines that have *RasK12* and find reduced proliferation, colony formation, and xenograft size. However, genes that act as tumor suppressors have loss-of-function phenotypes that allow for increased cell division. The *Drosophila uev1a* mutant is viable and fertile, suggesting that it is not a tumor suppressor in flies. Additionally, they do not deplete human *ubev1/2* from human cancer cell lines and assess whether this increases cell division, colony formation, and xenograph growth.

We apologize for our misleading description. In Figure 5, we aimed to demonstrate that *UBE2V1/2*, like Uev1A in *Drosophila* "*nos>Ras^{G12V}+bam-RNAi*" germline tumors (Figure 4), suppress oncogenic *KRAS*-driven overgrowth in diploid human cancer cells. Importantly, this function of Uev1A and *UBE2V1/2* is dependent on *Ras*-driven tumors; there is no evidence that they act as broad tumor suppressors in the absence of oncogenic *Ras*. *Drosophila uev1a* mutants were lethal, not viable (see Lines 131-133), and germline-specific knockdown of *uev1a* (*nos>uev1a-RNAi*) caused female sterility without inducing tumors. These findings suggest that Uev1A lacks tumor-suppressive activity in the *Drosophila* female germline in the absence of *Ras*-driven tumors. We will revise the manuscript to prevent misinterpretation. Furthermore, we will investigate whether depletion of *UBE2V1*, *UBE2V2*, or both promotes oncogenic *KRAS*-driven overgrowth in human cancer cells.

(4) A critical part of the model does not make sense. *CycA* is a key part of their model, but they do not show *CycA* protein expression in WT egg chambers or in their over-expression models (*nos.RasV12* or *bam>RasV12*). Based on Lilly and Spradling 1996, Cyclin A is not expressed in germ cells in region 2-3 of the germarium; whether *CycA* is expressed in

nurse cells in later egg chambers is not shown but is critical to document comprehensively.

We appreciate this critical comment. CycA is a key cyclin that partners with Cdk1 to promote cell division (Edgar and Lehner, 1996). Notably, nurse cells are post-mitotic endocycling cells (Hammond and Laird, 1985) and typically do not express CycA (Lilly and Spradling, 1996) (see the last sentence, page 2518, paragraph 3). However, their death induced by oncogenic *Ras*^{G12V} is significantly suppressed by monoallelic deletion of either *cycA* or *cdk1* (Zhang et al., 2024). Conversely, ectopic CycA expression in nurse cells triggers their death (Figure 2C, 2D). These findings suggest that polyploid nurse cells exhibit high sensitivity to aberrant division-promoting stress, which may represent a distinct form of cellular stress unique to polyploid cells. To further test our model, we will compare CycA expression levels in normal nurse cells versus those undergoing oncogenic *Ras*^{G12V}-induced cell death.

(5) The authors should provide more information about the knowledge base of uev1a and its homologs in the introduction.

Thanks for this suggestion. We will include this information in the introduction of the revised manuscript.

Reviewer #2 (Public review):

Summary:

*The authors performed a genetic screen using deficiency lines and identified Uev1a as a factor that protects nurse cells from *Ras*^{G12V}-induced cell death. According to a previous study from the same lab, this cell death is caused by aberrant mitotic stress due to CycA upregulation (Zhang et al.). This paper further reveals that Uev1a forms a complex with APC/C to promote proteasome-mediated degradation of CycA.*

*In addition to polyploid nurse cells, the authors also examined the effect of *Ras*^{G12V}-overexpression in diploid germline cells, where *Ras*^{G12V}-overexpression triggers active proliferation, not cell death. Uev1a was found to suppress its overgrowth as well.*

Finally, the authors show that the overexpression of the human homologs, UBE2V1 and UBE2V2, suppresses tumor growth in human colorectal cancer xenografts and cell lines. Notably, the expression of these genes correlates with the survival of colorectal cancer patients carrying the Ras mutation.

Strength:

This paper presents a significant finding that UBE2V1/2 may serve as a potential therapy for cancers harboring Ras mutations. The authors propose a fascinating mechanism in which Uev1a forms a complex with APC/C to inhibit aberrant cell cycle progression.

We greatly appreciate these comments.

Weakness:

The quantification of some crucial experiments lacks sufficient clarity.

Thanks for highlighting this issue. We will provide requested details regarding these quantification data in the revised manuscript.

References

Edgar, B.A., and Lehner, C.F. (1996). Developmental control of cell cycle regulators: a fly's perspective. *Science* 274, 1646-1652.

Hammond, M.P., and Laird, C.D. (1985). Chromosome structure and DNA replication in nurse and follicle cells of *Drosophila melanogaster*. *Chromosoma* 91, 267-278.

Lilly, M.A., and Spradling, A.C. (1996). The *Drosophila* endocycle is controlled by Cyclin E and lacks a checkpoint ensuring S-phase completion. *Genes Dev* 10, 2514-2526.

Zhang, Q., Wang, Y., Bu, Z., Zhang, Y., Zhang, Q., Li, L., Yan, L., Wang, Y., and Zhao, S. (2024). Ras promotes germline stem cell division in *Drosophila* ovaries. *Stem Cell Reports* 19, 1205-1216.

<https://doi.org/10.7554/eLife.107104.1.sa0>