

Spatially defined multicellular functional units in colorectal cancer revealed from single cell and spatial transcriptomics

Reviewed Preprint

v2 • September 26, 2025

Revised by authors

Reviewed Preprint

v1 • March 11, 2025

Inbal Avraham-Davidi , Simon Mages, Johanna Klughammer, Noa Moriel, Shinya Imada, Matan Hofree, Evan Murray, Jonathan Chen, Karin Pelka, Arnav Mehta, Genevieve M Boland, Toni Delorey, Leah Caplan, Danielle Dionne, Robert Strasser, Jana Lalakova, Anezka Niesnerova, Hao Xu, Morgane Rouault, Itay Tirosh, Nir Hacohen, Fei Chen, Omer Yilmaz, Jatin Roper , Orit Rozenblatt-Rosen , Mor Nitzan , Aviv Regev 

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States • Gene Center and Department of Biochemistry, Ludwig-Maximilians-Universität München, Munich, Germany • School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel • Department of Biology at MIT, Koch Institute for Integrative Cancer Research at MIT, Cambridge, United States • Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States • Massachusetts General Hospital (MGH) Cancer Center, Harvard Medical School (HMS), Boston, United States • Department of Pathology, Massachusetts General Hospital, Boston, United States • Gladstone-UCSF Institute of Genomic Immunology, Gladstone Institutes, San Francisco, United States • Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, United States • Dana Farber Cancer Institute, Boston, United States • Department of Surgery, Massachusetts General Hospital, Boston, United States • 10xGenomics, Solnavägen 3H, Stockholm, Sweden • Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel • Harvard Stem Cell and Regenerative Biology, Cambridge, United States • Department of Pharmacology and Cancer Biology and Department of Medicine, Division of Gastroenterology, Duke University, Durham, United States • Genentech, South San Francisco, United States • Racah Institute of Physics and Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel • Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This work presents a **valuable** resource combining scRNA-seq and spatial transcriptomics studies to map mouse pre-clinical models of colorectal cancer, identifying distinct cellular programs and microenvironments that could enhance patient stratification and therapeutic approaches in colorectal cancer. While the evidence provided in the manuscript are not fully validated, these **solid** data were collected and analyzed using a validated methodology that will be of interest to the community in future studies.

<https://doi.org/10.7554/eLife.104815.2.sa2>

Abstract

While advances in single cell genomics have helped to chart the cellular components of tumor ecosystems, it has been more challenging to characterize their specific spatial organization and functional interactions. Here, we combine single cell RNA-seq, spatial transcriptomics by Slide-seq, and *in situ* multiplex RNA analysis, to create a detailed spatial map of healthy and dysplastic colon cellular ecosystems and their association with disease progression. We profiled inducible genetic CRC mouse models that recapitulate key features of human CRC, assigned cell types and epithelial expression programs to spatial tissue locations in tumors, and computationally used them to identify the regional features spanning different cells in the same spatial niche. We find that tumors were organized in cellular neighborhoods, each with a distinct composition of cell subtypes, expression programs, and local cellular interactions. Comparing to scRNA-seq and Slide-seq data from human CRC, we find that both cell composition and layout features were conserved between the species, with mouse neighborhoods correlating with malignancy and clinical outcome in human patient tumors, highlighting the relevance of our findings to human disease. Our work offers a comprehensive framework that is applicable across various tissues, tumors, and disease conditions, with tools for the extrapolation of findings from experimental mouse models to human diseases.

Introduction

The spatial organization of diverse cells in the tumor ecosystem impacts and drives interactions between malignant cells and neighboring immune and stromal cells, either promoting or suppressing tumor growth¹. Recent studies have shown that systematic understanding of the spatial organization of tumors can shed light on disease progression and response to therapy, with specific features correlated with tumor subtypes^{2–4}, cancer prognosis^{5–7}, or response to treatment^{8,9}.

However, genome-scale, high-resolution dissection of the spatial organization of tumors and its functional implications remains challenging, largely due to technical limitations. Methods such as fluorescent *in situ* hybridization (FISH) and immunohistochemistry can only measure a handful of pre-selected transcripts or proteins, whereas single cell RNA-seq (scRNA-seq) does not directly capture spatial relations. Recent advances in spatial genomics and proteomics allow multiplexed or genome-scale measurements *in situ*^{10–17}, but with a trade-off between genomic scale and spatial resolution¹⁸. As a result, data from different experimental methods need to be integrated for a comprehensive view of the tissue biology. Many analytical tools have been developed to integrate some crucial aspects of the data^{19–23}, but it can be challenging to deploy them and distill answers to specific disease biology questions. This leaves open many fundamental questions about tissue organization and collective function, including whether there are canonical functional units in tumors, what may be their organization in the tumor landscape, and what role does each play in tumor progression.

A case in point is colorectal cancer (CRC), where initial lesions (adenomatous polyps) progress over time to carcinoma and eventually to metastatic disease. While the mutations that drive this process were extensively studied^{24–27}, and the cellular ecosystem of CRC has now been deeply charted^{28,29}, the spatial landscape is less well-characterized. In a recent study of human CRC², we statistically associated cell profiles across tumors and showed that they map to different cellular communities, reside in different locations in the tumor and reflect different tumor subtypes. However, absent genome wide *in situ* measurements, these statistical inferences

do not yet reflect the full spatial organization of the tumor. Moreover, it is important to relate such patterns to those in animal models used in mechanistic studies and as pre-clinical models, to understand the relation between lab models and human tumors.

Here, we deciphered the spatial and cellular organization of colorectal cancer (CRC) by combining scRNA-seq, spatial transcriptomics by Slide-seq, and *in situ* RNA multiplex analysis, using a novel computational framework. We first profiled two inducible genetic mouse models of colorectal cancer that recapitulate key features of human CRC, before and after tumor initiation. We integrated the spatial and cell profiles to create a spatial cell map of the tumor landscape, revealing dysplasia-specific cellular layout and potential physical interactions. We found that the tumor landscape is organized in cell neighborhoods, each with distinct epithelial, immune, and stromal cell compositions, and governed by different gene programs. Three of the cell neighborhoods are associated with tumor progression, each activating different biological pathways but all active simultaneously albeit in different parts of the tumor. We devised a computational framework, based on the TACCO²¹ method, extending it to compare single cell and spatial features of tumors between species and applied it to scRNA-seq data from human CRC. Multiple features were conserved between tumors in the mouse model and the human patients, and the mouse cellular neighborhoods correlated with malignancy and clinical outcome (progression-free intervals (PFI) and overall survival (OS)) in human patient tumors. Our work provides a general approach that can be applied to other tissues, tumors and disease conditions.

Results

A cell atlas of genetic models of colorectal cancer

To chart the cell ecosystem of CRC and how it changes during tumor progression, we studied two genetic mouse models of CRC, one with inactivation of *Apc* and another in which *Apc* inactivation is accompanied by an oncogenic *Kras*^{G12D/+} mutation and inactivation of *Trp53*³⁰⁻³² (Figure 1A). In the AV model, *Apc*^{fl/fl}*Villin*^{CreERT2} mice are injected with 4-hydroxytamoxifen to the submucosal layer of the colon, inducing the deletion of *Apc* specifically in epithelial cells within the injection site³⁰. In the AKPV model (*Apc*^{fl/fl}; *LSL-Kras*^{G12D}; *Trp53*^{fl/fl}; *Rosa*^{26LSL-tdTomato/+}; *Villin*^{CreERT2} mice, **Methods**), 4-hydroxytamoxifen injection also induces an oncogenic *Kras*^{G12D/+} mutation and then inactivation of *Trp53*. In both cases, 4-hydroxytamoxifen injection leads to the formation of local lesions that resemble human dysplastic lesions³⁰.

We first generated a single-cell atlas of the models consisting of 48,115 high quality scRNA-seq profiles from normal colon, AV (3 weeks after 4-hydroxytamoxifen induction) and AKPV (3 and 9 weeks after induction) tissues. We captured a diverse cell census (Figure 1B,C, **Methods**), with 35 clusters annotated *post hoc* by the expression of known marker genes (Figure 1B, Figure S1A,B, **Methods**), across epithelial, immune and stromal cell compartments (Figure S1C). For validation purposes, we also generated multiplex *in situ* RNA profiles (with the Cartana³³ method) in 6 sections each from normal and AV conditions, using a panel of 66-180 marker genes chosen to best represent the cell types and programs found in the tissues (Figure S1D, **Methods**).

Tumorigenic genotypes cause shifts in the composition of epithelial cell populations and their microenvironment in AV and AKPV lesions

Dysplastic lesions exhibited shifts in proportions of immune and stromal cells, including changes in subsets pre-existing in normal tissue, as well as infiltration of new cell subsets (Figure 1C,D and Figure S1D,E,F). This resulted in both increase in cells of existing populations (e.g., some T cell subsets (TNK05 (GdT/Il17+), TNK06 (Treg)) and emergence of new dysplasia-associated cells




Figure 1.

A single cell atlas of healthy colon and dysplastic lesions in mouse.

A. Study overview. **B.** Major cell subsets of healthy colon and dysplastic lesions. 2D embedding of 48,115 single cell profiles colored by cluster (top, legend) or annotated cell type (bottom, legend). **C,D.** Changes in cell composition in dysplastic tissues. **C.** 2D embedding of single cell profiles, showing only the cells in each condition state, subsampled to equal numbers of cells per condition state, colored by cluster (same legend as in B (top)). **D.** Proportion of cells (y axis) of each cell type in each sample (x axis). **E.** multiplex RNA *in situ* analysis. Representative images of Cartana analysis of normal colon (left) and AV lesions (right) colored by cell type assignment (same as **Supplementary item 2A**).

(e.g., granulocytes (Gran01, Gran02) and monocytes (Mono02, Mono03)) mirroring observations in human CRC², breast cancer³⁴, and non-small cell lung cancer³⁵ (**Figure 1C,D** and **Figure S1D,E** and **Figure S2 A-C**). We validated these patterns using multiplex *in situ* RNA analysis (**Figure 1E**, **Figure S1F-H** and **Supplementary item 1** and **2**). Infiltration is likely to underlie many of these changes as many of the increasing cell subsets (granulocytes, monocytes, mast cells) expressed genes, such as *Sell* and *Ccr2*, indicating tissue recruitment, and as the cells dramatically increase in proportion despite negligible signals of proliferation programs.

Two of four monocyte subsets, Mono02 and Mono03, were unique dysplasia-associated cells (**Figure S2D-F**) and were respectively enriched for general inflammatory response genes (FDR=5.7 10⁻³⁰, two-sided Fisher's exact test in GO term enrichment) and interferon beta and gamma response genes (FDR=3.5 10⁻¹¹, 1.0 10⁻¹³). T cell subsets showed the expected diversity across nine subsets (**Figure S2G-I**), with a significant decrease (out of all T cells) in TNK01 (GdT/Cd8) in the dysplastic microenvironment and an increase in TNK05 (GdT/Il17+) (**Figure S2J**). This is consistent with the T cell composition in tumors from mismatch repair proficient (MMRp) CRC patients (**Figure S2K**). RNA velocity analysis³⁷ of T cells from normal and AV tissue (**Figure S3A**) showed a change in inferred cellular relationships with TNK03 (naïve T) and TNK02 (Th1/Th17) preceding TNK06 (Treg), consistent with the promotion of an immunosuppressive microenvironment, and TNK08 (proliferating T) also preceding TNK02 (Th1/Th17), TNK04 (GdT/Cd8) and TNK05 (GdT/Il17+) populations.

Within the five subsets of stromal cells (including vascular endothelial and lymphatic endothelial cells and three fibroblast subsets, **Figure S2L-N** and **Methods**), Endo01 (vascular) were enriched in dysplastic lesions compared to normal colon (FDR=1.8 10⁻³, two-sided Welch's t test on CLR transformed compositions; **Figure S2O**). Angiogenesis-related pathways, such as angiogenesis (FDR=1.0 10⁻¹¹) and positive regulation of angiogenesis (FDR=3.4 10⁻⁴) as well glycolytic process (FDR=1.4 10⁻⁴) were enriched in Endo01 (vascular) from lesions compared to normal colon (Table S1). This is in line with vascular adaptation to the tumor's growing needs for nutrients and oxygen³⁸ and with the increased expression of the vascular growth factor *Vegf-A* in both monocytes and macrophages (**Figure S2P**).

Cell-intrinsic expression shifts in different sub-lineages in the dysplastic epithelium

Epithelial cells showed dramatic cell-intrinsic changes between normal tissues and either AV or AKPV lesions, such that the cell profiles of dysplastic epithelial cells in both models were highly distinct from normal epithelial cells (and similar to each other) (**Figure 1C**, **Figure S4A** and **Methods**). Epithelial cell profiles from normal mice (41% of cells) separated from most of those from AV and AKPV models (59% of cells) (**Figure 2A** and **Figure S4A,B**), suggesting a common shift in all dysplastic cells from the normal state. Notably, 11% of epithelial cells from AV/AKPV mice were classified as non-dysplastic healthy cells, indicating that normal, non-dysplastic, cells may be present in or adjacent to the lesion microenvironment (**Figure 2B** and **Figure S4A**), although we cannot firmly rule out adjacent tissue contamination. We annotated two cell clusters – Epi01 (dysplastic stem-like) and Epi05 (dysplastic secretory-like) – as dysplastic, due to their virtually exclusive presence in AV and AKPV models and because they expressed high levels of *Apc* target genes (e.g., *Axin2*, *Ascl2*, *Myc*, *Ccnd1*, *Lgr5*) and were enriched in tdTomato⁺ cells from AKPVT mice (**Figure 2C** and **Figure S4C,D**) while also spanning the enterocyte to secretory continuum with healthy cells (**Figure S4B**, PC2).

Interestingly, Epi05 (dysplastic secretory-like) had distinguishing markers (e.g., *Ccl9*, *Mmp7*, *Ifitm3*) from their counterparts in normal tissue, Epi04 (secretory) (**Figure 2A-C** and **Figure S4A,C,E**). *Mmp7* and *Ifitm3* are known to promote metastasis in human CRC^{39,40}, and *Ccl9* expression by epithelial cells promotes tumor invasion through recruitment of Ccr1⁺ myeloid cells to the tumor's invasive front in a mouse model of CRC⁴¹. Notably, *Ccr1* is expressed by newly recruited




Figure 2.

Composition and cell intrinsic expression program changes in dysplastic epithelial cells.

A-C. Compositional changes in epithelial cells in dysplastic tissue. **A.** 2D embedding of epithelial cell profiles colored by clusters (legend). Cluster Epi06: potential doublets (**Methods**). **B.** Proportion of cells out of all epithelial cells (y axis) of each epithelial cell subset in each sample (x axis). **C.** Fraction of expressing cells (dot size) and mean expression in expressing cells (dot color) of marker genes (columns) for each cluster (rows). **D-E.** Use of epithelial cell programs changes in dysplastic tissue. **D.** Weights (x axis) of each of the 20 top ranked genes (y axis) for each program. **E.** Proportion of program weights summed over all epithelial cells (y axis) in each sample (x axis). **F.** Stem cell program 16 is induced in epithelial cells in dysplastic tissue. Scaled log-normalized expression (color bar) of the top 100 genes differentially expressed between cells from normal colon and from dysplastic (AV and AKPV) across the 10,812 cells that accounted for 90% of program 16's expression across all epithelial cells (columns). Selected program genes are marked.

monocytes, macrophages and granulocytes in our model, suggesting a potential mechanism for tumor infiltration and invasion (**Figure S4E**). Thus, dysplastic secretory epithelial cells may perform additional functions that differ from those of their healthy counterparts.

RNA velocity³⁷ analysis of the epithelial cell compartment predicted that in the dysplastic epithelium (**Figure S3B, right**) a proliferating stem-cell like dysplastic subset is a direct source to both a massively expanded and heterogeneous non-proliferating population of dysplastic stem-like cells (expressing WNT signaling and angiogenesis programs) and to dysplastic cells of different ‘differentiation states’ (MHCII expressing stem/progenitors leading to enterocytes and secretory-like cells) (**Figure S3B,C**). Conversely, in the normal epithelium (**Figure S3B, left**), the ‘root’ is placed in a much smaller population of proliferating intestinal stem cells, and the stem cell compartment is overall much more modest. The proliferation and differentiation path of previously identified normal epithelial cells remains intact (*i.e.*, reminiscent of the one in normal samples) even in the dysplastic lesions (**Figure S3B**). While our RNA velocity analysis provides insights into potential cellular trajectories, further experimental validation is required to confirm these findings.

Expression programs for stem-like functions, Wnt signaling, angiogenesis and inflammation are activated in dysplastic epithelial cells

Both normal and dysplastic epithelial cells varied along a continuum, as expected and previously observed in the ongoing differentiation in the colon epithelium^{2,42–44} and our RNA Velocity analysis (**Figure S3**). Using non-negative matrix factorization (iNMF from LIGER⁴⁵, **Methods**), we recovered 20 expression programs spanning the different epithelial functions, and annotated them by Gene Ontology terms enriched in their top 100 weighted genes (**Figure 2D,E**, **Figure S3C**, **S4F-L**, **Methods**).

The programs enriched in different dysplastic cells highlighted key processes that play a role in tumor promotion, including stem cell programs, Wnt signaling, angiogenesis, and inflammation and innate immunity, including interferon alpha, beta and gamma pathways (**Figure 2D,E**, **Figure S3C**). In particular, the stem cell program (#16) detected in some cells across all conditions, was enriched in dysplastic samples ($\text{FDR}=5.6 \times 10^{-10}$, two-sided Welch’s t-test on CLR transformed compositions), reminiscent of a recently described population in human^{28, 29}. Comparing cells from dysplastic and normal samples that express the stem cell program, the dysplastic cells had distinct expression profiles with induction of negative regulators of Wnt signaling ($\text{FDR}=4.5 \times 10^{-5}$, two-sided Fisher’s exact test in GO term enrichment, *e.g.*, *Notum*, Wnt inhibitory factor 1 (*Wif1*) and *Nkd1*) and genes that are related to cellular response to interferon-gamma ($\text{FDR}=1.7 \times 10^{-6}$, *e.g.*, *Ccl9*, *Ccl6*) and immune system process ($\text{FDR}=6.4 \times 10^{-4}$, *e.g.*, *Ifitm1* and *Ifitm3*) (**Figure 2E,F**). This is consistent with recent studies showing that *Apc*-mutant stem cells secrete negative regulators of Wnt signaling to induce the differentiation of the WT stem cells in their proximity, thereby outcompeting them and promoting tumor formation^{46, 47}. Thus, stem cells from dysplastic lesions may have non-canonical function and regulation. In addition, the programs for Wnt signaling (expressing both positive and negative regulators; #4, $\text{FDR}=2.8 \times 10^{-6}$), angiogenesis (#14, $\text{FDR}=1.2 \times 10^{-9}$), inflammatory response (#6, $\text{FDR}=1.4 \times 10^{-6}$), and innate immune response and interferon response (#7, $\text{FDR}=1.2 \times 10^{-2}$) were all predominantly expressed or enriched in AV/AKPV epithelium (all with two-sided Welch’s t-test on CLR transformed compositions, **Figure 2E** and **Figure S4I-M**). These results are consistent with the known role of the Wnt signaling pathway in CRC, and of angiogenesis, response to hypoxia and inflammation in tumor progression^{48–50}.

Malignant-like tissue programs and composition are conserved between mouse and human tumors

To evaluate the relevance of our findings to human colorectal cancer, we compared them to a scRNA-seq atlas we recently generated from tumor and adjacent normal tissue from 62 patients with either MMRp or MMRd CRC². We compared mouse and human tumors in terms of their epithelial expression programs, cellular composition, and cell associations in multicellular hubs². While our mouse model should more closely resemble MMRp tumors, we compared to both classes separately and together to identify any shared features.

To assess the similarity between mouse and human programs we controlled for overall cross-species and batch differences by normalizing program-specific expression profiles with species-specific background profiles (**Methods**). The Pearson correlation coefficients of these normalized scores between the human and mouse programs indicate some overlap in the programs defined on mouse and human data (**Figure S5A**). Epithelial cells from human and mouse tumors expressed many programs highly correlated between the species (**Figure S5B**, **Methods**), including for cell cycle, inflammation, epithelial secretory, angiogenesis, Wnt signaling, and normal colon functions.

Co-variation in cell proportions across samples (by scRNA-seq) was also conserved between human and mouse tumors, suggesting broad conservation of tumor composition. For example, in both species the proportion of endothelial cells and fibroblasts correlated across samples, as did T and B cell proportions in human tumors and mouse dysplastic lesions (**Figure S5C**). Moreover, when we transferred epithelial program annotations from mouse to human scRNA-seq and calculated their co-variation across samples in each species, programs 11 (proliferation), 14 (angiogenesis) and 16 (stem cells), co-varied both across dysplastic mouse tumors and across human MMRp and MMRd tumors (**Figure S5D** and **Methods**), suggesting a conserved dysplastic tissue architecture.

Integrated spatial and single-cell atlas of mouse CRC tumors

To comprehensively decipher the distribution of cells and programs in the tumor spatial niche, we next used Slide-seqV2¹⁵ for genome-wide spatial RNA-seq at 10 µm resolution. We sectioned and profiled frozen tissues from four normal colon and four AV lesions using 10 Slide-seqV2 pucks (**Methods**), recovering 221,936 high quality beads (**Figure S6A-C**, **Methods**). We then integrated the single cell census and spatial profiles using TACCO, which allowed us to annotate each bead with compositions of discrete cell types (from epithelial, immune and stromal compartments) and to further annotate the epithelial fraction of each bead with a composition of epithelial program activity (**Figure 1A** “annotation”).

We first used TACCO to annotate every bead in the Slide-seq data with a composition of discrete cell subtypes for every puck separately, using its matching single-cell reference (normal or disease; **Figure 1A**, **Methods**). To this end, TACCO iteratively solved optimal transport problems to assign cell subtypes to fractions of reads of the beads. TACCO relies on unbalanced optimal transport to allow for shifts in the frequency of cell subtypes in the pucks vs. the single-cell dataset, while using the reference cellular frequencies as prior knowledge (**Figure S6E,G**). TACCO’s cell type mapping recapitulated the muscularis layer in its expected tissue location based on the inferred cellular composition pattern (**Figure 3A**, **4A** and **S6D**). This illustrates that TACCO’s compositional annotations align well with biological patterns.

Next, we used TACCO to map the epithelial gene programs (defined above), focusing on transcript counts that are inferred as derived from epithelial cells. TACCO partitioned the read count matrices for each puck, assigning counts to epithelial cells based on the mapped per-bead cell subtype annotations (from the first step) and the expression profiles associated with each subtype




Figure 3.

Altered cell type neighborhood in CRC.

A. Cell type distributions *in situ*. Slide-seq pucks of dysplastic (top) and normal (bottom) tissue colored by TACCO assignment of cell labels (legend, light grey: low quality beads) (x and y axis: spatial coordinates in μm). **B,C.** Cell type neighborhoods in normal and dysplastic colon tissue. Short-range (up to $20\mu\text{m}$) neighborhood enrichment (Z score, color bar) vs. a background of spatially random annotation assignments for each pair of cell annotations (rows, columns) in normal (A) and dysplastic (B) tissue.

(**Methods**). It then summed all epithelial contributions into an epithelial-only spatial count matrix, followed by optimal transport to assign epithelial program contributions to individual beads, based on epithelial cell-only read signals. As for cell type mapping, the proportional contribution of the programs largely recapitulated their contributions in scRNA-seq (**Figure S6F,H** [↗](#)).

Altered and less ordered local cellular organization of dysplastic lesions

We assessed the local cellular architecture in terms of the preferential proximity of cells of certain type or expressing particular epithelial programs, within a fixed-sized neighborhood, by adapting an earlier method. We defined a z-score as significance of the observed neighborhood relations compared to the null for neighborhoods of 20, 40 or 60 μm diameter (**Figure 3B,C** [↗](#) and **Figure S7A,B** [↗](#)). This z-score is defined with respect to a population of random cell type annotations generated by random permutations of the cell type annotations between the beads, where in our case we permute fractional cell type contributions.

Cell proximity preferences in the normal colon tissue are consistent with the expected morphology, validating our approach (**Figure 3A,B** [↗](#)). Epithelial cells were organized such that the differentiated Epi02 (Enterocytes) are excluded from the stem cell niche (**Figure 3B** [↗](#) and **Figure S7C** [↗](#)), and endothelial cells and fibroblasts were also spatially co-located in a focused region (**Figure 3B** [↗](#)), with T cells in their vicinity (**Figure 3B** [↗](#)). Our multiplex *in situ* RNA analysis validated the exclusion of enterocytes from the stem cell niche, as also seen in Slide-seq data (**Figure S7C,D** [↗](#)).

While some normal tissue features are preserved in dysplastic samples, including co-location of cells of the same lineage⁵ [↗](#), ¹⁰ [↗](#) (**Figure 3A-C** [↗](#)), there were notable changes, and more disorder. Cell types were more randomly distributed in AV lesion vs. normal tissue, reflected in lower z-scores ($p=1.6 \cdot 10^{-37}$, one-sided Mann-Whitney U test; **Figure S7E** [↗](#)). At short distances, all epithelial cells (normal and dysplastic) were preferentially located close to cells from the same subtype (**Figure 3C** [↗](#)) and even to cells with similar functions: epithelial cells expressing programs associated with malignant-like function (e.g., program 4 (Wnt signaling), 14 (angiogenesis) and 16 (stem cells)) resided close to each other and were spatially distant from cells expressing programs that are related to normal epithelial functions (e.g., program 5 (basolateral plasma membrane), 8 (apical plasma membrane) and 10 (oxidation-reduction process)), supporting a model where tumor progression is structured and compartmentalized (**Figure S7F** [↗](#)). Immune and stromal cells were generally excluded from epithelial cell neighborhoods. Granulocytes aggregated together (self-proximal) (**Figure 3C** [↗](#)) and were relatively close to endothelial cells and dysplasia-associated monocytes (Mono02, Mono03), consistent with their recruitment from the blood through the vessels (**Figure 3C** [↗](#) and **Figure S7G** [↗](#)). Our multiplex *in situ* RNA analysis validated the spatial enrichment of monocytes and granulocytes near vessels (**Figure S7H** [↗](#)).

Epithelial regional analysis recovers canonical structures in normal colon

To detect distinctive tissue regions in tumors, which lack traditional tissue references, we identified cellular neighborhoods with both similar epithelial program activity and a particular composition of immune and stromal cells. Specifically, we first identified “epithelial program regions” as areas of distinct epithelial program activity, and then found immune or stromal cells associated with each region (**Figure 1A** [↗](#) “annotation”). Intuitively, we defined “regions” based on both the similarity in epithelial expression program activity and proximity in space. To do this, after assigning epithelial programs to epithelial beads, we clustered the beads based on a weighted sum of spatial proximity and expression program similarity. This results in spatially contiguous annotation of beads with distinct epithelial program activity, which, together with the immune and stromal cells in their proximity, compose the “region”. Specifically, using TACCO, we defined

epithelial program regions by Leiden clustering of the weighted sum of neighborhood graphs for spatial bead proximity and epithelial expression program similarity, such that transcriptionally similar epithelial beads on different pucks can be connected (despite “infinite” spatial distance, **Methods**). We then used this single framework for region annotation across all pucks (**Figure 4A**), to determine the distinctive composition of additional cell types in the same set of spatial regions (**Figure 4B-D** and **Figure S8A**).

In the normal colon, the regional analysis (**Figure 4A**, bottom) robustly recovered the expected spatial organization of the healthy colon across five regions and their cellular composition and sublayers (**Figure 1A**), from luminal/apical to basal. Four regions recovered by TACCO corresponded to different layers of the mucosa (**Figure 4A** and **4E,F**): a luminal layer with reads found beyond the cellular layer and likely representing cellular debris trapped in the mucus; three apical layers expressing programs related to normal epithelial function (transmembrane transport, oxidation-reduction process) with gradual transition from apical to basal features; and a basal-most layer, enriched for the deep crypt, proliferation (G1/S,G2/M), MHCII and basolateral plasma membrane programs, all common features of the deep crypt area. Finally, region 2, enriched with fibroblasts, myofibroblasts and endothelial cells, and located in the most basal side of the tissue, captured the submucosal and muscularis propria layers, which are predominantly comprised of fibroblasts and muscle, respectively, alongside blood and lymphatic vessels, nerves and immune cells. Overall, TACCO recovered the known organization of the colon, showing the power of our unsupervised mapping approach and shedding light on expression programs that are required for the maintenance of normal colon homeostasis.

Dysplastic lesions maintain some of the programs of the corresponding regions in healthy tissue

AV lesions did not maintain the robust organization of normal tissues, and reflected the expected histopathology of high grade dysplasia, when dysplastic cells are confined to the mucosal layer and do not invade the submucosa⁵¹ (**Figure 1A** and **4A**, top). Specifically, the submucosal and muscularis propria layers from both normal and AV lesions were assigned to region 2 (**Figure 4A**).

Despite the altered morphology, some of the disrupted regions also expressed programs characteristic of their normal healthy function, suggesting that tumor progression is spatially structured and compartmentalized. For example, the region above the submucosa, captured as region 1 in AV lesions and region 5 in normal colon (**Figure 4A**), had similar features in both AV lesions and normal samples. Thus, although the overall spatial organization was disrupted in the lesion, region 1 in AV lesions expressed programs that are reminiscent of the normal deep crypt region 5, and was enriched for deep crypt cells and programs that are related to proliferation and MHC II⁴³ (**Figure 4A,C**). These included proliferation programs 3 and 11 and both normal stem cells (Epi03) and dysplastic secretory-like cells (Epi05), as well as dysplastic stem cells (Epi01, though to a lesser extent than some other regions), so it may reflect one of the proliferative stem cell (and dysplastic secretory-like) niches in AV models (**Figure 4B,C**). Other regions in the AV lesions also contained some epithelial cells with normal profiles, expressing programs that should allow them to maintain their capacity to perform normal tasks. For example, region 3 expressed apical plasma membrane functions and region 10 was enriched with oxidation-reduction functions (**Figure 4C**).

To learn about the spatial distribution of the dysplastic regions, we measured their distance from region 2 (muscularis), which is a stable landmark in the lesions. Remnants of the layered structure of the healthy tissue were still observed in the AV tissue, especially at relatively low distances from the muscularis. For example, healthy region 5 – characteristically located at distances of about 150-200µm from the muscularis – is replaced by dysplastic region 1, peaking at 200µm. All malignant-like regions (6/8/11) were spatially associated at ~300-700µm from the muscularis (**Figure S8C**),




Figure 4.

Three cellular neighborhoods associated with tumor progression.

A. Spatial regions. Slide-seq pucks of AV (top) and normal (bottom) mouse colon colored by TACCO regions (legend, light gray: low quality beads) (x and y axis are spatial coordinates in μm). **B, C.** Enrichment and depletion of cell subsets and epithelial programs across different regions. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of specific cell subsets (rows, B) or epithelial cell programs (rows, C) in the different regions defined by TACCO (columns) as well as all normal ("N (ref.)", leftmost column) and AV ("AV (ref.)", leftmost column) samples. **D.** TACCO defined regions preferentially relate to normal or AV tissue. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each TACCO defined region (rows) in normal ("N vs. rest") and AV ("AV vs. N") samples (columns). **E.** TACCO reveals normal colon architecture. Left: Slide-seq puck of normal mouse colon colored by TACCO region annotations (legend) (x and y axis: spatial coordinates (μm)). Right: Main epithelial expression programs enriched in each region (FDR < 6.3×10^{-4} , two-sided Welch's t test on CLR-transformed compositions) except region 2 (muscularis), which is characterized by non-epithelial (stromal) cell types. **F.** Expression signatures of cells in normal regions 3, 5, 10 and 12. Scaled log-normalized expression of the top 20 differentially expressed genes (rows) for each bead (columns) in the region. **G, H.** Malignant-like regions. G. Slide-seq pucks of two AV lesions colored by TACCO annotations of malignant-like regions 6, 8 and 11. H. Scaled log-normalized expression of the top 20 differentially expressed genes (rows) of each bead (left, columns) in the region; or epithelial (middle left), immune (middle right) or stromal (right) fractions of beads (columns) in regions 6, 8 and 11 in dysplastic lesions. **I, J.** Epithelial cell subsets and programs associated with "malignant-like", "normal-like" and normal tissues. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of epithelial cell programs (I, rows) or epithelial, immune and stromal cell subsets (J, rows) in different tissue types (columns) based on Slide-seq or scRNA-seq ("sc") samples. **K.** Inferred interaction pathways. Enrichment (FDR) in AV vs. normal tissue of corresponding "sender" (x axis) and "receiver" (y axis) (aggregated over ligand-receptor pairs) pathways (dots). Red/blue: pathways significantly enriched in AV/normal samples and (light red: only for either sender or receiver). The top 5 enriched pathways (in each direction) are labeled.

located ~100-400µm apart from each other (**Figure S8D** [↗](#)). We further validated this result at the protein level, by staining for b-catenin, showing an (inactive) cytoplasmic localization in the region adjacent to the muscularis, and mostly nuclear (active) localization in distal regions, near the lumen (**Figure S8B** [↗](#)).

Three spatially and functionally distinct tumor regions enriched in AV lesions

Three regions – 6, 8, and 11 – had epithelial composition and programs that suggested advanced malignant-like characteristics, each highlighting a potentially different mechanism for tumor progression (**Figure 4G,H** [↗](#)). These three ‘malignant-like regions’, were enriched (vs. all other regions) in stem cell, Wnt signaling and angiogenesis programs (16, #4 and #14; FDR=9.6 10⁻²¹, 8.1 10⁻¹⁰, 3.7 10⁻¹⁰, two-sided Welch’s t-test on CLR transformed compositions) and depleted of normal epithelial programs (#5, #8, and #10; FDR=5.5 10⁻¹², 1.6 10⁻¹², 4.2 10⁻¹⁰, **Figure 4I** [↗](#)). Furthermore, the malignant-like regions were enriched in immune cells, including monocytes-macrophages (FDR≤1.5 10⁻⁵; excluding Mac02 (Lyve1+)), T cell subsets TNK02 (Th1/Th17), TNK05 (GdT/IL17+), TNK06 (Treg), TNK08 (proliferating T) (FDR=2.4 10⁻³, 2.2 10⁻⁴, 1.7 10⁻³, 9.6 10⁻⁴; two-sided Welch’s t-test on CLR transformed compositions), infiltrating granulocytes (FDR≤9.7 10⁻³), and mast cells (FDR=1.4 10⁻²), suggesting an ongoing immune response (**Figure 4J** [↗](#)). However, each one of the three regions had a different epithelial program composition, suggesting that in each type of region there is a different dominant pathway/feature that may drive tumor progression (**Figure 4C** [↗](#), Table S2).

Region 6 was characterized by an inflammatory and angiogenic multicellular community, with epithelial and immune cells expressing inflammatory programs, endothelial cells and monocytes connected in a pro-angiogenic circuit, and pro-invasive genes expressed by both endothelial and immune cells (**Figure 4B,C** [↗](#)). Specifically, region 6 was distinctly enriched for proliferation (programs 3 and 11; FDR=2.2 10⁻¹⁴, 1.2 10⁻¹⁵, two-sided Welch’s t-test on CLR transformed compositions) and inflammatory epithelial programs (programs 6 and 7; FDR=9.0 10⁻⁷, 2.0 10⁻¹¹), and its non-epithelial compartment was correspondingly enriched for genes from inflammatory pathways, including the response to TNF, IL-1 and IFN γ (FDR=3.1 10⁻⁴, 2.8 10⁻³, 4.9 10⁻⁶, two-sided Fisher’s exact test in GO term enrichment), and chemotaxis of monocytes, neutrophils, and lymphocytes (FDR=1.5 10⁻³, 3.2 10⁻¹⁰, 1.0 10⁻², Table S2), suggesting recruitment of inflammatory cells from the circulation or other parts of the tissue. Region 6 was also enriched for collagen binding genes and collagen-containing extracellular matrix (ECM) genes (FDR=1.4 10⁻², 7.6 10⁻⁵, two-sided Fisher’s exact test in GO term enrichment, Table S2), which are important for migration and invasiveness⁵² [↗](#). These include *Sparc*, expressed mainly by endothelial cells and fibroblasts in our data, known to promote colorectal cancer invasion⁵³ [↗](#), and *Ctss*, a peptidase expressed by T cells and monocytes-macrophages that promotes CRC neovascularization and tumor growth⁵⁴ [↗](#). Finally, gene expression patterns in endothelial cells and monocytes in region 6 suggested active angiogenesis through a multi-cellular feedback loop, with enriched numbers of vascular and lymphatic endothelial cells expressing immune-attracting chemokines (*Cxcl9*) and adhesion molecules (e.g. *Chd5*, *Mcam*), monocytes expressing proangiogenic factors that induce proliferation of endothelial cells (e.g., *Mmp12*), and monocytes and macrophages expressing *Ctsd*, which increases tumorigenesis in CRC models⁵⁵ [↗](#) (**Figure S8E** [↗](#)).

Region 8 was enriched for deep crypt cells (program 13; FDR=1.7 10⁻¹⁴, two-sided Welch’s t-test on CLR transformed compositions), reminiscent of the normal stem cell niche in normal colon, an epithelial innate immune program (program 1; FDR=3.4 10⁻¹⁰⁰) expressed by secretory cells in AV and AKPV lesions, and plasma and B cell activity. Unlike the canonical (normal) deep crypt region (region 5), which is enriched for MHCII expression (program 18; FDR=8.6 10⁻²⁸), this region was depleted for the program’s expression (FDR=2.1 10⁻²⁶), which may indicate an earlier stem cell-like state⁴³ [↗](#), or a decoupling of the cell cycle and MHCII programs (which are coupled in normal ISC differentiation, and allow a cross talk with T cells to modulate T cell differentiation) (**Figure**

4B,C). The region's non-epithelial compartment was enriched for B cell activation and BCR signaling genes ($FDR=4.0 \cdot 10^{-4}$, $1.9 \cdot 10^{-3}$, two-sided Fisher's exact test in GO term enrichment, Table S2). This may be related to B cell function in protection from lumen antigens⁵⁶ or to tertiary lymphoid structures (TLS), which is correlated with clinical benefits in cancer patients⁵⁷. Notably, Epi05 (dysplastic secretory-like) enriched in Region 8 (**Figure 4B**) expressed higher levels of inflammatory genes and immune chemokines (*e.g.*, *Ccl9*, *Ifitm3*) compared to normal counterparts, Epi04 (secretory) (**Figure 2C** and **Figure S4C,E**), and may thus promote the formation of this region. We validated the presence of TLS-like structures in association with deep crypt secretory cells in AV lesions using multiplex RNA analysis (Cartana) (**Supplementary item 3**), showing that the dominant population of B cells is accompanied by monocyte-macrophages and T cells characteristic of TLSs, as well as the expression of *Reg4* and *Muc2*, deep crypt goblet/secretory cell markers.

Region 11 was populated by cells expressing the Wnt signaling pathway program (4, $FDR=9.3 \cdot 10^{-13}$, two-sided Welch's t-test on CLR transformed compositions), with several lines of evidence supporting an active epithelial to mesenchymal transition (EMT) in this region. Epithelial cells in region 11 were enriched for the expression of mesenchymal genes, including Vimentin⁵⁸ (*Vim*, $FDR=7.4 \cdot 10^{-245}$, one-sided Fisher's exact test), *Prox1*⁵⁹ ($FDR=3.7 \cdot 10^{-153}$), and *Sox11*⁶⁰ ($FDR=7.7 \cdot 10^{-224}$) (**Figure S8F**), as well as for EMT signatures from a mouse model of lung adenocarcinoma⁶¹ ($FDR=1.5 \cdot 10^{-142}$, two-sided Mann-Whitney U test) and from human head and neck squamous cell carcinoma tumors⁶² ($FDR=6.9 \cdot 10^{-55}$, two-sided Mann-Whitney U test). This is consistent with the role of Wnt signaling in promoting EMT and a mesenchymal phenotype in CRC, breast cancer and other epithelial tumors^{63, 64}. Region 11 non-epithelial cells also expressed genes encoding MHC-I binding proteins ($FDR=4.6 \cdot 10^{-2}$, two-sided Fisher's exact test in GO term enrichment, Table S2) and actin cytoskeleton, filament and binding proteins ($FDR=3.1 \cdot 10^{-6}$, $7.7 \cdot 10^{-3}$, $2.7 \cdot 10^{-10}$). Organization of the cytoskeleton affects migration, adherence, and interaction of lymphocytes with antigen presenting cells⁶⁵. Notably, region 11 also concentrated at a more distal part of the tissue at $\sim 900\mu m$ from the muscularis suggesting an outgrowth of the tissue towards the lumen (**Figure S8C**).

Non-epithelial cells formed two cellular hubs in the malignant-like regions (6,8,11) (**Figure S8G**): An endothelial-fibroblast hub, detected in all three regions, and an immune hub with B cells, TNK cells, monocytes, and macrophages, which was prominent in inflammatory region 6, weaker (less spatially correlated) in region 8 (but validated *in situ*), and not correlated in region 11. Thus, activation of an immune response is reflected by close proximity between immune cells. We further characterized the organization of the vascular niche using our multiplex *in situ* RNA data, finding that while neighbors of the *Pdgfrb*-expressing pericytes are mainly other *Pdgfrb*-expressing pericytes and endothelial cells, *Pecam1*-expressing endothelial cells appear self-enriched next to themselves at cellular scale distances and close to *Pdgfrb*-expressing pericytes for larger distances (**Figure S8H**).

Overall, three multicellular community regions were enriched in AV lesions: (1) inflammatory epithelial regions with endothelial cells and monocytes expressing angiogenesis, inflammation and invasion programs; (2) epithelial stem-like regions, associated with plasma and B cell activity; and (3) regions with epithelial to mesenchymal transition (EMT) and Wnt signaling dysplastic cells. Each region highlights different processes that modulate tumorigenesis or invasion, and the three regions co-exist in the same tumor at different spatial locations.

Cell-cell interactions are rewired in AV lesions

To identify cell-cell signaling mechanisms that may underly these regional associations, we used COMMOT⁶⁶, a computational framework that uses Optimal Transport to infer cell-cell communication from receptor-ligand expression patterns in spatially resolved data. We used COMMOT's bead-wise communication 'output' and devised a method to address p-value inflation in statistical enrichment testing, using spatially-informed data aggregation (**Methods**).

We observed stronger and distinct ligand-receptor interactions in AV vs. normal samples, reflecting the activated state in the dysplastic tissue (**Figure 4K** [↗](#), **Supplementary item 4A** [↗](#)). In particular, while interactions enriched in AV lesions involved immune, epithelial and stromal signaling, those enriched in normal tissue involved neuropeptides, such as NPY and GCG (glucagon). Moreover, malignant-like regions 6 and 11 as well as region 2 (muscularis) were particularly enriched for active communication pathways (**Figure S8I** [↗](#) and **Supplementary item 4B** [↗](#)). This included the WNT signaling pathway, angiogenesis (VEGF, PDGF, FGF) and the OSM pathway.

Similar spatial patterns in human and mouse tumors

The overall spatial distribution of cell types and epithelial profiles was conserved between mouse and human tumors, when comparing to scRNA-seq^{2, 28, 29, 67–70}. We examined mouse-defined regions in human tumors, using TACCO to map the expression profiles associated with the epithelial, immune and stroma compartments in each of the TACCO-identified mouse regions to scRNA-seq profiles from human CRC, and probabilistically annotated region-specific expression profiles for each scRNA-seq profile from the human samples. This identified two main “meta compartments”, with epithelial, stromal and immune profiles from human MMRp and MMRd tumors associated with regions 6, 8 and 11 that were enriched in AV lesions (as well as 0 and 2), while those from normal human tissue were associated with normal regions (e.g., 5, 10, 12) (**Figure 5A** [↗](#) and **Figure S9A** [↗](#), **Methods**).

Malignant-like regions are associated with tumor progression in human colorectal tumors

We next assessed if the regional epithelial programs that we spatially identified in mouse are conserved in human. To this end, we constructed pseudo-bulk profiles from epithelial cells for our mouse samples and for recently published human samples profiled along different stages of malignant transformation, from normal tissue to polyp to CRC^{2, 28, 29, 67–70}. We scored each epithelial pseudo-bulk profile with the differentially expressed genes between the epithelial parts of the regions and computed the principal components of these scores across all human and mouse samples (**Methods**). The first principal component (PC1) captured features that are related to malignancy, with higher values for human tumors vs. polyps (**Figure 5B** [↗](#) and **Figure S9B** [↗](#)). In addition, malignant-like region (6/8/11) scores were higher in dysplastic vs. normal samples (**Figure 5C** [↗](#)). Thus, the spatial region profiles defined in mouse capture features that correlate with malignant transformation in human.

We further classified each full pseudo-bulk profile from the dysplastic human samples into one of the four groups in the CMS expression-based classification⁷¹ (**Methods**) and compared the mouse region scores for each class of samples (**Figure S9C** [↗](#), **Methods**). CMS2 classified samples were most closely related to our dysplastic mouse models: all of the dysplasia-associated regions were enriched in CMS2 tumors while most normal regions were depleted, relative to the other CMS classes.

Finally, we found that expression of the malignant like regions (6/8/11) in tumors was associated with clinical outcome. We scored each tumor based on genes that were differentially expressed between the full expression profile of malignant-like regions (6/8/11) and compared the progression-free interval (PFI) and overall survival (OS) for patients in TCGA whose RNA-seq profiles were in the top and bottom quartile of malignant-like region scores (**Methods**). High scores for malignant-like region 11 (EMT) were correlated with shorter PFI, while those for malignant-like region 6 (inflammation) correlate with longer PFI and longer OS, (**Figure 5D,E** [↗](#)). These associations were driven primarily by MMRp tumors (**Figure S9D,E** [↗](#)). This suggests that




Figure 5.

Mouse tumor regions associated with tumor progression in human colorectal tumors.

A. Expression profiles characterizing mouse regions are recapitulated in human tumors. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of region-associated epithelial, immune or stromal profiles (rows) compared between normal, MMRp, or MMRd samples (columns). **B,C.** Mouse regions capture malignant features in human tumors. **B.** Top left: First (PC1, x axis) and second (PC2, y axis) principal components of mouse region scores of mouse and human epithelial pseudo-bulk samples. Top right: PC1 loadings (x axis) of each mouse region score (y axis). Bottom: PC1 values (box plots show mean, quartiles, and whiskers for the full data distribution except for outliers outside 1.5 times the interquartile range (IQR)) for each type of mouse or human sample (x axis). **C.** Significance (FDR, color bar, two-sided Welch's t-test) of enrichment (red) or depletion (blue) of region-associated profile scores (rows) in normal and dysplastic samples (columns) in human or mouse. **D,E.** Expression of malignant like regions 6 and 11 in tumors are associated with PFI (D) and OS (E) in human patients. Kaplan-Meier PFI (e, n = 662⁹⁵) or OS (f, n = 662⁹⁵) analysis of human bulk RNA-seq cohort stratified by malignant-like region profile scores.

region 11 is associated with pro-tumorigenic properties in human patients, while region 6 might be associated with tumor controlling properties. This highlights the importance of multicellular functional tissue modules in the CRC tumor ecosystem.

Discussion

Here, we systematically charted the spatial organization of cellular expression in dysplastic tissue of the colon, to help identify putative functional units in the tumor. We used TACCO²¹ to integrate scRNA-seq and Slide-seq data, not only by mapping cell types to their positions, but also distinguishing different cell programs, the regions that they dominate, and their characteristic microenvironments. This allowed us to overcome technical limitations, such as lack of spatial context in scRNA-seq and sparse readout in Slide-seq, and to generate a high-resolution spatial map of the dysplastic landscape transcending beyond the mapping of individual cells to spatial positions. We used this map to show correlation with clinical outcome in human patient tumors.

Our scRNA-seq analysis revealed profound enrichment of a stem cell program in dysplastic tissues. The profiles of dysplastic cells expressing this program are distinct from normal stem cells and enriched with expression of negative regulators of the WNT signaling pathway and inflammation, suggesting a non-canonical function. The abundance of these cells with stemness potential across all our malignant-like regions, points to a dynamic population that can affect the cells in its proximity, by secretion of negative regulators of the WNT signaling and inflammatory function but may also adopt various functions depending on the environmental cues and dysplasia-associated cells in its proximity. A similar population, designated “high-plasticity cell state”, was previously described in a mouse model of lung adenocarcinoma and in human patients, where it was correlated with resistance to chemotherapy⁶¹. Whether these cells can be manipulated to take on specific phenotypes or even to differentiate into normal-like enterocytes given the appropriate signal from the microenvironment, remain as open questions.

Within the dysplastic lesions, alongside malignant-like regions, we found regions with normal features (Regions 3,4,9,10), comparable to regions found in the normal colon, most likely representing compartments driven by clones that were not affected by the genetic perturbation. One of these regions, region 4, contained mainly goblet cells with normal expression profiles. Whether this neighborhood represents normal cells that reside alongside malignant cells or a cancer transition state, it may modify tumor progression, by recruiting immune cells or by secreting factors that affect epithelial proliferation in adjacent regions. For example, region 4 in dysplastic lesions is enriched with chemokine activity genes relative to region 4 in normal colon suggesting a possible role in recruitment of immune cells to the dysplastic landscape. Further work is required to understand the role of these regions (expressing normal features) in tumor progression.

While the malignant-like regions were identified as discrete spatial entities, each with coordinated features across epithelial, immune, and stromal cells, these regions are adjacent to each other. Thus, they may still influence one another through signaling or by utilizing branches of the same main vessels. For example, *Osm* is expressed by cells in region 6, whereas its receptor is expressed on fibroblasts and endothelial cells enriched in region 11. *OSMR* was previously shown to be expressed by inflammatory fibroblasts^{44, 72} and its activation in malignant cells promotes EMT in breast cancer and pancreatic cancer^{73, 74} and a mesenchymal state in glioblastoma⁷⁵. Future studies can help determine if these regions are functionally inter-dependent and if they evolved from the same clones and can inter-convert, or whether they developed independently.

Because animal models complement cell and tissue atlases of human colorectal cancer^{28, 29}, by allowing experimental manipulation for mechanistic studies^{76, 77}, it is important to relate between models and patients. By studying genetically engineered mouse models using high resolution single cell spatial genomics we can help determine to what extent they recapitulate the cellular and spatial organization of human disease, in the context of two distinct genetic states that represent human CMS-2 lesions. To this end, we developed several approaches to allow cross species comparison of tumors at the single cell and spatial level, despite the high level of both intra- and inter-individual variation within each species. Comparing to human CRC, our analysis suggests that the CRC landscape is organized in similar multicellular functional tissue modules between human and mouse, and disease subtypes (e.g., MMRp and MMRd). Future studies applying our approaches to patient cohorts could help understand whether the expression of different tissue modules may contribute to the partial response to immunotherapy reported for MMRd patients⁷⁸, and to define specific tissue modules predictive of response to therapy. Notably, while our study focused on the tumor landscape, its findings may be relevant for tissue response to other challenges (e.g., inflammation, fibrosis, wound healing), which involve activation of similar functional tissue modules, a result of collective function of parenchymal, immune and stromal cells.

Taken together, our integrative approach facilitates spatial analysis with high resolution, constructing regional neighborhoods and their spatial layout at both high cellular resolution and genomic scale. Our work is an important step toward a systematic understanding of the organization of dysplastic tissue with the potential to contribute to improved patient stratification by the multicellular functional units in the tumor landscape.

Methods

Human subjects

The MGH Institutional Review Board approved protocols for tissue collection used for sequencing (Protocol 02-240). Informed consent was obtained from all subjects prior to collection. Only patients with primary treatment-naïve colorectal cancer were included in this study.

Mice

Mice were housed in the animal facility at the Koch Institute for Integrative Cancer Research at MIT. All animal studies described in this study were approved by the MIT Institutional Animal Care and Use Committee (Protocol 1213-106-16). *Apc*^{fl/fl} mice⁷⁹ were obtained from NCI mouse repository, *Kras*^{LSL-G12D/+} Ref.⁸⁰, *Rosa26*^{LSL-tdTomato} Ref.⁸¹ and *Trp53*^{fl/fl} Ref.⁸² mice obtained from Jackson, *Villin*^{CreERT2} Ref.⁸³ mice were a gift from Dr. Sylvie Robine. All mice were maintained on C57BL/6J genetic background. Approximately equal numbers of male and female mice of 6–10 weeks of age were used for all experiments. Where indicated, mice were injected to the submucosal layer of the colon with 4-hydroxytamoxifen (EMD Millipore # 579002) dissolved in ethanol at a concentration of 100 μ M (for the mice that were kept for 3 weeks after injection) or 30 μ M (for the mice that were kept for 9 weeks after injection). Tumors were resected at either 3 or 9 weeks after 4-hydroxytamoxifen injection. Colonoscopy and colonoscopy-guided injection methods were previously described in detail.^{30, 31}

Tissue processing for scRNA-seq

Single-cell suspensions from healthy colon or dysplastic lesions were processed using a modified version of a previously published protocol⁴⁴. Tissue samples were rinsed in 30ml of ice-cold PBS (ThermoFisher 10010-049), chopped to small pieces and washed twice in 25 ml PBS, 5mM EDTA (ThermoFisher AM9261), 1%FBS (ThermoFisher 10082-147). To prime tissue for enzymatic digestion, samples were incubated for 10 minutes at 37°C, placed on ice for 10 minutes before

shaking vigorously 15 times followed by supernatant removal. Tissues were placed into a large volume of ice-cold PBS to rinse prior to transferring to 5ml of enzymatic digestion mix (Base: RPMI1640, 10 mM HEPES (ThermoFisher 15630-080), 2% FBS), freshly supplemented immediately before use with 100 mg/mL of Liberase TM (Roche 5401127001) and 50 mg/mL of DNase I (Roche 10104159001), and incubated at 37°C with 120 rpm rotation for 30 minutes. After 30 minutes, enzymatic dissociation was quenched by addition of 1ml of 100% FBS and 10mM EDTA. Samples were then filtered through a 40 mM cell strainer into a new 50 mL conical tube and rinsed with PBS to 30 mL total volume. Tubes were spun down at 400 g for 7 minutes, at 4°C. Resulting cell pellets were resuspended in 1ml PBS, placed on ice and counted.

Cell hashing

Cell hashing was performed based on the published protocol⁸⁴ as summarized below. Dissociated cells were resuspended in 1ml of Cell Hashing Staining Buffer (1× PBS with 2% BSA (New England Biolabs, B9000S) and 0.02% Tween (Tween®-20 Solution, 10%, Teknova, VWR-100216-360) and counted. 500,000 cells were resuspended in 100 µL of Cell Hashing Staining Buffer and incubated for 30 minutes on ice, with 2 µL of the appropriate BioLegend TotalSeq™ Hashing antibody (a 1:50 dilution, using a total of 1 µg of antibody per cell suspension). TotalSeq™-A anti-mouse Hashtag antibodies #1-8 (catalog numbers:155801, 155803, 155805, 155807, 155809, 155811, 155813, 155815) were used. Cells were washed three times with 0.5 mL of Cell Hashing Staining Buffer and filtered through low-volume 40-µm cell strainers. All cell suspensions were recounted to achieve a uniform concentration of 7,000 cells per microliter before pooling for capture by 10x Chromium controller following the manufacturer protocol for the v2 or v3 3' kit (10X Genomics, Pleasanton, CA).

Hashtag oligo (HTO) library preparation

Separation of hashtag oligo (HTO)-derived cDNAs (<180 bp) and mRNA-derived cDNAs (>300 bp) was done after whole-transcriptome amplification by performing 0.6× SPRI bead purification (Agencourt) on cDNA reactions as described in 10x Genomics protocol. Briefly, supernatant from 0.6× SPRI purification contains the HTO fraction, which was subsequently purified using 1.4 and 2× SPRI purifications per the manufacturer's protocol (Agencourt). HTOs were eluted by resuspending SPRI beads in 15 µL TE. Purified HTO sequencing libraries were then amplified by PCR (1µL clean HTO cDNA, 25µL 2X NEBNext Master Mix (NEB #M0541)), 10 µM SI-PCR and D701 or D704 primers performed dial out PCR (98°C (10 sec), (98°C for 2 sec, 72°C for 15 sec) x 12/18 then 72°C for 1 min) for 12 and 18 cycles, and used the 18 cycles product for sequencing. PCR reactions were purified using another 2× SPRI clean up and eluted in 15 µL of 1× TE. HTO libraries were quantified by Qubit High sensitivity DNA assay (ThermoFisher) and loaded onto a BioAnalyzer high sensitivity DNA chip (Agilent).

SI-PCR: **AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGC***T*C

D701: CAAGCAGAAGACGGCATACGAGATCGAGTAATGTGACTGGAGTTCAGACGTGTGCG

D704 : CAAGCAGAAGACGGCATACGAGATGGAATCTCGTGACTGGAGTTCAGACGTGTGCG

Sequencing

Samples were sequenced using HiSeq X (Illumina). Hashing libraries were sequenced with spike-ins of 2.5%.

Tissue processing for Slide-seq

Colons were flushed with cold PBS and a segment that includes the lesion and surrounding tissue (or a respective healthy segment from normal mice) was dissected. Samples were then mounted in cold OCT, flash frozen on dry ice covered with ETOH and long-term stored in -80°C.

Slide-seq

For mouse and human experiments, 10 mm sections were cut and the Slide-seq V2 protocol was used as previously described¹⁵. For mouse experiments, four and six arrays were collected from normal colons and AV lesions respectively. The muscularis was fit onto the array of both healthy and dysplastic lesions to allow appropriate orientation.

Multiplex *in situ* RNA Analysis

Multiplex *in situ* RNA analysis was performed with Cartana³³ technology (a newer version is now available as Xenium (10x Genomics)). In total we measured 3 samples with one section per sample in each state (normal/AV) and probe set (V1/V2), with an additional replicate section for one of the samples in normal V2.

Fresh Frozen OCT-embedded tissues from normal colon and AV lesions were cryosectioned as 10 µm sections and placed onto SuperFrost Plus glass slides (ThermoFisher) and further stored at -80°C before experiments. Samples were fixed (with 4% formaldehyde) for 5 minutes and permeabilized for 5 minutes (with 0,1 mg/mL pepsin in 0.1M HCl (P7012 Sigma-Aldrich)) prior to library preparation.

For library preparation, chimeric padlock probes (directly targeting RNA and containing an anchor sequence as well as a gene-specific barcode) for a custom panel of 66 (V1) or 180 (V2) genes (Table S3, see below) were hybridized overnight at 37°C, then ligated before the rolling circle amplification was performed overnight at 30°C using the HS Library Preparation kit for CARTANA technology and following manufacturer's instructions. All incubations were performed in SecureSealTM chambers (Grace Biolabs). Note that prior to final library preparation, optimal RNA integrity and assay conditions were assessed using Malat1 and Rplp0 housekeeping genes only using the same protocol.

To prevent tissue sections detachment, an additional baking step of 30 minutes at 37°C was performed before mounting. To quench autofluorescence background, TrueView (SP-8400 VectorLabs) was used for 1 minute at room temperature. For tissue sections mounting, Slow Fade Antifade Mountant (Thermo Fisher) was used for optimal handling and imaging.

Quality control of library preparation was performed by applying anchor probes to detect simultaneously all rolling circle amplification products from all genes in all panels. Anchor probes are labeled probes with Cy5 fluorophore (excitation at 650 nm and emission at 670 nm). All samples passed quality control and went through *in situ* barcode sequencing, imaging and data processing. Briefly, adapter probes and sequencing pools (containing 4 different fluorescent labels: Alexa Fluor® 488, Cy3, Cy5 and Alexa Fluor® 750) were hybridized to the padlock probes to detect the gene-specific barcodes, through a sequence specific signal for each gene specific rolling circle amplification product. This was followed by imaging and performed 6 times in a row to allow for the decoding of all genes in the panel. To reduce lipofuscin autofluorescence, 1X Lipofuscin Autofluorescence Quencher (Promocell) was applied for 30 seconds before fluorescence labeling.

Raw data consisting of 20x or 40x images from 5 fluorescent channels (DAPI, Alexa Fluor® 488, Cy3, Cy5 and Alexa Fluor® 750) were each taken as z-stack and flattened to 2D using maximum intensity projection. After image processing and decoding, the results were summarized in a csv file and gene plots were generated using MATLAB⁸⁵.

scRNA-seq pre-processing and quality control filtering

Count matrices for scRNA-seq were generated using the Cumulus feature barcoding workflow v0.2.0⁸⁶ with CellRanger v3.1.0 and the mm10_v3.0.0 mouse genome reference. Cell profiles were quality filtered by requiring between 1,000 and 50,000 counts, and between 500 and 7,000 genes, less than 20% mitochondrial counts, and less than 10% hemoglobin counts. Cell profiles that did not meet all these criteria were discarded. The top 5,000 highly variable genes were annotated on the remaining cells after normalization to 10,000 counts and log1p transform using Scanpy's "highly_variable_genes" function⁸⁷ and providing the chemistry (v2/v3) by hashing (True/False) combination as batch-annotation. Putative doublets were removed using Scrublet⁸⁸ with default parameters.

Selection of variable genes, dimensionality reduction and clustering

A preliminary clustering using the Leiden algorithm with resolution 1.0 was performed after normalization to 10,000 counts, log1p transform, correction for number of counts and percentage of mitochondrial genes, scaling with a max_value of 10, and generating a *k*-nearest neighbors (*k*-NN) graph with 15 neighbors on a PCA of the previously annotated 5000 highly variable genes with 50 components using Scanpy⁸⁷. The single cell profiles were provisionally annotated with SingleR⁸⁹ cell-wise (i.e. without using clustering information) using the SingleR built-in MouseRNAseqData and an intestine specific dataset from Tabula Muris⁹⁰ (<https://figshare.com/download/files/13092143>). For further processing, the dataset was then split into the three compartments, epithelial, immune and stromal, using the provisional SingleR annotations.

For each compartment, the top 5,000 highly variable genes were annotated using Scanpy's "highly_variable_genes" function on cells normalized to 10,000 counts after log1p-transformation and providing the chemistry (v2/v3) by hashing (True/False) combination as batch-annotation.

Expression programs and batch correction

For the dataset of each compartment separately (generated as described above), an integrative NMF was performed (using a part of the LIGER⁴⁵ implementation) with *k*=20 and lambda=5 to identify 20 programs and their respective weights per cell. This iNMF factorization represents the single cell expression matrix as a weighted sum of profiles such that both the weights and programs contain only non-negative numbers, while allowing for and separating out batch-only contributions. The same approach was also used with a higher *k* (epithelial and immune: 200, stromal: 50) to yield a detailed and batch corrected decomposition of expression which was then combined to obtain a count-like corrected expression matrix for the top 5,000 highly variable genes. For each compartment separately, these batch-corrected data were normalized to 10,000 counts, log1p transformed, corrected for number of counts and percentage of mitochondrial genes by linear regression, scaled with a max_value of 10, followed by a PCA of the previously annotated 5,000 highly variable genes. A *k*-nearest neighbors (*k*-NN) graph was constructed from the top 50 PCs, with *k*=15 neighbors using Scanpy, and clustered using a compartment-specific Leiden resolution parameter (epithelial: 0.2, immune: 0.4, stromal: 0.1). This clustering was used as the cluster level annotation of the mouse scRNA-seq data for the epithelial and stromal compartment. Separately per compartment the data were annotated with SingleR using the cluster information. The same per-compartment batch-corrected and preprocessed data from the Leiden clustering was used to create UMAP embeddings with PAGA initialization using Scanpy.

To improve the clustering and annotation in the immune compartment and to filter out additional doublets not detected by Scrublet, the immune data were separately filtered and clustered using information from the compartment level clustering and annotation. To that end, myeloid and T/NK

cells were partitioned separately and further processed, and additional likely doublet cells were labeled and removed by the following procedure:

1. Cells were labeled as doublets based on higher number of UMIs of marker genes for other compartments than the 95th percentile observed in this immune partition (i.e., Epcam and Cdh1 to remove immune-epithelial doublets and Cav1 and Kdr to remove immune-stromal doublets) and other immune partitions (i.e., Cd3d, Cd3e, and Cd3g to remove myeloid-lymphoid doublets from the myeloid cells). This type of filter criterion for lowly expressed genes (“larger than some percentile” on integer counts) also allows to keep more than 95% of the cells if, for example, all cells of this partition happened to have 0 UMIs of a particular marker gene.
2. Cells were labeled as doublets if they had inconsistent cell-wise and cluster-wise SingleR annotations.
3. Cells were labeled as doublets if they had significantly (Benjamini-Hochberg FDR=0.05, one-sided Fisher’s exact test) more neighbors in the k-NN graph from the immune compartment that were already marked as doublets.
4. All cells labeled as doublets were removed.

After filtering, the count matrices were batch corrected as above using the integrative NMF from LIGER with $k=20$ and $\lambda=5$, and clustered like above with group specific Leiden resolution (myeloid: 0.2, TNK: 0.4). For myeloid and TNK cells, this clustering superseded the original clustering. The integrative NMF result here was only used for updating the clustering and not for generating an extra set of expression programs.

Note that cluster Epi06 shows a broad expression spectrum; while such pattern can best be explained by remaining doublets, they were not called by Scrublet and could also not be consistently removed by extra QC as was applied to the immune compartment. The interpretation of Epi06 as containing doublets is supported by its overrepresentation in the annotation of the Slide-seq data (**Figure S6E** [↗](#)), as both doublets and slide-seq beads represent compositions of different cells. Still, non-doublet explanations (e.g. undifferentiated cells) cannot be ruled out. Therefore, Epi06 cells are retained in the data but not interpreted biologically.

Marker selection for *in situ* RNA analysis (CARTANA)

CARTANA V1 markers were selected from genes differentially expressed between compartments, cell types and clusters (for the immune and stromal compartments), and highly ranked genes for programs (for the epithelial compartment) were first filtered by biological relevance and the literature, obtaining 87 genes. To further reduce the set to the available panel size (66 genes), genes were annotated into categories of potential redundancy. To choose between redundant genes, a global objective function was optimized over the gene selection, looping over all potentially redundant gene sets until convergence, exhaustively testing all choices within a gene set, and accepting the best choice for this gene set in terms of the global objective function. The global objective function was constructed as the mean 4-fold cross-validation scores (using “GroupKFold” and “cross_val_score” from sklearn⁹¹ [↗](#)) of multi-class logistic regression classification (using “LogisticRegression” from sklearn) for discriminating cell classes and of ridge regression of epithelial program weights (using “RidgeRegression” from sklearn) on the scRNA-seq data subsampled to the expected sparsity of CARTANA data. The cell classifications used were between the stromal, immune, and epithelial compartment, within the stromal compartment between Endo01, Endo02, and fibroblasts, within the immune compartment between myeloid and lymphoid lineage, within the lymphoid lineage between T cells and B cells, within the myeloid lineage between granulocytes, mast cells, and all monocytes and macrophages together, between monocytes and macrophages, and within monocytes between Mono01, Mono02, Mono03, and

Mono04. The programs used in ridge regression were programs 3, 4, 6, 7, 13, 14, 15, and 16. As the probes for Ly6c1 and Ly6c2 could not discriminate sufficiently between Ly6c1 and Ly6c2, we chose combined probes that measure both.

The CARTANA V2 panel, included 59 of the 66 genes in the V1 panel (the others had to be removed for technical reasons), another 113 genes from the standard fixed gene panel for CARTANA, and 8 selected genes from literature.

Analysis of CARTANA data

Each measured molecule was annotated with an originating cell type cluster label (using TACCO's "tc.tl.annotate_single_molecules", with RCTD¹⁹ as the core annotation method and parameters bin_size=20, n_shifts=3, assume_valid_counts=True) separately for each sample. For this, genes in the reference that would likely cross hybridize in the probe panel design were summed over (Ly6c1 and Ly6c2). TLS-like regions were annotated by visual inspection of the cell type cluster composition and morphology.

To assess cell type compositions of the full dataset, molecules with cell type cluster annotations were binned into 10µm bins (using TACCO's "tc.utils.bin" and "tc.utils.hash" functions) and cluster-level annotations were merged to cell-type level.

To assess the compositions of TLS-like regions, CARTANA v2 data were aggregated, conserving the categorical TLS annotation (using TACCO's "tc.utils.bin" and "tc.tl.dataframe2anndata" functions).

Comparison between experimental methods

To compare cell type composition between methods, CLR-transformed compositions of samples (or of spatially split samples for the spatial methods, see subsection "Ligand-Receptor Analysis and Spatially Informed Enrichment") were computed. Then, using a 100 bootstrapped means of the compositions, mean and standard error of the mean were calculated once for the AV and once for the normal samples. To compare gene expression, the mean difference of gene counts between normal and AV samples and its standard error were calculated using all pairwise differences between the bootstrapped normal and AV samples. For gene comparisons, the CLR-transformed composition over all genes that were measured in all of scRNA-seq, Slide-seq, CARTANA V1, and CARTANA V2 was used. Pearson correlation between mean compositions was calculated for each pair of methods.

RNA-velocity analysis

Splicing aware count matrices for scRNA-seq were generated using CellRanger v6.1.2 and velocity v0.17.17³⁷ with the ensembl v108 mouse genome reference. Scvelo v0.2.5 was used to infer velocity separately for the epithelial and TNK subsets (using the functions "scv.pp.filter_and_normalize", "scv.pp.moments", "scv.tl.velocity" (with mode='stochastic'), and "scv.tl.velocity_graph"). Scanpy and bbknn v1.5.1⁹² were used to generate batch-corrected UMAP embeddings for the two subsets for visualization with scvelo's "scv.pl.velocity_embedding_stream" function.

Selection of human single cell data for the comparison of cell type and epithelial program composition

ScRNA-seq data from Ref² was used as reference for human CRC. To avoid biases in cell type compositions, only the subset of the data where "PROCESSING_TYPE = unsorted" were used.

Comparison of human and mouse samples by cell type composition

To compare human and mouse samples by composition of T/NK cell subsets, T/NK annotations from mouse and human data²¹ were matched by TACCO, using optimal transport (OT). First, human expression data were mapped to mouse genes using MGI homology information [subsection “Mapping of mouse and human orthologs”]. Then, human cell cluster annotations (‘cl295v11SubFull’) were mapped from the subset of human cells annotated as T/NK/ILC to the subset of mouse cells annotated as T/NK using TACCO’s “annotate” function with OT as core method, basic platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter lambda of 0.1, and 4 iterations of bisectioning with a divisor of 3. Annotation with maximum probability per cell was used as the unique cluster level annotation for mouse T/NK cells. Annotations were aggregated per sample to yield a compositional annotation over the identical cluster annotation categories (from the human dataset) for the T/NK subsets of human and mouse samples. Annotation vectors were then processed using the `sc.pp.neighbors` and `sc.tl.umap` functions from Scanpy⁸⁷ to yield a 2D sample embedding with respect to T/NK cell composition. Using the coordinates in the UMAP in place of spatial coordinates, neighborhood enrichment z-scores were computed with TACCO’s `co_occurrence_matrix` function with `max_distance=2` and `n_permutation=100`.

Slide-seq compositional annotation

Slide-seq data were annotated with scRNA-seq reference annotations. First, Slide-seq and scRNA-seq data were filtered to retain only 15759 genes that were detected in both datasets and only beads and cells that had at least 10 counts across all these common genes.

Next, TACCO²¹ was used to perform compositional annotation of each bead, allowing the bead counts to be explained by fractional contributions. In its basic application, TACCO finds an “optimal” mapping between scRNA-seq annotation categories (*e.g.*, cell types) and beads by solving a variant of an entropically regularized Optimal Transport (OT) problem in expression space. In its iterative application, TACCO uses a bi-sectioning functionality iteratively, annotating only fractions of the counts in each round, and reserving the remainder for the next round to improve the sensitivity to sub-leading annotation contributions (that is, first capture a portion of the counts for the “top” cell types, but preserving others for other, more minor, cell types).

For the compositional annotation of Slide-seq beads with the categorical cell clusters from the single cell data, the “annotate” function of TACCO with OT was used as core annotation method per Slide-seq puck with the subset of the single cell data with matching disease state, with basic platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter lambda of 0.001, and 4 iterations of bisectioning with a divisor of 3.

For the compositional annotation of Slide-seq beads with the compositional epithelial programs, the annotated beads were split using the “split_observations” function of TACCO on the cluster-level annotation and then aggregated to compartment level using the “merge_observations” function keeping only beads for a compartment with at least 50 counts assigned to that compartment. The genes from the epithelial part were filtered to retain only those that were used to define the epithelial programs in scRNA-seq, and then annotated using again the “annotate” function with OT as core annotation method, basic platform normalization, entropy regularization parameter epsilon 0.01, and a marginal relaxation parameter lambda of 0.001.

Slide-seq region annotation

Region annotation was done for all pucks (normal and AV pucks) in one step to get comparable region annotations across pucks, such that beads that have similar epithelial program activity and are spatially close are called as one region. Because cell type composition can change drastically from one bead to its neighbor at the length scales of Slide-seq data, there is a need to compromise between optimizing the two similarities. This is done with the “find_regions” function of TACCO, which performs a Leiden clustering on the weighted sum of connectivity matrices derived from epithelial program similarity and spatial proximity, using a position weight of 0.7, a Leiden resolution of 1.3, and 15 nearest neighbors per bead in position space and epithelial program space. To determine the neighbors in epithelial program space, the square-roots of the program weights were used for neighbor finding which effectively uses Bhattacharyya coefficients as overlap in epithelial program space instead of the Euclidean scalar products used for position space. These regions are defined by construction only on beads with a large enough epithelial contribution (see above) and are then extended to all beads by assigning unannotated beads the region from the nearest bead with region annotation.

Submucosal and muscularis propria layers are predominantly comprised of fibroblasts and muscle cells, respectively, alongside blood and lymphatic vessels, nerves and immune cells. Our algorithm depends on the epithelial expression component in beads. Since these layers do not contain epithelial cells⁹³, mapping likely relied either on “noisy” signal from non-epithelial cells or from the basal-most epithelial layer.

To determine region composition at a certain distance of a reference region, TACCO’s “annotation_coordinate” function is used with max_distance=1000 and delta_distance=10.

Slide-seq quality filtering

For all downstream analyses, all beads with less than 100 reads were discarded.

Region- and cell type-characterizing genes in Slide-seq data

Genes to characterize regions on Slide-seq pucks irrespective of compartment composition were found using Scanpy’s rank_genes_groups function on the full bead expression profiles. To find them separately for each compartment, the compartment-level split beads [sub-section “Slide-seq annotation”] were used instead of the full beads. To compare gene expression between cell types on Slide-seq pucks, cluster-level split beads [sub-section “Slide-seq annotation”] were aggregated to cell type level.

EMT scoring

Malignant regions were scored for EMT signatures, using only counts attributed to the epithelial compartments within these regions and only genes expressed on at least 3 beads. Bead profiles were normalized to 10,000 counts, log1p transformed and scaled, and Scanpy’s “sc.tl.score_genes” function was used to score the top 50 genes in two EMT gene signatures^{61, 94}.

Cell-type neighborships in Slide-seq data

To evaluate the local cell-type neighborhood relations in the different disease states on the cluster level, the clusters were filtered per disease state to contain only clusters which account for at least 1% of the UMIs in that state. Then neighbourhood-enrichment z-scores were calculated using TACCO’s “co_occurrence_matrix” function with max_distance=20 and n_permutation=10. To evaluate the stability of the result, this is also repeated for (max_distance, n_permutation)=(40,10),

(60,10),(20,5), and (20,50). To get the significance of the overall change in z-scores between the states, a one-sided Mann-Whitney-U test was performed on the values of the upper triangular half of the matrix between the two disease states for (max_distance, n_permutation)=(20,10).

A similar neighborhood analysis was performed on the coarser cell-type level separately for the three malignant regions 6, 8, and 11, using TACCO's "co_occurrence_matrix" function with max_distance=20 and n_permutation=10.

Cell-type co-occurrence in Slide-seq data

Cell-type compositions relative to a spatial landmark, Region 2=muscularis, were evaluated using TACCO's "annotation_coordinate" function with max_distance=1000 and delta_distance=10. To reduce tissue structure bias from the muscularis, the distance dependency of cell-type frequency relations was evaluated only for beads deep in the "epithelial domain", defined as follows. The effective distance from stromal annotation was computed using TACCO's "annotation_coordinate" function (with max_distance=100, delta_distance=10, critical_neighbourhood_size=4.0) and only beads with a distance of at least 75µm were used. On these remaining beads, TACCO's "co_occurrence" function was used (with delta_distance=20, max_distance=1000) to compute cell types co-occurrence as a function of their distance.

Epithelial program neighborhoods in Slide-seq data

As for cell types above, neighborhood relations were evaluated for epithelial programs in the AV Slide-seq samples using TACCO's "co_occurrence_matrix" function with max_distance=20 and n_permutation=10, after selecting only the programs which make up at least 1% of the UMIs in the AV Slide-seq samples.

Ligand-Receptor analysis and spatially informed enrichment

For the ligand-receptor analysis on slide-seq data we used COMMOT v0.03⁶⁶[\[66\]](https://doi.org/10.5281/zenodo.7272562). COMMOT employs optimal transport to construct sender and receiver side receptor-ligand interactions for every bead in one run of COMMOT. After filtering to beads with at least 100 counts, we applied basic preprocessing (normalization, log1p-transformation) and load the CellChat database as done in the COMMOT tutorial (https://commot.readthedocs.io/en/latest/notebooks/Basic_usage.html[\[67\]](#)). We then follow the Slide-Seq v2 analysis from the COMMOT paper ("slideseqv2-mouse-hippocampus/1-lr_signaling.ipynb" from <https://doi.org/10.5281/zenodo.7272562>[\[68\]](#)) to filter the database and to reconstruct the spatial communication network on ligand-receptor pair and pathway level separately for every puck. In particular, we use the distance cutoff of 200µm for inference of ligand-receptor interactions. The resulting bead-wise sender- and receiver communication values were then used for enrichment analysis between disease states and between spatial regions.

Unlike the downstream Slide-Seq v2 analysis from the COMMOT paper ("slideseqv2-mouse-hippocampus/2-downstream_analysis.ipynb" from <https://doi.org/10.5281/zenodo.7272562>[\[68\]](#)), we do not treat each bead on the puck as statistically independent observation in statistical tests, which leads to unreliably small p-values. Instead, we split all pucks along their axis of greatest extent (defined by the first principal component axis of the distribution of the spatial measurements) into spatial patches discarding 400µm of boundary layer (twice the COMMOT distance cutoff, also co-occurrences have decayed strongly at this distance) in between the patches to reduce the correlation between the patches. This is done iteratively with the patches to get a set of weakly correlated patches. On these patches we calculate the mean of the communication values and treat them as statistically independent observations for statistical tests. We argue that multiple sufficiently separated spatial patches of single spatial samples can be seen as multiple spatial samples using a spatial method with a smaller measurement area, and therefore can be treated as replicates. Unlike splitting into patches without removing a boundary layer, this procedure does not converge to the case of treating each bead as an independent observation as

the number of iterations rises as it accounts for the spatial correlations between adjacent measurements. This gives a natural lower limit of p-values reachable with p-values rising again if too many splits are performed as too much data is lost to remove the correlations (**Supplementary item 4C,D**). We chose a number of iterations of 2 as a compromise between having more patches and not discarding much data. For the enrichment of the pathway communication values we used a two-sided Mann-Whitney U test across the patches and cite Benjamini-Hochberg FDR values. For the enrichment testing in a given group (e.g. an annotated spatial region) only those patches are used which have at least 100 beads on the patch.

Unless otherwise noted, for consistency, an analogous enrichment procedure was also used for COMMOT-unrelated quantities, like cell types, even though their spatial footprint is not as large as COMMOT's (distance cutoff of 200µm). For cluster-level cell type enrichment analyses on Slide-seq data, only the top 3 contributing subtypes were considered, to better reflect the expected properties of Slide-seq data and better compare with categorically annotated scRNA-seq data.

Mapping of mouse and human orthologs

We applied TACCO's functions "setup_orthology_converter" and "run_orthology_converter" with option "use_synonyms=True" to map human to mouse genes using the ortholog mapping from Mouse Genome Informatics (<http://www.informatics.jax.org/homology.shtml>). Specifically we used http://www.informatics.jax.org/downloads/reports/HOM_MouseHumanSequence.rpt (downloaded April 26th, 2021) for the analyses comparing T/NK compositions of human and mouse scRNAseq data, analyses comparing epithelial program similarity in human and mouse scRNAseq data, analyses involving the annotation of human scRNA-seq data with mouse regions, and analyses comparing the cell type and epithelial program correlations cross samples between human and mouse scRNA-seq data, and http://www.informatics.jax.org/downloads/reports/HOM_AllOrganism.rpt (downloaded on August 8th, 2022) for the analyses involving the scoring of mouse regions in mouse and human scRNA-seq data and leading to the CMS classification.

GO term enrichment analysis

We used TACCO's functions "setup_goa_analysis" and "run_goa_analysis" to perform GO terms enrichment. As "gene_info_file" we used https://ftp.ncbi.nih.gov/gene/DATA/GENE_INFO/Mammalia/Mus_musculus.gene_info.gz, as "GO_obo_file" <http://purl.obolibrary.org/obo/go/go-basic.obo>, and as "gene2GO_file" <https://ftp.ncbi.nih.gov/gene/DATA/gene2go.gz> (all downloaded on August 10th, 2022).

Comparison between mouse and human programs

To compare mouse and human epithelial expression programs², genes were mapped to mouse homologs using MGI homology information [subsection "Mapping of mouse and human orthologs"]. Mouse and human programs were then characterized by a single vector of mean expression per program in mouse gene space. Specifically, both mouse and human programs were defined such that their weighted sum approximates the expression profiles of the cells without any transformations. Programs and weights were normalized to sum to 1. To reduce batch-effects (including species-specific ones), a background expression profile was defined for each species dataset as the pseudo-bulk epithelial expression profile in the respective scRNAs-seq data. Program and background profiles were normalized to 10,000 counts and the log ratio of the normalized program and background expression vectors was used to define a vector for each species. Pearson correlation coefficients were calculated for each pair of program vectors (mouse vs. human).

Human expression program associations across mouse scRNA-seq

All sets of programs that were previously used to define human CRC tissue hubs² (epithelial, T/NK cells and myeloid cells) were mapped to mouse genes with MGI homology information [subsection “Mapping of mouse and human orthologs”] and then to mouse single cell data with TACCO, using TACCOs platform normalization to account for batch effects. The “annotate” function in TACCO was used with OT as the core annotation method, on the comparable subsets of cells from mouse and human single-cell datasets (*e.g.*, myeloid cells from mouse and human), with basic platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter lambda of 0.1, and 4 iterations of bisectioning with a divisor of 3, and flat annotation prior distribution. The resulting probabilistic per-cell program annotations were aggregated to get probabilistic per-sample program annotations for all dysplastic mouse samples and CLR-transformed. For each pair of programs, the Pearson correlation coefficient was calculated on these transformed values.

Annotating human scRNA-seq data with mouse-derived region information

Human scRNA-seq profiles² were mapped to mouse gene space using MGI homology information [subsection “Mapping of mouse and human orthologs”]. Working in the same expression space, the “annotate” function in TACCO with OT as core annotation method was used on the full human scRNA-seq and mouse Slide-seq dataset with basic platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter lambda of 0.1, and 7 iterations of bisectioning with a divisor of 3, and 10-fold sub-clustering of the region annotations. The region transfer is done separately per compartment, with the Slide-seq compartment split as described above and the human scRNA-seq data split using the cell type annotation of the data. For validation, mapping was also performed with mouse scRNA-seq data, as well as mapping the region information from the mouse pucks back to themselves.

To test for enrichments of region annotations across disease state, region composition was aggregated to sample-level (for Slide-seq to 4-way split pucks), CLR-transformed, and enrichment was calculated using a two-sided Welch’s t-test. This was done for region annotation on human and mouse scRNA-seq data, and on the original and mapped region annotation on the mouse Slide-seq data.

Cell-type associations across samples

To compare associations of cell types across samples in human and mouse scRNA-seq the “clMidwayPr” cell type annotation in the human data² was aggregated to the same level as mouse cell type annotation, and then aggregated per sample and CLR-transformed. Pearson correlation coefficients were calculated for every cell type pair for different subsets of samples: all samples, normal samples, dysplastic, and for human MMRd/MMRp samples.

Epithelial program associations across human samples

To determine epithelial program associations across human samples, TACCO’s “annotate” function was used to annotate human epithelial scRNA-seq (after mapping to mouse orthologs using MGI homology information [subsection “Mapping of mouse and human orthologs”]) with mouse epithelial programs from mouse scRNA-seq data using OT as core method, basic platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter lambda of 0.1, and 4 iterations of bisectioning with a divisor of 3. The remaining steps were performed as for cell-type association (subsection “Cell-type associations across samples”).

Scoring epithelial mouse regions in mouse and human epithelial pseudo-bulk data

The published processed and filtered count matrices were used (where available) or instead raw count matrices for single cell/nucleus RNA seq data from Pelka² (GEO accession number GSE178341; downloaded on August 8th, 2022), Chen²⁸ (Synapse IDs syn27056096, syn27056097, syn27056098, syn27056099; downloaded on August 5th, 2022), Khaliq⁶⁹ (GEO accession number GSE200997; downloaded on August 5th, 2022), Becker²⁹ (GEO accession number GSE201348; downloaded on August 5th, 2022), Zheng⁶⁸ (GEO accession number GSE161277; downloaded on August 5th, 2022; excluding ‘blood’ samples), Che⁶⁷ (GEO accession number GSE178318; downloaded on August 5th, 2022; only ‘CRC’ and ‘LM’ samples) and Joanito⁷⁰ (Synapse IDs syn26844072, syn26844073, syn26844078, syn26844087, syn26844111; downloaded on August 22nd, 2022; excluding the ‘LymphNode’ sample).

To subset the human single cell data to epithelial cells, the epithelial annotation was used where readily available², ²⁸. For the remaining datasets²⁹, ⁶⁷–⁷⁰, TACCOs `tc.tl.annotate` function was used with default parameters to transfer the ‘cl295v11SubShort’ annotation from Pelka², from which a compositional compartment annotation was constructed, and then a cell was assigned to the epithelial compartment if it had more than 95% epithelial fraction.

To correct for batch effects between the different data sources, first batches were defined by species times protocol: ‘mouse-10x3p’, ‘mouse-SlideSeq’, ‘human-10x3p’², ⁶⁷, ⁶⁸, ⁷⁰, ‘human-10x5p’⁶⁹, ⁷⁰, ‘human-inDrop’²⁸, and ‘human-snRNA’²⁹. Then TACCO’s “`tc.pp.normalize_platform`” function was used to determine per gene batch normalization factors using only the normal samples of one data source per batch (choosing the normal samples from Zheng for ‘human-10x3p’ and the normal 5’ samples from Joanito for ‘human-10x5p’). The resulting factors are then used to rescale the sample-by-gene count matrices for the full dataset per batch, i.e. including non-normal samples. The normalization factors are calculated with respect to an (arbitrarily chosen) normal reference dataset⁶⁸.

The epithelial mouse region score was defined as the mean of the CLR-transformed expression values in the pseudo-bulk expression profile of the epithelial part of a dataset using the top 200 differentially expressed genes between all regions by a one-sided Fisher’s exact test.

To account for species-specific biases (in-set vs. out-of-set prediction: the DEGs are calculated in mouse), the scores per region across samples were zero-centered and scaled to unit variance across all samples (including normal and non-normal samples and all batches) per species. A Principal Components Analysis (PCA) of the region scores across all species, batches and samples was conducted and the values for the first PC were compared between different conditions using a two-sided Mann-Whitney U test with Benjamini-Hochberg FDR.

Assessing the relationship between mouse regions and CMS tumor classification

We used the package ‘CMSclassifier’ (<https://github.com/Sage-Bionetworks/CMSclassifier>) referred to in Ref.⁷¹ to classify human pseudo-bulk CRC profiles from all samples which were not normal or unaffected from the human studies above into CMS classes. We determine the enrichment (Benjamini-Hochberg FDR, two-sided Welch’s t test) of the same mouse region scores in the CMS classes.

Assessing the relation between mouse regions and clinical endpoints in human bulk RNA-seq

Published RNA-seq data from the COAD and READ cohorts of TCGA PanCancerAtlas⁹⁵ were used. Mouse region scores were defined as the mean of the log1p-transformed, zero-centered and scaled expression values in the bulk expression profile using the top 200 differentially expressed genes between the malignant mouse regions (6, 8, 11) by a one-sided Fisher's Exact test (comparing each of the three regions to the other two). Scores were stratified into quartiles. PFI and OS was compared between patients with tumors whose scores were in the lowest and highest quartiles using the Logrank test as implemented in the lifelines package⁹⁶, followed by Benjamini-Hochberg FDR.

Compositional enrichment analyses

Enrichments on compositional data (cell type compositions, etc.) were evaluated with a two-sided Welch's t test on sample level using CLR-transformed compositions followed by Benjamini-Hochberg FDR. For the enrichment of tdTomato, counts and ALR-transformation were used instead with all non-tdTomato counts used as reference compartment. Enrichment analyses were performed using TACCO's "enrichments" function.

Supplementary figures




Figure S1.

Marker genes, cell states and cell types in the healthy and dysplastic mouse colon atlas.

A. Cell-type expression signatures. Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows) in cells (columns) from each cell type. **B,C.** Distinct condition states and compartments. 2D embedding of all single cell profiles (dots) colored by either condition state (B) or compartment (C). **D.** Changes in cell composition in dysplastic lesions. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each cell subset (rows) between samples from different conditions (columns). **E.** Changes in cell composition between healthy and dysplastic tissue. Proportion of cells (y axis) from each cell subset (x axis) out of all cells in each sample (x axis). **F-H.** Multiplex *in situ* RNA profiles (Cartana) reproduce scRNA-seq and Slide-Seq findings. F. Proportion of RNA molecules (y axis) attributed to each cell type in each sample (x axis) in two Cartana experiments, with different gene panels. G. Change in cell proportions in AV vs. normal samples (difference of CLR-transformed cell type fractions, x and y axes) for each cell type (dots, color) based on scRNA-seq, Slide-Seq or Cartana (axis labels). Error bars: bootstrapped standard error of the mean (normal n=8, 16, 12, 16 and AV n=11, 24, 12, 12 for scRNA-seq, Slide-seq, Cartana V1, Cartana V2). H. Change in marker gene expression (dots) in AV vs. normal samples (difference of CLR-transformed gene fractions, x and y axes) based on scRNA-seq, Slide-Seq or Cartana (axis labels), for genes measured by all 3 methods. Genes with the 8 maximal and minimal expression ranks across methods are labeled. Error bars: bootstrapped standard error of the mean (normal n=8, 16, 12, 16 and AV n=11, 24, 12, 12 for scRNA-seq, Slide-seq, Cartana V1, Cartana V2).




Figure S2.

Compositional and cell intrinsic changes in stromal and immune cells.

A-C. Immune cell subsets and composition. A. 2D embedding of immune cell profiles colored by clusters (as in B). B. Proportion of cells out of all immune cells (y axis) of each immune cell subset in each sample (x axis). C. Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows) in cells (columns) from each immune cell subset (color bar on top). **D-F.** Monocyte and macrophage cell subsets and composition. D. 2D embedding of monocyte and macrophage cell profiles colored by clusters (as in E). E. Proportion of cells out of all monocytes and macrophages (y axis) of each monocyte and macrophage cell subset in each sample (x axis). F. Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows) in cells (columns) from each monocyte and macrophage cell subset (color bar on top). **G-I.** T/NK cell subsets and composition. G. 2D embedding of T/NK cell profiles colored by clusters (as in H). H. Proportion of cells out of all T/NK cells (y axis) of each T/NK cell subset in each sample (x axis). I. Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows) in cells (columns) from each T/NK cell subset (color bar on top). **J.** Enrichment of IL17+ gdT cells (TNK05) and depletion of CD8+ gdT cells (TNK01) in dysplastic lesions. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each T/NK cell subset (rows) between samples from different conditions (columns). **K.** T/NK cell compositions is similar in human and mouse. Left: 2D embedding of T/NK cell composition profiles of human and mouse samples colored by sample type (legend) (**STAR Methods**). Right: Similarity of T/NK cell composition (enrichment z-scores) in the 2D embedding between each set of samples (rows, columns). **L-N.** Stromal cell subsets and composition. L. 2D embedding of stromal cell profiles colored by clusters (legend). M. Proportion of cells out of all stromal cells (y axis) of each stromal cell subset in each sample (x axis). N. Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows) in cells (columns) from each stromal cell subset (color bar on top). **O.** Enrichment of vascular endothelial cells (Endo01) and depletion of myofibroblasts (Fibro02) in dysplastic lesions. Significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each stromal cell subset (rows) between samples from different conditions (columns). **P.** Increased VEGFA expression in monocyte-macrophage populations with dysplasia. Distribution of expression (y axis, log1p(counts)) of *Vegfa* in monocytes and macrophages from different conditions (x axis).




Figure S3.

Changes in T/NK and epithelial cell differentiation in dysplastic lesions.

A. T/NK cell differentiation. 2D embedding of T/NK cell profiles from normal tissues (left) and dysplastic lesions (right), colored by cell subset. Streamlines: averaged and projected RNA velocities. **B,C.** Proliferating stem-like cells give rise to an expanded stem-like compartment and differentiated-like tumor cells in dysplastic lesions. 2D embedding of epithelial cell profiles from normal tissues (B, left and C, top) and dysplastic lesions (B, right, and C, bottom), colored by cell subset (B), or by expression of epithelial programs (C). Streamlines: averaged and projected RNA velocities. Outlined dots (B): Proliferative cells (cells with more than 50% program weight in the proliferation programs #3 and #11).




Figure S4.

Changes in cell composition and expression programs usage in dysplastic epithelium.

A-D. Changes in epithelial cell composition in dysplastic tissues. **A.** 2D embedding of all single cell epithelial profiles showing only the profiles (dots) of cells from each condition state, subsampled to equal numbers of cells per condition state, colored by cluster. **B.** PCA captures differentiation status and lineage of epithelial clusters. First (PC1, x axis) and second (PC2, y axis) principal components of the expression of epithelial clusters. **C.** Mean expression (dot color) and fraction of expressing cells (dot size) of epithelial cell marker genes (columns) in each cluster (rows). **D.** Significance (FDR, color bar, two-sided Welch's t test on ALR-transformed compositions with all non-tdTomato counts used as reference compartment) of enrichment (red) or depletion (blue) of tdTomato expression in cells from AKPV samples between every pair of epithelial clusters. **E.** Dysplastic secretory like (Epi05) and immune cells express tumor-related genes. Distribution of expression (y axis, log1p(counts)) of different marker genes in cells from each epithelial/immune cell cluster (x axis). **F-L.** Epithelial gene programs. 2D embedding of all epithelial cells colored by the weight of each program (color bar) and the expression of selected program genes (color bar). **M.** Epithelial program characteristics of normal and dysplastic colon. Heatmap significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each epithelial program (rows) between normal vs. dysplastic tissues (columns).




Figure S5.

Conservation of cellular composition and expression programs between mouse and human scRNA-seq data.

A,B. Conservation of general and subtype-specific human and mouse epithelial expression programs. A. Pearson correlation coefficients (color) between program-specific expression profiles (**Methods**) of human (all samples, left, MMRd, middle and MMRp, right)) (rows) and mouse programs (columns). B. Mouse epithelial program enrichments in mouse and human tumor and normal samples. Significance (FDR, color bar, two-sided Welch t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of mouse epithelial program expression (rows) in sample classes from human or mouse (columns).

C,D. Human-mouse conservation of cell type and program associations. Pearson correlation coefficients (color) of the CLR-transformed cell type (C) or epithelial program (D) compositions across samples in mouse (left) or human (right) single cell data. In (D), data are hierarchically clustered for the “not normal” mouse case (AV and AKPV) and this ordering is applied to all other panels.




Figure S6.

Spatial distributions of cells and programs across regions.

A-C. Slide-seq quality controls. Slide-seq pucks of AV (top) and normal (bottom) mouse colon colored by number of UMIs (A) or of genes (B) per bead. (x and y axis: spatial coordinates (μm)). **C.** H&E staining of sections adjacent to those used for Slide-seq. **D.** Selected marker gene expression. Slide-seq pucks from a normal (top) and AV (bottom) sample, colored by marker gene detected per bead. **E, F.** Spatial mapping of cell types and programs yields comparable composition to scRNA-seq. Distribution of the proportion (y axis) of contributions to each cell type (D) or program (E; based on fractional annotations) in cells (for scRNA-seq; "sc") or beads (for Slide-seq; "spatial"; based on fractional annotations) in samples from normal (N) or AV (AV) tissue. **G, H.** Distinct cell types and programs associated with AV and normal colon. Significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of cell types (F, rows) or epithelial programs (G, rows) in normal (N) or AV (AV) tissues based on Slide-seq ("spatial") data or scRNA-seq ("sc").




Figure S7.

Distinct cellular layout in normal and AV tissues.

A,B. Neighborhood analysis is robust to the distance and number of randomizations. Neighborhood enrichment (z-scores, color) vs. a background of spatially random annotation assignments for each pair of cell annotations (rows, columns) in normal (top) and AV (bottom) samples at varying distance (a, left: $\leq 40 \mu\text{m}$; right: $\leq 60 \mu\text{m}$) or number of permutations (B, left: 5, right: 50). **C,D.** Reproducible spatial arrangement of epithelial cells in normal colon in Slide-seq and Cartana V2. Co-occurrence (y axis) of normal epithelial cell types (color) at different distances (x axis) from different central normal epithelial cell type (annotated on top) in Slide-Seq (C) and Cartana (D) data. **E.** Decreased spatial order of cell types in AV vs. normal tissue. Distribution of enrichment z-values (x axis) of cluster-cluster interactions (as shown in **Figure 3B,C**) in AV lesions (orange) and normal colon (blue). **F.** Epithelial program neighborships in AV tissue. Short-range ($\leq 20 \mu\text{m}$) neighborhood enrichment z-scores (color) vs. a background of spatially random annotation assignments for each pair of epithelial program annotations (rows, columns) in AV lesions. **G,H.** Reproducible spatial arrangement of dysplasia-associated immune cells relative to endothelial cells in Slide-seq and Cartana V2. Co-occurrence (y axis) of dysplasia-associated monocyte or granulocyte cells (color) with endothelial cells at different distances (x axis) in Slide-Seq (G) and Cartana (H) data.




Figure S8.

Cellular neighborhoods.

A. Distinct characteristic region distributions in healthy and AV tissue. Proportion of UMIs assigned to each region (y axis) in each Slide-seq puck (x axis). **B.** β -catenin expression in AV lesions. Representative images (i) of β -catenin IHC on sections of AV lesions showing either nuclear (*) or cytoplasmic (**) staining pattern (x10 magnification), along with high magnification (x20, ii) of the region highlighted in (i), and H&E staining (iii) of a serial section showing a region proximal to the region in (i). (M-muscularis, L-lumen, dashed line: border between areas of nuclear and cytoplasmic staining of β -catenin). **C,D.** Spatial relations between the regions. Proportion of beads (y axis) of each region category (color code) at different distances (x axis) from region 2 (C, muscularis), 6 (D, left), 8 (D, middle) or 11 (D, right). **E,F.** Fraction of expressing cells (dot size) and mean expression per celltype (E, dot color) or per region (F, dot color) of marker genes (columns) across cell types in region 6 (E, rows) or regions 6, 8 and 11 (F, rows). **G.** Cell type neighborhood enrichment z-scores (color) vs. a background of spatially random annotation assignment for each pair of cell type annotations (rows, columns) in malignant-like regions 6, 8 and 11 within AV lesions. **H.** Spatial organization of endothelial cells and pericytes based on Cartana multiplex *in situ* RNA analysis. Co-occurrence (y axis) of Cdh5 (top) or Pecam1 (middle) -expressing (endothelial) cells or Pdgfrb (bottom) -expressing cells (pericytes) at different distances (x axis) from each central cell type (color code), based on expression of a specific gene (color bar). **I.** Cell-cell interaction pathways enriched in different regions. Enrichment (FDR) in regions 2 (muscularis, left), 6 (malignant-like, inflammation, middle) or 11 (malignant-like, EMT, right) of corresponding "sender" (x axis) and "receiver" (y axis) (aggregated over ligand-receptor pairs) pathways (dots). Red/blue: pathways significantly enriched/depleted in region (light red/blue: only enriched/depleted for either sender or receiver). The top 5 enriched pathways (in each direction) are labeled.





Figure S9.

Transfer of mouse spatial region expression profiles to human patient data.


A. Expression profiles characterizing mouse regions in human and mouse samples. Significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of mouse region-associated epithelial (left), immune (middle) or stromal profiles (right) in pucks from normal colon and dysplastic lesions, in the same pucks but after mapping the region annotation to itself (consistency check), in mouse single cell data after mapping region annotation from mouse pucks, and in human single cell data after mapping region annotation from mouse pucks. **B.** PC1 values (y axis; box plots show mean, quartiles, and whiskers for the full data distribution except for outliers outside 1.5 times the interquartile range (IQR)) in a PCA of region scores of mouse and human samples (x axis) sorted by malignant status and colored by status (top, legend) or study (bottom, legend). **C.** Spatial regions profiles associated with different CMS classes. Significance (Benjamini-Hochberg FDR, color bar, two-sided Welch's t test) of enrichment (red) or depletion (blue) of each region profile (rows) in pseudo bulk profiles of human tumor samples classified in each CMS class (columns). **D,E.** Expression of malignant-like regions 6 and 11 in tumors is associated with PFI and OS in MMRd and MMRp patients. Kaplan-Meier PFI (f, MMRd $n = 189^{95}$, MMRp $n = 447^{95}$) or OS (g, MMRd $n = 189^{95}$, MMRp $n = 447^{95}$) analysis of human bulk RNA-seq cohort stratified by malignant-like region profile scores.



Supplementary item 1.

Correspondence between scRNA-seq, Slide-seq, and Cartana data


A,B. Agreement in cell type proportions. Inferred cell type proportions (y and x axis; CLR-transformed cell type fractions) for each cell type (dot color) in normal (A) and AV (B) samples in scRNA-seq, Slide-Seq and Cartana data (axis labels). Pearson's r : top left corners. Error bars: bootstrapped standard error of the mean ((A) $n=8, 16, 12, 16$ and (B) $n=11, 24, 12, 12$ for scRNA-seq, Slide-seq, Cartana V1, Cartana V2). **C,D.** Agreement in gene expression levels. Measured expression (y and x axis; CLR-transformed gene fractions) for each gene (dot) measured by all three methods in normal (A) and AV (B) samples in scRNA-seq, Slide-Seq and Cartana data (axis labels). Pearson's r : top left corners. Error bars: bootstrapped standard error of the mean ((C) $n=8, 16, 12, 16$ and (D) $n=11, 24, 12, 12$ for scRNA-seq, Slide-seq, Cartana V1, Cartana V2). Genes with the 8 maximal and minimal expression ranks across methods are labeled.



Supplementary item 2.

Dysplasia-associated cells in the tumor microenvironment.


A,B. Representative images of multiplex RNA analysis in normal colon (left) and AV lesions (right) colored by cell type assignment (A, the same is [Figure 1E](#)) and expression of marker genes (B) for Mono02 (Arg1, Hlfpda, Nos2, Cd274, Vegfa, Trem1; cyan), Mono03 (Osm, Ifi204 and Thbs1; red), granulocytes (S100a8; yellow), as well as Sell (green), Ccr2 (magenta), Ptprc (blue) and Cd14 (gray).



Supplementary item 3.

TLS-like structures in AV lesions by Cartana multiplex *in situ* RNA profiles.

A. Proportion of RNA molecules (y axis) attributed to each cell type (top) or gene (bottom) in TLS-like regions (x axis) in Cartana V2 multiplex *in situ* RNA experiment. Genes which constitute at least 3% of all measured transcripts in any of the identified follicular structures are noted (bottom). **B.** Representative images from TLS-like regions showing the distribution of cell type (top row) and of cell type markers (bottom four rows; Cd14, Cd68 (monocytes, magenta), Cd3d, Cd3e, Cd3g (T cells, yellow), and Cd79, Jchain, Igkc (B cells, dark cyan)).



Supplementary item 4.

Ligand-receptor analysis.

A,B. Cell-cell interaction pathways enriched in normal and AV tissue and regions. Significance (FDR, color bar, two-sided Mann-Whitney U test) of enrichment (red) or depletion (blue) of expression of “sender” (left) and “receiver” (right) cell-cell communication pathways (rows) (aggregated from ligand-receptor pairs) in normal vs. AV samples (A, columns), or regions (B, columns). **C,D.** Split power analysis for cell-cell communication pathway analysis. Significance of enrichment (log₁₀(FDR), y-axis) for receiver (solid lines) and sender (dashed line) for each of the top 5 enriched communication pathways (color) in each iteration for the spatial split of the Slide-Seq pucks into spatially disconnected and separated patches (x axis), for either normal vs. AV tissue enrichment (C) or for specific regions (D).

Acknowledgements

We thank N. Friedman, I. Benhar, N. Habib, M. Biton, K. Geiger-Schuller, J.C. Hütter, B. Dumitrascu, E. Baker and A. Greenwald for helpful discussions. We thank C. McCabe, O. Kuksenko and I. Barrera for technical assistance. We thank P. Yadollahpour, E. Dhaval, G. Smith-Rosario, S. Vickovic, D. Schapiro, S. Farhi, D. Abbondanza, A. Segerstolpe and T. Biancalani for their important contribution to this project. We thank L. Gaffney and A. Hupalowska for help with figure preparation. S.M. was supported by a DFG research fellowship (MA 9108/1-1), J.K. was supported by a HFSP long term fellowship (LT000452/2019-L), A.R. was a Howard Hughes Medical Institute (HHMI) Investigator when conducting this work. Work was supported by the Klarman Cell Observatory, a CEGS grant (5RM1HG006193-09) from the NHGRI, the NIH/NIAID (grants 1U24 CA180922, 1U19 MH114821, 1RC2 DK114784), the MIT Ludwig Center, the Manton Family Foundation, and HHMI (A.R.); Azrieli Foundation Early Career Faculty Fellowship, and an ISF Research Grant (1079/21) (M.N.), the Center for Interdisciplinary Data Science Research at the Hebrew University of Jerusalem (N.M. and M.N.), the Israeli Council for Higher Education Ph.D. fellowship (N.M.), SU2C Peggy Prescott Early Career Scientist Award PA-6146, SU2C Phillip A. Sharp Award SU2C-AACR-PS-32 and NIH/NCI R00CA259511 (K.P.), NIH/NCI R01 CA208756; Arthur, Sandra, and Sarah Irving Fund for Gastrointestinal Immuno-Oncology (N.H.), NIH/NCI R01CA257523, MIT Stem Cell Initiative (Foundation MIT) (O.Y.), and NIH R37CA259363, R21CA256414, R21DK125911, R41EB032693, R01CA254108, R01CA256530, and R01CA244359; DOD W81XWH-20-1-0203; and a Duke-NC State Translational Research Grant (J.R.). The authors gratefully acknowledge LMU Klinikum for providing computing resources on their Clinical Open Research Engine (CORE) and the Bioinformatic Core Facility of the Biomedical Center Munich for providing computing resources on their HPC system.

Additional information

Author contributions

I.A-D. and A.R. conceived the study and designed experiments. I.A-D. J.R., S.Y., E.M., T.D., L.C. and D.D. performed experiments, with guidance from O.Y. and A.R.. S.M., J.K., and N.M. developed computational approaches and analyzed data with I.A-D., M.H., E.M. and R.S, and with guidance from M.N. and A.R.. M.H., E.M., J.C., K.P., A.M. and G.M.B. generated and analyzed human data, with guidance from N.H., I.T., N.H., F.C., O.Y., O.R-R, and A.R.. A.N. and H.X. generated and processed the multiplex *in-situ* RNA data, with guidance from J.L. and M.R.. M.N. and A.R. provided supervision and acquired funding. I.A-D., S.M., M.N. and A.R. wrote the manuscript with input from all authors.

Code availability

The analysis code is available on GitHub (<https://github.com/simonwm/mouseCRC> )

Data availability

All data generated in this project is deposited on the Single Cell Portal and is available under the accession SCP1891 (https://singlecell.broadinstitute.org/single_cell/study/SCP1891 ). The raw scRNA-seq and Slide-seq data is also deposited on GEO under the GSE260801 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE260801> ). The public human scRNA-seq datasets can be found on GEO under accession numbers GSE178341, GSE200997, GSE201348, GSE161277, GSE178318 and on Synapse under the IDs syn27056096, syn27056097, syn27056098, syn27056099, syn26844072, syn26844073, syn26844078, syn26844087, syn26844111.

Additional files

Supplementary table 1 [↗](#)

Supplementary table 2 [↗](#)

Supplementary table 3 [↗](#)

References

1. McAllister S.S., Weinberg R.A (2014) **The tumour-induced systemic environment as a critical regulator of cancer progression and metastasis** *Nat. Cell Biol* **16**:717–727 [Google Scholar](#)
2. Pelka K., Hofree M., Chen J.H., Sarkizova S., Pirl J.D., Jorgji V., Bejnood A., Dionne D., Ge W.H., Xu K.H., et al. (2021) **Spatially organized multicellular immune hubs in human colorectal cancer** *Cell* **184**:4734–4752 [Google Scholar](#)
3. Wagner J., Rapsomaniki M.A., Chevrier S., Anzeneder T., Langwieder C., Dykgers A., Rees M., Ramaswamy A., Muenst S., Soysal S.D., et al. (2019) **A Single-Cell Atlas of the Tumor and Immune Ecosystem of Human Breast Cancer** *Cell* **177**:1330–1345 [Google Scholar](#)
4. Hunter M.V., Moncada R., Weiss J.M., Yanai I., White R.M (2021) **Spatially resolved transcriptomics reveals the architecture of the tumor-microenvironment interface** *Nat. Commun* **12**:1–16 [Google Scholar](#)
5. Keren L., Bosse M., Marquez D., Angoshtari R., Jain S., Varma S., Yang S.-R., Kurian A., Van Valen D., West R., et al. (2018) **A Structured Tumor-Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging** *Cell* **174**:1373–1387 [Google Scholar](#)
6. Schürch C.M., Bhate S.S., Barlow G.L., Phillips D.J., Noti L., Zlobec I., Chu P., Black S., Demeter J., McIlwain D.R., et al. (2020) **Coordinated cellular neighborhoods orchestrate antitumoral immunity at the colorectal cancer invasive front** *Cell* **182**:1341–1359 [Google Scholar](#)
7. Jackson H.W., Fischer J.R., Zanutelli V.R.T., Ali H.R., Mechera R., Soysal S.D., Moch H., Muenst S., Varga Z., Weber W.P., et al. (2020) **The single-cell pathology landscape of breast cancer** *Nature* **578**:615–620 [Google Scholar](#)
8. Jerby-Arnon L., Shah P., Cuoco M.S., Rodman C., Su M.-J., Melms J.C., Leeson R., Kanodia A., Mei S., Lin J.-R., et al. (2018) **A Cancer Cell Program Promotes T Cell Exclusion and Resistance to Checkpoint Blockade** *Cell* **175**:984–997 [Google Scholar](#)
9. Grünwald B.T., Devisme A., Andrieux G., Vyas F., Aliar K., McCloskey C.W., Macklin A., Jang G.H., Denroche R., Romero J.M., et al. (2021) **Spatially confined sub-tumor microenvironments in pancreatic cancer** *Cell* **184**:5577–5592 [Google Scholar](#)
10. Goltsev Y., Samusik N., Kennedy-Darling J., Bhate S., Hale M., Vazquez G., Black S., Nolan G.P (2018) **Deep Profiling of Mouse Splenic Architecture with CODEX Multiplexed Imaging** *Cell* **174**:968–981 [Google Scholar](#)
11. Angelo M., Bendall S.C., Finck R., Hale M.B., Hitzman C., Borowsky A.D., Levenson R.M., Lowe J.B., Liu S.D., Zhao S., et al. (2014) **Multiplexed ion beam imaging of human breast tumors** *Nat. Med* **20**:436–442 [Google Scholar](#)
12. Giesen C., Wang H.A.O., Schapiro D., Zivanovic N., Jacobs A., Hattendorf B., Schüffler P.J., Grolimund D., Buhmann J.M., Brandt S., et al. (2014) **Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry** *Nat. Methods* **11**:417–422 [Google Scholar](#)

13. Ståhl P.L., Salmén F., Vickovic S., Lundmark A., Navarro J.F., Magnusson J., Giacomello S., Asp M., Westholm J.O., Huss M., et al. (2016) **Visualization and analysis of gene expression in tissue sections by spatial transcriptomics** *Science* **353**:78–82 [Google Scholar](#)
14. Rodriques S.G., Stickels R.R., Goeva A., Martin C.A., Murray E., Vanderburg C.R., Welch J., Chen L.M., Chen F., Macosko E.Z (2019) **Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution** *Science* **363**:1463–1467 [Google Scholar](#)
15. Stickels R.R., Murray E., Kumar P., Li J., Marshall J.L., Di Bella D.J., Arlotta P., Macosko E.Z., Chen F. (2021) **Highly sensitive spatial transcriptomics at near-cellular resolution with Slide-seqV2** *Nat. Biotechnol* **39**:313–319 [Google Scholar](#)
16. Marx V (2021) **Publisher Correction: Method of the Year: spatially resolved transcriptomics** *Nat. Methods* **18**:219 [Google Scholar](#)
17. Waylen L.N., Nim H.T., Martelotto L.G., Ramialison M (2020) **From whole-mount to single-cell spatial assessment of gene expression in 3D** *Commun Biol* **3**:602 [Google Scholar](#)
18. Palla G., Fischer D.S., Regev A., Theis F.J (2022) **Spatial components of molecular tissue biology** *Nat. Biotechnol* <https://doi.org/10.1038/s41587-021-01182-1> | [Google Scholar](#)
19. Cable D.M., Murray E., Zou L.S., Goeva A., Macosko E.Z., Chen F., Irizarry R.A (2021) **Robust decomposition of cell type mixtures in spatial transcriptomics** *Nat. Biotechnol* :1–10 [Google Scholar](#)
20. Kleshchevnikov V., Shmatko A., Dann E., Aivazidis A., King H.W., Li T., Elmentaite R., Lomakin A., Kedlian V., Gayoso A., et al. (2022) **Cell2location maps fine-grained cell types in spatial transcriptomics** *Nat. Biotechnol* **40**:661–671 [Google Scholar](#)
21. Mages S., Moriel N., Avraham-Davidi I., Murray E., Watter J., Chen F., Rozenblatt-Rosen O., Klughammer J., Regev A., Nitzan M (2023) **TACCO unifies annotation transfer and decomposition of cell identities for single-cell and spatial omics** *Nat. Biotechnol* :1–9 [Google Scholar](#)
22. Lopez R., Li B., Keren-Shaul H., Boyeau P., Kedmi M., Pilzer D., Jelinski A., Yofe I., David E., Wagner A., et al. (2022) **DestVI identifies continuums of cell types in spatial transcriptomics data** *Nat. Biotechnol* **40**:1360–1369 [Google Scholar](#)
23. Biancalani T., Scalia G., Buffoni L., Avasthi R., Lu Z., Sanger A., Tokcan N., Vanderburg C.R., Segerstolpe Å., Zhang M., et al. (2021) **Deep learning and alignment of spatially resolved single-cell transcriptomes with Tangram** *Nat. Methods* **18**:1352–1362 [Google Scholar](#)
24. Cancer Genome Atlas Network (2012) **Comprehensive molecular characterization of human colon and rectal cancer** *Nature* **487**:330–337 [Google Scholar](#)
25. Kwong L.N., Dove W.F (2009) **APC and its modifiers in colon cancer** *Adv. Exp. Med. Biol* **656**:85–106 [Google Scholar](#)
26. Fearon E.R (2011) **Molecular genetics of colorectal cancer** *Annu. Rev. Pathol* **6**:479–507 [Google Scholar](#)

27. Fearon E.R., Vogelstein B (1990) **A genetic model for colorectal tumorigenesis** *Cell* **61**:759–767 [Google Scholar](#)
28. Chen B., Scurrah C.R., McKinley E.T., Simmons A.J., Ramirez-Solano M.A., Zhu X., Markham N.O., Heiser C.N., Vega P.N., Rolong A., et al. (2021) **Differential pre-malignant programs and microenvironment chart distinct paths to malignancy in human colorectal polyps** *Cell* **184**:6262–6280 [Google Scholar](#)
29. Becker W.R., Nevins S.A., Chen D.C., Chiu R., Horning A.M., Guha T.K., Laquindanum R., Mills M., Chaib H., Ladabaum U., et al. (2022) **Single-cell analyses define a continuum of cell state and composition changes in the malignant transformation of polyps to colorectal cancer** *Nat. Genet* :1–11 [Google Scholar](#)
30. Roper J., Tammela T., Cetinbas N.M., Akkad A., Roghanian A., Rickelt S., Almqdadi M., Wu K., Oberli M.A., Sánchez-Rivera F.J., et al. (2017) **In vivo genome editing and organoid transplantation models of colorectal cancer and metastasis** *Nat. Biotechnol* **35**:569–576 [Google Scholar](#)
31. Roper J., Tammela T., Akkad A., Almqdadi M., Santos S.B., Jacks T., Yilmaz Ö.H (2018) **Colonoscopy-based colorectal cancer modeling in mice with CRISPR–Cas9 genome editing and organoid transplantation** *Nat. Protoc* **13**:217–234 [Google Scholar](#)
32. Golovko D., Kedrin D., Yilmaz Ö.H., Roper J (2015) **Colorectal cancer models for novel drug discovery** *Expert Opin. Drug Discov* **10**:1217–1229 [Google Scholar](#)
33. Gyllborg D., Langseth C.M., Qian X., Choi E., Salas S.M., Hilscher M.M., Lein E.S., Nilsson M (2020) **Hybridization-based in situ sequencing (HybISS) for spatially resolved transcriptomics in human and mouse brain tissue** *Nucleic Acids Res* **48**:e112 [Google Scholar](#)
34. Hagerling C., Gonzalez H., Salari K., Wang C.-Y., Lin C., Robles I., van Gogh M., Dejmek A., Jirstrom K., Werb Z. (2019) **Immune effector monocyte–neutrophil cooperation induced by the primary tumor prevents metastatic progression of breast cancer** *Proc. Natl. Acad. Sci. U. S. A* **116**:21704–21714 [Google Scholar](#)
35. Arenberg D.A., Keane M.P., DiGiovine B., Kunkel S.L., Strom S.R., Burdick M.D., Iannettoni M.D., Strieter R.M (2000) **Macrophage infiltration in human non-small-cell lung cancer: the role of CC chemokines** *Cancer Immunol. Immunother* **49**:63–70 [Google Scholar](#)
36. Smith P.M., Garrett W.S (2011) **The gut microbiota and mucosal T cells** *Front. Microbiol* **2**:111 [Google Scholar](#)
37. La Manno G., Soldatov R., Zeisel A., Braun E., Hochgerner H., Petukhov V., Lidschreiber K., Kastri M.E., Lönnerberg P., Furlan A., et al. (2018) **RNA velocity of single cells** *Nature* **560**:494–498 [Google Scholar](#)
38. Ziyad S., Iruela-Arispe M.L (2011) **Molecular mechanisms of tumor angiogenesis** *Genes Cancer* **2**:1085–1096 [Google Scholar](#)
39. Zeng Z.-S., Shu W.-P., Cohen A.M., Guillem J.G (2002) **Matrix metalloproteinase-7 expression in colorectal cancer liver metastases: evidence for involvement of MMP-7 activation in human cancer metastases** *Clin. Cancer Res* **8**:144–148 [Google Scholar](#)
40. Li D., Peng Z., Tang H., Wei P., Kong X., Yan D., Huang F., Li Q., Le X., Li Q., et al. (2011) **KLF4-mediated negative regulation of IFITM3 expression plays a critical role in colon cancer**

pathogenesis *Clin. Cancer Res* **17**:3558–3568 [Google Scholar](#)

41. Kitamura T., Kometani K., Hashida H., Matsunaga A., Miyoshi H., Hosogi H., Aoki M., Oshima M., Hattori M., Takabayashi A., et al. (2007) **SMAD4-deficient intestinal tumors recruit CCR1+ myeloid cells that promote invasion** *Nat. Genet* **39**:467–475 [Google Scholar](#)
42. Haber A.L., Biton M., Rogel N., Herbst R.H., Shekhar K., Smillie C., Burgin G., Delorey T.M., Howitt M.R., Katz Y., et al. (2017) **A single-cell survey of the small intestinal epithelium** *Nature* **551**:333–339 [Google Scholar](#)
43. Biton M., Haber A.L., Rogel N., Burgin G., Beyaz S., Schnell A., Ashenberg O., Su C.-W., Smillie C., Shekhar K., et al. (2018) **T Helper Cell Cytokines Modulate Intestinal Stem Cell Renewal and Differentiation** *Cell* **175**:1307–1320 [Google Scholar](#)
44. Smillie C.S., Biton M., Ordovas-Montanes J., Sullivan K.M., Burgin G., Graham D.B., Herbst R.H., Rogel N., Slyper M., Waldman J., et al. (2019) **Intra- and Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis** *Cell* **178**:714–730 [Google Scholar](#)
45. Welch J.D., Kozareva V., Ferreira A., Vanderburg C., Martin C., Macosko E.Z (2019) **Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity** *Cell* **177**:1873–1887 [Google Scholar](#)
46. van Neerven S.M., de Groot N.E., Nijman L.E., Scicluna B.P., van Driel M.S., Lecca M.C., Warmerdam D.O., Kakkar V., Moreno L.F., Vieira Braga F.A., et al. (2021) **Apc-mutant cells act as supercompetitors in intestinal tumour initiation** *Nature* **594**:436–441 [Google Scholar](#)
47. Flanagan D.J., Pentimikko N., Luopajarvi K., Willis N.J., Gilroy K., Raven A.P., McGarry L., Englund J.I., Webb A.T., Scharaw S., et al. (2021) **NOTUM from Apc-mutant cells biases clonal competition to initiate cancer** *Nature* **594**:430–435 [Google Scholar](#)
48. Clevers H (2006) **Wnt/beta-catenin signaling in development and disease** *Cell* **127**:469–480 [Google Scholar](#)
49. Folkman J. (2002) **Role of angiogenesis in tumor growth and metastasis. Semin Oncol** **29**:15–18 [Google Scholar](#)
50. Lasry A., Zinger A., Ben-Neriah Y (2016) **Inflammatory networks underlying colorectal cancer** *Nat. Immunol* **17**:230–240 [Google Scholar](#)
51. Fleming M., Ravula S., Tatishchev S.F., Wang H.L (2012) **Colorectal carcinoma: Pathologic aspects** *J. Gastrointest. Oncol* **3**:153–173 [Google Scholar](#)
52. Winkler J., Abisoye-Ogunniyan A., Metcalf K.J., Werb Z (2020) **Concepts of extracellular matrix remodelling in tumour progression and metastasis** *Nat. Commun* **11**:1–19 [Google Scholar](#)
53. Drev D., Harpain F., Beer A., Stift A., Gruber E.S., Klimpfinger M., Thalhammer S., Reti A., Kenner L., Bergmann M., et al. (2019) **Impact of Fibroblast-Derived SPARC on Invasiveness of Colorectal Cancer Cells** *Cancers* **11** <https://doi.org/10.3390/cancers11101421> | [Google Scholar](#)
54. Burden R.E., Gormley J.A., Jaquin T.J., Small D.M., Quinn D.J., Hegarty S.M., Ward C., Walker B., Johnston J.A., Olwill S.A., et al. (2009) **Antibody-mediated inhibition of cathepsin S blocks colorectal tumor invasion and angiogenesis** *Clin. Cancer Res* **15**:6042–6051 [Google Scholar](#)

55. Basu S., Cheriyaundath S., Gavert N., Brabletz T., Haase G., Ben-Ze'ev A (2019) **Increased expression of cathepsin D is required for L1-mediated colon cancer progression** *Oncotarget* **10**:5217–5228 [Google Scholar](#)
56. Spencer J., Sollid L.M (2016) **The human intestinal B-cell response** *Mucosal Immunol* **9**:1113–1124 [Google Scholar](#)
57. Sautès-Fridman C., Petitprez F., Calderaro J., Fridman W.H (2019) **Tertiary lymphoid structures in the era of cancer immunotherapy** *Nat. Rev. Cancer* **19**:307–325 [Google Scholar](#)
58. Mendez M.G., Kojima S.-I., Goldman R.D (2010) **Vimentin induces changes in cell shape, motility, and adhesion during the epithelial to mesenchymal transition** *FASEB J* **24**:1838–1851 [Google Scholar](#)
59. Lu M.-H., Huang C.-C., Pan M.-R., Chen H.-H., Hung W.-C (2012) **Prospero homeobox 1 promotes epithelial-mesenchymal transition in colon cancer cells by inhibiting E-cadherin via miR-9** *Clin. Cancer Res* **18**:6416–6425 [Google Scholar](#)
60. Oliemuller E., Newman R., Tsang S.M., Foo S., Muirhead G., Noor F., Haider S., Aurrekoetxea-Rodríguez I., Vivanco M. dM, Howard B.A. (2020) **SOX11 promotes epithelial/mesenchymal hybrid state and alters tropism of invasive breast cancer cells** *eLife* **9**:e58374 <https://doi.org/10.7554/eLife.58374> | [Google Scholar](#)
61. Marjanovic N.D., Hofree M., Chan J.E., Canner D., Wu K., Trakala M., Hartmann G.G., Smith O.C., Kim J.Y., Evans K.V., et al. (2020) **Emergence of a High-Plasticity Cell State during Lung Cancer Evolution** *Cancer Cell* **38**:229–246 [Google Scholar](#)
62. Puram S.V., Tirosh I., Parikh A.S., Patel A.P., Yizhak K., Gillespie S., Rodman C., Luo C.L., Mroz E.A., Emerick K.S., et al. (2017) **Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer** *Cell* **171**:1611–1624 [Google Scholar](#)
63. Schwab R.H.M., Amin N., Flanagan D.J., Johanson T.M., Phesse T.J., Vincan E (2018) **Wnt is necessary for mesenchymal to epithelial transition in colorectal cancer cells** *Dev. Dyn* **247**:521–530 [Google Scholar](#)
64. DiMeo T.A., Anderson K., Phadke P., Fan C., Perou C.M., Naber S., Kuperwasser C (2009) **A novel lung metastasis signature links Wnt signaling with cancer cell self-renewal and epithelial-mesenchymal transition in basal-like breast cancer** *Cancer Res* **69**:5364–5373 [Google Scholar](#)
65. Penninger J.M., Crabtree G.R (1999) **The Actin Cytoskeleton and Lymphocyte Activation** *Cell* **96**:9–12 [Google Scholar](#)
66. Cang Z., Zhao Y., Almet A.A., Stabell A., Ramos R., Plikus M.V., Atwood S.X., Nie Q (2023) **Screening cell-cell communication in spatial transcriptomics via collective optimal transport** *Nat. Methods* **20**:218–228 [Google Scholar](#)
67. Che L.-H., Liu J.-W., Huo J.-P., Luo R., Xu R.-M., He C., Li Y.-Q., Zhou A.-J., Huang P., Chen Y.-Y., et al. (2021) **A single-cell atlas of liver metastases of colorectal cancer reveals reprogramming of the tumor microenvironment in response to preoperative chemotherapy** *Cell Discovery* **7**:1–21 [Google Scholar](#)
68. Zheng X., Song J., Yu C., Zhou Z., Liu X., Yu J., Xu G., Yang J., He X., Bai X., et al. (2022) **Single-cell transcriptomic profiling unravels the adenoma-initiation role of protein tyrosine kinases**

during colorectal tumorigenesis *Signal Transduction and Targeted Therapy* **7**:1–14 [Google Scholar](#)

69. Khaliq A.M., Erdogan C., Kurt Z., Turgut S.S., Grunvald M.W., Rand T., Khare S., Borgia J.A., Hayden D.M., Pappas S.G., et al. (2022) **Correction: Refining colorectal cancer classification and clinical stratification through a single-cell atlas** *Genome Biol* **23**:156 [Google Scholar](#)
70. Joanito I., Wirapati P., Zhao N., Nawaz Z., Yeo G., Lee F., Eng C.L.P., Macalinalao D.C., Kahraman M., Srinivasan H., et al. (2022) **Single-cell and bulk transcriptome sequencing identifies two epithelial tumor cell states and refines the consensus molecular classification of colorectal cancer** *Nat. Genet* **54**:963–975 [Google Scholar](#)
71. Guinney J., Dienstmann R., Wang X., de Reyniès A., Schlicker A., Soneson C., Marisa L., Roepman P., Nyamundanda G., Angelino P., et al. (2015) **The consensus molecular subtypes of colorectal cancer** *Nat. Med* **21**:1350–1356 [Google Scholar](#)
72. West N.R., Hegazy A.N., Owens B.M.J., Bullers S.J., Linggi B., Buonocore S., Coccia M., Görtz D., This S., Stockenhuber K., et al. (2017) **Oncostatin M drives intestinal inflammation and predicts response to tumor necrosis factor-neutralizing therapy in patients with inflammatory bowel disease** *Nat. Med* **23**:579–589 [Google Scholar](#)
73. West N.R., Murray J.I., Watson P.H (2013) **Oncostatin-M promotes phenotypic changes associated with mesenchymal and stem cell-like differentiation in breast cancer** *Oncogene* **33**:1485–1494 [Google Scholar](#)
74. Smigiel J.M., Parameswaran N., Jackson M.W (2017) **Potent EMT and CSC phenotypes are induced by oncostatin-M in pancreatic cancer** *Mol. Cancer Res* **15**:478–488 [Google Scholar](#)
75. Hara T., Chanoch-Myers R., Mathewson N.D., Myskiw C., Atta L., Bussema L., Eichhorn S.W., Greenwald A.C., Kinker G.S., Rodman C., et al. (2021) **Interactions between cancer cells and immune cells drive transitions to mesenchymal-like states in glioblastoma** *Cancer Cell* **39**:779–792 [Google Scholar](#)
76. de Sousa e Melo F., Kurtova A.V., Harnoss J.M., Kljavin N., Hoeck J.D., Hung J., Anderson J.E., Storm E.E., Modrusan Z., Koeppen H., et al. (2017) **A distinct role for Lgr5+ stem cells in primary and metastatic colon cancer** *Nature* **543**:676–680 [Google Scholar](#)
77. Schepers A.G., Snippert H.J., Stange D.E., van den Born M., van Es J.H., van de Wetering M., Clevers H. (2012) **Lineage tracing reveals Lgr5+ stem cell activity in mouse intestinal adenomas** *Science* **337**:730–735 [Google Scholar](#)
78. André T., Shiu K.-K., Kim T.W., Jensen B.V., Jensen L.H., Punt C., Smith D., Garcia-Carbonero R., Benavides M., Gibbs P., et al. (2020) **Pembrolizumab in Microsatellite-Instability-High Advanced Colorectal Cancer** *N. Engl. J. Med* **383**:2207–2218 [Google Scholar](#)
79. Kuraguchi M., Wang X.-P., Bronson R.T., Rothenberg R., Ohene-Baah N.Y., Lund J.J., Kucherlapati M., Maas R.L., Kucherlapati R (2006) **Adenomatous polyposis coli (APC) is required for normal development of skin and thymus** *PLoS Genet* **2**:e146 [Google Scholar](#)
80. Johnson L., Mercer K., Greenbaum D., Bronson R.T., Crowley D., Tuveson D.A., Jacks T (2001) **Somatic activation of the K-ras oncogene causes early onset lung cancer in mice** *Nature* **410**:1111–1116 [Google Scholar](#)

81. Madisen L., Zwingman T.A., Sunkin S.M., Oh S.W., Zariwala H.A., Gu H., Ng L.L., Palmiter R.D., Hawrylycz M.J., Jones A.R., et al. (2009) **A robust and high-throughput Cre reporting and characterization system for the whole mouse brain** *Nat. Neurosci* **13**:133–140 [Google Scholar](#)
82. Marino S., Vooijs M., van Der Gulden H., Jonkers J., Berns A. (2000) **Induction of medulloblastomas in p53-null mutant mice by somatic inactivation of Rb in the external granular layer cells of the cerebellum** *Genes Dev* **14**:994–1004 [Google Scholar](#)
83. el Marjou F., Janssen K.-P., Chang B.H.-J., Li M., Hindie V., Chan L., Louvard D., Chambon P., Metzger D., Robine S. (2004) **Tissue-specific and inducible Cre-mediated recombination in the gut epithelium** *Genesis* **39**:186–193 [Google Scholar](#)
84. Stoeckius M., Zheng S., Houck-Loomis B., Hao S., Yeung B.Z., Mauck W.M., Smibert P., Satija R. (2018) **Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics** *Genome Biol* **19**:224 [Google Scholar](#)
85. Qian X., Harris K.D., Hauling T., Nicoloutsopoulos D., Muñoz-Manchado A.B., Skene N., Hjerling-Leffler J., Nilsson M. (2019) **Probabilistic cell typing enables fine mapping of closely related cell types in situ** *Nat. Methods* **17**:101–106 [Google Scholar](#)
86. Li B., Gould J., Yang Y., Sarkizova S., Tabaka M., Ashenberg O., Rosen Y., Slyper M., Kowalczyk M.S., Villani A.-C., et al. (2020) **Cumulus provides cloud-based data analysis for large-scale single-cell and single-nucleus RNA-seq** *Nat. Methods* **17**:793–798 [Google Scholar](#)
87. Wolf F.A., Angerer P., Theis F.J. (2018) **SCANPY: large-scale single-cell gene expression data analysis** *Genome Biol* **19**:15 [Google Scholar](#)
88. Wolock S.L., Lopez R., Klein A.M. (2019) **Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic Data** *Cell Syst* **8**:281–291 [Google Scholar](#)
89. Aran D., Looney A.P., Liu L., Wu E., Fong V., Hsu A., Chak S., Naikawadi R.P., Wolters P.J., Abate A.R., et al. (2019) **Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage** *Nat. Immunol* **20**:163–172 [Google Scholar](#)
90. Tabula Muris Consortium (2018) **Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris** *Nature* **562**:367–372 [Google Scholar](#)
91. Pedregosa F., Varoquaux G., Gramfort A., Michel V. (2011) **Scikit-learn: Machine Learning in Python** *Journal of Machine Learning Research* **12**:2825–2830 [Google Scholar](#)
92. Polański K., Young M.D., Miao Z., Meyer K.B., Teichmann S.A., Park J.-E. (2020) **BBKNN: fast batch alignment of single cell transcriptomes** *Bioinformatics* **36**:964–965 [Google Scholar](#)
93. Rao J.N., Wang J.-Y. (2010) **Intestinal Architecture and Development** Morgan & Claypool Life Sciences [Google Scholar](#)
94. Puram S.V., Parikh A.S., Tirosh I. (2018) **Single cell RNA-seq highlights a role for a partial EMT in head and neck cancer** *Mol Cell Oncol* **5**:e1448244 [Google Scholar](#)
95. Liu J., Lichtenberg T., Hoadley K.A., Poisson L.M., Lazar A.J., Cherniack A.D., Kovatich A.J., Benz C.C., Levine D.A., Lee A.V., et al. (2018) **An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics** *Cell* **173**:400–416 [Google Scholar](#)

96. Davidson-Pilon C. (2019) **lifelines: survival analysis in Python** *The Journal of Open Source Software* **4**:1317 [Google Scholar](#)
97. Fazilaty H., Brügger M.D., Valenta T., Szczerba B.M., Berkova L., Doumpas N., Hausmann G., Scharl M., Basler K. (2021) **Tracing colonic embryonic transcriptional profiles and their reactivation upon intestinal damage** *Cell Rep* **36**:109484 [Google Scholar](#)

Author information

Inbal Avraham-Davidi*

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States

For correspondence: inbalavr@gmail.com

*These authors contributed equally

Simon Mages*

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States, Gene Center and Department of Biochemistry, Ludwig-Maximilians-Universität München, Munich, Germany

*These authors contributed equally

Johanna Klughammer*

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States, Gene Center and Department of Biochemistry, Ludwig-Maximilians-Universität München, Munich, Germany

*These authors contributed equally

Noa Moriel

School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel

Shinya Imada

Department of Biology at MIT, Koch Institute for Integrative Cancer Research at MIT, Cambridge, United States

Matan Hofree

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States

Evan Murray

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States

Jonathan Chen

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States, Massachusetts General Hospital (MGH) Cancer Center, Harvard Medical School (HMS),

Boston, United States, Department of Pathology, Massachusetts General Hospital, Boston, United States

Karin Pelka

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States, Massachusetts General Hospital (MGH) Cancer Center, Harvard Medical School (HMS), Boston, United States, Gladstone-UCSF Institute of Genomic Immunology, Gladstone Institutes, San Francisco, United States, Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, United States

Arnav Mehta

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States, Massachusetts General Hospital (MGH) Cancer Center, Harvard Medical School (HMS), Boston, United States, Dana Farber Cancer Institute, Boston, United States

Genevieve M Boland

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States, Department of Surgery, Massachusetts General Hospital, Boston, United States

Toni Delorey

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States

Leah Caplan

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States

Danielle Dionne

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States

Robert Strasser

Gene Center and Department of Biochemistry, Ludwig-Maximilians-Universität München, Munich, Germany

Jana Lalakova

10xGenomics, Solnavägen 3H, Stockholm, Sweden

Anezka Niesnerova

10xGenomics, Solnavägen 3H, Stockholm, Sweden

Hao Xu

10xGenomics, Solnavägen 3H, Stockholm, Sweden

Morgane Rouault

10xGenomics, Solnavägen 3H, Stockholm, Sweden

Itay Tirosh

Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel

Nir Hacohen

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States, Massachusetts General Hospital (MGH) Cancer Center, Harvard Medical School (HMS), Boston, United States

Fei Chen

Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, United States, Harvard Stem Cell and Regenerative Biology, Cambridge, United States
ORCID iD: [0000-0003-2308-3649](https://orcid.org/0000-0003-2308-3649)

Omer Yilmaz

Department of Biology at MIT, Koch Institute for Integrative Cancer Research at MIT, Cambridge, United States, Department of Pathology, Massachusetts General Hospital, Boston, United States

Jatin Roper

Department of Biology at MIT, Koch Institute for Integrative Cancer Research at MIT, Cambridge, United States, Department of Pharmacology and Cancer Biology and Department of Medicine, Division of Gastroenterology, Duke University, Durham, United States

For correspondence: jatin.roper@duke.edu

Orit Rozenblatt-Rosen

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States, Genentech, South San Francisco, United States

For correspondence: orit.r.rosen@gmail.com

Mor Nitzan

School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel, Racah Institute of Physics and Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel, Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

For correspondence: mor.nitzan@mail.huji.ac.il

Aviv Regev

Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, United States, Department of Biology at MIT, Koch Institute for Integrative Cancer Research at MIT, Cambridge, United States, Genentech, South San Francisco, United States

For correspondence: aviv.regev.sc@gmail.com

Editors

Reviewing Editor

Florent Ginhoux

Singapore Immunology Network, Singapore, Singapore

Senior Editor

Aleksandra Walczak

CNRS, Paris, France

Reviewer #2 (Public review):

In their study, Avraham-Davidi et al. combined scRNA-seq and spatial mapping studies to profile two preclinical mouse models of colorectal cancer: *Apcfl/fl VillinCreERT2* (AV) and *Apcfl/fl LSL-KrasG12D Trp53fl/fl Rosa26LSL-tdTomato/+ VillinCreERT2* (AKPV). In the first part of the manuscript, the authors describe the analysis of the normal colon and dysplastic lesions induced in these models following tamoxifen injection. They highlight broad variations in immune and stromal cell composition within dysplastic lesions, emphasizing the infiltration of monocytes and granulocytes, the accumulation of IL-17+gdT cells and the presence of a distinct group of endothelial cells. A major focus the study is the remodeling of the epithelial compartment, where most significant changes are observed. Using non-negative matrix factorization, the authors identify molecular programs of epithelial cell functions, emphasizing stemness, Wnt signaling, angiogenesis and inflammation as major features associated with dysplastic cells. They conclude that findings from scRNA-seq analyses in mouse models are transposable to human CRC. In the second part of the manuscript, the authors aim to provide the spatial contexture for their scRNA-seq findings using Slide-seq and TACCO. They demonstrate that dysplastic lesions are disorganized and contain tumor-specific regions, which contextualize the spatial proximity between specific cell states and gene programs. Finally, they claim that these spatial organizations are conserved in human tumors and associate region-based gene signatures with patient outcome in public datasets. Overall, the data were collected and analyzed using solid and validated methodology to offer a useful resource to the community.

Main comments:

(1) Clarity. The manuscript would benefit from a substantial reorganization to improve clarity and accessibility for a broad readership. The text could be shortened and the number of figure panels reduced to emphasize the novel contributions of this work while minimizing extensive discussions on general and expected findings, such as tissue disorganization in dysplastic lesions. Additionally, figure panels are not consistently introduced in the correct order, and some are not discussed at all (e.g., Fig. S1D; Fig. 3C is introduced before Fig. 3A; several panels in Fig. 4 are not discussed). The annotation of scRNA-seq cell states is insufficiently explained, with no corresponding information about associated genes provided in the figures or tables. Multiple annotations are used to describe cell groups (e.g., TKN01 = $\gamma\delta$ T and CD8 T, TKN05 = $\gamma\delta$ T_IL17+), but these are not jointly accessible in the figures, making the manuscript challenging to follow. It is also not clear what is the respective value of the two mouse models and timepoints of tissue collection in the analysis.

(2) Novelty. While the study is of interest, it does not present major findings that significantly advance the field or motivate new directions and hypotheses. Many conclusions related to tissue composition and patient outcomes, such as the epithelial programs of Wnt signaling, angiogenesis, and stem cells, are well-established and not particularly novel. Greater exploration of the scRNA-seq data beyond cell type composition could enhance the novelty of the findings. For instance, several tumor microenvironment clusters uniquely detected in dysplastic lesions (e.g., Mono2, Mono3, Gran01, Gran02) are identified, but no further investigation is conducted to understand their biological programs, such as applying nNMF as was done for epithelial cells. Additional efforts to explore precise tissue localization and cellular interactions within tissue niches would provide deeper insights and go beyond the limited analyses currently displayed in the manuscript.

(3) Validation. Several statements made by the authors are insufficiently supported by the data presented in the manuscript and should be nuanced in the absence of proper validation. For example: 1.) RNA velocity analyses: The conclusions drawn from these analyses are speculative and need further support. 2.) Annotations of epithelial clusters as dysplastic: These annotations could have been validated through morphological analyses and staining

on FFPE slides. 3.) Conservation of mouse epithelial programs in human tumors: The data in Figure S5B does not convincingly demonstrate enrichment of stem cell program 16 in human samples. This should be more explicitly stated in the text, given the emphasis placed on this program by the authors. 4.) Figure S6E: Cluster Epi06 is significantly overrepresented in spatial data compared to scRNA-seq, yet the authors claim that cell type composition is largely recapitulated without further discussion, which reduces confidence in other conclusions drawn.

Furthermore, stronger validation of key dysplastic regions (regions 6, 8, and 11) in mouse and human tissues using antibody-based imaging with markers identified in the analyses would have considerably strengthened the study. Such validation would better contextualize the distribution, composition, and relative abundance of these regions within human tumors, increasing the significance of the findings and aiding the generation of new pathophysiological hypotheses.

Comments on revisions:

The authors have improved the clarity of the manuscript and responded adequately to all my initial comments.

I don't have any other comments. Congratulations to the authors on this work.

<https://doi.org/10.7554/eLife.104815.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

Summary:

The authors conducted a spatial analysis of dysplastic colon tissue using the Slide-seq method. Their main objective is to build a detailed spatial atlas that identifies distinct cellular programs and microenvironments within dysplastic lesions. Next, they correlated this observation with clinical outcomes in human colorectal cancer.

Strengths:

The work is a good example of utilising spatial methods to study different tumour models. The authors identified a unique stem cell program to understand tumours gently and improve patient stratification strategies.

Weaknesses:

However, the study's predominantly descriptive nature is a significant limitation. Although the spatial maps and correlations between cell states are interesting observations, the lack of functional validation-primarily through experiments in mouse models-weakens the causal inferences regarding the roles these cellular programs play in tumour progression and therapy resistance.

We thank the reviewer for this comment. Indeed, functional validation to pin down causal dependencies and a more thorough investigation of tumor progression and therapy resistance both in mouse model as well as human patients and/or patient derived samples would broaden the insights to be gained from this work. Unfortunately, this is beyond the scope of this study.

The authors also missed an opportunity to link the mutational status of malignant cells with the cellular neighbourhoods.

The data reported in this study only contains spatial data for one mouse model (AV). As spatial data for the other model (AKPV) is missing, it is not possible to link the mutational type of the model with the cellular neighborhoods. We did investigate whether there is extra somatic mutational heterogeneity in the AV data, both regarding single nucleotide variations (SNVs) and copy number variations (CNVs). But at the time when the mice were sacrificed (after 3 weeks) there was no significant mutational heterogeneity discoverable.

Overall, the study contributes to profiling the dysplastic colon landscape. The methodologies and data will benefit the research community, but further functional validation is crucial to validate the biological and clinical implications of the described cellular interactions.

Reviewer #2 (Public review):

In their study, Avraham-Davidi et al. combined scRNA-seq and spatial mapping studies to profile two preclinical mouse models of colorectal cancer: Apcfl/fl VillinCreERT2 (AV) and Apcfl/fl LSL-KrasG12D Trp53fl/fl Rosa26LSL-tdTomato/+ VillinCreERT2 (AKPV). In the first part of the manuscript, the authors describe the analysis of the normal colon and dysplastic lesions induced in these models following tamoxifen injection. They highlight broad variations in immune and stromal cell composition within dysplastic lesions, emphasizing the infiltration of monocytes and granulocytes, the accumulation of IL-17+gdT cells, and the presence of a distinct group of endothelial cells. A major focus of the study is the remodeling of the epithelial compartment, where the most significant changes are observed. Using non-negative matrix factorization, the authors identify molecular programs of epithelial cell functions, emphasizing stemness, Wnt signaling, angiogenesis, and inflammation as major features associated with dysplastic cells. They conclude that findings from scRNA-seq analyses in mouse models are transposable to human CRC. In the second part of the manuscript, the authors aim to provide the spatial context for their scRNA-seq findings using Slide-seq and TACCO. They demonstrate that dysplastic lesions are disorganized and contain tumor-specific regions, which contextualize the spatial proximity between specific cell states and gene programs. Finally, they claim that these spatial organizations are conserved in human tumors and associate region-based gene signatures with patient outcomes in public datasets. Overall, the data were collected and analyzed using solid and validated methodology to offer a useful resource to the community.

Main comments:

(1) Clarity

The manuscript would benefit from a substantial reorganization to improve clarity and accessibility for a broad readership. The text could be shortened and the number of figure panels reduced to emphasize the novel contributions of this work while minimizing extensive discussions on general and expected findings, such as tissue disorganization in dysplastic lesions. Additionally, figure panels are not consistently introduced in the correct order, and some are not discussed at all (e.g., Figure S1D; Figure 3C is introduced before Figure 3A; several panels in Figure 4 are not discussed). The annotation of scRNA-seq cell states is insufficiently explained, with no corresponding information about associated genes provided in the figures or tables. Multiple annotations are used to describe cell groups (e.g., TKN01 = $\gamma\delta$ T and CD8 T, TKN05 = $\gamma\delta$ IL17+), but these are not jointly accessible in the figures, making the manuscript challenging to follow. It is also

not clear what is the respective value of the two mouse models and time points of tissue collection in the analysis.

We thank the reviewer for this suggestion. We clarified and simplified the revised manuscript, however we believe that the current discussions are an important part of the manuscript and would be useful to readers. We reordered panels in Figures S1 and 3 to align with their appearance in the manuscript. We kept the order of other panels as it is to keep both context and coherence of those figures intact. We changed the way we reference cell clusters in the manuscript to better align with the naming scheme introduced in Figure 1B. The respective value of the two mouse models as well as the time points of tissue collection are described in lines 108-120 of the manuscript.

(2) Novelty

While the study is of interest, it does not present major findings that significantly advance the field or motivate new directions and hypotheses. Many conclusions related to tissue composition and patient outcomes, such as the epithelial programs of Wnt signaling, angiogenesis, and stem cells, are well-established and not particularly novel. Greater exploration of the scRNA-seq data beyond cell type composition could enhance the novelty of the findings. For instance, several tumor microenvironment clusters uniquely detected in dysplastic lesions (e.g., Mono2, Mono3, Gran01, Gran02) are identified, but no further investigation is conducted to understand their biological programs, such as applying nNMF as was done for epithelial cells. Additional efforts to explore precise tissue localization and cellular interactions within tissue niches would provide deeper insights and go beyond the limited analyses currently displayed in the manuscript.

We thank the reviewer for this comment. Our study aimed to spatially characterize the tumor microenvironment, with scRNA-seq analysis serving to support this spatial characterization.

Due to technical limitations—such as the number of samples and the limited capture efficiency of Slide-seq—the resolution of immune cell identification in our spatial analysis is constrained. Additionally, while immune and stromal cells formed distinct clusters, epithelial cells exhibited a continuum that was better captured using nNMF.

Lastly, our manuscript provides a general characterization of monocyte and granulocyte populations in scRNA-seq (line 144) and their spatial microenvironments (line 400). We believe that additional analyses of these populations would be beyond the scope of this study and could place an unnecessary burden on the reader. Instead, we suggest that such analyses be explored in future studies.

We remark that we analyzed tissue localization for two entirely different spatial transcriptomics assays (Slide-seq and Cartana) at the resolution of cell types and programs, which was feasible within the constraints of the sparsity, gene panel and sample size in the experiments. A future potential path to further increase the resolution of investigation in this dataset is to include other datasets, e.g. by the emerging transformer-based spatial transcriptomics integration methods.

We also remark that the manuscript already includes an investigation of cellular interactions within tissue niches based on COMMOT (Fig 4k, Fig S8i, Supp Item 4).

(3) Validation

Several statements made by the authors are insufficiently supported by the data presented in the manuscript and should be nuanced in the absence of proper validation. For example:

(a) RNA velocity analyses: The conclusions drawn from these analyses are speculative and need further support.

We thank the reviewer for this comment. We clarified that our conclusions from the RNA velocity analysis need further support by experimental validation (lines 223-225), which is outside the scope of the current study.

(b) Annotations of epithelial clusters as dysplastic: These annotations could have been validated through morphological analyses and staining on FFPE slides.

We thank the reviewer for this comment. While this could have been a possible approach, our study primarily relies on scRNA-seq, which does not preserve tissue morphology, and Slide-seq of fresh tissue, where such an analysis is particularly challenging.

(c) Conservation of mouse epithelial programs in human tumors: The data in Figure S5B does not convincingly demonstrate the enrichment of stem cell program 16 in human samples. This should be more explicitly stated in the text, given the emphasis placed on this program by the authors.


We thank the reviewer for pointing this out. We clarified the section about the stem cell program 16 and references to Figures S5A and S5B (lines 269-274): while we do see correlation in the definition of human programs with the mouse stem cell program (Figure S5A), we do not see a correlated expression of the stem cell program across human and mouse (Figure S5B).

(d) Figure S6E: Cluster Epi06 is significantly overrepresented in spatial data compared to scRNA-seq, yet the authors claim that cell type composition is largely recapitulated without further discussion, which reduces confidence in other conclusions drawn.

We thank the reviewer for this remark. Indeed, Epi06 was a cluster which drew our attention during early analyses for its mixed expression profiles with contributions of vastly different cell types. We concluded that this is best explained by doublets, but we cannot rule out (partial) non-doublet explanations (e.g. undifferentiated cells). As doublet detection with Scrublet did not flag those cells as doublets, we kept these cells in the workflow, but excluded them from further interpretation. While in the previous version of the manuscript we only shortly hinted to this in figure legend 2A ("Cluster Epi06: doublets (not called by Scrublet)"), we expanded on this in the methods section of the revised manuscript (lines 863-869). Given the doublet interpretation, the observation that this cluster is significantly overrepresented in the annotation of the spatial data is not surprising as this annotation comes from the decomposition of compositional data which contains contributions of multiple cells per Slide-seq bead which are structurally very similar to doublets. While Epi06 appears enriched in S6E when comparing Slide-Seq to scRNA-seq, there are multiple technical cross platform differences, including different per-gene sensitivities or capture biases for certain cell types (e.g. stromal cells suffering more from dissociation in scRNA compared to Slide-Seq). We believe that comparisons between disease states within a single platform are more biologically meaningful, like the comparison between normal and premalignant tissue, which is presented in Figure S6G. To increase confidence in the analysis and to assess whether intra-platform biological conclusions are affected by the inclusion/exclusion of Epi06, we recreated Figure S6G for a Slide-Seq cell type annotation without Epi06 in the reference (see Author response image 1). Even though Epi06 is missing in that annotation, the strong enrichments are consistently preserved between the two analysis variants, while as expected some less significant enrichments with larger FDR values are not preserved.

Author response image 1.

Significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of cell clusters (rows) in normal (N) or AV (AV) tissues based on Slide-seq ("spatial") data or scRNA-seq ("sc") including (A) or excluding (B) Epi06 in the reference for annotating the Slide-Seq data (A is identical to Figure S6G in the manuscript).



Furthermore, stronger validation of key dysplastic regions (regions 6, 8, and 11) in mouse and human tissues using antibody-based imaging with markers identified in the analyses would have considerably strengthened the study. Such validation would better contextualize the distribution, composition, and relative abundance of these regions within human tumors, increasing the significance of the findings and aiding the generation of new pathophysiological hypotheses.

We agree with the reviewer with their assessment that validation by antibody-based imaging (or other spatial proteomics data) would have been useful follow-up experiments, yet these are beyond the scope of the current study.

Reviewer #1 (Recommendations for the authors):

AV and AKPV have different oncogenic mutations, and their impact on spatial neighbourhoods is unclear. Can authors perform an analysis to understand the contribution of oncogenic mutations on the spatial landscape of CRC?

The data reported in this study only contains spatial data for one mouse model (AV). As spatial data for the other model (AKPV) is missing, it is not possible to comparatively link the mutational type of the model with the spatial landscape.

<https://doi.org/10.7554/eLife.104815.2.sa0>