

The *Drosophila* hematopoietic niche assembles through collective cell migration controlled by neighbor tissues and Slit-Robo signaling

Kara A Nelson, Kari F Lenhart, Lauren Anllo, Stephen DiNardo 

Department of Cell and Developmental Biology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, United States • Institute for Regenerative Medicine at the University of Pennsylvania, Philadelphia, United States • Department of Biology, Drexel University, Philadelphia, United States

 https://en.wikipedia.org/wiki/Open_access Copyright information

Abstract

Niches are often found in specific positions in tissues relative to the stem cells they support. Consistency of niche position suggests that placement is important for niche function. However, the complexity of most niches has precluded a thorough understanding of how their proper placement is established. To address this, we investigated the formation of a genetically tractable niche, the *Drosophila* Posterior Signaling Center (PSC), the assembly of which had not been previously explored. This niche controls hematopoietic progenitors of the lymph gland (LG). PSC cells were previously shown to be specified laterally in the embryo, but ultimately reside dorsally, at the LG posterior. Here, using live-imaging, we show that PSC cells migrate as a tight collective and associate with multiple tissues during their trajectory to the LG posterior. We find that Slit emanating from two extrinsic sources, visceral mesoderm and cardioblasts, is required for the PSC to remain a collective, and for its attachment to cardioblasts during migration. Without proper Slit-Robo signaling, PSC cells disperse, form aberrant contacts, and ultimately fail to reach their stereotypical position near progenitors. Our work characterizes a novel example of niche formation and identifies an extrinsic signaling relay that controls precise niche positioning.

Impact Statement

A new example of niche formation is provided which reveals the mode of niche cell migration, implicates extrinsic sources that deliver positional information, and uncovers the signaling pathway required for the precise, stereotypical positioning of the niche.

eLife assessment

This study presents **valuable** findings on the role of a well-studied signal transduction pathway, the Slit/Robo system, in the context of the assembly of the hematopoietic niche in the *Drosophila* embryo. The evidence supporting the claims of the authors is **solid**. However, one aspect that needs attention is whether the cells are migrating and not being pushed to a more dorsal position through dorsal closure and/or other similar large-scale embryo movement. This does not detract from the very interesting analysis of PSC morphogenesis and will interest developmental biologists working on molecular mechanisms of tissue morphogenesis.

<https://doi.org/10.7554/eLife.100455.1.sa4>

Introduction

Homeostasis and repair of many organs relies on a resident stem cell population. Stem cell behavior is often coordinated by a niche (Morrison & Spradling, 2008) —the specific microenvironment that contains stem cells—and abnormal regulation can severely impact health by leading to tissue atrophy or tumor formation (Chakkalakal et al., 2012; Ferraro et al., 2010; Kobiela et al., 2007; Walkley et al., 2007). Thus, it is essential to understand stem cell-niche interactions. An important step in understanding the behavior of a mature niche is understanding the particular attributes it acquires—such as cellular features and the positioning of its cells with respect to each other and stem cells—and how those attributes are regulated as the niche is built during development. Yet, there are very few studies of niche formation at the cellular level. The few well-studied niches consistently acquire tissue-specific positional and structural characteristics (Anllo et al., 2019; Biggs et al., 2018; Gordon et al., 2020; Sumigray et al., 2018). The reproducibility of these characteristics suggests that acquisition of a particular architecture is essential to niche function and emphasizes the importance of understanding how the architecture of a given niche is established.

Frequent barriers to studying niche formation include niche complexity and lack of niche cell markers (Heitman et al., 2018; Pinho & Frenette, 2019). Studying niche formation *in vivo* is further hindered by an inability to follow its constituent cells as they form the niche in real-time due to lack of tissue and organism transparency, and the existence of large-scale tissue movements (Bostock et al., 2020; Boulais & Frenette, 2015; Gregg & Butcher, 2012). The mammalian hematopoietic stem cell niche is a prime example of a difficult-to-study niche due to its location within opaque bone and its complexity—it is comprised of numerous cellular components and molecular factors (Boulais & Frenette, 2015; Calvi et al., 2003; Pinho & Frenette, 2019; Zhang et al., 2003).

Despite these limitations, some niches, such as the mammalian intestinal epithelium and hair follicle, are beginning to be studied at the cellular level (Anllo & DiNardo, 2022; Díaz-Torres et al., 2021; Gupta et al., 2019; Pentimikko et al., 2022; Sumigray et al., 2018). However, investigations about establishment of these well-characterized niches still suffer from challenges including limited real-time imaging and lack of tissue-specific manipulations.

To overcome these limitations and gain insight into the mechanisms that drive niche formation, we have investigated the development of the niche that supports the *Drosophila* larval hematopoietic organ: the lymph gland (LG). The cells that constitute this niche, called the posterior signaling center (PSC), and the markers that label it are known (Croizatier et al., 2004; Lebestky

et al., 2003 [\[link\]](#); Mandal et al., 2007 [\[link\]](#)). Additionally, the PSC resides under a thin epidermal covering, making it amenable to live-imaging with high spatial resolution, as we show here. The dynamic information afforded by live-imaging can reveal the mechanism of migration and implicate nearby tissues as a source for guiding signals. These advantages, paired with facile *Drosophila* genetics, make for a powerful experimental system in which PSC formation can be visualized *in vivo* and the underlying mechanisms probed.

Minimal information exists about the formation of the PSC, as most analyses of the lymph gland have focused on the steady-state operations of late larval stages when the gland is mature and most accessible due to its larger size. While the mature LGs are comprised of multiple pairs of lobes (Koranteng et al., 2022 [\[link\]](#); Shrestha & Gateff, 1982 [\[link\]](#)), we use ‘lymph gland’ to refer to only one of the pair of thoroughly characterized, bilaterally symmetric ‘primary’ lobes. In late larval stages the LG contains thousands of cells, organized into multiple zones: the medullary, intermediate, and cortical zones, and the PSC (Banerjee et al., 2019 [\[link\]](#); Jung et al., 2005 [\[link\]](#); Lanot et al., 2001 [\[link\]](#)). The PSC regulates the adjacent hematopoietic progenitors of the medullary zone; these progenitors are progressively differentiated from the innermost region of the LG, radially outward (Cho et al., 2020 [\[link\]](#); Jung et al., 2005 [\[link\]](#); Luo et al., 2020 [\[link\]](#); Mandal et al., 2007 [\[link\]](#)). The mature, terminally differentiated hemocytes reside at the outermost cortex of the gland, in the cortical zone, and between these zones are the cells of the intermediate zone (Jung et al., 2005 [\[link\]](#); Krzemien et al., 2010 [\[link\]](#); Spratford et al., 2021 [\[link\]](#)). Recent scRNA-seq analyses have identified new types of hemocytes in the LG (Cho et al., 2020 [\[link\]](#); Girard et al., 2021 [\[link\]](#)), but the three main types are analogous to vertebrate myeloid cells and include crystal cells, plasmacytes, and lamellocytes, which are responsible for wound healing and innate immunity (Banerjee et al., 2019 [\[link\]](#); Lanot et al., 2001 [\[link\]](#); Rizki & Rizki, 1992 [\[link\]](#)).

The larval PSC facilitates wound repair, immune response, and homeostasis and does so by performing two key functions: maintaining progenitors and inducing progenitor differentiation (Baldeosingh et al., 2018 [\[link\]](#); Crozatier et al., 2004 [\[link\]](#); Krzemień et al., 2007 [\[link\]](#); Mandal et al., 2007 [\[link\]](#); Ramesh et al., 2021 [\[link\]](#)). PSC positioning is such that it contacts both the least differentiated progenitors and mature hemocytes (Baldeosingh et al., 2018 [\[link\]](#); Jung et al., 2005 [\[link\]](#)), a seemingly prime location where the PSC is poised to both implement regulation and receive feedback, thereby maintaining homeostasis in the LG. For example, upon immune challenge, the PSC senses and responds to the threat by instructing differentiation of progenitors (Khadilkar et al., n.d.; Louradour et al., 2017 [\[link\]](#); Sinenko et al., 2012 [\[link\]](#)). The PSC engages multiple signaling pathways to execute its roles and relies on cell biological characteristics of its component cells, such as occluding or gap junctions, to do so (Baldeosingh et al., 2018 [\[link\]](#); Ho et al., 2023 [\[link\]](#); Khadilkar et al., 2017 [\[link\]](#); Mandal et al., 2007 [\[link\]](#); Sinenko et al., 2009 [\[link\]](#)).

PSC functionality is integral for organism health, as PSC loss can cause precocious differentiation of progenitors under homeostatic conditions and inability to produce lamellocytes under immune challenge conditions (Baldeosingh et al., 2018 [\[link\]](#); Krzemień et al., 2007 [\[link\]](#); Mandal et al., 2007 [\[link\]](#)). PSC-regulated homeostatic maintenance of the larval LG is crucial, as the LG will ultimately rupture, releasing its constituents into circulation to contribute to the hematopoietic pool in pupal and adult stages (Grigorian et al., 2011 [\[link\]](#); Holz et al., 2003 [\[link\]](#)). Furthermore, recent publications suggest that the PSC itself adopts new functions upon LG rupture – PSC cells become highly motile and phagocytic and are capable of transdifferentiating into lamellocytes or plasmacytes upon immune challenge (Boulet et al., 2021 [\[link\]](#); Hirschhäuser et al., 2023 [\[link\]](#)).

Though the late larval PSC is comprised of about 30-50 cells (Ho et al., 2021 [\[link\]](#); Morin-Poulard et al., 2016 [\[link\]](#); Tokusumi et al., 2015 [\[link\]](#)), it is initially specified as only about five cells (Mandal et al., 2007 [\[link\]](#)). The expansion of PSC cell number takes place only after the five cells become organized into the PSC at the posterior of the developing LG during late stages of embryogenesis (Mandal et al., 2007 [\[link\]](#)). How the PSC becomes consistently positioned after specification is unknown and is our focus here. The LG itself is specified about mid-way through embryogenesis, around stage 12,

from cardiogenic mesoderm as three cell clusters located laterally— one in each thoracic segment (schematized in **Figure 1A**, left; Crozatier et al., 2004; Mandal et al., 2004, 2007). The PSC is specified from the posterior-most LG cells by expression of *Antennapedia* and the silencing of *Homothorax* (Mandal et al., 2007). A transcription factor, *Collier*, the *Drosophila* ortholog of mammalian early B-cell factor, is initially expressed in the entire LG primordium but becomes restricted to the PSC and is necessary for its maintenance and function (Crozatier et al., 2004; Krzemień et al., 2007). From fixed preparations, it is known that the three LG clusters coalesce along the anterior-posterior axis before reaching the dorsal midline. At the end of embryogenesis, the cluster of PSC niche cells resides at the posterior of each bilaterally symmetric lymph gland; each gland flanks the dorsal vessel—the *Drosophila* heart—at the dorsal midline (schematized in **Figure 1A**, right; (Crozatier et al., 2004; Mandal et al., 2007). Apart from this description nothing is known about how the PSC is built, and no live-imaging of the PSC has been conducted during embryogenesis. Despite the probable importance of PSC positioning to later function, the mechanism of migration, the signals guiding migration, and interactions among constituent cells—each integral aspects of organogenesis—are entirely unknown.




Figure 1.

Live-imaging reveals dynamics of PSC migration and presence of nearby muscles.

All images are oriented with anterior to the left and posterior to the right. A) Schematic depicts prior knowledge of PSC migration. Prospective cells of lymph glands (hematopoietic progenitors, magenta, and PSC, green) and cardioblasts (blue) of the dorsal vessel are specified as distinct clusters prior to migration onset around stage 13 (left). Around stage 16 (right) lymph gland clusters have coalesced and flank the dorsal vessel at the dorsal midline; bilaterally symmetric counterparts are aligned. B-F) Fixed *w¹¹¹⁸* embryos stained for Antp and Odd to label the PSC. B) Stage 14 PSC (green) is lateral in embryo, tightly clustered, and flanks 2 Antp+ CBs (blue). C) Late stage 14 PSC is more dorsal, flanking 2 Antp+ CBs. D) The stage 15 PSC is even more dorsal, elongated, and flanks 4 Antp+ CBs; contralateral Antp+ CBs in view. E) Late stage 15 PSC is more medial and compact. F) Stage 16 LGs and CBs at final position at dorsal midline, right LG partially in view. PSC is compact and coalesced at LG posterior, flanking Antp+ CBs. Antp+ CBs (yellow brackets) are neatly aligned in stereotypical 2x4 organization. B'-F') Schematics depicting relevant cell types in B-F: hematopoietic progenitors (magenta), PSC (green), and Antp+ cardioblasts (blue). G-J) Live-imaging stills from Hand-RFP, Antp-GAL4, 10xUAS-myr::GFP embryos; RFP in magenta and GFP in white. G) Timelapse shows dorsolateral view of PSC (magenta and white cells above green underline and below string of CBs, arrows) migrating as a collective from migration onset (0m), when the PSC is lateral, to migration completion (around 2h50m), when the PSC is dorsal. Arrowhead indicates fixed positioning of the posterior-most PSC cell in the collective. H-H''') Live-imaging stills show PSC protrusions during mid-migration. H) Merge shows positional relation of PSC (white and magenta) and its protrusions to rest of lymph gland and CBs (along top of image). H'-H''') Only GFP channel. Progressively shallower z-slices of same PSC. H') Anterior protrusions; some short (arrowheads) and one long (bracket) that encases lymph gland. H'') Posterior protrusion (bracket). H''') Green dot indicates dorsal-most PSC cell with short, branching protrusions (arrowheads). Much of image contains Antp+ epidermis. I) Live-imaging stills of GFP channel only show PSC cells shifting position within collective. Initially connected yellow and cyan PSC cells (0m, arrowhead) become separated by intercalation of magenta and green PSC cells while the posterior-most, dorsal vessel-adjacent PSC cell (orange dot) remains at a fixed position in the collective. J) Same embryo as (G), GFP channel only. For the entire timelapse PSC cells migrate dorsally in close association with concurrently migrating neighbor tissues, CBs and Vm. Posterior-most PSC cell (arrowhead) maintains its dorsal vessel-adjacent position throughout. 0m inset shows PSC proximity to Vm and 2h50m inset shows PSC proximity to CBs. J') Schematics depict relevant cell types for each panel of J time series: cardioblasts (blue), PSC (green), and visceral mesoderm (orange). K) Live-imaging series from dorsal vantage point shows A1 AMs (magenta; labeled with *org-1-HN39::RFP*) contact PSCs (green outlines; labeled by *Antp>CD8::GFP*) throughout the timelapse as both tissues migrate to dorsal midline. L) Schematics depict the relative positioning of the tissues identified to be near the PSC at migration onset (left) and completion (right).

We have conducted the first *in vivo* live-imaging of PSC formation, visualizing migration of PSC cells from their origin, lateral in the embryo, to their final, dorsal position at the posterior of the LG. We find that a prominent, regulated feature of this niche is the migration of its constituent cells as a collective. Live-imaging also revealed neighboring tissues that could feasibly signal to influence PSC migration as well as its positioning in the LG. Using genetic ablation and mutant analyses, we have identified that both the dorsal vessel and visceral mesoderm are required for proper positioning of the PSC. In addition, mutant analyses and tissue-specific challenges revealed an intricate web of Slit-Robo signaling in and between these tissues that is essential to establish positioning of PSC cells as a coalesced, compact collective at the LG posterior. Finally, we use live-imaging to show that Slit signaling is essential to maintain collectivity of PSC cells and properly position these cells during their migration. Taken together, we have uncovered a new example of collective cell migration, revealed some of its mechanistic underpinnings, and implicated a signaling relay required to properly build and position this niche.

Results

Live-imaging reveals dynamics of PSC formation

While PSC formation is undoubtedly a dynamic process, current knowledge in the field is derived exclusively from static timepoints using fixed embryos (**Figure 1A** [↗](#)). Thus, we sought to visualize the entirety of PSC migration in real time. To identify and track PSC cells live, we first confirmed their location at various time points in fixed embryos using the accepted markers, *Antp* and *Odd*, where co-expression demarcates PSC cells. We characterized PSC positioning in stages ranging from migration onset, stage 14, until migration completion, stage 16 (schematized in **Fig 1B** [↗](#)–**F**). Prior to migration, the PSC appeared as a compact cluster of cells (**Figure 1B** [↗](#), green underline), it appeared more elongated through mid-migration (**Figure 1C** [↗](#)–**1E** [↗](#)), and ultimately it assumed a compact, clustered organization at the posterior of the lymph gland (**Figure 1F** [↗](#)). During migration and upon its completion, the PSC flanked *Antp*⁺ cardioblasts of the dorsal vessel (**Figure 1F** [↗](#), brackets). Though not explored further here, ECM components, Perlecan and Viking, were expressed around the PSC collective including between the PSC and the adjacent cardioblasts (Figure S1A and S1B).

To our knowledge, a sufficiently bright, stable marker that labels only PSC cells during migration stages does not exist. Thus, for live-imaging we combined Hand-RFP, which marks lymph gland (including the PSC) and cardioblasts, with a myristoylated GFP driven by *Antp*-GAL4 (*Antp* > myrGFP) to mark the PSC but not other lymph gland cells (**Figure 1G–J** [↗](#)). We performed timelapse imaging on embryos beginning after PSC specification (stage 13) until the PSC completed migration and morphogenesis (stage 17). We observed that the PSC moved as a collective for the entire migration (**Figure 1G** [↗](#)). During migration many PSC cells extended protrusions—some reached far anteriorly, encasing the lymph gland (**Figure 1H** [↗](#), bracket) while others were shorter or branched (**Figure 1H** [↗](#), arrowheads). The protrusions exhibited no apparent directional bias but were more common at non-dorsal vessel-adjacent surfaces of the collective. During migration some cells shifted positioning with respect to one another. For example, two lateral PSC cells that abut early on (**Figure 1I** [↗](#), 0m, arrowhead) become separated by interdigitating cells (**Figure 1I** [↗](#), 8–40m, magenta and green dots). In general, however, most cells remained at relatively fixed positions within the collective—particularly the posterior-most, dorsal vessel-adjacent PSC cells (**Figure 1I** [↗](#), orange dots and **Figure 1G** [↗](#), arrowheads). Live-imaging revealed the importance of collective migration in building this niche, even as some PSC cells exhibit cellular extensions and shift position within the collective.

Visceral, cardiac, and alary muscles are near the PSC throughout its migration

Crucially, live-imaging also uncovered that the PSC was within signaling distance of various muscles for the entirety of its migration. It is well-documented that muscles can supply positional information to nearby tissues (Anillo & DiNardo, 2022 [↗](#); Lucila Scimone et al., 2017 [↗](#); Witchley et al., 2013 [↗](#)), and our imaging indicated that visceral muscle (or, visceral mesoderm; Vm) and cardioblasts (CBs) of the dorsal vessel were each near the PSC. We live-imaged with a dorsolateral vantage point which revealed that the PSC migrated dorsally in synchrony with CBs and Vm (**Figure 1J** [↗](#)). The PSC remained laterally affixed to *Antp*⁺ CBs (**Figure 1J** [↗](#), blue) and slightly dorsal to *Antp*⁺ Vm (**Figure 1J** [↗](#), orange) throughout migration.

Based on segmental positioning, we suspected the PSC was also near alary muscle (AM)—segmentally repeating muscles that attach internal organs to the body wall. We live-imaged PSC and AM migration from a dorsal vantage point with *Antp*>CD8:GFP and *org-1*-HN39::RFP, a marker with expression restricted to AMs. Indeed, the first AM was immediately lateral to the PSC for the

entirety of their synchronized migration to the dorsal midline (**Figure 1K** [↗](#), PSCs outlined in green). We confirmed this proximity by fixing *org-1*-HN39::RFP embryos and staining for Antp, which revealed that at late stages, the A1 AM ensheathed the PSC (Figure S1D), with some fibers encasing individual PSC cells (Figure S1E).

Thus, live-imaging provided invaluable insights, advancing our knowledge of PSC formation from that attainable through analysis of fixed preparations (**Figure 1A** [↗](#)), by revealing the mode of PSC migration and the spatiotemporal dynamics of the PSC relative to nearby tissues (**Figure 1L** [↗](#)). The sustained proximity of alary muscles, visceral mesoderm, and cardioblasts to the PSC made each a strong candidate for influencing PSC positioning.

Vm and CBs are required for proper PSC positioning

Having identified multiple candidates, we next investigated whether any or all of these muscles were necessary for proper PSC positioning. To test a role for AM, we ablated it by expressing the pro-apoptotic factor, *grim*, using the AM-specific AME_r-GAL4. Ablation was successful, evidenced by lack of GFP-labeled AMs (**Figure 2B** [↗](#)) compared to controls (**Figure 2A** [↗](#)). Based on our characterization of the wildtype PSC (**Figure 1** [↗](#)), we developed criteria to score PSCs at the culmination of their migration as positioned ‘normally’ or ‘abnormally’. A ‘normal’ PSC must be 1) coalesced within one nuclear diameter of one another, 2) adjacent to the dorsal vessel, and 3) at the same dorsal-ventral position as the posterior-most progenitors of the lymph gland. Surprisingly, AM ablation had no impact on PSC positioning (**Figure 2C** [↗](#)) nor the total number of PSC cells (Figure S2A), indicating that AMs are not required for proper PSC formation.

To determine whether Vm was involved in forming the PSC, we examined *biniou* mutants in which the Vm is genetically ablated (Zaffran et al., 2001 [↗](#)). In controls, the Vm is apparent at stage 12 as columnar, Fas3⁺ cells (**Figure 2D** [↗](#), brackets). By stage 16, Vm has surrounded the gut and constricted it into sections (**Figure 2E** [↗](#), arrow is first Vm constriction that separates the first two gut sections). By contrast, in *bin* mutants, minimal Fas3⁺ Vm was detected at stage 12 (**Figure 2G** [↗](#)), and it was absent at stage 16 (**Figure 2H** [↗](#)). The majority of *bin*^{R22} mutants and *bin*^{R22}/*bin*^{S4} transheterozygous mutants had abnormal PSC positioning (**Figure 2I** [↗](#), arrowheads) compared to heterozygote controls with normal PSC positioning (**Figure 2F** [↗](#), outlines; quantified in 2J). We detected no Bin protein in PSC cells (Figure S2B), indicating PSC positioning defects in *bin* mutants did not originate PSC-intrinsically but rather were caused by lack of Vm.

To confirm a role for Vm in PSC formation, we ablated Vm by expressing the proapoptotic gene, *hid*, using a Vm-specific driver, *bap*-GAL4. Stage 11 sibling controls displayed characteristic organization of Vm precursors (**Figure 2K** [↗](#)): segmentally repeating mounds of fusion competent myoblasts (FCMs; between green and magenta lines) atop columnar founder cells (FCs; below magenta line). Stage 11 ablated embryos were largely missing FCMs (**Figure 2N** [↗](#), brackets). Though most FCs were present at this stage, they exhibited substantially reduced Bin expression compared to controls (Figure S2C and S2D, quantified in S2E), suggesting improper differentiation. By stage 16 no Vm was detectable (**Figure 2O** [↗](#)). Vm-ablated embryos had abnormal PSC positioning (**Figure 2P** [↗](#), arrowheads) more often than sibling controls (**Figure 2M** [↗](#), quantified in 2Q). Taken together, these analyses definitively establish that an extrinsic cue(s) provided by Vm governs PSC positioning.

To address when the cue might be delivered, we analyzed *jelly belly* mutants and *bagpipe* hypomorphs, *bap*²⁰⁸. In both backgrounds Vm precursors are present initially at stage 11 but do not differentiate or migrate properly from stages 13-16 (Figure S2F-S2J and Englund et al., 2003; Lee et al., 2003 [↗](#); Weiss et al., 2001 [↗](#)). In both mutants PSC positioning was unaffected (Figure S2K, N, and P; quantified in S2L and S2Q), revealing that early signaling from Vm is sufficient for PSC positioning.




Figure 2.

Visceral mesoderm and cardioblasts are required for PSC formation.

All PSCs, co-labeled by Antp and Odd, are viewed dorsally from st16 or 17 embryos unless otherwise noted. A and B) LGs with normally-positioned PSCs outlined in yellow from control (A) and AM-ablated (B) embryos with AMs labeled by AME-Gal4 driven mCD8:GFP. A' and B') GFP channel only with PSC outlines overlayed. C) PSC positioning quantification. D and E) *bin* heterozygotes labeled with Fas3, lateral views. D) green bracket indicates st12 Vm. E) st16 with green arrow indicating first Vm constriction that segregates two sections of gut. F) *bin* heterozygote LGs with normal PSC positioning. G and H) lateral views of *bin* mutants with minimal (G, st12) and absent (H, st16) Vm. I) *bin* mutant LGs with dispersed PSCs (cyan arrowheads). J) PSC positioning quantification including analysis of *bin*^{R22}/*bin*^{S4} transheterozygous mutants. K-P) control embryos (K-M) compared to Vm-ablated embryos (N-P). K) Lateral view of normal st11 Vm labeled by Fas3. Founder cells are below magenta line and fusion competent myoblasts are between magenta and green lines. L) Dorsal view of normal st16 Vm. M) Normal PSC positioning, yellow outlines, in control. N) Fusion competent myoblasts absent, green brackets, in st11 Vm ablated embryo. Most founder cells present with occasional gaps. O) Vm absent in st16 Vm ablated embryo. P) Abnormal PSC positioning, cyan arrowheads. Q) PSC positioning quantification. R and S) st15 LGs with cardioblasts labeled by Mef2 in control (R) and CB-ablated embryos (S). R) Normal PSC positioning, yellow outlines. S) CBs ablated on left side, bracket, and corresponding PSC is mis-positioned, arrowheads, while right side CBs are intact and the R PSC is positioned normally, yellow outline. R' and S') Mef2 channel only. T) Quantification comparing PSC positioning in control embryos to positioning of PSCs with ablated ipsilateral CBs. Scale bars as indicated. Ns = not significant, **p*<0.05, ***p*<0.01, ****p*<0.001, *****p*<0.0001, Fisher's Exact test. Sample sizes as indicated.

Finally, to assess whether cardioblasts of the dorsal vessel impact PSC positioning, we expressed *hid* or *grim* with the CB-specific driver, *tinC44-GAL4*. We used conditions that restricted defects to CBs and left Vm unaffected (see Materials and Methods). Whereas controls had uninterrupted strings of Mef2-labeled CBs (**Figure 2R**), the manipulated embryos had significant dorsal vessel ablation, evident by gaps in CBs (**Figure 2S**, bracket). We elected to analyze stage 14 and 15 embryos because these presented with appreciable ablation over larger distances compared to stage 16 onwards. Oftentimes both sides of the bilaterally symmetric dorsal vessel were ablated; sometimes only one side was ablated. For a given instance of dorsal vessel ablation, we documented whether the ipsilateral PSC was positioned normally. While the PSC occasionally appeared mis-positioned in non-ablated controls, for ablated dorsal vessel the ipsilateral PSC was mis-positioned more frequently (**Figure 2T**). These data suggest that dorsal vessel CBs influence PSC formation.

Slit-Robo signaling is required for proper PSC formation

To identify a cue necessary for positioning the PSC, we sought a signal that is expressed in early Vm and/or CBs. The secreted glycoprotein Slit is known to be expressed in both tissues and has been implicated as a positional cue in multiple developmental contexts (Anillo & DiNardo, 2022; Kidd et al., 1999; Kolesnikov & Beckendorf, 2005; MacMullin & Jacobs, 2006a; Rothberg et al., 1990). We confirmed that Slit is detectable in both Vm and CBs at relatively early stages of PSC migration (**Figure 3A** and **3B**). Importantly, *sli* mutants exhibited mis-positioned PSCs more frequently compared to controls (**Figure 3A** and **3B**, quantified in 3E), indicating a requirement for Slit in properly positioning the PSC.

To confirm that PSC cells can respond to Slit ligand, we examined the expression of the three Robo class receptors. We found evidence for only Robo1 and Robo2 expression in the PSC (**Figure S3A** and **S3B**). Thus, to test the requirement for Robo signaling, we analyzed single and double *robo1* and *robo2* mutants. For both single mutants, about 50% of embryos had abnormal PSC positioning (**Figure 3I** and **3J**, arrowheads). About 75% of double mutants had abnormal PSCs (**Figure 3K**), a frequency which matches that of *sli* mutants. These results demonstrate that Slit signaling through Robo1 and Robo2 is required for properly positioning the PSC.

Dorsal vessel-derived Slit signaling is required for PSC positioning

To test whether CBs were a source of Slit important for PSC positioning we used *tinC44-GAL4* to prevent either production or release of Slit specifically from dorsal vessel using RNAi or sequestration via overexpression of the Robo1 receptor, respectively. We confirmed that two independent Slit RNAi's eliminated nearly all Slit accumulation at CBs (**Figure 3O**' and **3Q**'), while Robo overexpression resulted in more pronounced Slit accumulation on CB membranes (**Figure 3S**') compared to control CBs (**Figure 3M**'). PSCs were abnormally positioned more frequently under all challenge conditions (**Figure 3N**, **3P**, **3R**) compared to controls (**Figure 3S**). Thus, Slit produced from CBs is essential to properly position the PSC.

Vm-derived Slit signaling is required for PSC positioning

The additional requirement for Vm in PSC positioning and the presence of Slit expression in Vm suggested that Vm-derived Slit might also regulate PSC formation. To block Slit produced by Vm from signaling to PSCs, we sequestered Slit specifically in Vm using *bap-GAL4* to overexpress Robo1. This manipulation prevents diffusion of Slit away from the tissue, as evidenced by distinct accumulation of Slit on Vm membranes (**Figure 3Z**) compared to diffuse Slit puncta in control Vm (**Figure 3W**). Under this condition, we indeed found that PSCs were mis-positioned significantly more often (**Figure 3AA**; quantified in **3BB**). Taken together, these data suggest that Vm and CB-derived Slit signaling are both required for PSC positioning.




Figure 3.

Slit from Vm and CBs signals through Robo for PSC positioning.

All PSCs, co-labeled by Antp and Odd, are viewed dorsally from st16 or 17 embryos. Normally coalesced and positioned PSC cells are outlined in yellow; dispersed PSC cells indicated by cyan arrowheads **A**) Schematic of dorsal st14 embryo. Relevant tissues are indicated; Vm is ventral to the rest. **B**) Slit expression in st14 CBs, blue outline, and Vm, orange outline; these tissues are in reversed orientation from the schematic due to embryo tilt, and the projection necessary to have both tissues in view. **C-E**) Analysis of PSC positioning in *slit²* heterozygotes (**C**, normal PSCs) and *slit²* mutants (**D**, dispersed PSCs), with frequency of positioning phenotype quantified in **E**. **F-L**) Analysis of PSC positioning under various *robo* depletion scenarios. *robo1* heterozygote (**F**) and *robo2* heterozygote (**G**) with normal PSCs. **H**) *robo1, robo2* double heterozygote with abnormal PSCs. *robo1* single (**I**), *robo2* single (**J**), and *robo1, robo2* double (**K**) mutants with abnormal PSCs; frequency of positioning phenotype quantified in **L** ('H' is heterozygote, 'M' is mutant). **M-U**) Analysis of PSC positioning when Slit signaling from CBs is compromised; frequency of PSC positioning phenotype quantified in **U**. **M, O, Q, S**) Slit expression in Mef2 labeled cardioblasts of controls (**M**) and *tinCΔ4*-GAL4 driven Slit RNAi knockdown (**O** and **Q**) or Robo1 overexpression (**S**) embryos. **M', O', Q', S')** Slit channel only. **N**) Control with normal PSCs. **P, R, T**) Dispersed PSCs upon compromised Slit signaling from CBs. **V-BB**) Analysis of PSC positioning when Slit signaling from Vm is compromised; frequency of PSC positioning phenotype quantified in **BB**. **V** and **Y**) Schematics of dorsolateral st14 control (**V**) or *bap*-GAL4 driven Robo1 overexpression (**Y**) embryos with relevant tissues and proteins indicated. **W**) Diffuse Slit expression in Fas3-labeled Vm of st14 control embryo. **Z**) Slit trapping at Fas3-labeled Vm membranes in *bap*-GAL4 driven Robo1 overexpression embryo. **W' and Z')** Slit channel only. **X**) Normally positioned PSCs and Antp+ CBs (brackets) in control. **AA**) Abnormally positioned PSC and Antp+ CBs (brackets) in *bap*-GAL4 driven Robo1 overexpression embryo. Scale bars as indicated. **BB**) Analysis of PSC positioning when Slit signaling from Vm is compromised; frequency of PSC positioning phenotype quantified in **BB**. **V** and **Y**) Schematics of dorsolateral st14 control (**V**) or *bap*-GAL4 driven Robo1 overexpression (**Y**) embryos with relevant tissues and proteins indicated. **W**) Diffuse Slit expression in Fas3-labeled Vm of st14 control embryo. **Z**) Slit trapping at Fas3-labeled Vm membranes in *bap*-GAL4 driven Robo1 overexpression embryo. **W' and Z')** Slit channel only. **X**) Normally positioned PSCs and Antp+ CBs (brackets) in control. **AA**) Abnormally positioned PSC and Antp+ CBs (brackets) in *bap*-GAL4 driven Robo1 overexpression embryo. Scale bars as indicated. **BB**) Analysis of PSC positioning when Slit signaling from Vm is compromised; frequency of PSC positioning phenotype quantified in **BB**.

ns = not significant, **p*<0.05, ***p*<0.01, ****p*<0.001, *****p*<0.0001, Fisher's Exact test. Sample sizes as indicated.

Vm influences dorsal vessel positioning

While the data thus far supported a simple model wherein Slit secreted from Vm and CBs acts directly on Robo receptors expressed by PSC cells to influence PSC positioning, further investigation revealed a more complex situation. Autocrine Slit-Robo signaling among dorsal vessel cells is important for proper polarity and organization of the vessel (Medioni et al., 2008a [↗](#); Qian et al., 2005 [↗](#); Santiago-Martinez et al., 2008 [↗](#)). Intriguingly, in experiments where Slit was sequestered in Vm, we noticed defects in the organization of the cardioblasts (**Figure 3AA** [↗](#), brackets; compare to control, 3X brackets). This suggested that signaling from Vm was required for organizing the dorsal vessel—a possibility that had not been previously explored. To test this, we examined the dorsal vessels of *bin* mutants which lack Vm (**Figure 2H** [↗](#)). We identified multiple phenotypes, two of which we termed ‘twisted’ and ‘sunken’; the dorsal vessel in **Figure 4B** [↗](#) exhibits both phenotypes. We scored a dorsal vessel as ‘twisted’ (**Figure 4B** [↗](#), bracket) when there was an apparent kink in the vessel such that the left or right side sat directly atop the other; thus, when viewing a single z-slice, a span of contralateral CBs appear to be missing since they are displaced above or below that focal plane. We defined a sunken dorsal vessel (**Figure 4B** [↗](#)’ and 4B”) as those cases where the whole dorsal vessel, or part of it, was displaced ventrally, deeper into the embryo. Such a case is evident in the **Figure 4B** [↗](#) mutant dorsal vessel, which spans 8μm of depth, compared to the sibling control dorsal vessel, which spans only 4μm. Both phenotypes were significantly more frequent in *bin* mutants compared to sibling controls (**Figure 4C** [↗](#)). Together, the dorsal vessel defects observed upon removal of the Vm or from trapping Slit on Vm cells reveal a previously unrecognized role for Vm in dorsal vessel formation.




Figure 4.

PSC positioning requires Vm-mediated dorsal vessel organization and Robo activation in PSC cells.

A-C) Analysis of dorsal vessels with and without Vm; phenotype frequencies quantified in (C). A and B) Mef2-labeled CBs and Slit-labeled lumens of dorsal vessels. A) dorsal vessel of sibling control, captured in 4.0µm projection, shows normal, neatly-aligned CB organization. B) *bin^{R22}* mutant with twisted dorsal vessel (bracket), fully captured only by projection of an 8.0µm deep stack. B') twisted phenotype evidenced by the absence of the sunken portion in the first 3.5µm; B'') 3.5-8µm projection. D) Schematics depicting three expectations for PSC and CB positioning. In controls (left), PSCs are coalesced and adjacent to CBs. In *robo2*, *robo1* mutants, either PSC cells are associated with mis-positioned CBs (middle), suggesting passive displacement of the PSCs; or displaced PSC cells are sometimes separate from mis-positioned CBs (right), suggesting the PSC cells themselves require activated Robo signaling. E-I) Analysis of CB and PSC positioning with additional CB marker, *svp-LacZ*. Brackets indicate normally-positioned CBs. Yellow outlines indicate normally-positioned PSC cells. Arrowheads indicate abnormally-positioned PSC cells; green arrowheads indicate passive mis-positioning and magenta arrowheads are displaced PSC cells without a similarly displaced CB nearby. Arrows indicate the next set of *svp+* CBs to the posterior. E) Control with normally-positioned PSCs and CBs; second set of *svp+* CBs (arrow) nearby. F) *robo2*, *robo1* mutant with abnormal CB positioning; Antp+ CBs are dispersed and the second set of *svp+* CBs (arrow) are displaced far posteriorly. PSCs display both passive mis-positioning posteriorly (green arrowheads) and mis-positioning without a similarly mispositioned CB nearby (magenta arrowheads; bottom PSC cell invaded midline and top PSC cell is displaced laterally). G) *robo2*, *robo1* mutant with left side CBs positioned normally (brackets) and the second set of *svp+* CBs in view (arrow); despite normal CB positioning, the ipsilateral PSC has a mis-positioned cell (magenta arrowhead). Right side CBs positioned abnormally (an Antp+ CB is displaced laterally and the second set of *svp+* CBs are displaced posteriorly, not in view); correspondingly, the ipsilateral PSC has two laterally displaced cells (green arrowheads). H) Frequency of mis-positioning of both a PSC and the ipsilateral CBs; this occurs more frequently in *robo2*, *robo1* mutants than heterozygotes. I) Frequency of correlated mis-positioning of PSCs and CBs. Both *robo2*, *robo1* heterozygotes and mutants have CB-independent instances of PSC mis-positioning. Sample sizes as indicated. **p<0.01, ****p<0.0001, Fisher's Exact test. Scale bars as indicated.

PSC positioning requires Robo signaling in CBs and in PSCs

Our results indicated that lack of Vm causes abnormal dorsal vessel positioning, and that the dorsal vessel is required for PSC positioning. In addition, our live-imaging revealed that some PSC cells are affixed to CBs of the dorsal vessel (Figure 1). Thus, it became important to test whether all PSC phenotypes might be 'passive', explained by PSC attachment to a malforming dorsal vessel. Alternatively, the PSC defects could reflect a requirement for Robo activation directly in PSC cells. Without a PSC-specific driver to directly test the latter, we re-examined *robo1*, *robo2* double mutants, this time including an additional marker that allowed us to observe PSC and CB defects independently and score for correlation between the two. If defective PSC positioning mostly correlated with defective CB positioning this would suggest a passive effect on PSCs by CBs of the dorsal vessel (Figure 4D, middle schematic). By contrast, if PSC cells were mis-positioned without a similarly mis-positioned CB nearby, this would suggest a requirement for intrinsic activation of Robo in PSC cells (Figure 4D, rightmost schematic).

The CB marker, *svp-lacZ*, labels two CBs in each hemisegment (Figure 4E, two hemisegments shown; Lo & Frasch, 2001). Normally, the PSC is compact and located adjacent to the compact Antp+ CBs (Figure 4E, PSCs outlined). *Svp* co-labels the last of the four strongly Antp+ CBs, and it labels the immediately posterior Antp-CB (Figure 4E, brackets). At the end of migration, *Svp-LacZ*+ CBs, like all CBs, are well-aligned with their bilaterally symmetric counterparts. Thus,

including Svp-LacZ in our analysis served as a useful registration marker that afforded the ability to detect bilateral matching, as well as shifting of CBs or the PSC along the anterior-posterior axis (**Figure 4E–4G**, arrows indicate second set of Svp+ CBs).

In *robo1*, *robo2* double mutants, we found that mis-positioning of CBs and the ipsilateral PSC occurred more frequently compared to double heterozygotes (**Figure 4H**). We then further analyzed those instances of mis-positioning for whether the position of PSCs correlated with position of CBs. In about two-thirds of cases mis-positioning was correlated (**Figure 4F** and **4G**, green arrowheads, quantified in 4I), suggesting that proper PSC positioning relies in part on proper CB positioning. However, in about one-third of cases PSC mis-positioning appeared independent of CB mis-positioning (**Figure 4F** and **4G**, magenta arrowheads). We also observed PSC mis-positioning (**Figure 4G**, magenta arrowhead) when ipsilateral CBs were positioned normally (**Figure 4G**, brackets). These instances of non-correlation strongly suggest that Robo activation is required in PSC cells for their proper positioning.

Finally, we live-imaged *slit* mutants with Hand-RFP, Antp>myrGFP to examine the dynamics of PSC migration when signaling was compromised. In contrast to the robust collectivity noted throughout the migration of control PSCs (**Figure 1G**), this imaging revealed progressive deterioration of PSC integrity during migration. For instance, throughout imaging, one PSC cell (**Figure 5A**, cyan dot) bridged the gap between the Antp+ CBs (**Figure 5A**, yellow brackets, 0 min) and posteriorly-displaced CBs. Ultimately this PSC cell made aberrant contact with a CB from the contralateral side (**Figure 5A**, yellow dot, 70min). This region had a persistent gap, possibly reflecting improper sealing of the dorsal vessel (**Figure 5A**, arrows, 112 min), as well as improper A-P alignment of Antp+ CBs with their contralateral counterparts (**Figure 5A**, offset brackets, 112 min). Another PSC cell (**Figure 5A**, cyan arrowhead, 0 – 35min) extended long protrusions along the peripheral edge of the PSC and was cleared by a macrophage (**Figure 5A**, yellow arrowhead, 42 – 49min). At imaging onset in the same embryo, 2 PSC cells were laterally displaced (**Figure 5B**, arrowheads, 0 – 28min) but connected to the main PSC collective by a thin protrusion (**Figure 5B**, green arrow, 0min). These 2 cells detached from the collective (7-28min), underwent cell shape changes (elongated at 42-56min, brackets; highly protrusive at 77min), and remained relatively stationary as the rest of the PSC and CBs migrated away (out of this focal plane).

Altogether, these data demonstrate that proper PSC formation requires 1) Slit signaling from the Vm and dorsal vessel where signaling is necessary for proper alignment and organization of the dorsal vessel, which, in turn, affects positioning of attached PSC cells; and 2) Robo activation in PSC cells for their persistent association with one another as a collective (**Figure 6**).

Discussion

This is the first work to examine how the PSC is positioned. We describe the steady-state, coalesced positioning of the cells comprising the PSC, characterize their migration, and reveal the signaling requirements that facilitate their association and recruitment to the lymph gland. Altogether, the evidence we provide culminates in a model in which the PSC migrates to the dorsal midline as a collective of cells—some adhere to the dorsal vessel and the rest adhere to each other. We show that association of PSC cells requires input from Vm and dorsal vessel, and we implicate Slit as a necessary signal. Though we have not ruled out direct Vm-to-PSC signaling, we find at a minimum that Vm affects PSC positioning indirectly by its novel role in forming the dorsal vessel. The intricate regulation described herein ensures that PSC cells achieve a precise steady-state positioning as a coalesced group at the lymph gland posterior.




Figure 5.

During migration *Slit* is required for proper PSC adhesions to CBs and for association of the PSC collective.

A and B) timelapse imaging of *sli* mutant embryo with LG and PSC labeled by Hand-RFP, Antp>myrGFP; dorsolateral view. A) The line of GFP+ cells within the brackets are Antp+ CBs; GFP+ cells below this are Antp+ PSC cells. Elongated PSC cell (cyan arrowhead; 0-35') cleared by macrophage (yellow arrowhead; 42-49'). Anterior and posterior edges of another PSC cell (cyan dot) bridges separated CBs (0-70'); same PSC cell aberrantly contacts a contralateral Antp+ CB (yellow dot; 49-70'). Persisting gap in CBs (arrows; 112') evident in same region. A') just GFP channel. Brackets indicate mis-aligned contralateral Antp+ CBs. B) Cells within the yellow brackets are Antp+ CBs. Same PSC as (A) contains two laterally displaced PSC cells (cyan arrowheads; 0-28') barely attached (green arrow; 0') to main cluster. These cells are disconnected from the main cluster (7') and remain stationary as the other PSC cells and CBs migrate away (7-77'). The disconnected PSC cell(s) change shape (cyan brackets) and develop membranous spikes (77' arrowhead). Scale bars as indicated.




Figure 6.

Model of PSC formation.

Collective cell migration of the PSC

Prior knowledge of the embryonic PSC has been focused on PSC cell specification (Crozatier et al., 2004 [DOI](#); Mandal et al., 2007 [DOI](#)). Thus, our live-imaging provides a substantial advancement by revealing PSC dynamics after specification, during migration. PSC formation is an example of collective cell migration, reminiscent of border cell migration in the *Drosophila* egg chamber and migration of the lateral line primordium in *Danio rerio*. These cell collectives respond to guidance cues, and their constituent cells are highly protrusive and shift position within the collective (Cliffe et al., 2017 [DOI](#); Dalle Nogare et al., 2020 [DOI](#); Haas & Gilmour, 2006 [DOI](#); Peercy & Starz-Gaiano, 2020 [DOI](#)). PSC migration is no different (**Figure 1** [DOI](#))—we observe extension and retraction of protrusions on all surfaces of the collective except those in contact with CBs. This suggests that the migrating PSC explores and responds to its environment. Some cells of the PSC maintain their position adjacent to the dorsal vessel throughout the entirety of migration, while other cells exhibit fluidity within the collective. The positional shifts suggest remodeling of adhesive contacts between cells—albeit in a coordinated manner to achieve unidirectional migration for the entire cohort. Interestingly, circulating hemocytes can be seen interacting with the PSC—hemocytes are known to deposit and remodel ECM (Bunt et al., 2010 [DOI](#)), which could serve as a scaffold for adhesion molecules and as a substrate during migration. Indeed, we identified ECM components surrounding the PSC and at its interface with the dorsal vessel. The characteristics noted above could be further explored to improve our understanding not only of how the PSC is built, but more broadly of collective cell migration.

PSC heterogeneity

For improved live-imaging we sought a PSC-specific marker but found most candidates to be expressed in only subsets of PSC cells (unpublished data). The lack of homogenous expression across all cells suggests PSC heterogeneity. The existence of fixed and fluid PSC cell positionings in the collective further hints at heterogeneity—perhaps there are different types and polarities of adhesive molecules. If PSC heterogeneity exists at this early stage, its purpose is unknown. One possibility is that the fixed, dorsal vessel-adjacent PSC cells act as “leaders” while the more fluid PSC cells are “followers”, a common phenomenon in collective cell migration (Qin et al., 2021 [DOI](#)). The dorsal vessel has recently been shown to serve as a second hematopoietic niche component (Destalminil-Letourneau et al., 2021 [DOI](#); Tian et al., 2023 [DOI](#)), so another possibility is that the fixed, dorsal vessel-adjacent PSC cells have a special function as intermediaries that coordinate communication between the niche components. That said, whether the niche components interact remains unknown. Finally, PSC heterogeneity in this early context could reflect a division of labor later, when the PSC regulates larval hematopoietic progenitors. Gene expression heterogeneity has been noted in the testis niche (Le Bras & Van Doren, 2006 [DOI](#); Raz et al., 2023 [DOI](#); Zheng et al., 2011 [DOI](#)), suggesting that this may be a conserved feature of niches which warrants further investigation. Our findings reported herein are foundational for future studies of PSC heterogeneity.

Coordinated input of extrinsic signals positions the PSC

We reveal that positioning the PSC requires input from different extrinsic sources. Organogenesis commonly relies on organ-extrinsic inputs. Like the PSC, the *Drosophila* excretory tubules require multiple inputs to navigate a complex signaling environment and achieve their final, precise positioning. To traverse the anterior-posterior axis, the tubules rely on alary muscles for posterior anchoring and on tissue sources located progressively more anteriorly for attractive BMP-dependent signaling (Weavers & Skaer, 2013 [DOI](#)). It is perhaps notable that the few well-studied examples of niche formation all involve extrinsic inputs. The mammalian intestine is organized into protrusive villi and invaginated crypts; Wnt and EphB3 signals emitted from the crypt base recruit niche cells from the crypt-villus interface to their final position in the compact troughs of the crypts (Batlle et al., 2002 [DOI](#); Holmberg et al., 2006 [DOI](#); van Es et al., 2005 [DOI](#)). The *C. elegans*

gonadal niche cell relies on adjacent germ cell proliferation for propulsion towards its steady-state position at the apex of the gonad (Agarwal et al., 2022). A microniche of the mammalian hair follicle, the dermal papilla, originates via extrinsic Fgf20 signaling to dermal fibroblasts. The fibroblasts are recruited into a condensate via directed migration, and then the condensate is segregated deeper into the skin via epidermal invagination driven by reciprocal signaling between dermis and epidermis (Biggs et al., 2018). The *Drosophila* testis niche cells require FGF and Slit signals from Vm to migrate through the testis and assemble as a cap that is further compacted via actomyosin contractility (Anllo & DiNardo, 2022; Warder et al., 2024). Together with our findings on PSC formation, a paradigm for niche formation is emerging that, subsequent to niche cell specification, extrinsic input segregates niche cells from other constituents, and positions the collective toward one end of the tissue. Oftentimes positioning is accompanied by or precedes compaction of the recruited cells into a final niche shape/architecture.

We show that extrinsic Slit-Robo signaling is integral in establishing PSC position. Removing Slit causes PSC defects at the same frequency as removal of both canonical Robo receptors. This argues that in PSC formation, as in the gonadal niche and dorsal vessel, Slit only signals through Robo, and not through the non-canonical receptor, Dscam1 (Anllo & DiNardo, 2022; Qian et al., 2005). We noted that the PSC positioning phenotype was about 75% penetrant, and the severity of the defects varied. These observations argue that PSC formation relies on an additional cue(s) yet to be discovered. In fact, our lab showed in addition to Slit, gonadal niche formation requires the FGF ligand, *pyr*, and its receptor, *htl* (Anllo & DiNardo, 2022).

. Evidence for an additional PSC coalescence cue exists in later larval stages, as manipulated Insulin Receptor signaling was shown to disrupt PSC coalescence (Tokusumi et al., 2015). Unfortunately, most pathways are difficult to test for a role in PSC formation due to lack of an embryonic PSC-specific tool with the necessary temporal control.

Haploinsufficiency of Slit and Robo in PSC positioning

In the nervous system, one copy of the normal (WT) allele of Slit or Robo are typically sufficient for proper development (Kidd et al., 1999; Rothberg et al., 1988). In contrast, we find that about half of *robo2, robo1* double heterozygotes or of *slit* heterozygotes have abnormal PSCs, suggesting haploinsufficiency in this process. This observation for embryonic stages is supported by previous findings at later larval stages where both *slit* or *robo2* heterozygotes have dispersed PSCs (Morin-Poulard et al., 2016). Perhaps haploinsufficiency in the PSC reflects different regulation and function of the Slit-Robo pathway from the nerve cord.

Alternatively, because we identified that Slit-Robo signaling is required within both CBs and PSC for proper development of both tissues, the haploinsufficiency may reflect a combinatorial defect caused by diminished signaling in both tissues. Furthermore, we are surprised that *robo2* and *robo1* single mutants have a similar frequency of abnormal PSCs because in the embryo Robo1 is expressed in all PSC cells whereas Robo2 appears to be expressed in only one PSC cell. However, it is possible that all PSC cells express Robo2 and our finding reflects limited detection ability with the Robo2-GFP reporter. An alternative, intriguing possibility is that the Robo2-expressing PSC cell is the “leader” in PSC migration, and without Robo2 signaling in the leader, the collective fails to migrate appropriately, or the followers fail to remain properly adherent.

Morin-Poulard et al. show that all larval PSC cells express Robo2, and it has a more prominent role than Robo1 in maintaining PSC coalescence (termed ‘clustering’ in their work) during larval stages (Morin-Poulard et al., 2016). In fact, they found Robo1 knockdown alone was insufficient to disrupt clustering. Taken together with our work, perhaps this indicates that Robo1 is more important for first establishing PSC coalescence and Robo2 is more important for maintaining it. Most manipulations occurred after the PSC had already formed, and therefore the changes reported cannot stem from defects during establishment but rather from dispersion of the cluster after it formed. One experiment without temperature control that could have diminished Robo

signaling in PSC and CBs early (Antp>Robo RNAi), at about the time of PSC specification, generated a more dispersed late larval PSC compared to Robo knockdown subsequent to PSC formation. One can imagine that aberrant embryonic PSC positioning would become exacerbated and increasingly catastrophic as the PSC and nearby progenitors proliferate throughout larval stages. The fact that earlier Robo knockdown worsened the larval PSC defect supports our finding that Slit-Robo signaling is vital to establish PSC coalescence.

Function of Slit-Robo in PSC positioning

Determining how Slit affects a given process is complex because it can act as an attractive or repulsive cue, or even affect cell adhesion, and in some instances the outcome relies on whether Slit is cleaved by Tok (Englund et al., 2002 [↗](#); Kellermeyer et al., 2020 [↗](#); Kolesnikov & Beckendorf, 2005 [↗](#); Kramer et al., 2001 [↗](#)). Thus, elucidating the role of the Slit-Robo pathway in PSC formation is difficult. Upon compromising the pathway, we observed PSC mis-positioning in all directions. Also, we found that Slit is required in both Vm and dorsal vessel, and the role for each tissue in PSC formation is intertwined. The existence and relative positioning of two Slit sources, the reliance on one for positioning of the other, and the inconsistent direction of PSC mis-positioning suggest that it is too simplistic to ask whether Slit is acting as a repulsive or attractive cue for PSC cells. Notably, mis-positioned PSC cells adopted aberrant contacts with CBs, LG, and pericardial cells of the dorsal vessel. When reported in the dorsal vessel, this inappropriate mixing of cells was attributed to improper polarity of cell adhesion molecules (Medioni et al., 2008b [↗](#); Qian et al., 2005 [↗](#); Santiago-Martínez et al., 2006 [↗](#)). Furthermore, Slit has been shown to be important for maintaining adhesion amongst larval PSC cells (Morin-Poulard et al., 2016 [↗](#)). Taken together with our data, we reason that during PSC migration Slit must affect the ability of PSC cells to properly adhere to the dorsal vessel and to one another. In this manner, Slit facilitates PSC cell association as a compact cluster.

The various effects we have documented for Slit signaling indicate that downstream regulation is likely complex and dependent on the specific context, in time and space. Morin-Poulard et al. show that the Slit-Robo pathway maintains clustering of larval PSC cells via DE-Cadherin and Cdc42 (Morin-Poulard et al., 2016 [↗](#)); however, our data indicate different effectors are at play during PSC formation, as we detect no DE-Cad in the embryonic PSC (data not shown). Additionally, whereas constitutively active Cdc42 in the larval PSC caused dispersion (Morin-Poulard et al., 2016 [↗](#)), our attempt to elicit PSC dispersion with this manipulation in the embryo yielded normally coalesced PSCs (data not shown). Therefore, we reason that different downstream effectors are engaged by the Slit-Robo pathway for initial positioning of the PSC.

This study was limited by the lack of a truly PSC-specific driver—a tool that would improve live-imaging and allow testing of potential Slit-Robo pathway effectors. We and others observe Fasciclin III, a homotypic cell adhesion molecule, in the PSC (unpublished data and Mandal et al., 2007), however, it is first detectable too late in embryogenesis to function during PSC formation. Thus, the particular adhesion molecules that maintain association of PSC cells as they migrate remain unknown. Elucidating this tool will facilitate interrogation of how Slit-Robo signaling impacts PSC adhesion to the dorsal vessel and amongst itself.

Passive and active roles for Slit-Robo signaling in PSC positioning

Due to the necessity of autocrine Slit-Robo signaling in dorsal vessel formation (MacMullin & Jacobs, 2006b [↗](#); Medioni et al., 2008a [↗](#); Qian et al., 2005 [↗](#); Santiago-Martínez et al., 2006 [↗](#), 2008 [↗](#)), it was challenging to distinguish whether effects on the PSC were secondary to mis-positioned CBs, or primary consequences of diminished Robo activation in PSC cells. Our analysis of *robo2, robo1* double mutants along with the registration marker, *svp-lacZ*, strongly suggests both passive mis-positioning of PSC cells via attachment to mis-positioned CBs and a requirement for Robo activation directly in PSC cells. A direct requirement for Robo in PSC cells is further suggested in *slit* mutant live-imaging. Here, the lateral-most PSC cells separated from the main

cluster even though no CBs were nearby (**Figure 5B** [↗](#)). Perhaps during normal migration these more ‘fluid’, lateral PSC cells possess a stronger requirement for Robo activation and require direct input to achieve that. In this view, those PSC cells that are passively mis-positioned in mutants are the same PSC cells that remain affixed to the dorsal vessel during normal migration. Alternatively, or additionally, passive mis-positioning of CB-adjacent PSC cells might occur due to improper polarization of adhesive molecules, whereas separation of lateral PSC cells from the collective might occur due to a total loss of adhesive molecules.

Identification of novel mechanisms controlling dorsal vessel development

Our investigation of PSC formation unexpectedly revealed new insights about the dorsal vessel. We uncover a novel role for Vm in positioning the dorsal vessel. In Vm manipulations we noted abnormal positioning of Antp+ CBs, and upon further investigation of mutants lacking Vm, we show that large portions of the dorsal vessel are malformed. The twisting phenotype we observe suggests luminal collapse, and, thus, Vm may contribute to lumen formation. Alternatively, Vm could affect CBs at the time of their specification. Both Vm and CBs originate from dorsal mesoderm, and we learned that Vm is able to act at this early timepoint via our finding that early Vm signals influence PSC positioning. Additionally, we were surprised in dorsal vessel ablation experiments to observe more apparent ablation in mid-stage embryos than late-stage. Note that the CBs should be post-mitotic ([Gajewski et al., 2000](#) [↗](#); [Han & Bodmer, 2003](#) [↗](#); [Ward & Skeath, 2000](#) [↗](#)). Thus, under challenge conditions CBs fill the gaps by either re-entering the cell cycle to produce new daughters, or, more likely, by elongating to fill gaps. Regardless of mechanism, this observation suggests plasticity of the dorsal vessel upon challenge.

Connecting niche structure, position, and function

The coalesced nature and posterior positioning of the PSC are its most prominent features. The consistency of these features suggested that they are under tight regulation, as we report here, and strongly indicates their relevance to function of the PSC. Though we have yet linked the precise architecture of the embryonic PSC to function, the specific architectures of multiple other niches have been shown to be functionally relevant. In the *Drosophila* gonad, a coalesced and compact niche is necessary for proper germline stem cell maintenance, niche signaling, and for orienting characteristically orthogonal germline stem cell divisions ([Anllo & DiNardo, 2022](#) [↗](#); [Warder et al., 2024](#) [↗](#)). In the mammalian hair follicle, niche position determines stem cell fate ([Rompolas et al., 2013](#) [↗](#)).

Drosophila neural stem cells are supported by glia, and the precise morphology of this encasing niche is required for newborn neuronal survival, and therefore homeostasis of the entire nerve cord ([Spéder & Brand, 2018](#) [↗](#)). Most pertinently, recent work has shown that the larval PSC communicates amongst itself via a calcium signaling network. The integrity of this network is gap junction-dependent and is required for the PSC to emit proper levels of a progenitor maintenance signal ([Ho et al., 2021](#) [↗](#), [2023](#) [↗](#)). We find it reasonable to postulate that a dispersed PSC would exhibit defective calcium signaling such that the PSC would lack coordinated maintenance of progenitors, thereby leading to an imbalance in the ratio of progenitors to differentiated hemocytes in the gland. Future experiments will determine how PSC coalescence is functionally relevant to its regulation of hematopoietic progenitors.

Acknowledgements

We thank the Bloomington *Drosophila* Stock Center (NIH P40OD018537), J. Skeath, J. Reinitz, E. Furlong, G. Bashaw, M. Crozatier, U. Banerjee, M. Frasch, G. Vogler, and DSHB for antibodies and stocks. We thank CDB microscopy core director Andrea Stout for advice on imaging. Thanks to B.




Figure S1.

ECM components surround the PSC and AM encases PSC cells.

A and B) ECM components Perlecan and Viking (Collagen IV subunit) surround migrating PSCs. A) *perlecan*-GFP reporter, white, surrounds migrating Antp-labeled PSC cells, cyan arrowheads, of st14 embryo. B) *viking*-GFP reporter, white, surrounds migrating PSC collective on embryo left side, cyan outline, and a single PSC cell on the right side, arrowhead. A' and B') GFP channel only. C-E) *org-1*-RFP embryos with PSCs labeled by Antp, yellow brackets or arrowheads. C) Alary muscle of first abdominal segment (A1 AM), cyan, contacts posterior of PSC collective in st15 embryo. D) A1 AMs contact entire PSC collectives of st17 embryo. E) st16 embryo with individual A1 AM fibers encasing two PSC cells, arrowheads. E') RFP channel alone. Scale bars as indicated.




Figure S2.

All PSC positioning analyses were completed on st16 or st17 embryos by co-labeling with Antp and Odd.

A) Quantitation of PSC cells in controls and AM-ablated embryos. Not significant (ns), Mann-Whitney test. B) lateral view of st13 embryo stained for Antp (white) and Bin (red). Antp labels PSC collective (arrowhead) and Bin is expressed in Vm (brackets), as expected. B') Magnified view of Antp+ collective (outline) shows no detectable Bin. B'') Bin channel only; PSC collective outlined. C and D) Fas3-labeled Vm (white) of st11 control (C) and FCM-ablated (D) embryos also stained for Bin (red). E) Bin expression is reduced in remaining FCs of Vm-ablated embryos. ****p<0.0001, Mann-Whitney test. F-K) Analysis of *bap*²⁰⁸ sibling controls (F-H) and hypomorphic mutants (I-K). F and I) lateral view of st11 Fas3-labeled Vm precursors of control (F) and mutant (I), which has almost normal Vm organization with occasional gaps. G) dorsolateral view of Fas3 stained st16 sibling control embryo with normal Vm morphology. H) Sibling control LGs with normal PSCs. J) Dorsolateral view of Fas3 stained late stage mutant embryo with disorganized Vm that is missing in the midline region where the PSC is normally located. K) *bap*²⁰⁸ mutant LGs with normal PSCs. L) Quantification of frequency of PSC positioning phenotype in *bap*²⁰⁸ sibling controls and mutants; ns, Fisher's Exact test. M-Q) analysis of PSC positioning for various *jeb* alleles; frequency of phenotypes quantified in Q (ns, Fisher's Exact test). M and N) LGs with normal PSC positioning from *jeb*^{weli} heterozygous (M) and homozygous (N) mutant. O and P) LG with normal PSC positioning from a sibling control (O) and a *jeb*^{weli}/*jeb* Df transheterozygous mutant (P). Scale bars as indicated. Sample sizes as indicated.




Figure S3.

Robo1 and Robo2 are expressed by PSC cells.

A) Robo1 expression (cyan) on PSC cell membranes (arrowheads) of st14 embryo stained for Antp; brackets are two of the eventual 4 Antp+ CBs. A') Robo1 channel alone. B) *robo2*-GFP reporter expression (cyan) on membrane of single PSC cell (arrowhead) of st15 embryo stained for Antp; brackets enclose Antp+ CBs. B') GFP channel alone.

Warder, G. Vida, G. Bashaw, and the K. Lenhart lab for input during this project. This work was supported by NIH grants T32 HD083185 and F31 HD111208 to KAN; R01 GM138705 to KFL; F32 GM125123 to LA; R35 GM136270 and R03 HD111973 to SD.

Materials and Methods

In this study we used FlyBase (releases 2020_06 - 2024_02; [Jenkins et al., 2022](#)) to find information on phenotypes, function, stocks, and gene expression. All data describe biological replicates. Each experiment was repeated at least once. The Mann-Whitney test was used for statistical comparison of two groups of unpaired numerical data with non-Gaussian distribution. Fisher's exact test was used for statistical comparison of two groups of categorical data. All corresponding data spreadsheets depicted by the graphs herein are available upon email request.

Drosophila genetics

Detailed information on the *Drosophila* strains used in this study is in the Key Resources Table. Controls were GAL4 only, a cross to *w¹¹¹⁸*, or a sibling control identified by a fluorescent balancer. This study generated the following embryo genotypes by combination into a stock (available upon email request) or obtained through a cross:

Hand-RFP, Antp-GAL4, UAS-myr:GFP
 UAS-CD8:GFP / org-1-HN39-RFP ; Antp-GAL4 / +
 UAS-CD8:GFP / + ; tupAME-GAL4 / +
 UAS-CD8:GFP / UAS-grim ; tupAME-GAL4 / TM6Hu or MKRS
bin^{R22} / +
bin^{R22} / *bin^{S4}*
 bap-GAL4 ; + / CyODfd-YFP ; bap-GAL4
 bap-GAL4 ; + / UAS-hid ; bap-GAL4
 UAS-grim ; tinCΔ4-GAL4
robo1^{GA285} / +
robo2¹ / +
robo1^{GA285}, *robo2¹²³* / +
 tinCΔ4-GAL4 / +
 UAS-sli RNAi / + ; tinCΔ4-GAL4 / UAS-dcr-2
 UAS-dcr-2 / + ; tinCΔ4-GAL4 / UAS-sli RNAi
 UAS-Robo1 OX / + ; tinCΔ4-GAL4 / +
 bap-GAL4 ; + / UAS-Robo1 OX ; bap-GAL4 / +
robo1^{GA285}, *robo2¹²³* / CyO, hindgut-LacZ ; *svp*-LacZ / TM6Dfd-GFP
sli² ; Hand-RFP, Antp-GAL4, UAS-myr:GFP
jeb^{weli} / +
jeb^{weli} / *jeb*[Df]

Embryo collection

Unless otherwise noted, embryos were collected on apple juice agar plates overnight at 25 degrees, unless an experiment involved a Gal4, in which case collections occurred at 29 degrees. The next morning embryos were dechorionated in 50% bleach.

Embryo fixation

Using a paintbrush, embryos were transferred from a collection basket to a 50/50 mixture by volume of heptane and 4% paraformaldehyde in Buffer B (16.7mM KPO₄, pH 6.8; 75mM KCl; 25mM NaCl; 3.3mM MgCl₂) (de Cuevas & Spradling, 1998 [\[4\]](#)). Embryos were fixed on a rocker for 15 minutes, then fixative was removed. An equal volume of MeOH was added and the vial was shaken vigorously for about 5 seconds to remove vitelline membranes. The heptane and MeOH mixture was discarded, and the embryos were rinsed three times with MeOH.

Embryo staging and genotyping

Embryos were staged according to Atlas of *Drosophila* Development (Hartenstein, 1993 [\[4\]](#)). Embryos were genotyped according to presence of a balancer chromosome: CyO, P{Dfd-EYFP}; or CyO, P{Wg-lacZ}; or CyO, hindgut-lacZ; or TM6, P{Dfd-EYFP}, Sb, Hu, e.

Immunostaining

Manipulations were room temperature unless otherwise noted. Embryos were rehydrated in 50%MeOH/50% PBS (10mM Na₂ HPO₄; 1.8mM KH₂PO₄; 2.7mM KCl; 137mM NaCl; pH 7.4), followed by 100% PBS. Embryos rocked for 5 minutes in PBS with 0.1% Triton X-100 (PBST), then 1 hour in 4% normal donkey serum in PBST.

Embryos were then transferred to a rocker at 4 degrees until the end of the day or began incubation in primary antibody solution. The next day embryos were rinsed 3 times in PBST then rocked in PBST for 1 hour, incubated in secondary antibody solution for 1 hour, rinsed 3 times in PBST, rocked in PBST for 1 hour, equilibrated in 50% glycerol/50% Ringer's solution (5mM HEPES, pH 7.3; 130mM NaCl; 5mM KCl; 2mM MgCl₂; 2mM CaCl₂) for 15 minutes or overnight at 4 degrees, then mounted in 2% nPropyl-gallate in 90% glycerol. Primary antibodies were diluted in normal donkey serum as follows: mouse antibody against Antp (1:50; DSHB, 8C11), rabbit antibody against Odd skipped (1:400; gift from James Skeath, Washington University School of Medicine) chick antibody against GFP (1:1500; Aves Labs, GFP-1020), rabbit antibody against RFP (1:1000; Abcam, ab62341), mouse antibody against Fasciclin 3 (1:50; DSHB, 7G10), rabbit antibody against Mef2 (1:1000; DSHB), guinea pig antibody against Odd skipped (1:1200; gift from John Reinitz, University of Chicago), rabbit antibody against Bin (1:100; gift from Eileen Furlong, EMBL), mouse antibody against Slit (1:200; gift from Greg Bashaw, University of Pennsylvania), mouse antibody against Robo1 (1:200; gift from Greg Bashaw, University of Pennsylvania), chick antibody against LacZ (1:1000; Abcam, ab9361-250). Secondary antibodies (Alexa Fluor 488, Cy3, Cy5, and Alexa Fluor 647; Molecular Probes or Jackson ImmunoResearch) were all used at 3.75ug/mL, for 1 hour.

Live-imaging

After dechorionation, embryos were selected by stage and transgene expression, using a stereofluorescent microscope, transferred to a piece of agar, and then oriented in a line with the ventral/ventrolateral surface facing up – hanging off the edge of the agar. A heptane-glue mixture was dried as a strip on a glass slide, and that surface touched to the embryos for transfer (now dorsal/dorsolateral surface of embryos faces up). 3uL of halocarbon oil was added atop the embryos. Bridging coverslips were glued to the slide on either side of the line of embryos, then the main coverslip was laid atop the bridging coverslips. We imaged every 5-10 minutes for 2-4.5 hours; Z-stacks spanned 25-40uM with 0.3-1.0uM step sizes.

Microscopy

Fixed embryos were imaged on a Zeiss Axio Imager with ApoTome using a 40x, 1.2 NA water immersion objective or a 20x, 0.8 NA objective; z-steps were 0.5 – 1.0 μ m. All live-imaging except for alary muscle imaging occurred with a CrestOptics X-Light V3 spinning disk confocal microscope using a 60x, 1.3NA silicone immersion objective; images were captured with two pco.edge 4.2 bi sCMOS cameras operated by VisiView (Visitron) software. Alary muscles were live-imaged with an IX7 Olympus spinning disk confocal using a 63x, NA 1.2 water immersion objective and captured with an EMCCD camera (Hamamatsu photonics, model C9100-13) controlled by MetaMorph software.

PSC positioning phenotypic characterization

We used immunostains for the accepted markers, Antp and Odd, to identify PSC cells. Sometimes images contained Antp⁺ and Odd⁺ cells in epidermal stripes anterior to the LG; these were not considered PSC cells. An embryo was scored as having ‘normal’ PSC positioning if both PSCs were 1) coalesced within one nuclear diameter of one another, 2) adjacent to the dorsal vessel, and 3) at the same dorsal-ventral position as the posterior-most cells of the lymph gland. Abnormal PSC positioning presented as a range of phenotypes including PSCs dispersed into multiple groups, and PSCs that were coalesced but not located at the LG posterior. Fisher’s Exact test was used for statistical analysis and $p < 0.05$ was considered statistically significant.

AM ablation PSC cell counting

We considered a cell to be a PSC cell if it was co-labeled by Antp and Odd and was not located in epidermis.

Number of PSC cells was recorded separately for the left and right PSCs of each embryo, and the total number of PSC cells per embryo was plotted. The Mann-Whitney test was used to determine a non-significant difference with $p > 0.05$.

Bin fluorescent intensity

For 9 sibling controls and 10 Vm-ablated embryos, a normalized Bin fluorescent intensity was calculated for 3 different regions of st11 Vm founder cells with Bin-stained nuclei and Fas3-stained membranes. For st11 embryos imaged laterally, Fas3 labels the 2 lateral surfaces of a given Vm founder cell membrane. ImageJ software was used to extract fluorescent intensity values for regions of interest including large non-positive background regions in the embryo, founder cell nuclei, and founder cell membranes. Bin intensity for a given region was calculated by normalizing the background subtracted Bin level of a single representative nucleus to the average background subtracted Fas3 level of at least the 2 lateral surfaces of that same cell; most often an average Fas3 level for a given region was calculated based on the membranes of 2-4 founder cells. The normalized Bin fluorescent intensity for a given region is plotted. The Mann-Whitney test was used for statistical analysis to determine a significant difference of $p < 0.05$.

Tissue ablations

AM ablation: the *tupAME-GAL4* driver was combined with *UAS-CD8:GFP* for AM visualization, and GFP-labeled AMs were clearly visible in st13 controls. This line was crossed to *UAS-grim* to generate AM-ablated embryos; AMs were ablated by st14.

CB ablation: flies with the *tinCΔ4-GAL4* driver were crossed to flies with either *UAS-grim* or *UAS-hid* and embryos were collected overnight at 29 degrees. Expression of either pro-apoptotic gene led to quite effective dorsal vessel ablation, but there were also defects in either Vm (for Grim) or in germband retraction (for Hid; data not shown), which could confound a PSC positioning

analysis. We circumvented this issue by lowering GAL4 activity—embryos from the cross to UAS-grim were collected at 25 degrees—which yielded relatively normal Vm but successfully ablated CBs beginning at st14.

Slit RNAi in dorsal vessel

Two independent UAS-controlled RNAi's targeting different regions of Exon 4 of Slit along with UAS-controlled Dcr2 were driven by tinCΔ4-GAL4. Knockdown was evident at st14, the same stage when Slit expression becomes discernable in control CBs.

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
genetic reagent (<i>D. melanogaster</i>)	Antp-GAL4	Emerald and Cohen, 2004	FLYB:FBal0155891	FlyBase symbol: GAL4 ^{Antp-21}
genetic reagent (<i>D. melanogaster</i>)	w ¹¹¹⁸	Bloomington <i>Drosophila</i> Stock Center	BDSC:3605; FLYB:FBal0018186; RRID:BDSC_3605	FlyBase symbol: w ¹¹¹⁸
genetic reagent (<i>D. melanogaster</i>)	Hand-RFP	other		Gift from Georg Vogler
genetic reagent (<i>D. melanogaster</i>)	UAS-myr::GFP	Bloomington <i>Drosophila</i> Stock Center	BDSC:32200; FLYB:FBti0131976; RRID:BDSC_32200	FlyBase symbol: P{10XUAS-IVS-myr::GFP}su(Hw)attP1
genetic reagent (<i>D. melanogaster</i>)	tupAME-GAL4	Bataillé et al., 2020	FLYB:FB _{tp} 0142468	FlyBase symbol: P{tup-GAL4.AME-R} Gift from J.L. Frendo
genetic reagent (<i>D. melanogaster</i>)	UAS-CD8::GFP	other		Gift from J.L. Frendo
genetic reagent (<i>D. melanogaster</i>)	org-1-HN39-RFP	Schaub and Frasch, 2015	FLYB:FBal0276776	FlyBase symbol: RFP ^{org-1.HN39}
genetic reagent (<i>D. melanogaster</i>)	UAS-grim	Hugo Bellen	FLYB:FBti0154788	Flybase symbol: Dmel\P{UAS-grim.Y}2

genetic reagent (<i>D. melanogaster</i>)	<i>bin</i> ^{R22}	Zaffran and Frasch, 2001	FLYB:FBal0043738	Flybase symbol: Dmel\bin ^{R22}
genetic reagent (<i>D. melanogaster</i>)	<i>bin</i> ^{S4}	Zaffran and Frasch, 2001	FLYB:FBal0043739	Flybase symbol: Dmel\bin ^{S4}
genetic reagent (<i>D. melanogaster</i>)	UAS-hid	Bloomington <i>Drosophila</i> Stock Center	BDSC:65403; FLYB:FBti0183136 RRID:BDSC_65403	Flybase symbol: Dmel\{P{UAS-hid.Z}2
genetic reagent (<i>D. melanogaster</i>)	bap-GAL4	Zaffran and Frasch, 2001	BDSC:91540; FLYB:FBti0214156	Flybase symbol: Dmel\{P{bap-GAL4.3}1.1
genetic reagent (<i>D. melanogaster</i>)	bap-GAL4	other		gift from Manfred Frasch; Chr: X
genetic reagent (<i>D. melanogaster</i>)	tinCΔ4-GAL4	Bloomington <i>Drosophila</i> Stock Center	BDSC:92965; FLYB:FBti0216630	Flybase symbol: Dmel\{P{tinC-Gal4.Δ4}12a
genetic reagent (<i>D. melanogaster</i>)	<i>slit</i> ²	Bloomington <i>Drosophila</i> Stock Center	BDSC:3266 FLYB:FBal0015700	Flybase symbol: Dmel\slit ²
genetic reagent (<i>D. melanogaster</i>)	<i>robo1</i> ^{GA285}	other	FLYB:FBal0032588	Gift from Greg Bashaw Flybase symbol: Dmel\robo1 ¹
genetic reagent (<i>D. melanogaster</i>)	<i>robo2</i> ¹	Rajagopalan and Dickinson, 2000	FLYB:FBal0121562	Gift from Greg Bashaw Flybase symbol: Dmel\robo2 ¹
genetic reagent (<i>D. melanogaster</i>)	<i>robo2</i> ¹²³	other	FLYB:FBal0123720	Gift from Greg Bashaw Flybase

				symbol:Dmel robo2 ^{X123}
genetic reagent (<i>D. melanogaster</i>)	UAS-Slit RNAi #1	Vienna <i>Drosophila</i> Stock Center	VDRC:v10885 3 FLYB:FBti015 9991	Flybase symbol: Dmel\{P{KK1 00803}\VIE- 260B
genetic reagent (<i>D. melanogaster</i>)	UAS-Slit RNAi #2	Bloomington <i>Drosophila</i> Stock Center	BDSC:31468 FLYB:FBal02 45521	Flybase symbol:Dmel slit ^{JF01229}
genetic reagent (<i>D. melanogaster</i>)	UAS-Robo1 OX	Evans and Bashaw, 2015	BDSC:97240 FLYB:FBal03 16479	Flybase symbol:Dmel robo1 $\Delta C.10\times UAS.Tag:$ HA, Tag:SS(wg)
genetic reagent (<i>D. melanogaster</i>)	UAS-dcr2	Bloomington <i>Drosophila</i> Stock Center	BDSC:24650 FLYB:FBti010 0275	Flybase symbol:Dmel P{UAS-Dcr- 2.D}2
genetic reagent (<i>D. melanogaster</i>)	UAS-dcr2	Bloomington <i>Drosophila</i> Stock Center	BDSC:24651 FLYB:FBti010 0276	Flybase symbol:Dmel P{UAS-Dcr- 2.D}10
genetic reagent (<i>D. melanogaster</i>)	svp-lacZ	Bloomington <i>Drosophila</i> Stock Center	BDSC:7314 FLYB:FBti000 2862	Flybase symbol: Dmel\{P{HZ}s vp ³
genetic reagent (<i>D. melanogaster</i>)	perlecan- GFP	Flytrap; GFP Protein Trap Database	FLYB:FBal02 43609	Flybase symbol:Dmel trol ^{ZCL1700}
genetic reagent (<i>D. melanogaster</i>)	viking-GFP	Buszczak and Spradling, 2007	FLYB:FBal02 11825	Flybase symbol:Dmel vkg ^{CC00791}
genetic reagent (<i>D. melanogaster</i>)	bap ²⁰⁸	Bloomington <i>Drosophila</i> Stock Center	BDSC:91539 FLYB:FBal00 34201	Flybase symbol:Dmel bap ²⁰⁸

genetic reagent (<i>D. melanogaster</i>)	<i>jeb</i> ^{weli}	Stute and Holz, 2004	FLYB:FBal0159133	Flybase symbol:Dmel\jeb ^{weli}
genetic reagent (<i>D. melanogaster</i>)	<i>jeb</i> Df	Bloomington <i>Drosophila</i> Stock Center	BDSC:26551 FLYB:FBab0045764	Flybase symbol:Df(2R)BSC699
genetic reagent (<i>D. melanogaster</i>)	robo2-GFP	Bloomington <i>Drosophila</i> Stock Center	BDSC:61774 FLYB:FBal0265307	Flybase symbol:Dmel\robo2 ^{MI04295}
antibody	anti-Antp (Mouse monoclonal)	Developmental Studies Hybridoma Bank	Cat#:8C11, RRID:AB_528083	IF(1:50)
antibody	anti-Odd skipped (Rabbit polyclonal)	Ward and Skeath, 2000		IF(1:400); gift from James Skeath
antibody	anti-GFP (Chick polyclonal)	Aves labs	Cat#:GFP-1020 RRID:AB_2307313	IF(1:1500)
antibody	anti-Fas3 (Mouse monoclonal)	Developmental Studies Hybridoma Bank	Cat#:7G10 RRID:AB_528238	IF(1:50)
antibody	anti-Mef2 (Rabbit polyclonal)	Developmental Studies Hybridoma Bank	Cat#:Mef2 RRID:AB_2892602	IF(1:1000)
antibody	anti-Slit (Mouse monoclonal)	Developmental Studies Hybridoma Bank	Cat#:C555.6D RRID:AB_528470	IF(1:200); gift from Greg Bashaw

antibody	anti-LacZ (Chick polyclonal)	Abcam	Cat#:ab9361 RRID:AB_307210	IF(1:1000)
antibody	anti-RFP (Rabbit polyclonal)	Abcam	Cat#:ab62341 RRID:AB_945213	IF(1:1000)
antibody	anti-Bin (Rabbit polyclonal)	other		IF(1:100); gift from Eileen Furlong
antibody	anti-Robo1 (Mouse monoclonal)	other		IF(1:200); gift from Greg Bashaw
antibody	anti-Odd skipped (Guinea pig polyclonal)	other		IF(1:1200); gift from John Reinitz
chemical compound, drug	Paraformaldehyde	Electron Microscopy Sciences	Cat#:15710	
chemical compound, drug	Propyl-gallate	Sigma Aldrich	PubChem Substance ID:24898394; SKU:P3130; CAS Number:121-79-9	
chemical compound, drug	Normal Donkey Serum	Jackson ImmunoResearch Labs Inc	Cat#:017-000-121 RRID:AB_2337258	
chemical compound, drug	Ringer's solution	other		Recipe from https://doi.org/10.1242/dev.125.15.2781

chemical compound, drug	Triton X-100	MilliporeSigma	CAS Number: 9036-19-5	
software, algorithm	FIJI	ImageJ	RRID:SCR_002285	http://fiji.sc
software, algorithm	Photoshop	Adobe	RRID:SCR_014199	https://www.adobe.com/products/photoshop.html
software, algorithm	Prism	Graphpad	RRID:SCR_002798	v9.0.0-v10.0.0
software, algorithm	Axio-Vision Imaging Software	Zeiss		v4.8.1
software, algorithm	VisiView	Visitron		
software, algorithm	Metamorph Microscopy Automation and Image Analysis Software	Leica		v7.8.40
other	63x / 1.2 NA water immersion objective	Leica		
other	60x / 1.3 NA silicone immersion objective	Olympus		
other	AxioCam HRm	Zeiss		
other	40x / 1.2 NA water immersion objective	Zeiss		
other	20x / 0.8 NA objective	Zeiss		

other	M165FC	Leica		
other	Achromat 1.6x objective	Leica		
other	GFP Filter set ET470/40x; ET525/50m	Leica		
other	mCherry Filter set ET560/40x; ET630/75m	Leica		
other	pco.edge 4.2 bi sCMOS	PCO		
other	Cell Center Stockroom (Penn)	other	RRID:SCR_0 22399	
other	CDB Microscopy Core (Penn)	other	RRID:SCR_0 22373	

References

- Agarwal P., Shemesh T., Zaidel-Bar R (2022) **Directed cell invasion and asymmetric adhesion drive tissue elongation and turning in *C. elegans* gonad morphogenesis** *Developmental Cell* **57**:2111–2126 <https://doi.org/10.1016/j.devcel.2022.08.003>
- Anllo L., DiNardo S (2022) **Visceral mesoderm signaling regulates assembly position and function of the *Drosophila* testis niche** *Developmental Cell* **57**:1009–1023 <https://doi.org/10.1016/j.devcel.2022.03.009>
- Anllo L., Plasschaert L. W., Sui J., DiNardo S (2019) **Live imaging reveals hub cell assembly and compaction dynamics during morphogenesis of the *Drosophila* testis niche** *Developmental Biology* **446**:102–118 <https://doi.org/10.1016/j.ydbio.2018.12.014>
- Baldeosingh R., Gao H., Wu X., Fossett N (2018) **Hedgehog signaling from the Posterior Signaling Center maintains U-shaped expression and a prohemocyte population in *Drosophila*** *Developmental Biology* **441**:132–145 <https://doi.org/10.1016/j.ydbio.2018.06.020>
- Banerjee U., Girard J. R., Goins L. M., Spratford C. M (2019) ***Drosophila* as a genetic model for hematopoiesis** *Genetics* **211**:367–417 <https://doi.org/10.1534/genetics.118.300223>
- Batlle E. *et al.* (2002) **β -Catenin and TCF Mediate Cell Positioning in the Intestinal Epithelium by Controlling the Expression of EphB/EphrinB** *Cell* **111**:251–263 [https://doi.org/10.1016/S0092-8674\(02\)01015-2](https://doi.org/10.1016/S0092-8674(02)01015-2)
- Biggs L. C., Mäkelä O. J. M., Myllymäki S. M., Das Roy R., Närhi K., Pispä J., Mustonen T., Mikkola M. L (2018) **Hair follicle dermal condensation forms via FGF20 primed cell cycle exit, cell motility, and aggregation** *ELife* **7**:1–33 <https://doi.org/10.7554/eLife.36468>
- Bostock M. P., Prasad A. R., Chaouni R., Yuen A. C., Sousa-Nunes R., Amoyel M., Fernandes V. M (2020) **An Immobilization Technique for Long-Term Time-Lapse Imaging of Explanted *Drosophila* Tissues** *Frontiers in Cell and Developmental Biology* **8** <https://doi.org/10.3389/fcell.2020.590094>
- Boulais P. E., Frenette P. S (2015) **Making sense of hematopoietic stem cell niches** *Blood* <https://doi.org/10.1182/blood-2014-09-570192>
- Boulet M., Renaud Y., Lapraz F., Benmimoun B., Vandel L., Waltzer L (2021) **Characterization of the *Drosophila* Adult Hematopoietic System Reveals a Rare Cell Population With Differentiation and Proliferation Potential** *Frontiers in Cell and Developmental Biology* **9** <https://doi.org/10.3389/fcell.2021.739357>
- Bunt S., Hooley C., Hu N., Scahill C., Weavers H., Skaer H (2010) **Hemocyte-secreted type IV collagen enhances BMP signaling to guide renal tubule morphogenesis in *Drosophila*** *Developmental Cell* **19**:296–306 <https://doi.org/10.1016/j.devcel.2010.07.019>
- Chakkalakal J. V., Jones K. M., Basson M. A., Brack A. S (2012) **The aged niche disrupts muscle stem cell quiescence** *Nature* **490**:355–360 <https://doi.org/10.1038/nature11438>

- Cho B. *et al.* (2020) **Single-cell transcriptome maps of myeloid blood cell lineages in *Drosophila*** *Nature Communications* **11** <https://doi.org/10.1038/s41467-020-18135-y>
- Cliffe A., Doupé D. P., Sung H., Lim I. K. H., Ong K. H., Cheng L., Yu W (2017) **Quantitative 3D analysis of complex single border cell behaviors in coordinated collective cell migration** *Nature Communications* **8** <https://doi.org/10.1038/ncomms14905>
- Crozatier M., Ubeda J. M., Vincent A., Meister M (2004) **Cellular immune response to parasitization in *Drosophila* requires the EBF orthologue collier** *PLoS Biology* **2** <https://doi.org/10.1371/journal.pbio.0020196>
- Dalle Nogare D. E., Natesh N., Vishwasrao H. D., Shroff H., Chitnis A. B (2020) **Zebrafish Posterior Lateral Line primordium migration requires interactions between a superficial sheath of motile cells and the skin** *ELife* **9** <https://doi.org/10.7554/eLife.58251>
- de Cuevas M., Spradling A. (1998) **Morphogenesis of the *Drosophila* fusome and its implications for oocyte specification** *Development* **125** <https://doi.org/10.1242/dev.125.15.2781>
- Destalminil-Letourneau M., Morin-Poulard I., Tian Y., Vanzo N., Crozatier M (2021) **The vascular niche controls *drosophila* hematopoiesis via fibroblast growth factor signaling** *ELife* **10**:1–21 <https://doi.org/10.7554/eLife.64672>
- Díaz-Torres A., Rosales-Nieves A. E., Pearson J. R., Santa-Cruz Mateos C., Marín-Menguiano M., Marshall O. J., Brand A. H., González-Reyes A (2021) **Stem cell niche organization in the *Drosophila* ovary requires the ECM component Perlecan** *Current Biology* **31**:1744–1753 <https://doi.org/10.1016/j.cub.2021.01.071>
- Englund C., Lorén C. E., Grabbe C., Varshney G. K., Deleuil F., Hallberg B., Palmer R. H (2003) **Jeb signals through the Alk receptor tyrosine kinase to drive visceral muscle fusion** *Nature* **425**:512–516 <https://doi.org/10.1038/nature01950>
- Englund C., Steneberg P., Falileeva L., Xylourgidis N., Samakovlis C (2002) **Attractive and repulsive functions of Slit are mediated by different receptors in the *Drosophila* trachea** *Development* **129**:4941–4951 <https://doi.org/10.1242/dev.129.21.4941>
- Ferraro F., Lo Celso C., Scadden D. (2010) **Adult stem cells and their niches** *Advances in Experimental Medicine and Biology* **695**:155–168 https://doi.org/10.1007/978-1-4419-7037-4_11
- Gajewski K., Choi C. Y., Kim Y., Schulz R. A (2000) **Genetically distinct cardiac cells within the *Drosophila* heart** *Genesis* **28**:36–43 [https://doi.org/10.1002/1526-968X\(200009\)28:1<36::AID-GENE50>3.0.CO;2-4](https://doi.org/10.1002/1526-968X(200009)28:1<36::AID-GENE50>3.0.CO;2-4)
- Girard J. R., Goins L. M., Vu D. M., Sharpley M. S., Spratford C. M., Mantri S. R., Banerjee U (2021) **Paths and pathways that generate cell-type heterogeneity and developmental progression in hematopoiesis** *eLife* **10** <https://doi.org/10.7554/eLife.67516>
- Gordon K. L., Zussman J. W., Li X., Miller C., Sherwood D. R (2020) **Stem cell niche exit in *C. Elegans* via orientation and segregation of daughter cells by a cryptic cell outside the niche** *ELife* **9**:1–56 <https://doi.org/10.7554/eLife.56383>
- Gregg C. L., Butcher J. T. (2012) **Quantitative in vivo imaging of embryonic development: Opportunities and challenges** *Differentiation* :149–162 <https://doi.org/10.1016/j.diff.2012.05.003>

Grigorian M., Mandal L., Hartenstein V (2011) **Hematopoiesis at the onset of metamorphosis: terminal differentiation and dissociation of the *Drosophila* lymph gland** *Development Genes and Evolution* **221**:121–131 <https://doi.org/10.1007/s00427-011-0364-6>

Gupta K. *et al.* (2019) **Single-Cell Analysis Reveals a Hair Follicle Dermal Niche Molecular Differentiation Trajectory that Begins Prior to Morphogenesis** *Developmental Cell* **48**:17–31 <https://doi.org/10.1016/j.devcel.2018.11.032>

Haas P., Gilmour D (2006) **Chemokine Signaling Mediates Self-Organizing Tissue Migration in the Zebrafish Lateral Line** *Developmental Cell* **10**:673–680 <https://doi.org/10.1016/j.devcel.2006.02.019>

Han Z., Bodmer R (2003) **Myogenic cells fates are antagonized by Notch only in asymmetric lineages of the *Drosophila* heart, with or without cell division** *Development* **130**:3039–3051 <https://doi.org/10.1242/dev.00484>

Hartenstein V. (1993) **Atlas of *Drosophila* Development**

Heitman N., Saxena N., Rendl M. (2018) **Advancing insights into stem cell niche complexities with next-generation technologies** *Current Opinion in Cell Biology* :87–95 <https://doi.org/10.1016/j.ceb.2018.06.012>

Hirschhäuser A., Molitor D., Salinas G., Großhans J., Rust K., Bogdan S (2023) **Single-cell transcriptomics identifies new blood cell populations in *Drosophila* released at the onset of metamorphosis** *Development* **150** <https://doi.org/10.1242/dev.201767>

Ho K. Y. L., An K., Carr R. L., Dvoskin A. D., Ou A. Y. J., Vogl W., Tanentzapf G. (2023) **Maintenance of hematopoietic stem cell niche homeostasis requires gap junction-mediated calcium signaling** *Proceedings of the National Academy of Sciences of the United States of America* **120** <https://doi.org/10.1073/pnas.2303018120>

Ho K. Y. L., Khadilkar R. J., Carr R. L., Tanentzapf G (2021) **A gap-junction-mediated, calcium-signaling network controls blood progenitor fate decisions in hematopoiesis** *Current Biology* **31**:4697–4712 <https://doi.org/10.1016/j.cub.2021.08.027>

Holmberg J., Genander M., Halford M. M., Annerén C., Sondell M., Chumley M. J., Silvary R. E., Henkemeyer M., Frisén J (2006) **EphB Receptors Coordinate Migration and Proliferation in the Intestinal Stem Cell Niche** *Cell* **125**:1151–1163 <https://doi.org/10.1016/j.cell.2006.04.030>

Holz A., Bossinger B., Strasser T., Janning W., Klapper R (2003) **The two origins of hemocytes in *Drosophila*** *Development* **130**:4955–4962 <https://doi.org/10.1242/dev.00702>

Jenkins V.K., Larkin A., Thurmond J., Consortium FlyBase (2022) **Using FlyBase: A Database of *Drosophila* Genes and Genetics** *Methods Mol. Biol* **2540**

Jung S. H., Evans C. J., Uemura C., Banerjee U (2005) **The *Drosophila* lymph gland as a developmental model of hematopoiesis** *Development* **132**:2521–2533 <https://doi.org/10.1242/dev.01837>

Kellermeyer R., Heydman L. M., Gillis T., Mastick G. S., Song M., Kidd T (2020) **Proteolytic cleavage of Slit by the Tolkin protease converts an axon repulsion cue to an axon growth cue in vivo** *Development* <https://doi.org/10.1242/dev.196055>

- Khadilkar R. J., Vogl W., Goodwin K., Tanentzapf G. (2017) **Modulation of occluding junctions alters the hematopoietic niche to trigger immune activation** *eLife* <https://doi.org/10.7554/eLife.28081>
- Khadilkar R. J., Vogl W., Goodwin K., Tanentzapf G (2017) **Modulation of occluding junctions alters the hematopoietic niche to trigger immune activation** *ELife* **6**:1–30 <https://doi.org/10.7554/eLife.28081>
- Kidd T., Bland K. S., Goodman C. S (1999) **Slit Is the Midline Repellent for the Robo Receptor in Drosophila** *Cell* **96**:785–794 [https://doi.org/10.1016/S0092-8674\(00\)80589-9](https://doi.org/10.1016/S0092-8674(00)80589-9)
- Kobielak K., Stokes N., De La Cruz J., Polak L., Fuchs E. (2007) **Kobielak, K., Stokes, N., De La Cruz, J., Polak, L., & Fuchs, E. (2007). Loss of a quiescent niche but not follicle stem cells in the absence of bone morphogenetic protein signaling.** www.pnas.org/cgi/content/full/
- Kolesnikov T., Beckendorf S. K (2005) **NETRIN and SLIT guide salivary gland migration** *Developmental Biology* **284**:102–111 <https://doi.org/10.1016/j.ydbio.2005.04.037>
- Koranteng F., Cho B., Shim J. (2022) **Intrinsic and Extrinsic Regulation of Hematopoiesis in Drosophila** *Molecules and Cells* **45**:101–108 <https://doi.org/10.14348/molcells.2022.2039>
- Kramer S. G., Kidd T., Simpson J. H., Goodman C. S (2001) **Switching Repulsion to Attraction: Changing Responses to Slit During Transition in Mesoderm Migration** *Science* **292**:737–740 <https://doi.org/10.1126/science.1058766>
- Krzemień J., Dubois L., Makki R., Meister M., Vincent A., Crozatier M (2007) **Control of blood cell homeostasis in Drosophila larvae by the posterior signalling centre** *Nature* **446**:325–328 <https://doi.org/10.1038/nature05650>
- Krzemien J., Oyallon J., Crozatier M., Vincent A (2010) **Hematopoietic progenitors and hemocyte lineages in the Drosophila lymph gland** *Developmental Biology* **346**:310–319 <https://doi.org/10.1016/j.ydbio.2010.08.003>
- Lanot R., Zachary D., Holder F., Meister M (2001) **Postembryonic hematopoiesis in Drosophila** *Developmental Biology* **230**:243–257 <https://doi.org/10.1006/dbio.2000.0123>
- Le Bras S., Van Doren M. (2006) **Development of the male germline stem cell niche in Drosophila** *Developmental Biology* **294**:92–103 <https://doi.org/10.1016/j.ydbio.2006.02.030>
- Lebestky T., Jung S. H., Banerjee U (2003) **A Serrate-expressing signaling center controls Drosophila hematopoiesis** *Genes and Development* **17**:348–353 <https://doi.org/10.1101/gad.1052803>
- Lee H. H., Norris A., Weiss J. B., Frasch M (2003) **Jelly belly protein activates the receptor tyrosine kinase Alk to specify visceral muscle pioneers** *Nature* **425**:507–512 <https://doi.org/10.1038/nature01916>
- Lo P. C. H., Frasch M (2001) **A role for the COUP-TF-related gene seven-up in the diversification of cardioblast identities in the dorsal vessel of Drosophila** *Mechanisms of Development* **104**:49–60 [https://doi.org/10.1016/S0925-4773\(01\)00361-6](https://doi.org/10.1016/S0925-4773(01)00361-6)

- Louradour I., Sharma A., Morin-Poulard I., Letourneau M., Vincent A., Crozatier M., Vanzo N (2017) **Reactive oxygen species-dependent Toll/NF-κB activation in the drosophila hematopoietic niche confers resistance to wasp parasitism** *ELife* **6**:1–22 <https://doi.org/10.7554/eLife.25496>
- Lucila Scimone M., Cote L. E., Reddien P. W (2017) **Orthogonal muscle fibres have different instructive roles in planarian regeneration** *Nature* **551**:623–628 <https://doi.org/10.1038/nature24660>
- Luo F., Yu S., Jin L. H. (2020) **The Posterior Signaling Center Is an Important Microenvironment for Homeostasis of the Drosophila Lymph Gland** *Frontiers in Cell and Developmental Biology* <https://doi.org/10.3389/fcell.2020.00382>
- MacMullin A., Jacobs J. R (2006) **Slit coordinates cardiac morphogenesis in Drosophila** *Developmental Biology* **293**:154–164 <https://doi.org/10.1016/j.ydbio.2006.01.027>
- MacMullin A., Jacobs J. R (2006) **Slit coordinates cardiac morphogenesis in Drosophila** *Developmental Biology* **293**:154–164 <https://doi.org/10.1016/j.ydbio.2006.01.027>
- Mandal L., Banerjee U., Hartenstein V (2004) **Evidence for a fruit fly hemangioblast and similarities between lymph-gland hematopoiesis in fruit fly and mammal aorta-gonadal-mesonephros mesoderm** *Nature Genetics* **36**:1019–1023 <https://doi.org/10.1038/ng1404>
- Mandal L., Martinez-Agosto J. A., Evans C. J., Hartenstein V., Banerjee U (2007) **A Hedgehog- and Antennapedia-dependent niche maintains Drosophila haematopoietic precursors** *Nature* **446**:320–324 <https://doi.org/10.1038/nature05585>
- Medioni C., Astier M., Zmojdzian M., Jagla K., Sémériva M (2008) **Genetic control of cell morphogenesis during Drosophila melanogaster cardiac tube formation** *Journal of Cell Biology* **182**:249–261 <https://doi.org/10.1083/jcb.200801100>
- Medioni C., Astier M., Zmojdzian M., Jagla K., Sémériva M (2008) **Genetic control of cell morphogenesis during Drosophila melanogaster cardiac tube formation** *Journal of Cell Biology* **182**:249–261 <https://doi.org/10.1083/jcb.200801100>
- Morin-Poulard I., Sharma A., Louradour I., Vanzo N., Vincent A., Crozatier M (2016) **Vascular control of the Drosophila haematopoietic microenvironment by Slit/Robo signalling** *Nature Communications* **7** <https://doi.org/10.1038/ncomms11634>
- Morrison S. J., Spradling A. C (2008) **Stem Cells and Niches: Mechanisms That Promote Stem Cell Maintenance throughout Life** *Cell* **132**:598–611 <https://doi.org/10.1016/j.cell.2008.01.038>
- Peercy B. E., Starz-Gaiano M (2020) **Clustered cell migration: Modeling the model system of Drosophila border cells** *Seminars in Cell & Developmental Biology* **100**:167–176 <https://doi.org/10.1016/j.semcdb.2019.11.010>
- Pentinmikko N. *et al.* (2022) **Cellular shape reinforces niche to stem cell signaling in the small intestine** *In Sci. Adv* **8**
- Pinho S., Frenette P. S. (2019) **Haematopoietic stem cell activity and interactions with the niche** *Nature Reviews Molecular Cell Biology* :303–320 <https://doi.org/10.1038/s41580-019-0103-9>

- Qian L., Liu J., Bodmer R (2005) **Slit and Robo control cardiac cell polarity and morphogenesis** *Current Biology* **15**:2271–2278 <https://doi.org/10.1016/j.cub.2005.10.037>
- Qin L., Yang D., Yi W., Cao H., Xiao G (2021) **Roles of leader and follower cells in collective cell migration** *Molecular Biology of the Cell* **32**:1267–1272 <https://doi.org/10.1091/mbc.E20-10-0681>
- Ramesh P., Dey N. S., Kanwal A., Mandal S., Mandal L (2021) **Relish plays a dynamic role in the niche to modulate drosophila blood progenitor homeostasis in development and infection** *ELife* **10** <https://doi.org/10.7554/eLife.67158>
- Raz A. A. *et al.* (2023) **Emergent dynamics of adult stem cell lineages from single nucleus and single cell RNA-Seq of Drosophila testes** *eLife* **12** <https://doi.org/10.7554/eLife.82201>
- Rizki T. M., Rizki R. M (1992) **Lamellocyte differentiation in Drosophila larvae parasitized by Leptopilina** *Developmental & Comparative Immunology* **16**:103–110 [https://doi.org/10.1016/0145-305X\(92\)90011-Z](https://doi.org/10.1016/0145-305X(92)90011-Z)
- Rompolas P., Mesa K. R., Greco V (2013) **Spatial organization within a niche as a determinant of stem-cell fate** *Nature* **502**:513–518 <https://doi.org/10.1038/nature12602>
- Rothberg J. M., Hartley D. A., Walther Z., Artavanis-Tsakonas S (1988) **slit: An EGF-homologous locus of D. melanogaster involved in the development of the embryonic central nervous system** *Cell* **55**:1047–1059 [https://doi.org/10.1016/0092-8674\(88\)90249-8](https://doi.org/10.1016/0092-8674(88)90249-8)
- Rothberg J. M., Jacobs J Roger, Goodman C. S., Artavanis-Tsakonas S. (1990) **slit: an extracellular protein necessary for development of midline glia and commissural axon pathways contains both EGF and LRR domains** *Genes Dev.* **4**:2169–87 <https://doi.org/10.1101/gad.4.12a.2169>
- Santiago-Martínez E., Soplop N. H., Kramer S. G. (2006) **Lateral positioning at the dorsal midline: Slit and Roundabout receptors guide Drosophila heart cell migration** *PNAS* **103**:12441–12446 <https://doi.org/10.1073/pnas.0605284103>
- Santiago-Martínez E., Soplop N. H., Patel R., Kramer S. G (2008) **Repulsion by Slit and Roundabout prevents Shotgun/E-cadherin-mediated cell adhesion during Drosophila heart tube lumen formation** *Journal of Cell Biology* **182**:241–248 <https://doi.org/10.1083/jcb.200804120>
- Shrestha R., Gateff E. (1982) **Ultrastructure and Cytochemistry of the Cell Types in the Larval Hematopoietic Organs and Hemolymph of Drosophila Melanogaster** *drosophila/hematopoiesis/blood cells/ultrastructure/cytochemistry Development, Growth & Differentiation* **24**:65–82 <https://doi.org/10.1111/j.1440-169X.1982.00065.x>
- Sinenko S. A., Mandal L., Martinez-Agosto J. A., Banerjee U (2009) **Dual Role of Wingless Signaling in Stem-like Hematopoietic Precursor Maintenance in Drosophila** *Developmental Cell* **16**:756–763 <https://doi.org/10.1016/j.devcel.2009.03.003>
- Sinenko S. A., Shim J., Banerjee U (2012) **Oxidative stress in the haematopoietic niche regulates the cellular immune response in Drosophila** *EMBO Reports* **13**:83–89 <https://doi.org/10.1038/embor.2011.223>
- Spéder P., Brand A. H (2018) **Systemic and local cues drive neural stem cell niche remodelling during neurogenesis in Drosophila** *ELife* <https://doi.org/10.7554/eLife.30413>

- Spratford C. M., Goins L. M., Chi F., Girard J. R., Macias S. N., Ho V. W., Banerjee U (2021) **Intermediate progenitor cells provide a transition between hematopoietic progenitors and their differentiated descendants** *Development* **148** <https://doi.org/10.1242/dev.200216>
- Sumigra y K. D., Terwilliger M., Lechler T (2018) **Morphogenesis and Compartmentalization of the Intestinal Crypt** *Developmental Cell* **45**:183–197 <https://doi.org/10.1016/j.devcel.2018.03.024>
- Tian Y., Morin-Poulard I., Liu X., Vanzo N., Crozatier M (2023) **A mechanosensitive vascular niche for Drosophila hematopoiesis** *Proceedings of the National Academy of Sciences of the United States of America* **120** <https://doi.org/10.1073/pnas.2217862120>
- Tokusumi T., Tokusumi Y., Hopkins D. W., Schulz R. A (2015) **Bag of Marbles controls the size and organization of the Drosophila hematopoietic niche through interactions with the Insulin-like growth factor pathway and Retinoblastoma-family protein** *Development* **142**:2261–2267 <https://doi.org/10.1242/dev.121798>
- van Es J. H. *et al.* (2005) **Wnt signalling induces maturation of Paneth cells in intestinal crypts** *Nature Cell Biology* **7**:381–386 <https://doi.org/10.1038/ncb1240>
- Walkley C. R., Olsen G. H., Dworkin S., Fabb S. A., Swann J., McArthur G. A. A., Westmoreland S. V., Chambon P., Scadden D. T., Purton L. E (2007) **A Microenvironment-Induced Myeloproliferative Syndrome Caused by Retinoic Acid Receptor γ Deficiency** *Cell* **129**:1097–1110 <https://doi.org/10.1016/j.cell.2007.05.014>
- Ward E. J., Skeath J. B (2000) **Characterization of a novel subset of cardiac cells and their progenitors in the Drosophila embryo** *Development* **127** <https://doi.org/10.1242/dev.127.22.4959>
- Warder B. N., Nelson K. A., Sui J., Anllo L., Dinardo S (2024) **An actomyosin network organizes niche morphology and responds to feedback from recruited stem cells** *BioRxiv* <https://doi.org/10.1101/2023.09.08.556877>
- Weavers H., Skaer H (2013) **Tip Cells Act as Dynamic Cellular Anchors in the Morphogenesis of Looped Renal Tubules in Drosophila** *Developmental Cell* **27**:331–344 <https://doi.org/10.1016/j.devcel.2013.09.020>
- Weiss J. B., Suyama K. L., Lee H. H., Scott M. P (2001) **Jelly belly: A Drosophila LDL receptor repeat-containing signal required for mesoderm migration and differentiation** *Cell* **107**:387–398 [https://doi.org/10.1016/S0092-8674\(01\)00540-2](https://doi.org/10.1016/S0092-8674(01)00540-2)
- Witchley J. N., Mayer M., Wagner D. E., Owen J. H., Reddien P. W (2013) **Muscle cells provide instructions for planarian regeneration** *Cell Reports* **4**:633–641 <https://doi.org/10.1016/j.celrep.2013.07.022>
- Zaffran S., K  chler A., Lee H. H., Frasch M (2001) **biniou (FoxF), a central component in a regulatory network controlling visceral mesoderm development and midgut morphogenesis in Drosophila** *Genes and Development* **15**:2900–2915 <https://doi.org/10.1101/gad.917101>
- Zhang J. *et al.* (2003) **Identification of the haematopoietic stem cell niche and control of the niche size** *Nature* **425**:836–841 <https://doi.org/10.1038/nature02041>

Zheng Q., Wang Y., Vargas E., DiNardo S (2011) **Magu is required for germline stem cell self-renewal through BMP signaling in the *Drosophila* testis** *Developmental Biology* **357**:202–210 <https://doi.org/10.1016/j.ydbio.2011.06.022>

Editors

Reviewing Editor

Lolitika Mandal

Indian Institute of Science Education and Research Mohali, Mohali, India

Senior Editor

Utpal Banerjee

University of California, Los Angeles, Los Angeles, United States of America

Reviewer #1 (Public Review):

Summary:

The study by Nelson et al. is focused on the formation of the *Drosophila* Posterior Signaling Center (PSC) which ultimately acts as a niche to support hematopoietic stem cells of the lymph gland (LG). Using a combination of genetics and live imaging, the authors show that PSC cells migrate as a tight collective and associate with multiple tissues during a trajectory that positions them at the posterior of the LG.

This is an important study that identifies Slit-Robo signaling as a regulator of PSC morphogenesis, and highlights the complex relationship of interacting cell types - PSC, visceral mesoderm (VM), and cardioblasts (CBs) - in the coordinated development of these three tissues during organ development. However, one point requiring clarification is the idea that PSC cells exhibit a collective cell migration; it is not clear that the cells are migrating rather than being pushed to a more dorsal position through dorsal closure and/or other similar large-scale embryo movement. This does not detract from the very interesting analysis of PSC morphogenesis as presented.

Strengths:

(1) Using the expression of Hid or Grim to ablate associated tissues, they find evidence that the VM and CB of the dorsal vessel affect PSC migration/morphology whereas the alary muscles do not. Slit is expressed by both VM and CBs, and therefore Slit-Robo signaling was investigated as PSCs express Robo.

(2) Using a combination of approaches, the authors convincingly demonstrate that Slit expression in the CBs and VM acts to support PSC positioning. A strength is the ability to knockdown slit levels in particular tissue types using the Gal4 system and RNAi.

(3) Although in the analysis of robo mutants, the PSC positioning phenotype is weaker in the individual mutants (robo1 and robo2) with only the double mutant (robo1,robo2) exhibiting a phenotype comparable to the slit RNAi. The authors make a reasonable argument that Slit-Robo signaling has an intrinsic effect, likely acting within PSCs because PSCs show a phenotype even when CBs do not (Figure 4G).

(4) New insight into dorsal vessel formation by VM is presented in Figure 4A, B, as loss of the VM can affect dorsal vessel morphogenesis. This result additionally points to the VM as important.

Weaknesses:

(1) The authors are cautioned to temper the result that Slit-Robo signaling is intrinsic to PSC since the loss of robo may affect other cell types (besides CBs and PSCs) to indirectly affect PSC migration/morphogenesis. In fact, in the robo2, robo1 mutant, the VM appears to be incorrectly positioned (Figure 4G).

(2) If possible, the authors should use RNAi to knockdown Robo1 and Robo2 levels specifically in the PSCs if a Gal4 is available; might Antp.Gal4 (Fig 1K) be useful? Even if knockdown is achieved in PSCs+CBs, this would be a better/complementary experiment to support the approach outlined in Figure 4D.

(3) Movies are hard to interpret, as it seems unclear that the PSCs actively migrate rather than being pushed/moved indirectly due to association with VM and CBs/dorsal vessel.

<https://doi.org/10.7554/eLife.100455.1.sa3>

Reviewer #2 (Public Review):

The paper by Nelson KA, et al. explored the collective migration, coalescence, and positioning of the posterior signaling center (PSC) cells in *Drosophila* embryo. With live imaging, the authors observed the dynamic progress of PSC migration. Throughout this process, visceral mesoderm (VM), alary muscles (Ams), and cardioblasts (CBs) are in proximity to PSC. Genetic ablation of these tissues reveals the requirement for VM and CBs, but not AMs in this process. Genetic manipulations further demonstrated that Slit-Robo signaling was critical during PSC migration and positioning. While the genetic mechanisms of positioning the PSC were explored in much detail, including using live imaging, the functional consequence of mispositioning or (partial) absence of PSC cells has not been addressed, but would much increase the relevance of their findings. A few additional issues need to be addressed as well in this otherwise well-done study.

Major points:

(1) The only readout in their experiments is the relative correctness of PSC positioning. Importantly, what is the functional consequence if PSC is not properly positioned? This would be particularly important with robo-sli manipulations, where the PSC is present but some cells are misplaced. What is the consequence? Are the LGs affected, like the specification of their cell types, structure, and function? To address this for at least the robo-sli requirement in the PSC, it may be important to manipulate them directly in the PSC with a split Gal4 system, using Antp and Odd promoters.

(2) The densely, parallel aligned fibers in the part of Figure 1J seemed to be visceral mesoderm, but further up (dorsally) that may be epidermis. It is possible that the PSC migrate together with the epidermis? This should be addressed.

(3) Although the authors described the standards of assessing PSC positioning as "normal" or "abnormal", it is rather subtle at times and variable in the mutant or KD/OE examples. The criteria should be more clearly delineated and analyzed double-blind, also since this is the only readout. Further examples of abnormal positioning in supplementary figures would also help.

(4) The Discussion is very lengthy and should be shortened.

<https://doi.org/10.7554/eLife.100455.1.sa2>

Reviewer #3 (Public Review):

Summary:

This work is a detailed and thorough analysis of the morphogenesis of the posterior signaling center (PSC), a hematopoietic niche in the *Drosophila* larva. Live imaging is performed from the stage of PSC determination until the appearance of a compact lymph gland and PSC in the stage 16 embryo. This analysis is combined with genetic studies that clarify the involvement of adjacent tissue, including the visceral mesoderm, alary muscle, and cardioblasts/dorsal vessels. Lastly, the Slit/Robo signaling system is clearly implicated in the normal formation of the PSC.

Strengths:

The data are clearly presented, well documented, and fully support the conclusions drawn from the different experiments. The manuscript differs in character from the mainstay of "big data" papers (for example, no sets of single-cell RNAseq data of, for instance, PSC cells with more or less Slit input, are offered), but what it lacks in this regard, it makes up in carefully planned and executed visualizations and genetic manipulations.

Weaknesses:

A few suggestions concerning improvement of the way the story is told and contextualized.

(1) The minute cluster of PSC progenitors (5 or so cells per side) is embedded (as known before and shown nicely in this study) in other "migrating" cell pools, like the cardioblasts, pericardial cells, lymph gland progenitors, alary muscle progenitors. These all appear to move more or less synchronously. What should also be mentioned is another tissue, the dorsal epidermis, which also "moves" (better: stretches?) towards the dorsal midline during dorsal closure. Would it be reasonable to speculate (based on previously published data) that without the force of dorsal closure, operating in the epidermis, at least the lateral>medial component of the "migration" of the PSC (and neighboring tissues) would be missing? If dorsal closure is blocked, do essential components of PSC and lymph gland morphogenesis (except for the coming-together of the left and right halves) still occur? Are there any published data on this?

(2) Along similar lines: the process of PSC formation is characterized as "migration". To be fair: the authors bring up the possibility that some of the phenotypes they observe could be "passive"/secondary: "Thus, it became important to test whether all PSC phenotypes might be 'passive', explained by PSC attachment to a malforming dorsal vessel. Alternatively, the PSC defects could reflect a requirement for Robo activation directly in PSC cells." And the issue is resolved satisfactorily. But more generally, "cell migration" implies active displacement (by cytoskeletal forces) of cells relative to a substrate or to their neighbors (like for example migration of hemocytes). This to me doesn't seem really clearly to happen here for the dorsal mesodermal structures. Couldn't one rather characterize the assembly of PSC, lymph gland, pericardial cells, and dorsal vessel in terms of differential adhesion, on top of a more general adhesion of cells to each other and the epidermis, and then dorsal closure as a driving force for cell displacement? The authors should bring in the published literature to provide a background that does (or does not) justify the term "migration".

(3) That brings up the mechanistic centerpiece of this story, the Slit/Robo system. First: I suggest adding more detailed data from the study by Morin-Poulard et al 2016, in the Introduction, since these authors had already implicated Slit-Robo in PSC function and offered a concrete molecular mechanism: "vascular cells produce Slit that activates Robo receptors in the PSC. Robo activation controls proliferation and clustering of PSC cells by regulating Myc, and small GTPase and DE-cadherin activity, respectively". As stated in the

Discussion: the mechanism of Slit/Robo action on the PSC in the embryo is likely different, since DE-cadherin is not expressed in the embryonic PSC; however, it maybe not be THAT different: it could also act on adhesion between PSC cells themselves and their neighbors. What are other adhesion proteins that appear in the late lateral mesodermal structures? Could DN-cadherin or Fasciclins be involved?

<https://doi.org/10.7554/eLife.100455.1.sa1>

Author response:

eLife assessment

This study presents valuable findings on the role of a well-studied signal transduction pathway, the Slit/Robo system, in the context of the assembly of the hematopoietic niche in the Drosophila embryo. The evidence supporting the claims of the authors is solid. However, one aspect that needs attention is whether the cells are migrating and not being pushed to a more dorsal position through dorsal closure and/or other similar large-scale embryo movement. This does not detract from the very interesting analysis of PSC morphogenesis and will interest developmental biologists working on molecular mechanisms of tissue morphogenesis.

We appreciate the thoughtful and quite useful comments provided by each of the referees. Our responses are noted below each referee's comment.

Public Reviews:

Reviewer #1 (Public Review):

Summary:

The study by Nelson et al. is focused on the formation of the Drosophila Posterior Signaling Center (PSC) which ultimately acts as a niche to support hematopoietic stem cells of the lymph gland (LG). Using a combination of genetics and live imaging, the authors show that PSC cells migrate as a tight collective and associate with multiple tissues during a trajectory that positions them at the posterior of the LG.

This is an important study that identifies Slit-Robo signaling as a regulator of PSC morphogenesis, and highlights the complex relationship of interacting cell types - PSC, visceral mesoderm (VM), and cardioblasts (CBs) - in the coordinated development of these three tissues during organ development. However, one point requiring clarification is the idea that PSC cells exhibit a collective cell migration; it is not clear that the cells are migrating rather than being pushed to a more dorsal position through dorsal closure and/or other similar large-scale embryo movement. This does not detract from the very interesting analysis of PSC morphogenesis as presented.

Since each referee asked for clarification concerning collective cell migration, we present a combined response further below, placed after the comments from Reviewer #3.

Strengths:

(1) Using the expression of Hid or Grim to ablate associated tissues, they find evidence that the VM and CB of the dorsal vessel affect PSC migration/morphology whereas the alary muscles do not. Slit is expressed by both VM and CBs, and therefore Slit-Robo signaling was investigated as PSCs express Robo.

(2) Using a combination of approaches, the authors convincingly demonstrate that Slit expression in the CBs and VM acts to support PSC positioning. A strength is the ability to knockdown slit levels in particular tissue types using the Gal4 system and RNAi.

(3) Although in the analysis of robo mutants, the PSC positioning phenotype is weaker in the individual mutants (robo1 and robo2) with only the double mutant (robo1,robo2) exhibiting a phenotype comparable to the slit RNAi. The authors make a reasonable argument that Slit-Robo signaling has an intrinsic effect, likely acting within PSCs because PSCs show a phenotype even when CBs do not (Figure 4G).

(4) New insight into dorsal vessel formation by VM is presented in Figure 4A, B, as loss of the VM can affect dorsal vessel morphogenesis. This result additionally points to the VM as important.

Weaknesses:

(1) The authors are cautioned to temper the result that Slit-Robo signaling is intrinsic to PSC since the loss of robo may affect other cell types (besides CBs and PSCs) to indirectly affect PSC migration/morphogenesis. In fact, in the robo2, robo1 mutant, the VM appears to be incorrectly positioned (Figure 4G).

We have reexamined our wording in the relevant Results section and, given that this referee agrees that we, “make a reasonable argument that Slit-Robo signaling has an intrinsic effect, likely acting within PSCs because PSCs show a phenotype even when CBs do not (Figure 4G)”, it was not clear how we might temper our conclusions more. Given that PSC cells express Robo1 and Robo2, and that the Vm does not contact the PSC, our ‘reasonable argument’ appears fair and parsimonious. Since we agree with the referee that a reader should be made as aware as possible of alternatives, we will add a comment to the Discussion, reminding the reader of the possibility of a secondary defect.

(2) If possible, the authors should use RNAi to knockdown Robo1 and Robo2 levels specifically in the PSCs if a Gal4 is available; might Antp.Gal4 (Fig 1K) be useful? Even if knockdown is achieved in PSCs+CBs, this would be a better/complementary experiment to support the approach outlined in Figure 4D.

While we agree that PSC-specific knockdown of Robo1 and Robo2 simultaneously would be ideal, this is not possible. First, the most-effective UAS-RNAi transgenes (that is, those in a Valium 20 backbone) are both integrated at the same chromosomal position; these cannot be simultaneously crossed with a GAL4 transgenic line to attempt double knock down. Additionally, as with all RNAi approaches that must rely on efficient knockdown over the rapid embryonic period, even having facile access to the above does not ensure the RNAi approach will cause as effective depletion as the genetic null condition that we use. Second, as the referee concedes, there is no embryonic PSC-specific GAL4. The proposed use of Antp-GAL4 would cause knockdown in many tissues (PSC, CB, Vm, epidermis and amnioserosa). This would lead to a reservation similar to that caused by our use of the straight genetic double mutant, as regards potential indirect requirement for Robo function.

(3) Movies are hard to interpret, as it seems unclear that the PSCs actively migrate rather than being pushed/moved indirectly due to association with VM and CBs/dorsal vessel.

First, the Vm does not directly contact the PSC, so it cannot be pushing the PSC dorsally. We will re-examine our text to be certain to make this clear. Second, in our analysis of bin mutants, which lack Vm, LGs and PSCs are able to reach the dorsal midline region in the absence of Vm. Finally, please see our response to Reviewer #3, point 2, for why we maintain that PSC cells are “migrating” even though some PSC cells are attached to CBs.

Reviewer #2 (Public Review):

The paper by Nelson KA, et al. explored the collective migration, coalescence, and positioning of the posterior signaling center (PSC) cells in Drosophila embryo. With live imaging, the authors observed the dynamic progress of PSC migration. Throughout this process, visceral mesoderm (VM), alary muscles (Ams), and cardioblasts (CBs) are in proximity to PSC. Genetic ablation of these tissues reveals the requirement for VM and CBs, but not AMs in this process. Genetic manipulations further demonstrated that Slit-Robo signaling was critical during PSC migration and positioning. While the genetic mechanisms of positioning the PSC were explored in much detail, including using live imaging, the functional consequence of mispositioning or (partial) absence of PSC cells has not been addressed, but would much increase the relevance of their findings. A few additional issues need to be addressed as well in this otherwise well-done study.

Major points:

(1) The only readout in their experiments is the relative correctness of PSC positioning. Importantly, what is the functional consequence if PSC is not properly positioned? This would be particularly important with robo-sli manipulations, where the PSC is present but some cells are misplaced. What is the consequence? Are the LGs affected, like the specification of their cell types, structure, and function? To address this for at least the robo-slit requirement in the PSC, it may be important to manipulate them directly in the PSC with a split Gal4 system, using Antp and Odd promoters.

We agree that the functional consequence of PSC mis-positioning is important and a relevant question to eventually address. However, virtually all markers and reagents used to assess the effect of the PSC on progenitor cells and their differentiated descendants are restricted to analyses carried out on the third larval instar - some three days after the experiments reported here. Most of the manipulated conditions in our work are no longer viable at this phase and, thus, addressing the functional consequences of a malformed PSC will require the field to develop new tools.

As we noted in the Introduction, the consistency with which the wildtype PSC forms as a coalesced collective at the posterior of the LG strongly suggests importance of its specific positioning and shape, as has now been found for other niches (citations in manuscript). Additionally, in the Discussion we mention the existence of a gap junction-dependent calcium signaling network in the PSC that is important for progenitor maintenance. Without continuity of this network amongst all PSC cells (under conditions of PSC mis-positioning), we strongly anticipate that the balance of progenitors to differentiated hemocytes will be mis-managed, either constitutively, and / or under immune challenge conditions.

Finally, to our knowledge, the tools do not exist to build a “split Gal4 system using Antp and Odd promoters”. The expression pattern observed using the genomic Antp-GAL4 line must be driven by endogenous enhancers—none of which have been defined by the field, and thus cannot be used in constructing second order drivers. Similarly, for odd skipped, in the embryo the extant Odd-GAL4 driver expresses only in the epidermis, with no expression in the embryonic LG. Thus, the cis regulatory element controlling Odd expression in the embryonic LG is unknown. In the future, the discovery of an embryonic PSC-specific driver will aid in addressing the specific functional consequences of PSC mis-positioning.

(2) The densely, parallel aligned fibers in the part of Figure 1J seemed to be visceral mesoderm, but further up (dorsally) that may be epidermis. It is possible that the PSC migrate together with the epidermis? This should be addressed.

See response to Reviewer #3.

(3) Although the authors described the standards of assessing PSC positioning as "normal" or "abnormal", it is rather subtle at times and variable in the mutant or KD/OE examples. The criteria should be more clearly delineated and analyzed double-blind, also since this is the only readout. Further examples of abnormal positioning in supplementary figures would also help.

We appreciate the Reviewer's concern and acknowledge that the phenotypes we observed were indeed variable, and, at times subtle. As we show and discuss in the paper, our results revealed that the signaling requirements for proper PSC positioning are complex; this was favorably commented upon by Reviewer #1 ("...highlights the complex relationship of interacting cell types - PSC, visceral mesoderm (VM), and cardioblasts (CBs) - in the coordinated development of these three tissues during organ development..."). We suspect the phenotypic variability is attributable to any number of biological differences such as heterogeneity of PSC cells and an accompanying difference in the timing of their competence to receive and respond to Slit-Robo signaling, the timing of release of Slit from CBs and Vm, number of cells in a given PSC, which PSC cells in the cluster respond to too little or too much signaling, and/or typical variability between organisms. Furthermore, PSC positioning analyses were conducted by two of the authors, who independently came to the same conclusions. For many of the manipulations double blinding was not possible since the genotype of the embryo was discernible due to the obvious phenotype of the manipulated tissue.

(4) The Discussion is very lengthy and should [be] shortened.

We will re-examine the prose and emphasize more conciseness, while maintaining clarity for the reader.

Reviewer #3 (Public Review):

Summary:

This work is a detailed and thorough analysis of the morphogenesis of the posterior signaling center (PSC), a hematopoietic niche in the Drosophila larva. Live imaging is performed from the stage of PSC determination until the appearance of a compact lymph gland and PSC in the stage 16 embryo. This analysis is combined with genetic studies that clarify the involvement of adjacent tissue, including the visceral mesoderm, alary muscle, and cardioblasts/dorsal vessels. Lastly, the Slit/Robo signaling system is clearly implicated in the normal formation of the PSC.

Strengths:

The data are clearly presented, well documented, and fully support the conclusions drawn from the different experiments. The manuscript differs in character from the mainstay of "big data" papers (for example, no sets of single-cell RNAseq data of, for instance, PSC cells with more or less Slit input, are offered), but what it lacks in this regard, it makes up in carefully planned and executed visualizations and genetic manipulations.

Weaknesses:

A few suggestions concerning improvement of the way the story is told and contextualized.

(1) The minute cluster of PSC progenitors (5 or so cells per side) is embedded (as known before and shown nicely in this study) in other "migrating" cell pools, like the

cardioblasts, pericardial cells, lymph gland progenitors, alary muscle progenitors. These all appear to move more or less synchronously. What should also be mentioned is another tissue, the dorsal epidermis, which also "moves" (better: stretches?) towards the dorsal midline during dorsal closure. Would it be reasonable to speculate (based on previously published data) that without the force of dorsal closure, operating in the epidermis, at least the lateral>medial component of the "migration" of the PSC (and neighboring tissues) would be missing? If dorsal closure is blocked, do essential components of PSC and lymph gland morphogenesis (except for the coming-together of the left and right halves) still occur? Are there any published data on this?

Each of the Reviewers is interested in our response to this very relevant question, and, thus, we will address the issue en bloc here. First, we will add a Supplementary Figure showing that LG and CBs are still able to progress medially towards the dorsal midline when dorsal closure stalls. This rules out any major effect for the most prominent “large-scale embryo cell sheet movement” in positioning the PSC. Second, published work by Haack et. al. and Balaghi et. al. shows that CBs and leading edge epidermal cells are independently migratory, and we will add this context to the manuscript for the reader.

(2) Along similar lines: the process of PSC formation is characterized as "migration". To be fair: the authors bring up the possibility that some of the phenotypes they observe could be "passive"/secondary: "Thus, it became important to test whether all PSC phenotypes might be 'passive', explained by PSC attachment to a malforming dorsal vessel. Alternatively, the PSC defects could reflect a requirement for Robo activation directly in PSC cells." And the issue is resolved satisfactorily. But more generally, "cell migration" implies active displacement (by cytoskeletal forces) of cells relative to a substrate or to their neighbors (like for example migration of hemocytes). This to me doesn't seem really clearly to happen here for the dorsal mesodermal structures. Couldn't one rather characterize the assembly of PSC, lymph gland, pericardial cells, and dorsal vessel in terms of differential adhesion, on top of a more general adhesion of cells to each other and the epidermis, and then dorsal closure as a driving force for cell displacement? The authors should bring in the published literature to provide a background that does (or does not) justify the term "migration".

Before addressing this specifically, we remind readers of our response above that states the rationale ruling out large, embryo-scale movements, such as epidermal dorsal closure, in driving PSC positioning. So, how are PSC cells arriving at their reproducible position? This manuscript reports the first live-imaging of the PSC as it comes to be positioned in the embryo. We interpret these movies to suggest strongly that these cells are a ‘collective’ that migrates. Neither the data, nor we, are asserting that each PSC cell is ‘individually’ migrating to its final position. Rather, our data suggest that the PSC migrates as a collective. The most paradigmatic example of directed, collective cell migration, is of *Drosophila* ovarian border cells. That cell cluster is surrounded at all times by other cells (nurse cells, in that case), and for the collective to traverse through the tissue, the process requires constant remodeling of associations amongst the migrating cells in the collective (the border cells), as well as between cells in the collective and those outside of it (the nurse cells). In fact, the nurse cells are considered the substrate upon which border cells migrate. Note also that in collective border cell migration cells within the collective can switch neighbors, suggesting dynamic changes to cell associations and adhesions.

In our analysis, the PSC cells exhibit qualities reminiscent of the border cells, and thus we infer that the PSC constitutes a migratory cell collective. We also show in Figure 1H that PSC cells exhibit cellular extensions, and thus have a very active, intrinsic actin-based cytoskeleton. In fact, in Figure 1I, we point out that PSC cells shift position within the collective, which is not only a direct feature of migration, but also occurs within the border

cell collective as that collective migrates. Additionally, the fact that the lateral-most PSC cells shift position in the collective while remaining a part of the collective—and they do this while executing net directional movement—makes a strong argument that the PSC is migratory, as no cell types other than PSCs are contacting the surfaces of those shifting PSC cells. Lastly, the Reviewer's supposition that, rather than migration, dorsal mesoderm structures form via “differential adhesion, on top of a more general adhesion of cells to each other” is, actually, precisely an inherent aspect of collective cell migration as summarized above for the ovarian border collective.

In our resubmission we will adjust text citing the existing literature to better put into context the reasoning for why PSC formation based on our data is an example of collective cell migration.

(3) That brings up the mechanistic centerpiece of this story, the Slit/Robo system. First: I suggest adding more detailed data from the study by Morin-Poulard et al 2016, in the Introduction, since these authors had already implicated Slit-Robo in PSC function and offered a concrete molecular mechanism: "vascular cells produce Slit that activates Robo receptors in the PSC. Robo activation controls proliferation and clustering of PSC cells by regulating Myc, and small GTPase and DE-cadherin activity, respectively". As stated in the Discussion: the mechanism of Slit/Robo action on the PSC in the embryo is likely different, since DE-cadherin is not expressed in the embryonic PSC; however, it maybe not be THAT different: it could also act on adhesion between PSC cells themselves and their neighbors. What are other adhesion proteins that appear in the late lateral mesodermal structures? Could DN-cadherin or Fasciclins be involved?

We agree with the Reviewer that Slit-Robo signaling likely acts in part on the PSC by affecting PSC cell adhesion to each other and/or to CBs (lines 428-435). As stated in the Discussion, we do not observe Fasciclin III expression in the PSC until late stages when the PSC has already been positioned, suggesting that Fasciclin III is not an active player in PSC formation. Assessing whether the PSC expresses any other of the suite of potential cell adhesion molecules such as DN-Cadherin or other Fasciclins, and then study their potential involvement in the Slit-Robo pathway in PSC cells, would be part of a follow-up study.

<https://doi.org/10.7554/eLife.100455.1.sa0>