

Global risk mapping of highly pathogenic avian influenza H5N1 and H5Nx in the light of epidemic episodes occurring from 2020 onward

Reviewed Preprint

v3 • November 13, 2025

Revised by authors

Reviewed Preprint

v2 • October 20, 2025

Reviewed Preprint

v1 • February 13, 2025

Marie-Cécile Dupas , Maria F Vincenti-Gonzalez, Madhur Dhingra, Claire Guinat, Timothée Vergne, William Wint, Guy Hendrickx, Cedric Marsboom, Marius Gilbert, Simon Dellicour

Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles (ULB), Brussels, Belgium • Data Science Institute, University of Hasselt, Hasselt, Belgium • Food and Agriculture Organization of the United Nations, Rome, Italy • Interactions Hôtes-Agents Pathogènes (IHAP), Université de Toulouse, INRAE, ENVT, Toulouse, France • Environmental Research Group Oxford Ltd, c/o Department of Biology, Oxford, United Kingdom • Avia-GIS research department, Avia-GIS, Zoersel, Belgium • Department of Microbiology, Immunology and Transplantation, Rega Institute, KU Leuven, Leuven, Belgium • Interuniversity Institute of Bioinformatics in Brussels, Université Libre de Bruxelles, Vrije Universiteit Brussel, Brussels, Belgium

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This global study compares environmental niche model outputs of avian influenza pathogen niche constructed for two distinct periods, and uses differences between those outputs to suggest that the changed case numbers and distribution relate to intensification of chicken and duck farming, and extensive cultivation. While a **useful** update to existing niche models of highly pathogenic avian influenza, the justification for the use of environmental niche models to explore correlative relationships between land cover change and changed case epidemiology is **incomplete**.

<https://doi.org/10.7554/eLife.104748.3.sa3>

Abstract

Avian influenza (AI) is a highly contagious viral disease affecting poultry and wild water birds, posing significant global challenges due to its high mortality rates and economic impacts. Highly pathogenic avian influenza (HPAI) outbreaks, particularly those caused by H5N1 and its variants, have surged since their first occurrence in 1959. The HPAI H5N1 clade 2.3.4.4b viruses have notably expanded their geographical reach, affecting numerous countries, diverse avian species, and now wild and domestic mammals. Using an ecological niche modelling approach, this study aims to elucidate the environmental factors associated with the increased HPAI H5 cases since 2020, investigate potential shifts in ecological niches, and predict new areas suitable for local viral circulation. Focusing on H5N1 and H5Nx strains, we have developed ecological niche models for HPAI cases in both wild and domestic birds while considering two distinct periods: 2015-2020 and 2020-2022. Key environmental predictors include chicken and duck population density, human density, distance to water

bodies, and several land cover variables. Post-2020, we observe a notable increase in the relative influence of some of these predictors, such as intensive chicken population density and cultivated vegetation. The resulting risk maps reveal notable ecological suitability for local HPAI H5 circulation in Europe, Asia, as well as North and South America, with notable expansions of the areas at risk post-2020. The spatial distribution of HPAI H5 occurrences in wild birds appears to be primarily correlated with urban areas and open water regions. Overall, global risk maps derived from our models identify regions at risk where surveillance and control measures should be prioritised. Finally, our analyses also highlight a potential shift in the diversity of wild bird species affected by HPAI outbreaks, with a higher variety of avian species, particularly sea birds, being impacted post-2020. This increased diversity could suggest that ecological shifts in HPAI H5 circulation may be accompanied by a broader range of susceptible species. Overall, these results further contribute to the understanding of HPAI epidemiology.

Introduction

Since 2020, an increase in both H5Nx and H5N1 cases has been observed (**Fig. 1**). A variant of the Gs/Gd H5N1 viruses belonging to the H5 clade 2.3.4.4b has led to an unprecedented number of deaths in wild birds and poultry in many countries in Africa, Asia, and Europe (Huang et al., 2023). In 2021, the virus spread to North America, and in 2022, to Central and South America (FAO, WHO and WOA, 2023) spilling over into poultry farms and infecting an alarming number of wild terrestrial, marine and domestic mammals (Elsmo et al., 2023; Leguia et al., 2023; Neumann and Kawaoka, 2024; Peacock et al., 2024; Plaza et al., 2024); marking an unprecedented expansion in the geography and impact of HPAI. In 2022, 67 countries across five continents reported H5N1 HPAI outbreaks in poultry and wild birds to the World Organisation for Animal Health (WOAH), resulting in over 131 million domestic poultry deaths or cullings (WOAH, 2023). In 2023, another 14 countries, mostly in the Americas, reported outbreaks. Several mass death events in wild birds were caused by influenza A(H5N1) clade 2.3.4.4b viruses (Klaassen and Wille, 2023; WOA, 2023).

In that epidemiological context, understanding the environmental factors associated with the risk of HPAI circulation remains crucial. Extensive exploration of risk factors for HPAI presence, spread, and persistence has been conducted in various countries and regions. These studies have identified domestic waterfowl, several anthropogenic variables (human population density, distance to roads) and indicators of water presence as important factors for risk of H5N1 local circulation (Gilbert and Pfeiffer, 2012). More recent work on global suitability for HPAI have pinpointed that host-related variables such as poultry density are the strongest contributors to HPAI persistence (Dhingra et al., 2016; Gierak and Śmietanka, 2021; Martin et al., 2011).

In addition, poultry intensification, international poultry trade, live bird markets, and wild bird migratory routes have been recognised as playing a key role in the transmission and spread of HPAI (Vandegrift et al., 2010). Lastly, evidence of climate change impacting the dynamics of HPAI has also been discussed (Prosser et al., 2023). For example, changes in rainfall alter the distribution, abundance, and quality of wetlands can impact waterfowl populations (Forcey et al., 2007; Gilbert et al., 2008; Vandegrift et al., 2010). However, recent outbreaks raise questions about whether earlier ecological niche models still accurately predict the current distribution of areas ecologically suitable for the local circulation of HPAI H5 viruses. Ecological niche model outputs for range-shifting pathogens must therefore be interpreted with caution (Elith et al., 2010). Despite this limitation, correlative ecological niche models remain useful for identifying broad-scale associations and potential shifts in distribution. To account for this, we analysed two distinct time periods (2015-2020 and 2020-2023).




Figure 1

Epidemic curves for both wild and domestic bird cases.

Orange and blue histograms report the weekly number of H5N1 cases and the weekly number of H5Nx cases, respectively.

Understanding the interplay between environmental variables and avian influenza becomes increasingly crucial to have an in-depth understanding of the risk factors that govern avian influenza viruses (AIVs) circulation and spread. Following a previous study dedicated to the ecological niche of H5Nx and H5N1 (Dhingra et al., 2016 [DOI](#)), we aimed to address the following questions: (i) Which factors are associated with the increase in H5Nx and H5N1 cases since 2020? (ii) Do we observe a change or an extension of the ecological conditions suitable for local HPAI circulation? (iii) Are we able to predict the areas at risk for local HPAI circulation based on ecological niche models trained on occurrence data before 2020? To this end, we used HPAI H5N1 and H5Nx occurrence data collected before 2020 to train ecological niche models and tested whether these models could accurately predict areas ecologically suitable for HPAI circulation post-2020. We developed separate ecological niche models for wild and domestic bird HPAI occurrences, as these two groups are influenced by different ecological processes, surveillance biases, and management contexts. These models thus predict the ecological suitability for the risk of local viral circulation leading to the detection of HPAI occurrences within each host group (rather than the niche of the virus or the host groups alone). Specifically, we employed a boosted regression trees (BRT) approach to estimate the HPAI ecological suitability given local environmental conditions. We also computed the diversity indices of wild bird species involved in HPAI occurrences, helping to understand how shifts in species diversity might correspond with the environmental factors considered in the ecological niche modelling analyses of HPAI in wild or domestic birds. Our results provide a better understanding of HPAI dynamics by identifying key environmental factors correlated with the increase in H5Nx and H5N1 cases in poultry and wild birds, investigating potential shifts in their ecological niches, and improving the prediction of at-risk areas. This knowledge might aid policymakers, epidemiologists, and wildlife conservationists in better targeting surveillance and control measures.

Results

In this study, we applied an ecological niche modelling approach using the boosted regression trees (BRT) method to evaluate the distribution of areas ecologically suitable for the risk of local circulation of HPAI H5Nx and H5N1 viruses in wild and domestic bird populations across two periods: 2015-2020 and 2020-2022. We trained separate ecological niche models for each combination of host groups (wild or domestic birds), virus strain (H5Nx or H5N1), and time period. In addition to evaluating each model separately, models trained on data from the 2015-2020 period were tested to predict occurrences in the 2020-2022 period, allowing us to evaluate the potential of pre-2020 models to predict a more recent distribution of H5N1 and H5Nx cases.

We assessed the relative influence (RI) of a broad range of spatial predictors (Elith et al., 2008), including anthropogenic, topographical, land cover, eco-climatic, and poultry density, on virus occurrences (**Fig. S1** [DOI](#)). We used occurrence records from the EMPRES-i database and sampled pseudo-absence points to account for the lack of virus detection data in some areas. These pseudo-absence points were distributed based on human population density, with more pseudo-absence points sampled in areas of higher population to reflect the greater surveillance efforts in those regions (**Fig. S2** [DOI](#)).

The accuracy of the models was validated through three types of cross-validation: (i) a standard cross-validation with a random and stratified divide between training and validation sets, (ii) a spatial cross-validation based on the approach used by Dhingra and colleagues (2016) [DOI](#) that clustered presence and pseudo-absence points into folds using reference presence points, and (iii) a second spatial cross-validation based on a blocks generation technique (Valavi et al., 2019 [DOI](#)). Models using spatial cross-validation techniques are less affected by spatial autocorrelation, as shown by higher spatial sorting bias (SSB) values (**Fig. S3** [DOI](#)). Among the seven datasets evaluated, four show better performance with the reference points-based spatial cross-validation method,

suggesting it was less impacted by spatial autocorrelation compared to the block generation technique. Consequently, we have proceeded with the reference points approach for consistency with [Dhingra et al. \(2016\)](#).

Using pre-2020 data, our models demonstrated a relatively robust capability to predict the distribution of post-2020 occurrence data. The Area Under the Curve (AUC) values for ecological niche models trained on occurrence data in domestic birds before 2020 ranged between 0.74 and 0.77 when evaluated against post-2020 data, indicating a relatively good predictive performance. As expected, ecological niche models trained on post-2020 data showed a slightly higher predictive performance when evaluated on data from the same period, with AUC values ranging from 0.78 to 0.83 (**Table S1**). These results suggest that pre-2020 models captured broad patterns of suitability for H5Nx and H5N1 outbreaks, while post-2020 models provided a closer fit to the more recent epidemiological situation.

Previous literature reviews ([Gilbert and Pfeiffer, 2012](#); [Dhingra et al., 2016](#)) have summarised the predictor variables commonly correlated with HPAI occurrences. As detailed in the Materials and Methods section and in the table in Supplementary Information Resources S1, we explored four sets of environmental variables considered by Dhingra and colleagues ([Dhingra et al., 2016](#)): host variables (set 1), land cover variables (set 2), eco-climatic variables (set 3), and a risk-based selection of variables performed by Dhingra and colleagues (set 4). Set 3 showed the lowest predictive performance for both domestic and wild bird cases during the two periods and, in contrast, sets 2 and 4 demonstrated the highest predictive accuracy for wild and domestic birds, respectively (**Fig. S3**). Set 4 combined host variables with cultivated and managed vegetation, open water areas, distance to water, and the annual mean of land surface temperature. For domestic birds, set 1 (host variables) performed well before 2020, but its predictive performance decreased after 2020 in favour of set 4. Ecological niche models trained on wild and domestic bird cases, respectively with the sets 2 and 4, are associated with AUC values exceeding 0.77, indicating a relatively strong performance in predicting H5 occurrences based on these sets of environmental variables.

Figure 2 displays the response curves of the most important variables with their respective RI (**Table S2**). In the ecological niche models for H5N1 and H5Nx in domestic birds, the densities of intensive chicken populations, domestic duck populations, and human populations emerged as significant predictors, each with RI values exceeding 5%. Notably, the RI for intensive chicken density in H5N1 models increased sharply from 8.5% to 30.4% since 2020. Similarly, the RI of cultivated and managed vegetation has doubled for both strains post-2020. The response curves showed a positive correlation, indicating that higher values of these predictors were linked with an increased likelihood of HPAI occurrences given local environmental conditions. These findings indicate a trend towards increased HPAI susceptibility in environments characterised by intensive agricultural and vegetation management practices.

In contrast, eco-climatic factors such as land surface temperature and precipitation (set 3) showed only moderate influence. We also observed a decrease in the importance of domestic duck population density for both subtypes after 2020, possibly due to the increasing diversity of bird species involved in the transmission dynamics of H5N1 and H5NX (**Fig. S4**). These findings highlight the crucial role of anthropogenic and host-related variables in accurately predicting HPAI occurrences.

In the ecological niche models trained for wild bird cases, urban and built-up areas were associated with H5N1 and H5Nx occurrences (**Fig. 2**), showing the highest relative influence (RI) prior to 2020. Specifically, RI values reached 54.5% before 2020 but decreased to 39.3% in the post-2020 period. This decline may indicate a reduced bias in observation data: typically, dead wild birds are more frequently found near human-populated areas due to opportunistic detections, whereas more recent surveillance efforts have become increasingly proactive ([Giacinti et al.,](#)




Figure 2

Response curves associated with the environmental variables included in the ecological niche models.

For the ecological niche models trained on wild bird infection records, we here only display the response curves estimated for the environmental variables associated with an averaged relative influence (RI) >4% for at least one of the considered occurrence datasets (thus not reporting the response curves obtained for the following variables: evergreen deciduous needleleaf trees, evergreen broadleaf trees, shrublands, and regularly flooded vegetation). Each curve was retrieved from a distinct boosted regression tree (BRT) model trained for a specific dataset of occurrence data. We also report the averaged RI (in %) of each environmental variable in the respective ecological models trained on a specific dataset of occurrence data (see [Table S1](#) for the complete list of RI estimates along with their first and third quartiles). Due to lack of data, the model was not trained for H5N1 in wild birds before 2020.

2024 [\[link\]](#)). After 2020, we observed an increase in the importance of habitats, such as deciduous broadleaf trees (6.1%), mixed and other tree regions (11.4%), and herbaceous vegetation for H5Nx (9.5%). However, response curves (**Fig. 2** [\[link\]](#)) indicated that H5N1 and H5Nx occurrences in wild birds decreased in areas with greater tree cover, while it increased in regions dominated by herbaceous vegetation. Open water areas consistently showed high RI values across all time periods and virus strains, particularly for H5Nx before 2020 (25.5%) and H5N1 after 2020 (22.0%), highlighting their significant role in the ecological models of HPAI outbreaks.

Figure 3 [\[link\]](#) displays maps of the predicted ecological suitability for the risk of local H5N1 and H5Nx circulation across two time periods, 2015-2020 and 2020-2022; these maps being also accessible on the following link for a dynamic visualisation of the results: <https://app.mood-h2020.eu/core> [\[link\]](#). All maps are in a WGS84 projection with a spatial resolution of 0.0833 decimal degrees (i.e. 5 arcmin, or approximately 10 km at the equator). For H5N1 and H5Nx in domestic birds, these maps reveal several regions associated with a relatively high ecological suitability, especially in Europe and Asia, in countries such as South Korea, Japan, Singapore, Malaysia, Vietnam, Cambodia, Thailand, and the Philippines, as well as the United Kingdom, France, the Netherlands, Germany, Italy, Ukraine, and Poland. Additionally, several areas in African nations, including Nigeria and South Africa, were identified as suitable environments for both H5N1 and H5Nx local circulation in scenarios before and after 2020. Regions in North America and South America — including Bolivia, Brazil, Colombia, Ecuador, Chile, Peru, and Venezuela — demonstrated ecological suitability for the risk of local circulation for H5Nx and H5N1, with a marked increase in the predicted suitability for H5N1 after 2020. For H5Nx post-2020, areas of high predicted ecological suitability, such as Brazil, Bolivia, the Caribbean islands, and Jilin province in China, likely result from spatial extrapolations, as these regions reported few or no outbreaks in the training data.

Our findings were overall consistent with those previously reported by Dhingra and colleagues (Dhingra et al., 2016 [\[link\]](#)), who used data from January 2004 to March 2015 for domestic poultry. However, some differences were noted: their maps identified higher ecological suitability for H5 occurrences before 2016 in North America, West Africa, eastern Europe, and Bangladesh, while our maps mainly highlight ecologically suitable regions in China, South-East Asia, and Europe (**Fig. S5** [\[link\]](#)). In India, analyses consistently identified high ecologically suitable areas for the risk of local H5Nx and H5N1 circulation for the three time periods (pre-2016, 2016-2020, and post-2020). Similar to the results reported by Dhingra and colleagues, we observed an increase in the ecological suitability estimated for H5N1 occurrence in South America's domestic bird populations post-2020. Finally, Dhingra and colleagues identified high suitability areas for H5Nx occurrence in North America, which are predicted to be associated with a low ecological suitability in the 2016-2020 models.

For H5Nx and H5N1 cases in wild birds, the estimated ecological niches were primarily associated with environmental factors such as open water and distance to water, which result in ecological suitability hotspots estimated near coasts and main rivers. For both H5Nx and H5N1, certain areas of predicted high ecological suitability appear spatially isolated, i.e. surrounded by regions of low predicted ecological suitability. These areas likely meet the environmental conditions associated with past HPAI occurrences, but their spatial isolation may imply a lower risk of actual occurrences, particularly in the absence of nearby outbreaks or relevant wild bird movements. Some of the areas with high predicted ecological suitability reflect the result of extrapolations. This is particularly the case in coastal regions of West and North Africa, the Nile Basin, Central Asia (Kyrgyzstan, Tajikistan, Uzbekistan), Brazil (including the Amazon and coastal areas), southern Australia, and the Caribbean, where ecological conditions are similar to those in areas where outbreaks are known to occur but where records of outbreaks are still rare.




Figure 3

areas ecologically suitable for local H5N1 or H5Nx circulation leading to infection cases in domestic bird populations.

We estimated the ecological suitability for two different time periods (2015-2020 and 2020-2022) and for both wild and domestic bird populations. Dynamic visualisations of the results are available here: <https://mood-platform.avia-gis.com/core>.

We observed that areas of relatively high ecological suitability have expanded after 2020. North America, particularly near the Great Lakes region, appeared more suitable for local H5Nx circulation. This expansion was also evident in Russia and South America, aligning with major bird migration routes. Notably, while H5Nx had not been reported in Australia, the models indicate potential ecological suitability there as well. Expanded ecological suitability was also observed in China, India, and European countries. However, models trained on pre-2020 data maintained reasonable predictive performance when tested on post-2020 data, suggesting that the overall ecological niche of HPAI did not drastically shift over time.

In **Table 1** [↗](#), we report the estimation of avian species diversity indices for species involved in HPAI outbreaks for the pre-2020 and post-2020 periods. Note that these indices reflect the diversity of bird species detected in outbreak records, not necessarily their abundance in the wild. We observed variations between these two periods both in the overall bird population and within specific wild bird species groups, including sea bird species (e.g., gulls, terns, boobies, gannets). For all birds, the Shannon index increased from 4.23 before 2020 to 5.03 after 2020, which might indicate a more diverse infected bird population in the latter period. The Simpson index however only rose from 0.97 to 0.98, but which could potentially suggest a slightly higher concentration of certain species post-2020. Sea birds exhibited a similar upward trend in diversity post-2020, with the Shannon index increasing from 2.19 to 2.26. This latter trend could potentially suggest a broader diversity of sea bird species involved in HPAI outbreaks.

When considering wild birds affected by the H5N1 strain of HPAI, there is a notable increase in the Shannon index from 2.00 before 2020 to 2.95 after 2020, accompanied by a rise in the Simpson index from 0.83 to 0.91. These increases could be indicative of greater diversity and a more even distribution of wild bird species among the reported HPAI H5N1 wild bird cases after 2020. In contrast, birds not affected by the H5N1 strain showed a slight decrease in diversity post-2020. The Shannon index decreases from 3.42 to 3.26, and the Simpson index decreases from 0.95 to 0.92. Given that HPAI particularly affected sea birds >2020, we further explore the yearly distribution of sea bird families. As depicted in Figure S4, positive cases among sea birds belonging to the families Laridae (gulls and terns), Sulidae (gannets and boobies), Ardeidae (herons), Pelecanidae (pelicans), Spheniscidae (penguins), Alcidae (auks), Phalacrocoracidae (cormorants) were detected more frequently from 2020 in comparison to before 2020. Notably, the occurrence of Laridae and Sulidae species among positive cases increased markedly from 2021 onwards.

To evaluate whether the post-2020 increase in species diversity estimated for infected wild birds could result from an increase in the number of tests performed on wild birds, we compared European annual surveillance test counts (EFSA et al., 2025 [↗](#), 2019 [↗](#)) before and after 2020 using a Wilcoxon rank-sum test. We relied on European data because it was readily accessible and offered standardised and systematically collected metrics across multiple years, making it suitable for a comparative analysis. Although borderline significant (p-value = 0.063), the Wilcoxon rank-sum test indeed highlighted a recent increase in the number of wild bird tests (on average >11,000/year pre-2020 and >22,000 post-2020), which indicates that the comparison of bird species diversity metrics should be interpreted with caution. However, such an increase in the number of tests conducted in the context of a passive surveillance framework would thus also be in line with an increase in the number of wild birds found dead and thus tested. Therefore, while the increase in the number of tests could indeed impact species diversity metrics such as the Shannon index, it can also reflect an absolute higher wild bird mortality in line with a broadened range of infected bird species.

	All birds		Sea birds		Wild birds		Domestic birds		All birds H5N1		All birds non-H5N1	
	<i>Shannon</i>	<i>Simpson</i>	<i>Shannon</i>	<i>Simpson</i>	<i>Shannon</i>	<i>Simpson</i>	<i>Shannon</i>	<i>Simpson</i>	<i>Shannon</i>	<i>Simpson</i>	<i>Shannon</i>	<i>Simpson</i>
< 2020	4.227	0.971	2.187	0.843	3.072	0.832	3.919	0.966	2.003	0.833	3.415	0.945
> 2020	5.033	0.982	2.255	0.831	5.066	0.983	2.967	0.897	2.952	0.907	3.254	0.914

Table 1.

Species diversity indices estimated from infected bird cases before and after 2020.

This table presents the Shannon and Simpson species diversity indices for various bird groups, comparing values before and after the year 2020. The indices are provided for all birds, sea birds, wild birds, domestic birds, as well as all birds affected by H5N1 and non-H5N1 strains.

Discussion

Using an ecological niche modelling approach, our study presents an analysis of the ecological suitability, interpreted here as the risk of local circulation, for H5Nx and H5N1 HPAI viruses across different temporal and ecological contexts. By conducting boosted regression tree (BRT) analyses and incorporating a diverse set of spatial predictors, our study provides insights into the environmental and anthropogenic factors that are correlated with the distribution of these viruses in both wild and domestic bird populations. For domestic birds, our results concur with those previously reported by Dhingra and colleagues (Dhingra et al., 2016), indicating that ecological niche models incorporating host variables (chicken and duck population densities) outperform those based on land-use or eco-climatic factors in terms of predictive performance. Contrariwise, in our analysis of H5N1 and H5Nx HPAI in wild birds, the most significant predictors were anthropogenic factors, notably urban areas and distance to open water. However, observation biases might affect these findings since outbreaks in wild birds are commonly reported near human populations.

In our analyses, intensive chicken population density emerges as a significant predictor, underscoring the strong association of poultry farming practices with HPAI spread (Dhingra et al., 2016) and revealing a complex interplay between ecological niches and the anthropogenic landscape. We note, however, that intensive poultry density may reflect both surveillance intensity and epidemiological risk, and its predictive role in our models should be interpreted in light of both processes. Similarly to the results of Dhingra and colleagues (Dhingra et al., 2016), land-use and climatic predictors do not play an important role in the niche ecological models, even for wild birds. This suggests that HPAI H5Nx is not as strongly environmentally constrained as vector-borne pathogens, for which clear eco-climatic boundaries (e.g., vector borne diseases) can be mapped (Dhingra et al., 2016). Additionally, we found that while domestic duck population density is a significant predictor for HPAI occurrence in domestic birds before 2020, its importance has decreased post-2020, particularly within the H5Nx ecological niche model. This observation is particularly noteworthy, as domestic ducks have historically been closely linked to areas of persistence and evolution of H5N1 HPAI (Gilbert and Pfeiffer, 2012). However, our findings, in line with earlier studies such as the one of Dhingra and colleagues (Dhingra et al., 2016), highlight a significant shift in the epidemiological dynamics of HPAI. Specifically, while the association between the H5Nx clade 2.3.4.4 and domestic ducks has for instance been documented in South Korea (Hill et al., 2015), it is becoming increasingly evident that HPAI is now more strongly associated with high-density chicken farming and anthropogenic factors. This shift suggests that the virus could now be circulating more among farms (farm to farm transmission) (FAO, 2023), with fewer introductions by wild birds — particularly wild ducks, which have previously acted as “Trojan horses” for H5N1 HPAI (Kim et al., 2014). However, this remains an interpretation, as the available data do not allow us to distinguish between index cases and secondary transmission events. Consequently, we could be witnessing a shift from the traditional duck/rice ecosystems towards a scenario where intensive chicken farming and a wider diversity of wild birds play a more prominent role in the transmission dynamics of HPAI. Therefore, monitoring areas with high intensive chicken densities and regular surveillance of wild birds remain crucial for the early detection and management of HPAI outbreaks.

A difference between the ecological niche models previously trained by Dhingra and colleagues (Dhingra et al., 2016) and our models is the pronounced increase in ecological suitability for H5N1 in domestic bird populations estimated in India and China. More generally, our models highlight the extensive spread of H5N1 and H5Nx HPAI among domestic birds in Asia previously identified as “reassortment sink areas” (Dhingra et al., 2018). Reassortment refers to the process where influenza viruses exchange genetic material, potentially leading to new, more virulent

strains. The extent of virus suitability in Asia highlights a potential change in the dynamics of virus spread, possibly driven by intensified poultry farming practices and increased human-wildlife interactions in these regions (Bahl et al., 2016 [↗](#); Gilbert et al., 2017 [↗](#)).

The recent outbreaks of infection in cattle within the United States have been partially delineated by our HPAI risk mapping approach, highlighting major hotspots predominantly in the Great Lakes region, including Michigan. From March to July 2024, approximately 26 herds in Michigan were affected (U.S. Department of Agriculture, 2024 [↗](#)). Additionally, our risk maps identify localised hotspots around urban centres in Colorado and Texas, where herds have also been infected. These outbreaks appear to stem from a single introduction event in Texas by wild birds (Worobey et al., 2024a [↗](#), 2024b [↗](#)). This trend may be indicative of an increasing risk of future introductions from wild birds, as we observed a significant rise in the diversity of infected bird species after 2020, particularly among sea birds.

The latest strain of H5N1 (H5 clade 2.3.4.4b), believed to have emerged in 2014, shows adaptations for infecting a wide range of wild birds and mammals (Caliendo et al., 2022 [↗](#); Graziosi et al., 2024 [↗](#)). By 2020, its spread among wild birds was three times faster than in farmed poultry due to mutations enabling the virus to infect diverse species (CDC, 2024; Xie et al., 2023). Unlike previous epidemic waves in Europe, outbreaks in the last years have also continued through summer, confirming that the virus's current circulation is no longer seasonal but nesting in the wild bird population (Sciensano, 2024 [↗](#)). Therefore, the virus is able to infect a wider range of species than previous forms, as it circulates all seasons (Sacristán et al., 2024 [↗](#)). The densely packed colonies of sea birds have also facilitated the rapid virus spread, resulting in high mortality in these sea bird colonies (Boulinier, 2023 [↗](#); Bregnballe et al., 2024 [↗](#)). For instance, the Netherlands lost up to 80% of its Sandwich terns in weeks, and the UK saw significant outbreaks, especially on Scottish islands (Knief et al., 2024 [↗](#)). In South America, similar trends have been observed, with notable die-offs in seabird populations along coastal regions, particularly impacting species such as penguins, pelicans and cormorants (Leguia et al., 2023 [↗](#)). Since then, in early 2023, the Peruvian outbreak had spread to marine mammals, particularly affecting the South American sea lion, which also began to experience a mass die-off (Campagna et al., 2024 [↗](#); Leguia et al., 2023 [↗](#)).

The increase in the number of wild bird species being infected by HPAI, particularly H5N1, could be attributed to several factors such as viral evolution, increased interaction between domestic and wild birds, climate and land-use changes, among others. Regarding land-use changes, large parts of Europe have become increasingly fragmented due to the expansion of urban and transport infrastructure (European Environment Agency (EEA), 2022 [↗](#)). This habitat loss and fragmentation can significantly impact water availability and trophic resources (Santos-Tovar et al., 2024 [↗](#)), which could in turn aid the spread and transmission of avian influenza viruses due to relocation of birds and outbreaks (Yin et al., 2022 [↗](#)). Climate change has also been reported as a factor that could affect wild bird distribution in different ways (Gilbert et al., 2008 [↗](#)). As global climate conditions change, avian migratory patterns and routes are also changing (Hitch and Leberg, 2007 [↗](#); Van Doren, 2022 [↗](#)). Furthermore, higher temperatures and extreme weather have resulted in large-scale population shifts in a range of temperate species (Dezfuli et al., 2022 [↗](#)). Overall, these elements therefore raise the question of the role of climate and land-use changes in the recent shifting epidemiology of HPAI in wild birds.

The increasing diversity of infected bird species is part of a broader resurgence of H5N1 globally. Phylogenetic analysis from Europe revealed significant genetic diversity in HPAI-H5N1 viruses since October 2020, likely due to multiple reassortment events with LPAI viruses. Notable introductions from Russia to Europe were observed through autumn migrations of wild birds (ECDC, 2021 [↗](#)). Additionally, the transatlantic transport of HPAI H5N1, documented in Canada in December 2021, suggests the virus can spread across continents, a route previously seen only with LPAI viruses (Caliendo et al., 2022 [↗](#); Dusek et al., 2014 [↗](#); Huang et al., 2014 [↗](#)). In November

2022, H5N1 (clade 2.3.4.4b) was introduced into South America, presumably through the movements of migratory wild birds travelling south during the boreal winter. The virus spread rapidly and caused significant mortality in wild birds and marine mammals across multiple countries (Leguia et al., 2023 [↗](#); (Banyard et al., 2023 [↗](#); Campagna et al., 2024 [↗](#); Gamarra-Toledo et al., 2023 [↗](#); WOA, 2024 [↗](#)).

Interestingly, our models spatially extrapolate ecological suitability for H5Nx in wild birds in areas where few or no outbreaks have been reported. This discrepancy may be explained by limited surveillance or underreporting in those regions. For instance, there is significant evidence that Kazakhstan and Central Asia play a role as a centre for the transmission of avian influenza viruses through migratory birds (Amirgazin et al., 2022 [↗](#); FAO, 2005 [↗](#); Sultankulova et al., 2024 [↗](#)). However, very few wild bird cases are reported in EMPRES-i. In contrast, Australia appears environmentally suitable in our models, yet no incursion of HPAI H5N1 2.3.4.4b has occurred despite the arrival of millions of migratory shorebirds and seabirds from Asia and North America. Extensive surveillance in 2022 and 2023 found no active infections nor evidence of prior exposure to the 2.3.4.4b lineage (Wille et al., 2024 [↗](#); Wille and Klaassen, 2023 [↗](#)).

Despite the robustness of our ecological niche models, some limitations must be acknowledged. Firstly, the accuracy of our models is inherently dependent on the quality and completeness of the occurrence data used, which is influenced by the different reporting systems in place in different countries. Secondly, the definitions of key explanatory variables, such as intensive versus extensive production, may vary across regions, adding complexity to global predictions. To address these limitations, future work could focus on developing country-level or regional models that incorporate localised data and context-specific definitions. Thirdly, underreporting and biases in outbreak reports, particularly in wild bird populations, can skew model predictions and underestimate areas at risk. Furthermore, another limitation of our models for wild bird populations is the lack of incorporation of migratory patterns and behaviours. Migratory birds have complex movement patterns influenced by seasonal changes, food availability, and habitat conditions. Our models, using static spatial predictors, do not capture these dynamic routes and stopover sites, leading to potential inaccuracies in predicting areas at risk for HPAI circulation. In future work, the integration of dynamic migratory data and environmental changes into our ecological niche models could provide a more comprehensive and accurate risk assessment for HPAI spread among wild bird populations. In addition, aligning outbreak occurrences with seasonally matched environmental variables could further refine predictions of HPAI risk linked to migratory dynamics. Finally, our models for domestic poultry do not distinguish between primary introduction events (e.g., spillover from wild birds) and secondary transmission between farms due to limitations in the available surveillance data. While environmental factors likely influence the risk of initial spillover events, secondary spread is more often driven by anthropogenic factors such as biosecurity practices and poultry trade, which are not included in our current modelling framework.

Despite these limitations, our study provides significant contributions to the understanding of H5N1 and H5Nx HPAI virus dynamics. By integrating a wide range of spatial predictors and using ecological niche modelling techniques, we provide a quantitative analysis on the factors driving virus distribution. Overall, our study underscores the importance of continuous surveillance and the need for adaptive management strategies to address the evolving threat of HPAI viruses globally.

Materials and Methods

Our approach is similar to the one implemented by Dhingra and colleagues (2016) [↗](#). While Dhingra et al. (2016) [↗](#) developed their models only for domestic birds over the 2003-2016 period, our models were developed for two host groups separately (wild and domestic birds) and for two

time periods (2016-2020 and 2020-2023).

HPAI occurrence data acquisition

The dataset containing highly pathogenic avian influenza (HPAI) occurrences was constructed by extracting relevant information from the EMPRES-i database spanning the period from January 5, 2015, to March 7, 2023. This dataset encompasses comprehensive details pertaining to H5N1 and H5Nx subtypes, including the family of birds involved in transmission, observation and reporting dates, location coordinates, and the respective countries where the reports originated. The dataset comprises a total of 17,410 H5Nx and 8,383 H5N1 reported cases. It is important to note that the EMPRES-i database does not distinguish between index cases (e.g., primary spillover from wild birds) and secondary farm-to-farm transmissions. As such, our ecological niche models are trained on confirmed HPAI outbreaks in poultry that may result from different transmission dynamics — including both initial introduction events influenced by environmental factors and subsequent spread within poultry systems.

Environmental data acquisition

Previous literature reviews (Gilbert and Pfeiffer, 2012; Dhingra et al., 2016) have summarised the predictor variables commonly linked to occurrences of HPAI. Here, we explored four sets of environmental variables considered by Dhingra and colleagues (Dhingra et al., 2016), and which are detailed in the table in Supplementary Information Resources S1: host variables (set 1), land cover variables (set 2), eco-climatic variables (set 3), and a risk-based selection of variables performed by Dhingra and colleagues (set 4).

Set 1 includes $\log_{10}$ -transformed extensive and intensive chicken density (Gilbert et al., 2015), domestic duck density (Robinson et al., 2014) and human population density from the Gridded Population of the World (Center For International Earth Science Information Network-CIESIN-Columbia University, 2017) database. The chicken density data includes GIS layers representing the global distribution of chickens, categorised into extensive and intensive systems. It uses the model published by Gilbert et al. (2018) applied to the Gridded Livestock of the World (GLW) version 3 chicken layers (Gilbert et al., 2022a). The global domestic duck distribution data were computed using the GLW version 4 (Gilbert et al., 2022b). Set 2 includes land cover information, which was obtained from the global 1-km Consensus Land Cover database by (Tuanmu and Jetz, 2014) Tuanmu and Jetz (2014), and identifies different land cover categories by providing a percentage of representation of each land cover category within raster cells of approximately one square kilometre (<http://www.earthenv.org/landcover.html>). Additionally, this dataset was supplemented by a layer about the distance of each spatial point to the open water. Set 3 encompasses seasonal and large-scale patterns of eco-climatic indices like daytime land surface temperature (day LST) and the normalised difference vegetation index (NDVI), as documented by Scharlemann and colleagues (Scharlemann et al., 2008). Finally, set 4 is a selection from the previous sets based on prior epidemiological knowledge (Gilbert and Pfeiffer, 2012; Dhingra et al., 2016).

Ecological niche modelling

The different ecological niche modelling analyses were performed with the boosted regression trees (BRT), available in the R package “dismo” (Hijmans et al., 2017). BRT is a machine learning approach that can be used to generate a collection of sequentially fitted regression trees optimising the predictive probability of occurrence given local environmental conditions (Elith et al., 2006). The predictive probability, interpreted as ecological suitability for local virus circulation, ranges from “0” (unsuitable environmental conditions) and “1” (highly suitable environmental conditions). BRT allows us to model complex non-linear relationships between the

response and various predictor variables (Elith et al., 2008). It has also been demonstrated that the BRT approach had a superior predictive performance compared to alternative modelling approaches (Elith et al., 2006 [↗](#)).

We selected 2020 as a cut-off point to reflect a well-documented shift in HPAI epidemiology, notably the emergence and global spread of clade 2.3.4.4b. This event marked a turning point in viral dynamics, influencing both the range of susceptible hosts and the geographical distribution of outbreaks. Based on this distinction, we performed ten replicated BRT analyses using each of the seven occurrence datasets: (i) H5Nx cases in wild birds < 2020 (01/01/2015 - 31/12/2019), (ii) H5N1 cases in wild birds > 2020 (01/01/2020 - 06/03/2023), (iii) H5Nx cases in wild birds > 2020 (01/01/2020 - 06/03/2023), (iv) H5Nx cases in domestic birds < 2020 (01/01/2015 - 31/12/2019), (v) H5N1 cases in domestic birds < 2020 (01/01/2015 - 31/12/2019), (vi) H5N1 cases in domestic birds > 2020 (01/01/2020 - 06/03/2023), and (vii) H5Nx cases in domestic birds > 2020 (01/01/2020 - 06/03/2023). Of note, there was not enough occurrence data to train ecological niche models on H5N1 cases in wild birds < 2020 (only 33 occurrence data from eight different countries — Bangladesh, Bulgaria, China, India, Kazakhstan, Nepal, Romania, and Russia — between the 01/01/2015 and 31/12/2019).

We ran BRT analyses using both occurrence and “pseudo-absence” data, the latter corresponding to randomly sampled absence points within the study area (Elith et al., 2006 [↗](#)). To account for heterogeneity in AIV surveillance and minimise the risk of sampling pseudo-absences in poorly monitored regions, we restricted our analysis to countries (or administrative level 1 units in China and Russia) with at least five confirmed outbreaks. Unlike Dhingra et al. (2016) [↗](#), who sampled pseudo-absences across a broader global extent, our sampling was limited to regions with demonstrated surveillance activity. In addition, we adjusted the density of pseudo-absence points according to the number of reported outbreaks in each country or admin-1 unit, as a proxy for surveillance effort — an approach not implemented in this previous study.

In addition, the sampling pseudo-absence points was based on the human population distribution to further account for potential within country differences in surveillance intensity (Dhingra et al., 2016 [↗](#)): the probability to sample a pseudo-absence from a given grid cell was proportional to the log-transformed human population density in that cell. The human population density raster was retrieved from the Gridded Population of the World (GPW) project, which was downsampled to a resolution of 5 arcminutes. Another approach to sampling pseudo-absences would have been to distribute them according to the density of domestic poultry. However, as pointed out by Dhingra and colleagues (2016) [↗](#), the locations of outbreaks in the EMPRES-i database are often georeferenced using place name nomenclatures due to a lack of accurate GPS data, which could introduce a spatial bias towards populated areas. Therefore, the use of human population density as an indicator of surveillance effort takes into account not only the probability of detection, but also the geo-referencing bias in household records. To further reduce bias from unsuitable regions where the pathogen may not have been introduced, we restricted pseudo-absence sampling to areas located at a minimum of 10 km and a maximum of 1,000 km from known presence points (Phillips et al., 2009; Dhingra et al., 2016 [↗](#)). Initially, ten times more pseudo-absences than presence points were simulated, but we ultimately retained three times more pseudo-absences than presences in each country/admin-1 unit.

Given that spatial data typically exhibits autocorrelation, employing a standard cross-validation procedure may result in an overestimation of the BRT model’s accuracy (Randin et al., 2006). To circumvent this issue, we employed a spatial cross-validation procedure to select the optimal number of trees in each BRT model. Specifically, we tested and compared two distinct spatial cross-validation procedures: (i) the procedure introduced by Dhingra and colleagues and based on (five) reference presence points used to cluster the presence and pseudo-absence points in (five) folds according to their nearest distance to a reference point, and (ii) the procedure introduced by Valavi and colleagues and based on a block generation method (Valavi et al., 2019 [↗](#)). In the first

spatial cross-validation procedure, the five reference presence points were randomly selected among all presence points. If each of the five generated spatial folds did not gather at least 10% of the presence points, the random selection of reference points was re-initiated. To compare the performance of the three cross-validation procedures applied in this study (the standard procedure and the spatial procedures based on the reference points or blocks generation), we estimated the spatial sorting bias (SSB) metric (Hijmans, 2012 [↗](#)) ranging from 0 to 1, an SSB near 0 and 1 indicating an extreme impact and the absence of an impact of spatial autocorrelation on the model training, respectively.

All BRT analyses were run and averaged over 10 cross-validated replicates, with a tree complexity of 4, a learning rate of 0.01, a tolerance parameter of 0.001, and while considering 5 spatial folds. Each model was initiated with 10 trees, and additional trees were incrementally added (in steps of 5) up to a maximum of 10,000, with the optimal number selected based on cross-validation tests. We evaluated the inferences using the area under the receiver operating characteristic (ROC) curve, also simply referred to as “area under the curve” (AUC). We further used the AUC metric to assess the capacity of our models trained on occurrence data < 2020 to predict > 2020 distribution of occurrence data. To assess how each environmental factor contributed to the different BRT models, we computed their relative influence (RI) in those models and their response curves showing how ecological suitability varies with one specific factor. RI values were obtained by evaluating the number of times a specific environmental factor was selected for splitting a tree, weighted by the squared improvement to the model as a result of each split, averaged over all trees (Elith et al., 2008).

We explored the four sets of environmental variables assembled by Dhingra and colleagues (Dhingra et al., 2016 [↗](#)) and described above in the “Environmental data acquisition” subsection. The BRT analyses presented here were eventually performed based on set 2 for wild bird cases and set 4 for domestic bird cases. This selection is aligned with the methodology of Dhingra and colleagues (2016 [↗](#)), prioritising variables which demonstrated the best predictive accuracy and clear epidemiological significance to minimise the potential for coincidental correlations. The RI values obtained when analysing each set of environmental variables separately are all reported in a table in Supplementary Information Resources S2.

Bird diversity indices calculation

Diversity index calculations were performed using the HPAI dataset. Cases lacking information on the date of collection, bird family, and subtype were excluded from the analyses. The diversity of HPAI cases across bird families was quantified using two indices: the Shannon diversity index and the Simpson diversity index. The Shannon index accounts for evenness of bird families (Shannon, 1948 [↗](#)), whereas the Simpson index provides a measure of dominance (Simpson, 1949 [↗](#)), emphasising the most abundant families in the dataset. Simpson index ranges between 0 and 1, where 1 represents infinite diversity and 0 no diversity. Calculations were done using two time points: before and after 2020, and for various data classifications: overall and annually, across all bird families and specifically sea bird families, as well as for all subtypes (HxNx) and those identified as H5N1 subtype, wild birds, and poultry. These indices were calculated using the R package “vegan” (Oksanen et al., 2023): the dataset was first aggregated by year, HPAI subtype and bird family, and we then used the “diversity” function to compute the Shannon and Simpson indices for each year.

Data and code accessibility

All data were retrieved from EMPRES-i database, and are publicly available. R scripts and related files needed to run the analyses, as well as Supplementary Information Resources S1 (description and source of each environmental variable included in the original sets of variables) and S2 (mean




Figure S1

Environmental factors included in the ecological niche modelling.

“LST” refers to “land surface temperature”, “NDVI” to the “normalised difference vegetation index”, and “EVI” to the “enhanced vegetation index”.




Figure S1 (continued)




Figure S2

Comparison between the presence/pseudo-absence data used to train the ecological niche models (1° column) and the resulting ecological niche suitability maps (2° column).

In the first column, presence and sampled pseudo-absence points are displayed as red and dark points, respectively. As detailed in the Materials and Methods section, pseudo-absence points were only sampled in countries in which there was at least one presence point (for either H5N1 or H5Nx, the considering time period, and bird populations). For Russia and China, we considered the admin-1 administrative areas instead of the country to define if pseudo-absence points should be sampled in the corresponding area. Specifically, for a given administrative region (i.e. a given country or admin-1 region in case of Russia or China) gathering at least one presence point, we sampled a number of pseudo-absence points equal to three times the number of presence points in that region. Furthermore, pseudo-absence points were sampled according to the log-transformed raster of human population density, which prevented the sampling of pseudo-absence in unpopulated areas non associated with any surveillance activity.




Figure S2 (continued)

Figure S3

boxplots reporting the sorting sampling bias (SSB) and area under the curve (AUC) of the receiving operator curve metrics associated with various ecological niche models trained for H5N1 and H5Nx.

For the SSB estimates, we report values obtained for the standard cross-validation (A), as well as spatial cross-validation procedures based on the reference points (B) and the blocks generation (C) approaches (see the Materials and Methods section for further detail). SSB estimates range from 0 to 1; values close to 0 indicate a strong impact of spatial autocorrelation on model training, while values close to 1 suggest little to no impact. The AUC metrics computed for the ecological niche models trained with the spatial cross-validation procedures based on the “epicentres”, and we here report the values obtained for the models trained on each of the four distinct sets of environmental factors.



Figure S4

distribution of H5Nx and H5N1 occurrence records in sea birds from 2015 to 2023, categorised by bird family.

The three panels successively show the total occurrence records for all H5Nx subtypes, occurrence records for non-H5N1 H5Nx subtypes, and all H5N1 occurrence records.




Figure S5

comparison between the ecological niche suitability estimated for H5N1 and H5Nx before 2016 and after 2020 for domestic bird populations.

Specifically, we here compare the ecological niche models trained by [Dhingra et al. \(2016\)](#) on occurrence records collected between January 2004 to March 2015 (< 2016) and the ecological niche models trained in the present study on occurrence records collected between January 2020 to March 2023 (> 2020).




Table S1

Assessment of the predictive performance of ecological niche models to predict present and past ecological suitability maps.

We here report, for each replicate analysis, a measure of the predictive performance of the trained ecological niche through the computation of the area under the receiver operating characteristic (ROC) curve, also simply referred to as “area under the curve” (AUC). (*) refers to AUC estimates obtained when assessing the capacity of models trained on occurrence data < 2020 to predict > 2020 distribution of occurrence data.

	Wild birds			Domestic birds					
	H5Nx			H5N1			H5Nx		
	< 2020	> 2020	>2020*	< 2020	> 2020	> 2020*	< 2020	> 2020	> 2020*
Replicate 1	0.899	0.793	0.700	0.832	0.812	0.769	0.833	0.791	0.755
Replicate 2	0.896	0.769	0.703	0.832	0.822	0.769	0.829	0.781	0.752
Replicate 3	0.895	0.796	0.707	0.834	0.820	0.770	0.818	0.784	0.743
Replicate 4	0.894	0.773	0.705	0.843	0.820	0.769	0.828	0.804	0.751
Replicate 5	0.893	0.797	0.706	0.837	0.816	0.770	0.828	0.780	0.752
Replicate 6	0.898	0.797	0.701	0.845	0.816	0.770	0.829	0.780	0.752
Replicate 7	0.897	0.798	0.700	0.835	0.816	0.769	0.827	0.775	0.750
Replicate 8	0.894	0.805	0.705	0.836	0.825	0.770	0.828	0.777	0.752
Replicate 9	0.894	0.770	0.702	0.834	0.813	0.769	0.828	0.779	0.750
Replicate 10	0.894	0.775	0.703	0.835	0.823	0.769	0.832	0.795	0.754

	Wild birds			
	H5N1		H5Nx	
	< 2020	> 2020	< 2020	> 2020
Evergreen deciduous needleleaf trees	-	1.6 % [1.5-1.6]	1.6 % [1.3-1.7]	0.4 % [0.4-0.5]
Evergreen broadleaf trees	-	1.3 % [1.2-1.4]	0.1 % [0.1-0.1]	1.0 % [0.8-1.1]
Deciduous broadleaf trees	-	4.1 % [4.0-4.2]	1.6 % [1.5-1.8]	6.1 % [5.9-6.2]
Mixed and other trees	-	11.5 % [11.4-11.7]	5.3 % [4.9-5.4]	11.4 % [11.3-11.8]
Shrublands	-	2.3 % [2.0-2.3]	3.2 % [3.0-3.3]	2.4 % [2.1-2.7]
Herbaceous vegetation	-	8.8 % [8.8-9.0]	2.8 % [2.5-3.0]	9.5 % [9.5-9.6]
Cultivated and managed vegetation	-	4.4 % [4.3-4.5]	4.8 % [4.6-4.9]	5.8 % [5.4-6.1]
Regularly flooded vegetation	-	0.3 % [0.3-0.3]	0.1 % [0.1-0.1]	0.2 % [0.1-0.3]
Urban and built-up areas	-	39.3 % [39.0-39.9]	54.5 % [53.5-55.0]	37.5 % [36.5-38.2]
Open water areas	-	22.0 % [21.9-22.3]	25.5 % [25.3-25.8]	23 % [22.8-23.3]
Distance to water	-	4.1 % [3.9-4.3]	0.9 % [0.7-1.0]	2.8 % [2.4-3.1]
	Domestic birds			
	H5N1		H5Nx	
	< 2020	> 2020	< 2020	> 2020
Duck pop. density (2010, log)	18.2 % [17.8-18.4]	15.7 % [13.3-16.4]	28.7 % [28.4-28.8]	12.9 % [12.9-13.5]
Extensive chicken pop. density (2015, log)	4.4 % [4.2-4.5]	2.3 % [0.5-2.6]	3.9 % [3.7-4.0]	2.7 % [2.5-3.4]
Intensive chicken pop. density (2015, log)	8.5 % [8.4-8.7]	30.4 % [29.6-36.2]	29.6 % [29.3-30.1]	43.1 % [40.5-43.4]
Human pop. density (2020, log)	54.9 % [53.9-55.1]	20.7 % [20.4-21.6]	15.5 % [15.3-15.7]	14.9 % [14.8-15.2]
Cultivated and managed vegetation	9.5 % [9.3-9.7]	16.6 % [16.3-18.5]	7.1 % [7.0-7.2]	13.2 % [12.9-13.3]
Open water areas	0.5 % [0.5-0.6]	1.4 % [0.3-1.5]	0.0 % [0.0-0.0]	0.6 % [0.4-0.8]
Day LST annual mean	4.1 % [4.0-4.2]	12.9 % [9.6-13.3]	15.4 % [15.1-15.6]	13.0 % [12.4-13.9]

Table S2

relative influence (RI, in %) of each environmental variable in the respective ecological models trained on a specific dataset of occurrence data.

A RI estimate was obtained for each replicate BRT analysis and we here report median as well as first and third quartile RI values.

relative influence computed for the environmental variables within each considered set of variables), are all available at https://github.com/sdellicour/h5nx_risk_mapping .

Acknowledgements

MFVG and SD acknowledge funding from the European Union Horizon 2020 project MOOD (grant agreement n°874850). MCD was supported by the UKRI GCRF One Health Poultry Hub (Grant No. B/S011269/1), one of twelve interdisciplinary research hubs funded under the UK government's Global Challenge Research Fund Interdisciplinary Research Hub initiative. MCD and SD are supported by the BELSPO project BE-PIN. SD also acknowledges support from the *Fonds National de la Recherche Scientifique* (F.R.S.-FNRS, Belgium; grant n°F.4515.22), from the Research Foundation - Flanders (*Fonds voor Wetenschappelijk Onderzoek - Vlaanderen*, FWO, Belgium; grant n°G098321N), from the European Union Horizon 2020 project LEAPS (grant agreement n°101094685), from the ImmunReach project funded by the *Institut d'Encouragement de la Recherche Scientifique et de l'Innovation de Bruxelles* (Innoviris, Belgium), and from the doctoral network VIVACE funded by the Marie Skłodowska-Curie Actions (MSCA) of the European Commission (grant agreement n°101167768).

Additional information

Funding

EC | Horizon 2020 Framework Programme (H2020) (874850)

- Maria F Vincenti-Gonzalez
- Simon Dellicour

UK Research and Innovation (UKRI) (B/S011269/1)

- Marie-Cécile Dupas

Fonds De La Recherche Scientifique - FNRS (FNRS) (F.4515.22)

- Simon Dellicour

Belgian Federal Science Policy Office (BELSPO) (TD/231/BE-PIN)

- Marie-Cécile Dupas
- Simon Dellicour

Fonds Wetenschappelijk Onderzoek (FWO) (G098321N)

- Simon Dellicour

EC | Horizon 2020 Framework Programme (H2020) (101094685)

- Simon Dellicour

References

- Amirgazin A, Shevtsov A, Karibayev T, Berdikulov M, Kozhakhmetova T, Syzdykova L, Ramankulov Y, Shustov AV (2022) **Highly pathogenic avian influenza virus of the A/H5N8 subtype, clade 2.3.4.4b, caused outbreaks in Kazakhstan in 2020** *PeerJ* **10**:e13038 <https://doi.org/10.7717/peerj.13038> | Google Scholar
- Bahl J, Pham TT, Hill NJ, Hussein ITM, Ma EJ, Easterday BC, Halpin RA, Stockwell TB, Wentworth DE, Kayali G, Krauss S, Schultz-Cherry S, Webster RG, Webby RJ, Swartz MD, Smith GJD, Runstadler JA (2016) **Ecosystem Interactions Underlie the Spread of Avian Influenza A Viruses with Pandemic Potential** *PLOS Pathogens* **12**:e1005620 <https://doi.org/10.1371/journal.ppat.1005620> | Google Scholar
- Banyard A, Begeman L, Black J, Breed A, Dewar M, Fijn R. (2023) **Continued expansion of high pathogenicity avian influenza H5 in wildlife in South America and incursions in to the Antarctic Region** OFFLU | Google Scholar
- Boulinier T (2023) **Avian influenza spread and seabird movements between colonies** *Trends in Ecology & Evolution* **38**:391–395 <https://doi.org/10.1016/j.tree.2023.02.002> | Google Scholar
- Bregnballe T, Herrmann C, Globig A, Günther A, Staubach C, Heise JN, Harder T, Beer M, Knief U, Heinicke T, Leivits M, Lundström K, Nurmoja I, Liang Y, Larsen LE, Hjulsager CK, Pohlmann A, Fox AD. (2024) **Outbreaks of highly pathogenic avian influenza (HPAI) epidemics in Baltic Great Cormorant *Phalacrocorax carbo* colonies in 2021 and 2022** *Bird Study* :1–14 <https://doi.org/10.1080/00063657.2024.2399168> | Google Scholar
- Caliendo V, Lewis NS, Pohlmann A, Baillie SR, Banyard AC, Beer M, Brown IH, Fouchier RAM, Hansen RDE, Lameris TK, Lang AS, Laurendeau S, Lung O, Robertson G, Van Der Jeugd H, Alkie TN, Thorup K, Van Toor ML, Waldenström J, Yason C, Kuiken T, Berhane Y. (2022) **Transatlantic spread of highly pathogenic avian influenza H5N1 by wild birds from Europe to North America in 2021** *Sci Rep* **12**:11729 <https://doi.org/10.1038/s41598-022-13447-z> | Google Scholar
- Campagna C, Uhart M, Falabella V, Campagna J, Zavattieri V, Vanstreels RET, Lewis MN (2024) **Catastrophic mortality of southern elephant seals caused by H5N1 avian influenza** *Marine Mammal Science* **40**:322–325 <https://doi.org/10.1111/mms.13101> | Google Scholar
- Center For International Earth Science Information Network-CIESIN-Columbia University (2017) **Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11** <https://doi.org/10.7927/H49C6VHW>
- Dezfuli A, Horton KG, Zuckerberg B, Schubert SD, Bosilovich MG (2022) **Continental Patterns of Bird Migration Linked to Climate Variability** *Bulletin of the American Meteorological Society* **103**:E536–E547 <https://doi.org/10.1175/BAMS-D-21-0220.1> | Google Scholar
- Dhingra MS, Artois J, Dellicour S, Lemey P, Dauphin G, Von Dobschuetz S, Van Boeckel TP, Castellan DM, Morzaria S, Gilbert M. (2018) **Geographical and Historical Patterns in the Emergences of Novel Highly Pathogenic Avian Influenza (HPAI) H5 and H7 Viruses in Poultry** *Frontiers in Veterinary Science* **5** | Google Scholar

- Dhingra MS, Artois J, Robinson TP, Linard C, Chaiban C, Xenarios I, Engler R, Liechti R, Kuznetsov D, Xiao X, Dobschuetz SV, Claes F, Newman SH, Dauphin G, Gilbert M (2016) **Global mapping of highly pathogenic avian influenza H5N1 and H5Nx clade 2.3.4.4 viruses with spatial cross-validation** *eLife* **5**:e19571 <https://doi.org/10.7554/eLife.19571> | Google Scholar
- Dusek RJ, Hallgrimsson GT, Ip HS, Jónsson JE, Sreevatsan S, Nashold SW, TeSlaa JL, Enomoto S, Halpin RA, Lin X, Fedorova N, Stockwell TB, Dugan VG, Wentworth DE, Hall JS (2014) **North Atlantic Migratory Bird Flyways Provide Routes for Intercontinental Movement of Avian Influenza Viruses** *PLOS One* **9**:e92075 <https://doi.org/10.1371/journal.pone.0092075> | Google Scholar
- ECDC (2021) **Avian influenza overview September – December 2021** <https://www.ecdc.europa.eu/en/publications-data/avian-influenza-overview-september-december-2021>
- EFSA, Abrahantes JC, Aznar I, Catalin I, Kohnle L, Mulligan KF, Mur L, Stoicescu A, van Houtum A, Zancanaro G. (2025) **Avian influenza annual report 2023** *Efs2* **23** <https://doi.org/10.2903/j.efsa.2025.9197> | Google Scholar
- EFSA, Brouwer A, Gonzales J, Huneau A, Mulatti P, Kuiken T, Staubach C, Stegeman A, Antoniou S, Baldinelli F, Van der Stede Y, Aznar I. (2019) **Annual Report on surveillance for avian influenza in poultry and wild birds in Member States of the European Union in 2018** *Efs2* **17** <https://doi.org/10.2903/j.efsa.2019.5945> | Google Scholar
- Elith J, Graham C H., Anderson R P., Dudík M, Ferrier S, Guisan A, Hijmans R J., Huettmann F, Leathwick J R., Lehmann A (2006) **Novel methods improve prediction of species' distributions from occurrence data** *Ecography* **29**:129–151 | Google Scholar
- Elsmo EJ, Wünschmann A, Beckmen KB, Broughton-Neiswanger LE, Buckles EL, Ellis J, Fitzgerald SD, Gerlach R, Hawkins S, Ip HS, Lankton JS, Lemley EM, Lenocho JB, Killian ML, Lantz K, Long L, Maes R, Mainenti M, Melotti J, Moriarty ME, Nakagun S, Ruden RM, Shearn-Bochsler V, Thompson D, Torchetti MK, Van Wettene AJ, Wise AG, Lim AL (2023) **Highly Pathogenic Avian Influenza A(H5N1) Virus Clade 2.3.4.4b Infections in Wild Terrestrial Mammals, United States, 2022** *Emerg Infect Dis* **29**:2451–2460 <https://doi.org/10.3201/eid2912.230464> | Google Scholar
- European Environment Agency (EEA) (2022) **Landscape fragmentation pressure in Europe** <https://www.eea.europa.eu/en/analysis/indicators/landscape-fragmentation-pressure-in-europe>
- FAO (2023) **Global consultation on highly pathogenic avian influenza (HPAI)** FAO <https://doi.org/10.4060/cc7302en> | Google Scholar
- FAO (2005) **Potential risk of Highly Pathogenic Avian Influenza (HPAI) spreading through wild water bird migration** <https://openknowledge.fao.org/items/d6185856-dae8-4e8a-a345-1eb34e591a36>
- FAO, WHO, WOA (2023) **Ongoing avian influenza outbreaks in animals pose risk to humans** AnimalHealth <https://www.fao.org/animal-health/news-events/news/detail/ongoing-avian-influenza-outbreaks-in-animals-pose-risk-to-humans/en>
- Forcey GM, Linz GM, Thogmartin WE, Bleier WJ (2007) **Influence of land use and climate on wetland breeding birds in the Prairie Pothole region of Canada** *Can J Zool* **85**:421–436 <https://doi.org/10.1139/Z07-005> | Google Scholar

Gamarra-Toledo V, Plaza PI, Gutiérrez R, Inga-Díaz G, Saravia-Guevara P, Pereyra-Meza O, Coronado-Flores E, Calderón-Cerrón A, Quiroz-Jiménez G, Martínez P, Huamán-Mendoza D, Nieto-Navarrete JC, Ventura S, Lambertucci SA (2023) **Mass Mortality of Marine Mammals Associated to Highly Pathogenic Influenza Virus (H5N1) in South America** *bioRxiv* <https://doi.org/10.1101/2023.02.08.527769> | Google Scholar

Giacinti JA, Signore AV, Jones MEB, Bourque L, Lair S, Jardine C, Stevens B, Bollinger T, Goldsmith D, British Columbia Wildlife AIV Surveillance Program (BC WASP), Pybus M, Stasiak I, Davis R, Pople N, Nituch L, Brook RW, Ojke D, Massé A, Dimitri-Masson G, Parsons GJ, Baker M, Yason C, Harms J, Jutha N, Neely J, Berhane Y, Lung O, French SK, Myers L, Provencher JF, Avery-Gomm S, Robertson GJ, Barychka T, Gurney KEB, Wight J, Rahman I, Hargan K, Lang AS, Montevecchi WA, Burt TV, Brown MGC, Pekarik C, Thompson T, McLaughlin A, Willie M, Wilson L, Flemming SA, Ross MV, Leafloor J, Baldwin F, Sharp C, Lewis H, Beaumont M, Hanson A, Ronconi RA, Reed E, Campbell M, Saunders M, Soos C. (2024) **Avian influenza viruses in wild birds in Canada following incursions of highly pathogenic H5N1 virus from Eurasia in 2021–2022** *mBio* **15**:e03203–23 <https://doi.org/10.1128/mbio.03203-23> | Google Scholar

Gierak A, Śmietanka K. (2021) **The impact of selected risk factors on the occurrence of highly pathogenic avian influenza in commercial poultry flocks in Poland** *Journal of Veterinary Research* **65**:45–52 <https://doi.org/10.2478/jvetres-2021-0013> | Google Scholar

Gilbert M, Cinardi G, Da Re D, Wint WGR, Wisser D, Robinson TP. (2022a) **Global chickens distribution in 2015 (5 minutes of arc)** *Harvard Dataverse* <https://doi.org/10.7910/DVN/SXHLF3>

Gilbert M, Cinardi G, Da Re D, Wint WGR, Wisser D, Robinson TP. (2022b) **Global ducks distribution in 2015 (5 minutes of arc)** *Harvard Dataverse* <https://doi.org/10.7910/DVN/S9ONXV>

Gilbert M, Nicolas G, Cinardi G, Van Boeckel TP, Vanwambeke SO, Wint G, Robinson TP. (2018) **Global distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens and ducks in 2010** *Scientific data* **5**:1–11 | Google Scholar

Gilbert M, Pfeiffer DU (2012) **Risk factor modelling of the spatio-temporal patterns of highly pathogenic avian influenza (HPAIV) H5N1: A review** *Spatial and Spatio-temporal Epidemiology* **3**:173–183 <https://doi.org/10.1016/j.sste.2012.01.002> | Google Scholar

Gilbert M, Slingenbergh J, Xiao X (2008) **Climate change and avian influenza** *Rev Sci Tech* **27**:459–466 | Google Scholar

Gilbert M, Xiao X, Robinson TP (2017) **Intensifying poultry production systems and the emergence of avian influenza in China: a ‘One Health/Ecohealth’ epitome** *Archives of Public Health* **75**:48 <https://doi.org/10.1186/s13690-017-0218-4> | Google Scholar

Graziosi G, Lupini C, Catelli E, Carnaccini S (2024) **Highly Pathogenic Avian Influenza (HPAI) H5 Clade 2.3.4.4b Virus Infection in Birds and Mammals** *Animals* **14**:1372 <https://doi.org/10.3390/ani14091372> | Google Scholar

Hijmans RJ (2012) **Cross-validation of species distribution models: removing spatial sorting bias and calibration with a null model** *Ecology* **93**:679–688 | Google Scholar

Hijmans RJ, Phillips S, Leathwick J, Elith J, Hijmans MRJ. (2017) **Package ‘dismo.’** *Circles* **9**:1–68 | Google Scholar

Hitch AT, Leberg PL (2007) **Breeding Distributions of North American Bird Species Moving North as a Result of Climate Change** *Conservation Biology* **21**:534–539 <https://doi.org/10.1111/j.1523-1739.2006.00609.x> | [Google Scholar](#)

Huang P, Sun L, Li J, Wu Q, Rezaei N, Jiang S, Pan C (2023) **Potential cross-species transmission of highly pathogenic avian influenza H5 subtype (HPAI H5) viruses to humans calls for the development of H5-specific and universal influenza vaccines** *Cell Discov* **9**:1–13 <https://doi.org/10.1038/s41421-023-00571-x> | [Google Scholar](#)

Huang Y, Wille M, Dobbin A, Walzthöni NM, Robertson GJ, Ojic D, Whitney H, Lang AS (2014) **Genetic Structure of Avian Influenza Viruses from Ducks of the Atlantic Flyway of North America** *PLOS One* **9**:e86999 <https://doi.org/10.1371/journal.pone.0086999> | [Google Scholar](#)

Klaassen M, Wille M. (2023) **Wild birds' plight and role in the current bird flu panzootic** *bioRxiv* [Google Scholar](#)

Knief U, Bregnballe T, Alfarwi I, Ballmann MZ, Brenninkmeijer A, Bzoma S, Chabrolle A, Dimmlich J, Engel E, Fijn R, Fischer K, Hälterlein B, Haupt M, Hennig V, Herrmann C, T Veld R, Kirchhoff E, Kristersson M, Kühn S, Larsson K, Larsson R, Lawton N, Leopold M, Lilipaly S, Lock L, Marty R, Matheve H, Meissner W, Morrison P, Newton S, Olofsson P, Packmor F, Pedersen KT, Redfern C, Scarton F, Schenk F, Scher O, Serra L, Sibille A, Smith J, Smith W, Sterup J, Stienen E, Strassner V, Valle RG, Van Bemmelen RSA, Veen J, Vervaeke M, Weston E, Wojcieszek M, Courtens W (2024) **Highly pathogenic avian influenza causes mass mortality in Sandwich Tern *Thalasseus sandvicensis* breeding colonies across north-western Europe** *Bird Conservation International* **34**:e6 <https://doi.org/10.1017/S0959270923000400> | [Google Scholar](#)

Leguia M, Garcia-Glaessner A, Muñoz-Saavedra B, Juárez D, Barrera P, Calvo-Mac C, Jara J, Silva W, Ploog K, Amaro Lady, Colchao-Claux P, Johnson CK, Uhart MM, Nelson MI, Lescano J (2023) **Highly pathogenic avian influenza A (H5N1) in marine mammals and seabirds in Peru** *Nat Commun* **14**:5489 <https://doi.org/10.1038/s41467-023-41182-0> | [Google Scholar](#)

Martin V, Pfeiffer DU, Zhou X, Xiao X, Prosser DJ, Guo F, Gilbert M (2011) **Spatial Distribution and Risk Factors of Highly Pathogenic Avian Influenza (HPAI) H5N1 in China** *PLoS Pathog* **7**:e1001308 <https://doi.org/10.1371/journal.ppat.1001308> | [Google Scholar](#)

Neumann G, Kawaoka Y (2024) **Highly pathogenic H5N1 avian influenza virus outbreak in cattle: the knowns and unknowns** *Nat Rev Microbiol* **22**:525–526 <https://doi.org/10.1038/s41579-024-01087-1> | [Google Scholar](#)

Peacock T, Moncla L, Dudas G, VanInsberghe D, Sukhova K, Lloyd-Smith JO, Worobey M, Lowen AC, Nelson MI (2024) **The global H5N1 influenza panzootic in mammals** *Nature* :1–2 <https://doi.org/10.1038/s41586-024-08054-z> | [Google Scholar](#)

Plaza PI, Gamarra-Toledo V, Euguí JR, Lambertucci SA (2024) **Recent Changes in Patterns of Mammal Infection with Highly Pathogenic Avian Influenza A(H5N1) Virus Worldwide** *Emerg Infect Dis* **30** <https://doi.org/10.3201/eid3003.231098> | [Google Scholar](#)

Prosser DJ, Teitelbaum CS, Yin S, Hill NJ, Xiao X (2023) **Climate change impacts on bird migration and highly pathogenic avian influenza** *Nat Microbiol* **8**:2223–2225 <https://doi.org/10.1038/s41564-023-01538-0> | [Google Scholar](#)

Sacristán C, Ewbank AC, Ibáñez Porras P, Pérez-Ramírez E, De La Torre A, Briones V, Iglesias I. (2024) **Novel Epidemiologic Features of High Pathogenicity Avian Influenza Virus A H5N1**

2.3.3.4b Panzootic: A Review *Transboundary and Emerging Diseases* **2024**:5322378 <https://doi.org/10.1155/2024/5322378> | [Google Scholar](#)

Santos-Tovar A, Ramírez-Bastida P, Navarro-Sigüenza AG, Paz H, Ruiz-Rodríguez A, Vázquez-Reyes LD (2024) **Habitat fragmentation erodes taxonomic and functional diversity of waterbird communities of the South Pacific coast of Mexico** *Ornithol Res* **32**:124–134 <https://doi.org/10.1007/s43388-024-00172-6> | [Google Scholar](#)

Scharlemann JP, Benz D, Hay SI, Purse BV, Tatem AJ, Wint GW, Rogers DJ (2008) **Global data for ecology and epidemiology: a novel algorithm for temporal Fourier processing MODIS data** *PloS one* **3**:e1408 [Google Scholar](#)

Sciensano (2024) **Avian Influenza: Numbers** *sciensano.be* <https://www.sciensano.be/en/health-topics/avian-influenza/numbers>

Shannon CE (1948) **A Mathematical Theory of Communication** *Bell System Technical Journal* **27**:379–423 <https://doi.org/10.1002/j.1538-7305.1948.tb01338.x> | [Google Scholar](#)

Simpson EH (1949) **Measurement of Diversity** *Nature* **163**:688–688 <https://doi.org/10.1038/163688a0> | [Google Scholar](#)

Sultankulova KT, Argimbayeva TU, Aubakir NA, Bopi A, Omarova ZD, Melisbek AM, Karamendin K, Kydyrmanov A, Chervyakova OV, Kerimbayev AA, Burashev YD, Kasymbekov YT, Orynbayev MB (2024) **Reassortants of the Highly Pathogenic Influenza Virus A/H5N1 Causing Mass Swan Mortality in Kazakhstan from 2023 to 2024** *Animals* **14**:3211 <https://doi.org/10.3390/ani14223211> | [Google Scholar](#)

Tuanmu M, Jetz W (2014) **A global 1-km consensus land-cover product for biodiversity and ecosystem modelling** *Global Ecology and Biogeography* **23**:1031–1045 [Google Scholar](#)

U.S. Department of Agriculture A and PHIS (2024) **HPAI Confirmed Cases in Livestock** <https://www.aphis.usda.gov/livestock-poultry-disease/avian/avian-influenza/hpai-detections/hpai-confirmed-cases-livestock>

Valavi R, Shafizadeh-Moghadam H, Matkan A, Shakiba A, Mirbagheri B, Kia SH (2019) **Modelling climate change effects on Zagros forests in Iran using individual and ensemble forecasting approaches** *Theor Appl Climatol* **137**:1015–1025 <https://doi.org/10.1007/s00704-018-2625-z> | [Google Scholar](#)

Van Doren BM (2022) **How migratory birds might have tracked past climate change** *Proceedings of the National Academy of Sciences* **119**:e2121738119 <https://doi.org/10.1073/pnas.2121738119> | [Google Scholar](#)

Vandegrift KJ, Sokolow SH, Daszak P, Kilpatrick AM (2010) **Ecology of avian influenza viruses in a changing world** *Annals of the New York Academy of Sciences* **1195**:113–128 <https://doi.org/10.1111/j.1749-6632.2010.05451.x> | [Google Scholar](#)

Wille M, Atkinson R, Barr IG, Burgoyne C, Bond AL, Boyle D, Christie M, Dewar M, Douglas T, Fitzwater T, Hassell C, Jessop R, Klaassen H, Lavers JL, Leung KK -S., Ringma J, Sutherland DR, Klaassen M. (2024) **Long- Distance Avian Migrants Fail to Bring 2.3.4.4b HPAI H5N1 Into Australia for a Second Year in a Row** *Influenza Resp Viruses* **18**:e13281 <https://doi.org/10.1111/irv.13281> | [Google Scholar](#)

Wille M, Klaassen M (2023) **No evidence for HPAI H5N1 2.3.4.4b incursion into Australia in 2022** *Influenza Resp Viruses* **17**:e13118 <https://doi.org/10.1111/irv.13118> | Google Scholar

WOAH (2024) **Practical guide for authorised field responders to HPAI outbreaks in marine mammals** <https://www.woah.org/app/uploads/2024/02/woah-practicalguide-forauthorisedfieldresponders-hpaimarinemammals-feb24.pdf>

(2023) **WOAH. . Ongoing avian influenza outbreaks in animals pose risk to humans2023. .** <https://www.who.int/news/item/12-07-2023-ongoing-avian-influenza-outbreaks-in-animals-pose-risk-to-humans> <https://www.who.int/news/item/12-07-2023-ongoing-avian-influenza-outbreaks-in-animals-pose-risk-to-humans>

Worobey M, Gangavarapu K, Pekar J E., Joy J B., Moncla L, Kraemer M U. G., Dudas G, Goldhill D, Ruis C, Malpica Serrano L, Ji X, Andersen K G., Wertheim J O., Lemey P, Suchard M A, Rasmussen A L., Chand M, Groves N, Pybus O G., Peacock T P., Rambaut A, Nelson M I. (2024a) **Preliminary report on genomic epidemiology of the 2024 H5N1 influenza A virus outbreak in U.S. cattle (Part 1 of 2) - Influenza virus / H5N1-global** *Virological* <https://virological.org/t/preliminary-report-on-genomic-epidemiology-of-the-2024-h5n1-influenza-a-virus-outbreak-in-u-s-cattle-part-1-of-2/970>

Worobey M, Gangavarapu K, Pekar J E., Joy J B., Moncla L, Kraemer M U. G., Dudas G, Goldhill D, Ruis C, Malpica Serrano L, Ji X, Andersen K G., Wertheim J O., Lemey P, Suchard M A, Rasmussen A L., Chand M, Groves N, Pybus O G., Peacock T P., Rambaut A, Nelson M I. (2024b) **Preliminary report on genomic epidemiology of the 2024 H5N1 influenza A virus outbreak in U.S. cattle (Part 2 of 2) - Influenza virus / H5N1-global** *Virological* <https://virological.org/t/preliminary-report-on-genomic-epidemiology-of-the-2024-h5n1-influenza-a-virus-outbreak-in-u-s-cattle-part-1-of-2/970>

Yin S, Xu Y, Xu M, de Jong MCM, Huisman MRS, Contina A, Prins HHT, Huang ZYX, de Boer WF (2022) **Habitat loss exacerbates pathogen spread: An Agent-based model of avian influenza infection in migratory waterfowl** *PLOS Computational Biology* **18**:e1009577 <https://doi.org/10.1371/journal.pcbi.1009577> | Google Scholar

Author information

Marie-Cécile Dupas*

Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles (ULB), Brussels, Belgium,
Data Science Institute, University of Hasselt, Hasselt, Belgium
ORCID iD: [0000-0002-8769-2243](https://orcid.org/0000-0002-8769-2243)

For correspondence: dupas.mc@gmail.com

*denotes equal contribution

Maria F Vincenti-Gonzalez*

Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles (ULB), Brussels, Belgium

*denotes equal contribution

Madhur Dhingra

Food and Agriculture Organization of the United Nations, Rome, Italy

Claire Guinat

Interactions Hôtes-Agents Pathogènes (IHAP), Université de Toulouse, INRAE, ENVT, Toulouse, France

Timothée Vergne

Interactions Hôtes-Agents Pathogènes (IHAP), Université de Toulouse, INRAE, ENVT, Toulouse, France

William Wint

Environmental Research Group Oxford Ltd, c/o Department of Biology, Oxford, United Kingdom

Guy Hendrickx

Avia-GIS research department, Avia-GIS, Zoersel, Belgium

Cedric Marsboom

Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles (ULB), Brussels, Belgium, Avia-GIS research department, Avia-GIS, Zoersel, Belgium

Marius Gilbert

Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles (ULB), Brussels, Belgium

Simon Dellicour

Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles (ULB), Brussels, Belgium, Department of Microbiology, Immunology and Transplantation, Rega Institute, KU Leuven, Leuven, Belgium, Interuniversity Institute of Bioinformatics in Brussels, Université Libre de Bruxelles, Vrije Universiteit Brussel, Brussels, Belgium
ORCID iD: [0000-0001-9558-1052](https://orcid.org/0000-0001-9558-1052)

Editors

Reviewing Editor

Joshua Schiffer

Fred Hutch Cancer Center, Seattle, United States of America

Senior Editor

Joshua Schiffer

Fred Hutch Cancer Center, Seattle, United States of America

Reviewer #1 (Public review):

The authors aim to predict ecological suitability for transmission of highly pathogenic avian influenza (HPAI) using ecological niche models. This class of models identify correlations between the locations of species or disease detections and the environment. These correlations are then used to predict habitat suitability (in this work, ecological suitability for disease transmission) in locations where surveillance of the species or disease has not been conducted. The authors fit separate models for HPAI detections in wild birds and farmed birds, for two strains of HPAI (H5N1 and H5Nx) and for two time periods, pre- and post-2020. The authors also validate models fitted to disease occurrence data from pre-2020 using post-2020 occurrence data.

<https://doi.org/10.7554/eLife.104748.3.sa2>

Reviewer #2 (Public review):**Summary:**

The geographic range of highly pathogenic avian influenza cases changed substantially around the period 2020, and there is much interest in understanding why. Since 2020 the pathogen irrupted in the Americas and the distribution in Asia changed dramatically. This study aimed to determine which spatial factors (environmental, agronomic and socio-economic) explain the change in numbers and locations of cases reported since 2020 (2020–2023). That's a causal question which they address by applying correlative environmental niche modelling (ENM) approach to the avian influenza case data before (2015–2020) and after 2020 (2020–2023) and separately for confirmed cases in wild and domestic birds. To address their questions they compare the outputs of the respective models, and those of the first global model of the HPAI niche published by Dhingra et al 2016.

ENM is a correlative approach useful for extrapolating understandings based on sparse geographically referenced observational data over un- or under-sampled areas with similar environmental characteristics in the form of a continuous map. In this case, because the selected covariates about land cover, use, population and environment are broadly available over the entire world, modelled associations between the response and those covariates can be projected (predicted) back to space in the form of a continuous map of the HPAI niche for the entire world.

Strengths:

The authors are clear about expected bias in the detection of cases, such geographic variation in surveillance effort (testing of symptomatic or dead wildlife, testing domestic flocks) and in general more detections near areas of higher human population density (because if a tree falls in a forest and there is no-one there, etc), and take steps to ameliorate those. The authors use boosted regression trees to implement the ENM, which typically feature among the best performing models for this application (also known as habitat suitability models). They ran replicate sets of the analysis for each of their model targets (wild/domestic x pathogen variant), which can help produce stable predictions. Their code and data is provided, though I did not verify that the work was reproducible.

The paper can be read as a partial update to the first global model of H5Nx transmission by Dhingra and others published in 2016 and explicitly follows many methodological elements. Because they use the same covariate sets as used by Dhingra et al 2016 (including the comparisons of the performance of the sets in spatial cross-validation) and for both time periods of interest in the current work, comparison of model outputs is possible. The authors further facilitate those comparisons with clear graphics and supplementary analyses and presentation. The models can also be explored interactively at a weblink provided in text, though it would be good to see the model training data there too.

The authors' comparison of ENM model outputs generated from the distinct HPAI case datasets is interesting and worthwhile, though for me, only as a response to differently framed research questions.

Weaknesses:

This well-presented and technically well-executed paper has one major weakness to my mind. I don't believe that ENM models were an appropriate tool to address their stated goal, which was to identify the factors that "explain" changing HPAI epidemiology.

Comments on the revised version from the editors:

We are extremely grateful to the authors for presenting a thoughtful and respectful point by point rebuttal to the prior reviewers' comments. After reading these comments carefully, we conclude that there is a straightforward strongly held disagreement between the authors and the reviewers as to the validity of the methods (Ecological Niche Modeling) for this particular dataset. Please note that the two reviewers have substantial expertise in the area of Ecologic Niche Modeling. We elected not to reach out to the reviewers for a third set of comments as we do not think their overall opinions will change, and wish to be respectful of their time.

To allow readers a balanced assessment of the paper, we intend to publish your rebuttal comments in full. It is our hope that interested readers can weigh both sides of this respectful and interesting debate in order to reach their own conclusions about the strength of evidence presented in your manuscript.

<https://doi.org/10.7554/eLife.104748.3.sa1>

Author response:

The following is the authors' response to the previous reviews

Public Reviews:

We thank the Reviewers for their thorough attention to our paper and the interesting discussion about the findings. Before responding to more specific comments, here some general points we would like to clarify:

(1) Ecological niche models are indeed correlative models, and we used them to highlight environmental factors associated with HPAI outbreaks within two host groups. We will further revise the terminology that could still unintentionally suggest causal inference. The few remaining ambiguities were mainly in the Discussion section, where our intent was to interpret the results in light of the broader scientific literature. Particularly, we will change the following expressions:

- “Which factors can explain...” to “Which factors are associated with...” (line 75);

- “the environmental and anthropogenic factors influencing” to “the environmental and anthropogenic factors that are correlated with” (line 273);

- “underscoring the influence” to “underscoring the strong association” (line 282).

(2) We respectfully disagree with the suggestion that an ecological niche modelling (ENM) approach is not appropriate for this work and the research question addressed therein. Ecological niche models are specifically designed to estimate the spatial distribution of the environmental suitability of species and pathogens, making them well suited to our research questions. In our study, we have also explicitly detailed the known limitations of ecological niche models in the Discussion section, in line with prior literature, to ensure their appropriate interpretation in the context of HPAI.

(3) The environmental layers used in our models were restricted to those available at a global scale, as listed in Supplementary Information Resources S1 (https://github.com/sdellicour/h5nx_risk_mapping/blob/master/Scripts/%26_data/SI_Resource/S1.xlsx). Naturally, not all potentially relevant environmental factors could be included, but the selected layers are explicitly documented and only these were assessed for their importance. Despite this limitation, the performance metrics indicate that the models performed well, suggesting that the chosen covariates capture meaningful associations with HPAI occurrence at a global scale.

Reviewer #1 (Public review):

The authors aim to predict ecological suitability for transmission of highly pathogenic avian influenza (HPAI) using ecological niche models. This class of models identify correlations between the locations of species or disease detections and the environment. These correlations are then used to predict habitat suitability (in this work, ecological suitability for disease transmission) in locations where surveillance of the species or disease has not been conducted. The authors fit separate models for HPAI detections in wild birds and farmed birds, for two strains of HPAI (H5N1 and H5Nx) and for two time periods, pre- and post-2020. The authors also validate models fitted to disease occurrence data from pre-2020 using post-2020 occurrence data. I thank the authors for taking the time to respond to my initial review and I provide some follow-up below.

Detailed comments:

In my review, I asked the authors to clarify the meaning of "spillover" within the HPAI transmission cycle. This term is still not entirely clear: at lines 409-410, the authors use the term with reference to transmission between wild birds and farmed birds, as distinct to transmission between farmed birds. It is implied but not explicitly stated that "spillover" is relevant to the transmission cycle in farmed birds only. The sentence, "we developed separate ecological niche models for wild and domestic bird HPAI occurrences ..." could have been supported by a clear sentence describing the transmission cycle, to prime the reader for why two separate models were necessary.

We respectfully disagree that the term “spillover” is unclear in the manuscript. In both the Methods and Discussion sections (lines 387-391 and 409-414), we explicitly define “spillover” as the introduction of HPAI viruses from wild birds into domestic poultry, and we distinguish this from secondary farm-to-farm transmission. Our use of separate ecological niche models for wild and domestic outbreaks reflects not only the distinction between primary spillover and secondary transmission, but also the fundamentally different ecological processes, surveillance systems, and management implications that shape outbreaks in these two groups. We will clarify this choice in the revised manuscript when introducing the separate models. Furthermore, on line 83, we will add “as these two groups are influenced by different ecological processes, surveillance biases, and management contexts”.

I also queried the importance of (dead-end) mammalian infections to a model of the HPAI transmission risk, to which the authors responded: "While spillover events of HPAI into mammals have been documented, these detections are generally considered dead-end infections and do not currently represent sustained transmission chains. As such, they fall outside the scope of our study, which focuses on avian hosts and models ecological suitability for outbreaks in wild and domestic birds." I would argue that any infections, whether they are in dead-end or competent hosts, represent the presence of environmental conditions to support transmission so are certainly relevant to a niche model and therefore within scope. It is certainly understandable if the authors have not been able to access data of mammalian infections, but it is an oversight to dismiss these infections as irrelevant.

We understand the Reviewer’s point, but our study was designed to model HPAI occurrence in avian hosts only. We therefore restricted our analysis to wild birds and domestic poultry, which represent the primary hosts for HPAI circulation and the focus of surveillance and control measures. While mammalian detections have been reported, they are outside the scope of this work.

Correlative ecological niche models, including BRTs, learn relationships between occurrence data and covariate data to make predictions, irrespective of correlations between covariates. I am not convinced that the authors can make any "interpretation" (line 298) that the covariates that are most informative to their models have any "influence" (line 282) on their response variable. Indeed, the observation that "land-use and climatic predictors do not play an important role in the niche ecological models" (line 286), while "intensive chicken population density emerges as a significant predictor" (line 282) begs the question: from an operational perspective, is the best (e.g., most interpretable and quickest to generate) model of HPAI risk a map of poultry farming intensity?

We agree that poultry density may partly reflect reporting bias, but we also assumed it a meaningful predictor of HPAI risk. Its importance in our models is therefore expected. Importantly, our BRT framework does more than reproduce poultry distribution: it captures non-linear relationships and interactions with other covariates, allowing a more nuanced characterisation of risk than a simple poultry density map. Note also that we distinguished in our models intensive and extensive chicken poultry density and duck density. Therefore, it is not a “map of poultry farming intensity”.

At line 282, we used the word “influence” while fully recognising that correlative models cannot establish causality. Indeed, in our analyses, “relative influence” refers to the importance metric produced by the BRT algorithm (Ridgeway, 2020), which measures correlative associations between environmental factors and outbreak occurrences. These scores are interpreted in light of the broader scientific literature, therefore our interpretations build on both our results and existing evidence, rather than on our models alone. However, in the next version of the paper, we will revise the sentence as: “underscoring the strong association of poultry farming practices with HPAI spread (Dhingra et al., 2016)”.

I have more significant concerns about the authors' treatment of sampling bias: "We agree with the Reviewer's comment that poultry density could have potentially been considered to guide the sampling effort of the pseudo-absences to consider when training domestic bird models. We however prefer to keep using a human population density layer as a proxy for surveillance bias to define the relative probability to sample pseudo-absence points in the different pixels of the background area considered when training our ecological niche models. Indeed, given that poultry density is precisely one of the predictors that we aim to test, considering this environmental layer for defining the relative probability to sample pseudo-absences would introduce a certain level of circularity in our analytical procedure, e.g. by artificially increasing to influence of that particular variable in our models." The authors have elected to ignore a fundamental feature of distribution modelling with occurrence-only data: if we include a source of sampling bias as a covariate and do not include it when we sample background data, then that covariate would appear to be correlated with presence. They acknowledge this later in their response to my review: "...assuming a sampling bias correlated with poultry density would result in reducing its effect as a risk factor." In other words, the apparent predictive capacity of poultry density is a function of how the authors have constructed the sampling bias for their models. A reader of the manuscript can reasonably ask the question: to what degree are is the model a model of HPAI transmission risk, and to what degree is the model a model of the observation process? The sentence at lines 474-477 is a helpful addition, however the preceding sentence, "Another approach to sampling pseudo-absences would have been to distribute them according to the density of domestic poultry," (line 474) is included without acknowledgement of the flow-on consequence to one of the key findings of the manuscript, that "...intensive chicken

population density emerges as a significant predictor..." (line 282). The additional context on the EMPRES-i dataset at line 475-476 ("the locations of outbreaks ... are often georeferenced using place name nomenclatures") is in conflict with the description of the dataset at line 407 ("precise location coordinates"). Ultimately, the choices that the authors have made are entirely defensible through a clear, concise description of model features and assumptions, and precise language to guide the reader through interpretation of results. I am not satisfied that this is provided in the revised manuscript.

We thank the Reviewer for this important point. To address it, we compared model predictive performance and covariate relative influences obtained when pseudo-absences were weighted by poultry density versus human population density (Author response table 1). The results show that differences between the two approaches are marginal, both in predictive performance (Δ AUC ranging from -0.013 to +0.002) and in the ranking of key predictors (see below Author response images 1 and 2). For instance, intensive chicken density consistently emerged as an important predictor regardless of the bias layer used.

Note: the comparison was conducted using a simplified BRT configuration for computational efficiency (fewer trees, fixed 5-fold random cross-validation, and standardised parameters). Therefore, absolute values of AUC and variable importance may differ slightly from those in the manuscript, but the relative ranking of predictors and the overall conclusions remain consistent.

Given these small differences, we retained the approach using human population density. We agree that poultry density partly reflects surveillance bias as well as true epidemiological risk, and we will clarify this in the revised manuscript by noting that the predictive role of poultry density reflects both biological processes and surveillance systems. Furthermore, on line 289, we will add "We note, however, that intensive poultry density may reflect both surveillance intensity and epidemiological risk, and its predictive role in our models should be interpreted in light of both processes".


Author response table 1.

Comparison of model predictive performances (AUC) between pseudo-absence sampling were weighted by poultry density and by human population density across host groups, virus types, and time periods. Differences in AUC values are shown as the value for poultry-weighted minus human-weighted pseudo-absences.

Dataset	AUC (poultry-weighted pseudo-absences)	AUC (human-weighted pseudo-absences)	AUC differences between the two approaches
H5N1, domestic birds, <2020	0.877	0.864	-0.013
H5N1, wild birds, >2020	0.837	0.829	-0.008
H5N1, domestic birds, >2020	0.831	0.828	-0.003
H5Nx, wild birds, <2020	0.856	0.858	0.002
H5Nx, domestic birds, <2020	0.853	0.855	0.002
H5Nx, wild birds, >2020	0.847	0.842	-0.005
H5Nx, domestic birds, >2020	0.835	0.833	-0.002


Author response image 1.

Comparison of variable relative influence (%) between models trained with pseudo-absences weighted by poultry density (red) and human population density (blue) for domestic bird outbreaks. Results are shown for four datasets: H5N1 (<2020), H5N1 (>2020), H5Nx (<2020), and H5Nx (>2020).



Author response image 2.

Comparison of variable relative influence (%) between models trained with pseudo-absences weighted by poultry density (red) and human population density (blue) for wild bird outbreaks. Results are shown for three datasets: H5N1 (>2020), H5Nx (<2020), and H5Nx (>2020).



The authors have slightly misunderstood my comment on "extrapolation": I referred to "environmental extrapolation" in my review without being particularly explicit about my meaning. By "environmental extrapolation", I meant to ask whether the models were predicting to environments that are outside the extent of environments included in the occurrence data used in the manuscript. The authors appear to have understood this to be a comment on geographic extrapolation, or predicting to areas outside the geographic extent included in occurrence data, e.g.: "For H5Nx post-2020, areas of high

predicted ecological suitability, such as Brazil, Bolivia, the Caribbean islands, and Jilin province in China, likely result from extrapolations, as these regions reported few or no outbreaks in the training data" (lines 195-197). Is the model extrapolating in environmental space in these regions? This is unclear. I do not suggest that the authors should carry out further analysis, but the multivariate environmental similarity surface (MESS; see Elith et al., 2010) is a useful tool to visualise environmental extrapolation and aid model interpretation.

On the subject of "extrapolation", I am also concerned by the additions at lines 362-370: "...our models extrapolate environmental suitability for H5Nx in wild birds in areas where few or no outbreaks have been reported. This discrepancy may be explained by limited surveillance or underreporting in those regions." The "discrepancy" cited here is a feature of the input dataset, a function of the observation distribution that should be captured in pseudo-absence data. The authors state that Kazakhstan and Central Asia are areas of interest, and that the environments in this region are outside the extent of environments captured in the occurrence dataset, although it is unclear whether "extrapolation" is informed by a quantitative tool like a MESS or judged by some other qualitative test. The authors then cite Australia as an example of a region with some predicted suitability but no HPAI outbreaks to date, however this discussion point is not linked to the idea that the presence of environmental conditions to support transmission need not imply the occurrence of transmission (as in the addition, "...spatial isolation may imply a lower risk of actual occurrences..." at line 214). Ultimately, the authors have not added any clear comment on model uncertainty (e.g., variation between replicated BRTs) as I suggested might be helpful to support their description of model predictions.

Many thanks for the clarification. Indeed, we interpreted your previous comments in terms of geographic extrapolations. We thank the Reviewer for these observations. We will adjust the wording to further clarify that predictions of ecological suitability in areas with few or no reported outbreaks (e.g., Central Asia, Australia) are not model errors but expected extrapolations, since ecological suitability does not imply confirmed transmission (for instance, on Line 362: "our models extrapolate environmental suitability" will be changed to "Interestingly, our models extrapolate geographical"). These predictions indicate potential environments favorable to circulation if the virus were introduced.

In our study, model uncertainty is formally assessed when comparing the predictive performances of our models (Fig. S3, Table S1), the relative influence (Table S3) and response curves (Fig. 2) associated with each environmental factor (Table S2). All the results confirming a good converge between these replicates. Finally, we indeed did not use a quantitative tool such as a MESS to assess extrapolation but did rely on qualitative interpretation of model outputs.

All of my criticisms are, of course, applied with the understanding that niche modelling is imperfect for a disease like HPAI, and that data may be biased/incomplete, etc.: these caveats are common across the niche modelling literature. However, if language around the transmission cycle, the niche, and the interpretation of any of the models is imprecise, which I find it to be in the revised manuscript, it undermines all of the science that is presented in this work.

We respectfully disagree with this comment. The scope of our study and the methods employed are clearly defined in the manuscript, and the limitations of ecological niche modelling in this context are explicitly acknowledged in the Discussion section. While we appreciate the Reviewer's concern, the comment does not provide specific examples of unclear or imprecise language regarding the transmission cycle, niche, or interpretation of the models. Without such examples, it is difficult to identify further revisions that would improve clarity.

Reviewer #2 (Public review):

The geographic range of highly pathogenic avian influenza cases changed substantially around the period 2020, and there is much interest in understanding why. Since 2020 the pathogen irrupted in the Americas and the distribution in Asia changed dramatically. This study aimed to determine which spatial factors (environmental, agronomic and socio-economic) explain the change in numbers and locations of cases reported since 2020 (2020--2023). That's a causal question which they address by applying correlative environmental niche modelling (ENM) approach to the avian influenza case data before (2015--2020) and after 2020 (2020--2023) and separately for confirmed cases in wild and domestic birds. To address their questions they compare the outputs of the respective models, and those of the first global model of the HPAI niche published by Dhingra et al 2016.

We do not agree with this comment. In the manuscript, it is well established that we are quantitatively assessing factors that are associated with occurrences data before and after 2020. We do not claim to determine the causality. One sentence of the Introduction section (lines 75-76) could be confusing, so we intend to modify it in the final revision of our manuscript.

ENM is a correlative approach useful for extrapolating understandings based on sparse geographically referenced observational data over un- or under-sampled areas with similar environmental characteristics in the form of a continuous map. In this case, because the selected covariates about land cover, use, population and environment are broadly available over the entire world, modelled associations between the response and those covariates can be projected (predicted) back to space in the form of a continuous map of the HPAI niche for the entire world.

We fully agree with this assessment of ENM approaches.

Strengths:

The authors are clear about expected bias in the detection of cases, such geographic variation in surveillance effort (testing of symptomatic or dead wildlife, testing domestic flocks) and in general more detections near areas of higher human population density (because if a tree falls in a forest and there is no-one there, etc), and take steps to ameliorate those. The authors use boosted regression trees to implement the ENM, which typically feature among the best performing models for this application (also known as habitat suitability models). They ran replicate sets of the analysis for each of their model targets (wild/domestic x pathogen variant), which can help produce stable predictions. Their code and data is provided, though I did not verify that the work was reproducible.

The paper can be read as a partial update to the first global model of H5Nx transmission by Dhingra and others published in 2016 and explicitly follows many methodological elements. Because they use the same covariate sets as used by Dhingra et al 2016 (including the comparisons of the performance of the sets in spatial cross-validation) and for both time periods of interest in the current work, comparison of model outputs is possible. The authors further facilitate those comparisons with clear graphics and supplementary analyses and presentation. The models can also be explored interactively at a weblink provided in text, though it would be good to see the model training data there too.

The authors' comparison of ENM model outputs generated from the distinct HPAI case datasets is interesting and worthwhile, though for me, only as a response to differently

framed research questions.

Weaknesses:

This well-presented and technically well-executed paper has one major weakness to my mind. I don't believe that ENM models were an appropriate tool to address their stated goal, which was to identify the factors that "explain" changing HPAI epidemiology.

Here is how I understand and unpack that weakness:

(1) Because of their fundamentally correlative nature, ENMs are not a strong candidate for exploring or inferring causal relationships.

(2) Generating ENMs for a species whose distribution is undergoing broad scale range change is complicated and requires particular caution and nuance in interpretation (e.g., Elith et al, 2010, an important general assumption of environmental niche models is that the target species is at some kind of distributional equilibrium (at time scales relevant to the model application). In practice that means the species has had an opportunity to reach all suitable habitats and therefore its absence from some can be interpreted as either unfavourable environment or interactions with other species). Here data sets for the response (N5H1 or N5Hx case data in domestic or wild birds) were divided into two periods; 2015--2020, and 2020--2023 based on the rationale that the geographic locations and host-species profile of cases detected in the latter period was suggestive of changed epidemiology. In comparing outputs from multiple ENMs for the same target from distinct time periods the authors are expertly working in, or even dancing around, what is a known grey area, and they need to make the necessary assumptions and caveats obvious to readers.

We thank the Reviewer for this observation. First, we constrained pseudo-absence sampling to countries and regions where outbreaks had been reported, reducing the risk of interpreting non-affected areas as environmentally unsuitable. Second, we deliberately split the outbreak data into two periods (2015-2020 and 2020-2023) because we do not assume a single stable equilibrium across the full study timeframe. This division reflects known epidemiological changes around 2020 and allows each period to be modeled independently. Within each period, ENM outputs are interpreted as associations between outbreaks and covariates, not as equilibrium distributions. Finally, by testing prediction across periods, we assessed both niche stability and potential niche shifts. These clarifications will be added to the manuscript to make our assumptions and limitations explicit.

Line 66, we will add: "Ecological niche model outputs for range-shifting pathogens must therefore be interpreted with caution (Elith et al., 2010). Despite this limitation, correlative ecological niche models remain useful for identifying broad-scale associations and potential shifts in distribution. To account for this, we analysed two distinct time periods (2015-2020 and 2020-2023)."

Line 123, we will revise "These findings underscore the ability of pre-2020 models in forecasting the recent geographic distribution of ecological suitability for H5Nx and H5N1 occurrences" to "These results suggest that pre-2020 models captured broad patterns of suitability for H5Nx and H5N1 outbreaks, while post-2020 models provided a closer fit to the more recent epidemiological situation".

(3) To generate global prediction maps via ENM, only variables that exist at appropriate resolution over the desired area can be supplied as covariates. What processes could influence changing epidemiology of a pathogen and are their covariates that represent them? Introduction to a new geographic area (continent) with naive population, immunity in previously exposed populations, control measures to limit spread such as

vaccination or destruction of vulnerable populations or flocks? Might those control measures be more or less likely depending on the country as a function of its resources and governance? There aren't globally available datasets that speak to those factors, so the question is not why were they omitted but rather was the authors decision to choose ENMs given their question justified? How valuable are insights based on patterns of correlation change when considering different temporal sets of HPAI cases in relation to a common and somewhat anachronistic set of covariates?

We agree that the ecological niche models trained in our study are limited to environmental and host factors, as described in the Methods section with the selection of predictors. While such models cannot capture causality or represent processes such as immunity, control measures, or governance, they remain a useful tool for identifying broad associations between outbreak occurrence and environmental context. Our study cannot infer the full mechanisms driving changes in HPAI epidemiology, but it does provide a globally consistent framework to examine how associations with available covariates vary across time periods.

*(4) In general the study is somewhat incoherent with respect to time. Though the case data come from different time periods, each response dataset was modelled separately using exactly the same covariate dataset that predated both sets. That decision should be understood as a strong assumption on the part of the authors that conditions the interpretation: the world (as represented by the covariate set) is immutable, so the model has to return different correlative associations between the case data and the covariates to explain the new data. While the world represented by the selected covariates *may* be relatively stable (could be statistically confirmed), what about the world not represented by the covariates (see point 3)?*

We used the same covariate layers for both periods, which indeed assumes that these environmental and host factors are relatively stable at the global scale over the short timeframe considered. We believe this assumption is reasonable, as poultry density, land cover, and climate baselines do not change drastically between 2015 and 2023 at the resolution of our analysis. We agree, however, that unmeasured processes such as control measures, immunity, or governance may have changed during this time and are not captured by our covariates.

Recommendations for the Authors:

Reviewer #1 (Recommendations for the authors):

- Line 400-401: "over the 2003-2016 periods" has an extra "s"; "two host species" (with reference to wild and domestic birds) would be more precise as "two host groups".

- Remove comma line 404

Many thanks for these comments, we have modified the text accordingly.

Reviewer #2 (Recommendations for the authors):

Most of my work this round is encapsulated in the public part of the review.

The authors responded positively to the review efforts from the previous round, but I was underwhelmed with the changes to the text that resulted. Particularly in regard to limiting assumptions - the way that they augmented the text to refer to limitations raised in review downplayed the importance of the assumptions they've made. So they acknowledge the significance of the limitation in their rejoinder, but in the amended text merely note the limitation without giving any sense of what it means for their interpretation of the findings of this study.

The abstract and findings are essentially unchanged from the previous draft.

I still feel the near causal statements of interpretation about the covariates are concerning. These models really are not a good candidate for supporting the inference that they are making and there seem to be very strong arguments in favour of adding covariates that are not globally available.

We never claimed causal interpretation, and we have consistently framed our analyses in terms of associations rather than mechanisms. We acknowledge that one phrasing in the research questions (“Which factors can explain...”) could be misinterpreted, and we are correcting this in the revised version to read “Which factors are associated with...”. Our approach follows standard ecological niche modelling practice, which identifies statistical associations between occurrence data and covariates. As noted in the Discussion section, these associations should not be interpreted as direct causal mechanisms. Finally, all interpretive points in the manuscript are supported by published literature, and we consider this framing both appropriate and consistent with best practice in ecological niche modelling (ENM) studies.

We assessed predictor contributions using the “relative influence” metric, the terminology reported by the R package “gbm” (Ridgeway, 2020). This metric quantifies the contribution of each variable to model fit across all trees, rescaled to sum to 100%, and should be interpreted as an association rather than a causal effect.

L65-66 The general difficulty of interpreting ENM output with range-shifting species should be cited here to alert readers that they should not blithely attempt what follows at home.

I believe that their analysis is interesting and technically very well executed, so it has been a disappointment and hard work to write this assessment. My rough-cut last paragraph of a reframed intro would go something like - there are many reasons in the literature not to do what we are about to do, but here's why we think it can be instructive and informative, within certain guardrails.

To acknowledge this comment and the previous one, we revised lines 65-66 to: “However, recent outbreaks raise questions about whether earlier ecological niche models still accurately predict the current distribution of areas ecologically suitable for the local circulation of HPAI H5 viruses. Ecological niche model outputs for range-shifting pathogens must therefore be interpreted with caution (Elith et al., 2010). Despite this limitation, correlative ecological niche models remain useful for identifying broad-scale associations and potential shifts in distribution.”

We respectfully disagree with the Reviewer’s statement that “there are many reasons in the literature not to do what we are about to do”. All modeling approaches, including mechanistic ones, have limitations, and the literature is clear on both the strengths and constraints of ecological niche models. Our manuscript openly acknowledges these limits and frames our findings accordingly. We therefore believe that our use of an ENM approach is justified and contributes valuable insights within these well-defined boundaries.

Reference: Ridgeway, G. (2007). Generalized Boosted Models: A guide to the gbm package. Update, 1(1), 2007.

<https://doi.org/10.7554/eLife.104748.3.sa0>