

Endophilin A1 facilitates organization of the GABAergic postsynaptic machinery to maintain excitation-inhibition balance

Reviewed Preprint

v2 • July 2, 2025

Revised by authors

Reviewed Preprint

v1 • November 12, 2024

Xue Chen, Deng Pan, Jia-Jia Liu , Yanrui Yang 

State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China • College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This study presents a **valuable** finding on the molecular mechanisms that govern GABAergic inhibitory synapse function. The authors propose that Endophilin A1 serves as a novel regulator of GABAergic synapses by acting as a component of the inhibitory postsynaptic density. The authors have added substantial new analyses that take a wide range of approaches to provide **solid** support for their conclusions, although one of the reviewers concludes that the premise that gephyrin and endophilin A1 interact requires more robust analysis. The findings are likely to interest a broad audience of scientists focusing on inhibitory synaptic transmission, the excitation-inhibition balance, and its disruption in disorders such as epilepsy.

<https://doi.org/10.7554/eLife.102792.2.sa4>

Abstract

The assembly and operation of neural circuits in the brain rely on the coordination and balance of excitatory and inhibitory activities. Inhibitory synapses are key regulators of the functional balance of neural circuits. However, due to the diversity of inhibitory presynaptic neurons, the complex composition of postsynaptic receptor subunits and the lack of typical postsynaptic dense structure, there are relatively few studies on the regulatory mechanisms for inhibitory synaptic structure and function, and insufficient understanding of the cellular and molecular abnormalities of inhibitory synapses in neurological and neuropsychiatric disorders. Here, we report a crucial role for endophilin A1 in inhibitory synapses. We show that endophilin A1 directly interacts with the inhibitory postsynaptic scaffold protein gephyrin in excitatory neurons, and promotes organization of the inhibitory postsynaptic density and synaptic recruitment/stabilization of the γ -aminobutyric acid type A receptors via its plasma membrane association and actin polymerization promoting activities. Loss of endophilin A1 by gene knockout in mouse hippocampal CA1 pyramidal cells weakens inhibitory synaptic transmission and causes imbalance in the excitatory/inhibitory function

of neural circuits, leading to increased susceptibility to epilepsy. Our findings identify endophilin A1 as an iPSD component and provide new insights into the organization and stabilization of inhibitory postsynapses to maintain E/I balance as well as the pathogenesis of epilepsy.

Introduction

The excitatory and inhibitory synaptic activities of neural circuits are tightly orchestrated in the central nervous system (Sakimoto et al., 2021 [DOI](#)). The imbalance between excitatory and inhibitory neuronal activity is implicated in many neurological and neuropsychiatric diseases including schizophrenia, autism spectrum disorder and epilepsy (Canitano and Pallagrosi, 2017 [DOI](#); Canitano and Palumbi, 2021 [DOI](#); Foss-Feig et al., 2017 [DOI](#); Fritschy, 2008 [DOI](#)). In the brain, in contrast to asymmetric excitatory synapses formed between axonal boutons and dendritic postsynaptic structures with thick disc-shaped PSD, inhibitory synapses are symmetric and are constituted of the presynaptic terminal from diverse interneurons and the postsynaptic plasma membrane with thin sheet-like PSD where receptor complexes are anchored (Bai et al., 2021 [DOI](#); Favuzzi et al., 2019 [DOI](#); Liu et al., 2020 [DOI](#); Mody and Pearce, 2004 [DOI](#); Tao et al., 2018 [DOI](#)). Although the formation, maturation and plasticity of both excitatory and inhibitory synapses are vitally important for neural circuit functioning, much more research efforts have been focused on the regulatory mechanisms governing excitatory synaptic structure and function. The mechanisms by which inhibitory synapses are constructed in the brain and fine-tuned in response to neural activity remain largely unexplored.

The γ -aminobutyric acid type A receptors (GABA_ARs), major targets of antiseizure drugs, are heteropentameric chloride channels that mediate fast synaptic transmission at inhibitory synapses in the brain (Scott and Aricescu, 2019 [DOI](#)). Several neuropathologies are characterized by alterations in GABA_A receptor trafficking, synaptic E/I ratio and neuronal excitability (Sakimoto et al., 2021 [DOI](#)). At the inhibitory postsynaptic site, synaptic localization and functioning of GABA_ARs requires trans-synaptic cell adhesion molecules, submembranous scaffold proteins and intracellular signaling proteins. Gephyrin is a master organizer of iPSD, which self-assembles into planar lattices beneath the plasma membrane and anchors GABA_ARs or glycine receptors (GlyRs) to the postsynaptic sites (George et al., 2021 [DOI](#); Kasaragod and Schindelin, 2019 [DOI](#); Kirsch et al., 1991 [DOI](#); Tyagarajan and Fritschy, 2014 [DOI](#)). Collybistin, a membrane lipid-binding protein, functions in recruitment of gephyrin and GABA_ARs to the plasma membrane by interacting with gephyrin, the synaptic adhesion molecule neuroligin-2 (NL2) and the $\alpha 2$ subunit of GABA_ARs (Kalscheuer et al., 2009 [DOI](#); Kins et al., 2000 [DOI](#); Pouloupoulos et al., 2009 [DOI](#); Saiepour et al., 2010 [DOI](#)). The transmembrane protein Shisa7 binds GABA_ARs and regulates their trafficking, synaptic abundance and functional properties via binding to protein(s) other than gephyrin (Han et al., 2019 [DOI](#); Liu et al., 2020 [DOI](#)). Nevertheless, mechanisms governing the assembly, recruitment and maintenance of various inhibitory synaptic components remain largely enigmatic.

The majority of inhibitory synapses are formed on the dendritic shaft, or around the cell body of excitatory principal neurons where both actin and microtubule cytoskeletons are present underneath the plasma membrane. Through direct interaction with polymerized tubulin, gephyrin organizes the iPSD by anchoring the GlyRs and GABA_ARs to the microtubule cytoskeleton (Parato and Bartolini, 2021 [DOI](#)). Notably, although small F-actin patches are observed within the dendritic shaft where the majority of inhibitory synapses are assembled (van Bommel et al., 2019 [DOI](#)), role(s) of the actin cytoskeleton in the organization and maintenance of inhibitory synapses is still unclear.

Endophilin A1 (EndoA1) is a member of the endophilin A protein family characterized by an amino (N)-terminal membrane-binding N-BIN/amphiphysin/Rvs (BAR) domain and a carboxyl (C)-terminal Src homology 3 (SH3) domain. The gene encoding endophilin A1 (*EndoA1*, gene name

sh3gl2) is almost exclusively expressed in the brain (Ringstad et al., 1997), and has been implicated in epilepsy, schizophrenia, depression, Parkinson's disease, and Alzheimer's disease (Chang et al., 2017; Corponi et al., 2019; Germer et al., 2019; Ren et al., 2008; Yu et al., 2018a; Yu et al., 2018b). In excitatory synapses, endophilin A1 functions in synaptic vesicle recycling and autophagosome formation at the axon terminal (Bademosi et al., 2023; Milosevic et al., 2011; Murdoch et al., 2016; Ringstad et al., 1999; Schuske et al., 2003; Soukup et al., 2016; Verstreken et al., 2003; Watanabe et al., 2018), and regulates the morphogenesis, maturation, and acute structural plasticity of dendritic spines by promoting actin polymerization via its effector protein p140Cap (Yang et al., 2021; Yang et al., 2018; Yang et al., 2015).

Previous studies reported that the double knockout (KO) mice of *EndoA1* and *EndoA2* exhibit spontaneous epileptic seizures (Milosevic et al., 2011). However, *EndoA1* depletion only causes a slight decrease in excitatory synaptic transmission in hippocampal CA1 neurons (Yang et al., 2021; Yang et al., 2018). As E/I imbalance is caused by changes in either excitatory or inhibitory activity, or both, we were prompted to investigate the biological function of endophilin A1 in the inhibitory synapses. In this study, we show that endophilin A1 interacts with gephyrin and localizes to GABAergic postsynapses. Via its membrane-binding and actin polymerization-promoting activities, endophilin A1 facilitates organization of the inhibitory postsynaptic machinery. Loss of endophilin A1 causes impairment of inhibitory synaptic transmission, E/I imbalance and aggravation of epileptic susceptibility.

Results

Ablation of endophilin A1 causes E/I imbalance and increases epilepsy susceptibility

Previous studies found that *EndoA1* and *EndoA2* double knockout (KO) mice exhibit spontaneous epileptic seizures (Milosevic et al., 2011), suggesting that endophilin A1 and A2 are involved in maintaining the E/I balance. However, another study reported that the protein levels of endophilin A1 were higher in temporal neocortex neurons of temporal lobe epilepsy (TLE) patients (Yu et al., 2018b). Moreover, in a chemical kindling mouse model of epilepsy, in which mice are chronically induced with the GABA_AR antagonist pentylenetetrazole (PTZ) (Dhir, 2012), endophilin A1 was also upregulated in the hippocampus and adjacent temporal cortex, and silencing its expression in the hippocampal CA3 region reduced PTZ-induced seizure susceptibility and severity (Yu et al., 2018b). To investigate the role of endophilin A1 in epilepsy, we determined the impact of *EndoA1* gene ablation on chemical kindling of pan-neural *EndoA1* KO mice (Yang et al., 2018) with PTZ at a sub-convulsive dose of 25 mg/kg. Compared with the wild-type, PTZ-induced epileptic behaviors were significantly exacerbated in *EndoA1* KO mice (**Figure 1A-C** and Video S1).




Figure 1.

Ablation of endophilin A1 in CA1 neurons disrupts the E/I balance and increases epilepsy susceptibility.

- (A) Experimental protocol of PTZ kindling. Saline serves as negative control for PTZ.
- (B) Representative images of convulsive behavior in each group.
- (C) Seizures were monitored and scored after each injection. Nine 10-week-old mice were used in each condition. Seizure scores are shown as means $\pm$ S.E.M. Statistical test: repeated measures ANOVA followed by Tukey post-hoc test at each time point. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, *EndoA1*^{-/-}_PTZ vs *EndoA1*^{+/+}_PTZ at each time point.
- (D) The representative mCherry-positive hippocampal CA1 neuron used for electrophysiological recording.
- (E-F) Representative traces of evoked excitatory postsynaptic currents (eEPSC) and quantification of mean eEPSC amplitude (E), and paired-pulse recording (F) from 10 control (Ctrl, uninfected) and 10 *EndoA1* KO (mCherry and Cre) pyramidal neurons of 4 independent animals.
- (G-H) Representative traces of evoked inhibitory postsynaptic currents (eIPSCs) and quantification of mean eIPSC amplitude (G), and paired-pulse recording (H) from 9 control and 9 *EndoA1* KO pyramidal neurons of 4 independent animals.
- (I) Effect of *EndoA1* KO on the E/I ratio from 9 control and 9 *EndoA1* KO pyramidal neurons of 4 independent animals.
- (J) Representative recordings of miniature excitatory postsynaptic currents (mEPSCs) traces (+10 mV) from control or *EndoA1* KO CA1 pyramidal cells.
- (K) Cumulative probability and quantification of mEPSC amplitude from 7 control and 7 *EndoA1* KO pyramidal neurons of 3 independent animals.
- (L) Cumulative probability and quantification of mEPSC frequency from 7 control and 7 *EndoA1* KO pyramidal neurons of 3 independent animals.
- (M) Representative recordings of miniature inhibitory postsynaptic currents (mIPSCs) traces (-70 mV) from control or *EndoA1* KO CA1 pyramidal cells.
- (N) Cumulative probability and quantification of mIPSC amplitude from 7 control and 7 *EndoA1* KO pyramidal neurons of 3 independent animals.
- (O) Cumulative probability and quantification of mIPSC frequency from 7 control and 7 *EndoA1* KO pyramidal neurons of 3 independent animals. Statistical test: two-tailed Student's t-test in E-I & K, L, N, O. ns, non-significant. All data shown are scatterplots with means $\pm$ S.E.M.

We previously found that *EndoA1* KO caused a slight decrease in the amplitude of evoked excitatory postsynaptic currents (eEPSC) and a significant attenuation of long-term potentiation (Yang et al., 2018). As imbalance between excitatory and inhibitory signals in neuronal circuits is implicated in epilepsy, we then determined whether loss of endophilin A1 in excitatory neurons affects the E/I ratio by electrophysiological analysis. To this end, we eliminated the *EndoA1* gene in pyramidal neurons by stereotaxic injection of adeno-associated virus (AAV) expressing CaMKII α promoter-driven Cre recombinase into the hippocampal CA1 region of *EndoA1*^{fl/fl} mice (Figure 1D). Whole-cell patch clamp recording of CA1 pyramidal cells indicated that, while the eEPSC

and its paired-pulse ratio were not significantly changed, the evoked inhibitory postsynaptic current (eIPSC), not its PPR, was decreased in the *EndoA1* KO neurons, resulting in a significant increase in the E/I ratio (**Figure 1E-I**). Of note, neither the frequency nor the amplitude of miniature EPSCs (mEPSC) or miniature IPSCs (mIPSC) was altered in *EndoA1* KO neurons (**Figure 1J-O**). Moreover, neither fast mIPSCs (perisomatic events: < 2.8 ms) nor slow mIPSCs (distal dendritic events: > 2.8 ms) (Miles et al., 1996) were affected by *EndoA1* KO (Figure S1), indicating that presynaptic quantal release of soma- or dendrite-targeting inhibitory synapses is normal. Taken together, these data indicate that endophilin A1 in hippocampal pyramidal neurons functions in inhibitory synaptic transmission to maintain E/I balance.

Endophilin A1 is required for the formation and stabilization of inhibitory synapses

Diminished eIPSC in *EndoA1*^{-/-} pyramidal neurons prompted us to determine the role of endophilin A1 in inhibitory synapses by immunofluorescence staining of brain sections from *EndoA1*^{-/-} mice. Analysis of confocal microscopy images revealed that, compared with the wild-type, there was a significant decrease in fluorescent signals of both VGAT and gephyrin, the inhibitory pre- and post-synaptic markers, or VGLUT1 and PSD95, the excitatory pre- and postsynaptic markers, in all subregions of the hippocampus of the *EndoA1*^{-/-} mice on postnatal day 21 (P21) (**Figure 2A,C,D,F**, and S2A,C,D,F), suggesting defects in synapse development. While a slight reduction in excitatory presynaptic protein was observed, the decreases in inhibitory but not excitatory postsynaptic proteins were also detected in the *EndoA1*^{-/-} mice at P60 (**Figure 2B,C,E,F**, and S2B,C,E,F), indicating that the assembly and/or stabilization of inhibitory postsynapses was impaired throughout brain development. Notably, gephyrin signals in all layers of the different hippocampal subregions were reduced in the *EndoA1*^{-/-} mice (Figure S2G,H). In line with the data from immunofluorescent labeling (**Figure 2** and S2), decreased endogenous PSD95 or gephyrin clusters at synapses were also visualized by the recombinant probes, PSD95.FingR-eGFP or mRuby2-gephyrin.FingR, respectively (Bensussen et al., 2020; Gross et al., 2013), in *EndoA1* KO neurons in the CA1 region at P21 (Figure S3A-D). As different types of interneurons synapse with hippocampal excitatory neurons at distinct subcellular compartments (soma, axon initial segment, proximal and distal dendrites) indicated by the layered structures of the hippocampus (Spruston, 2008), these data suggest that endophilin A1 functions in the assembly/stabilization of inhibitory synapses formed between various presynaptic neurons and postsynaptic neurons.

Next we analyzed the GABA_A receptors and found that, consistent with changes in inhibitory postsynaptic marker, signals of the GABA_A receptor γ2 subunit were decreased in the hippocampi of both juvenile and adult *EndoA1*^{-/-} mice (**Figure 2G-I**). These data collectively indicate that endophilin A1 orchestrates the assembly of inhibitory synapses, the architecture of inhibitory postsynapses in particular, and functioning of GABAergic synapses.

Endophilin A1 localizes to the iPSD and augments the organization of the inhibitory postsynaptic machinery

To substantiate the role of postsynaptic endophilin A1 in the assembly of inhibitory synapses and exclude the impacts of presynaptic endophilin A1 and its depletion on early embryonic development, we determined the effects of endophilin A1 KO on inhibitory and excitatory synapses in cultured hippocampal pyramidal neurons during late developmental stages. As the expression levels of GABA_A receptor α1 are highly heterogeneous in individual cultured neurons (Figure S4A), we determined the effect of Cre-mediated *EndoA1* KO on surface expression of GABA_A receptor γ2 in *EndoA1*^{fl/fl} hippocampal neurons by immunofluorescence staining. Whilst no changes in surface GluA1 was detected, the fluorescence signal intensity and number of surface GABA_A receptor γ2 clusters were largely lower in dendrites of *EndoA1* KO neurons, which was rescued by re-expression of endophilin A1 (**Figure 3A-D**). We further examined the total and surface levels of synaptic




Figure 2.

EndoA1 KO causes defects in the formation and stabilization of inhibitory synapses.

(A-B) Immunofluorescence staining and confocal microscopy of gephyrin (GPN) in the hippocampal region of 3-week-old (P21) and 10-week-old (P60) *EndoA1*^{+/+} or *EndoA1*^{-/-} mouse brains. Scale bar, 50 μ m.

(C) Quantification of the signal intensity of GPN in (A) and (B). $n = 6$ mice for each genotype. (D-E) Same as (A-B) except that slices were immunostained for VGAT, (D) for P21, (E) for P60. Scale bar, 50 μ m.

(F) Quantification of VGAT signal intensity in (D) and (E). $n = 6$ mice for each genotype.

(G-H) Same as (A-B) except that slices were immunostained for GABA_AR γ 2 (γ 2). (G) for P21, (H) for P60. Scale bar, 50 μ m.

(I) Quantification of γ 2 signal intensity in (G) and (H). $n = 6$ mice for each genotype. Statistical test: unpaired two-tailed Student's t-test. ns, non-significant. All data are scatterplots with means $\pm$ S.E.M. Abbreviations: CA1-3, cornus ammonis region 1-3; DG, dentate gyrus; DAPI staining of cell nuclei is shown in blue.

proteins by immunoblotting. Biotinylation assay of surface proteins revealed that, compared with wild-type, although there was no change in total levels of synaptic proteins in *EndoA1* KO pyramidal neurons, the surface expression of GABA_ARs but not NL2 or the NR1 subunit of the NMDA-type glutamate receptor was significantly reduced (**Figure 3E,F** [↗](#)). Collectively, these data indicate that postsynaptic endophilin A1 enhances the surface clustering of GABA_ARs and facilitates the assembly of inhibitory postsynapses in pyramidal neurons.




Figure 3.

Endophilin A1 is distributed in the iPSD and regulates the assembly of the inhibitory postsynaptic machinery.

(A) Cultured *EndoA1^{fl/fl}* hippocampal neurons were transfected with construct expressing GFP or GFP-2A-Cre on DIV12, fixed and immunostained for surface GluA1 (sGluA1) or GABA_AR γ 2 (sy2) on DIV16. Shown are representative confocal microscopy images. Lower panels are magnification of boxed regions in the upper panels. Scale bar, 10 μ m in the upper panels and 5 μ m in the lower panels.

(B) Quantification of the signal intensity (left) and cluster number (right) of GPN puncta in dendrites in (A). GFP: 15 neurons; Cre: 18 neurons, from 3 independent cultures. Statistical test: unpaired two-tailed Student's t-test. ns, non-significant.

(C-D) Cultured *EndoA1^{fl/fl}* hippocampal neurons were co-transfected with constructs for GFP or GFP-2A-Cre and Flag vector or Flag-endophilin A1 (Flag-EndoA1) on DIV12, and immunostained for surface γ 2 on DIV16. Shown are representative confocal images (C) and quantification of surface γ 2 in dendrites (D). Scale bar, 2 μ m. GFP: 18 neurons; Cre: 19 neurons; Cre + EndoA1: 18 neurons, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test.

(E-F) *EndoA1^{fl/fl}* hippocampal neurons were infected with adeno-associated virus (AAV) carrying mCherry or mCherry-2A-Cre on DIV3 and surface biotinylated on DIV16. Total and surface proteins were detected by SDS-PAGE and immunoblotting. N = 3. Statistical test: unpaired two-tailed t-test.

(G) Hippocampal neurons were transfected with construct expressing GFP to label neuronal morphology on DIV12, then immunostained with antibodies against EndoA1 and GPN on DIV16. Shown are representative z stack maximum projection images captured by structured illumination microscopy (SIM). Lower panels are insets of boxed regions in the upper panel. Dendrites and spines are outlined in white. Scale bars, 2 μ m in upper panels and 0.5 μ m in magnified images.

(H) Same as (G) except that neurons were immunostained for EndoA1 and surface γ 2.

(I) Same as (G) except that neurons were immunostained for EndoA1 and VGAT.

(J) Immunoprecipitation from the membrane fraction of mouse brain tissues with antibodies against GABA_AR α 1 (α 1) or EndoA1.

(K) Immunoprecipitation from *EndoA1^{+/+}* and *EndoA1^{-/-}* brain lysates with antibodies to GABA_AR α 1. Shown are representative immunoblot (left) and quantification of neuroligin 2 (NL2), GPN and GABA_AR γ 2 co-immunoprecipitated with α 1 (right). N = 6 independent experiments. Statistical test: unpaired two-tailed Student's t-test.

(L) PSD fractions were isolated by differential centrifugation from mouse brain lysates and analyzed by immunoblotting with antibodies to EndoA1 and synaptic proteins. S1, homogenates; PSD, postsynaptic density. SYP, synaptophysin. N = 7 independent experiments. Statistical test: unpaired two-tailed Student's t-test. All data represent scatter plots with means $\pm$ S.E.M.

Next we sought to determine the molecular function of endophilin A1 in inhibitory synapses at the cellular level. By immunofluorescence staining and super-resolution microscopy of hippocampal neurons in dissociated culture, we observed partial colocalization of endophilin A1 with gephyrin as well as surface GABA_AR γ 2 in dendrites of hippocampal neurons, and juxtaposition or partial colocalization of endophilin A1 and VGAT puncta (**Figure 3G-I** and Video S2-4). We further detected the localization of endophilin A1 to inhibitory synapses by co-immunostaining with both

pre- and post-synaptic markers (Figure. S4B and Video 5). Quantitative analysis of super-resolution localization maps revealed that $\sim 47\%$ of gephyrin or Bassoon were proximal to endophilin A1 (Figure. S4G, $n = 14$), with a mean distance between endophilin A1- and gephyrin-positive pixels of ~ 120 nm, or between endophilin A1- and Bassoon-positive pixels of ~ 130 nm (Figure. S4C-F). These results indicate that endophilin A1 localizes to not only ePSD but also iPSD. Therefore, we speculated that endophilin A1 acted as a component of iPSD. Previous studies reported that synaptic GABA_ARs contain the $\gamma 2$ subunit along with α and β subunits (e.g., $\alpha 1\beta 2\gamma 2$) (Scott and Aricescu, 2019 [\[1\]](#)). The $\gamma 2$ subunit is essential for receptor clustering at the synapse, mediated by interactions with scaffold proteins (Essrich et al., 1998 [\[2\]](#)). Indeed, co-immunoprecipitation (coIP) of endogenous proteins in membrane fractions from mouse brain lysates with antibodies against GABA_AR $\alpha 1$ detected not only the known iPSD components $\gamma 2$, NL2 and gephyrin, but also endophilin A1, and vice versa (Figure 3J [\[3\]](#)). Of note, loss of endophilin A1 caused decreases in levels of GABA_AR $\gamma 2$ and gephyrin but not NL2 in anti-GABA_AR $\alpha 1$ immunoprecipitates (Figure 3K [\[4\]](#)). Further, immunoblotting of synaptosome fractions revealed that the levels of GABA_AR $\gamma 2$ and gephyrin were reduced at synapses in adult *EndoA1* KO mice (Figure 3L [\[5\]](#)). Together, these data support that endophilin A1 functions at the iPSD to promote the assembly and/or stabilization of the GABAergic postsynaptic machinery.

Endophilin A1 directly interacts with the scaffold protein gephyrin and promotes its clustering and localization of GABA_ARs to iPSD

We subsequently sought to investigate the mechanistic role of endophilin A1 in the clustering of inhibitory postsynaptic proteins. To this end, we performed coIP from mouse brain lysates with antibodies against endophilin A1 (Figure 4A [\[6\]](#)) and analyzed the immunoprecipitates by mass spectrometry. Gene Ontology (GO) analysis revealed that, in addition to proteins participating in known cellular processes involving endophilin A1, such as synaptic vesicle recycling, actin cytoskeleton remodeling, endocytosis and long-term potentiation (Gad et al., 2000 [\[7\]](#); Ringstad et al., 1999 [\[8\]](#); Schuske et al., 2003 [\[9\]](#); Verstreken et al., 2003 [\[10\]](#); Yang et al., 2021 [\[11\]](#); Yang et al., 2018 [\[12\]](#); Yang et al., 2015 [\[13\]](#)), proteins related to not only glutamatergic but also dopaminergic and GABAergic synapses were also identified, including gephyrin (Figure 4B [\[14\]](#) and Table S1).

Gephyrin is the major scaffold protein which interacts with the intracellular domains of GABA_ARs and anchors them at the inhibitory postsynaptic membrane (Essrich et al., 1998 [\[2\]](#); Kneussel et al., 1999 [\[15\]](#); Mukherjee et al., 2011 [\[16\]](#); Tretter et al., 2008 [\[17\]](#)). The cell adhesion molecule NL2 drives postsynaptic assembly at perisomatic inhibitory synapses via interaction with gephyrin and the membrane lipid-binding protein collybistin (Pouloupoulos et al., 2009 [\[18\]](#)). We speculated that endophilin A1 functions in the assembly of iPSD via interaction with the scaffold and/or intercellular adhesion proteins. To determine which inhibitory synaptic protein(s) interacts with endophilin A1, we performed coIP from HEK293T cells transiently expressing endophilin A1 and NL2, or gephyrin, and detected interaction of endophilin A1 with gephyrin but not NL2 (Figure 4C [\[19\]](#) and 4D [\[20\]](#)). Moreover, coIP from mouse brain lysates corroborated interaction between endophilin A1 and gephyrin but not GABA_AR $\alpha 1$ or $\gamma 2$ (Figure 4E [\[21\]](#)). Further, neither endophilin A2 nor A3 interacted with gephyrin (Figure 4F [\[22\]](#)).

Notably, although the expression levels of gephyrin were not affected by ablation of *EndoA1*, its distribution in the PSD fraction was decreased (Figure 3L [\[5\]](#)). In dendrites of mature cultured hippocampal neurons following *EndoA1* depletion, the signal intensity and number of gephyrin.FingR-eGFP clusters were also significantly attenuated, with no obvious changes in PSD95.FingR-eGFP clusters (Figure S5A-D), which was consistent with the reduced gephyrin but unaltered PSD95 labeled by antibodies in brain slice of mature *EndoA1* KO mice (Figure 2 [\[23\]](#) and S2). Moreover, immunofluorescence staining revealed a decrease in the clustering of gephyrin across the soma, the axon initial segment (AIS), and dendrites (Figure S5E-H). The gephyrin clustering phenotype was rescued by expression of endophilin A1 but not endophilin A2 or A3 (Figure 4G,H [\[24\]](#)), suggesting that their functions at the inhibitory postsynapses are not redundant.




Figure 4.

Endophilin A1 interacts with gephyrin and promotes its clustering in postsynaptic neurons.

(A) Silver staining of proteins co-immunoprecipitated from *EndoA1*^{+/+} mouse brain lysates with antibodies to endophilin A1.

(B) Gene Ontology (GO) enrichment analysis of EndoA1-interacting proteins identified by mass spectrometry.

(C-D) HEK293T cells co-transfected with constructs expressing Myc-tagged GPN or HA-tagged NL2 and Flag-tagged EndoA1 for 48 h were lysed for Flag IP. Input and bound proteins were analyzed by immunoblotting with antibodies against Myc, HA and Flag.

(E) Co-IP from brain lysates with antibodies to EndoA1.

(F) HEK293T cells overexpressing Flag-tagged endophilin As and Myc-tagged GPN were lysed for co-immunoprecipitation.

(G) *EndoA1*^{fl/fl} hippocampal neurons were co-transfected with constructs expressing GFP and Flag vector, or GFP-2A-Cre and Flag vector, Flag-EndoA1, Flag-EndoA2 or Flag-EndoA3 on DIV12, fixed on DIV16 and immunostained for GPN, GFP and Flag. Shown are representative confocal microscopy images. Scale bar, 2 μ m.

(H) Quantification of GPN signal intensity (left) and cluster number (right) in dendrites in (G). 36 neurons for GFP + vector; 34 neurons for Cre + vector; 38 neurons for Cre + EndoA1; 29 neurons for Cre + EndoA2; 27 neurons for Cre + EndoA3, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test.

(I) Hippocampal neurons were co-transfected with constructs expressing GFP and vector or Flag-EndoA1 on DIV12, fixed and immunostained for GPN on DIV16. Shown are representative confocal microscopy images of GPN in the soma (left) and dendrites (right). Scale bars, 5 μ m in the upper panels and 2 μ m in the lower panels.

(J) Quantification of GPN signal intensity (left) and cluster number (right) in the soma and dendrites. 34 neurons for GFP + vector; 37 neurons for GFP + EndoA1, from 3 independent cultures. Statistical test: unpaired Student's two-tailed t-test. All data represent scatterplots with means $\pm$ S.E.M.

Conversely, overexpression of endophilin A1 caused an increase in the mean signal intensity and number of gephyrin puncta in the soma and dendrites (**Figure 4I,J** [↗](#)). Collectively, these data demonstrate that endophilin A1 interacts with the scaffold protein gephyrin and facilitates its clustering at inhibitory postsynapses.

At the submembraneous postsynaptic density, gephyrin self assembles via the N-terminal G and C-terminal E domains to provide scaffold for inhibitory postsynapses and anchors GABA_ARs to the postsynaptic sites. Consistent with the effect of *EndoA1* KO on the clustering of gephyrin (**Figure 4G,H** [↗](#)) and GABA_AR $\gamma 2$ (**Figure 3A-D** [↗](#)), immunofluorescence staining and confocal microscopy analysis showed that, depletion of endophilin A1 also caused a decrease in the localization of GABA_AR $\gamma 2$ to gephyrin puncta in *EndoA1* KO neurons (**Figure 5A-C** [↗](#)). Moreover, overexpression of endophilin A1 also enhanced clustering of GABA_AR $\gamma 2$ in the soma and dendrites (**Figure 5D,E** [↗](#)). Together these data indicate that endophilin A1 promotes anchoring of GABA_ARs to the iPSD.




Figure 5.

Both BAR and SH3 domains are required for endophilin A1-mediated organization of iPSD.

(A) *EndoA1^{fl/fl}* hippocampal neurons expressing GFP or Cre were immunostained for GPN, surface γ 2 and GFP on DIV16. Shown are representative confocal microscopy images. Scale bars, 5 μ m in left panels and 1 μ m in magnified images.

(B-C) Quantification of the mean intensity of GPN (green) and surface γ 2 (red) and their colocalization in (A). 21 neurons for GFP, 22 neurons for Cre, from 3 independent cultures. Statistical test: unpaired two-tailed Student's t-test. All data represent scatterplots with means $\pm$ S.E.M.

(D) DIV12 hippocampal neurons were co-transfected with constructs expressing GFP and vector or Flag-EndoA1, fixed on DIV16 and immunostained for surface γ 2. Shown are representative confocal microscopy images of surface γ 2 in the soma (left) and dendrites (right). Scale bars, 5 μ m in the upper panels and 2 μ m in the lower panels.

(E) Quantification of surface γ 2 mean intensity (left) and cluster number (right) in the soma and dendrites. Soma: 20 neurons for GFP + vector, 20 neurons for GFP + EndoA1; Dendrites: 35 neurons for GFP + vector, 30 neurons for GFP + EndoA1, from 3 independent cultures. Statistical test: unpaired two-tailed Student's t-test. All data represent scatterplots with means $\pm$ S.E.M.

(F) Schematic representation of EndoA1 and GPN domain structure and diagrams of fragments used for interaction mapping.

(G) GST-tagged EndoA1 fragments conjugated to Glutathione-Sepharose beads were incubated with lysates of HEK293T cells overexpressing Myc-tagged GPN. Input and bound proteins were analyzed by immunoblotting with antibodies against Myc. Bottom panel: Coomassie blue-stained SDS-PAGE gel showing GST fusion proteins. FL, full length.

(H) HEK293T cells co-transfected with constructs encoding Myc-tagged GPN and Flag-tagged EndoA1 fragments were lysed for Flag IP. Input and bound proteins were analyzed by immunoblotting with antibodies against Myc and Flag.

(I) Binding of His-tagged EndoA1 to GST-tagged GPN fragments in pull-down assay. Bound proteins were analyzed by SDS-PAGE and immunoblotting with anti-His antibodies. Input and bound proteins were analyzed by immunoblotting with antibodies against His. Bottom panel: Coomassie blue-stained SDS-PAGE gel showing GST fusion proteins. FL, full length. Right panel: Quantification of EndoA1 binding to GPN domain. N = 4 independent experiments. Statistical test: one-way ANOVA by Tukey post hoc test. All data represent scatterplots with means $\pm$ S.E.M.

(J) DIV12 *EndoA1^{fl/fl}* hippocampal neurons were co-transfected with constructs expressing GFP and Flag vector or GFP-2A-Cre, and Flag vector or Flag-EndoA1 FL, Flag-EndoA1 Δ SH3, Flag-EndoA1 aa172-352, fixed and immunostained for GPN on DIV16. Shown are representative confocal microscopy images of GPN. Scale bar, 2 μ m.

(K) Quantification of mean intensity (left) and cluster number (right) in dendrites are shown for GPN in (J). 41 neurons for GFP + vector; 44 neurons for Cre + vector; 35 neurons for Cre + FL; 24 neurons for Cre + Δ SH3; 27 neurons for Cre + aa172-352, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test. All data represent scatterplots with means $\pm$ S.E.M.

(L) Same as (I) except that neurons were immunostained for surface γ 2. Scale bar, 2 μ m.

(M) Quantification of mean intensity (left) and cluster number (right) in dendrites are shown for surface $\gamma 2$ in (K). 38 neurons for GFP + vector; 31 neurons for Cre + vector; 26 neurons for Cre + FL; 33 neurons for Cre + Δ SH3; 36 neurons for Cre + aa172-352, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test. All data represent scatterplots with means $\pm$ S.E.M.

To further understand the mechanism by which endophilin A1 regulates the assembly and function of inhibitory postsynapses, next we attempted domain mapping of the endophilin A1-gephyrin interaction. To this end, we generated a series of truncation and internal deletion mutants for endophilin A1 or gephyrin (**Figure 5F**). For those which did not express in mammalian cells, we determined their interaction with gephyrin by in vitro binding assays with recombinant proteins. None of the truncation or internal deletion mutants lost the ability to bind to gephyrin, and deletion of neither the BAR nor the SH3 domain weakened the binding (**Figure 5G,H**), suggesting that they are involved in the interaction with gephyrin. Pull-down assay showed both G and E domains but not C domain of gephyrin were involved in the interaction with endophilin A1 (**Figure 5I**). Indeed, the decreases in clustering of either gephyrin or GABA_A $\gamma 2$ in *EndoA1* KO neurons were not rescued by expression of endophilin A1 fragments (**Figure 5J-M**), indicating that all functional domains of endophilin A1 are required for the assembly of the GABAergic postsynaptic machinery.

Actin polymerization is required for stabilization of inhibitory postsynapses

Endophilin A1 functions in the morphogenesis and structural plasticity of dendritic spines by promoting actin polymerization underneath the plasma membrane (Yang et al., 2021; Yang et al., 2018; Yang et al., 2015). Of note, previous studies reported controversial results about the role of the actin and microtubule cytoskeleton in gephyrin clustering (Allison et al., 2000; Charrier et al., 2006; Kirsch and Betz, 1995).

To further investigate the mechanism underlying endophilin A1-mediated assembly of inhibitory postsynapses, we asked whether actin polymerization contributes to the stabilization of iPSD and GABA_A receptor clustering in dendrites. Consistent with previous studies (van Bommel et al., 2019), we observed large patches of F-actin labeled by LifeAct-mCherry in spines and small puncta in dendritic shafts that exhibited partial colocalization with or adjoined gephyrin in cultured hippocampal neurons by super-resolution microscopy (Figure S6A and Video S6). We then determined the impact of F-actin depolymerization on gephyrin-labeled iPSD and $\gamma 2$ -containing GABA_ARs in cultured hippocampal neurons. Treatment of mature neurons with nocodazole, a microtubule depolymerizing reagent, for one hour (short-term) or two hours (long-term), caused decreases in the number of both PSD95 and gephyrin puncta (**Figure 6A-D**), which was in good agreement with previous findings that microtubule dynamics regulates the spatial organization of both excitatory and inhibitory postsynaptic structures (Gu et al., 2008; Jaworski et al., 2009; Kirsch and Betz, 1995; Kirsch et al., 1991). Short-term or long-term incubation with latrunculin A, which disrupts actin polymerization, also caused decreases in the number of both ePSD and iPSD puncta (**Figure 6A-D**), indicating that actin polymerization also contributes to the stabilization of the iPSD. We then tested the effect of actin depolymerization on GABA_A receptor clustering and found that, long-term but not short-term latrunculin A application caused a reduction in the number of GABA_A $\gamma 2$ puncta in dendrites (**Figure 6E-H**), this finding supported the observation that the decrease in gephyrin cluster preceded the reduction in GABA_A $\gamma 2$ receptor cluster following latrunculin A treatment (**Figure 6I**). In addition, colocalization analysis revealed less recruitment of $\gamma 2$ to gephyrin puncta even after short-term drug application, without changes in $\gamma 2$ clustering (**Figure 6J,K**). Together, these data demonstrate that actin polymerization is essential for the stabilization of iPSD and anchoring of GABA_ARs to the postsynaptic sites.




Figure 6.

Actin polymerization is required for stabilization of inhibitory postsynapses.

(A-D) DIV12 cultured hippocampal neurons transiently expressing GFP were treated with DMSO (vehicle control), 1 μ M Latrunculin A (LatA) or 1 μ M Nocodazole (Noco) for 60 min (A) or 120 min (C) on DIV16, respectively. Changes in postsynaptic proteins GPN (red) and PSD95 (green) were detected by confocal microscopy. Quantitative data shown were the mean intensity (left) and cluster number (right) of GPN and PSD95 at 60 min (B) and 120 min (D). Ctrl: 20 neurons for 60 min; 29 neurons for 120 min; LatA: 20 neurons for 60 min; 33 neurons for 120 min; Noco: 23 neurons for 60 min; 22 neurons for 120 min, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test. Scale bar, 5 μ m.

(E-H) Same as (A-D) except that neurons were immunostained for GPN (red) and surface γ 2 (green). Changes in GPN and surface γ 2 at 60 min (E) or 120 min (G) were detected by confocal microscopy. Quantitative data shown were the mean intensity (left) and cluster number (right) of GPN and surface γ 2 at 60 min (F) and 120 min (H). Ctrl: 35 neurons for 60 min; 24 neurons for 120 min; LatA: 34 neurons for 60 min; 20 neurons for 120 min; Noco: 34 neurons for 60 min; 24 neurons for 120 min, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test. Scale bar, 5 μ m.

(I) Levels of synaptic proteins normalized to control neurons following LatA or Noco treatment in (A,C,E,G).

(J-K) The colocalization between surface γ 2 and GPN signals in dendrites in (E) and (G). Statistical test: one-way ANOVA by Tukey post hoc test. All data are scatterplots with means $\pm$ S.E.M.

Both the membrane-binding and p140Cap-binding capacities of endophilin A1 are indispensable for its function at the inhibitory postsynapses

At the excitatory synapse, the plasma membrane-associated endophilin A1 regulates the morphogenesis, maturation and structural plasticity of dendritic spines by promoting branched actin polymerization underneath the postsynaptic membrane via its interaction with the cytoskeleton regulator p140Cap (Yang et al., 2021 [DOI](#); Yang et al., 2018 [DOI](#); Yang et al., 2015 [DOI](#)). Having established the requirement for both endophilin A1 and the actin cytoskeleton in the assembly and stabilization of the inhibitory postsynaptic structure, we speculated that regulation of inhibitory postsynapse assembly by endophilin A1 necessitated its plasma membrane-binding and actin polymerization-promoting activities. Silencing p140Cap expression in postsynaptic CA1 pyramidal cells by shRNA-mediated knockdown caused decreases in the clustering of gephyrin puncta in dendrites, which was restored by overexpression of p140Cap (Figure S6B, C), suggesting that p140Cap-mediated actin polymerization is required for the assembly of iPSD in postsynaptic neurons. However, compared with mere depletion of endophilin A1, p140Cap knockdown did not further attenuate the clustering of gephyrin in *EndoA1* KO neurons (Figure S6D, E). Moreover, overexpression of p140Cap did not rescue the iPSD phenotype in *EndoA1* KO neurons (**Figure 7A,B** [DOI](#)), suggesting that p140Cap-enhanced actin polymerization promotes the organization of iPSD through endophilin A1-mediated pathway. Indeed, overexpression of neither the membrane-binding deficient (KKK-EEE) nor the p140Cap-binding deficient (Y343A) mutant of endophilin A1 (Yang et al., 2021 [DOI](#)) could rescue the iPSD defect (**Figure 7A,B** [DOI](#)). Consistently, neither p140Cap nor the endophilin A1 loss-of-function mutants could rescue the GABA_A γ 2 clustering phenotype in *EndoA1* KO neurons (**Figure 7C,D** [DOI](#)). Collectively, these data indicate that both plasma membrane association and actin polymerization-promoting activities are required for endophilin A1 to promote the assembly of the inhibitory postsynaptic machinery.




Figure 7.

Both the membrane-binding and p140Cap-binding capacities of endophilin A1 are required for iPSD assembly and E/I balance.

(A) Cultured hippocampal neurons from *EndoA1^{fl/fl}* mice were co-transfected with construct expressing GFP and Flag vector or GFP-2A-Cre and Flag vector, Flag-EndoA1 wild-type (WT), mutants (KKK-EEE or Y343A) or p140Cap, and analyzed on DIV16 by immunofluorescence staining for GPN (green) and GFP (red). Scale bar, 2 μ m.

(B) Quantification of GPN mean intensity (left) and cluster number (right) in (A). 35 neurons for GFP vector; 43 neurons for Cre + vector; 32 neurons for Cre + WT; 33 neurons for Cre + KKK-EEE; 36 neurons for Cre + Y343A; 33 neurons for Cre + p140Cap, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test.

(C) Same as (A) except that neurons were immunostained for surface γ 2 on DIV16. Scale bar, 2 μ m.

(D) Quantification of surface γ 2 mean intensity (left) and number (right) in (C). 44 neurons for GFP + vector; 42 neurons for Cre + vector; 32 neurons for Cre + WT; 35 neurons for Cre + KKK-EEE; 35 neurons for Cre + Y343A; 32 neurons for Cre + p140Cap, from 3 independent cultures. Statistical test: one-way ANOVA by Tukey post hoc test.

(E) Experimental scheme for seizure scoring of *EndoA1^{fl/fl}* mice stereotactically co-injected with AAVs carrying mCherry and Cre, and EGFP and EndoA1 or its mutants (KKK-EEE and Y343A) in the hippocampal CA1 region. Mice were intraperitoneally administered PTZ for 12 days after AAV injections, then seizures were scored.

(F) PTZ-induced seizures in mice injected with the indicated AAVs were scored for 30 min. Ctrl, 7 mice; Cre, 7 mice; Cre + WT, 7 mice; Cre + KKK-EEE, 6 mice; Cre + Y343A, 7 mice.; Statistical test: repeated measures ANOVA followed by Tukey post-hoc test at each time point.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, red color is for Cre vs Ctrl, green color is for Cre + WT vs Cre. All data are means $\pm$ S.E.M.

(G-K) AAVs expressing Cre (AAV-mCherry-2A-Cre) and GFP (AAV-EGFP) or Cre and EndoA1 WT or mutant (AAV-EGFP-2A-EndoA1, AAV-EGFP-2A-EndoA1 KKK-EEE, AAV-EGFP-2A-EndoA1 Y343A) were stereotactically injected into the CA1 regions of *EndoA1^{fl/fl}* mouse brain at P0. Acute hippocampal slices were prepared on P14-P21 for dual recording of evoked excitatory postsynaptic currents (eEPSC) and inhibitory postsynaptic currents (eIPSC). Shown are representative traces and pairwise comparisons of noninfected (Ctrl) and infected neurons in the same slice. Scale bar = 50/50, 50/50, 50/50, 50/50, 40/20 (pA/ms) in (G); Scale bar = 100/100, 100/400, 100/100, 100/200, 300/100 (pA/ms) in (H); Scale bar = 50/50, 50/50, 50/50, 50/50, 50/50 (pA/ms) in (I); Scale bar = 100/100, 200/100, 100/100, 100/200, 100/200 (pA/ms) in (J). Statistical test: unpaired two-tailed Student's t-test. ns, non-significant; All data are scatterplots with means $\pm$ S.E.M. All number of recording were from more than 7 pyramidal neurons of 3 independent animals.

Next, we determined the mechanistic role of endophilin A1 in vivo by rescuing the epilepsy susceptibility caused by *EndoA1* KO. Indeed, when AAVs expressing the Cre recombinase and endophilin A1 were co-injected into the hippocampal CA1 region of *EndoA1^{fl/fl}* mice, only the wild-type but not the membrane-binding or p140Cap-binding deficient mutant of endophilin A1 rescued the early onset and severity of PTZ-induced seizures (**Figure 7E,F** and Video S7). Consistently, electrophysiological recording in acute brain slices indicated that only wild-type endophilin A1 but not the mutants restored the amplitude of eIPSC and E/I balance in *EndoA1* KO neurons (**Figure 7G-K**). All together, these data indicate that endophilin A1 contributes to the structure and function of inhibitory synapses by promoting the submembrane assembly of iPSD and synaptic localization of the GABA_ARs, thereby maintaining the balance of excitatory and inhibitory activities in neural circuits (**Figure 8**).




Figure 8.

Model for the role of endophilin A1 at the inhibitory postsynapses and E/I balance.

(A) Endophilin A1 directly interacts with the inhibitory postsynaptic scaffold protein gephyrin to promote organization of iPSD and synaptic recruitment/stabilization of the GABA_ARs via its plasma membrane association and actin polymerization-promoting activities. **(B)** Loss of endophilin A1 impairs inhibitory synaptic neurotransmission and disrupts the balance of excitatory/inhibitory activities in the neural circuits, leading to enhanced epileptic seizure susceptibility.

Discussion

Although collybistin is implicated in plasma membrane recruitment and clustering of gephyrin and GABA_A α 2 (Saiepour et al., 2010 [DOI](#)), studies of collybistin KO mice showed that the GABA_A α 3 subunit promotes gephyrin-mediated formation of inhibitory synapses in the absence of collybistin (Wagner et al., 2021 [DOI](#)), raising the possibility that formation of a subset of GABAergic synapses requires gephyrin-interacting membrane-associated proteins other than collybistin. In the present study, we identify endophilin A1 as a novel binding partner for gephyrin and demonstrate that it promotes the assembly of inhibitory synapses and clustering of GABA_ARs at the iPSD. Loss of endophilin A1 activity causes impairment of inhibitory synaptic transmission in neuronal circuits and increases susceptibility to seizures. In the central nervous system, inhibitory interneurons are highly diverse in morphologies, molecular profiles and physiological functions. Presynaptic terminals from different types of interneurons target distinct subcellular compartments in postsynaptic pyramidal neurons where GABA_A receptors of specific subunit compositions are located. Parvalbumin-expressing (PV+) interneurons usually contact the soma and proximal dendrites, whereas somatostatin-expressing (SST+) interneurons mainly innervate the dendrites of pyramidal cells (Klausberger and Somogyi, 2008 [DOI](#)). Although different inhibitory synapses localized in different subcellular compartments of postsynaptic neurons have distinct properties and express a diversity of synaptic proteins, gephyrin remains a core postsynaptic component (Favuzzi et al., 2019 [DOI](#)). Our study revealed that endophilin A1 promotes the clustering of gephyrin in all subcellular compartments of pyramidal neurons, indicating that endophilin A1 functions in the organization of inhibitory postsynapses innervated by various presynaptic interneurons.

Among the endophilin A proteins, endophilin A1 is the most abundant in the brain (Milosevic et al., 2011 [DOI](#)). Although they have redundant roles in physiological or pathological functions (Milosevic et al., 2011 [DOI](#)), endophilin A1, but not A2 or A3, promotes spine morphogenesis and excitatory postsynaptic organization (Yang et al., 2018 [DOI](#); Yang et al., 2015 [DOI](#); Milosevic et al., 2011 [DOI](#)). Here, we uncover a previously unanticipated role for endophilin A1 at GABAergic synapses. By deleting endophilin As in specific neuron subtypes, we found that endophilin A1 regulates GABAergic postsynaptic compartments. Endophilin A1, rather than A2 or A3, interacts with gephyrin and modulates the inhibitory postsynaptic machinery. Although endophilin A1 might function redundantly with other endophilin As in synaptic vesicle recycling at the presynaptic site, it participates in the organization or reorganization of postsynaptic structures through specific binding to postsynaptic proteins, a function that cannot be fully replaced by other endophilin As at both excitatory and inhibitory postsynapses.

The membrane lipid-binding protein collybistin interacts with the trans-synaptic cell adhesion molecule NL-2 to recruit gephyrin and GABA_A α 2 to the postsynaptic membrane (Kalscheuer et al., 2009 [DOI](#); Kins et al., 2000 [DOI](#); Pouloupoulos et al., 2009 [DOI](#); Saiepour et al., 2010 [DOI](#)). Adamtsl3 secreted from presynapses binds to the transmembrane protein DCC (Deleted in Colorectal Cancer) and controls its synaptic localization, which is required for the maintenance of GABAergic synapses in the adult brain (Cramer et al., 2023 [DOI](#)). Anchoring of the α 5 subunit of GABA_A α 5 to the actin cytoskeleton through radixin, an actin-binding protein, facilitates its clustering (Loebrich et al., 2006 [DOI](#)). These findings underscore the necessity for investigation of mechanism(s) regulating the plasma membrane-iPSD-actin cytoskeleton interaction. As membrane-binding and actin polymerization-promoting activities of endophilin A1 are both required for its function in enhancing iPSD formation and γ 2-containing GABA_A α 5 clustering to iPSD, we propose that membrane-bound endophilin A1 promotes postsynaptic assembly by coordinating the plasma membrane tethering of the postsynaptic protein complex and its stabilization with the actin cytomatrix (**Figure 8** [DOI](#)). In contrast, clustering of β 2/3 subunit-containing GABA_ARs is independent of the actin or microtubule cytoskeleton in hippocampal neurons (Allison et al.,

2000 [\[10\]](#); Kasaragod et al., 2022 [\[11\]](#)). Given that synaptic and extrasynaptic GABA_ARs are composed of distinct subunits (Kasaragod et al., 2022 [\[11\]](#); Koh et al., 2023 [\[12\]](#)), these findings indicate that the actin cytoskeleton plays distinct roles in the clustering and possibly dynamic changes in the subunit composition and subcompartmental localization of different types of GABA_ARs. Of note, the subcellular distribution of gephyrin, which has a tubulin-binding motif, is not affected by depolymerization of F-actin or microtubule (Allison et al., 2000 [\[13\]](#)). Overall, the discrepancy in data about the involvement of the actin cytoskeleton in inhibitory synapses may be due to differences in drug concentration, treatment time, receptor subtypes and neuronal cell types (Mody and Pearce, 2004 [\[14\]](#); Sexton et al., 2021 [\[15\]](#)). In the present study, the impaired clustering of gephyrin and GABA_A γ2 by F-actin depolymerization underscores the essential role of F-actin in the assembly and stabilization of the inhibitory postsynaptic machinery. Membrane-bound endophilin A1 promotes F-actin polymerization beneath the plasma membrane through its interaction with p140Cap, an F-actin regulatory protein, thereby facilitating and/or stabilizing the clustering of gephyrin and γ2-containing GABA_A receptors at postsynapses.

The number of synaptic GABA_A receptors is subject to alterations not only during development and neuronal plasticity but also during pathological processes such as epilepsy (Nusser et al., 1998 [\[16\]](#)). Synaptic GABA_A receptors are recruited directly from extrasynaptic pools following exocytosis, and their endocytosis occurs exclusively at extrasynaptic sites (Bogdanov et al., 2006 [\[17\]](#)). Therefore, lateral diffusion of GABA_ARs between synaptic and extrasynaptic sites and their iPSD anchoring are very important for their synaptic distribution (Thomas et al., 2005 [\[18\]](#)). In good agreement with the finding that GABA_ARs stabilization at synapses depends on gephyrin clustering (Jacob et al., 2005 [\[19\]](#)), we found that decreases in gephyrin clustering precedes the reduction in surface GABA_ARs following F-actin depolymerization (**Figure 6** [\[20\]](#)). As the core scaffold protein in iPSD, gephyrin is highly enriched at the inhibitory postsynaptic sites and drives the formation of iPSD sheets via phase separation (Bai et al., 2021 [\[21\]](#)). As binding of gephyrin by its partner proteins inhibits the iPSD condensate formation via autoinhibition of the multivalent interactions between its E domain and GABA_ARs (Bai et al., 2021 [\[21\]](#); Hoffmann and Milovanovic, 2021 [\[22\]](#); Liu et al., 2020 [\[23\]](#)), it is conceivable that endophilin A1 binding not only tethers the scaffold protein to the plasma membrane, but also releases it from the autoinhibitory conformation to facilitate the phase separation of the gephyrin-receptor complexes. Whether the activity of endophilin A1 is regulated by neuronal activity to participate in the structural and functional remodeling of iPSD awaits further investigation.

At the presynaptic site of excitatory synapses, endophilin A1 functions in synaptic vesicle recycling by recruiting the clathrin-uncoating factor synaptojanin (Milosevic et al., 2011 [\[24\]](#); Schuske et al., 2003 [\[25\]](#); Verstreken et al., 2003 [\[26\]](#)). At the excitatory postsynapses, plasma membrane-associated endophilin A1 facilitates the morphogenesis, stabilization, maturation and plasticity of dendritic spines by promoting actin polymerization via its interaction with p140Cap (Yang et al., 2021 [\[27\]](#); Yang et al., 2018 [\[28\]](#); Yang et al., 2015 [\[29\]](#)). In this study, we demonstrate that endophilin A1 functions upstream of p140Cap in not only excitatory but also inhibitory postsynapses of excitatory neurons. Intriguingly, recent studies reported that p140Cap is present in the inhibitory presynaptic compartment and regulates GABAergic synaptogenesis and neurotransmission (Russo et al., 2019 [\[30\]](#)), raising the possibility that endophilin A1 and p140Cap also coordinate to regulate the organization and activity of GABAergic synapses at the inhibitory presynaptic site. Remodeling of inhibitory synapses and dendritic spines are spatially clustered by sensory inputs in the central nervous system (Chen et al., 2012 [\[31\]](#)). Notably, the turnover of inhibitory synapses outpace that of excitatory synapses (Chen et al., 2012 [\[31\]](#)). Therefore, it is conceivable that deficits in inhibitory synapses are much more severe in *EndoA1* KO mice, leading to perturbation of the E/I balance and increased epileptic vulnerability.

Materials and Methods

Ethics statement

All animal experiments were approved by and performed in accordance with the guidelines of the Animal Care and Use Committee of Institute of Genetics and Developmental Biology, Chinese Academy of Sciences (Approval code: AP2022008). All animals were housed in standard mouse cages at 22–24°C on a 12 h light/dark cycle with access to food and water freely.

Animals

Generation of *EndoA1^{fl/fl}* and *EndoA1^{-/-}* (KO first) mice on the C57BL/6J background were as previously described. *EndoA1^{+/+}* and *EndoA1^{-/-}* mice were bred as littermates from heterozygous breeders. Genotyping of mouse lines was performed by genomic PCR of tail prep DNA from offspring with the following primer pairs: loxPF/loxPR: 5'-CAAGGACTCCCAGAGACCTAGCATC-3' and 5'-GAGATGGCGCAACGCAATTAAT-3' (A PCR product of 375 base pairs in *EndoA1* KO first mice but none in wild-type mice). zptF/zptR: 5'-gtaagcggctctagcgcgtgtct-3' and 5'-GCAGGGGCATGTAGGTGGCTCAAC-3' (A PCR product of 466 base pairs in wild-type mice, none in *EndoA1^{-/-}* mice, and of 627 base pairs in *EndoA1^{fl/fl}* mice).

Constructs and viruses

pCMV-Tag2B-EndoA1, pCMV-Tag2B-p140Cap, pCMV-Tag2B-EndoA1 Y343A, pCMV-Tag2B-EndoA1 KKK-EEE, pCMV-Tag2B-EndoA1 (aa172-352), pGEX-4T-1-SH3 (aa295-352), pET-28a (+)-EndoA1 and LifeAct-mCherry constructs were described previously (Yang et al., 2021 [DOI](#); Yang et al., 2015 [DOI](#)). pAOV-CAMKII α -mCherry-2A-3FLAG were purchased from OBIO Technology (Shanghai) Corp. Ltd., (Shanghai, China). pNICE-NL2(-) was a gift from Peter Scheiffele (Addgene plasmid # 15246) (Chih et al., 2006 [DOI](#)). pCR3-FLAG-Gephyrin P1 was a gift from Shiva Tyagarajan (Addgene plasmid # 68816) (Tyagarajan et al., 2013 [DOI](#)). pCAG_PSD95.FingR-eGFP-CCR5TC and pCAG_GPN.FingR-eGFP-CCR5TC were gifts from Don Arnold (Addgene plasmid # 46295 & #46296) (Gross et al., 2013 [DOI](#)). The pCMV-Tag3B-GPN construct was generated by cloning gephyrin cDNA amplified from pCR3-FLAG-gephyrin into pCMV-Tag3B (Clontech Laboratories, Inc.). The pGEX-4T-1 EndoA1 Δ BAR (Δ aa6-242) and pGEX-4T-1 EndoA1 Δ SH3 (Δ aa1-352) constructs were generated by cloning fragments amplified from pCMV-Tag2B-EndoA1 into pGEX-4T-1. The pGEX-4T-1 GPN G (aa15-169), pGEX-4T-1 GPN C (aa170-326) and pGEX-4T-1 GPN E (aa327-733) constructs were generated by PCR amplification from full-length constructs and subcloned into pGEX-4T-1. All constructs were verified by DNA sequencing. Viral particles of adeno-associated virus (AAV) carrying mCherry, EGFP, the Cre recombinase, EndoA1 WT or mutants were purchased from OBIO Technology (Shanghai) Corp. Ltd., (Shanghai, China). Viral particles of rAAV-EF1a-PSD95.FingR-eGFP-CCR5TC and rAAV-EF1a-mRuby2-Gephyrin.FingR-IL2RGTC (Bensussen et al., 2020 [DOI](#)) were purchased from brain case Corp. Ltd., (Shenzhen, China).

Antibodies

The following antibodies were obtained from commercial sources: rabbit anti-endophilin A1 (159 002, Synaptic Systems GmbH, Germany), mouse anti-Gephyrin (147 111, Synaptic Systems), guinea pig anti-gephyrin (147 318, Synaptic Systems), mouse anti-neurologin 2 (129 511, Synaptic Systems), mouse anti-GluA1 (MAB2263, Millipore), rabbit anti-GABA_A α 1 (06-868, Millipore Sigma), rabbit anti-GABA_A γ 2 (224 003, Synaptic Systems), mouse anti-SYP (D-4) (sc-17750, Santa Cruz), rabbit anti-FLAG (F3165, Sigma-Aldrich), rabbit and mouse anti-RFP which recognizes DsRed, mCherry and mRuby2 (PM005 and M165-3, Medical & Biological Laboratories, Naka-ku Nagoya, Japan); mouse anti- α -tubulin (T9026, Sigma-Aldrich), mouse anti-PSD95 (610495, BD Biosciences, San Diego, CA) and mouse anti-GST (M071-3, MBL) for western blotting (WB) and rabbit anti-VGAT (131

004, Synaptic Systems), mouse anti-PSD95 (75-028, NeuroMab, Davis, CA), guinea pig anti-Ankyrin G (386 004, Synaptic Systems), guinea pig anti GABA_AR γ 2 (224 004, Synaptic Systems) and mouse anti-Bassoon (sc-58509, Santa Cruz) for immunofluorescence staining; Rabbit anti-p140Cap was described previously (Yang et al., 2015 [DOI](#)). Alexa Fluor dye-conjugated secondary antibodies for immunofluorescence staining were from Molecular Probes (Invitrogen, Carlsbad, CA, USA).

Pentylentetrazole (PTZ)-kindling and Seizure Behavior Scoring

To evaluate seizure susceptibility, 8-10-week-old male and female *EndoA1*^{+/+} or *EndoA1*^{-/-} littermates or *EndoA1*^{fl/fl} littermates were intraperitoneally administered with PTZ (25 mg/kg; K0250, Sigma-Aldrich) or saline (control) for 8-12 days, and the resulting seizure behaviors were video-recorded for the next 30 min. Seizure susceptibility was measured blindly by rating seizures on a scale of 0 to 7 as follows (Van Erum et al., 2019 [DOI](#)): no abnormal behavior (0), reduced motility and prostate position (1), facial jerking (Schulz et al.), neck ierks (Huo et al.), clonic seizure (sitting) (4), clonic, tonic-clonic seizure (lving on belly) (5), clonic, tonic-clonic seizure (lving on side) and wild jumping (6), tonic extension, possibly leading to respiratory arrest and death (7).

Stereotaxic injection

For virus injection, 8-9-week-old naive male and female littermates were anesthetized and stereotactically injected with viral particles in the hippocampal CA1 region as described (Yang et al., 2018 [DOI](#)). Briefly, mice were weighed and anesthetized with isoflurane (2% 2,2,2-tribromoethanol, Sigma), and secured in a stereotaxic apparatus. The scalp of mouse was cut along the midline between the ears after sterilized with iodophors and 75% (vol/vol) alcohol. Holes for inserting needle were drilled in the bilateral skull. Viral particles carrying pAOV-CAMKII α -MCS-mCherry-3FLAG or pAOV-CAMKII α -mCherry-2A-Cre and pAAV-CaMKII α -EGFP-2A-MCS-3FLAG or pAAV-CaMKII α -EGFP-2A-MCS-3FLAG-EndoA1 WT or mutants (1.2 μ L, 2.0×10^{12} viral genomes/mL) (OBiO Technology Corp., Ltd) were injected bilaterally into the hippocampal CA1 regions using the following coordinates relative to bregma: 2.0 mm posterior, 1.8 mm lateral, and 1.4 mm ventral at a rate of 0.125 μ L/min using a microinjection system (World Precision Instruments). The needle was kept in place for 2 min before withdrawal, and the skin was sutured with surgical line. The virus-injected mice were tested for behavior two weeks later.

Immunofluorescence staining of brain sections

Male and female littermates (P21 or P60) were anesthetized and immediately perfused, brains were dissected out, fixed overnight in 4% (vol/vol) paraformaldehyde, incubated overnight with 30% sucrose in phosphate-buffered saline (PBS), and then sliced into 30 μ m-thick coronal sections using a cryotome (Model CM-1950; Leica Biosystems). For immunostaining of brain sections, floating 30 μ m-thick slices were rinsed with PBS and permeabilized in 0.4% Triton X-100 in 0.01M PBS for 30 min. Cyrosections were blocked with 2% BSA in PBS containing 0.4% Triton X-100 for 2 h at room temperature (RT), then incubated with primary antibodies overnight at 4 °C. Secondary antibodies conjugated with Alexa Fluor 488 or 555 were used for detection. Sections were then incubated with DAPI (Roche, Grenzach-Wyhlen, Germany) for nuclear staining for 3 min at RT. Following rinsing, cyrosections were mounted on gelatin-coated slides and covered with coverslip with mounting medium. Confocal images were collected using the Spectral Imaging Confocal Microscope Digital Eclipse A1Si (Nikon, Tokyo, Japan) with a 40 $\times$ Plan Fluo (NA 1.30) oil objective (Yang et al., 2018 [DOI](#)).

Cell culture, transfection and drug treatment

Primary hippocampal neurons were cultured as previously described (Yang et al., 2021 [DOI](#)). Briefly, Hippocampi of female or male pups (P0) were rapidly dissected under sterile conditions, kept in cold DMEM (4°C) with high glucose. Tissue was then digested with 0.25% trypsin solution at 37°C for 18-20 min, triturated in DMEM supplemented with 10% F12 (11765054, Gibco) and 10% FBS (Gibco, Carlsbad, CA, USA). Then the tissue was mechanically dissociated into single cells by gentle

pipetting up and down. Neurons were plated at a density of $\sim 3.0\text{--}3.5 \times 10^5$ cells/well in poly-D-lysine (P6407, Sigma-Aldrich) -coated 6-well plate and at a density of $2.5\text{--}3.0 \times 10^4$ cells/well in 24-well plate. Four hours later, media was changed to Neurobasal A (NB-A) growth medium supplemented with 2% B27 supplement (17504-044, Gibco), 1% GlutaMAX (35050-061, Gibco) and 33 mM glucose. Culture media were changed by half volume with Neurobasal maintenance media once a week. The cultures were maintained at 37°C with 5% CO₂. Experiments were performed at DIV 14-16 unless otherwise stated. For neuronal transfection, hippocampal neurons (DIV11) were transfected using Lipofectamine-LTX (15338100, Invitrogen, ThermoFisher Scientific). For 12 mm coverslips in individual wells of a 24 well plate, 0.5 µg of DNA was mixed with 0.5 µL PLUS reagent in 50 µL of NB-A and incubated for 5 min at RT, then gently mixed with 1 µL Lipofectamine LTX in 50 µL NB-A and incubated for 20 min at RT. The DNA-Lipofectamine LTX mixture was then added to the neurons in NB-A at 37°C in 5% CO₂ for 1 h. Neurons were then rinsed with NB-A and incubated in the original medium at 37°C in 5% CO₂ for 3-4 days.

For HEK293T cell culture, DMEM supplemented with 10% FBS were used. Cell transfections were performed using Lipofectamine-2000 (11668019, Invitrogen, ThermoFisher Scientific) according to the manufacturer's instructions. Cells were harvested 24-48 h post transfection.

For inhibitor treatment, primary neurons cultured on coverslips were pre-incubated with Latrunculin A (1 µM, 76343-93-6, Sigma-Aldrich), Nocodazole (1 µM, 31430-18-9, Sigma-Aldrich) for 60 min or 120 min.

Immunofluorescence staining of cultured neurons

Hippocampal neurons were fixed in 4% paraformaldehyde/4% sucrose in PBS at RT for 15 min, then permeabilized in 0.4% Triton X-100 in PBS for 10 min. After blocking with 2% BSA in PBS containing 0.4% Triton X-100 for 30 min at RT, neurons were incubated with primary antibodies for 1 h at RT or overnight at 4°C, and appropriate secondary antibodies conjugated with Alexa Fluor 488, Alexa Fluor 555, or Alexa Fluor 647 were used for detection. The primary antibody was diluted in a PBS-2% BSA solution and incubated overnight at 4°C. After washing with PBS three times, samples were incubated with Alexa Fluor secondary antibodies (1:2000, Thermo Fisher) for 1 h at RT, and mounted without bubbles.

Biotinylation of cell surface proteins

Surface biotinylation experiments were performed as previously described (Guo et al., 2022 [DOI](#)). Briefly, Neurons were washed with ice cold extracellular solution (ECS) (in mM: 125 NaCl, 2 CaCl₂, 2.5 KCl, 5 HEPES, 33 glucose, pH, 8.0) and incubated for 30 min with 0.8 mg/ml NHS-SS-Biotin (21327, Thermo Fisher Scientific) in ECS buffer at 37°C. After incubation, cells were washed once with PBS and the unbound biotin was quenched via two 8 min incubations with quenching buffer (20 mM glycine in PBS). Lysis was performed by mechanical scraping in ice-cold lysis buffer (50 mM Tris-Cl pH 7.4, 150 mM NaCl, 1% TritonX-100, 0.1% SDS). After rotating for 30 min at 4°C, lysates were centrifuged for 10 min at 14,000× *g* at 4°C. Balanced Pierce Streptavidin Agarose beads (25 µL, 20353, Thermo Fisher Scientific) were added to the supernatant. The mixture of lysates and beads were rotated at 4°C overnight and then washed twice with ice-cold lysis buffer. Beads were incubated with 2× SDS sample buffer for 10 min at 37°C and biotinylated proteins were analyzed by immunoblotting.

Protein expression and purification

GST-tagged EndoA1 or gephyrin fragments was expressed in *E. coli* BL21 (DE3). Cells were grown at 37°C in LB (g/L: tryptone 10, yeast extract 5, NaCl 10) supplemented with ampicillin. Cells were induced at OD600 of ~ 0.6 with 0.4 mM isopropyl-D-1-thiogalactopyranoside (IPTG, A600168-0005,

Bio Basic Inc) for 4 h at 30°C or 16 h at 16°C. Cell pellets were resuspended in PBS supplemented with 0.2% Triton X-100 and 0.1 mM PMSF. Proteins were purified with Glutathione Sepharose 4B (GE17-0756-01, Sigma-Aldrich, St. Louis, MO, USA) according to the manufacturer's instructions.

GST-pull down, co-immunoprecipitation (IP) and immunoisolation

For GST-pull down assays, GST-fused protein EndoA1 and its fragments were expressed and purified from *E. coli*. Myc-tagged GPN constructs were expressed in HEK293 cells for 48 h and lysates were prepared in a buffer (in mM: Tris-HCl 50, pH 7.4, NaCl 150, 1% NP-40, EDTA 2) plus protease inhibitors cocktail. Cell lysates were centrifuged at 12,000× *g* for 15 min at 4°C and the supernatant was incubated with individual GST-fused proteins conjugated with Glutathione-Sepharose beads for overnight at 4°C. Beads were washed five times with lysis buffer and boiled in 2× SDS sample buffer. Bound proteins were analyzed by immunoblotting.

For co-IP from brain lysates, the whole brain from 8-10-week-old WT and KO littermates were dissected and homogenized in the pre-chilled lysis buffer (in mM: Tris-HCl 20, pH 7.4, NaCl 150, 1% NP-40, EDTA 5) plus protease inhibitor cocktail, followed by incubation on ice for 15 min. The lysates were then centrifuged at 12,000× *g* for 15 min and the supernatant was collected and incubated with anti-EndoA1 antibody (2 µg) or control IgG overnight at 4°C, followed by incubation with 20 µL Dynabeads Protein A (ThermoFisher, 10007D for rabbit antibody) for 2 h at 4°C. After washing the beads with the same lysis buffer four times, the bound proteins were incubated 2× SDS sample buffer at 37°C for 10 min and subjected to SDS-PAGE and Western blot with indicated antibodies.

For co-IP of proteins in HEK293T cells (Cat. No. 632180, Takara), cells were lysed with lysis buffer (in mM: 0.5% NP-40, Tris-HCl 50, pH 7.4, NaCl 150, EDTA 5) supplemented with protease inhibitors. Lysates were then centrifuged at 16,000× *g* for 15 min at 4°C. Cell lysates were incubated with anti-Flag Affinity Gel (A2220, Sigma-Aldrich) overnight at 4°C on a roller mixer. Immunoprecipitates were washed four times in lysis buffer and boiled in 2× SDS sample buffer, then subjected to SDS-PAGE and immunoblotting.

For immunoisolation of membrane proteins, hippocampi were homogenized with lysis buffer (in mM: Tris-HCl 20, pH 7.4, HEPES 10, pH 7.4, NaCl 150, sucrose 250, EDTA 0.5, pH 8.0) supplemented with protease inhibitors and centrifuged at 800× *g* for 15 min. The supernatants were collected and subjected to high-speed centrifugation at 100,000× *g* for 1 h (TLS-55 rotor, OptimaTMMAX Ultracentrifuge; Beckman Coulter, USA). The supernatants (S100) and pellets (P100, the membrane fraction) resuspended in lysis buffer were subjected to immunoisolation with Dynabeads Protein A (100-4D, Invitrogen, Carlsbad, CA, USA) coupled with 2 µg of rabbit anti-EndoA1 or rabbit anti-GABA_A α1 antibody. Bound proteins were eluted by incubation with 2× SDS loading buffer at 37°C for 10 min and subjected to SDS-PAGE and immunoblotting.

PSD fractionation of brain lysates

PSD fractions from adult mouse brain were prepared as previously described (Yang et al., 2015 [\[15\]](#)). In brief, hippocampi were homogenized on ice with 20 strokes of a Teflon-glass homogenizer in 1 mL of HEPES-buffered sucrose (0.32 M sucrose, 4 mM HEPES, pH 7.4) containing freshly added protease inhibitors, then centrifuged at 800× *g* at 4°C to remove the pelleted nuclear fraction (P1). Supernatant (S1) was centrifuged at 10,000× *g* for 15 min to yield the crude synaptosomal pellet (P2) and the pellet was washed once in 1 mL HEPES-buffered sucrose. P2 was lysed by hypoosmotic shock in 900 µL ice-cold 4 mM HEPES, pH 7.4 plus protease inhibitors, homogenized by pipetting and rotating for 30 min at 4 °C. The lysate was centrifuged at 25,000× *g* for 20 min to yield supernatant (S3, crude synaptic vesicle fraction) and pellet (P3, lysed synaptosomal membrane fraction). To prepare the PSD fraction, P3 was resuspended in 900 µL of ice-cold 50 mM HEPES, pH 7.4, 2 mM EDTA, plus protease inhibitors and 0.5% Triton X-100, rotated for 15 min at 4

°C and centrifuged at 32,000× *g* for 20 min to obtain the PSD pellet. PSD pellets were resuspended in 50 µL ice-cold 50 mM HEPES, pH 7.4, 2 mM EDTA plus protease inhibitors. Proteins were subjected to SDS-PAGE and immunoblotting.

Protein identification with mass spectrometry and pathway analysis

For identification of EndoA1-interacting proteins, mouse brain was homogenized with lysis buffer (in mM: Tris-HCl 20, pH 7.4, HEPES 10, pH 7.4, NaCl 150, EDTA 1, 1% NP-40) supplemented with protease inhibitors. Lysates were centrifuged at 12,000× *g* for 15 min. The supernatants were incubated with rabbit anti-EndoA1 antibody overnight at 4°C. Beads were rinsed with lysis buffer for five times and bound proteins were resolved by SDS-PAGE. Proteins in the gels were reduced with 10 mM DTT at 37°C for 1 h, then alkylated with 25 mM iodoacetamide at RT for 1 h in the dark before digestion with trypsin (T1426, Sigma-Aldrich; enzyme-to-substrate ratio 1:50) in 25 mM ammonium bicarbonate at 37°C overnight. Tryptic peptides were extracted from gel by sonication with a buffer containing 5% trifluoroacetic acid and 50% acetonitrile. The liquid was dried by SpeedVac, and peptides were resolubilized in 0.1% formic acid and filtered with 0.45 µm centrifugal filters before analysis with a TripleTOF 5600 mass spectrometer (AB SCIEX, Canada) coupled to an Eksigent nanoLC. Proteins were identified by searching the MS/MS spectra against the *Mus musculus* SwissProt database using the ProteinPilot 4.2 software. Carbamidomethylation of cysteine was set as the fixed modification. Trypsin was specified as the proteolytic enzyme with a maximum of 2 miss cleavages. Mass tolerance was set to 0.05 Da and the false discovery rates for both proteins and peptides were set at 1%. For enrichment pathway analysis: Only proteins with intensity to IgG ratio of > 1.5 were considered in the analysis. These genes were analyzed by Gene Ontology (GO) Enrichment Analysis (<http://bioinfo.org/kobas> ).

Electrophysiology

To knockout *EndoA1* in the brain, *EndoA1*^{fl/fl} mice were injected with high-titer AAV stock carrying pAOV-CaMKIIα-mCherry-2A-Cre (AAV-mCherry-2A-Cre) ($1\sim5 \times 10^{13}$ IU/ml) in the hippocampal CA1 region within 24 h after birth. Newborn *EndoA1*^{fl/fl} littermates (male or female) were anesthetized on ice for 4-5 min and then mounted in a custom ceramic mold to align the head along the X and Y axes, with lambda set as (X, Y) = (0, 0). The zero point of the Z axis was the position at which the injecting needle penetrated the skin. Approximately 10 nL of the viral solution was injected at each of the seven sites (coordinates in mm, [X, Y, Z] = [1.2, 1.2, 1.4/1.0/0.6] and [1.5, 1.0, 1.7/1.3/0.9/0.5]) targeting the hippocampus in each cerebral hemisphere using a microsyringe (Sutter Instrument, Novato, CA, USA) and beveled glass injection pipette (Yang et al., 2018 ). For the rescue experiment, P0 *EndoA1*^{fl/fl} mice were injected with a mixture of two high-titer AAV stocks: AAV-EGFP and AAV-mCherry or AAV-mCherry-2A-Cre; AAV-mCherry-2A-Cre and AAV-EGFP-2A-FLAG-EndoA1, AAV-EGFP-2A-FLAG-EndoA1 Y343A or AAV-EGFP-2A-FLAG-EndoA1 KKK-EEE. Approximately 20 nL of the viral solution was injected at each of the seven sites. The injected pups were returned to the home cage and used for recording 2 weeks later.

To prepare acute brain slices for electrophysiological analyses, mice were anesthetized with tribromoethanol (330-390 mg/kg) (T48402, Sigma-Aldrich, St. Louis, MO, USA) and decapitated. Whole brains were removed promptly and placed in ice-cold oxygenated (95% O₂/5% CO₂) low-Ca²⁺/high-Mg²⁺ high-sucrose cutting solution (in mM: KCl 2.5, NaHCO₃ 25, NaH₂PO₄ 1.25, glucose 10, sucrose 210, Na-Ascorbate 1.3, CaCl₂ 0.5, MgSO₄ 7). Coronal 300 µm slices were obtained using a vibrating tissue slicer (Leica VT1200S; Leica Biosystems, Wetzlar, Deutschland). Slices were immediately transferred to standard artificial cerebrospinal fluid (ACSF) (in mM: NaCl 119, KCl 2.5, MgSO₄ 1.3, CaCl₂ 2.5, glucose 11, NaHCO₃ 26.2, NaH₂PO₄ 1, pH 7.3-7.4) at 33°C and continuously bubbled with 95% O₂/5% CO₂. After 30 min of incubation, slices were transferred to a recording chamber with the same extracellular buffer at room temperature (RT, ~25°C).

For whole-cell patch clamp recordings, brain slices were continuously perfused with standard ACSF at RT under visual control with differential interference contrast illumination on an upright microscope (BX51WI, Olympus, Tokyo, Japan) with a 40× water-immersion objective and a digital camera. The mCherry- and/or EGFP-positive hippocampal CA1 pyramidal neurons were identified by an epifluorescence microscope (pE-300, CoolLED Ltd, Andover, UK). Whole-cell recordings were performed on an AAV-infected neuron and a neighboring uninfected neuron. Patch clamp data were obtained using a Multiclamp 700B amplifier (Molecular Devices, San Jose, CA, USA), and signals were low-pass filtered at 2 kHz and digitized at 10 kHz using an Axon Digidata 1550B analog-to-digital board (Molecular Devices, San Jose, CA, USA).

To analyze evoked excitatory postsynaptic currents (eEPSCs), evoked inhibitory postsynaptic currents (eIPSCs), and paired-pulse ratio, cells were recorded with 3–7 MΩ borosilicate glass patch pipettes filled with the intracellular solution (in mM: CsMeSO₃ 135, HEPES 10, EGTA 1, QX-314 3.3, MgATP 4, NaGTP 0.3, and Na₂-phosphocreatine 8, pH 7.3) (Horn and Nicoll, 2018), following stimulation of Schaffer collaterals with a bipolar stimulation electrode placed in the stratum radiatum of the CA1 region. EPSCs were measured at a holding potential of -70 mV, and IPSCs were recorded at +10 mV. Paired-pulse ratios of EPSCs were measured at a holding potential of -70 mV by applying two electrical stimulations at a 50 ms interval and calculating the ratio of the peak amplitudes. Paired-pulse ratios of IPSCs were measured at a holding potential of +10 mV by applying two electrical stimulations at a 200 ms interval and calculating the ratio of the peak amplitudes. E/I ratios were measured by applying the same stimulation current and then calculating the ratio of the absolute values of the peak amplitudes of EPSCs and IPSCs. The miniature EPSCs and IPSCs were measured in the ACSF bath solution containing 1 μM tetrodotoxin (TTX, MSS0002, BGB Analytik AG, BL, Switzerland) at -70 mV and +10 mV, respectively, in the voltage-clamp mode. Data were collected and analyzed using the pClamp 10.6 software (Molecular Devices, San Jose, CA, USA).

All electrophysiological recordings were carried out blindly and all n numbers in the data figures indicate the number of recorded neurons. All chemicals were purchased from Sigma-Aldrich except otherwise noted.

Confocal and three-dimensional structured illumination microscopy (3D-SIM) image acquisition and analysis

Confocal images were collected using the Spectral Imaging Confocal Microscope Digital Eclipse A1Si (Nikon, Tokyo, Japan) with a 100× Plan Apochromat VC (NA 1.40) oil objective (4× optical zoom). Images were z projections of images taken at 0.25-μm step intervals. The number of planes, typically 5–7, was chosen to cover the entire dendrite from top to bottom. Images were captured using a 1024×1024 pixels screen for neuronal cells. Laser power, digital gain, and offset settings were identical in each experiment. Images were processed with the NIH ImageJ software and exported into Adobe Photoshop. Only brightness and contrast were adjusted for the whole frame, and no part of a frame was enhanced or modified in any way. Control and experimental group neurons which were to be directly compared were imaged with the same acquisition parameters.

For 3D-SIM, secondary antibodies conjugated with Alexa Fluor 568 were used and images were captured as described by Niu et al (Niu et al., 2017) on the DeltaVision OMX V4 imaging system (Applied Precision Inc, USA) with a 100× 1.4 oil objective (Olympus UPlanApo), solid state multimode lasers (488, 593 and 642 nm) and electron-multiplying charge-coupled device (CCD) cameras (Evolve 512× 512, Photometrics, USA). Serial Z-stack sectioning was done at 125 nm intervals. The microscope is routinely calibrated with 100 nm fluorescent spheres to calculate both the lateral and axial limits of image resolution. SIM image stacks were reconstructed using softWoRx 5.0 (Applied Precision) with the following settings: pixel size 39.5 nm; channel-specific optical transfer functions; Wiener filter 0.001000; discard Negative Intensities background; drift correction with respect to first angle; custom K0 guess angles for camera positions. Pixel

registration was corrected to be less than 1 pixel for all channels using 100 nm Tetraspeck beads. Three dimensional images were reconstructed from z stacks. For clarity of display, small linear changes of brightness and contrast were performed on three-dimensional reconstructions.

Quantification of immunostaining were carried out blindly. For quantitative analysis of immunofluorescent signals in neurons, maximal projection images were created with the ImageJ software (<https://imagej.net>) from 6-7 serial optical sections. Background was subtracted by using the “subtract background” function and the background level was held identical for all cells within each experiment. To examine the number and fluorescence intensity of signal puncta in soma or dendrites, the EGFP- or DsRed-labeled soma or dendrites were outlined automatically by using the “Wand Tool” function. The region of interest (ROI) in cultured neurons was defined along a segment of the dendrite according to the fluorescence signal distinguished from the background. Average values of fluorescence intensities in ROI (the total fluorescent intensity divided by the total area of a dendritic segment) were calculated by Analyze > measure. Gephyrin clusters were thresholded and confirmed visually to select appropriate clusters following a minimal size cut-off by using the “Analyze Particles” plugin of the ImageJ software. For colocalization quantifications, NIS-Elements Software was used as follows: region of interest (ROI) of dendritic segments was manually chosen, and thresholded using the “Threshold” function to remove background signals. Pearson’s correlation coefficient and Manders overlap coefficient were analyzed by using the “Analyze Control > colocalization” function.

For spatial correlation/colocalization analysis in super-resolution images, we using the method described in the reference (McCall, 2024). Briefly, channels in 3D images were split and followed by Gaussian blur performed on both GPN and Bassoon channels. Next masks of all possible localizations for the signal in GPN and Bassoon image channel were obtained. Then, the input images of the three channels were deconvolved using 10 iterations of the Richardson-Lucy algorithm in Deconvolution Lab 2 (version 2.1.2), with theoretical point spread functions (PSFs) generated by the PSF Generator plugin. PSFs were generated using the Born & Wolf model, with parameters matching the acquisition parameters: 1.518 immersion refractive index, 1.4 NA, 40 nm pixel size and 125 nm z-step, 512 × 512 × 14 output diameters, and 520 nm, 592 nm and 670 nm wavelengths. Finally, select the two images (GPN and EndoA1 or Bassoon and EndoA1) and the mask images, then perform a cross-correlation analysis (CCC) between the two images using the CCC plugin (https://github.com/andmccall/Colocalization_by_Cross_Correlation).

Statistical analysis

For all biochemical, cell biological and electrophysiological recordings, at least three independent experiments were performed (independent cultures, transfections or different mice). Statistical analysis was performed in GraphPad Prism 9.0 (GraphPad Software, La Jolla, CA). Data sets were tested for normality and direct comparisons between two groups were made using two-tailed Student’s t test (t test, for normally distributed data) as indicated. To evaluate statistical significance of three or more groups of samples, one-way ANOVA analysis with a Tukey test was used or repeated measures ANOVA analysis with a Tukey test was used in behavior assays. Statistical parameters are reported in the figures and the corresponding legends. The statistical significance was defined as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, respectively. All data are presented as scatterplots with mean ± S.E.M.

Acknowledgements

We thank Drs Shiva Tyagarajan (Addgene #68816), Peter Scheiffele (Addgene # 15246) and Don Arnold (Addgene # 46295 & #46296) for constructs. This work was supported by funding from the National Natural Science Foundation of China (NSFC 32271005 and 32070785 to Y. Yang, 31921002,

32270730 and 91954126 to J-J. Liu) and the State Key Laboratory of Molecular Developmental Biology (2022-MDB-KF-13).

The authors declare no competing financial interests.

Additional information

Author contributions

J-J. Liu and Y. Yang conceived the project. Y. Yang and J-J. Liu designed and supervised research. X. Chen, D. Pan and Y. Yang performed experiments and analyzed the data. Y. Yang and J-J. Liu wrote the manuscript with help from all other authors.

Funding

National Natural Science Foundation of China (32271005)

National Natural Science Foundation of China (32070785)

National Natural Science Foundation of China (31921002)

National Natural Science Foundation of China (32270730)

National Natural Science Foundation of China (91954126)

State Key Laboratory of Molecular Developmental Biology

Additional files

Figure S1. Loss of endophilin A1 in vivo does not affect fast mIPSCs (perisomatic events: < 2.8 ms) and slow mIPSCs (distal dendritic events: > 2.8 ms). Related to Figure 1. **(A)** Quantification of somatic and dendritic amplitudes in Ctrl and EndoA1 KO neuron. $n = 7$ Ctrl and 9 EndoA1 KO neurons from 3 independent mice. Data are scatterplots with mean $\pm$ S.E.M. **(B)** Frequency of somatic and dendritic events from Ctrl and EndoA1 KO mice. Ctrl: 7 neurons; EndoA1 KO: 9 neurons from 3 independent mice. Data are scatterplots with mean $\pm$ S.E.M. [↗](#)

Figure S2. Effects of EndoA1 knockout on excitatory and inhibitory synapses in the hippocampus. Related to Figure 2. **(A-B)** Immunofluorescence analysis of PSD95 clusters in the hippocampal region of 3-week-old (P21) (left) and 10-week-old (P60) (right) EndoA1^{+/+} and EndoA1^{-/-} mouse brains. Scale bar, 50 μ m. **(A)** **(C)** Quantification of the mean intensity of PSD95 in the hippocampal region of 3-week-old (P21) and 10-week-old (P60) EndoA1^{+/+} and EndoA1^{-/-} mouse brains. $n = 6$ mice for each genotype. Statistical test: unpaired two-tailed Student's t-test. Data are scatterplots with mean $\pm$ S.E.M. **(D-E)** Immunofluorescent signals of VGLUT1 in the hippocampal region of 3-week-old (P21) (left) and 10-week-old (P60) (right) EndoA1^{+/+} and EndoA1^{-/-} mouse brains. Scale bar, 50 μ m. **(A)** **(F)** Quantification of the mean intensity of VGLUT1 in the hippocampal region of 3-week-old (P21) and 10-week-old (P60) EndoA1^{+/+} and EndoA1^{-/-} mouse brains. $n = 6$ mice for each genotype. Statistical test: unpaired two-tailed Student's t-test. Data are scatterplots with mean $\pm$ S.E.M. **(B)** **(G)** Immunofluorescent signals of GPN in distinct layers of P60 mouse hippocampus. Scale bar, 50 μ m. **(C)** **(H)** Quantification of gephyrin (GPN) mean intensity in distinct layers of hippocampus. $n = 6$ mice for each genotype. Abbreviations: CA1-3, cornu Ammonis region 1-3; DG, dentate gyrus; slm, stratum lacunosum-moleculare; so, stratum oriens; sp, stratum pyramidale; sr, stratum radiatum. DAPI staining of cell nuclei is shown in blue. Statistical test: two-tailed Student's t-test. ns, non-significant. Data are scatterplots with mean $\pm$ S.E.M. [↗](#)

Figure S3. Changes of endogenous postsynaptic proteins visualized by recombinant probes after depletion of EndoA1. Related to Figure 4. **(A)** AAVs expressing GFP (AAV-GFP) or GFP and Cre (AAV-GFP-2A-Cre) and mRuby2-GPN.FingR were stereotactically injected into the CA1 regions of EndoA1^{fl/fl} mouse brain at P0. Brain sections from P21 mice were fixed and immunostained for GFP and mRuby. Representative images at 40× magnification with 4× digital zoom showed the expression patterns of mRuby2-GPN.FingR. Scale bars, 10 μm. **(B)** Quantification of signal intensity of mRuby2-GPN.FingR in (C). n = 8 mice for each genotype. **(C)** AAVs expressing mCherry (AAV-mCherry) or mCherry and Cre (AAV-mCherry-2A-Cre) and PSD95.FingR-eGFP were stereotactically injected into the CA1 regions of EndoA1^{fl/fl} mouse brain at P0. Brain sections from P21 mice were fixed and immunostained for mCherry and GFP. Representative images at 40× magnification with 4× digital zoom showed the expression patterns of PSD95.FingR-eGFP. Scale bars, 10 μm. **(D)** Quantification of signal intensity of PSD95.FingR-eGFP in (A). n = 6 mice for each genotype. 

Figure S4. Expression of GABA_A α1 in cultured hippocampal neurons and the localization of EndoA1 in inhibitory pre- and postsynaptic sites. Related to Figure 3. **(A)** Cultured hippocampal neurons were fixed on DIV16 and immunostained for surface GABA_A α1. Shown are representative confocal microscopy images. Scale bar, 10 μm. **(B)** Hippocampal neurons were fixed and immunostained with antibodies against EndoA1, Bassoon and GPN on DIV16. Shown are representative z stack maximum projection images captured by structured illumination microscopy (SIM). Lower panels are insets of boxed regions in the upper panel. Lower panels are insets of boxed regions in the upper panel. Scale bars, 5 μm in upper panels and 0.5 μm in magnified images. **(C-D)** Color composite of the two input images (left) and composite image of the data that contributed to the Gaussian result (right) for spatial colocalization analysis in super-resolutive images. Scale bars, 5 μm. **(E-F)** The blue line is the radial profile of original cross-correlation, the green line is the radial profile after the pixel randomization procedure, and the red dashed line is the fit Gaussian curve. CCC generates four statistical parameters (μ (the mean distance of the measured spatial correlation), σ (standard deviation distance for the spatial correlation), Confidence and R-squared) that was used to evaluate spatial correlation of GPN and EndoA1 in (E), and of Bassoon and EndoA1 in (F). **(G)** The ratio of GPN or Bassoon cross-correlated with EndoA1 to total GPN or Bassoon, respectively. n = 14, from 3 independent cultures. All data are scatterplots with means ± S.E.M. 

Figure S5. Effect of EndoA1 knockout on gephyrin clustering in cultured hippocampal neurons. Related to Figure 4. **(A)** EndoA1^{fl/fl} hippocampal neurons were co-transfected with constructs expressing mCherry and GPN.FingR-eGFP or mCherry-2A-Cre and GPN.FingR-eGFP on DIV12, then fixed and immunostained for eGFP and mCherry on DIV16. Shown are representative confocal microscopy images of GPN.FingR-eGFP. The right panels are magnification of soma or boxed regions in the left panels. Scale bar, 10 μm in the left panels and 5 μm in the right panels. **(B)** Quantification of GPN.FingR-eGFP mean intensity in the soma (left), mean intensity (center) and cluster number (right) in dendrites. Scale bar, 5 μm. 20 neurons for mCherry + GPN.FingR-eGFP; 20 neurons for Cre + GPN.FingR-eGFP. Statistical test: unpaired two-tailed Student's t-test. All data are scatterplots with means ± S.E.M. **(C)** Same as (B) except that neuron were co-transfected with PSD.FingR-eGFP. **(D)** Quantification of PSD95.FingR-eGFP mean intensity in the soma (left), mean intensity (center) and cluster number (right) in dendrites. Scale bar, 5 μm. 14 neurons for mCherry + PSD95.FingR-eGFP; 13 neurons for Cre + PSD95.FingR-eGFP. Statistical test: unpaired two-tailed Student's t-test. All data are scatterplots with means ± S.E.M. **(E-G)** EndoA1^{fl/fl} hippocampal neurons were transfected with construct expressing mCherry or mCherry-2A-Cre on DIV12 and immunostained for GPN and mCherry on DIV16. Shown are representative confocal microscopy images for GPN in the soma, axon initial segment (AIS) and dendrites 30 μm, 60 μm, 90 μm or 120 μm from the soma, respectively. Scale bars, 5 μm. **(H)** Quantification of GPN mean intensity normalized to the mCherry group in the soma (A), AIS (B) and dendrites at various distances from the soma (C). Number: 40 (mCherry), 47 (Cre) for soma; 26 (mCherry), 25 (Cre) for AIS; 55 (mCherry), 52 (Cre) for dendrites 30 μm; 52 (mCherry), 50 (Cre) for dendrites 60 μm; 48 (mCherry), 47 (Cre) for dendrites 90 μm; 49 (mCherry), 41 (Cre) for dendrites 120 μm. Ankyrin-G: Axon initial segment (AIS) labeling marker. Statistical test: unpaired two-tailed Student's t-test. All data are scatterplots with means ± S.E.M for each group. 

Figure S6. p140Cap promotes gephyrin clustering via endophilin A1-mediated pathway. Related to Figure 6 and 7. **(A)** Cultured hippocampal neurons transiently expressing GFP and LifeAct-mCherry were immunostained for GPN, GFP and mCherry on DIV16. Shown are representative z stack maximum projection images captured by structured illumination microscopy (SIM). Right panels are insets of boxed regions in the left panel. Scale bars, 2 μm in left panels and 0.5 μm in magnified images. **(B)** Cultured hippocampal neurons were co-transfected with Ctrl shRNA and vector or p140Cap shRNA and vector or RNAi-resistant Flag-tagged p140Cap on DIV 11, followed by immunostaining with antibodies to GPN, Flag (blue) and RFP on DIV 16. Scale bar, 5 μm . **(C)** Quantification of GPN mean intensity and puncta number in (B). 18 neurons for shCtrl + vector; 20 neurons for shp140Cap + vector; 22 neurons for shp140Cap + p140Cap, from 3 independent cultures. Statistical test: One-way ANOVA followed by Tukey post-hoc test. All data shown are scatterplots with means $\pm$ S.E.M. **(D)** EndoA1^{f/f} hippocampal neurons coexpressing GFP and Ctrl shRNA or p140Cap shRNA, or GFP-2A-Cre and Ctrl shRNA or p140Cap shRNA were immunostained for GPN on DIV16. Shown are representative confocal microscopy images. Scale bar, 5 μm . **(E)** Quantification of GPN mean intensity and puncta number in (D). 27 neurons for GFP + shCtrl; 26 neurons for GFP + shp140Cap; 23 neurons for Cre + shCtrl; 26 neurons for Cre + shp140Cap, from 3 independent cultures. Statistical test: One-way ANOVA followed by Tukey post-hoc test. All data shown are scatterplots with means $\pm$ S.E.M. [↗](#)

Table S1. Proteins identified by mass spec analysis of endophilin A1 co-immunoprecipitates and their abundance. Related to Figure 4 [↗](#)

Video S1. PTZ kindling behaviors of EndoA1^{+/+} and EndoA1^{-/-} mice. Related to Figure 1 [↗](#)

Video S2. The spatial relationship between EndoA1 and GPN. Related to Figure 3G. 3D-SIM movie of an enlarged region of interest (ROI) from a hippocampal neuron dendrite shows the association of EndoA1 (red) and GPN (green). [↗](#)

Video S3. The spatial relationship between endophilin A1 and surface GABA_A γ 2. Related to Figure 3H. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron dendrite shows the association of EndoA1 (red) and surface GABA_A γ 2 (green). [↗](#)

Video S4. The spatial relationship between endophilin A1 and VGAT. Related to Figure 3I. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron dendrite shows the signals of EndoA1 (red) and VGAT (green). [↗](#)

Video S5. The spatial relationship between Bassoon, gephyrin and endophilin A1. Related to Figure S3B. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron dendrite shows the signals of Bassoon (blue), GPN (green) and EndoA1 (red). [↗](#)

Video S6. The spatial relationship between gephyrin and LifeAct. Related to Figure 6 and Figure S5A. 3D-SIM movie of an enlarged region of interest from a hippocampal neuron dendrite shows the signals of LifeAct (red) and GPN (green). [↗](#)

Video S7. PTZ kindling behaviors of EndoA1^{f/f} mice injected with AAVs expressing Cre and EndoA1 WT or mutants. Related to Figure 7 [↗](#)

References

Allison D.W., Chervin A.S., Gelfand V.I., Craig A.M (2000) **Postsynaptic scaffolds of excitatory and inhibitory synapses in hippocampal neurons: maintenance of core components independent of actin filaments and microtubules** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **20**:4545–4554 [Google Scholar](#)

Bademosi A.T., Decet M., Kuenen S., Calatayud C., Swerts J., Gallego S.F., Schoovaerts N., Karamanou S., Louros N., Martin E., et al. (2023) **EndophilinA-dependent coupling between activity-induced calcium influx and synaptic autophagy is disrupted by a Parkinson-risk mutation** *Neuron* [Google Scholar](#)

Bai G., Wang Y., Zhang M (2021) **Gephyrin-mediated formation of inhibitory postsynaptic density sheet via phase separation** *Cell research* **31**:312–325 [Google Scholar](#)

Bensussen S., Shankar S., Ching K.H., Zemel D., Ta T.L., Mount R.A., Shroff S.N., Gritton H.J., Fabris P., Vanbenschoten H., et al. (2020) **A Viral Toolbox of Genetically Encoded Fluorescent Synaptic Tags** *iScience* **23**:101330 [Google Scholar](#)

Bogdanov Y., Michels G., Armstrong-Gold C., Haydon P.G., Lindstrom J., Pangalos M., Moss S.J (2006) **Synaptic GABAA receptors are directly recruited from their extrasynaptic counterparts** *The EMBO journal* **25**:4381–4389 [Google Scholar](#)

Canitano R., Pallagrosi M (2017) **Autism Spectrum Disorders and Schizophrenia Spectrum Disorders: Excitation/Inhibition Imbalance and Developmental Trajectories** *Frontiers in psychiatry* **8**:69 [Google Scholar](#)

Canitano R., Palumbi R (2021) **Excitation/Inhibition Modulators in Autism Spectrum Disorder: Current Clinical Research** *Frontiers in neuroscience* **15**:753274 [Google Scholar](#)

Chang D., Nalls M.A., Hallgrímsdóttir I.B., Hunkapiller J., van der Brug M., Cai F., Kerchner G.A., Ayalaon G., Bingol B., Sheng M., et al. (2017) **A meta-analysis of genome-wide association studies identifies 17 new Parkinson's disease risk loci** *Nature genetics* **49**:1511–1516 [Google Scholar](#)

Charrier C., Ehrensperger M.V., Dahan M., Lévi S., Triller A (2006) **Cytoskeleton regulation of glycine receptor number at synapses and diffusion in the plasma membrane** *J Neurosci* **26**:8502–8511 [Google Scholar](#)

Chen J.L., Villa K.L., Cha J.W., So P.T., Kubota Y., Nedivi E (2012) **Clustered dynamics of inhibitory synapses and dendritic spines in the adult neocortex** *Neuron* **74**:361–373 [Google Scholar](#)

Chih B., Gollan L., Scheiffele P (2006) **Alternative splicing controls selective trans-synaptic interactions of the neuroligin-neurexin complex** *Neuron* **51**:171–178 [Google Scholar](#)

Corponi F., Bonassi S., Vieta E., Albani D., Frustaci A., Ducci G., Landi S., Boccia S., Serretti A., Fabbri C (2019) **Genetic basis of psychopathological dimensions shared between schizophrenia and bipolar disorder** *Prog Neuropsychopharmacol Biol Psychiatry* **89**:23–29 [Google Scholar](#)

Cramer T.M.L., Pinan-Lucarre B., Cavaccini A., Damilou A., Tsai Y.C., Bhat M.A., Panzanelli P., Rama N., Mehlen P., Benke D., et al. (2023) **Adamtsl3 mediates DCC signaling to selectively promote GABAergic synapse function** *Cell reports* **42**:112947 [Google Scholar](#)

Dhir A (2012) **Pentylentetrazol (PTZ) kindling model of epilepsy** *Curr Protoc Neurosci* **9**:Unit9.37 [Google Scholar](#)

Essrich C., Lorez M., Benson J.A., Fritschy J.M., Luscher B (1998) **Postsynaptic clustering of major GABAA receptor subtypes requires the gamma 2 subunit and gephyrin** *Nature neuroscience* **1**:563–571 [Google Scholar](#)

Favuzzi E., Deogracias R., Marques-Smith A., Maeso P., Jezequel J., Exposito-Alonso D., Balia M., Kroon T., Hinojosa A.J., et al. (2019) **Distinct molecular programs regulate synapse specificity in cortical inhibitory circuits** *Science* **363**:413–417 [Google Scholar](#)

Foss-Feig J.H., Adkinson B.D., Ji J.L., Yang G., Srihari V.H., McPartland J.C., Krystal J.H., Murray J.D., Anticevic A (2017) **Searching for Cross-Diagnostic Convergence: Neural Mechanisms Governing Excitation and Inhibition Balance in Schizophrenia and Autism Spectrum Disorders** *Biol Psychiatry* **81**:848–861 [Google Scholar](#)

Fritschy J.M (2008) **Epilepsy, E/I Balance and GABA(A) Receptor Plasticity** *Frontiers in molecular neuroscience* **1**:5 [Google Scholar](#)

Gad H., Ringstad N., Low P., Kjaerulff O., Gustafsson J., Wenk M., Di Paolo G., Nemoto Y., Crun J., Ellisman M.H., et al. (2000) **Fission and uncoating of synaptic clathrin-coated vesicles are perturbed by disruption of interactions with the SH3 domain of endophilin** *Neuron* **27**:301–312 [Google Scholar](#)

George S., Chiou T.T., Kanamalla K., De Blas A.L. (2021) **Recruitment of Plasma Membrane GABA-A Receptors by Submembranous Gephyrin/Collybistin Clusters** *Cellular and molecular neurobiology* [Google Scholar](#)

Germer E.L., Imhoff S., Vilariño-Güell C., Kasten M., Seibler P., Brüggemann N., Klein C., Trinh J (2019) **The Role of Rare Coding Variants in Parkinson’s Disease GWAS Loci** *Frontiers in neurology* **10**:1284 [Google Scholar](#)

Gross G.G., Junge J.A., Mora R.J., Kwon H.B., Olson C.A., Takahashi T.T., Liman E.R., Ellis-Davies G.C., McGee A.W., Sabatini B.L., et al. (2013) **Recombinant probes for visualizing endogenous synaptic proteins in living neurons** *Neuron* **78**:971–985 [Google Scholar](#)

Gu J., Firestein B.L., Zheng J.Q (2008) **Microtubules in dendritic spine development** *J Neurosci* **28**:12120–12124 [Google Scholar](#)

Guo Z., Jiang C.H., Tong C., Yang Y., Wang Z., Lam S.M., Wang D., Li R., Shui G., Shi Y.S., et al. (2022) **Activity-dependent PI4P synthesis by PI4KIIIα regulates long-term synaptic potentiation** *Cell Rep* **38**:110452 [Google Scholar](#)

Han W., Li J., Pelkey K.A., Pandey S., Chen X., Wang Y.X., Wu K., Ge L., Li T., Castellano D., et al. (2019) **Shisa7 is a GABA(A) receptor auxiliary subunit controlling benzodiazepine actions** *Science* **366**:246–250 [Google Scholar](#)

Hoffmann C., Milovanovic D (2021) **Gephyrin: a scaffold that builds a phase at the inhibitory postsynapses** *Cell research* **31**:245–246 [Google Scholar](#)

Horn M.E., Nicoll R.A (2018) **Somatostatin and parvalbumin inhibitory synapses onto hippocampal pyramidal neurons are regulated by distinct mechanisms** *Proceedings of the National Academy of Sciences* **115**:589–594 [Google Scholar](#)

Huo J.Z., Cortez M.A., Snead O.C. (2009) **GABA receptor proteins within lipid rafts in the AY-9944 model of atypical absence seizures** *Epilepsia* **50**:776–788 [Google Scholar](#)

Jacob T.C., Bogdanov Y.D., Magnus C., Saliba R.S., Kittler J.T., Haydon P.G., Moss S.J (2005) **Gephyrin regulates the cell surface dynamics of synaptic GABAA receptors** *J Neurosci* **25**:10469–10478 [Google Scholar](#)

Jaworski J., Kapitein L.C., Gouveia S.M., Dortland B.R., Wulf P.S., Grigoriev I., Camera P., Spangler S.A., Di Stefano P., Demmers J., et al. (2009) **Dynamic microtubules regulate dendritic spine morphology and synaptic plasticity** *Neuron* **61**:85–100 [Google Scholar](#)

Kalscheuer V.M., Musante L., Fang C., Hoffmann K., Fuchs C., Carta E., Deas E., Venkateswarlu K., Menzel C., Ullmann R., et al. (2009) **A balanced chromosomal translocation disrupting ARHGEF9 is associated with epilepsy, anxiety, aggression, and mental retardation** *Human mutation* **30**:61–68 [Google Scholar](#)

Kasaragod V.B., Mortensen M., Hardwick S.W., Wahid A.A., Dorovych V., Chirgadze D.Y., Smart T.G., Miller P.S (2022) **Mechanisms of inhibition and activation of extrasynaptic alphabeta GABA(A) receptors** *Nature* **602**:529–533 [Google Scholar](#)

Kasaragod V.B., Schindelin H (2019) **Structure of Heteropentameric GABAA Receptors and Receptor-Anchoring Properties of Gephyrin** *Frontiers in molecular neuroscience* **12**:191 [Google Scholar](#)

Kins S., Betz H., Kirsch J (2000) **Collybistin, a newly identified brain-specific GEF, induces submembrane clustering of gephyrin** *Nat Neurosci* **3**:22–29 [Google Scholar](#)

Kirsch J., Betz H (1995) **The postsynaptic localization of the glycine receptor-associated protein gephyrin is regulated by the cytoskeleton** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **15**:4148–4156 [Google Scholar](#)

Kirsch J., Langosch D., Prior P., Littauer U.Z., Schmitt B., Betz H (1991) **The 93-kDa glycine receptor-associated protein binds to tubulin** *The Journal of biological chemistry* **266**:22242–22245 [Google Scholar](#)

Klausberger T., Somogyi P (2008) **Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations** *Science* **321**:53–57 [Google Scholar](#)

Kneussel M., Brandstätter J.H., Laube B., Stahl S., Müller U., Betz H (1999) **Loss of Postsynaptic GABAAR Clustering in Gephyrin-Deficient Mice** *J Neurosci* **19**:9289–9297 [Google Scholar](#)

Koh W., Kwak H., Cheong E., Lee C.J. (2023) **GABA tone regulation and its cognitive functions in the brain** *Nature reviews Neuroscience* **24**:523–539 [Google Scholar](#)

Liu Y.T., Tao C.L., Zhang X., Xia W., Shi D.Q., Qi L., Xu C., Sun R., Li X.W., Lau P.M., et al. (2020) **Mesophasic organization of GABAA receptors in hippocampal inhibitory synapses** *Nat*

Neurosci **23**:1589–1596 [Google Scholar](#)

Loebrich S., Bähring R., Katsuno T., Tsukita S., Kneussel M (2006) **Activated radixin is essential for GABAA receptor alpha5 subunit anchoring at the actin cytoskeleton** *The EMBO journal* **25**:987–999 [Google Scholar](#)

McCall A.D (2024) **Colocalization by cross-correlation, a new method of colocalization suited for super-resolution microscopy** *BMC bioinformatics* **25**:55 [Google Scholar](#)

Miles R., Tóth K., Gulyás A.I., Hájos N., Freund T.F (1996) **Differences between somatic and dendritic inhibition in the hippocampus** *Neuron* **16**:815–823 [Google Scholar](#)

Milosevic I., Giovedi S., Lou X., Raimondi A., Collesi C., Shen H., Paradise S., O'Toole E., Ferguson S., Cremona O., et al. (2011) **Recruitment of endophilin to clathrin-coated pit necks is required for efficient vesicle uncoating after fission** *Neuron* **72**:587–601 [Google Scholar](#)

Mody I., Pearce R.A (2004) **Diversity of inhibitory neurotransmission through GABA(A) receptors** *Trends in neurosciences* **27**:569–575 [Google Scholar](#)

Mukherjee J., Kretschmannova K., Gouzer G., Maric H.M., Ramsden S., Tretter V., Harvey K., Davies P.A., Triller A., Schindelin H., et al. (2011) **The residence time of GABA(A)Rs at inhibitory synapses is determined by direct binding of the receptor $\alpha 1$ subunit to gephyrin** *J Neurosci* **31**:14677–14687 [Google Scholar](#)

Murdoch J.D., Rostovsky C.M., Gowrisankaran S., Arora A.S., Soukup S.F., Vidal R., Capece V., Freytag S., Fischer A., Verstreken P., et al. (2016) **Endophilin-A Deficiency Induces the Foxo3a-Fbxo32 Network in the Brain and Causes Dysregulation of Autophagy and the Ubiquitin-Proteasome System** *Cell reports* **17**:1071–1086 [Google Scholar](#)

Niu Y., Dai Z., Liu W., Zhang C., Yang Y., Guo Z., Li X., Xu C., Huang X., Wang Y., et al. (2017) **Ablation of SNX6 leads to defects in synaptic function of CA1 pyramidal neurons and spatial memory** *eLife* **6** [Google Scholar](#)

Nusser Z., Hajos N., Somogyi P., Mody I (1998) **Increased number of synaptic GABA(A) receptors underlies potentiation at hippocampal inhibitory synapses** *Nature* **395**:172–177 [Google Scholar](#)

Parato J., Bartolini F (2021) **The microtubule cytoskeleton at the synapse** *Neuroscience letters* **753**:135850 [Google Scholar](#)

Poulopoulos A., Aramuni G., Meyer G., Soykan T., Hoon M., Papadopoulos T., Zhang M., Paarman I., Fuchs C., Harvey K., et al. (2009) **Neuroigin 2 drives postsynaptic assembly at perisomatic inhibitory synapses through gephyrin and collybistin** *Neuron* **63**:628–642 [Google Scholar](#)

Ren Y., Xu H.W., Davey F., Taylor M., Aiton J., Coote P., Fang F., Yao J., Chen D., Chen J.X., et al. (2008) **Endophilin I expression is increased in the brains of Alzheimer disease patients** *The Journal of biological chemistry* **283**:5685–5691 [Google Scholar](#)

Ringstad N., Gad H., Low P., Di Paolo G., Brodin L., Shupliakov O., De Camilli P. (1999) **Endophilin/SHP3p4 is required for the transition from early to late stages in clathrin-mediated synaptic vesicle endocytosis** *Neuron* **24**:143–154 [Google Scholar](#)

- Ringstad N., Nemoto Y., De Camilli P. (1997) **The SH3p4/Sh3p8/SH3p13 protein family: binding partners for synaptojanin and dynamin via a Grb2-like Src homology 3 domain** *Proc Natl Acad Sci U S A* **94**:8569–8574 [Google Scholar](#)
- Russo I., Gavello D., Menna E., Vandael D., Veglia C., Morello N., Corradini I., Focchi E., Alfieri A., Angelini C., et al. (2019) **p140Cap Regulates GABAergic Synaptogenesis and Development of Hippocampal Inhibitory Circuits** *Cerebral Cortex* **29**:91–105 [Google Scholar](#)
- Saiepour L., Fuchs C., Patrizi A., Sassoè-Pognetto M., Harvey R.J., Harvey K (2010) **Complex role of collybistin and gephyrin in GABAA receptor clustering** *J Biol Chem* **285**:29623–29631 [Google Scholar](#)
- Sakimoto Y., Oo P.M., Goshima M., Kanehisa I., Tsukada Y., Mitsushima D (2021) **Significance of GABAA Receptor for Cognitive Function and Hippocampal Pathology** *Int J Mol Sci* **22** [Google Scholar](#)
- Schulz A., Miehl C., Berry M.J., Gjorgjieva J. (2021) **The generation of cortical novelty responses through inhibitory plasticity** *eLife* **10** [Google Scholar](#)
- Schuske K.R., Richmond J.E., Matthies D.S., Davis W.S., Runz S., Rube D.A., van der Bliek A.M., Jorgensen E.M. (2003) **Endophilin is required for synaptic vesicle endocytosis by localizing synaptojanin** *Neuron* **40**:749–762 [Google Scholar](#)
- Scott S., Aricescu A.R (2019) **A structural perspective on GABA(A) receptor pharmacology** *Current opinion in structural biology* **54**:189–197 [Google Scholar](#)
- Sexton C.A., Penzinger R., Mortensen M., Bright D.P., Smart T.G (2021) **Structural determinants and regulation of spontaneous activity in GABAA receptors** *Nature communications* **12**:5457 [Google Scholar](#)
- Soukup S.F., Kuenen S., Vanhauwaert R., Manetsberger J., Hernandez-Diaz S., Swerts J., Schoovaerts N., Vilain S., Goukko N.V., Vints K., et al. (2016) **A LRRK2-Dependent EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Terminals** *Neuron* **92**:829–844 [Google Scholar](#)
- Spruston N (2008) **Pyramidal neurons: dendritic structure and synaptic integration** *Nature reviews Neuroscience* **9**:206–221 [Google Scholar](#)
- Tao C.L., Liu Y.T., Sun R., Zhang B., Qi L., Shivakoti S., Tian C.L., Zhang P., Lau P.M., Zhou Z.H., et al. (2018) **Differentiation and Characterization of Excitatory and Inhibitory Synapses by Cryo-electron Tomography and Correlative Microscopy** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **38**:1493–1510 [Google Scholar](#)
- Thomas P., Mortensen M., Hosie A.M., Smart T.G (2005) **Dynamic mobility of functional GABAA receptors at inhibitory synapses** *Nature neuroscience* **8**:889–897 [Google Scholar](#)
- Tretter V., Jacob T.C., Mukherjee J., Fritschy J.M., Pangalos M.N., Moss S.J (2008) **The clustering of GABA(A) receptor subtypes at inhibitory synapses is facilitated via the direct binding of receptor alpha 2 subunits to gephyrin** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **28**:1356–1365 [Google Scholar](#)

Tyagarajan S.K., Fritschy J.M (2014) **Gephyrin: a master regulator of neuronal function?** *Nat Rev Neurosci* **15**:141–156 [Google Scholar](#)

Tyagarajan S.K., Ghosh H., Yévenes G.E., Imanishi S.Y., Zeilhofer H.U., Gerrits B., Fritschy J.M. (2013) **Extracellular signal-regulated kinase and glycogen synthase kinase 3 β regulate gephyrin postsynaptic aggregation and GABAergic synaptic function in a calpain-dependent mechanism** *The Journal of biological chemistry* **288**:9634–9647 [Google Scholar](#)

van Bommel B., Konietzny A., Kobler O., Bar J., Mikhaylova M. (2019) **F-actin patches associated with glutamatergic synapses control positioning of dendritic lysosomes** *The EMBO journal* **38**:e101183 [Google Scholar](#)

Van Erum J., Van Dam D., De Deyn P.P. (2019) **PTZ-induced seizures in mice require a revised Racine scale** *Epilepsy & behavior : E&B* **95**:51–55 [Google Scholar](#)

Verstreken P., Koh T.W., Schulze K.L., Zhai R.G., Hiesinger P.R., Zhou Y., Mehta S.Q., Cao Y., Roos J., Bellen H.J (2003) **Synaptojanin is recruited by endophilin to promote synaptic vesicle uncoating** *Neuron* **40**:733–748 [Google Scholar](#)

Wagner S., Lee C., Rojas L., Specht C.G., Rhee J., Brose N., Papadopoulos T (2021) **The $\alpha 3$ subunit of GABA(A) receptors promotes formation of inhibitory synapses in the absence of collybistin** *J Biol Chem* **296**:100709 [Google Scholar](#)

Watanabe S., Mamer L.E., Raychaudhuri S., Luvsanjav D., Eisen J., Trimbuch T., Sohl-Kielczynski B., Fenske P., Milosevic I., Rosenmund C., et al. (2018) **Synaptojanin and Endophilin Mediate Neck Formation during Ultrafast Endocytosis** *Neuron* **98**:1184–1197 [Google Scholar](#)

Yang Y., Chen J., Chen X., Li D., He J., Wang S., Zhao S., Yang X., Deng S., Tong C., et al. (2021) **Endophilin A1 drives acute structural plasticity of dendritic spines in response to Ca²⁺/calmodulin** *The Journal of cell biology* **220** [Google Scholar](#)

Yang Y., Chen J., Guo Z., Deng S., Du X., Zhu S., Ye C., Shi Y.S., Liu J.J (2018) **Endophilin A1 Promotes Actin Polymerization in Dendritic Spines Required for Synaptic Potentiation** *Frontiers in molecular neuroscience* **11**:177 [Google Scholar](#)

Yang Y., Wei M., Xiong Y., Du X., Zhu S., Yang L., Zhang C., Liu J.J (2015) **Endophilin A1 regulates dendritic spine morphogenesis and stability through interaction with p140Cap** *Cell research* **25**:496–516 [Google Scholar](#)

Yu Q., Wang Y., Du F., Yan S., Hu G., Origlia N., Rutigliano G., Sun Q., Yu H., Ainge J., et al. (2018a) **Overexpression of endophilin A1 exacerbates synaptic alterations in a mouse model of Alzheimer's disease** *Nature communications* **9**:2968 [Google Scholar](#)

Yu X., Xu T., Ou S., Yuan J., Deng J., Li R., Yang J., Liu X., Li Q., Chen Y (2018b) **Endophilin A1 mediates seizure activity via regulation of AMPARs in a PTZ-kindled epileptic mouse model** *Experimental neurology* **304**:41–57 [Google Scholar](#)

Author information

Xue Chen

State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China, College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China

Deng Pan

State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China, College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China

Jia-Jia Liu

State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China, College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China
ORCID iD: [0000-0002-6099-1059](https://orcid.org/0000-0002-6099-1059)

For correspondence: jjliu@genetics.ac.cn

Yanrui Yang

State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China
ORCID iD: [0000-0001-5194-1239](https://orcid.org/0000-0001-5194-1239)

For correspondence: yryang@genetics.ac.cn

Editors

Reviewing Editor

Inna Slutsky

Tel Aviv University, Tel Aviv, Israel

Senior Editor

Lu Chen

Stanford University, Stanford, United States of America

Reviewer #1 (Public review):

Summary:

In the present study, Chen et al. investigate the role of Endophilin A1 in regulating GABAergic synapse formation and function. To this end, the authors use constitutive or conditional knockout of Endophilin A1 (EEN1) to assess the consequences on GABAergic synapse composition and function, as well as the outcome for PTZ-induced seizure susceptibility. The authors show that EEN1 KO mice show a higher susceptibility to PTZ-induced seizures, accompanied by a reduction in the GABAergic synaptic scaffolding protein gephyrin as well as specific GABAAR subunits and eIPSCs. The authors then investigate the underlying mechanisms, demonstrating that Endophilin A1 binds directly to gephyrin and GABAAR subunits, and identifying the subdomains of Endophilin A1 that contribute to this effect. Overall, the authors state that their study places Endophilin A1 as a new regulator of GABAergic synapse function.

Strengths:

Overall, the topic of this manuscript is very timely, since there has been substantial recent interest in describing the mechanisms governing inhibitory synaptic transmission at GABAergic synapses. The study will therefore be of interest to a wide audience of neuroscientists studying synaptic transmission and its role in disease. The manuscript is well written and contains a substantial quantity of data. In the revised version of the manuscript, the authors have increased the number of samples analyzed and have significantly improved the statistical analysis, thereby substantially strengthening the conclusions of their study.

Comments on revised version:

The authors have addressed all of my concerns, and the manuscript has been substantially improved.

<https://doi.org/10.7554/eLife.102792.2.sa3>

Reviewer #2 (Public review):

Summary:

The function of neural circuits relies heavily on the balance of excitatory and inhibitory inputs. Particularly, inhibitory inputs are understudied when compared to their excitatory counterparts due to the diversity of inhibitory neurons, their synaptic molecular heterogeneity, and their elusive signature. Thus, insights into these aspects of inhibitory inputs can inform us largely on the functions of neural circuits and the brain.

Endophilin A1, an endocytic protein heavily expressed in neurons, has been implicated in numerous pre- and postsynaptic functions, however largely at excitatory synapses. Thus, whether this crucial protein plays any role in inhibitory synapse, and whether this regulates functions at the synaptic, circuit, or brain level remains to be determined.

New Findings:

- (1) Endophilin A1 interacts with the postsynaptic scaffolding protein gephyrin at inhibitory postsynaptic densities within excitatory neurons.
- (2) Endophilin A1 promotes the organization of the inhibitory postsynaptic density and the subsequent recruitment/stabilization of GABA A receptors via Endophilin A1's membrane binding and actin polymerization activities.
- (3) Loss of Endophilin A1 in CA1 mouse hippocampal pyramidal neurons weakens inhibitory input and leads to susceptibility to epilepsy.
- (4) Thus the authors propose that via its role as a component of the inhibitory postsynaptic density within excitatory neurons, Endophilin A1 supports the organization, stability, and efficacy of inhibitory input to maintain the excitatory/inhibitory balance critical for brain function.
- (5) The conclusion of the manuscript is well supported by the data but will be strengthened by addressing our list of concerns and experiment suggestions.

Comments on revised version:

The authors addressed the concerns adequately. The three remaining concerns are:

- (1) The use of one-way ANOVA is not well justified.

(2) The use of superplots to show culture to culture variability would make it more transparent.

(3) Change EEN1 in Figure 8B to EndoA1.

<https://doi.org/10.7554/eLife.102792.2.sa2>

Reviewer #3 (Public review):

Summary:

The authors investigated a possible role of Endophilin A1 in the inhibitory postsynaptic density.

Strengths:

The authors used a broad array of experimental approaches to investigate this, including tests of seizure susceptibility, electrophysiology, biochemistry, neuronal culture and image analysis.

Weaknesses:

Many results are difficult to interpret, and data quality is not always convincing, unfortunately. The basic premise of the study, that gephyrin and endophilin A1 interact, requires more robust analysis to be convincing.

Specific comments:

The authors have made a substantial effort to improve their manuscript. A number of issues, related to numbers of observations mentioned by the reviewers, are clarified in the revised manuscript. The authors have also clarified some of the other questions from the reviewers. The long list of issues brought up by the reviewers and the many corrections needed still raise questions about data quality in this manuscript.

In response to my comments (Point 2), the added experiment with PSD95.FingR and GPN.FingR in cultured neurons (Fig. S5A-D) is a good addition; the in vivo data using FingRs in Figure S3 look less convincing however. In response to my Point 5, the authors have added a cell-free binding assay (Figure 5I). This is a useful addition, but to convincingly make the point of interaction between Gephyrin and EndoA1, more rigorous biophysical quantitation of binding is needed. The legend in Figure 5I states that 4 independent experiments were performed, but the graph only shows 3 dots. This needs to be corrected.

<https://doi.org/10.7554/eLife.102792.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

Summary:

In the present study, Chen et al. investigate the role of Endophilin A1 in regulating GABAergic synapse formation and function. To this end, the authors use constitutive or conditional knockout of Endophilin A1 (EEN1) to assess the consequences on GABAergic

synapse composition and function, as well as the outcome for PTZ-induced seizure susceptibility. The authors show that EEN1 KO mice show a higher susceptibility to PTZ-induced seizures, accompanied by a reduction in the GABAergic synaptic scaffolding protein gephyrin as well as specific GABAAR subunits and eIPSCs. The authors then investigate the underlying mechanisms, demonstrating that Endophilin A1 binds directly to gephyrin and GABAAR subunits, and identifying the subdomains of Endophilin A1 that contribute to this effect. Overall, the authors state that their study places Endophilin A1 as a new regulator of GABAergic synapse function.

Strengths:

Overall, the topic of this manuscript is very timely, since there has been substantial recent interest in describing the mechanisms governing inhibitory synaptic transmission at GABAergic synapses. The study will therefore be of interest to a wide audience of neuroscientists studying synaptic transmission and its role in disease. The manuscript is well-written and contains a substantial quantity of data.

Weaknesses:

A number of questions remain to be answered in order to be able to fully evaluate the quality and conclusions of the study. In particular, a key concern throughout the manuscript regards the way that the number of samples for statistical analysis is defined, which may affect the validity of the data analysed. Addressing this weakness will be essential to providing conclusive results that support the authors' claims.

We would like to thank the reviewer for appreciation of the value of our study and careful critics to help us improve the manuscript. We will correct the way that the number of samples for statistical analysis is defined throughout the manuscript as suggested and update figures, figure legends, and Materials and Methods accordingly. For example, we will average the values for all dendritic segments from one neuron, so that each data point represents one neuron in the graphs.

Reviewer #2 (Public review):

Summary:

The function of neural circuits relies heavily on the balance of excitatory and inhibitory inputs. Particularly, inhibitory inputs are understudied when compared to their excitatory counterparts due to the diversity of inhibitory neurons, their synaptic molecular heterogeneity, and their elusive signature. Thus, insights into these aspects of inhibitory inputs can inform us largely on the functions of neural circuits and the brain.

Endophilin A1, an endocytic protein heavily expressed in neurons, has been implicated in numerous pre- and postsynaptic functions, however largely at excitatory synapses. Thus, whether this crucial protein plays any role in inhibitory synapse, and whether this regulates functions at the synaptic, circuit, or brain level remains to be determined.

New Findings:

(1) Endophilin A1 interacts with the postsynaptic scaffolding protein gephyrin at inhibitory postsynaptic densities within excitatory neurons.

(2) Endophilin A1 promotes the organization of the inhibitory postsynaptic density and the subsequent recruitment/stabilization of GABA A receptors via Endophilin A1's membrane binding and actin polymerization activities.

(3) Loss of Endophilin A1 in CA1 mouse hippocampal pyramidal neurons weakens inhibitory input and leads to susceptibility to epilepsy.

(4) Thus the authors propose that via its role as a component of the inhibitory postsynaptic density within excitatory neurons, Endophilin A1 supports the organization, stability, and efficacy of inhibitory input to maintain the excitatory/inhibitory balance critical for brain function.

(5) The conclusion of the manuscript is well supported by the data but will be strengthened by addressing our list of concerns and experiment suggestions.

We would like to thank the reviewer for their favorable impression of manuscript. We also appreciate the great experiment suggestions to help us improve the manuscript.

Weaknesses:

Technical concerns:

(1) Figure 1F and Figure 1H, Figures 7H,J:

Can the authors justify using a paired-pulse interval of 50 ms for eEPSCs and an interval of 200 ms for eIPSCs? Otherwise, experiments should be repeated using the same paired pulse interval.

We apologize for the confusion. As illustrated by the schematic current traces, the decay time constants of eEPSCs and eIPSCs in hippocampal CA1 neurons are different. The eEPSCs exhibit a faster channel closing rate, corresponding to a smaller time constant Tau. Thus, a shorter inter-stimulus interval (50 ms) was chosen for paired-pulse ratio recordings. In contrast, the eIPSCs display a slower channel closing rate, with a Tau value larger than that of eEPSCs, so a longer inter-stimulus interval (200 ms) was used for PPR. This protocol has been long-established and adopted in previous studies (please see below for examples).

Contractor, A., Swanson, G. & Heinemann, S. F. Kainate receptors are involved in short- and long-term plasticity at mossy fiber synapses in the hippocampus. *Neuron* 29, 209-216, doi:10.1016/s0896-6273(01)00191-x (2001).

Babiec, W. E., Jami, S. A., Guglietta, R., Chen, P. B. & O'Dell, T. J. Differential Regulation of NMDA Receptor-Mediated Transmission by SK Channels Underlies Dorsal-Ventral Differences in Dynamics of Schaffer Collateral Synaptic Function. *Journal of neuroscience* 37, 1950-1964, doi:10.1523/JNEUROSCI.3196-16.2017 (2017).

(2) Figures 3G,H,I:

While 3D representations of proteins of interest bolster claims made by superresolution microscopy, SIM resolution is unreliable when deciphering the localization of proteins at the subsynaptic level given the small size of these structures (<1 micrometer). In order to determine the actual location of Endophilin A1, especially given the known presynaptic localization of this protein, the authors should complete SIM experiments with a presynaptic marker, perhaps an active zone protein, so that the relative localization of Endophilin A1 can be gleaned. Currently, overlapping signals could stem from the presynapse given the poor resolution of SIM in this context.

Thanks for your suggestions. It is certainly preferable to investigate the relative localization of endophilin A1 using both presynaptic and postsynaptic markers. For SIM imaging in Figure 3G-I, to visualize neuronal morphology, we immunostained GFP as cell fill, leaving two other channels for detection of immunofluorescent signals of endophilin A1 and another protein.

We will try co-immunostaining of endophilin A1, the active zone protein bassoon (presynaptic marker) and gephyrin without morphology labeling. Alternatively, we will do co-staining of endophilin A1 and bassoon in GFP-expressing neurons. We agree that overlapping signals or proximal localization of presynaptic endophilin A1 with gephyrin or GABA_AR γ 2 could not be ruled out. To note, if image resolution is improved with the use of a more advanced imaging system, the overlap between two proteins will become smaller or even disappear. With the ~110 nm lateral resolution of SIM microscopy, the degree of overlap between the two proteins of interest is much lower than in confocal microscopy. Given the presynaptic localization of endophilin, most likely we will observe a small overlap (presynaptic) or proximal localization (postsynaptic) of endophilin A1 with bassoon. Nevertheless, we will complete the SIM experiments as suggested to improve the manuscript.

Manuscript consistency:

(1) Figure 2:

The authors looked at VGAT and noticed a reduction of signals in hippocampal regions in their P21 slices, indicating that the proposed postsynaptic organization/stabilization functions of Endophilin A1 extend to the inhibitory presynapse, perhaps via Neuroligin 2-Neurexin. Simultaneously, hippocampal regions in P21 slices showed a reduction in PSD-95 signals, indicating that excitatory synapses are also affected. It would be crucial to also look at excitatory presynapses, via VGLUT staining, to assess whether EndoA1^{-/-} also affects presynapses. Given the extensive roles of Endophilin A1 in presynapses, especially in excitatory presynapses, this should be investigated.

Thanks for the thoughtful comments. Given that the both VGAT and PSD95 signals are reduced in hippocampal regions in P21 slices, it is conceivable that the proposed postsynaptic organization/stabilization functions of endophilin A1 extend to the inhibitory presynapse via Neuroligin-2-Neurexin and the excitatory presynapse as well during development. Of note, endophilin A1 knockout did not impair the distribution of Neuroligin-2 in inhibitory postsynapses (immunoisolated with anti-GABA_AR α 1) in mature mice (Figure 3K), and endophilin A1 did not bind to Neuroligin-2 (Figure 4D), suggesting that endophilin A1 might function via other mechanisms. Nevertheless, as functions of endophilin A family members at the presynaptic site are well-established, the reduction of presynaptic signals in developmental hippocampal regions of EndoA^{-/-} mice might result from the depletion of presynaptic endophilin A1. The presynaptic deficits can be compensatory by other mechanisms as neurons mature. Certainly, we will do VGLUT staining of EndoA1^{-/-} brain slices as suggested to assess the role of endophilin A1 in excitatory presynapses in vivo.

(2) Figure 7C:

The authors do not assess whether p140Cap overexpression rescues GABA_AR receptor loss exhibited in Endophilin A1 KO, as they did for Gephyrin. This would be an important data point to show, as p140Cap may somehow rescue receptor loss by another pathway. In fact, it is mentioned in the text that this experiment was done, "Consistently, neither p140Cap nor the endophilin A1 loss-of-function mutants could rescue the GABA_AR clustering phenotype in EEN1 KO neurons (Figure 7C, D)" yet the data for p140Cap overexpression seem to be missing. This should be remedied.

Thanks a lot for the thoughtful comment. We will determine whether p140Cap overexpression also rescues the GABA_AR clustering phenotype in EndoA1^{-/-} neurons by surface GABA_AR γ 2 staining in our revised manuscript.

Reviewer #3 (Public review):

Summary:

Chen et al. identify endophilin A1 as a novel component of the inhibitory postsynaptic scaffold. Their data show impaired evoked inhibitory synaptic transmission in CA1 neurons of mice lacking endophilin A1, and an increased susceptibility to seizures. Endophilin can interact with the postsynaptic scaffold protein gephyrin and promote assembly of the inhibitory postsynaptic element. Endophilin A1 is known to play a role in presynaptic terminals and in dendritic spines, but a role for endophilin A1 at inhibitory postsynaptic densities has not yet been described.

Strengths:

The authors used a broad array of experimental approaches to investigate this, including tests of seizure susceptibility, electrophysiology, biochemistry, neuronal culture, and image analysis.

Weaknesses:

Many results are difficult to interpret, and the data quality is not always convincing, unfortunately. The basic premise of the study, that gephyrin and endophilin A1 interact, requires a more robust analysis to be convincing.

We greatly appreciate the positive comment on our study and the very valuable feedback for us to improve the manuscript. We will conduct additional experiments to improve our data quality and strengthen our evidences according to these great constructive suggestions. To gain strong evidence for the interaction between endophilin A1 and gephyrin, we will perform in vitro pull-down assay with recombinant proteins from bacterial expression system.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

(1) For all of the electrophysiology experiments, only the number of neurons recorded is stated, but not the number of independent animals that these neurons were obtained from. The number of independent animals used should be stated for each panel. At least 3 independent animals should be used in each group, otherwise, more data needs to be added.

We apologize for missing the information in the original manuscript. For all electrophysiological experiments, data were obtained from more than 3 experimental animals. The figure legends were updated to include the number of independent animals used for each panel.

(2) For the cell culture experiments analyzing dendritic puncta at GABAergic synapses, the number of data points analysed appears to be the number of dendritic segments quantified, regardless of whether they originate from the same neuron or not. This analysis method is not valid, since dendritic segments from the same neuron cannot be counted as statistically independent samples. The authors need to average the values for all dendritic segments from one neuron, such that one neuron equals one data point. This alteration should be made for Figures 2B, 2D, 4H, 4J, 5B, 5C, 5E, 5J, 5L, 6B, 6D, 6F, 6H, 6J, 6K, 7B, and 7D. In addition, the number of independent cultures from which the neurons were obtained should be stated for each panel. At least 3 independent cultures should be used in each group, otherwise, more data need to be added.

Thanks for the criticism. We reanalyzed the data throughout the manuscript as suggested and updated the figure legends accordingly. Moreover, we increased the number of neurons from independent experiments to further confirm the results in our revised manuscript.

In the revised manuscript, we averaged the values for all dendritic segments from a single neuron and updated the data in Figure 3B, 3D, 4H, 4J, 5B, 5C, 5E, 5K, 5M, 6B, 6D, 6F, 6H, 6J, 6K, 7B, and 7D.

Neurons analyzed in each group were derived from at least 3 independent cultures. Due to very low efficiency of sparse transfection in primary cultured hippocampal neurons, multiple experimental repetitions were necessary to obtain the sufficient number of neurons for analysis. We described statistical analysis in “Material and Methods” section in the original manuscript as follows:

“For all biochemical, cell biological and electrophysiological recordings, at least three independent experiments were performed (independent cultures, transfections or different mice).”

(3) Individual data points should be shown on all graphs, particularly in Figures 2C, 2F, 2I, 3F, 3K, and 3L.

Thank you for the suggestion. We replaced the original graphs with scatterplots and mean $\pm$ S.E.M. in new Figures.

(4) For each experiment, the authors should state explicitly in the methods section whether that experiment was conducted blind to genotype.

Thank you for the suggestion. We have modified the description of blind analysis for each experiment in methods section to “Seizure susceptibility was measured blindly by rating seizures on a scale of 0 to 7 as follows...”, “Quantification of immunostaining were carried out blindly...” in our revised manuscript.

(5) For each experiment, the authors should state whether they used male or female mice, and what age the mice were at the time of the experiment

Thanks a lot for the suggestion. We usually use male and female mice for neuron culture and behavioral test. We observed no sex-related differences in PTZ-induced behaviors, so the results were pooled together.

For mice ages, P0 pups were used for hippocampal neuron cultures and virus injection in electrophysiological recording assays or FingR probes assays. P14-21 mice were used for electrophysiological recording, immunofluorescent staining and FingR probes detection in brain slice, while adult mice (P60) for behavioral tests, immunofluorescent staining in brain slice and biochemical assays. We have modified the description in genders and ages of mice in methods section to “To evaluate seizure susceptibility, 8-10-week-old male and female *EndoA1*^{+/+} or *EndoA1*^{-/-} littermates or *EndoA1*^{fl/fl} littermates were intraperitoneally administered...”, “For virus injection, 8-9-week-old naive male and female littermates were anesthetized...”, “Male and female littermates (P21 or P60) were anesthetized and immediately perfused...”, “Hippocampi of female or male pups (P0) were rapidly dissected under sterile conditions...”, “PSD fractions from adult mouse brain were prepared as previously described...”, “Newborn *EndoA1*^{fl/fl} littermates (male or female) were anesthetized on ice for 4-5 min...” in our revised manuscript.

(6) For each experiment involving WT and KO mice, please state whether WTs and KOs were bred as littermates from heterozygous breeders

Sorry for the confusion. In our study, *EndoA1*^{+/+} and *EndoA1*^{-/-} mice were bred as littermates from heterozygous breeders. We added the information in methods section as follows in our revised manuscript, “*EndoA1*^{+/+} and *EndoA1*^{-/-} mice were bred as littermates from heterozygous breeders...”, “To evaluate seizure susceptibility, 8-10-week-old male and female *EndoA1*^{+/+} or *EndoA1*^{-/-} littermates or *EndoA1*^{fl/fl} littermates...”, “For virus injection, 8-9-week-old naive male and female littermates were anesthetized...”, “Male and female littermates (P21 or P60) were anesthetized and immediately perfused...”, “For co-IP from brain lysates, the whole brain from 8-10-week-old WT and KO littermates were dissected...”, “Newborn *EndoA1*^{fl/fl} littermates (male or female) were anesthetized on ice for 4-5 min...”.

(7) For experiments comparing three or more groups, the authors claim in the methods section to have used a one-way ANOVA for statistical analysis. However, no ANOVA values are given, only the post-hoc tests. Please add the ANOVA values for each experiment before stating the values of the post-hoc analysis.

Sorry for the missing information. We used one-way ANOVA for comparing three or more groups in the original manuscript and have changed to two-way ANOVA for behavior data analysis in our revised manuscript as suggested in Recommendations (18). We added the ANOVA values (F & p values) for each experiment in new figures. For example, see Figure 1C.

(8) In Figure 1A-C, seizure susceptibility was compared in *EEN*^{+/+} and *EEN*^{-/-} mice, but the methods section states that seizure susceptibility was evaluated in 8-10-week-old male C57BL/6N mice (line 513). Was this meant to indicate that the *EEN*^{+/+} and *EEN*^{-/-} mice were on a C57BL/6N background? How does this match with the statement that *EEN1*^{-/-} mice were generated on a C57BL/6J background (line 467)?

We apologize for the mistake. In our study, *EEN1*^{-/-} mice were generated on a C57BL/6J background, as stated in our previously published papers (Yang et al., 2021; Yang et al., 2018) and in “Animals” in Material and Methods of our original manuscript. We had corrected the statement to “To evaluate seizure susceptibility, 8-10-week-old male and female *EndoA1*^{+/+} or *EndoA1*^{-/-} littermates...” in Material and Methods of the revised manuscript.

(9) In the electrophysiology experiments in Figure 1E-O, it is not clear to me which neurons were recorded in the control group. The methods section states that “Whole-cell recordings were performed on an AAV-infected neuron and a neighboring uninfected neuron” (line 736). However, the figure legends states that recordings were obtained from “10 control (Ctrl, mCherry alone) and 10 *EEN1* KO (mCherry and Cre) pyramidal neurons” (line 1079), which would indicate that the controls are not uninfected neurons from the same animal, but AAV-mCherry infected neurons from a different animal. Please clarify which of the two descriptions is accurate.

Thanks for catching the error! In all electrophysiological experiments, a neighboring uninfected neuron was used as the control in Figure 1E-O. This was incorrectly stated in the figure legend of the original manuscript. In the revised manuscript, the information has been corrected in figure legends of new Figure 1 (E-F).

(10) The authors show that in Endophilin A1 KO animals, eIPSCs are reduced, but mIPSC frequency and amplitude are unaltered. How do they explain this finding in the context of the fact that gephyrin and GABAAR1.

We apologize for the confusion about the data of electrophysiological recording. Compared with eIPSC, which are recorded in the presence of electrically evoked action potential that elicited a substantial release of neurotransmitter, mIPSCs are small, spontaneous currents

recorded in the presence of TTX during patch-clamp experiments, resulting from the release of neurotransmitters from presynaptic terminals in the absence of action potential. The amplitude of mIPSCs typically reflects the quantal release of neurotransmitters, while their frequency can vary depending on synaptic activity and the state of the neuron.

A number of molecules fine-tune presynaptic neurotransmitter release and functions of inhibitory postsynaptic receptors. In our study, inhibitory postsynapses were partially affected in endophilin A1 knockout neurons, while presynaptic endophilin A1 remained intact during electrophysiological recordings. Conceivably, the observed deficits in endophilin A1 knockout mice were mild. Following endophilin A1 depletion, inhibitory postsynaptic receptors appeared sufficient to respond to spontaneous neurotransmitter release but may be inadequate to large amounts of neurotransmitter release evoked by action potential. Meanwhile, spontaneous synaptic activity and the state of the neuron were not obviously affected under basic state by endophilin A1 depletion during postnatal stages. Consequently, mIPSC frequency and amplitude remain unaltered but eIPSCs were reduced compared to the control neurons. This finding was consistent with behavioral experiments, where aggressive epileptic behaviors were induced by PTZ rather than spontaneous epilepsy in endophilin A1 knockout mice.

(11) Distribution of gephyrin, VGAT, and GABAAR α 2 differs substantially between the different layers of hippocampal area CA1, and the same goes for the other regions of the hippocampus. However, in Figure 2, it is not clear to me from the sample images which layers of each subregion the authors quantified, or indeed whether they paid attention to which layers they included in their analysis. This can lead to a substantial skewing of the data if different layers were preferentially included in the two genotypes. Please clarify which layers were analysed, and how comparability between WT and KO was ensured. This is particularly important given the authors' claim that Endophilin A1 acts equally at all subtypes of GABAergic synapses (lines 373- 376).

Thanks for the cautiousness! We distinguished each hippocampal subregion based on the anatomical structure in brain slices. Quantification of fluorescent mean intensity of each synaptic protein in all layers of each subregion, as shown in new Figure 2 and Figure S2A-F, revealed that GABAergic synaptic proteins were impaired in both P21 and P60 KO mice.

We further analyzed the fluorescent signal of core postsynaptic component, gephyrin, in individual layers of each subregion in the hippocampus of mature WT and KO mice, as presented in new Figures S2G-H. Our findings demonstrated a decrease in gephyrin levels across all layers of each subregion in KO mice. Additionally, we examined gephyrin clustering across the soma, axon initial segment (AIS), and dendrites in cultured mature endophilin A1 knockout hippocampal neurons, as shown in new Figure S5E-H. The results showed that gephyrin was affected in all subcellular regions following endophilin A1 knockout.

Collectively, these data suggest that endophilin A1 functions across all subtypes of GABAergic postsynapses.

(12) In Figure 3E-F, the authors state that there was no change in the total level of synaptic neurons in EEN1 KO neurons (line 188). However, there is no quantification of the total level of synaptic neurons shown, and based on the immunoblot in Figure 3E, it looks like there is a substantial reduction in NR1, NL2, and g2. The authors should present a quantification of the total levels of these proteins and adjust their statement accordingly if necessary.

Thanks a lot for your comments. We quantified the total protein levels in Figure 3E and added the result to new Figure 3F, showing that total protein levels were not obviously affected in cultured KO neurons. When normalized to total protein levels, the surface levels

of GABA_A receptors were significantly compromised compared to surface GluN1 and NL2. Furthermore, the total protein levels were not affected in brains of KO mice, as shown in Figures 3K (input) and 3L (S1). Collectively, there was no change in the total level of synaptic proteins in KO neurons.

(13) In Figure 3G-I, the authors claim, based on super-resolution images as presented here, that Endophilin A1 colocalizes with gephyrin and g2. However, no quantification of this colocalization is presented. The authors should add this quantification to support their claim and indicate how many GABAergic synapses contain Endophilin A1.

Thank you for the thoughtful comments. The resolution of the images is significantly improved by super-resolution microscopy. As a result, the overlap between the two proteins will become smaller or even disappear. Since no two proteins can occupy the same physical space, they would show lower colocalization and instead exhibit proximal localization. As expected, in Figures 3G and 3H, we observed only small overlap or proximal localization of endophilin A1 with gephyrin or GABA_A R γ 2. To further confirm the localization of endophilin A1 in inhibitory synapses, we co-stained endophilin A1 with both pre- and post-synaptic proteins, gephyrin and Bassoon. Then we quantified the colocalization of endophilin A1 with gephyrin or with Bassoon using the method for super-resolution images described in the reference (Andrew D. McCall. Colocalization by cross-correlation, a new method of colocalization suited for super-resolution microscopy. McCall BMC Bioinformatics (2024) 25:55). The percentage of gephyrin or Bassoon puncta that were in close proximity with endophilin A1 was also calculated, as shown in new video 5 and new Figure S4B-G. These data have been added in the revised manuscript as follows, “We further detected the localization of endophilin A1 to inhibitory synapses by co-immunostaining with both pre- and post-synaptic markers (Figure. S4B and Video 5). Quantitative analysis of super-resolution localization maps revealed that ~ 47 % puncta of gephyrin or Bassoon were proximal to endophilin A1 (Figure. S4G, $n = 14$), with a mean distance between endophilin A1- and gephyrin-positive pixels of ~ 120 nm, or between endophilin A1- and Bassoon-positive pixels of ~ 130 nm (Figure. S4C-F).”

(14) In the quantification shown in Figure 3K-L, there are no error bars in the WT data sets. This presumably means that all values were normalized to WT. However, since this artificially eliminates the variance in the WT group, a t-test is no longer valid, since this assumes a normal distribution and normal variance, which are no longer given. The authors should either change the way they normalize their data to maintain the variance in the WT group or perform a different statistical test that can account for the artificial lack of variance in one of the groups.

Thank you for the suggestions! We modified our analysis approach. Specifically, we used mean value of WTs to normalize data to preserve the variance in the WT group and performed unpaired t-tests to assess statistical significance in Figure 3K-L. Additionally, we replaced the bar graphs with modified graphs showing individual data points. Please see Response to Recommendation (12).

(15) What is the difference between the coIP experiment in Figure 4E and 3J, right panel? In both cases, an Endophilin A1 IP is performed, and gephyrin, GABAAR γ 2, and GABAAR α 1 are assessed. However, Figure 3J's right panel indicates that Endophilin A1 does interact with the GABAAR subunits, whereas Figure 4E shows that it does not. How do the authors explain this discrepancy? Were these experiments performed more than once?

Sorry for the confusion. Figure 3J and Figure 4E show data from immunoprecipitation assay and conventional co-immunoprecipitation (co-IP), respectively. Immunoprecipitation allows for the

rapid and efficient separation of subcellular membrane compartments using antibodies conjugated to magnetic beads. In Figure 3J, we used antibodies against GABA_A α 1 subunit or endophilin A1 to isolate the inhibitory postsynaptic membranes or endophilin A1-associated membranous compartments. In contrast, co-immunoprecipitation detects direct protein-protein interactions in detergent-solubilized lysates. For Figure 4E, we applied antibodies against endophilin A1 to precipitate its interaction partners. The results in Figure 3J and Figure 4E demonstrate that endophilin A1 is localized in the inhibitory postsynaptic compartment and directly interacts with gephyrin, but not with GABA_ARs. Detailed information regarding the methods used for co-IP and immunoisolation can be found in “GST-pull down, co-immunoprecipitation (IP), and immunoisolation” in the “Material and Methods” section of original manuscript.

These experiments were repeated multiple times to ensure reliability. In fact, consistent data showing endophilin A1 localization in the inhibitory postsynaptic compartment were observed in Figure 3K, showing the quantified data as well.

(16) For the colocalization analysis in Figure 5A-C, what percentage of gephyrin puncta contain g2 in the WT and Endophilin A1 KO? Currently, only a correlation coefficient is provided, but not the degree of overlap. Please add this information to the figure.

Thanks for the comments on the colocalization analysis. We analyzed the percentage of gephyrin puncta overlapping with GABA_A γ 2 and added the graphs in new Figure 5C.

(17) Figure 6 investigates how actin depolarization affects GABAergic synapse function, but does not assess how Endophilin A1 contributes to this process. The authors then provide an extremely short statement in the discussion, stating that their data are contradictory to a previous study (lines 412 - 417). This section of the discussion should be expanded to address the specific role of Endophilin A1 in the consequences of actin depolymerization.

Thanks a lot for the advice. In the original manuscript, we discussed the specific role of endophilin A1 at inhibitory postsynapses as follows in Discussion:

“As membrane-binding and actin polymerization-promoting activities of endophilin A1 are both required for its function in enhancing iPSD formation and γ 2-containing GABA_A clustering to iPSD, we propose that membrane-bound endophilin A1 promotes postsynaptic assembly by coordinating the plasma membrane tethering of the postsynaptic protein complex and its stabilization with the actin cytomatrix”

Following your advice, we added a statement in the revised manuscript addressing the role of endophilin A1 in actin polymerization at inhibitory postsynapses, shown as follows, “In the present study, the impaired clustering of gephyrin and GABA_A γ 2 by F-actin depolymerization underscores the essential role of F-actin in the assembly and stabilization of the inhibitory postsynaptic machinery. Membrane-bound endophilin A1 promotes F-actin polymerization beneath the plasma membrane through its interaction with p140Cap, an F-actin regulatory protein, thereby facilitating and/or stabilizing the clustering of gephyrin and γ 2-containing GABA_A receptors at postsynapses.”

(18) Which statistical analysis was conducted in Figure 7F? Given the nature of the data, a repeated measures ANOVA would be necessary to accurately assess the statistical accuracy.

Sorry for the confusion. We conducted one-way ANOVA followed by Tukey post hoc test at each time point in original Figure 7F. We have employed the method of repeated measures ANOVA followed by Tukey post hoc test as suggested in new Figure 7F. Meanwhile, we

reanalyzed data in new Figure 1C with the same method. We also modified the description in “Statistical analysis” and Figure legends for new Figure 1C and 7F in revised manuscript.

Reviewer #2 (Recommendations for the authors):

Data presentation:

(1) Figures 2A, B, D, E, G, H. Figures S2A, B, D:

Add P21 or P60 labels to these figures so that the difference between similarly stained samples (e.g. Figures 2A, B) is obvious to the reader.

Thanks! We added “P21” or “P60” labels in new Figure 2 and Figure S2 as suggested.

(2) Figures 4C, D:

The authors must make their coIP data annotation consistent. In Figure 4C, they use actual microgram amounts when, e.g., describing how much input was present, yet in Figure 4D they use + and -. The authors should pick one.

Thanks for the comments. We labeled the consistent data annotation in new Figure 4C and 4D, we also changed the label in 4F for the consistent data annotation.

(3) Figure 5A

GFP is gray in this figure, but in all other figures, it is blue. Consider changing for presentation reasons.

Thanks a lot for pointing out the problem. We replaced gray with blue color to indicate GFP in new Figure 5A.

(4) Figures 6A, C, E, G

Label graphs as either short-term or long-term drug treatment.

Thanks for the suggestion. We labeled the graphs as 60 min for short-term or 120 min for long-term drug treatment in new Figure 6A, C, E, G for convenient reading.

Annotation, grammar, spelling, typing errors:

(1) Figure 4G:

Merge and GFP labels are seemingly swapped.

Thanks a lot for sharp eye. We corrected the labels in new Figure 4G.

(2) Fig 4I:

The authors use "Gephyrin" instead of GPN. They should be consistent and choose one.

Sorry for the mistake. We changed the label consistent with other figures in new Figure 4I and rearranged the images in figures for good looking.

(3) "One-hour or two-hour treatment of mature neurons with nocodazole..."

Thanks for your advice. We modified the sentence to “Treatment of mature neurons with nocodazole, a microtubule depolymerizing reagent, for one hour (short-term) or two hours

(long-term), caused...”.

(4) The authors should indicate that one-hour is their short-term treatment and that two-hour is their long-term treatment so that when these terms are used later to describe LatA experiments, it is clearer to the reader.

Thanks for your comments. We modified the statement as seen in Response to Recommendation (3), it is clearer to the reader.

(5) EEA1. The authors should use a more conventional term EndoA1 so that the manuscript can be searched easily.

Thanks a lot for the suggestion. We replaced all of the term “EEN1” with “EndoA1” in the revised manuscript.

Reviewer #3 (Recommendations for the authors):

Major Points

(1) The number of observations for the electrophysiology experiments in Figure 1 (dots are neurons) is very low and it is not clear whether the data shown is derived from different mice. The same criticism applies to the data shown in Figures 7G-K.

We apologize for the low neuron number in electrophysiology experiments. In the patch-clamp experiments, the number of neurons recorded was higher than what is shown in the figures. However, neurons with a membrane resistance (R_m) below 500 M Ω , indicating unstable seals or poor conditions, were excluded from the analysis. Additionally, we added the number of mice from which the data derived in each group in the figure legends for Figure 1, 7 and S1, this point was also raised by Reviewer #1 (Please see Response to Recommendation (1)).

(2) Images in Figure 2 are shown at low magnification, statements on changes in intensity of inhibitory synaptic markers in the hippocampal region are impossible to interpret. Analysis of inhibitory synapses in vivo would require sparse neuronal labeling and 3D reconstruction, for instance using gephyrin-FingRs (Gross et al., Neuron 2013).

Thanks for your insightful suggestion. We obtained pCAG_PSD95.FingR-eGFP-CCR5TC and pCAG_GPN.FingR-eGFP-CCR5TC constructs from Addgene (plasmid # 46295 & #46296). We attempted in utero electroporation (IUE) to introduce the DNAs into cortical neurons or hippocampal neurons at E14.5, unfortunately with no success. Following the repetitive operation for numerous times, we could eventually obtain newborn pups of ICR mice after IUE. However, we failed to obtain any newborn pups of C57BL/6J mice due to abortion following the procedure. Furthermore, pregnant C57BL/6J mice (WTs or KOs) did not survive or remained in a poor state of health after surgery. Therefore, we were unable to analyze synapses through sparse labeling and 3D reconstruction by IUE. Alternatively, we obtained commercial AAVs carrying rAAV-EF1a-PSD95.FingR-eGFP-CCR5TC and rAAV-EF1a-mRuby2-Gephyrin.FingR-IL2RGTC, then injected into the CA1 region of *EndoA1^{fl/fl}* mice at P0. Mice were fixed and detected the fluorescent signals in CA1 regions at P21. Consistent with immunostaining with antibodies, decreased mRuby2-Gephyrin.FingR or PSD95.FingR-eGFP was observed in dendrites of KO neurons at P21, as shown in new Figure S3. In combination with electrophysiological recording, PSD fractionation and immunoisolation from brains, these data support our conclusion regarding the effects of endophilin A1 knockout on the inhibitory synapses.

Additionally, we transfected DIV12 cultured hippocampal neurons with pCAG_PSD95.FingR-eGFP-CCR5TC or pCAG_GPN.FingR-eGFP-CCR5TC and observed fluorescent signals on DIV16. Both the signal intensity and number of GPN.FingR-eGFP clusters were also significantly attenuated, with no obvious changes in PSD95.FingR-eGFP clusters in dendrites of mature neurons, as shown in new Figure S5A-D. We are very pleased that the result further strengthened our original conclusion. We have added the new pieces of data in our revised manuscript.

(3) Figure 3: surface labeling of GluA1 or the GABAAR gamma 2 subunit is difficult to interpret: the patterns are noisy and the numerous puncta appear largely non-synaptic although this is difficult to judge in the absence of additional synaptic markers. It appears statistics are done on dendritic segments rather than the number of neurons. The legend does not mention how many independent cultures this data is derived from. In their previous study (Yang et al., Front Mol Neurosci 2018), the authors noted a decrease in surface GluA1 levels in the absence of endophilin A1. How do they explain the absence of an effect on surface GluA1 levels in the current study?

Sorry for the concern and thanks for your comments. First, we assessed changes in the surface levels of excitatory and inhibitory receptors by co-immunostaining in cultured WT and KO hippocampal neurons. Given the very low transfection efficiency of neurons in high density culture, numerous puncta of receptors from adjacent non-transfected neurons were also detected. This approach may contribute to the noisy pattern observed in Figure 3A. Besides, the projections of z-stack for higher magnified dendrites may likely introduced higher background signals. We have now replaced the original images with the newest repeat in new Figure 3A. Moreover, we confirmed a decrease in the surface expression of GABA_AR $\gamma 2$ by the biotinylation assay, as shown in Figure 3E. Indeed, we agree that some puncta for surface labeling of receptors seemed to be non-synaptic localization. In order to reflect the decrease in synaptic proteins at synapses, we isolated PSD fraction by biochemical assay and found that gephyrin and GABA_AR $\gamma 2$, two major inhibitory postsynaptic components, were reduced in the PSD fraction from KO brains, as shown in Figure 3L. Their colocalization was also attenuated in the absence of endophilin A1, as shown in Figure 5A-C. Combined with electrophysiological recording, these data from multiple assays indicate GluA1 at synapses was not obviously affected but GABA_AR $\gamma 2$ at synapses was impaired in endophilin A1 KO neurons in the present study.

We have corrected the way that the number of samples is defined for statistical analysis as suggested. This point was also raised by Reviewer #1 (Recommendation (2)). We averaged the values from all dendritic segments of a single neuron, such that one neuron equaled one data point. We had replaced the original Figure 3B and 3D (please see Response to Recommendation (2) by Reviewer #1). Additionally, we added the number of independent cultures these data were derived from to figure legends in revised manuscript.

Previously, we observed a small decrease in surface GluA1 levels in spines under basal conditions and a more pronounced suppression of surface GluA1 accumulation in spines upon chemical LTP in endophilin A1 KO neurons from *EndoA1*^{-/-} mice that knockout endophilin A1 since embryonic development stages (Figure 5C,H. Yang et al., Front Mol Neurosci, 2018). In Figure 3A and B in current study, we analyzed surface receptor levels in GFP-positive dendrites, rather than spines, under basal conditions when endophilin A1 was depleted at the later developmental stage. We found a decrease in surface GABA_AR $\gamma 2$ levels but no significant effects on surface GluA1 levels in dendrites. These findings indicate that endophilin A1 primarily affects excitatory synaptic proteins in spines during synaptic plasticity and inhibitory synaptic proteins in dendrites under basal conditions in mature neurons.

(4) Super-resolution images in Figure 3G, H, I: endophilin A1 puncta look different in panel 3I compared to 3G and 3H, which are very noisy. It is difficult to interpret how specific these EEN1 puncta are. Previous images showing EEN1 distribution in dendrites look different (Yang et al., Front Mol Neurosci 2018); is the same KO-verified antibody being used here? Colocalization of EEN1 with gephyrin or the GABAAR gamma 2 subunit is difficult to interpret; gephyrin mostly does not seem to colocalize with EEN1 in the example shown.

Sorry for your concerns. As stated previously in Major Points (3), transfection efficiency was very low in cultured neurons and our cultured neurons were at relative high density. As a result, numerous puncta of proteins located in the adjacent non-transfected neurons were also detected, which may contribute to noisy signals observed in Figure 3G-I.

In our previous paper, we confirmed the specificity of the antibody against endophilin A1 (5A,B. Yang et al., Front Mol Neurosci, 2018). We used the same antibody (rabbit anti-endophilin A1, Synaptic Systems GmbH, Germany) in the current study. While the previous images were obtained using confocal microscopy, the current images in Figures 3G, H, and I were acquired using super-resolution microscopy (SIM). The different patterns observed in the dendrites may be attributed to the difference in image resolution, antibodies dilution and reaction time.

Reviewer #1 also points out the quantification of colocalization of gephyrin and GABA_AR γ 2 with endophilin A1. Please see Response to Recommendation (13) by Reviewer #1.

(5) The interaction of gephyrin and endophilin A1 is based on coIP experiments in cells and brain tissue. To convincingly demonstrate that these proteins interact, biophysical experiments with purified proteins are necessary.

Thanks a lot for your great suggestions on the interaction of endophilin A1 with gephyrin. To convincingly demonstrate their interaction, we performed pull-down assay with purified recombinant proteins and the result shows that both G and E domains of gephyrin were involved in the interaction with endophilin A1. The data has been added to the revised manuscript as new Figure 5I. We also modified the statement about the data and figure legends in the revised manuscript.

(6) Figure 4G: the gephyrin images are not convincing; the inhibitory postsynaptic element typically looks somewhat elongated; these puncta are very noisy and do not appear to represent iPSDs. The same criticism applies to the images shown in Figures 5 and 7.

Thanks for the comment. The gephyrin puncta in our images exhibited heterogeneous shapes and sizes, with some appearing somewhat elongated. To address this, we compared the puncta pattern of gephyrin with that shown in the reference. As illustrated in the figure from the reference, gephyrin puncta also displayed distinct shapes and sizes, Figure 3A-F, Neuron 78, 971–985, June 19, 2013). Please note that the images were z-stack projections at higher magnification, as described in the "Materials and Methods" section. This approach may likely introduce higher background signals and may contribute to the much more heterogeneous appearance of the puncta in Figures 4, 5, and 7. As mentioned previously, the numerous gephyrin puncta located in the adjacent non-transfected neurons may also contribute to some of the noisy signals observed. We have replaced the original images with new images in new Figure 4G, 5 and 7.

Moreover, in order to confirm the effects of endophilin A1 KO on the gephyrin clustering, we also detected the endogenous clusters of gephyrin or PSD95 visualized by GPN.FingR-eGFP or

PSD95.FingR-eGFP in cultured mature neurons. The results were consistent with immunostaining with antibodies against gephyrin. Please see Response to Recommendation (2)

(7) Figure 7E, F: the rescue (Cre + WT) appears to perform better than the control (mCherry + GFP) in the PTZ condition; how do the authors explain this? Mixes of viral vectors were injected, would this approach achieve full rescue?

Thanks for the thoughtful comment. Mixed viruses were injected bilaterally into the hippocampal CA1 regions. The results showed a full rescue effect by WT endophilin A1 in knockout mice during the early days, with even a little bit better rescue effect than the control group in the later days under the PTZ condition, as shown in Figures 7E and 7F. In the current study, overexpression of endophilin A1 increased the clustering of gephyrin and GABA_A γ 2 in cultured neurons, as shown in Figures 4I-J and 5D-E. Presumably, the slightly better rescue effects observed in the behavioral tests was likely attributed to the enhanced clustering and/or stabilization of gephyrin/GABA_A γ 2 by WT endophilin A1 expression in KO neurons *in vivo*. Moreover, the electrophysiological recording also showed full rescue effects on eIPSC by WT endophilin A1 in KO neurons (Figure 7G-K).

Minor Points

(1) The authors mention that they previously found a decrease in eEPSC amplitude in EEN1 KO mice (Yang et al., Front Mol Neurosci 2018). The data in Fig. 1E suggests a decrease in eEPSC amplitude but is not significant here, likely due to the small number of observations. If both eEPSC and iEPSC amplitude are reduced in the absence of EEN1. Would the E/I ratio still be significantly changed?

We apologize for the confusion. In our previous study, AMPAR-mediated excitatory postsynaptic currents (eEPSCs) were found to be slightly but significantly reduced compared to the control group, while NMDAR-mediated excitatory postsynaptic currents showed no significant difference (Figure 4N,O. Yang et al., Front Mol Neurosci, 2018). In the current study, we adopted a different recording protocol, simultaneously measuring eEPSCs and eIPSCs from the same neuron to calculate the E/I ratio. Unlike previous studies, we did not use inhibitors to suppress GABA receptor activity. As a result, the recorded signals did not distinguish AMPAR-mediated or NMDAR-mediated excitatory postsynaptic currents to reflect total eEPSCs, which may explain the non-significant reduction observed compared to control neurons in this study.

It is possible that the eEPSC amplitude would show a significant reduction if a larger number of neurons were recorded. Nevertheless, the larger suppression of eIPSCs in the absence of endophilin A1 indicates that the E/I ratio is significantly altered.

(2) Page 7: the authors mention they aim to exclude effects on presynaptic terminals of deleting endophilin A1 in cultured neurons, is this because of a sparse transfection approach?

Please clarify.

Sorry for the confusion. In cultured neurons, we always observed sparse transfection due to the very low transfection efficiency (~ 0.5%). Therefore, we could examine the effects of endophilin A1 knockout specifically in the specific CamKII α promoter-driven Cre-expressing postsynaptic neurons, while endophilin A1 remained intact in the non-transfected presynaptic neurons.

(3) The representative blot of the surface biotinylation experiment (Figure 3E) suggests that loss of endophilin A1 also affects GluN1 and Nlgn2 levels, and error bars in panel 3F (lacking individual data points) suggest these experiments were highly variable.

Sorry for the confusion. Reviewer #1 also raised the question and we quantified the total level of GluN1 and NL2 in Figure 3E. And we replaced the original graphs with scatterplots and means $\pm$ S.E.M. Please see the Response to Recommendation (3) & (12) by Reviewer #1.

(4) Have other studies analyzing inhibitory synapse composition identified endophilin A1 as a component? The rationale for this study seems to be primarily based on the presence of epileptic seizures and E/I imbalance.

Thank you for your questions. To date, no other studies investigated endophilin A1 as an inhibitory postsynaptic component. We observed the proximal localization of endophilin A1 with inhibitory postsynaptic proteins using super-resolution microscopy (SIM) and quantification results showed $\sim 47\%$ puncta of gephyrin correlated with endophilin A1 (Figure 3G-I and S4B-G). We further immunisolated the inhibitory postsynaptic fraction using GABA_A receptors and found that endophilin A1 was present in the isolated fraction, and vice versa (Figure 3J). Additionally, we demonstrated that endophilin A1 directly interacted with gephyrin through co-IP and pull-down assays (Figure 5J-I). Together with data from immunolabeling, biochemical assays, electrophysiological recordings, and behavioral tests, these results identified endophilin A1 as an inhibitory postsynaptic component.

(5) Figure 3J: what are S100 and P100 labels? Is Nlgn2 part of the EEN1 complex? If it is, why are Nlgn2 surface levels not affected by EEN1 loss (Figure 3E, F, K)? Why does EEN1 not interact with Nlgn2 in HEK cells (Figure 4D)?

Sorry for the confusion. The detailed information regarding S100 and P100 can be found in the “GST-pull down, co-immunoprecipitation (IP), and immunoisolation” in the “Materials and Methods” section. S100 contains soluble proteins, while P100 refers to the membrane fraction after high speed (100,000xg) centrifugation.

Figures 3J-K and 4C-F showed the data from immunoisolation and conventional co-immunoprecipitation assays, respectively. Immunoisolation, which uses antibodies coupled to magnetic beads, allows for the rapid and efficient separation of subcellular membrane compartments. In Figure 3J-K, we used antibodies against GABA_A $\alpha 1$ to isolate membrane protein complexes from the inhibitory postsynaptic fraction. In contrast, co-immunoprecipitation typically detects direct interactions between proteins solubilized by detergent treatment. For Figure 4C-F, FLAG beads were used in HEK293 lysates, or antibodies against endophilin A1 were employed in brain lysates to precipitate direct interaction partners. Combined with the results from Figure 3J-L, the data in 4C-F indicated that endophilin A1 was localized in the inhibitory postsynaptic compartment and directly bound to gephyrin but not to either GABA_A receptors or Nlgn2 (NL2). This binding promoted the clustering of gephyrin and GABA_A $\gamma 2$ at synapses, facilitating GABA_A assembly.

Nlgn2 (NL2) is a key inhibitory postsynaptic component but does not directly bind to endophilin A1. Consequently, endophilin A1 failed to co-immunoprecipitate with NL2 in the presence of detergent in HEK293 cell lysates (Figure 4D). Furthermore, the surface levels of NL2 or its distribution in PSD fraction were unaffected by the loss of endophilin A1 (Figure 3E, F, K, L). This suggests that mechanisms independent of endophilin A1 orchestrate the surface expression and synaptic distribution of NL2.

(6) How do the authors interpret the finding that endophilin A1, but not A2 or A3, binds gephyrin? What could explain these differences?

Thanks for the thoughtful comment. Endophilin As contain BAR and SH3 domains. While the amino acid sequences in the BAR and SH3 domains are highly conserved, the intrinsically disordered loop region between BAR and SH3 domains is highly variable. A study by the Verstreken lab revealed that a human mutation in the unstructured loop region of endophilin A1 increases the risk of Parkinson's disease. They also demonstrated that the disordered loop region controls protein flexibility, which fine-tunes protein-protein and protein-membrane interactions critical for endophilin A1 function (Bademosi et al., Neuron 111, 1402–1422, May 3, 2023). Our previous study showed that endophilin A1 and A3, but not A2, bind to p140Cap through their SH3 domains, despite the high sequence homology in the SH3 domains among these proteins (Figure2A,B. Yang et al., Cell Research, 2015). These findings indicate that each endophilin A likely interacts with specific partners due to distinct key amino acids.

Additionally, endophilin A1 is expressed at much higher levels than A2 and A3 in neurons, with distinct distribution of them across different brain regions. Our lab demonstrated that the function of A1 at postsynapses (both excitatory and inhibitory synapses) cannot be compensated by A2 or A3. Therefore, it is reasonable that endophilin A1, rather than A2 or A3, binds to gephyrin, even though the underlying mechanisms remain unclear.

| (7) Figure 4G: panels are mislabeled (GFP vs merge).

Thanks for careful reading and sorry for the mistake. We corrected the label in new Figure 4G. Please see Response to Annotation, grammar, spelling, typing errors:(1) by Reviewer #2.

<https://doi.org/10.7554/eLife.102792.2.sa0>