

Assemblies, synapse clustering and network topology interact with plasticity to explain structure-function relationships of the cortical connectome

Reviewed Preprint

v2 • May 22, 2025

Revised by authors

Reviewed Preprint

v1 • November 7, 2024

András Ecker, Daniela Egas Santander, Marwan Abdellah, Jorge Blanco Alonso, Sirio Bolaños-Puchet, Giuseppe Chindemi, Dhuruva Priyan Gowri Mariyappan, James B Isbister, James Gonzalo King, Pramod Kumbhar, Ioannis Magkanaris, Eilif B Muller, Michael W Reimann 

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech, Geneva, Switzerland • Department of Basic Neurosciences, University of Geneva, Geneva, Switzerland • Department of Neurosciences, Faculty of Medicine, Université de Montréal, Montréal, Canada • Centre de Recherche Azrieli du CHU Sainte-Justine, Montréal, Canada • Mila Quebec AI Institute, Montréal, Canada

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **useful** study presents a biologically realistic, large-scale cortical model of the rat's non-barrel somatosensory cortex, investigating synaptic plasticity of excitatory connections under varying patterns of external activations and characterizing relations between network architecture and plasticity outcomes. The model offers an impressive level of biological detail, addressing many aspects of the cellular and network anatomy and properties, and investigating their relationships to the biologically plausible plasticity. The numerical simulations appear to be well executed and documented, providing an excellent resource to the community. The evidence supporting the main conclusions is **solid** with results being more observational in nature, and minor weaknesses relating to the lack of explanatory power of causal relationships and mechanisms.

<https://doi.org/10.7554/eLife.101850.2.sa4>

Abstract

Synaptic plasticity underlies the brain's ability to learn and adapt. While experiments in brain slices have revealed mechanisms and protocols for the induction of plasticity between pairs of neurons, how these synaptic changes are coordinated in biological neuronal networks to ensure the emergence of learning remains poorly understood. Simulation and modeling have emerged as important tools to study learning in plastic networks, but have yet to achieve a scale that incorporates realistic network structure, active dendrites, and multi-synapse interactions, key determinants of synaptic plasticity. To rise to this challenge, we endowed an existing large-scale cortical network model, incorporating data-constrained

dendritic processing and multi-synaptic connections, with a calcium-based model of functional plasticity that captures the diversity of excitatory connections extrapolated to *in vivo*-like conditions. This allowed us to study how dendrites and network structure interact with plasticity to shape stimulus representations at the microcircuit level. In our exploratory simulations, plasticity acted sparsely and specifically, firing rates and weight distributions remained stable without additional homeostatic mechanisms. At the circuit level, we found plasticity was driven by co-firing stimulus-evoked functional assemblies, spatial clustering of synapses on dendrites, and the topology of the network connectivity. As a result of the plastic changes, the network became more reliable with more stimulus-specific responses. We confirmed our testable predictions in the MICrONS datasets, an openly available electron microscopic reconstruction of a large volume of cortical tissue. Our results quantify at a large scale how the dendritic architecture and higher-order structure of cortical microcircuits play a central role in functional plasticity and provide a foundation for elucidating their role in learning.

1 Introduction

Learning and memory are orchestrated by synaptic plasticity, the ability of synapses to change their *efficacy* in an activity-dependent manner. Donald O. Hebb's postulate about how synaptic plasticity might manifest was paraphrased with the well known mantra: “*cells that fire together, wire together*” (Hebb, 1949 [↗](#); Shatz, 1992 [↗](#)). The first proof of coincident pre- and postsynaptic population activity leading to *potentiation* (an increase in efficacy) came from pathway stimulation in hippocampal slices (Bliss and Lømo, 1973 [↗](#)). It was later confirmed at the neuron pair level (Markram et al., 1997 [↗](#); Bi and Poo, 1998), and spike-time dependent plasticity (STDP) became a quintessential protocol to study Hebbian plasticity *in vitro*. In the early 2000s a plethora of cortical pathways were studied and plasticity proved to be synapse location- and therefore pathway-dependent (Sjöström and Häusser, 2006 [↗](#); Letzkus et al., 2006 [↗](#); Froemke et al., 2010 [↗](#)). The molecular substrate of Hebbian coincidence detection is the N-methyl-D-aspartate (NMDA) receptor, which upon removal of the Mg^{2+} block by depolarization, conducts Ca^{2+} (Mayer et al., 1984 [↗](#)). The calcium-control hypothesis, put forward by Lisman (1989 [↗](#)) postulates that prolonged, moderate amounts of Ca^{2+} lead to *depression* (a decrease in efficacy) while large transients of Ca^{2+} lead to potentiation. By putting these together, it became evident that it is not necessarily the timing of the postsynaptic spike, but rather the depolarization induced entrance of calcium into the postsynaptic dendrite that is important to evoke changes in synaptic efficacy (Goldberg et al., 2002 [↗](#); Lisman and Spruston, 2005 [↗](#)).

In parallel with slice electrophysiology, Hebbian plasticity was also studied through its effect on behavior via fear conditioning experiments (McKernan and Shinnick-Gallagher, 1997 [↗](#)) and this line of research led to numerous new techniques for tagging and re-activating cells that participate in newly formed memories (Tonegawa et al., 2015 [↗](#)). While these studies highlighted the need to study plasticity at the network level, most changes are expected to happen at the synapse level. Therefore, high-throughput methods tracking synaptic proteins like PSD95 (Ray et al., 2023 [↗](#)) and α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA) subunit GluA1 (Graves et al., 2021 [↗](#); Kim et al., 2023 [↗](#)) are currently being developed. While readily applicable to monitor synaptic efficacy *in vivo*, currently these techniques cannot be supplemented with recordings of neural activity, thus the reason for the changes in efficacy can only be speculated.

The bridge between *in vitro* pairs of neurons and *in vivo* behavior is often provided by complementary, simulation based-approaches. Early theoretical work explored the potential link between memories and cells that fire and therefore wire together, concentrating on the storage and retrieval of memories in strongly recurrent networks (Hopfield, 1982 [↗](#)), which remained an active topic of research (Fusi and Abbott, 2007 [↗](#); Krotov and Hopfield, 2016 [↗](#); Widrich et al., 2020 [↗](#)). Inspired by STDP experiments, modelers have postulated diverse *plasticity rules* that fit

the most recent experimental findings (Gerstner et al., 1996; Kempter et al., 1999; Song et al., 2000; Pfister and Gerstner, 2006; Clopath et al., 2010). Models that take into account the crucial role of calcium in driving plasticity outcomes have also been proposed (Shouval et al., 2002; Graupner and Brunel, 2012; Rubin et al., 2005; Jedrzejewska-Szmek et al., 2017; Rodrigues et al., 2023). The calcium-based model of Graupner and Brunel (2012) describes the evolution of intracellular calcium concentration ($[Ca^{2+}]_i$) given the pre- and postsynaptic spike trains and updates the efficacy of the synapse, upon $[Ca^{2+}]_i$ crossing thresholds for depression and potentiation. Bio-plausible plasticity rules have been shown to bring about Hebbian cell assemblies, i.e., groups of neurons that fire together to enable long-term memory storage (Litwin-Kumar and Doiron, 2014; Zenke et al., 2015; Fauth and Van Rossum, 2019; Kossio et al., 2021). A common theme in these models is the necessity for homeostatic plasticity to keep the networks stable. While experimental evidence exists for homeostatic plasticity (Turrigiano and Nelson, 2004), this has been shown to be too slow to induce stability (Zenke et al., 2017a) and alternative mechanism have been proposed (Vogels et al., 2011; Delattre et al., 2015). While these studies provide mechanistic explanations of learning and memory, they used point-neuron models, therefore neglecting the structural and functional importance of dendrites in plasticity (with the exception of Bono et al. (2017); Kastellakis and Poirazi (2019)). The compartmentalized nature of dendritic trees gives rise to spatial clustering of synapses (Kastellakis and Poirazi, 2019; Farinella et al., 2014; Iacaruso et al., 2017; Tazerart et al., 2020) and local, non-linear voltage events (Poirazi et al., 2003; Stuart and Spruston, 2015) both of which are thought to contribute to the removal of the Mg^{2+} block from NMDA receptors and therefore gate plasticity.

To investigate the effect of dendrites and multi-synaptic interactions for plasticity at the network level, in this study we equipped the biophysically detailed, large-scale cortical network model of Isbister et al. (2023) with our recently developed model of functional plasticity (Chindemi et al., 2022) between excitatory cells (Figure 1). At this level of detail, a calcium-based plasticity model is the natural choice since calcium dynamics are already part of the simulation. Moreover, it allowed us to model *in vivo*-like conditions, specifically the low extracellular calcium concentration ($[Ca^{2+}]_o$, Chindemi et al., 2022), which has been experimentally shown to reduce plasticity (Inglebert et al., 2020, Figure 1E). As we had access to the pre- and postsynaptic activity and efficacy of millions of synapses including their dendritic location we could characterize the rules governing plasticity at the microcircuit level. During simulated evoked activity, plasticity acted sparsely and specifically, although it was still able to reorganize the network's dynamics, manifesting in more pattern-specificity after plasticity. Potentiation dominated in amplitude and depression counteracted it in frequency, which led to stable firing rates without explicitly introducing any homeostatic terms (Zenke et al., 2017a; Turrigiano and Nelson, 2004). We found plasticity was driven by the network topology beyond pairwise connectivity. Specifically, by co-firing cell assemblies, spatial clustering of synapses on dendrites, and the centrality of a connection in the network. Furthermore, we confirmed our testable predictions in the MICrONS (2021) dataset, an openly available electron microscopic reconstruction.

2 Results

To study how plasticity and various network level features interact to explain structure-function relationships, we used a bio-realistic, large-scale cortical model of the rat non-barrel somatosensory cortex (nbS1). By doing so, we had a continuous readout of both the activity of all neurons and the efficacy of the synapses between them under *in vivo*-like, low $[Ca^{2+}]_o$ conditions (1 mM instead of 2.5 mM *in vitro*). The current model improves on Markram et al. (2015) in terms of both anatomical, e.g., atlas-based cell composition and placement (described in Reimann et al. (2024a)), and physiological properties, e.g., improved single cell models, multi-vesicular synaptic release, and layer-wise compensation for missing synapses (described in Isbister et al.




Figure 1

Overview of the cortical network model and calcium-based plasticity rule.

A: Rendering of the seven column subvolume of rat nbS1. (10% of the cells are shown). **B:** Representative morphologies for the 18 excitatory morphological types (m-types) and their typical firing pattern (top left). Neurite diameters are scaled (x2) for better resolution. **C:** Exemplary connections to layer 5 (thick-tufted pyramidal cells, L5_TPC:A and L5_TPC:B) L5 TTPCs (top) and their short-term dy-namics (bottom). Neurite diameters are uniform for better visibility. On the bottom: thin gray lines represent 20 individual trials, while the thicker black ones their means. **D:** Bouton density profiles of thalamocortical fibers, and locations of VPM (black) and POM (purple) synapses on neurons (in a 5 μm radius subvolume). The spatial scale bar on B applies to the whole figure. Panels adapted from Isbister et al. (2023) [\[1\]](#); Chindemi et al. (2022) [\[2\]](#); Reimann et al. (2024a) [\[3\]](#). **E:** Main variables of the plasticity model during coincident activation of the pre-and postsynaptic neurons for an exemplary synapse. Left: under *in vitro*-like conditions (adapted from Chindemi et al. (2022) [\[2\]](#)). Right: same pair of neurons under *in vivo*-like conditions. Schematics on their lefts illustrate the difference between typical *in vitro* and *in vivo* protocols.

(2023) [\[1\]](#). For this study, we used a seven column subvolume comprising 211,712 neurons in 2.4 mm^3 of tissue (**Figure 1A** [\[1\]](#)). In line with the biological variability, excitatory cells are modeled as a di-verse set of morphologies (Reimann et al., 2024a [\[2\]](#); Kanari et al., 2019 [\[3\]](#)) (**Figure 1B** [\[1\]](#)) equipped with conductances distributed across all electrical compartments (Reva et al., 2023 [\[4\]](#)) (Supplementary Figure S1A). The connectivity and synaptic physiology of these cells were extensively validated (Isbister et al., 2023 [\[5\]](#); Reimann et al., 2024a [\[2\]](#)) (**Figure 1C** [\[1\]](#); Supplementary Figure S1C). To deliver input with spatio-temporal precision to the circuit, we used thalamic fibers from the ventral posteromedial nucleus of the thalamus (VPM) and the high-order posteromedial nucleus of the thalamus (POm; **Figure 1D** [\[1\]](#); Meyer et al., 2010 [\[6\]](#)).

To simulate long-term plasticity, we integrated our recently published calcium-based plasticity model that was used to describe functional long-term potentiation and depression between pairs of pyramidal cells (PCs) (Chindemi et al., 2022 [\[7\]](#)). In short, the model follows the formalism of Graupner and Brunel (2012) [\[8\]](#), where pre-and postsynaptic spikes led to changes in synaptic $[\text{Ca}^{2+}]_i$ (**Figure 1E** [\[1\]](#)). Calcium entering through NMDA receptors and voltage-dependent calcium channels (VDCCs) contributes to $[\text{Ca}^{2+}]_i$ (Equation (2) [\[1\]](#) in Methods). When the integrated calcium trace of a synapse crosses the threshold for depression (θ_d) or the higher one for potentiation (θ_p), synaptic efficacy (ρ) is updated (**Figure 1E** [\[1\]](#) left; Equation (1) [\[1\]](#) in Methods). Changes in ρ are then converted into changes in the utilization of synaptic efficacy (U_{SE}), a variable of the Tsodyks-Markram model of short-term plasticity describing the baseline probability of vesicle release (Tsodyks and Markram, 1997 [\[9\]](#)), and the peak AMPA receptor conductance ($\hat{g}_{AMP A}$; Equations (7 [\[1\]](#)) and (8 [\[1\]](#)) in Methods respectively). As a result of updating U_{SE} , short-and long-term plasticity are tightly coupled in the model (Markram and Tsodyks, 1996 [\[10\]](#); Costa et al., 2015 [\[11\]](#); Deperrois and Graupner, 2020 [\[12\]](#)). In the absence of $[\text{Ca}^{2+}]_i$ influx ρ exhibits bistable dynamics (Lisman, 1985 [\[13\]](#); Graupner and Brunel, 2012 [\[8\]](#)) and at initialization, synapses are assumed to be at one of the two fixed points (fully depressed ones at $\rho = 0$ and fully potentiated ones at $\rho = 1$) and their assignment to these states is pathway-specific (Supplementary Figure S1C3).

We calibrated layer-wise spontaneous firing rates and evoked activity to brief VPM inputs matching *in vivo* data from Reyes-Puerta et al. (2015) [\[14\]](#) (Supplementary Figure S3). Spontaneous activity was driven by somatic injection of a layer-and cell type-specific noisy conductance signal (Isbister et al., 2023 [\[5\]](#), see Methods). Evoked activity was driven by a thalamocortical input stream already described in Ecker et al. (2024) [\[15\]](#). In short, 10 VPM input patterns were repeatedly presented in random order with a 500 ms inter-stimulus interval, together with a non-specific POm input. These VPM patterns were defined with varying degrees of overlap in the sets of activated fibers (**Figure 2A** [\[1\]](#), Supplementary Figure S4; see Methods). An exemplary raster plot, characterizing the evoked state of the plastic network is shown in **Figure 2B** [\[1\]](#).

2.1 Plasticity changes promote stable network activity

In vivo, plasticity does not seem to create macroscopic network instability. Therefore, it is a critical first step to validate our model in this regard. Previous theoretical work has shown that while embedding simple STDP rules in spiking networks without fast homeostatic plasticity leads to exploding firing rates (Morrison et al., 2007 [\[16\]](#); Zenke et al., 2017a [\[17\]](#)), this is not necessarily the case for calcium-based rules (Higgins et al., 2014 [\[18\]](#); Graupner et al., 2016 [\[19\]](#); Chindemi, 2018 [\[20\]](#); Wang and Aljadeff, 2022 [\[21\]](#)). To assess the stability of our network while undergoing plasticity with our specific choice of calcium-based rule, we simulated 10 minutes of biological time of *in vivo*-like, stimulus-evoked network activity and verified that excitatory firing rates remained stable throughout the entire length of the simulation (**Figure 2C** [\[1\]](#)).

When comparing the amount of changes in ρ across time steps, we found that most of the plastic changes happened in the first 1-2 minutes of the simulation, after which they stabilized (**Figure 2D1** [\[1\]](#)). While small changes were still apparent towards the end of the simulation, by visualizing individual synaptic efficacy traces we confirmed that most of them oscillated around a dynamic




Figure 2

Changes in synaptic efficacy stabilize and they promote macroscale stable dy-namics.

A: Top-down view of the spatial arrangement of the VPM fiber centers associated with the 10 input patterns. Bottom row 3^{rd} : pyramid-like overlap setup of VPM patterns, 4^{th} (bottom right) POM fibers (high-order thalamic input) associated with all stimuli. **B:** Raster plot of the microcircuit's activity and the population firing rates below. The y-axis shows cortical depth. As cortical layers do not have the same cell density, the visually densest layer is not necessarily the most active. Panel adapted from Ecker et al. (2024). **C:** Firing rate of excitatory cells during the 10-minutes-long simulation. **D:** Plastic evolution of synaptic efficacy (p): L2 norm of changes in p across time. Inset shows the distribution of p values in the beginning (green; strictly 0s and 1s, see Methods) and end (purple) of the simulation. **E:** Evolution of mean p (aggregated over multiple synapses under a single connections). **E1:** L2 norm of changes in mean p (black) across time against STDP control in pink (inset, see Methods). **E2:** L2 norm of changes in p (similar to E1, but) compared to $p_{t=5\text{minutes}}$ in black, STDP in pink, and their random walk controls (see Methods) with the same colors but dashed lines. **F:** Firing rates of excitatory cells from a simulation with higher $[\text{Ca}^{2+}]_o$ and thus population bursts (see Supplementary Figure S6, Markram et al., 2015, and Methods) in gray and the baseline one in red as on panel C. **G:** Grand average mean p values across time for the synchronous simulation in gray and for the corresponding STDP control in pink. **H:** Same as E2 but for the synchronous simulation (calcium-based plasticity rule in gray instead of black).

fixed-point and the amount of changes in the second half of the simulation were negligible (Chindemi, 2018 [↗](#), Supplementary Figure S5). The results were similar (Figure 2E1 [↗](#), black line) when we averaged ρ values within a connection, i.e., all synapses between a pair of neurons (on average 4.1 2.3 synapses per connection; Supplementary Figure S1C1). Considering changes at the connection level allowed us to compare our results against a traditional spike pair-based STDP rule (Gerstner et al., 1996 [↗](#); Kempter et al., 1999 [↗](#); Song et al., 2000 [↗](#), see Methods).

As it is evident from previous theoretical work that simply equipping networks with pair-based STDP rules leads to exploding firing rates due to a positive feedback loop (Morrison et al., 2007 [↗](#); Zenke et al., 2017a [↗](#)), we turned to a stricter control and executed the STDP rule on the 35 M excitatory spikes from our simulation. We observed that the STDP rule kept inducing the same magnitude of changes throughout all 10 minutes of the simulation instead of stabilizing after a transient phase (Figure 2E [↗](#), pink line). To further assess whether the remaining changes in mean ρ were in the form of fluctuations around a dynamic fixed-point, we compared the changes in the second half of the simulation to a random walk with the same step size (see Methods). The magnitude of changes were well below the random walk control for our simulation, while the changes from the STDP rule were above both, but below a random walk with a step size fitted to the changes induced by the STDP (Figure 2E2 [↗](#), see Methods). Consequently, the lifetime of synaptic efficacy configurations in the network are longer when using a calcium-based rule than an STPD rule or a random walk. This demonstrates that not only the scale of the simulated network, its biorealistic connection probabilities and the low, *in vivo*-like rates contribute to the stability of changes observed in our simulation, but also the use of a calcium-based plasticity model (Chindemi et al., 2022 [↗](#)) in line with previous modeling insights (Higgins et al., 2014 [↗](#)).

Next, we went beyond keeping a stable network stable and investigated if an unstable network exhibiting synchronous activity (Figure 2F [↗](#), left) could be brought back to a stable, asynchronous regime by the calcium-based plasticity rule. The synchronized activity state was the result of simulating a higher $[Ca^{2+}]_o$ and thus higher release probability (U_{SE} , see Methods). Surprisingly, the firing rate flipped from synchronous to asynchronous (Figure 2F [↗](#), right) and it did so within a minute of biological time (Supplementary Figure S6B). This was likely driven by a sudden decrease in the mean ρ values over synapses in the network, which decrease could not be observed for the STDP rule (Figure 2G [↗](#)). As before, both rules led to smaller changes than their corresponding random walk controls (Figure 2H [↗](#)). Taken together, our results indicate that the calcium-based rule evolves an over-excitable network towards lower excitability, which in this case is a role that a homeostatic rule would play (Turrigiano and Nelson, 2004 [↗](#); Zenke et al., 2017a [↗](#)).

In summary, the calcium-based plasticity rule promotes stability in our network. Indeed, when applied to asynchronous activity, it generates changes in synaptic efficacy that stabilize after a transient phase without leading to exploding firing rates. Furthermore, when applied to an unstable synchronous activity regime, the mean synaptic efficacy of the whole network consistently decreases making the network less excitable and fosters stability in the macroscopic activity.

2.2 Plasticity induces sparse and specific changes driven by the stimulus and the network's topology

We have shown that the changes in synaptic efficacy stabilize after a transient period, which raises the question: What predictions does the model make about the structure of the plastic changes *in vivo*? First of all, the changes were sparse, implying that the configuration of synaptic efficacy of the overall network remained largely unchanged. Indeed, we found that the number of connections changing was below 18% across the whole network and remained low when restricted to individual layers (Figure 3A1 [↗](#)). Moreover, the propensity of changes increased as the pairwise firing rates increased (Figure 3A2 [↗](#)), in line with previous modeling insights (Litwin-

Kumar and Doiron, 2014 [\[14\]](#); Graupner et al., 2016 [\[15\]](#)). On the level of individual synapses changes were even sparser (**Figure 3B1** [\[16\]](#)), and in both cases depression was more common than potentiation. Layer 5 (L5) PCs contributed mostly to changes on the basal dendrites, while apical changes happened mostly on L6 PCs (**Figure 3B2** [\[17\]](#)).

So far, we characterized plastic change in terms of ρ , as this parameter is bounded to the $[0, 1]$ interval and thus easy to interpret across pathways. Another parameter affecting the function of the synaptic connection is $\hat{g}_{AMP A}$. We found a minimal decrease (-2 pS, **Figure 3C1** [\[18\]](#)) in its mean value due to plasticity, which was explained by frequent depression and a heavy tail of potentiation amplitudes (**Figure 3C2** [\[19\]](#)). The distribution of $\hat{g}_{AMP A}$ remained lognormal, in line with biology (Buzsáki and Mizuseki, 2014 [\[20\]](#); Rößler et al., 2023 [\[21\]](#), **Figure 3C1** [\[22\]](#), bottom). The plasticity model employed was bistable around $\rho_* = 0.5$, i.e., in the absence of further Ca^{2+} influx, values above 0.5 would converge towards 1 and values below towards 0 (see Graupner and Brunel, 2012 [\[23\]](#) and Methods). The fraction of changes in ρ that crossed this threshold was slightly higher for depression (26%) than for potentiation (23%) (compare **Figure 3D1** [\[24\]](#) to B1). Furthermore, the vast majority of these crossings occurred early, with 60% of potentiation and 40% of depression happening within the first minute (**Figure 3D2** [\[25\]](#)).

Additionally, we hypothesized that both the amount, i.e., the level of sparsity, and the structure of plastic changes are non-random and instead shaped by the stimuli and the underlying network's topology. We confirmed our hypothesis first for the amount of plastic changes. We ran control simulations where we delivered random Poisson spikes on the same VPM fibers with the same mean rate, but without the spatio-temporal structure of the stimuli. We found that this reduced the number of connections undergoing plasticity by 25%, demonstrating the importance of stimulus structure over simple firing of pre- and postsynaptic neurons (Supplementary Figure S7). Additionally, without synaptic transmission, plasticity was reduced even further: In simulations of stimulus streams where intrinsic connectivity between the simulated neurons was cut we found occasional changes in ρ , but the number of changing synapses was an order of magnitude below baseline conditions (Supplementary Figure S7). Thus, our plasticity model is not strictly Hebbian, since the effect of postsynaptic firing alone could change synaptic efficacy, although presynaptic release was required for most of the observed changes (Graupner and Brunel, 2012 [\[23\]](#); Graupner et al., 2016 [\[15\]](#)). When the external stimuli were also left out, negligible plastic changes occurred (Supplementary Figure S7).

We then confirmed that the network's topology shaped plastic changes as well. In particular, we found layer-to-layer pathway specificity in the amount of plastic changes observed. That is, the amount of changes between layers differed from those expected in a random control with the same pre- and postsynaptic populations (**Figure 4A2** [\[26\]](#) vs. A3). We further quantified the difference in the subnetwork of changing connections from the random control by counting *directed simplices* (**Fig 4B** [\[27\]](#)). These are motifs that have previously shown to be linked to network function (Reimann et al., 2017 [\[28\]](#)) as well as quantify the complexity of the network's topology (Kahle, 2009 [\[29\]](#); Bobrowski and Kahle, 2018 [\[30\]](#)). Succinctly, a directed simplex of dimension k , is a motif on $k + 1$ neurons with all-to-all feed-forward connectivity (**Figure 4A1** [\[31\]](#) right, see Methods). We found strong overexpression of simplices in the subgraph of changing connections compared to the control (**Figure 4B** [\[32\]](#)). Furthermore, the maximal simplex dimension found in the subgraph exceeded the value expected by chance by two. This result shows that the connections changing are not determined only by their pre- and postsynaptic populations but also depend on their embedding in the whole network. We will explore this further in **Section 2.5** [\[33\]](#).

Finally, we showed that the input stimuli shaped plastic changes by running 2-minute-long plastic simulations in which we only presented a single pattern (several times, with the same 500 ms inter-stimulus interval as before) and compared the changes in mean ρ matrices. By using an input - output distance correlation analysis, we found that the distances between these mean ρ matrices were highly correlated with the input distances of the associated VPM fiber locations ($r =$




Figure 3

Changes in synaptic efficacy are sparse.

A1: Propensity of changes in connections (mean p values) across layers. Connections are grouped into layers based on the soma location of the postsynaptic cell. Percentages are calculated within layers, i.e., 7.4% depression (in blue) for layer 2 (L2) means that among all connections that have the postsynaptic cell in L2, 7.4% depressed. **A2:** Plastic changes in mean p vs. mean pairwise firing rates of the pre- and postsynaptic cell of a given connection. **B1:** Same as A1 but for p (changes in synapses rather than connections). **B2:** Layer- and neurite type-wise (apical and basal dendrites) distribution of Δp . Only synapses that underwent any plastic change (15M) are shown. **C1:** Distribution of $\hat{g}_{AMPA}$ in the beginning (in green) and end (in purple) of the simulation. **C2:** Plastic changes in $\hat{g}_{AMPA}$ leading to the slight shift in the distribution on C1. **D1:** Same as B1, i.e., propensity of changes at the synapse level, but for synapses that cross the $p_{*} = 0.5$ unstable fixed-point. **D2:** Cumulative histogram of p crossing the unstable fixed-point against time for depressing synapses in blue and potentiating ones in red.




Figure 4

Changes in synaptic efficacy are non-random and shaped by the networks topology and the input stimuli.

A: Layer-wise distribution of changing connections. **A1:** Left: schematic of the *changing subgraph*. In black, connections whose efficacies are changing and their pre-and postsynaptic populations. In grey, connections whose efficacies do not change and neurons not partaking in a changing connection. Right: random control of the changing subgraph, generated by randomly selecting the same number of connections as the changing connections (red edges) between the same pre-and postsynaptic populations (black nodes). **A2:** Sankey plot of layer-wise distribution of changing connections. Thickness of lines is proportional to the number of changing connections between pre-and post node populations. **A3:** As A2, but for the random control (A1 right). **B:** Directed simplex counts in the changing subgraph and its random control. **B1:** Schematic of directed k-simplices and their counts in the schematic graphs on A1. Note that by construction the control must have the same number of 0-and 1-simplices (see Methods) which correspond to the number of cells and connections in the subnetwork. **B2:** Directed simplex counts across dimensions in the changing subgraph (black) and its random control (red). **C:** Input - output distance correlations. **C1:** Input distances as the Earth mover's distance of the VPM fiber locations (see Figure 2A). Inset shows the overlap (based on Hamming distance, see Methods) of pattern fibers. **C2:** Output distances as Euclidean distance of mean p matrices in the 2-minute-long single pattern simulations. To its right: correlation of input and output distances. **C3:** Same as C2 but with Hamming distance. **C4:** Same as C2 but with Earth mover's distance (on the output side as well).

0.716, $p < 0.0001$, **Figure 4C2** [↗](#), Supplementary Figure S8A). Furthermore, this required using Euclidean distance between the ρ matrices on the output side, implying that both the mean ρ values themselves and their position in the network are relevant. When either Hamming distance (taking only the identity of changing connection into account) or Earth mover's distance (taking only the distribution of mean ρ values into account) were used instead, we found weaker and non-significant correlations respectively (**Figure 4C3 and C4** [↗](#) respectively).

In summary, we observed that 7% of synapses undergo long-term plasticity under realistic *in vivo*-like conditions in 10 minutes of biological time, and most of these synapses are on L5 PC's basal dendrites. Moreover, the changes are not random, but depend not only on the firing rates but also on the recurrent connectivity and input stimuli. Potentiation dominated in amplitude, while depression counteracted it in frequency, keeping the network stable without needing to model homeostatic plasticity (Zenke et al., 2017a [↗](#); Turrigiano and Nelson, 2004 [↗](#)).

2.3 More frequent plastic changes within and across cell assemblies

We have shown that plastic changes are sparse and highly specific. We therefore tried to understand the rules determining which synapses changed. From the parametrization of the plasticity model, we learned that presynaptic spikes contribute orders of magnitude higher $[Ca^{2+}]_i$ than postsynaptic ones if the NMDA receptors are fully unblocked (Supplementary Figure S2B). Thus, in order to effectively depolarize the dendrites and unblock NMDA receptors, spikes at low, *in vivo*-like rates must be synchronized in time. Therefore, we specifically hypothesized that plasticity of connections may be structured by the membership of participating neurons in Hebbian cell assemblies (Hebb, 1949 [↗](#); Harris, 2005 [↗](#)). In our previous analysis (Ecker et al., 2024 [↗](#)), and in line with experimental results (Harris, 2005 [↗](#); Song et al., 2005 [↗](#); Perin et al., 2011 [↗](#)), we observed that the number of afferents from an assembly, is a great predictor of a neuron's membership in an assembly. Strong positive interactions were also found *across assemblies*, but only when the direction of innervation reflected the temporal order of assembly activation. These results, combined with the biophysics of the plasticity model, suggest that connections within an assembly and the ones between temporarily ordered assemblies, are expected to undergo plastic changes with a higher probability.

To test this hypothesis, we detected cell assemblies, based on their co-firing function, from the *in silico* spiking activity of the 10-minute-long plastic simulation using methods established by experimentalists (Carrillo-Reid et al., 2015 [↗](#); Herzog et al., 2021 [↗](#), see Methods). In short, spikes were binned and bins with significantly high firing rates were hierarchically clustered (**Figure 5A** [↗](#)). These clusters correspond to the functional assemblies of neurons (**Figure 5C** [↗](#)), with a neuron being considered a member if its spiking activity correlates with the activity of an assembly significantly stronger than chance level. Since time bins and not neurons, were clustered in the first place, this method yields one assembly per time bin and single neurons can be part of several assemblies (**Figure 5B** [↗](#), D). To foster effective discussions in the field Miehle et al. (2023) [↗](#) urged to call co-firing neurons, detected from purely functional activity *ensembles*, and strongly interconnected ones - usually by design in modeling studies - *assemblies*. While we agree with the distinction, we prefer to call the groups of neurons used for analysis in the study *functional assemblies* as a mix of the two terms, since we have shown in our previous investigation using the same circuit model that they not only co-fire but are interconnected more than expected (Ecker et al., 2024 [↗](#)). Note, that in contrast to classical modeling studies, where assemblies are usually defined based on their strong internal connectivity (i.e., their structure Litwin-Kumar and Doiron, 2014 [↗](#); Zenke et al., 2015 [↗](#); Fauth and Van Rossum, 2019 [↗](#); Kossio et al., 2021 [↗](#)) our functional assemblies are detected from the activity of the network only, eliminating a potential bias arising from links between structural properties of changing synapses and the assembly detection method.




Figure 5

Changes are more frequent within and across cell assemblies.

A: Schematics of the assembly detection pipeline. Significant time bins are clustered by cosine similarity and these clusters define functional assemblies: olive, light blue, steel blue (see Methods). Bottom right, visual summary of the organization of pattern-specific assembly sequences of the two repetitions of pattern A. **B:** Activation of functional assembly sequences for multiple repetitions of the patterns presented (as in A bottom right). Each row within the 10 matrices corresponds to a single repetition of a given pattern. White: non-significant time bins. **C:** Number and location of neurons in each cell assembly. Constituent neurons are those for which the correlation of their spiking activity within the corresponding time bins is significantly stronger than chance (see Methods). Top: top-down view, bottom: depth-profile. **D:** Jaccard similarity (intersection over union) between cell assemblies. **E:** Initial mean p of within and cross-assembly synapses. **F:** Propensity of depression and potentiation of within and cross-assembly synapses in blue on the left and in red on the right respectively. Since assemblies are overlapping (see D) some synapses are counted in multiple pre- and postsynaptic assembly pairings. **G:** Firing rate-based random controls of changes within and across cell assemblies. **G1:** Firing rate distributions of cell assemblies and their corresponding firing rate matching random controls. Last gray box depicts rates of non-assembly (non-A) neurons. **G2:** Same as F above, but for rate-based controls of assemblies. Color bars are the same as in F for easier comparison.

Functional assemblies were activated in all stimulus repetitions and a series of three to four assemblies remained active for 190–30 ms (**Figure 5B**). Pattern C elicited the strongest response, while pattern H the weakest, and the responses of patterns H and I were the most similar to each other, as expected, since they share 66% of the VPM fibers (**Figure 2A**). Functional assembly activations had a well-preserved temporal order - with some of them always appearing early during a stimulus, while others later - and from now on we will refer to them as *early*, *middle*, and *late assemblies*, and will order them in the figures accordingly.

When checking the propensity of changes within and across functional assemblies, we indeed found more synapses undergoing long-term plasticity (**Figure 5F**). While only 5% of synapses depressed in the whole dataset, we found up to 13.8% when restricting the analysis to assemblies. Similarly, compared to 2% of all synapses potentiating, we observed up to 5.2% when restricting to functional assemblies. Interestingly, large values were found in the off-diagonal entries (**Figure 5F**), i.e., synapses across assemblies underwent more plastic changes than the synapses within these assemblies. In our model, the initial ρ values are pathway-specific and highest in L4 pathways (Brémaud et al., 2007, Supplementary Figure S1C3). Therefore, early assemblies, with large numbers of L4 cells have a higher than average initial ρ (**Figure 5C and E** respectively), thus their synapses are more likely to depress (**Figure 5F** left). As early assemblies are stimulus specific, and thus not part of the same assembly sequences, synaptic depression between these cells can be seen as some kind of stimulus separation. On the other hand, late assemblies, that are predominantly composed of cells from the deep layers, have a low initial ρ (**Figure 5E**; Supplementary Figure S1C3) and synapses towards them are more likely to potentiate (**Figure 5F** right). These functional assemblies are mostly non-specific and participate in all assembly-sequences, thus the potentiation of their efferents means a general strengthening of the stimulus response as a whole.

We have shown above that connections between pairs of neurons with high firing rate are more likely to undergo plastic changes (**Figure 3A2**). Thus, we were wondering if firing rate alone explains the increased propensity of changes we observed when restricting our analysis to assembly synapses. To this end, for each assembly, we randomly sampled the same number of neurons with the same firing rate distribution from the whole circuit (**Figure 5G1**) and studied the changes of their synapses. We observed fewer synapses undergoing plastic changes and also a loss of specificity in these rate-based controls (**Figure 5G2**, Supplementary Figure S9A). In conclusion, co-firing in a Hebbian cell assembly is a better predictor of plastic changes than just firing at a high rate.

Together these results indicate that, in line with 70-years-old predictions, cells that fire together wire together (Hebb, 1949). Our contribution lies in making the qualitative statement above into a quantitative one: Under *in vivo*-like conditions, cells that fire together more than expected have three times higher chances of changing the efficacy of their synapses.

2.4 Synapse clustering interacts with assembly formation and plastic changes in complex ways

In addition to co-firing, a group of presynaptic neurons is more effective in depolarizing a given dendritic branch if they all innervate the same branch, i.e., they form a spatial synapse cluster (Kastellakis and Poirazi, 2019; Farinella et al., 2014; Iacaruso et al., 2017; Tazerart et al., 2020; Kirchner and Gjorgjieva, 2021, 2022). To quantify this, we selected the 10 most innervated L5 TTPCs (thick-tufted pyramidal cells) within a cell assembly and then detected spatial clusters of synapses, defined as at least 10 synapses within a 20 μm stretch of a single dendritic branch (Methods; 15k clustered synapses per assembly on average). Next, we grouped all synapses on these 10 selected neurons per assembly into four categories based on assembly membership of the presynaptic neuron and whether the synapse was part of a cluster or not (see exemplary clustered assembly synapses in **Figure 6A**). While some of the ρ values of clustered synapses

within an assembly, underwent small constant changes, many of them changed at the same time (vertical stripes on **Figure 6B**); indicating the importance of spatial co-localization for plasticity. To systematically quantify this effect, we calculated the likelihood of plastic change in each category: (not) in an assembly and (non-)clustered, by contrasting the conditional probability of observing it in a given category with the probability of observing any change irrespective of the category (see **Equation (15)** in Methods; **Figure 6C**). Surprisingly, clustered synapses within an assembly were only likely to undergo plastic changes in the larger, but less general late assembly, A11, but not in early and middle assemblies (**Figure 6C** black arrows top). In fact, the only category that was likely to change across all assemblies is that of non-clustered within-assembly synapses. Therefore, we concluded that synapses within the other three categories tend to keep their initial arrangement of ρ values. To study the initial arrangements as well, we repeated the same analysis on the probability of the initial value of ρ to be either 0 or 1. Initial ρ values in early and middle assembly synapses, especially the clustered ones, were very likely to be initialized as fully potentiated, while synapses within the late assemblies were likely to be initialized in the fully depressed state (**Figure 6D** black arrows bottom). However, when comparing not the likelihood, but the amplitude of changes across categories with a 2-way ANOVA, we found that clustered, within-assembly synapses depress to a significantly smaller degree than all other categories (Supplementary Figure S10A). In contrast, non-clustered, within-assembly synapses (the ones that are most likely to change) depress the most. On the other hand, non-assembly synapses usually tended to be initialized depressed and to stay that way. Thus the picture emerging is as follows: early and middle assemblies are partially defined by clustered (both spatial and functional) synapses that are initialized as fully potentiated. These synapses are unlikely to change, but when they do, they depress less than the others, and would converge back to $\rho = 1$ in absence of activity, as they do not cross the $\rho_* = 0.5$ unstable fixed-point.

These stable early assemblies can therefore function as a stable backbone amid ongoing synaptic plasticity.

In our previous investigation, we found that most changes happened across assemblies (**Figure 5F**), so we extended the analysis described above to cross-assembly synapses (Supplementary Figure S10B). Here, the picture was reversed: cross-assembly synapses that were part of a spatial cluster were likely to be initiated as fully depressed and then had a high chance of undergoing potentiation (Supplementary Figure S10C black arrow). Together with the previous results, this suggests that synapses between assemblies are more likely to change, which is even more pronounced if these synapses form a cluster on the postsynaptic dendrite.

2.5 Network-level metrics predict plasticity and separate potentiation from depression

So far, we have found co-firing and synapse clustering to be good predictors of plastic changes, in line with previous work (Zenke et al., 2015; Kastellakis and Poirazi, 2019; Harris, 2005). Additionally, we have shown that changes are also driven by the underlying network's connectivity (**Figure 4A**, B). We now unfold the effect of network connectivity on plastic changes further, by studying the effect of the presynaptic population of the source (presynaptic) and target (postsynaptic) neurons of a connection on the ρ dynamics. Note first, that the indegree of a neuron, i.e., the size of its source population, is weakly correlated to its average firing rate (**Figure 7A**; $r = 0.13$). Furthermore, the indegree of the pre- and more strongly of the postsynaptic neurons of a connection are predictive of plastic changes (**Figure 7B1**). However, just like for assemblies, a purely rate-based control highlighted that the picture is more complex, firing rate alone is not the best predictor of plastic changes (Supplementary Figure S9B).

Moreover, the correlation of the pre- and postsynaptic indegree ($r = 0.18$) complicates the picture (**Figure 7B2**). Thus, we took the natural choice of considering common innervation (see below), instead of making a choice of how to combine independent innervation of both neurons mediating the




Figure 6

Interactions between plastic changes and synaptic clusters.

A: Changes in clustered assembly synapse on an exemplary neuron from assembly 11 (A11). Neurite diameters are scaled (x2) for better resolution. (Synapse diameters are arbitrary.) **B:** Temporal evolution of the synapses on basal dendrites shown on A. Synapses are grouped by their dendritic sections (single stretches of non-branching dendrites) and ordered by their mean distance to the soma (closest on the bottom and furthest on top). **C:** Michelson contrast (Equation (15) in Methods) of the likelihood of plastic changes of synapses grouped into four categories: (non)-assembly and (non)-clustered for each assembly. Negative values indicate that synapses belonging to the given category are unlikely to change. Depression on top (in blue) and potentiation below (in red). Black arrows indicate clustered assembly synapses. **D:** Same as C but for initial p values (see Methods).




Figure 7

Changes are stronger in central connections in the network.

A: Firing rate vs. indegree of all excitatory neurons ($r = 0.13$). **B:** Probability of plastic change vs. indegree. **B1:** Separately for indegree of the pre- and postsynaptic neurons. **B2:** Joint probability of change against the indegrees of both partners of the connection. **C1:** Schematics of k-edge indegree across dimensions ($k = 0, 1, 2$) and its difference from the classical (node) indegree. TODO: add something about colors. **C2:** Probability of changes (i.e., either depression or potentiation) vs. k-edge indegree in high-dimensional simplices (see C and Methods). **C3:** Probability of depression (left) and potentiation (right) conditioned on plastic change (in any direction) vs. k-edge indegree. **D:** Comparison with electron microscopy data. Total AMPA conductance ($\hat{g}_{AMP A}$) against k-edge indegree across dimensions in our nbS1 model on the left. Total synapse volume (see Methods) against k-edge indegree across dimensions in the MICRONS (2021) dataset on the right. Shaded areas indicate SEM. **E1:** Schematics of k-edge indegree in high-dimensional simplices in an exemplary “full” network on the top in black and in (assembly) “sub”-graphs on the bottom. Table summarizes k-edge indegree values for the connection from node u to v across dimensions for all three cases. **E2:** Probability of changes against k-edge indegree in assembly subgraphs (see C2 bottom). Left, probability of changes in (2-minutes-long) pattern A simulations against k-edge indegree in assembly subgraphs. Right, summary of the maximum probability values across patterns on the right. Arrow indicates the row shown in detail on its left and white crosses indicate the pattern-specific early assemblies (see Figure 5B).

connection. Additionally, it has been shown that not only the size of the presynaptic population but also its internal connectivity affect co-firing and assembly formation [Ecker et al. \(2024\)](#). We explore these effects using *k*-edge indegree, a novel yet simple network metric of edge centrality that considers how the edges (connections) are embedded in the entire network. More precisely, the *k*-edge indegree of a connection is the number of *k*-simplices that the connection is innervated by, where we call *k* the dimension (**Figure 7C1**, see Methods). In particular, for *k* = 0 it is the number of neurons innervating both the pre- and post-synaptic neurons of the connection, i.e., a common neighbor as in [Perin et al. \(2011\)](#). Indeed, for our circuit, in the lower dimensions, *k*-edge indegree is correlated to the indegree of its source and target populations (Supplementary Figure S11A; *r* = 0.48, 0.35 for *k* = 0 and *r* = 0.35, 0.27 for *k* = 1). We expected these motifs to have an effect on plasticity for two main reasons. First, pairs of connected neurons have higher spike correlations if the connection has higher *k*-edge indegree ([Reimann et al., 2017](#), [2024b](#)) and the strength of this correlation depends on the overall higher-order structure of the network ([Nolte et al., 2020](#)). Second, simplex motifs innervating a cell have been shown to be a good predictor of assembly membership ([Ecker et al., 2024](#)), which in turn have an effect on plasticity as demonstrated above.

Across dimensions, as *k*-edge indegree increased so did the probability of undergoing plastic change (**Figure 7C2**). Furthermore, among the connections that change, the ones with the highest *k*-edge indegree tend to get potentiated while those with the lowest favor depression (**Figure 7C3**). Thus, we hypothesized *k*-edge indegree to be a good predictor of synaptic strength. We verified this both in our model as well as in the MICrONS (2021) mm³ dataset, an electron microscopic reconstruction of cortical tissue that combines synaptic resolution with the scale needed to calculate meaningful *k*-edge indegree. Both networks had similar maximal dimension of simplices as well as ranges of *k*-edge indegree values (Supplementary Figure S11B), making the results of this analysis comparable. For our model we used the sum of $\hat{g}_{AMP A}$ values as a proxy for connection strength. For MICrONS, we used total synapse volume of a connection (see Methods), as a measure of connection strength ([Harris and Stevens, 1989](#)). Indeed, high *k*-edge indegree led to higher connection strength in MICrONS (2021) as predicted by our modeling insights; moreover, the curves across dimensions are qualitatively the same for both datasets (**Figure 7D**).

To gain insights into pattern-specificity, we analyzed the 2-minutes-long single pattern simulations and considered *k*-edge indegree within the assembly-specific subgraphs (**Figure 7E1-Sub**). When doing so, we found an even higher predictive power, which was additionally pattern specific (**Figure 7E2**). More precisely, the probability of changes only increased with *k*-edge indegree in the assembly sub-graph associated with the presented pattern or to a late assembly and that it was the strongest for the former (see **Figure 7E2** left for pattern A for an example, right for a summary of the maxima attained for all patterns, and Supplementary Figure S8B for consistency across repetitions of the same simulation).

Finally, *pattern-indegree*, i.e., the number of VPM fibers belonging to a pattern that innervate a neuron is also a good predictor of the probability of a connection to change i.e., an increase of either the pre- or postsynaptic pattern-indegree of a connection leads to more frequent plastic changes (Supplementary Figure S11C1). However, given the highly non-random structure of the full network, the values of pre- and postsynaptic pattern-indegree are not independent and thus their effect in driving plastic changes is hard to disentangle (Supplementary Figure S11C2, D; *r* = 0.230. on average).

In summary, *k*-edge indegree in the full network allowed us to distinguish potentiation from depression given change, as well as predict connection strength, which we confirmed in the MICrONS (2021) dataset. Moreover, *k*-edge indegree within assembly subgraphs is a pattern specific predictor of plastic changes.

2.6 Increased stimulus-specificity and more reliable assembly sequences characterize the network after plasticity

As the changes are sparse, one might wonder if they cause any functional change. To study this, we compared the network's activity *before* and *after* plasticity (see Methods). Firing rates decreased slightly as a result of plasticity (**Figure 8A** [left](#)), while spike correlations remained stable, in line with recent findings (Oby et al., 2019 [left](#); Feulner et al., 2022 [left](#), **Figure 8A** [right](#)). Both of these effects were more pronounced in the deeper layers (Supplementary Figure S12A). Correlations of connected pairs of neurons decreased slightly after plasticity following the decrease of firing rate (Supplementary Figure S12A). However, the highest correlations both before and after plasticity were found in the most central edges in the network i.e., those with highest edge participation in high dimensional simplices (**Figure 8B1** [left](#)). We further verified this prediction in the MICrONS dataset (**Figure 8B2** [left](#), see Methods). It has been previously shown that assemblies have many edges that are central in the network, therefore, we hypothesized that they remain functional and reliably active (Fauth and Van Rossum, 2019 [left](#); Kossio et al., 2021 [left](#); Pérez-Ortega et al., 2021 [left](#)). We showed this on the level of assembly sequences: plastic changes promote stimulus specific and reliable responses. While we found considerable overlap between assemblies detected before and after plasticity, an early assembly responding to patterns B and E before plasticity, split into two stimulus-specific assemblies (**Figure 8C** [left](#)). Note that this is not an artifact of the thresholding applied during assembly detection as forcing the pipeline to yield the same number of assemblies instead splits a middle assembly first (Supplementary Figure S12E). Moreover, the assembly sequences responding to repetitions of a given pattern generally became more reliable after plasticity (Supplementary Figure S12B). To quantify this, we measured the Hamming similarity of assembly sequences between repetitions and found significant increases for most patterns (see Methods, **Figure 8D** [left](#)). Interestingly, this increase on the assembly level lied in contrast with a slight decrease of the reliability of the neuronal activity on the single cell level (**Figure 8E** [left](#)).

The plastic changes we observed at the network-level are in contrast with *in vitro* observations at the connection level. In particular, with the redistribution of synaptic efficacy towards earlier spikes during high-frequency firing caused by changes in U_{SE} (Chindemi et al., 2022 [left](#); Markram and Tsodyks, 1996 [left](#); Costa et al., 2015 [left](#); Selig et al., 1999 [left](#); Sjöström et al., 2003 [left](#)), which under our *in vivo*-like, low firing rates and $[Ca^{2+}]_o$ is not in the ranges where this connection-level redistribution is relevant. Specifically, at *in vitro* levels of $[Ca^{2+}]_o$ the potentiated U_{SE} shifts the connection further into the depressing regime, causing the redistribution (**Figure 8F** [left](#)). *In vivo* the lower $[Ca^{2+}]_o$ counters this by moving the connection into the pseudo-linear regime; additionally, the low firing rates make this phenomenon less relevant in general (**Figure 8F** [right](#) and **G** respectively). Thus, while Markram and Tsodyks (1996) [left](#) showed a redistribution of synaptic efficacy after plasticity at the single connection level *in vitro* (**Figure 8F** [left](#)), we found a redistribution at the network level under *in vivo*-like conditions (**Figure 8A-D** [left](#), Supplementary Figure S12A).

In summary, we observed a network-level redistribution of synaptic efficacy that maintained spike correlations globally, while enhancing stimulus-specificity and the reliability of the subpopulations in terms of assembly activation. Furthermore, we derived and confirmed in the MICrONS (2021) [left](#) dataset a prediction stating that the most central connections of the network drive the correlated activity.




Figure 8

Changes promote stimulus specificity.

A: Firing rates and pairwise spike correlations extracted from non-plastic simulations *before* plasticity, i.e., in the naive circuit vs. *after* the 10-minute-long plastic simulation. **B1:** Spike correlation of connected cells vs. k-edge indegree in high dimensional simplices (see Methods) before and after plasticity on the left and right respectively. **B2:** Spike correlation of connected cells vs. k-edge indegree in the MICrONS (2021) dataset (see Methods). Shaded areas indicate SEM. **C:** Jaccard similarity of assemblies detected before vs. after plasticity on the left. Shared early assembly of patterns B and E splitting into two stimulus-specific ones after plasticity on the right. (Indicated by red arrow and rectangle on its left). See more detailed plots in Supplementary Figure S12B. **D:** Reliability of assembly sequences (see Methods and Supplementary Figure S12B). Significance of increases was determined with Kruskal-Wallis test: *: p 0.05, **: p 0.01, ***: p 0.001, ****: p 0.0001. **E:** Spike time reliability (see Methods) of single cells to the different patterns before and after plasticity. (The lack of stars means no significant increases after plasticity.) **F:** Short-term dynamics of an exemplary potentiated L5 TTPC connection before (in green) and after plasticity (in purple). At *in vitro* $[Ca^{2+}]_o$ on the left, and *in vivo* on the right. Thin lines represent 20 individual trials, while the thicker ones their means. **G:** Interspike interval (ISI) distribution of all excitatory neurons before and after plasticity on the left. Zoom in on low ISIs (≤ 50 ms) on the right.

3 Discussion

In this work, we aimed to understand how network structure, function, and dendritic processing interact to drive synaptic plasticity at the population level in the neocortex. Specifically, we used a plasticity model, whose parameters were determined by pairwise STDP experiments but modeled at the level of dendrites. We studied its effect at the population level under *in vivo*-like conditions to determine network-level rules that best predict plastic changes by simulating it in a biophysically detailed model of a cortical microcircuit that was subjected to 10 different stimuli under a variety of protocols. Our principal observations are as follows: (1) Plastic changes were sparse, affecting only 7% of the synapses. A balance between occasional, large-amplitude potentiation and more frequent depression stabilized the network without explicitly modeling homeostatic plasticity. Moreover, the changes were non-random and stimulus specific.

(2) Plastic changes were largely determined by the anatomical structure of synaptic connectivity and its relation to functional units, i.e., changes were most likely between co-firing cell assemblies, at clustered synapses, between neurons highly innervated by the same input, and in central edges in the network. (3) While assemblies remained fairly stable, they become more stimulus specific and reliable which has important implications for coding and learning.

The first observation (1) is quite significant considering that we did not design the learning rule to be sparse and stable. In previous models of plastic networks, changes were not sparse and additional mechanisms were needed to keep the network's activity stable (Litwin-Kumar and Doiron, 2014; Delattre et al., 2015; Zenke et al., 2015; Fauth and Van Rossum, 2019; Kossio et al., 2021; Zenke et al., 2017a; Turrigiano and Nelson, 2004). The machine learning community is also aware of the importance of sparse changes, as in continual learning one has to balance plasticity and stability of existing weights to avoid catastrophic forgetting (McCloskey and Cohen, 1989; Ratcliff, 1990). In recent years, they have come up with impressive techniques that mask or slow down weight updates (e.g., by a regularizer on the loss) to improve the performance of deep neural networks (Zenke et al., 2017b; Kirkpatrick et al., 2017; Mallya and Lazebnik, 2018; Frankle and Carbin, 2019), whereas in our model it emerged without any mechanisms explicitly enforcing it. Note however, that despite the comparison to machine learning, our model is not designed to solve any specific task. Instead it is a model of sensory cortex which only learns to represent stimulus features in an unsupervised manner. Nonetheless, demonstrating the stimulus-specificity of changes as well as an increase of the reliability of assembly sequences, was a crucial validation that our results are not just a description of a biophysical process, i.e., plasticity, but have implications for learning in general.

For our second observation (2), we described three different kinds of plasticity rules that emerge from the underlying plasticity model based on: activity, subcellular structure, and network structure. While all of these are powerful predictors of plastic changes, none of them fully determines them. It is rather the interplay between them (and potentially additional unknown rules) that brings about non-homogeneous changes across the full microcircuit. The first two can be explained from the biophysics of the plasticity model and links our results to the classical work of Hebb (1949) as well as the recent literature on synapse clustering (Kastellakis and Poirazi, 2019; Farinella et al., 2014; Iacaruso et al., 2017; Tazerart et al., 2020). With respect to synapse clustering, we would highlight that our synapses are stochastic and the release probability between PCs is 0.04 at the simulated low $[Ca^{2+}]_o = 1.05$ mM (Markram et al., 2015; Ecker et al., 2020; Jones and Keep, 1988; Borst, 2010). Therefore, care should be taken when comparing our results with either glutamate uncaging experiments, which bypass the presynaptic machinery (Pettit et al., 1997; Losonczy and Magee, 2006), or with other modeling studies that use deterministic synapses (Farinella et al., 2014; Poirazi et al., 2003; Ujfalussy and Makara, 2020). With respect to network-based rules, previous simulation approaches have characterized the

learned connectomes as having an overexpression of reciprocal connectivity and densely connected neu-ron motifs (Brunel, 2016 [↗](#); Zhang et al., 2019 [↗](#)). We expanded on this body of previous work, by introducing cell and connection-specific metrics that directly yield the probability of observ-ing depression or potentiation of any given connection. Most of our predictors behaved similarly with respect to depression and potentiation. Only a metric based on the embedding of individual edges in the global (or assembly-specific) network was able to separate them. This edge-based prediction was confirmed by analyzing an electron microscopic reconstruction of cortical tissue in a comparable way (MICrONS, 2021 [↗](#)).

Our third observation (3) that assembly sequence become more reliable and stimulus specific may sound counter-intuitive considering that we observed reduced firing rates and less reliable single neurons. This shows that plasticity mechanism can contribute to the emergence of a robust population code with unreliable neurons. Crucially, this means that plasticity can make circuit responses more distinct (between stimuli by increasing specificity) but also less distinct (for a given stimulus by increasing reliability). This has important computational implications that may be explored in a future study.

To facilitate further research, we are open-sourcing our model alongside detailed instructions to launch simulations and analyze the results (see Data and code availability). Simulating the model requires a performant hardware and software infrastructure. With respect to the latter we are continuously improving the performance and efficiency of the simulator (Kumbhar et al., 2019 [↗](#)). The model has several assumptions and limitations, which will ideally be improved upon iteratively and in a community-driven manner. First, by using the circuit model of Isbister et al. (2023) [↗](#), we inherit all assumptions listed therein; one of the most important one being on source data. A common limitation of data-driven modeling is the lack of high-quality source data, thus while we refer to our model as juvenile rat non-barrel somatosensory cortex, sometimes we used data from the barral cortex, from adult animals and from mouse. Second, for our simulations we used the model of Chindemi et al. (2022) [↗](#) with parameters obtained in a previous version of the circuit. Due to various improvements of synapse parameters (e.g. U_{SE} of L5 PCs) since fitting plasticity-specific parameters, the results for pairing protocols at various frequencies shift slightly while maintaining the presence of STDP (Supplementary Figure S2A). When exploring slightly different parameters of the plasticity model, we found that while the exact numerical values changed, our main conclusions about nonrandom changes and varios features predicting them were not affected (Supplementary Figure S13). Third, the extracellular magnesium con-centration of 1 mM used in *in vitro* preparations was assumed to be representative of the *in vivo* level. Fourth, experimental evidence indicates that firing bursts affects plasticity (Letzkus et al., 2006 [↗](#); Williams and Stuart, 1999 [↗](#); Larkum, 2013 [↗](#)). However, L5 TTPC bursts were rare in our simulations, as the model is based on an early developmental stage (P14-16; juvenile rats) and firing bursts only becomes prominent as the animals mature (Zhu, 2000 [↗](#)). Further-more, bursts can be triggered by synaptic input onto apical dendrites (Larkum, 2013 [↗](#); Hay et al., 2011 [↗](#)). Inputs to apical dendrites from high-order thalamus have been shown to gate plasticity in L23 PCs *in vivo*, via dis-inhibiting the distal dendrites (Gambino et al., 2014 [↗](#); Williams and Holtmaat, 2019 [↗](#)). While POm input was present in our model and targeted apical dendrites, we activated them in random and non-specific ways. On the other hand, top-down inputs rep-resent context or brain state and are thought to serve as an error or target signal for learning and thus are likely very specific (Makino, 2019 [↗](#)). Feedback signals from high-order cortical areas also innervate apical dendrites (Harris et al., 2019 [↗](#)), but as our model is of a single cortical area only, these inputs were absent. Missing excitatory input from other areas were compensated by somatic conductance injections (Isbister et al., 2023 [↗](#)), which could be replaced in the future by simulating additional dendritic synapses instead. Finally, learning in the brain is orches-trated by several mechanisms in unison and some of these are beyond the scope of this paper, e.g., inhibitory plasticity, metabotropic glutamate receptors, neuromodulation, slow homeostatic plasticity, structural plasticity, and calcium induced calcium release (Magee and Grienberger, 2020 [↗](#)).

Bottom-up modeling is often criticized for being arbitrarily complex. In our simulations, the mean value of synaptic efficacies of an over excited network reduced, comparable to [Turrigiano and Nelson \(2004\)](#) [without introducing rate homeostasis explicitly](#). In the machine learning field, mechanisms are added to classical algorithms to make updates sparser, overcome catastrophic forgetting, and allow the learning of tasks sequentially ([Kirkpatrick et al., 2017](#) [Zenke et al., 2017b](#)). The calcium-based plasticity rule employed here results in sparse updates without additional mechanisms, unlike STDP rules, which update efficacies at each spike arrival. Thus, we showed that adding biological complexity allowed us to reduce the number of mechanisms explicitly modeled. Calcium-based plasticity models existed for a long time ([Shouval et al., 2002](#) [Rubin et al., 2005](#)), have been shown to be stable ([Higgins et al., 2014](#) [Graupner et al., 2016](#) [Wang and Aljadeff, 2022](#)), have been modified to mimic *in vivo*-like low $[Ca^{2+}]_o$ ([Graupner et al., 2016](#) [Chindemi et al., 2022](#)), and versions more complex than ours exist ([Mäki-Marttunen et al., 2020](#) [Rodrigues et al., 2023](#)). Our contribution lies in integrating all these different aspects in network simulations at scale with subcellular resolution. Using such simulations allowed us to explore not only the temporal aspects of plasticity but also their link to spatial and network aspects, such as clusters of synapses and cell assemblies. Future work could explore the constellation of changing connections when stimuli change or cease, providing a more concrete examination of continuous learning paradigms within this model. Furthermore, the analysis methods developed on large scale functional and structural data empowered us to make and test novel predictions in the [MICrONS \(2021\)](#) dataset, which while pushing the boundaries of big data neuroscience, was so far only analyzed with focus on either single cells or first-order network metrics (e.g., node degree) ([Ding et al., 2023](#) [Wang et al., 2023](#)). We extend this work by considering the higher-order structure of the network.

4 Methods

4.1 Calcium-based plasticity model

The calcium-based long-term plasticity model used for E to E connections is fully described in [Chindemi et al. \(2022\)](#), but a minimal description of it can be found below. Synaptic efficacy (ρ) is based on the [Graupner and Brunel \(2012\)](#) formalism, which exhibits a bistable dynamics ($\rho = 0$ fully depressed, $\rho = 1$ fully potentiated, and $\rho_* = 0.5$ unstable fixed-point) described as:

$$\tau \frac{d\rho}{dt} = -\rho(1 - \rho)(\rho_* - \rho) + \gamma_p(1 - \rho)\Theta(Ca^* - \theta_p) - \gamma_d\rho\Theta(Ca^* - \theta_d) \quad (1)$$

where $\tau = 70$ s is the time constant of convergence, θ_d and θ_p are depression and potentiation thresholds, $\gamma_d = 101.5$ and $\gamma_p = 216.2$ are depression and potentiation rates and Θ is the Heaviside function. Ca^* is linked to the dynamics of $[Ca^{2+}]_i$ in spines (see below), which was modeled as:

$$\frac{d[Ca^{2+}]_i}{dt} = (I_{NMDA}^* + I_{VDCC}) \frac{\eta}{2FX} - \frac{[Ca^{2+}]_i - [Ca^{2+}]_i^{(0)}}{\tau_{Ca}} \quad (2)$$

where I^* and I_{VDCC} are calcium currents through NMDA receptors and VDCCs, $\eta = 0.04$ is the fraction of unbuffered calcium, F is the Faraday constant, X is the spine volume, $[Ca^{2+}]_i^{(0)} = 70$ pM is the resting value of $[Ca^{2+}]_i$, and $\tau_{Ca} = 12$ ms is the time constant of free (unbuffered) calcium clearance. I_{NMDA}^* depends on the state of the Mg^{2+} block as follows:

$$I_{NMDA}(t) = g_{NMDA}m(V - E_{NMDA}) \quad (3)$$

where g_{NMDA} and $E_{NMDA} = 3$ mV are the conductance and the reversal potential of the NMDA receptor, V is the local voltage, and m describes the nonlinear voltage dependence due to the Mg^{2+} block following the [Jahr and Stevens \(1990\)](#) formalism:

$$m = \frac{1}{1 + [Mg^{2+}]_o / \theta e^{-\kappa V}} \quad (4)$$

where θ is a scaling factor of the extracellular magnesium concentration $[Mg^{2+}]_o$, and κ is the slope of the voltage dependence. Parameters $\theta = 2.552$ and $\kappa = 0.072$ were obtained by refitting the model to cortical recordings from [Vargas-Caballero and Robinson \(2003\)](#) (as opposed to the original parameters fit to hippocampal ones [Jahr and Stevens, 1990](#)). Spines were assumed to be separate biochemical compartments, i.e., $[Ca^{2+}]_i$ of the dendrites does not influence that of the synapses.

Inspired by previous theoretical insights ([Rubin et al., 2005](#)), a leaky integrator of $[Ca^{2+}]_i$ was introduced (Ca^*) to slow down its time course instead of modeling enzymes downstream of calcium (e.g. CamKII as others did ([Mäki-Marttunen et al., 2020](#); [Rodrigues et al., 2023](#))):

$$\frac{dCa^*}{dt} = -\frac{Ca^*}{\tau^*} + [Ca^{2+}]_i - [Ca^{2+}]_i^{(0)} \quad (5)$$

where $\tau^* = 278.318$ ms is the time constant of the integrator. Updates in ρ were linked to this Ca^* variable crossing θ_d and/or θ_p (see equation (1)). The two synapse-specific threshold were derived based on peaks in $[Ca^{2+}]_i$ caused by pre- and postsynaptic spikes, c_{pre} and c_{post} respectively. To measure these parameters for all E to E 312,709,576 synapses, simulations of single cells were run, in which either the pre- or the postsynaptic excitatory cell was made to fire a single action potential and the local $[Ca^{2+}]_i$ was monitored in each synapse. Presynaptically evoked $[Ca^{2+}]_i$ peaks were three orders of magnitude larger, than the ones evoked by postsynaptic spikes (Supplementary Figure S2B). Postsynaptically evoked $[Ca^{2+}]_i$ peaks had a multimodal distribution in the apical dendrites (Supplementary Figure S2B right), in line with [Landau et al. \(2022\)](#). Since 8% of L6 PCs could not be made to fire a single action potential (only bursts), synapses on those cells (10,995,513 in total) were assumed to be non-plastic, i.e., their thresholds were set to a negative value that could not be crossed. Similarly, as the plasticity of connections between L4 spiny stellate cells was shown to be non-NMDA dependent ([Chindemi et al., 2022](#); [Egger et al., 1999](#)) those connections were made non-plastic. For the remainder of cells, θ_d and θ_p were derived as follows:

$$\begin{bmatrix} \theta_d \\ \theta_p \end{bmatrix} = \begin{bmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \end{bmatrix} \times \begin{bmatrix} c_{pre} \\ c_{post} \end{bmatrix} \text{ or } \begin{bmatrix} b_{00} & b_{01} \\ b_{10} & b_{11} \end{bmatrix} \times \begin{bmatrix} c_{pre} \\ c_{post} \end{bmatrix} \quad (6)$$

where a_{ij} and b_{ij} are constants optimized during model fitting for apical and basal dendrites respectively. Changes in ρ were then converted by low-pass filtering into changes U_{SE} and $\hat{g}_{AMPA}$ as follows:

$$\frac{dU_{SE}}{dt} = \frac{\bar{U}_{SE} - U_{SE}}{\tau_{change}} \quad \text{where} \quad \bar{U}_{SE} = U_{SE}^{(d)} + \rho(U_{SE}^{(p)} - U_{SE}^{(d)}) \quad (7)$$

$$\frac{d\hat{g}_{AMPA}}{dt} = \frac{\bar{g}_{AMPA} - \hat{g}_{AMPA}}{\tau_{change}} \quad \text{where} \quad \bar{g}_{AMPA} = \hat{g}_{AMPA}^{(d)} + \rho(\hat{g}_{AMPA}^{(p)} - \hat{g}_{AMPA}^{(d)}) \quad (8)$$

where $U_{SE}^{(d)}$, $U_{SE}^{(p)}$, $\hat{g}_{AMPA}^{(d)}$, and $\hat{g}_{AMPA}^{(p)}$ are the fully depressed (d) and fully potentiated (p) values of the given variables in-between which they evolve and $\tau_{change} = 100$ s is the time constant of the conversion of changes in ρ into changes in U_{SE} and $\hat{g}_{AMPA}$.

In the nbS1 model U_{SE} is also modulated by $[Ca^{2+}]_o$, where a reduction in $[Ca^{2+}]_o$ leads to a pathway-specific, non-linear reduction in U_{SE} (Markram et al., 2015 [↗](#); Ecker et al., 2020 [↗](#)).

When extracting the network's state after plasticity, not only the U_{SE} and $\hat{g}_{AMP A}$ values, but also the peak NMDA conductances ($\hat{g}_{NMDA}$) were updated according the ρ values in the last time step of the simulation.

Model parameter optimization (Chindemi et al., 2022 [↗](#)) and the derivation of thresholds from c_{pre} and c_{post} measurements was done with *plastyfire*.

4.2 *In vivo*-like spontaneous and evoked activity

The calibration process that leads to the *in vivo*-like spontaneous activity is fully described in Isbister et al. (2023) [↗](#), but a minimal description and a list of the parameters used in this article can be found below. As extracellular recordings are known to overestimate firing rates (Wohrer et al., 2013 [↗](#)), a spectrum of spontaneous states at fixed percentage of the rates reported in Reyes-Puerta et al. (2015) [↗](#) were calibrated (Isbister et al., 2023 [↗](#)). Matching specific firing rates *in silico* was achieved by iterative adjustments of layer and cell-type (excitatory/inhibitory) specific somatic conductance injection (following an Ornstein-Uhlenbeck process (Destexhe et al., 2001 [↗](#))). As in Isbister et al. (2023) [↗](#), extracellular recordings were assumed to have the same bias across layers and neuron populations. Furthermore it was assumed, that different inhibitory subpopulations require the same amount of input compensation. By introducing plasticity at all E to E synapses, an additional depolarizing current from VDCCs was added to the model, which made the network more active than its non-plastic counterpart (Supplementary Figure S3A). This required an algorithmic lowering of the amplitude of injected conductances from Isbister et al. (2023) [↗](#) to achieve the same *in vivo*-like layer-wise spontaneous firing rates (Supplementary Figure S3B). The spontaneous state used in the article is characterized by the parameters: $[Ca^{2+}]_o = 1.05$ mM (Jones and Keep, 1988 [↗](#)), percentage of reported firing rates = 40%, the coefficient of variation (CV; std/mean) of the noise process = 0.4. As in Markram et al. (2015) [↗](#), synapse dynamics (also known as short-term plasticity) of all synaptic pathways (not only E to E) played a pivotal role in achieving an *in vivo*-like asynchronous firing regime. In short, the modulation of the baseline release probability (U_{SE}) by $[Ca^{2+}]_o$ is pathway-dependent in the model, effecting E to E synapses more than I to E ones. This way, the I to E synapses have a smaller reduction in their efficacy at low $[Ca^{2+}]_o$ and therefore put the network in an E/I balanced state (Markram et al., 2015 [↗](#)).

The thalamic input patterns, and the spike trains delivered on them are fully described in Ecker et al. (2024) [↗](#), but a minimal description, highlighting the changes applied in this study, can be found below. First, the flatmap location (Bolaños-Puchet et al., 2024 [↗](#)) of VPM fibers avoiding the boundaries of the network were clustered with k-means to form 100 bundles of fibers. Second, the four base patterns (A, B, C, and D) were formed by randomly selecting four non-overlapping groups of bundles, each containing 12% of them. The remaining six patterns were derived from these base patterns with various degrees of overlap: three patterns as combinations of two of the base ones (E, F, G), two patterns as combinations of three of the base ones (H, I), and one pattern as a combination of all four base ones (J). Third, the input stream was defined as a random presentation of these 10 patterns, in a balanced way. Last, for each pattern presentation, unique spike times were generated for its corresponding fibers following a 100 ms-long inhomogeneous adapting Markov process (Muller et al., 2007 [↗](#)). The maximal rate of the VPM fibers was set to 17.5 Hz (compared to 30 Hz for the non-plastic circuit in Ecker et al., 2024 [↗](#)) and half of that for POM. The overlap of the patterns is clearly visible in the firing pattern of each group of fibers corresponding to them (Supplementary Figure S4).

4.3 Network simulations

Simulations were run using the NEURON simulator as a core engine with the Blue Brain Project's collection of hoc and NMODL templates for parallel execution on supercomputers (Kumbhar et al., 2019; Hines and Carnevale, 1997; Awile et al., 2022) (see Data and code availability). Simulating 10 minutes of biological time with reporting the state of all synapses (in every second) took 2,350,000 core hours (4x more than the corresponding non-plastic circuit without report-ing), on our HPE based supercomputer, installed at CSCS, Lugano. Although deep profiling of all campaigns was not done, an estimated 12M core hours were required to run all simulations presented in the manuscript. Simulations were always repeated at least three times to assess the qualitative consistency of the results (Supplementary Figure S8).

4.4 Control STDP rule

To compare the amount of changes induced by the calcium-based model of Chindemi et al. (2022) with classical plasticity rules, the 35,264,818 excitatory spikes from the 10-minute-long simulation were evaluated with a pair-based STDP rule (Gerstner et al., 1996; Kempter et al., 1999; Song et al., 2000). Synaptic weights evolved as follows under the STDP rule:

$$\Delta w_+ = A_+ \exp\left(-\frac{\Delta t}{\tau_+}\right) \text{ at } t_{post} \text{ if } t_{pre} < t_{post} \quad (9)$$

$$\Delta w_- = A_- \exp\left(\frac{\Delta t}{\tau_-}\right) \text{ at } t_{pre} \text{ if } t_{pre} > t_{post} \quad (10)$$

where t_{pre} and t_{post} are the times of pre-and postsynaptic spikes, $\Delta t = t_{post} - t_{pre}$ is the difference between them; $A_{\pm} = 0.05$ describe the weight update, which decayed exponentially with time constants $\tau_{\pm} = 20$ ms. The same parameters were used for the comparison with the 10-minute-long synchronous (unstable) simulation (76,025,009 excitatory spikes). The STDP rule was implemented in Brian2 (Stimberg et al., 2019).

4.5 Random walk control

Changes in ρ induced by the calcium-based model of Chindemi et al. (2022) and the STDP rule described above, were compared to a random control that takes into account global and noise-induced random changes in the network. To that end, a control model was used, where K connections of the network either increase or decrease by a noise value ϵ with equal probability in each time step. Note that, in this control, the size of the change vector at each time step is given by:

$$\|\rho(t + \Delta t) - \rho(t)\|_2 = \epsilon\sqrt{K}. \quad (11)$$

This process was modeled with a *random walk* in K -dimensional space of step size l , which at step N was given by:

$$\vec{R} = \sum_{i=0}^N \vec{r}_i, \quad (12)$$

where each $\vec{r}_i$ is a random variable in $\mathbb{R}^K$, where K is the number of changing connections of the network and such that the step size is given by $|\vec{r}_i| = \epsilon\sqrt{K} =: l$. The changes induced by the random walk at step N in the network is given by its distance from the origin at that time step, which for large enough N and K is approximately:

$$\|\vec{R}\| \approx l\sqrt{N}. \quad (13)$$

The step size l was determined as the average step size in the data after the 5 minute mark, when the transient activity has long passed. That is, l is given by:

$$\text{mean}_{t > 5 \text{ minutes}} \{ \|\rho(t + \Delta t) - \rho(t)\|_2 \}. \quad (14)$$

4.6 Cell assembly detection

The combination of methods from Carrillo-Reid et al. (2015) [\[15\]](#) and Herzog et al. (2021) [\[21\]](#) yielding the assembly detection pipeline is fully described in Ecker et al. (2024) [\[24\]](#), but a minimal description, highlighting the changes applied in this study, can be found below. First, spikes of excitatory cells were binned using 20 ms time bins (Harris et al., 2003 [\[3\]](#)). Second, time bins with significantly high firing rates were determined as crossing a threshold defined as the mean activity level plus the 95th percentile of the standard deviation of 100 shuffled controls. These shuffled controls were less strict than in Ecker et al. (2024) [\[24\]](#). Unlike in the original study, where spikes were only shifted by one time bin forward or backward (Carrillo-Reid et al., 2015 [\[15\]](#)), spikes were shifted by any amount. This change was introduced because the network's response to the same patterns was more variable in the plastic simulations, and to not miss any of them, a lower threshold was more fitting. Third, based on the cosine similarity of activation vectors, i.e., vectors of spike counts of all neurons in the given significant time bins, a similarity matrix was built (Carrillo-Reid et al., 2015 [\[15\]](#)). Fourth, this similarity matrix was hierarchically clustered using Ward's linkage (Pérez-Ortega et al., 2021 [\[25\]](#); Montijn et al., 2016 [\[16\]](#)). Like for any other unsupervised clustering method, the number of optimal clusters cannot be known beforehand, thus potential number of clusters were scanned between five and twenty. The number of clusters with the lowest Davis-Bouldin index was chosen, which maximizes the similarity within elements of the cluster while minimizing the between cluster similarity (Davies and Bouldin, 1979 [\[4\]](#)). Fifth, neurons were associated to these clusters based on their spiking activity, and it was determined whether they formed a cell assembly or not. The correlations between the spike trains of all neurons and the activation sequences of all clusters were computed and the ones with significant correlation selected to be part of the potential assemblies. Significance was determined based on exceeding the 95th percentile of correlations of shuffled controls (1000 controls with spikes of individual cells shifted by any amount as above; Herzog et al., 2021 [\[21\]](#); Montijn et al., 2016 [\[16\]](#)). Finally, it was required that the mean pairwise correlation of the spikes of the neurons with significant correlations was higher than the mean pairwise correlation of neurons in the whole dataset (Herzog et al., 2021 [\[21\]](#)). Clusters passing this last criterion were considered to be functional assemblies and the neurons with significant correlations their constituent cells.

The reliability of assembly sequences was defined as the Hamming similarity over the repetitions of a single pattern, which attains a value of 1 if two assembly sequences are identical and 0 if they are completely different. Assemblies of neurons were compared using their Jaccard similarity.

Assemblies were detected using the assemblyfire package.

4.7 Synapse clusters and likelihood of plastic changes within them

To quantify the importance of co-localization of synapses for plastic changes, clusters of synapses were detected. To be part of a synapse cluster, a synapse was required to have at least nine other synapses on the same dendritic branch, i.e., between two branching points of the dendrite, with 10 μm (Euclidean) distance. Significance of spatial clustering was determined similarly to Druckmann et al. (2014) [\[26\]](#). To that end, the distribution of synapse neighbor distances of the 10 selected synapses were compared with a Poisson model, assuming exponentially distributed inter-synapse distances, based on all (same branch) synapse neighbor distances on the given neuron. Clusters were merged in a subsequent step, thus synapse clusters with more than 10 synapses, spanning more than 20 μms were also feasible.

Plastic changes in synapse clusters were only analyzed for a small subpopulation of neurons in an assembly (10 L5 TTPCs per assembly), which were selected based on maximizing two connectivity features: assembly-indegree and synaptic clustering coefficient (SCC). The first one is the number of connections from a presynaptic assembly, while the second one quantifies the co-localization of synapses on the dendrites of a neuron from its presynaptic assembly. Maximizing these two metrics, which were introduced (and are described in detail) in [Ecker et al. \(2024\)](#), allows one to select subpopulations with high probability of finding synapse clusters.

Control synapse clusters, originating from non-assembly neurons were also detected on the same postsynaptic neurons.

The normalized likelihood of changes, conditioned on the four *categories* a synapse could fall into (assembly clustered, assembly non-clustered, non-assembly cluster, non-assembly non-clustered) were quantified using the Michaelson contrast, defined as:

$$\frac{Pr(changed | category) - Pr(changed)}{Pr(changed | category) + Pr(changed)} \quad (15)$$

where changed was split to be either potentiated ($\Delta\rho > 0$) or depressed ($\Delta\rho < 0$). Note that a nonzero value for one category always has to be compensated by a nonzero value with the opposite sign in another. For the normalized likelihood of initial ρ values, the same equation was used but $Pr(\Delta\rho > 0 | category)$ was replaced with $Pr(\rho_0 = 1 | category)$ for potentiated and $Pr(\Delta\rho < 0 | category)$ with $Pr(\rho_0 = 0 | category)$ for depressed synapses respectively.

4.8 Topological metrics

For quantifying the non-random nature of changing connections, directed simplex counts were used. A k -simplex in a network G is a set of $k + 1$ nodes (neurons) of G that are all-to-all connected in a feedforward fashion. That is, there is a numbering of the nodes $v_0, v_1, \dots, v_k$, such that for every $i < j$ there is an edge from v_i to v_j , and k is called the dimension of the simplex. In particular, 0-simplices are the nodes of the network, 1-simplices directed edges (connections) and 2-simplices are also known as transitive triads. Random controls were defined as the same number of edges between the same nodes, resulting in the same 0- and 1-simplex counts. Given an edge in G , a notion of its edge-centrality in the network is its *k-edge indegree*, which is the number of k -simplices the edge is innervated by. Note that a k -simplex innervating an edge is equivalent to it being an edge in a $k + 2$ -simplex going from the next to last to the last node of the simplex. This extends the classic notion of node indegree from nodes to edges. This value can be computed either for the simplices in the entire network or in an specified subnetwork e.g., the subnetwork on neurons belonging to an assembly.

Simplex counts and k -edge indegree values were computed using the connalysis package ([Egas Santander et al., 2024](#)), based on a fast C++ implementation from flagsercount ([Lüt-gehetmann et al., 2020](#)).

4.9 MICrONS dataset

For the comparison of two of our findings with a rodent electron microscopy dataset with co-registered calcium-imaging traces, the *minnie65_public* release of the MICrONS (2021) dataset was used. Synapses from sources other than one of the 60,048 classified neurons inside the bigger (*minnie65*) volume were discarded. The 117 version of the structural dataset was used for these analysis and made freely available in SONATA format ([Dai et al., 2020](#)) at <https://doi.org/10.5281/zenodo.8364070>. For comparable analysis we restricted the volume to its central part (650,000 x 950,000 and 700,000 z 1,000,000) and considered only E to E connections. Synapse volume was defined as the number of voxels painted by the automatic cleft segmentation (from the

corresponding synapses_pni_2 table) times the 4×4×40 nm voxel size. Total synapse volume was defined as the sum across all synapses mediating a connection. For the functional dataset, eight calcium-imaging sessions from the 661 version were used.

These sessions were selected by the same criteria as in Reimann et al. (2024b) [\[1\]](#); Egas Santander et al. (2024) [\[2\]](#), namely at least 1000 neurons were scanned in each session and at least 85% of them were co-registered in the structural data. Neurons with non-unique identifiers were dropped, and the remaining ones restricted to the central part of the structural volume described above. Deconvolved spike trains of neurons (243 neurons/session on average, 1817 unique neurons in total, 2% and 15% of all cells respectively) were used to calculate Pearson correlations. To average spike correlations from different session without any bias, they were first z-scored within each session.

4.10 Spike time reliability

Spike time reliability was defined as the mean of the cosine similarities of a given neuron's mean centered, smoothed spike times across all pairs of repetitions of a single pattern (Schreiber et al., 2003 [\[3\]](#); Cutts and Egle, 2014 [\[4\]](#)). To smooth the spike times, they were first binned to 1 ms time bins, and then convolved with a Gaussian kernel with a standard deviation of 10 ms as in Ecker et al. (2024) [\[5\]](#); Egas Santander et al. (2024) [\[2\]](#).

4.11 Visualization

Rendering of cells from the cortical network on **Figure 1A** [\[6\]](#) was done with Brayns, while the rendering of selected cells with their thalamo-cortical synapses on D with BioExplorer. Morphologies on **Figures 1** [\[7\]](#) and **6** [\[8\]](#) were rendered with NeuroMorphoVis (Abdellah et al., 2018 [\[9\]](#)).

Data and code availability

The 2.4 mm³ subvolume of the juvenile rat somatosensory cortex, containing 211,712 neurons and 312,709,576 plastic synapses in SONATA format (Dai et al., 2020 [\[10\]](#)) is freely available at: <https://doi.org/10.5281/zenodo.8158471> [\[11\]](#). It can be loaded and instantiated in CoreNEURON (Kumbhar et al., 2019 [\[12\]](#)) with neurodamus. The circuit and the simulations can be analyzed using Blue Brain SNAP and ConnectomeUtilities built on top of it. Cell assemblies were detected and can be analyzed with assemblyfire. Exemplary Jupyter notebooks using the packages above were deposited in the same repository on Zenodo.

Supplementary Figures




Figure S1

Physiology of excitatory cells and E to E connections.

A: Distribution of ion-channel densities in the excitatory (cADpyr) electrical type (etype). **B:** Validation of dendritic physiology of the cADpyr etype on L5 TTPC mtypes. **B1:** Validation of back-propagating action potential (bAP) amplitude for basal (teal) and apical (blue) dendrites. Reference data (in orange) comes from Stuart and Sakmann (1994) [Stuart and Sakmann \(1994\)](#); Larkum et al. (2001) [Larkum et al. \(2001\)](#) (apical) and Nevian et al. (2007) [Nevian et al. \(2007\)](#) (basal). Lines show exponential fits for the *in silico* (teal and blue) and *in vitro* (orange) data. Color bar indicates dendritic diameter. **B2:** Validation of EPSP attenuation. Reference data comes from Berger et al. (2001) [Berger et al. \(2001\)](#) (apical) and Nevian et al. (2007) [Nevian et al. \(2007\)](#) (basal). Lines and color bar same as in B2. Data taken from (and partially shown in) Reva et al. (2023) [Reva et al. \(2023\)](#). **C:** Anatomy and physiology of E to E connections. **C2:** Connection probability and number of synapses per connections for all E to E connections. White boxes indicate non-feasible connections, or on the left panel: no pairs found within the 200 μm intersomatic distance used. **C2:** Mean (over 100 pairs) PSP amplitude (left) and CV (std/mean on the right) of all E to E connections. Data taken from (and shown in) Isbister et al. (2023) [Isbister et al. \(2023\)](#). **C3:** Initial synaptic physiology parameters. From left to right: p , g_{AMPA} , and U_{SE} .




Figure S2

STDP and synapse-specific parameters of the plasticity model.

A: Spike timing, pairing frequency, and pathway dependence of plasticity. **B:** Layer- and neurite type-wise distribution of measured $[Ca^{2+}]_i$ peaks (used to derive parameters of the plasticity model shown in C). Synapses are grouped based on the soma location of the postsynaptic cell. 10% of all synapses are shown. Schematics on their lefts illustrate the measurement protocols. **C:** Layer- and neurite type-wise distribution of depression and potentiation thresholds (θ_p and θ_d) of the plasticity model. Synapses grouped and shown as in B. **D:** Correlations of the parameters shown in B and C.




Figure S3

Calibration of the *in vivo*-like network state.

A1: Same as [Figure 2B](#) (i.e., raster plots of the microcircuit's activity) under different synapse setups. The microcircuit equipped with the plasticity model of [Chindemi et al. \(2022\)](#) only resembles that of the non-plastic network's of [Isbister et al. \(2023\)](#) when VDCCs (voltage-dependent calcium channels) are blocked (last row). **A2:** Layer-wise (absolute) firing rates of excitatory (E) and inhibitory (I) subpopulations under stimulus-evoked conditions. Values correspond to the setup marked with an arrow on A1 (the setup used throughout the manuscript). Legend on its left applies to the three panels of B below as well. **B:** Re-calibration of the *in vivo*-like spontaneous state using the plasticity model. **B1:** Left: Euclidean distance of the measured percentages of firing rates (P_{FR} s) from the target ones in different iterations of the calibration process. Right: Validation of network states after the final (4th) iteration. Dashed gray line along the diagonal indicated perfect match. **B2** Left: Injected Ornstein-Uhlenbeck (OU) conductances in the non-plastic model of [Isbister et al. \(2023\)](#) vs. the plastic one for $P_{FR} = 40\%$ (the state used in the current article). Dashed gray line along the diagonal indicated perfect match. Right: Same as A2 but in the spontaneous state ($P_{FR} = 40\%$).




Figure S4

Activity of the thalamic fibers.

Raster plots of VPM fibers forming each of the 10 input patterns ([Figure 2A](#)) for the stimulus stream (i.e., from pattern A at 2000 ms to pattern J at 6500 ms). Bottom row shows the same for non-specific POM fibers. (Adapted from [Ecker et al. \(2024\)](#).)




Figure S5

Changes in synaptic efficacy during plasticity.

Individual p traces (10 per panel) during plasticity. Sampled traces have $|\Delta p| \geq 0.1$ and mean over the 10 minutes within $[0.2, 0.8]$.




Figure S6

Synchronous (unstable) network activity.

A: Same as **Figure 2B** (i.e., raster plots of the microcircuit's activity) under a different network state, characterized by higher $[\text{Ca}^{2+}]_o$ and thus population bursts (see Markram et al. (2015)). **A1** in the beginning of the 10-minute-long simulation, while **A2** at the end. **B:** Firing rate of excitatory cells (zoom-ins on the beginning and end (corresponding to the raster plots above) are shown in **Figure 2F**). **C** and **D:** Same as **Figure 2D** and E1 (i.e., L2 norm of changes in p (synapses) and mean p (connections) across time).




Figure S7

Changing connections in plastic control simulations.

A: Same as [Figure 2B](#) and [3A2](#) (i.e., raster plots of the microcircuit's activity and plastic changes in mean p vs. firing rates under different conditions). The last row of A2 is not an exact replica of [Figure 3A2](#) as these simulations were run for 2 minutes. **B:** Similar to [Figure 3A1](#) (i.e., layer-wise propensity of changes in mean p) but split across conditions. **C:** Similar to [Figure 2E1](#) (i.e., L2 norm of changes in mean p values for all conditions.)

Figure S8

Analysis results across 3 repetitions of the same simulations.

A: Same as [Figure 4C2](#), i.e., distances of mean ρ matrices in the 2-minute-long single pattern simulations. Each simulation was repeated three times, and simulations with different random seeds are along the diagonal (highlighted with black squares). **B:** Same as [Figure 7F](#), i.e., maximum probability of plastic changes against k -edge indegree in assembly subgraphs on top, and the bottom (pattern A) rows in more detail on the bottom. Arrows on top correspond to the detailed plots shown in the bottom. The first seed is the exact same as the left-hand side of [Figure 7F](#).





Figure S9

Rate-based controls.

A: Difference of propensity of changes in assemblies and their rate-based controls in [Figure 5F](#) and G2. **B:** Rate-based control of probability of changes vs. indegree bins ([Figure 7B1](#)) on the left. In particular, for each indegree bin, the same number of neurons with the same firing rate distribution were sampled at random. The probability of change was calculated for these random samples instead of the original ones. The y-axis is the same as of the main Figure for easier visual comparison. On the right: difference of probability of change vs. indegree and its rate-based control on its left.






Figure S10

Changes in cross-assembly synapse clusters.

A: Distribution of within-assembly Δp s across the four categories introduced in **Figure 6** (non-)assembly and (non-)clustered). Black arrow indicates clustered assembly synapses. Boxes show all values, while black dots are 1000 samples from each. Significance test was run on the balanced samples (1000 each): 2-way ANOVA and post-hoc Tukey's test: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$. **B:** Schematics of cross-assembly synapse clusters. Postsynaptic neuron from the green assembly, presynaptic ones from the orange assembly. **C:** As **Figure 6** C and D, i.e., Michelson contrast (*Equation (15)* in Methods) of the likelihood of plastic changes of synapses grouped into categories (not only four as there are several presynaptic assemblies) on the left, and initialization of p values on the right. Depression in blue and potentiation in red. Black arrow indicates clustered assembly synapses. Assembly 12 (A12) was chosen as the postsynaptic assembly as it appeared the latest in the sequence (**Figure 5B**) and thus had the most presynaptic assemblies which were active before it in time. **D:** Same as A, but for cross-assembly synapses (data from several postsynaptic assemblies, not only A12 shown on C).




Figure S11

Correlation of indegree and k-edge indegree, structural comparison with MICrONS and pattern-indegree analysis.

A: Pre- and postsynaptic (node) indegree vs. k-edge indegree across dimensions ($k = 0, 1$). R-values from left to right: 0.48, 0.35, 0.39, 0.27 **B:** Image taken from MICrONS (2021) (under CC-BY 4.0 license). Distributions of edge participation in high dimensional sim-plices (see Methods) across dimensions in the MICrONS (2021) dataset and in our nbS1 model (Reimann et al., 2024a). **C:** Probability of change vs. pattern-indegree (number of connections from VPM patterns). **C1:** Probability of changes (in the 2-minute-long single pattern simulations) vs. pattern-indegree of the presynaptic (left) and postsynaptic (right) neurons. **C2:** Probability of change (i.e., either depression or potentiation) against the pattern-indegree of both pre- and postsynaptic neurons on the left (for pattern A only). Probability of potentiation conditioned on plastic change (in any direction) against pattern-indegrees on the right. **D:** Pattern-indegree of pre- and postsynaptic cells (for the four base patterns only as the remaining ones are based on these four). R-values from left to right: 0.27, 0.23, 0.22, 0.19




Figure S13

Main results with slightly different plasticity model parameters.

Caption on the next page. The second subpanel, highlighted with red rectangle, is always the same as the first one, but with simulations ran with slightly different parameters of the plasticity model. Results on the first subpanels are replications of panels shown on other Figures. The following parameters of *equation (1)* differed between the simulations: $y_d = 100$ (instead of 101.5), $y_p = 450$ (instead of 216.2), and $\tau^* = 200$ ms (instead of 278.318 ms). **A1**: Same as **Figure 4C2** - extended to three repetitions as in Supplementary Figure S8A, i.e., input-output distance correlation. **B1**: Same as **Figure 5F**, i.e., propensity of depression and potentiation within and across cell assemblies. **C1**: Same as **Figure 6C**, i.e., Michaelson contrast of plastic changes of synapses grouped into four categories. **D1**: Same as **Figure 7C** and E2 - extended to three repetitions as in Supplementary Figure S8B, i.e., probability of change against k -edge indegree, and its restriction to assembly subgraphs.

Acknowledgements

The authors thank Nicolas Ninin for his involvement in the early stage of this project, Elvis Boci and Cyrille Favreau for their help with visualizations, Michael Gevaert, Joni Herttuainen and Thomas Delemontex for their assistance with software engineering, and Alberto Antonietti, Christoph Pokorny, Kathryn B. Hess, Ran Levi, Wulfram Gerstner, Roberto Araya and Henry Markram for discussions.

Additional information

Funding

This study was supported by funding to the Blue Brain Project, a research center of the École polytechnique fédérale de Lausanne (EPFL), from the Swiss government's ETH Board of the Swiss Federal Institutes of Technology. E.B.M. and D.P. were further supported by funding from the Institute for Data Valorization (IVADO), the CHU Sainte-Justine Research Center (CHUSJRC), Fonds de Recherche du Québec-Santé (FRQS), the Canada CIFAR AI Chairs Program, the Quebec Institute for Artificial Intelligence (Mila), and Google. Their compute infrastructure was supported through a grant of computing time awarded to E.B.M. from the Digital Research Alliance of Canada.

Author contributions

- Conceptualization: A.E., D.E.S., G.C., E.B.M., M.W.R.
- Formal analysis: A.E., D.E.S., M.W.R.
- Investigation: A.E., S.B.-P., G.C.
- Methodology: A.E., D.E.S., G.C., J.B.I., E.B.M., M.W.R.
- Project administration: M.W.R.
- Software: A.E., D.E.S., S.B.-P., G.C., J.K., P.K., M.W.R.
- Resources: J.B.A., I.M.
- Supervision: M.W.R.
- Validation: A.E., D.E.S., D.P.G.M., S.B.-P., J.B.I., M.W.R.
- Visualization: A.E., M.A.
- Writing - original draft: A.E., D.E.S., M.W.R.
- Writing - review & editing: A.E., D.E.S., S.B.-P., G.C., J.B.I., E.B.M., M.W.R.

References

- Abdellah M., Hernando J., Eilemann S., Lapere S., Antille N., Markram H., Schürmann F (2018) **NeuroMorphoVis: A collaborative framework for analysis and visualization of neuronal morphology skeletons reconstructed from microscopy stacks** *Bioinformatics* **34**:i574–i582 [Google Scholar](#)
- Awile O., Kumbhar P., Cornu N., Dura-Bernal S., King J. G., Lupton O., Magkanaris I., McDougal R. A., Newton A. J., Pereira F., Săvulescu A., Carnevale N. T., Lytton W. W., Hines M. L., Schürmann F (2022) **Modernizing the NEURON Simulator for Sustainability, Portability, and Performance** *Frontiers in Neuroinformatics* **16**:884046 [Google Scholar](#)
- Berger T., Larkum M. E., Lüscher H. R (2001) **High Ih channel density in the distal apical dendrite of layer V pyramidal cells increases bidirectional attenuation of EPSPs** *Journal of Neurophysiology* **85**:855–868 [Google Scholar](#)
- Bi, G.-q. and Poo, M.-m (1998) **Synaptic Modifications in Cultured Hippocampal Neurons: Dependence on Spike Timing, Synaptic Strength, and Postsynaptic Cell Type** *The Journal of Neuroscience* **18**:10464–10472 [Google Scholar](#)
- Bliss T. V. P., Lømo T (1973) **Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path** *The Journal of Physiology* **232**:331–356 [Google Scholar](#)
- Bobrowski O., Kahle M (2018) **Topology of random geometric complexes: a survey** *Journal of Applied and Computational Topology* **1**:331–364 [Google Scholar](#)
- Bolaños-Puchet S., Teska A., Hernando J. B., Lu H., Romani A., Schürmann F., Reimann M. W (2024) **Enhancement of brain atlases with laminar coordinate systems: Flatmaps and barrel column annotations** *Imaging Neuroscience* **2** [Google Scholar](#)
- Bono J., Wilmes K. A., Clopath C (2017) **Modelling plasticity in dendrites: from single cells to networks** *Current Opinion in Neurobiology* **46**:136–141 [Google Scholar](#)
- Borst J. G. G (2010) **The low synaptic release probability in vivo** *Trends in Neurosciences* **33**:259–266 [Google Scholar](#)
- Brémaud A., West D. C., Thomson A. M (2007) **Binomial parameters differ across neocortical layers and with different classes of connections in adult rat and cat neocortex** *PNAS* **104**:14134–14139 [Google Scholar](#)
- Brunel N (2016) **Is cortical connectivity optimized for storing information?** *Nature Neuroscience* **19**:749–755 [Google Scholar](#)
- Buzsáki G., Mizuseki K (2014) **The log-dynamic brain: How skewed distributions affect network operations** *Nature Reviews Neuroscience* **15**:264–278 [Google Scholar](#)
- Carrillo-Reid L., Miller J. E. K., Hamm J. P., Jackson J., Yuste R (2015) **Endogenous sequential cortical activity evoked by visual stimuli** *Journal of Neuroscience* **35**:8813–8828 [Google Scholar](#)

Chindemi G. (2018) **Towards a unified understanding of synaptic plasticity: parsimonious modeling and simulation of the glutamatergic synapse life-cycle** EPFL [Google Scholar](#)

Chindemi G., Abdellah M., Amsalem O., Benavides-Piccione R., Delattre V., Doron M., Ecker A., Jaquier A. T., King J., Kumbhar P., Monney C., Perin R., Rössert C., Tuncel M. A., van Geit W., DeFelipe J., Graupner M., Segev I., Markram H., Muller E. B. (2022) **A calcium-based plasticity model predicts long-term potentiation and depression in the neocortex** *Nature Communications* **13** [Google Scholar](#)

Clopath C., Büsing L., Vasilaki E., Gerstner W (2010) **Connectivity reflects coding: A model of voltage-based STDP with homeostasis** *Nature Neuroscience* **13**:344–352 [Google Scholar](#)

Costa R. P., Froemke R. C., Sjöström P. J., van Rossum M. C. (2015) **Unified pre-and postsynaptic long-term plasticity enables reliable and flexible learning** *eLife* **4**:e09457 <https://doi.org/10.7554/eLife.09457> | [Google Scholar](#)

Cutts C. S., Eglen X. S. J (2014) **Detecting pairwise correlations in spike trains: An objective comparison of methods and application to the study of retinal waves** *Journal of Neuroscience* **34**:14288–14303 [Google Scholar](#)

Dai K., Hernando J., Billeh Y. N., Gratiy S. L., Planas J., Davison A. P., Dura-Bernal S., Gleeson P., Devresse A., Dichter B. K., Gevaert M., King J. G., van Geit W. A., Povolotsky A. V., Muller E., Courcol J. D., Arkhipov A. (2020) **The SONATA data format for efficient description of large-scale network models** *PLoS Computational Biology* **16**:e1007696 [Google Scholar](#)

Davies D. L., Bouldin D. W (1979) **A Cluster Separation Measure**. *IEEE Transactions on Pattern Analysis and Machine Learning Pami* **1**:224–227 [Google Scholar](#)

Delattre V., Keller D., Perich M., Markram H., Muller E. B (2015) **Network-timing-dependent plasticity** *Frontiers in Cellular Neuroscience* **9**:220 [Google Scholar](#)

Deperrois N., Graupner M (2020) **Short-term depression and long-term plasticity together tune sensitive range of synaptic plasticity** *PLoS Computational Biology* **16**:e100826 [Google Scholar](#)

Destexhe A., Rudolph M., Fellous J. M., Sejnowski T. J (2001) **Fluctuating synaptic conductances recreate in vivo-like activity in neocortical neurons** *Neuroscience* **107**:13–24 [Google Scholar](#)

Ding Z., Fahey P. G., Papadopoulos S., et al. (2023) **Functional connectomics reveals general wiring rule in mouse visual cortex** *bioRxiv* [Google Scholar](#)

Druckmann S., Feng L., Lee B., Yook C., Zhao T., Magee J. C., Kim J (2014) **Structured Synaptic Connectivity between Hippocampal Regions** *Neuron* **81**:629–640 [Google Scholar](#)

Ecker A., Romani A., Sáray S., Káli S., Migliore M., Falck J., Lange S., Mercer A., Thomson A. M., Muller E., Reimann M. W., Ramaswamy S (2020) **Data-driven integration of hippocampal CA1 synaptic physiology in silico** *Hippocampus* **30**:1129–1145 [Google Scholar](#)

Ecker A., Santander D. E., Bolaños-Puchet S., Isbister J. B., Reimann M. W (2024) **Cortical cell assemblies and their underlying connectivity: an in silico study** *PLoS Computational Biology* **20**:e1011891 [Google Scholar](#)

Egas Santander D., Pokorny C., Ecker A., Lazovskis J., Santoro M., Smith J. P., Hess K., Levi R., Reimann M. W. (2024) **Efficiency and reliability in biological neural network architectures** *bioRxiv* [Google Scholar](#)

Egger V., Feldmeyer D., Sakmann B (1999) **Coincidence detection and changes of synaptic efficacy in spiny stellate neurons in rat barrel cortex** *Nature Neuroscience* **2**:1098–1105 [Google Scholar](#)

Farinella M., Ruedt D. T., Gleeson P., Lanore F., Silver R. A (2014) **Glutamate-Bound NM-DARs Arising from In Vivo-like Network Activity Extend Spatio-temporal Integration in a L5 Cortical Pyramidal Cell Model** *PLoS Computational Biology* **10**:4 [Google Scholar](#)

Fauth M. J., Van Rossum M. C. (2019) **Self-organized reactivation maintains and reinforces mem-ories despite synaptic turnover** *eLife* **8**:e43717 <https://doi.org/10.7554/eLife.43717> | [Google Scholar](#)

Feulner B., Perich M. G., Chowdhury R. H., Miller L. E., Gallego J. A., Clopath C (2022) **Small, correlated changes in synaptic connectivity may facilitate rapid motor learning** *Nature Communications* **13** [Google Scholar](#)

Frankle J., Carbin M. (2019) **The lottery ticket hypothesis: Finding sparse, trainable neural networks** *Iclr* [Google Scholar](#)

Froemke R. C., Letzkus J. J., Kampa B. M., Hang G. B., Stuart G. J (2010) **Dendritic synapse location and neocortical spike-timingdependent plasticity** *Frontiers in Synaptic Neuroscience* **2**:29 [Google Scholar](#)

Fusi S., Abbott L. F (2007) **Limits on the memory storage capacity of bounded synapses** *Nature Neuroscience* **10**:485–493 [Google Scholar](#)

Gambino F., Pagès S., Kehayas V., Baptista D., Tatti R., Carleton A., Holtmaat A. (2014) **Sensory-evoked LTP driven by dendritic plateau potentials in vivo** *Nature* **515**:116–119 [Google Scholar](#)

Gerstner W., Kempter R., Hemmen J. L. V., Wagnert H (1996) **A neuronal learning rule for sub-millisecond temporal coding** *Nature* **383**:76–78 [Google Scholar](#)

Goldberg J., Holthoff K., Yuste R (2002) **A problem with Hebb and local spikes** *Trends in Neurosciences* **25**:433–435 [Google Scholar](#)

Graupner M., Brunel N (2012) **Calcium-based plasticity model explains sensitivity of synaptic changes to spike pattern, rate, and dendritic location** *PNAS* **109**:3991–3996 [Google Scholar](#)

Graupner M., Wallisch P., Ostojic S (2016) **Natural firing patterns imply low sensitivity of synaptic plasticity to spike timing compared with firing rate** *Journal of Neuroscience* **36**:11238–11258 [Google Scholar](#)

Graves A. R., Roth R. H., Tan H. L., Zhu Q., Bygrave A. M., Lopez-Ortega E., Hong I., Spiegel A. C., Johnson R. C., Vogelstein J. T., Tward D. J., Miller M. I., Hugarir R. L (2021) **Visualizing synaptic plasticity in vivo by large-scale imaging of endogenous AMPA receptors** *eLife* **10**:e66809 <https://doi.org/10.7554/eLife.66809> | [Google Scholar](#)

- Harris J. A., Mihalas S., et al. (2019) **Hierarchical organization of cortical and thalamic connectivity** *Nature* **575**:195–202 [Google Scholar](#)
- Harris K. D (2005) **Neural signatures of cell assembly organization** *Nature Reviews Neuroscience* **6**:399–407 [Google Scholar](#)
- Harris K. D., Csicsvari J., Hirase H., Dragoi G., Buzsáki G (2003) **Organization of cell assemblies in the hippocampus** *Nature* **424**:552–556 [Google Scholar](#)
- Harris K. M., Stevens J. K (1989) **Dendritic spines of CA1 pyramidal cells in the rat hippocampus: Serial electron microscopy with reference to their biophysical characteristics** *Journal of Neuroscience* **9**:2982–2997 [Google Scholar](#)
- Hay E., Hill S., Schürmann F., Markram H., Segev I (2011) **Models of Neocortical Layer 5b Pyramidal Cells Capturing a Wide Range of Dendritic and Perisomatic Active Properties** *PLoS Computational Biology* **7**:e1002107 [Google Scholar](#)
- Hebb D. O (1949) **The Organization of Behavior; A Neuropsychological Theory** New York: John Wiley & Sons, Inc. [Google Scholar](#)
- Herzog R., Morales A., Mora S., Araya J., Escobar M. J., Palacios A. G., Cofré R (2021) **Scalable and accurate method for neuronal ensemble detection in spiking neural networks** *PLoS ONE* **16**:e0251647 [Google Scholar](#)
- Higgins D., Graupner M., Brunel N (2014) **Memory Maintenance in Synapses with Calcium-Based Plasticity in the Presence of Background Activity** *PLoS Computational Biology* **10**:e1003834 [Google Scholar](#)
- Hines M. L., Carnevale N. T (1997) **The NEURON simulation environment** *Neural computation* **9**:1179–1209 [Google Scholar](#)
- Hopfield J. J (1982) **Neural networks and physical systems with emergent collective computational abilities** *PNAS* **79**:2554–2558 [Google Scholar](#)
- Iacaruso M. F., Gasler I. T., Hofer S. B (2017) **Synaptic organization of visual space in primary visual cortex** *Nature* **547**:449–452 [Google Scholar](#)
- Inglebert Y., Aljadeff J., Brunel N., Debanne D (2020) **Synaptic plasticity rules with physiological calcium levels** *PNAS* **117**:33639–33648 [Google Scholar](#)
- Isbister J. B., Ecker A., Pokorny C., Bolaños-Puchet S., Egas Santander D., et al. (2023) **Modeling and Simulation of Neocortical Micro-and Mesocircuitry. Part II: Physiology and Experimentation** *bioRxiv* [Google Scholar](#)
- Jahr C. E., Stevens C. F (1990) **Voltage dependence of NMDA-activated macroscopic conductances predicted by single-channel kinetics** *The Journal of neuroscience* **10**:3178–3182 [Google Scholar](#)
- Jedrzejewska-Szmek J., Damodaran S., Dorman D. B., Blackwell K. T. (2017) **Calcium dynamics predict direction of synaptic plasticity in striatal spiny projection neurons** *European Journal of Neuroscience* **45**:1044–1056 [Google Scholar](#)
- Jones H., Keep R (1988) **Brain Fluid Calcium Concentration and Response To Acute Hypercal-caemia During Development in the Rat** *Journal of Physiology* **402**:579–593 [Google](#)

Scholar

- Kahle M (2009) **Topology of random clique complexes** *Discrete Mathematics* **309**:1658–1671 [Google Scholar](#)
- Kanari L., Ramaswamy S., Shi Y., Morand S., Meystre J., Perin R., Abdellah M., Wang Y., Hess K., Markram H (2019) **Objective Morphological Classification of Neocortical Pyramidal Cells** *Cerebral Cortex* **29**:1719–1735 [Google Scholar](#)
- Kastellakis G., Poirazi P (2019) **Synaptic Clustering and Memory Formation** *Frontiers in Molecular Neuroscience* **12**:300 [Google Scholar](#)
- Kempler R., Gerstner W., van Hemmen J. L. (1999) **Hebbian learning and spiking neurons** *Physical Review* **59**:4498–4514 [Google Scholar](#)
- Kim D., Park P., Li X., Wong-Campos J. D., Tian H., Moulton E. M., Grimm J. B., Lavis L., Cohen A. E (2023) **Mapping memories: pulse-chase labeling reveals AMPA receptor dynamics during memory formation** *bioRxiv* [Google Scholar](#)
- Kirchner J. H., Gjorgjieva J (2021) **Emergence of local and global synaptic organization on cortical dendrites** *Nature Communications* **12** [Google Scholar](#)
- Kirchner J. H., Gjorgjieva J (2022) **Emergence of synaptic organization and computation in dendrites** *Neuroforum* **28**:21–30 [Google Scholar](#)
- Kirkpatrick J., Pascanu R., Rabinowitz N., Veness J., Desjardins G., Rusu A. A., Milan K., Quan J., Ramalho T., Grabska-Barwinska A., Hassabis D., Clopath C., Kumaran D., Hadsell R (2017) **Overcoming catastrophic forgetting in neural networks** *PNAS* **114**:3521–3526 [Google Scholar](#)
- Kossio Y. F. K., Goedeke S., Klos C., Memmesheimer R. M (2021) **Drifting assemblies for persistent memory: Neuron transitions and unsupervised compensation** *PNAS* **118**:e2023832118 [Google Scholar](#)
- Krotov D., Hopfield J. J (2016) **Dense associative memory for pattern recognition** *Advances in Neural Information Processing Systems* **29**:1172–1180 [Google Scholar](#)
- Kumbhar P., Hines M., Fouriaux J., Ovcharenko A., King J., Delalondre F., Schürmann F (2019) **CoreNEURON: An Optimized Compute Engine for the NEURON Simulator** *Frontiers in Neuroinformatics* **13**:63 [Google Scholar](#)
- Landau A. T., Park P., Wong-Campos J. D., Tian H., Cohen A. E., Sabatini B. L (2022) **Dendritic branch structure compartmentalizes voltage-dependent calcium influx in cortical layer 2/3 pyramidal cells** *eLife* **11**:e76993 <https://doi.org/10.7554/eLife.76993> | [Google Scholar](#)
- Larkum M. E (2013) **A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex** *Trends in Neurosciences* **36**:141–151 [Google Scholar](#)
- Larkum M. E., Zhu J. J., Sakmann B (2001) **Dendritic mechanisms underlying the coupling of the dendritic with the axonal action potential initiation zone of adult rat layer 5 pyramidal neurons** *Journal of Physiology* **533**:447–466 [Google Scholar](#)
- Letzkus J. J., Kampa B. M., Stuart G. J (2006) **Learning rules for spike timing-dependent plasticity depend on dendritic synapse location** *Journal of Neuroscience* **26**:10420–

10429 [Google Scholar](#)

Lisman J (1989) **A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory** *PNAS* **86**:9574–9578 [Google Scholar](#)

Lisman J. E (1985) **A mechanism for memory storage insensitive to molecular turnover: A bistable autophosphorylating kinase** *PNAS* **82**:3055–3057 [Google Scholar](#)

Lisman J. E., Spruston N (2005) **Postsynaptic depolarization requirements for LTP and LTD: A critique of spike timing-dependent plasticity** *Nature neuroscience* **8**:839–841 [Google Scholar](#)

Litwin-Kumar A., Doiron B (2014) **Formation and maintenance of neuronal assemblies through synaptic plasticity** *Nature Communications* **5** [Google Scholar](#)

Losonczy A., Magee J. C (2006) **Integrative Properties of Radial Oblique Dendrites in Hippocampal CA1 Pyramidal Neurons** *Neuron* **50**:291–307 [Google Scholar](#)

Lütgehetmann D., Govc D., Smith J. P., Levi R (2020) **Computing persistent homology of directed flag complexes** *Algorithms* **13**:19 [Google Scholar](#)

Magee J. C., Grienberger C (2020) **Synaptic Plasticity Forms and Functions** *Annual Review of Neuroscience* **43** [Google Scholar](#)

Mäki-Marttunen T., Iannella N., Edwards A. G., Einevoll G. T., Blackwell K. T (2020) **A unified computational model for cortical post-synaptic plasticity** *eLife* **9**:e55714 <https://doi.org/10.7554/eLife.55714> | [Google Scholar](#)

Makino H (2019) **Top-down control: A unified principle of cortical learning** *Neuroscience Research* **141**:23–28 [Google Scholar](#)

Mallya A., Lazebnik S (2018) **PackNet: Adding Multiple Tasks to a Single Network by Iterative Pruning** In: *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition* [Google Scholar](#)

Markram H., Lübke J., Frotscher M., Sakmann B (1997) **Regulation of Synaptic Efficacy by Coincidence of Postsynaptic APs and EPSPs** *Science* **275**:213–215 [Google Scholar](#)

Markram H., Muller E. B., Ramaswamy S., Reimann M. W., et al. (2015) **Reconstruction and Simulation of Neocortical Microcircuitry** *Cell* **163**:456–492 [Google Scholar](#)

Markram H., Tsodyks M (1996) **Redistribution of synaptic efficacy between neocortical pyramidal neurons** *Letters to Nature* **382**:807–810 [Google Scholar](#)

Mayer M. L., Westbrook G. L., Guthrie P. B (1984) **Voltage-dependent block by Mg²⁺ of NMDA responses in spinal cord neurones** *Nature* **309**:261–263 [Google Scholar](#)

McCloskey M., Cohen N. J (1989) **Catastrophic Interference in Connectionist Networks: The Sequential Learning Problem** *The Psychology of Learning and Motivation* **24**:109–165 [Google Scholar](#)

- McKernan M. G., Shinnick-Gallagher P (1997) **Fear conditioning induces a lasting potentiation of synaptic currents in vitro** *Nature* **390**:607–611 [Google Scholar](#)
- Meyer H. S., Wimmer V. C., Hemberger M., Bruno R. M., De Kock C. P., Frick A., Sakmann B., Helmstaedter M. (2010) **Cell type-specific thalamic innervation in a column of rat vibrissa cortex** *Cerebral Cortex* **20**:2287–2303 [Google Scholar](#)
- MICrONS (2021) **Functional connectomics spanning multiple areas of mouse visual cortex** *bioRxiv* [Google Scholar](#)
- Miehl C., Onasch S., Festa D., Gjorgjieva J (2023) **Formation and computational implications of assemblies in neural circuits** *The Journal of Physiology* **601**:3071–3090 [Google Scholar](#)
- Montijn X. J. S., Olcese U., Pennartz X. C. M. A (2016) **Visual Stimulus Detection Correlates with the Consistency of Temporal Sequences within Stereotyped Events of V1 Neuronal Population Activity** *The Journal of Neuroscience* **36**:8624–8640 [Google Scholar](#)
- Morrison A., Aertsen A., Diesmann M (2007) **Spike-timing-dependent plasticity in balanced random networks** *Neural Computation* **19**:1437–1467 [Google Scholar](#)
- Muller E., Buesing L., Schemmel J., Meier K (2007) **Spike-Frequency Adapting Neural Ensembles: Beyond Mean Adaptation and Renewal Theories** *Neural Computation* **19**:2958–3010 [Google Scholar](#)
- Nevian T., Larkum M. E., Polsky A., Schiller J (2007) **Properties of basal dendrites of layer 5 pyramidal neurons: A direct patch-clamp recording study** *Nature Neuroscience* **10**:206–214 [Google Scholar](#)
- Nolte M., Gal E., Markram H., Reimann M. W (2020) **Impact of higher order network structure on emergent cortical activity** *Network Neuroscience* **4**:292–314 [Google Scholar](#)
- Oby E. R., Golub M. D., Hennig J. A., Degenhart A. D., Tyler-Kabara E. C., Yu B. M., Chase S. M., Batista A. P (2019) **New neural activity patterns emerge with long-term learning** *PNAS* **116**:15210–15215 [Google Scholar](#)
- Pérez-Ortega J., Alejandre-García T., Yuste R (2021) **Long-term stability of cortical ensembles** *eLife* **10**:e64449 <https://doi.org/10.7554/eLife.64449> | [Google Scholar](#)
- Perin R., Berger T. K., Markram H (2011) **A synaptic organizing principle for cortical neuronal groups** *PNAS* **108**:5419–5424 [Google Scholar](#)
- Pettit D. L., Wang S. S., Gee K. R., Augustine G. J (1997) **Chemical two-photon uncaging: A novel approach to mapping glutamate receptors** *Neuron* **19**:465–471 [Google Scholar](#)
- Pfister J.-P., Gerstner W (2006) **Triplets of Spikes in a Model of Spike Timing-Dependent Plasticity** *Journal of Neuroscience* **26**:9673–9682 [Google Scholar](#)
- Poirazi P., Brannon T., Mel B. W (2003) **Pyramidal Neuron as Two-Layer Neural Network** *Neuron* **37**:989–999 [Google Scholar](#)
- Ratcliff R (1990) **Connectionist models of recognition memory: constraints imposed by learning and forgetting functions** *Psychological review* **97**:285–308 [Google Scholar](#)

- Ray A., Christian J. A., Mosso M. B., Park E., Wegner W., Willig K. I., Barth A. L (2023) **Quantitative Fluorescence Analysis Reveals Dendrite-Specific Thalamocortical Plasticity in L5 Pyra-midal Neurons during Learning** *Journal of Neuroscience* **43**:584–600 [Google Scholar](#)
- Reimann M. W., Bolaños-Puchet S., Courcol J.-D., Egas Santandre D., et al. (2024a) **Modeling and Simulation of Rat Non-Barrel Somatosensory Cortex. Part I: Modeling Anatomy** *eLife* **13**:RP99688 <https://doi.org/10.7554/eLife.99688> | [Google Scholar](#)
- Reimann M. W., Egas Santander D., Ecker A., Muller E. B (2024b) **Specific inhibition and disinhibition in the higher-order structure of a cortical connectome** *Cerebral Cortex* **34**:bhae433 [Google Scholar](#)
- Reimann M. W., Nolte M., Scolamiero M., Turner K., Perin R., Chindemi G., Dłotko P., Levi R., Hess K., Markram H (2017) **Cliques of neurons bound into cavities provide a missing link between structure and function** *Frontiers in Computational Neuroscience* **11**:48 [Google Scholar](#)
- Reva M., Rössert C., Arnaudon A., Damart T., Mandge D., Tuncel A., Ramaswamy S., Markram H., Werner V. G (2023) **A universal workflow for creation, validation and generalization of detailed neuronal models** *Patterns* **4**:100855 [Google Scholar](#)
- Reyes-Puerta V., Sun J. J., Kim S., Kilb W., Luhmann H. J (2015) **Laminar and Columnar Structure of Sensory-Evoked Multineuronal Spike Sequences in Adult Rat Barrel Cortex in Vivo** *Cerebral Cortex* **25**:2001–2021 [Google Scholar](#)
- Rodrigues Y. E., Tigaret C. M., Marie H., O'donnell C., Veltz R (2023) **A stochastic model of hippocampal synaptic plasticity with geometrical readout of enzyme dynamics** *eLife* **12**:e80152 <https://doi.org/10.7554/eLife.80152> | [Google Scholar](#)
- Rößler N., Jungenitz T., Sigler A., Bird A., Mittag M., Rhee J. S., Deller T., Cuntz H., Brose N., Schwarzacher S. W., Jedlicka P (2023) **Skewed distribution of spines is independent of presynaptic transmitter release and synaptic plasticity and emerges early during adult neurogenesis** *Open Biology* **13**:230063 [Google Scholar](#)
- Rubin J. E., Gerkin R. C., Bi G. Q., Chow C. C (2005) **Calcium time course as a signal for spike-timing-dependent plasticity** *Journal of Neurophysiology* **93**:2600–2613 [Google Scholar](#)
- Schreiber S., Fellous J. M., Whitmer D., Tiesinga P., Sejnowski T. J (2003) **A new correlation-based measure of spike timing reliability** *Neurocomputing* [Google Scholar](#)
- Selig D. K., Nicoll R. A., Malenka R. C (1999) **Hippocampal long-term potentiation preserves the fidelity of postsynaptic responses to presynaptic bursts** *Journal of Neuroscience* **19**:1236–1246 [Google Scholar](#)
- Shatz C. J (1992) **The Developing Brain** *Scientific American* **267**:60–67 [Google Scholar](#)
- Shouval H. Z., Bear M. F., Cooper L. N (2002) **A unified model of NMDA receptor-dependent bidirectional synaptic plasticity** *PNAS* **99**:10831–10836 [Google Scholar](#)

- Sjöström P. J., Häusser M (2006) **A Cooperative Switch Determines the Sign of Synaptic Plasticity in Distal Dendrites of Neocortical Pyramidal Neurons** *Neuron* **51**:227–238 [Google Scholar](#)
- Sjöström P. J., Turrigiano G. G., Nelson S. B (2003) **Neocortical LTD via Coincident Activation of Presynaptic NMDA and Cannabinoid Receptors** *Neuron* **39**:641–654 [Google Scholar](#)
- Song S., Miller K. D., Abbott L. F (2000) **Competitive Hebbian learning through spike-timing-dependent synaptic plasticity** *Nature Neuroscience* **3**:919–926 [Google Scholar](#)
- Song S., Sjöström P. J., Reigl M., Nelson S., Chklovskii D. B (2005) **Highly nonrandom features of synaptic connectivity in local cortical circuits** *PLoS Biology* **3**:e68 [Google Scholar](#)
- Stimberg M., Brette R., Goodman D. F (2019) **Brian 2, an intuitive and efficient neural simulator** *eLife* **8**:e47314 <https://doi.org/10.7554/eLife.47314> | [Google Scholar](#)
- Stuart G. J., Sakmann B (1994) **Active propagation of somatic action potentials into neocortical pyramidal cell dendrites** *Nature* **367**:69–72 [Google Scholar](#)
- Stuart G. J., Spruston N (2015) **Dendritic integration: 60 years of progress** *Nature Neuroscience* **18**:1713–1721 [Google Scholar](#)
- Tazerart S., Mitchell D. E., Miranda-Rottmann S., Araya R (2020) **A spike-timing-dependent plasticity rule for dendritic spines** *Nature Communications* **11** [Google Scholar](#)
- Tonegawa S., Liu X., Ramirez S., Redondo R (2015) **Memory Engram Cells Have Come of Age** *Neuron* **87**:918–931 [Google Scholar](#)
- Tsodyks M., Markram H (1997) **The neural code between neocortical pyramidal neurons depends on neurotransmitter release probability** *PNAS* **94**:719–723 [Google Scholar](#)
- Turrigiano G. G., Nelson S. B (2004) **Homeostatic plasticity in the developing nervous system** *Nature Reviews Neuroscience* **5**:97–107 [Google Scholar](#)
- Ujfalussy B. B., Makara J. K (2020) **Impact of functional synapse clusters on neuronal response selectivity** *Nature Communications* **11** [Google Scholar](#)
- Vargas-Caballero M., Robinson H. P (2003) **A slow fraction of Mg²⁺ unblock of NMDA receptors limits their contribution to spike generation in cortical pyramidal neurons** *Journal of Neurophysiology* **89**:2778–2783 [Google Scholar](#)
- Vogels T. P., Spekreijse H., Zenke F., Clopath C., Gerstner W (2011) **Inhibitory Plasticity Balances Excitation and Inhibition in Sensory Pathways and Memory Networks** *Science* **334**:1569–1573 [Google Scholar](#)
- Wang B., Aljadeff J (2022) **Multiplicative Shot-Noise: A New Route to Stability of Plastic Networks** *Physical Review Letters* **129**:068101 [Google Scholar](#)
- Wang E. Y., Fahey P. G., Ponder K., Ding Z., Chang A., Muhammad T., Patel S., Ding Z., Tran D., Fu J., Papadopoulos S., Franke K., Ecker A. S., Reimer J., Pitkow X., Sinz F. H., Tolias A. S (2023) **Towards a Foundation Model of the Mouse Visual Cortex** *bioRxiv* [Google Scholar](#)
- Widrich M., Schäfl B., Pavlovic M., Ramsauer H., Gruber L., Holzleitner M., Brandstetter J., Sandve G. K., Greiff V., Hochreiter S., Klambauer G (2020) **Modern hopfield networks and**

attention for immune repertoire classification *Advances in Neural Information Processing Systems* **33** [Google Scholar](#)

Williams L. E., Holtmaat A (2019) **Higher-Order Thalamocortical Inputs Gate Synaptic Long-Term Potentiation via Disinhibition** *Neuron* **101**:91–102 [Google Scholar](#)

Williams S. R., Stuart G. J (1999) **Mechanisms and consequences of action potential burst firing in rat neocortical pyramidal neurons** *Journal of Physiology* **521**:467–482 [Google Scholar](#)

Wohrer A., Humphries M. D., Machens C. K (2013) **Population-wide distributions of neural activity during perceptual decision-making** *Progress in Neurobiology* **103**:156–193 [Google Scholar](#)

Zenke F., Agnes E. J., Gerstner W (2015) **Diverse synaptic plasticity mechanisms orchestrated to form and retrieve memories in spiking neural networks** *Nature Communications* **6** [Google Scholar](#)

Zenke F., Gerstner W., Ganguli S (2017a) **The temporal paradox of Hebbian learning and homeostatic plasticity** *Current Opinion in Neurobiology* **43**:166–176 [Google Scholar](#)

Zenke F., Poole B., Ganguli S. (2017b) **Continual learning through synaptic intelligence** *Icml* [Google Scholar](#)

Zhang D., Zhang C., Stepanyants A (2019) **Robust associative learning is sufficient to explain the structural and dynamical properties of local cortical circuits** *Journal of Neuroscience* **39**:6888–6904 [Google Scholar](#)

Zhu J. J (2000) **Maturation of layer 5 neocortical pyramidal neurons: amplifying salient layer 1 and layer 4 inputs by Ca²⁺ action potentials in adult rat tuft dendrites** *Journal of Physiology* **526**:571–587 [Google Scholar](#)

Author information

András Ecker[†]

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech, Geneva, Switzerland
ORCID iD: [0000-0001-9635-4169](#)

[†]Co-lead authors

Daniela Egas Santander[†]

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech, Geneva, Switzerland
ORCID iD: [0000-0001-9838-7992](#)

[†]Co-lead authors

Marwan Abdellah

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0000-0003-0751-2300](https://orcid.org/0000-0003-0751-2300)

Jorge Blanco Alonso

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0009-0000-4872-3855](https://orcid.org/0009-0000-4872-3855)

Sirio Bolaños-Puchet

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0000-0003-4049-6488](https://orcid.org/0000-0003-4049-6488)

Giuseppe Chindemi

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland, Department of Basic Neurosciences, University of Geneva, Geneva,
Switzerland
ORCID iD: [0000-0001-6872-2366](https://orcid.org/0000-0001-6872-2366)

Dhuruva Priyan Gowri Mariyappan

Department of Neurosciences, Faculty of Medicine, Université de Montréal, Montréal, Canada,
Centre de Recherche Azrieli du CHU Sainte-Justine, Montréal, Canada, Mila Quebec AI
Institute, Montréal, Canada

James B Isbister

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0000-0002-1013-3013](https://orcid.org/0000-0002-1013-3013)

James Gonzalo King

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0000-0003-0906-8389](https://orcid.org/0000-0003-0906-8389)

Pramod Kumbhar

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0000-0002-1756-801X](https://orcid.org/0000-0002-1756-801X)

Ioannis Magkanaris

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland
ORCID iD: [0000-0003-0726-1417](https://orcid.org/0000-0003-0726-1417)

Eilif B Muller

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland, Department of Neurosciences, Faculty of Medicine, Université de
Montréal, Montréal, Canada, Centre de Recherche Azrieli du CHU Sainte-Justine, Montréal,
Canada, Mila Quebec AI Institute, Montréal, Canada

ORCID iD: [0000-0003-4309-8266](https://orcid.org/0000-0003-4309-8266)

Michael W Reimann

Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech,
Geneva, Switzerland

ORCID iD: [0000-0003-3455-2367](https://orcid.org/0000-0003-3455-2367)

For correspondence: mwr@reimann.science

Editors

Reviewing Editor

Julijana Gjorgjieva

Technical University of Munich, Freising, Germany

Senior Editor

Panayiota Poirazi

FORTH Institute of Molecular Biology and Biotechnology, Heraklion, Greece

Reviewer #1 (Public review):

This paper investigates the dynamics of excitatory synaptic weights under a calcium-based plasticity rule, in long (up to 10 minutes) simulations of a 211,000-neuron biophysically detailed model of a rat cortical network.

Strengths

- (1) A very detailed network model, with a large number of neurons, connections, synapses, etc., and with a huge number of biological considerations implemented in the model.
- (2) A carefully developed calcium-based plasticity rule, which operates with biologically relevant variables like calcium concentration and NMDA conductances.
- (3) The study itself is detailed and thorough, covering many aspects of the cellular and network anatomy and properties and investigating their relationships to plasticity.
- (4) The model remains stable over long periods of simulations, with the plasticity rule maintaining reasonable synaptic weights and not pushing the network to extremes.
- (5) The variety of insights the authors derive in terms of relationships between the cellular and network properties and dynamics of the synaptic weights are potentially interesting for the field.
- (6) Sharing the model and the associated methods and tools is a big plus.

Weaknesses

- (1) Conceptually, there seems to be a missed opportunity here in that it is not clear what the network learns to do. The authors present 10 different input patterns, the network does some plasticity, which is then analyzed, but we do not know whether the learning resulted in anything functionally significant. Did the network learn to discriminate the patterns much better than at the beginning, to capture or anticipate the timing of pattern presentation, detect similarities between patterns, etc.? This is important to understand if one wants to assess the significance of synaptic changes due to plasticity. For example, if the network did not learn much new functionally, relative to its initial state, then the observed plasticity could be considered minor and possibly insufficient. In that case, were the network to learn

something substantial, one would potentially observe much more extensive plasticity, and the results of the whole study could change, possibly including the stability of the network. While this could be a whole separate study, this issue is of central importance, and it is hard to judge the value of the results when we do not know what the network learned to do, if anything.

(2) In this study, plasticity occurs only at E-to-E connections but not at others. However, it is well known that inhibitory connections in the cortex exhibit at the very least a substantial short-term plasticity. One would expect that not including these phenomena would have substantial consequences on the results.

(3) Lines 134-135: "We calibrated layer-wise spontaneous firing rates and evoked activity to brief VPM inputs matching in vivo data from Reyes-Puerta et al. (2015)."

(4) Can the authors show these results? It is an important comparison, and so it would be great to see firing rates (ideally, their distributions) for all the cell types and layers vs. experimental data, for the evoked and spontaneous conditions.

(5) That being said, the Reyes-Puerta et al. paper reports firing rates for the barrel cortex, doesn't it? Whereas here, the authors are simulating a non-barrel cortex. Is such a comparison appropriate?

(6) Comparison with STDP on pages 5-7 and Figure 2: if I got this right, the authors applied STDP to already generated spikes, that is, did not run a simulation with STDP. That seems strange. The spikes they use here were generated by the system utilizing their calcium-based plasticity rule. Obviously, the spikes would be different if STDP was utilized instead. The traces of synaptic weights would then also be different. The comparison therefore is not quite appropriate, is it?

(7) Section 2.3 and Figure 5: I am not sure this analysis adds much. The main finding is that plasticity occurs more among cells in assemblies than among all cells. But isn't that expected given what was shown in the previous figures? Specifically, the authors showed that for cells that fire more, plasticity is more prominent. Obviously, cells that fire little or not at all won't belong to any assemblies. Therefore, we expect more plasticity in assemblies.

(8) Section 2.4 and Figure 6: It is not clear that the results truly support the formulation of the section's title ("Synapse clustering contributes to the emergence of cell assemblies, and facilitates plasticity across them") and some of the text in the section. What I can see is that the effect on ρ is strong for non-clustered synapses (Figure 6C and Figure S8A). In some cases, it is substantially higher than what is seen for clustered synapses. Furthermore, the wording "synapse clustering contributes to the emergence of cell assemblies" suggests some kind of causal role of clustered synapses in determining which neurons form specific cell assemblies. I do not see how the data presented supports that. Overall, it appears that the story about clustered synapses is quite complicated, with both clustered and non-clustered synapses driving changes in ρ across the board.

(9) Section 2.5 and Figure 7: Can we be certain that it is the edge participation that is a particularly good predictor of synaptic changes and/or strength, as opposed to something simpler? For example, could it be the overall number of synapses, excitatory synapses, or something along these lines, that the source and/or target neurons receive, that determine the ρ dynamics? And then, I do not understand the claim that edge participation allows one to "delineate potentiation from depression". The only related data I can find is in Figure 7A3, about which the authors write "this effect was stronger for potentiation than depression". But I don't see what they mean. For both depression and facilitation, the changes observed are in the range of ~12% of probability values. And even if the effect is stronger, does it mean one can "delineate" potentiation from depression better? What does it mean, to "delineate"? If it is some kind of decoding based on the edge participation, then the authors did not show that.

(10) "test novel predictions in the MICrONS (2021) dataset, which while pushing the boundaries of big data neuroscience, was so far only analyzed with single cells in focus instead of the network as a whole (Ding et al., 2023; Wang et al., 2023)." That is incorrect. For example, the whole work of Ding et al. analyzes connectivity and its relation to the neuron's functional properties at the network level.

Comments on revisions:

The authors addressed all my concerns from the previous review, primarily via textual changes such as improved Discussion. Thus, most of the weaknesses raised in the original review are not eliminated - in particular, points 1, and 5-9 - but they are acknowledged and described better. This remains a useful study that should be of interest to researchers in the field.

<https://doi.org/10.7554/eLife.101850.2.sa3>

Reviewer #2 (Public review):

Summary:

This paper aims at understanding the effects of plasticity in shaping dynamics and structure of cortical circuits, as well as on how that depends on aspects as network structure and dendritic processing.

Strengths:

The level of biological detail included is impressive, and the numerical simulations appear to be well executed. Additionally, they have done a commendable job in open-sourcing the model.

Weaknesses (after revision):

- As noted in my initial review, the observation that network activity remains stable without an explicit homeostatic mechanism-while acknowledged by the authors as consistent with previous findings (e.g., Higgins et al., 2014)-is not clearly framed as a replication or validation step in the current manuscript. For instance, the abstract states: "In our exploratory simulations, plasticity acted sparsely and specifically, firing rates and weight distributions remained stable without additional homeostatic mechanisms," without noting that this outcome has been previously reported, albeit in models with different levels of biological detail. Furthermore, in the general response to reviewers, the authors list this as the first item in their summary of phenomena accounted for by the model, which gives the impression that it is being presented as a primary result.

If this finding is instead meant to serve as a necessary validation that prior results continue to hold under the authors' extended modeling framework-including multicompartmental neurons, stochastic synaptic transmission, and a modified calcium-based plasticity rule-this should be made more explicit in both the abstract and main text. Unless there were specific reasons to suspect that these model extensions might disrupt previously observed stability, the conceptual contribution of this validation step remains unclear.

I would encourage the authors to revise the manuscript to clarify the role and novelty of this result in the context of existing literature and to briefly motivate why confirming this property in their model was an important step.

- While the revised manuscript includes improvements in the discussion of the generality and specificity of the findings, it still offers limited interpretability and mechanistic insight. As it stands, the simulations provide limited understanding of the underlying principles or

mechanisms at play, which constrains the broader conclusions that can be drawn from the work.

- In my first review, I suggested that the comparison with the MICrONS dataset could be made more informative-specifically by showing the same quantification of Figure 7D (7B in the previous version) in a version of the model without plasticity and clarifying the interpretation of Figure 8B, where the data appears to align closely with the model before plasticity.

In their response, the authors explain that several of these features remain largely unchanged before and after plasticity. For example, they note that total g_{AMPA} increases with k -edge indegree even in the initial model configuration. I appreciate this clarification, but it highlights a conceptual point that should be more clearly addressed in the manuscript. If the aspects of the model that align with MICrONS data are already present before plasticity, then these similarities reflect properties of the initial network architecture or baseline dynamics, rather than outcomes shaped by the plasticity process itself. If this interpretation is correct, it represents an interesting and potentially important finding. However, it is not currently articulated in the text. The manuscript places strong emphasis on the role of plasticity in shaping network structure and dynamics, yet the comparisons with MICrONS data appear to reflect features that do not depend on plasticity. Clarifying this distinction would help readers better appreciate the implications of the model-data comparison and discern which conclusions are genuinely supported by the data.

<https://doi.org/10.7554/eLife.101850.2.sa2>

Reviewer #3 (Public review):

Summary:

Ecker et al. utilized a biologically realistic, large-scale cortical model of the rat's non-barrel somatosensory cortex, incorporating a calcium-dependent plasticity rule to examine how various factors influence synaptic plasticity under in vivo-like conditions. Their analysis characterized the resulting plastic changes and revealed that key factors, including the co-firing of stimulus-evoked neuronal ensembles, the spatial organization of synaptic clusters, and the overall network topology, play an important role in affecting the extent of synaptic plasticity.

Strengths:

The detailed, large-scale model employed in this study enables the evaluation of diverse factors across various levels that influence the extent of plastic changes. Specifically, it facilitates the assessment of synaptic organization at the subcellular level, network topology at the macroscopic level, and the co-activation of neuronal ensembles at the activity level. Moreover, modeling plasticity under in vivo-like conditions enhances the model's relevance to experiments.

Weaknesses:

The paper lacks mechanistic insights into the observed phenomena, particularly regarding aspects that are typically inaccessible in traditional simplified models, such as layer-specific and layer-to-layer pathway-specific plasticity changes.

<https://doi.org/10.7554/eLife.101850.2.sa1>

Author response:

The following is the authors' response to the original reviews

General response

Our modeling study integrates recent experimental advances on dendritic physiology, biophysical plasticity rules, and network connectivity motifs into a single model, aiming to clarify their hypothesized inseparable functional roles in neocortical learning. By modelling excitatory plasticity in multi-synaptic connections on dendrites within a network with biologically constrained higher-order structure, we show these aspects are sufficient to account for a wide range of interesting phenomena: First, the calcium-based plasticity rule acted sparsely and specifically, keeping the network stable without requiring homeostatic mechanisms or inhibitory plasticity, as usually employed for models based on STDP rules. Most importantly, simulations of the network initiated in a recurrent-excitation induced synchronous state transitioned to an in vivo-like asynchronous state, and remained there. Second, plastic changes were stimulus-dependent and could be predicted by neurons' membership in functional assemblies, spatial clustering of synapses on dendrites, and the topology of the network's connectivity. Several of our predictions could be confirmed by comparison to the MICrONS dataset.

Our study thus aims to provide a first broad exploration of these phenomena and their interactions in a model, as well as a foundation for future studies that examine specific aspects more deeply. Specific concerns of the reviewers about parameter choices (reviewer 2's 2nd point - 2.2), claims about stability (2.1 and 3.1), the STDP control (1.5), and the motivation behind network metrics (1.8, 2.3) are addressed in detail below and in the revised manuscript.

Reviewer #1 (Public review):

This paper investigates the dynamics of excitatory synaptic weights under a calcium-based plasticity rule, in long (up to 10 minutes) simulations of a 211,000-neuron biophysically detailed model of a rat cortical network.

Strengths

- (1) A very detailed network model, with a large number of neurons, connections, synapses, etc., and with a huge number of biological considerations implemented in the model.*
- (2) A carefully developed calcium-based plasticity rule, which operates with biologically relevant variables like calcium concentration and NMDA conductances.*
- (3) The study itself is detailed and thorough, covering many aspects of the cellular and network anatomy and properties and investigating their relationships to plasticity.*
- (4) The model remains stable over long periods of simulations, with the plasticity rule maintaining reasonable synaptic weights and not pushing the network to extremes.*
- (5) The variety of insights the authors derive in terms of relationships between the cellular and network properties and dynamics of the synaptic weights are potentially interesting for the field.*
- (6) Sharing the model and the associated methods and tools is a big plus.*

We thank the reviewer for their comments.

Weaknesses

(1) Conceptually, there seems to be a missed opportunity here in that it is not clear what the network learns to do. The authors present 10 different input patterns, the network does some plasticity, which is then analyzed, but we do not know whether the learning resulted in anything functionally significant. Did the network learn to discriminate the patterns much better than at the beginning, to capture or anticipate the timing of pattern presentation, detect similarities between patterns, etc.? This is important to understand if one wants to assess the significance of synaptic changes due to plasticity. For example, if the network did not learn much new functionally, relative to its initial state, then the observed plasticity could be considered minor and possibly insufficient. In that case, were the network to learn something substantial, one would potentially observe much more extensive plasticity, and the results of the whole study could change, possibly including the stability of the network. While this could be a whole separate study, this issue is of central importance, and it is hard to judge the value of the results when we do not know what the network learned to do, if anything.

(1.1) The reviewer raises a very interesting point of discussion. As they remarked, it is very hard to judge what the network learned to do. However, our model was not designed to solve a specific task and even defining precisely what "learning" entails in a primary sensory region is still an open question. As many before us, we hypothesized that one of the roles of the primary somatosensory cortex would be to represent stimuli features and that most of the learning process would happen in an unsupervised manner. This is indeed what we have demonstrated by showing the stimulus-specificity of changes as well as an increase of reliability of assembly sequences between repetitions after plasticity. We have added this to the Discussion in lines 523-525.

(2) In this study, plasticity occurs only at E-to-E connections but not at others. However, it is well known that inhibitory connections in the cortex exhibit at the very least a substantial short-term plasticity. One would expect that not including these phenomena would have substantial consequences on the results.

(1.2) This is indeed well known. Please consider that we do have short-term plasticity (called synapse dynamics in the manuscript) at all connections, including inhibitory ones. We thank the reviewer for pointing out this potential confusion in the wording. We have now clarified this in the Methods in lines: 691-697. Furthermore, we have listed not having long-term plasticity at inhibitory connections in the limitations part of the Discussion in line: 593.

(3) Lines 134-135: "We calibrated layer-wise spontaneous firing rates and evoked activity to brief VPM inputs matching in vivo data from Reyes-Puerta et al. (2015)."

(4) Can the authors show these results? It is an important comparison, and so it would be great to see firing rates (ideally, their distributions) for all the cell types and layers vs. experimental data, for the evoked and spontaneous conditions.

(1.3) The layer- and cell type specific spontaneous firing rates were indeed hidden in the Methods and on Supplementary Figure S3. We now reference that figure in the Results in line: 136. Furthermore, we have amended Supplementary Figure S3 (panel A2), to show these rates in the evoked state as well.

(5) That being said, the Reyes-Puerta et al. paper reports firing rates for the barrel cortex, doesn't it? Whereas here, the authors are simulating a non-barrel cortex. Is such a comparison appropriate?

(1.4) As correctly pointed out by the reviewer, we made the assumption that these rates would generalize to the whole S1 because of the sparsity of experimental data. This assumption is discussed in length in Isbister et al. (2023) and now in the limitations part of the Discussion in lines: 564-568.

(6) Comparison with STDP on pages 5-7 and Figure 2: if I got this right, the authors applied STDP to already generated spikes, that is, did not run a simulation with STDP. That seems strange. The spikes they use here were generated by the system utilizing their calcium-based plasticity rule. Obviously, the spikes would be different if STDP was utilized instead. The traces of synaptic weights would then also be different. The comparison therefore is not quite appropriate, is it?

(1.5) Yes, the reviewer's understanding is correct. However, considering the findings of Morrison et al. 2007 [PMID: 17444756], and Zenke et al. 2017 [PMID: 28431369] (cited in the manuscript in lines: 165-166), running STDP in a closed loop simulation would most likely make the network “blow up” because of the positive feedback loop. Thus, we argue that our comparison is more conservative, since by using pre-generated spikes, we opened the loop and avoided positive feedback. This is now further explained in lines: 166-167.

(7) Section 2.3 and Figure 5: I am not sure this analysis adds much. The main finding is that plasticity occurs more among cells in assemblies than among all cells. But isn't that expected given what was shown in the previous figures? Specifically, the authors showed that for cells that fire more, plasticity is more prominent. Obviously, cells that fire little or not at all won't belong to any assemblies. Therefore, we expect more plasticity in assemblies.

(1.6) We thank the reviewer for this comment. We added additional panels (G1 and G2) to Figure 5 (and describe their content in lines: 329-337) showing that this is not the case. Firing-rate alone is indeed predictive of plastic changes, but co-firing in assemblies is even more so.

(8) Section 2.4 and Figure 6: It is not clear that the results truly support the formulation of the section's title ("Synapse clustering contributes to the emergence of cell assemblies, and facilitates plasticity across them") and some of the text in the section. What I can see is that the effect on rho is strong for non-clustered synapses (Figure 6C and Figure S8A). In some cases, it is substantially higher than what is seen for clustered synapses. Furthermore, the wording "synapse clustering contributes to the emergence of cell assemblies" suggests some kind of causal role of clustered synapses in determining which neurons form specific cell assemblies. I do not see how the data presented supports that. Overall, it appears that the story about clustered synapses is quite complicated, with both clustered and non-clustered synapses driving changes in rho across the board.

(1.7) We agree with the reviewer, it is “quite complicated” and we also see that the writing could have been better/more precise and supported by the data shown on the Figure. We updated both the section title and a big chunk of the text to take the suggestions into account in lines: 361-373.

(9) Section 2.5 and Figure 7: Can we be certain that it is the edge participation that is a particularly good predictor of synaptic changes and/or strength, as opposed to something simpler? For example, could it be the overall number of synapses, excitatory synapses, or something along these lines, that the source and/or target neurons receive, that determine the rho dynamics? And then, I do not understand the claim that edge participation allows one to "delineate potentiation from depression". The only related

data I can find is in Figure 7A3, about which the authors write "this effect was stronger for potentiation than depression". But I don't see what they mean. For both depression and facilitation, the changes observed are in the range of ~12% of probability values. And even if the effect is stronger, does it mean one can "delineate" potentiation from depression better? What does it mean, to "delineate"? If it is some kind of decoding based on the edge participation, then the authors did not show that.

(1.8) We thank the reviewer for this comment. We have included an analysis of the predictive power of indegree of the pre and postsynaptic neuron of a connection on the rho dynamics in Figure 7 (panel B). Please consider, that the rho dynamics are described on the level of connections, while properties like indegree are on the level of nodes. Any procedure transferring a node based property to an edge based property involves choices e.g., should the values be added, multiplied, should one be preferential over the other, or should they be considered independently? As edge-based metrics avoid these arbitrary choices, we would argue that they are - ultimately - the simpler and more natural choice in this context.

Though we believe that the metric of edge participation is simple, we recognize it is perhaps not common. Thus, we have switched to using a version of it that is perhaps more intuitive for the community at large i.e., as a metric of common innervation. Moreover, we have changed the name “(k+2) edge participation” to “(k)-edge indegree”, to make it even more accessible. For k=0, this is the number of neurons that commonly innervate the connection, i.e., a common neighbour. And for k=1, this is the number of connections that commonly innervate the connection. This is equivalent to edge participation from the next to last to the last neuron in a simplex. Furthermore, in lines: 391-418 we have added additional text and references explaining the intuition of why we think this metric is relevant, as it has been shown to affect correlated activity of pairs of neurons, as well as assembly formation.

Furthermore, we have clarified the language referring to potentiation and depression in lines: 420-422 and 448.

(10) "test novel predictions in the MICrONS (2021) dataset, which while pushing the boundaries of big data neuroscience, was so far only analyzed with single cells in focus instead of the network as a whole (Ding et al., 2023; Wang et al., 2023)." That is incorrect. For example, the whole work of Ding et al. analyzes connectivity and its relation to the neuron's functional properties at the network level.

(1.9) We thank the reviewer for pointing this out. Indeed, the sentence was improperly worded. We have appropriately changed this phrasing in lines: 616-618.

Reviewer #2 (Public review):

Summary:

This paper aims to understand the effects of plasticity in shaping the dynamics and structure of cortical circuits, as well as how that depends on aspects such as network structure and dendritic processing.

Strengths:

The level of biological detail included is impressive, and the numerical simulations appear to be well executed. Additionally, they have done a commendable job in open-sourcing the model.

We thank the reviewer for their comments.

Weaknesses:

The main result of this work is that activity in their network model remains stable without the need for a homeostatic mechanism. However, as the authors acknowledge, this has been demonstrated in previous studies (e.g., Higgins et al. 2014). In those studies, stability was attributed to calcium-based rules combined with calcium concentrations at in vivo levels and background neuronal activity. Since the authors use the same calcium-based rule, it is unclear what new result, if any, is being presented. If the authors are suggesting that the mechanism in their simulations differs, that should be stated clearly, and evidence supporting that claim should be provided.

(2.1) We do not see this as the main result of our study, but rather a critical validation step, since our calcium rule, while similar to previous ones, is not exactly the same (see equations (1) and especially (2) in Methods). This has been clarified in the text in lines: 150-151. Note in particular, that one of the main differences is the stochastic synaptic transmission and the role of calcium concentration on the release probability. Furthermore, our model involves multicompartmental neurons instead of point neuron models, which to our knowledge was never tested before with calcium-based plasticity rules at the network level. Moreover, determining the time required for stability to be reached is a necessary step to set up the simulation parameters to test the main hypotheses about rules governing the plastic changes.

The other findings discussed in the paper are related to a characterization of the dependency of plastic changes on network structure. While this analysis is potentially interesting, it has the following limitations.

First, I believe the authors should include an analysis of the generality and specificity of their results. All the findings seem to be derived from a single run of the simulation. How do the results vary with different network initializations, simulation times, or parameter choices?

(2.2) All simulations were run with 3 different random seeds (mentioned in the Methods) and now shown in Supplementary Figure S8 for some selected analyses. The maximum duration of our simulations were limited by our hardware constraints. However, from the long (10 minutes) simulation we concluded that most changes happen within the first minute. This is how we determined 2 minutes as the simulation time for all other experiments. Parameters determining both the spontaneous and evoked network state are discussed in length in Isbister et al. (2023) and while we acknowledge that they are only shown in Supplementary Figure S3, we did not want to lengthen the manuscript with redundant details but rather refer to reader to the manuscript where this is discussed at large.

Crucially, we tried slightly different parameters of the plasticity model in the early phases of the research, and while they changed the exact numerical values of our results, the main trends (i.e., stabilization time, assemblies, synapse clustering, and network topology influencing plastic changes) remained unchanged. This is now shown in Supplementary Figure S13 and referenced in the Discussion in lines: 572-575.

Second, the presentation of the results is difficult to follow. The characterization comes across as a long list of experiments, making it hard to identify a central message or distinguish key findings from minor details. The authors provide little intuition about why certain outcomes arise, and the complexity of the simulation makes it challenging - if not impossible - to determine which model elements are essential for specific results and which mechanisms drive emergent properties. Additionally, the text often lacks crucial details. For instance, the description of k-edge participation should be expanded, and an explanation of what this method quantifies should be included. Overall, I believe the authors should focus on a smaller set of significant results and provide a more in-depth discussion.

(2.3) We acknowledge the complexity of these large-scale simulations and the interpretation of their results. We appreciate the reviewer's feedback on the areas that needed more detail. To address this, we have extended the Results section describing k-edge indegree with more background and intuition in lines: 391-418. See also our reply to reviewer 1 (1.8) above.


While the manuscript may appear to be "a long list of experiments," it is actually guided by the following logic: We choose a calcium-based rule because it was the natural choice in a multicompartimental model which already included calcium dynamics and NMDA receptors. After setting up the main network state, verifying stability (Figure 2), doing traditional basic analysis (Figure 3), and verifying that the changes are non-random (Figure 4); we elaborated on long-standing ideas about co-firing in cell assemblies (Figure 5) and spatial clustering of synapse on dendrites (Figure 6) interacting with plasticity. Finally as we had access to the network's non-random connectivity we tried to link the network's topology to the observed plastic changes. This was done with a higher order perspective, given that there was previous evidence for the relevance of these structures on cofiring and correlated activity.

While we understand the frustration, we would highlight that the study is the first of its kind at this scale and level of biological detail. Our goal was to offer a broad exploration of the factors influencing plasticity and their interactions at this scale. Thus, laying the groundwork for future studies to investigate specific aspects more deeply.

The comparison of the model with the MICrONS dataset could be improved. In Figure 7B, the authors should show how the same quantification looks in a network model without plasticity. In Figure 8B, the data aligns with the model before plasticity, so it's unclear how this serves as a verification of the theoretical predictions.

(2.4) Our only claim is that by being used to working with both functional and structural data we were able to develop a metric (k-edge indegree) that could be utilized to study the non-random, high-order topology of the MICrONS connectivity as well. On Figure 8, spike correlations in MICrONS more or less align with both cases (before vs. after plasticity); the only difference is that spike correlations looked different enough in the model so we thought they are worth showing for both cases. Moreover, as the changes are sparse (Figure 2 and 3) the synapse strength panel of Figure 7(D) looks almost exactly the same before plasticity (see first two panels of Author response image 1). In line with our results, the small and significant changes increase as k-edge indegree increases (last panel of Author response image 1). As the first two panels look almost the same and the third one is shown in a slightly different way (Figure 7C2) we would prefer not to include this in the manuscript, but only in our response.

Author response image 1.



Reviewer #3 (Public review):

Summary:

Ecker et al. utilized a biologically realistic, large-scale cortical model of the rat's non-barrel somatosensory cortex, incorporating a calcium-dependent plasticity rule to examine how various factors influence synaptic plasticity under in vivo-like conditions. Their analysis characterized the resulting plastic changes and revealed that key factors, including the co-firing of stimulus-evoked neuronal ensembles, the spatial organization of synaptic clusters, and the overall network topology, play an important role in affecting the extent of synaptic plasticity.

Strengths:

The detailed, large-scale model employed in this study enables the evaluation of diverse factors across various levels that influence the extent of plastic changes. Specifically, it facilitates the assessment of synaptic organization at the subcellular level, network topology at the macroscopic level, and the co-activation of neuronal ensembles at the activity level. Moreover, modeling plasticity under in vivo-like conditions enhances the model's relevance to experiments.

We thank the reviewer for their comments.

Weaknesses:

(1) The authors claimed that, under in vivo-like conditions and in the presence of plasticity, firing rates and weight distributions remain stable without additional homeostatic mechanisms during a 10-minute stimulation period. However, the weights do not reach the steady state immediately after the 10-minute stimulation. Therefore, extended simulations are necessary to substantiate the claim.

(3.1) We thank the reviewer for this comment, as it gave us the opportunity to clarify in the text our stabilization criteria. Indeed, the dynamical system of weight changes has not reached a zero-change steady state because the changes, while small, are non-zero. However, in a stochastic system with ongoing activity (stimulus- or noise-driven), non-zero changes are expected. Thus, we consider the system to be at steady state when changes become negligible relative to a null model given by a random walk. Our results show that this condition is met around the 2-minute mark, with negligible changes in the subsequent 8 minutes.

Moreover, for spontaneous activity, we showed that an unstable network exhibiting synchronous activity can be stabilized into an asynchronous regime by the calcium-based plasticity rule within 10 minutes. These results show that the system reaches a stochastic steady state within 10 minutes without requiring homeostatic mechanisms. Our work reveals that incorporating more biological detail (i.e. calcium-based plasticity), reduces the need for additional mechanisms to stabilize network activity (e.g. fast homeostatic mechanisms).

Interestingly, one might argue that after 10 minutes of stimulation the network might transition to a different weight configuration if the stimuli change or cease. We agree this is an intriguing question, which we added to the Discussion in lines 611-613. However, this scenario concerns continuous learning, not the system's steady-state dynamics.

(2) Another major limitation of the paper lies in its lack of mechanistic insights into the observed phenomena (particularly on aspects that are typically impossible to assess in traditional simplified models, like layer-specific and layer-to-layer pathways-specific

plasticity changes), as well as the absence of discussions on the potential computational implications of the corresponding observed plastic changes.

(3.2) Our study integrates recent experimental advances aiming to clarify their hypothesized inseparable functional roles in neocortical learning. In particular, we study three different kinds of mechanistic insight: co-firing in assemblies (Figure 5), synapse clustering on postsynaptic dendrites (Figure 6), and high-order network topology (Figure 7). Furthermore, layer specificity is shown (Figure 3A1, B1, B2, D1) and so is layer-to-layer specificity (Figure 4A2). In addition we also describe synapse clustering on postsynaptic dendrites (Figure 6) which is not available in simplified models either.

As such, the mechanistic insights provided in our work are integrative in nature and aim to provide a first broad exploration of these phenomena and their interactions-which are rarely considered together in experimental or modelling studies. This foundation paves the way for future studies that examine specific aspects more deeply in this level of biological detail.

Reviewer #1 (Recommendations for the authors):

(1) I would suggest the authors explain more explicitly that their study uses plasticity for E-to-E connections and not others. Doing so in multiple places in the paper, but certainly in Methods and early in Results, would be helpful. This is stated in lines 117-119 ("To simulate long-term plasticity, we integrated our recently published calcium-based plasticity model that was used to describe functional long-term potentiation and depression between pairs of pyramidal cells"), but could be highlighted more.

We have added it to several lines in the Methods: 621, 648, 649.

(2) "Simulations were always repeated at least three times to assess the consistency of the results." This sounds important. How is this used for the analysis? Do the results reported combine the data from the 3 simulations? How did the authors check the "consistency of the results"? Did they run any statistical tests comparing the results between the 3 simulations or was it more of a visual check?

The reported results come from a single simulation. Three simulations were run to check that no obvious qualitative differences could be found, such as a change of network regime, association between stimuli and assemblies. No statistical tests can be run with samples of size three. These are now shown in Supplementary Figure S8, and additional clarifying text has been added in Methods line: 722.

(3) "We needed 12M core hours to run the simulation presented in this manuscript." The Methods section mentions ~2.4 M core hours for a 10-minute simulation, which may be confusing. It might be helpful to provide a table with all the simulations run for this study.

We wanted to provide a rough estimate of the runtime, but did not run a deep profiling of all campaigns. The results depend on the actual hardware and configurations used (e.g., temporal resolution of synapse reporting). We understand the potential source of confusion and have clarified this in the Methods in lines 719-721 (and took it out from the Discussion).

Reviewer #2 (Recommendations for the authors):

(1) I found the paper somewhat challenging to follow, as there are many small points, making it unclear what the main message is. It sometimes feels like a list of 'we did this and found that.' It might be helpful if the authors focused on a smaller number of key results with more in-depth discussion. For instance, the discussion of network topology

on page 9 is intriguing but condensed into a single, dense paragraph that is hard to follow. Clarifying how the random control is generated would also be beneficial.

See our response to the public review's third point (2.3).

(2) Line 245: typo? "Furthermore, the maximal simplex dimension found in the subgraph was two higher than expected by chance."

We changed the grammar in line: 249.

(3) Line 410: typo? "It has been previously shown before that assemblies have many edges".

Noted and fixed in line: 463.

Reviewer #3 (Recommendations for the authors):

(1) The authors claimed that plasticity operates in a sparse and specific manner, with firing rates and weight distributions remaining stable without additional homeostatic mechanisms. However, as shown in Figure 2D inset, the weights do not reach their steady-state values immediately after the 10-minute stimulation. A similar issue is observed in Figure 2G. It would be necessary to show the claim is indeed true as the weights reach the steady states.

See our response to the public review's first point (3.1).

(2) In the model, synapses undergo both short- and long-term plasticity, but the contribution of short-term plasticity to the stated claim is unclear. It would be helpful to demonstrate how the results of Figure 2 are affected when short-term plasticity is excluded.

STP is needed to achieve the asynchronous in vivo-like firing state in our model (and is intimately linked to the fitting procedure of the plasticity rules - mean-field approximation is not possible due to the important role of synaptic failures in thresholded plasticity outcomes), thus it cannot be excluded. We have added this to the Methods in lines: 691-697.

(3) It would be helpful to include a supplementary plot, similar to Figure 2F, illustrating the corresponding results for STDP.

This is not possible as we did not run a different simulation with STDP, only evaluated the changes in connections with an STDP model using spikes from our simulation. We did not incorporate the STDP equations into our detailed network, as there is no canonical or unambiguous way for doing so (e.g., one would need to handle the fact the connections are multi-synaptic). Note however, that considering the findings of Morrison et al. 2007 [PMID: 17444756], and Zenke et al. 2017 [PMID: 28431369] (cited in the manuscript in lines: 165-166), running STDP in a closed loop simulation would most likely make the network "blow up" because of the positive feedback loop.

(4) It would be helpful to provide mechanistic insights into the current observations and to discuss the potential computational implications of the observed plastic changes. Particularly on aspects that are typically impossible to examine in traditional models, like layer-specific plastic changes presented in Fig. 3A1, B1, B2, D1, and layer-to-layer pathways-specific plastic changes illustrated in Figure 4A2.

See our response to the public review's second point (3.2).

(5) The use of the term 'assembly' in most places of the manuscript may cause confusion. To enhance clarity and foster effective discussions in the field, I would recommend replacing it with 'ensemble,' as suggested in Miehl et al. (2023), 'Formation and computational implications of assemblies in neural circuits' (The Journal of Physiology, 601(15), 3071-3090), which should also be cited.

We read the mentioned manuscript when it was published (and appreciated it a lot), now reference it, and explain why we did not exactly follow the suggestion in lines: 293-299.

(6) The title of Figure 5 is not directly supported by the current figure. To strengthen the alignment, it would be helpful to present the results from lines 303-306 in bar plots and incorporate them into Figure 5 to better substantiate the figure title.

While the mentioned lines compare maximum values to those within the whole dataset, we think those 2*12*12 values are better presented in condensed matrices than bar plots (while the maximum values are still easily grasped from the colorbars). We have added panel G2 to the figure to address a comment by reviewer 1 (1.7), we believe that this further supports the title of the Figure.

(7) Line 326, cite "Kirchner, J. H., & Gjorgjieva, J. (2021). Emergence of local and global synaptic organization on cortical dendrites. Nature Communications, 12(1), 4005." and "Kirchner, J. H., & Gjorgjieva, J. (2022). Emergence of synaptic organization and computation in dendrites. Neuroforum, 28(1), 21-30."

Although we were aware of the mentioned manuscripts, we did not include them originally because they are models of a different species. However, we have now cited these in line: 347.

(8) The contrast results for ensembles 11 and 12 do not appear to support the claims made in lines 339-341. Clarification on this point would be helpful.

The reviewer is right, we have updated lines: 360-361, to clarify the difference between the two late assemblies.

(9) For Figure 6C and 6D in Section 2.4, rather than presenting the results for individual ensembles (which could be moved to the supplementary materials), it would be easier if the authors could summarize the results by grouping them into three categories: early, middle, and late ensembles.

We agree with the reviewer's suggestion and tried it before, but as the results slightly depend on functional assembly size as well (not only temporal order) averaging them loses information (see different xlims of the panels). Given that the issue is complex we decided to show all the data on the Figure, but we have revised the text now to provide a more high-level interpretation.

<https://doi.org/10.7554/eLife.101850.2.sa0>