

Reviewed Preprint

v2 • August 29, 2025

Revised by authors

Reviewed Preprint

v1 • May 30, 2023

Viral genome sequence datasets display pervasive evidence of strand-specific substitution biases that are best described using non-reversible nucleotide substitution models

Rita Sianga-Mete , Penelope Hartnady, Wimbai Caroline Mandikumba, Kayleigh Rutherford, Christopher Brian Currin, Florence Phelanyane, Sabina Stefan, Steven Weaver, Sergei L Kosakovsky Pond, Darren Patrick Martin

Division of Computational Biology, Institute of Infectious Diseases and Molecular Medicine, Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa • Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa • Centre for Infectious Disease and Epidemiology Research, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa • Centre for Biomedical Engineering, School of Engineering, Brown University, Providence, United States • institute for Genomics and Evolutionary Medicine, Department of Biology, Temple University, Pennsylvania, United States • Wellcome Center for Infectious Diseases Research in Africa, Institute of Infectious Disease and Molecular Medicine and Department of Medicine, University of Cape Town, Cape Town, South Africa

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **valuable** study revisits the effects of substitution model selection on phylogenetics by comparing reversible and non-reversible DNA substitution models. The authors provide **solid** evidence that 1) it can be beneficial to include non-time-reversible models in addition to general time-reversible models when inferring phylogenetic trees out of simulated viral genome sequence data sets, and that 2) non time-reversible models may fit the real data better than the reversible substitution models commonly used in phylogenetics, a finding consistent with previous work.

<https://doi.org/10.7554/eLife.87361.2.sa3>

Abstract

Most phylogenetic trees are inferred using time-reversible evolutionary models that assume that the relative rates of substitution for any given pair of nucleotides are the same regardless of the direction of the substitutions. However, there is no reason to assume that the underlying biochemical mutational processes that cause substitutions are similarly symmetrical. We consider two non-reversible nucleotide substitution models: (1) a 6-rate non-reversible model (NREV6) that is applicable to analyzing mutational processes in double-

stranded genomes in that complementary substitutions occur at identical rates; and (2) a 12-rate non-reversible model (NREV12) that is applicable to analyzing mutational processes in single-stranded (ss) genomes in that all substitution types are free to occur at different rates. Using likelihood ratio and Akaike Information Criterion-based model tests, we show that, surprisingly, NREV12 provided a significantly better fit than the General Time Reversible (GTR) and NREV6 models to 21/31 dsRNA and 20/30 dsDNA datasets. As expected, however, NREV12 provided a significantly better fit to 24/33 ssDNA and 40/47 ssRNA datasets. We tested how non-reversibility impacts the accuracy with which phylogenetic trees are inferred. As simulated degrees of non-reversibility (DNR) increased, the tree topology inferences using both NREV12 and GTR became more accurate, whereas inferred tree branch lengths became less accurate. We conclude that while nonreversible models should be helpful in the analysis of mutational processes in most virus species, there is no pressing need to use these models for routine phylogenetic inference.

Introduction

Modelling the nucleotide substitution processes that underly the diversification of virus genome sequences lies at the heart of many viral evolutionary analyses. The most widely used nucleotide substitution models belong to the general time reversible (GTR) family (Tavare, 1986) and assume that the Markov process of evolution is time-reversible (Hoff, Orf, Riehm, Darriba, & Stamatakis, 2016), (Lio & Goldman, 1998), (Tavare, 1986).

The GTR model is defined by its instantaneous rate matrix Q_{ij} equation (1), where Q_{ij} defines the instantaneous rate of change from nucleotide $i \in \{A, C, G, T\}$ to nucleotide j ; subject to the detailed balance condition: $q_{ji}\pi_i = q_{ij}\pi_j$, with rates q and equilibrium frequencies π (Squartini & Arndt, 2008) (Posada, 2003). The instantaneous rate matrix of the GTR model includes six rate parameters (a, b, c, d, e, and f). Because only products of substitution rates and evolutionary times can be estimated, one of the rate parameters is set to 1 (e.g. b), or the entire matrix is normalized to yield one expected substitution per unit time.

$$Q = \{q_{ij}\} = \begin{pmatrix} - & a\pi_C & b\pi_G & d\pi_T \\ a\pi_A & - & c\pi_G & e\pi_T \\ b\pi_A & c\pi_C & - & f\pi_T \\ d\pi_A & e\pi_C & f\pi_G & - \end{pmatrix} \quad (1)$$

The rate matrix in equation (1) is symmetrical in that, for example, the relative rate at which A changes to G is the same as the relative rate at which G changes to A.

Time reversible nucleotide substitution models such as GTR form the basis of almost all nucleotide sequence-focused evolutionary analyses (including those involving eukaryotes, prokaryotes, and viruses) (Lefort, Longueville, & Gascuel, 2017), (Posada & Crandall, 2001), (Minin, Abdo, Joyce, & Sullivan, 2003).

The reliability of a phylogenetic tree constructed using a particular nucleotide sequence dataset should be maximized when the evolutionary models used to construct the tree accurately reflect the important aspects of the evolutionary process (Buckley & Cunningham, 2002), (Ripplinger & Sullivan, 2008), (Hoff, Orf, Riehm, Darriba, & Stamatakis, 2016). The suitability of different models for describing the evolution of DNA or RNA sequences is, therefore, expected to depend to some degree on the biological and environmental contexts of the sequences being analyzed.

Mutations in viral genomes arise due to diverse biotic (such as replication enzyme infidelities, RNA/DNA editing enzymes) and abiotic (such as ionizing radiation, inorganic oxidizers and chemical mutagens) factors (Sanjuan & Domingo-Calap, 2016). Mutagenic chemical reactions or types of radiation that, for example, cause G to A or C to U mutations in DNA or RNA, are not the same as those that cause A to G or U to C mutations (Cheng, Cahill, Kasai, Nishimura, & Loeb, 1992), (Nguyen, et al., 1992), (Chelico, Pham, Calabrese, & Goodman, 2006), (Sharma, Patnaik, Taggart, & Baysal, 2016). It should not be expected, therefore, that the relative rates of G to A substitution will equal the relative rates of A to G substitution. Instead, in evolving double-stranded (ds) DNA and dsRNA molecules where both strands of the genome are in existence for similar amounts of time, both G to A and C to T substitutions should occur at relatively similar rates. Therefore, for nucleotide sequence datasets derived from any organisms with dsDNA or dsRNA genomes, a non-reversible nucleotide substitution model with a different relative substitution rate category for each of the six possible pairs of complementary nucleotide substitutions (e.g. NREV6 in equation (2)), with $q_{AC} = q_{TG}$, $q_{AG} = q_{TC}$, $q_{AT} = q_{TA}$, $q_{CG} = q_{GC}$, $q_{CT} = q_{GA}$, $q_{GT} = q_{CA}$, might plausibly provide a better description of mutational processes than GTR (Baele, Van de Peer, & Vansteelandt, 2010), (Wickner, 1993).

$$Q = \{q_{ij}\} = \begin{pmatrix} - & a\pi_C & b\pi_G & c\pi_T \\ f\pi_A & - & d\pi_G & e\pi_T \\ e\pi_A & d\pi_C & - & f\pi_T \\ c\pi_A & b\pi_C & a\pi_G & - \end{pmatrix} \quad (2)$$

In the case of single-stranded (ss) RNA viruses, ssDNA viruses, retroviruses, and dsRNA/dsDNA viruses where the two complementary genome strands do not exist for equal amounts of time (Yu, et al., 2004), a model where all twelve different substitutions occur at different rates might be best. Specifically, with ssRNA viruses, ssDNA viruses, and retroviruses, only one of the genome strands (called the virion strand) is packaged into viral particles for transmission and, in many dsRNA viruses, the genome strand that is translated into proteins (called the + strand) exists for longer during the life cycle than does the complementary (or -) strand (Bruslind, 2020), (Onwubiko, et al., 2020). In all these viruses, some degree of strand-specific substitution bias is expected to occur (Van Der Walt, Martin, Varsani, Polston, & Rybicki, 2008), (Polak & Arndt, 2008) such that NREV6 might be anticipated to provide a poorer description of mutational processes than a model such as NREV12 (equation (3)) where each of the twelve different types of substitution has a separate rate (Baele, Van de Peer, & Vansteelandt, 2010).

$$Q = \{q_{ij}\} = \begin{pmatrix} - & a\pi_A & b\pi_A & c\pi_A \\ g\pi_C & - & d\pi_C & e\pi_C \\ h\pi_G & i\pi_G & - & f\pi_G \\ j\pi_T & k\pi_T & l\pi_T & - \end{pmatrix} \quad (3)$$

Because non-reversible models consider the directionality of evolution, they could, in some cases, be used to identify root nodes of phylogenetic trees (Yap & Speed, 2005), (Boussau & Gouy, 2006). It is, however, unclear whether non-reversible models might, in certain situations at least, perform better than reversible models in the context of phylogenetic inference. Although it is possible to use non-reversible nucleotide substitution models such as NREV6 and NREV12 during maximum likelihood-based phylogenetic inference with computer programs such as IQ-TREE (Nguyen, Schmidt, Von Haeseler, & Minh, 2015), these models are not routinely used for phylogenetic inference. This is in part because non-reversible models render several commonly used algorithmic techniques for efficient likelihood computation inapplicable making inference

slower. It is also in part because it remains undetermined whether, under conditions where strand-specific substitution biases are evident, non-reversible models consistently yield substantially more accurate phylogenetic trees than reversible models.

Here we present evidence that strand-specific nucleotide substitution biases are common within virus genomic sequence datasets such that NREV12 generally provides a significantly better fit than both GTR and NREV6 for such datasets. We then use simulations to demonstrate that whereas strand-specific nucleotide substitution biases reduce the accuracy of phylogenetic inference under both GTR or NREV12, when these biases become extreme use of NREV12 can yield significantly more accurate phylogenetic trees than GTR.

Results and Discussion

Non-reversible nucleotide substitution models generally provide a better fit than reversible models to virus sequence datasets

We tested for evidence of non-reversibility of the nucleotide substitution process in 141 virus sequence datasets (33 ssDNA virus datasets, 30 dsDNA virus datasets, 31 dsRNA virus datasets, and 47 ssRNA virus datasets, all consisting of either full genome sequences (for unsegmented viruses), or complete genome component sequences (for viruses with segmented genomes). Specifically, for each dataset, we compared the goodness-of-fit of the GTR+G, NREV6+G, and NREV12+G models (where G represents gamma-distributed nucleotide substitution rates among sites (Yang, 1994)).

Given that dsDNA viruses such as adenoviruses, papillomaviruses and herpesviruses have both their DNA strands in existence for similar amounts of time before DNA-dependant-DNA polymerase enzymes copy both their + and - DNA strands during replication (Hanson, 2009), we had anticipated that the best fitting substitution model for sequence datasets of these viruses would be NREV6. Using weighted AIC-c scores to reveal trends of model support (Figure 1), it is surprising that NREV12 was overall the best-supported model (illustrated by the redder hues around the top corner of the dsDNA plot in Figure 2). Out of the 30 dsDNA datasets considered, we found that NREV6 provided the best fit to five datasets (HPV18, HPV45, HPV16, BPV, and SV40) and GTR provided the best fit to five (Alphapapilloma virus 6, JC polyomavirus, DPV, RTBV, and DBAV). NREV12 was the best fitting model for the remaining 20 datasets (Table 1). Further, LRTs revealed strong overall support for NREV12, with this model providing a significantly better fit ($p < 0.05$) than NREV6 for 25/30 of the dsDNA datasets and a significantly better fit than GTR for 24/30 of the datasets.

As NREV6 was not the best fitting model for most of the dsDNA virus datasets we infer that, in most dsDNA virus species, strand-specific substitution biases are not ignorable. Further, the datasets where NREV6 was not the best fit are from species in families containing other species where NREV6 was the best fit, indicating that such strand-specific substitution biases are unlikely to be a consequence of some broadly conserved feature of viral life cycles in these families (such as ssDNA replicative intermediates). It is instead plausible that these differences may relate to:

1. differences in replication fidelity and/or proofreading efficiency on the leading and trailing DNA strands in some dsDNA virus species (Grigoriev, 1999). These differences are common in eukaryotes (Youri, Newlon, & KunkelThomas, 2002) (Furusawa, 2012) and prokaryotes (Fijałkowska, Jonczyk, Tkaczyk, Białoskorska, & Schaaper, 1998) and, considering that the replication processes of dsDNA viruses analyzed here mirror those of their eukaryote hosts, it is perhaps unsurprising that most of these viruses also display some evidence of strand-specific substitution biases.




Figure 1

Ternary plots illustrating the relative fit of the NREV12, NREV6, and GTR nucleotide substitution models based on weighted AIC-c scores for 30 dsDNA, 31 dsRNA, 33 ssDNA, and 47 ssRNA virus nucleotide sequence datasets.

These plots were produced using the Akaike weights function with an overlaid density function (implemented in the qpcR package of RStudio (Ritz & Spiess, 2008 <https://doi.org/10.1002/bio.1448>)) to indicate point densities. Each model is represented by a corner of the triangles, and each circle represents the relative fit of each of the three models to a single nucleotide sequence dataset. The sides of the triangle represent model support axes ranging from 0-100%, with the position of a circle in relation to each of the sides of the triangle indicating the probability of models best describing the nucleotide sequence dataset that is represented by that point. Red colours represent a very high density of nucleotide sequence datasets that favour a particular model, blue colours indicate a lower, but still substantial, density of datasets that favour a particular model.




Figure 2

Weighted Robinson-Foulds distances between inferred and true phylogenetic trees for datasets simulated with different degrees of nucleotide substitution non-reversibility and different average pairwise sequence identities (APIs) (~75%, ~80%, ~85%, ~90% and ~95%).

“ns” above a pair of box and whisker plots indicates a paired t-test adjusted p-value of greater than or equal to 0.05 and “*” indicate a paired t test adjusted p-value of <0.05.

Virus Family	Dataset	AIC Score	AIC Score	AIC Score	P-Value	P-Value	DNR
		GTR	NREV-6	NREV-12	GTR vs NREV-12	NREV-6 vs NREV-12	
Papillomaviridae	APPV 6	35099.5	35108.0	35102.2	>0.05	0.007	0.089
	HPV18_2	25202.9	25174.6	25179.2	<0.001	>0.05	0.323
	HPV45_2	23600.6	23599.0	23602.9	>0.05	>0.05	0.285
	HPV16_2	29734.0	29664.5	29665.4	<0.001	>0.05	0.371
	HPV31	24681.4	24677.3	24672.8	0.002	0.01	0.165

Table 1

AIC Scores and LRT results for double-stranded DNA virus datasets.

The lowest AIC-c scores indicating the bestfitting models are in bold.

	HPV6_1	31199.1	31150.0	31141.2	<0.001	<0.001	0.451
	LPV	67165.7	67188.1	67145.5	<0.001	<0.001	0.42
	DPV	69829.7	69889.2	69835.1	>0.05	<0.001	0.056
	XPV	95455.6	95617.1	95452.2	<0.001	<0.001	0.072
	BATV	134821	134511	133322	<0.001	<0.001	0.402
Polyomaviridae	JC_2	51806.7	51819.6	51812.0	>0.05	0.003	0.089
	BK_2	21472.6	21472.7	21471.1	0.03	0.03	0.244
	SV40	16859.8	16858.0	16858.4	0.037	>0.05	0.567
	BPV	148614.9	148573.8	148585.2	<0.001	>0.05	0.064
Caulimoviridae	CMV	124083.9	124221.0	123888.6	<0.001	<0.001	0.351
	CSSV	145327.0	146575	145202	<0.001	<0.001	0.158
	SVBV	138575	138488.1	138464.7	<0.001	<0.001	0.174
	DBAV	46495.5	46514.1	46502.0	>0.05	<0.001	0.0335
	RTBV	54987.9	55350.1	54991	>0.05	<0.001	0.082
	BDV	376325.2	376647.6	376029.9	<0.001	<0.001	0.140
Siphoviridae	CLV	237362.3	237351.8	237348.6	<0.001	<0.01	0.070
<u>Tectiviridae</u>	TTIV	913864.9	913915.4	913773.1	<0.001	<0.001	0.279
Adenoviridae	FAV_C	3074086.7	3074207.5	3073739.1	<0.001	<0.001	0.169
	FAV_E	103482.3	103222.7	102636.7	<0.001	<0.001	0.357
	FAV_D	2326925.6	2325719.4	2324784.5	<0.001	<0.001	0.551
	FAV_A	705328.5	705436..5	705197.8	<0.001	<0.001	0.645
	HMAV_B	103796.7	103937.44	103753.8	<0.001	<0.001	10.890
	HMAV_D	1748635.2	1749769	1748119.1	<0.001	<0.001	0.646
	HMAV_C	2851144.5	2851357.1	2851133	0.006	<0.001	0.0225
	HMAV_E	1915044.8	1915065.3	1914998	<0.001	<0.001	0.049

Table 1 (continued)

2. extra exposure to DNA damage of displaced template strands during unidirectional rolling circle replication in some papillomavirus species such as HPV16 could be a contributor to strand-specific nucleotide substitution biases (Kusumoto-Matsuo, Kanda, & Kukimoto, 2011 [\[link\]](#)).
3. extra time spent by non-coding strands in single-stranded dissociated states during RNA transcription in some papillomavirus and polyomavirus species (Fernandes & de Medeiros Fernandes, 2012 [\[link\]](#)): during transcription processes, the dissociated non-coding strand is transiently more exposed to damage than the coding strand (Wei, et al., 2010 [\[link\]](#)) which might also contribute to strand-specific substitution biases.

Similarly, and equally surprising, we found that NREV12 was overall the best supported model for dsRNA viruses (illustrated by the redder hues around the top corner of the dsRNA plot in **Figure 1** [\[link\]](#)).

NREV6 fit only two of the 31 dsRNA datasets better than both NREV12 and GTR (Human rotavirus A set H and Fiji virus). NREV12 was found to be the best fitting model for 21/31 datasets and GTR was the best fitting of 8/31 (**Table 2** [\[link\]](#)). In all three Birnaviridae family datasets (which contains virus species with two genome segments) and in 17/22 of Reoviridae family datasets (which contain virus species with 10-12 genome segments) the NREV12 model provided the best fit. Based on the LRTs, strong overall support for NREV12 was found, with this model providing a significantly better fit ($p < 0.05$) in 27/31 dsRNA virus datasets relative to NREV6, and 23/31 datasets relative to GTR.

We anticipated that NREV12 might fit many of these dsRNA datasets better than NREV6 simply because, during their infection cycles, the coding +strand of dsRNA viruses (the one from which protein translation occurs) tends to exist for longer periods within an infected cell than the noncoding -strand. Specifically, there are two main steps during double-stranded RNA virus replication (Wickner, 1993 [\[link\]](#)). Firstly, synthesis of the viral +strands from a dsRNA template occurs in the cytoplasm within viral particles. These +strands exist within the cell for prolonged periods in the absence of complementary -strands and are used as templates for translation of viral proteins. In the second step the +strands remaining after translation act as templates for -strand synthesis resulting in the formation of new dsRNA molecules. The +strands of dsRNA viruses are therefore likely more impacted by mutational processes, which in turn could explain the pervasive strand specific substitution biases seen in this group of viruses.

For the ssRNA and ssDNA viruses where one genome strand exists during the virus life cycle for far longer periods of time than the other such that complementary substitutions would not be expected to occur at similar rates, we anticipated that NREV12 should provide a better fit than both NREV6 and GTR. Indeed, for ssRNA viruses, NREV12 was a better fit than NREV6 and GTR for 40/47 of the ssRNA datasets and 24/33 of the ssDNA virus datasets (**Figure 1** [\[link\]](#)). Of the nine ssDNA virus datasets where NREV12 was not the best fitting model, GTR fit 7/9 better and NREV6 fit 2/9 better (**Table 3** [\[link\]](#)). Of the seven ssRNA datasets where NREV12 was not the best fitting model, GTR fit 6/7 better and NREV6 fit 1/7 better.

Based on the LRTs, strong overall support for NREV12 was found with this model providing a significantly better fit ($p < 0.05$) than NREV6 for 45/47 of the ssRNA virus datasets (**Table 4** [\[link\]](#)) and 31/33 of the ssDNA virus datasets (**Error! Reference source not found.**). Similarly, based on LRTs, NREV12 provided a significantly better fit than GTR for 27/33 of the ssDNA virus datasets (**Error! Reference source not found.**) and 40/47 of the ssRNA virus datasets (**Table 4** [\[link\]](#)).

We found that DNR estimates alone did not cleanly differentiate between datasets for which NREV12 was or was not best supported (**Tables 1** [\[link\]](#), **2** [\[link\]](#), **3** [\[link\]](#) and **4** [\[link\]](#)). For the 107 nucleotide sequence datasets with a model preference of NREV12, ten had estimated DNRs that were greater than 0.5, 13 had DNRs between 0.25 and 0.5 and 84 had DNRs between 0.0225 and 0.25. For the ten

Virus Family	Dataset	AIC score GTR	AIC score NREV-6	AIC score NREV-12	GTR vs NREV-12	NREV-6 vs NREV-12	DNR
Birnaviridae	AQBV	31754.9	31853.3	31721.9	<0.001	<0.001	0.219
	GBV_A	47176.9	47347.2	47154.8	<0.001	<0.001	0.142
	IPNV	79186.2	79221.9	79182.4	0.0145	<0.001	0.162
	GBV_B	39313.7	39062.8	38938.7	<0.001	<0.001	0.201
Reoviridae	BTv_A	34803.5	34895.1	34801.3	0.03	<0.001	0.042
	BTv_B	48849.9	48893.	48837.1	<0.001	<0.001	0.043
	BTv_C	28350.9	28386.5	28350.8	>0.05	<0.001	0.061
	BTv_D	24969.1	24947.3	24894.0	<0.001	<0.001	0.191
	BTv_F	20622.7	20708.5	20610.2	<0.001	<0.001	0.067
	BTv_G	63349.9	63485.0	63345.9	0.00426	<0.001	0.040
	BTv_H	20596.7	20685.5	20586.1	<0.001	<0.001	0.118
	BTv_I	17592.7	17622.5	17588.8	0.01	<0.001	0.095
	BRVA_C	41206.7	41187.4	41137.1	<0.001	<0.001	0.128
	HRVA_A	17030.5	17043.2	17035.5	>0.05	0.003	0.036
	HRVA_B	8275.1	8280.3	8281.7	>0.05	>0.05	0.087
	HRVA_C	12815.1	12842.6	12807.6	0.003	<0.001	0.132
	HRVA_D2	8036.8	8041.0	8043.7	>0.05	>0.05	0.057
	HRVA_E	7045.9	7056.1	7053.3	>0.05	0.02	0.102

Table 2

AIC Scores and LRT results for double-stranded RNA datasets.

The lowest AIC-c scores indicating the best fitting models are in bold.

	HRVA_F	7046.0	7056.7	7053.4	>0.05	0.02	0.0710
	HRVA_G	18424.2	18434.0	18425.1	>0.05	<0.001	0.123
	HRVA_H	20431.4	20413.87	20420.5.6	0.002	>0.05	0.163
	PRVA_A	28540.7	28441.9	28398.7	<0.001	<0.001	0.204
	PRVA_B	14757.7	14775.5	14732.6	<0.001	<0.001	0.351
	HRVC_A	6713.2	6718.2	6712.3	0.045	0.007	0.124
	PTOV	202011.3	202106.5	201878.5	<0.001	<0.001	0.039
	FJV_B	9274.1	9250.0	9250.9	<0.001	>0.05	0.194
Endornaviridae	EDV	1771992.8	1772689.1	1771950.6	<0.001	<0.001	0.121
	BPAV	70386.5	70540.2	70390.7	>0.05	0.00	0.047
Totiviridae	TTV	617302.6	617462.6	617172.9	<0.001	<0.001	0.052
	GDV	80435.8	80396.5	80387.7	<0.001	0.002	0.109
Hypoviridae	HPV	66859.8	66899.8	66857.8	0.03	<0.001	0.057

Table 2 (continued)

Virus Family	Dataset	AIC Score GTR	AIC Score NREV-6	AIC Score NREV-12	P-Value GTR vs NREV-12	P-Value NREV-6 vs NREV-12	DNR
Nanoviridae	BBTV M	15044.3	15207.9	14984.4	<0.001	<0.001	0.662
	BBTV N	10605.6	10686.2	10595.2	<0.001	<0.001	0.533
	BBTV R	18484.5	18544	18480.8	>0.05	<0.001	0.609
	BBTV S	12718.9	12757.2	12707.3	<0.001	<0.001	0.728
	CCDV	38622.7	38632.0	38630.5	>0.05	0.03	0.050
	MDV	36232.8	36063	36064	<0.001	>0.05	0.142
	PYDV	56138.4	56076.6	56056.4	<0.001	<0.001	0.187
	FBNS	100153.6	100135.6	100120.5	<0.001	<0.001	0.098

Table 3

AIC-c Scores and LRT results for single-stranded DNA datasets.

The lowest AIC-c scores indicating the best fitting models are in bold.

Geminiviridae	Begomo 5	28192.1	28311.9	28192.5	>0.05	<0.001	0.1995
	Begomo 6	16743.0	16722.6	16724.1	<0.001	>0.05	0.214
	Begomo 9	8517.6	8540.8	8515.6	0.03	<0.001	0.312
	Dicot 1	44730.7	44594.3	44583.3	<0.001	<0.001	0.200
	Dicot 2	39909.9	39919.8	39917.9	>0.05	<0.001	0.100
	MSV	252645.3	254347.5	254347.5	<0.001	<0.001	0.144
	PanSV	94601.2	94600.3	94593.7	<0.001	<0.001	0.182
	WDV	35301.7	35313.2	35253.8	<0.001	<0.001	0.1033
Circoviridae	BFDV	17256.7	17262.7	17246.7	<0.001	<0.001	0.224
	DG_CV	12754.8	12779.5	12758.3	>0.05	<0.001	0.116
	PiCV	19180.5	19192.5	19191.0	>0.05	0.04	0.117
	CCCC	84435.7	84377.4	84315.3	<0.001	<0.001	0.132
	BTC	262910.4	262060.1	261985.4	<0.001	<0.001	0.178
	POCV2	24940.9	24953.8	24915.8	<0.001	<0.001	0.162
	CCV	90307.9	90301.5	90285.9	<0.001	<0.001	0.114
Anelloviridae	TTV_1	825811	826800	825292	<0.001	<0.001	0.513
	TTSV	332287.9	332397.4	332258.2	<0.001	<0.001	1.560
Parvoviridae	MVM	26756.3	26743.9	26686.9	<0.001	<0.001	0.148
	HPV	67051.2	67080.1	67001.8	<0.001	<0.001	0.235
	CPV	85731	85695	85689.3	<0.001	0.007	0.062
	PPV	163006.8	163090.7	162995.9	<0.001	<0.001	0.143
	CAV_P	37073.3	37115.5	37065.7	<0.001	<0.001	0.162
Microviridae	BMV	31175.3	31164.8	31147.3	<0.001	<0.001	0.188
Pleolipoviridae	APV	85700.2	85617.4	85402.8	<0.001	<0.001	0.204
	BPV	204797.5	204802.3	204796.7	0.04	0.007	0.064

Table 3 (continued)

Virus Family	Dataset	AIC Score GTR	AIC Score NREV-6	AIC Score NREV-12	P-Value GTR vs NREV-12	P-Value NREV-6 vs NREV-12	DNR
Astroviridae	HAV	94580.7	94926.3	94548.1	<0.001	<0.001	0.096
	BAV	188307.1	188572.9	188144.9	<0.001	<0.001	0.108
	MMV	281072.2	281094.5	281076.9	>0.05	<0.001	0.072
	PAV	150626.88	150827.6	150609.5	<0.001	<0.001	0.069
	CKV	90902.3	91233.1	90873.0	<0.001	<0.001	0.083

Table 4

AIC-c Scores and LRT results for single-stranded RNA datasets.

The lowest AIC-c scores indicating the best fitting models are in bold.

	GA	64998.5	65223.9	64975.9	<0.001	<0.001	0.110
	CAV_A	85558.8	85617.4	85547.3	<0.01	<0.001	0.076
Bromoviridae	CMV RNA1	34197.5	34198.8	34147.7	<0.001	<0.001	0.124
	CMV RNA2	31398.2	31455.9	31388.7	<0.001	<0.001	0.091
	CMV RNA3	24337.2	24360.3	24343.9	>0.05	<0.001	0.073
	AMS	24337.2	24360.3	24343.9	>0.05	<0.001	0.073
	PSV	67707	67786.5	67691	<0.001	<0.001	0.048
Caliciviridae	LAV	73042.8	73102.4	72984.6	<0.001	<0.001	0.120
	NoV	207667.2	207777.5	207660	<0.001	<0.001	0.047
	VSV	235936.4	236051.4	235913.3	<0.001	<0.001	0.046
Closteroviridae	CTV	30062.2	29980.4	29960.1	<0.001	<0.001	0.272
Flaviviridae	DGV_T1	69771.9	70030.5	69776.2	>0.05	<0.001	0.063
	JEV	146920.8	148101.5	146885.5	<0.001	<0.001	0.091
Hepeviridae	HPVE1	200439.5	200863.8	200179.8	<0.001	<0.001	0.073
	HPVE2	155709.1	155983.8	155518.6	<0.001	<0.001	0.088
Picornaviridae	ENV_A	552287.9	553535.5	551794.1	<0.001	<0.001	0.061
	HRV_A	102218.7	102267.0	101550.7	<0.001	<0.001	0.285
	AIV	101073.1	101136.7	101052.2	<0.001	<0.001	0.093
	AHP	139635.7	140119.6	139506.9	<0.001	<0.001	0.170
	ECV	82078.9	82181.0	82065.8	<0.001	<0.001	0.066
	CDV	130551.3	130896.7	130478.3	<0.001	<0.001	0.086
	TCV	53027.3	53029	53023	0.0151	0.0422	0.033
	FMDV	455180.6	455582.6	454806.1	<0.001	<0.001	0.117
Fusariviridae	FRV	52413.1	52470.6	52418.4	>0.05	<0.001	0.076
Retroviridae	HIV1_setA	344014.4	344295.1	343669.7	<0.001	<0.001	0.237
	HIV1_M	80764.1	80829.5	80668.1	<0.001	<0.001	0.442
	HIV1_setC	180575.0	180702.3	180494.4	<0.001	<0.001	0.107

Table 4 (continued)

	HIV1_setD	298489.9	298695.3	298260.6	<0.001	<0.001	0.133
	HIV1_setE	289111.3	289292.1	288941.9	<0.001	<0.001	0.112
	HIV1_setF	214375.9	214692.2	214289.4	<0.001	<0.001	0.148
	EIV	126149	126365.4	125300	<0.001	<0.001	0.192
	BIV	24505.2	24506.9	24513.2	>0.05	>0.05	0.15
	FIV	164542.1	164487.9	164260.4	<0.001	<0.001	0.114
	CAV	351329.9	351871.5	350721.9	<0.001	<0.001	0.174
	SIV	110731.2	110816	110663.3	<0.001	<0.001	0.144
Filoviridae	Ebola_2	53147.3	53143.0	53149.9	>0.05	>0.50	0.264
Orthomyxo-viridae	Flu A 2	82872.8	83010.2	82849.7	<0.001	<0.001	0.27
	Flu B 1	50090.4	50144.1	50060.9	<0.001	<0.001	0.311
Coronaviridae	SARS-COV1	214715.3	214968.5	214644.39	<0.001	<0.001	0.198
	SARS-COV2	15715.4.2	15715.6	15696.7	<0.001	<0.001	1.536
	SARB	573966.3	573815.1	572517.0	<0.001	<0.001	0.301
	MERS-COV	516683.2	516983.4	516608.9	<0.001	<0.001	0.169

Table 4 (continued)

nucleotide sequence datasets with a model preference of NREV6, one had an estimated DNR greater than 0.5, four had estimated DNRs between 0.25 and 0.5 and five had estimated DNRs between 0.064 and 0.25 (**Figure 1** [↗](#)). For the 24 nucleotide sequence datasets with a model preference of GTR, none had estimated DNRs greater than 0.5, one had an estimated DNR between 0.25 and 0.5 and the remainder had estimated DNRs between 0.0335 and 0.25.

The dsDNA virus dataset with the highest DNR was Human mastadenovirus D (DNR = 0.646), the dsRNA virus dataset with the highest estimated DNR was Porcine_rotavirus_B (0.351), the ssRNA virus dataset with the highest DNR was SARS-CoV-2 (DNR = 1.536) and the ssDNA virus dataset with the highest DNR was Torque teno sus virus (DNR = 1.56).

Therefore, while NREV12 appears to be generally more appropriate than either NREV6 or GTR for describing mutational processes in ssRNA, ssDNA, dsDNA, and dsRNA viruses, this might only be particularly relevant from a practical perspective when datasets of these viruses yield DNR estimates that are greater than 0.25. For such datasets, NREV12 (and possibly NREV 6 in some instances) might be especially useful for both determining the direction of evolution across phylogenetic trees (i.e., it could potentially be used to root these trees) and for quantifying genomic strand-specific nucleotide substitution biases ([Harkins et al., 2009](#) [↗](#)).

Assessing the impacts of model misspecification on phylogenetic tree inference

To determine whether it is worthwhile to use NREV12 rather than GTR for phylogenetic inference when NREV12 is the best fitting nucleotide substitution model, we used simulated datasets to compare the accuracy of phylogenetic trees inferred using these models.

We found that, regardless of dataset diversity and the nucleotide substitution model used, phylogenetic inference tended to become less accurate (i.e. wRF scores increased) as DNR increased (**Figure 2** [↗](#)). This tendency was, however, more pronounced when using a (mis-specified) GTR model than when using a (correctly specified) NREV12 model with, for any given dataset having DNR > 0, the use of NREV12 tending to yield more accurate phylogenetic trees than when GTR was used. There were, however, only statistically significant improvements ($p < 0.05$ paired t-test) in the accuracy of phylogenetic trees inferred using NREV12 relative to those inferred using GTR in lower diversity datasets (i.e. those with APIs of 85%, 90% and 95%); and then only for DNR > 8. It is noteworthy that the highest estimated DNR in any of the empirical datasets that we analysed was 1.56 more than four-fold lower than the point where statistically significant differences in phylogenetic-inference accuracies became apparent in the simulated datasets.

Materials and Methods

Virus sequence datasets and phylogenetic trees

We obtained viral nucleotide sequences from the National Centre for Biotechnology Information Taxonomy database (<http://www.ncbi.nlm.nih.gov/taxonomy> [↗](#)) and the Los Alamos National Laboratory HIV sequence database (<https://www.hiv.lanl.gov/content/index> [↗](#)). These included gene and whole-genome sequences for viruses with ssRNA, ssDNA, dsRNA, and dsDNA genomes (datasets are summarized in **Error! Reference source not found.**). An outgroup sequence from a closely related virus species was added to each dataset to help root phylogenetic trees. The sequences in each of the datasets were aligned using MUSCLE ([Edgar, 2004](#) [↗](#)) implemented in Aliview ([Larsson, 2014](#) [↗](#)), and maximum likelihood phylogenetic trees were constructed from each alignment using RAXML v8.2 ([Stamatakis, 2016](#) [↗](#)).

Model Testing

We evaluated the fit of NREV12, NREV6, and GTR to the 141 individual sequence datasets using a previously published model test (Harkins, et al., 2009 [↗](#)) implemented as a HyPhy package module (Pond, Frost, & Muse, 2005 [↗](#)). This script (<https://github.com/veg/hyphy-analyses/tree/master/NucleotideNonREV> [↗](#)) and is also available as a module in the Datamonkey web server (Weaver, et al., 2018) which takes as input a rooted maximum likelihood phylogenetic tree (minus the rooting sequence) and its corresponding nucleotide sequence alignment. The three models described above: GTR, NREV6, and NREV12 are then fitted to the data using maximum likelihood (ML). The equilibrium frequencies (EF) of the GTR model match those empirically observed in the alignment, while EFs for NREV6 and NREV12 are induced by the model, satisfying the condition $\pi Q = 0$. An additional model NREV12 +F is estimated, where the distribution of nucleotides at the root of the tree is estimated by ML, instead of being set to the empirical frequencies. Nested models were compared by the likelihood ratio test (LRT) with significance evaluated using the χ^2_d distribution with d = difference in degrees of freedom. For all models we also computed the small sample corrected Akaike Information Criterion (AIC-c) score.

Quantification of non-reversibility

We further defined the degree of non-reversibility (DNR) as the absolute difference between the relative rate differences of two nucleotide pairs; i.e. for two nucleotides, x and y, there exists a relative rate of x to y substitutions that we will refer to as m , and a relative rate y to x substitutions that we will refer to as n . Under the NREV12 model, the degree of non-reversibility (DNR) between x and y is defined simply as the absolute difference between m and n : $|m - n|$ and will hereby be referred to as ij_{DNR} where i and j are two nucleotides. We use DNR as a mathematical representation of the degree to which the rates of all pairs of reverse substitutions differ from one another. For each of the 140 individual viral alignments we calculated the average DNR of the six ij_{DNR} estimates inferred using the NREV12 model.

Simulations for testing the impact of non-reversible evolution on phylogenetic inference

We tested the accuracy of phylogenetic tree inference under reversible and non-reversible models using simulated datasets with varying average pairwise nucleotide sequence identities (APIs) evolved under the NREV12 model with different degrees of non-reversibility (DNR). The goal of these tests was not to exhaustively evaluate model misspecification issues during phylogenetic tree inference, but rather to check, in instances where viral taxa are known to be evolving in a detectably non-reversible manner (i.e. where NREV12 or NREV6 fits the data better than GTR), whether not accounting for this might decrease the accuracy of phylogenetic inference. Using IQTREE, a phylogenetic inference program that has the option to apply an NREV12-like model (referred to in IQTREE as the UNREST model), a phylogenetic tree was inferred from an alignment of real sequences (Avian Leukosis virus) (**Figure 3** [↗](#)) with an average sequence identity (API) of ~90%. The branch lengths on this tree were then scaled to create four other phylogenetic trees representing sequences with approximately 95%, 85%, 80% and 75% API. These five trees are hereafter referred to as “true” trees and each individual tree was used as the starting point of a different set of simulations.

Phylogenetic trees were inferred from these 5500 simulated datasets and compared to the phylogenetic trees used to simulate the datasets (i.e., the true trees) using wRF distances to assess the impact of varying DNR on the accuracy of phylogenetic inference. We further tested whether the accuracy of phylogenetic inference could be improved for sequences that had evolved under $DNR > 0$ by using NREV12 instead of GTR. Specifically, for every simulated dataset a phylogenetic tree was inferred using GTR and another using NREV12 and the wRF distances of each of these trees to the true tree was determined. For each of the analysed DNRs a paired t-test was then used

to compare the wRF scores of trees inferred using GTR and NREV12. We were particularly interested in determining whether trees inferred using a mis-specified model (i.e. GTR in this case) would be less accurate than trees inferred with a correctly specified model (i.e. NREV12).

To test whether failure to account for non-reversibility might decrease the accuracy of phylogenetic inference we simulated the evolution of 5,500 nucleotide sequence alignments evolved non-reversibly under varying DNR along the five true phylogenetic trees: 100 datasets per true tree per simulated degree of non-reversibility (DNR). Specifically, simulations were done using HyPhy (Pond, Frost, & Muse, 2005 [\[4\]](#)) with relative rates ranging from a completely reversible matrix (equation (4) [\[4\]](#))

$$Q = \{q_{ij}\} = \begin{pmatrix} - & 0.166 & 1 & 0.14 \\ 0.166 & - & 0.131 & 1.101 \\ 1 & 0.131 & - & 0.188 \\ 0.14 & 1.101 & 0.188 & - \end{pmatrix} \quad (4)$$

representing DNR=0 - through matrices with DNR=2, 4, 6, 8, 10, 12, 14, 16, 18 and 20 (Table 5 [\[4\]](#)). These baselines simulated substitution rates are reflective of those seen in empirical viral nucleotide sequence datasets.

At each DNR, the relative rates used conformed to standard measures of non-reversibility under the Kolmogorov conditions according to which non-reversibly evolving sequence datasets should yield three irreversibility indices (IRI1, IRI2 and IRI3) that are all non-zero (Squartini & Arndt, 2008 [\[4\]](#)). It should be noted that all simulations under NREV12 were performed under the stationarity criterion: $\pi e^{Qt} = \pi$ (where Q is the rate matrix and π is the nucleotide frequency distribution and $t \geq 0$).

Quantifying the accuracy of phylogenetic inferences

We used the weighted Robinson-Foulds metric (wRF; implemented in the R phangorn package (Schliep, 2011 [\[4\]](#))) to quantify differences between the true trees used to simulate datasets and the trees inferred from these datasets using the GTR or NREV12 models. wRF considers differences between both the topology and branch lengths of actual and inferred trees (Kuhner, 2015 [\[4\]](#)), (Robinson & Foulds, 1981 [\[4\]](#)).

Conclusion

The non-reversible nucleotide substitution model, NREV12, provides a substantially better fit to most virus nucleotide sequence datasets than does the widely used reversible substitution model, GTR. NREV12 also provides a better fit to most virus nucleotide sequence datasets than does NREV6; a non-reversible model that would be expected to best describe the evolution of double stranded genome sequences that display no strand-specific nucleotide substitution biases. This suggests that, contrary to our expectations, substantial strand-specific nucleotide substitution biases (i.e. estimated DNRs > 0.25) are common during viral evolution irrespective of genome type. Such biases should be expected for any viruses where one genome strand either is in existence for substantially longer periods of time than the other, or is more exposed to mutagenic processes than the other during transmission, replication, or gene expression.

We had anticipated that, given evidence of sequences evolving both non-reversibly and with strand-specific substitution biases, inferring trees using a model such as NREV12 that appropriately accounts for this might: (1) minimize the impact of increasing DNR on the accuracy

Table 5

Relative rate change for C to A, G to A, A to T, G to C, T to G and C to T mutations under the 11 degrees of non-reversibility alongside the maintained rates for A to C, A to G, T to A, C to G, G to T, and T to C.

Degree of Non-Reversibility (DNR)	Relative rates of different nucleotide substitution types (from-to)											
	C-A	A-C	G-A	A-G	A-T	T-A	G-C	C-G	T-G	G-T	C-T	T-C
0	0.166	0.166	1	1	0.14	0.14	0.131	0.131	0.118	0.118	1.101	1.101
2	2.166	0.166	3	1	2.14	0.14	2.131	0.131	2.118	0.118	3.101	1.101
4	4.166	0.166	5	1	4.14	0.14	4.131	0.131	4.118	0.118	5.101	1.101
6	6.166	0.166	7	1	6.14	0.14	6.131	0.131	6.118	0.118	7.101	1.101
8	8.166	0.166	9	1	8.14	0.14	8.131	0.131	8.118	0.118	9.101	1.101
10	10.166	0.166	11	1	10.14	0.14	10.131	0.131	10.118	0.118	11.101	1.101
12	12.166	0.166	13	1	12.14	0.14	12.131	0.131	12.118	0.118	13.101	1.101
14	14.166	0.166	15	1	14.14	0.14	14.131	0.131	14.118	0.118	15.101	1.101
16	16.166	0.166	17	1	16.14	0.14	16.131	0.131	16.118	0.118	17.101	1.101
18	18.166	0.166	19	1	18.14	0.14	18.131	0.131	18.118	0.118	19.101	1.101
20	20.166	0.166	21	1	20.14	0.14	20.131	0.131	20.118	0.118	21.101	1.101

of phylogenetic inference (i.e. wRF scores presented in blue in **Figure 2** [↗](#) might have been expected to not increase with increasing DNR); and (2) yield significantly more accurate phylogenetic inferences than when using GTR for all datasets where NREV12 was the most appropriate model and DNRs were greater than zero. However, increasing DNR clearly decreased the accuracy of phylogenetic inference even when using NREV12, and, for datasets where DNRs were greater than zero, using GTR did not consistently yield significantly less accurate phylogenetic inferences than those attained using NREV12. From a practical perspective, choosing a non-reversible nucleotide substitution model to construct phylogenetic trees from virus genome sequences that display strand-specific nucleotide substitution biases is not guaranteed to yield more accurate phylogenetic trees. Nevertheless, in instances where strand-specific substitution biases are higher than ~0.5 (such as are found in our SARS-CoV-2, Torque teno sus virus, and Banana bunchy top virus datasets) it may be prudent to select a model such as NREV12 (such as is implemented in programs like IQTREE) over GTR as the better of two suboptimal choices.

The lack of available data regarding the proportions of viral life cycles during which genomes exist in single and double stranded states makes it difficult to rationally predict the situations where the use of models such as GTR, NREV6 and NREV12 might be most justified: particularly in light of the poor over-all performance of NREV6 and GTR relative to NREV12 with respect to describing mutational processes in viral genome sequence datasets. We therefore recommend case-by-case assessments of NREV12 vs NREV6 vs GTR model fit when deciding whether it is appropriate to consider the application of non-reversible models for phylogenetic inference and/or phylogenetic model-based analyses such as those intended to test for evidence of natural selection or the existence of molecular clocks.

Data availability

All sequence alignments used during model testing in this study are included in the manuscript supplementary table one. These datasets were obtained from nucleotide sequences from the National Centre for Biotechnology Information Taxonomy database (<http://www.ncbi.nlm.nih.gov/taxonomy>) and also obtained from the Los Alamos National Laboratory HIV sequence database (<https://www.hiv.lanl.gov/content/index>). These sequence alignments have been made available as supporting files datasets.

Supplementary Files

Genome Type	Virus Family	Virus Genus	Virus Species	Dataset Name
ssDNA	Circoviridae	Circovirus	Beak and feather disease virus	BFDV
			Duck circovirus, Goose circovirus	DG_CV
			Columbine circovirus	PICV
			Circovirus	CCCC
			Bat circovirus	BTC
			Porcine Circovirus 2	POCV2
			Cyclovirus	CCV
	Geminiviridae	Begomovirus	East Africa cassava mosaic virus, South African cassava mosaic virus	Begomo6
			Tomato yellow leaf curl virus	Begomo5
			Malvastrum yellow vein Yunnan virus, Cotton leaf curl Multan virus, Bhendi yellow vein India virus	Begomo9
			Tobacco yellow dwarf virus, Chickpea chlorosis virus, Chickpea yellows virus	Dicot_1
		Mastrevirus	Chickpea chlorotic dwarf virus	Dicot_2
			Maize streak virus	MSV
			Panicum streak virus	PanSV
			Wheat dwarf virus	WDV
	Anelloviridae	Anellovirus	Torque teno virus 1	TTV_1
			Torque teno sus virus	TTSV
	Parvoviridae	Aneptorquevirus	minute virus of mice, MVM	MVM
		Protoparvovirus	Human parvovirus	HPV
			Canine parvovirus	CPV
		Amdoparvovirus	porcine parvovirus	PPV
			Carnivore amdoparvovirus	CAV_P
	Nanoviridae	Babuvirus	Banana bunchy top virus	BBTV_M
				BBTV_N
				BBTV_R
				BBTV_S
		Nanovirus	Coconut foliar decay virus	CCDV
			Milk vetch dwarf virus full genome	MDV
			Pea necrotic yellow dwarf virus	PYDV
			Faba bean necrotic stunt virus	FBNS
	Microviridae	Microviruses	Microvirus	BMV
	Pleolipoviridae	Pleolipoviruses	Betapleolipovirus	BPV
			Alphapleolipovirus	APV
ssRNA	Astroviridae	Astroviruses	Human astrovirus	HAV
			Bovine astrovirus	BAV

Table S1.

Full details of the datasets used in the study.

			Mamastrovirus	MMV
			porcine astrovirus	PAV
			chicken astrovirus	CKV
			Goose astrovirus	GA
			Canine astrovirus	CAV_A
	Bromoviridae	Cucumovirus	Cucumber mosaic virus	CMV_RNA1
				CMV_RNA2
				CMV_RNA3
		Alphamovirus	Alphalfa mosaic virus	AMS
		Cucumovirus	Peanut stunt virus	PSV
	Caliciviridae	Lagovirus	Lagovirus	LAV
		coroviruses	Norovirus	NoV
		Vesivirus	Vesivirus	VSV
	Closteroviridae	Closterovirus	Citrus tristeza virus	CTV
	Flaviviridae	Flavivirus	Dengue virus	DGV_T1
			Japanese encephalitis virus	JEV
	Hepeviridae	Hepevirus	Hepatitis E virus	HPVE1
			Hepatitis E2 virus	HPVE2
	Picornaviridae	Enterovirus	Human Rhinovirus A	HRV_A
			Enterovirus A	ENV_A
		Teschovirus	Techovirus	TCV
		Aichivirus	Aichivirus	AiV
		Aphthovirus	Foot and mouth disease virus	FMDV
			Avihepatovirus	AHP
		Cardiovirus	Encephalomyo carditis virus	ECV
			Cardiovirus	CDV
	Fusariviridae	Fusarivirus	Fusariviruses	FRV
	Retroviridae	Lentivirus	Human immuno-deficiency virus 1	HIV1_setA
				HIV1_M
				HIV1_setC
				HIV1_setD
				HIV1_setE
				HIV1_setF
			Simian immuno-deficiency virus	SIV
			Bovine immunodeficiency	BIV
			Feline immunodeficiency	FIV
			equine infectious anemia virus	EIV
			caprine arthritis encephalitis virus	CAV
	Orthomyxo-viridae	Influenzavirus	Influenza virus A	FluA_2
			Influenza virus B	FluB_1
	Filoviridae	Ebolavirus	Ebola virus	Ebola_2
	Coronaviridae	Merbecovirus	Middle East respiratory syndrome	MERS-COV

Table S1. (continued)

		Sarbecoviruses	Severe acute respiratory syndrome coronavirus 1	SARS-COV1
			Severe acute respiratory syndrome coronavirus 2	SARS-COV2
			sarbecoviruses	SARB
dsDNA	Papillomaviridae	Alphapapillomavirus	Alphapapillomavirus 6	APPV 6
			Alphapapillomavirus 7	HPV18_2
				HPV45_2
			Alphapapillomavirus 9	HPV16_2
				HPV31
			Alphapapillomavirus 10	HPV6_1
			Bovine papillomavirus	BPV
			Lambdaapapillomavirus	LPV
	Polyomaviridae	Polyomavirus	Deltapapillomavirus	DPV
			Xipapillomavirus	XPV
			BK polyomavirus	BK_2
			JC polyomavirus	JC_2
	Caulimoviridae	Caulimovirus	Bat polyomavirus	BPV
			Simian virus 40	SMV_40
			cauliflower mosaic virus	CMV
			Cacao swollen shoot virus	CSSV
			Strawberry vein banding virus	SVBV
			Dioscorea bacilliform AL virus	DBAV
			Rice tungro bacilliform virus	RTBV
	Siphoviridae	Escherichia virus Lambda	badnavirus	BDV
	Tectiviridae	Tectivirus	coliphage lambda	CLV
	Adenoviridae	Aviadenovirus	Tectivirus	TTIV
			Fowl aviadenovirus C	FAV_C
			Fowl aviadenovirus E	FAV_E
			Fowl aviadenovirus A	FAV_A
		Mastadenovirus	Fowl aviadenovirus D	FAV_D
			Human mastadenovirus B	HMAV_B
			Human mastadenovirus D	HMAV_D
			Human mastadenovirus C	HMAV_C
			Human mastadenovirus E	HMAV_E
DsRNA	Birnaviridae	Avibirnavirus	Gumburo virus_setA	GBV_A
		Aquabirnavirus	Gumburo virus_setB	GBV_B
			Infectious pancreatic necrosis virus	IPNV
	Reoviridae	Orbivirus	Aquabirnavirus	AQBV
			Bluetongue_virus_setA	BTv_A
			Bluetongue_virus_setB	BTv_B
			Bluetongue_virus_setC	BTv_C
			Bluetongue_virus_setD	BTv_D

Table S1. (continued)

			Bluetongue_virus_setF	BTV_F
			Bluetongue_virus_setG	BTV_G
			Bluetongue_virus_setH	BTV_H
			Bluetongue_virus_setI	BTV_I
		Rotavirus	Bovine_rotavirus_A_setC	BRVA_C
			Human_rotavirus_A_setA	HRVA_A
			Human_rotavirus_A_setB	HRVA_B
			Human_rotavirus_A_setC	HRVA_C
			Human_rotavirus_A_setD2	HRVA_D2
			Human_rotavirus_A_setE	HRVA_E
			Human_rotavirus_A_setF	BRVA_F
			Human_rotavirus_A_setG	HRVA_G
			Human_rotavirus_A_setH	HRVA_H
			Porcine_rotavirus_A_setA	PRVA_A
			Porcine_rotavirus_A_setB	PRVA_B
			Human_rotavirus_C_setA	HRVC_A
		Orthoreovirus	Pteropine orthoreovirus	PTOV
		Fijivirus	Fijivirus_setB	FJV_B
	Totiviridae	Totivirus	Totivirus	TTV
		Giardiavirus	Giardiavirus	GDV
	Hypoviridae	Hypovirus	Hypovirus	HPV
	Endornaviridae	Endornavirus	Endornavirus	EDV
		Alphaendornavirus	Bell pepper alphaendornavirus	BPAV

Table S1. (continued)

Acknowledgements

The authors wish to thank the South African National Research Foundation (NRF) for funding this research under the South African Centre for Epidemiological Modelling and Analysis (SACEMA) bursary. SLKP and SW were supported in part by the U.S. National Institutes of Health (R01 AI134384 and AI140970, GM144468). DPM is supported by the Wellcome Trust (222574/Z/21/Z).

Additional information

Ethics

The University of Cape Town ethics committee declared that this research did not need ethics approval due to the use of freely accessible nucleotide sequences obtained from the National Centre for Biotechnology Information Taxonomy database (<http://www.ncbi.nlm.nih.gov/taxonomy>) and the Los Alamos National Laboratory HIV sequence database (<https://www.hiv.lanl.gov/content/index>).

Abbreviations

- API: Average pairwise nucleotide sequence identity
- NREV6: 6 Rate Non-Reversible Model
- NREV12: 12 Rate Non-Reversible Model
- GTR: General time Reversible Model
- dsDNA: Double-stranded deoxyribonucleic acid
- dsRNA: Double-stranded ribonucleic acid
- ssDNA: Single-stranded deoxyribonucleic acid
- ssRNA: Single-stranded ribonucleic acid
- DNR: Degrees of Non-Reversibility
- wRF: Weighted Robinson-Foulds distance
- LRT: likelihood ratio test
- AIC: Akaike information criterion test

References

- Anisimova M., Bielawski J. P., Yang Z. (2001) **Accuracy and power of the likelihood ratio test in detecting adaptive molecular evolution** *Molecular biology and evolution* **18**:1585–1592 [Google Scholar](#)
- Baele G., Van de Peer Y., Vansteelandt S. (2010) **Using non-reversible context-dependent evolutionary models to study substitution patterns in primate non-coding sequences** *Journal of molecular evolution* **17**:34–50 [Google Scholar](#)
- Boussau B., Gouy M. (2006) **Efficient likelihood computations with nonreversible models of evolution** *Systematic biology* **55**:756–768 [Google Scholar](#)
- Bruslind L. (n.d.) [pressbooks https://open.oregonstate.edu/generalmicrobiology/chapter/the-viruses/](https://open.oregonstate.edu/generalmicrobiology/chapter/the-viruses/)
- Buckley T. R., Cunningham C. W. (2002) **The effects of nucleotide substitution model assumptions on estimates of nonparametric bootstrap support** *Molecular Biology and Evolution* **19**:394–405 [Google Scholar](#)
- Chelico L., Pham P., Calabrese P., Goodman M. F. (2006) **APOBEC3G DNA deaminase acts processively 3' → 5' on single-stranded DNA** *Nature structural & molecular biology* **13**:392–399 [Google Scholar](#)
- Cheng K. C., Cahill D., Kasai H., Nishimura S., Loeb L. A. (1992) **8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes GT and AC substitutions** *Biological Chemistry* **267**:166–172 [Google Scholar](#)
- Edgar R. C. (2004) **MUSCLE: multiple sequence alignment with high accuracy and high throughput** *Nucleic acids research* **32**:1792–1797 [Google Scholar](#)
- Fernandes J. V., de Medeiros Fernandes T. A. (2012) **Human papillomavirus: biology and pathogenesis. In Human Papillomavirus and Related Diseases-From Bench to Bedside-A Clinical Perspective** *IntechOpen* [Google Scholar](#)
- Fijalkowska I. J., Jonczyk P., Tkaczyk M. M., Bialoskorska M., Schaaper R. M. (1998) **Unequal fidelity of leading and lagging strand DNA replication on the Escherichia coli chromosome** *Proceedings of the National Academy of Science* **95** **17**:10020–10025 [Google Scholar](#)
- Furusawa M. (2012) **Implications of fidelity difference between the leading and the lagging strand of DNA for the acceleration of evolution** *Frontiers in oncology* **2**:144 [Google Scholar](#)
- Grigoriev A. (1999) **Strand-specific compositional asymmetries in double-stranded DNA viruses** *Virus research* **60**:1–19 [Google Scholar](#)
- Hanson L. (2009) **2 Isolation of Viral DNA from Cultures** *Handbook of Nucleic Acid Purification* :23 [Google Scholar](#)

- Harkins G., Delport W., Duffy S., Wood N., Monjane A. L., Owor B. E., Varsani A. (2009) **Experimental evidence indicating that mastreviruses probably did not co-diverge with their hosts** *Virology Journal* **6**:1–14 [Google Scholar](#)
- Hoff M., Orf S., Riehm B., Darriba D., Stamatakis A. (2016) **Does the choice of nucleotide substitution models matter topologically?** *BMC bioinformatics* **17**:1–13 [Google Scholar](#)
- Kuhner M. K., Yamato J. (2015) **Practical performance of tree comparison metrics** *Systematic Biology* **64**:205–214 [Google Scholar](#)
- Kusumoto-Matsuo R., Kanda T., Kukimoto I. (2011) **Rolling circle replication of human papillomavirus type 16 DNA in epithelial cell extracts.** *Genes Cells Genes to Cells* :2333 <https://doi.org/10.1111/j.1365-2443.2010.01458.x> | [PubMed](#) | [Google Scholar](#)
- Larsson A. (2014) **AliView: a fast and lightweight alignment viewer and editor for large datasets** *Bioinformatics* **30**:3276–3278 [Google Scholar](#)
- Lefort V., Longueville J. E., Gascuel O. (2017) **SMS: smart model selection in PhyML** *Molecular biology and evolution* **34**:2422–2424 [Google Scholar](#)
- Lio P., Goldman N. (1998) **Models of molecular evolution and phylogeny** *Genome research* **8**:1233–1244 [Google Scholar](#)
- Minin V., Abdo Z., Joyce P., Sullivan J. (2003) **Performance-based selection of likelihood models for phylogeny estimation** *Systematic biology* **52**:674–683 [Google Scholar](#)
- Nguyen L. T., Schmidt H. A., Von Haeseler A., Minh B. Q. (2015) **IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies** *Molecular biology and evolution* **32**:268–274 [Google Scholar](#)
- Nguyen T., Brunson D., Crespi C., Penman B., Wishnok J., Tannenbaum S. (1992) **DNA damage and mutation in human cells exposed to nitric oxide in vitro** *National Academy of Science* **89**:3030–3034 [Google Scholar](#)
- Onwubiko O., Borst A., Diaz A., Passkowski K., Scheffel F., Tessmer I., Nasheuer H. (2020) **SV40 T antigen interactions with ssDNA and replication protein A: a regulatory role of T antigen monomers in lagging strand DNA replication** *National Library of medicine* **48**:3657–3677 <https://doi.org/10.1093/nar/gkaa138> | [Google Scholar](#)
- Polak P., Arndt P. F. (2008) **Transcription induces strand-specific mutations at the 5' end of human genes** *Genome Research* **18**:1216–1223 [Google Scholar](#)
- Pond S. L., Muse S. V. (2005) **HyPhy: hypothesis testing using phylogenies.** In *Statistical methods in molecular evolution* Springer :125–181 [Google Scholar](#)
- Posada D. (2003) **Using MODELTEST and PAUP to select a model of nucleotide substitution** :6–5
- Posada D., Buckley T. R. (2004) **Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and Bayesian approaches over likelihood ratio tests** *Systematic biology* **53**:793–808 [Google Scholar](#)
- Posada D., Crandall K. A. (2001) **Selecting models of nucleotide substitution: an application to human immunodeficiency virus 1 (HIV-1)** *Molecular biology and evolution* **18**:897–

906 [Google Scholar](#)

Posada D., Crandall K. A. (2001) **Selecting the best-fit model of nucleotide substitution** *Systematic biology* **50**:580–601 [Google Scholar](#)

Ripplinger J., Sullivan J. (2008) **Does choice in model selection affect maximum likelihood analysis?** *Systematic biology* **57**:76–85 [Google Scholar](#)

Ritz C., Spiess A. N. (2008) **qpcR: an R package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis** *Bioinformatics* **24**:1549–1551 [Google Scholar](#)

Robinson D. F., Foulds L. R. (1981) **Comparison of phylogenetic trees** *Mathematical biosciences* **53**:131–147 [Google Scholar](#)

Sanjuan R., Domingo-Calap P. (2016) **Mechanisms of viral mutation** *Cellular and molecular life sciences* **73**:4433–4448 [Google Scholar](#)

Schliep K. P. (2011) **phangorn: phylogenetic analysis in R** *Bioinformatics* **27**:592–593 [Google Scholar](#)

Sharma S., Patnaik S. K., Taggart R. T., Baysal B. E. (2016) **The double-domain cytidine deaminase APOBEC3G is a cellular site-specific RNA editing enzyme** *Scientific reports* **6**:1–12 [Google Scholar](#)

Squartini F., Arndt P. F. (2008) **Quantifying the stationarity and time reversibility of the nucleotide substitution process** *Molecular biology and evolution* **25**:2525–2535 [Google Scholar](#)

Stamatakis A. (2016) **Retrieved from Heidelberg Institute for Theoretical Studies** *The RAxML v8. 2. X Manual* <https://cme.h-its.org/exelixis/resource/download/NewManual.pdf>

Tavare S. (1986) **Some probabilistic and statistical problems in the analysis of DNA sequences** *Lectures on mathematics in the life sciences* **17**:57–86 [Google Scholar](#)

Van Der Walt E., Martin D. P., Varsani A., Polston J. E., Rybicki E. P. (2008) **Experimental observations of rapid Maize streak virus evolution reveal a strand-specific nucleotide substitution bias** *Virology journal* **5**:1–11 [Google Scholar](#)

Wei S., Shi M., Chen X., Sharkey M., van Achterberg C., Ye G., He J. (2010) **New views on strand asymmetry in insect mitochondrial genomes** *PLoS One* **5** [Google Scholar](#)

Wickner R. B. (1993) **Double-stranded RNA virus replication and packaging** *The Journal of biological chemistry* **268**:3797–3800 [Google Scholar](#)

Yap V. B., Speed T. (2005) **Rooting a phylogenetic tree with nonreversible substitution models** *BMC Evolutionary Biology* **5**:1–8 [Google Scholar](#)

Youri P., Newlon C. S., KunkelThomas A. (2002) **Yeast origins establish a strand bias for replicational mutagenesis** *Molecular cell* **10** **1**:207–213 [Google Scholar](#)

Yu Q., König R., Pillai S., Chiles K., Kearney M., Palmer S., Landau N. R. (2004) **Single-strand specificity of APOBEC3G accounts for minus-strand deamination of the HIV genome** *Nature structural & molecular biology* **11**:435–442 [Google Scholar](#)

Author information

Rita Sianga-Mete

Division of Computational Biology, Institute of Infectious Diseases and Molecular Medicine, Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

For correspondence: rita@aims.ac.za

Penelope Hartnady

Division of Computational Biology, Institute of Infectious Diseases and Molecular Medicine, Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

Wimbai Caroline Mandikumba

Division of Computational Biology, Institute of Infectious Diseases and Molecular Medicine, Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

Kayleigh Rutherford

Division of Computational Biology, Institute of Infectious Diseases and Molecular Medicine, Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

Christopher Brian Currin

Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

Florence Phelanyane

Centre for Infectious Disease and Epidemiology Research, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa

Sabina Stefan

Centre for Biomedical Engineering, School of Engineering, Brown University, Providence, United States

Steven Weaver

institute for Genomics and Evolutionary Medicine, Department of Biology, Temple University, Pennsylvania, United States

Sergei L Kosakovsky Pond

institute for Genomics and Evolutionary Medicine, Department of Biology, Temple University, Pennsylvania, United States

Darren Patrick Martin

Division of Computational Biology, Institute of Infectious Diseases and Molecular Medicine, Department of Integrative Biomedical Sciences, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa, Wellcome Center for Infectious Diseases Research in Africa, Institute of Infectious Disease and Molecular Medicine and Department of Medicine, University of Cape Town, Cape Town, South Africa

Editors

Reviewing Editor

Antonis Rokas

Vanderbilt University, Nashville, United States of America

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #1 (Public Review):

The study by Sianga-Mete et al revisits the effects of substitution model selection on phylogenetics by comparing reversible and non-reversible DNA substitution models. This topic is not new, previous works already showed that non-reversible, and also covarion, substitution models can fit the real data better than the reversible substitution models commonly used in phylogenetics. In this regard, the results of the present study are not surprising.

<https://doi.org/10.7554/eLife.87361.2.sa2>

Reviewer #2 (Public Review):

The authors evaluate whether non time reversible models fit better data presenting strand-specific substitution biases than time reversible models. Specifically, the authors consider what they call NREV6 and NREV12 as candidate non time-reversible models. On the one hand, they show that AIC tends to select NREV12 more often than GTR on real virus data sets. On the other hand, they show using simulated data that NREV12 leads to inferred trees that are closer to the true generating tree when the data incorporates a certain degree of non time-reversibility. Based on these two experimental results, the authors conclude that "We show that non-reversible models such as NREV12 should be evaluated during the model selection phase of phylogenetic analyses involving viral genomic sequences". This is a valuable finding, and I agree that this is potentially good practice. However, I miss an experiment that links the two findings to support the conclusion: in particular, an experiment that solves the following question: does the best-fit model also lead to better tree topologies?

[Editors' note: the reviewers were sent the revised submission and rebuttal and based on their response, an amended eLife Assessment has been formulated.]

<https://doi.org/10.7554/eLife.87361.2.sa1>

Author response:

The following is the authors' response to the original reviews

eLife Assessment

This valuable study revisits the effects of substitution model selection on phylogenetics by comparing reversible and non-reversible DNA substitution models. The authors provide evidence that 1) non time-reversible models sometimes perform better than general time-reversible models when inferring phylogenetic trees out of simulated viral genome sequence data sets, and that 2) non time-reversible models can fit the real data better than the reversible substitution models commonly used in phylogenetics, a finding

consistent with previous work. However, the methods are incomplete in supporting the main conclusion of the manuscript, that is that non time-reversible models should be incorporated in the model selection process for these data sets.

The non-reversible models should be incorporated in the selection model process not because they significantly perform better but only because they do not perform worse than the reversible models and that true biochemical processes of nucleotide substitution does support the science of non-reversibility.

Reviewer #1 (Public Review):

The study by Sianga-Mete et al revisits the effects of substitution model selection on phylogenetics by comparing reversible and non-reversible DNA substitution models. This topic is not new, previous works already showed that non-reversible, and also covarion, substitution models can fit the real data better than the reversible substitution models commonly used in phylogenetics. In this regard, the results of the present study are not surprising. Specific comments are shown below.

True

It is well known that non-reversible models can fit the real data better than the commonly used reversible substitution models, see for example,

<https://academic.oup.com/sysbio/article/71/5/1110/6525257>

<https://onlinelibrary.wiley.com/doi/10.1111/jeb.14147?af=R>

The manuscript indicates that the results (better fitting of non-reversible models compared to reversible models) are surprising but I do not think so, I think the results would be surprising if the reversible models provide a better fitting.

I think the introduction of the manuscript should be increased with more information about non-reversible models and the diverse previous studies that already evaluated them. Also I think the manuscript should indicate that the results are not surprising, or more clearly justify why they are surprising.

The surprise in the findings is in NREV12 performing better than NREV6 for double stranded DNA viruses as it was expected that NREV6 would perform better given the biochemical processes discussed in the introduction.

In the introduction and/or discussion I missed a discussion about the recent works on the influence of substitution model selection on phylogenetic tree reconstruction. Some works indicated that substitution model selection is not necessary for phylogenetic tree reconstruction,

<https://academic.oup.com/mbe/article/37/7/2110/5810088>

<https://www.nature.com/articles/s41467-019-08822-w>

<https://academic.oup.com/mbe/article/35/9/2307/5040133>

While others indicated that substitution model selection is recommended for phylogenetic tree reconstruction,

<https://www.sciencedirect.com/science/article/pii/S0378111923001774>

<https://academic.oup.com/sysbio/article/53/2/278/1690801>

<https://academic.oup.com/mbe/article/33/1/255/2579471>

The results of the present study seem to support this second view. I think this study could be improved by providing a discussion about this aspect, including the specific contribution of this study to that.

In our conclusion we have stated that:

The lack of available data regarding the proportions of viral life cycles during which genomes exist in single and double stranded states makes it difficult to rationally predict the situations where the use of models such as GTR, NREV6 and NREV12 might be most justified: particularly in light of the poor over-all performance of NREV6 and GTR relative to NREV12 with respect to describing mutational processes in viral genome sequence datasets. We therefore recommend case-by-case assessments of NREV12 vs NREV6 vs GTR model fit when deciding whether it is appropriate to consider the application of non-reversible models for phylogenetic inference and/or phylogenetic model-based analyses such as those intended to test for evidence of natural selection or the existence of molecular clocks.

The real data was downloaded from Los Alamos HIV database. I am wondering if there were any criterion for selecting the sequences or if just all the sequences of the database for every studied virus category were analysed. Also, was any quality filter applied? How gaps and ambiguous nucleotides were considered? Notice that these aspects could affect the fitting of the models with the data.

We selected varying number of sequences of the database for every studied virus type. Using the software aliview we did quality filter by re-aligning the sequences per virus type.

How the non-reversible model and the data are compared considering the non-reversible substitution process? In particular, given an input MSA, how to know if the nucleotide substitution goes from state x to state y or from state y to state x in the real data if there is not a reference (i.e., wild type) sequence? All the sequences are mutants and one may not have a reference to identify the direction of the mutation, which is required for the non-reversible model. Maybe one could consider that the most abundant state is the wild type state but that may not be the case in reality. I think this is a main problem for the practical application of non-reversible substitution models in phylogenetics.

True

Reviewer #1 (Recommendations for the authors):

The reversible and non-reversible models used in this study assume that all the sites evolve under the same substitution matrix, which can be unrealistic. This aspect could be mentioned.

Done

The manuscript indicates that "a phylogenetic tree was inferred from an alignment of real sequences (Avian Leukosis virus) with an average sequence identity (API) of ~90%.". I was wondering under which substitution model that phylogenetic tree reconstruction was performed? could the use of that model bias posterior results in terms of favoring results based on such a model?

We have stated that the GTR+G model was used to reconstruct the tree. The use of the GTR+G model could yes bias the posterior results as we have stated in the paper too.

I was wondering which specific R function was used to calculate the weighted Robinson-Foulds metric. I think this should be included in the manuscript.

We stated that We used the weighted Robinson-Foulds metric (wRF; implemented in the R phangorn package (Schliep, 2011))

Despite a minority, several datasets fitted better with a reversible model than with a non-reversible model. I think that should be clearly indicated. In addition, in my opinion the AIC does not enough penalizes the number of parameters of the models and favors the non-reversible models over the reversible models, but this is only my opinion based on the definition of AIC and it is not supported. Thus, I think the comparison between phylogenetic trees reconstructed under different substitution models was a good idea (but see also my second major comment).

Noted

When comparing phylogenetic trees I was wondering if one should consider the effect of the estimation method and quality of the studied data? For example, should bootstrap values be estimated for all the ancestral nodes and only ancestral nodes with high support be evaluated in the comparison among trees?

Yes the estimation method and quality of the studied data should be considered. When using RF unlike wRF this will not matter but for weighted RF it does. When building the trees, using RaxML only high support nodes are added to the tree.

In Figure 3, I do not see (by eye) significant differences among the models. I see in the legend that the statistical evaluation was based on a t test but I am not much convinced. Maybe it is only my view. Exactly, which pairs of datasets are evaluated with the t test? Next, I would expect that the influence of the substitution model on the phylogenetic tree reconstruction is higher at large levels of nucleotide diversity because with more substitution events there is more information to see the effects of the model. However, the t test seems to show that differences are only at low levels of nucleotide diversity (and large DNR), what could be the cause of this?

The paired T-tests compares the wRF distances of the inferred tree real tree and the trees simulated using the GTR model verses the wRF distances of the inferred true tree from the trees simulated using the NREV12 model.

The reason why the influence of the NREV12 model on the tree reconstructed is not significantly higher at large levels of nucleotide diversity could be because at a certain level the DNR are simply unrealistic.

Can the user perform substitution model selection (i.e., AIC) among reversible and non-reversible substitution models with IQTREE? If yes, then doing that should be the recommendation from this study, correct?

But, can DNR be estimated from a real dataset? DNR seems to be the key factor (Figure 3) for the phylogenetic analysis under a proper model.

Substitution model selection can be performed among reversible and non-reversible using both HyPhy and IQTREE. And we have recommended that model tests should be done as a first step before tree building. Estimating DNR from real datasets requires a substitution rate matrix of a non-reversible.

The manuscript has many text errors (including typos and incorrect citations). For example, many citations in page 20 show "Error! Reference source not found.". I think authors should double check the manuscript before submitting. Also, some text is not formally written. For example, "G represents gamma-distributed rates", rates of what? The text should be clear for readers that are not familiar with the topic (i.e., G represents gamma-distributed substitution rates among sites). In general, I recommend a detailed revision of the whole text of the manuscript.

Done

Reviewer #2 (Public Review):

The authors evaluate whether non time reversible models fit better data presenting strand-specific substitution biases than time reversible models. Specifically, the authors consider what they call NREV6 and NREV12 as candidate non time-reversible models. On the one hand, they show that AIC tends to select NREV12 more often than GTR on real virus data sets. On the other hand, they show using simulated data that NREV12 leads to inferred trees that are closer to the true generating tree when the data incorporates a certain degree of non time-reversibility.

Based on these two experimental results, the authors conclude that "We show that non-reversible models such as NREV12 should be evaluated during the model selection phase of phylogenetic analyses involving viral genomic sequences". This is a valuable finding, and I agree that this is potentially good practice.

However, I miss an experiment that links the two findings to support the conclusion: in particular, an experiment that solves the following question: does the best-fit model also lead to better tree topologies?

By NREV12 leading to inferred trees that are closer to the true generating tree as compared to GTR, it then shows that the best-fit model in this case being NREV12 leads to better tree topologies.

On simulated data, the significance of the difference between GTR and NREV12 inferences is evaluated using a paired t test. I miss a rationale or a reference to support that a paired t test is suitable to measure the significance of the differences of the wRF distance. Also, the results show that on average NREV12 performs better than GTR, but a pairwise comparison would be more informative: for how many sequence alignments does NREV12 perform better than GTR?

We have used the popular paired t-test as it is the most widely used when comparing means values between two matched samples where the difference of each mean pair is normally distributed. And the wRF distances do match the guidelines above.

The paired t-test contains the pairwise comparison and the boxplots side by side show the pairwise wRF comparisons.

Reviewer #2 (Recommendations for the authors):

The authors reference Baele et al., 2010 for describing NREV6 and NREV12. I suggest using the same name used in the referenced paper: GNR-SYM and GNR respectively. Although I do not think there is a standard name for these models, I would use a previously used one.

We have built studies based on the names NREV6 and NREV12. We would like to keep the naming as standard for our studies.

GTR and NREV12 models are already described in many other papers. I do not see the need to include such an extensive description. Also, a reference should be included to the discrete Gamma rate categories [1]

We included the extensive description to enable other readers who are not super familiar with these models better understanding since we have given the models our own naming different from those used in other papers.

We have added referencing for the discrete gamma rate as recommended. (Yang, 1994)

To evaluate the exhaustiveness and correctness of the results, I would recommend publishing as supplementary material the simulated data sets or the scripts for generating the data set, the scripts or command lines for the analysis, and the versions of the software used (e.g., IQTREE). Also, to strongly support the main conclusion of the manuscript, I suggest adding to the simulations section results the RF-distances of the best-fit selected model under AIC, AICc, and BIC as well.

We can go ahead and submit all the needed datasets. The simulated data RF-Distances results are available and will be submitted. We cannot however add them to the main document as this will create very long data tables.

In some instances, it is mentioned that the selection criterion used is AIC, while in others, AIC-c is referenced. Even in the table captions, both terms are mixed. It should be made clearer which criterion is being employed, as AIC is not suitable for addressing the overparameterization of evolutionary models, given that it does not account for the sample size. A previous pre-print of this article [2] does not mention AIC-c, but also explicitly includes the formulas for AIC that do not take the sample size into account, and reports the same results as this manuscript, what indicates that AIC and not AIC-c was used here. This should be clarified. It is recommended to use AIC-c instead of AIC, especially if the sample size to model parameters ratio is low [3]. Two things may be appointed here: some authors consider tree branch lengths as model free parameters and others do not. In this paper it is not specified how the model parameters are counted. AIC tends to select more parameterized models than AIC-c, and overparameterization can lead to different tree inferences, as evidenced in Hoff et al., 2016. Therefore, it is expected that NREV12 is more frequently selected than NREV6 and GTR.

In my opinion, a pairwise comparison between GTR and NREV12 performance is of great interest here, and the whiskers plots are not useful. Scatterplots would display the results better.

Boxplots are meant to offer a simplified view of the results as the paired t-tests does all of the comparisons. We shall provide the scatter plots as supplementary information so that readers can get full detailed plots as recommended.

Some references are missing.

Missing references added

<https://doi.org/10.7554/eLife.87361.2.sa0>