

Cntnap2 loss drives striatal neuron hyperexcitability and behavioral inflexibility

Reviewed Preprint

v2 • May 2, 2025

Revised by authors

Reviewed Preprint

v1 • July 29, 2024

Katherine R Cording, Emilie M Tu, Hongli Wang, Alexander HCW Agopyan-Miu, Helen S Bateup 

Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, United States • Department of Neuroscience, University of California, Berkeley, Berkeley, United States • Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, United States

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

This **important** and well-executed study describes how deleting the autism spectrum disorder risk gene CNTNAP2 in mice increases dorsolateral striatal projection neuron excitability and promotes repetitive behaviors and cognitive inflexibility. The evidence supporting this claim is **convincing**. The study provides a potential cellular explanation for the repetitive and inflexible behavior in *Cntnap2* knockout mice and CNTNAP2 disorder in humans, which would interest both basic and translational neuroscientists.

<https://doi.org/10.7554/eLife.100162.2.sa4>

Abstract

Autism spectrum disorder (ASD) is a neurodevelopmental disorder characterized by two major diagnostic criteria – persistent deficits in social communication and interaction, and the presence of restricted, repetitive patterns of behavior (RRBs). Evidence from both human and animal model studies of ASD suggest that alteration of striatal circuits, which mediate motor learning, action selection, and habit formation, may contribute to the manifestation of RRBs. *CNTNAP2* is a syndromic ASD risk gene, and loss of function of *Cntnap2* in mice is associated with RRBs. How loss of *Cntnap2* impacts striatal neuron function is largely unknown. In this study, we utilized *Cntnap2*^{-/-} mice to test whether altered striatal neuron activity contributes to aberrant motor behaviors relevant to ASD. We find that *Cntnap2*^{-/-} mice exhibit increased cortical drive of direct pathway striatal projection neurons (dSPNs). This enhanced drive is likely due to increased intrinsic excitability of dSPNs, which make them more responsive to cortical inputs. We find that *Cntnap2*^{-/-} mice exhibit spontaneous repetitive behaviors, increased motor routine learning, perseveration, and cognitive inflexibility. Increased corticostriatal drive of the direct pathway may therefore contribute to the acquisition of repetitive, inflexible behaviors in *Cntnap2* mice.

Introduction

Autism spectrum disorder (ASD) is characterized by alterations in social communication and interaction, as well as the presence of restricted, repetitive, inflexible behaviors (APA, 2022). Given that ASD has high heritability (Sandin et al., 2017), much work has been done in the last 30 years to identify genes that confer risk of developing ASD (De Rubeis et al., 2014; Iossifov et al., 2014; Sanders et al., 2015). Through this, hundreds of high-confidence risk genes have been identified, varying greatly in the proteins for which they code (Satterstrom et al., 2020). These include transcriptional and translational regulators, ion channels, receptors, cell adhesion molecules, and others (De Rubeis et al., 2014; Delorme et al., 2013; Ebert & Greenberg, 2013). Recent work has focused on identifying brain regions and circuits that may be commonly affected by ASD-related mutations. The basal ganglia, in particular the striatum, represents one such commonly altered brain region and prior studies have demonstrated changes in striatal function and striatum-associated behaviors in mice with mutations in ASD risk genes (Benthall et al., 2021; Fuccillo, 2016; Peca et al., 2011; Peixoto et al., 2016; Platt et al., 2017; Rothwell et al., 2014; Wang et al., 2016). However, whether basal ganglia circuits are convergently changed in ASD mouse models is an open question. Here we investigated whether loss of function of the syndromic ASD risk gene *Cntnap2* alters striatal physiology and basal ganglia-dependent behaviors in mice.

Cntnap2 codes for a neurexin-like cell adhesion molecule called Contactin-associated protein-like 2 (Caspr2) (Poliak et al., 1999; Poliak et al., 2003). In mice, Caspr2 is expressed in several cortical and subcortical regions, including the striatum, from embryonic day 14 (E14) onward into adulthood (Penagarikano et al., 2011). Caspr2 is primarily localized at the juxtaparanodes of myelinated axons and is involved in the clustering of potassium channels (Poliak et al., 2003; Scott et al., 2019). In mice, *in vitro* studies suggests that Caspr2 may also play a role in AMPAR trafficking and cell morphology (Anderson et al., 2012; Gdalyahu et al., 2015; Varea et al., 2015), and *ex vivo* experiments indicate that it can control cell excitability and circuit synchronicity (Martin-de-Saavedra et al., 2022). Caspr2 is important during neurodevelopment and has been implicated in neuronal migration (Penagarikano et al., 2011), the maturation and function of parvalbumin-positive GABAergic interneurons (Penagarikano et al., 2011; Scott et al., 2019; Vogt et al., 2018), and the timing of myelination (Scott et al., 2019). *CNTNAP2* mutations in people lead to a neurodevelopmental syndrome that can include language disorders, epilepsy, obsessive compulsive disorder, and ASD (Penagarikano & Geschwind, 2012; Rodenas-Cuadrado et al., 2014). A mouse model of this syndrome, *Cntnap2*^{-/-}, has been shown to exhibit good face validity for ASD-relevant social and motor behavior alterations (Brunner et al., 2015; Dawes et al., 2018; Penagarikano et al., 2011; Scott et al., 2019). However, the impact of *Cntnap2* loss on striatal physiology and corticostriatal-dependent behaviors has not been comprehensively assessed.

The striatum is primarily composed of GABAergic striatal projection neurons (SPNs), which make up two functionally distinct output pathways: the D1-receptor expressing cells of the direct pathway (dSPNs), which project to substantia nigra pars reticulata (SNr), and the D2-receptor expressing cells of the indirect pathway (iSPNs), which project to external globus pallidus (GPe) (Calabresi et al., 2014; Gerfen & Surmeier, 2011; Kravitz et al., 2010; Tai et al., 2012). The two types of SPNs are intermixed throughout the striatum and receive excitatory glutamatergic inputs from cortex and thalamus, as well as dopaminergic input from the midbrain (Ding et al., 2008; Doig et al., 2010; Gerfen & Surmeier, 2011). Coordinated activity between the populations of SPNs in response to these inputs mediates action selection, motor learning and habit formation (Hawes et al., 2015; Santos et al., 2015; Yin et al., 2005, 2006). Although SPNs comprise upwards of 95% of the cells in striatum, there are distinct types of GABAergic interneurons that contribute significantly to the inhibitory circuitry of the striatum. Parvalbumin

(PV) interneurons, which make up ~2% of the cells in striatum, provide the largest feedforward inhibition onto SPNs (Burke et al., 2017 [DOI](#)). Changes in the number and/or function of PV interneurons have been identified in several ASD mouse models, including *Cntnap2*^{-/-} mice, implicating PV circuitry as a potential common alteration across ASD mouse models (Filice et al., 2020 [DOI](#); Juarez & Martinez Cerdeno, 2022 [DOI](#)).


To determine how loss of *Cntnap2* affects striatal function, we assessed the physiology of SPNs and PV-interneurons in the dorsolateral striatum and utilized a range of assays to assess striatum-associated behaviors in *Cntnap2*^{-/-} mice. We find that SPNs of the direct pathway exhibit increased corticostriatal drive, despite unchanged excitatory cortical input. Although decreased inhibitory function has been identified in other brain regions in *Cntnap2*^{-/-} mice, we find no deficit in broad or PV-specific inhibitory input onto either SPN subtype in the case of *Cntnap2* loss. Instead, we identify a significant increase in the intrinsic excitability of dSPNs in *Cntnap2*^{-/-} mice, driven by a reduction in Kv1.2 channel function. Behaviorally, we find that *Cntnap2*^{-/-} mice exhibit RRB-like behaviors including increased grooming, marble burying and nose poking in the holeboard assay. *Cntnap2*^{-/-} mice also exhibit increased motor routine learning in the accelerating rotarod, and cognitive inflexibility in an odor-based reversal learning task. Taken together, these findings suggest that enhanced direct pathway drive may play a role in the spontaneous and learned repetitive behaviors exhibited by these mice.

Results


Cntnap2^{-/-} dSPNs exhibit increased cortical drive

Emerging evidence indicates that corticostriatal synapses are a common site of alteration in mouse models of ASD (Li & Pozzo-Miller, 2020 [DOI](#)). To test whether mice with loss of *Cntnap2* exhibit changes in corticostriatal connectivity, we crossed *Cntnap2*^{-/-} mice to *Thy1*-ChR2-YFP mice, which express channelrhodopsin in a subset of layer V pyramidal neurons (Fig. 1A [DOI](#)) (Arenkiel et al., 2007 [DOI](#); Poliak et al., 1999 [DOI](#)). These mice were crossed to a D1-tdTomato reporter line to visually identify dSPNs (Ade et al., 2011 [DOI](#)). We recorded from SPNs in the dorsolateral striatum (DLS), as this sensorimotor striatal subregion is implicated in the acquisition of habitual and procedural behaviors (Packard & Knowlton, 2002 [DOI](#)). Changes in physiological function in this area may be connected to the acquisition of repetitive motor behaviors in ASD (APA, 2022 [DOI](#); Fuccillo, 2016 [DOI](#)). To simulate a train of cortical inputs, we applied ten pulses of blue light over the recording site in DLS and measured the number of action potentials (APs) fired by SPNs in the absence of synaptic blockers (Fig. 1A [DOI](#)). We altered the light intensity to vary the probability of eliciting subthreshold depolarizations or AP firing. dSPNs were identified using tdTomato fluorescence, and tdTomato negative neurons were designated putative iSPNs.

A *Cntnap2*;D1-tdTomato;Thy1-ChR2



Corticostriatal excitability



Corticostriatal synapses




Figure 1.

***Cntnap2*^{-/-} dSPNs exhibit increased cortical drive.**

(A) Left: schematic of the corticostriatal connectivity experiments. For corticostriatal excitability, cortical terminals expressing ChR2 were stimulated with 10 pulses of blue light at 10 Hz and responses were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. For corticostriatal synaptic strength, cortical terminals expressing ChR2 were stimulated with blue light at increasing intensity and synaptic currents were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. Center: 10x confocal image of the striatum from a *Cntnap2*^{+/+};D1-tdTomato;Thy1-ChR2 mouse. Right: 20x confocal image of dorsolateral striatum from a *Cntnap2*^{+/+};D1-tdTomato;Thy1-ChR2 mouse. YFP (green) labels cell bodies and axons of a subset of layer V pyramidal neurons, tdTomato (red) labels dSPNs, and DAPI stained nuclei are in blue.

(B) Example single traces of action potentials (APs) in dSPNs evoked by cortical terminal stimulation at 20% light intensity for the indicated genotypes.

(C) Quantification (mean ± SEM) of the number of APs evoked in dSPNs at different light intensities. *Cntnap2*^{+/+} n=9 mice, 24 cells, *Cntnap2*^{-/-} n=10 mice, 22 cells. Repeated measures two-way ANOVA p values are shown; g x s F (12, 204) = 1.935, geno F (1, 17) = 2.034, stim F (1.931, 32.83) = 86.12.

(D) Example single traces of action potentials (APs) in iSPNs evoked by cortical terminal stimulation at 20% light intensity for the indicated genotypes.

(E) Quantification (mean ± SEM) of the number of APs evoked in iSPNs at different light intensities. *Cntnap2*^{+/+} n=9 mice, 23 cells, *Cntnap2*^{-/-} n=10 mice, 21 cells. Repeated measures two-way ANOVA p values are shown; g x s F (12, 216) = 0.5012, geno F (1, 18) = 0.9989, stim F (2.331, 41.96) = 60.62.

(F) Average EPSC traces from example dSPNs of each genotype induced by optogenetic cortical terminal stimulation at 14% light intensity.

(G) Quantification of EPSC amplitude evoked in dSPNs at different light intensities (line represents the mean, dots/squares represent average EPSC amplitude for each mouse). *Cntnap2*^{+/+} n=8 mice, 17 cells, *Cntnap2*^{-/-} n=5 mice, 13 cells. Repeated measures two-way ANOVA p values are shown; g x s F (2, 22) = 0.2154, geno F (1, 11) = 0.2585, stim F (1.053, 11.58) = 49.68.

(H) Average EPSC traces from example iSPNs of each genotype induced by optogenetic cortical terminal stimulation at 14% light intensity.

(I) Quantification of EPSC amplitude evoked in iSPNs at different light intensities (line represents mean, dots/squares represent average for mouse). *Cntnap2*^{+/+} n=6 mice, 13 cells, *Cntnap2*^{-/-} n=5 mice, 11 cells. Repeated measures two-way ANOVA p values are shown; g x s F (2, 18) = 0.4428, geno F (1, 9) = 0.5635, stim F (1.095, 9.851) = 23.82.

(J) Quantification (mean ± SEM) of AMPA:NMDA ratio in dSPNs evoked by 20% light intensity (dots/squares represent average AMPA:NMDA ratio for each mouse). *Cntnap2*^{+/+} n=5 mice, 22 cells, *Cntnap2*^{-/-} n=5 mice, 22 cells, p=0.8413, Mann-Whitney test.

(K) Example traces show pairs of EPSCs evoked by optogenetic corticostriatal stimulation (20% light intensity) recorded at +40 mV (top traces) and -70 mV (bottom traces) from *Cntnap2*^{+/+} and *Cntnap2*^{-/-} dSPNs.

(L) Quantification (mean ± SEM) of AMPA:NMDA ratio in iSPNs evoked by 20% light intensity (dots/squares represent average AMPA:NMDA ratio for each mouse). *Cntnap2*^{+/+} n=5 mice, 21 cells, *Cntnap2*^{-/-} n=5 mice, 21 cells, p=0.3095, Mann-Whitney test.

(M) Example traces show pairs of EPSCs evoked by optogenetic corticostriatal stimulation (20% light intensity) recorded at +40 mV (top traces) and -70 mV (bottom traces) from *Cntnap2*^{+/+} and *Cntnap2*^{-/-} iSPNs.

We quantified the number of evoked APs at different light intensities and found that dSPNs in young adult *Cntnap2*^{-/-} mice exhibited increased spike probability compared to wild-type (WT) dSPNs (Fig. 1B-C). The interaction effect of genotype and stimulation intensity in these cells suggests increased corticostriatal drive, consistent with findings in another mouse model with loss of function of the ASD-risk gene *Tsc1* (Benthall et al., 2021). *Cntnap2*^{-/-} iSPNs had subtly increased cortically-evoked AP firing compared to WT iSPNs, although this was not statistically significant (1D-E). To test whether the enhanced spiking probability of *Cntnap2*^{-/-} dSPNs was due to excitatory synaptic changes, we applied blue light of varying intensity over the recording site in DLS while holding cells at -70 mV to evoke AMPAR-driven excitatory postsynaptic currents (EPSCs). We found that the average optically-evoked EPSC amplitude was not significantly different across a range of light intensities in *Cntnap2*^{-/-} dSPNs or iSPNs compared to WT (Fig. 1F-I). In an additional group of mice, we measured the ratio of AMPAR currents recorded at -70 mV to NMDAR currents recorded at +40 mV (at 20% blue light intensity) and found no significant differences in AMPA:NMDA ratio in *Cntnap2*^{-/-} dSPNs or iSPNs (Fig. 1J-M).

To further assess synaptic inputs, we measured the number of dendritic spines in *Cntnap2*^{-/-} and WT SPNs, which are typically the sites of cortical synaptic innervation (Bouyer et al., 1984; Xu et al., 1989). To visualize spines, we injected neonate *Cntnap2*;*Drd1a*-tdTomato mice with AAV5-Syn1-GFP virus to sparsely label dSPNs and iSPNs in the DLS (Fig. S1) (Keaveney et al., 2018). We found that *Cntnap2*^{-/-} SPNs in adult mice had similar spine density as WT (Fig. S1), suggesting no overall change in synapse number. Together, these results show that dSPNs in *Cntnap2*^{-/-} mice exhibit enhanced cortically-driven spiking. However, this is not due to a change in corticostriatal synaptic strength or overall synapse density.

Inhibition is unchanged in *Cntnap2*^{-/-} SPNs

Previous work has indicated a reduction in the number and/or function of fast-spiking parvalbumin-expressing (PV) interneurons across multiple brain regions in *Cntnap2*^{-/-} mice (Ahmed et al., 2023; Antoine et al., 2019; Jurgensen & Castillo, 2015; Paterno et al., 2021; Penagarikano et al., 2011; Vogt et al., 2018). While inhibitory deficits have been identified in the cortex and hippocampus (Antoine et al., 2019; Jurgensen & Castillo, 2015), and the number of PV interneurons has been reported to be decreased in striatum (Penagarikano et al., 2011), a comprehensive assessment of inhibitory synaptic function has yet to be completed in the striatum of *Cntnap2*^{-/-} mice. We first determined whether there were global deficits in inhibition onto SPNs in *Cntnap2*^{-/-} mice using intrastriatal electrical stimulation to evoke inhibitory postsynaptic currents (IPSCs) (Fig. 2A). In *Cntnap2*^{-/-} dSPNs and iSPNs, the average amplitude of electrically-evoked IPSCs across a range of stimulation intensities was not different from WT (Fig. 2B-E).

There are many sources of inhibition in the striatum (Burke et al., 2017), which can all be activated with electrical stimulation. To assess whether inhibition from PV interneurons specifically is altered in *Cntnap2*^{-/-} mice, we crossed *Cntnap2*^{-/-};D1-tdTomato mice to *Pvalb*-Cre; RCL-ChR2-H134R-EYFP (Ai32) mice to express channelrhodopsin in PV interneurons (Fig. 2F) (Hippenmeyer et al., 2005; Madisen et al., 2010). We applied a blue light pulse of varying intensity over the recording site to evoke PV interneuron-specific IPSCs in SPNs, in the presence of excitatory synaptic blockers (Fig. 2F). We found that the average amplitude of optically-evoked IPSCs did not differ significantly in *Cntnap2*^{-/-} dSPNs or iSPNs compared to WT controls (Fig. 2G-J).




Figure 2.

Inhibition is not altered in *Cntnap2*^{-/-} SPNs.

(A) Top: schematic of the experiment. A bipolar stimulating electrode was placed approximately 200 μ m from the recording site. A range of electrical stimulation intensities was applied to the tissue while IPSCs were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. Bottom: 20x confocal image of dorsolateral striatum from a *Cntnap2*^{+/+};D1-tdTomato mouse. tdTomato (red) labels dSPNs, DAPI stained nuclei are in blue.

(B) Average IPSC traces from example dSPNs of each genotype evoked by electrical stimulation at 1.5 (x0.01 mA) intensity for the indicated genotypes.

(C) Quantification (mean $\pm$ SEM) of IPSC amplitude in dSPNs at different stimulation intensities. *Cntnap2*^{+/+} n=17 cells from 9 mice, *Cntnap2*^{-/-} n=16 cells from 9 mice. Repeated measures two-way ANOVA p values are shown; g x s F (7, 217) = 1.080, geno F (1, 31) = 0.0007751, stim F (1.815, 56.28) = 54.92.

(D) Average IPSC traces from example iSPNs of each genotype evoked by electrical stimulation at 1.5 (x0.01 mA) intensity for the indicated genotypes.

(E) Quantification (mean $\pm$ SEM) of IPSC amplitude in iSPNs at different stimulation intensities. *Cntnap2*^{+/+} n=16 cells from 9 mice, *Cntnap2*^{-/-} n=16 cells from 10 mice. Repeated measures two-way ANOVA p values are shown; g x s F (7, 210) = 0.8741, geno F (1, 30) = 1.746, stim F (1.591, 47.73) = 45.66.

(F) Top: schematic of the experiment. PV interneuron terminals expressing Chr2 were stimulated with blue light at a range of intensities and optically-evoked IPSCs were recorded from dSPNs (red) and iSPNs (grey) in dorsolateral striatum. Bottom: 20x confocal image of dorsolateral striatum from a *Cntnap2*^{+/+};D1-tdTomato;PV-Cre;Ai32 mouse. YFP (green) labels PV interneurons, tdTomato (red) labels dSPNs, DAPI stained nuclei are in blue.

(G) Average IPSC traces from example dSPNs of each genotype evoked by optogenetic PV interneuron stimulation at 30% light intensity.

(H) Quantification (mean $\pm$ SEM) of IPSC amplitude in dSPNs at different light intensities. *Cntnap2*^{+/+} n=29 cells from 15 mice, *Cntnap2*^{-/-} n=23 cells from 11 mice. Repeated measures two-way ANOVA p values are shown; g x s F (7, 441) = 0.2566, geno F (1, 63) = 0.001322, stim F (1.433, 90.25) = 32.57.

(I) Average IPSC traces from example iSPNs of each genotype evoked by optogenetic PV interneuron stimulation at 30% light intensity.

(J) Quantification (mean $\pm$ SEM) of IPSC amplitude in iSPNs at different light intensities. *Cntnap2*^{+/+} n=24 cells from 14 mice, *Cntnap2*^{-/-} n=27 cells from 13 mice. Repeated measures two-way ANOVA p values are shown; g x s F (7, 343) = 1.441, geno F (1, 49) = 0.4771, stim F (1.622, 79.46) = 38.49.

To directly measure PV neuron function, we assessed the intrinsic excitability of PV interneurons in *Cntnap2*^{-/-} mice. To visualize PV interneurons for recordings, we crossed *Cntnap2*^{-/-} mice to *Pvalb*-Cre;RCL-tdT (Ai9) mice (Fig. S2A). Plotting the number of APs fired as a function of current step amplitude indicated that there were no significant differences in the intrinsic excitability of PV interneurons in *Cntnap2*^{-/-} mice compared to controls (Fig. S2B,C). There were also no changes in intrinsic cell properties such as rheobase, membrane resistance, capacitance, resting membrane potential, or AP shape in *Cntnap2*^{-/-} PV interneurons (Fig. S2D-K).

Given prior reports of altered PV cell number in *Cntnap2*^{-/-} mice (Paterno et al., 2021 [↗](#); Penagarikano et al., 2011 [↗](#); Vogt et al., 2018 [↗](#)), we counted PV-expressing cells in the striatum, using immunohistochemistry and fluorescent in situ hybridization. We found no significant difference in the number of PV-positive cells in the dorsal striatum of *Cntnap2*^{-/-} mice compared to WT (Fig. S3A-F and K-N). We also found no changes in the average single cell or total level of PV protein expression in the dorsal striatum (Fig. 3G-J [↗](#)). Overall, we did not observe significant changes in PV interneuron number, PV expression, or PV interneuron-mediated inhibition in the adult *Cntnap2*^{-/-} striatum compared to WT controls.

dSPN intrinsic excitability is increased in *Cntnap2*^{-/-} mice

Given that the increased cortical drive onto *Cntnap2*^{-/-} dSPNs could not be explained by changes in excitatory or inhibitory synaptic function, we tested whether it could be due to a change in intrinsic excitability. To measure this, we recorded from dSPNs and iSPNs in *Cntnap2*;D1-tdTomato mice and injected current steps of increasing amplitude. We found that *Cntnap2*^{-/-} dSPNs had significantly increased intrinsic excitability compared to WT dSPNs (Fig. 3A-B [↗](#)). *Cntnap2*^{-/-} dSPNs also had reduced rheobase current (Fig. 3C [↗](#)), the minimum current required to evoke an AP, as well as increased membrane resistance (Fig. 3D [↗](#)). While there was a trend towards increased excitability in *Cntnap2*^{-/-} iSPNs, this effect was not statistically significant (Fig. 3G-H [↗](#)), and these cells did not exhibit changes in rheobase current (Fig. 3I [↗](#)) or membrane resistance (Fig. 3J [↗](#)). Membrane capacitance (Fig. 3E,K [↗](#)), resting membrane potential (Fig. 3F,L [↗](#)), and AP shape (Fig. S4) were not significantly changed in *Cntnap2*^{-/-} SPNs, although latency to first spike was reduced in both *Cntnap2*^{-/-} dSPNs and iSPNs (Fig. S4C, I). Given the lack of synaptic changes observed in *Cntnap2*^{-/-} dSPNs, the increase in dSPN intrinsic excitability likely underlies their enhanced corticostriatal drive (see Fig. 1 [↗](#)).

The effects of Kv1.2 blockade are occluded in *Cntnap2*^{-/-} dSPNs

Caspr2 is known to be involved in the clustering of voltage-gated potassium channels (Inda et al., 2006 [↗](#); Poliak et al., 1999 [↗](#); Poliak et al., 2003 [↗](#)), particularly Kv1.2 channels (Scott et al., 2019 [↗](#)). These channels play an important role in regulating the intrinsic excitability of SPNs (Nisenbaum et al., 1994 [↗](#)). Blockade of these channels with the drug α-Dendrotoxin (α-DTX), results in increased AP frequency, decreased rheobase current, decreased first AP latency, and decreased AP threshold (Shen et al., 2004 [↗](#)), particularly in dSPNs (Lahiri & Bevan, 2020 [↗](#)). Given that *Cntnap2*^{-/-} dSPNs exhibited increased AP frequency (Fig. 3B [↗](#)), decreased rheobase current (Fig. 3C [↗](#)), decreased first AP latency (Fig. S4C), and a trend towards decreased AP threshold (p=0.0516; Fig. S4B), we hypothesized that loss of function of Kv1.2 channels could be the mechanism. To test this, we recorded from dSPNs and iSPNs in *Cntnap2*;Drd1a-tdTomato mice and injected current steps of increasing amplitude in the absence or presence of α-DTX (100nM). We found that α-DTX enhanced AP firing and decreased AP threshold in WT dSPNs (Fig. 4A-C [↗](#)), but not in *Cntnap2*^{-/-} dSPNs (Fig. 4D-F [↗](#)). AP width was also significantly increased by α-DTX in WT but not *Cntnap2*^{-/-} dSPNs (Fig. S5H,Q), while AP fast afterhyperpolarization and AP latency were significantly decreased in both WT and *Cntnap2*^{-/-} dSPNs (Fig. S5E, I, N, R). Together these data show that the effects of α-DTX on *Cntnap2*^{-/-} dSPNs are largely occluded, indicating that Kv1.2 channels are basally altered by loss of *Cntnap2*. This likely accounts for the enhanced intrinsic excitability of *Cntnap2*^{-/-} dSPNs.




Figure 3.

Intrinsic excitability is increased in *Cntnap2*^{-/-} dSPNs.

(A) Example AP traces in dSPNs evoked by a 200 pA current step for the indicated genotypes.

(B) Quantification (mean $\pm$ SEM) of the number of APs evoked in dSPNs at different current step amplitudes. *Cntnap2*^{+/+} n=22 cells from 8 mice, *Cntnap2*^{-/-} n=23 cells from 8 mice. Repeated measures two-way ANOVA p values are shown; $g \times c F(12, 528) = 2.649$, $geno F(1, 44) = 107.5$, $current F(1.974, 86.86) = 147.5$.

(C) Quantification (mean $\pm$ SEM) of the rheobase current in dSPNs. Dots/squares represent the rheobase current for each neuron. n is the same as in panel B. **p=0.0016, two-tailed unpaired t test.

(D-F) Quantification (mean $\pm$ SEM) of dSPN membrane resistance (D), *p=0.0328, Mann-Whitney test; membrane capacitance (E), p=0.2182, Mann-Whitney test; and resting membrane potential (F), p=0.9914, two-tailed unpaired t test. Dots/squares represent the average value for each neuron. n is the same as in panel B.

(G) Example AP traces in iSPNs evoked by a 200 pA current step for the indicated genotypes.

(H) Quantification (mean $\pm$ SEM) of the number of APs evoked in iSPNs at different current step amplitudes. *Cntnap2*^{+/+} n=22 cells from 8 mice, *Cntnap2*^{-/-} n=21 cells from 8 mice. Repeated measures two-way ANOVA p values are shown; $g \times c F(12, 516) = 1.569$, $geno F(1, 43) = 1.042$, $current F(2.041, 87.78) = 284.7$.

(I) Quantification (mean $\pm$ SEM) of the rheobase current in iSPNs. Dots/squares represent the rheobase current for each neuron. n is the same as in panel H. p=0.0923, two-tailed unpaired t test.

(J-L) Quantification (mean $\pm$ SEM) of iSPN membrane resistance (J), p=0.8193, Mann-Whitney test; membrane capacitance (K), p=0.6886, two-tailed unpaired t test; and resting membrane potential (L), p=0.4859, two-tailed unpaired t test. Dots/squares represent the average value for each neuron. n is the same as in panel H.




Figure 4.

The effects of Kv1.2 blockade are occluded in *Cntnap2*^{-/-} dSPNs.

- (A) Example AP traces from *Cntnap2*^{+/+} dSPNs evoked by a 200 pA current step in the absence or presence of α -Dendrotoxin (α -DTX).
- (B) Quantification (mean $\pm$ SEM) of the number of APs evoked in *Cntnap2*^{+/+} dSPNs in the absence or presence of α -DTX at different current step amplitudes. *Cntnap2*^{+/+} n=17 cells from 8 mice, *Cntnap2*^{+/+} + α -DTX n=21 cells from 8 mice. Repeated measures two-way ANOVA p values are shown; drug x c F (13, 468) = 3.097, drug F (1, 36) = 2.399, current F (2.091, 75.28) = 172.0.
- (C) Quantification (mean $\pm$ SEM) of the AP threshold in *Cntnap2*^{+/+} dSPNs in the absence or presence of α -DTX. Dots/triangles represent the threshold for each neuron. n is the same as in panel B. **p=0.0010, Mann-Whitney test.
- (D) Example AP traces from *Cntnap2*^{-/-} dSPNs evoked by a 200 pA current step in the absence or presence of α -DTX.
- (E) Quantification (mean $\pm$ SEM) of the number of APs evoked in *Cntnap2*^{-/-} dSPNs. *Cntnap2*^{-/-} n=21 cells from 9 mice, *Cntnap2*^{-/-} + α -DTX n=20 cells from 9 mice. Repeated measures two-way ANOVA p values are shown; drug x c F (13, 507) = 0.6054, drug F (1, 39) = 0.03352, current F (2.156, 84.07) = 247.8.
- (F) Quantification (mean $\pm$ SEM) of the AP threshold in *Cntnap2*^{-/-} dSPNs. Squares/triangles represent the threshold for each neuron. n is the same as in panel E. p=0.1348, Mann-Whitney test.
- (G) Example AP traces in *Cntnap2*^{+/+} iSPNs evoked by a 200 pA current step in the absence or presence of α -DTX.
- (H) Quantification (mean $\pm$ SEM) of the number of APs evoked in *Cntnap2*^{+/+} iSPNs. *Cntnap2*^{+/+} n=17 cells from 9 mice, *Cntnap2*^{+/+} + α -DTX n=17 cells from 9 mice. Repeated measures two-way ANOVA p values are shown; drug x c F (13, 416) = 0.3719, drug F (1, 32) = 2.161, current F (1.955, 62.57) = 215.9.
- (I) Quantification (mean $\pm$ SEM) of the AP threshold in *Cntnap2*^{+/+} iSPNs. Dots/triangles represent the threshold for each neuron. n is the same as in panel H. p=0.5401, Mann-Whitney test.
- (J) Example AP traces in *Cntnap2*^{-/-} iSPNs evoked by a 200 pA current step in the absence or presence of α -DTX.
- (K) Quantification (mean $\pm$ SEM) of the number of APs evoked in *Cntnap2*^{-/-} iSPNs. *Cntnap2*^{-/-} n=18 cells from 10 mice, *Cntnap2*^{-/-} + α -DTX n=19 cells from 10 mice. Repeated measures two-way ANOVA p values are shown; drug x c F (13, 455) = 2.623, drug F (1, 35) = 0.01734, current F (1.721, 60.23) = 227.4.
- (L) Quantification (mean $\pm$ SEM) of the AP threshold in *Cntnap2*^{-/-} iSPNs. Squares/triangles represent the threshold for each neuron. n is the same as in panel K. p=0.4250, two-tailed unpaired t test.

We tested the effects of α -DTX on WT and *Cntnap2*^{-/-} iSPNs and found minimal changes (Fig. 4G-K, Fig. S6), with no alterations in AP threshold (Fig. 4I,L) and only a small shift in the input-output relationship of *Cntnap2*^{-/-} iSPNs (Fig. 4K). The effects of α -DTX on iSPN excitability have been less well documented and our data suggest a difference in the sensitivity of dSPNs and iSPNs to Kv1 blockade.

Cntnap2^{-/-} mice display increased repetitive behaviors

RRBs comprise one of the primary symptom domains of ASD (APA, 2022). Alterations in striatal circuits are thought to be involved in the manifestation of RRBs, given the striatum's role in action selection and motor control (Estes et al., 2011; Fuccillo, 2016; Hollander et al., 2005; Langen et al., 2014). To determine whether changes in motor behavior accompanied the altered striatal physiology in *Cntnap2*^{-/-} mice, we assessed locomotor activity and spontaneous repetitive behaviors using the open field, marble burying, and holeboard assays (Fig. 5A-D). In the open field, we found no significant difference in the total distance traveled, average speed, or number of rears in *Cntnap2*^{-/-} mice compared to WT controls (Fig. 5E-G). We did find that *Cntnap2*^{-/-} mice made significantly more entries into the center of the open field arena than WT mice, which may reflect a reduction in avoidance behavior in these mice (Fig. 5H). Manually scored grooming behavior revealed that *Cntnap2*^{-/-} mice initiated more grooming bouts than WT controls in the open field (Fig. 5I), consistent with prior reports (Penagarikano et al., 2011).

To further assess motor behaviors in *Cntnap2*^{-/-} mice, we utilized the marble burying assay (Fig. 5C), which takes advantage of a mouse's natural tendency to dig or bury. The number of marbles buried is used as a measure of persistent or repetitive behavior (Angoa-Perez et al., 2013). We found that *Cntnap2*^{-/-} mice buried significantly more marbles on average than WT controls (Fig. 5J). Another measure of repetitive behavior, which is based on the natural exploratory behavior of mice is the holeboard assay (Fig. 5D). In this test, the number of nose pokes made into unbaited holes is recorded. *Cntnap2*^{-/-} mice made significantly more nose pokes within a 10-minute period than WT mice (Fig. 5K). This was largely due to increased poking during the last 5 minutes of the test (Fig. 5L-M), indicating persistent poking behavior in *Cntnap2*^{-/-} mice. Together, the increased grooming, marble burying and nose poking indicate an increase in RRBs in *Cntnap2*^{-/-} mice. A summary of behavior test results by genotype and sex is shown in Table S1.

To gain further insight into the spontaneous behavior profile of *Cntnap2*^{-/-} mice, we utilized a combination of DeepLabCut and Keypoint-MoSeq to perform unbiased, machine learning-based assessment of general locomotion and behavior in an additional cohort of *Cntnap2*^{-/-} mice (Fig. 5N-P, Fig. S7) (Mathis et al., 2018; Wiltchko et al., 2020). Again, we found that *Cntnap2*^{-/-} mice did not exhibit changes in general locomotor activity compared to WT littermates (Fig. S7A). Analysis of movement “syllables” using Keypoint-MoSeq revealed that across the 25 most frequent syllables, two syllables associated with grooming were performed with significantly increased frequency in *Cntnap2*^{-/-} mice (Fig. 5N). *Cntnap2*^{-/-} mice also had an increase in the total number of grooming bouts (Fig. S7B), replicating the findings in the manually scored cohort (see Fig. 5I). While syllable usage was generally similar between WT and *Cntnap2*^{-/-} mice, transitions between syllables differed between the groups (Fig. S7C). A measure of the entropy of syllable transitions revealed that *Cntnap2*^{-/-} mice exhibited less entropy, suggesting less variability in the transition from one movement syllable to the next (Fig. 5O). This rigidity in motor sequence may be indicative of more restricted motor behavior overall. Finally, we tested whether a trained decoder could accurately distinguish WT and *Cntnap2*^{-/-} mice using information about movement, syllable usage, or syllable transitions. The decoding models performed significantly better than chance at identifying WT and *Cntnap2*^{-/-} mice based on their syllable usage and transitions, but not general locomotor activity (Fig. 5P). Together, this analysis demonstrates that while overall locomotor activity is not strongly affected in *Cntnap2*^{-/-} mice, the behavior patterns of these mice are distinct from WT, reflecting enhanced repetitive behaviors.




Figure 5.

Cntnap2^{-/-} mice have increased repetitive behaviors.

(A-D) Schematics of the behavioral assays used to measure repetitive behaviors in *Cntnap2*^{+/+} and *Cntnap2*^{-/-} mice, created with BioRender.com. Male and female mice were used for all tests.

(E-H) Quantification (mean ± SEM) of open field activity over 60 minutes. (E) total distance traveled, $p = 0.3538$, Mann-Whitney test; (F) average speed, $p = 0.3832$, Mann-Whitney test; (G) number of rears, $p = 0.1892$, two-tailed unpaired t test; (H) number of center entries, $*p = 0.0101$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n = 41$ mice, *Cntnap2*^{-/-} $n = 34$ mice.

(I) Quantification (mean ± SEM) of the number of manually scored grooming bouts in the first 20 minutes of the open field test, $**p = 0.0034$, Mann-Whitney test, *Cntnap2*^{+/+} $n = 41$ mice, *Cntnap2*^{-/-} $n = 34$ mice.

(J) Quantification (mean ± SEM) of total marbles buried in the marble burying assay. *Cntnap2*^{+/+} $n = 33$ mice and *Cntnap2*^{-/-} $n = 33$ mice, $*p = 0.0396$, two-tailed unpaired t test.

(K-M) Quantification (mean ± SEM) of performance in the holeboard assay. (K) Total number of nose pokes in 10 minutes, $*p = 0.0212$, Mann-Whitney test; (L) nose pokes in the first five minutes, $p = 0.4811$, two-tailed unpaired t test; and (M) nose pokes in the last five minutes, $*p = 0.0116$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n = 25$ mice, *Cntnap2*^{-/-} $n = 22$ mice.

(N) Quantification (mean ± SEM) of the frequency of movement syllables (top 25 most frequent syllables) in the open field assay defined by Keypoint-MoSeq. *Cntnap2*^{+/+} $n = 13$ mice and *Cntnap2*^{-/-} $n = 11$ mice, $**p = 0.0013$ for grooming 6, $**p = 0.0013$ for grooming 7, Kruskal-Wallis test with Dunn's correction for multiple comparisons.

(O) Quantification (mean ± SEM) of the entropy of syllable transitions in the open field assay. *Cntnap2*^{+/+} $n = 13$ mice and *Cntnap2*^{-/-} $n = 11$ mice, $*p = 0.0236$, two-tailed unpaired t test.

(P) Accuracy of a Random Forest decoder trained on DeepLabCut basic locomotor data (Movement), Keypoint-MoSeq syllable usage data (Syllable usage), or Keypoint-MoSeq syllable transition data (Syllable transition) in distinguishing between *Cntnap2*^{+/+} and *Cntnap2*^{-/-} mice. Dotted line represents chance performance.

For panels E-M and O, dots/squares represent the value for each mouse.

***Cntnap2*^{-/-} mice exhibit enhanced motor learning**

The accelerating rotarod is a striatal-dependent measure of motor coordination and learning that has been used across a range of ASD mouse models (Cording & Bateup, 2023 [↗](#)). Changes in corticostriatal circuits have been identified in mouse models of ASD with altered performance in the task (Cording & Bateup, 2023 [↗](#)). Given the altered corticostriatal drive in *Cntnap2*^{-/-} mice, we tested whether motor coordination and learning were affected in these mice. In the rotarod test, mice learn to walk and then run to stay on a rotating rod as it increases in speed over the course of five minutes. Mice perform three trials a day for four days. In trials one through six, the rod increases in speed from five to 40 revolutions per minute (RPM), while in trials seven through 12 the rod increases from 10 to 80 RPM (Fig. 6A [↗](#)). Learning occurs over trials within a day, as well as across days, as the mouse develops and hones a stereotyped motor pattern to stay on the rod for increasing amounts of time (Rothwell et al., 2014 [↗](#); Yin et al., 2009 [↗](#)). We found that *Cntnap2*^{-/-} mice performed significantly better than WT mice in this task, particularly in the later trials when the rod rotates at the faster 10 to 80 RPM speed (Fig. 6B [↗](#)).

Initial performance (terminal velocity on trial one) was not significantly different between WT and *Cntnap2*^{-/-} mice (Fig. 6C [↗](#)), but the learning rate from trial one to trial 12 was increased in *Cntnap2*^{-/-} mice (Fig. 6D [↗](#)). These findings expand upon previous work indicating increased performance on both steady-state and accelerating rotarod tasks utilizing slower speeds in *Cntnap2*^{-/-} mice (Dawes et al., 2018 [↗](#); Penagarikano et al., 2011 [↗](#)). These results also align with the increased rotarod performance seen in other ASD mouse models exhibiting enhanced corticostriatal drive (Benthall et al., 2021 [↗](#); Cording & Bateup, 2023 [↗](#)).

***Cntnap2*^{-/-} mice exhibit cognitive inflexibility**

RRBs include not just stereotyped movements, but also insistence on sameness and perseverative interests (APA, 2022 [↗](#)). Cognitive inflexibility, a deficit in the ability to flexibly adapt to changes in the environment and update behavior, is a manifestation of ASD and some other psychiatric disorders, which are associated with striatal dysfunction (Fuccillo, 2016 [↗](#)). Indeed, in individuals with ASD, the severity of RRBs is associated with measures of cognitive inflexibility, and evidence from imaging studies suggests that altered corticostriatal connectivity may be present in the case of both repetitive behaviors and cognitive inflexibility (Uddin, 2021 [↗](#)). To assess cognitive flexibility in *Cntnap2*^{-/-} mice, we utilized a four-choice odor-based reversal learning assay (Johnson et al., 2016 [↗](#); Lin et al., 2022 [↗](#)). Briefly, mice were trained to dig for a reward in one of four pots containing scented wood shavings (Fig. 7A [↗](#)). On the first day of the task (acquisition), the rewarded pot was scented with odor one (O1). Mice reached criterion when they chose O1 for at least eight of 10 consecutive trials. On day two, mice were given a recall test in which the rewarded pot was again scented with O1. After reaching criterion, the reversal trials began, and the rewarded pot was scented with the previously unrewarded odor two (O2). To reach criterion, mice must learn the new association of O2 and reward and choose O2 for eight of 10 consecutive trials.

During acquisition, *Cntnap2*^{-/-} mice performed similarly to WT controls, not differing in the average number of trials needed to reach criterion, the number of quadrant entries made before making a choice, or the latency to choose a pot (Fig. 7B-D [↗](#)). On day two, *Cntnap2*^{-/-} mice performed similarly to controls during recall, demonstrating successful consolidation of the odor-reward pairing (Fig. 7E [↗](#)). However, we found that *Cntnap2*^{-/-} mice had a deficit in reversal learning, requiring significantly more trials on average than WTs to reach criterion once the odor-reward pairing was changed (Fig. 7F [↗](#)). Interestingly, during reversal, *Cntnap2*^{-/-} mice made fewer quadrant entries before making a digging choice and had significantly decreased latency to make a choice compared to controls (Fig. 7G,H [↗](#)). Even after the first correct choice of O2 during reversal, *Cntnap2*^{-/-} mice took more trials to reach criterion than WTs (Fig. 7I [↗](#)). In terms of




Figure 6.

Cntnap2^{-/-} mice exhibit enhanced motor learning.

(A) Schematic of the rotarod apparatus (top), and design of the task (bottom), created with *BioRender.com*. Mice walk to stay on the rotating rod for three 5-minute trials a day for two days at 5-40 RPM acceleration over 5 minutes, followed by three trials a day for two days at 10-80 RPM.

(B) Quantification (mean ± SEM) of accelerating rotarod performance across 12 trials for the indicated genotypes. *Cntnap2*^{+/+} n=30 mice, *Cntnap2*^{-/-} n=29 mice. Repeated measures two-way ANOVA p values are shown; g x t F (11, 616) = 4.935, geno F (1, 56) = 15.29, trial F (7.245, 405.7) = 108.4.

(C) Quantification (mean ± SEM) of rotarod performance on trial 1 quantified as terminal speed. Dots/squares represent the performance of individual mice. n is same as in panel B, p=0.1518, Mann-Whitney test.

(D) Quantification (mean ± SEM) of learning rate (RPM/day) calculated as the slope of the line of performance from the first trial (1) to the last trial (12) for each mouse. Dots/squares represent the learning rate for individual mice. n is same as in panel B, ***p=0.0002, two-tailed unpaired t test.




Figure 7.

Cntnap2^{-/-} mice demonstrate cognitive inflexibility.

(A) Schematic of the four-choice odor-based reversal learning task, created with *BioRender.com*. Colored circles represent pots with different scented wood shavings. Yellow ring represents the food reward. Red circle in the Day 2 Reversal panel indicates a novel odor.

(B-D) Quantification of parameters during acquisition learning. Mean $\pm$ SEM number of trials to reach criterion (B, at least 8 out of last 10 trials correct), *p*=0.5397, Mann-Whitney test; number of quadrant entries before making a choice (C), *p*=0.9118, Mann-Whitney test; and latency to make a choice (D), *p*=0.7224, two-tailed unpaired t test. Dots/squares represent the value for each mouse.

(E) Quantification (mean $\pm$ SEM) of the number of trials to reach criterion (at least 8 out of last 10 trials correct) during the recall test on day 2, *p*=0.3737, Mann-Whitney test.

(F-I) Quantification of parameters during reversal learning. Mean $\pm$ SEM number of trials to reach criterion (F, at least 8 out of last 10 trials correct), ***p*=0.0048, two-tailed unpaired t test; number of quadrant entries before making a choice (G), **p*=0.0158, two-tailed unpaired t test; latency to make a choice (H), ***p*=0.0013, two-tailed unpaired t test; and number of trials to reach criterion after the first correct choice (I), **p*=0.0183, two-tailed unpaired t test.

(J) Quantification (mean $\pm$ SEM) of the total number of errors made during reversal learning, ***p*=0.0034, Mann-Whitney test.

(K) Quantification (mean $\pm$ SEM) of the different error types made during reversal learning. Perseverative errors, ****p*=0.0005, regressive errors ***p*=0.0068, novel errors, *p*=0.9955, irrelevant errors *p*=0.9988, omissions, *p*>0.9999, repeated measures two-way ANOVA with Šidák's multiple comparisons test; *g x e F* (4, 72) = 4.292, *geno F* (1, 18) = 10.91, *error F* (4, 72) = 53.49.

(L) Quantification (mean $\pm$ SEM) of the percent of choices made for each odor, binned across three trials, during reversal learning. Odor 1 was rewarded during acquisition learning. Odor 2 was rewarded during reversal learning. Odor 3 was never rewarded (irrelevant). Odor 4 was a novel odor introduced during the reversal learning phase.

For panels B-L, *n*=10 *Cntnap2*^{+/-} mice and 10 *Cntnap2*^{-/-} mice.

errors, *Cntnap2*^{-/-} mice made more reversal errors than WT mice (**Fig. 7J**), in particular perseverative (continuing to choose O1) and regressive (choosing O1 after correctly choosing O2 once) errors (**Fig. 7K**). *Cntnap2*^{-/-} mice did not differ from WT controls in choices of the novel (newly introduced during reversal) or irrelevant (never rewarded) odors, or in the number of omitted trials (timing out without making a choice) (**Fig. 7K**). Instead, the persistence in choosing O1, even after at least one correct choice of O2, drove the cognitive inflexibility in these mice (**Fig. 7L**). This persistence in choice may be reflective of the broader scope of RRBs in *Cntnap2*^{-/-} mice.

Discussion

In this study we tested whether loss of the ASD risk gene *Cntnap2* altered striatal physiology or striatal-dependent behaviors. We found that direct pathway SPNs exhibited enhanced cortical drive in *Cntnap2*^{-/-} mice. This change was not due to differences in excitatory or inhibitory synapses, as cortical inputs onto SPNs were unchanged and there were no significant deficits in inhibition onto SPNs in these mice. Instead, loss of *Cntnap2* resulted in a significant increase in the excitability of dSPNs, likely driven by altered contribution of Kv1.2 channels to intrinsic firing properties. At the behavioral level, *Cntnap2*^{-/-} mice exhibited repetitive behaviors including increased grooming, nose poking, and marble burying. These mice also had enhanced motor learning, performing significantly better than controls in the accelerating rotarod task. Finally, *Cntnap2*^{-/-} mice exhibited cognitive inflexibility in the four-choice reversal learning assay.

Cellular phenotypes of *Cntnap2* loss

The loss of *Caspr2* has a variable impact on intrinsic excitability across brain regions and cell types. The increased intrinsic excitability that we identified in dSPNs has also been observed in Purkinje cells of the cerebellum (Fernandez et al., 2021) and pyramidal cells of the cortex (Antoine et al., 2019; Cifuentes-Diaz et al., 2023). However, we note hypoactivity (Brumback et al., 2018) or unchanged excitability (Lazaro et al., 2019) of pyramidal cells in some cortical regions in *Cntnap2*^{-/-} mice. *Caspr2* is involved in the clustering of voltage-gated potassium channels, in particular at the juxtaparanodes of myelinated axons (Poliak et al., 1999; Poliak et al., 2003) and the axon initial segment (Inda et al., 2006). Indeed, there are profound deficits in the clustering of Kv1-family channels in *Cntnap2*^{-/-} mice, particularly Kv1.2 channels (Scott et al., 2019). These channels play an important role in regulating the intrinsic excitability of SPNs (Nisenbaum et al., 1994), in particular dSPNs (Lahiri & Bevan, 2020), and when blocked, result in increased excitability (Shen et al., 2004). The loss of *Caspr2* in *Cntnap2*^{-/-} mice may result in improper localization of Kv1.2 channels, and thus alter their contribution to intrinsic excitability. Indeed, we found that the effects of Kv1.2 blockade were largely occluded in *Cntnap2*^{-/-} dSPNs, indicating that Kv1.2 loss of function is likely the mechanism driving the change in excitability. Interestingly, we find that blocking Kv1.2 channels has less of an effect on the excitability of iSPNs, which may account for the greater impact of *Cntnap2* loss on dSPNs.

Prior studies of *Cntnap2*^{-/-} mice have identified changes in the number of PV-expressing interneurons in the cortex (Penagarikano et al., 2011; Vogt et al., 2018), hippocampus (Paterno et al., 2021; Penagarikano et al., 2011) and striatum (Penagarikano et al., 2011). However, this finding is inconsistent across studies, as others have reported no change in the number of PV interneurons in these regions (Ahmed et al., 2023; Lauber et al., 2018; Scott et al., 2019). One possible explanation for this disparity is altered PV protein expression in *Cntnap2*^{-/-} mice such that immunoreactivity varies in cell counting assessments. This is supported by the finding that the number of Vicia Villosa Agglutinin-positive (VVA+) perineuronal nets that preferentially surround PV cells is unchanged in *Cntnap2*^{-/-} mice, even when PV immunoreactivity varies (Hartig et al., 1992; Haunso et al., 2000; Lauber et al., 2018). Parvalbumin, a Ca²⁺ buffer, plays an important role in the intrinsic fast-spiking properties of PV

interneurons, such that a reduction in PV protein expression is known to change PV intrinsic function (Orduz et al., 2013). However, altered intrinsic properties of PV interneurons has also been variably reported across brain regions and studies of *Cntnap2*^{-/-} mice, with subtle changes in PV firing properties reported in the developing striatum (Ahmed et al., 2023) and adult cortex (Vogt et al., 2018), but unchanged in the hippocampus (Paterno et al., 2021) and medial prefrontal cortex (Lazaro et al., 2019). In this study, we find no significant change in the number of PV interneurons or the striatal expression of PV protein in *Cntnap2*^{-/-} mice. Consistent with this, we find no deficits in PV-mediated inhibition onto SPNs. Together, this suggests that primary changes in PV interneurons are unlikely to account for altered striatal circuit function in *Cntnap2*^{-/-} mice.

Loss of *Cntnap2* alters striatal-dependent behaviors

The striatum can be separated into functionally distinct subregions. We focused on the dorsal striatum in this study because of its role in controlling motor and cognitive functions (Voorn et al., 2004), which are relevant to ASD (Fuccillo, 2016; Subramanian et al., 2017). The dorsal striatum can be further subdivided into the dorsomedial striatum (DMS) and the dorsolateral striatum (DLS), with the former considered an associative region involved in goal-directed action-outcome learning and the latter implicated in the acquisition of habitual or procedural behaviors (Packard & Knowlton, 2002). We focused on cellular properties in the DLS as stereotyped, perseverative or persistent behaviors likely recruit DLS circuitry (Evans et al., 2024; Fuccillo, 2016). In the accelerating rotarod assay, learning and performance in the task has been associated with changes in the DLS. Positive modulation of the firing rate of DLS neurons occurs during rotarod training, in particular in later trials of the task, and synaptic potentiation of DLS SPNs in late training is necessary for intact performance (Yin et al., 2009). In line with this, lesions of the DLS impair both early and late rotarod learning (Yin et al., 2009). We found that *Cntnap2*^{-/-} mice had increased rotarod performance, most notably at the later stages when DLS function is strongly implicated. Functionally, we also found increased cortical drive of DLS dSPNs in these mice, a change that was sufficient to increase rotarod performance in another mouse model with disruption of an ASD-risk gene (Benthall et al., 2021). Together, this supports a connection between the change observed in DLS SPN physiology and the increased motor routine learning in *Cntnap2*^{-/-} mice, although this idea remains to be causally tested.

In terms of restricted, repetitive behaviors, we replicated prior studies showing increased spontaneous grooming in *Cntnap2*^{-/-} mice (Penagarikano et al., 2011). Early evidence implicates the striatum in the control of the syntax or sequence of movements in a rodent grooming bout, such that very small lesions of DLS are capable of disrupting grooming (Cromwell & Berridge, 1996). However, recent work has also outlined roles for cellular modulation in DMS and ventral striatal Islands of Calleja in the control of grooming behavior (Ramirez-Armenta et al., 2022; Zhang et al., 2021). *Cntnap2*^{-/-} mice also exhibited increased marble burying and nose poking.

The precise neurobiological substrates of these behaviors are yet unclear, but evidence linking increases in these behaviors to changes in cortico-striatal and amygdala-striatal function supports the notion that these behaviors may fit into a broader basal ganglia-associated RRB-like domain (Albelda & Joel, 2012; Lee et al., 2024).

In the four-choice reversal learning task, *Cntnap2*^{-/-} mice showed no differences during the acquisition phase, suggesting that there were no broad deficits in learning. However, in the reversal stage of the task, *Cntnap2*^{-/-} mice took significantly more trials to learn a new odor-reward pairing, owing primarily to continued choice of the previously rewarded odor. The DMS and ventral striatum (nucleus accumbens) have been shown to play an important role in reversal learning (Izquierdo et al., 2017), and in the four choice task specifically (Delevich et al., 2022). Additionally, decreased dopamine release in the DLS is associated with deficits in reversal learning in this task (Kosillo et al., 2019; Lin et al., 2022). Together, the learning phenotypes seen in

Cntnap2^{-/-} mice in the accelerating rotarod and reversal learning assay share an underlying rigidity in behavioral choice. In both cases, changes in striatal circuits likely underlie the repetitive, stereotyped behaviors.

In summary, our results fit into a model whereby divergent cellular changes in the striatum driven by a functionally diverse set of ASD risk genes similarly enhance corticostriatal drive, in particular, of the direct pathway. This in turn may facilitate striatal-dependent motor routine learning and behavioral perseveration. We speculate that a shared gain-of-function in striatal circuits may play a role in the formation of perseverative or repetitive behaviors in a sub-set of ASDs more broadly.


Limitations and future directions

This study characterized multiple striatal cell types and synapses as well as striatum-associated behaviors in *Cntnap2*^{-/-} mice for the first time. However, there remain open questions as to the impact that *Cntnap2* loss has on striatal function. Although we did not identify excitatory or inhibitory synaptic changes in *Cntnap2*^{-/-} SPNs in this study, we only focused on a subset of these connections. While cortical inputs are a major source of excitation onto SPNs, there are other excitatory inputs onto these cells, such as from thalamus, that were not assessed in this study (Ding et al., 2008 [DOI](#); Doig et al., 2010 [DOI](#); Gerfen & Surmeier, 2011 [DOI](#)). In addition, while all intrastriatal GABAergic interneurons would have been sampled in the electrical stimulation experiments in this study, it is possible that interrogation of individual interneuron subtypes would reveal changes in specific inhibitory connections. Finally, major modulators of SPN activity such as cholinergic interneurons and dopaminergic inputs were not assessed here. These inputs have been implicated in several ASD mouse models (Kosillo & Bateup, 2021 [DOI](#); Paval & Miclutia, 2021 [DOI](#); Rapanelli et al., 2017 [DOI](#)), and changes in striatal cholinergic interneuron function have been identified in young (P21) *Cntnap2*^{-/-} mice, making further study of this circuit particularly cogent (Ahmed et al., 2023 [DOI](#)).

The major cellular phenotype we observed was enhanced intrinsic excitability of *Cntnap2*^{-/-} dSPNs. Given that Kv1.2 channels are known to be organized in part by Caspr2 (Poliak et al., 1999 [DOI](#); Poliak et al., 2003 [DOI](#); Scott et al., 2019 [DOI](#)) and the fact that blockade of Kv1.2 did not affect the excitability of *Cntnap2*^{-/-} dSPNs, we conclude that loss of *Cntnap2* leads to the improper clustering, number, or function of Kv1.2. However, more direct measurement of the number and/or localization of these channels through imaging, or the function of these channels through voltage clamp measurement of potassium currents, would bolster this conclusion. Further, assessing whether similar occlusion of the effects of blocking Kv1.2 channels occurs in the cortical and cerebellar cell types that have also been shown to be hyperexcitable in *Cntnap2*^{-/-} mice would support a more holistic understanding of the impact of *Cntnap2* loss on neuronal function (Antoine et al., 2019 [DOI](#); Cifuentes-Diaz et al., 2023 [DOI](#); Fernandez et al., 2021 [DOI](#)). Considering that these cell types also exhibit α-DTX-sensitive currents that boost AP firing, it is possible that the mechanism by which *Cntnap2* loss increases excitability may be shared (Guan et al., 2007 [DOI](#); Haghdoust et al., 2007 [DOI](#); Khavandgar et al., 2005 [DOI](#)).

Finally, while we did identify changes in several striatum-associated behaviors in *Cntnap2*^{-/-} mice, a causative relationship between the physiological and behavioral changes that we identified has not been established. Given that hyperexcitability of the SPNs of the movement-initiating direct pathway is the primary physiological change we identified, testing whether this change is necessary (i.e. by decreasing dSPN activity in *Cntnap2*^{-/-} mice through cell type-specific expression of inward rectifying potassium channel Kir2.1) or sufficient (i.e. by increasing dSPN activity in WT mice using cell type-specific G_q-coupled (hM2Dq) DREADD activation) to alter behavior could illuminate the relationship between striatal function and behavior in *Cntnap2*^{-/-} mice.

Supplementary Figures and Table



Supplementary Figure 1.


***Cntnap2*^{-/-} SPNs do not have altered spine density.**

(A) Representative deconvoluted images of dendritic spines from dSPNs for the indicated genotypes.

(B) Quantification (mean ± SEM) of dendritic spine density per 10 μm of dendrite in *Cntnap2*^{+/+} and *Cntnap2*^{-/-} dSPNs. Dots/squares represent the average spine density per neuron. *Cntnap2*^{+/+} n=9 neurons (15 dendrites) from 6 mice, *Cntnap2*^{-/-} n=8 neurons (15 dendrites) from 6 mice. p=0.9964, two-tailed unpaired t test.

(C) Representative deconvoluted images of dendritic spines from iSPNs for the indicated genotypes.

(D) Quantification (mean ± SEM) of dendritic spine density per 10 μm of dendrite in *Cntnap2*^{+/+} and *Cntnap2*^{-/-} iSPNs. Dots/squares represent the average spine density per neuron. *Cntnap2*^{+/+} n=7 neurons (15 dendrites) from 4 mice, *Cntnap2*^{-/-} n=10 neurons (15 dendrites) from 7 mice. p=0.5362, Mann-Whitney test.



Supplementary Figure 2.

PV interneuron intrinsic excitability is unchanged in *Cntnap2*^{-/-} mice.

(A) Top: schematic of the experiment. tdTomato-expressing PV interneurons were recorded in the dorsolateral striatum and injected with current steps of increasing magnitude to evoke firing. Bottom: 20x confocal image of dorsolateral striatum from a *Cntnap2*^{+/+};PV-Cre;Ai9 mouse. tdTomato (red) labels PV interneurons, DAPI stained nuclei are in blue.

(B) Example traces of APs in PV interneurons evoked by a 200 pA current step for the indicated genotypes.


(C) Quantification (mean $\pm$ SEM) of the number of APs evoked in PV interneurons at different current step amplitudes. *Cntnap2*^{+/+} n=28 cells from 6 mice, *Cntnap2*^{-/-} n=23 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; $g \times c \ F(40, 1960) = 0.1251$, $geno \ F(1, 49) = 0.06951$, $current \ F(2, 033, 99.60) = 107.5$.

(D) Quantification (mean $\pm$ SEM) of the rheobase current. Dots/squares represent the rheobase current for each neuron. n is the same as in panel C. $p=0.8852$, two-tailed unpaired t test.

(E-G) Quantification (mean $\pm$ SEM) of the membrane resistance (E), $p=0.3422$, Mann-Whitney test; membrane capacitance (F), $p=0.9055$, two-tailed unpaired t test; and resting membrane potential (G), $p=0.3141$, two-tailed unpaired t test. Dots/squares represent the average value for each neuron. n is the same as in panel C.

(H) Phase plane plot of a single AP evoked in PV interneurons from *Cntnap2*^{+/+} (black) and *Cntnap2*^{-/-} (red) mice.

(I-K) Quantification (mean $\pm$ SEM) of the AP threshold (I), $p=0.6025$, two-tailed unpaired t test; AP height (J), $p=0.2850$, two-tailed unpaired t test; and AP width at half max (K), $p=0.2561$, two-tailed unpaired t test. Dots/squares represent the value for the first spike evoked at rheobase for each neuron. n is the same as in panel C.




Supplementary Figure 3.

***Cntnap2*^{-/-} mice do not exhibit changes in PV-positive cell number or expression.**

- (A) Representative confocal images of dorsolateral striatum labeled with an antibody against Parvalbumin (PV) protein for the indicated genotypes, mice aged postnatal day (P) 21.
- (B) Quantification (mean ± SEM) of the number of PV positive cells per ROI at P21, $p=0.2554$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n=8$ sections from 4 mice (2 sections imaged per mouse), *Cntnap2*^{-/-} $n=8$ sections from 4 mice (2 sections imaged per mouse).
- (C) Representative confocal images of dorsolateral striatum labeled with an antibody against PV for the indicated genotypes, mice aged P40-50.
- (D) Quantification (mean ± SEM) of the number of PV positive cells per ROI, $p=0.5144$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n=24$ sections from 12 mice (2 sections imaged per mouse), *Cntnap2*^{-/-} $n=24$ sections from 12 mice (2 sections imaged per mouse).
- (E) Representative confocal images of dorsal striatum labeled with an antibody against PV for the indicated genotypes, mice aged P40-50.
- (F) Quantification (mean ± SEM) of the number of PV positive cells in the whole dorsal striatum at P40-50, $p=0.4201$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n=8$ sections from 4 mice (2 sections imaged per mouse), *Cntnap2*^{-/-} $n=8$ sections from 4 mice (2 sections imaged per mouse).
- (G) Quantification (mean ± SEM) of the average single cell fluorescence of PV antibody staining in dorsal striatum, $p=0.9996$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n=8$ sections from 4 mice (2 sections imaged per mouse), *Cntnap2*^{-/-} $n=8$ sections from 4 mice (2 sections imaged per mouse).
- (H) Quantification (mean ± SEM) of the average bulk fluorescence of PV antibody staining in dorsal striatum per section, $p=0.4878$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n=8$ sections from 4 mice (2 sections imaged per mouse), *Cntnap2*^{-/-} $n=8$ sections from 4 mice (2 sections imaged per mouse).
- (I) Representative western blots for CASPR2, PV, and GAPDH protein in dorsal striatal tissue punches. 10 independent samples per genotype are shown.
- (J) Quantification (mean ± SEM) of PV protein expression normalized to GAPDH. Data are presented as a percentage of *Cntnap2*^{+/+} levels, $p=0.1485$, two-tailed unpaired t test. *Cntnap2*^{+/+} $n=10$ mice, *Cntnap2*^{-/-} $n=10$ mice.
- (K) Representative confocal images of in situ hybridization for *Pvalb* in dorsolateral striatum for the indicated genotypes, mice aged P40-50.
- (L-N) Quantification (mean ± SEM) of the number of *Pvalb* positive cells in the dorsolateral striatum per section (L), $p>0.9999$, Mann-Whitney test; dorsomedial striatum (M), $p=0.7429$, Mann-Whitney test; and central striatum (N), $p=0.6286$, Mann-Whitney test. *Cntnap2*^{+/+} $n=4$ sections from 2 mice (2 sections imaged per mouse), *Cntnap2*^{-/-} $n=4$ sections from 2 mice (2 sections imaged per mouse).

Experimental procedures



Supplementary Figure 4.

Latency to spike is reduced in *Cntnap2*^{-/-} SPNs.

(A) Phase plane plot of a single AP evoked in dSPNs from *Cntnap2*^{+/+} (black) and *Cntnap2*^{-/-} (orange) mice.

(B) Quantification (mean ± SEM) of the dSPN AP threshold. Dots/squares represent the threshold for each neuron. *Cntnap2*^{+/+} *n* = 22 cells from 8 mice, *Cntnap2*^{-/-} *n* = 23 cells from 8 mice, *p* = 0.0516, two-tailed unpaired *t* test.

(C) Quantification (mean ± SEM) of dSPN AP latency to first spike plotted by number of APs fired in the current pulse. Shaded area around line represents SEM. Mixed-effects model with Geisser-Greenhouse correction *p* values are shown; *g* x APs *F* (19, 387) = 7.870, *geno* *F* (1, 43) = 24.98, APs *F* (1.740, 35.44) = 913.5.


(D-F) Quantification (mean ± SEM) of the dSPN AP fast afterhyperpolarization (D), *p* = 0.4664, two-tailed unpaired *t* test; AP height (E), *p* = 0.0833, two-tailed unpaired *t* test; and AP width at half max (F), *p* = 0.3792, two-tailed unpaired *t* test. Dots/squares represent the value for the first spike evoked at rheobase for each neuron, *n* same as in panel B.

(G) Phase plane plot of a single AP evoked in iSPNs in *Cntnap2*^{+/+} (green) and *Cntnap2*^{-/-} (blue) mice.

(H) Quantification (mean ± SEM) of the iSPN AP threshold. Dots/squares represent the threshold for each neuron. *Cntnap2*^{+/+} *n* = 22 cells from 8 mice, *Cntnap2*^{-/-} *n* = 21 cells from 8 mice, *p* = 0.1495, two-tailed unpaired *t* test.

(I) Quantification (mean ± SEM) of iSPN AP latency to first spike plotted by number of APs fired in the current pulse. Shaded area around line represents SEM. Mixed-effects model with Geisser-Greenhouse correction *p* values are shown; *g* x APs *F* (19, 343) = 3.300, *geno* *F* (1, 41) = 45.55, APs *F* (1.817, 32.81) = 680.4.

(J-L) Quantification (mean ± SEM) of iSPN AP fast afterhyperpolarization (J), *p* = 0.3281, two-tailed unpaired *t* test; AP height (K), *p* = 0.1794, Mann-Whitney test; and AP width at half max (L), *p* = 0.2409, two-tailed unpaired *t* test. Dots/squares represent the value for the first spike evoked at rheobase for each neuron, *n* same as in panel H.



Supplementary Figure 5.

Inhibition of Kv1.2 differentially impacts action potential properties in *Cntnap2*^{+/+} and *Cntnap2*^{-/-} dSPNs.

(A) Phase plane plot of a single AP evoked in *Cntnap2*^{+/+} dSPNs in the absence (solid black) or presence (dotted black) of α -Dendrotoxin (α -DTX).

(B-C) Quantification (mean $\pm$ SEM) of the *Cntnap2*^{+/+} dSPN membrane resistance (B), $p=0.2681$, Mann-Whitney test; and membrane capacitance (C), $p=0.9886$, two-tailed unpaired t test. Dots/triangles represent the average value for each neuron, *Cntnap2*^{+/+} $n=17$ cells from 8 mice, *Cntnap2*^{+/+} + α -DTX $n=21$ cells from 8 mice.

(D) Quantification (mean $\pm$ SEM) of the rheobase current in *Cntnap2*^{+/+} dSPNs. Dots/triangles represent the rheobase current for each neuron. n is the same as in panels B-C. $p=0.2703$, two-tailed unpaired t test.

(E-H) Quantification (mean $\pm$ SEM) of the *Cntnap2*^{+/+} dSPN fast afterhyperpolarization (E), $*p=0.0274$, two-tailed unpaired t test; resting membrane potential (F), $p=0.1207$, two-tailed unpaired t test; AP height (G), $p=0.3037$, two-tailed unpaired t test; and AP width at half max (H), $**p=0.0084$, two-tailed unpaired t test. Dots/triangles represent the value for the first spike evoked at rheobase for each neuron. n is the same as in panels B-C.

(I) Quantification (mean $\pm$ SEM) of *Cntnap2*^{+/+} dSPN AP latency to first spike, plotted by number of APs fired in the current pulse. Shaded area around line represents SEM. Mixed-effects model with Geisser-Greenhouse correction p values are shown; $d \times$ APs $F(19, 126) = 4.825$, drug $F(1, 36) = 23.33$, APs $F(19.00, 126.0) = 106.7$.


(J) Phase plane plot of a single AP evoked in *Cntnap2*^{-/-} dSPNs in the absence (solid orange) or presence (dotted orange) of α -Dendrotoxin (α -DTX).

(K-L) Quantification (mean $\pm$ SEM) of the *Cntnap2*^{-/-} dSPNs membrane resistance (K), $*p=0.0452$, Mann-Whitney test; and membrane capacitance (L), $p=0.4950$, two-tailed unpaired t test. Squares/triangles represent the average value for each neuron, *Cntnap2*^{-/-} $n=21$ cells from 9 mice, *Cntnap2*^{-/-} + α -DTX $n=20$ cells from 9 mice.

(M) Quantification (mean $\pm$ SEM) of the rheobase current in *Cntnap2*^{-/-} dSPNs. Squares/triangles represent the rheobase current for each neuron. n is the same as in panels K-L. $p=0.6239$, Mann-Whitney test.

(N-Q) Quantification (mean $\pm$ SEM) of the *Cntnap2*^{-/-} dSPN fast afterhyperpolarization (N), $**p=0.0017$, two-tailed unpaired t test; resting membrane potential (O), $p=0.5681$, two-tailed unpaired t test; AP height (P), $p=0.0652$, two-tailed unpaired t test; and AP width at half max (Q), $p=0.3400$, two-tailed unpaired t test. Squares/triangles represent the value for the first spike evoked at rheobase for each neuron. n is the same as in panels K-L.

(R) Quantification (mean $\pm$ SEM) of *Cntnap2*^{-/-} dSPN AP latency to first spike, plotted by number of APs fired in the current pulse. Shaded area around line represents SEM. Mixed-effects model with Geisser-Greenhouse correction p values are shown; $d \times$ APs $F(19, 135) = 3.362$, drug $F(1, 39) = 30.66$, APs $F(19.00, 135.0) = 123.8$.



Supplementary Figure 6.

Inhibition of Kv1.2 does not strongly affect the excitability of iSPNs.

(A) Phase plane plot of a single AP evoked in *Cntnap2*^{+/+} iSPNs in the absence (solid green) or presence (dotted green) of α-Dendrotoxin (α-DTX).

(B-C) Quantification (mean ± SEM) of the *Cntnap2*^{+/+} iSPN membrane resistance (B), p=0.6624, two-tailed unpaired t test; and membrane capacitance (C), p=0.6444, two-tailed unpaired t test. Dots/triangles represent the average value for each neuron, *Cntnap2*^{+/+} n=17 cells from 9 mice, *Cntnap2*^{+/+} + α-DTX n=17 cells from 9 mice.

(D) Quantification (mean ± SEM) of the rheobase current in *Cntnap2*^{+/+} iSPNs. Dots/triangles represent the rheobase current for each neuron. n is the same as in panels B-C. p=0.5267, two-tailed unpaired t test.

(E-H) Quantification (mean ± SEM) of the *Cntnap2*^{+/+} iSPN fast afterhyperpolarization (E), *p=0.0238, Mann-Whitney test; resting membrane potential (F), p=0.6832, Mann-Whitney test; AP height (G), p=0.6448, two-tailed unpaired t test; and AP width at half max (H), p=0.3070, two-tailed unpaired t test. Dots/triangles represent the value for the first spike evoked at rheobase for each neuron. n is the same as in panels B-C.

(I) Quantification (mean ± SEM) of *Cntnap2*^{+/+} iSPN AP latency to first spike, plotted by number of APs fired in the current pulse. Shaded area around line represents SEM. Mixed-effects model with Geisser-Greenhouse correction p values are shown; d x APs F (17, 82) = 2.375, drug F (1, 32) = 6.104, APs F (17.00, 82.00) = 40.22.


(J) Phase plane plot of a single AP evoked in *Cntnap2*^{-/-} iSPNs in the absence (solid blue) or presence (dotted blue) of α-Dendrotoxin (α-DTX).

(K-L) Quantification (mean ± SEM) of the *Cntnap2*^{-/-} iSPNs membrane resistance (K), p=0.1746, two-tailed unpaired t test; and membrane capacitance (L), p=0.7511, two-tailed unpaired t test. Squares/triangles represent the average value for each neuron, *Cntnap2*^{-/-} n=18 cells from 10 mice, *Cntnap2*^{-/-} + α-DTX n=19 cells from 10 mice.

(M) Quantification (mean ± SEM) of the rheobase current in *Cntnap2*^{-/-} iSPNs. Squares/triangles represent the rheobase current for each neuron. n is the same as in panels K-L. p=0.6589, Mann-Whitney test.

(N-Q) Quantification (mean ± SEM) of the *Cntnap2*^{-/-} iSPN fast afterhyperpolarization (N), p=0.0538, two-tailed unpaired t test; resting membrane potential (O), p=0.9461, two-tailed unpaired t test; AP height (P), p=0.5920, two-tailed unpaired t test; and AP width at half max (Q), *p=0.0335, two-tailed unpaired t test, in the absence or presence of α-DTX. Squares/triangles represent the value for the first spike evoked at rheobase for each neuron. n is the same as in panels K-L.

(R) Quantification (mean ± SEM) of *Cntnap2*^{-/-} iSPN AP latency to first spike in the absence or presence of α-DTX, plotted by number of APs fired in the current pulse. Shaded area around line represents SEM. Mixed-effects model with Geisser-Greenhouse correction p values are shown; d x APs F (17, 68) = 3.181, drug F (1, 35) = 20.66, APs F (17.00, 68.00) = 82.73.



Supplementary Figure 7.

DeepLabCut and Keypoint-MoSeq analysis of *Cntnap2*^{-/-} mice.

(A) Quantification (mean ± SEM) of the total distance traveled in the open field measured by DeepLabCut, $p=0.8201$, Mann-Whitney test; *Cntnap2*^{+/+} $n=13$ mice and *Cntnap2*^{-/-} $n=11$ mice.

(B) Quantification (mean ± SEM) of the number of grooming bouts in the open field assessed by Keypoint-MoSeq, calculated as the number of instances of all grooming syllables. *Cntnap2*^{+/+} $n=13$ mice and *Cntnap2*^{-/-} $n=11$ mice, $*p = 0.0352$, Mann-Whitney test.

(C) Map depicting the difference in transitions between syllables (top 25 most frequent syllables) for *Cntnap2*^{+/+} and *Cntnap2*^{-/-} mice (represented as *Cntnap2*^{-/-} minus *Cntnap2*^{+/+}). A red circle around a syllable number indicates increased usage of that syllable in *Cntnap2*^{-/-} mice, a blue circle indicates decreased usage of that syllable in *Cntnap2*^{-/-} mice. A red line between syllable numbers indicates increased usage of that syllable transition in *Cntnap2*^{-/-} mice, a blue line indicates decreased usage of that syllable transition in *Cntnap2*^{-/-} mice. The thickness of the line scales with the size of the difference and the size of the circle surrounding the syllable number scales with usage of the syllable.

Group	N (mice)	Mean	Std. deviation	Std. error of the mean
Open field				
<i>Cntnap2</i> ^{+/+} male distance traveled	22	127.7 m	54.34	11.58
<i>Cntnap2</i> ^{-/-} male distance traveled	16	156 m	28.36	7.091
<i>Cntnap2</i> ^{+/+} female distance traveled	19	139.2 m	48.71	11.17
<i>Cntnap2</i> ^{-/-} female distance traveled	18	131.7 m	57.52	13.56
<i>Cntnap2</i> ^{+/+} male avg. speed	22	0.03536 m/s	0.01500	0.003198
<i>Cntnap2</i> ^{-/-} male avg. speed	16	0.04331 m/s	0.007964	0.001991
<i>Cntnap2</i> ^{+/+} female avg. speed	19	0.03868 m/s	0.01341	0.003076
<i>Cntnap2</i> ^{-/-} female avg. speed	18	0.03644 m/s	0.01590	0.003749
<i>Cntnap2</i> ^{+/+} male rears	22	557.6 rears	201.8	43.03
<i>Cntnap2</i> ^{-/-} male rears	16	462.8 rears	173.0	43.25
<i>Cntnap2</i> ^{+/+} female rears	19	361.2 rears	188.3	43.19
<i>Cntnap2</i> ^{-/-} female rears	18	355.1 rears	158.8	37.42
<i>Cntnap2</i> ^{+/+} male center entries	22	254.3 entries	123.3	26.29
<i>Cntnap2</i> ^{-/-} male center entries	16	350.8 entries	102.4	25.59
<i>Cntnap2</i> ^{+/+} female center entries	19	204.8 entries	71.51	16.41
<i>Cntnap2</i> ^{-/-} female center entries	18	247.6 entries	88.30	20.81
<i>Cntnap2</i> ^{+/+} male number of grooming bouts (20 min.)	22	14.91 bouts	8.602	1.834
<i>Cntnap2</i> ^{-/-} male number of grooming bouts (20 min.)	16	19.56 bouts	11.52	2.879
<i>Cntnap2</i> ^{+/+} female number of grooming bouts (20 min.)	19	13.37 bouts	6.353	1.457
<i>Cntnap2</i> ^{-/-} female number of grooming bouts (20 min.)	18	28.44 bouts	18.99	4.475

Supplementary Table 1.

Summary of behavior data by sex and genotype

Marble burying				
<i>Cntnap2^{+/+}</i> male marbles buried	16	8.125 marbles	6.206	1.552
<i>Cntnap2^{-/-}</i> male marbles buried	17	10.82 marbles	5.187	1.258
<i>Cntnap2^{+/+}</i> female marbles buried	17	8.235 marbles	5.641	1.368
<i>Cntnap2^{-/-}</i> female marbles buried	16	11.25 marbles	5.310	1.328
Holeboard				
<i>Cntnap2^{+/+}</i> male 10 min. nose pokes	11	156.5 pokes	21.18	6.385
<i>Cntnap2^{-/-}</i> male 10 min. nose pokes	11	192.3 pokes	38.79	11.70
<i>Cntnap2^{+/+}</i> female 10 min. nose pokes	14	175.9 pokes	37.51	10.03
<i>Cntnap2^{-/-}</i> female 10 min. nose pokes	11	180.5 pokes	47.50	14.32
<i>Cntnap2^{+/+}</i> male first 5 min. nose pokes	11	105.4 pokes	16.26	4.901
<i>Cntnap2^{-/-}</i> male first 5 min. nose pokes	11	114.5 pokes	30.25	9.121
<i>Cntnap2^{+/+}</i> female first 5 min. nose pokes	14	110.5 pokes	22.93	6.128
<i>Cntnap2^{-/-}</i> female first 5 min. nose pokes	11	112.3 pokes	30.09	9.073
<i>Cntnap2^{+/+}</i> male last 5 min. nose pokes	11	51.18 pokes	10.09	3.042
<i>Cntnap2^{-/-}</i> male last 5 min. nose pokes	11	77.73 pokes	14.80	4.462
<i>Cntnap2^{+/+}</i> female last 5 min. nose pokes	14	65.36 pokes	20.82	5.565
<i>Cntnap2^{-/-}</i> female last 5 min. nose pokes	11	68.27 pokes	20.15	6.077
Rotarod				
<i>Cntnap2^{+/+}</i> male learning rate	15	1.839 RPM/day	1.416	0.3655
<i>Cntnap2^{-/-}</i> male learning rate	14	2.856 RPM/day	1.166	0.3115
<i>Cntnap2^{+/+}</i> female learning rate	15	2.162 RPM/day	1.083	0.2798
<i>Cntnap2^{-/-}</i> female learning rate	15	3.607 RPM/day	1.103	0.2849
Four choice reversal learning				

Supplementary Table 1. (continued)

<i>Cntnap2</i> ^{+/+} male acquisition trials	5	20.80 trials	6.834	3.056
<i>Cntnap2</i> ^{-/-} male acquisition trials	5	20 trials	7.106	3.178
<i>Cntnap2</i> ^{+/+} female acquisition trials	5	17.20 trials	4.550	2.035
<i>Cntnap2</i> ^{-/-} female acquisition trials	5	19.60 trials	2.302	1.030
<i>Cntnap2</i> ^{+/+} male recall trials	5	11 trials	1.871	0.8367
<i>Cntnap2</i> ^{-/-} male recall trials	5	11 trials	4.243	1.897
<i>Cntnap2</i> ^{+/+} female recall trials	5	10.60 trials	1.140	0.5099
<i>Cntnap2</i> ^{-/-} female recall trials	5	10.20 trials	1.095	0.4899
<i>Cntnap2</i> ^{+/+} male reversal trials	5	18.40 trials	1.817	0.8124
<i>Cntnap2</i> ^{-/-} male reversal trials	5	30.40 trials	4.037	1.806
<i>Cntnap2</i> ^{+/+} female reversal trials	5	26.40 trials	9.737	4.354
<i>Cntnap2</i> ^{-/-} female reversal trials	5	32.60 trials	4.827	2.159
<i>Cntnap2</i> ^{+/+} male reversal errors	5	7.400 errors	0.8944	0.4000
<i>Cntnap2</i> ^{-/-} male reversal errors	5	17.60 errors	2.608	1.166
<i>Cntnap2</i> ^{+/+} female reversal errors	5	12.20 errors	6.380	2.853
<i>Cntnap2</i> ^{-/-} female reversal errors	5	15.20 errors	1.789	0.8000
DeepLabCut Keypoint-MoSeq Open Field				
<i>Cntnap2</i> ^{+/+} male distance traveled	7	272 m	6112	2310
<i>Cntnap2</i> ^{-/-} male distance traveled	5	232 m	4876	2181
<i>Cntnap2</i> ^{+/+} female distance traveled	6	202 m	1604	654.9
<i>Cntnap2</i> ^{-/-} female distance traveled	6	227 m	4651	1899
<i>Cntnap2</i> ^{+/+} male number of grooming bouts (60 min.)	7	324.7 bouts	166.1	62.77

Supplementary Table 1. (continued)

<i>Cntnap2</i> ^{-/-} male number of grooming bouts (60 min.)	5	411.6 bouts	131.6	58.84
<i>Cntnap2</i> ^{+/+} female number of grooming bouts (60 min.)	6	361.0 bouts	92.18	37.63
<i>Cntnap2</i> ^{-/-} female number of grooming bouts (60 min.)	6	534.0 bouts	206.9	84.47

Supplementary Table 1. (continued)

Mice

All animal procedures were conducted in accordance with protocols approved by the University of California, Berkeley Institutional Animal Care and Use Committee (IACUC) and Office of Laboratory Animal Care (OLAC) (AUP-2016-04-8684-3). *Cntnap2*^{-/-} mice and littermate *Cntnap2*^{+/+} controls with the following alleles were utilized for each experiment.

Mice were group housed on a 12 h light/dark cycle (dark cycle 9:00 AM – 9:00 PM) and given ad libitum access to standard rodent chow and water. Both male and female animals were used for experimentation. The ages, sexes, and numbers of mice used for each experiment are indicated in the respective method details and figure legends. All mice used for experiments were heterozygous or hemizygous for the *Drd1a*-tdTomato, *Thy1*-Chr2-YFP, PV-Cre, Ai32, or Ai9 transgenes to avoid potential physiological or behavioral alterations.

Electrophysiology

Mice (P50-60) were briefly anesthetized with isoflurane and perfused transcardially with ice-cold ACSF (pH = 7.4) containing (in mM): 127 NaCl, 25 NaHCO₃, 1.25 NaH₂PO₄, 2.5 KCl, 1 MgCl₂, 2 CaCl₂, and 25 glucose, bubbled continuously with carbogen (95% O₂ and 5% CO₂). Brains were rapidly removed and coronal slices (275 μm) were cut on a VT1000S vibratome (Leica) in oxygenated ice-cold choline-based external solution (pH = 7.8) containing (in mM): 110 choline chloride, 25 NaHCO₃, 1.25 NaHPO₄, 2.5 KCl, 7 MgCl₂, 0.5 CaCl₂, 25 glucose, 11.6 sodium ascorbate, and 3.1 sodium pyruvate. Slices were recovered in ACSF at 36°C for 15 min and then kept at room temperature (RT) before recording. Recordings were made with a MultiClamp 700B amplifier (Molecular Devices) at RT using 3-5 MΩ glass patch electrodes (Sutter, #BF150-86-7.5). Data were acquired using ScanImage software, written and maintained by Dr. Bernardo Sabatini (<https://github.com/bernardosabatini/SabalabAcq>). Traces were analyzed in Igor Pro (Wavemetrics). Recordings with a series resistance > 25 MΩ or holding current above 200 pA were rejected. Passive properties were calculated using the double exponential curve fit of the average of five -5 mV, 100 ms long pulse steps applied at the beginning of every experiment.

Current-clamp recordings

Current clamp recordings were made using a potassium-based internal solution (pH = 7.4) containing (in mM): 135 KMeSO₄, 5 KCl, 5 HEPES, 4 Mg-ATP, 0.3 Na-GTP, 10 phosphocreatine, and 1 EGTA. For corticostriatal excitability experiments, optogenetic stimulation consisted of a full-field pulse of blue light (470 nm, 0.5 ms pulse width, CoolLED) through a 63x objective (Olympus, LUMPLFLN60XW). Light power was linear over the range of intensities tested. No synaptic blockers were included. For intrinsic excitability experiments (SPN, PV interneuron, and SPN + α-DTX experiments), NBQX (10 μM, Tocris, #1044), CPP (10 μM, Tocris, #0247) and picrotoxin (50 μM, Abcam, #120315) were added to the external solution to block synaptic transmission. For Kv1.2 inhibition experiments, α-DTX (100nM, Alomone Labs, #D-350) was added to the external solution. Control recordings in the absence of α-DTX were performed on slices prior to drug application or on fresh slices after drug washout in alternating order across recording days. Bovine serum albumin (BSA, 0.005%, Sigma, #A7030) was included in both control and α-DTX-containing external solutions to minimize nonspecific binding. One second depolarizing current steps were applied to induce action potentials. No holding current was applied to the membrane.

Voltage-clamp recordings

Voltage-clamp recordings were made using a cesium-based internal solution (pH = 7.4) containing (in mM): 120 CsMeSO₄, 15 CsCl, 10 TEA-Cl, 8 NaCl, 10 HEPES, 1 EGTA, 5 QX-314, 4 Mg-ATP, and 0.3 Na-GTP. Recordings were acquired with the amplifier Bessel filter set at 3 kHz. Corticostriatal synaptic stimulation experiments to measure evoked EPSCs were performed in picrotoxin (50 μM), and optogenetic stimulation consisted of a full-field pulse of blue light (470 nm, 0.15 ms pulse

Experiment	Mouse line	Allele 1 (reference; JAX strain #)	Allele 2 (reference; JAX strain #)	Allele 3 (reference; JAX strain #)	Allele 4 (reference; JAX strain #)
Corticostriatal transmission (Fig. 1)	<i>Cntnap2</i> ;D1-tdTomato;Thy1-ChR2	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Drd1a</i> -tdTomato (Ade et al., 2011); #016204)	<i>Thy1</i> -ChR2-YFP (Arenkiel et al., 2007); #007612)	
Broad inhibition (Fig. 2A-E)	<i>Cntnap2</i> ;D1-tdTomato	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Drd1a</i> -tdTomato (Ade et al., 2011); #016204)		
PV-specific inhibition (Fig. 2F-J)	<i>Cntnap2</i> ;D1-tdTomato;PV-Cre;Ai32	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Drd1a</i> -tdTomato (Ade et al., 2011); #016204)	<i>Pvalb</i> -Cre (Hippenmeyer et al., 2005); #017320)	Ai32 (Madisen et al., 2012); #012569)
SPN intrinsic excitability (Fig. 3, S4)	<i>Cntnap2</i> ;D1-tdTomato	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Drd1a</i> -tdTomato (Ade et al., 2011); #016204)		
SPN intrinsic excitability in the presence of α -DTX (Fig. 4, S5, S6)	<i>Cntnap2</i> ;D1-tdTomato	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Drd1a</i> -tdTomato (Ade et al., 2011); #016204)		
Behavior experiments* (Fig. 5-7, S7)	<i>Cntnap2</i> ;D1-tdTomato	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Drd1a</i> -tdTomato (Ade et al., 2011); #016204)		
Spine analysis, PV cell counting	<i>Cntnap2</i> ;D1-tdTomato	<i>Cntnap2</i>	<i>Drd1a</i> -tdTomato		

Table 1.

Summary of mouse lines used.

(Fig. S1, S3A-H)		(Poliak et al., 2003); #017482)	(Ade et al., 2011); #016204)		
PV intrinsic excitability (Fig. S2)	<i>Cntnap2</i> ;PV-Cre;Ai9	<i>Cntnap2</i> (Poliak et al., 2003); #017482)	<i>Pvalb</i> -Cre (Hippenmeyer et al., 2005); #017320)	Ai9 (Madisen et al., 2010); #007909)	
PV in situ, western blot (Fig. S3I-N)	<i>Cntnap2</i>	<i>Cntnap2</i> (Poliak et al., 2003); #017482)			

*Littermate animals both positive and negative for D1-tdTomato were used in behavior experiments

Table 1. (continued)

width) through a 63x objective. To record AMPAR-mediated EPSCs, cells were held at -70mV ; to record NMDAR-mediated EPSCs, cells were held at $+40\text{mV}$. Synaptic stimulation experiments to measure evoked IPSCs were performed in NBQX ($10\text{ }\mu\text{M}$) and CPP ($10\text{ }\mu\text{M}$). For electrically evoked IPSCs, a concentric bipolar stimulating electrode (FHC, #30214) was placed in dorsal striatum, roughly $200\text{ }\mu\text{m}$ medial to the recording site in dorsolateral striatum, and a 0.15 ms stimulus was applied. For PV-interneuron optically evoked IPSCs, a full-field pulse of blue light (470 nm , 0.15 ms pulse width) was applied through a 63x objective at the recording site. All evoked IPSC experiments were recorded with cells held at -70mV .

Dendritic imaging and spine analysis

Neonatal (P1-3) *Cntnap2*^{-/-};D1-tdT and *Cntnap2*^{+/-};D1-tdT mice were cryoanesthetized and injected bilaterally with 200 nL AAV1.hSyn.eGFP.WPRE.bGH (Penn Vector Core, #p1696 (Keaveney et al., 2018)), diluted 1:75 in saline to achieve sparse transduction. Injections were targeted to the dorsal striatum, with coordinates approximately 1.3 mm lateral to midline, 2.0 mm posterior to bregma, and 1.5 mm ventral to the head surface. At P50-60, mice were anesthetized with isoflurane and transcardial perfusion was performed with 10 mL of 1x PBS followed by 10 mL of ice cold 4% PFA (EMS, #15710-S) in 1x PBS. Brains were post-fixed in 4% PFA in 1x PBS overnight at 4°C . $80\text{ }\mu\text{m}$ coronal sections were made using a freezing microtome (American Optical, AO 860) and stored in 1x PBS at 4°C . Sections were blocked for 1 hour at RT in BlockAid (ThermoFisher, #B10710) and incubated for 48 hours with gentle shaking at 4°C with antibodies against GFP (1:2500, Abcam, #13970) and RFP (1:1000, Rockland (VWR, #600-401-379) diluted in PBS-Tx (1x PBS with 0.25% Triton X-100 (Sigma, #T8787). Sections were washed $3 \times 10\text{ min}$ in PBS-Tx and incubated with gentle shaking for 1 hour at RT with Alexa Fluor 488 and 546 secondary antibodies (1:500, Invitrogen, #A11039, #A11035). Sections were washed $3 \times 10\text{ min}$ in 1x PBS and mounted onto SuperFrost slides (VWR, #48311-703) using VECTASHIELD HardSet Antifade Mounting Medium (Vector Laboratories, #H-1400-10). Z stack images of individual dendrites were taken on a confocal microscope (Olympus FLUOVIEW FV3000) with a 60x oil immersion objective (Olympus #1-U2B832) at 2.5x zoom with a step size of $0.4\text{ }\mu\text{m}$ and deconvoluted using Olympus CellSens software. To quantify spine density, dendrites and spines were reconstructed using the FilamentTracer module in Imaris software (Oxford Instruments). The spine density of each dendrite was calculated using Imaris. Dendrites analyzed varied in total length, but excluded the most proximal and distal portions of the dendrite.

Brain sectioning and immunohistochemistry

Adult mice were perfused as above and brains were post-fixed with 4% paraformaldehyde overnight, then sectioned coronally at $30\text{ }\mu\text{m}$. For immunohistochemistry, individual wells of sections were washed for $3 \times 5\text{ min}$ with 1x PBS, then blocked for 1 hour at RT with BlockAid blocking solution. Primary antibodies diluted in PBS-Tx were added and tissue was incubated for 48 hours with gentle shaking at 4°C . Sections were then washed $3 \times 10\text{ min}$ with PBS-Tx. Secondary antibodies diluted 1:500 in PBS-Tx were added and incubated with gentle shaking for 1 hour at RT. Sections were washed $3 \times 10\text{ min}$ in 1x PBS. Sections were mounted onto SuperFrost slides (VWR, #48311-703) and coverslipped with VECTASHIELD HardSet with DAPI (Vector Laboratories, #H-1500-10) or VECTASHIELD HardSet Antifade Mounting Medium (Vector Laboratories, #H-1400-10). The following antibodies were used: mouse anti-PV (1:1000, Sigma, #P3088), rabbit anti-PV (1:1000, Abcam, #11427), anti-RFP (1:500, Rockland, #600-401-379), Alexa Fluor 405, 488 and 546 conjugated secondary antibodies (1:500, Invitrogen, #A-31553, #A-11001, #A-11003, and #A-11035).

PV cell counting

To count PV+ interneurons, Z-stack images of immunostained striatal sections were taken on a confocal microscope (Olympus FLUOVIEW FV3000) with a 10x or 20x objective (Olympus #1-U2B824 or Olympus #1-U2B825) and step size of $2\text{ }\mu\text{m}$. For quantification, image stacks were Z-

projected to maximum intensity using Fiji (ImageJ) and cropped to a 400 μm x 400 μm image in anatomically matched sections of the DLS. All PV-expressing cells within this region were counted using the ROI manager tool in ImageJ. Designation of a cell as PV positive was determined by the experimenter and consistently maintained across animals. Experimenter was blind to genotype and ROIs were made on the DAPI channel to avoid selecting regions based on PV expression. To quantify bulk PV fluorescence, ROIs were manually defined in ImageJ using the Freehand tool to cover as much of the DLS as possible, and mean fluorescence intensity was measured. To quantify individual cell PV fluorescence, ROIs were manually defined around every PV positive cell in the previously drawn DLS ROI using the Freehand tool, and mean fluorescence intensity was measured.

Western Blot

Adult mice (P48-55) were deeply anesthetized with isoflurane and decapitated. Brains were rapidly dissected and 1.5 mm dorsal striatum punches (Biopunch, Ted Pella, #15111-15) were collected from both hemispheres, flash-frozen in liquid nitrogen, and stored at -80°C . On the day of analysis, frozen samples were sonicated (QSonica Q55) until homogenized in 200 μl lysis buffer containing 1% SDS in 1x PBS with Halt phosphatase inhibitor cocktail (Thermo Fisher Scientific, #PI78420) and Complete mini EDTA-free protease inhibitor cocktail (Roche, #4693159001). Sample homogenates were then boiled on a heat block at 95°C for 5 min and allowed to cool to RT. Total protein content was determined using a BCA assay (Thermo Fisher Scientific, #23227). Following the BCA assay, protein homogenates were mixed with 4x Laemmli sample buffer (BioRad, #161-0747). 12.5 μg of total protein per sample were then loaded onto 12% Criterion TGX gels (BioRad, #5671044) and run at 65 V. Proteins were transferred to a PVDF membrane (BioRad, #1620177) at 11 V for 14 hours at 4°C using the BioRad Criterion Blotter (BioRad, #1704070). Membranes (BioRad, #1620177) were briefly reactivated in methanol and rinsed in water 3x. After rinsing, membranes were blocked in 5% milk in 1x TBS with 1% Tween (TBS-Tween) for 1 hour at RT before being incubated with primary antibodies diluted in 5% milk in TBS-Tween overnight at 4°C . The following day, after 3 x 10 min washes with TBS-Tween, membranes were incubated with secondary antibodies for 1 hour at RT. Following 6 x 10 min washes, membranes were incubated with chemiluminescence substrate (PerkinElmer #NEL105001EA) for 1 min and exposed to Amersham Hyperfilm ECL (VWR, #95017-661).

Bands were quantified by densitometry using ImageJ software. GAPDH was used to normalize protein content and data are expressed as a percentage of control within a given experiment. The following antibodies were used: anti-Caspr2 (1:5000, Abcam, #153856), anti-PV (1:2500, Abcam, #11427), anti-GAPDH (1:5000, Cell Signaling, #51745S), and anti-rabbit goat HRP conjugate (1:5000, BioRad, #1705046).

In situ hybridization

Fluorescent in situ hybridization was performed to quantify *Pvalb* mRNA expression in the striatum of *Cntnap2*^{+/+} and *Cntnap2*^{-/-} mice. Mice were briefly anesthetized with isoflurane and brains were harvested, flash-frozen in OCT mounting medium (Thermo Fisher Scientific, #23-730-571) on dry ice and stored at -80°C for up to 6 months. 16 μm sections were collected using a cryostat (Thermo Fisher Scientific, Microm HM 550), mounted directly onto Superfrost Plus glass slides (VWR, #48311-703) and stored at -80°C for up to 6 months. In situ hybridization was performed according to the protocols provided with the RNAscope Multiplex Fluorescent Reagent Kit (ACD, #323100). *Drd1a* mRNA was visualized with a probe in channel 2 (ACD, #406491-C2) and *Pvalb* mRNA in channel 3 (ACD, #421931-C3). After incubation, sections were secured on slides using ProLong Gold Antifade Mountant with DAPI (Invitrogen, P36935) and 60 x 24 mm rectangular glass coverslips (VWR, #16004-096). Sections were imaged on an Olympus FluoView 3000 confocal microscope using a 10x objective with 1.5x zoom and a step size of 2 μm . *Pvalb*-

expressing cells were quantified across the entire striatum using the ROI manager tool in ImageJ. A cell was considered *Pvalb* positive if over 50% of the cell contained fluorescent puncta when compared to the DAPI channel. Experimenter was blind to genotype.

Behavioral analysis

All behavior studies were carried out in the dark phase of the light cycle under red lights (open field) or white lights (marble burying, holeboard, rotarod, and four choice reversal learning). Mice were habituated to the behavior testing room for at least 30 min prior to testing. Mice were given at least one day between different tests. All behavior equipment was cleaned between each trial and mouse with 70% ethanol and rinsed in diluted soap followed by water at the end of the day. If male and female mice were to be tested on the same day, male mice were run first then returned to the housing room, after which all equipment was thoroughly cleaned prior to bringing in female mice for habituation. Behavioral tests were performed with young adult male and female mice (7-11 weeks old). The experimenter was blind to genotype throughout the testing and scoring procedures.

Open field assay

Exploratory behavior in a novel environment and general locomotor activity were assessed by a 60 min session in an open field chamber (40 cm L x 40 cm W x 34 cm H) made of transparent plexiglass. Horizontal infrared photobeams (Stoelting, 60001-02A) were positioned to detect rearing. The mouse was placed in the bottom right-hand corner of the arena and behavior was recorded using an overhead camera and analyzed using ANY-maze software (Stoelting). An observer manually scored self-grooming behavior during the first 20 minutes of the test. A grooming bout was defined as an unbroken series of grooming movements, including licking of body, paws, or tail, as well as licking of forepaws followed by rubbing of face with paws.

Open field assay with DeepLabCut Keypoint-MoSeq analysis

Mice were placed in the open field arena and video recorded with a monochrome camera (FLIR Grasshopper 3, GS3-U3-41C6NIR-C) and a 16 mm wide angle lens (Kowa, LM16HC) placed above the arena from a height of 50 cm. To extract the body part (keypoint) coordinates from the video recordings, DeepLabCut (DLC) 2.3.4 (Mathis, et al. 2018 [DOI](#); Nath, et al. 2019) was used. Fourteen body parts including nose, head, left ear, right ear, left forelimb, right forelimb, spine 1, spine 2, spine 3, left hindlimb, right hindlimb, tail 1, tail 2, and tail 3 were manually labeled on a small subset of the video frames. A DLC model was then trained using the annotated frames to label those 14 body parts for all videos recorded. The total distance traveled, and number of center entries were calculated using the coordinate of bodypart tail 1.

Discrete behavior syllables were extracted using Keypoint-MoSeq 0.4.4 (Weinreb, et al. 2023). Syllable usage and transition data were obtained using built-in functions of the Keypoint-MoSeq package. Decoding and entropy analysis were performed using customized Python 3.9 script. Code available upon request in Github. Entropy was calculated using the following equation, where u_i denotes the frequency of the syllable i and $p_{i,j}$ denotes the transition probability from syllable i to syllable j :

$$Entropy = - \sum_{i,j} u_i \times p_{i,j} \times \log_2 p_{i,j}$$

Marble burying assay

The marble burying assay was used to test for repetitive behavior. 20 black marbles were arranged in an orderly 4 x 5 grid on top of 5 cm of clean corn cob bedding in a standard mouse cage. Overhead room lights were on and white noise was played to induce mild stress. Mice were

placed in the cage with the marbles for 30 minutes. The number of unburied marbles (>50% exposed) was recorded after the session.

Holeboard assay

The holeboard assay was used to measure exploratory and repetitive behavior. The holeboard apparatus consisted of a smooth, flat, opaque gray plastic platform, suspended 10 cm from the base by four plastic pegs in each corner. The board contained 16 evenly spaced 2 cm diameter holes and was surrounded by a 30 cm high clear plastic square encasing. During testing, mice were placed into the center of the holeboard. Mice explored the board for 10 minutes while video was recorded from both an above and side-view camera. Videos were used post-hoc to manually count and map the number of nose pokes made during the task. Nose pokes were defined as the mouse's nose passing through the board barrier when viewed through the side-view camera.

Accelerating rotarod assay

The accelerating rotarod test was used to examine motor coordination and learning.

Mice were trained on a rotarod apparatus (Ugo Basile, #47650) for four consecutive days. Three trials were completed per day with a 5 min break between trials. The rotarod was accelerated from 5-40 revolutions per minute (rpm) over 300 s for trials 1-6 (days 1 and 2), and from 10-80 rpm over 300 s for trials 7-12 (days 3 and 4). On the first testing day, mice were first acclimated to the apparatus by being placed on the rotarod rotating at a constant 5 rpm for 60 s and returned to their home cage for 5 min prior to starting trial 1. Latency to fall, or to rotate off the top of the rotarod barrel, was measured by the rotarod stop-trigger timer.

Four choice odor-based reversal learning test

The four-choice odor-based reversal learning test was used to assess learning and cognitive flexibility. Animals were food restricted for 6 days in total, with unrestricted access to drinking water, and maintained at 90-95% of ad lib feeding body weight. Food was given at the end of the day once testing was completed. Food restriction and introduction to Froot Loop cereal piece (Kellogg's, Battle Creek, MI) began 48 hours before pre-training. The four-choice test was performed in a custom-made square box (30.5 cm L × 30.5 cm W × 23 cm H) constructed of clear acrylic. Four internal walls 7.6 cm wide partially divided the arena into four quadrants. A 15.2 cm diameter removable cylinder fit in the center of the maze and was lowered between trials (after a digging response) to isolate the mouse from the rest of the maze. Odor stimuli were presented mixed with wood shavings in white ceramic pots measuring 7.3 cm in diameter and 4.5 cm deep. All pots were sham baited with a piece of Froot Loop cereal secured underneath a mesh screen at the bottom of the pot. This was to prevent mice from using the odor of the Froot Loop to guide their choice. The apparatus was cleaned with 2.5% acetic acid followed by water and the pots were cleaned with 70% ethanol followed by water between mice. The apparatus was cleaned with diluted soap and water at the end of each testing day.

On the first habituation day of pre-training (day 1), animals were allowed to freely explore the testing arena for 30 min and consume small pieces of Froot Loops placed inside empty pots positioned in each of the four corners. On the second shaping day of pre-training (day 2), mice learned to dig to find cereal pieces buried in unscented coarse pine wood shavings (Harts Mountain Corporation, Secaucus, NJ). A single pot was used and increasing amounts of unscented wood shavings were used to cover each subsequent cereal reward. The quadrant containing the pot was alternated on each trial and all quadrants were rewarded equally. Trials were untimed and consisted of (in order): two trials with no shavings, two trials with a dusting of shavings, two trials with the pot a quarter full, two trials with the pot half full, and four trials with the cereal piece completely buried by shavings. The mouse was manually returned to the center cylinder between trials.

On the days for odor discrimination (day 3, acquisition) and reversal (day 4), wood shavings were freshly scented on the day of testing. Anise extract (McCormick, Hunt Valley, MD) was used undiluted at 0.02 ml/g of shavings. Clove, litsea, and eucalyptus oils (San Francisco Massage Supply Co., San Francisco, CA) were diluted 1:10 in mineral oil and mixed at 0.02 ml/g of shavings. Thymol (thyme; Alfa Aesar, A14563) was diluted 1:20 in 50% ethanol and mixed at 0.01 ml/g of shavings. During the discrimination phase (day 3), mice had to discriminate between four pots with four different odors and learn which one contained a buried food reward. Each trial began with the mouse confined to the start cylinder. Once the cylinder was lifted, timing began, and the mouse could freely explore the arena until it chose to dig in a pot. Digging was defined as purposefully moving the shavings with both front paws. A trial was terminated if no choice was made within 3 min and recorded as omission. Criterion was met when the animal completed eight out of ten consecutive trials correctly. The spatial location of the odors was shuffled on each trial. The rewarded odor during acquisition was anise.

The first four odor choices made during acquisition were analyzed to determine innate odor preference by the percentage of choices for a given odor: *Cntnap2*^{+/+} mice: 60% thyme, 25% anise, 12.5% clove, and 2.5% litsea. *Cntnap2*^{-/-} mice: 47.5% thyme, 45% anise, 7.5% clove, 0% litsea. We note that both *Cntnap2*^{+/+} and *Cntnap2*^{-/-} mice exhibited the strongest innate preference for thyme, an unrewarded odor. There were no significant differences in innate odor preference.

The reversal phase of the task was carried out on day 4. Mice first performed the task with the same rewarded odor as the discrimination day to ensure they learned and remembered the task. After reaching criterion on recall (eight out of ten consecutive trials correct), the rewarded odor was switched, and mice underwent a reversal learning test in which a previously unrewarded odor (clove) was rewarded. A novel odor (eucalyptus) was also introduced, which replaced thyme. Perseverative errors were choices to dig in the previously rewarded odor that was no longer rewarded. Regressive errors were choosing the previously rewarded odor after the first correct choice of the newly rewarded odor. Novel errors were choices to dig in the pot with the newly introduced odor (eucalyptus). Irrelevant errors were choices to dig in the pot that had never been rewarded (litsea). Omissions were trials in which the mouse failed to make a digging choice within 3 min from the start of the trial. Total errors were the sum of perseverative, regressive, irrelevant, novel, and omission errors. Criterion was met when the mouse completed eight out of ten consecutive trials correctly. The spatial location of the odors was shuffled on each trial.

Quantification and statistical analysis

Experiments were designed to compare the main effect of genotype. The sample sizes were based on prior studies and are indicated in the figure legend for each experiment. Whenever possible, quantification and analyses were performed blind to genotype. GraphPad Prism version 10 was used to perform statistical analyses. The statistical tests and outcomes for each experiment are indicated in the respective figure legend. Two-tailed unpaired t tests were used for comparisons between two groups. For data that did not pass the D'Agostino & Pearson normality test, a Mann-Whitney test was used. Two-way ANOVAs or mixed effects models were used to compare differences between groups for experiments with two independent variables. Statistical significance was defined in the figure panels as follows: **p* < 0.05, ***p* < 0.01, ****p* < 0.001.

Acknowledgements

This work was supported by Simons Foundation Autism Research Initiative (SFARI) research grant #514428 to H.S.B. and NIH fellowship #F31NS124499 to K.R.C.

References

1. Ade K. K., Wan Y., Chen M., Gloss B., Calakos N (2011) **An Improved BAC Transgenic Fluorescent Reporter Line for Sensitive and Specific Identification of Striatonigral Medium Spiny Neurons** *Front Syst Neurosci* **5**:32 <https://doi.org/10.3389/fnsys.2011.00032>
2. Ahmed N. Y., Knowles R., Liu L., Yan Y., Li X., Schumann U., Wang Y., Sontani Y., Reynolds N., Natoli R., Wen J., Del Pino I., Mi D., Dehorter N (2023) **Developmental deficits of MGE-derived interneurons in the Cntnap2 knockout mouse model of autism spectrum disorder** *Front Cell Dev Biol* **11**:1112062 <https://doi.org/10.3389/fcell.2023.1112062>
3. Albelda N., Joel D (2012) **Animal models of obsessive-compulsive disorder: exploring pharmacology and neural substrates** *Neurosci Biobehav Rev* **36**:47–63 <https://doi.org/10.1016/j.neubiorev.2011.04.006>
4. Anderson G. R., Galfin T., Xu W., Aoto J., Malenka R. C., Sudhof T. C (2012) **Candidate autism gene screen identifies critical role for cell-adhesion molecule CASPR2 in dendritic arborization and spine development** *Proc Natl Acad Sci U S A* **109**:18120–18125 <https://doi.org/10.1073/pnas.1216398109>
5. Angoa-Perez M., Kane M. J., Briggs D. I., Francescutti D. M., Kuhn D. M (2013) **Marble burying and nestlet shredding as tests of repetitive, compulsive-like behaviors in mice** *J Vis Exp* **82** <https://doi.org/10.3791/50978>
6. Antoine M. W., Langberg T., Schnepel P., Feldman D. E (2019) **Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models** *Neuron* **101**:648–661 <https://doi.org/10.1016/j.neuron.2018.12.026>
7. APA (2022) **Diagnostic and Statistical Manual of Mental Disorders** <https://doi.org/10.1176/appi.books.9780890425787>
8. Arenkiel B. R., Peca J., Davison I. G., Feliciano C., Deisseroth K., Augustine G. J., Ehlers M. D., Feng G (2007) **In vivo light-induced activation of neural circuitry in transgenic mice expressing channelrhodopsin-2** *Neuron* **54**:205–218 <https://doi.org/10.1016/j.neuron.2007.03.005>
9. Benthall K. N., Cording K. R., Agopyan-Miu A., Wong C. D., Chen E. Y., Bateup H. S (2021) **Loss of Tsc1 from striatal direct pathway neurons impairs endocannabinoid-LTD and enhances motor routine learning** *Cell Rep* **36**:109511 <https://doi.org/10.1016/j.celrep.2021.109511>
10. Bouyer J. J., Park D. H., Joh T. H., Pickel V. M (1984) **Chemical and structural analysis of the relation between cortical inputs and tyrosine hydroxylase-containing terminals in rat neostriatum** *Brain Res* **302**:267–275 [https://doi.org/10.1016/0006-8993\(84\)90239-7](https://doi.org/10.1016/0006-8993(84)90239-7)
11. Brumback A. C., Ellwood I. T., Kjaerby C., Iafrati J., Robinson S., Lee A. T., Patel T., Nagaraj S., Davatolhagh F., Sohal V. S (2018) **Identifying specific prefrontal neurons that contribute to autism-associated abnormalities in physiology and social behavior** *Mol Psychiatry* **23**:2078–2089 <https://doi.org/10.1038/mp.2017.213>

12. Brunner D., Kabitzke P., He D., Cox K., Thiede L., Hanania T., Sabath E., Alexandrov V., Saxe M., Peles E., Mills A., Spooren W., Ghosh A., Feliciano P., Benedetti M., Luo Clayton A., Biemans B (2015) **Comprehensive Analysis of the 16p11.2 Deletion and Null Cntnap2 Mouse Models of Autism Spectrum Disorder** *PLoS One* **10**:e0134572 <https://doi.org/10.1371/journal.pone.0134572>
13. Burke D. A., Rotstein H. G., Alvarez V. A (2017) **Striatal Local Circuitry: A New Framework for Lateral Inhibition** *Neuron* **96**:267–284 <https://doi.org/10.1016/j.neuron.2017.09.019>
14. Calabresi P., Picconi B., Tozzi A., Ghiglieri V., Di Filippo M. (2014) **Direct and indirect pathways of basal ganglia: a critical reappraisal** *Nat Neurosci* **17**:1022–1030 <https://doi.org/10.1038/nn.3743>
15. Cifuentes-Diaz C., Canali G., Garcia M., Druart M., Manett T., Savariradjane M., Guillaume C., Le Magueresse C., Goutebroze L. (2023) **Differential impacts of Cntnap2 heterozygosity and Cntnap2 null homozygosity on axon and myelinated fiber development in mouse** *Front Neurosci* **17**:1100121 <https://doi.org/10.3389/fnins.2023.1100121>
16. Cording K. R., Bateup H. S (2023) **Altered motor learning and coordination in mouse models of autism spectrum disorder** *Front Cell Neurosci* **17**:1270489 <https://doi.org/10.3389/fncel.2023.1270489>
17. Cromwell H. C., Berridge K. C (1996) **Implementation of action sequences by a neostriatal site: a lesion mapping study of grooming syntax** *J Neurosci* **16**:3444–3458 <https://doi.org/10.1523/JNEUROSCI.16-10-03444.1996>
18. Dawes J. M., Weir G. A., Middleton S. J., Patel R., Chisholm K. I., Pettingill P., Peck L. J., Sheridan J., Shakir A., Jacobson L., Gutierrez-Mecinas M., Galino J., Walcher J., Kuhnemund J., Kuehn H., Sanna M. D., Lang B., Clark A. J., Themistocleous A. C., ..., Bennett D. L (2018) **Immune or Genetic-Mediated Disruption of CASPR2 Causes Pain Hypersensitivity Due to Enhanced Primary Afferent Excitability** *Neuron* **97**:806–822 <https://doi.org/10.1016/j.neuron.2018.01.033>
19. De Rubeis S., He X., Goldberg A. P., Poultney C. S., Samocha K., Cicek A. E., Kou Y., Liu L., Fromer M., Walker S., Singh T., Klei L., Kosmicki J., Shih-Chen F., Aleksic B., Biscaldi M., Bolton P. F., Brownfeld J. M., Cai J., ..., Buxbaum J. D. (2014) **Synaptic, transcriptional and chromatin genes disrupted in autism** *Nature* **515**:209–215 <https://doi.org/10.1038/nature13772>
20. Delevich K., Hoshal B., Zhou L. Z., Zhang Y., Vedula S., Lin W. C., Chase J., Collins A. G. E., Wilbrecht L. (2022) **Activation, but not inhibition, of the indirect pathway disrupts choice rejection in a freely moving, multiple-choice foraging task** *Cell Rep* **40**:111129 <https://doi.org/10.1016/j.celrep.2022.111129>
21. Delorme R., Ey E., Toro R., Leboyer M., Gillberg C., Bourgeron T (2013) **Progress toward treatments for synaptic defects in autism** *Nat Med* **19**:685–694 <https://doi.org/10.1038/nm.3193>
22. Ding J., Peterson J. D., Surmeier D. J (2008) **Corticostriatal and thalamostriatal synapses have distinctive properties** *J Neurosci* **28**:6483–6492 <https://doi.org/10.1523/JNEUROSCI.0435-08.2008>
23. Doig N. M., Moss J., Bolam J. P (2010) **Cortical and thalamic innervation of direct and indirect pathway medium-sized spiny neurons in mouse striatum** *J Neurosci* **30**:14610–14618 <https://doi.org/10.1523/JNEUROSCI.1623-10.2010>

24. Ebert D. H., Greenberg M. E (2013) **Activity-dependent neuronal signalling and autism spectrum disorder** *Nature* **493**:327–337 <https://doi.org/10.1038/nature11860>
25. Estes A., Shaw D. W., Sparks B. F., Friedman S., Giedd J. N., Dawson G., Bryan M., Dager S. R (2011) **Basal ganglia morphometry and repetitive behavior in young children with autism spectrum disorder** *Autism Res* **4**:212–220 <https://doi.org/10.1002/aur.193>
26. Evans M. M., Kim J., Abel T., Nickl-Jockschat T., Stevens H. E (2024) **Developmental Disruptions of the Dorsal Striatum in Autism Spectrum Disorder** *Biol Psychiatry* **95**:102–111 <https://doi.org/10.1016/j.biopsych.2023.08.015>
27. Fernandez M., Sanchez-Leon C. A., Llorente J., Sierra-Arregui T., Knafo S., Marquez-Ruiz J., Penagarikano O (2021) **Altered Cerebellar Response to Somatosensory Stimuli in the Cntnap2 Mouse Model of Autism** *eNeuro* **8**:5 <https://doi.org/10.1523/ENEURO.0333-21.2021>
28. Filice F., Janickova L., Henzi T., Bilella A., Schwaller B (2020) **The Parvalbumin Hypothesis of Autism Spectrum Disorder** *Front Cell Neurosci* **14**:577525 <https://doi.org/10.3389/fncel.2020.577525>
29. Fuccillo M. V (2016) **Striatal Circuits as a Common Node for Autism Pathophysiology** *Front Neurosci* **10**:27 <https://doi.org/10.3389/fnins.2016.00027>
30. Gdalyahu A., Lazaro M., Penagarikano O., Golshani P., Trachtenberg J. T., Geschwind D. H (2015) **Correction: The Autism Related Protein Contactin-Associated Protein-Like 2 (CNTNAP2) Stabilizes New Spines: An In Vivo Mouse Study** *PLoS One* **10**:e0129638 <https://doi.org/10.1371/journal.pone.0129638>
31. Gerfen C. R., Surmeier D. J (2011) **Modulation of striatal projection systems by dopamine** *Annu Rev Neurosci* **34**:441–466 <https://doi.org/10.1146/annurev-neuro-061010-113641>
32. Guan D., Lee J. C., Higgs M. H., Spain W. J., Foehring R. C (2007) **Functional roles of Kv1 channels in neocortical pyramidal neurons** *J Neurophysiol* **97**:1931–1940 <https://doi.org/10.1152/jn.00933.2006>
33. Haghdoust H., Janahmadi M., Behzadi G (2007) **Physiological role of dendrotoxin-sensitive K⁺ channels in the rat cerebellar Purkinje neurons** *Physiol Res* **56**:807–813 <https://doi.org/10.33549/physiolres.931041>
34. Hartig W., Brauer K., Bruckner G (1992) **Wisteria floribunda agglutinin-labelled nets surround parvalbumin-containing neurons** *Neuroreport* **3**:869–872 <https://doi.org/10.1097/00001756-199210000-00012>
35. Haunso A., Ibrahim M., Bartsch U., Letiembre M., Celio M. R., Menoud P (2000) **Morphology of perineuronal nets in tenascin-R and parvalbumin single and double knockout mice** *Brain Res* **864**:142–145 [https://doi.org/10.1016/s0006-8993\(00\)02173-9](https://doi.org/10.1016/s0006-8993(00)02173-9)
36. Hawes S. L., Evans R. C., Unruh B. A., Benkert E. E., Gillani F., Dumas T. C., Blackwell K. T (2015) **Multimodal Plasticity in Dorsal Striatum While Learning a Lateralized Navigation Task** *J Neurosci* **35**:10535–10549 <https://doi.org/10.1523/JNEUROSCI.4415-14.2015>

37. Hippenmeyer S., Vrieseling E., Sigrist M., Portmann T., Laengle C., Ladle D. R., Arber S (2005) **A developmental switch in the response of DRG neurons to ETS transcription factor signaling** *PLoS Biol* **3**:e159 <https://doi.org/10.1371/journal.pbio.0030159>
38. Hollander E., Anagnostou E., Chaplin W., Esposito K., Haznedar M. M., Licalzi E., Wasserman S., Soorya L., Buchsbaum M (2005) **Striatal volume on magnetic resonance imaging and repetitive behaviors in autism** *Biol Psychiatry* **58**:226–232 <https://doi.org/10.1016/j.biopsych.2005.03.040>
39. Inda M. C., DeFelipe J., Munoz A (2006) **Voltage-gated ion channels in the axon initial segment of human cortical pyramidal cells and their relationship with chandelier cells** *Proc Natl Acad Sci U S A* **103**:2920–2925 <https://doi.org/10.1073/pnas.0511197103>
40. Iossifov I., O’Roak B. J., Sanders S. J., Ronemus M., Krumm N., Levy D., Stessman H. A., Witherspoon K. T., Vives L., Patterson K. E., Smith J. D., Paepier B., Nickerson D. A., Dea J., Dong S., Gonzalez L. E., Mandell J. D., Mane S. M., Murtha M. T., ..., Wigler M (2014) **The contribution of de novo coding mutations to autism spectrum disorder** *Nature* **515**:216–221 <https://doi.org/10.1038/nature13908>
41. Izquierdo A., Brigman J. L., Radke A. K., Rudebeck P. H., Holmes A (2017) **The neural basis of reversal learning: An updated perspective** *Neuroscience* **345**:12–26 <https://doi.org/10.1016/j.neuroscience.2016.03.021>
42. Johnson C. M., Peckler H., Tai L. H., Wilbrecht L (2016) **Rule learning enhances structural plasticity of long-range axons in frontal cortex** *Nat Commun* **7**:10785 <https://doi.org/10.1038/ncomms10785>
43. Juarez P., Martinez Cerdeno V (2022) **Parvalbumin and parvalbumin chandelier interneurons in autism and other psychiatric disorders** *Front Psychiatry* **13**:913550 <https://doi.org/10.3389/fpsy.2022.913550>
44. Jurgensen S., Castillo P. E (2015) **Selective Dysregulation of Hippocampal Inhibition in the Mouse Lacking Autism Candidate Gene CNTNAP2** *J Neurosci* **35**:14681–14687 <https://doi.org/10.1523/JNEUROSCI.1666-15.2015>
45. Keaveney M. K., Tseng H. A., Ta T. L., Gritton H. J., Man H. Y., Han X (2018) **A MicroRNA-Based Gene-Targeting Tool for Virally Labeling Interneurons in the Rodent Cortex** *Cell Rep* **24**:294–303 <https://doi.org/10.1016/j.celrep.2018.06.049>
46. Khavandgar S., Walter J. T., Sageser K., Khodakhah K (2005) **Kv1 channels selectively prevent dendritic hyperexcitability in rat Purkinje cells** *J Physiol* **569**:545–557 <https://doi.org/10.1111/jphysiol.2005.098053>
47. Kosillo P., Bateup H. S (2021) **Dopaminergic Dysregulation in Syndromic Autism Spectrum Disorders: Insights From Genetic Mouse Models** *Front Neural Circuits* **15**:700968 <https://doi.org/10.3389/fncir.2021.700968>
48. Kosillo P., Doig N. M., Ahmed K. M., Agopyan-Miu A., Wong C. D., Conyers L., Threlfell S., Magill P. J., Bateup H. S (2019) **Tsc1-mTORC1 signaling controls striatal dopamine release and cognitive flexibility** *Nat Commun* **10**:5426 <https://doi.org/10.1038/s41467-019-13396-8>
49. Kravitz A. V., Freeze B. S., Parker P. R., Kay K., Thwin M. T., Deisseroth K., Kreitzer A. C (2010) **Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry** *Nature* **466**:622–626 <https://doi.org/10.1038/nature09159>

50. Lahiri A. K., Bevan M. D (2020) **Dopaminergic Transmission Rapidly and Persistently Enhances Excitability of D1 Receptor-Expressing Striatal Projection Neurons** *Neuron* **106**:277–290 <https://doi.org/10.1016/j.neuron.2020.01.028>
51. Langen M., Bos D., Noordermeer S. D., Nederveen H., van Engeland H., Durston S. (2014) **Changes in the development of striatum are involved in repetitive behavior in autism** *Biol Psychiatry* **76**:405–411 <https://doi.org/10.1016/j.biopsych.2013.08.013>
52. Lauber E., Filice F., Schwaller B (2018) **Dysregulation of Parvalbumin Expression in the Cntnap2-/-Mouse Model of Autism Spectrum Disorder** *Front Mol Neurosci* **11**:262 <https://doi.org/10.3389/fnmol.2018.00262>
53. Lazaro M. T., Taxidis J., Shuman T., Bachmutsky I., Ikrar T., Santos R., Marcello G. M., Mylavaram A., Chandra S., Foreman A., Goli R., Tran D., Sharma N., Azhdam M., Dong H., Choe K. Y., Penagarikano O., Masmanidis S. C., Racz B., ..., Golshani P (2019) **Reduced Prefrontal Synaptic Connectivity and Disturbed Oscillatory Population Dynamics in the CNTNAP2 Model of Autism** *Cell Rep* **27**:2567–2578 <https://doi.org/10.1016/j.celrep.2019.05.006>
54. Lee I. B., Lee E., Han N. E., Slavuj M., Hwang J. W., Lee A., Sun T., Jeong Y., Baik J. H., Park J. Y., Choi S. Y., Kwag J., Yoon B. J (2024) **Persistent enhancement of basolateral amygdala-dorsomedial striatum synapses causes compulsive-like behaviors in mice** *Nat Commun* **15**:219 <https://doi.org/10.1038/s41467-023-44322-8>
55. Li W., Pozzo-Miller L (2020) **Dysfunction of the corticostriatal pathway in autism spectrum disorders** *J Neurosci Res* **98**:2130–2147 <https://doi.org/10.1002/jnr.24560>
56. Lin W. C., Liu C., Kosillo P., Tai L. H., Galarce E., Bateup H. S., Lammel S., Wilbrecht L (2022) **Transient food insecurity during the juvenile-adolescent period affects adult weight, cognitive flexibility, and dopamine neurobiology** *Curr Biol* **32**:3690–3703 <https://doi.org/10.1016/j.cub.2022.06.089>
57. Madisen L., Mao T., Koch H., Zhuo J. M., Berenyi A., Fujisawa S., Hsu Y. W., Garcia A. J., Gu X., Zanella S., Kidney J., Gu H., Mao Y., Hooks B. M., Boyden E. S., Buzsaki G., Ramirez J. M., Jones A. R., Svoboda K., ..., Zeng H. (2012) **A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing** *Nat Neurosci* **15**:793–802 <https://doi.org/10.1038/nn.3078>
58. Madisen L., Zwingman T. A., Sunkin S. M., Oh S. W., Zariwala H. A., Gu H., Ng L. L., Palmiter R. D., Hawrylycz M. J., Jones A. R., Lein E. S., Zeng H (2010) **A robust and high-throughput Cre reporting and characterization system for the whole mouse brain** *Nat Neurosci* **13**:133–140 <https://doi.org/10.1038/nn.2467>
59. Martin-de-Saavedra M. D., Dos Santos M., Culotta L., Varea O., Spielman B. P., Parnell E., Forrest M. P., Gao R., Yoon S., McCoig E., Jalloul H. A., Myczek K., Khalatyan N., Hall E. A., Turk L. S., Sanz-Clemente A., Comoletti D., Lichtenthaler S. F., Burgdorf J. S., ..., Penzes P (2022) **Shed CNTNAP2 ectodomain is detectable in CSF and regulates Ca(2+) homeostasis and network synchrony via PMCA2/ATP2B2** *Neuron* **110**:627–643 <https://doi.org/10.1016/j.neuron.2021.11.025>
60. Mathis A., Mamidanna P., Cury K. M., Abe T., Murthy V. N., Mathis M. W., Bethge M (2018) **DeepLabCut: markerless pose estimation of user-defined body parts with deep learning** *Nat Neurosci* **21**:1281–1289 <https://doi.org/10.1038/s41593-018-0209-y>

61. Nisenbaum E. S., Xu Z. C., Wilson C. J (1994) **Contribution of a slowly inactivating potassium current to the transition to firing of neostriatal spiny projection neurons** *J Neurophysiol* **71**:1174–1189 <https://doi.org/10.1152/jn.1994.71.3.1174>
62. Orduz D., Bischof D. P., Schwaller B., Schiffmann S. N., Gall D (2013) **Parvalbumin tunes spike-timing and efferent short-term plasticity in striatal fast spiking interneurons** *J Physiol* **591**:3215–3232 <https://doi.org/10.1113/jphysiol.2012.250795>
63. Packard M. G., Knowlton B. J (2002) **Learning and memory functions of the Basal Ganglia** *Annu Rev Neurosci* **25**:563–593 <https://doi.org/10.1146/annurev.neuro.25.112701.142937>
64. Paterno R., Marafija J. R., Ramsay H., Li T., Salvati K. A., Baraban S. C (2021) **Hippocampal gamma and sharp-wave ripple oscillations are altered in a Cntnap2 mouse model of autism spectrum disorder** *Cell Rep* **37**:109970 <https://doi.org/10.1016/j.celrep.2021.109970>
65. Paval D., Miclutia I. V (2021) **The Dopamine Hypothesis of Autism Spectrum Disorder Revisited: Current Status and Future Prospects** *Dev Neurosci* **43**:73–83 <https://doi.org/10.1159/000515751>
66. Peca J., Feliciano C., Ting J. T., Wang W., Wells M. F., Venkatraman T. N., Lascola C. D., Fu Z., Feng G (2011) **Shank3 mutant mice display autistic-like behaviours and striatal dysfunction** *Nature* **472**:437–442 <https://doi.org/10.1038/nature09965>
67. Peixoto R. T., Wang W., Croney D. M., Kozorovitskiy Y., Sabatini B. L (2016) **Early hyperactivity and precocious maturation of corticostriatal circuits in Shank3B(-/-) mice** *Nat Neurosci* **19**:716–724 <https://doi.org/10.1038/nn.4260>
68. Penagarikano O., Abrahams B. S., Herman E. I., Winden K. D., Gdalyahu A., Dong H., Sonnenblick L. I., Gruver R., Almajano J., Bragin A., Golshani P., Trachtenberg J. T., Peles E., Geschwind D. H (2011) **Absence of CNTNAP2 leads to epilepsy, neuronal migration abnormalities, and core autism-related deficits** *Cell* **147**:235–246 <https://doi.org/10.1016/j.cell.2011.08.040>
69. Penagarikano O., Geschwind D. H (2012) **What does CNTNAP2 reveal about autism spectrum disorder?** *Trends Mol Med* **18**:156–163 <https://doi.org/10.1016/j.molmed.2012.01.003>
70. Platt R. J., Zhou Y., Slaymaker I. M., Shetty A. S., Weisbach N. R., Kim J. A., Sharma J., Desai M., Sood S., Kempton H. R., Crabtree G. R., Feng G., Zhang F (2017) **Chd8 Mutation Leads to Autistic-like Behaviors and Impaired Striatal Circuits** *Cell Rep* **19**:335–350 <https://doi.org/10.1016/j.celrep.2017.03.052>
71. Poliak S., Gollan L., Martinez R., Custer A., Einheber S., Salzer J. L., Trimmer J. S., Shrager P., Peles E (1999) **Caspr2, a new member of the neurexin superfamily, is localized at the juxtaparanodes of myelinated axons and associates with K⁺ channels** *Neuron* **24**:1037–1047 [https://doi.org/10.1016/s0896-6273\(00\)81049-1](https://doi.org/10.1016/s0896-6273(00)81049-1)
72. Poliak S., Salomon D., Elhanany H., Sabanay H., Kiernan B., Pevny L., Stewart C. L., Xu X., Chiu S. Y., Shrager P., Furley A. J., Peles E (2003) **Juxtaparanodal clustering of Shaker-like K⁺ channels in myelinated axons depends on Caspr2 and TAG-1** *J Cell Biol* **162**:1149–1160 <https://doi.org/10.1083/jcb.200305018>
73. Ramirez-Armenta K. I., Alatraste-Leon H., Verma-Rodriguez A. K., Llanos-Moreno A., Ramirez-Jarquín J. O., Tecuapetla F (2022) **Optogenetic inhibition of indirect pathway neurons in the**

dorsomedial striatum reduces excessive grooming in Sapap3-knockout mice

Neuropsychopharmacology **47**:477–487 <https://doi.org/10.1038/s41386-021-01161-9>

74. Rapanelli M., Frick L. R., Pittenger C (2017) **The Role of Interneurons in Autism and Tourette Syndrome** *Trends Neurosci* **40**:397–407 <https://doi.org/10.1016/j.tins.2017.05.004>
75. Rodenas-Cuadrado P., Ho J., Vernes S. C (2014) **Shining a light on CNTNAP2: complex functions to complex disorders** *Eur J Hum Genet* **22**:171–178 <https://doi.org/10.1038/ejhg.2013.100>
76. Rothwell P. E., Fuccillo M. V., Maxeiner S., Hayton S. J., Gokce O., Lim B. K., Fowler S. C., Malenka R. C., Sudhof T. C (2014) **Autism-associated neuroligin-3 mutations commonly impair striatal circuits to boost repetitive behaviors** *Cell* **158**:198–212 <https://doi.org/10.1016/j.cell.2014.04.045>
77. Sanders S. J., He X., Willsey A. J., Ercan-Sencicek A. G., Samocha K. E., Cicek A. E., Murtha M. T., Bal V. H., Bishop S. L., Dong S., Goldberg A. P., Jinlu C., Keaney J. F., Klei L., Mandell J. D., Moreno-De-Luca D., Poultney C. S., Robinson E. B., Smith L., ..., State M. W. (2015) **Insights into Autism Spectrum Disorder Genomic Architecture and Biology from 71 Risk Loci** *Neuron* **87**:1215–1233 <https://doi.org/10.1016/j.neuron.2015.09.016>
78. Sandin S., Lichtenstein P., Kuja-Halkola R., Hultman C., Larsson H., Reichenberg A (2017) **The Heritability of Autism Spectrum Disorder** *JAMA* **318**:1182–1184 <https://doi.org/10.1001/jama.2017.12141>
79. Santos F. J., Oliveira R. F., Jin X., Costa R. M (2015) **Corticostriatal dynamics encode the refinement of specific behavioral variability during skill learning** *eLife* **4**:e09423 <https://doi.org/10.7554/eLife.09423>
80. Satterstrom F. K., Kosmicki J. A., Wang J., Breen M. S., De Rubeis S., An J.-Y., Peng M., Collins R., Grove J., Klei L., Stevens C., Reichert J., Mulhern M. S., Artomov M., Gerges S., Sheppard B., Xu X., Bhaduri A., Norman U., ..., Walters R. K. (2020) **Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism** *Cell* **180**:568–584 <https://doi.org/10.1016/j.cell.2019.12.036>
81. Scott R., Sanchez-Aguilera A., van Elst K., Lim L., Dehorter N., Bae S. E., Bartolini G., Peles E., Kas M. J. H., Bruining H., Marin O. (2019) **Loss of Cntnap2 Causes Axonal Excitability Deficits, Developmental Delay in Cortical Myelination, and Abnormal Stereotyped Motor Behavior** *Cereb Cortex* **29**:586–597 <https://doi.org/10.1093/cercor/bhx341>
82. Shen W., Hernandez-Lopez S., Tkatch T., Held J. E., Surmeier D. J (2004) **Kv1.2-containing K⁺ channels regulate subthreshold excitability of striatal medium spiny neurons** *J Neurophysiol* **91**:1337–1349 <https://doi.org/10.1152/jn.00414.2003>
83. Subramanian K., Brandenburg C., Orsati F., Soghomonian J. J., Hussman J. P., Blatt G. J (2017) **Basal ganglia and autism – a translational perspective** *Autism Res* **10**:1751–1775 <https://doi.org/10.1002/aur.1837>
84. Tai L. H., Lee A. M., Benavidez N., Bonci A., Wilbrecht L (2012) **Transient stimulation of distinct subpopulations of striatal neurons mimics changes in action value** *Nat Neurosci* **15**:1281–1289 <https://doi.org/10.1038/nn.3188>
85. Uddin L. Q (2021) **Cognitive and behavioural flexibility: neural mechanisms and clinical considerations** *Nat Rev Neurosci* **22**:167–179 <https://doi.org/10.1038/s41583-021-00428-w>

86. Varea O., Martin-de-Saavedra M. D., Kopeikina K. J., Schurmann B., Fleming H. J., Fawcett-Patel J. M., Bach A., Jang S., Peles E., Kim E., Penzes P (2015) **Synaptic abnormalities and cytoplasmic glutamate receptor aggregates in contactin associated protein-like 2/Caspr2 knockout neurons** *Proc Natl Acad Sci U S A* **112**:6176–6181 <https://doi.org/10.1073/pnas.1423205112>
87. Vogt D., Cho K. K. A., Shelton S. M., Paul A., Huang Z. J., Sohal V. S., Rubenstein J. L. R (2018) **Mouse Cntnap2 and Human CNTNAP2 ASD Alleles Cell Autonomously Regulate PV+ Cortical Interneurons** *Cereb Cortex* **28**:3868–3879 <https://doi.org/10.1093/cercor/bhx248>
88. Voorn P., Vanderschuren L. J., Groenewegen H. J., Robbins T. W., Pennartz C. M (2004) **Putting a spin on the dorsal-ventral divide of the striatum** *Trends Neurosci* **27**:468–474 <https://doi.org/10.1016/j.tins.2004.06.006>
89. Wang X., Bey A. L., Katz B. M., Badea A., Kim N., David L. K., Duffney L. J., Kumar S., Mague S. D., Hulbert S. W., Dutta N., Hayrapetyan V., Yu C., Gaidis E., Zhao S., Ding J. D., Xu Q., Chung L., Rodriguiz R. M., ..., Jiang Y. H (2016) **Altered mGluR5-Homer scaffolds and corticostriatal connectivity in a Shank3 complete knockout model of autism** *Nat Commun* **7**:11459 <https://doi.org/10.1038/ncomms11459>
90. Wiltchko A. B., Tsukahara T., Zeine A., Anyoha R., Gillis W. F., Markowitz J. E., Peterson R. E., Katon J., Johnson M. J., Datta S. R (2020) **Revealing the structure of pharmacobehavioral space through motion sequencing** *Nat Neurosci* **23**:1433–1443 <https://doi.org/10.1038/s41593-020-00706-3>
91. Xu Z. C., Wilson C. J., Emson P. C (1989) **Restoration of the corticostriatal projection in rat neostriatal grafts: electron microscopic analysis** *Neuroscience* **29**:539–550 [https://doi.org/10.1016/0306-4522\(89\)90129-2](https://doi.org/10.1016/0306-4522(89)90129-2)
92. Yin H. H., Knowlton B. J., Balleine B. W (2005) **Blockade of NMDA receptors in the dorsomedial striatum prevents action-outcome learning in instrumental conditioning** *Eur J Neurosci* **22**:505–512 <https://doi.org/10.1111/j.1460-9568.2005.04219.x>
93. Yin H. H., Knowlton B. J., Balleine B. W (2006) **Inactivation of dorsolateral striatum enhances sensitivity to changes in the action-outcome contingency in instrumental conditioning** *Behav Brain Res* **166**:189–196 <https://doi.org/10.1016/j.bbr.2005.07.012>
94. Yin H. H., Mulcare S. P., Hilário M. R. F., Clouse E., Holloway T., Davis M. I., Hansson A. C., Lovinger D. M., Costa R. M (2009) **Dynamic reorganization of striatal circuits during the acquisition and consolidation of a skill** *Nature Neuroscience* **12**:333–341 <https://doi.org/10.1038/nn.2261>
95. Zhang Y. F., Vargas Cifuentes L., Wright K. N., Bhattarai J. P., Mohrhardt J., Fleck D., Janke E., Jiang C., Cranfill S. L., Goldstein N., Schreck M., Moberly A. H., Yu Y., Arenkiel B. R., Betley J. N., Luo W., Stegmaier J., Wesson D. W., Spehr M., ..., Ma M (2021) **Ventral striatal islands of Calleja neurons control grooming in mice** *Nat Neurosci* **24**:1699–1710 <https://doi.org/10.1038/s41593-021-00952-z>

Author information

Katherine R Cording

Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, United States,
Department of Neuroscience, University of California, Berkeley, Berkeley, United States

Emilie M Tu

Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, United States

Hongli Wang

Department of Neuroscience, University of California, Berkeley, Berkeley, United States

Alexander HCW Agopyan-Miu

Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, United States

Helen S Bateup

Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, United States,
Department of Neuroscience, University of California, Berkeley, Berkeley, United States,
Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, United States

ORCID iD: [0000-0002-0135-0972](https://orcid.org/0000-0002-0135-0972)

For correspondence: bateup@berkeley.edu

Editors

Reviewing Editor

Jun Ding

Stanford University, Stanford, United States of America

Senior Editor

Kate Wassum

University of California, Los Angeles, Los Angeles, United States of America

Reviewer #1 (Public review):

Summary:

Cording et al. investigated how deletion of CNTNAP2, a gene associated with autism spectrum disorder, alters corticostriatal engagement and behavior. Specifically, the authors present slice electrophysiology data showing that striatal projection neurons (SPNs) are more readily driven to fire action potentials in response to stimulation of corticostriatal afferents, and this is due to increases in SPN intrinsic excitability rather than changes in excitatory or inhibitory synaptic inputs. Specifically, these changes seem to be due to preferential reduction of Kv1.2 in dSPNs. The authors separately show that CNTNAP2 mice display repetitive behaviors, enhanced motor learning and cognitive inflexibility. Overall, the authors' conclusions are supported by their data, but a few claims could use some more evidence to be convincing.

Strengths:

The use of multiple behavioral techniques, both traditional and cutting-edge machine learning-based analyses, provides a powerful means of assessing repetitive behaviors and behavioral transitions/rigidity. Characterization of both excitatory and inhibitory synaptic

responses in slice electrophysiology experiments offers a broad survey of the synaptic alterations that may lead to increased corticostriatal engagement of SPNs.

Weaknesses:

As it stands, the reported changes in dorsolateral striatum SPN excitability are only correlative with reported changes in repetitive behaviors, motor learning and cognitive flexibility. The authors do broach this in the text (particularly in "Limitations and future directions").

<https://doi.org/10.7554/eLife.100162.2.sa3>

Reviewer #2 (Public review):

Summary:

This is an important study characterizing striatal dysfunction and behavioral deficits in *Cntnap2*^{-/-} mice. There is growing evidence suggesting that striatal dysfunction underlies core symptoms of ASD but the specific cellular and circuit level abnormalities disrupted by different risk genes remain unclear. This study addresses how deletion of *Cntnap2* affects the intrinsic properties and synaptic connectivity of striatal spiny projection neurons (SPN) of the direct (dSPN) and indirect (iSPN) pathways. Using Thy1-ChR2 mice and optogenetics the authors found increased firing of both types of SPNs in response to cortical afferent stimulation. However, there was no significant difference in the amplitude of optically-evoked excitatory postsynaptic currents (EPSCs) or spine density between *Cntnap2*^{-/-} and WT SPNs, suggesting that the increased corticostriatal coupling might be due to changes in intrinsic excitability. Indeed, the authors found *Cntnap2*^{-/-} SPNs, particularly dSPNs, exhibited higher intrinsic excitability, reduced rheobase current and increased membrane resistance compared to WT SPNs. The enhanced spiking probability in *Cntnap2*^{-/-} SPNs is not due to reduced inhibition. Despite previous reports of decreased parvalbumin-expressing (PV) interneurons in various brain regions of *Cntnap2*^{-/-} mice, the number and function (IPSC amplitude and intrinsic excitability) of these interneurons in the striatum were comparable to WT controls.

This study also includes a comprehensive behavioral analysis of striatal related behaviors. *Cntnap2*^{-/-} mice demonstrated increased repetitive behaviors (RRBs), including more grooming bouts, increased marble burying, and increased nose poking in the holeboard assay. MoSeq analysis of behavior further showed signs of altered grooming behaviors and sequencing of behavioral syllables. *Cntnap2*^{-/-} mice also displayed cognitive inflexibility in a four-choice odor-based reversal learning assay. While they performed similarly to WT controls during acquisition and recall phases, they required significantly more trials to learn a new odor-reward association during reversal, consistent with potential deficits in corticostriatal function.

Strengths:

This study provides significant contributions to the field. The finding of altered SPN excitability, the detailed characterization of striatal inhibition, and the comprehensive behavioral analysis are novel and valuable to understand the pathophysiology of *Cntnap2*^{-/-} mice.

Weaknesses:

All my concerns were addressed in the revised version of the manuscript

<https://doi.org/10.7554/eLife.100162.2.sa2>

Reviewer #3 (Public review):

Summary:

The authors analyzed *Cntnap2* KO mice to determine whether loss of the ASD risk gene *CNTNAP2* alters the dorsal striatum's function.

Strengths:

The results demonstrate that loss of *Cntnap2* results in increased excitability of striatal projection neurons (SPNs) and altered striatal-dependent behaviors, such as repetitive, inflexible behaviors. Unlike other brain areas and cell types, synaptic inputs onto SPNs were normal in *Cntnap2* KO mice. The experiments are well-designed, and the results support the authors' conclusions.

Weaknesses:

The mechanism underlying SPN hyperexcitability was not explored, and it is unclear whether this cellular phenotype alone can account for the behavioral alterations in *Cntnap2* KO mice. No clear explanation emerges for the variable phenotype in different brain areas and cell types.

Comments on revisions:

The authors have appropriately addressed all my comments. In my opinion, no further changes are required.

<https://doi.org/10.7554/eLife.100162.2.sa1>

Author response:

The following is the authors' response to the original reviews

We thank the reviewers for their careful review of our manuscript and the constructive comments. We have addressed the majority of comments with either new experiments, analyses, and/or text revisions. A summary of the major changes is listed below, followed by our point-by-point responses to the reviewer comments.

Major changes:

(1) We sought to gain insight into the potential mechanistic cause of the increased intrinsic excitability of *Cntnap2*^{-/-} dSPNs. Given that Kv1.1 and 1.2 potassium channels are known to interact with Caspr2 (the protein encoded by *Cntnap2*), we hypothesized that altered number, location, and/or function of these channels may underlie the excitability change in these cells. To investigate this, we performed new analyses of the initial dataset to assess action potential (AP) properties known to be impacted by potassium channel function. Indeed, we found that AP frequency was increased, and rheobase current, AP latency and AP threshold were decreased in *Cntnap2*^{-/-} dSPNs, suggestive of altered Kv1.2 function. These data are in the new Supplemental Fig. 4. We also performed new electrophysiology experiments in which we pharmacologically blocked Kv1.1 and 1.2 to assess whether the effects of blocking these channels would be occluded in *Cntnap2*^{-/-} dSPNs. We found that 1) WT dSPNs responded to blockade of Kv1.1/1.2 channels by increasing their excitability but *Cntnap2*^{-/-} dSPNs did not and 2) Kv1.1/1.2 channels were more important contributors to the excitability of dSPNs compared to iSPNs. These new data are presented in the revised Fig. 4 and Supplemental Figs. 5 and 6.

(2) We performed additional experiments to assess excitatory synaptic properties, specifically AMPA/NMDA receptor ratio. This has been added to Fig. 1.

(3) We performed more rigorous statistical analyses of the initial physiology datasets to align with the statistics performed for the revision experiments. This applies to Fig. 1, Fig. 2, Fig. 3, Fig. 5, and Supp. Fig. 2.

(4) In the discussion section, we now highlight potential limitations of the study and further discuss the variable impact that *Cntnap2* loss has on different cell types and brain regions.

Reviewer #1 (Public Review):

Summary:

Cording et al. investigated how deletion of CNTNAP2, a gene associated with autism spectrum disorder, alters corticostriatal engagement and behavior. Specifically, the authors present slice electrophysiology data showing that striatal projection neurons (SPNs) are more readily driven to fire action potentials in response to stimulation of corticostriatal afferents, and this is due to increases in SPN intrinsic excitability rather than changes in excitatory or inhibitory synaptic inputs. The authors show that CNTNAP2 mice display repetitive behaviors, enhanced motor learning, and cognitive inflexibility. Overall the authors' conclusions are supported by their data, but a few claims could use some more evidence to be convincing.

Strengths:

The use of multiple behavioral techniques, both traditional and cutting-edge machine learning-based analyses, provides a powerful means of assessing repetitive behaviors and behavioral transitions/rigidity.

Characterization of both excitatory and inhibitory synaptic responses in slice electrophysiology experiments offers a broad survey of the synaptic alterations that may lead to increased corticostriatal engagement of SPNs.

Weaknesses:


(1) The authors conclude that increased cortical engagement of SPNs is due to changes in SPN intrinsic excitability rather than synaptic strength (either excitatory or inhibitory). One weakness is that only AMPA receptor-mediated responses were measured. Though the holding potential used for experiments in Figure 1F wasn't clear, recordings were presumably performed at a hyperpolarized potential that limits NMDA receptor-mediated responses. Because the input-output experiments used to conclude that corticostriatal engagement of SPNs is elevated (Figure 1B-E) were conducted in the current clamp, it is possible that enhanced NMDA receptor engagement contributed to increased SPN responses to cortical stimulation. Confirming that NMDA receptor-mediated EPSC components are not altered would strengthen the main conclusion.

The reviewer is correct, the initial optically-evoked EPSC assessments were performed at a hyperpolarized potential (-70mV), thus measuring primarily AMPAR-mediated currents. We agree that assessing potential changes in the NMDAR-mediated EPSC component is important and we have completed new experiments to assess this. We find no differences in NMDAR-mediated EPSCs assessed at +40mV or the AMPA:NMDA ratio.

These results have been added to Fig. 1. An expanded analysis of these results is shown in Author response image 1. We note that the previous AMPAR-mediated EPSC results have been replicated in this additional experiment, again showing no change in *Cntnap2*^{-/-} SPNs.

Author response image 1.

AMPA and NMDA receptor-mediated EPSCs are unchanged in *Cntnap2*^{-/-} SPNs. (A) Quantification (mean ± SEM) of AMPA:NMDA ratio per cell for *Cntnap2*^{+/+} and *Cntnap2*^{-/-} dSPNs, p=0.9537, Mann-Whitney test. (B) dSPN AMPA current per cell, p=0.6172, Mann-Whitney test. (C) dSPN NMDA current per cell, p=0.6009, Mann-Whitney test. (D) dSPN AMPA:NMDA ratio averaged by animal, p=0.8413, Mann-Whitney test. (E) dSPN AMPA current averaged by animal, p>0.9999, Mann-Whitney test. (F) dSPN NMDA current averaged by animal, p=0.6905, Mann-Whitney test. (G) Quantification (mean ± SEM) of AMPA:NMDA ratio per cell for *Cntnap2*^{+/+} and *Cntnap2*^{-/-} iSPNs, p=0.4104, Mann-Whitney test. (H) iSPN AMPA current per cell, p=0.9010, Mann-Whitney test. (I) iSPN NMDA current per cell, p=0.9512, two-tailed unpaired t test. (J) iSPN AMPA:NMDA averaged by animal, p=0.3095, Mann-Whitney test. (K) iSPN AMPA current averaged by animal, p>0.9999, Mann-Whitney test. (L) iSPN NMDA current averaged by animal, p=0.8413, Mann-Whitney test. All values were recorded using 20% blue light intensity. For dSPNs: *Cntnap2*^{+/+} n=22 cells from 5 mice, *Cntnap2*^{-/-} n=22 cells from 5 mice. For iSPNs: *Cntnap2*^{+/+} n=21 cells from 5 mice, *Cntnap2*^{-/-} n=21 cells from 5 mice.



(2) Data clearly show that SPN intrinsic excitability is increased in knockout mice. Given that CNTNAP2 has been linked to potassium channel regulation, it would be helpful to

show and quantify additional related electrophysiology data such as negative IV curve responses and action potential hyperpolarization.

We appreciate this suggestion. As indicated by the reviewer, Caspr2, has previously been shown to control the clustering of Kv1-family potassium channels in axons isolated from optic nerve and corpus callosum (PMIDs: 10624965, 12963709, 29300891). In particular, Caspr2 is known to associate directly with Kv1.2 (PMID: 29300891). To assess a potential contribution of Kv1.2 to the excitability phenotype, we performed additional analyses of our original dataset to quantify AP properties known to be impacted by changes in Kv1.2 function (i.e. latency to fire and AP threshold, new Supp. Fig. 4). We identified several changes in *Cntnap2*^{-/-} dSPNs resembling those that occur in wild-type cells when Kv1.2 is blocked (i.e. reduced threshold and reduced latency to fire, Supp. Fig. 4).

We then performed a pharmacological experiment, blocking Kv1.2 using α -dendrotoxin (α -DTX) while recording intrinsic excitability to assess whether the effects of this drug on dSPN excitability were occluded in *Cntnap2*^{-/-} cells. Indeed, we found that while blocking Kv1.2 in wild-type dSPNs significantly reduced threshold and increased intrinsic excitability, these effects were not seen in *Cntnap2*^{-/-} dSPNs (new Fig. 4). We believe that this suggests an altered contribution of Kv1.2 to the intrinsic excitability of mutant dSPNs, owing to a change in the clustering, number, or function of these channels. Therefore, loss-of-function of Kv1.2 is a likely explanation for the enhanced intrinsic excitability of *Cntnap2*^{-/-} dSPNs. Interestingly, we found that α -DTX had only subtle effects on iSPNs (*Cntnap2* WT or mutant), suggesting a lesser contribution of this channel in controlling the excitability of indirect pathway cells. This finding can account for the relatively stronger effect of *Cntnap2* loss on dSPN physiology. The results of these new experiments and analyses are presented in the new Fig. 4, Supp. Fig. 5 and Supp. Fig. 6.

(3) As it stands, the reported changes in dorsolateral striatum SPN excitability are only correlative with reported changes in repetitive behaviors, motor learning, and cognitive flexibility.

We agree that we have not identified a causative relationship between the change in dorsolateral dSPN excitability and the behaviors that we measured in *Cntnap2*^{-/-} mice. That said, in a previous study, we showed that selective deletion of the autism spectrum disorder (ASD) risk gene *Tsc1* from dorsal striatal dSPNs resulted in increased corticostriatal drive and this was sufficient to increase rotarod motor learning (PMID: 34380034). Therefore, while we have not demonstrated causality in this study, we hypothesize that changes in dSPN excitability are likely to contribute to the behavioral phenotypes observed in *Cntnap2*^{-/-} mice.

Reviewer #2 (Public Review):

Summary:

*This is an important study characterizing striatal dysfunction and behavioral deficits in *Cntnap2*^{-/-} mice. There is growing evidence suggesting that striatal dysfunction underlies core symptoms of ASD but the specific cellular and circuit level abnormalities disrupted by different risk genes remain unclear. This study addresses how the deletion of *Cntnap2* affects the intrinsic properties and synaptic connectivity of striatal spiny projection neurons (SPN) of the direct (dSPN) and indirect (iSPN) pathways. Using *Thy1-ChR2* mice and optogenetics the authors found increased firing of both types of SPNs in response to cortical afferent stimulation. However, there was no significant difference in the amplitude of optically-evoked excitatory postsynaptic currents (EPSCs) or spine density between *Cntnap2*^{-/-} and WT SPNs, suggesting that the increased corticostriatal coupling might be due to changes in intrinsic excitability. Indeed, the authors found *Cntnap2*^{-/-} SPNs, particularly dSPNs, exhibited higher intrinsic excitability, reduced rheobase*

current, and increased membrane resistance compared to WT SPNs. The enhanced spiking probability in *Cntnap2*^{-/-} SPNs is not due to reduced inhibition. Despite previous reports of decreased parvalbumin-expressing (PV) interneurons in various brain regions of *Cntnap2*^{-/-} mice, the number and function (IPSC amplitude and intrinsic excitability) of these interneurons in the striatum were comparable to WT controls.

This study also includes a comprehensive behavioral analysis of striatal-related behaviors. *Cntnap2*^{-/-} mice demonstrated increased repetitive behaviors (RRBs), including more grooming bouts, increased marble burying, and increased nose poking in the holeboard assay. MoSeq analysis of behavior further showed signs of altered grooming behaviors and sequencing of behavioral syllables. *Cntnap2*^{-/-} mice also displayed cognitive inflexibility in a four-choice odor-based reversal learning assay. While they performed similarly to WT controls during acquisition and recall phases, they required significantly more trials to learn a new odor-reward association during reversal, consistent with potential deficits in corticostriatal function.

Strengths:

This study provides significant contributions to the field. The finding of altered SPN excitability, the detailed characterization of striatal inhibition, and the comprehensive behavioral analysis are novel and valuable to understanding the pathophysiology of *Cntnap2*^{-/-} mice.

Weaknesses:

(1) The approach based on Thy-ChR2 mice has the advantage of overcoming issues caused by injection efficiency and targeting variability. However, the spread of oEPSC amplitudes across mice shown in panels of Figure 1 G/I is very high with almost one order of magnitude difference between some mice. Given this is one of the most important points of the study it will be important to further analyze and discuss what this variability might be due to. Typically, in acute slice recordings, the within-animal variability is larger than the variability across animals. From the sample sizes reported it seems the authors sampled a large number of animals, but with a relatively low number of neurons per animal (per condition). Could this be one of the reasons for this variability?


We agree with the reviewer that the variability in these experiments is quite large. We have replicated these experiments in the process of performing AMPA:NMDA ratio recordings (see above response to Reviewer 1's comment). We again find no differences in AMPAR-mediated EPSC amplitude between WT and mutant SPNs (Author response image 2). Notably, these experiments also demonstrate a large amount of variability. In the original dataset, a small number of cells were collected from each animal (~1-3 cells/mouse). However, the variability remains in the new dataset, in which more cells were collected from each animal (~4-6 cells/mouse). We find both withinanimal and between-animal variability, as can be seen in Author response image 2 (recordings made from the same animal are color-coordinated). Potential sources of variability in this experiment include: 1) variable expression of ChR2 per mouse, 2) variable innervation of ChR2-expressing terminals onto any given recorded cell, and/or 3) differences in prior plasticity state between cells (i.e. some neurons may have recently undergone corticostriatal LTP or LTD).

Author response image 2.

Optically-evoked AMPAR EPSCs exhibit within- and between-animal variability.

(A) Quantification of EPSC amplitude evoked in dSPNs at different light intensities from the original dataset, plotted by cell (line represents the mean, dots/squares represent average EPSC amplitude for each recorded cell). *Cntnap2*^{+/+} n=17 cells from 8 mice, *Cntnap2*^{-/-} n=13

cells from 5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 56) = 0.3879, geno F (1, 28) = 0.8098, stim F (1.047, 29.32) = 76.56. (B) Quantification of EPSC amplitude evoked in dSPNs, averaged by mouse (line represents the mean, dots/squares represent average EPSC amplitude for each mouse). *Cntnap2*^{+/+} n=8 mice, *Cntnap2*^{-/-} n=5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 22) = 0.2154, geno F (1, 11) = 0.2585, stim F (1.053, 11.58) = 49.68. (C) Quantification of EPSC amplitude in dSPNs from the revision dataset, plotted by cell (line represents the mean, dots/squares represent average EPSC amplitude for each recorded cell). *Cntnap2*^{+/+} n=22 cells from 5 mice, *Cntnap2*^{-/-} n=22 cells from 5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 84) = 0.01885, geno F (1, 42) = 0.002732, stim F (1.863, 78.26) = 20.93. (D) Quantification of EPSC amplitude in dSPNs from the revision dataset, averaged by mouse (line represents the mean, dots/squares represent average EPSC amplitude for each mouse). *Cntnap2*^{+/+} n=5 mice, *Cntnap2*^{-/-} n=5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 16) = 0.06288, geno F (1, 8) = 0.006548, stim F (1.585, 12.68) = 16.97. (E) Quantification of EPSC amplitude evoked in iSPNs from the original dataset, plotted by cell (line represents the mean, dots/squares represent average EPSC amplitude for each recorded cell). *Cntnap2*^{+/+} n=13 cells from 6 mice, *Cntnap2*^{-/-} n=11 cells from 5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 44) = 0.9414, geno F (1, 22) = 1.333, stim F (1.099, 24.18) = 52.26. (F) Quantification of EPSC amplitude evoked in iSPNs from original dataset, averaged by mouse (line represents the mean, dots/squares represent average EPSC amplitude for each mouse). *Cntnap2*^{+/+} n=6 mice, *Cntnap2*^{-/-} n=5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 18) = 0.4428, geno F (1, 9) = 0.5635, stim F (1.095, 9.851) = 23.82. (G) Quantification of EPSC amplitude evoked in iSPNs from the revision dataset, plotted by cell (line represents the mean, dots/squares represent average EPSC amplitude for each recorded cell). *Cntnap2*^{+/+} n=21 cells from 5 mice, *Cntnap2*^{-/-} n=21 cells from 5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 80) = 0.04134, geno F (1, 40) = 0.007025, stim F (1.208, 48.31) = 102.9. (H) Quantification of EPSC amplitude evoked in iSPNs from the revision dataset, averaged by mouse (line represents the mean, dots/squares represent average EPSC amplitude for each mouse). *Cntnap2*^{+/+} n=5 mice, *Cntnap2*^{-/-} n=5 mice. Repeated measures two-way ANOVA p values are shown; $g \times s$ F (2, 16) = 0.001865, geno F (1, 8) = 0.1004, stim F (1.179, 9.433) = 61.31.



(2) This is particularly important because the analysis of corticostriatal evoked APs in panels C and E is performed on pooled data without considering the variability in evoked current amplitudes across animals shown in G and I. Were the neurons in panels C/E recorded from the same mice as shown in G/I? If so, it would be informative to regress AP firing data (say at 20% LED) to the average oEPSC amplitude recorded on those mice at the same light intensity. However, if the low number of neurons recorded per mouse is due to technical limitations, then increasing the sample size of these experiments would strengthen the study.

We appreciate this point; however, the evoked AP experiment and the evoked EPSC experiment were performed on different mice, so it is not possible to correlate the data across experiments. While the evoked AP experiments were performed using potassium-based internal, we used a cesium-based internal to measure AMPAR-mediated EPSCs to more accurately detect synaptic currents. We note that the evoked AP experiments share a similar amount of variability as the evoked EPSC experiments, again possibly owing to variable expression of channelrhodopsin per mouse, variable innervation of ChR2-positive terminals onto individual cells, and/or differences in prior plasticity status between cells.

(3) On a similar note, there is no discussion of why iSPNs also show increased corticostriatal evoked firing in Figure 1E, despite the difference in intrinsic excitability shown in Figure 3. This suggests other potential mechanisms that might underlie altered corticostriatal responses. Given the role of Caspr2 in clustering K channels in axons, altered presynaptic function or excitability could also contribute to this phenotype, but potential changes in PPR have not been explored in this study.

We have now performed more rigorous statistics on the data in Fig. 1 (repeated measures two-way ANOVA) such that the difference in corticostriatal evoked firing in *Cntnap2*^{-/-} iSPNs no longer reaches statistical significance. This is consistent with the modest but statistically non-significant effect of *Cntnap2* loss on iSPN intrinsic excitability. We agree with the reviewer that presynaptic alterations could potentially contribute to the changes in cortically-driven action potentials, especially as this experiment was performed without any synaptic blockers present, and *Cntnap2* is deleted from all cells. That said, if changes in presynaptic release probability accounted for the increased corticostriatal drive, we would expect to see differences in cortically-evoked EPSCs onto SPNs.

While we can't rule out the possibility of pre-synaptic changes, a straightforward explanation for our findings is that loss or alteration of Kv1.2 channel function is responsible for the increased excitability of *Cntnap2*^{-/-} dSPNs, resulting in enhanced spiking in response to cortical input. Given the fact that Kv1.2 channels appear less important for regulating iSPN excitability (see new Fig. 4 and Supp. Fig. 6), this can explain the greater impact of *Cntnap2* loss on dSPN physiology.

(4) Male and female SPNs have different intrinsic properties but the number and/or balance of M/F mice used for each experiment is not reported.

We agree that this is an important consideration. Author response table 1 provides the sex breakdown for the intrinsic excitability experiments. While we did not explicitly power the experiments to test for sex differences, Author response image 3 shows the data separated by sex and genotype for the intrinsic excitability experiments. Within genotype, we find no significant differences between males and females, except for *Cntnap2*^{-/-} iSPNs which showed a significant interaction between sex and current step (Author response image 3F). Interestingly, while present in both sexes, the excitability shift of *Cntnap2*^{-/-} dSPNs may be

slightly more pronounced in females compared to males (Author response image 3C and D). However, this result would require further validation with a greater sample size.

Author response table 1.


Numbers of male and female mice used for the intrinsic excitability experiments.

Cell Type	Genotype	Sex	Cells (Animals)
dSPN	<i>Cntnap2</i> ^{+/+}	Male	12 (4)
		Female	8 (4)
	<i>Cntnap2</i> ^{-/-}	Male	12 (4)
		Female	11 (4)
iSPN	<i>Cntnap2</i> ^{+/+}	Male	10 (4)
		Female	12 (4)
	<i>Cntnap2</i> ^{-/-}	Male	12 (4)
		Female	9 (4)

Author response image 3.

Enhanced excitability of *Cntnap2*^{-/-} dSPNs is present in both males and females. (A) Quantification (mean ± SEM) of the number of APs evoked in dSPNs in *Cntnap2*^{+/+} males and females at different current step amplitudes. *Cntnap2*^{+/+} males n=12 cells from 4 mice, *Cntnap2*^{+/+} females n=8 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; s x c F (28, 560) = 0.8992, sex F (1, 20) = 0.3754, current F (1.279, 25.57) = 56.85. (B) Quantification (mean ± SEM) of the number of APs evoked in dSPNs in *Cntnap2*^{-/-} males and females at different current step amplitudes. *Cntnap2*^{-/-} males n=12 cells from 4 mice, *Cntnap2*^{-/-} females n=11 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; s x c F (28, 588) = 0.6752, sex F (1, 21) = 0.04534, current F (2.198, 46.15) = 78.89. (C) Quantification (mean ± SEM) of the number of APs evoked in dSPNs in *Cntnap2*^{+/+} males and *Cntnap2*^{-/-} males at different current step amplitudes. *Cntnap2*^{+/+} males n=12 cells from 4 mice, *Cntnap2*^{-/-} males n=12 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; g x c F (28, 672) = 2.233, geno F (1, 24) = 3.746, current F (1.708, 40.98) = 79.82. (D) Quantification (mean ± SEM) of the number of APs evoked in dSPNs in *Cntnap2*^{+/+} females and *Cntnap2*^{-/-} females at different current step amplitudes. *Cntnap2*^{+/+} females n=8 cells from 4 mice, *Cntnap2*^{-/-} females n=11 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; g x c F (28, 476) = 1.547, geno F (1, 17) = 5.912, current F (1.892, 32.17) = 58.76. (E) Quantification (mean ± SEM) of the number of APs evoked in iSPNs in *Cntnap2*^{+/+} males and females at different current step amplitudes. *Cntnap2*^{+/+} males n=10 cells from 4 mice, *Cntnap2*^{+/+} females n=12 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; s x c F (28, 560) = 1.236, sex F (1, 20) = 1.074, current F (2.217, 44.34) = 179.6. (F) Quantification (mean ± SEM) of the number of APs evoked in iSPNs in *Cntnap2*^{-/-} males and females at different current step amplitudes. *Cntnap2*^{-/-} males n=12 cells from 4 mice, *Cntnap2*^{-/-} females n=9 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; s x c F (28, 532) = 2.513, sex F (1, 19) = 2.639, current F (1.858, 35.31) = 152.5. (G) Quantification (mean ± SEM) of the number of APs evoked in iSPNs in *Cntnap2*^{+/+} males and *Cntnap2*^{-/-} males at different current step amplitudes. *Cntnap2*^{+/+} males n=10 cells from 4

mice, *Cntnap2*^{+/+} females n=12 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; s x c F (28, 560) = 1.236, sex F (1, 20) = 1.074, current F (2.217, 44.34) = 179.6. (F) Quantification (mean ± SEM) of the number of APs evoked in iSPNs in *Cntnap2*^{-/-} males and females at different current step amplitudes. *Cntnap2*^{-/-} males n=12 cells from 4 mice, *Cntnap2*^{-/-} females n=9 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; s x c F (28, 532) = 2.513, sex F (1, 19) = 2.639, current F (1.858, 35.31) = 152.5. (G) Quantification (mean ± SEM) of the number of APs evoked in iSPNs in *Cntnap2*^{+/+} males and *Cntnap2*^{-/-} males at different current step amplitudes. *Cntnap2*^{+/+} males n=10 cells from 4 mice, *Cntnap2*^{-/-} males n=12 cells from 4 mice. Repeated measures twoway ANOVA p values are shown; g x c F (28, 560) = 0.4723, geno F (1, 20) = 0.5675, current F (2.423, 48.47) = 301.7. (H) Quantification (mean ± SEM) of the number of APs evoked in iSPNs in *Cntnap2*^{+/+} females and *Cntnap2*^{-/-} females at different current step amplitudes. *Cntnap2*^{+/+} females n=12 cells from 4 mice, *Cntnap2*^{-/-} females n=9 cells from 4 mice. Repeated measures two-way ANOVA p values are shown; g x c F (28, 532) = 1.655, geno F (1, 19) = 0.2322, current F (2.081, 39.55) = 99.45.



(5) There is no mention of how membrane resistance was calculated, and no I/V plots are shown.

Passive properties were calculated from the average of five -5 mV, 100 ms long test pulse steps applied at the beginning of every experiment. Membrane resistance was calculated from the double exponential curve fit. This has now been added to the methods section.

(6) It would be interesting to see which behavior transitions most contribute to the decrease in entropy. Are these caused by repeated or perseverative grooming bouts? Or is this inflexibility also observed across other behaviors? The transition map in Figure S5 shows the overall number of syllables and transitions but not their sequence during behavior. Can this be analyzed by calculating the ratio of individual $u_i \times p_{i,j} \times \log_2 p_{i,j}$ factors across genotypes?

We thank the reviewer for raising an insightful question. Here we use a finite state Markov chain model to describe the syllable transitions in animal behavior. To quantify the randomness in the system, we calculated the entropy of the Markov chain (see methods section). The reviewer suggested calculating the partial entropy of the transition matrix, which would allow us to estimate the contribution of a subset of states to the entropy of the whole system, given by the equation:

$$\text{Entropy}(S) = - \sum_{i \in S} u_i \sum_{j \in S} p_{i,j} \log \log(p_{i,j})$$


The partial equation can indeed quantify the stochasticity, or “flexibility” in our context, of the sub-system containing only a subset of the behavior syllables. However, there are two main limitations to this approach:

- (1) The partial entropy fails to account for the transitions connecting the subset with the rest of the states in the system
- (2) The stationary distribution may not reflect the actual probabilities in the isolated sub-system S.

Consequently, the partial entropy cannot be directly interpreted as the fraction of contributions from specific syllable pairs or sub-system to the entropy of the whole system. To be more specific, while a significant difference between the same sub-system in WT and KO groups could indicate that the sub-system contributes significantly to the difference of overall entropy, a non-significant result does not mean that the sub-system does not contribute to overall entropy difference, as interactions between the sub-system and other notconsidered states are not accounted for.

Author response image 4.

Grooming syllables contribute to some but not all differences in syllable transitions in *Cntnap2^{-/-}* mice. We calculated the entropy of each syllable pair using $u_i \times p_{i,j} \times \log_2 p_{i,j}$ for every syllable pair and every animal. We then statistically tested the difference between genotypes for each syllable pair using Mann-Whitney tests. This plot displays those adjusted p-values for each syllable pair between WT and KO groups. The significant p-values suggest that the transitions to syllables 24 and 25 are different between genotypes (note that these correspond to grooming syllables, see Fig. 5N). However, since the overall entropy is a summation of every pair, it is difficult to conclude that syllables 24 and 25 are the sole contributors to the different entropy we observed.



Reviewer #3 (Public Review):

Summary:

The authors analyzed Cntnap2 KO mice to determine whether loss of the ASD risk gene CNTNAP2 alters the dorsal striatum's function.

Strengths:

The results demonstrate that loss of Cntnap2 results in increased excitability of striatal projection neurons (SPNs) and altered striatal-dependent behaviors, such as repetitive, inflexible behaviors. Unlike other brain areas and cell types, synaptic inputs onto SPNs were normal in Cntnap2 KO mice. The experiments are well designed, and the results support the authors' conclusions.

Weaknesses:

The mechanism underlying SPN hyperexcitability was not explored, and it is unclear whether this cellular phenotype alone can account for the behavioral alterations in Cntnap2 KO mice. No clear explanation emerges for the variable phenotype in different brain areas and cell types.

We agree that identifying the mechanism by which *Cntnap2* loss affects intrinsic excitability is interesting and important. We have added experiments to address this and conclude that the improper clustering, number, or function of Kv1.2 channels in *Cntnap2*^{-/-} dSPNs is likely responsible for their increased excitability. These channels are known to be clustered/organized in part by Caspr2 (PMIDs: 10624965, 12963709, 29300891), and Kv1.2 channels are known to play an important role in regulating excitability in SPNs (PMIDs: 13679409, 32075716). In the case of dSPNs, blocking these channels with α -DTX significantly increased the excitability of WT cells (as has been previously reported); however, this effect was occluded in mutant cells, perhaps owing to a decreased contribution of Kv1.2 channels to excitability in *Cntnap2*^{-/-} dSPNs. In addition, we found that blockade of these channels with α -DTX only modestly affected the excitability of iSPNs. Therefore, this can explain why loss of *Cntnap2* more strongly affects the excitability of dSPNs. Please see new Fig. 4, Supp. Fig. 5 and Supp. Fig. 6 for these new data.

We agree with the reviewer that we have not identified a causative relationship between the change in dSPN excitability and the behavioral alterations in *Cntnap2*^{-/-} mice. This is a limitation of the study.

It is interesting to speculate on the root of the varying impacts to excitability that occur across different brain regions and cell types in *Cntnap2*^{-/-} mice. Increased excitability, as we see in dSPNs, has been identified in cerebellar Purkinje cells and L2/3 pyramidal neurons in somatosensory cortex in the context of *Cntnap2* loss (PMIDs: 34593517, 30679017, 36793543). However, other cell types in *Cntnap2*^{-/-} mice have exhibited no change in excitability (mPFC, L2/3 pyramidal neurons, PMID: 31141683) or hypoexcitability (subset of L5/6 pyramidal neurons, PMID: 29112191). While all of these cell types express Kv1.2 channels, they fundamentally vary in their intrinsic properties, owing to the role that other ion channels play in membrane excitability. As a result, loss of *Cntnap2* is expected to have a variable effect on excitability depending on the cell type and the complement of other ion channels that are present. In addition, an initial change in excitability may drive secondary, potentially compensatory, changes in other channels that lead to a different excitability state. These changes are also expected to be cell type-specific. We do note that both of the cell types that show increased excitability in the context of *Cntnap2* loss have been shown to exhibit an α -DTX-sensitive Kv1 channel current, such that application of α -DTX results in increased firing of these cells (cerebellar Purkinje cells; PMIDs: 17087603, 16210348 and L2/3 pyramidal neurons in somatosensory cortex; PMID: 17215507). These findings are consistent with our results in *Cntnap2*^{-/-} dSPNs.


Reviewer #1 (Recommendations For The Authors):

More thorough analysis of some of the manually quantified behaviors would be helpful. For example, only the grooming bout number was presented- what about the duration of bouts and total time grooming? Similarly, for the open field the number of center entries was reported but what about the total time in the center?

We have quantified the time spent grooming and total time spent in the center during the open field test from our original data (Author response image 5). These data were not originally included in the manuscript because they were recorded for only a subset of the total animals. For each of these measures we find trend level changes, which are consistent with the primary measures reported in the main manuscript.

Author response image 5.

Time in center and time spent grooming trend towards an increase in *Cntnap2*^{-/-} mice. (A) Quantification (mean $\pm$ SEM) of total time spent in the center of the open field during a 60 minute test, $p=0.0656$, Mann-Whitney test. (B) Time spent grooming during the first 20 minutes of the open field test, $p=0.0611$, Mann-Whitney test. For both measurements, *Cntnap2*^{+/+} $n=18$ mice, *Cntnap2*^{-/-} $n=19$ mice.



Reviewer #3 (Recommendations For The Authors):

What accounts for the hyperexcitability observed in Cntnap2-deficient SPNs? The authors noted that excitability is reportedly increased, reduced, or unchanged in different brain areas. What accounts for this disparity? Is it about the subcellular localization of Kv1 channels? The authors may want to test this possibility experimentally. At least, they may want to test whether Kv1 channels are mislocalized in SPNs.

We agree that this is an important point, and we have performed additional experiments to address this. We find that the Kv1.2 blocker a-DTX significantly increases the excitability of WT dSPNs but not *Cntnap2*^{-/-} dSPNs. This suggests that the mechanism underlying dSPN hyperexcitability in *Cntnap2* mutants is the improper clustering, number, or function of Kv1.2 channels. These channels are known to be clustered and organized in part by Caspr2 (PMIDs: 10624965, 12963709, 29300891) and have been shown to play an important role in regulating the excitability of SPNs (PMIDs: 13679409, 32075716). Interestingly, we find that a-DTX has less of an effect on the excitability of iSPNs, which may account for the greater impact of *Cntnap2* loss on dSPNs. Please see new Fig. 4, Supp. Fig. 5 and Supp. Fig. 6 for these added data and analyses.

Please see above response to Reviewer #3 for our speculation on the variable impact of *Cntnap2* loss on different cell types and brain regions.

We agree with the reviewer that assessing potential differences in subcellular localization of Kv1 channels in our model would bolster the conclusion that these channels are mislocalized in the *Cntnap2*^{-/-} striatum. We piloted these experiments using immunohistochemistry to stain for Kv1.1 and 1.2 but found that without very high-resolution imaging, it would be challenging to accurately quantify Kv1 puncta in a cell type-specific manner. We instead chose to investigate the functional contribution of Kv1 channels to the dSPN hyperexcitability phenotype through the a-DTX experiments outlined above. a-DTX strongly inhibits Kv1.2 channels, but also Kv1.1 channels to some extent (PMIDs: 12042352, 13679409). We find that the effects of a-DTX on SPN excitability are occluded in *Cntnap2*^{-/-} dSPNs; therefore, we conclude that Kv1.2 (and possibly Kv1.1) channels have reduced function in these cells. Further work will be needed to determine if this is a result of channel mislocalization or another type of alteration.

The authors did not detect synaptic changes in Cntnap-deficient SPNs. This important observation should be briefly discussed in the context of previous work in other brain

regions and cell types. For example, some studies reported structural and functional changes at excitatory synapses. The variable impact on synapses suggests distinct compensatory mechanisms in different brain areas.

Given the prior literature showing effects of *Cntnap2* loss on synapses in other brain regions, we were surprised that striatal synapses were not impacted in our model. We agree with the reviewer that the variable changes in synaptic properties across brain regions in *Cntnap2* mutant mice is likely a result of distinct compensatory changes in these regions. Differences may also arise depending on whether the synaptic changes originate from the post-synaptic cell or from pre-synaptic changes. An interesting direction for future studies would be to explore the developmental trajectory of excitability and synaptic changes to determine which may be initial perturbations versus those that are secondary and potentially compensatory.

Line 138: "synaptic excitability". How is this term defined? Consider "synaptic changes" instead.

"Synaptic excitability" was used to mean a change in the number and/or function of glutamate receptors. We have now changed this term to "excitatory synaptic changes."

*Consider a short paragraph to highlight some limitations of this study. For example, it is unclear whether SPN hyperexcitability results from a compensatory change in *Cntnap2* KO mice and whether the behavioral phenotype is solely due to this cellular phenotype. The study focuses on cortical projections onto SPNs, but these cells receive inputs from other brain areas that were not explored. Lastly, no clear explanation emerges for the variable phenotype in different brain areas and cell types.*

We thank the reviewer for this suggestion and have added several paragraphs to the discussion highlighting some limitations of this study.

We hypothesize that the dSPN hyperexcitability in *Cntnap2*^{-/-} mice is a primary change, due to the direct relationship between Caspr2 and Kv1.2 channels. The results of our $\square$ -DTX experiments suggest that the function and/or contribution of these channels to excitability is altered in *Cntnap2*^{-/-} dSPNs. However, it is possible that there are additional changes in dSPNs that occur as a result of *Cntnap2* loss and contribute to the hyperexcitability of these cells. Rather surprisingly, we don't find evidence for altered excitatory (specifically from cortical inputs) or inhibitory synaptic function, suggesting lack of engagement of homeostatic mechanisms at the synaptic level.

We have not yet determined whether there is a causative relationship between the change in dSPN excitability and the behavioral alterations in *Cntnap2*^{-/-} mice. This is a limitation of the current study. In our discussion section, we highlight that the dSPN changes we observe in dorsolateral striatum (DLS) are known to be sufficient to enhance rotarod learning in other mouse models and thus supports a connection between this cellular change and behavior. For the other behaviors we measured, we acknowledge that both DLS and other striatal or extra-striatal brain regions have been implicated in these behaviors, and therefore less of a direct connection can be made.

In terms of the inputs, we focused on cortical inputs given their known role in mediating motor and habit learning (PMID: 15242609, 16237445, 19198605). Notably, corticostriatal synapses have been shown to be altered across a variety of mouse models with mutations in ASD risk genes and therefore may be a point of convergence for disparate genetic insults (PMID: 31758607). We agree that the striatum receives inputs from a variety of brain regions, notably the thalamus, which we did not explore in this study. This would be an interesting area for future studies.

Finally, it is difficult to speculate on the root of the varying impacts to excitability that occur across different brain regions and cell types in *Cntnap2*^{-/-} mice. Please see above response to Reviewer #3 for some speculation on this point in regard to the potential involvement of Kv1.2 in the excitability changes in various *Cntnap2*^{-/-} cell types. To expand upon this, it is known that ASD-associated mutations can have varying impacts on cell function even across similar cell types within a given brain region – we have seen this between dSPNs and iSPNs (this study, PMIDs: 34380034, 39358043), as have other groups studying ASD risk gene mutations in striatum (PMID: 24995986). This differential impact of the same mutation on intrinsic and/or synaptic physiology across cell types has been identified in other brain regions as well (PMID: 22884327, 26601124). Differences in transcriptional programs, protein expression, neuronal morphology, synaptic inputs and plasticity state make up a non-exhaustive set of variables that will impact the physiological function of a neuron, both in terms of the direct but also indirect consequences of an ASD risk gene mutation. To better address this important question, future studies would benefit from a systematic approach to assessing physiological changes in a given ASD mouse model, both across development and across brain regions.

<https://doi.org/10.7554/eLife.100162.2.sa0>