

Investments in photoreceptors compete with investments in optics to determine eye design

Reviewed Preprint

v2 • July 22, 2025

Revised by authors

Reviewed Preprint

v1 • June 13, 2024

Francisco JH Heras, Simon B Laughlin 

Department of Zoology, University of Cambridge, Cambridge, United Kingdom

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This paper makes a **valuable** contribution to our understanding of the tradeoffs in eye design - specifically between improvements in optics and in photoreceptor performance. The authors successfully build a formal theory that enables comparisons across a wide range of species and eye types. One notable example is that how space should be allocated to optics and photoreceptors depends on eye type - with particularly notable differences between compound and simple eyes. The framework introduced to compare different design properties is **convincing** and provides a nice example of how to study tradeoffs in seemingly disparate design properties.

<https://doi.org/10.7554/eLife.96517.2.sa4>

Abstract

When an animal invests space, materials and energy in an eye to meet behavioural needs, the eye's optics and photoreceptor array compete for these resources to improve the eye's performance. To discover how this competition influences eye design, we introduce a new and superior measure of cost, specific volume in $\mu\text{m}^3 \text{sr}^{-1}$, that depends on the dimensions of the eye's components, applies to both optics and photoreceptor array, accounts for space, materials and energy (including photoreceptors' high metabolic rates), and links investments to an eye's performance via optical, physiological and geometrical constraints. Specific volume enables us to construct a performance surface across the morphospace of an eye of given type and total cost by modelling all of its configurations and determining each model's information capacity. We model three eye types, neural superposition and fused-rhabdom apposition compound eyes and a simple (camera type) eye, across a 10^5 -fold range of total cost. Performance surfaces are flat-topped, therefore the optimum configuration lies in a broad high-efficiency zone within which eyes adapted for specific tasks lose <5% of information. This robust region will increase adaptability by reducing loss of function. Comparing optimised models: simple eye information capacity increases as $(\text{total cost})^{0.8}$ and $(\text{total cost})^{0.55}$ in apposition eyes and simple eyes are x10 to x100 more efficient than apposition eyes of the same total cost. In both eye types 30%-80% of total cost is invested in photoreceptor arrays, optimum photoreceptor length increases with total cost and is reduced

by photoreceptor energy consumption. Simple eyes' photoreceptors are much shorter than apposition eyes' and their length more sensitive to energy consumption. We analyse published data that cover the same range of total specific volumes. The apposition eyes of fast-flying diurnal insects follow three trends predicted by our models: photoreceptor arrays are allocated 40% - 80% of total specific volume, spatial resolution and photoreceptor length increase with increasing specific volume, and apposition photoreceptors are much longer than simple. We conclude that photoreceptor costs are considerable and often exceed optical costs. Thus, competition between optics and photoreceptors for resources helps determine eye design, photoreceptor energy cost plays a major role in determining an eye's efficiency and design, and matching investments in optics and photoreceptors to improve efficiency is a design principle. Our new methodology can be developed to view the adaptive radiation of eyes through a cost-benefit lens.

Introduction

The design of eyes has long fascinated biologists with its variations, innovations, numerous adaptations of form to function, and insights into the evolution of organs, both rudimentary and perfectly contrived (Darwin, 1859 [↗](#); Land & Fernald, 1992 [↗](#); Nilsson & Land, 2012 [↗](#); Oakley & Speiser, 2015 [↗](#); Walls, 1942 [↗](#)). We address a question of eye design that has not been considered before, but is familiar to people who construct imaging systems on tight budgets. How should the resources invested in an eye be divided between an eye's two major components, the optical system that forms images and the photoreceptor array that captures images?

The benefits of enlarging the dioptric apparatus and the photoreceptor array are well understood, consequently numerous studies relate measures of eye morphology (radius, width, length, lens diameter, pupil area, and photoreceptor length) to optical, geometrical and physiological constraints on performance to show how morphology adapts an eye to lifestyle and habitat (Barlow, 1952 [↗](#); Kirschfeld, 1976 [↗](#); Labhart & Nilsson, 1995 [↗](#); Land, 1981 [↗](#); Land, 1997 [↗](#); Nilsson & Land, 2012 [↗](#); Niven et al., 2007 [↗](#); Snyder, 1979 [↗](#); K. N. Thomas et al., 2020 [↗](#)). However, the fact that an eye's performance depends on two systems that effectively compete for the resources invested in an eye, the dioptric apparatus and the photoreceptor array, is rarely considered (Kirschfeld, 1976 [↗](#); Kröger & Biehlmaier, 2009 [↗](#)). We investigate how trade-offs between dioptrics and photoreceptor array determine an eye's performance and efficiency by developing a new cost-benefit approach, which uses measures of eye morphology to calculate costs, and relates these measures to optical, geometrical and physiological constraints to calculate benefits.

We consider three basic eye types, two types of apposition compound eye, neural superposition and fused rhabdom, and a simple eye (**Figure 1** [↗](#)). We establish and road test our cost-benefit approach using apposition compound eyes because they offer a unique combination of advantages. The optical and physiological constraints are well documented, their effects on performance are measured and modelled, (Hardie & Postma, 2008 [↗](#); Heras & Laughlin, 2017 [↗](#); Howard et al., 1987 [↗](#); Snyder, 1979 [↗](#); Song et al., 2012 [↗](#); Stavenga, 2004b [↗](#)) and their influence on eye design is well established. For example, the compound eyes of housefly, praying mantis, dragonfly and robber fly have foveas for detecting, locating and catching flying mates or prey. Here the facet lens are enlarged to reduce the diffraction limit to optical resolving power, the ommatidia are more densely packed to increase angular resolution, and photoreceptors are elongated to increase signal to noise ratio by transducing more photons (B. G. Burton & Laughlin, 2003 [↗](#); Hardie, 1985 [↗](#); Labhart & Nilsson, 1995 [↗](#); Rossel, 1979 [↗](#); Wardill et al., 2017 [↗](#)). In addition, there are clear indications that photoreceptors are costly and investments in optics and photoreceptors are matched. Apposition eyes' photoreceptor arrays obviously fill a substantial fraction of eye volume because photoreceptors construct long photoreceptive waveguides (rhabdomeres, rhabdoms), substantial photoreceptor energy costs increase with rhabdomere

length (Niven et al., 2007 [↗](#)), and there is a precise, fine scale covariation of acuity and photoreceptor length, measured in two apposition eyes (Labhart & Nilsson, 1995 [↗](#); Rossel, 1979 [↗](#)) and obvious in many others (e.g. **Figure 1b** [↗](#)).

Our new cost-benefit approach establishes that photoreceptor cost is a major factor in eye design that influences optics by competing for the resources invested in an eye. Consequently matching investments in optics and photoreceptors to improve efficiency is a principle of eye design. This principle can explain why apposition eyes have longer photoreceptors than simple eyes and photoreceptor length increases with spatial acuity in apposition and simple eyes. We suggest that our approach can advance our understanding of visual ecology by costing the benefits of the adaptations observed in eyes (Cronin et al., 2014 [↗](#); Nilsson & Land, 2012 [↗](#)). Moreover, because the allocation of resources to optics and photoreceptors both promotes cost-effectiveness, and depends upon developmental and physiological mechanisms (Casares & McGregor, 2021 [↗](#); Niven & Laughlin, 2008 [↗](#)), our cost-benefit approach may well contribute to our understanding of how and why eyes evolve (Nilsson & Pelger, 1994 [↗](#)).

Symbols used in the Results and the Discussion

symbol	description	units
D	lens diameter	μm
f	lens focal length, in air	μm
f'	focal distance	μm
F	F-number	
d_{rh}	diameter of rhabdom (fused-rhabdom eye) or rhabdomere (NS eye)	μm
$\Delta\phi$	interommatidial or interreceptor angle	rad (Equations) deg (Figures)
R	radius of eye or locally spherical eye region	μm
p	eye parameter	
L	length of rhabdom or rhabdomere, also depth of photoreceptor array	μm
λ	wavelength of light	nm
η_i	refractive index of eye's internal medium	
$\Delta\rho_l$	half width of lens point-spread-function	rad
$\Delta\rho_{rh}$	half-width of rhabdom(ere) acceptance angle	rad
$\Delta\rho_{ph}$	half-width of photoreceptor acceptance angle	rad
V_o	volume of optics	$\mu\text{m}^3 \text{sr}^{-1}$
V_{ph}	volume of photoreceptor array	$\mu\text{m}^3 \text{sr}^{-1}$
V_{tot}	total eye volume	$\mu\text{m}^3 \text{sr}^{-1}$
C_o	cost of optics	$\mu\text{m}^3 \text{sr}^{-1}$
C_{ph}	costs of photoreceptor array	$\mu\text{m}^3 \text{sr}^{-1}$
C_{tot}	total cost of eye	$\mu\text{m}^3 \text{sr}^{-1}$
S_E	photoreceptor energy surcharge	$\mu\text{m}^3 \text{sr}^{-1}$
K_E	photoreceptor energy tariff	$\mu\text{m}^3 \text{sr}^{-1}$ per microvillus
N_{vil}	number of microvilli	
ν	number of microvilli per unit length of rhabdom or rhabdomere	μm^{-1}
ψ	transduction rate	
SNR_{ph}	photoreceptor signal to noise ratio	per unit contrast
H	spatio-temporal information capacity	bits $\text{sr}^{-1} \text{s}^{-1}$




Figure 1.

Eye structure and geometry defines resolution and costs.

a) Fused rhabdom apposition eye, photoreceptors coding a pixel form fused rhabdom and send axons to a single neural module; b) neural superposition (NS) apposition eye, photoreceptor forms its own rhabdomere, photoreceptors with same optical axes code single pixel and send axons to single neural module; c) simple eye, as in a camera each photoreceptor codes a single pixel. d) Gradient of investment in spatial acuity: apposition eye, honeybee drone, *Apis mellifera*. From ventral to dorsal, lens diameter D increases and interommatidial angle $\Delta\phi$ decreases to increase spatial resolution, and rhabdom length, L , increases to increase SNR_{ph} . $10 \mu\text{m}$ thick longitudinal section, (DA) dorsal eye area (VA) ventral area. BM - retina's basement membrane; E - equator separating dorsal and ventral regions. From Menzel et al., 1991 [\[1\]](#), original micrograph, courtesy of Doeke Stavenga. e) Schematic section of locally spherical apposition eye region. Volumes of optics and photoreceptor array are determined by dimensions that constrain the quality of the spatial image coded by photoreceptors: lens diameter D , focal distance f , interommatidial angle $\Delta\phi = D/R$ where R is eye radius, and rhabdom(ere) length L .

Results

We start with the neural superposition (NS) eyes of flies. This type of apposition compound eye provides both the detailed analyses of optical and physiological constraints required to develop a model that relates investments in optics and photoreceptor arrays to performance, and the published data on dioptrics and photoreceptor arrays required to test our model. Moreover, because fly NS eyes vary in size between species while retaining similar structure and function we simplify our study by developing and using a generic fly NS model. We then model and examine fused rhabdom apposition eyes to identify general principles of apposition eye design. Finally we model a simple (camera) eye to demonstrate that changing eye type changes the distribution of costs. In all three types, we consider diurnal eyes operating in daylight when the number of available photons is not limiting. Although this restriction eliminates the many ways in which eyes are designed to operate efficiently at lower light levels (Nilsson & Land, 2012), it makes it easier to establish points of principle by reducing the numbers of variables in our models.

Calculating the benefits and costs of investing in optics

In daylight, when the number of available photons is not limiting, investments in optics buy increased optical contrast and spatial resolving power by reducing optical and geometrical constraints, as follows. An apposition eye forms and captures images with an array of ommatidia (Figure 1). Each ommatidium uses a facet lens of diameter, D , and focal length (in air), f , to focus light onto the entrance aperture of photoreceptive waveguides, a single fused rhabdom or several rhabdomeres (Figure 1) of diameter d_{rh} . Increasing D sharpens a photoreceptor's angular sensitivity by reducing the blur produced by the Airy disk diffraction pattern, whose half-width $\Delta\rho_L = \lambda/D$ radians, where λ is the wavelength of light, taken to be 500 nm. Increasing f and reducing d_{rh} also sharpen angular sensitivity by reducing the angular subtense of the photoreceptor entrance aperture, d_{rh}/f . Sharpening angular sensitivity has two benefits: it increases spatial resolving power and increases image contrast.

We calculate the effects of D , d_{rh} and f on the half-width of a fly photoreceptor's angular sensitivity, $\Delta\rho$, using two formulae, Snyder's simplification, his Convolution of Gaussians (CoG) $\Delta\rho^2 = \Delta\rho_L^2 + \Delta\rho_{rh}^2$, where $\Delta\rho_L = \lambda/D$ and $\Delta\rho_{rh} = d_{rh}/f$; and Stavenga's better approximation, $\Delta\rho = 1.26\Delta\rho_L$, obtained using a comprehensive wave-optics model (WOM) of flies' NS eyes operating in bright light (see Methods for more details). For CoG we fix $d_{rh} = 1.9 \mu\text{m}$ because this value is typical of fly R1-6 (Hardie, 1985). CoG provides a lower bound to optical resolving power and WOM a more efficient upper bound that, in bright light, makes better use of investments in optics by operating closer to the diffraction limit.

The benefits of investing in optics come at a cost: increasing D and lengthening f to reduce $\Delta\rho$ expands the volume of the dioptric apparatus, V_o , which in apposition eyes is easily calculated from eye geometry (Figure 1e). D and $\Delta\Phi$ define the radius $R = D/\Delta\Phi$ of a locally spherical eye region, in which the dioptric apparatus (lens, cone and screening pigment) is a shell of thickness f' , the focal distance from lens front surface to focal plane. Thus

$$V_o = \frac{1}{3}[R^3 - (R - f')^3] \mu\text{m}^3 \text{sr}^{-1} \quad (1)$$

Because V_o is a specific measure, volume per unit solid angle of visual space, we can compare investments in eye regions that differ in radius and angular extent. Thus V_o handles the regional variations in R observed in many compound eyes.

To simplify our expression for V_o (Equation 1) we fix lens F-number, $F = f/D$ and apply the formula for image formation by a convex cornea, $f' = f \cdot n_i$ where n_i is the refractive index of the internal medium (Nilsson & Land, 2012). This formula describes several apposition eyes, including fly NS eyes (Rossel, 1979; Stavenga, 2003a; Stavenga et al., 1990; Zeil, 1983). Thus

$$V_o = \frac{1}{3}[R^3 - (R - F n_i D)^3] \mu\text{m}^3 \text{sr}^{-1} \quad (2)$$

We use values measured in blowfly, $F = 2.0$ (Stavenga, 2003a) and $n_i = 1.34$, (Seitz, 1968).

Fixing F both simplifies our models by reducing the number of free parameters and accounts for the dependence of the photon flux entering an ommatidium on D by making image brightness (the flux per unit area of rhabdomere entrance aperture) independent of D . We use the specific volume of optics, V_o , as our measure of the cost of optics, $C_o \mu\text{m}^3 \text{sr}^{-1}$. This usage assumes that the cost of materials, the cost of metabolic energy for maintaining function, and the energy cost of carriage all increase in equal proportion to volume, in all parts of the dioptric apparatus, namely the corneal lens, cone, and surrounding pigment cells. Given the lack of definitive data on composition and costs, this assumption is a reasonable starting point (Discussion).

Calculating the benefits and costs of investing in photoreceptors

Investing in photoreceptors buys photoreceptor signal to noise ratio, SNR_{ph} , by reducing the effects of an optical constraint, the Poisson statistics of photon absorption (Nilsson & Land, 2012), and at higher light levels a physiological constraint, the saturation of a photoreceptor's transduction units - its light sensitive microvilli (Heras & Laughlin, 2017; Howard et al., 1987; Song et al., 2012). A microvillus contains rhodopsin molecules, the intermediate molecules of the phototransduction cascade and the ion channels they activate, and it uses these to generate a brief pulse of depolarising current, a quantum bump, following the activation of a single rhodopsin molecule by an absorbed photon. This all-or-none response takes time to complete, during which the microvillus cannot respond to the absorption of another photon (Hardie & Postma, 2008; Song & Juusola, 2014). Therefore, as light intensity rises to daylight levels, SNR_{ph} falls below the Poisson limit because a photoreceptor's transduction units saturate (Howard et al., 1987). At any given time a significant fraction of microvilli are failing to respond to the absorption of a photon by rhodopsin because they are already engaged in processing a bump. Consequently the transduction rate ψ no longer increases linearly with the absorption rate, and bump statistics change from Poisson to binomial. In daylight, flies activate a photomechanical response, a longitudinal pupil, to prevent excessive saturation and maintain SNR_{ph} close to the maximum permitted by binomial statistics,

$$SNR_{ph} = \frac{1}{2} \sqrt{N_{vil}} \quad (3)$$

where SNR_{ph} is defined with respect to a signal of unit contrast (Howard et al., 1987). Note that by operating at this upper limit the fly obtains maximum benefit from investing in N_{vil} . Thus the benefit of investing in a longer rhabdomere is an increase in SNR_{ph} in bright light, according to

$$SNR_{ph} = \frac{1}{2} \sqrt{\nu L} \quad (4)$$

where L is rhabdomere length and ν is the number of microvilli per unit length. We assume ν is constant along the rhabdomere and from published results we estimate $\nu = 230 \mu\text{m}^{-1}$ (Methods). We acknowledge that diameter, cross section and taper often vary within and among rhabdomeres. We are fixing ν at a plausible value for a generic model.

The cost of increasing SNR_{ph} by increasing L is easily calculated. In fly NS eyes, L is also the depth of the photoreceptor array because the rhabdomeres of the six largest photoreceptors, R1-6, stretch from the focal plane of the lens to the basement membrane (Hardie, 1985). Thus in a locally spherical eye region (Figure 1) the photoreceptor array's specific volume is

$$V_{ph} = \frac{1}{3}[(R - f')^3 - (R - f' - L)^3] \mu\text{m}^3 \text{sr}^{-1} \quad (5)$$

Substituting $f' = F.n_i.D$

$$V_{ph} = \frac{1}{3}[(R - F.n_i.D)^3 - (R - F.n_i.D - L)^3] \mu\text{m}^3 \text{sr}^{-1} \quad (6)$$

According to eye geometry (Figure 1), the total specific volume of an eye region is

$$V_{tot} = V_o + V_{ph} \mu\text{m}^3 \text{sr}^{-1} \quad (7)$$

Observe that because both V_o (Equations 1 & 2) and V_{ph} (Equations 5 & 6) depend upon R and $R = D/\Delta\Phi$, both V_o and V_{ph} increase with the number of ommatidia per unit solid angle.

For the photoreceptor array we assume that the costs of space, materials and energy for carriage increase in proportion to volume with the same constants of proportionality as optics. Again, in the absence of measurements this assumption is a reasonable starting point (Discussion). However, this assumption breaks down when we consider specific metabolic rates. To enable and power phototransduction, a photoreceptor has an exceptionally high specific metabolic rate (energy consumed per gram, and hence unit volume, per second) (Laughlin et al., 1998; Niven et al., 2007; Pangršič et al., 2005). We account for this extra cost by applying an energy surcharge, S_E . To equate with our other estimates of cost this surcharge has units of $\mu\text{m}^3 \text{sr}^{-1}$, and we assume that it increases in proportion to N_{vil} because N_{vil} determines the magnitude of the photoreceptor's light-gated conductance (Hardie & Postma, 2008; Heras & Laughlin, 2017). We satisfy these two requirements by defining

$$S_E = K_E N_{vil} \mu\text{m}^3 \text{sr}^{-1} \quad (8)$$

where K_E , the photoreceptor energy tariff, converts the energy consumed by a photoreceptor per microvillus into an equivalent volume.

We estimate K_E (Methods) by adopting the method used to estimate the energy cost of weapons carried by flying beetles (Goyens et al., 2015). We divide the energy consumed per microvillus by the animal's mass specific metabolic rate, SMR, to obtain an equivalent body mass which, assuming a density of 1.0, is our volume equivalent, $K_E \mu\text{m}^3$ per microvillus. K_E is at present poorly determined: it depends upon several ecological, physiological and behavioural factors, it will vary among species and among individuals of the same species, and there is barely enough data (Methods). Given this uncertainty, we model a range of values from $K_E = 0$ to $K_E = 0.64$ that covers the range we estimate for blowfly, $K_E = 0.13$ to $K_E = 0.52$ (Methods).

To summarise costs, for the photoreceptor array

$$C_{ph} = V_{ph} + S_E \mu\text{m}^3 \text{sr}^{-1} \quad (9)$$

and for the optics

$$C_o = V_o \mu\text{m}^3 \text{sr}^{-1} \quad (10)$$

Because C_o and C_{ph} have the same units and are specific (per unit solid angle), we can transfer resources between optics and photoreceptors within the constraint of total specific cost

$$C_{tot} = C_o + C_{ph} \text{ } \mu\text{m}^3 \text{ sr}^{-1} \quad (11)$$

Modelling the effects of resource allocation on NS eye performance

When C_{tot} is fixed, allocating a smaller proportion to optics and a larger proportion to photoreceptors trades the benefits of investing in optics for the benefits of investing in photoreceptors (**Figure 2** [↗](#)).

Having used constraints to link investments in dioptrics and photoreceptor array to three major determinants of the quality of the achromatic images captured by fly photoreceptors R1-6, namely $\Delta\rho$, $\Delta\Phi$ and SNR_{ph} , we can now model how trade-offs between optics and photoreceptors change an eye's ability to support vision. Several measures of visual performance depend on $\Delta\rho$, $\Delta\Phi$ and SNR_{ph} . To establish proof of principle we use a general measure that embraces a wide variety of resolvable achromatic image details - spatio-temporal information capacity in bits $\text{sr}^{-1} \text{ s}^{-1}$. Information capacity has proved useful for measuring the performance of compound eyes (de Ruyter van Steveninck & Laughlin, 1996 [↗](#)), discovering design principles (Howard & Snyder, 1983 [↗](#); Snyder et al., 1977 [↗](#); van Hateren, 1992 [↗](#)), and illuminating the evolution of simple eyes (Nilsson & Pelger, 1994 [↗](#)). The measure is particularly relevant for a fly NS eye because coding by second order neurons is adapted to maximise information capacity (Laughlin, 1981 [↗](#); van Hateren, 1992 [↗](#)).

To model the effects of resource allocation on information capacity, we fix C_{tot} and generate pairs of D and L that cover the range of values permitted by eye geometry. Each D , L pair specifies an investment in optics, $C_o(D, L)$, an investment in photoreceptors $C_{ph}(D, L)$ and hence the three determinants of image quality, $\Delta\rho(D, L)$, $\Delta\Phi(D, L)$, and $SNR_{ph}(D, L)$, that specify spatio-temporal information capacity $H(D, L)$ bits $\text{sr}^{-1} \text{ s}^{-1}$.

We calculate $H(D, L)$ (Methods) in the frequency domain (Howard & Snyder, 1983 [↗](#); van Hateren, 1992 [↗](#)). In brief, a 2-D power spectrum typical of natural scenes is low-pass filtered by the photoreceptor angular sensitivity function, $\Delta\rho(D, L)$, and sampled by an hexagonal lattice of photoreceptors specified by $\Delta\Phi(D, L)$. Movement of the retinal image modulates the flux of photons entering a photoreceptor by converting spatial frequencies into temporal frequencies, according to the distribution of angular velocities generated during behaviour. Then during transduction, the photoreceptors low-pass filter these temporal frequencies and add the noise generated by sampling the photon flux stochastically with transduction units (microvilli). We use van Hateren's (1992) formulae for the $1/f^2$ power spectrum of natural scenes and the distribution of image velocities appropriate for blowfly, and we temporally low pass filter according to the measured properties of a fully light-adapted blowfly photoreceptor R1-6 (Methods). We also modify van Hateren's method to take account of the spatial aliasing that occurs when an array of ommatidia undersamples an image (Methods).

By calculating information capacities for each D , L pair we define the performance surface $H(D, L)$ that covers the morphospace of model eyes of given Fnumber, F , and specific cost, C_{tot} . The performance surface is a flat-topped ridge with steep flanks (**Figure 3** [↗](#)). Atop the ridge, a single point of maximum capacity, $H(D_{opt}, L_{opt})$, sits in an extensive zone within which capacity, and hence efficiency, is $> 95\%$ maximum (red zone in **Figure 3** [↗](#)). With efficiency dropping steeply away from the ridge, approximately two thirds of D , L combinations have efficiencies $< 70\%$. The shape of the performance surface depends upon photoreceptor energy cost (**Figure 3** [↗](#)). Increasing the energy tariff K_E reduces L_{opt} from the maximum permitted by eye geometry to one quarter of maximum. However, because the ridge is flattopped, L_{opt} still lies in an extensive high-




Figure 2.

The effect of trading investment in photoreceptor array for investment in dioptric apparatus.

Schematic shows two eye regions of equal specific volume, the left investing more heavily in photoreceptors, the right more heavily on optics. The images they capture show that transferring resources from photoreceptor array to optics increases image sharpness and contrast at the expense of increasing noise.

efficiency zone, within which L can be changed three-fold and D by 40%, while maintaining $> 95\%$ efficiency (**Figure 3**). This broad high-efficiency zone is bad news for investigators who set out to establish that eyes are optimised, but good news for insects. An eye region can be adapted for a specific purpose, e.g. by increasing L to increase SNR , without sacrificing more than 5% of a measure of general-purpose vision, information capacity. Thus this robust region of design space can facilitate evolution by reducing loss of function.

Patterns of investment in optimised NS eye models

We next establish how, in theory, eye structure and investments in optics and photoreceptor array vary with total investment when efficiency is optimised by maximising information capacity. We run our fly NS eye model with different combinations of C_{tot} and K_E and for each combination find the values of L_{opt} , D_{opt} and $\Delta\Phi_{opt}$ that maximise information capacity and the corresponding investments in optics and photoreceptor array, C_o and C_{ph} . For each combination of C_{tot} and K_E we run two versions of our NS model, one calculates $\Delta\rho$ using CoG (Snyder, 1979) and the other WOM, which produces a narrower $\Delta\rho$ that comes closer to the lens diffraction limit (Stavenga, 2004a). The two versions give similar patterns of investment (**Figure 4**).

Rhabdomere length is matched to acuity and rhabdomeres are long

As C_{tot} increases, D_{opt} widens to increase lens contrast transfer and resolving power by reducing $\Delta\rho$, $\Delta\Phi_{opt}$ narrows to increase spatial resolution, and L_{opt} lengthens to improve SNR_{ph} , reaching several hundred microns in larger eyes (**Figure 4a,b,c**). Thus efficient resource allocation can, in theory, explain why NS eye rhabdomeres are long and rhabdomere length increases with spatial acuity. Note that improving the performance of the optical system by approaching the lens diffraction limit to resolving power improves an eye's performance in full daylight. With the WOM approximation for $\Delta\rho$ the lens diameter D required to achieve a given $\Delta\rho$ is decreased, and this allows $\Delta\Phi$ to decrease (**Figure 4b**), thereby increasing the spatial resolution achieved by an efficient eye of given total cost.

Photoreceptor energy cost changes the structure of an efficient eye

Increasing the photoreceptor energy tariff K_E has little effect on the optimum values of D and $\Delta\Phi$, but it reduces the optimum value of L , and hence the depth of the photoreceptor array, by as much as 90% (**Figure 4c**). The optimum value of L is most sensitive to K_E in the range 0.04 to 0.16, and tends jump up and down as C_{tot} increases. Careful interrogation of our MATLAB code suggests that coding and rounding errors are not responsible for these jumps. We attribute the jumps to interactions between competing non-linear terms (e.g. the dependence on contrast signal on D , noise on $L^{0.5}$, information on $\log(SNR)$, effective spatial bandwidth on $R^{0.5}$, volume on R^3) that shift the optimum across a flat-topped performance surface. This explanation is supported by the observation that these large jumps in optimum L have very little effect on the optimum information rate (**Figure 4d**). The effects of K_E on eye structure are also sensitive to optical performance: increasing K_E has a greater effect on L_{opt} in the more sharply focussed WOM model at all but the highest values of C_{tot} (**Figure 4c**). As expected, increasing K_E reduces the spatial resolution achieved at a given C_{tot} by reducing the resources available for investment in the volume costs V_o and V_{ph} .

Photoreceptor arrays are allocated more resources than optics

When our NS eye models are optimised for information capacity, more than 50% of C_{tot} is allocated to photoreceptor arrays over the range $C_{tot} = 2 \times 10^6$ to $3 \times 10^{11} \mu m^3 sr^{-1}$, irrespective of optical effectiveness (CoG c.f. WOM) and photoreceptor energy tariff, K_E (**Figure 4e**). At lower




Figure 3.

The performance surface $H(D, L)$ describes how information capacity changes across all geometrically permissible configurations of an eye of fixed cost - the eye's morphospace.

Red area: high-performance zone in which capacity is $> 95\%$ maximum. Surfaces plotted for model fly neural superposition (NS) eyes, with F-number, $F = 2$ and total cost, $C_{tot} = 4 \times 10^9 \mu m^3 sr^{-1}$ at 3 values of photoreceptor energy tariff K_E . Acceptance angle $\Delta\rho$ calculated using Snyder's (1979) CoG approximation.




Figure 4.

When model fly NS eyes are optimised for information capacity, eye structure, eye performance and the division of resources between optics and photoreceptor array depend on total investment, C_{tot} and photoreceptor energy tariff, K_E .

a) lens diameter D ; b) interommatidial angle, $\Delta\phi$; c) rhabdomere length, L . Note that sudden jump in the curve for CoG $K_E = 0.12$. d) optimum information capacity H and rhabdomere length L at CoG $K_E = 0.12$ plotted over the range of C_{tot} within which L jumps upward. The eight-fold increase in L has very little effect on the rate of increase of H . e) % C_{tot} allocated to photoreceptor array; f) photoreceptor energy cost as % C_{tot} . CoG, acceptance angle approximated by convolving Gaussians (Snyder, 1979 [\[1\]](#)); WOM, acceptance angle approximated according to wave optics model (Stavenga, 2004a [\[2\]](#)). Key in b) also applies to a).

values of K_E , the optimum allocation to photoreceptor arrays is strongly dependent on C_{tot} , increasing from 45% to 90% C_{tot} over the range $C_{tot} = 1 \times 10^6 \mu\text{m}^3 \text{sr}^{-1}$ to $1 \times 10^{10} \mu\text{m}^3 \text{sr}^{-1}$, and then jumping down to approximately 60% between $C_{tot} = 1 \times 10^{10} \mu\text{m}^3 \text{sr}^{-1}$ and $C_{tot} = 1 \times 10^{11} \mu\text{m}^3 \text{sr}^{-1}$ (Figure 4e). At higher values of K_E , the optimum photoreceptor allocation is relatively insensitive to C_{tot} and lies between 60% C_{tot} and 65% C_{tot} . We conclude that, in theory, the cost of the photoreceptor array is as significant as the cost of optics. Therefore to fully understand eye design both costs must be taken into account.

The relative contributions of photoreceptor array volume, V_{ph} , and photoreceptor energy cost, S_E , to total array cost, C_{ph} , depends on total investment, C_{tot} (Figure 4f). At our lowest C_{tot} , i.e. in the lowest acuity eyes regions, the energy cost is 25% to 50% of C_{tot} , depending on K_E , but as eye size increases the costs associated with array volume come to dominate and photoreceptor energy cost drops to 5% of C_{tot} .

Flies invest efficiently in costly photoreceptor arrays

Our new method for calculating costs enables us to estimate, for the first time, the total investment an animal makes in an eye or eye region, and break this down into its two major components, optics and photoreceptor array. Using published measurements of D , $\Delta\Phi$ and L we determine the specific volumes of dioptric apparatus, V_o , and photoreceptor array, V_{ph} in fly neural superposition (NS) eyes. Although numerous publications give values of D and $\Delta\Phi$ (Feller et al., 2021; Land, 1997), very few include L . We have values for just 12 eye regions, taken from 7 species of flies (Table 2). For each region we insert the values, L , D , $\Delta\Phi$ and f' , into Equations 1 and 5 to calculate V_o , V_{ph} , and V_{tot} . Where focal distance f' is not reported (Table 2), we follow our models (Equation 2) and calculate $f' = F D n_i$ with $F = 2$ and $n_i = 1.34$.

The 12 eye regions provide a well-distributed set of empirical values of L , D , $\Delta\Phi$, V_o , V_{ph} and V_{tot} that spans almost five orders of magnitude of V_{tot} , from $1.4 \times 10^6 \mu\text{m}^3 \text{sr}^{-1}$ and $2.1 \times 10^6 \mu\text{m}^3 \text{sr}^{-1}$ for the low acuity eye region of the March fly *Dilophus febrilis* and the eye of *Drosophila melanogaster* respectively, to $9 \times 10^{10} \mu\text{m}^3 \text{sr}^{-1}$ for the remarkable fovea of the small robber fly *Holcocephala fusca* (Wardill et al., 2017).

We compare the empirical values obtained from flies with theoretical curves, obtained from model NS eyes that are optimised for information capacity (Figure 5). Note that for our theoretical curves we plot D , $\Delta\Phi$, L and V_{ph} against total specific cost, $C_{tot} \mu\text{m}^3 \text{sr}^{-1}$ (Figure 4). This total includes the surcharge for photoreceptor energy consumption $S_E \mu\text{m}^3 \text{sr}^{-1}$, which increases in proportion to the number of microvilli and the photoreceptor energy tariff, K_E (Equation 8). However, we cannot estimate C_{tot} in flies because their values of K_E are not well-determined (Methods). Although K_E will vary considerably between species, we only have enough published data to calculate its order of magnitude and likely range (Methods). Furthermore, K_E is species-specific and, because we have no more than two data points per species and we are using a generic model that does not account for differences in d_{rh} and F among species (Table 2), we cannot estimate K_E by fitting our model to the empirical data. For these reasons we compare theoretical findings to empirical data by plotting them against a measure of eye size that is both calculated by our models and estimated from data, total specific volume, V_{tot} . Because K_E is uncertain, we plot theoretical curves calculated with values of K_E from within our likely range.

Looking across 12 fly eye regions, D increases and $\Delta\Phi$ decreases as V_{tot} increases. D lies within, or close to, the theoretical envelope defined by CoG models optimised with $K_E = 0$ and $K_E = 0.32$ (Figure 5a), and tends to slightly exceed optimum values at lower specific volumes. $\Delta\Phi$ is approximately 25% wider than optimum (Figure 5b) in the four eye regions with lowest specific volume and close to optimum elsewhere. The eye parameter, $p = D\Delta\Phi$, (Figure 5c) indicates the degree to which an apposition compound eye sacrifices spatial resolution by sampling below the lens diffraction limit. Only *Holcocephala* fovea, $p = 0.37$, comes close to the diffraction limit $p = 0.29$ (Snyder, 1979). As observed in a number of diurnal apposition eyes

species	eye region	D μm	$\Delta\phi$ deg	L μm	f μm	f' μm	F	source
<i>Calliphora vicina</i>	male acute zone	37	1.07	340	74	99	2	1,2,3,4
<i>Calliphora vicina</i>	female acute zone	29	1.28	280	58	78	2	2
<i>Musca domestica</i>	male love spot	36	1.75	250	72	96	2	2,5
<i>Musca domestica</i>	male peripheral	20	3.5	140	40	54	2	2
<i>Drosophila melanogaster</i>	average	16.5	5	83	20	27	1.25	1,2,6
<i>Coenesia attenuata</i>	female frontal	20	2.3	140	25	28	1.25	7
<i>Bibio marci</i>	male dorsal	33	1.6	230	70	94	2.1	8
<i>Bibio marci</i>	male ventral and female	21	3.7	110	36	48	1.7	8
<i>Dilophus febrilis</i>	male dorsal	24	2.2	180	40	54	1.7	8
<i>Dilophus febrilis</i>	male ventral and female	15	5.1	60	16	21	1.1	8
<i>Holcocephala fusca</i>	acute zone, max. acuity	75	0.28	230	160	176	2.53	9
<i>Holcocephala fusca</i>	peripheral	21	3.5	123	33	45	1.6	9

Table 1

Measurements of Dipteran neural superposition (NS) eyes, extracted from published sources.

1. Land 1997 [\[1\]](#); 2. Hardie 1985 [\[2\]](#); 3. Stavenga et al. 1990 [\[3\]](#); 4. Stavenga 2003a [\[4\]](#); 5. Land and Eckert 1985 [\[5\]](#); 6. Stavenga 2003b [\[6\]](#); 7. Gonzalez-Bellido et al. 2011 [\[7\]](#); 8. Zeil 1983 [\[8\]](#); 9. Wardill et al. 2017 [\[9\]](#). Further details of the measurements used and results obtained are given in supplementary table S1.

species	eye region	D μm	$\Delta\phi$ deg	L μm	f μm	f' μm	F	source
<i>Tenodora australasiae</i>	eye coordinates z=15;x=20 (fovea)	47.5	0.64	500	205	445	8	1
	z=15;x=30	51.5	0.8	535	335	405	6.5	1
	z=15;x=10	41	1	415	305	375	7.4	1
	z=15;x=0	35	1.1	285	205	275	5.9	1
	z=15;x=60	42	1.6	390	150	212	3.6	1
	z=15;x=100	33	2.5	330	110	153	3.3	1
<i>Sympetrum spp.</i>	ventral -60 deg	31	1.85	461	124	166	4	2
	ventral -5 deg	27	1.75	461	108	148		2
	dorso-ventral border	31	1.69	516	124	167		2
	dorsal +50	53	1.82	627	92	123		2
	dorsal +70	61.5	1.78	886	246	330	4	2
	dorsal fovea	71	0.35	1107	305	410	4.4	2
<i>Apis mellifera</i> drone	dorsal eye	40	1	500	205	275	5.1	3
	top ventral	28	2.1	400	145	194	5.1	3
	mid ventral	25	2.8	280	82	110	3.3	3
	basal ventral	18	4.4	150	54	72	3	3
<i>Apis mellifera</i> worker	max. acuity	25	1.6	350	100	134	5.6	4,5

Table 2

Measurements of fused-rhabdom apposition eyes, extracted from published sources.

1. Rossel 1979 [link](#); 2. Labhart and Nilsson, 1995 [link](#); 3. Menzel et al 1991 [link](#); 4. Varela and Wiitanen, 1970 [link](#); 5. Kelber and Somanathan, 2019 [link](#). Details of measurements are given in supplementary table S1.




Figure 5.

Parameters defining the configurations of 12 fly NS eye regions, taken from 7 species (Table 2), compared to model NS eyes optimised for information capacity at given values of photoreceptor energy tariff K_E , and total specific volume V_{tot} .

a) lens diameter D , b) interommatidial angle $\Delta\phi$ and c) eye parameter $p = D\Delta\phi$ (keys as in a), d) rhabdome length L . e) Log-log plot of L vs D , dashed line shows slope for isomorphic scaling, $L \propto D$. f) Specific volume of photoreceptor array, V_{ph} expressed as % total specific volume of eye, $\%V_{tot}$. Models use Snyder's CoG approximation for photoreceptor acceptance angle (Snyder, 1979).

(Wehner, 1981 [↗](#)), the remaining 11 fly eye regions in our data set undersample by a factor of 2 or more, with p decreasing with increasing V_{tot} from 1.4 to 0.65 (**Figure 5c** [↗](#)). Our optimised models also undersample, with p decreasing with increasing V_{tot} from 0.89 to 0.65 (**Figure 5c** [↗](#)). Note that like D and $\Delta\Phi$, p is relatively insensitive to K_E . We conclude that in most eye regions p is not optimised for spatio-temporal information capacity, at least according to our model, because it is higher than predicted at all but the highest values of V_{tot} .

Rhabdomere length, L , increases with V_{tot} from 60 μm to 340 μm (**Figure 5d** [↗](#); **Table 2** [↗](#)), within or close to the theoretical envelope defined by $K_E = 0$ and $K_E = 0.16$. There is a robust trend: the empirical values of $\log L$ increase with $\log V_{tot}$, and hence with increasing D and decreasing $\Delta\Phi$ (**Figure 5a,b** [↗](#)), along a trajectory that resembles theoretical curves for optimum $\log L$. However, the data do not follow the theoretical curve for any particular value of K_E , and there is some scatter among the data points (**Figure 5d** [↗](#)). We have three reasons to attribute most of this scatter to unknown differences in K_E . First, the optimum value of L is sensitive to K_E (**Figure 4c** [↗](#)); second, K_E will vary between species (Methods); and third, the empirical values of D and $\Delta\Phi$, which are in theory relatively insensitive to K_E , are less scattered and lie closer to our theoretical predictions (**Figure 5a,b** [↗](#)). The one obvious outlier, the shorter than predicted rhabdomeres in *Holcocephala* fovea (**Figure 5d** [↗](#)), is easily explained: the optics and photoreceptor array of this forward pointing fovea, together with underlying optic lobe, are tightly packed into a head that is less than 600 μm from front to back ([Wardill et al., 2017](#) [↗](#)), and this lack of headroom limits the depth of the eye, $(L + f)$.

The log-log plot of L versus D (**Figure 5e** [↗](#)) shows that the increase in L with V_{tot} is not explained by proportional scaling because the empirical values do not lie on a straight line of slope 1, $L = kD$. As expected of a physiological system subject to several competing constraints ([G. K. Taylor & Thomas, 2014](#) [↗](#)), the theoretical curve continuously changes slope, and this is reflected in the data. Thus the matching of L to D and $\Delta\Phi$ observed in the data is not the consequence of a simple developmental constraint, it is the product of developmental mechanisms that allocate resources efficiently to optics and photoreceptor array.

We conclude that the eye regions we analyse are, according to our generic model, not optimised for spatio-temporal information capacity because the data points that are least sensitive to the photoreceptor energy tariff K_E , namely D , $\Delta\Phi$ and p , are generally larger than predicted. Nonetheless, we argue on two grounds that the matching of rhabdomere length to spatial resolution seen in flies promotes the efficiency of their total investments in eye. L lies within, or close to, the theoretical envelope predicted by models optimised for efficiency, and because the theoretical performance surface has a wide flat high-efficiency zone (**Figure 3** [↗](#)), large deviations from optimum have relatively little impact on efficiency.

To accommodate photoreceptors' lengthy rhabdomeres, flies allocate between 56% and 78% of eye volume, V_{tot} , to photoreceptor arrays (**Figure 5f** [↗](#)). The smaller remainder, between 44% and 22%, is allocated to optics. There is a slight trend for the percentage allocated to photoreceptors to increase with V_{tot} . With the exception of *Holcocephala*'s fovea, where the depth of the photoreceptor array is much less than predicted for optimum allocation, the volumes allocated to photoreceptor arrays lie within or close to the theoretical envelope defined by $K_E = 0$ and $K_E = 0.16$. On this basis we conclude that our diurnal flies allocate a larger fraction of eye volume to photoreceptors than to optics, and this allocation promotes the efficient use of resources. Thus photoreceptor costs are playing a major role in determining the design and efficiency of flies' NS eyes.

Fused rhabdom apposition eyes show similar patterns of investment

We investigate diurnal apposition eyes with fused rhabdoms by modifying our NS eye model and comparing theoretical results with empirical data using the procedures developed for NS eyes. The modifications are straightforward. We increase lens F-number to 5.5 and reduce d_{rh} to 1.8 μm . These values are in the middle of the wide range found in our sample of apposition eye regions (Table 3). We also adjust the relationship between L and the number of microvilli N_{vil} to account for a fused rhabdom (Methods). For simplicity we assume that all photoreceptors contribute equal numbers of microvilli along the full length of the rhabdom so that N_{vil} increases in proportion to the depth of the photoreceptor array.

We then compare the models' predictions of the optimum values of L , D , $\Delta\Phi$, V_o , V_{ph} and V_{tot} with empirical values extracted from published measurements of D , $\Delta\Phi$, L and f' (Table 3). The necessary data is only available from studies of four species: the mantid *Tenodera australis* (Rossel, 1979), the honeybee *Apis mellifera* (Kelber & Somanathan, 2019; Menzel et al., 1991; Varela & Wiitanen, 1970), and two species of the Libellulid dragonfly *Sympetrum* that are so similar that their data are lumped together (Labhart & Nilsson, 1995). The studies provide empirical values for 16 eye regions, which span a range of V_{tot} , from $4.4 \times 10^7 \mu\text{m}^3 \text{sr}^{-1}$ for *Apis* worker ventral eye to $1.8 \times 10^{11} \mu\text{m}^3 \text{sr}^{-1}$ for *Sympetrum* dorsal fovea (Figure 6).

Looking across our limited sample of fused rhabdom apposition eyes, D increases with V_{tot} , $\Delta\Phi$ decreases, and L increases from 280 μm to 1100 μm (Figure 6a,b,c). These empirical values are more widely scattered than those for NS eyes (Figure 5a,b,d), and the discrepancies between empirical and predicted values are larger. We expect a wider scatter because our sample of fused rhabdom eyes is more diverse than our sample of NS eyes, both structurally and phylogenetically. In addition, the scatter is dominated by regional differences within eyes (Figure 6), several of which adapt a region for a specific task. For example, the 1100 μm rhabdomeres of the dragonfly *Sympetrum*, which are the longest photoreceptive waveguides found in any eye, adapt the fovea for detecting small prey against a bright blue sky by enhancing SNR_{ph} (Labhart & Nilsson, 1995). Despite the scatter in data points, fused rhabdom eyes follow the trends seen in NS eyes (Figure 6 c.f. Figure 5). In most eye regions, D and $\Delta\Phi$ are larger than predicted by models optimised for efficiency (Figure 6a,b), so therefore is the eye parameter p (values not plotted). At low V_{tot} , L lies within the theoretical envelope defined by models run with plausible values of photoreceptor energy tariff, K_E . However, at higher specific volumes L falls increasingly short (Figure 6c). As with flies' NS eyes, large investments are made in photoreceptor arrays. The volume fractions of photoreceptor arrays increase from 33% V_{tot} to 75% V_{tot} as V_{tot} increases, and the majority of values lie between 40% V_{tot} and 60% V_{tot} (Figure 6d). Our information maximisation models predict an increase from 33% V_{tot} to 85% V_{tot} .

Our models also give the relative contribution of photoreceptor cost to total cost when fused rhabdom apposition eyes are optimised for spatial information capacity. The percentage of total investment devoted to the photoreceptor array increases with the energy tariff K_E and climbs steadily with increasing C_{tot} , from 43% to more than 80% (Figure 6e). As in NS eyes, the contribution of photoreceptor energy consumption to array cost is progressively marginalised by array volume cost as C_{tot} increases, falling to less than 20% in the largest eyes (Figure 6f).

We conclude that the 16 eye regions in our sample of fused rhabdom apposition eyes are not optimised for spatio-temporal information capacity, at least according to our models. However, the regions exhibit three trends that are associated with efficient resource allocation - rhabdomeres are relatively long, length increases with acuity, and the photoreceptor array occupies a large

Table 3


Dependence of energy surcharge K_E on time spent flying T_F and hours of daylight D_L , calculated for blowfly.

T_F hrs	D_L hrs	K_E $\mu\text{m}^3/\text{microvillus}$
12	12	0.13
12	16	0.15
2	12	0.44
2	16	0.52

Figure 6.

Theoretical results from apposition eye models optimised for information capacity compared to empirical data from 16 apposition eye regions taken from 3 species (Table 3).

a) D vs specific volume V_{tot} ; b) $\Delta\Phi$ vs V_{tot} ; c) L vs V_{tot} ; d) photoreceptor array specific volume V_{ph} as % V_{tot} vs V_{tot} . (e & f) Models demonstrate impact of photoreceptor costs and its dependence on total cost, C_{tot} . e) Photoreceptor cost, C_{ph} , as % C_{tot} ; f) Photoreceptor energy cost as % C_{ph} . Models run with F-number $F = 5.5$, using COG approximation for acceptance angle (Snyder, 1979) and values of energy tariff K_E given in keys.



fraction of eye volume. These trends suggest that, as in NS eyes, resources are being divided between optics and photoreceptor array to promote efficiency, consequently photoreceptor costs play a major role in the design of fused-rhabdom apposition eyes.

Patterns of efficient investment are different in simple eyes

To further investigate how investment patterns differ according to eye type, we construct a basic hemispherical model of a simple eye with rhabdomeric photoreceptors (Methods; [Figure 7a](#)). We then follow the procedures described above to see how optimum patterns of investment depend on total investment, C_{tot} , and photoreceptor energy tariff, K_E . We consider the same 5 orders of magnitude of total investment, $C_{tot} = 1 \times 10^6$ to $2.5 \times 10^{11} \mu\text{m}^3 \text{sr}^{-1}$, and for simple eye models this encompasses a 100 fold range of lens diameters, $D = 29 \mu\text{m}$ to 2.9 mm , and focal distances, $f' = 79 \mu\text{m}$ to 7.9 mm .

For each C_{tot} there is an optimum division of resources between optics and photoreceptors that maximises information capacity. L_{opt} increases with V_{tot} , scaling close to $(C_{tot})^{1/3}$ ([Figure 7b](#)). Increasing the energy tariff, K_E , greatly reduces L_{opt} without changing this scaling exponent. At any given C_{tot} and K_E , L_{opt} is much shorter in simple eyes and the difference increases with C_{tot} from at least 50% shorter at $C_{tot} = 1 \times 10^6 \mu\text{m}^3 \text{sr}^{-1}$ to over 90% shorter at $C_{tot} = 1 \times 10^9 \mu\text{m}^3 \text{sr}^{-1}$. We conclude that, in theory, efficient resource allocation explains why photoreceptors in diurnal simple eyes have much shorter light-sensitive waveguides than photoreceptors in diurnal apposition eyes.

Patterns of investment differ from apposition eyes in other respects ([Figure 7c](#)). In simple models the optimum volume fraction of the photoreceptor array is insensitive to C_{tot} whereas in apposition models it increases with C_{tot} . In addition, a simple eye is much more sensitive to photoreceptor energy consumption. The small energy tariff, $K_E = 0.04$, reduces simple eye array volume from 35% V_{tot} to 12% V_{tot} , but has a negligible effect on an NS eye. Raising the tariff to $K_E = 0.32$, reduces the simple eye's array volume to just 3% V_{tot} but in apposition models it is always >30% V_{tot} . Despite this ten-fold reduction in % V_{tot} the total resources allocated to photoreceptor array increases from 35% to 57% ([Figure 7d](#)) because the relative contribution of photoreceptor energy consumption increases from 40% C_{tot} to 55% C_{tot} . Finally, in simple eyes the relative contribution of photoreceptor energy consumption is insensitive to C_{tot} , but in apposition models it falls steadily with increasing C_{tot} to around 5% C_{tot} ([Figure 7d](#)). This difference between eye types can be attributed to differences in eye geometry ([Figure 7a](#)). As C_{tot} increases, the simple eye maintains its dense packing of photoreceptors but an apposition eye's photoreceptors become increasingly widely spaced as larger lenses force the boundaries between ommatidia further apart. We conclude that when resources are efficiently allocated, the differences in geometry that define eye type can profoundly influence both the impact of photoreceptor energy consumption on eye morphology and the distribution of costs within an eye.

Information capacity and investment

In all optimised eye models, apposition and simple, the spatio-temporal information capacity, H_{opt} , increases sub-linearly with total investment, C_{tot} ([Figure 8a](#)), making a bit of information more expensive in a larger eye. In simple eyes, H_{opt} increases as $(C_{tot})^{0.8}$ across the full range of C_{tot} while in apposition models the exponent is lower, decreasing from 0.6 to 0.5 as C_{tot} increases. Thus the simple eye model is 10 times more efficient when $C_{tot} = 1 \times 10^6 \mu\text{m}^3 \text{sr}^{-1}$ and approximately 100 times more efficient when $C_{tot} = 1 \times 10^{11} \mu\text{m}^3 \text{sr}^{-1}$ ([Figure 8b](#)). Increasing the photoreceptor energy tariff reduces the efficiency of all eye types, but this effect declines as apposition eye models increase in size ([Figure 8b](#)) because, for reasons of geometry, volume costs marginalise photoreceptor energy cost.




Figure 7.

Resource allocation in model simple eyes optimised for information capacity compared to optimised model neural superposition apposition eyes (NS).

a) Schematic showing apposition eye and simple eye with identical spatial resolution, as defined by lens diameter D , focal distance f , rhabdom(ere) diameter d_{rh} , and rhabdom(ere) length, L (after Kirschfeld, 1976). Note the denser packing of photoreceptors inside the simple eye. (b - e) Properties of simple and NS eye models optimised for information capacity in full daylight. b) Photoreceptor length, L vs specific volume V_{tot} ; c) photoreceptor specific volume V_{ph} expressed as $\% V_{tot}$ vs V_{tot} ; d) Photoreceptor investment C_{ph} as $\% C_{tot}$ vs C_{tot} ; e) photoreceptor energy cost as $\% C_{tot}$ vs C_{tot} . Models use Snyder's (1979) CoG approximation for acceptance angle.




Figure 8.

Information rate, H v.s. total cost C_{tot} for simple and apposition model eyes optimised to maximise information capacity in full daylight.

a) Rates are higher in simple eyes and more sensitive to photoreceptor energy tariff, K_E b) Efficiency, bits per unit specific volume per second, falls less steeply with increasing C_{tot} in simple eyes. Models use Snyder's CoG approximation (1979) for acceptance angle.

Discussion

We initiate a new approach to understanding eye design by investigating how the division of resources between an eye's dioptric apparatus and photoreceptor array influences an eye's performance and efficiency. To achieve this we formulate a new measure of cost, specific volume, that is defined by an eye's shape and the dimensions of its components, can be adjusted to account for the high energy cost of photoreceptors, and relates investments in optics and investments in photoreceptors to three basic determinants of image quality, interreceptor angle $\Delta\Phi$, photoreceptor acceptance angle $\Delta\rho$, and photoreceptor signal to noise ratio SNR_{ph} , via optical, geometrical and physiological constraints. These advantages allow us to model how an eye's performance changes when resources are transferred between optics and photoreceptors. Thus we can, for the first time, model performance across the morphospace of an eye of given type and total cost. Using a general-purpose measure of performance, spatio-temporal information capacity (Howard & Snyder, 1983 [\[1\]](#); van Hateren, 1992 [\[2\]](#)), we find that the surface for an apposition compound eye (**Figure 3** [\[3\]](#)) is flat topped - the optimum configuration that maximises performance sits in a broad high efficiency zone in which performance over 95% of maximum. Within this robust region of morphospace an eye, or eye region, can be specialised for a specific task with little loss of information, and this can facilitate evolution by reducing loss of function. Our models predict that efficiently configured diurnal apposition eyes invest heavily in deep photoreceptor arrays with long rhabdomeres and rhabdoms, and match investments in optics and photoreceptor arrays so that rhabdom(ere) length increases with spatial resolution (**Figure 4** [\[4\]](#)). By applying our new method of costing investments to published data, we show that fast-flying diurnal insects conform to these trends and are, therefore, configured to be efficient (**Figures 5** [\[5\]](#), **6** [\[6\]](#)). Our models also demonstrate that photoreceptor energy consumption is an important factor in an apposition eye's design and performance: it reduces photoreceptor length by as much as 90% and information capacity by as much as 60%. To see how resource allocation and performance depend on eye type we model a simple (camera type) eye (**Figure 7** [\[7\]](#)). When optimised for efficiency, a simple eye has much shorter photoreceptors and its structure and function are more sensitive to photoreceptor energy demands: photoreceptors length is greatly reduced by and information capacity decreases by as much as 90% (**Figures 7** [\[7\]](#), **8** [\[8\]](#)). We conclude that when apposition and simple eyes are designed to use resources efficiently, the cost of the photoreceptor array is as significant as the cost of optics, and matching investments in optics and photoreceptor arrays to increase efficiency is a principle of eye design. Our analysis, which is the first to account for the costs of both optics and photoreceptor array, indicates that one benefit of investing in an eye, the ability to gather more information follows the law of diminishing returns. As expected from its superior geometry (Kirschfeld, 1976 [\[9\]](#)), a simple eye is in theory a better investment: its information capacity increases as (total cost)^{0.8}, compared with (total cost)^{0.55} for an apposition eye, and the simple eye is 10 to 100 times more efficient at gathering information (**Figure 8** [\[8\]](#)). We now discuss the validity of our methodology, the robustness of our findings, and their contributions to our understanding of eye design and evolution.

Specific volume is a useful measure of cost

This new measure of cost has several advantages. First, specific volume is defined by dimensions and angles that are measured anatomically and optically (**Equations 1** [\[1\]](#), **2**, **5**, **6**) and relates directly to benefits via optical, geometrical and physiological constraints on spatial acuity and sensitivity. Thus, like previous indicators of cost - eye radius, eye length, pupil area and corneal area (Brandon et al., 2015 [\[10\]](#); Brandon & Dudycha, 2014 [\[11\]](#); Snyder et al., 1977 [\[12\]](#); K. N. Thomas et al., 2020 [\[13\]](#); R. J. Thomas et al., 2004 [\[14\]](#)) - specific volume can evaluate investments and trade-offs that adapt eyes for specific purposes. Second, specific volume is more flexible and far reaching than previous indicators of cost because it can be applied to eyes that are not well approximated by solid spheres. Third, it applies to small eye regions. Fourth, it provides a common currency for

comparing investments in different eyes, eye regions, and components within an eye (**Figures 5**, **6**, **7**). This facility enables one to investigate how and why resources are distributed across visual fields and, as we demonstrate here, evaluate trade-offs between investments in an eye's components. Given that biological processes and systems evolve to become more cost-efficient because costs depress fitness (Alexander, 1996; Niven & Laughlin, 2008; G. K. Taylor & Thomas, 2014), we suggest a fifth advantage: specific volume will help us understand how eyes are adapted and evolve.

How well does specific volume represent costs? Volume accurately measures one limiting resource, space, but because there are insufficient data, we have to make assumptions to calculate the costs of energy and materials. Assuming that the densities of an eye's cells, extracellular materials and fluid-filled cavities are close to 1, space is a reasonable proxy for three costs that increase in proportion to mass: the energy expended carrying an eye (carriage cost), materials, and cellular energy consumption (Witter & Cuthill, 1993).

Equating mass and carriage cost is a reasonable starting point. Experiments on individual animals - a human walker carrying a backpack (Bastien et al., 2005), a flightless ant carrying nectar (Duncan & Lighton, 1994), and a flying beetle carrying a heavy weapon (Goyens et al., 2015) - show that the cost of carrying an additional load is the load's mass times the individual's mass specific metabolic rate when moving unloaded.

Turning to the costs of the materials and energy used within the eye, there is no data for an apposition eye's dioptric apparatus. Nonetheless, its corneal lenses, pigment cells, and cones cells are densely packed with macromolecules, and because their optical function is passive (they refract and absorb) they contain very few, if any, mitochondria. Indeed, the corneal cuticle and the fluid filled pseudo-cone of fly NS eye are extracellular (Hardie, 1985). Thus, the assumption that the costs of materials and energy for the dioptric apparatus increase in fixed proportion to volume is a reasonable starting point.

Cost estimates can be adjusted for situations in which costs per unit volume are not equal, as illustrated by our treatment of photoreceptor energy consumption. To support transduction the photoreceptor array has an exceptionally high metabolic rate (Laughlin et al., 1998; Niven et al., 2007; Pangršič et al., 2005). We account for this higher energy cost by using the animal's specific metabolic rate (power per unit mass and hence power per unit volume) to convert an array's power consumption into an equivalent volume (Methods). Photoreceptor ion pumps are the major consumers of energy and the smaller contribution of pigmented glia (Coles, 1989) is included in our calculation of the energy tariff K_E (Methods). The higher costs of materials and their turnover in the photoreceptor array can be added to the energy tariff K_E but given the magnitude of the light-gated current (Laughlin et al., 1998) the relative increase will be very small. Thus, for our intents and purposes the effects of these additional costs are covered by our models. For want of sufficient data, K_E is uncertain, therefore we model a reasonable range of values (Methods). Because the optimum eye configuration is most sensitive to K_E in the lower half of this range (**Figures 4**, **7**), we are unlikely to seriously underestimate the effects of photoreceptor costs on eye morphology and efficiency.

Matching photoreceptor length to spatial acuity

The systematic increase in rhabdomere or rhabdom length, L , with spatial acuity observed in apposition eyes (**Figures 1**, **5**, **6**) and simple eyes (see below), and predicted by our models (**Figures 4**, **7**), is an obvious manifestation of efficient resource allocation. Two studies of apposition eyes, one of dragonfly and the other of praying mantis (Labhart & Nilsson, 1995; Rossel, 1979), map the covariation of lens diameter, D , interommatidial angle, $\Delta\Phi$, and rhabdom length, L , across an entire eye. Both show that when acuity is increased by enlarging the dioptric apparatus and increasing ommatidial density, i.e. by reducing $\Delta\Phi$, L is increased to improve SNR_{ph} . In both insects L peaks in a fovea that is used to detect and capture prey, and in dragonfly dorsal

fovea the 1.1 mm rhabdomeres (the longest light sensitive waveguides reported in any eye, diurnal or nocturnal) are adapted to improve the detection and localisation of small flying prey (Labhart & Nilsson, 1995; Rigosi et al., 2017). However, specialisation for prey capture is unlikely to explain why L varies systematically with acuity across an entire eye.

We find that the trend to increase rhabdomere length with acuity is more generally advantageous. The trend is widespread: looking across 28 eye regions drawn from 10 species of fast flying insect, L increases with acuity over a 25-fold range of interommatidial angle (Figures 5, 6). Modelling shows that matching L to acuity improves a general measure of performance, spatial information capacity and the widespread covariation of L and acuity observed in apposition eyes follows the trend predicted by modelling (Figures 5, 6). Finally, because information capacity captures the interplay between three basic determinants of image quality, $\Delta\rho$, $\Delta\Phi$ and SNR_{ph} , several tasks could benefit from an increase in L with acuity.

Other factors can link L to acuity. One possibility, developmental constraints impose inflexible scaling relationships on apposition eyes, can be dismissed. Scaling changes across apposition compound eyes (Perl & Niven, 2016; Scales & Butler, 2016; G. J. Taylor et al., 2019), and our empirical log-log plots of L v.s. eye volume and L v.s. D continuously change slope (Figure 5d,e), as required for the efficient allocation of resources (Figure 4c). Head size is an obvious constraint: smaller insects have smaller heads, consequently they will tend to have smaller eyes with lower acuity and shorter rhabdom(ere)s. The strongest evidence that lack of headroom reduces the length of photoreceptors comes from the fovea of a small insect, *Holcocephala* (Wardill et al., 2017), that is adapted for exceptionally high acuity (Table 2; Figure 5d). One must also consider the costs associated with enlarging the head by adding more eye, such as mechanical support, wind resistance, and conspicuousness. We suggest that more comparative studies are required to establish the influence of head size on eye design.

Then there is the difference in L that is so obvious and widespread that it has been accepted without question - L is generally much longer in diurnal apposition eyes than in diurnal simple eyes. Comparing eyes with rhabdomeric photoreceptors, L ranges from 60 μm to 1100 μm in apposition eyes (Tables 2 & 3), and from 10 μm to 90 μm , in spiders' simple eyes (Land, 1985). Efficient resource allocation provides an explanation: optimised simple eye models predict that L is approximately x10 shorter than in apposition compound eyes of the same cost (Figure 7b). L also increases with acuity (increasing D and decreasing $\Delta\Phi$) in optimised simple eye models and there is evidence that this matching occurs in diurnal spiders. L increases with increasing D when comparing eyes in different species (Land, 1985), different eyes in the same individual, and the same eye increasing in size and acuity as a juvenile spider grows to adulthood (Göte et al., 2019). However, more accurate models of these spiders' eyes are needed prove that this co-variation is increasing efficiency. The degree to which the lengthening of cone outer segments in primate fovea (Provis et al., 2013) increases the efficiency of resource utilisation is also an open question. A critical ingredient, a physiological constraint that links investments in outer segments to noise levels in full daylight, has yet to be identified in ciliary photoreceptors. In addition, the economics of vertebrate simple eyes differs from invertebrate simple eyes in two ways. First, a vertebrate outer segment is almost 10 times more effective at absorbing photons because its folded sheets of membrane pack visual pigment molecules more densely than arrays of tubes (microvilli) (Fain et al., 2010). Second, photoreceptors compete with neural circuits for resources within the eye (Kröger & Biehlmaier, 2009; Sterling & Laughlin, 2015).

Photoreceptor costs are a major factor in eye design

In apposition eye models optimised for spatio-temporal information capacity more than 40% of total investment, C_{tot} , is allocated to photoreceptor arrays (Figures 4e, 6e) and, in line with this theoretical finding, our analysis of data from diurnal insects shows that arrays occupy 30% to 75% of eye volume (Figures 5f, 6d). For simple eye models the efficient allocation is 30% to 60% of C_{tot} (Figure 7d). Our modelling identifies roles for two types of array cost, the rarely considered

costs associated with array volume (Baden & Nilsson, 2022 [↗](#); Kirschfeld, 1976 [↗](#); Kröger & Biehlmaier, 2009 [↗](#)), and the well-known energy costs associated with phototransduction (Ames, 2000 [↗](#); Fain & Sampath, 2021 [↗](#); Laughlin et al., 1998 [↗](#); Okawa et al., 2008 [↗](#); Pangršič et al., 2005). We find that this energy cost greatly influences the configuration of an efficient eye: increasing the energy tariff K_E reduces array depth, L , as much as tenfold (Figures 4c [↗](#), 5d, 6c, 7b). Reducing L reduces SNR_{ph} (Equation 4 [↗](#)), however the SNR_{ph} that remains is worth paying for because the % C_{tot} allocated to photoreceptor energy consumption increases (Figures 4f [↗](#), 7e). We conclude that because SNR_{ph} is valuable, photoreceptor costs are always significant and this necessitates efficient resource allocation. It follows that photoreceptor costs influence the morphologies of both photoreceptor array and dioptric apparatus because optics and array compete for the resources invested in an eye.

The impact of photoreceptor costs depends on eye type

Photoreceptor energy cost has more impact on both the total cost and the design of a simple eye than an equivalent apposition compound eye because the simple eye has more photoreceptors per unit volume (Figure 7a [↗](#)). Applying our lowest energy tariff, $K_E = 0.04 \mu\text{m}^3$ per microvillus, to an optimised apposition eye model has almost no effect on the depth of the photoreceptor array, L , but in the optimised simple eye model it reduces L by more than 80%, and array volume by more than 60% (Figure 7c [↗](#)). Despite this shrinkage, investment in the simple eye model's photoreceptor array increases from 35% C_{tot} to 50% C_{tot} because energy costs have risen from 30% C_{tot} to 55% C_{tot} (Figure 7d [↗](#)). Also, in simple eye models the percentages of C_{tot} allocated to the photoreceptor array volume and energy change very little as C_{tot} increases, whereas in apposition eyes the relative contribution of photoreceptor energy consumption falls steeply as photoreceptors become increasingly widely spaced (Figure 7a [↗](#)). Thus the distribution of costs within an efficiently configured eye changes with eye type, according to the eye's typical geometry

Efficiency depends on eye type and photoreceptor energy cost

In both apposition and simple eye models, efficiency decreases with increasing C_{tot} (Figure 8b [↗](#)), making a bit of information more expensive in a larger eye. The decline in efficiency with increasing eye size is steeper in apposition models, as expected from the well-known scaling relationships enforced by lens diffraction and eye geometry, spatial resolution increases as eye radius, R , in a simple eye and as $\sqrt{R}$ in an apposition eye (Kirschfeld, 1976 [↗](#)). However, using R as a measure of investment fails to identify the contribution of the photoreceptor array and fails to account for the fact that an apposition eye is not a solid sphere. When our more realistic measure of cost, specific volume, is used a *Drosophila* size apposition eye is 10 times less efficient than a simple eye of the same cost and a dragonfly size eye is 100 times less efficient (Figure 8b [↗](#)). In both simple and apposition eyes efficiency is reduced by increasing the photoreceptor energy tariff K_E . This effect is much greater in simple eyes (Figure 8b [↗](#)) thus, as also found for the reduction of photoreceptor length (Figure 7b [↗](#)), K_E has more impact on the design of simple eyes.

Expanding our understanding of apposition eye design

We build on theoretical studies that show how constraints imposed by apposition optics, eye geometry and eye size determine spatial resolution and information capacity when eye radius R , and hence corneal area, represent the limiting resource (Howard & Snyder, 1983 [↗](#); Kirschfeld, 1976 [↗](#); Snyder et al., 1977 [↗](#)). Howard & Snyder (1983) [↗](#) introduced the physiological constraint, transduction unit saturation, and showed that it increases the optimum value of eye parameter, p , in bright light by placing a fixed ceiling on SNR_{ph} that is equivalent to operating at a lower light level. We extend Howard and Snyder's (1983) approach by using a measure of cost, specific volume, that allows eye radius R to vary while total cost remains constant. This added realism allows us to raise and lower the SNR ceiling imposed by transduction units by trading investments in rhabdomere length against investments in dioptric apparatus. In so doing we discover that it is

advantageous for photoreceptors to lengthen rhabdomeres so as to reduce shot noise. Indeed, in daylight *Calliphora vicina* R1-6 achieve the lowest level of photoreceptor noise reported to date, an equivalent contrast of 0.0084 (Anderson & Laughlin, 2000 [↗](#)). This extreme specialisation resembles Autrum's proposition that apposition eye photoreceptors compensate for poor spatial resolution by increasing temporal resolution (Autrum, 1958 [↗](#)), thereby achieving the highest flicker fusion frequencies known, in excess of 400 Hz (Tatler et al., 2000 [↗](#)). Perhaps enhancing photoreceptor performance to compensate for optical inefficiency is a principle of apposition eye design?

Increasing temporal resolution increases photoreceptor energy consumption, so to improve efficiency apposition eye photoreceptors match temporal bandwidth to the properties of the images they form (Laughlin & Weckström, 1993 [↗](#); Niven et al., 2007 [↗](#)). Eye regions that have lower spatial resolution and encounter more slowly moving images save energy by reducing photoreceptor bandwidth (B. G. Burton et al., 2001 [↗](#)). Although this matching makes large energy savings it is not included in our models because, to reduce the burden of establishing proof of principle, we fix photoreceptor bandwidth at the high value measured in a fully light-adapted blowfly. Nonetheless, our models can accommodate the costs and benefits of changing temporal bandwidth because they operate in the frequency domain (Methods). Boosting temporal resolution will be more cost-effective in large apposition eyes because photoreceptor energy costs are marginalised by costs associated with array volume (**Figure 4e,f** [↗](#)), therefore investments of energy that increase temporal bandwidth are increasing the return on the much larger investment in array volume (e.g. carriage cost). Some fast flying diurnal insects with large facet lenses reduce the carriage cost of the photoreceptor array by surrounding each ommatidium with a palisade of air-filled tracheae (Horridge, 1969 [↗](#); Laughlin & McGinness, 1978 [↗](#); Smith & Butler, 1991 [↗](#); Wardill et al., 2017 [↗](#)). The volumes and savings involved can be large. Published micrographs of dragonfly dorsal eye (Horridge, 1969 [↗](#)) suggest that air fills at least 50% of array cross section.

How efficiently do insects allocate resources within their apposition eyes?

In principle one can establish the efficiency of an insect's investments in optics and photoreceptors by computing the performance surface for a model eye region that has the same parameters as the insect's eye region, then observing how closely the insect's eye region approaches the optimum configuration. We cannot do this at present because a critical parameter, the photoreceptor energy tariff K_E , has not been determined: the best we can do is use data from blowflies to estimate K_E 's likely range (Methods). Not knowing K_E is a major impediment to obtaining exact fits of our models to data because our modelling has discovered that the morphology of an efficient eye is sensitive to photoreceptor energy costs: in apposition eyes, increasing K_E across its likely range reduces the depth of the photoreceptor array by as much as 90% (**Figure 4c** [↗](#)). Nor can we use our models to infer K_E by adjusting it to fit the data. Changing K_E has very little effect on D and $\Delta\Phi$, therefore we must fit the model to L , and this requires several data points from eyes of the same species because K_E will be species specific (Methods). In our most consistent data set, the NS eyes (**Figure 5** [↗](#); **Table 2** [↗](#)), it is impossible to obtain a satisfactory fit: the 12 data points come from 7 species and there are never more than 2 points per species. There are several data points per species for fused rhabdom apposition eyes (**Figure 6** [↗](#); **Table 3** [↗](#)), but in each of the four species the data points are too widely scattered to obtain a useful fit. This scatter in values of L is not surprising. Our generic fused rhabdom model glosses over the large differences in F-number and rhabdom diameter seen within the eye of each species (**Table 3** [↗](#)), and these differences probably reflect the specialisation of eye regions for different purposes, e.g. in the fovea the detection, tracking and interception of mates and prey, and in the periphery the monitoring of optic flow and the detection of potential prey and predators (Labhart & Nilsson, 1995 [↗](#); Menzel et al., 1991 [↗](#); Straw et al., 2006 [↗](#); van Hateren et al., 1989 [↗](#)). This proposition is supported by the observation that the empirical data for D and $\Delta\Phi$ (**Figure 6a,b** [↗](#))

are also more widely scattered than the data for the more homogeneous group, fly NS eyes (**Figure 5a,b**). Because our method calculates three basic determinants of the quality of the captured image, $\Delta\Phi$, $\Delta\rho$ and SNR_{ph} , it can be adapted to compute performance surfaces for these specialised tasks, however the necessary theory has not been developed.

Having raised the question, efficient for what, we must examine our assumption that every eye region of a diurnal insect is adapted for vision in full daylight. The fact that insects in our data sets have larger lenses, interommatidial angles and eye parameters than our models predict (**Figure 5**) suggests that their eye regions are adapted for lower light levels (Snyder, 1979). We can think of two mutually inclusive reasons why a diurnal eye will not be adapted for vision in full daylight. The eye might be adapted for the time of day when the behaviour it supports is most profitable, e.g. insects that are most active early and/or late in the day to avoid predators and desiccation, and insects that prey on these crepuscular insects. Second, because an eye's performance surface is flat-topped, an eye region optimized for an intermediate light intensity might, when averaged over the daylight hours, perform better than an eye adapted for bright light. Our models can be developed to investigate how costs and benefits influence the design of eyes adapted for lower light levels (see below), and modified to investigate two further reasons for increasing the eye parameter p , higher image speeds (Snyder, 1979) and increasing D in "bright zones" to improve the detection of small targets and small movements (Straw et al., 2006; van Hateren et al., 1989).

In summary, for want of both data and theory, we are currently unable to quantify the efficiency with which an apposition eye region performs the function for which it is adapted. Nonetheless, we have two reasons to conclude that flying diurnal insects allocate resources to the optics and photoreceptor arrays of their apposition eyes efficiently. First, the empirical values we extract from published data lie within or close to the boundaries defined by our theoretical curves for eyes that maximise a general-purpose measure of performance, information capacity (**Figures 5,6**). Second, the flat-topped performance surfaces have extensive high-efficiency zones within which an eye's morphology can be changed while retaining >95% of the maximum information capacity (**Figure 3**). Thus eyes that are not optimised to maximise information capacity can still be gathering information with high efficiency. This observation speaks to the value of deriving performance surfaces: one can consider competing demands, evaluate loss of function and identify compromises.

Efficient resource allocation is a general principle

Efficient resource allocation is observed at many levels of biological organization, from single molecules to life histories (Bigman & Levy, 2020; Garland et al., 2022; Nijhout & Emlen, 1998; Stearns, 1989). Eyes are no exception. In an apposition eye, optical radius, R , represents corneal surface area per unit solid angle, therefore models that maximise spatial acuity or information capacity at constant R show that corneal area can be allocated to facet lenses to maximise spatial acuity and information capacity by trading spatial resolution for SNR (Snyder, 1977; Snyder et al., 1977). In the POL region of fly NS eye a fixed resource, the number of microvilli in the central rhabdom, is allocated to a pair of photoreceptors, R7 and R8, to increase the efficiency of polarisation coding (Heras & Laughlin, 2017), and in primate eye retinal area is allocated to cone spectral types to increase coding efficiency (Garrigan et al., 2010; Zhang et al., 2022).

Matching the sampling density of the photoreceptor array to lens resolving power (Nilsson & Land, 2012) also allocates the resources invested in photoreceptors more efficiently by avoiding wasteful overcapacity, according to the principle of symmorphosis (Piersma & van Gils, 2011; C. Taylor & Weibel, 1981; Weibel, 2000). However, unlike the matching of L to $\Delta\Phi$ that allocates resources efficiently to optics and photoreceptors, the matching of capacities does not consider the individual cost-benefit functions of a system's components. When these are taken into account, the

allocation of investment that maximises efficiency can instal over-capacity in components with more favourable cost-benefit ratios, as demonstrated for bones in racehorse forelegs (Alexander, 1997) and chains of signalling molecules (Schreiber et al., 2002).

Understanding the adaptive radiation of eyes

The method we introduce here can be developed to view several aspects of eye adaptation and evolution through a cost-benefit lens because it is flexible. Our approach is not wedded to information capacity: it supports other measures of visual performance because it relates investments to fundamental determinants of image quality. Moreover, the method can be applied to several trade-offs that influence eye design by relaxing the simplifying assumptions we make here. To give a concrete example, the trade-offs between acuity and sensitivity that adapt eyes for nocturnal or diurnal vision can be viewed through a cost-benefit lens by allowing F-number and rhabdom(ere) diameter to vary, and by adopting a method that calculates the mean and variance of photon transduction rate on N_{vil} in rhabdomeres of any given length and variable cross section, at any biologically plausible light intensity (Heras & Laughlin, 2017).

One can also relax the restrictions we place on eye geometry. Our approach does not depend on spherical geometry, it depends on being able to relate the shapes, sizes and constituents of an eye's components to their costs and performance. Advances in scanning, imaging and digital reconstruction offer opportunities to determine shapes, volumes and constituents directly, and advances in modelling optical and physiological processes are extending our ability to relate these properties to performance (Kim, 2014; Song et al., 2012; Sumner-Rooney et al., 2019; G. J. Taylor et al., 2019). These powerful new methods can take us closer to the goal of understanding of how costs, benefits and innovative mechanisms drive eye evolution (Sumner-Rooney, 2018), and our flexible cost-benefit approach can support this endeavour.

Our approach also identifies factors that contribute to selective pressure, and gauges their effects. For example, the pressure to reduce photoreceptor energy cost depends on a photoreceptor's impact on an animal's energy budget. This effect is captured by our photoreceptor energy tariff, K_E , because it relates a photoreceptor's energy consumption to the animal's metabolic rate (Methods). We also find that photoreceptor energy costs are marginalised when apposition eye size increases and this encourages the use of wide bandwidth, high SNR photoreceptors with high metabolic rates. Most interestingly, the derivation of performance surfaces (Figure 3) reveals a large area, the robust high-performance zone within which an eye's morphology can be adapted to a particular task while retaining more than 95% of a measure of general-purpose capability, information capacity. This low "loss of opportunity" cost helps to explain the remarkable degree of fine tuning observed in apposition compound eyes (B. G. Burton et al., 2001; Land, 1997; Nilsson & Land, 2012; Sumner-Rooney et al., 2019; G. J. Taylor et al., 2019), in line with suggestions that robustness increases evolvability (Drack & Betz, 2022).

Two key steps in the early evolution of eyes were the stacking of photoreceptive membranes to absorb more photons, and the formation of optics to intercept more photons and concentrate them according to angle of incidence (Nilsson, 2013, 2021). Our modelling of well-developed image forming eyes shows that to improve performance stacked membranes (rhabdomeres) compete with optics for the resources invested in an eye, and this competition profoundly influences both form and function. Thus it is likely that competition between optics and photoreceptors was shaping eyes as lenses evolved to support low resolution spatial vision. If so, the developmental mechanisms that allocate resources within extant high resolution eyes (Casares & McGregor, 2021), by controlling cell size and shape and, as our study emphasises, setting up gradients of shape and size across the eye, will have analogues or homologues in ancient eyes. Their discovery will not only tell us how Nature constructs perfectly contrived instruments, it will illuminate the evolutionary pathway to perfection.

Methods

Calculating a photoreceptor array's information rate

Our calculations of the rate at which an eye's photoreceptor array codes achromatic information, in bits per steradian per second, take account of five factors. 1) The statistics of the signals presented to an eye under natural conditions; 2) the blurring of the spatial image by the eye's dioptric apparatus; 3) the sampling of the spatial image by the photoreceptor array; 4) the introduction of noise by the stochastic activation of a finite population of transduction units and 5) the temporal smoothing of signals and noise during phototransduction. We calculate information rates in the frequency domain using methods developed for apposition eyes (de Ruyter van Steveninck & Laughlin, 1996 [\[1\]](#); Howard & Snyder, 1983 [\[2\]](#); Snyder et al., 1977 [\[3\]](#); van Hateren, 1992 [\[4\]](#)).

Natural intensity signals

We calculate the spatial and temporal power spectra of naturalistic signals using the model established for blowfly (van Hateren, 1992 [\[4\]](#)). An eye images an ensemble of natural scenes in which power decreases as the square of spatial frequency, as measured and predicted theoretically (G. J. Burton & Moorhead, 1987 [\[5\]](#); Ruderman, 1997 [\[6\]](#); van Hateren, 1992 [\[4\]](#)). Consequently the two dimensional power spectrum of the spatial frequencies viewed by the eye is

$$S_{xy} = \frac{c_c}{f_x^2 + f_y^2} \quad (12)$$

The proportionality constant c_c is chosen to set the total contrast of the image formed by the eye (defined as the square root of the ratio between the power of the signal and the square of the mean level of signal) to an appropriate value. This contrast depends on the eye that observes the scene. For images spatially filtered by a blowfly compound eye the average contrast, calculated locally over intervals of 25-50 degrees, is 0.4 (Laughlin, 1981 [\[7\]](#)). Accordingly we use a proportionality constant that scales the power of frequencies between one cycle per 50 degrees and the upper cut-off frequency dictated by sampling, one cycle per 1.5 degrees, to be $(0.4)^2 = 0.16$.

To add the temporal dimension van Hateren (1992) [\[4\]](#) modelled how the movement of an eye through an ensemble of uniformly distributed objects generates temporal structure. For the case of an eye moving in a straight line, the structure is adequately described by a random distribution of angular speeds

$$a_v(v) = \frac{c_v}{(|v| + \sigma_v)^2} \quad (13)$$

where $a_v(v)$ is the probability of encountering image speed v , c_v is the normalization constant that sets the sum of probabilities over the distribution to 1, and σ_v is a constant that regulates the width of the distribution. Local movements produced by objects moving within a scene, although acknowledged to be biologically significant, have little impact on the statistics of the signal encountered by an eye moving in a straight line (van Hateren, 1992 [\[4\]](#)).

Distributions of this form peak at lower speeds and have a long tail that extends to speeds faster than σ_v . The exact value of σ_v will depend on the behaviour of the animal. For *Calliphora vicina* yaw, pitch and roll movements rarely exceed 3.5 rad s^{-1} between saccades (van Hateren & Schilstra, 1999 [\[8\]](#)). Assuming a similar limit in translational movements, we take σ_v to be 1 rad s^{-1} . In his earlier 1992 paper van Hateren assumed σ_v is 0.29 rad s^{-1} and noted that the information capacities he calculated are not particularly sensitive to σ_v .

The three-dimensional power density of naturalistic signals generated by the aforementioned distributions is

$$S_{xyt}(f_x, f_y, f_t) = \frac{\pi}{2f_r^3} c_c a_v \left(\frac{\pi}{2} \frac{f_t}{f_r} \right) \quad (14)$$

where $f_r = \sqrt{f_x^2 + f_y^2}$ and c_c is the proportionality constant used in Equation 12. We checked that this distribution retains the local contrast value of 0.4 for different values of the parameter $|\sigma_v|$.

Blurring by the optics

The lens point spread function and the angular sensitivity of the photoreceptor entrance aperture combine to determine photoreceptor angular sensitivity, which in the eyes we consider approximates a Gaussian function of half-width $\Delta\rho$. The transfer function of this spatial filter is

$$m(f_x, f_y) = \exp \left(-\frac{\pi^2}{4 \ln 2} (f_x^2 + f_y^2) \Delta\rho^2 \right) \quad (15)$$

The lens is diffraction limited and the photoreceptive rhabdomeres and rhabdoms are waveguides, consequently calculating the acceptance angle of this coupled system is a complicated problems of wave optics, especially when, with narrow d_{rh} , waveguide effects are significant (Stavenga, 2003a, 2003b, 2004a, 2004b). We employ two expressions for photoreceptor acceptance angle. One is Snyder's (1979) convolution of Gaussians (CoG) approximation

$$\Delta\rho^2 = (\lambda/D)^2 + \Delta\rho_r^2 \quad (16)$$

where λ is the wavelength of light, D is lens diameter, λ/D is the half-width of the Airy disk and $\Delta\rho_r$ is the angular acceptance of the rhabdomere, taken to be its angular subtense, $\Delta\rho_r = d_{rh}/f$.

The other is Stavenga's more principled expression, obtained using his wave optics model (WOM) of larger fly NS eyes (Stavenga, 2004a)

$$\Delta\rho = 1.26\lambda/D \quad (17)$$

Stavenga's WOM shows that the coefficient in Equation 17 reduces progressively as the fly's longitudinal pupil closes, reaching a minimum of 1.14 when fully closed in full daylight. We use an intermediate value, 1.26, because the pupil closes progressively over the upper two log units of daylight intensity, while holding SNR_{ph} close to maximum.

Temporal filtering during phototransduction

A photoreceptor's elementary response to a single photon, a quantum bump, is generally fitted with functions of the type

$$h(t) \propto \left(\frac{t}{\tau} \right)^{\alpha-1} e^{-t/\tau} \quad (18)$$

whose Fourier transform is

$$H(f) = \frac{1}{(1 + i2\pi\tau f)^\alpha} \quad (19)$$

Thus transduction's first temporal low pass filter has the form

$$|m_t(f_t)|^2 = \frac{1}{(1 + (2\pi\tau f)^2)^\alpha} \quad (20)$$

We use the fit of $\alpha = 3.12$ measured in *Musca* (B. G. Burton, 2006 [\[1\]](#)) and $\tau = 0.001$ s obtained in *Calliphora* at the highest light intensities (Juusola et al., 1994 [\[2\]](#)). According to these values photon shot noise power falls to half-maximum at around 170 Hz.

Following the activation of a microvillus by an absorbed photon the quantum bump is produced with a variable delay. This latency dispersion low-pass filters signals that are composed of many bumps by widening the impulse response. We use the low-pass filter

$$|m_d(f_t)|^2 = \frac{1}{(1 + (2\pi\tau_d f)^2)^{\alpha_d}} \quad (21)$$

where $\alpha_d = 2$ (Wong et al., 1980 [\[3\]](#)). With $\tau_d = 0.0014$ s this filter produces a signal corner frequency close to the 55 Hz measured in *Calliphora vicina* (Anderson & Laughlin, 2000 [\[4\]](#)). The trends we observe in our study are not critically dependent on the exact values of these bump parameters.

Sampling

For one dimensional receptor arrays, frequencies will not be correctly sampled above the Nyquist limit (Snyder, 1979 [\[5\]](#))

$$f_{Ny} = \frac{1}{2\Delta\phi} \quad (22)$$

where Φ , as defined above, is the sampling angle of the array.

For a two dimensional hexagonal lattice, only frequencies up to $\frac{1}{\sqrt{3}\Delta\phi}$ can be sampled independently of the orientation of the array (Snyder et al., 1977 [\[6\]](#)). These authors also noted that frequencies up to $\frac{1}{\Delta\phi}$ can be sampled for special orientations of the array, however we will assume that the highest spatial frequencies sampled by an array is $\frac{1}{\sqrt{3}\Delta\phi}$.

Deriving the signal coded by the array

To obtain the power spectral density of the signal coded by a photoreceptor the natural image signal derived above is spatially filtered by the dioptric apparatus, converted to a stream of shot events, quantum bumps, with mean rate ψ , and temporally filtered during phototransduction. We start with van Hateren's (1992) expression for the signal coded by a photoreceptor that is transducing photons at a rate ψ

$$(\Delta S)^2(f_x, f_y, f_t) = \psi^2 S_{xyt}(f_x, f_y, f_t) |m(f_x, f_y)|^2 |m_t(f_t)|^2 |m_d(f_t)|^2 \quad (23)$$

This expression assumes that the photoreceptor signal increases linearly with photon capture rate, as happens at low light levels when the saturation of transduction units is negligible, and signal and noise follow Poisson statistics.

In brighter conditions the saturation of microvilli changes the relationship between transduction rate, ψ and photoreceptor signal ΔS (Howard et al., 1987 [\[7\]](#)). The probability that a microvillus generates a signal is given by

$$p = \frac{I}{N_{vil}/\tau_r + I} \quad (24)$$

where I , the effective intensity, is the rate at which microvilli are absorbing photons, and the cycling time τ_r is the average time taken to reset a microvillus to its receptive state, following activation by a photon capture (Howard et al., 1987 [\[3\]](#)).

The transduction rate is given by

$$\psi = pN_{vil}/\tau_r \quad (25)$$

There are some limits to this treatment. Because light intensity decreases along the rhabdom(ere), microvilli in the distal part will, in principle, absorb more photons than microvilli in the proximal part, making it impossible for all microvilli to have the same probability of being active. However, adaptations have been described that partially compensate for this effect by reducing the quantum capture efficiency of distal microvilli (Labhart & Nilsson, 1995 [\[3\]](#)).

According to the saturation equation (24), the light intensity that activates half the microvilli, $I_{50} = N_{vil}/\tau_r$. Therefore the dependence of transduction rate on I is given by

$$\psi = pN_{vil}/\tau_r = \frac{I_{50}I}{I_{50} + I} \quad (26)$$

and

$$\frac{d\psi}{dI} = \frac{I_{50}^2}{(I_{50} + I)^2} \quad (27)$$

$\Delta\psi$, the change in ψ produced by a change in intensity, ΔI , is given by

$$\Delta\psi = \frac{d\psi}{dI} \Delta I = \frac{d\psi}{dI} cI \quad (28)$$

where c is the stimulus contrast (Howard et al., 1987 [\[3\]](#))

Under the condition we model, half-saturation of transduction units, $I = 2\psi$ and $\frac{d\psi}{dI} = \frac{1}{4}$. Consequently the photoreceptor signal is

$$(\Delta S)^2(f_x, f_y, f_t) = \left(\frac{\psi}{2}\right)^2 S_{xyt}(f_x, f_y, f_t) |m(f_x, f_y)|^2 |m_t(f_t)|^2 |m_d(f_t)|^2 \quad (29)$$

where

$$\psi = \frac{N_{vil}}{2\tau_r} \quad (30)$$

In the fly NS eye 6 photoreceptors sample the same point in space, therefore the signal transmitted for each pixel in the array is

$$(\Delta S)^2(f_x, f_y, f_t) = (3\psi)^2 S_{xyt}(f_x, f_y, f_t) |m(f_x, f_y)|^2 |m_t(f_t)|^2 |m_d(f_t)|^2 \quad (31)$$

Noise

We consider shot noise generated at the first step in the generation of a quantum bump, the absorption of a photon by a rhodopsin molecule in a microvillus that is not currently processing a bump. Noise generated elsewhere in photoreceptors is not incorporated in our model. Assuming that the photon capture rate ψ is never far from its average value, the shot noise power in a single photoreceptor is given by:

$$\sigma_{sn}^2(f_t) = \frac{\psi}{2} |H(f_t)|^2 \quad (32)$$

where $|H(f_t)|$ is the Fourier transform of the quantum bump and the factor 2 takes account of the fact that at half-saturation the statistics of microvillus activation are binomial (Howard et al., 1987 [\[1\]](#)).

Substituting from Equation 20 [\[2\]](#)

$$\sigma_{sn}^2(f_t) = \frac{\psi}{2} |m_t(f_t)|^2 \quad (33)$$

We assume that the shot noise in different photoreceptors is uncorrelated, therefore the shot noise generated by the 6 photoreceptors coding a pixel is

$$\sigma_{sn}^2(f_t) = 3\psi |m_t(f_t)|^2 \quad (34)$$

Because we assume that the shot noise in different photoreceptors is uncorrelated the power spectrum of shot noise, is on average, the same across the photoreceptor array

$$\sigma_{sn}^2(f_x, f_y, f_t) \propto \begin{cases} 3\psi |H(f_t)|^2 & \text{if } (f_x, f_y) \text{ is sampled} \\ 0 & \text{if } (f_x, f_y) \text{ is not sampled} \end{cases} \quad (35)$$

with the proportionality constant being the inverse of the area in frequency space that is sampled by the photoreceptor array.

Aliasing

When the image projected to the photoreceptor array contains spatial frequencies that are beyond the spatial resolving power of the array, these are aliased to lower frequencies, thereby distorting the sampled image (Snyder et al., 1994 [\[3\]](#)). We assume that all aliased spatial frequencies act as noise, shared equally between the correctly sampled spatial frequencies:

$$\sigma_{al}^2(f_t) = \psi^2 k(f_t) \quad (36)$$

Calculating Information

We calculate the spatial information gathered by the photoreceptor array by modifying the formula for a square spatial sampling lattice (Howard & Snyder, 1983 [\[4\]](#)). For a hexagonal array, the equation for spatial information has the form

$$H = A \int_0^{\mu_s} f_r \log_2[1 + (S/N)(f_r)] df_r \quad (37)$$

where the array cut-off frequency is $\mu_s = \frac{1}{\sqrt{3}\Delta\phi}$. To obtain the factor A , we consider the extreme case where the signal-to-noise ratio is constant across all spatial frequencies, in which case

$$H = A \frac{1}{3(\Delta\phi)^2} \log_2[(N_i)] \quad (38)$$

where $N_i = \sqrt{1 + S/N}$ is the number of discriminable levels of contrast coded by each ommatidium. The factor multiplying $\log_2[N_i]$ is the number of ommatidia in a steradian, which for a hexagonal lattice is $1/S_0 = \frac{2\sqrt{3}}{3(\Delta\phi)^2}$. Thus $A = 2\sqrt{3}$. Therefore spatial information capacity per steradian is

$$H = 2\sqrt{3} \int_0^{\mu_s} f_r \log_2[1 + (S/N)(f_r)] df_r \quad (39)$$

We then extend this equation for spatial information capacity to the temporal domain (van Hateren, 1992). For a spatio-temporal signal the information capacity is

$$H = 2\sqrt{3} \int_{-\infty}^{\infty} \int_0^{\mu_s} f_r \log_2[1 + (S/N)(f_r, f_t)] df_t df_r \quad (40)$$

Inserting signal and noise

$$H = 2\sqrt{3} \int_{-\infty}^{\infty} \int_0^{\mu_s} f_r \log_2\left(1 + \frac{(\Delta S)^2(f_r, f_t)}{\sigma_{sn}^2(f_r, f_t) + \sigma_{al}^2(f_t)}\right) df_t df_r \quad (41)$$

Ballpark estimate of the photoreceptor energy tariff

Energy consumed per microvillus

Our estimates are based on high resolution measurements of the oxygen consumed by isolated *Calliphora vicina* compound eyes, made using a magnetic diver balance (Pangršič et al., 2005). In bright light, an eye consumes 1.8×10^{-9} litres of oxygen per second which, converting to the units commonly used to express metabolic rates, is 6.48 mm^3 of oxygen per hour. With 5000 ommatidia, 60,000 microvilli in the rhabdomere of a photoreceptor R1-6, and taking the central rhabdom formed by R7 and R7 to be equivalent to a single R1-6, the eye contains 2×10^9 microvilli, giving an oxygen consumption per microvillus in bright light $O_{vil}^L = 3.24 \times 10^{-9} \text{ mm}^3$ oxygen per microvillus per hour. In darkness the eye's oxygen consumption drops to one third, giving $O_{vil}^D = 1.08 \times 10^{-9} \text{ mm}^3$ oxygen per microvillus per hour.

Because these estimates are based on measurements from whole eyes they include consumption by all cell types. There are three reasons why photoreceptors dominate total consumption. First, the values of an eye's oxygen consumption in light and in darkness are close to the total photoreceptor consumption predicted from electrophysiological estimates of the ATP consumed by single photoreceptors (Laughlin et al., 1998; Pangršič et al., 2005). Second, photoreceptors are the compound eye's most active cells because transducing photons at high light intensities necessarily requires large fluxes of ions and chemical intermediates. Third, to support their activity photoreceptors contain most of a compound eye's mitochondria.

The rate of oxygen consumption averaged over the course of 24 hrs $\bar{O}_{vil}$, depends on the hours of daylight, DL

$$\bar{O}_{vil} = \frac{DL * O_{vil}^L + (24 - DL) * O_{vil}^D}{24} \text{ mm}^3 \text{ oxygen per mg per hour} \quad (42)$$

Energy consumed by the fly

To convert the energy consumption of the eye per microvillus into an equivalent volume we should ideally use the animal's average specific metabolic rate, measured under natural conditions, but to the best of our knowledge this field metabolic rate has not been measured in blowflies. Therefore, we combine a value of the specific metabolic rate in continuous flight that is towards the high end of the range measured in *Calliphora*, $sFMR = 30 \text{ mm}^3$ oxygen per mg per hour (Yurkiewicz & Smyth, 1966), with the resting metabolic rate measured in the blowfly *Phormia*, $RMR = 2.6 \text{ mm}^3$ oxygen per mg per hour (Keister & Buck, 1961; Niven & Scharlemann, 2005), according the number of hours per day spent flying, T_F , to estimate the daily average specific metabolic rate

$$\overline{SMR} = \frac{T_F * sFMR + (24 - T_F) * RMR}{24} \text{ mm}^3 \text{ oxygen per mg per hour.} \quad (43)$$

Estimating the energy tariff, K_E

Assuming that the density of the eye is 1

$$K_E = \frac{\bar{O}_{vil}}{\overline{SMR}} \text{ mm}^3 \text{ of eye per microvillus} \quad (44)$$

To obtain a plausible range of values we calculate K_E in four conditions (Table 4)

Note that, as expected, our measure of the impact of photoreceptor energy costs, K_E , depends on both the energy consumption per microvillus and the animal's metabolic rate: K_E is increased by increasing daily consumption by photoreceptors and decreased by increasing the animal's overall metabolic rate by increasing time flying, T_F .

Preliminary model of a simple eye

Our generic simple eye model has the same F-number $F = 2$ as our generic fly NS model, and its rhabdom diameter equals the rhabdomere diameter in fly $d_{rh} = 1.9 \mu\text{m}$. The simple eye is hemispherical (Figure 7a), therefore the specific volume of optics is given by

$$V_o = \frac{\pi}{3} f^3 = \frac{\pi}{3} (DF)^3 \mu\text{m}^3 \text{ sr}^{-1} \quad (45)$$

and with a photoreceptor array with rhabdoms of length L , the total specific volume of the eye is given by

$$V_{tot} = \frac{\pi}{3} (DF + L)^3 \quad (46)$$

and the specific volume of the photoreceptor array by

$$V_{ph} = V_{tot} - V_o \quad (47)$$

The receptor spacing, $\Delta\phi$, depends on the lens focal length, f and d_{rh} ,

$$\Delta\phi = \frac{d_{rh}}{f} = \frac{d_{rh}}{DF} \quad (48)$$

This is an important difference between simple eyes and apposition eyes. In apposition eyes model $\Delta\phi$ is can be varied at constant f and D by changing the optical radius R . However, in a simple eye with a diffraction limited lens the relationship between the optical resolving power of the lens and

the anatomical resolving power of the photoreceptor mosaic depends on F-number, F , as follows (Snyder, 1979 [↗](#)). For a hexagonal mosaic of photoreceptors the diffraction limit is reached when

$$\lambda/D = \sqrt{3} \frac{d_{rh}}{f} = \sqrt{3} \frac{d_{rh}}{DF} \quad (49)$$

When λ , the wavelength of light is 500 nm, and $d_{rh} = 2 \mu\text{m}$, then $F = 6.9$. It follows that our model simple eye with $F = 2$ is undersampling by a factor of approximately 3.5. Running our simple eye model with $F = 4$ had very little effect on the optimum L (data not shown), suggesting that our conclusion that efficient simple eyes have much shorter rhabdoms than efficient apposition eyes is robust with respect to undersampling.

Calculating N_{vil} as a function of L

We estimate v , the number of microvilli per unit length of rhabdomere, for our NS eye model using findings from photoreceptors R1-6 in female blowflies. Dividing the number of microvilli in a rhabdomere, 6×10^4 (Howard et al., 1987 [↗](#)), by $L = 260 \mu\text{m}$, which is towards the upper end of the range $L = 220 \mu\text{m} - 280 \mu\text{m}$ reported for blowfly (Hardie, 1985 [↗](#)), gives $v = 230 \mu\text{m}^{-1}$. Although fly rhabdomeres vary in both diameter and degree of taper according to eye region, sex and species, and this sculpting adapts photoreceptor arrays to visual ecology, e.g. (Gonzalez-Bellido et al., 2011 [↗](#)), we fix v to simplify the derivation of points of principle.

For our generic model of a fused rhabdom apposition eye we consider a column of six photoreceptors that, like the short visual fibres R1-6 in fly, terminate in the first optic neuropile. Their six rhabdomeres form the triradiate rhabdom described in locust (Wilson et al., 1978 [↗](#)). This fused rhabdom has a cross-section that approximates an equilateral triangle in which each side is constructed by a pair photoreceptors whose parallel microvilli are equivalent to a single fly R1-6 rhabdomere. Thus each pixel is coded by a set of six rhabdomeres that are equivalent to three fly R1-6 rhabdomeres. Therefore, the number of transduction units coding a pixel is half that in fly and the resulting changes in signal and noise are simply accounted for by replacing the factor $(3\psi)^2$ in Equation 29 [↗](#) with $(\frac{3\psi}{2})^2$, and 3ψ in Equations 33 [↗](#) and 34 [↗](#) with $(\frac{3\psi}{2})$.

For our generic model of a diurnal simple eye we assume that, as in spiders, the rhabdomeres of adjacent photoreceptors abut; i.e. there are no gaps between photoreceptors, therefore a photoreceptor's rhabdomeres project from a central column of cytoplasm. To compare with our apposition models we assume that the diameter of the photoreceptor entrance aperture, $d_{rh} = 1.9 \mu\text{m}$ and contains microvilli projecting in three directions. In this case, the simple eye photoreceptor is equivalent to the triradiate fused rhabdom.

Acknowledgements

We are grateful to Doekele Stavenga for his expert advice on optics, to Jeremy Niven for his advice during the project and his comments on the manuscript, and to Daniel Osorio and James Herbert-Read for their comments on the manuscript.

Additional information

Authors' contributions

FJHH and SBL conceived of the project, FJHH developed, constructed, programmed and ran the models, FJHH and SBL applied the models, curated and analysed published data, and wrote the paper. FJHH's contributions were made while he was a member of the University of Cambridge.

Funding

Francisco Heras was supported by a La Caixa Postgraduate Scholarship and a Caja-Madrid Postgraduate Scholarship. Simon Laughlin was supported by an honorarium from H. Britton Sanderford. The funders had no role in study design, data collection and analysis, the decision to publish, and the preparation of the manuscript.

Additional files

Literature data Table S1 [↗](#)

References

- Alexander R. M. (1997) **A theory of mixed chains applied to safety factors in biological systems** *Journal of Theoretical Biology* **184**:247–252 <https://doi.org/10.1006/jtbi.1996.0270> | [Google Scholar](#)
- Alexander R. M. (1996) **Optima for animals** Princeton University Press [Google Scholar](#)
- Ames A. (2000) **CNS energy metabolism as related to function** *Brain Research Reviews* **34**:42–68 [https://doi.org/10.1016/S0165-0173\(00\)00038-2](https://doi.org/10.1016/S0165-0173(00)00038-2) | [Google Scholar](#)
- Anderson J. C., Laughlin S. B. (2000) **Photoreceptor performance and the coordination of achromatic and chromatic inputs in the fly visual system** *Vision Research* **40**:13–31 [https://doi.org/10.1016/S0042-6989\(99\)00171-6](https://doi.org/10.1016/S0042-6989(99)00171-6) | [Google Scholar](#)
- Autrum H. (1958) **Electrophysiological analysis of the visual systems in insects** *Experimental Cell Research, Supplement* **5**:426–439 [Google Scholar](#)
- Baden T., Nilsson D.-E. (2022) **Is our retina really upside down?** *Current Biology* **32**:R300–R303 <https://doi.org/10.1016/j.cub.2022.02.065> | [Google Scholar](#)
- Barlow H. B. (1952) **The size of ommatidia in apposition eyes** *Journal of Experimental Biology* **29**:667–674 [Google Scholar](#)
- Bastien G. J., Willems P. A., Schepens B., Heglund N. C. (2005) **Effect of load and speed on the energetic cost of human walking** *European Journal of Applied Physiology* **94**:76–83 <https://doi.org/10.1007/s00421-004-1286-z> | [Google Scholar](#)
- Bigman L. S., Levy Y. (2020) **Proteins: Molecules defined by their trade-offs** *Current Opinion in Structural Biology* **60**:50–56 <https://doi.org/10.1016/j.sbi.2019.11.005> | [Google Scholar](#)
- Brandon C. S., James T., Dudycha J. L. (2015) **Selection on incremental variation of eye size in a wild population of *Daphnia*** *Journal of Evolutionary Biology* **28**:2112–2118 <https://doi.org/10.1111/jeb.12711> | [Google Scholar](#)
- Brandon C. S., Dudycha J. L. (2014) **Ecological constraints on sensory systems: Compound eye size in *Daphnia* is reduced by resource limitation** *Journal of Comparative Physiology A* **200**:749–758 <https://doi.org/10.1007/s00359-014-0918-y> | [Google Scholar](#)
- Burton B. G. (2006) **Adaptation of single photon responses in photoreceptors of the housefly, *Musca domestica*: A novel spectral analysis** *Vision Research* **46**:622–635 <https://doi.org/10.1016/j.visres.2005.09.020> | [Google Scholar](#)
- Burton B. G., Laughlin S. B. (2003) **Neural images of pursuit targets in the photoreceptor arrays of male and female houseflies *Musca domestica*** *Journal of Experimental Biology* **206**:3963–3977 <https://doi.org/10.1242/jeb.00600> | [Google Scholar](#)
- Burton B. G., Tatler B. W., Laughlin S. B. (2001) **Variations in photoreceptor response dynamics across the fly retina** *Journal of Neurophysiology* **86**:950–960 <https://doi.org/10.1152/jn.2001.86.2.950> | [Google Scholar](#)

- Burton G. J., Moorhead I. R. (1987) **Color and spatial structure in natural scenes** *Applied Optics* **26**:157–170 <https://doi.org/10.1364/AO.26.000157> | Google Scholar
- Casares F., McGregor A. P. (2021) **The evolution and development of eye size in flies** *WIREs Developmental Biology* **10**:e380 <https://doi.org/10.1002/wdev.380> | Google Scholar
- Coles J. A. (1989) **Functions of glial cells in the retina of the honeybee drone** *Glia* **2**:1–9 <https://doi.org/10.1002/glia.440020102> | Google Scholar
- Cronin T. W., Johnsen S., Marshall N. J., Warrant E. J. (2014) **Visual ecology** Princeton University Press [Google Scholar](https://doi.org/10.1017/9781107301510)
- Darwin C. (1859) **The origin of species by means of natural selection** John Murray [Google Scholar](https://doi.org/10.1017/9781107301510)
- van Steveninck de Ruyter, Laughlin S. B. (1996) **The rate of information transfer at graded-potential synapses** *Nature* **379**:642–645 <https://doi.org/10.1038/379642a0> | Google Scholar
- Drack M., Betz O. (2022) **A technomorphic conceptualisation of biological ‘constructions’ and their evolution** *Vertebrate Zoology* **72**:839–855 <https://doi.org/10.3897/vz.72.e86968> | Google Scholar
- Duncan F. D., Lighton J. R. B. (1994) **The burden within: The energy cost of load carriage in the honeypot ant, *Myrmecocystus*** *Physiological Zoology* **67**:190–203 <https://doi.org/10.1086/physzool.67.1.30163842> | Google Scholar
- Fain G. L., Hardie R., Laughlin S. B. (2010) **Phototransduction and the evolution of photoreceptors** *Current Biology* **20**:R114–R124 <https://doi.org/10.1016/j.cub.2009.12.006> | Google Scholar
- Fain G. L., Sampath A. P. (2021) **Light responses of mammalian cones** *Pflügers Archiv - European Journal of Physiology* **473**:1555–1568 <https://doi.org/10.1007/s00424-021-02551-0> | Google Scholar
- Feller K. D., Sharkey C. R., McDuffee-Altekruse A., Bracken-Grissom H. D., Lord N. P., Porter M. L., Schweikert L. E. (2021) **Surf and turf vision: Patterns and predictors of visual acuity in compound eye evolution** *Arthropod Structure and Development* **60**:101002 <https://doi.org/10.1016/j.asd.2020.101002> | Google Scholar
- Garland T., Downs C. J., Ives A. R. (2022) **Trade-offs (and constraints) in organismal biology** *Physiological and Biochemical Zoology* **95**:82–112 <https://doi.org/10.1086/717897> | Google Scholar
- Garrigan P., Ratliff C. P., Klein J. M., Sterling P., Brainard D. H., Balasubramanian V. (2010) **Design of a trichromatic cone array** *PLOS Computational Biology* **6**:e1000677 <https://doi.org/10.1371/journal.pcbi.1000677> | Google Scholar
- Gonzalez-Bellido P. T., Wardill T. J., Juusola M. (2011) **Compound eyes and retinal information processing in miniature dipteran species match their specific ecological demands** *Proceedings of the National Academy of Sciences* **108**:4224–4229 <https://doi.org/10.1073/pnas.1014438108> | Google Scholar
- Gote J. T., Butler P. M., Zureka D. B., Buschbeck E. K., Morehouse N. (2019) **Growing tiny eyes: How juvenile jumping spiders retain high visual performance in the face of size**

limitations and developmental constraints *Vision Research* **160**:24–36 <https://doi.org/10.1016/j.visres.2019.04.006> | [Google Scholar](#)

Goyens J., Van Wassenbergh S., Dirckx J., Aerts P. (2015) **Cost of flight and the evolution of stag beetle weaponry** *Journal of The Royal Society Interface* **12**:20150222 <https://doi.org/10.1098/rsif.20150222> | [Google Scholar](#)

Hardie R. C. (1985) **Functional organization of the fly retina** In: Autrum H., Ottoson D., Perl E. R., Schmidt R. F., Shimazu H., Willis W. D., editors. *Progress in sensory physiology* Springer Berlin Heidelberg pp. 1–79 https://doi.org/10.1007/978-3-642-70408-6_1 | [Google Scholar](#)

Hardie R. C., Postma M. (2008) **Phototransduction in microvillar photoreceptors of Drosophila and other invertebrates** In: Masland R., Albright T. D., editors. *The senses, a comprehensive reference. volume 1, vision 1* Academic Press pp. 77–130 [Google Scholar](#)

Heras F. J. H., Laughlin S. B. (2017) **Optimizing the use of a sensor resource for opponent polarization coding** *PeerJ* **5**:e2772 <https://doi.org/10.7717/peerj.2772> | [Google Scholar](#)

Horridge G. A. (1969) **Unit studies on retina of dragonflies** *Zeitschrift Fur Vergleichende Physiologie* **62**:1–37 <https://doi.org/10.1007/BF00298040> | [Google Scholar](#)

Howard J., Blakeslee B., Laughlin S. B. (1987) **The intracellular pupil mechanism and photoreceptor signal:noise ratios in the fly *Lucilia cuprina*** *Proc R Soc Lond B* **231**:415–435 <https://doi.org/10.1098/rspb.1987.0053> | [Google Scholar](#)

Howard J., Snyder A. W. (1983) **Transduction as a limitation on compound eye function and design** *Proc R Soc Lond B* **217**:287–307 <https://doi.org/10.1098/rspb.1983.0011> | [Google Scholar](#)

Juusola M., Kouvalainen E., Järvilehto M., Weckström M. (1994) **Contrast gain, signal-to-noise ratio, and linearity in light-adapted blowfly photoreceptors** *Journal of General Physiology* **104**:593–621 <https://doi.org/10.1085/jgp.104.3.593> | [Google Scholar](#)

Keister M., Buck J. (1961) **Respiration of *Phormia regina* in relation to temperature and oxygen** *Journal of Insect Physiology* **7**:51–72 [https://doi.org/10.1016/0022-1910\(61\)90044-0](https://doi.org/10.1016/0022-1910(61)90044-0) | [Google Scholar](#)

Kelber A., Somanathan H. (2019) **Spatial vision and visually guided behavior in Apidae** *Insects* **10**:418 <https://doi.org/10.3390/insects10120418> | [Google Scholar](#)

Kim J. (2014) **Absorption-assisted mode transformation in butterfly compound eyes** *Scientific Reports* **4**:6291 <https://doi.org/10.1038/srep06291> | [Google Scholar](#)

Kirschfeld K. (1976) **The resolution of lens and compound eyes** In: Zettler F., Weiler R., editors. *Neural principles in vision* Springer pp. 354–370 https://doi.org/10.1007/978-3-642-66432-8_19 | [Google Scholar](#)

Kröger R. H., Biehlmaier O. (2009) **Space-saving advantage of an inverted retina** *Vision Research* **49**:2318–2321 <https://doi.org/10.1016/j.visres.2009.07.001> | [Google Scholar](#)

- Labhart T., Nilsson D.-E. (1995) **The dorsal eye of the dragonfly *Sympetrum*: specializations for prey detection against the blue sky** *Journal of Comparative Physiology A* **176**:437–453 <https://doi.org/10.1007/BF00196410> | [Google Scholar](#)
- Land M. F. (1981) **Optics and vision in invertebrates** In: Autrum H., editors. *Handbook of sensory physiology* Springer-Verlag pp. 471–593 [Google Scholar](#)
- Land M. F. (1985) **The morphology and optics of spider eyes** In: Barth F. G., editors. *Neurobiology of arachnids* Springer pp. 53–78 https://doi.org/10.1007/978-3-642-70348-5_4 | [Google Scholar](#)
- Land M. F., Eckert H. (1985) **Maps of the acute zones of fly eyes** *Journal of Comparative Physiology A* **156**:525–538 <https://doi.org/10.1007/BF00613976> | [Google Scholar](#)
- Land M. F., Fernald R. D. (1992) **The evolution of eyes** *Annual Review of Neuroscience* **15**:1–29 <https://doi.org/10.1146/annurev.ne.15.030192.000245> | [Google Scholar](#)
- Land M. F. (1997) **Visual acuity in insects** *Annual Review of Entomology* **42**:147–177 <https://doi.org/10.1146/annurev.ento.42.1.147> | [Google Scholar](#)
- Laughlin S. B. (1981) **A simple coding procedure enhances a neuron's information capacity** *Zeitschrift für Naturforschung C* **36**:910–912 [Google Scholar](#)
- Laughlin S. B., McGinness S. (1978) **The structures of dorsal and ventral regions of a dragonfly retina** *Cell and Tissue Research* **188**:427–447 [Google Scholar](#)
- Laughlin S. B., van Steveninck R. R. R., Anderson J. C. (1998) **The metabolic cost of neural information** *Nature Neuroscience* **1**:36–41 <https://doi.org/10.1038/236> | [Google Scholar](#)
- Laughlin S. B., Weckström M. (1993) **Fast and slow photoreceptors—a comparative study of the functional diversity of coding and conductances in the Diptera** *Journal of Comparative Physiology A* **172**:593–609 <https://doi.org/10.1007/BF00213682> | [Google Scholar](#)
- Menzel J. G., Wunderer H., Stavenga D. G. (1991) **Functional morphology of the divided compound eye of the honeybee drone (*Apis mellifera*)** *Tissue and Cell* **23**:525–535 [https://doi.org/10.1016/0040-8166\(91\)90010-Q](https://doi.org/10.1016/0040-8166(91)90010-Q) | [Google Scholar](#)
- Nijhout H. F., Emlen D. J. (1998) **Competition among body parts in the development and evolution of insect morphology** *Proceedings of the National Academy of Sciences* **95**:3685–3689 <https://doi.org/10.1073/pnas.95.7.3685> | [Google Scholar](#)
- Nilsson D.-E. (2013) **Eye evolution and its functional basis** *Visual Neuroscience* **30**:5–20 <https://doi.org/10.1017/S0952523813000035> | [Google Scholar](#)
- Nilsson D.-E. (2021) **The diversity of eyes and vision** *Annual Review of Vision Science* **7**:19–41 <https://doi.org/10.1146/annurev-vision-121820-074736> | [Google Scholar](#)
- Nilsson D.-E., Land M. F. (2012) **Animal eyes** Oxford University Press [Google Scholar](#)
- Nilsson D.-E., Pelger S. (1994) **A pessimistic estimate of the time required for an eye to evolve** *Proceedings of the Royal Society B: Biological Sciences* **256**:53–58 <https://doi.org/10.1098/rspb.1994.0048> | [Google Scholar](#)

- Niven J. E., Anderson J. C., Laughlin S. B. (2007) **Fly photoreceptors demonstrate energy-information trade-offs in neural coding** *PLoS Biology* **5**:828–840 <https://doi.org/10.1371/journal.pbio.0050116> | [Google Scholar](#)
- Niven J. E., Laughlin S. B. (2008) **Energy limitation as a selective pressure on the evolution of sensory systems** *Journal of Experimental Biology* **211**:1792–1804 <https://doi.org/10.1242/jeb.017574> | [Google Scholar](#)
- Niven J. E., Scharlemann J. P. (2005) **Do insect metabolic rates at rest and during flight scale with body mass?** *Biology Letters* **1**:346–349 <https://doi.org/10.1098/rsbl.2005.0311> | [Google Scholar](#)
- Oakley T. H., Speiser D. I. (2015) **How complexity originates: The evolution of animal eyes** *Annual Review of Ecology, Evolution, and Systematics* **46**:237–260 <https://doi.org/10.1146/annurev-ecolsys-110512-135907> | [Google Scholar](#)
- Okawa H., Sampath A. P., Laughlin S. B., Fain G. L. (2008) **ATP consumption by mammalian rod photoreceptors in darkness and in light** *Current Biology* **18**:1917–1921 <https://doi.org/10.1016/j.cub.2008.10.029> | [Google Scholar](#)
- Pangršić T., Stušek P., Belušić G., Zupancić G. (2005) **Light dependence of oxygen consumption by blowfly eyes recorded with a magnetic diver balance** *Journal of Comparative Physiology A* **191**:75–84 <https://doi.org/10.1007/s00359-004-0571-y> | [Google Scholar](#)
- Perl C. D., Niven J. E. (2016) **Differential scaling within an insect compound eye** *Biology Letters* **12**:20160042 <https://doi.org/10.1098/rsbl.2016.0042> | [Google Scholar](#)
- Piersma T., van Gils J. A. (2011) **The flexible phenotype** Oxford University Press [Google Scholar](#)
- Provis J. M., Dubis A. M., Maddess T., Carroll J. (2013) **Adaptation of the central retina for high acuity vision: Cones, the fovea and the avascular zone** *Progress in Retinal and Eye Research* **35**:63–81 <https://doi.org/10.1016/j.preteyeres.2013.01.005> | [Google Scholar](#)
- Rigosi E., Wiederman S. D., O’Carroll D. C. (2017) **Photoreceptor signalling is sufficient to explain the detectability threshold of insect aerial pursuers** *The Journal of Experimental Biology* **220**:4364–4369 <https://doi.org/10.1242/jeb.166207> | [Google Scholar](#)
- Rossel S. (1979) **Regional differences in photoreceptor performance in the eye of the praying mantis** *Journal of Comparative Physiology* **131**:95–112 <https://doi.org/10.1007/BF00619070> | [Google Scholar](#)
- Ruderman D. L. (1997) **Origins of scaling in natural images** *Vision Research* **37**:3385–3398 [https://doi.org/10.1016/S0042-6989\(97\)00008-4](https://doi.org/10.1016/S0042-6989(97)00008-4) | [Google Scholar](#)
- Scales J. A., Butler M. A. (2016) **The relationship between microhabitat use, allometry and functional variation in the eyes of Hawaiian Megalagrion damselflies** *Functional Ecology* **30**:356–368 <https://doi.org/10.1111/1365-2435.12479> | [Google Scholar](#)

- Schreiber S., Machens C. K., Herz A. V. M., Laughlin S. B. (2002) **Energy-efficient coding with discrete stochastic events** *Neural Computation* **14**:1323–1346 <https://doi.org/10.1162/089976602753712963> | [Google Scholar](#)
- Seitz G. (1968) **Der Strahlengang in Appositionsaugen von Calliphora erythrocephala (Meig.)** *Zeitschrift für Vergleichende Physiologie* **59**:205–231 [Google Scholar](#)
- Smith W. C., Butler J. F. (1991) **Ultrastructure of the Tabanidae compound eye: Unusual features for Diptera** *Journal of Insect Physiology* **37**:287–296 [https://doi.org/10.1016/0022-1910\(91\)90063-6](https://doi.org/10.1016/0022-1910(91)90063-6) | [Google Scholar](#)
- Snyder A. W. (1977) **Acuity of compound eyes - physical limitations and design** *Journal of Comparative Physiology* **116**:161–182 <https://doi.org/10.1007/BF00605401> | [Google Scholar](#)
- Snyder A. W. (1979) **Physics of vision in compound eyes** In: Autrum H., editors. *Handbook of sensory physiology* Springer pp. 225–313 [Google Scholar](#)
- Snyder A. W., Bossomaier T. J., Hughes A. (1994) **The theory of comparative eye design** In: Blakemore C., editors. *Vision: Coding and efficiency* pp. 45–52 [Google Scholar](#)
- Snyder A. W., Stavenga D. G., Laughlin S. B. (1977) **Spatial information capacity of compound eyes** *Journal of Comparative Physiology* **116**:183–207 [https://doi.org/10.1016/0042-6989\(77\)90151-1](https://doi.org/10.1016/0042-6989(77)90151-1) | [Google Scholar](#)
- Song Z., Juusola M. (2014) **Refractory sampling links efficiency and costs of sensory encoding to stimulus statistics** *The Journal of Neuroscience* **34**:7216–7237 <https://doi.org/10.1523/JNEUROSCI.4463-13.2014> | [Google Scholar](#)
- Song Z., Postma M., Billings S. A., Coca D., Hardie R. C., Juusola M. (2012) **Stochastic, adaptive sampling of information by microvilli in fly photoreceptors** *Current Biology* **22**:1371–1380 <https://doi.org/10.1016/j.cub.2012.05.047> | [Google Scholar](#)
- Stavenga D. G. (2003a) **Angular and spectral sensitivity of fly photoreceptors. I. Integrated facet lens and rhabdomere optics** *Journal of Comparative Physiology A* **189**:1–17 <https://doi.org/10.1007/s00359-002-0370-2> | [Google Scholar](#)
- Stavenga D. G. (2003b) **Angular and spectral sensitivity of fly photoreceptors. II. Dependence on facet lens F-number and rhabdomere type in Drosophila** *Journal of Comparative Physiology A* **189**:189–202 <https://doi.org/10.1007/s00359-003-0390-6> | [Google Scholar](#)
- Stavenga D. G. (2004a) **Angular and spectral sensitivity of fly photoreceptors. III. Dependence on the pupil mechanism in the blowfly Calliphora** *Journal of Comparative Physiology A* **190**:115–129 <https://doi.org/10.1007/s00359-003-0477-0> | [Google Scholar](#)
- Stavenga D. G. (2004b) **Visual acuity of fly photoreceptors in natural conditions - dependence on UV sensitizing pigment and light-controlling pupil** *Journal of Experimental Biology* **207**:1703–1713 <https://doi.org/10.1242/jeb.00949> | [Google Scholar](#)
- Stavenga D. G., Kruizinga R., Leertouwer H. (1990) **Dioptrics of the facet lenses of male blowflies Calliphora and Chrysomya** *Journal of Comparative Physiology A* **166**:365–371 <https://doi.org/10.1007/BF00204809> | [Google Scholar](#)

- Stearns S. C. (1989) **Trade-offs in life-history evolution** *Functional Ecology* **3**:259–268 <https://doi.org/10.2307/2389364> | [Google Scholar](#)
- Sterling P., Laughlin S. (2015) **Principles of neural design** MIT Press [Google Scholar](#)
- Straw A. D., Warrant E. J., O’Carroll D. C. (2006) **A ‘bright zone’ in male hoverfly *Eristalis tenax* eyes and associated faster motion detection and increased contrast sensitivity** *Journal of Experimental Biology* **209**:4339–4354 <https://doi.org/10.1242/jeb.02517> | [Google Scholar](#)
- Sumner-Rooney L. (2018) **The kingdom of the blind: Disentangling fundamental drivers in the evolution of eye loss** *Integrative and Comparative Biology* **58**:372–385 <https://doi.org/10.1093/icb/icy047> | [Google Scholar](#)
- Sumner-Rooney L., Kenny N. J., Ahmed F., Williams S. T. (2019) **The utility of micro-computed tomography for the non-destructive study of eye microstructure in snails** *Scientific Reports* **9**:15411 <https://doi.org/10.1038/s41598-019-51909-z> | [Google Scholar](#)
- Tatler B., O’Carroll D. C., Laughlin S. B. (2000) **Temperature and the temporal resolving power of fly photoreceptors** *Journal of Comparative Physiology A* **186**:399–407 <https://doi.org/10.1007/s003590050439> | [Google Scholar](#)
- Taylor C., Weibel E. (1981) **Design of the mammalian respiratory system. I. problem and strategy** *Respiratory Physiology* **44**:1–10 [Google Scholar](#)
- Taylor G. K., Thomas A. L. R. (2014) **Evolutionary biomechanics: Selection, phylogeny and constraint** Oxford University Press [Google Scholar](#)
- Taylor G. J., Tichit P., Schmidt M. D., Bodey A. J., Rau C., Baird E. (2019) **Bumblebee visual allometry results in locally improved resolution and globally improved sensitivity** *eLife* **8**:e40613 <https://doi.org/10.7554/eLife.40613> | [Google Scholar](#)
- Thomas K. N., Gower D. J., Bell R. C., Fujita M. K., Schott R. K., Streicher J. W. (2020) **Eye size and investment in frogs and toads correlate with adult habitat, activity pattern and breeding ecology** *Proceedings of the Royal Society B: Biological Sciences* **287**:20201393 <https://doi.org/10.1098/rspb.2020.1393> | [Google Scholar](#)
- Thomas R. J., Kelly D. J., Goodship N. M. (2004) **Eye design in birds and visual constraints on behavior** *Ornitologia Neotropicana* **15**:243–250 [Google Scholar](#)
- van Hateren J. H. (1992) **Theoretical predictions of spatiotemporal receptive fields of fly LMCs, and experimental validation** *Journal of Comparative Physiology A* **171**:157–170 <https://doi.org/10.1007/BF00188924> | [Google Scholar](#)
- van Hateren J. H., Hardie R. C., Rudolph A., Laughlin S. B., Stavenga D. G. (1989) **The bright zone, a specialized dorsal eye region in the male blowfly *Chrysomya megacephala*** *Journal of Comparative Physiology A* **164**:297–308 [Google Scholar](#)
- van Hateren J. H., Schilstra C. (1999) **Blowfly flight and optic flow. II. head movements during flight** *Journal of Experimental Biology* **202**:1491–1500 [Google Scholar](#)

- Varela F., Wiitanen W. (1970) **Optics of the compound eye of the honeybee *Apis mellifera*** *Journal of General Physiology* **55**:336–358 <https://doi.org/10.1085/jgp.55.3.336> | [Google Scholar](#)
- Walls G. L. (1942) **The vertebrate eye and its adaptive radiation** Cranbrook Institute of Science [Google Scholar](#)
- Wardill T. J., Fabian S. T., Pettigrew A. C., Stavenga D. G., Nordström K., Gonzalez-Bellido P. T. (2017) **A novel interception strategy in a miniature robber fly with extreme visual acuity** *Current Biology* **27**:854–859 <https://doi.org/10.1016/j.cub.2017.01.050> | [Google Scholar](#)
- Wehner R. (1981) **Spatial vision in arthropods** In: Autrum H., editors. *Handbook of sensory physiology* Springer-Verlag pp. 287–616 [Google Scholar](#)
- Weibel E. (2000) **Symmorphosis: On form and function shaping life** Harvard University Press [Google Scholar](#)
- Wilson M., Garrard P., McGinness S. (1978) **The unit structure of the locust compound eye** *Cell and Tissue Research* **195**:205–226 <https://doi.org/10.1007/BF00236720> | [Google Scholar](#)
- Witter M., Cuthill I. (1993) **The ecological costs of avian fat storage** *Philosophical Transactions of the Royal Society B-Biological Sciences* **340**:73–92 <https://doi.org/10.1098/rstb.1993.0050> | [Google Scholar](#)
- Wong F., Knight B. W., Dodge F. A. (1980) **Dispersion of latencies in photoreceptors of limulus and the adapting-bump model** *Journal of General Physiology* **76**:517–537 <https://doi.org/10.1085/jgp.76.5.517> | [Google Scholar](#)
- Yurkiewicz W. J., Smyth T. (1966) **Effects of temperature on oxygen consumption and fuel utilization by the sheep blowfly** *Journal of Insect Physiology* **12**:403–408 [https://doi.org/10.1016/0022-1910\(66\)90003-5](https://doi.org/10.1016/0022-1910(66)90003-5) | [Google Scholar](#)
- Zeil J. (1983) **Sexual dimorphism in the visual system of flies: The compound eyes and neural superposition in Bibionidae (Diptera)** *Journal of Comparative Physiology A* **150**:379–393 <https://doi.org/10.1007/BF00605027> | [Google Scholar](#)
- Zhang L.-Q., Cottaris N. P., Brainard D. H. (2022) **An image reconstruction framework for characterizing initial visual encoding** *eLife* **11**:e71132 <https://doi.org/10.7554/eLife.71132> | [Google Scholar](#)

Author information

Francisco JH Heras

Department of Zoology, University of Cambridge, Cambridge, United Kingdom
ORCID iD: [0000-0001-8124-2359](https://orcid.org/0000-0001-8124-2359)

Simon B Laughlin

Department of Zoology, University of Cambridge, Cambridge, United Kingdom
ORCID iD: [0000-0003-4659-6543](https://orcid.org/0000-0003-4659-6543)

For correspondence: sl104@cam.ac.uk

Editors

Reviewing Editor

Fred Rieke

University of Washington, Seattle, United States of America

Senior Editor

Lois Smith

Boston Children's Hospital, Boston, United States of America

Reviewer #1 (Public review):

Summary:

Two important factors in visual performance are the resolving power of the lens and the signal-to-noise ratio of the photoreceptors. These both compete for space: a larger lens has improved resolving power over a smaller one, and longer photoreceptors capture more photons and hence generate responses with lower noise. The current paper explores the tradeoff of these two factors, asking how space should be allocated to maximize eye performance (measured as encoded information).

The revisions, to my read, have greatly improved the paper. Most of this was due to setting clear expectations from the start of the paper. Nice work!

<https://doi.org/10.7554/eLife.96517.2.sa3>

Reviewer #2 (Public review):

Summary:

In short, the paper presents a theoretical framework that predicts how resources should be optimally distributed between receptors and optics in eyes.

After revision of an already excellent contribution, the manuscript is now even better. The authors have responded carefully to all reviewer comments.

Strengths:

The authors build on the principle of resource allocation within an organism and develop a formal theory for optimal distribution of resources within an eye between the receptor array and the optics. Because the two parts of eyes, receptor arrays and optics, share the same role of providing visual information to the animal it is possible to isolate these from resource allocation in the rest of the animal. This allows for a novel and powerful way of exploring the principles that govern eye design. By clever and thoughtful assumptions/constraints, the authors have built a formal theory of resource allocation between the receptor array and the optics for two major types of compound eye as well as for camera-type eyes. The theory is formalized with variables that are well characterized in a number of different animal eyes, resulting in testable predictions.

The authors use the theory to explain a number of design features that depend on different optimal distribution of resources between the receptor array and the optics in different types of eye. As an example, they successfully explain why eye regions with different spatial resolution should be built in different ways. They also explain differences between different types of eye, such as long photoreceptors in apposition compound eyes and much shorter receptors in camera type eyes. The predictive power in the theory is impressive.

To keep the number of parameters at a minimum, the theory was developed for two types of compound eye (neural superposition, and apposition) and for camera-type eyes. It is possible to extend the theory to other types of eye, although it would likely require more variables and assumptions/constraints to the theory. It is thus good to introduce the conceptual ideas without overdoing the applications of the theory.

The paper extends a previous theory, developed by the senior author, that develops performance surfaces for optimal cost/benefit design of eyes. By combining this with resource allocation between receptors and optics, the theoretical understanding of eye design takes a major leap and provides entirely new sets of predictions and explanations for why eyes are built the way they are.

The paper is well written and even though the theory development in the Results may be difficult to take in for many biologists, the Discussion very nicely lists all the major predictions under separate headings, and here the text is more tuned for readers that are not entirely comfortable with the formalism of the Results section. I must point out though that the Results section is kept exemplary concise. The figures are excellent and help explain concepts that otherwise may go above the head of many biologists.

<https://doi.org/10.7554/eLife.96517.2.sa2>

Reviewer #3 (Public review):

Summary:

This is a proposal for a new theory for the geometry of insect eyes. The novel cost-benefit function combines the cost of the optical portion with the photoreceptor portion of the eye. These quantities are put on the same footing using a specific (normalized) volume measure, plus an energy factor for the photoreceptor compartment. An optimal information transmission rate then specifies each parameter and resource allocation ratio for a variable total cost. The elegant treatment allows for comparison across a wide range of species and eye types. Simple eyes are found to be several times more efficient across a range of eye parameters than neural superposition eyes. Some trends in eye parameters can be explained by optimal allocation of resources between the optics and photoreceptors compartments of the eye.

Strengths:

Data from a variety of species roughly align with rough trends in the cost analysis, e.g. as a function of expanding the length of the photoreceptor compartment.

New data could be added to the framework once collected, and many species can be compared.

Eyes of different shapes are compared.

Weaknesses:

Detailed quantitative conclusions are not possible given the approximations and simplifying assumptions in the models and weak accounting for trends in the data across eye types.

Comments on revisions:

I have no additional comments for the authors and appreciate the revisions and corrections implemented - I think those changes have improved the clarity of the manuscript and expanded the potential readership for the paper.

Author Response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

Summary:

Two important factors in visual performance are the resolving power of the lens and the signal-to-noise ratio of the photoreceptors. These both compete for space: a larger lens has improved resolving power over a smaller one, and longer photoreceptors capture more photons and hence generate responses with lower noise. The current paper explores the tradeoff of these two factors, asking how space should be allocated to maximize eye performance (measured as encoded information).

Your summary is clear, concise and elegant. The competition is not just for space, it is for space, materials and energy. We now emphasise that we are considering these three costs in our rewrites of the Abstract and the first paragraph of the Discussion.

Strengths:

The topic of the paper is interesting and not well studied. The approach is clearly described and seems appropriate (with a few exceptions - see weaknesses below). In most cases, the parameter space of the models are well explored and tradeoffs are clear.

Weaknesses:

Light level

The calculations in the paper assume high light levels (which reduces the number of parameters that need to be considered). The impact of this assumption is not clear. A concern is that the optimization may be quite different at lower light levels. Such a dependence on light level could explain why the model predictions and experiment are not in particularly good agreement. The paper would benefit from exploring this issue.

Thank you for raising this point. We briefly explained in our original Discussion, under Understanding the adaptive radiation of eyes (Version 1, lines 756 – 762), how our method can be modified to investigate eyes adapted for lower light levels. We have some thoughts on how eyes might be adapted. In general, transduction rates are increased by increasing D , reducing f , increasing d_{rh} and increasing L . In addition, d_{rh} is increased to allow for a larger D within the constraint of eye radius/corneal surface area, and to avoid wasteful oversampling (the changes in D , f and d_{rh} increase acceptance angle $\Delta\rho$). We suspect that in eyes optimised for the efficient use of space, materials and energy the increases in L will be relatively small, first because increasing D , reducing f and increasing d_{rh} are much more effective at increasing transduction rate than increasing L . Second, increasing sensitivity by reducing f decreases the cost V_0 whereas increasing sensitivity by increasing L increases the cost V_{ph} . This disadvantage, together with exponential absorption, might explain why L is only 10% - 20% longer in the apposition eyes of nocturnal bees (Somanathan et al, J. comp. Physiol. A195, 571583, 2009). Because this line of argument is speculative and enters new territory, we have not included it in our revised version. We already present a lot of new material for readers to digest, and we agree with referee 2 that “It is possible to extend the theory to other types of eyes, although it would likely require more variables and assumptions/constraints to the theory. It is thus good to introduce the conceptual ideas without overdoing the applications of the theory”. Nonetheless, we take your point that some

of the eyes in our data set might be adapted for lower light levels, and we have rewritten the Discussion section, How efficiently do insects allocate resources within their apposition eyes accordingly. On line 827 – 843 we address the assumption that eyes are adapted for full daylight, and also take the opportunity to mention two more reasons for increasing the eye parameter p : namely increasing image velocity (Snyder, 1979), and constructing bright zones that increase the detectability of small targets (van Hateren et al., 1989; Straw et al., 2006).

Discontinuities

The discontinuities and non-monotonicity of the optimal parameters plotted in Figure 4 are concerning. Are these a numerical artifact? Some discussion of their origin would be quite helpful.

Good points, we now address the discontinuities in the Results, where they are first observed (lines 311 - 319)

Discrepancies between predictions and experiment

As the authors clearly describe, experimental measurements of eye parameters differ systematically from those predicted. This makes it difficult to know what to take away from the paper. The qualitative arguments about how resources should be allocated are pretty general, and the full model seems a complex way to arrive at those arguments. Could this reflect a failure of one of the assumptions that the model rests on - e.g. high light levels, or that the cost of space for photoreceptors and optics is similar? Given these discrepancies between model and experiment, it is also hard to evaluate conclusions about the competition between optics and photoreceptors (e.g. at the end of the abstract) and about the importance for evolution (end of introduction).

Your misgivings boil down to two issues: what use is a model that fails to fit the data, and do we need a complicated model to show something that seems to be intuitively obvious? Our study is useful because it introduces new approaches, methods, factors and explanations which advance our analysis and understanding of eye design and evolution. Your comments make it clear that we failed to get this message across and we have revised the manuscript accordingly. We have rewritten the Abstract and the first paragraph of the Discussion to emphasise the value of our new measure of cost, specific volume, by including more of its practical advantages. In particular, our use of specific volume 1) opens the door to the morphospace of all eyes of given type and cost. 2) This allows one to construct performance surfaces across morphospace that not only identify optima, but by evaluating the sub-optimal cast light on efficiency and adaptability. 3) Shows that photoreceptor energy costs have a major impact on design and efficiency, and 4) allows us to calculate and compare the capacities and efficiencies of compound eyes and simple eyes using a superior measure of cost. It is also possible that your dissatisfaction was deepened by disappointment. The first sentence of our original Abstract said that the goal of design is to maximize performance, so you might have expected to see that eyes are optimised. Given that optimization provides cast iron proof that a system is designed to be efficient, and previous studies of coding by fly LMCs (Laughlin, 1981; Srinivasan et al., 1982 & van Hateren 1992) validated Barlow's Efficient Coding Hypothesis by showing that coding is optimised, your expectation is reasonable. However, our investigation of how the allocation of resources to optics and photoreceptors affects an eye's performance, efficiency and design does not depend *a priori* on finding optima, therefore we have removed the "maximized". Our revised Abstract now says, "to improve performance".

In short, our study illustrates an old adage in statistics "All models fail to fit, but some are useful". As is often the case, the way in which our model fails is useful. In the original version of the Results and Discussion, we argued that the allocation of resources is efficient, and

identified factors that can, in principle, explain the scattering of data points. Indeed, our modelling identifies two of these deficiencies; a lack of data on species-specific energy usage, and the need for models that quantify the relationship between the quality of the captured image and the behavioural tasks for which an eye might be specialised. Thus, by examining the model's failings we identify critical factors and pose new questions for future research. We have rewritten the Discussion section How efficiently do insects allocate resources.... to make these points. We hope that these revisions will convince you that we have established a starting point for definitive studies, invented a vehicle that has travelled far enough to discover new territory, and shown that it can be modified to cope with difficult terrain.

Turning to the need for a complicated model, because the costs and benefits depend on elementary optics and geometry, we too thought that there ought to be a simple model. However, when we tried to formulate a simple set of equations that approximate the definitive findings of our more complicated model we discovered that this is not as straightforward as we thought. Many of the parameters in our model interact to determine costs and benefits, and many of these interactions are non-linear (e.g. the volumes of shells in spheres involve quadratic and cubic terms, and information depends on the log of a square root). So, rather than hold back publication of our complicated model, we decided to explain how it works as clearly as we can and demonstrate its value.

In response to your final comment, "it is hard to evaluate conclusions about the competition between optics and photoreceptors (e.g. at the end of the abstract) and about the importance for evolution (end of introduction)", we stand by our original argument. There must be competition in an eye of fixed cost, and because competition favours a heavy investment in photoreceptors, both in theory and in practice, it is a significant factor in eye design. A match between investments in optics and photoreceptors is predicted by theory and observed in fly NS eyes, therefore this is a design principle. As for evolution, no one would deny that it is important to view the adaptive radiation of eyes through a cost-benefit lens. Our lens is the first to view the whole eye, optics and photoreceptor array, and the first to treat the costs of space, materials and energy. Although the view through our lens is a bit fuzzy, it reveals that costs, benefits and trade-offs are important. Thus we have established a promising starting point for a new and more comprehensive cost-benefit approach to understanding eye design and evolution. As for the involvement of genes, when there are heritable changes in phenotype genes must be involved and if, as we suggest, efficient resource allocation is beneficial, the developmental mechanisms responsible for allocating resources to optics and photoreceptor array will be playing a formative role in eye evolution.

Reviewer #2 (Public Review):

Summary:

In short, the paper presents a theoretical framework that predicts how resources should be optimally distributed between receptors and optics in eyes.

Strengths:

The authors build on the principle of resource allocation within an organism and develop a formal theory for optimal distribution of resources within an eye between the receptor array and the optics. Because the two parts of eyes, receptor arrays and optics, share the same role of providing visual information to the animal it is possible to isolate these from resource allocation in the rest of the animal. This allows for a novel and powerful way of exploring the principles that govern eye design. By clever and thoughtful assumptions/constraints, the authors have built a formal theory of resource allocation between the receptor array and the optics for two major types of compound eye as well as for camera-type eyes. The theory is formalized with variables that are well characterized in a number of different animal eyes, resulting in testable predictions.

The authors use the theory to explain a number of design features that depend on different optimal distribution of resources between the receptor array and the optics in different types of eyes. As an example, they successfully explain why eye regions with different spatial resolution should be built in different ways. They also explain differences between different types of eyes, such as long photoreceptors in apposition compound eyes and much shorter receptors in camera type eyes. The predictive power in the theory is impressive.

To keep the number of parameters at a minimum, the theory was developed for two types of compound eye (neural superposition, and apposition) and for camera-type eyes. It is possible to extend the theory to other types of eyes, although it would likely require more variables and assumptions/constraints to the theory. It is thus good to introduce the conceptual ideas without overdoing the applications of the theory.

The paper extends a previous theory, developed by the senior author, that develops performance surfaces for optimal cost/benefit design of eyes. By combining this with resource allocation between receptors and optics, the theoretical understanding of eye design takes a major leap and provides entirely new sets of predictions and explanations for why eyes are built the way they are.

The paper is well written and even though the theory development in the Results may be difficult to take in for many biologists, the Discussion very nicely lists all the major predictions under separate headings, and here the text is more tuned for readers that are not entirely comfortable with the formalism of the Results section. I must point out though that the Results section is kept exemplary concise. The figures are excellent and help explain concepts that otherwise may go above the head of many biologists.

We are heartened by your appreciation of our manuscript - it persuaded us not to undertake extensive revisions – thank you.

Reviewer #3 (Public Review):

Summary:

This is a proposal for a new theory for the geometry of insect eyes. The novel costbenefit function combines the cost of the optical portion with the photoreceptor portion of the eye. These quantities are put on the same footing using a specific (normalized) volume measure, plus an energy factor for the photoreceptor compartment. An optimal information transmission rate then specifies each parameter and resource allocation ratio for a variable total cost. The elegant treatment allows for comparison across a wide range of species and eye types. Simple eyes are found to be several times more efficient across a range of eye parameters than neural superposition eyes. Some trends in eye parameters can be explained by optimal allocation of resources between the optics and photoreceptors compartments of the eye.

Strengths:

Data from a variety of species roughly align with rough trends in the cost analysis, e.g. as a function of expanding the length of the photoreceptor compartment.

New data could be added to the framework once collected, and many species can be compared.

Eyes of different shapes are compared.

Weaknesses:

Detailed quantitative conclusions are not possible given the approximations and simplifying assumptions in the models and poor accounting for trends in the data across eye types.

Reviewer #1 (Recommendations For The Authors):

Figure 1: Panel E defines the parameters described in panel d. Consider swapping the order of those panels (or defining D and Delta Phi in the figure legend for d). Order follows narrative, eye types then match

We think that you are referring to Figure 1. We modified the legend.

Lines 143-145: How does a different relative cost impact your results?

Thank you for raising this question. Because our assumption that relative costs are the same is our starting point, and for optics it is not an obvious mistake, we do not raise your question here. We address your question where you next raise it because, for photoreceptors the assumption is obviously wrong. We now emphasise that our method for accounting for photoreceptor energy costs can be applied to other costs.

Lines 187-190: Same as above - how do your results change if this assumption is not accurate?

We have revised our manuscript to emphasise that we are dealing with the situation in which our initial assumption (costs per unit volume are equal) breaks down. On (lines 203 - 208) we write “ However, this assumption breaks down when we consider specific metabolic rates. To enable and power phototransduction, photoreceptors have an exceptionally high specific metabolic rate (energy consumed per gram, and hence unit volume, per second) (Laughlin et al., 1998; Niven et al., 2007; Pangršič et al., 2005). We account for this extra cost by applying an energy surcharge, S_E . To equate....

We also revised part of the Discussion section, Specific volume is a useful measure of cost to make it clear that we are able take account for situations in which the costs per unit volume are not equal, and we give our treatment of photoreceptor energy costs as an example of how this is done. On lines 626 - 640 we say

Cost estimates can be adjusted for situations in which costs per unit volume are not equal, as illustrated by our treatment of photoreceptor energy consumption. To support transduction the photoreceptor array has an exceptionally high metabolic rate (Laughlin et al., 1998; Niven et al., 2007; Pangršič et al., 2005). We account for this higher energy cost by using the animal's specific metabolic rate (power per unit mass and hence power per unit volume) to convert an array's power consumption into an equivalent volume (Methods). Photoreceptor ion pumps are the major consumers of energy and the smaller contribution of pigmented glia (Coles, 1989) is included in our calculation of the energy tariff K_E . (Methods) The higher costs of materials and their turnover in the photoreceptor array can be added the energy tariff K_E but given the magnitude of the light-gated current (Laughlin et al., 1998) the relative increase will be very small. Thus for our intents and purposes the effects of these additional costs are covered by our models. For want of sufficient data...”.

Reviewer #2 (Recommendations For The Authors):

A few comments for consideration by the authors:

(1) In the abstract, Maybe give another example explaining why other eyes should be different to those of fast diurnal insects.

This worthwhile extrapolation is best kept to the Discussion.

(2) Would it be worthwhile mentioning that the photopigment density is low in rhabdoms compared to vertebrate outer segments? This will have major effects on the relative size of retina and optics.

Thank you, we now make this good point in the Discussion (lines 698-702).

(3) It took me a while to understand what you mean by an energy tariff. For the less initiated reader many other variables may be difficult to comprehend. A possible remedy would be to make a table with all variables explained first very briefly in a formal way and then explained again with a few more words for readers less fluent in the formalism.

A very useful suggestion. We have taken your advice (p.4).

(4) The "easy explanation" on lines 356-357 need a few more words to be understandable.

We have expanded this argument, and corrected a mistake, the width of the head front to back is not 250 μm , it is 600 μm (lines 402-407)

(5) Maybe devote a short paragraph in the Discussion to other types of eye, such as optical superposition eyes and pinhole eyes. This could be done very shortly and without formalism. I'm sure the authors already have a good idea of the optimal ratio of receptor arrays and optics in these eye types.

We do not discuss this because we have not found a full account of the trade-offs and their effects on costs and benefits. We hope that our analysis of apposition and simple eyes will encourage people to analyse the relationships between costs and benefits in other eye types. To this end we pointed out in the Discussion that recent advances in imaging and modelling could be helpful.

(6) Could the sentence on lines 668-671 be made a little clearer?

"Efficiency is also depressed by increasing the photoreceptor energy tariff K_E , and in line with the greater impact of photoreceptor energy costs in simple eyes, the reduction in efficiency is much greater in simple eyes (Figure 8b).0.

We replaced this sentence with "In both simple and apposition eyes efficiency is reduced by increasing the photoreceptor energy tariff K_E . This effect is much greater in simple eyes, thus as found for reductions in photoreceptor length (Figure 7b), K_E has more impact on the design of simple eyes" (lines 749 – 752).

(7) I have some reservations about the text on lines 789-796. The problem is that optics can do very little to improve the performance of a directional photoreceptor where Δr should optimally be very wide. Here, membrane folding is the only efficient way to improve performance (SNR). The option to reduce Δr for better performance comes later when simultaneous spatial resolution (multiple pixels) is introduced.

Yes, we have been careless. We have rewritten this paragraph to say (lines 920-931)

"Two key steps in the evolution of eyes were the stacking of photoreceptive membranes to absorb more photons, and the formation of optics to intercept more photons and concentrate them according to angle of incidence to form an image (Nilsson, 2013, 2021). Our modelling of

well-developed image forming eyes shows that to improve performance stacked membranes (rhabdomeres) compete with optics for the resources invested in an eye, and this competition profoundly influences both form and function. It is likely that competition between optics and photoreceptors was shaping eyes as lenses evolved to support low resolution spatial vision. Thus the developmental mechanisms that allocate resources within modern high resolution eyes (Casares & MacGregor, 2021), by controlling cell size and shape, and as our study emphasises, gradients in size and shape across an eye, will have analogues or homologues in more ancient eyes. Their discovery....” (lines 920-931

Reviewer #3 (Recommendations For The Authors):

Suggestions for major revisions:

While the approach is novel and elegant, the results from the analysis of insect morphology do not broadly support the optimization argument and hardly constrain parameters, like the energy tariff value, at all. The most striking result of the paper is the flat plateau in information across a broad range of shape parameters and the length, and resolution trend in Figure 5.

At no point in the Results and Discussion do we argue that resource allocation is optimized. Indeed, we frequently observe that it is not. Our mistake was to start the Abstract by observing that animals evolve to minimise costs. We have rewritten the Abstract accordingly.

The information peaks are quite shallow. This might actually be a very important and interesting result in the paper - the fact that the information plateaus could give the insect eye quite a wide range of parameters to slide between while achieving relatively efficient sensing of the environment. Instead of attempting to use a rather ad hoc and poorly supported measure of energetics in PR cost, perhaps the pitch could focus on this flexibility. K_E does not seem to constrain eye parameters and does not add much to the paper.

We agree, being able to construct performance surfaces across morphospace is an important advance in the field of eye design and evolution, and the performance surface’s flat top has interesting implications for the evolution of adaptations. Encouraged by your remarks, we have rewritten the Abstract and the introductory paragraph of the Discussion to draw attention to these points.

We are disappointed that we failed to convince you that our energy tariff, K_E , is no better than a poorly supported ad hoc parameter that does not add much to the paper. In our opinion a resource allocation model that ignores photoreceptor energy consumption is obviously inadequate because the high energy cost of phototransduction is both wellknown and considered to be a formative factor in eye evolution (Niven and Laughlin, 2008). One of the advantages of modelling is that one can assess the impact of factors that are known to be present, are thought to be important, but have not been quantified. We followed standard modelling practice by introducing a cost that has the same units as the other costs and, for good physiological reasons, increases linearly with the number of microvilli, according to K_E . We then vary this unknown cost parameter to discover when and why it is significant. We were pleased to discover that we could combine data on photoreceptor energy demands and whole animal metabolic rates to establish the likely range of K_E . This procedure enabled us to unify the cost-benefit analyses of optics and photoreceptors, and to discover that realistic values of K_E have a profound impact on the structure and performance of an efficient eye. We hope that this advance will encourage people to collect the data needed to evaluate K_E . To emphasise the importance of K_E and dispel doubts associated with the failure of the model to fit the data, we have revised two sections: Flies invest efficiently in costly photoreceptor arrays in the Results, and How efficiently do insects allocate resources within their apposition

eyes? in the Discussion. These rewrites also explain why it is impossible for us to infer K_E by adjusting its value so that the model's predictions fit the data.

The graphics after Figure 3 are quite dense and hard to follow. None of the plateau extent shown in Fig 3 is carried through to the subsequent plots, which makes the conclusions drawn from these figures very hard to parse. If the peak information occurs on a flat plateau, it would be more helpful to see those ranges of parameters displayed in the figures.

Ideally one should do as you suggest and plot the extent of the plateau, but in our situation this is not very helpful. In the best data set, flies, optimised models predict D well, get close to Δp in larger eyes, and demonstrate that these optimum values are not very sensitive to K_E . L is a different matter, it is very sensitive to K_E L which, as we show (and frequently remind) is poorly constrained by experimental data. The best we can do is estimate the envelope of L vs C_{tot} curves, as defined by a plausible range of K_E L . Because most of the plateau boundaries you ask for will fall within this envelope, plotting them does little to clear the fog of uncertainty. We note that all three referees agree that our model can account for two robust trends, i) in apposition eyes L increase with optical resolving power and acuity, both within individual eyes and among eyes of different sizes, and ii) L is much longer in apposition eyes than in simple eyes. Nonetheless, the scatter of data points and their failure to fit creates a bad impression. We gave a number of reasons why the model does not fit the data points, but these were scattered throughout the Results and Discussion and, as referees 1 and 3 point out, this makes it difficult to draw convincing conclusions. To rectify this failing, we have rewritten two sections, in the Results Flies invest efficiently in costly photoreceptor arrays and in the Discussion, How efficiently do insects allocate resources within their apposition eyes?, to discuss these reasons en bloc, draw conclusions and suggest how better data and refinements to modelling could resolve these issues.

Throughout the figures, the discontinuities in the optimal cuts through parameter space are not sufficiently explained.

We added a couple of sentences that address the “jumps” (lines 313 – 318)

None of the data seems to hug any of the optimal lines and only weakly follow the trends shown in the plots. This makes interpretation difficult for the reader and should be better explained. The text can be a little telegraphic in the Results after roughly page 10, and requires several readings to glean insight into the manuscript's conclusions.

We revised the Results section in which we compare the best data set, flies' NS eyes with theoretical predictions, Flies invest efficiently in costly photoreceptor arrays, to expand our interpretation of the data and clarify our arguments. The remaining sections have not been expanded. In the next section, which is on fused rhabdom apposition eyes, our interpretation of the scattering of data points follows the same line of argument. The remaining Results sections are entirely theoretical.

Overall, the rough conclusions outlined in the Results seem moderately supported by the matches of the data to the optimal information transmission cuts through parameter space, but only weakly.

We agree, more data is required to test and refine our theoretical predictions.

The Discussion is long and well-argued, and contains the most cogent writing in the manuscript.

Thank you: this is most pleasing. We submitted our study to eLife because it allows longer Discussions, but we worried that ours was too long. However, we felt that our extensive Discussion was necessary for two reasons. First, we are introducing a new approach to understanding of eye design and evolution. Second, because the data on eye morphology and costs are limited, we had to make a number of assumptions and by discussing these, warts and all, we hoped to encourage experimentalists to gather more data and focus their efforts on the most revealing material.

Minor comments:

We have acted upon most of your minor comments and we confine our remarks to our disagreements. We are grateful for your attention to details that we \textshould have picked up on.

It's a more standard convention to say "cost-benefit" rather than with a colon.

"equation" should be abbreviated "eq" or "eqn", never with a "t"

when referring to the work of van Hateren, quote the paper and the database using "van Hateren" not just "Hateren"

small latex note: use "\textit{SNR}" to get the proper formatting for those letters when in the math environment

Line 100-110: "f" is introduced, but only f' is referenced in the figure. This should be explained in order. d_{rh} is not included in the figure. Also in this section, d_{rh}/f is also referenced before $\Delta \rho_{rh}$, which is the same quantity, without explanation.

Figure 1 shows eye structure and geometry. f' is a lineal dimension of the eye but f is not, so f is not shown in Fig 1e. We eliminated the confusion surrounding $\Delta \rho_{rh}$ by deleting "and changing the acceptance angle of the photoreceptive waveguide $\Delta \rho_{rh}$ (Snyder, 1979)".

Fig 1 caption: this says "From dorsal to ventral," then describes trends that run ventral to dorsal, which is a confusing typo.

Fig 3 - adding some data points to these plots might help the reader understand how (or if) K_E is constrained by the data.

It is not possible to add data points because to total cost, C_{tot}, is unknown.

Fig 4c (and in other subplots): the jumps in L with C_{tot} could be explained better in the text - it wasn't clear to this reviewer why there are these discontinuities.

Dealt with in the revised text (lines 310-318).

Fig 4d: The caption for this subplot could be more clearly written.

We have rewritten the subscript for subplot 4d.

Fig 5 and other plots with data: please indicate which symbols are samples from the same species. This info is hard to reconstruct from the tables.

We have revised Figure 5 accordingly. Species were already indicated in Figure 6.

Line 328: missing equation number

<https://doi.org/10.7554/eLife.96517.2.sa0>