

Reviewed Preprint

v1 • January 14, 2026

Not revised

✉ For correspondence:

ajith.vasanthakumar@petermac.org

rallan@wehi.edu.au

* These authors contributed equally.

† These authors contributed equally.

Competing interests: No

competing interests declared

Funding: See [page 14](#)

Reviewing editor: Florent Ginhoux,
Singapore Immunology Network,
Singapore

© 2026, Quon et al. This article is
distributed under the terms of the
[Creative Commons Attribution](#)
[License](#), which permits unrestricted
use and redistribution provided that
the original author and source are
credited.

The long non-coding RNA *Dreg1* is required for optimal ILC2 development

Sara Quon^{1,2,*}, Adelynn Tang^{3,4,5,*}, Nadia Iannarella^{1,2}, Kael Schoffer^{1,2}, Wing Fuk Chan^{1,2},

Timothy M Johanson^{1,2}, Ajithkumar Vasanthakumar^{3,4,5,†} ✉, Rhys S Allan^{1,2,†} ✉

¹The Walter and Eliza Hall Institute of Medical Research, Parkville, Australia • ²Department of Medical Biology, The University of Melbourne, Parkville, Australia • ³Olivia Newton-John Cancer Research Institute, Heidelberg, Australia • ⁴La Trobe University, Bundoora, Australia • ⁵Peter MacCallum Cancer Centre, Melbourne, Australia

eLife Assessment

This is a **useful** study that investigates the role of the long non-coding RNA *Dreg1* for the development, differentiation, or maintenance of group 2 ILC (ILC2). The authors generate *Dreg1*^{-/-} mice and show a reduction of group 2 innate lymphoid cells (ILC2). However, the strength of evidence supporting the impact of *Dreg1* on *Gata3* expression, a transcription factor required for ILC2 cell fate decisions, and the cell-intrinsic requirement of *Dreg1* for ILC2 remain **incomplete**. This study will be of interest to immunologists.

<https://doi.org/10.7554/eLife.109408.1.sa2>

Abstract

Gata3 is an essential transcription factor for the development of several distinct immune cell lineages such as T cells, natural killer (NK) cells and innate lymphoid cells (ILC). As such, the levels and timing of *Gata3* expression are critical for directing lineage fate decisions. The *Gata3* locus has a complex and dynamic distal regulatory enhancer landscape. Recently we identified a non-coding RNA, *Dreg1*, located immediately upstream of the classic +280kb T/NK cell enhancer (Tce1). To test its function, we excised the *Dreg1* locus in mice and observed a selective reduction of group 2 ILCs (ILC2) across multiple tissues, but mature T, NK and other ILC lineages remained unchanged. In bone marrow, common innate lymphoid cell progenitors (ILCP) increased while ILC2 progenitors (ILC2P) decreased, with a modest reduction of *Gata3* in upstream progenitors consistent with an early developmental bottleneck. Chromatin profiling showed the *Dreg1* locus is accessible in early lymphoid progenitors and became decorated with H3K27ac in ILCP in a Tcf1-dependent manner. Furthermore, Tcf1-deficient cells did not express *Dreg1* and showed alterations in the epigenetic landscape of the *Dreg1* locus. Finally, we discovered that potential homologues of *Dreg1* harboured in a syntenic enhancer of *GATA3* are also highly expressed in human ILC2. Taken together we conclude that *Dreg1* is a Tcf1-dependent non-coding RNA critical for fine tuning the high level of *Gata3* required for the optimal development of the ILC2 lineage.

Introduction

The immune system comprises a collection of functionally distinct cell types which act in concert to protect the body from infection and malignancy. These diverse cell populations develop from common haematopoietic progenitors in a highly controlled fashion driven by a network of transcription factors that control lineage-specific gene expression programs. GATA-binding protein 3 (*Gata3*) is one of the master transcription factors that regulates the development of immune cells such as T cells, Natural Killer (NK) cells, and innate lymphoid cells (ILCs) (1, 2). *Gata3* is also critical for the development of CD4⁺ T helper 2 (Th2) cells (3) which produce cytokines such as IL4, 5 and 13 that are critical for protection against extracellular pathogens. It has recently emerged that a population of ILCs known as group 2 ILCs (ILC2) with similar characteristics to Th2 cells reside in non-lymphoid tissues such as the small intestine, visceral adipose tissue and lung and have been

shown to play a non-redundant role in immunity to helminths and drive allergic immune responses(4). ILC2s develop in the bone marrow via common ILC progenitor (CILP)(5), common helper ILC progenitor (CHILP)(6), innate lymphoid cell progenitor (ILCP) (7) and finally ILC2 progenitor (ILC2P)(8). Gata3 is critical to the development of ILCs (8, 9) and high levels are required for the development of ILC2P in the bone marrow (8, 10).

Dosage of Gata3 plays a critical role in T, NK and ILC development and peripheral immunity (8, 10, 11). To support this *Gata3* expression is spatiotemporally tuned by several distal enhancers that dictate *Gata3* activity in these individual lineages. Previous research identified a 7.1kb region ~280kb from the *Gata3* gene that is important for its expression in NK and T cell development, known as T cell enhancer 1 (Tce1) (12, 13). Subsequent work found that this region (also termed *Gata3* +278/285) also impacts ILC2 development but not Th2 cell differentiation (14). ILC2-specific effects have been attributed to additional distal regulatory regions, such as *Gata3* +674/762 (14) and the *Gata3* tandem super enhancers (+500/764)(15). In addition, a Th2 specific enhancer (*Gata3* +906/935) has recently been identified (16). Overall, this complex regulatory landscape mechanistically fit with a dosage-threshold model controlling *Gata3* expression.

Recently we discovered a long non-coding RNA gene that we named *Distal regulatory enhancer of Gata3 1 (Dreg1)* which lies ~0.6kb upstream of Tce1 enhancer (17, 18). Expression of *Dreg1* is high in early T cell progenitors and Th2 cells, mirroring *Gata3* expression (17). Overexpression studies indicated that it may be involved in the establishment but not maintenance of *Gata3* expression (17, 18), but it remains unclear how the loss of *Dreg1* affects immune system development. Here, we explore the role of *Dreg1* in the regulation of immune cell differentiation. We find that deletion of the *Dreg1* locus leads to a specific loss of ILC2 progenitors and their downstream progeny in non-lymphoid tissues but not T or NK cells. Chromatin profiling reveals that the regulatory regions around the *Dreg1* gene are open in haematopoietic progenitors and are bound by the transcription factor TCF1 in early innate lymphoid progenitors and TCF1-deficient mice lack *Dreg1* expression. Finally, we discovered potential homologues of *Dreg1* that highly expressed in human ILC2 emanating from a syntenic enhancer of GATA3.

Results

Deletion of the *Dreg1* locus results in a specific reduction of ILC2

To understand how *Dreg1* contributes to the development of the immune system we generated *Dreg1*-deficient mice by CRISPR-Cas9-mediated excision of the complete 3kb of the *Dreg1* gene (Figure 1A). We confirmed that the *Dreg1* gene was removed from the germline and found that these mice reproduced at Mendelian frequencies and exhibited no overt developmental defects suggesting that *Dreg1* is not required for embryogenesis.

Given our previous finding of the relatively high levels of *Dreg1* expression in T cells (17), and of a critical role of the Tce1 (*Gata3* +278/285) enhancer (12, 14), we expected that the deletion of *Dreg1* would impact T cell and NK cell development. We therefore performed an analysis of the different populations of T cells. Mature CD4⁺ and CD8⁺ T cells in the spleen appeared unaltered in frequency (Supplementary Figure 1). Moreover, both populations had similar proportions of naïve, effector, and memory cells (Supplementary Figure 1B & C). This suggests that *Dreg1* is not required for conventional T cell development or maintenance. In addition, the loss of *Dreg1* did not affect the development of NK cells and γδ T cells as normal proportions were observed in the spleen (Supplementary Figure 1D). Overall, this analysis suggests that *Dreg1* is dispensable for NK and T cell development.

Gata3 is also critical for the development of all innate lymphoid cells (ILCs)(9) and the levels of Gata3 are critical in directing ILC2 lineage development (8, 19). Therefore, we next explored peripheral sites known to harbour substantial numbers of these cell types such as visceral adipose tissue (VAT), the small intestine lamina propria (SI LP) and the lung. While these tissues had relatively normal proportions of NK, ILC1 and ILC3 (Figure 1B, C & E) we noted a specific




Figure 1. Dreg1 deletion results in a specific reduction in peripheral ILC2 cells.

(A) Deletion of Dreg1 in mice using CRISPR-Cas9. (B-E) Representative FACS plots and % of CD45⁺ of NK and ILC populations from the visceral adipose tissue (VAT), small intestinal lamina propria (SI LP), and lung from wild type (WT) or Dreg1^{-/-} mice. Shown is one representative experiment of two with n=4 mice/group. Mean and SEM together with individual data points are shown. Data were statistically analyzed by Student *t* test.

reduction in the population of ILC2 in each of the tissues we examined (Figure 1D [↗](#)). Taken together these results show that the loss of *Dreg1* results in the specific loss of ILC2 in different peripheral tissues.

Dreg1 deficiency results in the reduction of ILC2 progenitors

We next investigated the origins of the ILC2 defect in the *Dreg1*-deficient mice. It is known that expression of *Gata3* in the early ILC progenitor stage is required for development of mature ILC2s ([8](#), [10](#), [19](#)). Indeed, analysis of publicly available RNAseq data ([20](#)) demonstrated that *Gata3* expression was very high in ILC2P and ILC2 compared to other populations, whereas the highest expression of *Dreg1* was found in ILC2P (Figure 2A [↗](#)). We examined ILC development in the bone marrow (Figure 2B-G [↗](#), gating strategy in Supplementary Figure 2A [↗](#)). We began by examining the common lymphoid (CLP) and lymph-myeloid primed progenitors (LMPP) and found no overt changes in proportions or total number of these populations (Figure 2B [↗](#) & C). However, we found increased CILP, CHILP and ILCP in the *Dreg1*-deficient mice (reaching significance in the CILPs) (Figure 2D-F [↗](#)). We observed a substantial reduction of the ILC2P population suggesting that the loss of *Dreg1* leads to a block in ILC development (Figure 2G [↗](#)). This suggests that the loss of *Dreg1* results in a bottleneck at the early stages of ILC development. We next examined the levels of *Gata3* in these populations (Figure 2H [↗](#)). We observed a reduction in *Gata3* signal in the upstream CHILP and the ILCP from the *Dreg1*-deficient mice, but interestingly *Gata3* levels were normal in the ILC2P even though this population is reduced in the *Dreg1*-deficient mice. This suggests that *Dreg1* acts to regulate *Gata3* prior to the ILC2P stage which leads to a block in their development, however, if cells can overcome this perturbation, they express relatively normal levels of *Gata3*.

Dynamic epigenetic regulation of the *Dreg1* locus in a Tcf1-dependent manner during ILC development

To gain a greater understanding of how the expression of *Dreg1* is regulated we next studied the chromatin state of the *Dreg1* locus during early ILC development via analysis of ATAC-Seq data generated on progenitors and their downstream ILC progeny (Figure 3A [↗](#)). Of note, the original Refseq annotation of *Dreg1* was missing the first exon and TSS which we identified ([17](#)) and show in Figure 3A [↗](#). We observed three significant accessible chromatin “peaks”, with the predominant one at the promoter which was open at the CLP stage remained so in ILCP and ILC2P. One region downstream became more accessible in ILCP and ILC2P in line with that observed by others ([14](#)). Overall, we find the promoter of *Dreg1* is open in early lymphoid progenitors and the acquisition of chromatin accessibility downstream correlates with increased *Dreg1* expression in ILC2 progenitors.

To understand the transcription factors that might be involved in regulating *Dreg1* expression we examined the motifs at the *Dreg1* promoter and downstream accessible region. We observed a specific enrichment of motifs for Tcf1, Lef1 and Ets1 suggesting that these transcription factors may be involved in regulating *Dreg1* expression. As Tcf1 (encoded by the *Tcf7* gene) has been implicated in ILC2 development ([21](#), [22](#)) we focussed on its role in regulating *Dreg1* expression. Firstly, we examined publicly available Tcf1 CUT&Run data ([23](#)) and found that indeed Tcf1 was specifically bound to the accessible sites of the *Dreg1* locus in early innate lymphoid progenitors (EILP, which are a population that sit between CILP and CHILP) (Figure 3B [↗](#)). We next examined ChIC-Seq data ([24](#)) of active (H3K27ac) or repressive (H3K27me3) histone modifications at different stages of ILC development (Figure 3D [↗](#)). Firstly, we found no evidence of H3K27me3 in this region in any of the cell types examined. However, there was a modest accumulation of H3K27ac around the accessible regions in the *Dreg1* locus at the EILPs which was noticeably increased at the ILCP stage. Of note, Tcf1-deficient (via deletion of the *Tcf7* gene) EILP showed lower levels of H3K27ac suggesting that Tcf1 is likely involved in creating a permissive chromatin landscape at the *Dreg1* locus during ILC development. Indeed, the expression of *Dreg1* was dependent on the presence of




Figure 2. *Dreg1* deletion results in a bottleneck in ILC development and a reduction in ILC2P.

(A) Boxplots showing expression of *Gata3* and *Dreg1* from RNA-Seq (GSE77695). (B-G) Quantification of ILC progenitor subsets with percentages (upper) and total numbers (below). (H) Mean fluorescence intensity (MFI) of *Gata3* measured on the CHILP, ILCP and ILC2P. Shown is one representative experiment of two with $n=3-4$ mice/group. Mean and SEM together with individual data points are shown. Data were statistically analyzed by Student *t* test.




Figure 3. *Dreg1* locus is dynamically regulated in a Tcf1-dependent manner during ILC2 development.

(A) Chromatin accessibility around *Dreg1* in ILC progenitors. (B) Motifs enriched in accessible regions around *Dreg1*. (C) Tcf1 binding around *Dreg1* locus in EILP. (D) Histone marks around *Dreg1* at different stages of ILC development and in Tcf7^{-/-} EILP. (E) *Gata3* and *Dreg1* expression quantified from RNA-Seq data in ILC progenitors and Tcf7^{-/-} EILP.

Tcf1 as *Tcf7*^{-/-} EILPs lacked *Dreg1* expression (Figure 3E). Taken together this data suggests that the alterations to the chromatin state *Dreg1* locus and its subsequent transcriptional activity during ILC development is dependent on Tcf1.

Identification of a human GATA3 enhancer harbouring *Dreg1* homologues that are active in ILC2s

Previously we identified a region syntenic to the mouse *Dreg1* locus on human chromosome 10 that is a similar distance from the *GATA3* gene and forms an 3D interaction in T cells but not B cells suggesting it may represent an enhancer element (17). Interestingly, bidirectional transcription initiates from the sequence-conserved *cis* element to produce two potential human homologs of *Dreg1*, the lncRNAs CAT00000105356.1 (we named *DREG1.1*) and CAT00000117261.1 (*DREG1.2*). To explore whether these transcripts are also expressed in ILC populations we examined publicly available RNAseq data from healthy human blood (25)(Figure 4A). We found that *GATA3* was highly expressed in human ILCP, ILC2 and TH2 cells. We also observed this pattern for *DREG1.1* and to an even greater extent for *DREG1.2* suggesting that the syntenic region harbours transcripts which reflect the same expression pattern as murine *Dreg1*.

Finally, we investigated the functional role of this potential enhancer of *GATA3* by examining a recently published tiling CRISPR deletion screen searching for regulatory elements that control *GATA3* expression in human Th2 cells (26). We focused on the syntenic region which was defined as functional sequence (FS) 23 (Figure 4B). Specifically, guides that deleted conserved sequences FS23-4 and FS23-5 led to a significant downregulation of *GATA3* expression (26) suggesting that this region represents a distal enhancer of *GATA3* (Figure 4B). ATACseq data from either cultured (pcILC2) or fresh ILC2 from healthy human blood (27) revealed a clear accessible region in these cells that overlaps the *DREG1.1/2* and FS23 (Figure 4C). Overall, we find that this syntenic region in humans represents an enhancer that is important for high levels of *GATA3* expression and contains two non-coding RNA genes that show a similar expression pattern to murine *Dreg1*.

Discussion

Here we show that germline excision of the *Dreg1* locus resulted in a specific reduction in ILC2 cells owing to a developmental bottleneck during ILC development in the bone marrow. These results are in line with the fact that very high levels of *Gata3* are required for ILC2 development (8, 10). Given its proximity to the *Tcf1* enhancer, a surprising finding was that the deletion of *Dreg1* did not affect T or NK cells lineages. While it is tempting to speculate that the *Dreg1* locus represents a ILC2-specific enhancer element we favour the idea that *Dreg1* likely contributes to dosage tuning of *Gata3* during ILC2 commitment and that ILC2s because of their higher *Gata3* threshold are more sensitive to modest reductions, whereas T and NK cells are buffered by redundant modules.

How *Dreg1* is involved in regulating *Gata3* levels mechanistically is yet to be determined. *Dreg1* may act as an enhancer RNA that cooperates with chromatin modifying proteins (28) to reinforce local H3K27ac and enhancer-promoter communication. Future experiments that selectively suppress *Dreg1* transcription for example via antisense oligonucleotides or CRISPRi at the *Dreg1* promoter will enable the discrimination of RNA-dependent from DNA element-dependent effects.

In humans, we identify two noncoding transcripts within the syntenic *GATA3* distal enhancer that are highly expressed in ILCP, ILC2 and TH2 cells. Perturbations within this conserved block in an independent tiling-deletion screen resulted in reduced *GATA3* expression in T cells. While these experiments were performed in T cells, we expect that this enhancer region will likely also play an important role in the regulation of ILC2 development. Given that ILC2 have been implicated in the development of allergic reactions (4) the noncoding transcripts we identified may represent targets for the modulation of *GATA3* levels and ILC2 function in disease.




Figure 4. A syntenic region in humans represents a *GATA3* enhancer element with a transcriptional profile akin to murine *Dreg1*.

(A) *GATA3* and *DREG1.1* and *DREG1.2* expression quantified from RNA-Seq data in ILC and T helper populations from isolated from healthy human blood. (B) Alignment of Functional Sequence 23 (FS23) that overlaps the syntenic region revealed as regulating *GATA3* expression in a CRISPR deletion screen of human TH2 cells (26). (C) Chromatin accessibility around FS23 guides from the CRISPR deletion screen (26).

Methods

Mice

The *Dreg1* knockout mouse line was generated at the Melbourne Advanced Genome Editing Centre (MAGEC) laboratory using CRISPR-Cas9 through microinjection of 2 sgRNAs (CTACTTGCTGACAAGTCGTC and TCTAGTAAGTCCAGTTGCTT) to target the 3,594 bp genomic region around *Dreg1*. The F0 offspring were validated for *Dreg1* deletion with forward primer GACCAGATATGGAGACGTGCA and reverse primer TCTTTGCCATCTTCTGTGTGC and were backcrossed with wild-type C57BL6/J mice.

Cells and tissues were obtained from littermate control mice aged at least 6 weeks unless otherwise specified. Due to sex differences in ILC populations (29), only female mice were used. All mice were maintained at the WEHI Animal Facility under specific pathogen-free conditions.

Cell isolation

Thymus and spleen

The spleen and thymus were made into single-cell suspension by mashing through a 70µm mesh. Splenic cells were treated with red cell lysis buffer.

Bone marrow

Both pairs of hips, femurs, and tibia bones were grounded in a mortar and pestle to collect the bone marrow. The cell suspension was filtered through a 70µm cell strainer and negatively enriched for ILC cells with the mouse lineage cell depletion kit (Miltenyi).

Extraction of lymphocytes from non-lymphoid tissues

Lymphocytes from the VAT and lung were finely chopped in 3-5mL digestion buffer (RPMI and 2% FCS with 2mg/mL Collagenase IV). The suspension was digested for 40 min at 37°C and shaken at 180 rpm. Digested VAT samples were topped up to 30ML with FACS buffer (1xPBS with 2% FCS) and spun at 800g for 15 min at 4°C. Lungs were mashed through a 70µm filter and topped to 30mL FACS buffer and pelleted at 1700rpm for 5min. Both VAT and lung pellets were treated with RBC lysis buffer.

Isolation of lymphocytes from small intestinal lamina propria

Mesentery and Peyer's patches were removed longitudinally, and faeces removed. Intestine was chopped into 0.5cm pieces into 1xPBS and vortexed. Intestinal pieces were placed into 20mLs of dissociation buffer (1xHBSS without Ca²⁺ and Mg²⁺ with 2% FCS, 10mM HEPES and 5 mM EDTA) and digested for 40min at 37°C at 180rpm. Digested pieces were washed with FACS buffer and then placed in 5-8mL of digestion buffer (1xHBSS with indicator containing 2% FCS, 10mM HEPES, 2mg/mL Collagenase IV and 2mg/mL DNase I) and incubated for 30-40min at 37°C at 180rpm. Digested tissue was mashed through a 70 µm cell strainer and pelleted. A 40/80% percoll gradient was performed to enrich for lymphocytes.

Flow cytometry and antibodies

Fluorophore conjugated antibodies against mouse antigens were used for flow cytometry. Antibodies, clone names and manufactures: **Biological**- CD4 (GK1.5) APCFire750, CD8a (53- 6.7) BV570, NKp46 (29A1.4) PEcy7, ICOS (7E.17G9) BV421, Ki67 (11F6) BV650, CD62L (MEL-14) APC and T-bet (4B10) APC. **BD**- CD19 (1D3) BV786, TCRb (H57-597) BV750, TCRgd (GL3) BV711, B220 (RA3-6B2) BV786, GATA3 (L50-823) BUV395, RORgt (Q31-378) BV421, RORgt (Q31-378) PerCPcy5.5, KLRG1 (2F1) BUV615, CD49a (Ha31/8), CD25 (PC61) BV480, NK1.1 (PK136) BUV805, NK1.1 (PK136) BB700, CD44 (IM7) BUV496. **Invitrogen/bioscience**- CD11c (N418) FITC, FOXP3 (FJK-16s) PE, IL-33R (ST2) (RMST2-2) PerCP-eFLuor710, Streptavidin BUV563. **WEHI**-CD45.2 (104) AF700. Live dead staining was performed using Fixable viability stain 440UV diluted in 1xPB at 1:1000 dilution. Surface staining of antibodies was for 30min on ice at 1/200, 1/300 or 1/400 in FACS buffer. Intracellular staining of transcription factors was overnight at 4°C using the Foxp3/Transcription factor staining

buffer set from eBioscience overnight. Cell numbers for specific cell populations were calculated using the volume of the sample that was analyzed, the cell counts per sample, and the percentage of the specific cell population.

RNA-Seq Analysis

Publicly available RNA-Seq data were downloaded from SRA and re-analysed. Briefly fastq files were analyzed with FASTQC (0.12.1) and adapters were removed using trimmomatic(30) (0.36). The reads were aligned to GRCm39 for mouse samples and GRCh38 for human samples using hisat2(31) (2.0.5). Gene counts were quantified using Rsubread(32) featureCounts (1.6.3) with annotation from gencode(33) (vM33 for mouse and v42 for human), which had the annotation for the *Dreg1* gene or human homologs of the *Dreg1* gene added, respectively. Samples were processed using edgeR(34) (3.42.4). Lowly expressed genes were removed, and samples were normalized according to library size. Gene expression was plotted as logCPM using ggplot2 (3.4.4).

ATAC-Seq Analysis

Publicly available ATAC-Seq data were downloaded from SRA and re-analysed. Processing was based on the ENCODE ATAC-Seq pipeline. Briefly fastq files were analyzed with FASTQC and adapters were removed using trimmomatic (0.36). The reads were aligned to the GRCm39 genome for mouse samples and GRCh38 genome for human samples using bowtie2 (2.4.4). Duplicate reads were marked using Picard tools (2.26.11) and reads were filtered using samtools(35) (1.9) with -F 1804 -f 2. Fold-change coverage tracks were created using macs2 (2.2.7.1) callpeak with a smooth window of 150 and a shift size of -75 and bdgcmp. The bedgraph was converted to a bigwig using ucsc tools (331) bedGraphToBigWig(36) and visualized using IGV(37) (2.17.4) with group scaling among samples from the same experiment.

Motif Scanning


The regions of interest as identified through peak calling of ATAC-Seq data were scanned for motifs using FIMO(38) from the MEME suite (5.0.5) using the Hocomoco(39) (H12) core motif database. The resulting list was filtered to remove factors that were not expressed in ILC2P cells as determined through bulk RNA-Seq (GSE77695) with a cutoff of ≥ 1 LogCPM.

ChIC and CUT&Run Analysis


Publicly available ChIP-Seq or CUT&Run data were downloaded from SRA and re-analysed. Briefly fastq files were analyzed with FASTQC and adapters were removed using trimmomatic (0.36). The reads were aligned to the GRCm39 genome for mouse samples using bowtie2 (2.4.4) with the options -q -5 0 -3 0. Duplicate reads were marked using Picard tools (2.26.11) and reads were filtered using samtools (1.9) with -F 1804 -f 2. Fold-change coverage tracks were created using macs2 (2.2.7.1) callpeak with a smooth window of 50 and a shift size of -25 for transcription factors and 150 and -75 for histone marks along with macs2 bdgcmp. The bedgraph was converted to a bigwig using ucsc tools (331) bedGraphToBigWig and visualized using IGV (2.17.4) with group scaling among samples from the same experiment.

CRISPR Guides

The guide counts were downloaded from GSE190860 and were used to create a bed file to identify the target regions. The bed file was visualized in IGV (2.17.4).



Supplemental Figure 1. *Dreg1* deletion in does not affect T or NK cells. (A-C) Representative FACS plots with quantification (right) showing expression of (A) CD8 α and CD4 in splenic T cells. (B) CD44 and CD62L in splenic CD8⁺ T cells (C) CD44 and CD62L in splenic CD4⁺ T cells. (D) Quantification of splenic $\gamma\delta$ T cells and NK cells. Shown is one representative experiment of two with n=3 mice/group. Mean and SEM together with individual data points are shown. Data were statistically analyzed by Student *t* test.



Supplemental Figure 2. Gating of ILC populations in peripheral tissues and bone marrow.

(A) Gating scheme for bone marrow ILC progenitors corresponding to Figure 2 [link](#).

Data availability

All data generated or analyzed during this study are included in the manuscript and supporting files; source data files have been provided for all figures.

Dataset type	Accession Number
Bulk RNA-Seq (Mouse-ILC)(20)	GSE77695
Bulk RNA-Seq (Mouse-ILC w TCF1KO)(23)	GSE113767
Bulk RNA-Seq (Human)(25)	PRJEB35186
Bulk ATAC-Seq (ILCP)(14)	GSE169542
Bulk ATAC-Seq (HumanMouseILC2)(27)	GSE231999
Bulk ChIP-Seq (Histone)(40)	GSE85156
Bulk CUT&Run (EILP TCF1)(23)	GSE128483
Bulk ChIC-Seq (Histone Tcf1 KO)(24)	GSE142468
CRISPR Screen(26)	GSE190860

Acknowledgements

The authors thank the members of the Allan and Vasanthakumar labs for their input into this project. The authors also gratefully acknowledge the WEHI Flow Cytometry and Bioservices Facilities for their support and assistance in this work. We also thank the Melbourne Advanced Genome Editing Centre (MAGEC) laboratory for generation of the *Dreg1*^{-/-} mice.

Additional information

Funding

This work was supported by the NHMRC, Australia Ideas Grant 2001131 and a Stafford Fox Medical Foundation Research Grant to R.A and an NHMRC Investigator Grant 2009336 to A.V. This work was made possible through Victorian State Government Operational Support Program and the Australian Government NHMRC IRISS.

Authorship contributions

Sara Quon: Conceptualization, Investigation, Formal Analysis, Writing-Original Draft, Review & Editing. **Adelynn Tang:** Conceptualization, Investigation, Formal Analysis, Writing-Review & Editing. **Nadia Iannarella:** Investigation. **Kael Schoffer:** Investigation. **Wing Fuk Chan:** Conceptualization, Methodology. **Timothy Johanson:** Investigation, Supervision. **Ajith Vasanthakumar:** Conceptualization, Supervision, Funding Acquisition, Writing-Review & Editing. **Rhys Allan:** Conceptualization, Supervision, Funding Acquisition, Writing-Original Draft, Review & Editing.

Funding

Funder	Grant reference number	Author
Australian NHMRC	2001131	Rhys Allan
Stafford Fox Medical Research Foundation		Rhys Allan
Australian NHMRC	2009336	Ajithkumar Vasanthakumar

Author ORCID iDs

Sara Quon: <https://orcid.org/0000-0003-3860-3491>

Adelynn Tang: <https://orcid.org/0000-0001-8256-6013>

Kael Schoffer: <https://orcid.org/0000-0002-5683-9935>

Wing Fuk Chan: <https://orcid.org/0000-0002-8719-7787>

Timothy M Johanson: <https://orcid.org/0000-0003-4605-6416>

Ajithkumar Vasanthakumar: <https://orcid.org/0000-0002-1620-7781>

Rhys S Allan: <https://orcid.org/0000-0003-0906-2980>

Note

This reviewed preprint has been updated to correct author email address.

References

1. **Harly C.**, Cam M., Kaye J., Bhandoola A. (2018) Development and differentiation of early innate lymphoid progenitors. *J Exp Med* **215**:249-262
2. **Ho I. C.**, Tai T. S., Pai S. Y. (2009) GATA3 and the T-cell lineage: essential functions before and after T-helper-2-cell differentiation. *Nat Rev Immunol* **9**:125-135
3. **Zheng W.**, Flavell R. A. (1997) The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* **89**:587-596
4. **Jarick K. J.**, Topczewska P. M., Jakob M. O., Yano H., Arifuzzaman M., Gao X., Boulekou S., Stokic-Trtica V., Leclerc P. S., Preusser A., et al. (2022) Non-redundant functions of group 2 innate lymphoid cells. *Nature* **611**:794-800
5. **Yu X.**, Wang Y., Deng M., Li Y., Ruhn K. A., Zhang C. C., Hooper L. V. (2014) The basic leucine zipper transcription factor NFIL3 directs the development of a common innate lymphoid cell precursor. *eLife* **3**
6. **Klose C. S. N.**, Flach M., Mohle L., Rogell L., Hoyler T., Ebert K., Fabiunke C., Pfeifer D., Sexl V., Fonseca-Pereira D., et al. (2014) Differentiation of type 1 ILCs from a common progenitor to all helper-like innate lymphoid cell lineages. *Cell* **157**:340-356
7. **Constantinides M. G.**, McDonald B. D., Verhoef P. A., Bendelac A. (2014) A committed precursor to innate lymphoid cells. *Nature* **508**:397-401
8. **Hoyler T.**, Klose C. S., Souabni A., Turqueti-Neves A., Pfeifer D., Rawlins E. L., Voehringer D., Busslinger M., Diefenbach A. (2012) The transcription factor GATA-3 controls cell fate and maintenance of type 2 innate lymphoid cells. *Immunity* **37**:634-648
9. **Yagi R.**, Zhong C., Northrup D. L., Yu F., Bouladoux N., Spencer S., Hu G., Barron L., Sharma S., Nakayama T., et al. (2014) The transcription factor GATA3 is critical for the development of all IL-7R α -expressing innate lymphoid cells. *Immunity* **40**:378-388
10. **Klein Wolterink R. G.**, Serafini N., van Nimwegen M., Vosshenrich C. A., de Bruijn M. J., Fonseca Pereira D., Veiga Fernandes H., Hendriks R. W., Di Santo J. P. (2013) Essential, dose-dependent role for the transcription factor Gata3 in the development of IL-5 $^{+}$ and IL-13 $^{+}$ type 2 innate lymphoid cells. *Proc*

Natl Acad Sci U S A **110**:10240-10245

11. **Scripture-Adams D. D.**, Damle S. S., Li L., Elihu K. J., Qin S., Arias A. M., Butler R. R., Champhekar A., Zhang J. A., Rothenberg E. V. (2014) GATA-3 dose-dependent checkpoints in early T cell commitment. *J Immunol* **193**:3470-3491
12. **Hosoya-Ohmura S.**, Lin Y. H., Herrmann M., Kuroha T., Rao A., Moriguchi T., Lim K. C., Hosoya T., Engel J. D. (2011) An NK and T cell enhancer lies 280 kilobase pairs 3' to the gata3 structural gene. *Mol Cell Biol* **31**:1894-1904
13. **Ohmura S.**, Mizuno S., Oishi H., Ku C. J., Hermann M., Hosoya T., Takahashi S., Engel J. D. (2016) Lineage-affiliated transcription factors bind the Gata3 Tce1 enhancer to mediate lineage-specific programs. *J Clin Invest* **126**:865-878
14. **Kasal D. N.**, Liang Z., Hollinger M. K., O'Leary C. Y., Lisicka W., Sperling A. I., Bendelac A. (2021) A Gata3 enhancer necessary for ILC2 development and function. *Proc Natl Acad Sci U S A* **118**
15. **Furuya H.**, Toda Y., Iwata A., Kanai M., Kato K., Kumagai T., Kageyama T., Tanaka S., Fujimura L., Sakamoto A., *et al.* (2024) Stage-specific GATA3 induction promotes ILC2 development after lineage commitment. *Nat Commun* **15**:5610
16. **Kumagai T.**, Iwata A., Furuya H., Kato K., Okabe A., Toda Y., Kanai M., Fujimura L., Sakamoto A., Kageyama T., *et al.* (2024) A distal enhancer of GATA3 regulates Th2 differentiation and allergic inflammation. *Proc Natl Acad Sci U S A* **121**:e2320727121
17. **Chan W. F.**, Coughlan H. D., Iannarella N., Smyth G. K., Johanson T. M., Keenan C. R., Allan R. S. (2021) Identification and characterization of the long noncoding RNA Dreg1 as a novel regulator of Gata3. *Immunol Cell Biol* **99**:323-332
18. **Chan W. F.**, Coughlan H. D., Chen Y., Keenan C. R., Smyth G. K., Perkins A. C., Johanson T. M., Allan R. S. (2022) Activation of stably silenced genes by recruitment of a synthetic de-methylating module. *Nat Commun* **13**:5582
19. **Zhong C.**, Zheng M., Cui K., Martins A. J., Hu G., Li D., Tessarollo L., Kozlov S., Keller J. R., Tsang J. S., *et al.* (2020) Differential Expression of the Transcription Factor GATA3 Specifies Lineage and Functions of Innate Lymphoid Cells. *Immunity* **52**:83-95.
20. **Shih H. Y.**, Sciume G., Mikami Y., Guo L., Sun H. W., Brooks S. R., Urban J. F., Davis F. P., Kanno Y., O'Shea J. J. (2016) Developmental Acquisition of Regulomes Underlies Innate Lymphoid Cell Functionality. *Cell* **165**:1120-1133
21. **Mielke L. A.**, Groom J. R., Rankin L. C., Seillet C., Masson F., Putoczki T., Belz G. T. (2013) TCF-1 controls ILC2 and NKp46+RORgammat+ innate lymphocyte differentiation and protection in intestinal inflammation. *J Immunol* **191**:4383-4391
22. **Yang Q.**, Li F., Harly C., Xing S., Ye L., Xia X., Wang H., Wang X., Yu S., Zhou X., *et al.* (2015) TCF-1 upregulation identifies early innate lymphoid progenitors in the bone marrow. *Nat Immunol* **16**:1044-1050
23. **Harly C.**, Kenney D., Ren G., Lai B., Raabe T., Yang Q., Cam M. C., Xue H. H., Zhao K., Bhandoola A. (2019) The transcription factor TCF-1 enforces commitment to the innate lymphoid cell lineage. *Nat Immunol* **20**:1150-1160
24. **Ren G.**, Lai B., Harly C., Baek S., Ding Y., Zheng M., Cao Y., Cui K., Yang Y., Zhu J., *et al.* (2022) Transcription factors TCF-1 and GATA3 are key factors for the epigenetic priming of early innate lymphoid progenitors toward distinct cell fates. *Immunity* **55**:1402-1413.
25. **Ercolano G.**, Wyss T., Salome B., Romero P., Trabanelli S., Jandus C. (2020) Distinct and shared gene expression for human innate versus adaptive helper lymphoid cells. *J Leukoc Biol* **108**:723-737
26. **Chen H. V.**, Lorenzini M. H., Lavalley S. N., Sajeev K., Fonseca A., Fiaux P. C., Sen A., Luthra I., Ho A. J., Chen A. R., *et al.* (2023) Deletion mapping of regulatory elements for GATA3 in T cells reveals a distal enhancer involved in allergic diseases. *Am J Hum Genet* **110**:703-714

27. Laurie S. J., Foster J. P., Bruce D. W., Bommasamy H., Kolupaev O. V., Yazdimamaghani M., Pattenden S. G., Chao N. J., Sarantopoulos S., Parker J. S., *et al.* (2024) Type II innate lymphoid cell plasticity contributes to impaired reconstitution after allogeneic hematopoietic stem cell transplantation. *Nat Commun* **15**:6000
28. Statello L., Guo C. J., Chen L. L., Huarte M. (2021) Gene regulation by long non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol* **22**:96-118
29. Kadel S., Ainsua-Enrich E., Hatipoglu I., Turner S., Singh S., Khan S., Kovats S. (2018) A Major Population of Functional KLRG1(-) ILC2s in Female Lungs Contributes to a Sex Bias in ILC2 Numbers. *Immunohorizons* **2**:74-86
30. Bolger A. M., Lohse M., Usadel B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**:2114-2120
31. Kim D., Paggi J. M., Park C., Bennett C., Salzberg S. L. (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**:907-915
32. Liao Y., Smyth G. K., Shi W. (2019) The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res* **47**:e47
33. Mudge J. M., Carbonell-Sala S., Diekhans M., Martinez J. G., Hunt T., Jungreis I., Loveland J. E., Arnan C., Barnes I., Bennett R., *et al.* (2025) GENCODE 2025: reference gene annotation for human and mouse. *Nucleic Acids Res* **53**:D966-D975
34. Robinson M. D., McCarthy D. J., Smyth G. K. (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139-140
35. Danecek P., Bonfield J. K., Liddle J., Marshall J., Ohan V., Pollard M. O., Whitwham A., Keane T., McCarthy S. A., Davies R. M., *et al.* (2021) Twelve years of SAMtools and BCFtools. *Gigascience* **10**
36. Kent W. J., Zweig A. S., Barber G., Hinrichs A. S., Karolchik D. (2010) BigWig and BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**:2204-2207
37. Thorvaldsdottir H., Robinson J. T., Mesirov J. P. (2013) Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* **14**:178-192
38. Grant C. E., Bailey T. L., Noble W. S. (2011) FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**:1017-1018
39. Kulakovskiy I. V., Vorontsov I. E., Yevshin I. S., Sharipov R. N., Fedorova A. D., Rumynskiy E. I., Medvedeva Y. A., Magana-Mora A., Bajic V. B., Papatsenko D. A., *et al.* (2018) HOCOMOCO: towards a complete collection of transcription factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res* **46**:D252-D259
40. Gury-BenAri M., Thaïss C. A., Serafini N., Winter D. R., Giladi A., Lara-Astiaso D., Levy M., Salame T. M., Weiner A., David E., *et al.* (2016) The Spectrum and Regulatory Landscape of Intestinal Innate Lymphoid Cells Are Shaped by the Microbiome. *Cell* **166**:1231-1246.

Peer reviews

Reviewer #1 (Public review):

Summary:

This study examines the role of the long non-coding RNA Dreg1 in regulating Gata3 expression and ILC2 development. Using Dreg1-deficient mice, the authors show a selective loss of ILC2s but not T or NK cells, suggesting a lineage-specific requirement for Dreg1. By integrating public chromatin and TF-binding datasets, they propose a Tcf1-Dreg1-Gata3 regulatory axis. The topic is relevant for understanding epigenetic regulation of ILC differentiation.

Strengths:

- (1) Clear in vivo evidence for a lineage-specific role of Dreg1.
- (2) Comprehensive integration of genomic datasets.
- (3) Cross-species comparison linking mouse and human regulatory regions.

Weaknesses:

- (1) Mechanistic conclusions remain correlative, relying on public data.
- (2) Lack of direct chromatin or transcriptional validation of Tcf1-mediated regulation.
- (3) Human enhancer function is not experimentally confirmed.
- (4) Insufficient methodological detail and limited mechanistic discussion.

<https://doi.org/10.7554/eLife.109408.1.sa1>

Reviewer #2 (Public review):

The authors investigate the role of the long non-coding RNA Dreg1 for the development, differentiation, or maintenance of group 2 ILC (ILC2). Dreg1 is encoded close to the Gata3 locus, a transcription factor implicated in the differentiation of T cells and ILC, and in particular of type 2 immune cells (i.e., Th2 cells and ILC2). The center of the paper is the generation of a Dreg1-deficient mouse. While Dreg1^{-/-} mice did not show any profound ab T or gd T cell, ILC1, ILC3, and NK cell phenotypes, ILC2 frequencies were reduced in various organs tested (small intestine, lung, visceral adipose tissue). In the bone marrow, immature ILC2 or ILC2 progenitors were reduced, whereas a common ILC progenitor was overrepresented, suggesting a differentiation block. Using ATAC-seq, the authors find that the promoter of Dreg1 is open in early lymphoid progenitors, and the acquisition of chromatin accessibility downstream correlates with increased Dreg1 expression in ILC2 progenitors. Examining publicly available Tcf1 CUT&Run data, they find that Tcf1 was specifically bound to the accessible sites of the Dreg1 locus in early innate lymphoid progenitors. Finally, the syntenic region in the human genome contains two non-coding RNA genes with an expression pattern resembling mouse Dreg1.

The topic of the manuscript is interesting. However, there are various limitations that are summarized below.

- (1) The authors generated a new mouse model. The strategy should be better described, including the genetic background of the initially microinjected material. How many generations was the targeted offspring backcrossed to C57BL/6J?
- (2) The data is obtained from mice in which the Dreg1 gene is deleted in all cells. A cell-intrinsic role of Dreg1 in ILC2 has not been demonstrated. It should be shown that Dreg1 is required in ILC2 and their progenitors.
- (3) The data on how Dreg1 contributes to the differentiation and or maintenance of ILC2 is not addressed at a very definitive level. Does Dreg1 affect Gata3 expression, mRNA stability, or turnover in ILC2? Previous work of the authors indicated that knockdown of Dreg1 does not affect Gata3 expression (PMID: 32970351).
- (4) How Dreg1 exactly affects ILC2 differentiation remains unclear.

<https://doi.org/10.7554/eLife.109408.1.sa0>