

Bactabolize: A tool for high-throughput generation of bacterial strain-specific metabolic models

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

September 25, 2023 (this version)

Reviewed preprint version 1

May 30, 2023

Sent for peer review

March 7, 2023

Posted to bioRxiv

February 27, 2023

Ben Vezina , Stephen C. Watts, Jane Hawkey, Helena B. Cooper, Louise M. Judd, Adam W. J. Jenney, Jonathan M. Monk, Kathryn E. Holt, Kelly L. Wyres 

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia • Microbiology Unit, Alfred Health, Melbourne, Victoria, Australia • Department of Bioengineering, University of California, San Diego, CA, United States of America • Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Metabolic capacity can vary substantially within a bacterial species, leading to ecological niche separation, as well as differences in virulence and antimicrobial susceptibility. Genome-scale metabolic models are useful tools for studying the metabolic potential of individuals, and with the rapid expansion of genomic sequencing there is a wealth of data that can be leveraged for comparative analysis. However, there exist few tools to construct strain-specific metabolic models at scale.

Here we describe Bactabolize (github.com/kelwyres/Bactabolize), a reference-based tool which rapidly produces strain-specific metabolic models and growth phenotype predictions. We describe a pan reference model for the priority antimicrobial-resistant pathogen, *Klebsiella pneumoniae* (github.com/kelwyres/KpSC-pan-metabolic-model), and a quality control framework for using draft genome assemblies as input for Bactabolize.

The Bactabolize-derived model for *K. pneumoniae* reference strain KPPR1 performed comparatively or better than currently available automated approaches CarveMe and gapseq across 507 substrate and 2317 knockout mutant growth predictions. Novel draft genomes passing our systematically-defined quality control criteria resulted in models with a high degree of completeness ($\geq 99\%$ genes and reactions captured compared to models derived from matched complete genomes) and high accuracy (mean 0.97, $n=10$).

We anticipate the tools and framework described herein will facilitate large-scale metabolic modelling analyses that broaden our understanding of diversity within bacterial species and inform novel control strategies for priority pathogens.

eLife assessment

This study presents Bactabolize, a **valuable** tool for the rapid genome-scale reconstruction of bacteria and the prediction of growth phenotypes. Using validated methodology, the tool relies on a reference pan-genome model to create strain-specific draft metabolic models, as demonstrated in this study using *Klebsiella pneumoniae*. While the evidence in this specific case is **solid**, validation across diverse bacterial species is yet to be confirmed.

Introduction

Bacteria exhibit metabolic diversity and can utilise a broad range of substrates for growth. It has become clear amongst pathogens that there is an intertwined relationship between metabolism and nutrient usage with virulence and antimicrobial resistance (1–7). Comparative analyses of metabolic profiles (e.g. substrate usage) are key to fully understanding these relationships. Traditionally, these profiles have been assessed via phenotypic growth on a limited number of substrates, such as those used to delineate between species (8–10) which form the basis of a number of commercial products for species identification. However, these methods are not sufficiently discriminatory for in-depth comparisons within species, and alternative approaches such as the Omnilog Phenotype MicroArray system (Biolog) are too expensive and/or labour intensive for application to large numbers of isolates. Similarly, probing of essential metabolism-associated genes via transposon mutant libraries (e.g. to identify novel virulence factors and therapeutic targets) (4, 11, 12) cannot be easily scaled across diverse bacterial populations.

Genome-scale metabolic models or metabolic reconstructions are a computational approach to analysing the metabolic potential of an organism, within which the entire biochemical network is represented as a stoichiometric matrix (13). Metabolic models are constructed programmatically, but typically informed and at least partially validated using phenotypic growth data (14–16). Once constructed, they can be run through simulations and analysed under various contexts, such as *in silico* growth experiments (Flux Balance Analysis [FBA]) to predict substrate usage profiles (17), evaluate the impact of single gene knockouts on growth (14, 18), and identify metabolic chokepoints for drug targets (19), among others. Traditionally, metabolic models are strain-specific (i.e. each model represents a unique individual <http://bigg.ucsd.edu/models>) and may not be applicable to other isolates due to unrepresented genetic diversity.

We recently described 37 curated strain-specific models for the *Klebsiella pneumoniae* Species Complex (KpSC) (14) comprised of *K. pneumoniae* and its close relatives (20). These organisms are a common cause of healthcare-associated infections world-wide, and among the World Health Organization's priority antimicrobial resistant pathogens (21). KpSC are highly diverse and gene content can differ substantially between strains (22, 23). Accordingly, our models varied in terms of gene and reaction content, resulting in variable growth substrate usage profiles and metabolic redundancy (14). Similar variation has also been described in other key bacterial pathogens e.g. *Escherichia coli* (24), *Salmonella enterica* (25), *Staphylococcus aureus* (26) and *Pseudomonas aeruginosa* (27). This is highly relevant to the use of metabolic models for the exploration of virulence and antimicrobial resistance, and for the identification of novel drug targets. Therefore, such works should seek to include multiple strain-specific models, and in some cases 100s-1000s of models may be required to accurately represent population diversity (22, 28, 29).

There are several open source tools currently available that can rapidly produce strain-specific metabolic models, including CarveMe (30), gapseq (31), ModelSEED (32) and KBase (33) (see the recent review by Mendoza and colleagues for comparative descriptions (34)), as well as a recently published modelling and analysis pipeline, ChiMera, which leverages CarveMe for model construction (35). In their systematic analysis Mendoza *et al.* indicated CarveMe and ModelSEED to be of particular interest for large-scale studies due to their speed and model quality (34) (note that gapseq was published later and therefore not included in this review). KBase implements ModelSEED (32) as a web interface application, limiting its utility for high-throughput analysis of 100s – 1000s of bacterial genomes. CarveMe is a command line application; it is open source but is dependent on commercial solvers such as CPLEX (free for academic use). However, its use of a universal reference model may limit specificity of strain-specific models (36), and result in overestimation of model genes. These limitations can be overcome by manual curation of the output models, but such curation is highly labour intensive and not suitable for high-throughput analyses. Furthermore, the CarveMe database (BiGG universal_model) appears to be no longer actively maintained, meaning that there is no opportunity to integrate novel structural and/or biochemical data as these become available in the literature (as discussed in COBRA community forums). gapseq is a recently published command line tool which leverages an independent universal database (31). In their comparative analyses, the gapseq authors demonstrate superior accuracy to both CarveMe and ModelSEED; however the concerns about the specificity of universal models remain and it has been reported that model construction takes considerable time (several hours in some cases (37)).

Here, we present Bactabolize (available at <https://github.com/kelwyres/Bactabolize>), an easy-to-use tool which allows scalable production of strain-specific draft metabolic models and prediction of growth phenotypes in under three minutes per input genome. Bactabolize builds upon the reference-based model reconstruction approach described by Norsigian *et al.* (36), leveraging the COBRApy framework (38) and BiGG nomenclature (39). We present a pan-metabolic reference model for the *KpSC* (derived from our 37 curated strain-specific models (14)), and describe an exemplar quality control framework for the application of Bactabolize to *KpSC* draft genome assemblies. We show that Bactabolize can rapidly produce strain-specific models from draft genomes with a high degree of completeness (as compared to models generated from completed genome assemblies), resulting in highly accurate growth predictions that match or exceed the accuracy of models from CarveMe and manual curation efforts.

Results

Description of Bactabolize

Bactabolize is written in Python 3 and utilises the metabolic modelling library COBRApy (38). Bactabolize has four main commands:

- i) Draft model generation (`draft_model` command), which generates a strain-specific draft metabolic reconstruction ('model') using the approach outlined previously (36), and uses gap-filling to identify any missing reactions required to simulate growth in the user-specified conditions (`draft_model` command), which generates a strain-specific draft metabolic reconstruction ('model') using the approach outlined previously (36), and uses gap-filling to identify any missing reactions required to simulate growth in the user-specified conditions
- ii) Patching incomplete models (`patch_model` command) by the addition of missing reactions e.g. those identified by the automated gap-filling process (`patch_model` command) by the addition of missing reactions e.g. those identified by the automated gap-filling process

- iii) Substrate usage analysis via Flux Balance Analysis (FBA) (*fba* command) to predict growth outcomes for a specified range of substrates supported by the model(s) **Figure 1** ([fba](#) command) to predict growth outcomes for a specified range of substrates supported by the model(s) **Figure 1** ([fba](#) command).

Additional processing scripts are provided alongside Bactabolize to improve model metadata annotation (*improve_model_annotations.py*), convert models generated using KBase and ModelSEED to Bactabolize/BiGG-compatible format (*SEED_to_BiGG_model_convert.sh*), generate network graph files from models (*model_to_network_graph.py*) and merging output FBA profiles (*merge_fba_profiles_longtable.sh*). Full documentation, including example code and test data are available at the Bactabolize code repository (<https://github.com/kelwyres/Bactabolize>).

For draft model construction, Bactabolize requires users to provide an input assembly (annotated or unannotated FASTA or Genbank format respectively), a reference model (JSON format) and the corresponding reference sequence data (gene and protein sequences in two separate multi-fasta files or a single Genbank annotation in a .gbk file) (**Figure S1** [fba](#) command). For optimum results we suggest using a pan-model that captures as much diversity as possible for the target species or group of interest, because Bactabolize's reconstruction method is reductive i.e. each output strain-specific model will include only genes, reactions and metabolites that are present in the reference or a subset thereof (although novel genes, reactions and metabolites can be added via manual curation).

If the input assembly is unannotated, Bactabolize will identify coding sequences using Prodigal ([40](#) [fba](#)) but will otherwise honour the existing coding sequence (CDS) notations and optionally use Prodigal to search for additional CDS. Draft genome-scale metabolic models are output in both SBML v3.1 ([41](#) [fba](#)) and JSON formats (one pair of files for each independent strain-specific model), along with an optional MEMOTE quality report ([42](#) [fba](#)). Bactabolize will identify orthologs in the input genome(s) compared to the reference sequence data using Bi-directional BLAST ([43](#) [fba](#)) Best Hits (BBH) ([44](#) [fba](#)) using BLAST+ ([36](#) [fba](#)). Users can parameterise the ortholog finding settings (coverage and identity thresholds) for BBH. Alternatively, there is the option of using protein similarity to identify orthologs instead of identity.

Once a draft model has been constructed, it is validated via a simulated growth experiment on user-input choice of media and atmosphere (aerobic or anerobic). Predefined media include BG11 (Gibco), M9 + glucose ([36](#) [fba](#)), nutrient media ([45](#) [fba](#)), Luria-Bertani (LB) ([45](#) [fba](#)), Tryptic Soy (TSA) ([45](#) [fba](#)), TSA + sheep blood ([45](#) [fba](#)), LB as specified by the CarveMe developers ([30](#) [fba](#)), Chemically Defined Medium (CDM)-like ([34](#) [fba](#)), Plantarum Minimal Medium (PMM) PMM5-like ([34](#) [fba](#)) and PMM7-like ([34](#) [fba](#)). Users can also define custom media as Bactabolize supports several complex media ingredients, including peptone (peptic digest of bovine and porcine tissue) ([46](#) [fba](#)-[48](#) [fba](#)), tryptone (pancreatic digest of casein) ([46](#) [fba](#), [48](#) [fba](#), [49](#) [fba](#)), soy peptone/soytone (digest of soy meal) ([46](#) [fba](#), [48](#) [fba](#), [50](#) [fba](#), [51](#) [fba](#)), yeast extract ([52](#) [fba](#)-[57](#) [fba](#)) and beef extract ([46](#) [fba](#), [48](#) [fba](#)). If the model fails to simulate growth, gap-filling is performed to indicate missing reactions. Users can add these reactions to a patch JSON file and optionally use the *patch_model* command to correct the model (**Figure S2** [fba](#)). Bactabolize uses a conservative gap-filling approach that only adds the minimum number of reactions to enable growth under the chosen conditions. We recommend testing the models in minimal media and atmosphere expected to support growth for all isolates of the species of interest, unless the user has access to matched phenotypic data demonstrating growth for individual isolates in specific conditions. Aggressive gap-filling will effectively homogenise the models and should be avoided if the goal is to understand the underlying strain diversity.

Substrate usage analysis (the *fba* command) is performed iteratively for each possible carbon, nitrogen, sulfur and phosphorus substrate supported by the model(s) (**Figure S3** [fba](#)), by replacing the default substrate in the user specified growth medium (specified in the *fba_spec* JSON file). For example, in M9 media the default substrates are glucose (carbon), ammonia (nitrogen), sulphate




Figure 1

Simplified overview of Bactabolize's main commands. In pink is the *draft_model* command, which builds a draft strain-specific metabolic model using an input reference model and an input target assembly (approach adapted from (36)). If the model fails to simulate growth, Bactabolize will attempt automated gap-filling and produce a model patch file. The *patch_model* command (orange) allows the addition of missing reactions to produce a valid draft model that can simulate growth in a user-specified growth environment. A functioning model can be passed to the *fba* command (yellow), which performs Flux Balance Analysis to simulate growth in the user specified conditions, across all carbon, nitrogen, phosphorus and sulfur metabolite sources supported by the model under aerobic and anerobic conditions. The *sgk* command (blue) shows the Single Gene Knockout analysis, which outputs a predicted phenotype. User inputs and outputs are shown in white boxes while Bactabolize commands are shown inside the grey box.

(sulfur) and phosphate (phosphor). Each substrate can be tested in aerobic and/or anaerobic conditions. Growth prediction output is recorded in a tab delimited file (one per strain). The `merge_fba_profiles_longtable.sh` helper script will combine the outputs for multiple strains into a single file for downstream analysis.

The growth impacts of single-gene knockout mutations can be simulated via the `sgk` command (Figure S4). Bactabolize will iterate through every gene in the model, temporarily removing it and its associated reactions (unless they are also associated with another gene) and running FBA to simulate growth in the user-specified conditions. The output is comparable to single-gene knockout studies such as transposon mutagenesis and can be used to probe gene essentiality.

KpSC pan-metabolic reference model

We constructed a species complex-specific pan-metabolic reference model by combining a collection of 37 manually curated models for which we have previously demonstrated high accuracy (range 88.3%–96.8% for prediction of 94 distinct growth phenotypes (14)). These models represent a diverse collection of *KpSC* (14) (including at least one each of the seven major taxa in the complex; *K. pneumoniae*, *Klebsiella variicola* subsp *variicola*, *Klebsiella variicola* subsp *tropica*, *Klebsiella quasipneumoniae* subsp *quasipneumoniae*, *Klebsiella quasipneumoniae* subsp *similipneumoniae*, *Klebsiella quasivariicola*, *Klebsiella africana*). The combined pan-model, known as *KpSC*-pan v1, comprises a total of 1265 distinct genes, 2319 reactions and 1696 metabolites, and is available at github.com/kelwyres/KpSC-pan-metabolic-model.

Performance comparison

We compared the output and performance of Bactabolize to the two previously published tools that can support high-throughput analyses i.e. CarveMe (30) and gapseq (31). To aid interpretation in the context of community standard approaches, we also include a comparison to the popular web-based reconstruction tool, KBase (ModelSEED), and a manually curated metabolic reconstruction of *K. pneumoniae* strain KPPR1 (also known as VK055 and ATCC 43816, metabolic model named iKp1289) (15). This isolate was chosen as there is a completed genome sequence (Genbank accession: CP009208), single-source growth phenotype (15) and single-gene knockout growth essentiality data available (58). *De novo* draft models for strain KPPR1 were built using; i) Bactabolize with the *KpSC* pan v1 reference; ii) CarveMe, with its universal reference model (CarveMe universal); iii) CarveMe, with *KpSC*-pan v1 reference (CarveMe *KpSC* pan); iv) gapseq; and v) KBase (ModelSEED). Importantly, neither *K. pneumoniae* KPPR1 nor its genetic lineage (7 gene multi-locus sequence type, ST493), are represented in the *KpSC* pan-reference model, meaning these benchmarking comparisons were on equal footing. Subsequently, each model was used to predict growth phenotypes; i) in M9 minimal media with different sole sources of carbon, nitrogen, phosphorus and sulfur; and ii) for all possible single gene knockouts in LB under aerobic conditions. The predicted phenotypes were compared directly to the published phenotype data.

Among the high-throughput approaches, the Bactabolize draft model captured fewer genes and reactions ($n = 1233$ and 2307 , respectively) than the gapseq ($n = 1489$ and 3186 , respectively) and CarveMe universal models ($n = 1960$ and 2857 , Figure 2A). The Bactabolize and CarveMe universal models contained similar numbers of metabolites (1696 vs 1737 , respectively), while the gapseq model contained many more ($n = 2519$). Notably, the initial gapseq model was gap-filled by simulation of growth in M9 minimal media with glucose (consistent with Bactabolize and CarveMe media recipes), which resulted in the addition of 31 reactions to the draft model. This is in contrast to CarveMe and Bactabolize-generated models which produced biomass in M9 minimal media plus glucose without additional gap-filling. The CarveMe *KpSC* pan model captured considerably more genes than any of the other models ($n = 2407$), but these were associated with many fewer unique reactions and metabolites (1206 and 825 , respectively). Upon further investigation we determined

that this method resulted in the over prescription of gene reaction rules (GPRs) to multiple reactions (mean 2.2 GPRs per reaction when compared to Bactabolize using the same pan reference model: 1.94 GPRs per reaction).

Figures S5 [↗](#) and **S6** show the overlaps of metabolites and reactions between the high-throughput reconstruction methods after processing with MetaNetX ([59](#) [↗](#)) to standardise the reaction and metabolite nomenclatures (excluding CarveMe pan for simplicity and given the likely problems of reaction oversubscription). The majority of the reactions included in the Bactabolize model were conserved in either the CarveMe universal model ($n = 1225$, 53.2%), gapseq model ($n = 54$, 2.3%) or both ($n = 665$, 28.9%). The reaction overlap was skewed to the CarveMe universal model which shared 1225 reactions that were conserved in the Bactabolize model but absent from the gapseq model. Notably, the gapseq model contained a large number (2200) of unique reactions (70.4% of those in the model). Similarly, the vast majority of metabolites in the Bactabolize model were conserved in one or both of the other models ($n = 917$, 85.6%). However, it is likely that true overlaps between methods are underrepresented due to the different reaction identifiers and chemical synonyms used within the BiGG (Bactabolize, CarveMe) vs ModelSEED nomenclatures (gapseq), which are difficult to harmonise in an automated manner even after the application of MetaNetX.

In comparison to the low-throughput reconstruction approaches, the Bactabolize model contained a similar number of genes, reactions and metabolites to the manually curated model ($n = 1289$, 2484 and 1827, respectively) and many more than the KBase model (72, 544 and 534, respectively). This is likely due to the low number of metabolism-associated genes identified by KBase, which has impacted the associated reaction and metabolite data.

MEMOTE scores, (produced by the MEMOTE report ([42](#) [↗](#))) indicate the quality of the model metadata annotations, with the scores ranging between 0 – 100%. These provide a measure of model portability and the level of connected databases available to support the metabolite, reaction and genetic information represented in the model, but bear no reflection on model accuracy. Bactabolize performs on the lower end, with CarveMe universal and gapseq performing the best (**Figure 2B** [↗](#)). The KBase model appears to perform well in this regard, however this is due to the low number of genes, reactions and metabolites included in the model. Bactabolize using the *KpSC*-pan model outperforms the model propagation mode of CarveMe using the same reference model (**Figure 2B** [↗](#)). Work is ongoing to improve the metadata annotations in the *KpSC*-pan reference model, to support large-scale model propagation.

We assessed the performance of each model for *in silico* prediction of growth phenotypes compared to the previously published experimental data ([15](#) [↗](#)). Accuracy, sensitivity, specificity, precision and F1 scores were calculated ([60](#) [↗](#)). Note that the specific set of growth substrates and gene knockouts that can be simulated is determined by the sets of genes and metabolites captured by each model and is therefore model-dependent (**Data S1 and S2**). Among those with matched experimental phenotype data, the Bactabolize and CarveMe universal models were able to predict growth for a greater number of carbon, nitrogen, phosphorous and sulfur substrates than gapseq, CarveMe *KpSC* pan, KBase and iKp1289 models (**Figure 2C** [↗](#), **Data S1**). While the CarveMe universal model had the highest number of true-positive growth predictions overall ($n = 132$ of 617 total predictions), it also had a comparably high number of false-positive predictions ($n = 39$ of 617 total predictions, **Figure 2D** [↗](#)). Similarly, the gapseq and iKp1289 models resulted in 31 (262 total predictions) and 50 (513 total predictions) false-positive predictions, respectively. In contrast, the Bactabolize model had fewer false-positive predictions ($n = 21$ of 505 total predictions) alongside a high number of true-positive predictions ($n = 117$ of 505 total predictions), resulting in the highest overall accuracy metrics (**Figure 2E** [↗](#), **Data S1**). The KBase model was a notable outlier, associated with a high number of false-negative predictions ($n = 31$ of 103 total




Figure 2

K. pneumoniae KPPR1 metabolic model benchmarking comparisons. **A)** Counts of model features; genes, metabolites and reactions captured by each model. Exchanges refers to number of exchange reactions, a subset of reactions involved in substrate uptake, which determine the number of distinct growth substrates for which phenotypes can be predicted with the model. **B)** MEMOTE scores indicating the richness of annotations and metadata for metabolic model features according to database outlinks. SBO refers to score of Systems Biology Ontology (SBO), a controlled vocabulary for systems biology. Consistency refers to the score of stoichiometric consistency and chemical formulae annotation. Total refers to total MEMOTE score, as a combination of all previous scores, and is shown in bold. **C)** Counts of carbon, nitrogen, phosphorus and sulfur growth substrates that can be simulated by models and for which matched phenotypes were available for comparison (15). Hatched columns indicate the total number of substrates for which phenotypic data for *K. pneumoniae* KPPR1 were described (15). **D)** and **E)** Accuracy metrics for predicted to true phenotypes for the growth substrates shown in D and E, respectively. False-negatives, true-negatives, false-positives and true-positives are coloured as shown in legend. **F)** and **G)** Accuracy metrics for the KPPR1 single-gene knockout mutant library described in (58) shown in F and G, respectively. Numbers of true positives and false positives are shown to the left of the respective columns.

predictions) and low false-positive predictions ($n = 3$ of 103 total predictions), presumably resulting from the very low number of genes and reactions included in the model, driving low sensitivity and accuracy.

The gene essentiality results showed that gapseq produced the highest absolute number of true-positive gene essentiality predictions ($n = 79$ of 1403), followed by Bactabolize *KpSC* pan ($n = 44$ of 1220 total predictions), then CarveMe universal ($n = 39$ of 1951 total predictions). CarveMe universal had the largest number of true-negatives by a wide margin ($n = 1599$ of 1951 total predictions), followed by gapseq ($n = 1085$ of 1403 total predictions), then Bactabolize *KpSC* pan ($n = 939$ of 1220 total predictions), driving their high accuracies (83.96%, 82.96% and 80.57%, respectively). The Bactabolize model was associated with the greatest overall precision and specificity (**Figures 2F & 2G**) while the gapseq model resulted in the highest F1-score and sensitivity.

While model features and accuracy are essential metrics for comparison, computation time is also a key consideration for high-throughput analyses. We recorded the time required for each tool to build draft models for 10 of the completed *KpSC* genomes used in the quality control framework (see below) on a high-performance computing cluster (Intel Xeon Gold 6150 CPU @ 2.70GHz and 155 GB of requested memory on a CentOS Linux release 7.9.2009 environment). CarveMe *KpSC* pan was the fastest with a mean of 20.04 (range 19.90 - 20.18) seconds, followed by CarveMe universal at 30.28 (range 29.20 - 31.80) seconds, then Bactabolize *KpSC* pan at 98.05 (range 92.19 - 100.4) seconds. KBase took 183.50 (range 120.00 - 338.00) seconds per genome via batch analysis, including genome upload time and queuing. gapseq took 5.46 (range 4.55 - 6.28) hours to produce draft models (not including the required gap-filling), consistent with previous reports (37).

Quality control framework for input genome assemblies

There are now thousands of bacterial genomes available in public databases, the majority of which are in draft form, comprising 10s to 1000s of assembly contigs. This fragmentation of the genome is caused by repetitive sequences that cannot be resolved by the assembly algorithm and/or sequence drop-out. The latter can result in the loss of genetic information such that some portion of genes present in the underlying genome are lost from the genome assembly (either completely or partially). This in turn, poses a limitation for the reconstruction of metabolic models using these assemblies, since most published approaches use sequence searches to predict the presence/absence of genes and their associated enzymatic reactions. Therefore, if we are to use public genome data for high-throughput metabolic modelling studies, it is essential to evaluate the expected model accuracies and understand the minimum input genome quality requirements.

Here we performed a systematic analysis leveraging our published curated *KpSC* models ($n=37$, (14)), which were generated using completed genome sequences and were therefore considered to represent ‘complete’ models for which the underlying genome sequence contains all genes that are truly present in the genome (note the biological accuracy of these models was reported previously (14) and is not the subject of the current study). We randomly subsampled the corresponding Illumina read sets to various depths (10 – 100x, increments of 10) in triplicate and generated draft assemblies that were passed to Bactabolize for generation of draft metabolic models (**Data S3**). Due to low read depth ($\leq 30x$), two isolates were removed from this analysis. Additionally, ten 10x depth read samples failed to produce assemblies, leaving 1040 draft genomes for analysis. The resulting draft metabolic models were compared to the complete models to; i) determine the proportions of complete model genes and reactions captured in the draft models; and ii) compare 846 *in silico* aerobic growth predictions in M9 minimal media, where growth on 266 carbon, 153 nitrogen, 59 phosphorus and 25 sulfur sources were examined. Substrates containing multiple elements were tested as sole sources of each element independently and in combination, e.g. 1,5-Diaminopentane was tested as a sole carbon, sole nitrogen and sole carbon plus nitrogen source.

As expected, assembly quality generally increased with increasing sequencing depth i.e. assemblies generated from higher depth read sets were associated with higher N50 values, fewer contigs and fewer assembly graph dead-ends, although the rate of improvement drastically declined beyond 40-50x depth (**Figure S7** [↗](#), **Data S3**). We noted that it was rare for draft models to capture 100% of model genes and reactions (just 420 of all 1040 draft assemblies were associated with models that captured 100% model genes) (**Data S3**, **Figure S8** [↗](#)), even when using the highest quality draft genomes. However, $\geq 99\%$ of genes and reactions were commonly captured, which plateaued from 40x depth onwards (**Figure S8** [↗](#)). Therefore, we sought to evaluate whether $\geq 99\%$ model capture would produce functionally accurate models.

We used FBA to simulate substrate growth profiles for the 40x depth assemblies, representing a sequencing depth that can be routinely achieved with standard Illumina library preparations. All but one assembly triplicate set (isolate SB4767 98% gene capture, 99% reaction capture) captured $\geq 99\%$ but $\leq 100\%$ model genes and/or reactions. The substrate growth profiles were then compared to those of the complete models. The vast majority of draft models produced accurate growth predictions; 102 of 108 models resulted in predictions with 100% concordance to those from the corresponding complete models. Three models for *K. quasipneumoniae similipneumoniae* isolate SB164 resulted in predictions with a mean of 99.8% concordance. The remaining three models were for isolate SB4767 and resulted in mean of 80.4% concordance. Notably, these models were those representing $< 99\%$ gene capture. Together, these data suggest that draft models capturing $\geq 99\%$ of the complete model genes/reactions generate highly accurate growth predictions and that these capture rates can be readily achieved from draft genome assemblies.

We investigated the relationships between assembly quality metrics and model gene/reaction capture in more detail. Variation in assembly graph dead-ends accounted for the greatest amount of variation in model capture, closely followed by raw contig counts (cubic polynomial fit, R^2 of ≥ 0.98 for graph dead-ends, R^2 of ≥ 0.9 for contig count). A segmented linear model was fitted to N50 length ($R^2 \geq 0.83$), producing a breakpoint at 25153 bp (**Figure 3** [↗](#)).

To further explore the optimum thresholds for assembly metrics, we tallied the number of draft assemblies resulting in $\geq 99\%$ and $< 99\%$ gene and reaction capture at increasing graph dead-end and contig count cut-offs, and decreasing N50 cut-offs. Draft models that captured $\geq 99\%$ of the complete model genes/reactions were considered ‘good’ models, whereas draft models that captured $< 99\%$ of complete model genes/reactions were considered ‘bad’ models. The optimum threshold for assembly graph dead end was determined to be ≤ 200 . At this value, 94.44% of ‘good’ models were captured, and 0% ‘bad’ models. The optimum threshold for contig counts was determined as ≤ 130 contigs at which 67.92% of ‘good’ and 0% ‘bad’ models were captured (**Figure 4** [↗](#)). The optimum threshold for N50 was determined to be ≥ 65000 , at which 94.97% of ‘good’ and 1.71% of ‘bad’ models were captured. The assembly graph dead-end threshold results in comparatively higher sensitivity (i.e. a higher proportion of ‘good’ models pass the threshold) than contig count and comparatively better specificity (i.e. lower proportion of ‘bad’ models pass the threshold) than N50, but the underlying metric information is not universally available because many isolate genomes are deposited in public databases only as assemblies without the associated assembly graph. We therefore recommend a three-tier approach, whereby the assembly graph dead-end criterion is preferenced if available, followed by N50 and then contig count.

Impact of gap-filling models

Of the 901 draft genome assemblies which passed our QC criteria (≤ 200 assembly graph dead ends), 23 of the resulting draft models failed to simulate growth in M9 minimal media with glucose (despite capturing $\geq 99\%$ of the genes and reactions in the corresponding complete models). It is expected that all *KpSC* models should be able to simulate growth on M9 media with glucose as a sole carbon source, as this central metabolism is universal amongst *KpSC*. To replace missing, critical reactions required for growth on M9 with glucose, we investigated model gap-filling using




Figure 3

Scatterplots showing distribution of best performing assembly metrics 'assembly graph dead ends', 'contigs' and 'N50' against model feature capture (genes and reactions). Each point represents the mean values from a single genome (technical triplicate) and is coloured by model quality. 'Good' models capture $\geq 99\%$ of the model metric as compared to the corresponding complete model (shown at each facet), 'Bad' models capture $< 99\%$. Cubic polynomial line plotted for assembly 'graph dead ends', 'contigs', while a segmented linear model was plotted for 'N50'. R^2 is shown on each panel.




Figure 4

Line graphs showing the impact of assembly metric cut-off thresholds on model feature capture ($n = 1040$). 'Good' models which captured $\geq 99\%$ of model features are shown in green, while 'bad' models captured $< 99\%$ model features are shown in gold. The blue dotted line shows the metric cut-off thresholds, to minimize the number of models that capture $< 99\%$ model features and maximise models that capture $\geq 99\%$. Metric cut-off statistics are calculated in intervals of 10 for assembly graph dead ends and contigs, and every 5000 for N50.

the *patch_model* command of Bactabolize. We then assessed the accuracy of the gap-filled models for prediction of growth on the full range of substrates, as compared to the predictions from the corresponding complete models.

Gap filling added 1 – 3 missing reactions to each model, with a median of one, fully restoring biomass production in M9 media with glucose in all but two of the 23 failed models. The missing reactions appeared to be random genes across these 23 genomes, likely due to missing information in these assemblies.

Substrate usage predictions from the 21 successfully gap-filled models were highly accurate, with 18/21 having a prediction concordance of $\geq 99\%$ across all 846 growth conditions (12/21 had 100% concordance) (**Figure S9** [↗](#)). We therefore conclude that models generated for genome assemblies passing our QC criteria, which have been gap-filled to successfully simulate growth on minimal media plus glucose, are suitable for the prediction of growth across a range of substrates.

Predictive accuracy of draft models

We assessed the accuracy of Bactabolize for the construction of draft models for 10 novel *KpSC* clinical isolates, representing five of the major taxa in the complex. We included five isolates for which the associated STs were represented in the *KpSC*-pan v1 model and five isolates with STs that were not represented. Whole genome sequence data were generated on the Illumina platform and draft assemblies generated *de novo*. The resultant assemblies had 0-4 graph dead-ends, N50s of 151958-388486 bp and 83-187 contigs (**Data S4**), within the tiered threshold values.

FBA was performed, and the predicted growth profiles compared to matched phenotypic growth data for 16 carbon sources derived from Vitek GN ID cards. Though the number of tested carbon sources was limited, all were associated with high accuracy metrics (**Figure 5** [↗](#), **Data S4**). As expected, models for isolates with STs represented in the *KpSC*-pan v1 reference performed slightly better (mean accuracy = 0.98) than those for non-represented STs (mean accuracy = 0.95).

Discussion

In this work we described Bactabolize, a pipeline for rapid and scalable production of accurate bacterial strain-specific metabolic models and growth phenotype predictions. We describe a pan-reference model for the *KpSC* and demonstrate that a draft strain-specific model generated *de novo* via Bactabolize using the *KpSC*-pan v1 reference was highly accurate for growth phenotype prediction (85.79% accuracy for substrate usage across 190 substrates, and 80.57% for gene essentiality across 1220 genes). Importantly, we also described a quality control framework for the use of draft genome assemblies as input for metabolic reconstructions. We used a systematic analysis to: i) evaluate the proportion of gene and reaction capture compared to the corresponding ‘completed’ models; ii) define quality control thresholds for input assemblies (three tier approach for *KpSC*; ≤ 200 assembly graph dead ends, followed by ≥ 65000 N50, followed by ≤ 130 contigs); and iii) estimate the accuracy of the resultant growth predictions. While the quality control thresholds and accuracy estimates are specific to *KpSC*, the conceptual framework can be applied to any organism and is essential to support the confident application of metabolic modelling for large-scale genome datasets. We appreciate that assembly graphs may not be available for dead end count, e.g. for draft genome assemblies accessed via public repositories, however we encourage users to include this information in their quality control procedures wherever possible (e.g. using the recently published counter tool available at <https://github.com/rrwick/GFA-dead-end-counter> [↗](#)), because these counts represent a direct reflection of the completeness of the genome assembly. In contrast, contig counts and N50 are influenced by biological features such as repeat copy numbers as well as the underlying sequence




Figure 5

A) Comparisons of predicted to true phenotypes for 16 carbon source substrates. False-negatives, true-negatives, false-positives and true-positives are coloured as shown in legend. Each column represents a different isolate, separated by ST representation in the *KpSC*-pan model. **B)** Accuracy metrics for predicted vs phenotypic growth comparisons shown in A. Each column represents a different isolate, coloured by taxa and separated by ST representation in the *KpSC*-pan v1 model.

data quality e.g. a bacterial genome harbouring many insertion sequence insertions will result in a draft assembly with a high number of contigs regardless of the sequence data quality and completeness.

Traditionally, genome-scale metabolic reconstruction approaches have relied upon significant manual curation efforts. While there will always remain a need for high quality curated models, such resource intensive approaches preclude their application at scale, and have therefore limited analyses to small numbers of individual strains (15, 16). However, automated reconstruction approaches can support the generation and comparison of multiple strain-specific draft models from which meaningful biological insights can be derived (61). Additionally, the quality of curated models is likely to vary depending on their age, level and type of curation, as well as the approach used for preliminary drafting. Indeed it is possible for automated approaches to outperform manually curated models; a draft model for *K. pneumoniae* KPPR1 generated using Bactabolize with the KpSC pan-v1 reference model outperformed the manually curated iKp1289 model representing the same strain (15). iKp1289 was published in 2017 (6 years prior to this study) and was initially drafted via the KBase pipeline (33), which uses RAST to annotate the sequences with Enzyme Commission numbers. It has been demonstrated several times that the Enzyme Commission scheme has systematic errors (62, 63), leading to a loss in accuracy when compared to the ortholog identification methods used by automated approaches. Consistent with this assertion, our draft KPPR1 model constructed with KBase (without manual curation) was an outlier in terms of the very low number of genes, reactions and metabolites that were included.

CarveMe with universal model (30) and gapseq (31) are the current gold standard automated approaches for model reconstruction, and we show that a draft KpSC model generated by Bactabolize with the KpSC pan v1 reference resulted in similar or better accuracy for phenotype prediction (Figure 2). Both the CarveMe universal and gapseq models resulted in high numbers of true-positive and true-negative growth predictions. However, these were also accompanied by comparatively higher numbers of false-positive predictions that resulted in a lower overall accuracy for substrate usage analysis compared to Bactabolize with the KpSC-pan v1 reference (Figure 2), and comparatively lower precision and specificity for the gene essentiality analysis. False-positive predictions may indicate that the relevant metabolic machinery are present in the cell but were not active during the growth experiments (e.g. due to lack of gene expression). In this regard, false-positives are not always a sign of model inaccuracy. However, false-positive predictions can also occur from incorrect gene annotations e.g. due to reduced specificity of ortholog assignment resulting from the use of the universal model without manual curation. Given a key objective here is to facilitate high-throughput analysis for large numbers of genomes, it is not feasible to expect that all models will be manually curated, and therefore we believe that identifying fewer genes with lower overall error rates provides greater confidence in the resulting draft models. We also note that the BiGG universal reference model which CarveMe leverages is no longer being actively maintained. In contrast, user defined reference models can be iteratively curated and updated to incorporate new knowledge and data as they become available.

Bactabolize's reference-based reconstruction approach is reductive, meaning the resultant draft models will comprise only the genes, reactions and metabolites present in the reference, or a subset thereof, and will not include novel reactions unless they are manually identified and curated by the user. This is an important caveat that should be considered carefully for application of Bactabolize to large genome data sets, particularly for genetically diverse organisms such as those in the KpSC. For optimum results we suggest using a curated pan-model that captures as much diversity as possible for the target species or group of interest. While we acknowledge that a reasonable resource investment is required to generate a high-quality reference, we have shown that a pan-model derived from just 37 representative strains can be sufficient to support the generation of highly-accurate draft models (Figure 2 and 5). Additionally, we note that it is possible to use a single strain reference model, which should ideally represent the same or closely related species to that of the input genome assemblies, in order to

facilitate accurate identification of gene orthologs. It is technically possible to use an unrelated reference model, but this is expected to result in inaccurate and/or incomplete outputs and has not been tested in this study. In circumstances where no high quality closely-related reference model is available, we recommend alternative reconstruction approaches that leverage universal databases e.g. CarveMe (30) or gapseq (31). However, gapseq's long compute time makes it inappropriate for application to datasets comprising 100s-1000s of genomes (such as have become increasingly common in the bacterial population biology literature) (64, 65).

Bactabolize and the *KpSC* pan v1 model are freely available under open source licenses and satisfy the four features of the FAIR research principles (findability, accessibility, interoperability and reusability) (66). In addition to the *KpSC* pan-reference described here, a pan-reference model has been described previously for *Salmonella enterica* (representing 410 strains) (25), *Bacillus subtilis* (67) (representing 183 strains) and *Escherichia coli* (68) (representing 222 strains). We are actively working to expand and improve the *KpSC* pan reference model and welcome similar efforts to generate high quality references for other organisms. Together these resources will facilitate population wide metabolic analyses for global priority pathogens, which can be used to understand how they transmit, cause disease and evolve drug-resistance, and to identify novel therapeutic targets.

Methods

Bactabolize pipeline

Bactabolize utilises the existing metabolic modelling library COBRApy (38) and Python 3 (69). All code is freely available and open source at GitHub (www.github.com/kelwyres/Bactabolize (<http://www.github.com/kelwyres/Bactabolize>)) under a GNU General Public License v3.0. Users should additionally cite COBRApy (38) if Bactabolize is used.

Klebsiella pneumoniae Species Complex-pan metabolic model

The 37 metabolic models from a previous study (14) were combined with the iY1228 model using the `create_master_model.py` script (available at 10.6084/m9.figshare.21728717). Briefly, all GPRs from the iYL1228 model and the associated sequences were included, as well as new GPRs identified from the 36 additional strains by manual curation following comparison to the matched phenotype data (as described in (14)). Additionally, orthologous sequence variants with <75% nucleotide identity to gene sequences associated with these gene reaction rules (GPRs) were added if there was phenotype data supporting the reaction. The biomass reaction was updated, removing the metabolites `udpgal_c` and `udpgal_c` as their production was strain-specific.

Metadata annotations were improved using the `improve_model_annotations.py` script (also available in the Bactabolize code repository) resulting in the *KpSC* pan v1 used in this study, available at www.github.com/kelwyres/KpSC-pan-metabolic-model (<http://www.github.com/kelwyres/KpSC-pan-metabolic-model>).

Draft model generation

The annotated and unannotated genome of *K. pneumoniae* KPPR1 were obtained from Genbank accession number: CP009208.

Bactabolize draft models were generated using the `draft_model` command in Bactabolize v1 with the *KpSC*-pan v1 model as a reference, the annotated *K. pneumoniae* KPPR1 as input and the following options:

```
--min_coverage 25 --min_pident 80 --media M9 --atmosphere aerobic
```

A draft model for *K. pneumoniae* KPPR1 was also generated via CarveMe version 1.5.1 using the universal reference, the annotated *K. pneumoniae* KPPR1 as input, with the following commands:

‘-g M9 -i M9’. Subsequently, the --universe-file mode was also used, so the *KpSC*-pan model could be used as a reference, with the previously described command.

A draft model was generated using gapseq version 1.2 with the ‘doall’ command using the unannotated genome (as gapseq does not take annotated input files). Gap-filling was subsequently performed using the ‘fill’ command and a custom M9 media file to match the nutrient list found in Bactabolize (https://github.com/kelwyres/Bactabolize/blob/main/data/media_definitions/m9_media.json).

Finally, a draft model was constructed using the annotated genbank *K. pneumoniae* KPPR1 file and the KBase narrative (15) (<https://narrative.kbase.us/narrative/ws.14145.obj.1>).

Speed calculations

Modelling methods were timed via a script using the date +%s.%N command run before and after command on the MASSIVE computing cluster (Intel Xeon Gold 6150 CPU @ 2.70GHz and 155 GB of memory, CentOS Linux release 7.9.2009 environment). 10 individual complete *KpSC* genomes used in the quality control framework were tested for each method and the mean and range reported using R version 4.0.3 (70).

Performance comparisons

The annotated and unannotated genome of *K. pneumoniae* KPPR1 were obtained from Genbank under the accession: CP009208, and draft metabolic models were generated using Bactabolize, CarveMe, ModelSEED and gapseq as described above. The previously described, manually curated model for KPPR1 (iKp1289) was also included for comparison (15). MEMOTE version 0.13.0 was used to collate basic model statistics. The following KPPR1 phenotype data were retrieved from published studies: BIOLOG Phenotypic Microarray data (15) and single gene knockout data inferred from the outputs of a TraDIS transposon mutagenesis library (58).

A list of BIOLOG growth substrates for plates PM1, PM2A, PM3B, and PM4A (71) were converted where possible to BiGG and SEED IDs by manual search of the BiGG (bigg.ucsd.edu) and SEED websites (<https://modelseed.org/biochem/compounds>). An updated BiGG to SEED dictionary can be found in **Data S1**. A total of 143 of 190 carbon, 82 of 95 nitrogen, 46 of 59 phosphor and 26 of 35 sulfur substrates were successfully matched to BiGG and SEED IDs (**Data S1**). These growth data were compared to *in silico* predictions generated via FBA using the *fba* command from Bactabolize to optimise the biomass objective function with the following options:

```
--fba_spec_name m9 --fba_open_value -20
```

Gene essentiality was inferred from single gene knockout growth predictions using the *sgk* command from Bactabolize with the following options to mirror the growth conditions of the TraDIS library (LB media grown aerobically):

```
--media_type lb --atmosphere aerobic
```

In all cases, an objective value cut-off of $\geq 10^{-4}$ was used to indicate binarised growth as per previous studies (14, 72).

In silico predictions were compared to matched phenotype data and the following accuracy metrics were calculated:

$$\text{Precision} = \frac{TP}{TP + FP}$$

$$\text{Sensitivity/recall} = \frac{TP}{TP + FN}$$

$$\text{Specificity} = \frac{TN}{TN + FP}$$

$$\text{Accuracy} = \frac{TP + TN}{TP + FP + TN + FN}$$

$$F1 - \text{score} = 2 \times \frac{\text{Precision} \times \text{Sensitivity}}{\text{Precision} + \text{Sensitivity}}$$

Model metabolite and reaction IDs were harmonized for overlap comparisons using the ‘Import model’ function from *MetaNetX.org* (<http://metanetx.org/>) (59) after import and export via the `write_sbml_model` function from COBRApy.

Quality control framework

Illumina read sets (250 bp paired end) and completed genome sequences for 37 *KpSC* isolates were described previously (14). Here we randomly subsampled the Illumina reads at various depths (10 – 100, by increments of 10) using *rasusa* version 0.3.0 (73) in technical triplicate. Reads were then trimmed using *TrimGalore* version 0.5.0 (74) and assembled *de novo* with *Unicycler* version 0.4.7 (75), default parameters. Assembly statistics and assembly graph dead ends were calculated using the *GFA-dead-end-counter* version 1.0.0 (<https://github.com/rrwick/GFA-dead-end-counter>) (76). Draft metabolic models were generated with *Bactabolize* using the *KpSC*-pan v1 reference, and growth substrate profiles were predicted as described above. We compared the outputs from models generated for draft genome assemblies to those generated for the corresponding completed genomes. Where necessary models were gap-filled via the *patch_model* command.

Predictive accuracy of draft models

Novel growth phenotype data were generated for 10 *KpSC* clinical isolates from our in house collection using the VITEK 2 GN ID card system as described previously (14). Briefly, isolates were grown on Tryptic Soy (OXOID) agar plates overnight at 37°C, then analysed using VITEK 2 GN ID cards (bioMérieux) and read on the VITEK 2 Compact (bioMérieux) as per manufacturer’s instructions using software version 8.0. DNA was extracted for whole-genome sequencing via Genfind v3 extraction kit, library preparation performed using Nextera Flex (Illumina) using ¼ reagents. Paired-end read data (300 bp) were generated on an Illumina NovaSeq6000 SP v1.0 and have been deposited in the European Nucleotide Archive under Bioproject PRJNA777643 (individual read accession numbers are given in **Data S4**). Draft genome assemblies were generated with *Unicycler*, and draft metabolic models and growth predictions were generated with *Bactabolize* as described above.

Statistics and visualisation

Statistical analysis and graphical visualisation were performed using R version 4.0.3 (70), RStudio version 1.3.1093 (77), with the following software packages: *tidyverse* version 1.3.1 (78), *viridis* version 0.5.1 (79), *RColorBrewer* version 1.1-2 (80), *ggpubr* version 0.4.0 (81), *ggpmisc* version 0.4.4 (82), *aplot* version 0.1.6 (83), *colorspace* version 2.0-2 (84), *ggpattern* version 0.4.3-3 (85), *ggtext* version 0.1.1 (86) and *glue* version 1.4.2 (87).

Linear regression analysis was performed in R using the `lm` function and a third degree polynomial model was fitted to plots with the following equation: $y \sim \text{poly}(x, 3, \text{raw} = \text{TRUE})$. The segmented linear model was fitted using segmented version 1.6-2 (88 [↗](#)).

All code used to generate results can be found as supplemental material, <https://github.com/kelwyres/Bactabolize> [↗](#), <https://github.com/kelwyres/KpSC-pan-metabolic-model> [↗](#) and on Figshare (10.6084/m9.figshare.21728717).

Logo

The Bactabolize logo was constructed in Inkscape version 1.0.1 (89 [↗](#)). The font used is Proportional TFB (90 [↗](#)) and Element (91 [↗](#)).

Author contributions

Conceptualization: BV, JH, JMM, KEH, K LW

Methodology: SCW, BV, LMJ, JH, JMM, K LW

Software: SCW, BV

Validation: BV, HBC, K LW

Formal analysis: BV, K LW

Investigation: BV, K LW

Resources: AJ

Writing - Original Draft: BV, K LW

Writing - Review & Editing: All authors

Visualization: BV, K LW

Supervision: JMM, KEH, K LW

Project administration: K LW

Funding acquisition: JMM, KEH, K LW

All authors contributed to and approve of the manuscript in its current form.

Acknowledgements

We thank Sylvain Brisse for the isolates used in this study and review of the manuscript.

Funding

This work was funded by the Australian Research Council Discovery Project DP200103364.

Data availability

All genomes used are publicly available and generated by referenced studies. All Bactabolize code and data generated for this study can be found at <https://github.com/kelwyres/Bactabolize>. The KpSC-pan v1 reference model is freely available at <https://github.com/kelwyres/KpSC-pan-metabolic-model>.

Conflict of Interest

The authors declare that they have no competing or conflicts of interest.

Supplementary figures




Figure S1

Flow diagram showing the overview of the draft_model module from Bactabolize, which produces draft metabolic models. Input and output files are shown in light pink while Bactabolize processes are shown in dark pink. Third-party dependencies are indicated within the white boxes.




Figure S2

Flow diagram showing the overview of the patch_model module from Bactabolize, which patches metabolic models that do not simulate growth. Input and output files are shown in light orange while Bactabolize processes are shown in dark orange. Third-party dependencies are indicated within the white boxes.




Figure S3

Flow diagram showing the overview of the fba module from Bactabolize, which performs growth simulations using Flux Balance Analysis. Input and output files are shown in light yellow while Bactabolize processes are shown in dark yellow. Third-party dependencies are indicated within white boxes. C, carbon; N, nitrogen, P, phosphorus; S, sulphur; O₂, oxygen.




Figure S4

Flow diagram showing the overview of the sgk module from Bactabolize, which performs Single Gene Knockout analysis. Input and output files are shown in light blue while Bactabolize processes are shown in dark blue. Third-party dependencies are indicated within white boxes.




Figure S5

Venn diagram comparing the metabolite output of the best-performing tools for *K. pneumoniae* KPPR1.




Figure S6

Venn diagram comparing the reactions output of the best-performing tools for *K. pneumoniae* KPPR1.




Figure S7

Raincloud plot showing distributions of assembly metrics across various read subsampling depths (10x increments).




Figure S8

Line graph showing the capture of model features of draft assemblies (short read only) at various depths, compared to the corresponding completed genome (long-read + short read assemblies).




Figure S9

Faceted graphs showing the number of substrate usage (fba module) discrepancies of gap-filled models (patch_model module) which initially did not produce biomass (models which failed to simulate growth). The dots indicate percentage concordance with the completed genome model, while the columns indicate number of substrates with discrepancies (no simulated growth in patched model, but growth in completed genome model).

References

1. Su Q, Guan T, He Y, Lv H (2016) **Siderophore Biosynthesis Governs the Virulence of Uropathogenic Escherichia coli by Coordinately Modulating the Differential Metabolism** *Journal of Proteome Research* **15**:1323–32
2. Wu Y, Meng Y, Qian L, Ding B, Han H, Chen H, et al. **The Vancomycin Resistance-Associated Regulatory System VraSR Modulates Biofilm Formation of Staphylococcus epidermidis in an ica-Dependent Manner** *mSphere* **6**:e00641–21
3. Mir M, Priscic S, Kang C-M, Lun S, Guo H, Murry JP, et al. (2014) **Mycobacterial gene *cuvA* is required for optimal nutrient utilization and virulence** *Infection and immunity* **82**:4104–17
4. Vornhagen J, Sun Y, Breen P, Forsyth V, Zhao L, Mobley HLT, et al. (2019) **The Klebsiella pneumoniae citrate synthase gene, *gltA*, influences site specific fitness during infection** *PLOS Pathogens* **15**
5. Eberl C, Weiss AS, Jochum LM, Durai Raj AC, Ring D, Hussain S, et al. (2021) **E. coli enhance colonization resistance against Salmonella Typhimurium by competing for galactitol, a context-dependent limiting carbon source** *Cell Host & Microbe*
6. Jenior ML, Dickenson ME, Papin JA (2021) **Genome-scale metabolic modeling reveals increased reliance on valine catabolism in clinical isolates of Klebsiella pneumoniae** *bioRxiv*
7. Hudson AW, Barnes AJ, Bray AS, Zafar MA (2022) **Klebsiella pneumoniae L-Fucose metabolism promotes gastrointestinal colonization and modulates its virulence determinants** *bioRxiv*
8. Rodrigues C, Passet V, Rakotondrasoa A, Diallo TA, Criscuolo A, Brisse S (2019) **Description of Klebsiella africanensis sp. nov Klebsiella variicola subsp. tropicalensis subsp. nov. and Klebsiella variicola subsp. variicola subsp. nov.** *Res Microbiol* **170**:165–70
9. Blin C, Passet V, Touchon M, Rocha EPC, Brisse S (2017) **Metabolic diversity of the emerging pathogenic lineages of Klebsiella pneumoniae** *Environmental Microbiology* **19**:1881–98
10. Brisse S, Fevre C, Passet V, Issenhuth-Jeanjean S, Tournebize R, Diancourt L, et al. (2009) **Virulent Clones of Klebsiella pneumoniae: Identification and Evolutionary Scenario Based on Genomic and Phenotypic Characterization** *PLOS ONE* **4**
11. Mobegi FM, van Hijum SAFT, Burghout P, Bootsma HJ, de Vries SPW, van der Gaast-de Jongh CE, et al. (2014) **From microbial gene essentiality to novel antimicrobial drug targets** *BMC Genomics* **15**
12. Hogan AM, Scoffone VC, Makarov V, Gislason AS, Tesfu H, Stietz MS, et al. (2018) **Competitive Fitness of Essential Gene Knockdowns Reveals a Broad-Spectrum Antibacterial Inhibitor of the Cell Division Protein FtsZ** *Antimicrobial Agents and Chemotherapy* **62**:e01231–18
13. Edwards JS, Palsson BO (1999) **Systems Properties of the Haemophilus influenzae Rd Metabolic Genotype** *Journal of Biological Chemistry* **274**:17410–6

14. Hawkey J, Vezina B, Monk JM, Judd LM, Harshegyi T, López-Fernández S, et al. (2022) **A curated collection of *Klebsiella* metabolic models reveals variable substrate usage and gene essentiality** *Genome Research*
15. Henry CS, Rotman E, Lathem WW, Tyo KEJ, Hauser AR, Mandel MJ (2017) **Generation and Validation of the iKp1289 Metabolic Model for *Klebsiella pneumoniae* KPPR1** *The Journal of Infectious Diseases* **215**
16. Liao Y-C, Huang T-W, Chen F-C, Charusanti P, Hong JSJ, Chang H-Y, et al. (2011) **An Experimentally Validated Genome-Scale Metabolic Reconstruction of *Klebsiella pneumoniae* MGH 78578, iYL1228** *Journal of Bacteriology* **193**
17. Orth JD, Thiele I, Palsson BØ (2010) **What is flux balance analysis?** *Nature Biotechnology* **28**:245–8
18. Stanway RR, Bushell E, Chiappino-Pepe A, Roques M, Sanderson T, Franke-Fayard B, et al. (2019) **Genome-Scale Identification of Essential Metabolic Processes for Targeting the *Plasmodium* Liver Stage** *Cell* **179**:1112–28
19. Ramos PIP (2018) **Fernández Do Porto D, Lanzarotti E, Sosa EJ, Burguener G, Pardo AM et al. An integrative, multi-omics approach towards the prioritization of *Klebsiella pneumoniae* drug targets.** *Scientific Reports* **8**
20. Wyres KL, Lam MMC, Holt KE (2020) **Population genomics of *Klebsiella pneumoniae*** *Nature Reviews Microbiology* **18**:344–59
21. WHO (2017) **WHO publishes list of bacteria for which new antibiotics are urgently needed**
22. Gorrie CL, Mirčeta M, Wick RR, Judd LM, Lam MMC, Gomi R, et al. (2022) **Genomic dissection of *Klebsiella pneumoniae* infections in hospital patients reveals insights into an opportunistic pathogen** *Nature Communications* **13**
23. Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, et al. (2015) **Genomic analysis of diversity, population structure, virulence, and antimicrobial resistance *Klebsiella pneumoniae*, an urgent threat to public health** *Proceedings of the National Academy of Sciences* **112**
24. Monk JM, Charusanti P, Aziz RK, Lerman JA, Premyodhin N, Orth JD, et al. (2013) **Genome-scale metabolic reconstructions of multiple *Escherichia coli* strains highlight strain-specific adaptations to nutritional environments** *Proceedings of the National Academy of Sciences* **110**
25. Seif Y, Kavvas E, Lachance J-C, Yurkovich JT, Nuccio S-P, Fang X, et al. (2018) **Genome-scale metabolic reconstructions of multiple *Salmonella* strains reveal serovar-specific metabolic traits** *Nature Communications* **9**
26. Bosi E, Monk JM, Aziz RK, Fondi M, Nizet V, Palsson BØ (2016) **Comparative genome-scale modelling of *Staphylococcus aureus* strains identifies strain-specific metabolic capabilities linked to pathogenicity** *Proceedings of the National Academy of Sciences* **113**:E3801–E9
27. Bartell JA, Blazier AS, Yen P, Thøgersen JC, Jelsbak L, Goldberg JB, et al. (2017) **Reconstruction of the metabolic network of *Pseudomonas aeruginosa* to interrogate virulence factor synthesis** *Nature Communications* **8**

28. Croucher NJ, Coupland PG, Stevenson AE, Callendrello A, Bentley SD, Hanage WP (2014) **Diversification of bacterial genome content through distinct mechanisms over different timescales** *Nature Communications* **5**
29. Cummins EA, Hall RJ, Connor C, McInerney JO, McNally A (2022) **Distinct evolutionary trajectories in the Escherichia coli pangenome occur within sequence types** *Microbial Genomics* **8**
30. Machado D, Andrejev S, Tramontano M, Patil KR (2018) **Fast automated reconstruction of genome-scale metabolic models for microbial species and communities** *Nucleic Acids Res* **46**:7542–53
31. Zimmermann J, Kaleta C (2021) **Waschina S. gapseq: informed prediction of bacterial metabolic pathways and reconstruction of accurate metabolic models** *Genome Biology* **22**
32. Seaver SMD, Liu F, Zhang Q, Jeffries J, Faria JP, Edirisinghe JN, et al. (2021) **The ModelSEED Biochemistry Database for the integration of metabolic annotations and the reconstruction, comparison and analysis of metabolic models for plants, fungi and microbes** *Nucleic Acids Res* **49**:D575–D88
33. Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, et al. (2018) **KBase: The United States Department of Energy Systems Biology Knowledgebase** *Nature Biotechnology* **36**:566–9
34. Mendoza SN, Olivier BG, Molenaar D, Teusink B (2019) **A systematic assessment of current genome-scale metabolic reconstruction tools** *Genome Biology* **20**
35. Tamasco G, Kumar M, Zengler K (2022) **Silva-Rocha R, da Silva RR. ChiMera: an easy to use pipeline for bacterial genome based metabolic network reconstruction, evaluation and visualization** *BMC Bioinformatics* **23**
36. Norsigian CJ, Fang X, Seif Y, Monk JM, Palsson BO (2020) **A workflow for generating multi-strain genome-scale metabolic models of prokaryotes** *Nature Protocols* **15**:1–14
37. Zimmermann J, Kaleta C, Waschina S (2021) **37. Zimmermann J, Kaleta C, Waschina S. gapseq GitHub issue #77: Speed improvement suggestions 2021 [Available from: <https://github.com/jotech/gapseq/issues/77>]**
38. Ebrahim A, Lerman JA, Palsson BO, Hyduke DR (2013) **COBRApy: CONstraints-Based Reconstruction and Analysis for Python** *BMC Systems Biology* **7**
39. Schellenberger J, Park JO, Conrad TM, Palsson BØ (2010) **BiGG: a Biochemical Genetic and Genomic knowledgebase of large scale metabolic reconstructions** *BMC Bioinformatics* **11**
40. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ (2010) **Prodigal: prokaryotic gene recognition and translation initiation site identification** *BMC Bioinformatics* **11**
41. Keating SM, Waltemath D, König M, Zhang F, Dräger A, Chaouiya C, et al. (2020) **SBML Level 3: an extensible format for the exchange and reuse of biological models** *Molecular Systems Biology* **16**
42. Lieven C, Beber ME, Olivier BG, Bergmann FT, Ataman M, Babaei P, et al. (2020) **MEMOTE for standardized genome-scale metabolic model testing** *Nature Biotechnology* **38**:272–6

43. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. (2009) **BLAST+: architecture and applications** *BMC Bioinformatics* **10**
44. Hernández-Salmerón JE, Moreno-Hagelsieb G (2020) **Progress in quickly finding orthologs as reciprocal best hits: comparing blast, last, diamond and MMseqs2** *BMC Genomics* **21**
45. BD (2009) **BD. Difco™ & BBL™ Manual, 2nd Edition.** In: BD, editor. 2009. *Difco™ & BBL™ Manual, 2nd Edition.* In: BD, editor
46. Biosciences BD (2015) **BD Bionutrients Technical Manual: BD Biosciences – Advanced Bioprocessing** *BD=Biosciences*
47. Loginova LI, Manuilova VP, Tolstikov VP (1974) **Content of free amino acids in peptone and the dynamics of their consumption in the microbiological synthesis of dextran** *Pharmaceutical Chemistry Journal* **8**:249–51
48. ThermoFisherScientific (2019) **Technical guide to peptones, supplements, and feeds: Enhancing performance of mammalian and microbial bioprocesses** *ThermoFisherScientific*
49. Clausen E, Gildberg A, Raa J (1985) **Preparation and testing of an autolysate of fish viscera as growth substrate for bacteria** *Applied and Environmental Microbiology* **50**:1556–7
50. Hagely KB, Palmquist D, Bilyeu KD (2013) **Classification of Distinct Seed Carbohydrate Profiles in Soybean** *Journal of Agricultural and Food Chemistry* **61**:1105–11
51. Choct M, Dersjant-Li Y, McLeish J, Peisker M (2010) **Soy Oligosaccharides and Soluble Non-starch Polysaccharides: A Review of Digestion, Nutritive and Anti-nutritive Effects in Pigs and Poultry** *Asian-Australasian Journal of Animal Sciences* **23**
52. Tomé D (2021) **Yeast Extracts: Nutritional and Flavoring Food Ingredients** *ACS Food Science & Technology* **1**:487–94
53. Plata MR, Koch C, Wechselberger P, Herwig C, Lendl B (2013) **Determination of carbohydrates present in *Saccharomyces cerevisiae* using mid-infrared spectroscopy and partial least squares regression** *Anal Bioanal Chem* **405**:8241–50
54. Liu Y, Huang G, Lv M (2018) **Extraction, characterization and antioxidant activities of mannan from yeast cell wall** *International Journal of Biological Macromolecules* **118**:952–6
55. Blagović B (2001) **Lipid Composition of Brewer's Yeast** *Food Technology and Biotechnology* **39**:175–81
56. Blagović B, Mesarić M, Marić V, Rupčić J (2005) **Characterization of lipid components in the whole cells and plasma membranes of baker's Yeast** *Croatica Chemica Acta* **78**:479–84
57. Avramia I, Amariei S (2021) **Spent Brewer's Yeast as a Source of Insoluble β -Glucans** *International Journal of Molecular Sciences* **22**
58. Francesca L Short, Di Sario G, Reichmann Nathalie T, Kleanthous C, Parkhill J, Taylor Peter W, et al. (2020) **Genomic Profiling Reveals Distinct Routes To Complement Resistance in *Klebsiella pneumoniae*** *Infection and Immunity* **88**:e00043–20

59. Moretti S, Van Du T Tran, Mehl F, Ibberson M, Pagni M (2021) **MetaNetX/MNXref: unified namespace for metabolites and biochemical reactions in the context of metabolic models** *Nucleic Acids Res* **49**:D570–D4
60. Powers DMW (2020) **Evaluation: from precision, recall and F-measure to ROC, informedness, markedness and correlation** *arXiv*
61. Magnúsdóttir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, et al. (2017) **Generation of genome-scale metabolic reconstructions for 773 members of the human gut microbiota** *Nature Biotechnology* **35**:81–9
62. Green ML, Karp PD (2005) **Genome annotation errors in pathway databases due to semantic ambiguity in partial EC numbers** *Nucleic Acids Res* **33**:4035–9
63. Rembeza E (2021) **Engqvist MKM Experimental and computational investigation of enzyme functional annotations uncovers misannotation in the EC 1.1.3.15 enzyme class.** *PLOS Computational Biology* **17**
64. Ludden C, Moradigaravand D, Jamrozy D, Gouliouris T, Blane B, Naydenova P, et al. (2020) **A One Health Study of the Genetic Relatedness of Klebsiella pneumoniae and Their Mobile Elements in the East of England** *Clinical Infectious Diseases* **70**:219–26
65. Dyson ZA, Holt KE (2021) **Five Years of GenoTyphi: Updates to the Global Salmonella Typhi Genotyping Framework** *The Journal of Infectious Diseases* **224**
66. Wilkinson MD, Dumontier M, Aalbersberg IJ, Appleton G, Axton M, Baak A, et al. (2016) **The FAIR Guiding Principles for scientific data management and stewardship** *Scientific Data* **3**
67. Blázquez B, San León D, Rojas A, Tortajada M, Nogales J (2023) **New Insights on Metabolic Features of Bacillus subtilis Based on Multistrain Genome-Scale Metabolic Modeling** *International Journal of Molecular Sciences* [Internet] **24**
68. Monk JM (2022) **Genome-scale metabolic network reconstructions of diverse Escherichia strains reveal strain-specific adaptations** *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**
69. Van Rossum G, Drake F. L (2009) **Van Rossum G, & Drake, F. L. Python 3 Reference Manual. Scotts Valley, CA2009.** *Python 3 Reference Manual. Scotts Valley, CA*
70. R-Core-Team (2020) **R-Core-Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing; 2020. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing**
71. BIOLOG (2020) **BIOLOG. Phenotype MicroArrays. In: BIOLOG, editor. 2020. Phenotype MicroArrays. In: BIOLOG, editor**
72. Norsigian CJ, Attia H, Szubin R, Yassin AS, Palsson BØ, Aziz RK, et al. (2019) **Comparative Genome-Scale Metabolic Modeling of Metallo-Beta-Lactamase–Producing Multidrug-Resistant Klebsiella pneumoniae Clinical Isolates** *Frontiers in Cellular and Infection Microbiology* **9**
73. Hall MB (2019) **Hall MB. Rasusa: Randomly subsample sequencing reads to a specified coverage 2019 [Available from: <https://github.com/mbhall88/rasusa>.**

74. Krueger F (2012) **Krueger F. Trim Galore GitHub2012** [Available from: <https://github.com/FelixKrueger/TrimGalore>].
75. Wick RR, Judd LM, Gorrie CL, Holt KE (2017) **Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads** *PLOS Computational Biology* **13**
76. Wick RR (2023) **Wick RR. Dead-end count for QC of short-read assemblies. 2023.** *Dead-end count for QC of short-read assemblies*
77. RStudio-Team (2020) **RStudio: Integrated Development for R. Boston MA RStudio**
78. Wickham H, Averick M, Bryan J, Chang W, McGowan LDA, François R, et al. (2019) **Welcome to the Tidyverse** *Journal of Open Source Software* **4**
79. Garnier S (2018) **Garnier S. viridis: Default Color Maps from 'matplotlib'. 2018.** *viridis: Default Color Maps from 'matplotlib'*
80. Neuwirth E (2014) **Neuwirth E. RColorBrewer: ColorBrewer Palettes. 1.1–2 ed2014.** *RColorBrewer: ColorBrewer Palettes* **1:1–2**
81. Kassambara A (2022) **Kassambara A. ggpubr: 'ggplot2' Based Publication Ready Plots. 0.5.0 ed2022.** *ggpubr: 'ggplot2' Based Publication Ready Plots. 0.5.0*
82. Aphalo PJ, Slowikowski K, Mouksassi S. (2021) **Aphalo PJ, Slowikowski K, Mouksassi S. ggpmisc: Miscellaneous Extensions to 'ggplot2'. 0.4.1 ed2021.** *ggpmisc: Miscellaneous Extensions to 'ggplot2'. 0.4.1 ed*
83. Yu G (2020) **Yu G. aplot: Decorate a 'ggplot' with Associated Information. 0.0.6 ed2020.** *aplot: Decorate a 'ggplot' with Associated Information. 0.0.6 ed*
84. Zeileis A, Fisher JC, Hornik K, Ihaka R, McWhite CD, Murrell P, et al (2020) . **colorspace: A Toolbox for Manipulating and Assessing Colors and Palettes** *Journal of Statistical Software* **96:1–49**
85. Davis TL F.C. M (2022) **F.C. M, Davis TL. Package 'ggpattern'. 2022.** *Package 'ggpattern'*
86. Wilke CO (2020) **Wilke CO. ggtext: Improved Text Rendering Support for 'ggplot2'. 0.1.1 ed2020.** *ggtext: Improved Text Rendering Support for 'ggplot2'. 0.1.1 ed*
87. Hester J, Bryan J (2022) **Hester J, Bryan J, RStudio. glue: Interpreted String Literals. 1.6.2 ed2022.** *RStudio. glue: Interpreted String Literals. 1.6.2 ed*
88. Muggeo VMR (2022) **Muggeo VMR. segmented: Regression Models with Break-Points / Change-Points (with Possibly Random Effects) Estimation. 1.6-2 ed2022.** *segmented: Regression Models with Break-Points / Change-Points (with Possibly Random Effects) Estimation. 1.6-2 ed*
89. The-Inkscape-Team (2020) **The-Inkscape-Team. Inkscape. 1.0.1 (3bc2e813f5, 2020-09-07) ed2020.** *Inkscape. 1.0.1 (3bc2e813f5, 2020-09-07)*
90. zanatlja (2012) **zanatlja. Proportional TFB. 2012.** *Proportional TFB*
91. weknow (2015) **weknow. Element. 2015.** *Element*

Article and author information

Ben Vezina

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia

For correspondence: benjamin.vezina@monash.edu

ORCID iD: [0000-0003-4224-2537](https://orcid.org/0000-0003-4224-2537)

Stephen C. Watts

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia

ORCID iD: [0000-0001-7084-635X](https://orcid.org/0000-0001-7084-635X)

Jane Hawkey

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia

ORCID iD: [0000-0001-9661-5293](https://orcid.org/0000-0001-9661-5293)

Helena B. Cooper

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia

ORCID iD: [0000-0002-7096-0010](https://orcid.org/0000-0002-7096-0010)

Louise M. Judd

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia

ORCID iD: [0000-0003-3613-4839](https://orcid.org/0000-0003-3613-4839)

Adam W. J. Jenney

Microbiology Unit, Alfred Health, Melbourne, Victoria, Australia

ORCID iD: [0000-0002-5237-2506](https://orcid.org/0000-0002-5237-2506)

Jonathan M. Monk

Department of Bioengineering, University of California, San Diego, CA, United States of America

ORCID iD: [0000-0002-3895-8949](https://orcid.org/0000-0002-3895-8949)

Kathryn E. Holt

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia, Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK

ORCID iD: [0000-0003-3949-2471](https://orcid.org/0000-0003-3949-2471)

Kelly L. Wyres

Department of Infectious Diseases, Central Clinical School, Monash University, Melbourne, Victoria, Australia

For correspondence: kelly.wyres@monash.edu

ORCID iD: [0000-0002-4033-6639](https://orcid.org/0000-0002-4033-6639)

Copyright

© 2023, Vezina et al.

This article is distributed under the terms of the [Creative Commons Attribution License \(https://creativecommons.org/licenses/by/4.0/\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Marisa Nicolás

Laboratório Nacional de Computação Científica, Brazil

Senior Editor

Christian Landry

Université Laval, Canada

Reviewer #1 (Public Review):

In this work, Vezina et al. present Bactabolize, a rapid reconstruction tool for the generation of strain-specific metabolic models. Similar to other reconstruction pipelines such as CarveMe, Bactabolize builds a strain-specific draft reconstruction and subsequently gap-fills it. The model can afterwards be used to predict growth on carbon sources. The authors constructed a pan-model of the *Klebsiella pneumoniae* species complex (KpSC) and used it as input for Bactabolize to construct a genome-scale reconstruction of *K. pneumoniae* KPPR1. They compared the generated reconstruction with a reconstruction built through CarveMe as well as a manually curated reconstruction for the same strain. They then compared predictions of carbon, nitrogen, phosphorus, and sulfur sources and found that the Bactabolize reconstruction had the overall highest accuracy. Finally, they built draft reconstructions for 10 clinical isolates of *K. pneumoniae* and evaluated their predictive performance. Overall, this is a useful tool, the data is well-presented, and the paper is well-written.

Reviewer #2 (Public Review):

The authors present a pipeline for generating strain-specific genome-scale metabolic models for bacteria using *Klebsiella* spp. as the demonstrative data. This paper claims to provide a high-throughput tool for generating strain-specific models for bacteria. However, in reality, the tool requires a reference pan-genome-based complete model to generate the strain-specific model of the species of interest, which in this study is *Klebsiella pneumoniae*. This requirement renders the tool redundant for high-throughput purposes since the process of building or generating the pan-genome reference model is performed separately. Additionally, the quality of the newly built strain-specific model will depend on the reference model used. Therefore, this tool, on its own, can only work specifically with the available pan-genome model of reference, which in this case is only applicable to *Klebsiella pneumoniae*. Its effectiveness with other bacteria has not been proven. I would suggest that the authors either reframe the performance and results to be applicable only to *Klebsiella* or consider adding more reference pan-genome models for the study.

Author Response

The following is the authors' response to the original reviews.

Reviewer #1

1. Overall, this is a useful tool, the data is well-presented, and the paper is well-written. However, the predictions are only compared with two existing reconstruction tools though more have been recently published

The aim of this work was to facilitate high-throughput generation of strain-specific metabolic models e.g. at the scale of 100s -1000s as indicated throughout the introduction (see lines 74-82, 91-94), and therefore we only compared tools which were capable of high-throughput analysis via command line and excluded others (e.g. merlin). We have now tested this against the other recent command line tool, gapseq which had escaped our gaze. Thank you for bringing this to our attention. Additionally, we have included KBase (ModelSEED, a web-based app that does not support high-throughput analysis) to allow for readers to interpret the results in the context of community standard approaches, since KBase is a popular tool.

We have added an explicit statement about the choice of approaches, now at lines 194-199 as follows:

“We compared the output and performance of Bactabolize to the two previously published tools that can support high-throughput analyses i.e. CarveMe (30) and gapseq (31). To aid interpretation in the context of community standard approaches, we also include a comparison to the popular web-based reconstruction tool, KBase (ModelSEED), and a manually curated metabolic reconstruction of *K. pneumoniae* strain KPPR1 (also known as VK055 and ATCC 43816, metabolic model named iKp1289) (15).”

The methods section was updated accordingly (now lines 552-558):

“A draft model was generated using gapseq version 1.2 with the ‘doall’ command using the unannotated genome (as gapseq does not take annotated input files). Gap-filling was subsequently performed using the ‘fill’ command and a custom M9 media file to match the nutrient list found in Bactabolize (https://github.com/kelwyres/Bactabolize/blob/main/data/media_definitions/m9_media.json).

Finally, a draft model was constructed using the annotated genbank *K. pneumoniae* KPPR1 file and the KBase narrative (15) (<https://narrative.kbase.us/narrative/ws.14145.obj.1>).”

The results section (now lines 193-308) and Figure 2 have also been updated / restructured to reflect the new analyses, and include a comparison of the relative compute times for the construction of models (lines 281-291) as follows:

“While model features and accuracy are essential metrics for comparison, computation time is also a key consideration for high-throughput analyses. We recorded the time required for each tool to build draft models for 10 of the completed KpSC genomes used in the quality control framework (see below) on a high-performance computing cluster (Intel Xeon Gold 6150 CPU @ 2.70GHz and 155 GB of requested memory on a CentOS Linux release 7.9.2009 environment. CarveMe KpSC pan was the fastest with a mean of 20.04 (range 19.90 - 20.18) seconds, followed by CarveMe universal at 30.28 (range 29.20 - 31.80) seconds, then Bactabolize KpSC pan at 98.05 (range 92.19 - 100.4) seconds. KBase took 183.50 (range 120.00 - 338.00) seconds per genome via batch analysis, including genome upload time and queuing. gapseq took 5.46 (range 4.55 - 6.28) hours to produce draft models (not including the required gap-filling), consistent with previous reports (37).”

Finally, the whole discussion has been updated and substantially restructured (lines 472-474, 475-493, 494-512). Specific mentions to the new analyses are at lines:

472-474: “Consistent with this assertion, our draft KPPR1 model constructed with KBase (without manual curation) was an outlier in terms of the very low number of genes, reactions and metabolites that were included.”

475-493: “CarveMe with universal model (30) and gapseq (31) are the current gold standard automated approaches for model reconstruction, and we show that a draft KpSC model generated by Bactabolize with the KpSC pan v1 reference resulted in similar or better accuracy for phenotype prediction (Figure 2). Both the CarveMe universal and gapseq models resulted in high numbers of true-positive and true-negative growth predictions. However, these were also accompanied by comparatively higher numbers of false-positive predictions that resulted in a lower overall accuracy for substrate usage analysis compared to Bactabolize with the KpSC-pan v1 reference (Figure 2), and comparatively lower precision and specificity for the gene essentiality analysis. False-positive predictions may indicate that the relevant metabolic machinery are present in the cell but were not active during the growth experiments (e.g. due to lack of gene expression). In this regard, false-positives are not always a sign of model inaccuracy. However, false-positive predictions can also occur from incorrect gene annotations e.g. due to reduced specificity of ortholog assignment resulting from the use of the universal model without manual curation. Given a key objective here is to facilitate high-throughput analysis for large numbers of genomes, it is not feasible to expect that all models will be manually curated, and therefore we believe that identifying fewer genes with lower overall error rates provides greater confidence in the resulting draft models. We also note that the BiGG universal reference model which CarveMe leverages is no longer being actively maintained. In contrast, user defined reference models can be iteratively curated and updated to incorporate new knowledge and data as they become available.”

510-512: “However, gapseq’s long compute time makes it inappropriate for application to datasets comprising 100s-1000s of genomes (such as have become increasingly common in the bacterial population biology literature).”

1. *My understanding is that the tool requires a set of reference reconstructions for other strains of the target species. If no reference reconstruction is available for another strain of the target species, can this species not be reconstructed?*

Any input reference can be used to generate models however, single strain models matching the target species, or ideally a species-specific panreference, are recommended for best results. We have added a discussion on these points at lines 128-133:

“For optimum results we suggest using a pan-model that captures as much diversity as possible for the target species or group of interest, because Bactabolize’s reconstruction method is reductive i.e. each output strainspecific model will include only genes, reactions and metabolites that are present in the reference or a subset thereof (although novel genes, reactions and metabolites can be added via manual curation).”

We expand on these points further in the discussion:

494-512: “Bactabolize’s reference-based reconstruction approach is reductive, meaning the resultant draft models will comprise only the genes, reactions and metabolites present in the reference, or a subset thereof, and will not include novel reactions unless they are manually identified and curated by the user. This is an important caveat that should be considered carefully for application of Bactabolize to large genome data sets, particularly for genetically diverse organisms such as those in the KpSC. For optimum results we suggest using a curated

pan-model that captures as much diversity as possible for the target species or group of interest. While we acknowledge that a reasonable resource investment is required to generate a high-quality reference, we have shown that a pan-model derived from just 37 representative strains can be sufficient to support the generation of highly accurate draft models (Figure 2 and 5). Additionally, we note that it is possible to use a single strain reference model, which should ideally represent the same or closely related species to that of the input genome assemblies, in order to facilitate accurate identification of gene orthologs. It is technically possible to use an unrelated reference model, but this is expected to result in inaccurate and/or incomplete outputs and has not been tested in this study. In circumstances where no high quality closely-related reference model is available, we recommend alternative reconstruction approaches that leverage universal databases e.g. CarveMe (30) or gapseq (31). However, gapseq's long compute time makes it inappropriate for application to datasets comprising 100s-1000s of genomes (such as have become increasingly common in the bacterial population biology literature)."

1. *How do the reconstructions generated by Bactabolize compare to those generated by other reconstruction tools besides CarveMe and ModelSEED, e.g., gapseq (Zimmermann et al, Genome Biology 2021. 22:81) or merlin (Capela et al, Nucleic Acids Res 2022, 50(11):6052-6066)?*

See response to rev 1 point 1.

1. *How are the accuracy, specificity, and sensitivity of the pan-models calculated? Is the compared experimental data on the species level?*

We used the pan-model as a reference from which we generated a strain-specific model for *K. pneumoniae* KPPR1 (using Bactabolize and CarveMe). This strain-specific metabolic model was then used to simulate growth phenotypes and compared to published experimental data for KPPR1. This was described in the methods section, including the calculations for the metrics (lines 589-593); however, we have also expanded the description within the results section to clarify the approach (lines 201-209):

"De novo draft models for strain KPPR1 were built using; i) Bactabolize with the KpSC pan v1 reference; ii) CarveMe, with its universal reference model (CarveMe universal); iii) CarveMe, with KpSC-pan v1 reference (CarveMe KpSC pan); iv) gapseq; and v) KBase (ModelSEED). Subsequently, each model was used to predict growth phenotypes; i) in M9 minimal media with different sole sources of carbon, nitrogen, phosphorus and sulfur; and ii) for all possible single gene knockouts in LB under aerobic conditions. The predicted phenotypes were compared directly to the published phenotype data." [Note the published data are cited in the previous manuscript sentence, not shown here].

1. The link <https://github.com/rwick/GFA-dead-end-counter>, in line 286 does not work.

Link regenerated – now at line 451-452 and 604

Reviewer #2

1. *KpSC pan-metabolic reference model is provided. Are they required as input for Bactabolize? Are the gene, metabolite information open accessible by users? o See response to reviewer 1 point 2 above and;*

All data for the KpSC pan-model described in this work are accessible in the model files and amino acid + nucleotide files + data table at <https://github.com/kelwyres/KpSC-pan-metabolic-model>. This is also linked in the manuscript at line 631 and in the Data availability statement at line 661.

1. In the results section "description of Bactabolize", the authors present technical details on how to generate a metabolic model. For the input and output, please provide concrete examples to show the functionality of Bactabolize.

Detailed instructions, example code and example input/output files are available via the Bactabolize GitHub repository: <https://github.com/kelwyres/Bactabolize>.

Instructions and example code can be found on the wiki:

<https://github.com/kelwyres/Bactabolize/wiki> Test data and example files are at:

https://github.com/kelwyres/Bactabolize/tree/main/data/test_data

The Github repository is linked in the manuscript at lines 95, 124, 552, and 667, and we have added a further reference at line 124, which mentions the example code/data: "Full documentation, including example code and test data are available at the Bactabolize code repository (<https://github.com/kelwyres/Bactabolize>)."

1. To generate metabolic models, the authors present comparison results with other methods. However, the authors only present the numbers in genes, metabolites and substrates. Since the interactions between gene, metabolite, and substrate are also critical, if possible, please provide the coverage details about these interactions. Venn diagram is recommended to compare these coverage differences.

Two additional supplementary figures have been generated (Figures S5 and 6) showing Venn diagrams of metabolites and reactions for the highthroughput analysis approaches that are most relevant to this work (see also response to rev 1, point 1). These are discussed at lines 224-237:

"Figures S5 and S6 show the overlaps of metabolites and reactions between the high-throughput reconstruction methods after processing with MetaNetX (59) to standardise the reaction and metabolite nomenclatures (excluding CarveMe pan for simplicity and given the likely problems of reaction oversubscription). The majority of the reactions included in the Bactabolize model were conserved in either the CarveMe universal model (n = 1225, 53.2%), gapseq model (n = 54, 2.3%) or both (n = 665, 28.9%). The reaction overlap was skewed to the CarveMe universal model which shared 1225 reactions that were conserved in the Bactabolize model but absent from the gapseq model. Notably, the gapseq model contained a large number (2200) of unique reactions (70.4% of those in the model). Similarly, the vast majority of metabolites in the Bactabolize model were conserved in one or both of the other models (n = 917, 85.6%). However, it is likely that true overlaps between methods are underrepresented due to the different reaction identifiers and chemical synonyms used within the BiGG (Bactabolize, CarveMe) vs ModelSEED nomenclatures (gapseq), which are difficult to harmonise in an automated manner even after the application of MetaNetX."

Figure 2 shows not only the model numbers but also includes benchmarking to real phenotypic data in 2DEFG as the key mode of comparison between models. This encompasses meaningful interactions between gene, metabolic and substrate. The results are discussed at length in text at lines 253-271:

"We assessed the performance of each model for in silico prediction of growth phenotypes compared to the previously published experimental data (15). Accuracy, sensitivity, specificity, precision and F1 scores were calculated (60). Note that the specific set of growth

substrates and gene knockouts that can be simulated is determined by the sets of genes and metabolites captured by each model and is therefore model-dependent (Data S1 and S2). Among those with matched experimental phenotype data, the Bactabolize and CarveMe universal models were able to predict growth for a greater number of carbon, nitrogen, phosphorous and sulfur substrates than gapseq, CarveMe KpSC pan, KBase and iKp1289 models (Figure 2C, Data S1). While the CarveMe universal model had the highest number of true-positive growth predictions overall ($n = 132$ of 617 total predictions), it also had a comparably high number of false-positive predictions ($n = 39$ of 617 total predictions, Figure 2D). Similarly, the gapseq and iKp1289 models resulted in 31 (262 total predictions) and 50 (513 total predictions) false-positive predictions, respectively. In contrast, the Bactabolize model had fewer false-positive predictions ($n = 21$ of 505 total predictions) alongside a high number of true-positive predictions ($n = 117$ of 505 total predictions), resulting in the highest overall accuracy metrics (Figure 2E, Data S1). The KBase model was a notable outlier, associated with a high number of false-negative predictions ($n = 31$ of 103 total predictions) and low false-positive predictions ($n = 3$ of 103 total predictions), presumably resulting from the very low number of genes and reactions included in the model, driving low sensitivity and accuracy.”

Lines 272-280:

“The gene essentiality results showed that gapseq produced the highest absolute number of true-positive gene essentiality predictions ($n = 79$ of), followed by Bactabolize KpSC pan ($n = 44$ of 1220 total predictions), then CarveMe universal ($n = 39$ of 1951 total predictions). CarveMe universal had the largest number of true-negatives by a wide margin ($n = 1599$ of 1951 total predictions), followed by gapseq ($n = 1085$ of 1403 total predictions), then Bactabolize KpSC pan ($n = 939$ of 1220 total predictions), driving their high accuracies (83.96%, 82.96% and 80.57%, respectively). The Bactabolize model was associated with the greatest overall precision and specificity (Figures 2F & 2G) while the gapseq model resulted in the highest F1-score and sensitivity.”

1. *Are quality control and gap-filling needed to be processed when constructing a new metabolic model?*

Our goal here was to implement an approach to support high-throughput analyses (see response to rev 1 point 1), including leveraging draft genome assemblies as the bases for the construction of strain-specific metabolic models. As part of this work, we have described a robust quality control (QC) framework for screening draft *K. pneumoniae* genomes i.e. to identify genome assemblies that should not be used. We developed this framework by comparison to models generated for matched completed genomes. Our analyses demonstrate the importance of applying QC to the input draft genome assemblies. When appropriate QC is applied to the input genomes, the resultant draft models show a high degree of completeness compared to the matched models derived from complete genomes. The draft models can also be used to simulate growth phenotypes with high accuracy as compared to those simulated for the matched complete genome models.

No specific QC was applied to the draft models themselves, other than confirmation of positive growth prediction in m9 minimal media plus glucose (which is expected to support growth of all *K. pneumoniae*). In cases where the input assembly passed our QC criteria but the resultant model was unable to simulate growth in m9 minimal media plus glucose, gap-filling may be optionally applied. Again, by comparison to the simulated phenotypes from matched complete genome models, we show that these gap-filled draft models can produce accurate phenotype predictions. See lines 396-404:

“Of the 901 draft genome assemblies which passed our QC criteria (≤ 200 assembly graph dead ends), 23 of the resulting draft models failed to simulate growth in M9 minimal media with

glucose (despite capturing $\geq 99\%$ of the genes and reactions in the corresponding complete models). It is expected that all KpSC models should be able to simulate growth on M9 media with glucose as a sole carbon source, as this central metabolism is universal amongst KpSC. To replace missing, critical reactions required for growth on M9 with glucose, we investigated model gap-filling using the `patch_model` command of Bactabolize. We then assessed the accuracy of the gap-filled models for prediction of growth on the full range of substrates, as compared to the predictions from the corresponding complete models.” Lines 409-413: “Substrate usage predictions from the 21 successfully gap-filled models were highly accurate, with 18/21 having a prediction concordance of $\geq 99\%$ across all 846 growth conditions (12/21 had 100% concordance) (Figure S9). We therefore conclude that models generated for genome assemblies passing our QC criteria, which have been gap-filled to successfully simulate growth on minimal media plus glucose, are suitable for the prediction of growth across a range of substrates.”

1. *Are there any visualization results to check the status of the generated draft model?*

No. This is a tool for large-scale and rapid production of metabolic models, and phenotype prediction and we have not included visualisation tools. Third party tools are available e.g. <https://fluxer.umbc.edu/>. We do provide optional generation of MEMOTE reports at lines 136-138:

“Draft genome-scale metabolic models are output in both SMBL v3.1 (41) and JSON formats (one pair of files for each independent strain-specific model), along with an optional MEMOTE quality report (42)”.

Reviewer #3

1. *The justification and evaluation of the generated models are inadequate and onedimensional. The authors only focus on statistics such as the number of reactions and genes in the models, which does not accurately depict the completeness of the model.*

The reviewer has misunderstood how we have used ‘completeness’ in this manuscript. In the section describing our novel QC framework, we use this term to refer to the relative completeness of draft models generated from draft genome assemblies as compared to curated models generated from complete genome assemblies for the same strains. The latter were considered as the ‘complete’ models for this purpose. We are not referring to any measure of network or metabolic pathway completeness. We specifically refer to gene and reaction capture compared to the ‘complete’ models because these features directly reflect the problem we are trying to address i.e. that draft genome assemblies may not contain the complete set of genes that are truly present in the underlying genome. We have updated the manuscript text to further clarify the problem we aim to address in this section and justify the use of gene and reaction capture metrics:

Lines 310-319: “There are now thousands of bacterial genomes available in public databases, the majority of which are in draft form, comprising 10s to 1000s of assembly contigs. This fragmentation of the genome is caused by repetitive sequences that cannot be resolved by the assembly algorithm and/or sequence drop-out. The latter can result in the loss of genetic information such that some portion of genes present in the underlying genome are lost from the genome assembly (either completely or partially). This in turn, poses a limitation for the reconstruction of metabolic models using these assemblies, since most published approaches use sequence searches to predict the presence/absence of genes and their associated enzymatic reactions. Therefore, if we are to use public genome data for high-throughput

metabolic modelling studies, it is essential to evaluate the expected model accuracies and understand the minimum input genome quality requirements.”

The biological accuracy of the curated ‘complete’ models has been described previously, and this is now noted in the text at lines 320-324:

“Here we performed a systematic analysis leveraging our published curated KpSC models (n=37, (14)), which were generated using completed genome sequences and were therefore considered to represent ‘complete’ models for which the underlying genome sequence contains all genes that are truly present in the genome (note the biological accuracy of these models was reported previously (14) and is not the subject of the current study).”

Throughout the manuscript we not only compare models in terms of the numbers of genes and reactions, but through comparison of binary growth predictions. Specifically, in the Performance Comparison section (Bactabolize vs other approaches) we use comparison of predicted to experimental phenotypes for strain KPPR1 (see response to rev 1 point 4 for details). In the QC Framework section we compare the predictions derived from draft models generated from draft genome assemblies to those derived from the matched ‘complete’ models, and report the concordance as a measure of impact of input assembly quality (lines 309-394). In the final results section (Predictive accuracy of draft models), we generate 10 additional models and compare the growth predictions to matched experimental data (lines 414-433). We view these phenotype prediction comparisons as the ultimate measure of ‘completeness’ with which to assess our models, because these data have direct biological meaning.

1. The authors have not provided evidence or discussion on the accuracy of any metabolic fluxes, which are considered to be crucial for reconstructing metabolic models. Additionally, the authors have not mentioned the importance of non-growth associated maintenance and the criticality of biomass composition analysis, both of which significantly determine the fluxes in the system.

We acknowledge the importance of flux calculations and accurate biomass compositions when using genome-scale models to quantitatively predict growth rates. However, at this stage, the reconstructions developed using Bactabolize are intended for binary predictions and comparisons of growth capabilities on various substrates. The accuracies we report are based on measures of network completion (presence/absence of relevant reactions leading to growth or no-growth phenotypes) rather than specific growth rates. Thus, the models generated by Bactabolize can be used to explore diversity at the strain level in terms of growth capabilities and can serve as a scaffold for building detailed (customized biomass), strain-specific models. Measuring biomass composition and metabolic flux analysis require significant experimental comparisons that are outside the scope of the current study but could be performed for target strains based on reconstructions developed using Bactabolize.

1. It would be interesting to compare the accuracy of the models generated using Bactabolize with those manually curated.

We did exactly this. We compared the manually curated model iKp1289 as part of our benchmarking. Lines 194 – 199:

“We compared the output and performance of Bactabolize to the two previously published tools that can support high-throughput analyses i.e. CarveMe (30) and gapseq (31). To aid interpretation in the context of community standard approaches, we also include a comparison to the popular web-based reconstruction tool, KBase (ModelSEED), and a manually curated metabolic reconstruction of *K. pneumoniae* strain KPPR1 (also known as VK055 and ATCC 43816, metabolic model named iKp1289) (15).”

Unfortunately, as far as we aware there are currently no other published manually curated models for strains with matched phenotype data that are also not included as part of our pan-reference model (the latter is a key point to ensure a fair comparison of models generated using our pan-reference vs those generated with a universal reference).

1. *The authors have not provided evidence or discussion on the accuracy of any metabolic fluxes, which are considered to be crucial for reconstructing metabolic models.*

See response to rev 3, point 2.

1. *The justification regarding the completeness of the models requires further discussion.*

See response to rev 3, point 1.

1. *A detailed discussion on the importance of manually curated models would significantly enhance the quality of the manuscript.*

This has been added at lines 458-474:

“Traditionally, genome-scale metabolic reconstruction approaches have relied upon significant manual curation efforts. While there will always remain a need for high quality curated models, such resource intensive approaches preclude their application at scale, and have therefore limited analyses to small numbers of individual strains (15, 16). However, automated reconstruction approaches can support the generation and comparison of multiple strain-specific draft models from which meaningful biological insights can be derived (61). Additionally, the quality of curated models is likely to vary depending on their age, level and type of curation, as well as the approach used for preliminary drafting. Indeed it is possible for automated approaches to outperform manually curated models; a draft model for *K. pneumoniae* KPPR1 generated using Bactabolize with the KpSC pan-v1 reference model outperformed the manually curated iKp1289 model representing the same strain (15). iKp1289 was published in 2017 (6 years prior to this study) and was initially drafted via the KBase pipeline (33), which uses RAST to annotate the sequences with Enzyme Commission numbers. It has been demonstrated several times that the Enzyme Commission scheme has systematic errors (62, 63), leading to a loss in accuracy when compared to the ortholog identification methods used by automated approaches. Consistent with this assertion, our draft KPPR1 model constructed with KBase (without manual curation) was an outlier in terms of the very low number of genes, reactions and metabolites that were included.”