

OsNF-YB7 inactivates OsGLK1 to inhibit chlorophyll biosynthesis in rice embryo

Reviewed Preprint

v2 • August 29, 2024

Revised by authors

Reviewed Preprint

v1 • March 22, 2024

Zongju Yang, Tianqi Bai, Zhiguo E, Baixiao Niu, Chen Chen ✉

Jiangsu Key Laboratory of Crop Genomics and Molecular Breeding/ Zhongshan Biological Breeding Laboratory, Agricultural College of Yangzhou University, Yangzhou, China • Jiangsu Co-Innovation Center for Modern Production Technology of Grain Crops/ Key Laboratory of Plant Functional Genomics of the Ministry of Education, Agricultural College of Yangzhou University, Yangzhou, China • State Key Laboratory of Rice Biology and Breeding, China National Rice Research Institute, Hangzhou, China

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

Abstract

As a master regulator of seed development, Leafy Cotyledon 1 (LEC1) promotes chlorophyll (Chl) biosynthesis in *Arabidopsis*, but the mechanism underlying this remains poorly understood. Here, we found that loss of function of *OsNF-YB7*, a *LEC1* homolog of rice, leads to chlorophyllous embryo, indicating that *OsNF-YB7* plays an opposite role in Chl biosynthesis in rice compared with that in *Arabidopsis*. *OsNF-YB7* regulates the expression of a group of genes responsible for Chl biosynthesis and photosynthesis by directly binding to their promoters. In addition, *OsNF-YB7* interacts with Golden 2-Like 1 (OsGLK1) to inhibit the transactivation activity of OsGLK1, a key regulator of Chl biosynthesis. Moreover, *OsNF-YB7* can directly repress *OsGLK1* expression by recognizing its promoter *in vivo*, indicating the involvement of *OsNF-YB7* in multiple regulatory layers of Chl biosynthesis in rice embryo. We propose that *OsNF-YB7* functions as a transcriptional repressor to regulate Chl biosynthesis in rice embryo.

eLife assessment

This paper provides **important** insights into the role of rice *OsNF-YB7*, an ortholog of *Arabidopsis* LEC1, in chlorophyll biosynthesis, uncovering the genetic and molecular basis for negative regulation of chlorophyll production in the rice embryo. Mutational analysis, gene expression profiles and protein interaction combine for **convincing** evidence that *OsNF-YB7* represses chlorophyll biosynthesis.

<https://doi.org/10.7554/eLife.96553.2.sa4>

Introduction

Angiosperms can be divided into chloroembryophytes and leucoembryophytes, depending on the presence or absence of chlorophyll (Chl) in the embryo, respectively (Puthur et al., 2013; Smolikova and Medvedev, 2016). Some plant species such as *Arabidopsis* (*A. thaliana*) produce chloroembryos, which have photochemically active chloroplasts capable of producing assimilates that are further converted into reserve biopolymers (Simkin et al., 2020). Grass species, such as rice (*Oryza sativa*), are unable to produce Chl during embryo development. To the best of our knowledge, it remains completely unclear what determines the ability to biosynthesize Chl in plant embryo. However, studies have identified several genes that contribute to Chl degradation for chloroembryos, the mutation of which can lead to a stay-green phenotype in mature seeds such as Mendel's green pea (Sato et al., 2007; Delmas et al., 2013; Wang et al., 2018; Li et al., 2017; Thomas and Ougham, 2014).

Golden 2-Like (GLK) genes encode GARP-type transcription factors (TFs), which are key components regulating chloroplast development and Chl biosynthesis in plants (Rossini et al., 2001; Fitter et al., 2002; Nakamura et al., 2009). GLK can recognize the CCAATC cis-element of the Chl biosynthesis- and photosynthesis-associated nuclear genes to trigger their expression (Waters et al., 2009). The G-box (CACGTG) is also enriched in the GLK-targeted genes in *Arabidopsis*, possibly due to GLK being able to interact with certain G-box binding factors (Tamai et al., 2002). A genetic study showed that GLK activates Chl biosynthesis in roots, in a manner dependent on the G-box binding TF Elongated Hypocotyl 5 (HY5) (Kobayashi et al., 2012).

There are two copies of *GLK* in the rice genome, designated as *OsGLK1* and *OsGLK2* (Fitter et al., 2002). They redundantly regulate a set of genes, such as rice *Chlorophyllide A Oxygenase* (*OsCAO*) and *Protochlorophyllide Oxidoreductase A* (*OsPORA*) responsible for Chl biosynthesis, and *Light Harvesting Complex B1* (*OsLHCB1*) and *OsLHCB4* responsible for photosynthesis (Nakamura et al., 2009; Wang et al., 2013; Sakuraba et al., 2017). *OsGLK1* overexpression in rice leads to green calli and chloroplast development in the vascular bundles (Nakamura et al., 2009; Wang et al., 2017). Organelle development in rice vascular sheath cells is induced by ectopically expressed maize (*Zea mays*) *GLK* genes, mimicking a key step in the evolutionary transition from C3 to C4 plants (Wang et al., 2017). In accordance with this, the ectopic expression of maize *GLKs* in rice can boost biomass and grain yield by facilitating chloroplast development and photosynthesis (Li et al., 2020; Yeh et al., 2022). Moreover, the overexpression of maize *GLKs* in calli was shown to improve the ability of rice and maize to regenerate (Luo et al., 2022). A recent study found that *OsGLK1* also participates in tapetum plastid development and programmed cell death, consequently affecting pollen fertility in rice (Zheng et al., 2022). The findings indicate that GLK plays multiple roles in relation of plant development.

Leafy Cotyledon 1 (LEC1), a member of the nuclear factor Y (NF-Y) TF family, is a central regulator controlling many aspects of seed development, such as Chl accumulation in the embryo (Meinke et al., 1994; Pelletier et al., 2017). LEC1 can also act as a pioneer TF to regulate flowering by reprogramming the embryonic chromatin state (Tao et al., 2017). Previous studies have reported that *lec1* mutants have paler green embryos than wild-type (WT) at maturation in *Arabidopsis* (Meinke et al., 1994; West et al., 1994). LEC1 transcriptionally regulates the expression of genes that encode light-reaction components of photosystems I and II, as well as the expression of genes involved in chloroplast biogenesis in *Arabidopsis* and soybean embryos (Pelletier et al., 2017; Jo et al., 2020). These findings suggest that LEC1 is important for photosynthesis and chloroplast development during seed development. However, the underlying molecular mechanisms remain largely unclear.

It is still a mystery why plant species such as rice cannot synthesize chlorophyll in embryos, while species such as *Arabidopsis* can. There are two LEC1 homologs, *OsNF-YB7* and *OsNF-YB9*, encoded by the rice genome (E et al., 2018). *OsNF-YB7* is restricted to the embryo, and defective *OsNF-YB7* may result in seed lethality (Niu et al., 2021b; Zhang and Xue, 2013). Here, we found that *OsNF-YB7* acts as a key inhibitor of Chl biosynthesis in rice embryo. It inactivates the transactivation activity of *OsGLK1*, at multiple regulatory layers, to inhibit Chl accumulation in the embryo of rice, explaining the achlorophyllous embryo produced in rice.

Results

Loss of function of *OsNF-YB7* leads to chloroembryos

By observing seeds produced by the loss-of-function mutant of *OsNF-YB7*, we surprisingly found that the *osnf-yb7* embryo was greenish (Fig. 1A and B), suggesting that *OsNF-YB7* plays a negative role in chloroplast biogenesis or Chl biosynthesis, or both, during embryogenesis. We therefore examined the phenotype of green embryos at various seed development stages. The results showed that WT had an achlorophyllous embryo throughout embryonic development, whereas *osnf-yb7* embryo turned green at 5 days after fertilization (DAF) and the chloroembryo remained green until maturity (Fig. S1A–C). In support of this, transmission electron microscopy (TEM) showed well-developed chloroplast in the scutellum tip of the mutant, but this was not seen in the WT sections (Fig. 1C). By measuring Chl contents in the developing and mature embryos, we found that total Chl content in *osnf-yb7* was consistently higher than that in the WT, both at 10 DAF and at maturation (30 DAF) (Fig. 1D). As revealed by confocal laser scanning microscopy (CLSM), Chl autofluorescence was detectable as early as 5 DAF in *osnf-yb7*, indicating the initiation of Chl accumulation at this stage (Fig. 1E). This is consistent with our previous finding that *OsNF-YB7* was highly activated at 5 DAF in rice embryo (Niu et al., 2021b).

Light is a critical signal triggering Chl biosynthesis (Wietrzynski and Engel, 2021). To determine whether the Chl biosynthesis in *osnf-yb7* is induced by light, we investigated the embryo phenotype of WT and *osnf-yb7* in the dark, using aluminum foil to cover rice panicles prior to flowering. The result showed that, similar to WT, *osnf-yb7* embryo was achlorophyllous in the dark, although the embryogenesis defects, such as degenerated epiblast and coleorhiza, and maldeveloped coleoptile, were still observable (Fig. S2A–H). This indicated that Chl biosynthesis in the mutant is light dependent. Only a small amount of light can be perceived by rice embryos because the external hulls block light penetration (Simkin et al., 2020). We removed the hulls to directly expose the embryo to light; however, Chl accumulation still failed to occur in the WT embryo (Fig. S2I–L), suggesting that there are internal signals that repress Chl biosynthesis in rice embryo. We suspected that the *OsNF-YB7* mutation may attenuate the activity of such inhibitors; alternatively, *OsNF-YB7* itself could be an inhibitor.

OsNF-YB7 negatively regulates photosynthesis- and Chl biosynthesis-related genes

Using 5- and 10-DAF-old WT and *osnf-yb7* embryos, we previously performed deep sequencing of the transcriptome (RNA-seq) to identify possible downstream genes of *OsNF-YB7* (Niu et al., 2021a). As revealed by Mapman analysis, photosynthesis-related pathways, such as photosystem I, photosystem II, and the light reaction, were significantly enriched for the differentially expressed genes (DEGs) in the mutant embryos (Fig. 1F, Fig. S3A and B). Moreover, 96.9% (95/98) and 96.4% (106/110) of the photosynthesis-related DEGs that identified from 5- and 10-DAF-old embryos, respectively, were upregulated in the mutant (Fig. 1G). To confirm this finding, we next examined the expression of *OsLHCAs* and *OsLHCBs*, which are primarily associated with photosystems I and II, respectively, via quantitative real-time PCR (RT-qPCR). It showed higher expression of all of the studied *OsLHCAs* and *OsLHCBs* in the *osnf-yb7*




Figure 1.

OsNF-YB7 negatively regulates Chlorophyll (Chl) biosynthesis in embryo.

A, B. Morphologies of wild-type (WT) and *osnf-yb7* detached embryos at 10 days after fertilization (DAF) (**A**) and longitudinally dissected embryos at maturation (**B**). Scale bars = 200 μ m.

C. Transmission electron microscopy images of embryos from WT and *osnf-yb7* at 10 DAF. Scale bars = 5 μ m.

D. Chl levels in WT and *osnf-yb7* embryos at 10 DAF and maturation. Data are means $\pm$ SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

E. Chl autofluorescence of WT and mutant embryos at 5 DAF. BF, bright field; Chl, Chl autofluorescence. Scale bars = 100 μ m.

F. Photosynthesis-related pathways enriched among the differentially expressed genes (DEGs) identified from the 5- and 10-DAF-old *osnf-yb7* embryos compared to WT, respectively.

G. Most of the photosynthesis-related genes were upregulated in the mutant embryos.

H. A heat map shows the expression of the Chl biosynthesis-related genes in the WT and *osnf-yb7* embryos at 5 and 10 DAF. Reads per kilobase per million mapped reads (RPKM) is used to indicate the expression level.

embryo at 10 DAF (**Fig. S4A**). Likewise, many of the genes participating in Chl biosynthesis, including rice *Genomes Uncoupled 4* (*OsGUN4*), *Mg-Chelatase H Subunit* (*OsCHLH*), *OsCHLI*, *OsCHLD*, *Copper Response Defect 1* (*OsCRD1*), *OsPORA*, *OsPORB*, and *Divinyl Reductase* (*OsDVR*), were significantly activated in the mutant (**Fig. 1H** and **Fig. S4B**). The findings indicated that OsNF-YB7 might act as a repressor of Chl biosynthesis and photosynthesis, which is opposite to the role of its homologue LEC1 in *Arabidopsis*.

Since OsNF-YB7 is a TF, we assumed that it may directly regulate the expression of genes related to Chl biosynthesis and photosynthesis, such as *OsPORA*, and *OsLHCB4*, which were significantly activated in the *osnf-yb7* embryo (**Fig. 2A and B**). We first generated transgenic lines that overexpressed *OsNF-YB7*, tagged with either green fluorescent protein (*NF-YB7-GFP*) or 3× Flag (*NF-YB7-Flag*) in the Zhonghua11 (ZH11, *O. sativa* ssp. *gengljaponica*) or Kitaake (*O. sativa* ssp. *gengljaponica*) background, respectively. As expected, *OsPORA* and *OsLHCB4* were significantly downregulated in leaves of the *OsNF-YB7* overexpressors (**Fig. 2C and D**, and **Fig. S4C**). Similar to previously reported (Zhang and Xue, 2013; Ito et al., 2011), the *OsNF-YB7* overexpression lines displayed severe reproductive development defects, which prevented us from obtaining sufficient embryo tissues for subsequent experiments. Instead, using the *NF-YB7-Flag* seedling, we conducted a chromatin immunoprecipitation assay coupled with quantitative PCR (ChIP-qPCR). The results showed that OsNF-YB7 was highly enriched in the promoter regions of *OsPORA* and *OsLHCB4* harboring the G-box motif (**Fig. 2E and F**), a putative binding site of OsNF-YB7 (Guo et al., 2022). To confirm the ability of OsNF-YB7 to bind to the *OsPORA* and *OsLHCB4* promoters, we next performed electrophoretic mobility-shift assays (EMSAs), using recombinant OsNF-YB7-His protein, and biotin-labeled subfragments of the *OsPORA* or *OsLHCB4* promoters containing G-boxes. The results showed that OsNF-YB7-His was able to bind to the labeled probes, and the shifted band signals were substantially weakened upon application of the unlabeled cold probes or hot probes with a mutated G-box (**Fig. 2G and H**).

To confirm that the binding of OsNF-YB7 represses target gene expression, dual luciferase (LUC) reporter (DLR) assays were performed. We first generated LUC reporters driven by the *OsPORA* or *OsLHCB4* promoters (designated as *proOsPORA:LUC* and *proOsLHCB4:LUC* hereafter) (**Fig. 2I**). When the reporters were coexpressed with OsNF-YB7, which acted as an effector, in rice protoplasts, we found that OsNF-YB7 significantly repressed the activity of both *proOsPORA:LUC* and *proOsLHCB4:LUC* (**Fig. 2J**). Taken together, these results suggested that OsNF-YB7 directly binds to the promoters of photosynthesis- and Chl biosynthesis-related genes and represses their transcription.

OsNF-YB7 represses *OsGLK1* in the embryo

Several TFs that regulate Chl biosynthesis or chloroplast development have been identified in plants (Jarvis and López-Juez, 2013). Some of these, such as rice *OsGLK1*, *HY5-like 1* (*OsHYSL1*), *PIF-like 14* (*OsPIL14*), and *GATA Nitrate-inducible Carbon-metabolism-involved* (*OsGNC*), were found to be upregulated in the embryo of *osnf-yb7* (**Fig. 3A**).

OsGLK1 was the most strikingly activated TF (**Fig. 3A and B**), while it was significantly repressed in *NF-YB7-GFP* (**Fig. 3C**). To test the idea that *OsGLK1* is a direct downstream target of OsNF-YB7, a ChIP-qPCR assay was first carried out using *NF-YB7-Flag* transgenic seedlings. The results showed that the promoter segments P3 and P4 were significantly enriched in the immunoprecipitated chromatin (**Fig. 3D**), suggesting that OsNF-YB7 was able to bind to the *OsGLK1* promoter *in vivo*. However, we failed to validate the binding ability *in vitro* using the EMSA assay (**Fig. S5**), suggesting that OsNF-YB7 requires other NF-Ys or TFs to form a TF complex to recognize the promoter, as many NF-Y members do for function (Laloum et al., 2013). A DLR assay was next performed to investigate the negative regulation of OsNF-YB7 on *OsGLK1* transcription using rice protoplasts. The *OsGLK1* promoter was inserted upstream of the 5× upstream activating sequence (UAS) as a reporter; OsNF-YB7 was fused C-terminally to the DNA binding domain (BD) of the yeast GAL4 and the herpes virus VP16 transactivation domain




Figure 2.

OsNF-YB7 binds to the promoters of *OsPORA* and *OsLHCB4* to regulate their expression.

A, B. Quantitative real-time PCR (RT-qPCR) analysis of the transcription levels of *OsPORA* (**A**) and *OsLHCB4* (**B**) in the embryos of WT and *osnf-yb7* at 10 DAF. Data are means $\pm$ SD ($n = 3$). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

C, D. Expression of *OsPORA* (**C**) and *OsLHCB4* (**D**) in the leaves of WT and *OsNF-YB7*-overexpressing transgenic plants (*NF-YB7-GFP*). Data are means $\pm$ SD ($n = 3$). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

E, F. Chromatin immunoprecipitation assay coupled with quantitative PCR (ChIP-qPCR) analyses showing the enrichment of *OsNF-YB7* at the *OsPORA* (**E**) and *OsLHCB4* (**F**) promoters in 14-day-old *OsNF-YB7-Flag* seedlings. Precipitated DNA was quantified by qPCR and DNA enrichment is displayed as a percentage of input DNA. Data are means $\pm$ SD ($n = 3$). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene. Diagrams in the upper panel showing the promoter structures of *OsPORA* and *OsLHCB4*, and the PCR amplicons used for ChIP-qPCR.

G, H. Electrophoretic mobility-shift assays (EMSAs) showing that *OsNF-YB7* directly binds to the promoters of *OsPORA* (**G**) and *OsLHCB4* (**H**). Hot probes were biotin-labeled. The hot mProbes contain mutant nucleic acid from CACATG to AAAAAA. The arrow heads indicate the shift bands.

I. Schematic diagram displaying the constructs used in the dual luciferase reporter (DLR) assays of **J**. LUC, firefly luciferase; REN, *Renilla* luciferase.

J. DLR assays showing that *OsNF-YB7* directly represses the promoter activities of *OsPORA* and *OsLHCB4*. Data are means $\pm$ SD ($n = 3$). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis.




Figure 3.

OsNF-YB7 associates with the promoter of *OsGLK1* and represses its expression.

A. A heat map showing the expression of transcription factors associated with Chl biosynthesis and chloroplast development in the 5- and 10-DAF-old embryos of WT and *osnf-yb7*. The colored dots indicate $\log_2$ (RPKM mean) of the genes in three biological replicates.

B. RT-qPCR analysis of *OsGLK1*, *OsGNC*, and *OsPIL14* expression levels in the embryos from WT and *osnf-yb7* at 5- and 10 DAF. Numbers represent fold changes of expression. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

C. RT-qPCR analysis of *OsGLK1* expression levels in leaves from WT and *NF-YB7-GFP*. Data are means $\pm$ SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

D. ChIP-qPCR analysis showing the enrichment of OsNF-YB7 at the *OsGLK1* promoter in 14-day-old *OsNF-YB7-Flag* seedling. Chromatin of each sample was immunoprecipitated using anti-Flag or IgG antibodies. Precipitated DNA was quantified by qPCR and DNA enrichment is displayed as a percentage of input DNA. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene. The experiment was performed three times with similar results. The diagram in the upper panel showing the promoter structure of *OsGLK1* and PCR amplicons (P1, P2, P3, P4, and P5) used for ChIP-qPCR.

E. Schematic diagram displaying the constructs used in the DLR assays of **F**. LUC, firefly luciferase; REN, *Renilla* luciferase; UAS, upstream activating sequence.

F. DLR assays showing that OsNF-YB7 represses the promoter activity of *OsGLK1*. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; Student's *t*-test was used for statistical analysis.

(VP16:OsNF-YB7), as an effector (Fig. 3E). Coexpression of VP16:OsNF-YB7 with the reporter in rice protoplasts significantly decreased the transcriptional activity of VP16 (Fig. 3F), indicating that OsNF-YB7 represses *OsGLK1* promoter activity.

OsGLK1 is involved in OsNF-YB7-regulated Chl biosynthesis in embryo

To confirm the contribution of *OsGLK1* for the production of chloroembryo in *osnf-yb7*, we first generated *OsGLK1*-overexpressing lines (*OsGLK1-OX*) driven by the rice ubiquitin promoter. In association with over-accumulated Chl in the glume and seed coat of the transformant, RT-qPCR and Western blot assays confirmed *OsGLK1* activation in *OsGLK1-OX* (Fig. S6A–C). As observed in *osnf-yb7*, green embryos were produced (Fig. 4A). The Chl content in *OsGLK1-OX* were higher than that in WT (Fig. 4B and C), suggesting that *OsGLK1* overexpression in rice embryo induces Chl biosynthesis.

Using a clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated 9 (Cas9) cassette including three tandemly arrayed guide RNAs targeting *OsNF-YB7*, *OsGLK1*, and *OsGLK2*, respectively, we successfully obtained the *osnf-yb7;osglk2* double mutant and the *osnf-yb7;osglk1;osglk2* triple mutant (Fig. S7A and B). By analyzing the embryos that the mutants produced, we found that significantly less Chl accumulated in *osnf-yb7;osglk1;osglk2* than in *osnf-yb7* and *osnf-yb7;osglk2* (Fig. 4D–G and Fig. S8). In comparison to the achlorophyllous embryo of WT, the *osnf-yb7;osglk1;osglk2* triple mutant still showed somewhat green coloration in the apical part of the embryos (Fig. 4G). We hypothesized that this was at least partially due to that genes like *OsLHCB4* and *OsPORA* can be induced by the mutation of *OsNF-YB7*, given the fact that OsNF-YB7 represses the genes' expression independent of *OsGLK1* (Fig. 2J).

OsNF-YB7 and OsGLK1 regulate a common set of genes in the embryo

We next performed RNA-seq analysis to explore the transcriptomic changes in the chloroembryo of *OsGLK1-OX* at 10 DAF. More than 64.4% of the DEGs identified in *OsGLK1-OX* were overlapped with the ones identified in *osnf-yb7*, the vast majority of which were either activated (60.6%) or repressed (27.6%) in both *OsGLK1-OX* and *osnf-yb7* (Fig. 4H and I, and Table S1 and S2). As revealed by the Gene Ontology (GO) analysis, genes involved in Chl biosynthesis and photosynthesis, such as *OsPORA* and *OsLHCB4*, were markedly enriched among the common DEGs (Fig. 4J and Fig. S9), implying that OsNF-YB7 and *OsGLK1* antagonistically regulate a common set of genes for Chl biosynthesis and photosynthesis.

The ChIP-qPCR results showed that *OsGLK1* associated with the regions of the *OsPORA* and *OsLHCB4* promoters to which OsNF-YB7 binds (Fig. 5A and B). In agreement with this, the EMSA results suggested that *OsGLK1* directly binds to the same DNA probes of *OsPORA* and *OsLHCB4* *in vitro* (Fig. 5C and D). In opposite to OsNF-YB7, when we coexpressed the *OsGLK1* effector vector with the reporter vector *proOsLHCB4:LUC* or *proOsPORA:LUC* in rice protoplasts, *OsGLK1* showed significant transactivation activity on *OsPORA* and *OsLHCB4* (Fig. S10A and B).

Recently, the putative binding sites of OsNF-YB7 and *OsGLK1* were investigated at the whole genome scale (Guo et al., 2022; Tu et al., 2022). This allows us to test our hypothesis that OsNF-YB7 and *OsGLK1* can target a common set of genes involved in Chl biosynthesis and photosynthesis. By reanalyzing the ChIP-seq data, we found that 91.4% (235/257) and 90.7% (167/184) of the *OsGLK1*-binding genes overlapped with the OsNF-YB7-binding genes in two replicates (Fig. 5E and Table S3), although the peak number of OsNF-YB7 was much greater than that of *OsGLK1* (Guo et al., 2022; Tu et al., 2022). A large number of the common targets were activated in the embryos of *osnf-yb7* and *OsGLK1-OX* at 10 DAF (Fig. S11A). GO analysis suggested that most of the common targets were genes involved in Chl biosynthesis or




Figure 4.

Chl biosynthesis in *osnf-yb7* embryo requires active *OsGLKs*.

- A.** Embryo morphologies of WT and *OsGLK1-OX* detached embryos at 10 DAF. Scale bars = 2 mm.
- B.** Chl autofluorescence of the WT and *GLK-OX* embryos at 10 DAF. BF, bright field; Chl, Chl autofluorescence. Scale bars = 100 μ m.
- C.** Chl levels in WT and *OsGLK1-OX* embryos at maturation. Data are means $\pm$ SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.
- D-G.** Morphologies of the embryos produced by WT (**D**), *osnf-yb7* (**E**), *osnf-yb7;osglk2* double mutant (**F**) and *osnf-yb7;osglk1;osglk2* triple mutant (**G**). Scale bars = 1 mm.
- H.** A Venn diagram showing overlaps of the DEGs identified from the embryos of *osnf-yb7* and *OsGLK1-OX* at 10 DAF.
- I.** A pie chart showing similar transcriptional changes of the common DEGs identified from *osnf-yb7* and *OsGLK1-OX*.
- J.** Gene Ontology (GO) analysis of the common DEGs identified from *osnf-yb7* and *OsGLK1-OX*.




Figure 5.

OsNF-YB7 and OsGLK1 regulate a common set of genes in the embryo.

A, B. ChIP-qPCR analysis showing the enrichment of OsGLK1 on the *OsPORA* (**A**) and *OsLHCB4* (**B**) promoters. PCR amplicons used for ChIP-qPCR are indicated in the schematic diagrams. OsGLK1-GFP was transiently expressed in protoplasts isolated from green tissues of the 14-day-old WT seedling. Chromatin of each sample was immunoprecipitated using anti-GFP or IgG antibodies. Precipitated DNA was quantified by qPCR and DNA enrichment is displayed as a percentage of input DNA. Data are means $\pm$ SD ($n = 3$). *, $p < 0.05$; **, $p < 0.01$; Student's t -test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene. The experiment was performed three times with similar results.

C, D. EMSA assays showing that OsGLK1 directly binds to the promoters of *OsPORA* (**C**) and *OsLHCB4* (**D**). The arrow heads indicate the shift bands.

E. A Venn diagram showing overlaps of the target genes between OsGLK1 and OsNF-YB7.

F. Distribution of the distance between the OsNF-YB7 and OsGLK1 summits showing that OsNF-YB7 and OsGLK1 bind to proximal regions of their common targets.

photosynthesis (**Fig. S11B**). By retrieving the sequences of the OsNF-YB7 and OsGLK1 binding peaks in the common targets for MEME analysis, we found that the TFs probably recognize similar DNA motifs. A short sequence containing the G-box motif was the most significantly enriched (**Fig. S12A and B**). There was also enrichment of another sequence containing a CCAAT motif recognized by the NF-Y TF complexes and a CCAATC motif recognized by GLKs, as previously reported (Waters et al., 2009 [↗](#); Pelletier et al., 2017 [↗](#)) (**Fig. S12A and B**). The results suggested that OsNF-YB7 and OsGLK1 can bind to the same region of their common target. In agreement with this, approximately 75.0% and 76.5% of the peak summits of OsGLK1 were located proximally to the summit of OsNF-YB7 in two replicates, within an adjacent region no more than 100 bp away (**Fig. 5F [↗](#) and Table S3**). Consistent with our biochemical evidence, the ChIP-seq results showed that OsNF-YB7 does bind to the same regions of the *OsPORA* and *OsLHCB4* promoters as OsGLK1 binds to (**Fig. S12C**).

OsNF-YB7 physically interacts with OsGLK1 to inhibit its transcriptional activity

To resolve how OsNF-YB7 and OsGLK1 bind to the same regions to regulate their common targets, we speculated that the TFs probably form a dimer in rice in order to exert their functions. We therefore carried out a yeast two-hybrid (Y2H) assay by transforming yeast cells with a bait construct expressing OsNF-YB7 fused with the GAL4 DNA-binding domain (BD), together with a prey construct expressing OsGLK1 fused with the yeast GAL4 activation domain (AD). The results showed that OsNF-YB7 interacts with OsGLK1 in yeast (**Fig. 6A and B [↗](#)**). Furthermore, to determine the functional domains required for the interaction, we generated two truncated versions of OsNF-YB7, which contained the N- or C-terminal, and three truncated versions of OsGLK1, which contained the N-terminal, DNA-BD, or GCT box domain, as previously reported (Fitter et al., 2002 [↗](#); Zhang et al., 2021b [↗](#)) (**Fig. 6A [↗](#)**). We found that the C-terminus of OsNF-YB7 was sufficient for the interaction (**Fig. 6B [↗](#)**). In addition, the GCT box domain of OsGLK1 strongly interacted with the full length or C-terminus of OsNF-YB7, while the DNA-BD of OsGLK1 showed a weak interaction (**Fig. 6B [↗](#)**).

A split complementary LUC assay further confirmed the interaction between OsNF-YB7 and OsGLK1 in the epidermal cells of *Nicotiana benthamiana* (**Fig. 6C [↗](#)**). As suggested by the bimolecular fluorescence complementation (BiFC) analysis, the interaction occurred exclusively in the nuclei of *N. benthamiana* epidermal cells (**Fig. 6D [↗](#)**). Moreover, we transiently coexpressed OsNF-YB7 tagged with GFP (OsNF-YB7-GFP) and OsGLK1 tagged with 3× Flag (OsGLK1-Flag) in rice protoplasts, and co-immunoprecipitation (Co-IP) analysis showed that OsGLK1-Flag could be immunoprecipitated by the anti-GFP antibody (**Fig. 6E [↗](#)**), indicating that the interactions do occur *in vivo*. In addition, both Y2H and split complementary LUC assays showed that OsGLK2 could interact with OsNF-YB7 (**Fig. S13A and B**). These findings indicated that OsNF-YB7 interacts with OsGLKs, explaining why OsNF-YB7 and OsGLKs share a common set of targets in rice.

To explore the biological meaning of the interaction, we next performed DLR assays using rice protoplasts. Transient expression of OsGLK1 in rice protoplasts substantially activated the reporters, driven by either the *OsPORA* or the *OsLHCB4* promoter; however, when we coexpressed OsNF-YB7 with OsGLK1, the transactivation ability of OsGLK1 was significantly repressed in rice protoplasts (**Fig. 6F [↗](#)**). These findings suggested that OsNF-YB7/OsGLK1 dimerization reduces the transactivation ability of OsGLK1 for fine-tuning the Chl biosynthetic and photosynthetic genes, such as *OsPORA* and *OsLHCB4*. The EMSA assay showed that OsGLK1-MBP recombinant proteins could bind to the promoter of *OsPORA*; however, when we incubated the probe with OsGLK1-MBP and OsNF-YB7-His together, the binding ability of OsGLK1 substantially decreased (**Fig. 6G [↗](#)**). As a control, when the probe was incubated with OsGLK1-MBP and GST-His, the binding ability of OsGLK1 remained unchanged (**Fig. 6G [↗](#)**). These findings implied that the reduced transactivity of OsGLK1 is likely due to the formation of OsNF-YB7/OsGLK1 heterodimers inhibiting the binding of OsGLK1 to its downstream genes.




Figure 6.

OsNF-YB7 interacts with OsGLK1 to regulate the expression of *OsPORA* and *OsLHCB4*.

A. Schematic diagrams showing the protein structures of OsGLK1 and OsNF-YB7.

B. Yeast two-hybrid (Y2H) assays showing the interaction between OsNF-YB7 and OsGLK1. AD and BD indicate the activation domain and binding domain of GAL4, respectively. The full length or truncated OsNF-YB7 and OsGLK1 were fused with BD and AD, respectively. The indicated combinations of constructs were cotransformed into yeast cells and grown on the nonselective medium SD/-L-W and selective medium SD/-L-W-H.

C. A split complementary luciferase (LUC) assay confirmed the interaction between OsNF-YB7 and OsGLK1. Coexpression of the fusion of OsGLK1 and the N-terminal half of LUC (nLUC-OsGLK1) and the fusion of the C-terminal half of LUC and OsNF-YB7 (cLUC-OsNF-YB7) in the epidermal cells of *N. benthamiana* leaves induced LUC activities, whereas the epidermal cells coexpressing nLUC-OsGLK1 and cLUC, nLUC and cLUC-OsNF-YB7, or nLUC and cLUC did not show LUC activities.

D. Bimolecular fluorescence complementation (BiFC) assays showed interactions between OsNF-YB7 and OsGLK1 in the nucleus. OsNF-YB7 was fused with the N-terminal of yellow fluorescent protein (nYFP-OsNF-YB7); OsGLK1 was fused with the C-terminal of yellow fluorescent protein (cYFP-OsGLK1). The indicated combinations of constructs were transiently coexpressed in the leaf epidermal cells of *N. benthamiana*. Scale bar = 10 μ m.

E. Co-immunoprecipitation (Co-IP) assays showing that OsNF-YB7 interacts with OsGLK1 *in vivo*. 35S::OsNF-YB7:GFP (OsNF-YB7-GFP) and 35S::OsGLK1:3 $\times$ Flag (OsGLK1-Flag) were coexpressed in rice protoplasts and were immunoprecipitated with an anti-GFP antibody, and the immunoblots were probed with anti-GFP and anti-Flag antibodies. 35S::GFP (GFP) was used as a negative control.

F. DLR assays showing that OsNF-YB7 represses the transactivation activity of OsGLK1 on *OsPORA* and *OsLHCB4*. Protoplasts isolated from etiolated seedlings were used for the analyses. EV, empty vector. Data are means $\pm$ SD ($n = 3$). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

G. EMSA assay indicated that OsNF-YB7 inhibits the DNA binding of OsGLK1 to the promoter of *OsPORA*. The black and white arrow heads indicate the OsGLK1- and OsNF-YB7-bound probes, respectively; "+" and "++" indicate that 2- and 4 μ M recombinant proteins were used for the reactions. The GST-His was used as a negative control.

Discussion

In crops such as soybean and canola, the presence of green embryos is considered as a valuable trait due to its association with increased photosynthetic capacity, which consequently promotes fatty acid biosynthesis (Ruuska et al., 2004 [↗](#)). This highlights the potential value of the green embryo. Nevertheless, the degradation of chlorophyll in mature seeds is necessary to prevent adverse effects on seed viability and meal quality (Chung et al., 2006 [↗](#)).

By surveying 1,094 species from 666 genera and 182 families, Yakovlev and Zhukova (1980) [↗](#) found that 428 angiosperms produce embryos with the presence of chlorophyll. Seeds with chlorophyllous embryo are scattered throughout the angiosperms and have evolved in many unrelated groups, but they are more prevalent in non-endospermous seeds (Dahlgren, 1980 [↗](#)). Currently, the evolutionary force driving the divergence of chloroembryophytes and leucoembryophytes remains largely unknown. Embryo photosynthesis contributes a large amount of oxygen to fuel energy-generating pathways in seed (Simkin et al., 2020 [↗](#)). The presence of chlorophyll in the embryo facilitates photosynthesis at early developmental stages, potentially leading to improved seedling growth and vigor (Smolikova and Medvedev, 2016 [↗](#)). In many chloroembryophytes, such as *Arabidopsis*, the embryo occupies a large proportion of the seed. From an evolutionary perspective, the presence of chlorophyll in the embryo may promote adaptation in such chloroembryophytes, as more reserves can be accumulated in the seed through active photosynthesis, better supporting the embryo development and subsequent seedling growth (Sela et al., 2020 [↗](#)). On the other hand, some leucoembryophytes, such as rice, have a persistent endosperm rich in storage reserves to nourish embryo development (Liu et al., 2022 [↗](#)). The acquisition of the ability to accumulate chlorophyll in the embryo is not necessary in such species. Although chlorophyllous embryos are rare among the ‘primitive’ angiosperm superorders, they are observed in some (but not all) Nymphaeales species that are surrounded by some amount of endosperm (Yakovlev and Zhukova, 1980 [↗](#)). As there is a paucity of knowledge regarding the evolution of chlorophyllous embryos, further comprehensive studies are necessary.

Some algae and gymnosperm species have evolved an ability to synthesize Chl in the dark (Myers, 1940 [↗](#); Ranade et al., 2019 [↗](#); Bogorad, 1950 [↗](#)). However, it has remained largely unclear whether access to light accounts for the induction of chloroembryos (Dahlgren, 1980 [↗](#); Periasamy and Vivekanandan, 1981 [↗](#); Liu et al., 2017 [↗](#)). Here, we found that removing the tissues covering an embryo failed to produce chlorophyllous embryos in the WT; however, light avoidance inhibited Chl accumulation in *osnf-yb7* embryos (Fig. S2A–H). These results suggest that light is necessary but insufficient to trigger chloroplast biogenesis in rice. Because light itself does not induce Chl accumulation in rice embryo (Fig. S2I–L), we inferred that there are intrinsic cues to repress chloroplast development in rice embryo. However, to the best of our knowledge, the underlying mechanism that determines an embryo’s ability to synthesize Chl is completely unknown. Here, we showed that OsNF-YB7, a LEC1 homolog of rice, acts as an inhibitor to repress Chl accumulation in the embryo. In line with this, a recent independent study also showed that the *OsNF-YB7* null mutant accumulates Chl in the embryo, although the cause of this remains unresolved (Guo et al., 2022 [↗](#)).

Several TFs involved in chloroplast development and photomorphogenesis in rice have been identified (Nakamura et al., 2009 [↗](#); Li et al., 2019 [↗](#); Bai et al., 2019 [↗](#); Hudson et al., 2013 [↗](#)). Interestingly, we found that many of them were upregulated in the *osnf-yb7* embryos. For example, *OsGLK1* was shown to be the most activated (Fig. 3A and B [↗](#)). Our genetic and biochemical evidence suggests that *OsGLK1* is involved in the OsNF-YB7-mediated repression of Chl biosynthesis in rice embryo. *OsGLK1* overexpression mimicked the chloroembryo phenotype in WT, while knockout of *OsGLK1* and *OsGLK2*, simultaneously, suppressed Chl accumulation in *osnf-yb7* embryo (Fig. 4A–G [↗](#)). The biochemical experiments suggested that OsNF-YB7 associates

with the promoter of *OsGLK1*, *in vivo*, to transcriptionally inactivate *OsGLK1* (Fig. 3D–F), indicating that *OsGLK1* is a downstream target of OsNF-YB7. However, OsNF-YB7 alone failed to bind to the promoter of *OsGLK1* *in vitro* (Fig. S5), presumably due to some as-yet-unidentified TFs being recruited by OsNF-YB7, assisting in recognizing *OsGLK1* for transcriptional regulation. An NF-Y TF complex usually consists of three subunits (NF-YA/B/C) in order to exert its function (Laloum et al., 2013). NF-YA is responsible for DNA binding, while NF-YB and NF-YC are primarily responsible for transactivation (Chaves-Sanjuan et al., 2021; Gnesutta et al., 2017). OsNF-YB7 possibly forms a heterotrimer with NF-YA and NF-YC in the embryo to recognize *OsGLK1*. In addition, previous studies showed that *Arabidopsis* LEC1 can interact with different TFs for developmental regulation (Huang et al., 2015b; Yamamoto et al., 2009; Fathihi et al., 2016; Boulard et al., 2018; Huang et al., 2015a). Identifying such TFs may reveal how OsNF-YB7 recognizes *OsGLK1* in order to exert its function.

OsNF-YB7 could recognize a common set of genes involved in Chl biosynthesis and photosynthesis that are recognized by *OsGLK1* (Fig. 2E–H and Table S3). The biochemical results suggested that OsNF-YB7 can directly repress *OsPORA* and *OsLHCB4*, which are activated by *OsGLK1* (Fig. 2J, and Fig. S10). OsNF-YB7 probably hinders *OsGLK1* to access the target genes by forming an *OsGLK1*/OsNF-YB7 heterodimer (Fig. 6A–E), or by occupying the motif *OsGLK1* recognized in the promoter, given the binding sites of *OsGLK1* and OsNF-YB7 are likely overlapped (Fig. 5F, Fig. S12 and Table S3). The first of these hypotheses is more plausible because, upon co-incubation of *OsGLK1* and OsNF-YB7 with the *OsPORA* promoter *in vitro*, the shifted band signal of *OsGLK1* was substantially decreased (Fig. 6G), indicating that the protein-protein interactions overwhelm the protein-DNA interactions. Thus, when OsNF-YB7 is expressed, it interacts with *OsGLK1*, then *OsGLK1* is less available to activate downstream targets. The findings suggested that OsNF-YB7 plays a dual role in regulating Chl biosynthesis in rice embryo: first, it represses the downstream genes, achieved via its function as a transcriptional inactivator; second, OsNF-YB7 can interact with *OsGLKs* to disturb their abilities to transactivate genes related to Chl biosynthesis and photosynthesis (Fig. S14). In addition to OsNF-YB7, a recent study showed that Deep Green Panicle 1, a plant-specific protein with a conserved TIGR01589 domain, can interact with *OsGLKs* to suppress *OsGLK*-mediated transcription (Zhang et al., 2021a). The findings suggested that, as the central regulator responsible for chloroplast development, *GLK* is tightly regulated at the post-translational level to fine-tune Chl biosynthesis in plants.

Previous studies suggested that *Arabidopsis* LEC1 is a positive regulator of Chl biosynthesis (Pelletier et al., 2017), given that the mature embryos of *lec1* were paler than the WT (Meinke, 1992; West et al., 1994). However, the null mutation of *OsNF-YB7* activated Chl biosynthesis, implying that the LEC1-type gene acts as a negative regulator in rice. By surveying the literature, we noted that Meinke reported that the cotyledons of the *lec1* mutant remained green unusually late in development (Meinke, 1992). Although there was no significant difference in Chl content upon using whole seeds for quantification, Parcy et al. did observe that the tip of *lec1* cotyledons accumulated more Chl (Parcy et al., 1997). Moreover, the *lec1;abi3* double mutant embryos produced much more Chl than the *abi3* single mutant (Parcy et al., 1997). Pelletier et al. recently reported that a cluster of LEC1 targets is enriched in genes related to photosynthesis and chloroplast development, and many of which are down-regulated in the embryo of *lec1* (Pelletier et al., 2017); however, we noticed that in their dataset, the photosynthesis and Chl biosynthesis related genes were more strikingly enriched in the up-regulated genes of *lec1*, at either the mature green or postmature green stage. These observations challenge the concept that LEC1 positively regulates Chl biosynthesis and photosynthesis in *Arabidopsis*. Most studies in *Arabidopsis* have emphasized the importance of LEC1 in embryo development at the maturation stage. However, LEC1 is activated within 24 h after fertilization (Lotan et al., 1998), but its role in the early embryo developmental stages for Chl biosynthesis is still unknown. Previously, we have expressed *OsNF-YB7* in the *lec1-1* background, driven by the native promoter of *Arabidopsis* LEC1 (Niu et al., 2021b). Given that *OsNF-YB7* could rescue embryo morphogenesis defects in *Arabidopsis* (Niu et al., 2021b), we postulated that *OsNF-YB7* in rice plays a similar role to that of LEC1 in

Arabidopsis. In order to ascertain whether *LEC1* can fully restore *osnf-yb7*, it is necessary to ectopically express *LEC1* driven by the native *OsNF-YB7* promoter in the *osnf-yb7* background in the future, due to the possibility of functional divergence between the genes with regard to the regulation of chlorophyll biosynthesis in the embryo. It would also be worthwhile to carry out similar tests in other grass organisms to better understand the regulatory mechanism in rice.

In addition to producing chlorophyllous embryo, the *osnf-yb7* mutants display an array of developmental defects, including abnormal embryogenesis, reduced dormancy, and desiccation intolerance, similar to those found in *Arabidopsis lec1* mutants (Niu et al., 2021a [↗](#)). Mutation of *osglk1* and *osglk2* in *osnf-yb7* could recover the chloroembryo phenotype, but did not alleviate the other embryo defects (Fig. 4D—G [↗](#)). The findings suggest that OsGLKs specifically function in Chl biosynthesis, but that OsNF-YB7 is responsible for many aspects of embryo development. In agreement with this, the number of OsNF-YB7-targeted genes is far greater than that of OsGLK-targeted genes, and the DEGs in *osnf-yb7* embryo is far greater than that in *OsGLK1-OX* embryo (Fig. 4H [↗](#) and Fig. 5E [↗](#)). The mechanisms underlying the OsNF-YB7-regulated multiple embryo developmental processes require further investigation.

Methods and Materials

Plant materials and growth conditions

The *osnf-yb7* mutant lines used in this study were previously generated in our laboratory (Niu et al., 2021b [↗](#)). Rice cultivars Zhonghua11 (ZH11) and Kitaake (Kit) were used for gene transformation. Rice plants were grown in a paddy field in Yangzhou, Jiangsu Province, China, or in a climate-controlled room under long-day conditions with a photocycle of 14 h of light (32°C) and 10 h of darkness (28°C), at 50% humidity. The *N. benthamiana* plants were grown in a growth chamber at 22°C with long-day conditions (16 h light/8 h dark). To determine whether the chloroembryo of *osnf-yb7* is light-dependent, the emerging panicles of the mutants were covered with aluminum foil until seed maturation. The lemmas of the WT were carefully removed with forceps at 1–2 DAF to expose the WT embryo to light in a climate-controlled room.

Vector construction and plant transformation

To generate the overexpression plants, full-length coding sequences (CDSs) of *OsNF-YB7* or *OsGLK1* (the stop codon removed) were cloned into pCambia1300-Flag or pUN1301-GFP under the control of *ubiquitin* promoter, using the ClonExpress® II One Step Cloning Kit (Vazyme). The constructs were transformed into rice calli through an *Agrobacterium*-mediated strategy, as described previously (Chen et al., 2016 [↗](#)). The higher-order mutants of *OsNF-YB7*, *OsGLK1* and *OsGLK2* mutants, were generated using a previously described method for multiple gene editing in ZH11 (Cheng et al., 2021 [↗](#)). The primers used for vector construction are listed in Table S4.

RNA extraction and RT-qPCR

Embryos or seedlings were collected by flash freezing in liquid nitrogen and stored at –80 °C until processing. Samples were finely ground using a mortar and pestle with liquid nitrogen. Total RNA was isolated using the RNA-easy Isolation Reagent (Vazyme, R701-01). The experiments were performed with at least three biological replicates. The relative expression levels of the tested genes were normalized using the rice *Actin* gene and calculated by the $2^{-\Delta C_t}$ method. The primers used for the RT-qPCR are listed in Table S4.

RNA-sequencing and data analysis

Ten-DAF-old embryos of the Wild-type (WT) and *OsGLK1-OX* were used for RNA-sequencing. Two biological replicates were set. RNA extraction, library preparation, and high-throughput sequencing of the collected samples were outsourced to BGI Genomics Co., Ltd., Shenzhen, China. CLC Genomics Workbench 12.0 software was used for RNA-seq data analysis, as previously reported (Xu et al., 2021 [DOI](#)). The thresholds of fold change > 2 and Bonferroni-corrected *FDR* < 0.05 were used for defining a DEG. The previously generated ChIP-seq data of OsNF-YB7 and OsGLK1 (Guo et al., 2022 [DOI](#); Tu et al., 2022 [DOI](#)) were reanalyzed using CLC Genomics Workbench 12.0 software for peak calling. Enriched motifs were identified by the online tool MEME-ChIP (<https://meme-suite.org/meme/tools/meme-chip> [DOI](#)) with default parameter set. The software Mapman was used for pathway analysis (Usadel et al., 2009 [DOI](#)). The online tool AgriGO 2.0 was used for GO analysis (Du et al., 2010 [DOI](#)). The Venn diagrams were drawn using an online tool (<https://bioinfogp.cnb.csic.es/tools/venny/index.html> [DOI](#)). TBtools was used for heat map generation (Chen et al., 2020 [DOI](#)).

Dual luciferase reporter (DLR) assays

The *OsGLK1*, *OsLHCB4* (2 kb upstream of translation start site), and *OsPORA* (1.5 kb upstream of translation start site) promoter sequences were amplified from ZH11 genomic DNA and cloned into the vector pGreenII 0800-LUC (Hellens et al., 2005 [DOI](#)), as reporters; the 35S::*OsGLK1* and 35S::*OsNF-YB7* constructs were used as effectors. The reporters and effectors were transfected into rice etiolated protoplasts in different combinations and incubated overnight. Firefly LUC and *Renilla* luciferase (REN) activities were measured using the DLR Assay Kit (Vazyme), following the manufacturer's instructions, and the LUC:REN ratios were calculated for analysis. The primers used for generating these constructs are listed in **Table S4**.

Chromatin immunoprecipitation (ChIP) assays

The CDS (stop codons removed) of *OsGLK1* was cloned into the pJIT163-GFP vector driven by a 35S promoter to generate the *OsGLK1*-GFP construct. Protoplasts transformed with *OsGLK1*-GFP and *OsNF-YB7*-Flag transgenic lines were used for the ChIP assays, in accordance with a previously described method (Zhao et al., 2020 [DOI](#)). Briefly, protoplasts or 0.2 g of 14-day-old seedlings were harvested and crosslinked with 1% formaldehyde for 15 min, followed by neutralization using 0.125 M glycine for an additional 5 min. The seedlings were then ground into powder in liquid nitrogen. The nuclei were isolated and lysed using Buffer S [50 mM HEPES-KOH (pH 7.5), 150 mM NaCl, 1 mM ethylenediaminetetraacetic acid (EDTA), 1% Triton X-100, 0.1% sodium deoxycholate, 1% SDS] and Buffer F [50 mM HEPES-KOH (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate]. The chromatin was then sonicated with the segment size ranging from 200 to 600 bp. The lysates were then immunoprecipitated by anti-GFP (abcam no. ab290) and anti-Flag (Sigma no. F3165) antibodies, respectively. Immunocomplexes were washed with low-salt ChIP buffer (50 mM HEPES-KOH, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS), high-salt ChIP buffer (low-salt ChIP buffer but replacing 150 mM NaCl with 350 mM NaCl), ChIP wash buffer (10 mM Tris-HCl pH 8.0, 250 mM LiCl, 0.5% NP-40, 1 mM EDTA, 0.1% sodium deoxycholate), and TE buffer (10 mM Tris-HCl, pH 8.0, and 1 mM EDTA). The protein-DNA complexes were eluted from beads using ChIP Elution buffer (50 mM Tris-HCl pH 7.5, 10 mM EDTA, 1% SDS) for 15 min at 65 °C and the crosslinking was reversed by incubation overnight with proteinase K. The fragment DNA was extracted with phenol:chloroform:isoamyl alcohol (25:24:1), precipitated with ethanol, and resuspended in TE buffer. The immunoprecipitated DNA was used as a template for qPCR. The primers used here are listed in **Table S4**.

Yeast two-hybrid assays

The CDSs of *OsGLK1/2* and *OsNF-YB7* were cloned into pGADT7 and pGBKT7, respectively. The constructs were cotransformed into yeast strain AH109 using Frozen-EZ Yeast Transformation II kit (Zymo), in accordance with the manufacturer's instructions. The empty pGADT7 and pGBKT7 vectors were cotransformed in parallel as negative controls. The transformants were first selected on synthetic dropout medium (SD/-Leu-Trp) plates. We tested protein-protein interactions using selective SD/-Leu-Trp-His dropout medium. Interactions were observed after 3 days of incubation at 28 °C. The primers used for generating these constructs are listed in **Table S4**.

Split complementary LUC assays

Split complementary LUC assays were performed as previously described (Niu et al., 2020 [\[1\]](#)). The CDSs of *OsGLK1/2* and *OsNF-YB7* were cloned into JW771 and JW772 vectors to generate nLUC-*OsGLK1/2* and cLUC-*OsNF-YB7*, respectively. The constructs were introduced into *Agrobacterium tumefaciens* strain GV3101 and then co-infiltrated into *N. benthamiana* leaves, and the LUC activities were analyzed after 48 h of infiltration using Tanon Imaging System (5200 Multi; Tanon). The primers used for vector construction are shown in **Table S4**.

Bimolecular fluorescence complementation assays

The CDSs of *OsNF-YB7* and *OsGLK1* were cloned into the vector pSPYNE (nYFP) and pSPYCE (cYFP), respectively. The prepared plasmids were transformed into *Agrobacterium* strain GV3101, and the indicated transformant pairs were infiltrated into *N. benthamiana* leaves. Forty-eight hours after infiltration, the fluorescence signal of yellow fluorescent protein (YFP) was observed with a confocal laser scanning microscope (CLSM) (Carl Zeiss, LSM 710). Images were captured at 514 nm laser excitation and 519–620 nm emission for YFP. The primers used for vector construction are shown in **Table S4**.

Co-immunoprecipitation (Co-IP) assays

Co-IP assays were performed using rice protoplast as described previously (Zhang et al., 2011 [\[2\]](#)). The CDSs (stop codons removed) of *OsGLK1* and *OsNF-YB7* were cloned into the vectors pUC19-35S-FLAG-RBS and pJIT163-GFP driven by a 35S promoter, respectively. Ten micrograms of plasmid DNA (*OsGLK1*-GFP, GFP, and *OsNF-YB7*-Flag) was transformed or cotransformed into 200 µl of protoplasts and incubated in WI buffer (0.5 M mannitol, 20 mM KCl, and 4 mM MES at pH 5.7) for 12 h. The protoplasts were collected and lysed in 500 µl of lysis buffer [50 mM Tris-HCl, 150 mM NaCl, 5 mM EDTA (pH 8.0), 1% NP-40, 0.1 mM PMSF]. The extracts were incubated with GFP-Trap agarose beads at 4 °C for 3 h and washed three times with washing buffer. Samples were boiled in SDS protein loading buffer. Immunoblots were detected using corresponding primary antibodies (anti-GFP, ABclonal no. AE012; anti-Flag, Sigma no. F3165). The primers used for vector construction are shown in **Table S4**.

Chl measurement and confocal imaging

Approximately one hundred micrograms of embryos of the indicated genotypes were extracted in 1 ml of 100% dimethyl sulfoxide (DMSO) with incubation at 65 °C for 1 h. Then, the absorbance values at wavelengths of 648.2 and 664.9 nm were measured by spectrophotometry and total Chl content was calculated as reported previously (Barnes et al., 1992 [\[3\]](#)). Chl autofluorescence signal was detected by CLSM (Carl Zeiss, LSM 710), with excitation and emission wavelengths of 633 and 625–730 nm, respectively.

Electrophoretic mobility-shift assays (EMSAs)

OsNF-YB7 CDS was amplified by PCR and cloned into pET-28a vector to generate the *OsNF-YB7-His* construct. The full-length CDS of *OsGLK1* was cloned into pMAL-c5X vector to generate *OsGLK1-MBP* construct. All constructs were transformed into *E. coli* strain BL21 to produce recombinant proteins. The promoter subfragments of *OsPORA* (42 bp, from -278 to -237) and *OsLHCB4* (38 bp, from -259 to -222) were PCR amplified and labeled with biotin at the 3' hydroxyl end of the double strands using EMSA Probe Biotin Labeling Kit (Beyotime, GS008). EMSA was performed using EMSA/Gel-Shift kit (Beyotime, GS009), in accordance with the manufacturer's instructions. The labeled probes were detected in accordance with the instructions provided with the EMSA/Gel-Shift kit. All oligonucleotides used to generate the biotin-labeled probes are listed in **Table S4**.

Transmission electron microscopy (TEM)

TEM analysis was performed as described previously (Cheng et al., 2021 [DOI](#)). Briefly, embryos of WT and *osnf-yb7* were fixed overnight at 4°C in 2.5% glutaraldehyde and 0.1 M PBS. The samples were subsequently washed three times with 0.1 M PBS and then fixed with 1% osmic acid for 4 h. The samples were dehydrated in a series of ethanol and embedded in acrylic resin at 37 °C for 12 h and at 60 °C for 48 h. The samples were sectioned at 100 nm and observed by TEM (TECNAI 12).

Accession Numbers

Gene sequence data of this article can be found in the Rice Annotation Project Database (RAP-DB) under the following accession numbers: *OsNF-YB7* (Os02g0725700), *OsGLK1* (Os06g0348800), *OsGLK2* (Os01g0239000), *OsLHCA1* (Os06g0320500), *OsLHCA2* (Os07g0577600), *OsLHCA3* (Os02g0197600), *OsLHCA4* (Os08g0435900), *OsLHCA5* (Os02g0764500), *OsLHCA6* (Os09g0439500), *OsLHCB1.1* (Os01g0720500), *OsLHCB2* (Os03g0592500), *OsLHCB3* (Os07g0562700), *OsLHCB4* (Os07g0558400), *OsLHCB5* (Os11g0242800), *OsLHCB6* (Os04g0457000), *OsGUN4* (Os11g0267000), *OsCHLH* (Os03g0323200), *OsCHLI* (Os03g0563300), *OsCHLD* (Os03g0811100), *OsCRD1* (Os01g0363900), *OsDVR* (Os03g0351200), *OsPORA* (Os04g0678700), *OsGNC* (Os06g0571800), *OsPIL14* (Os07g0143200).

Acknowledgements

We thank Prof. Hengxiu Yu and Dr. Chao Zhang of Yangzhou University for kindly providing the positive and negative control vectors for dual luciferase reporter assays.

Funding

This research was supported by grants from the National Natural Science Foundation of China (32170344, 32300689), the Project of Zhongshan Biological Breeding Laboratory (BM2022008-02), the Jiangsu Province Government (JBGS001), the Natural Science Foundation of the Higher Education Institutions of Jiangsu Province (21KJB21003), the China Postdoctoral Science Foundation (2022M712702), the Independent Scientific Research Project Funds of the Jiangsu Key Laboratory of Crop Genomics and Molecular Breeding (PLR202101), the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

Author contributions

C.C. conceived the project. C.C., Z.Y. and T.B. designed the research. T.B., Z.Y. and B.N performed the experiments. Z.Y., T.B., Z.E. and C.C. analyzed the data. Z.Y. and C.C. wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

References

- Bai B., Lu N., Li Y., Guo S., Yin H., He Y., Sun W., Li W., Xie X (2019) **OsBBX14 promotes photomorphogenesis in rice by activating OsHY5L1 expression under blue light conditions** *Plant Sci* **284**:192–202
- Barnes J.D., Balaguer L., Manrique E., Elvira S., Davison A.W (1992) **A reappraisal of the use of DMSO for the extraction and determination of chlorophylls a and b in lichens and higher plants** *Environ. Exp. Bot* **32**:85–100
- Bogorad L (1950) **Factors Associated with the Synthesis of Chlorophyll in the Dark in Seedlings of Pinus jeffreyi** *Bot. Gaz* **111**:221–241
- Boulard C., Thévenin J., Tranquet O., Laporte V., Lepiniec L., Dubreucq B (2018) **LEC1 (NF-YB9) directly interacts with LEC2 to control gene expression in seed** *Biochim. Biophys. Acta - Gene Regul. Mech* **1861**:443–450
- Chaves-Sanjuan A., Gnesutta N., Gobbini A., Martignago D., Bernardini A., Fornara F., Mantovani R., Nardini M (2021) **Structural determinants for NF-Y subunit organization and NF-Y/DNA association in plants** *Plant J* **105**:49–61
- Chen C., Begcy K., Liu K., Folsom J.J., Wang Z., Zhang C., Walia H (2016) **Heat stress yields a unique MADS box transcription factor in determining seed size and thermal sensitivity** *Plant Physiol* **171**:606–622
- Chen C., Chen H., Zhang Y., Thomas H.R., Frank M.H., He Y., Xia R (2020) **TBtools: An Integrative Toolkit Developed for Interactive Analyses of Big Biological Data** *Mol. Plant* **13**:1194–1202
- Cheng X., Pan M., E, Z, Zhou Y., Niu B., Chen C. (2021) **The maternally expressed polycomb group gene OsEMF2a is essential for endosperm cellularization and imprinting in rice** *Plant Commun* **2**
- Chung D.W., Pružinská A., Hörtensteiner S., Ort D.R (2006) **The role of pheophorbide a oxygenase expression and activity in the canola green seed problem** *Plant Physiol* **142**:88–97
- Dahlgren R (1980) **The taxonomic significance of chlorophyllous embryos in angiosperm seeds** *Bot. Not* **133**:337–341
- Delmas F., Sankaranarayanan S., Deb S., Widdup E., Bournonville C., Bollier N., Northey J.G.B., McCourt P., Samuel M.A (2013) **ABI3 controls embryo degreening through Mendel's i locus** *Proc. Natl. Acad. Sci. U. S. A* **110**:1–7
- Du Z., Zhou X., Ling Y., Zhang Z., Su Z (2010) **agriGO: a GO analysis toolkit for the agricultural community** *Nucleic Acids Res* **38**:W64–W70
- E, Z, Li T., Zhang H., Liu Z., Deng H., Sharma S., Wei X., Wang L., Niu B., Chen C (2018) **A group of nuclear factor Y transcription factors are sub-functionalized during endosperm development in monocots** *J. Exp. Bot* **69**:2495–2510

- Fatihi A., Boulard C., Bouyer D., Baud S., Dubreucq B., Lepiniec L. (2016) **Deciphering and modifying LAFL transcriptional regulatory network in seed for improving yield and quality of storage compounds** *Plant Sci* **250**:198–204
- Fitter D.W., Martin D.J., Copley M.J., Scotland R.W., Langdale J.A. (2002) **GLK gene pairs regulate chloroplast development in diverse plant species** *Plant J* **31**:713–727
- Gnesutta N., Saad D., Chaves-Sanjuan A., Mantovani R., Nardini M (2017) **Crystal Structure of the Arabidopsis thaliana L1L/NF-YC3 Histone-fold Dimer Reveals Specificities of the LEC1 Family of NF-Y Subunits in Plants** *Mol. Plant* **10**:645–648
- Guo F., et al. (2022) **Rice LEAFY COTYLEDON1 Hinders Embryo Greening During the Seed Development** *Front. Plant Sci* **13**:1–18
- Hellens R.P., Allan A.C., Friel E.N., Bolitho K., Grafton K., Templeton M.D., Karunairetnam S., Gleave A.P., Laing W.A (2005) **Transient expression vectors for functional genomics, quantification of promoter activity and RNA silencing in plants** *Plant Methods* **1**
- Huang M., Hu Y., Liu X., Li Y., Hou X. (2015) **Arabidopsis LEAFY COTYLEDON1 controls cell fate determination during post-embryonic development** *Front. Plant Sci* **6**
- Huang M., Hu Y., Liu X., Li Y., Hou X (2015) **Arabidopsis LEAFY COTYLEDON1 mediates postembryonic development via interacting with PHYTOCHROME-INTERACTING FACTOR4** *Plant Cell* **27**:3099–3111
- Hudson D., Guevara D.R., Hand A.J., Xu Z., Hao L., Chen X., Zhu T., Bi Y.M., Rothstein S.J (2013) **Rice cytokinin GATA transcription factor1 regulates chloroplast development and plant architecture** *Plant Physiol* **162**:132–144
- Ito Y., Thirumurugan T., Serizawa A., Hiratsu K., Ohme-Takagi M., Kurata N (2011) **Aberrant vegetative and reproductive development by overexpression and lethality by silencing of OsHAP3E in rice** *Plant Sci* **181**:105–110
- Jarvis P., López-Juez E (2013) **Biogenesis and homeostasis of chloroplasts and other plastids** *Nat. Rev. Mol. Cell Biol* **14**:787–802
- Jo L., Pelletier J.M., Hsu S., Baden R., Goldberg R.B., Harada J.J (2020) **Combinatorial interactions of the LEC1 transcription factor specify diverse developmental programs during soybean seed development** *Proc. Natl. Acad. Sci* **117**:1223–1232
- Kobayashi K. et al. (2012) **Regulation of root greening by light and auxin/cytokinin signaling in Arabidopsis** *Plant Cell* **24**:1081–1095
- Laloum T., De Mita S., Gamas P., Baudin M., Niebel A. (2013) **CCAAT-box binding transcription factors in plants: Y so many?** *Trends Plant Sci* **18**:157–166
- Li X., Wang P., Li J., Wei S., Yan Y., Yang J., Zhao M., Langdale J.A., Zhou W (2020) **Maize GOLDEN2-LIKE genes enhance biomass and grain yields in rice by improving photosynthesis and reducing photoinhibition** *Commun. Biol* **3**
- Li Z., Mo W., Jia L., Xu Y.C., Tang W., Yang W., Guo Y.L., Lin R. (2019) **Rice FLUORESCENT1 Is Involved in the Regulation of Chlorophyll** *Plant Cell Physiol* **60**:2307–2318

- Li Z., Wu S., Chen J., Wang X., Gao J., Ren G., Kuai B. (2017) **NYEs/SGRs-mediated chlorophyll degradation is critical for detoxification during seed maturation in Arabidopsis** *Plant J* **92**:650–661
- Liu H., Wang X., Ren K., Li K., Wei M., Wang W., Sheng X. (2017) **Light deprivation-induced inhibition of chloroplast biogenesis does not arrest embryo morphogenesis but strongly reduces the accumulation of storage reserves during embryo maturation in arabidopsis** *Front. Plant Sci* **8**:1–14
- Liu J., Wu M., Liu C (2022) **Cereal Endosperms: Development and Storage Product Accumulation** *Annu. Rev. Plant Biol* **73**:255–291
- Lotan T., Ohto M., Yee K.M., West M.A.L., Lo R., Kwong R.W., Yamagishi K., Fischer R.L., Goldberg R.B., Harada J.J (1998) **Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells** *Cell* **93**:1195–1205
- Luo W. *et al.* (2022) **Overexpression of maize GOLDEN2 in rice and maize calli improves regeneration by activating chloroplast development** *Sci. China Life Sci*
- Meinke D.W (1992) **A Homoeotic Mutant of Arabidopsis thaliana with Leafy Cotyledons** *Science* **258**:1647–50
- Meinke D.W., Franzmann L.H., Nickle T.C., Yeung E.C (1994) **Leafy cotyledon mutants of Arabidopsis** *Plant Cell* **6**:1049–1064
- Myers J (1940) **A Study Of The Pigments Produced In Darkness By Certain Green Algae** *Plant Physiol* **15**:575–588
- Nakamura H., Muramatsu M., Hakata M., Ueno O., Nagamura Y., Hirochika H., Takano M., Ichikawa H (2009) **Ectopic overexpression of the transcription factor osglk1 induces chloroplast development in non-green rice cells** *Plant Cell Physiol* **50**:1933–1949
- Niu B., Deng H., Li T., Sharma S., Yun Q., Li Q., E, Z, Chen C (2020) **OsZIP76 interacts with OsNF-YBs and regulates endosperm cellularization in rice (Oryza sativa)** *J. Integr. Plant Biol* **62**:1983–1996
- Niu B. *et al.* (2021) **Rice LEAFY COTYLEDON1 regulates embryonic envelope development and chlorophyll biogenesis in embryo** *Research Square* <https://doi.org/10.21203/rs.3.rs-888827/v1>
- Niu B., Zhang Z., Zhang J., Zhou Y., Chen C. (2021) **The rice LEC1-like transcription factor OsNF-YB9 interacts with SPK, an endosperm-specific sucrose synthase protein kinase, and functions in seed development** *Plant J* **106**:1233–1246
- Parcy F., Valon C., Kohara A., Miséra S., Giraudat J (1997) **The ABSCISIC ACID-INSENSITIVE3, FUSCA3, and LEAFY COTYLEDON1 loci act in concert to control multiple aspects of arabidopsis seed development** *Plant Cell* **9**:1265–1277
- Pelletier J.M. *et al.* (2017) **LEC1 sequentially regulates the transcription of genes involved in diverse developmental processes during seed development** *Proc. Natl. Acad. Sci* **114**:E6710–E6719
- Periasamy K., Vivekanandan M (1981) **Photosynthesis in the Chloroembryo of Cyamopsis tetragonaloba** *Taub Ann. Bot* **47**:793–797

- Puthur J.T., Shackira A.M., Saradhi P.P., Bartels D (2013) **Chloroembryos: A unique photosynthesis system** *J. Plant Physiol* **170**:1131–1138
- Ranade S.S., Delhomme N., García-Gil M.R (2019) **Global gene expression analysis in etiolated and de-etiolated seedlings in conifers** *PLoS One* **14**
- Rossini L., Cribb L., Martin D.J., Langdale J.A (2001) **The maize Golden2 gene defines a novel class of transcriptional regulators in plants** *Plant Cell* **13**:1231–1244
- Ruuska S.A., Schwender J., Ohlrogge J.B (2004) **The capacity of green oilseeds to utilize photosynthesis to drive biosynthetic processes** *Plant Physiol* **136**:2700–2709
- Sakuraba Y., Kim E.Y., Han S.H., Piao W., An G., Todaka D., Yamaguchi-Shinozaki K., Paek N.C (2017) **Rice Phytochrome-Interacting Factor-Like1 (OsPIL1) is involved in the promotion of chlorophyll biosynthesis through feed-forward regulatory loops** *J. Exp. Bot* **68**:4103–4114
- Sato Y., Morita R., Nishimura M., Yamaguchi H., Kusaba M (2007) **Mendel's green cotyledon gene encodes a positive regulator of the chlorophyll-degrading pathway** *Proc. Natl. Acad. Sci. U. S. A* **104**:14169–14174
- Sela A., Piskurewicz U., Megies C., Mene-Saffrane L., Finazzi G., Lopez-Molina L (2020) **Embryonic photosynthesis affects post-germination plant growth** *Plant Physiol* **182**
- Simkin A.J., Faralli M., Ramamoorthy S., Lawson T (2020) **Photosynthesis in non-foliar tissues: implications for yield** *Plant J* **101**:1001–1015
- Smolikova G.N., Medvedev S.S (2016) **Photosynthesis in the seeds of chloroembryophytes** *Russ. J. Plant Physiol* **63**:1–12
- Tamai H., Iwabuchi M., Meshi T (2002) **Arabidopsis GARP transcriptional activators interact with the Pro-rich activation domain shared by G-box-binding bZIP factors** *Plant Cell Physiol* **43**:99–107
- Tao Z., Shen L., Gu X., Wang Y., Yu H., He Y (2017) **Embryonic epigenetic reprogramming by a pioneer transcription factor in plants** *Nature* **551**:124–128
- Thomas H., Ougham H (2014) **The stay-green trait** *J. Exp. Bot* **65**:3889–3900
- Tu X., et al. (2022) **Limited conservation in cross-species comparison of GLK transcription factor binding suggested wide-spread cistrome divergence** *Nat. Commun* **13**
- Usadel B., Poree F., Nagel A., Lohse M., Czedik-Eysenberg A., Stitt M (2009) **A guide to using MapMan to visualize and compare Omics data in plants: A case study in the crop species, Maize** *Plant, Cell Environ* **32**:1211–1229
- Wang M., et al. (2018) **Parallel selection on a dormancy gene during domestication of crops from multiple families** *Nat. Genet* **50**:1435–1441
- Wang P. et al. (2013) **Evolution of GOLDEN2-LIKE gene function in C3 and C4 plants** *Planta* **237**:481–495
- Wang P., Khoshravesh R., Karki S., Tapia R., Balahadia C.P., Bandyopadhyay A., Quick W.P., Furbank R., Sage T.L., Langdale J.A (2017) **Re-creation of a Key Step in the Evolutionary Switch from C3 to C4 Leaf Anatomy** *Curr. Biol* **27**:3278–3287

- Waters M.T., Wang P., Korkaric M., Capper R.G., Saunders N.J., Langdale J.A (2009) **GLK transcription factors coordinate expression of the photosynthetic apparatus in Arabidopsis** *Plant Cell* **21**:1109–1128
- West M.A.L., Yee K.M., Danao J., Zimmerman J.L., Fischer R.L., Goldberg R.B., Harada J.J (1994) **LEAFY COTYLEDON1 is an essential regulator of late embryogenesis and cotyledon identity in Arabidopsis** *Plant Cell* **6**:1731–1745
- Wietrzynski W., Engel B.D (2021) **Chlorophyll biogenesis sees the light** *Nat. Plants* **7**:380–381
- Xu X., E, Z, Zhang D., Yun Q., Zhou Y., Niu B., Chen C (2021) **OsYUC11-mediated auxin biosynthesis is essential for endosperm development of rice** *Plant Physiol* **185**:934–950
- Yakovlev M.S., Zhukova G.Y. (1980) **Chlorophyll in embryos of angiosperm seeds, a review** *Bot. Not* **133**:323–336
- Yamamoto A., Kagaya Y., Toyoshima R., Kagaya M., Takeda S., Hattori T (2009) **Arabidopsis NF-YB subunits LEC1 and LEC1-LIKE activate transcription by interacting with seed-specific ABRE-binding factors** *Plant J* **58**:843–856
- Yeh S.Y., et al. (2022) **Maize Golden2-like transcription factors boost rice chloroplast development, photosynthesis, and grain yield** *Plant Physiol* **188**:442–459
- Zhang C., Zhang J., Tang Y., Liu K., Liu Y., Tang J., Zhang T., Yu H (2021) **DEEP GREEN PANICLE1 suppresses GOLDEN2-LIKE activity to reduce chlorophyll synthesis in rice glumes** *Plant Physiol* **185**:469–477
- Zhang D., Tan W., Yang F., Han Q., Deng X., Guo H., Liu B., Yin Y., Lin H (2021) **A BIN2-GLK1 Signaling Module Integrates Brassinosteroid and Light Signaling to Repress Chloroplast Development in the Dark** *Dev. Cell* **56**:310–324
- Zhang J.-J., Xue H.-W (2013) **OsLEC1/OsHAP3E participates in the determination of meristem identity in both vegetative and reproductive developments of rice** *J. Integr. Plant Biol* **55**:232–49
- Zhang Y. et al. (2011) **A highly efficient rice green tissue protoplast system for transient gene expression and studying light/chloroplast-related processes** *Plant Methods* **7**
- Zhao L., et al. (2020) **Integrative analysis of reference epigenomes in 20 rice varieties** *Nat. Commun* **11**
- Zheng S. et al. (2022) **A cytosolic pentatricopeptide repeat protein is essential for tapetal plastid development by regulating OsGLK1 transcript levels in rice** *New Phytol* **234**:1678–1695

Editors

Reviewing Editor

Paula Casati

Center of Photosynthetic And Biochemical Studies (CEFOBI), Santa Fe, Argentina

Senior Editor

Jürgen Kleine-Vehn

University of Freiburg, Freiburg, Germany

Reviewer #1 (Public Review):

Summary:

This manuscript investigates the regulation of chlorophyll biosynthesis in rice embryos, focusing on the role of OsNF-YB7. The rigorous experimental approach, combining genetic, biochemical, and molecular analyses, provides a robust foundation for these findings. The research achieves its objectives, offering new insights into chlorophyll biosynthesis regulation, with the results convincingly supporting the authors' conclusions.

Strengths:

The major strengths include the detailed experimental design and the findings regarding OsNF-YB7's inhibitory role.

Weaknesses:

However, the manuscript's discussion on the practical implications for agriculture and the evolutionary analysis of regulatory mechanisms could be expanded.

<https://doi.org/10.7554/eLife.96553.2.sa3>

Reviewer #2 (Public Review):

Summary:

The authors set out to establish the role of the rice LEC1 homolog OsNF-YB7 in embryo development, especially as it pertains to the development of photosynthetic capacity, with chlorophyll production as a primary focus.

Strengths:

The results are well-supported and each approach used complements each other. There are no major questions left unanswered and the central hypothesis is addressed in every figure.

Weaknesses:

There are a handful of sections which could use clarifying for readers, but overall this is a solidly composed manuscript.

The authors clearly achieved their aims; the results compellingly establish a disparity between how this system operates in rice and Arabidopsis. Conclusions are thoroughly supported by the provided data and interpretations. This work will force a reconsideration of the value of Arabidopsis as a model organism for embryo chlorophyll biosynthesis and possibly photosynthesis during embryo maturation more broadly, as rice is a major crop organism and it very clearly does not follow the Arabidopsis model. It will thus be useful to carry out similar tests in other organisms rather than relying on Arabidopsis and attempt to more fully establish the regulatory mechanism in rice.

<https://doi.org/10.7554/eLife.96553.2.sa2>

Reviewer #3 (Public Review):**Summary:**

In this study, the authors set out to understand the mechanisms behind chlorophyll biosynthesis in rice, focusing in particular on the role of OsNF-YB7, an ortholog of Arabidopsis LEC1, which is a positive regulator of chlorophyll (Chl) biosynthesis in Arabidopsis. They showed that OsNF-YB7 loss-of-function mutants in rice have chlorophyll-rich embryos, in contrast to Arabidopsis LEC1 loss-of-function mutants. This contrasting phenotype led the authors to carry out extensive molecular studies on OsNF-YB7, including in vitro and in vivo protein interaction studies, gene expression profiling and protein-DNA interaction assays. The evidence provided well supported the core arguments of the authors, emphasising that OsNF-YB7 is a negative regulator of Chl biosynthesis in rice embryos by mediating the expression of OsGLK1, a transcription factor that regulates downstream Chl biosynthesis genes. In addition, they showed that OsNF-YB7 interacts with OsGLK1 to negatively regulate the expression of OsGLK1, demonstrating the broad involvement of OsNF-YB7 in rice Chl biosynthetic pathways.

Strengths:

This study clearly demonstrated how OsNF-YB7 regulates its downstream pathways using several in vitro and in vivo approaches. For example, gene expression analysis of OsNF-YB7 loss-of-function and gain-of-function mutants revealed the expression of selected downstream chl biosynthetic genes. This was further validated by EMSA on the gel. The authors also confirmed this using luciferase assays in rice protoplasts. These approaches were used again to show how the interaction of OsNF-YB7 and OsGLK1 regulates downstream genes. The main idea of this study is very well supported by the results and data.

Weaknesses:

It would be interesting to see how two similar genes have come to play opposite roles in Arabidopsis and rice. Interspecies complementation might help to understand this point.

<https://doi.org/10.7554/eLife.96553.2.sa1>

Author response:

The following is the authors' response to the original reviews.

eLife assessment

This is an important study on the regulation of chlorophyll biosynthesis in rice embryos. It provides insights into the genetic and molecular interactions that underlie chlorophyll accumulation, highlighting the inhibition of OsGLK1 by OsNF-YB7 and the broader implications for understanding chloroplast development and seed maturation in angiosperms. The results presented, including mutation analysis, gene expression profiles, and protein interaction studies, provide convincing evidence for the function of OsNF-YB7 as a repressor in the chlorophyll biosynthesis pathway.

Thank you very much for your positive assessment of our manuscript. We have carefully revised the manuscript according to the reviewers' valuable suggestions and comments. For more details, please see the point-to-point response to the reviewers below.

Public Reviews:

Reviewer #1 (Public Review):

Summary:

This manuscript investigates the regulation of chlorophyll biosynthesis in rice embryos, focusing on the role of OsNF-YB7. The rigorous experimental approach, combining genetic, biochemical, and molecular analyses, provides a robust foundation for these findings. The research achieves its objectives, offering new insights into chlorophyll biosynthesis regulation, with the results convincingly supporting the authors' conclusions.

Strengths:

The major strengths include the detailed experimental design and the findings regarding OsNF-YB7's inhibitory role.

Weaknesses:

However, the manuscript's discussion on the practical implications for agriculture and the evolutionary analysis of regulatory mechanisms could be expanded.

Thank you for your insightful comments and suggestions. In the revised manuscript, we discussed the potential application of the chlorophyllous embryo (please see line 270-274). The presence of chlorophyll in the embryo facilitates photosynthesis at early developmental stages, potentially leading to improved seedling growth and vigor (Smolikova and Medvedev, 2016). In crops such as soybean and canola, green embryo is considered as a valuable trait due to its association with enhanced photosynthetic capacity, which consequently promotes fatty acid biosynthesis (Ruuska et al., 2004). However, chlorophyll degradation must be carefully managed during seed maturation to avoid negative effects on seed viability and meal quality (Chung et al., 2006). Interestingly, the green embryo of lotus (*Nelumbo nucifera*) is widely used as a food ingredient in Asian, Australia, and North America. It is employed in herbal medicine to treat nervous disorders, insomnia, and other conditions (Zhu et al., 2017; Ha et al., 2022), highlighting the significant potential value of the green embryo.

In many chloroembryophytes, such as Arabidopsis, the embryo occupies a large proportion of the seed. From an evolutionary perspective, the presence of chlorophyll in the embryo may promote adaptation in such chloroembryophytes because more reserves can be accumulated in the seed through active photosynthesis, better supporting the embryo development and subsequent seedling growth (Sela et al., 2020). On the other hand, some leucoembryophytes, such as rice, have persistent endosperm rich in storage reserves to nourish embryo development (Liu et al., 2022). Gaining the ability to accumulate chlorophyll in the embryo is unnecessary for such species. In agreement with this hypothesis, chlorophyllous embryos are more prevalent in non-endospermous seeds (Dahlgren, 1980). However, we would like to emphasize that the evolutionary force driving the divergence of chloroembryophytes and leucoembryophytes is currently almost completely unknown and deserves in-depth investigation in the future. We discussed the possible evolution of the ability to accumulate chlorophyll in the embryo, please find the details in Line 276-295.

Reviewer #2 (Public Review):

Summary:

The authors set out to establish the role of the rice LEC1 homolog OsNF-YB7 in embryo development, especially as it pertains to the development of photosynthetic capacity,

with chlorophyll production as a primary focus.

Strengths:

The results are well-supported and each approach used complements each other. There are no major questions left unanswered and the central hypothesis is addressed in every figure.

Weaknesses:

There are a handful of sections that could use clarifying for readers, but overall this is a solidly composed manuscript.

The authors clearly achieved their aims; the results compellingly establish a disparity between how this system operates in rice and Arabidopsis. Conclusions are thoroughly supported by the provided data and interpretations. This work will force a reconsideration of the value of Arabidopsis as a model organism for embryo chlorophyll biosynthesis and possibly photosynthesis during embryo maturation more broadly, as rice is a major crop organism and it very clearly does not follow the Arabidopsis model. It will thus be useful to carry out similar tests in other organisms rather than relying on Arabidopsis and attempting to more fully establish the regulatory mechanism in rice.

Thank you very much for your positive comments. We have carefully revised the manuscript according to your and the other reviewers' comments and suggestions. Particularly, we emphasized the necessary to carry out similar tests in other organisms rather than relying on Arabidopsis to better understand the regulatory mechanism in rice.

Reviewer #3 (Public Review):

Summary:

In this study, the authors set out to understand the mechanisms behind chlorophyll biosynthesis in rice, focusing in particular on the role of OsNF-YB7, an ortholog of Arabidopsis LEC1, which is a positive regulator of chlorophyll (Chl) biosynthesis in Arabidopsis. They showed that OsNF-YB7 loss-of-function mutants in rice have chlorophyll-rich embryos, in contrast to Arabidopsis LEC1 loss-of-function mutants. This contrasting phenotype led the authors to carry out extensive molecular studies on OsNF-YB7, including in vitro and in vivo protein interaction studies, gene expression profiling, and protein-DNA interaction assays. The evidence provided well supported the core arguments of the authors, emphasising that OsNF-YB7 is a negative regulator of Chl biosynthesis in rice embryos by mediating the expression of OsGLK1, a transcription factor that regulates downstream Chl biosynthesis genes. In addition, they showed that OsNF-YB7 interacts with OsGLK1 to negatively regulate the expression of OsGLK1, demonstrating the broad involvement of OsNF-YB7 in rice Chl biosynthetic pathways.

Strengths:

This study clearly demonstrated how OsNF-YB7 regulates its downstream pathways using several in vitro and in vivo approaches. For example, gene expression analysis of OsNF-YB7 loss-of-function and gain-of-function mutants revealed the expression of selected downstream chl biosynthetic genes. This was further validated by EMSA on the gel. The authors also confirmed this using luciferase assays in rice protoplasts. These approaches were used again to show how the interaction of OsNF-YB7 and OsGLK1 regulates downstream genes. The main idea of this study is very well supported by the results and data.

Weaknesses:

From an evolutionary perspective, it is interesting to see how two similar genes have come to play opposite roles in Arabidopsis and rice. It would have been more interesting if the authors had carried out a cross-species analysis of AtLEC1 and OsNF-YB7. For example, overexpressing AtLEC1 in an osnf-yb7 mutant to see if the phenotype is restored or enhanced. Such an approach would help us understand how two similar proteins can play opposite roles in the same mechanism within their respective plant species.

We appreciate your insightful comments and suggestions. It is a very interesting question whether *AtLEC1* can fully restore *osnf-yb7*, given the possible functional divergence between the genes in terms of regulation of chlorophyll biosynthesis in the embryo. We have previously expressed *OsNF-YB7* in the *lec1-1* background in Arabidopsis, driven by the native promoter of *LEC1* (Niu et al., 2021). We found that *OsNF-YB7* could almost completely rescue the embryo defects in Arabidopsis, indicating that *OsNF-YB7* plays a resemble role in rice as the *LEC1* does in Arabidopsis (Niu et al., 2021). We sought to determine whether *AtLEC1* can complement the chlorophyll defect in *osnf-yb7*. However, given the fact that *osnf-yb7* shows severe callus induction defect, which is not surprising, because many studies have shown that *LEC1* is indispensable for somatic embryo development in various plant species, we are struggling to obtain the genetic materials for analysis. We have to transform *OsNF-YB7pro::AtLEC1* into the WT background first, and then cross the transformant with the *osnf-yb7* mutant. This is a time-consuming process in rice, but hopefully we will be able to isolate a line expressing *OsNF-YB7pro::AtLEC1* in the *osnf-yb7* background from the resulting segregating population.

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

A minor comment regarding the chlorophyll contents quantification in the study. Line 87: "The results showed that WT had an achlorophyllous embryo throughout embryonic development,...." In the TEM result, chloroplast was not observed in the WT embryo sections, indicating a lack of chlorophyll-containing structures, contrary to what was found in the osnf-yb7 embryos where chloroplasts were observed.

The authors stated that the embryo morphologies and Chl autofluorescence data showed that WT had an achlorophyllous embryo throughout embryonic development. However, the quantification of Chl levels in Figure 1D and Figure 4C showed that WT does produce some chlorophylls, albeit at lower levels than osnf-yb7 or OSLK1-OX embryos (WT values in the two figures are slightly different). This discrepancy warrants clarification to ensure consistency and accuracy in the manuscript's findings.

We re-evaluated the Chl content in the embryos of WT and *OsGLK1-OX* mature seeds. The result confirmed our previous finding that WT embryos produce a small amount of chlorophyll (please see the updated Fig. 4C). Notably, we observed that the dark-grown etiolated plants still have measurable chlorophyll content as reported in many studies (for example, Wang et al., 2017; Yoo et al., 2019), suggesting that there is potential bias in measuring chlorophyll content using an absorbance-based approach. We assume this possibly explains the concern you have raised.

Reviewer #2 (Recommendations For The Authors):

Mild editing for grammar is needed throughout, e.g. line 73, "It is still a mysterious why plant species".

We have carefully edited the grammar.

As a minor point, the placement of figure panels, such as in Figure 1, is not always intuitive.

Thank you for your suggestion. This figure has been revised as suggested. Please see the updated Fig. 1.

What is the significance of the two GFP mutants in Figures 2C and 2D? Is one of those the mislabeled Flag mutant?

The lines showed in Fig. 2C and D were not mislabeled. They were two independent transgenic events, both of which showed that OsNF-YB7 inhibited the expression of *OsPORA* and *OsLHCB4* in rice. The transgenic lines overexpressing OsNF-YB7 tagging with the 3× Flag (*NF-YB7-Flag*) were also used for this experiment. In agreement, *OsPORA* and *OsLHCB4* were significantly downregulated in the three independent *NF-YB7-Flag* lines (Fig. S4C), confirming the results showed in Fig. 2C and D.

In Figures 2G and 2H, what is that enormous band at the bottom of the gel?

The bands at the bottom of the gel were free probes. We indicated this in the revised figure.

Not until the Materials and Methods section did I realize that any of this study was being done in tobacco; the Introduction implies it's rice vs. Arabidopsis and it might be a good idea to mention the organism of study somewhere before Figure 6.

We apologize for any confusion caused by our previous writing. While the majority of this study was performed with rice plants or protoplasts, the split complementary LUC assays and BiFC assays were performed with tobacco. We have specified these in the revised manuscript as suggested.

Reviewer #3 (Recommendations For The Authors):

It would be nice if the author could show what the phenotype is in AtLEC1 OX in osnf-yb7 and also OsNF-YB7 OX in atlec1 mutants.

Thank you for your suggestion. We have previously expressed *OsNF-YB7* in the *lec1-1* background of Arabidopsis, driven by the native promoter of Arabidopsis *LEC1* (Niu et al., 2021). Since *OsNF-YB7* could rescue the embryo morphogenesis defects in Arabidopsis (Niu et al., 2021), we assumed that *OsNF-YB7* plays a similar role in rice as the *LEC1* does in Arabidopsis. However, it remains unknown whether expression of *LEC1* in *osnf-yb7* may restore the chlorophyllous embryo phenotype in rice. As the generation of genetic material is time-consuming, and especially given the fact that *osnf-yb7* has a severe callus induction defect, we are struggling to obtain the complementary line for analysis. We have to transform *OsNF-YB7pro::AtLEC1* in a WT background first, and then cross the transformant with the *osnf-yb7* mutant. Hopefully, we will be able to isolate a line expressing *OsNF-YB7pro::AtLEC1* in *osnf-yb7* background, from the derived segregating population. We discussed the reviewer's concern in the revised manuscript, please see Line 369-376.

Line 46, I think it is vague to mention that 'Like most plant species'. Some species might have different copy numbers, for example, a single GLK in liverwort M. polymorpha.

The statement has been revised. Please see Line 46.

Figures 2F and 5B, why was only one promoter region used for OsLHCB4? It would be better to have more regions like OsPORA.

Thank you for your comments. Here, we have examined more promoter regions (P1, P2 and P3) in the revised manuscript as suggested, among which, the previously selected promoter region (P3) contains both the G-box and CCAATC motifs that can be potentially recognized by GLK1. Consistent to our previous report, the results showed that OsNF-YB7 (left) and OsGLK1 (right) were associated with the P3 region, but showed no significant differences in the other probes. Please see the results in Fig. 2F and Fig. 5B of the revised manuscript.

Legend of Figures 2G, H, OsPORA (I), and OsLHCB (J) should be (G) and (H) respectively.

Corrected.

References

- Chung, D.W., Pruzinska, A., Hortensteiner, S., and Ort, D.R. (2006). The role of pheophorbide a oxygenase expression and activity in the canola green seed problem. *Plant Physiol* 142, 88-97.
- Ha, T., Kim, M.S., Kang, B., Kim, K., Hong, S.S., Kang, T., Woo, J., Han, K., Oh, U., Choi, C.W., and Hong, G.S. (2022). Lotus Seed Green Embryo Extract and a Purified Glycosyloxyflavone Constituent, Narcissoside, Activate TRPV1 Channels in Dorsal Root Ganglion Sensory Neurons. *J Agric Food Chem* 70, 3969-3978.
- Liu, J., Wu, M.W., and Liu, C.M. (2022). Cereal Endosperms: Development and Storage Product Accumulation. *Annu Rev Plant Biol* 73, 255-291.
- Niu, B., Zhang, Z., Zhang, J., Zhou, Y., and Chen, C. (2021). The rice LEC1-like transcription factor OsNF-YB9 interacts with SPK, an endosperm-specific sucrose synthase protein kinase, and functions in seed development. *Plant J* 106, 1233-1246.
- Ruuska, S.A., Schwender, J., and Ohlrogge, J.B. (2004). The capacity of green oilseeds to utilize photosynthesis to drive biosynthetic processes. *Plant Physiol* 136, 2700-2709.
- Sela, A., Piskurewicz, U., Megies, C., Mene-Saffrane, L., Finazzi, G., and Lopez-Molina, L. (2020). Embryonic Photosynthesis Affects Post-Germination Plant Growth. *Plant Physiol* 182, 2166-2181.
- Smolikova, G.N., and Medvedev, S.S. (2016). Photosynthesis in the seeds of chloroembryophytes. *Russ J Plant Physiol* 63, 1-12.
- Wang, Z., Hong, X., Hu, K., Wang, Y., Wang, X., Du, S., Li, Y., Hu, D., Cheng, K., An, B., and Li, Y. (2017). Impaired Magnesium Protoporphyrin IX Methyltransferase (ChlM) Impedes Chlorophyll Synthesis and Plant Growth in Rice. *Front Plant Sci* 8, 1694.
- Yoo, C.Y., Pasoreck, E.K., Wang, H., Cao, J., Blaha, G.M., Weigel, D., and Chen, M. (2019). Phytochrome activates the plastid-encoded RNA polymerase for chloroplast biogenesis via nucleus-to-plastid signaling. *Nat Commun* 10, 2629.
- Zhu, M., Liu, T., Zhang, C., and Guo, M. (2017). Flavonoids of Lotus (*Nelumbo nucifera*) Seed Embryos and Their Antioxidant Potential. *J Food Sci* 82, 1834-1841.

<https://doi.org/10.7554/eLife.96553.2.sa0>