

Reviewed Preprint

v1 • October 2, 2025

Not revised

Reviewed Preprint

v2 • December 5, 2025

Revised by authors

For correspondence:

michor@jimmy.harvard.edu

Author notes: See [page 28](#)

Reviewing editor: Anna Panchenko,
Queen's University, Canada

© 2025, Graser et al. This article is
distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted
use and redistribution provided that
the original author and source are
credited.

Punctuated mutagenesis promotes multi-step evolutionary adaptation in human cancers

Christopher Graser^{1,2,3}, Wenbo Wu¹, Cole Christini^{1,4}, Mia Petljak^{5,6}, Franziska Michor^{1,2,3,7,8,9}✉

¹Department of Data Science, Dana-Farber Cancer Institute, Boston, United States • ²Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, United States • ³Department of Stem Cell and Regenerative Biology, Harvard University, Cambridge, United States • ⁴School of Computer Science, Carnegie Mellon University, Pittsburgh, United States • ⁵Department of Pathology, New York University School of Medicine, New York, United States • ⁶Laura and Isaac Perlmutter Cancer Center, New York University, New York, United States • ⁷Center for Cancer Evolution, Dana-Farber Cancer Institute, Boston, United States • ⁸The Eli and Edythe L. Broad Institute, Cambridge, United States • ⁹Ludwig Center at Harvard, Harvard Medical School, Boston, United States

eLife Assessment

This **valuable** study presents a theoretical model of how punctuated mutations influence multistep adaptation, supported by empirical evidence from some TCGA cancer cohorts. This **solid** model points to the case of possible punctuated evolution rather than gradual genomic change. There was some disagreement amongst the reviewers in terms of how closely the theoretical results apply to the phenomena examined empirically, and alternative explanations should be considered in the future.

<https://doi.org/10.7554/eLife.108058.2.sa3>

Abstract

The rate of acquisition of genomic changes in cancer has been the topic of much discussion, with several recent investigations finding evidence of punctuated evolution instead of gradual accumulation of such changes. Despite forays into the description and quantification of these punctuated events, the effects of such changes on subsequent cancer evolution remain incompletely understood. Here we investigate how non-gradual mutagenesis affects the ability of tumor cells to acquire and retain fitness-enhancing adaptations. We find that punctuated mutagenesis significantly facilitates adaptation in scenarios where adaptation requires crossing a fitness valley, i.e. when multiple mutations are required which individually are maladaptive but jointly confer a fitness advantage. By increasing the probability that multiple mutations occur in close succession, punctuation increases the chance that mutants in a fitness valley mutate further to reach a fitness peak before going extinct. Analyzing data from The Cancer Genome Atlas, we find that tumors with signatures of APOBEC mutagenesis, which has been shown to proceed in episodic bursts, exhibit patterns consistent with higher rates of crossing fitness valleys. Lastly, we characterize how the interplay between this enhanced ability to cross fitness valleys and adaptation-limiting effects of clonal interference affects overall adaptability in complex fitness landscapes.

Introduction

Tumors evolve via the acquisition of randomly arising genetic and/or epigenetic alterations and their selection¹. The rate at which tumor cells accumulate such genomic changes to produce potentially adaptive innovation is thus an important determinant of the capacity of tumor cells to adapt to diverse selection pressures^{2,3}, affecting their propensity to progress locally, to

metastasize, or to evolve resistance to therapeutic intervention. Several methods to quantify mutation rates have been developed^{4–6} and proxies for mutation rates such as tumor mutation burden often feature in prediction models of therapeutic outcomes^{7,8}.

Most existing methods deal with constant mutation rates^{6,9}, assuming gradual evolutionary change over time. However, accumulating evidence suggests that mutagenesis in cancer is fluctuating. Indeed, a recent study¹⁰ demonstrated in vitro that mutations associated with DNA-editing activity of APOBEC cytidine deaminases¹¹ occur in episodic bursts, with more than 100-fold differences in the rate of APOBEC-associated mutations across otherwise identical cell culture replicates (Fig 1A [↗](#)). This episodicity observed in vitro aligned with patterns in previously published in vivo data investigating APOBEC mutagenesis in lung cancer^{12,13} and data from intestinal crypts¹⁴ later confirmed explicitly that this episodic pattern also occurs in patients. Other recent studies using genomic data to reconstruct evolutionary histories of tumors have found evidence of punctuated evolution across several cancer types^{15–17}. These studies have demonstrated the existence of distinct phases of mutation bursts^{17,18} (Fig 1B [↗](#)), challenging the prevailing paradigm of gradual emergence of mutant lineages (Fig 1C [↗](#)). Various mutagenic processes such as chromothripsis^{19,20}, chromoplexy²¹ and other drivers of chromosomal instability²² have been implicated in causing such punctuated patterns²³.

How punctuated mutagenesis affects the evolutionary dynamics of tumor cell adaptation across complex fitness landscapes remains incompletely understood. Addressing this question with existing experimental data is inconclusive, as only short time horizons are observed (single expansions¹⁰; time to a most recent common ancestor^{17,18}), and as fitness differences between observed cell lineages are incompletely characterized. Mathematical modelling of human cancer genomics data, however, can elucidate the dynamics of adaptation during punctuated tumor evolution. Here, we set out to systematically investigate the evolutionary consequences of punctuated mutation acquisition during tumorigenesis using mathematical modeling and analysis of genomic data from The Cancer Genome Atlas (TCGA).

Results

Temporal clustering facilitates multi-step adaptation

We set out to investigate how *temporal clustering* of mutagenic events into distinct episodes affects the ability of a cell population to achieve multi-step evolutionary adaptations. We considered two scenarios: tumor evolution via a single advantageous step, such as an oncogenic adaptation²⁴ (“1-step adaptations”, Fig 1D [↗](#)), and accumulation of multiple mutations with synergistic fitness effects (“multi-step adaptations”, Fig 1E [↗](#)). Indeed, the two hit-hypothesis²⁵ for tumor suppressor genes (TSGs) suggests that inactivation of one copy of a TSG can be inconsequential or maladaptive, while bi-allelic inactivation confers a selective advantage²⁶. While some TSGs exhibit (context-dependent) haploinsufficiency^{27–29}, synergistic fitness effects for successive (epigenetic^{27,28,30}) alterations in the same gene remain the prominent feature of TSGs. Analogously, there are ample examples of successive alterations across different genes acting synergistically^{31–33}. High mutation rates are known to limit the rate of retained 1-step adaptations per emerging 1-step adaptation due to clonal interference^{34–36} (Fig 1F [↗](#)), which has been highlighted as a potential consequence of punctuated cancer evolution³⁷. The dynamics of multi-step adaptation have been studied extensively^{26,38–44}, elucidating two modes of evolution (Fig 1G [↗](#)): ‘sequential fixation’ refers to the scenario in which the second mutation only emerges after cells harboring the first mutant have taken over the population, while ‘stochastic tunneling’ refers to situations in which the second mutation arises sooner. In large populations, mutants with a selective disadvantage become vanishingly unlikely to reach fixation, so that stochastic tunneling becomes the dominant mode of evolution for multi-step adaptations.

To investigate how the rate of successful stochastic tunneling events – enabling multistep adaptation – is affected by punctuated mutagenesis, we simulated selection dynamics of a population of cells which proliferate according to a Wright-Fisher process^{45,46} and accumulate mutations according to a constant (Fig 1H [↗](#)) or temporally clustered (Fig 1I [↗](#)) rate. The Wright-




Figure 1. Population dynamics under gradual vs punctuated mutagenesis.

(A) Evidence for punctuated APOBEC mutagenesis (Petljak et al. 2019) from successive cell culture expansions, seeded with single progenitors from the preceding expansion. Sequencing of the expanded populations revealed large fluctuations in APOBEC-associated mutagenic signatures SBS2 and SBS13. (B) Evidence for punctuated copy-number evolution (Gao et al. 2017, Minussi et al. 2021). Patterns in branch lengths of reconstructed phylogenetic trees reveal fluctuations in the rates of copy number alterations. Dynamics prior to most recent common ancestor (MRCA) are unidentifiable. (C) Gradual vs. punctuated evolution. In gradual evolution novelty emerges and spreads at a constant rate over time. In punctuated evolution novelty emerges and spreads during distinct burst phases. (D) Fitness schematic of two 1-step adaptations which each independently confer a fitness advantage. (E) Fitness schematic for a 2-step adaptation, in which carrying one mutation confers a fitness disadvantage but carrying two mutations confers a fitness advantage. (F,G) Schematic of possible evolutionary dynamics for two 1-step adaptations (F) and of modes of valley-crossing with or without prior fixation of the first mutant (G). Sequential fixation occurs at low mutation rates where emerging mutant lineages are likely to have fixed or gone extinct before the next mutation occurs. At higher mutation rates, the second mutation can occur in a multi-clonal population. (H) Sketch of failing two-step adaptation under a uniform (time-invariant) mutation rate. (I) Sketch of a successful two-step adaptation via stochastic tunneling under a punctuated mutation process with distinct clusters of high mutation rates.

Fisher process models evolution as successive, non-overlapping generations of constant size. Each new generation is populated by drawing with replacement from the cells in the previous generation, with probabilities proportional to their fitness (Fig 2A). Cells can accumulate mutations that change their fitness, i.e. increase their probability to survive to the next generation. We considered fitness landscapes in which two-step adaptations are the only way by which cells can increase their fitness, with fixed fitness ratios across subsequent two-step adaptations (Fig 2B,C). We also confirmed that the findings observed for the Wright-Fisher process, described below, are qualitatively consistent for expanding populations that proliferate according to a branching process model (Methods, Fig S1).

Interestingly, for two-step adaptations for which the intermediate mutants that carry only one of the two required mutations have a strong selective disadvantage (Fig 2B), we found that the population undergoes substantially (3.55 times) more two-step adaptations when accumulating mutations in a temporally clustered rather than in a uniform way (Fig 2D). This effect arises because intermediate mutants in “fitness valleys” have a high chance of going extinct before acquiring the next mutation. During mutation bursts, acquiring the next mutation in time before the disadvantageous mutant goes extinct becomes more likely. Similarly, intermediate mutants are more likely to emerge in a mutation burst. This temporal clustering of the likelihood of two succeeding mutation events increases the rate of successful two-step adaptations.

This effect also emerges for two-step adaptations that do not constitute proper fitness valleys, i.e. for which the intermediate mutant is not maladaptive (Fig 2E; 1.15-fold increase). Under stochastic selection dynamics, even mutants with a slight selective advantage (Fig 2C) have a high chance of going extinct due to random drift. Lineages acquiring sets of synergistic mutations, thus, often do so without prior fixation of each intermediate mutant – via stochastic tunneling²⁶. In those regimes, temporal clustering of mutations therefore also increases the speed of adaptation, as confirmed with simulations (Fig S2A-D). Moreover, the proportion of such sets of synergistic mutations acquired via stochastic tunneling rather than sequential fixation also increases with temporal clustering (Fig S2E).

Our results generalize to multi-step adaptations with arbitrarily many steps (SI1). To demonstrate this result, we compared a uniform mutation process with mutation rate μ to temporally clustered mutation processes in which mutations emerge exclusively during burst phases that constitute a fraction $\frac{1}{k}$ of the time, but at a k -fold increased rate, $k\mu$ (Fig S2A). We showed analytically that in the limit of the average mutation rate μ going to zero, the rate f_k of successful $(n + 1)$ -step adaptations is k^n times higher in the temporally clustered mutation process than in the uniform mutation process. We validated this result with simulations of a Wright-Fisher process for two-step adaptations, $n = 1$, and a range of small values for μ (Fig S2F,G), and with numerical calculations of f_k in a branching process (see SI2-SI3 and Fig S3-S4 for complete characterizations of parameter dependencies). As the mutation rate increases, this fold-increase $\frac{f_k}{f_1}$ becomes smaller. However, since the absolute rates f_k and f_1 are proportional to μ^{n+1} (SI1), the absolute effect of temporal clustering on the rate of fixations of mutants with multi-step adaptations increases in μ (Fig S2F). Analogously, we found that $\frac{f_k}{f_1}$ increases with the selective disadvantage that cells in a fitness valley have, while $f_k - f_1$ decreases with this selective disadvantage (SI2). In contrast, for low μ , and particularly for parameter choices in biologically relevant ranges ($\mu \leq 10^{-7}$), the fold-change becomes insensitive to the magnitude of the selective disadvantage so that ever smaller fitness differences suffice to reach high fold-changes, $\frac{f_k}{f_1} \approx k^n$ (SI2). For instance, in a population of constant size that evolves according to a branching process with a birth-death ratio of 1, if intermediate mutations in a multi-step adaptation sequence reduce the birth-death ratio to 0.98, we have $\frac{f_{50}}{f_1} = 49.4 = 0.988k$ for 2-step adaptations, and $\frac{f_{50}}{f_1} = 2440.6 = 0.976k^2$ for 3-step adaptations. In light of the large fold-changes in mutation rates during a burst and before or after a burst^{10,15,17,18}, our findings suggest that the resulting effect on the dynamics of multi-step adaptation is substantial.




Figure 2. Simulation results: valley-crossing under uniform vs. temporally clustered mutation rates.

(A) Schematic of a Wright Fisher Process. (B)-(C) Fitness landscapes used for the simulations in Panel (D) and (E) respectively. Mutations move a cell from left to right through the landscape. Each 2-step adaptation corresponds to a fitness increase by a factor of 1.5. Having an odd number of mutations comes at a multiplicative fitness disadvantage of 0.5 in (B) and an advantage of 1.01 in panel (D). (D)-(E) Heatmaps of simulation results for a Wright-Fisher process with 50 cells. The mutation rate trajectories in each panel are chosen such that the total expected number of mutations under the uniform trajectory is identical to that under the temporally clustered trajectory. Each column of pixels represents a population after a specific number of generations. Homogeneity in pixel-color per column thus indicates that cells with a given number of adaptations have fixated in the population, whereas persistent diversity of pixel colors (see Fig S1) indicates coexistence of lineages with different numbers of adaptations.

Temporal clustering in an exploration-exploitation setting

The increased rate of valley crossing is driven by phases of high mutation rates, when the fold-increase in the chance of valley-crossing is larger than the fold-increase in the mutation rate. This effect improves adaptability while the fitness landscape offers scope for multi-step adaptation. However, in fitness landscapes which have global maxima, as the population adapts there is ever less scope for (further) multi-step adaptations, and excessive exploration of the landscape becomes costly.

To investigate the effect of temporal clustering in such an exploration-exploitation setting, we constructed a two-dimensional fitness landscape (Fig 3A) which is randomly re-drawn at regular intervals, mimicking exposure to novel physical environments or drugs during cancer evolution and treatment (Methods). Between subsequent re-drawings, the population may move to a global fitness maximum where any further mutation decreases fitness. We considered mutation processes with different average mutation rates μ and different values of the clustering parameter k (Fig 3B) and investigated the average fitness of the cell population over long time horizons. As before, the clustering parameter k denotes the fold-increase of the mutation rate in burst phases relative to the average mutation rate μ . However, rather than fixing the out-of-burst mutation rate at zero, the burst duration is now held constant (Fig 3B) to control for differences in waiting time until a burst occurs.

We found that, in this scenario, relative to the uniform mutation process ($k = 1$) increasing k initially increases the average fitness of the population for any fixed μ (Fig 3C). However, at high k , marginal increases of k reduce fitness, since high in-burst mutation rates in small populations increase the chance of leaving fitness peaks due to stochastic drift, and low mutation rates outside of bursts delay the acquisition of one-step adaptations. Similarly, for a fixed k , increasing μ when starting at low μ -values increases the average fitness. However, moving further in either of these directions in the parameter space – towards increasing μ or towards increasing k – eventually brings about a decrease in average fitness reflecting a tradeoff between exploration and exploitation (see SI4 and Fig 3D-F for detailed descriptions and plots of the dynamics). For any fixed k , average fitness peaks at intermediate μ , and the highest average fitness on the k - μ -plane is achieved by a $k > 1$. Maximizing average fitness, thus, involves temporal clustering even when there is a trade-off between exploration and exploitation.

Proxies for valley crossing and for temporal clustering found in patient data

We then sought evidence of the modeling prediction that temporal clustering facilitates multi-step adaptation in patient data, leveraging whole exome sequencing (WES) data from TCGA. We reasoned that among tumors in which similar numbers of mutations had emerged, those tumors with more temporally clustered mutation processes in their evolutionary history would be more likely to have undergone a larger number of multi-step adaptations. Generalizing this idea to enable comparison of tumors which may differ in the number of mutations they have acquired, we reasoned further that tumors with high levels of temporal clustering would have a high fraction of successful multi-step adaptations per attempted multi-step adaptation, i.e. per emergence of a first mutant in a multi-step adaptation sequence. We would, thus, expect analogous ratios of detectable quantities that scale with the numbers of successful or attempted multi-step adaptations to correlate with indicators of fluctuations in the mutation rate history of a tumor such as those found for APOBEC mutagenesis. In particular, we considered the frequency of observing two deactivating mutations in a TSG as an indicator for successful multi-step adaptation, and the total number of observed mutations in a tumor as a quantity that scales with attempted multi-step adaptations.

Before considering the TCGA data, to confirm *in silico* that such a correlation would indeed emerge under biologically realistic parameters, we performed large-scale simulations of tumor growth from a single cell to up to realistically detectable cell numbers of 10^6 – 10^7 cells⁴⁷ (Methods). We recorded the fraction of mutations that emerged during burst phases, constructed a TSG




Figure 3. Exploration and exploitation with temporally clustered mutation rates.

(A) Cells move through two-dimensional fitness landscapes. These fitness landscapes are randomly redrawn every 50714 divisions. (B) Mutation rate trajectories are parameterized with a mean mutation rate μ , and a clustering parameter k . (C) Average fitness in simulations of a population of 20 cells in a Wright Fisher Process. Simulations were run for 10^8 division events. (D)-(F) Average fitness trajectories for representative snippets of the simulations in (C).

deactivation score (Fig 4A,B) by counting the number of TSGs (modelled as mutations with synergistic fitness effects) with at least two mutations across the population and divided by the total number of mutations across the genome (Methods). Our simulation results confirmed that these two readouts are correlated (Pearson correlation 0.42, $p < 0.01$; Fig 4C), showing that in silico predictions regarding the likelihood of acquiring 2-step adaptations under mutation processes with more vs. less temporal clustering are well reflected in our score for TSG deactivation.

Next, to construct similar readouts from the TCGA data, we first performed SBS signature decomposition on each sample and computed the relative contribution of APOBEC-associated mutation signatures (SBS2 and SBS13) vs all other mutation signatures⁴⁸ (Methods). Given the evidence for episodic APOBEC mutagenesis^{10,14}, we used this relative contribution as a score for temporal clustering. As a proxy for multi-step adaptations, a list of known TSGs (the TSGene 2.0 database⁴⁹) was considered, and for each TCGA sample, we counted the number of TSGs in this list which harbored at least two single nucleotide substitutions classified as a missense or nonsense mutation (Methods) normalized by the total number of single nucleotide substitutions in that sample. The TCGA WES data did not enable us to determine whether such mutations indeed deactivated different alleles of a TSG, rather than appearing on the same allele. If the likelihood of appearing on the same allele varied with the contribution of the APOBEC signature, this association could confound our results. Since mutations that are interdependently generated on the same copy of a gene would likely appear in close spatial proximity^{11,50,51}, we thus verified in a robustness check that our results remain unchanged when filtering out mutations with close-by genomic coordinates (Methods).

We then ranked the different tumor types in TCGA by the average contribution of SBS2 and SBS13 relative to all detected mutation signatures, and for each tumor type, investigated the correlation between our proxies for multi-step adaptations and for temporal clustering. For the top four categories with the highest mean contribution of APOBEC-driven mutagenesis to all mutations, we observed a positive correlation between the two proxies. This correlation was significant at the 5% level (t-test on the Pearson correlation coefficient) for three out of four of these top four categories (Fig 4D, for results for categories with lower APOBEC signature contribution see Fig S5).

Consistent with the hypothesis that this correlation arises because TSG deactivation occurs more readily with APOBEC mutagenesis, we found that mutations which contribute to the TSG-deactivation score on average have a strongly increased likelihood of being caused by APOBEC (Methods, Fig 4E). This finding is consistent across cancer types and independent of whether probabilities are averaged over samples or over individual mutations, suggesting that this result is not driven by outlier samples with many TSG inactivations and many APOBEC-associated mutations (Methods, Fig 4F).

An alternative score for multi-step adaptations, in which the list of TSGs is replaced with a list of pairs of genes with synergistic fitness effects³¹ (Methods) again exhibits a positive correlation with the relative contribution of APOBEC-associated mutation signatures in all categories; this correlation is significant for two of these categories (Fig S5). Additionally, we verified that our results are robust to the choice of correlation metric: using Spearman instead of Pearson correlations yields significant positive correlations across all top four categories discussed above (Fig 4D, S7).

Taken together, our analyses of TCGA data show that tumors with a larger share of APOBEC-associated mutations are enriched for pairs of mutations with synergistic fitness effects. While we cannot exclude the influence of unobserved confounders, given the limitations of the available data, we found that the patterns from our analysis are consistent with the theoretical predictions of our model.




Figure 4. Proxies for valley crossing and temporal clustering in simulations and in TCGA data.

(A) Sketch of the simulation analysis workflow. **(B)** Distributions of the fraction of mutations acquired during mutation bursts and the TSG deactivation score in simulations. **(C)** Joint distribution of the quantities in **(B)** indicates strong correlation. **(D)** Schematic of analysis workflow for TCGA data (methods) and results for the four cancer types with highest APOBEC signature contribution. **(E)** Probabilities that single base substitutions (SBSs) in samples of the cancer types in **(D)** were caused by APOBEC associated signatures SBS2 and SBS13. On the left, these probabilities are averaged over all SBSs. On the right, only those SBSs are considered which are classified as “nonsense” or “missense”, and which appear in a TSG with at least 2 such deactivating SBSs. The average probabilities between both groups are significantly different (t-test, $p < 0.001$). **(F)** Average probabilities analogous to those in panel **(D)** were constructed for each individual sample and then averaged across samples. Samples without deactivated TSGs were excluded. The average probabilities between groups are significantly different for the four cancer types (*, ** and *** respectively indicate that the p-value in a t-test lies below 0.05, 0.01 and 0.001).

Stochastic Tunneling vs Clonal Interference during Punctuated Evolution

In tumors evolving at high mutation rates, clonal interference may limit the rate at which a cell population manages to acquire and retain 1-step adaptations (Fig 1F). Previous literature has quantified this limiting effect as a function of the mutation rate and the distribution of fitness values of emerging mutants^{52–54}. However, the effects of non-constant mutation rates in this setting have not been elucidated. We thus set out to investigate clonal interference in the setting of temporally clustered mutation rates (Fig 5A).

We first considered a scenario in which cells can only acquire 1-step adaptations and measured the rate at which novel 1-step adaptations reach fixation in the population across simulations with different clustering parameters k (Fig 5B). We found that fixation rates quickly decrease as k increases (Fig 5C). This pattern emerges because the extent to which clonal interference reduces fixation rates disproportionately increases with the mutation rate; for a fixed average mutation rate, temporal clustering thus increases the effects of clonal interference.

These findings stand in contrast to our results for 2-step adaptations whose rate of fixation increases when temporal clustering is introduced. However, if in-burst mutation rates are sufficiently large such that multiple clones in the population may acquire a 2-step adaptation independently before one of these clones has fixated, effects of clonal interference also play a role for 2-step adaptations. Performing simulations for 2-step adaptations, we found that fixation rates are non-monotone in k . While at low k increasing k leads to a steep increase in the fixation rate, this trend eventually levels off and becomes negative, with further increases in k leading to a decrease in the fixation rate (Fig 5C).

Having observed that temporal clustering increases effects of clonal interference but also facilitates stochastic tunneling, we next set out to investigate the relative contribution of these two effects on overall adaptation rates. We considered a setting in which cells can increase their fitness through both 1-step and 2-step adaptations (Fig 5D). We found that 1-step adaptations are acquired more quickly under the uniform mutation process ($k=1$), while 2-step adaptations are acquired more quickly under a temporally clustered mutation process ($k=5$; Fig 5E). The relative magnitude of both effects varies with time. As 1-step adaptations spread more readily in the population, differences in the speed at which they fixate manifest early in the dynamics. Over time, cells gradually exhaust the possibilities for 1-step adaptations and the difference between the adaptation rate in both processes becomes dominated by differences in the speed of acquiring two-step adaptations. In this latter phase we observe large differences between the two mutation processes in the expected time until any given proportion of the possible two-step adaptations have spread in the population. Analogous time differences for reaching fixed proportions of one-step adaptations in the initial phase of the dynamics are much smaller (Fig 5E).

In these analyses, parameters such as mutation rates, fitness effects and population sizes were chosen so that simulations remained computationally manageable. Decreasing mutation rates toward more biologically realistic ranges decreases the likelihood that multiple lineages with adaptations emerge simultaneously and therefore weaken the effects of clonal interference. In contrast, larger population sizes and smaller fitness differences increase the likelihood of co-existence between competing lineages as each emerging lineage requires more time to fixate in the population, which strengthens the effects of clonal interference.

We also investigated how this pattern depends on the ruggedness of the fitness landscape (Fig S9A). To this end, we varied the relative proportions of loci with and without fitness valleys and quantified adaptability differences. We found that the duration of the initial phase in which the uniform process yields higher adaptability varies with the ruggedness of the fitness landscape: this phase is roughly three times as long when the proportion of loci with fitness valleys is 5% compared to when it is 95% (Fig S9B–D). However, over this range of proportions, we consistently observed that this initial phase only accounts for a small fraction of the time it takes to acquire all adaptations.




Figure 5. Stochastic tunneling vs clonal interference under temporally clustered mutation rates:

(A) Schematic of the simulation approach. **(B)** Sketch of the mutation process. **(C)** Simulation results: measuring fixations of higher-fitness mutations per generation as a function of the clustering parameter and of the fitness landscape. In the 1-step adaptation setting fitness is defined as $1.5^{\text{\#mutations}}$. Fitness in the 2-step adaptation setting is defined as in Fig 2B. Results are averaged over simulation runs with 10^7 generations. **(D)** Sketch of fitness landscape for simulations shown in (E). Cells start with an unmutated genome of 200 loci. Mutations on each locus have independent multiplicative effects on cell fitness. In the first 100 loci a single mutation confers a multiplicative fitness change of 1.5. In the remaining 100, 2 mutations are required to reach this multiplicative fitness change of 1.5, with the first mutation conferring a multiplicative fitness change of 0.5. **(E)** Simulation results for adaptation with genome sketched in (D). Results are averaged over 100 simulation runs per value of k . Shaded regions indicate 10th and 90th percentiles. Smaller plots (left) are zoomed in versions of the first 1200 generations of the larger plots (right), with identical color-coding.

Discussion

Accumulating evidence suggests that mutagenesis in tumor cell populations proceeds in punctuated bursts rather than gradually; however, the effects of such punctuation on the ability of a tumor cell population to acquire and retain fitness-enhancing adaptations remains incompletely understood. Here we set out to investigate the evolutionary dynamics of these processes using mathematical modeling and genomics data analysis of human tumors.

We found that when a population acquires multi-step adaptations via stochastic tunneling, the rate of adaptation is substantially enhanced by punctuation. Stochastic tunneling is the predominant mode of evolution in large populations that traverse fitness valleys. However, stochastic tunneling also occurs when the necessary mutation steps to reach a substantial fitness advantage are not individually maladaptive, i.e. if steps do not constitute a fitness valley but confer a slight fitness advantage. Punctuated evolution therefore facilitates the accumulation of sets of mutations that jointly and synergistically confer a fitness advantage, while the fitness effect of carrying only a subset of these mutations can range from being strongly maladaptive to being moderately adaptive. For the limit of low average mutation rates we show analytically that the rate of stochastic tunneling under a temporally clustered process is k^n times larger than under the time-invariant process, where k is the fold-increase in the mutation rate relative to the average mutation rate during mutation bursts in the temporally clustered process, and n is the number of mutation steps that is required to exit the fitness valley. Moving towards higher average mutation rates, this fold-change in the stochastic tunneling rates decreases, but the absolute difference between the rates increases. Investigating these relative and absolute effects in a branching process as a function of the birth-death ratio elucidated a similar pattern: While the fold-change is highest if lineages in a fitness valley have a low birth-death ratio, we observed the largest difference between rates of stochastic tunneling at birth-death ratios close to one.

Applied to cancer evolution where cells' birth-death ratios are influenced by a variety of internal and external factors, these findings highlight that for any given multi-step adaptation sequence, the magnitude of the facilitating effects described here likely varies depending on whether the tumor is in a phase of constant or growing population size.

While temporal clustering facilitates multi-step adaptations, it also impedes 1-step adaptations due to clonal interference. We examined the interplay between these contrasting effects in a setting in which cells can acquire both types of adaptations. We showed that the relative importance of both effects varies over time. Since 1-step adaptations tend to be acquired more readily, clonal interference matters most in the early phase of the process. As the share of possible 2-step adaptations relative to 1-step adaptations increases, differences in the tunneling rate become the more relevant determinant for the speed of adaptation. A uniform mutation rate, thus, makes a population more efficient at finding local fitness maxima. However, temporal clustering allows the population to more quickly move between local maxima, thus speeding up the search for a global maximum.

While our theoretical results are easily verified in simulation settings where we can choose a fitness landscape, applying these results to real data remains challenging because of the inherent complexities in determining fitness effects in biological systems. Estimating fitness effects of individual mutations requires either intricate experimental approaches⁵⁵ or large patient cohorts to control for differences in the (epi-)genetic background against which these mutations emerge; such estimations become considerably harder when considering joint fitness effects of sets of multiple mutations.

To circumvent these challenges, we focused on TSGs as a representative set of gene pairs with synergistic effects and on one mutation process known to tend to fluctuate over time – APOBEC-driven mutagenesis. Moreover, we restricted attention to single-nucleotide substitutions as the principal mode of APOBEC-driven mutagenesis and the context in which mutation signatures are best characterized, and we did not consider other classes of mutations such as copy number changes or epigenomic modifications. Consequently, we observed only a subset of groups of mutations with synergistic fitness effects in the data and possibly only a small fraction of the

variability in punctuation between different samples. Nevertheless, we found that these scores significantly correlate, which aligns with our model predictions. However, given this narrow scope, the generalization of our model towards a full quantitative estimate of the relevance of punctuation to adaptation dynamics in cancer requires additional investigations.

The observed empirical patterns may in part reflect the influence of unmeasured confounders. Rates of multi-step adaptation are determined not only by temporal clustering, but also by factors such as average mutation rates, fitness landscapes, and population-size trajectories. Structural differences in these factors between tumors with high or low APOBEC signature contributions therefore represent potential sources of confounding (for a review of known APOBEC-related biological mechanisms and clinical associations, see⁶⁰). Additionally, we cannot measure rates of multi-step adaptation directly, and therefore need to consider a proxy; to this end, we chose a score that relates deactivating TSG mutations to the absolute mutation count. Structural differences affecting the relative rates of deactivating TSG mutations, such as APOBEC-related enrichment for nonsense mutations⁵⁹, compared to other mutations can therefore also influence our empirical findings.

Our results have implications for both mechanistic and statistical modeling of tumor evolution. A key quantity in mechanistic models is the rate at which different types of adaptations arise; our results show that this rate depends on the temporal dynamics of the mutation process. Accounting for this effect might be particularly important when modelling the emergence of treatment resistance in models used to optimize treatment schedules⁵⁶. There is evidence linking APOBEC mutagenesis to resistance in breast cancer⁶¹. Our results suggest that APOBEC, and other punctuated mutation processes, are likely to be relevant more generally to settings in which developing resistance requires multi-step adaptation. Analogously, in statistical models, incorporating measures of punctuated mutagenesis may improve the ability to predict treatment outcomes by accounting for how these temporal dynamics shape adaptability.

Methods

Simulations of a Wright-Fisher and a branching process in an unbounded fitness landscape

In the simulations presented in Fig 2⁴ and Fig S1⁴, mutations occur between selection steps and are independent of divisions. In the Wright-Fisher process simulations (Fig 2 C⁴) with a temporally clustered mutation rate, the out-of-burst mutation probability per cell and per update step (between two divisions) was set to 0.01 and was increased to 0.1 in bursts. For the corresponding uniform mutation rate trajectory, this probability was simply set equal to the total amount of mutations that occurred in the simulations under the temporally clustered mutation rate, divided by the total number of update steps.

Mutation rates in the branching model simulations (Fig S1⁴) were chosen analogously. However, to achieve temporally equidistant mutation bursts in the branching process simulations, we scaled the burst duration and the time between bursts by the population size.

Simulations Wright-Fisher Process in exploration/exploitation setting

To produce the results shown in Fig 3⁴, we performed agent-based stochastic simulations of a Wright-Fisher Process with 20 cells. Cells were characterized by their location on a discrete fitness landscape (a 30 by 30 two-dimensional lattice). The fitness values f associated with the positions on the lattice were drawn independently at random as $f = 0.5 + x^4$ where $x \sim U[0,1]$ is distributed uniformly between zero and one. The lower bound of 0.5 ensures that cells at all positions have a non-negligible probability of dividing.

Raising x to the fourth power creates a landscape with few peak-positions surrounded by many valley-positions with little variation in fitness.

In each selection step, one cell was randomly chosen to divide with probability proportional to its fitness, and replaced a cell which was drawn uniformly at random. Mutations occurred between consecutive selection steps, and the mutating cell was chosen uniformly at random amongst all cells in the population (independently of the preceding selection step). If a cell mutated, it would move to a location in the fitness landscape chosen uniformly at random from the set of (at most 8) locations in the Moore neighborhood of its current location.

The rate at which mutation events occurred was governed by the parameters μ and k . Mutation bursts lasted 10^2 division events and started every 10^3 division events. Fitness landscapes were redrawn every 50714 ($\approx 50000 + \frac{10^3}{7}$) division events, to periodically vary the relative timing of mutation bursts and re-drawings of the fitness landscape. In this manner, we avoid artifacts in our results caused by phase-alignment of bursts and re-drawings.

Larger-scale simulations

We simulated tumor evolution as a branching process, starting from a single unmutated cell, up to a randomly drawn target cell number between 10^6 and 10^7 cells. We assume a constant death rate, so that over the course of a simulation the probability that the next event is a death event rather than a division event remains fixed (at a value of 0.35). In case of a death event, one cell picked uniformly at random is removed from the population. In case of a division event, one cell is chosen to divide with probabilities proportional to its fitness.

We assume that the number of mutations per division follows a binomial distribution $\text{Bin}(10^5, \mu)$, and for each simulation we randomly draw a baseline mutation probability μ uniformly between 0 and $3 \cdot 10^{-5}$. In mutation bursts, this probability gets multiplied by a factor of 20 or 50, again chosen randomly for each simulation. The population enters a burst phase with probability $\frac{n}{10}$ per division, where n is the current population size, and exits a burst phase with probability $\frac{n}{3}$, so that in expectation bursts phases last 3 generations, and start every 10 generations. The first cell in a simulation is initialized to be in a burst with probability $\frac{3}{13}$ consistent with the expected time spent in bursts given those parameters.

The fitness effect of mutations is modeled as follows. The starting cell has a fitness of one. Each new mutation has an additive effect on the cell's fitness. We assume that mutations occur uniformly at random across the genome and that multi-step adaptations can occur in a fraction $\theta = 0.01$ of the genome, roughly aligning with the ratio of the number of TSGs vs the total number of genes in the human genome. We subdivide this part of the genome into 200 TSGs which are all hit by a mutation with equal probability. For each individual TSG, the first mutation reduces a cell's fitness by 0.05. The second mutation increases fitness by 0.15, and all further mutations are fitness neutral. For the remaining $(1 - \theta)$ fraction of the genome, we assume that mutations are fitness-neutral with a probability of 0.9, and that fitness effects are otherwise drawn from a Gaussian with mean equal to -0.005 and standard deviation equal to 0.005.

To produce the results in Fig 4A-B, we keep track of all mutations that arise in a population, and remove all mutations with less than 1% variant allele frequency from the output, as those would be unlikely to be detected in WES. Moreover, we keep track of the fraction of mutations that a sample acquired during a mutation burst. For each simulation, we then count the number of TSGs for which there are at least two mutations found in the population and divide this by the total number of mutations in the sample.

Analysis of TCGA WES data

Whole-exome sequencing data was acquired from The Cancer Genome Atlas. We performed mutation signature decomposition using the `cosmic_fit()` function in python from the package SigProfilerAssignment⁵⁷ version 0.1.8 with cosmic version 3.4. Moreover, we used the mutation-level signature probabilities generated via SigProfilerAssignment to compute mean probabilities of signatures SBS2 and SBS13 for individual SBSs (Fig 4D,E).

To compute our TSG deactivation score, we filtered the SNV data for missense and nonsense mutations in genes belonging to the TSGene 2.0 database⁴⁹. For each sample we calculated the number of TSGs with at least two mutations and divided by the total number of SNVs.

Analogously, to construct our synergistic mutations score, we filtered for missense and nonsense mutations in frequently co-mutated gene pairs identified by Gu et al.³¹, and for each sample divided the number of pairs in this list with mutations in both genes by the number of SNVs.²⁹ This enrichment for co-occurrences of mutations in both genes in a pair suggests that mutations in the two genes tend to have a synergistic fitness effect. For each sample in the TCGA WES data we thus computed the number of gene-pairs from this list for which there is at least one non-synonymous single nucleotide substitution mutation in each of the two genes, and divided this number by the total number of single nucleotide substitutions in the sample.

As a robustness-check, we investigated whether the results change if we require a minimum distance between the genomic locations of the mutations in TSGs that we count to our TSG deactivation score. APOBEC has been linked to clustered mutagenesis through mutation processes of kataegis and omikili.^{11,50,51} If samples with higher APOBEC activity have higher rates of having multiple close-by mutations on the same allele, and if such a sample has multiple deactivating mutations in a TSG, these inactivating mutations might be more likely to be localized on the same allele compared to samples with lower APOBEC activity. Since we use appearances of more than one inactivating mutation as a proxy for bi-allelic deactivation of TSGs, such a pattern would bias our results. To account for that, we explored filtering TSG mutations in the data based different minimum-distance-thresholds ranging from 1 to 100 bp and found no impact on our results. We did not perform any analyses with allele-specific mutation information, as determining allele-specific TSG inactivation relies on informative co-occurring mutations. For any given mutation, tumors with lower mutation burden are less likely to harbor such co-occurring mutations and preferentially filtering out observations in tumors with lower mutation burden would have biased our findings.

Data availability

The results shown here are in part based upon data generated by the TCGA Research Network: <https://www.cancer.gov/tcga>.

Supplementary Information

SI1 Derivations of results on multi-step adaptation rates in the limit of infrequent mutations

In the main text, we discussed differences in the rate of multi-step adaptations f_k as a function of a temporal clustering parameter k . Here we formalize this discussion and derive the result presented in the main text. We consider processes in which multi-step adaptations occur via stochastic tunneling rather than sequential fixation, i.e. processes in which mutants with a selective disadvantage are negligibly unlikely to reach fixation. We assume that mutations happen sufficiently infrequently, so that we can neglect valley crossing scenarios in which the same mutation in a multi-step adaptation sequence occurs multiple times. Additionally, we require that valley crossing happens solely via stochastic tunneling, i.e. that the probability that a maladaptive mutant fixates is vanishingly low.

We denote the average rate at which cells acquire mutations by μ , and compare dynamics under different mutation processes which we index by a temporal clustering parameter $k \geq 1$. Starting every d units of time, a process with parameter k undergoes a mutation burst lasting $\frac{d}{k}$ units of time, during which mutations occur at a rate $k\mu$. Outside of bursts, the mutation rate is 0. We assume that the time $\frac{(k-1)d}{k}$ between subsequent burst phases is sufficiently long such that we can restrict attention to multi-step adaptations that occur during a single burst, and that the duration

of each burst $\frac{d}{k}$ is sufficiently long relative to the time that it takes to cross a fitness valley, so that valley crossings, which fail because the burst phase ended but which would have been successful, have a negligible effect on f_k .

As stated in the main text, with this setup we can show that for any population dynamics process for which the above limits can be motivated, the fold-increase in the rate f_k at which $(n + 1)$ -step adaptations occur in a process with clustering parameter k relative to the uniform process approaches k^n .

$$\lim_{d \rightarrow \infty} \lim_{\mu \rightarrow 0} \frac{f_k}{f_1} = \lim_{\mu \rightarrow 0} \lim_{d \rightarrow \infty} \frac{f_k}{f_1} = k^n$$

Deriving this result for any selection process indeed becomes straightforward, if it can be motivated that the mutation rate does not affect the fate of any mutant lineage once the first mutant in this lineage has emerged.

Somewhat more formally, we index the steps in an n -step adaptation sequence by $I \in \{1, \dots, n\}$, and denote the expected size of the lineage descending from mutant i by $Y_i(t)$. This quantity $Y_i(t)$ of course depends on the specificities of the selection process, but for our derivations it suffices to only require that $Y_i(t)$ does not depend on the mutation rate.

Since all intermediate mutants have a selective disadvantage, and since we assume that we are in a parameter regime in which the likelihood that a disadvantageous mutant fixates is negligibly low, it follows that the integral $\int_0^\infty Y_i(t) dt$ converges.

$$\int_0^\infty Y_i(t) dt < \infty$$

In the limit of a low (and for now time-invariant) mutation rate μ , the probability that the i 'th mutant produces a further mutant $P(i \rightarrow i + 1)$ is simply the product of the mutation rate and this integral.

$$P(i \rightarrow i + 1) = \mu \int_0^\infty Y_i(t) dt$$

The probability that the first mutant spawns a sequence of $n+1$ mutants can, thus, be written as follows.

$$\begin{aligned} P(1 \rightarrow n + 1) &= \prod_{i=1}^n \left(\mu \int_0^\infty Y_i(t) dt \right) \\ &= \mu^n \prod_{i=1}^n \left(\int_0^\infty Y_i(t) dt \right) \end{aligned}$$

Lastly, we use $\tilde{\mu}$ to denote the rate at which new mutants with only one mutation emerge. For a given selection process, this rate may vary with the population size. For the Wright Fisher models of constant population size N , this expression simplifies to $\tilde{\mu} = N\mu$. We arrive at the following formulation for the rate at which new advantageous mutants emerge f_1 .

$$\begin{aligned} f_1 &= \tilde{\mu} \cdot P(1 \rightarrow n + 1) \\ &= \tilde{\mu} \cdot \mu^n \prod_{i=1}^n \left(\int_0^\infty Y_i(t) dt \right) \end{aligned}$$

Finally, we can introduce our clustering parameter k to scale both μ and μ^n , and multiply by a factor of $\frac{1}{k}$ to arrive at the average rate of valley-crossing in the temporally clustered process with parameter k .

$$\begin{aligned} f_k &= \frac{1}{k} \cdot (k \cdot \tilde{\mu}) \cdot (k \cdot \mu)^n \prod_{i=1}^n \left(\int_0^\infty Y_i(t) dt \right) \\ &= k^n \cdot \tilde{\mu} \cdot \mu^n \prod_{i=1}^n \left(\int_0^\infty Y_i(t) dt \right) \\ &= k^n \cdot f_1 \end{aligned}$$

One consequence of our assumption that none of the maladaptive intermediate mutants reaches fixation is that the composition of the population once the final mutant emerges in the limit of $\mu \rightarrow 0$ does not depend on the mutation rate. For population dynamics models with a constant population size in which the probability that an emerging mutant (absent further mutation events) reaches fixation only depends on this composition, such as the Wright-Fisher process, we can interpret $\frac{f_k}{f_1}$ therefore also as the ratio of the rates at which mutants with $n+1$ mutations fixate.

Analogously, in certain models of branching evolution in which the prospects of one branch do not depend on other co-evolving branches, such as the Galton-Watson process⁵⁸, whether the lineage of an emerging mutant with $(n + 1)$ mutations survives is independent of the mutation rate in the limit of $\mu \rightarrow 0$. The ratio $\frac{f_k}{f_1}$ therefore also reflects the ratio of the rates at which surviving lineages with $(n+1)$ adaptations arise in such models. We provide thorough treatment of dynamics in such processes below (SI2).

The above result suggests that the effect of temporal clustering on relative rates of valley crossing is substantial, and gets exponentially stronger the wider the fitness valley is (n). We validated this result with simulations of a Wright-Fisher process for a range of small values for μ (Fig S2 B-E [↗](#)). As we move away from the limit of rare mutations, the relative effect of temporal clustering $\frac{f_k}{f_1}$ gets smaller, as will be discussed below. However, as the absolute rates f_k and f_1 are proportional to μ^{n+1} , the absolute effect of temporal clustering on the rate of fixations of mutants with multi-step adaptations initially increases in μ (Fig S2 D [↗](#)).

SI2 Dynamics in a Branching Process

In the Galton-Watson process, a branching process with constant birth and death rates, it is possible to derive explicit formulas for the probability that a lineage will acquire a mutation before going extinct (often termed ‘evolutionary rescue’⁶²). Formulating this probability for the lineage of the first mutant in a 2-step adaptation sequence thus allows us to explore numerically how the magnitude of the effects that we describe behaves across a wide range of parameter combinations, which is especially attractive for parameter regions for which simulations become prohibitively expensive.

We thus investigated this probability of evolutionary rescue, which translates to f_k/μ in our model, as a function of k , μ and the birth/death ratio (Fig S3A-C [↗](#)). Note that the birth/death ratio that we considered here is that of the first mutant, not that of the wild type cell. Unsurprisingly, we found that as μ decreases, rescue probabilities for any k converge to zero for subcritical processes (birth/death ratios < 1). For supercritical processes (birth/death ratios > 1), rescue probabilities converge to one minus the extinction probability of the lineage – rescue only happens if the process expands indefinitely. Throughout, rescue probabilities of course increase in the expected time that the lineage persists, i.e. they increase with the birth death ratio. However, this increase is most steep for large μ and flattens for the subcritical range as μ decreases.

For the ratio of these escape probabilities for different k relative to that of the uniform process, i.e. for $\frac{f_k}{f_1} = \frac{f_k}{f_1}$, we found that in the subcritical regime, as μ decreases $\frac{f_k}{f_1}$ approaches k , consistent with our derivations in the previous section (SI1). In the supercritical regime, the condition $\int_0^\infty Y(t)dt < \infty$ used in our derivations above is violated – lineages have a non-zero chance of branching out indefinitely. Here we saw that as μ decreases towards zero, $\frac{f_k}{f_1}$ approaches 1 (Fig S3D-F [↗](#)).

Moreover, we found that $\frac{f_k}{f_1}$ consistently decreases with increasing birth-death ratio. As a result of the interplay between this decrease in $\frac{f_k}{f_1}$ and increase in the absolute rescue probability, we found that the absolute effect, $f_k - f_1$, is highest for birth/death ratios close to one (Fig S3G [↗](#)). These results remain as μ decreases (Fig S3H-I [↗](#)) but the magnitude of the absolute effect of course shrinks, as rescue of lineages that would otherwise go extinct (all subcritical lineages, and those lineages in the supercritical regime that go extinct by chance) generally becomes unlikely at small values of μ .

SI3 Dynamics as a function of time between bursts

In our derivations, we assumed that bursts are sufficiently spaced out in time, and last sufficiently long so that we can restrict attention to mutant lineages that emerge and go extinct within the duration of a single burst ($d \rightarrow \infty$). To investigate how effect sizes behave away from this limit, as a function of the time between bursts, we numerically approximated rescue probabilities for finite d .

Specifically, we approximated the Galton Watson process by a Markov process with finitely many states $s \in \{0, 1, \dots, N, \text{rescue}\}$, starting with $s(0) = 1$, where $s \in \{0, 1, \dots, N\}$ corresponds to the number of viable cells. We considered 0, N and *rescue* as absorbing states, and otherwise modelled transitions to states $s - 1$ or $s + 1$ as occurring at rates $s \cdot r_d$ and $s \cdot r_b$ respectively with constant birth and death rates r_d and r_b , and transitions to the *rescue* state as occurring at rate $s \cdot k \cdot \mu$ within bursts and at rate zero otherwise. We iteratively solved the corresponding forward equations of this continuous time Markov process numerically for subsequent cycles of in-burst and out-of-burst periods, until the summed mass on states $\{1, \dots, N - 1\}$ fell below a tolerance threshold, and we considered the joint mass on states N and *rescue* as approximation for the rescue probability. For our results in Fig S4 [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)

Generally, we showed that differences in multi-step adaptation rates emerge because lineages with a first mutation experience higher mutation rates under a temporally clustered mutation process compared to a uniform mutation process (SI1). As the burst duration and the time-interval between subsequent bursts decrease, this expected difference in mutation rates for mutant lineages diminishes because of the increased likelihood that a mutant lineage that emerged in a burst also experiences a non-burst phase. Consequently, we found that lowering d diminishes the effect of temporal clustering on the probability of producing a second mutant (Fig S4 A-C [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)). This pattern manifests both in the relative effects, f_k/f_1 (Fig S4 D-F [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)) and in the absolute effects $f_k - f_1$ (Fig S4 H-J [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)). However, for the absolute differences $f_k - f_1$, we found that the strength of this dependency varies non-monotonously with the birth-death ratio. While absolute differences under large d peak close to a birth-death ratio of one (Fig S3 [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#), Fig S4 [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)), we also observed the strongest decrease in absolute differences at these values for the birth-death ratio, so that under low d this peak turns into a local minimum of the absolute-difference curve (Fig S4 H, I [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)). This effect arises because the expected lifetime of a mutant lineage before going extinct increases steeply when moving from a subcritical process (birth-death ratios < 1) to a critical process (birth-death ratio = 1), making such lineages with higher birth-death ratios more sensitive to changes in mutation rates that occur long after the lineage first emerged.

SI4 Dynamics in exploration-exploitation setting


Our simulations show that for given μ the effect of increasing the clustering parameter k on the average fitness becomes negative at some k (Fig 3C [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)). In the simulations in this section, this effect is driven by two main factors. Firstly, after a re-drawing, the population often is not in a local maximum of the fitness landscape, and hence may be able to increase its fitness by single mutations (1-step adaptations), without tunneling. Waiting for a burst to occur and thereby delaying such local explorations decreases average fitness. Second, once the population has found a peak in the fitness landscape, having very high mutation rates in bursts temporarily scatters cells to lower points in the landscape, and may even cause drift to points of lower fitness. Both of these effects become apparent when considering a representative snapshot of the simulations (Fig 3D [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)). The misalignment of re-drawing and burst causes the population to remain in a valley for several divisions until the first burst occurs. Moreover, relative to similar snapshots of simulations with lower k (Fig 3E, F [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)), once the population has reached a peak, the scattering during bursts at higher k leads to sharper decreases in fitness, and the population may even remain in a lower fitness point after a burst (Fig 3D [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)).

This increased scattering can also be seen when going from $k = 1$ to $k = 5$ (Fig 3E, F [we used \$N=300\$ and a tolerance threshold of \$10^{-10}\$.](#)). However, at $k = 1$, the population spends much time in local maxima, as large jumps in fitness only occur right after re-drawings, whereas at $k = 5$ jumps occur also long after the landscape was re-drawn and


the population found a local maximum. These later jumps to higher points in the landscape correspond to tunneling events.

SI5 TSG-deactivation and ROS-associated mutagenesis


Following a reviewer suggestion, we also explored the contribution of mutation signatures attributed to reactive oxygen species (ROS; Fig S8 [↗](#)). However, we found that these signatures account for only small fractions of the total observed mutation signatures, suggesting that these mutagenic processes would not account for much of the fluctuation behavior of overall mutation rates and we saw no consistent correlations with TSG inactivation.




Supplementary Figure 1. Simulation results: valley-crossing under uniform vs. temporally clustered mutation rates in branching process. (A) Schematic of fitness landscape as in Fig 2. (B) Schematic of Branching Process model. (C) Simulation results for the Branching Process model. The mutation rate trajectories in (C) are chosen such that the total expected number of mutations under the uniform trajectory is identical to that under the temporally clustered trajectory.




Supplementary Figure 2. Effect of temporal clustering on stochastic tunneling rates as a function of the mutation rate. (A) Sketch of the mutation process with clustering parameter k . (B) Sketch of the fitness landscape as in Fig 2. (C) Description of the simulations. (D)-(E) Simulation results of a Wright-Fisher process with 100 cells over 10^6 generations with a fixed average mutation rate of 10^{-5} per cell per generation and varying temporal clustering k . The fitness is as in (B), with subsequent peaks differing in fitness by a factor of 1.5. However, on the y-axis in (F) and (G) we vary the fitness of the valleys relative to the preceding peak from 0.5 (as in (B)) to 1.5. Panel (D) shows the number of valleys crossed per generation (color bar in log10 values), where contrary to the previous panels it is no longer true that every valley crossing leads to a fixation – the next mutant might emerge before that. Panel (E) the share of these valley crossings that occur without fixation of the first mutant. (F) Simulation results of the process described in (A)-(C). Each simulation ran for $10^5/(k\mu)$ generations. The y-axis measures fixations of adaptive mutations per generation (see Fig 5D). (G) Fixations per generation relative to the uniform mutation process ($k=1$) for $k=5$ and $k=10$.




Supplementary Figure 3. Effect sizes for 2-step adaptations in a Galton-Watson process as a function of the mutation rate and the birth-death ratio. Each row corresponds to a unique value for μ . **(A)-(C)** Probabilities that a mutant lineage will give rise to a second mutant. **(D)-(F)** Ratios, f_k/f_1 , of the rates at which lineages with one mutation produce a lineage with both mutations for processes with different clustering parameters k . **(G)-(I)** Differences, $f_k - f_1$, of the rates at which lineages with one mutation produce a lineage with both mutations for processes with different clustering parameters k .




Supplementary Figure 4. Effect sizes for 2-step adaptations in a Galton-Watson process as a function of the distance d between subsequent burst phases. Each row corresponds to a unique value for d . **(A)-(C)** Probabilities that a mutant lineage will give rise to a second mutant. **(D)-(F)** Ratios, f_k/f_i , of the rates at which lineages with one mutation produce a lineage with both mutations for processes with different clustering parameters k . **(G)-(I)** Differences, $f_k - f_i$, of the rates at which lineages with one mutation produce a lineage with both mutations for processes with different clustering parameters k .




Supplementary Figure 5. TSG deactivation scores in TCGA. Distribution and correlation of APOBEC signature contribution and TSG deactivation scores as in Fig 4, sorted by cancer type.




Supplementary Figure 6. Synergistic mutation scores in TCGA. Distribution and correlation of APOBEC signature contribution and synergistic mutation scores, sorted by cancer type.



Supplementary Figure 7. Robustness to choice of correlation coefficient. Distribution and Spearman correlation of APOBEC signature contribution and TSG deactivation score (top) or synergistic mutation scores (bottom), sorted by cancer type. Confidence intervals for Spearman correlation coefficients were computed using a percentile bootstrap with 1000 resamples at the 95% confidence level.



Supplementary Figure 8. Robustness check: TSG deactivation and ROS in TCGA. Distribution and correlation of TSG deactivation score and ROS signature contribution measured as SBS18 (top) or as the joint contribution of SBS17a, SBS17b, SBS18 and SBS36. SBS18 only accounts for a small proportion of signature contributions and most correlations are close to zero. Including SBS17a, SBS17b and SBS36 which are less strongly associated with ROC does not change this correlation pattern relevantly.



Supplementary Figure 9. Temporal clustering and adaptation in fitness landscapes with varying ruggedness. (A) Sketch of the setup. Like in Fig 5, we assume that cells have two types of loci on which one-step or two-step adaptations are possible respectively. Fitness effects per locus are as in Fig 5. Here, we run simulations in which we vary the relative shares of the two types of loci. (B) Simulation results over time, averaged for 100 simulation replicates per row and clustering parameter k . Colors indicate the fitness ratio between simulations with $k=5$ and $k=1$ on a log10 scale. (C)-(D) show the corresponding absolute numbers of loci with adaptive mutations (i.e. loci in the black and blue states in panel (A)).

Additional information

Author ORCID iDs

Franziska Michor: <http://orcid.org/0000-0003-4869-8842>

Author notes

Competing interests: M.P. is a shareholder in Vertex Pharmaceuticals and a compensated consultant for the GLG Network. F.M. is a co-founder and consultant of Harbinger Health, a consultant for Zephyr AI, and is also on the board of directors of Recursion Pharmaceuticals. She declares that none of these relationships are directly or indirectly related to the content of this manuscript. The other authors declare no conflicts of interest.

References

1. Merlo LMF, Pepper JW, Reid BJ, Maley CC (2006) Cancer as an evolutionary and ecological process. *Nature Reviews Cancer* 2006 6:12 6:924-935 <https://doi.org/10.1038/nrc2013>

2. Yates LR, Campbell PJ (2012) Evolution of the cancer genome. *Nature Reviews Genetics* 2012 13:11 13:795-806 <https://doi.org/10.1038/nrg3317>
3. Neinavaie F, Ibrahim-Hashim A, Kramer AM, Brown JS, Richards CL (2021) The Genomic Processes of Biological Invasions: From Invasive Species to Cancer Metastases and Back Again. *Front Ecol Evol* 9:681100 <https://doi.org/10.3389/FEVO.2021.681100>
4. Araten DJ, Golde DW, Zhang RH, et al. (2005) A Quantitative Measurement of the Human Somatic Mutation Rate. *Cancer Res* 65:8111-8117 <https://doi.org/10.1158/0008-5472.CAN-04-1198>
5. Williams MJ, Werner B, Heide T, et al. (2018) Quantification of subclonal selection in cancer from bulk sequencing data. *Nature Genetics* 2018 50:6 50:895-903 <https://doi.org/10.1038/s41588-018-0128-6>
6. Werner B, Case J, Williams MJ, et al. (2020) Measuring single cell divisions in human tissues from multi-region sequencing data. *Nature Communications* 2020 11:1 11:1-9 <https://doi.org/10.1038/s41467-020-14844-6>
7. Galuppini F, Dal Pozzo CA, Deckert J, Loupakis F, Fassan M, Baffa R. (2019) Tumor mutation burden: From comprehensive mutational screening to the clinic. *Cancer Cell Int* 19:1-10 <https://doi.org/10.1186/S12935-019-0929-4>
8. Sha D, Jin Z, Budczies J, Kluck K, Stenzinger A, Sinicrope FA (2020) Tumor mutational burden as a predictive biomarker in solid tumors. *Cancer Discov* 10:1808-1825 <https://doi.org/10.1158/2159-8290.CD-20-0522>
9. Altrock PM, Liu LL, Michor F. (2015) The mathematics of cancer: integrating quantitative models. *Nature Reviews Cancer* 2015 15:12 15:730-745 <https://doi.org/10.1038/nrc4029>
10. Petljak M, Alexandrov LB, Brummel JS, et al. (2019) Characterizing Mutational Signatures in Human Cancer Cell Lines Reveals Episodic APOBEC Mutagenesis. *Cell* 176:1282-1294.e20. <https://doi.org/10.1016/j.CELL.2019.02.012>
11. Petljak M, Dananberg A, Chu K, et al. (2022) Mechanisms of APOBEC3 mutagenesis in human cancer cells. *Nature* 2022 607:7920 607:799-807 <https://doi.org/10.1038/s41586-022-04972-y>
12. Jamal-Hanjani M, Wilson GA, McGranahan N, et al. (2017) Tracking the Evolution of Non-Small-Cell Lung Cancer. *New England Journal of Medicine* 376:2109-2121 <https://doi.org/10.1056/NEJMOA1616288>
13. Lee JK, Lee J, Kim S, et al. (2017) ClonalHistory & genetic predictors of transformation into small-cell carcinomas from lung adenocarcinomas. *Journal of Clinical Oncology* 35:3065-3074 <https://doi.org/10.1200/JCO.2016.71.9096>
14. Wang Y, Robinson PS, Coorens THH, et al. (2023) APOBEC mutagenesis is a common process in normal human small intestine. *Nature Genetics* 2023 55:2 55:246-254 <https://doi.org/10.1038/s41588-022-01296-5>
15. Navin N, Kendall J, Troge J, et al. (2011) Tumor Evolution Inferred by Single Cell Sequencing. *Nature* 472:90 <https://doi.org/10.1038/NATURE09807>
16. Sottoriva A, Kang H, Ma Z, et al. (2015) A Big Bang model of human colorectal tumor growth. *Nat Genet* 47:209 <https://doi.org/10.1038/NG.3214>
17. Gao R, Davis A, McDonald TO, et al. (2016) Punctuated copy number evolution and clonal stasis in triple-negative breast cancer. *Nature Genetics* 2016 48:10 48:1119-1130 <https://doi.org/10.1038/ng.3641>
18. Minussi DC, Nicholson MD, Ye H, et al. (2021) Breast Tumors Maintain a Reservoir of Subclonal Diversity During Expansion. *Nature* 592:302 <https://doi.org/10.1038/S41586-021-03357-X>
19. Stephens PJ, Greenman CD, Fu B, et al. (2011) Massive genomic rearrangement acquired in a single catastrophic event during cancer development. *Cell* 144:27-40 <https://doi.org/10.1016/j.CELL.2010.11.055>
20. Zhang CZ, Spektor A, Cornils H, et al. (2015) Chromothripsis from DNA damage in micronuclei. *Nature* 2015 522:7555 522:179-184 <https://doi.org/10.1038/nature14493>

21. Baca SC, Prandi D, Lawrence MS, et al. (2013) Punctuated evolution of prostate cancer genomes. *Cell* **153**:666-677 <https://doi.org/10.1016/j.CELL.2013.03.021>
22. Drews RM, Hernando B, Tarabichi M, et al. (2022) A pan-cancer compendium of chromosomal instability. *Nature* 2022 606:7916 **606**:976-983 <https://doi.org/10.1038/s41586-022-04789-9>
23. Davis A, Gao R, Navin N. (2017) Tumor evolution: Linear, branching, neutral or punctuated?. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* **1867**:151-161 <https://doi.org/10.1016/j.BBCAN.2017.01.003>
24. Chakravarty D, Gao J, Phillips S, et al. (2017) OncoKB: A Precision Oncology Knowledge Base. *JCO Precis Oncol* **2017**:1-16 <https://doi.org/10.1200/PO.17.00011>
25. Knudson AG (1971) Mutation and Cancer: Statistical Study of Retinoblastoma. *Proceedings of the National Academy of Sciences* **68**:820-823 <https://doi.org/10.1073/PNAS.68.4.820>
26. Iwasa Y, Michor F, Nowak MA (2004) Stochastic tunnels in evolutionary dynamics. *Genetics* **166**:1571-1579 <https://doi.org/10.1534/GENETICS.166.3.1571>
27. Paige AJW (2003) Redefining tumour suppressor genes: exceptions to the two-hit hypothesis. *Cell Mol Life Sci* **60**:2147 <https://doi.org/10.1007/S00018-003-3027-6>
28. Wang LH, Wu CF, Rajasekaran N, Shin YK (2019) Loss of Tumor Suppressor Gene Function in Human Cancer: An Overview. *Cellular Physiology and Biochemistry* **51**:2647-2693 <https://doi.org/10.1159/000495956>
29. Park S, Supek F, Lehner B. (2021) Higher order genetic interactions switch cancer genes from two-hit to one-hit drivers. *Nature Communications* 2021 12:1 **12**:1-10 <https://doi.org/10.1038/s41467-021-27242-3>
30. Issa JP (2022) The Two-Hit Hypothesis Meets Epigenetics. *Cancer Res* **82**:1167-1169 <https://doi.org/10.1158/0008-5472.CAN-22-0405>
31. Gu Y, Yang D, Zou J, et al. (2010) Systematic interpretation of comutated genes in large-scale cancer mutation profiles. *Mol Cancer Ther* **9**:2186-2195 <https://doi.org/10.1158/1535-7163.MCT-10-0022>
32. Hsu MS, Yu JC, Wang HW, et al. (2010) Synergistic effects of polymorphisms in DNA repair genes and endogenous estrogen exposure on female breast cancer risk. *Ann Surg Oncol* **17**:760-771 <https://doi.org/10.1245/S10434-009-0802-0/METRICS>
33. Deming DA, Leystra AA, Nettekoven L, et al. (2013) PIK3CA and APC mutations are synergistic in the development of intestinal cancers. *Oncogene* 2014 33:17 **33**:2245-2254 <https://doi.org/10.1038/onc.2013.167>
34. Gerrish PJ, Lenski RE (1998) The fate of competing beneficial mutations in an asexual population. *Genetica* 1998 102:0 **102**:127-144 <https://doi.org/10.1023/A:1017067816551>
35. Wilke CO (2004) The Speed of Adaptation in Large Asexual Populations. *Genetics* **167**:2045-2053 <https://doi.org/10.1534/GENETICS.104.027136>
36. Fogle CA, Nagle JL, Desai MM (2008) Clonal Interference, Multiple Mutations and Adaptation in Large Asexual Populations. *Genetics* **180**:2163 <https://doi.org/10.1534/GENETICS.108.090019>
37. Sun R, Hu Z, Curtis C. (2018) Big Bang Tumor Growth and Clonal Evolution. *Cold Spring Harb Perspect Med* **8** <https://doi.org/10.1101/CSHPERSPECT.A028381>
38. Komarova NL, Sengupta A, Nowak MA (2003) Mutation-selection networks of cancer initiation: tumor suppressor genes and chromosomal instability. *J Theor Biol* **223**:433-450 [https://doi.org/10.1016/S0022-5193\(03\)00120-6](https://doi.org/10.1016/S0022-5193(03)00120-6)
39. Nowak MA, Michor F, Komarova NL, Iwasa Y. (2004) Evolutionary dynamics of tumor suppressor gene inactivation. *Proceedings of the National Academy of Sciences* **101**:10635-10638 <https://doi.org/10.1073/PNAS.0400747101>
40. Weinreich DM, Chao L. (2005) Rapid evolutionary escape by large populations from local fitness peaks is likely in nature. *Evolution (N Y)* **59**:1175-1182 <https://doi.org/10.1111/j.0014-3820.2005.TB01769.X>

41. Weissman DB, Desai MM, Fisher DS, Feldman MW (2009) The rate at which asexual populations cross fitness valleys. *Theor Popul Biol* **75**:286-300 <https://doi.org/10.1016/j.TPB.2009.02.006>
42. Proulx SR (2011) The rate of multi-step evolution in Moran and Wright–Fisher populations. *Theor Popul Biol* **80**:197-207 <https://doi.org/10.1016/j.TPB.2011.07.003>
43. Ashcroft P, Michor F, Galla T. (2015) Stochastic Tunneling and Metastable States During the Somatic Evolution of Cancer. *Genetics* **199**:1213-1228 <https://doi.org/10.1534/GENETICS.114.171553>
44. Van Egeren D, Madsen T, Michor F. (2018) Fitness variation in isogenic populations leads to a novel evolutionary mechanism for crossing fitness valleys. *Communications Biology* **2018** 1:1 1:1-9 <https://doi.org/10.1038/s42003-018-0160-1>
45. Wright S. (1931) Evolution in Mendelian Populations. *Genetics* **16**:97 <https://doi.org/10.1093/GENETICS/16.2.97>
46. Fisher R. (1922) On the dominance ratio. *Proc R Soc Edinb* **42**:321-341
47. Fischer BM, Olsen MWB, Ley CD, et al. (2006) How few cancer cells can be detected by positron emission tomography? A frequent question addressed by an in vitro study. *Eur J Nucl Med Mol Imaging* **33**:697-702 <https://doi.org/10.1007/S00259-005-0038-6/METRICS>
48. Alexandrov LB, Nik-Zainal S, Wedge DC, et al. (2013) Signatures of mutational processes in human cancer. *Nature* **2013** 500:7463 **500**:415-421 <https://doi.org/10.1038/nature12477>
49. Zhao M, Kim P, Mitra R, Zhao J, Zhao Z. (2016) TSGene 2.0: an updated literature-based knowledgebase for tumor suppressor genes. *Nucleic Acids Res* **44**:D1023-D1031 <https://doi.org/10.1093/NAR/GKV1268>
50. Nik-Zainal S, Alexandrov LB, Wedge DC, et al. (2012) Mutational processes molding the genomes of 21 breast cancers. *Cell* **149**:979-993 <https://doi.org/10.1016/j.CELL.2012.04.024>
51. Mas-Ponte D, Supek F. (2020) DNA mismatch repair promotes APOBEC3-mediated diffuse hypermutation in human cancers. *Nat Genet* **52**:958-968 <https://doi.org/10.1038/S41588-020-0674-6>
52. Campos PR, Adami C, Wilke CO (2004) Modelling Stochastic Clonal Interference. In: Ciobanu G, Rozenberg G (Eds). *Modelling in Molecular Biology* pp. 21-38 https://doi.org/10.1007/978-3-642-18734-6_2
53. Campos PRA, De Oliveira VM (2004) Mutational effects on the clonal interference phenomenon. *Evolution* **58**:932-937 <https://doi.org/10.1111/J.0014-3820.2004.TB00427.X>
54. Park SC, Krug J. (2007) Clonal interference in large populations. *Proc Natl Acad Sci U S A* **104**:18135-18140 <https://doi.org/10.1073/PNAS.0705778104>
55. Salehi S, Kabeer F, Ceglia N, et al. (2021) Clonal fitness inferred from time-series modelling of single-cell cancer genomes. *Nature* **595**:585-590 <https://doi.org/10.1038/S41586-021-03648-3>
56. McDonald TO, Cheng YC, Graser C, Nicol PB, Temko D, Michor F. (2023) Computational approaches to modelling and optimizing cancer treatment. *Nature Reviews Bioengineering* **2023** 1:10 **1**:695-711 <https://doi.org/10.1038/s44222-023-00089-7>
57. Díaz-Gay M, Vangara R, Barnes M, et al. (2023) Assigning mutational signatures to individual samples and individual somatic mutations with SigProfilerAssignment. *Bioinformatics* **39** <https://doi.org/10.1093/BIOINFORMATICS/BTAD756>
58. Watson HW, Galton F. (1875) On the Probability of the Extinction of Families. *The Journal of the Anthropological Institute of Great Britain and Ireland* **4**:138 <https://doi.org/10.2307/2841222>
59. Adler N, Bahcheli AT, Cheng KC, et al. (2023) Mutational processes of tobacco smoking and APOBEC activity generate protein-truncating mutations in cancer genomes. *Science Advances* **9** <https://doi.org/10.1126/sciadv.adh3083>
60. Butler K, Banday AR (2023) APOBEC3-mediated mutagenesis in cancer: causes, clinical significance and therapeutic potential. *Journal of Hematology & Oncology* **16**:31 <https://doi.org/10.1186/s13045-023-01425-5>

61. Gupta A, Gazzo A, Selenica P, et al. (2025) APOBEC3 mutagenesis drives therapy resistance in breast cancer. *Nature Genetics* **57** <https://doi.org/10.1101/2024.04.29.591453>
62. Azevedo RBR, Olofsson P (2021) A branching process model of evolutionary rescue. *Mathematical Biosciences* **341** <https://doi.org/10.1016/j.mbs.2021.108708>

Peer reviews

Reviewer #1 (Public review):

Summary:

Grasper et al. present a combined analysis of the role of temporal mutagenesis in cancer, which includes both theoretical investigation and empirical analysis of point mutations in TCGA cancer patient cohorts. They find that temporal elevated mutation rates contribute to cancer fitness by allowing fast adaptation when the fitness drops (due to previous deleterious mutations). This may be relevant in the case of tumor suppressor genes (TSG), which follow the 2-hit hypothesis (i.e., biallelic 2 mutations are necessary to deactivate TS), and in cases where temporal mutagenesis occurs (e.g. high APOBEC, ROS). They provide evidence that this scenario is likely to occur in patients, in some cancer types. This is an interesting and potentially important result that merits the attention of the target audience. Nonetheless, I have some questions (detailed below) regarding the design of the study, the tools and parametrization of the theoretical analysis and the empirical analysis - that I think if addressed would make the paper more solid and the conclusion more substantiated.

Strengths:

Combined theoretical investigation with empirical analysis of cancer patients

Weaknesses:

Parametrization and systematic investigation of theoretical tools and their relevant to tumor evolution

Comments on revisions:

The authors have adequately addressed my suggestions. I think some of the details provided in some of the replies to my comments (specifically with regard to my points 1, 4, 6ii; minor point 6) could be integrated into relevant text in the introduction, discussion and methods, to help the readers follow better the model and its interpretation - but this is up to the authors to decide what to emphasize.

<https://doi.org/10.7554/eLife.108058.2.sa2>

Reviewer #2 (Public review):

This work presents theoretical results concerning the effect of punctuated mutation on multistep adaptation along with empirical analysis of multistep adaptation in cancer. The empirical results are claimed to demonstrate the acceleration of multistep adaptation predicted theoretically. However, there is an important disconnect between the theoretical results and the empirical observations, such that it is not clear that punctuated mutation can produce the phenomena observed empirically. Furthermore, there are other plausible explanations for the empirical observations.

The theoretical work emphasizes the positive effect of punctuated mutation on the rate of crossing a "fitness valley", i.e., multistep adaptation where the first mutation is deleterious. The empirical work, however, focuses on inactivation of both alleles of a tumor suppressor

gene (TSG), for which the first mutation--inactivation of one gene copy--is expected to be neutral or slightly advantageous, not maladaptive as suggested by the authors. Pairs of genes with putative synergistic effects were also analyzed, but there is no indication that these generally involve fitness valleys either.

This disconnect is most glaring in Figure 4, in which the simulations are supposed to confirm that punctuated mutation can produce the empirical phenomena reported for TSG inactivation. If this is the case, it should be possible to produce such results in simulations in which inactivation of just one allele is neutral. Instead, simulations assuming a substantial fitness penalty (0.05) for the first mutation are presented. Contrary to what is claimed in the text (line 212), this is not a "biologically realistic" parameter value for TSG inactivation. The insensitivity of results to the size the fitness penalty is irrelevant: a substantial fitness penalty is qualitatively different from no penalty at all.

The paper does report a small (15%) effect of punctuation on the rate of multistep adaptation in the absence of a fitness valley. This effect is much smaller than the fourfold increase in the presence of a fitness valley. The results presented--a single stochastic run for each condition--are insufficient to establish that there is any effect at all: if we assume that the number of pairs of fixations (about 150-180 in each simulation) is Poisson distributed, the 15% difference is not statistically significant.

Assuming that this effect is genuine, it is likely due to a mutation rate that is unrealistically high (considering that "rescue" requires inactivation of a particular gene). Theoretical considerations suggest that punctuated mutation has little or no effect in the absence of a fitness valley in the limit of low mutation rate:

(A1) The authors' theoretical results for a Galton-Watson process (SI2) imply that there is no effect without a fitness valley in that limit. This is so because there is no effect in the "supercritical" regime. Cancer cells must be supercritical (otherwise there would be no net growth), and a neutral or advantageous mutant would remain in the supercritical regime.

(A2) Fig. S2D indicates, as far as I can tell from the colors, that punctuation makes little or no difference to the rate of adaptation in the absence of a fitness valley, i.e., for vertical axis values of 1 or more. I am not sure why the authors (line 129) point to this figure as evidence that punctuation speeds two-step adaptation when the first mutation is not maladaptive; the figure appears to say that it does not. The fraction of events due to "stochastic tunneling" of course increases with punctuation, but that does not change the fact that adaptation is no faster.

(A3) The authors' verbal argument to the contrary (line 124ff) is flawed. Despite the fact that even a mildly advantageous mutant is likely to go extinct, its expected frequency only increases with time, and that of a neutral allele remains constant over time. Thus, the average number of opportunities for a second mutation does not decrease with time since the first mutation, as it does when the first mutation is deleterious.

(A4) I ran some simulations for a Wright-Fisher population, and they seem to confirm the lack of an effect in the low mutation rate limit.

Thus, it is unclear whether punctuated mutation can explain the reported phenomena or should be expected to have major effects on the rate or nature of cancer cell adaptation.

I would also note that routes to inactivation of both copies of a TSG that are not accelerated by punctuation will dilute any effects of punctuation. An example is a single somatic mutation followed by loss of heterozygosity. Such mechanisms are not included in the theoretical analysis nor assessed empirically. If, for example, 90% of double inactivations were the result of such mechanisms with a constant mutation rate, a factor of two effect of punctuated mutagenesis would increase the overall rate by only 10%. Consideration of the

rate of apparent inactivation of just one TSG copy and of deletion of both copies would shed some light on the importance of this consideration.

Several factors besides the effects of punctuated mutation might explain or contribute to the empirical observations. Though these are now mentioned in the paper, I will list them here for clarity:

(B1) High APOBEC3 activity can select for inactivation of TSGs (references in Butler and Banday 2023, PMID 36978147). This could explain the empirical correlations.

(B2) Without punctuation, the rate of multistep adaptation is expected to rise more than linearly with mutation rate. Thus, if APOBEC signatures are correlated with a high mutation rate due to the action of APOBEC, this alone could explain the correlation with TSG inactivation.

(B3) The nature of mutations caused by APOBEC might explain the results. Notably, one of the two APOBEC mutation signatures, SBS13, is particularly likely to produce nonsense mutations. The authors count both nonsense and missense mutations, but nonsense mutations are more likely to inactivate the gene, and hence to be selected.

<https://doi.org/10.7554/eLife.108058.2.sa1>

Author response:

The following is the authors' response to the original reviews

eLife Assessment

his valuable study presents a theoretical model of how punctuated mutations influence multistep adaptation, supported by empirical evidence from some TCGA cancer cohorts. This solid model is noteworthy for cancer researchers as it points to the case for possible punctuated evolution rather than gradual genomic change. However, the parametrization and systematic evaluation of the theoretical framework in the context of tumor evolution remain incomplete, and alternative explanations for the empirical observations are still plausible.

We thank the editor and the reviewers for their thorough engagement with our work. The reviewers' comments have drawn our attention to several important points that we have addressed in the updated version. We believe that these modifications have substantially improved our paper.

There were two major themes in the reviewers' suggestions for improvement. The first was that we should demonstrate more concretely how the results in the theoretical/stylized modelling parts of our paper quantitatively relate to dynamics in cancer.

To this end, we have now included a comprehensive quantification of the effect sizes of our results across large and biologically-relevant parameter ranges. Specifically, following reviewer 1's suggestion to give more prominence to the branching process, we have added two figures (Fig S3-S4) quantifying the likelihood of multi-step adaptation in a branching process for a large range of mutation rates and birth-death ratios. Formulating our results in terms of birth-death ratios also allowed us to provide better intuition regarding how our results manifest in models with constant population size vs models of growing populations. In particular, the added figure (Fig S3) highlights that the effect size of temporal clustering on the probability of successful 2-step adaptation is very sensitive to the probability that the lineage of the first mutant would go extinct if it did not acquire a second mutation. As a result, the phenomenon we describe is biologically likely to be most effective in those phases during tumor evolution in which tumor growth is constrained. This important pattern had

not been described sufficiently clearly in the initial version of our manuscript, and we thank both reviewers for their suggestions to make these improvements.

The second major theme in the reviewers' suggestions was focused on how we relate our theoretical findings to readouts in genomic data, with both reviewers pointing to potential alternative explanations for the empirical patterns we describe.

We have now extended our empirical analyses following some of the reviewers' suggestions. Specifically, we have included analyses investigating how the contribution of reactive oxygen species (ROS)-related mutation signatures correlates with our proxies for multi-step adaptation; and we have included robustness checks in which we use Spearman instead of Pearson correlations. Moreover, we have included more discussion on potential confounds and the assumptions going into our empirical analyses as well as the challenges in empirically identifying the phenomena we describe.

Below, we respond in detail to the individual comments made by each reviewer.

Public Reviews:

Reviewer #1 (Public review):

Summary:

Grasper et al. present a combined analysis of the role of temporal mutagenesis in cancer, which includes both theoretical investigation and empirical analysis of point mutations in TCGA cancer patient cohorts. They find that temporally elevated mutation rates contribute to cancer fitness by allowing fast adaptation when the fitness drops (due to previous deleterious mutations). This may be relevant in the case of tumor suppressor genes (TSG), which follow the 2-hit hypothesis (i.e., biallelic 2 mutations are necessary to deactivate TS), and in cases where temporal mutagenesis occurs (e.g., high APOBEC, ROS). They provide evidence that this scenario is likely to occur in patients with some cancer types. This is an interesting and potentially important result that merits the attention of the target audience. Nonetheless, I have some questions (detailed below) regarding the design of the study, the tools and parametrization of the theoretical analysis, and the empirical analysis, which I think, if addressed, would make the paper more solid and the conclusion more substantiated.

Strengths:

Combined theoretical investigation with empirical analysis of cancer patients.

Weaknesses:

Parametrization and systematic investigation of theoretical tools and their relevance to tumor evolution.

We sincerely thank Reviewer 1 for their comments. As communicated in more detail in the point-by-point replies to the "Recommendations for the authors", we have revised the paper to address these comments in various ways. To summarize, Reviewer 1 asked for (1) more comprehensive analyses of the parameter space, especially in ranges of small fitness effects and low mutation rates; (2) additional clarifications on details of mechanisms described in the manuscript; and (3) suggested further robustness checks to our empirical analyses. We have addressed these points as follows: we have added detailed analyses of dynamics and effect sizes for branching processes (see Sections SI2 and SI3 in the Supplementary Information, as well as Figures S3 and S4). As suggested, these additions provide characterizations of effect sizes in biologically relevant parameter ranges (low mutation rates and smaller fitness effect sizes), and extend our descriptions to processes with dynamically changing population sizes. Moreover, we have added further clarifications at suggested

points in the manuscript, e.g. to elaborate on the non-monotonicities in Fig 3. Lastly, we have undertaken robustness checks using Spearman rather than Pearson correlation coefficients to quantify relations between TSG deactivation and APOBEC signature contribution, and have performed analyses investigating dynamics of reactive oxygen species-associated mutagenesis instead of APOBEC.

Reviewer #2 (Public review):

This work presents theoretical results concerning the effect of punctuated mutation on multistep adaptation and empirical evidence for that effect in cancer. The empirical results seem to agree with the theoretical predictions. However, it is not clear how strong the effect should be on theoretical grounds, and there are other plausible explanations for the empirical observations.

Thank you very much for these comments. We have now substantially expanded our investigations of the parameter space as outlined in the response to the “eLife Assessment” above and in the detailed comments below (A(1)-A(3)) to convey more quantitative intuition for the magnitude of the effects we describe for different phases of tumor evolution. We agree that there could be potential additional confounders to our empirical investigations besides the challenges regarding quantification that we already described in our initial version of the manuscript. We have thus included further discussion of these in our manuscript (see replies to B(1)-B(3)), and we have expanded our empirical analyses as outlined in the response to the “eLife Assessment”.

For various reasons, the effect of punctuated mutation may be weaker than suggested by the theoretical and empirical analyses:

(A1) The effect of punctuated mutation is much stronger when the first mutation of a two-step adaptation is deleterious (Figure 2). For double inactivation of a TSG, the first mutation--inactivation of one copy--would be expected to be neutral or slightly advantageous. The simulations depicted in Figure 4, which are supposed to demonstrate the expected effect for TSGs, assume that the first mutation is quite deleterious. This assumption seems inappropriate for TSGs, and perhaps the other synergistic pairs considered, and exaggerates the expected effects.

Thank you for highlighting this discrepancy between Figure 2 and Figure 4. For computational efficiency and for illustration purposes, we had opted for high mutation rates and large fitness effects in Figure 2; however, our results are valid even in the setting of lower mutation rates and fitness effects. To improve the connection to Figure 4, and to address other related comments regarding parameter dependencies, we have now added more detailed quantification of the effects we describe (Figures SF3 and SF4) to the revised manuscript. These additions show that the effects illustrated in Figure 2 retain large effect sizes when going to much lower mutation rates and much smaller fitness effects. Indeed, while under high mutation rates we only see the large relative effects if the first mutation is highly deleterious, these large effects become more universal when going to low mutation rates.

In general, it is correct that the selective disadvantage (or advantage) conveyed by the first mutation affects the likelihood of successful 2-step adaptations. It is also correct that the magnitude of the ‘relative effect’ $R_e = \alpha R_f + (1 - \alpha) R_g$, of temporal clustering on valley-crossing is highest if the lineage with only the first of the two mutations is vanishingly unlikely to produce a second mutant before going extinct. If the first mutation is strongly deleterious, the lineage of such a first mutant is likely to quickly go extinct – and therefore also more likely to do so before producing a second mutant.

However, this likelihood of producing the second mutant is also low if the mutation rate is low. As our added figure (Figure SF3) illustrates, at low mutation rates appropriate for cancer cells, $R_g = \alpha R_f + (1 - \alpha)R_s$, is insensitive to the magnitude of the fitness disadvantage for large parts of the parameter space. Especially in populations of constant size (approximated by a birth/death ratio of 1), the relative effects $R_g = \alpha R_f + (1 - \alpha)R_s$, for first mutations that reduce the birth rate by 0.5 or by 0.05 are indistinguishable (Figure SF3f).

Moreover, the absolute effect $(f_k - f_1)$, as we discuss in the paper (Figures SF2 and SF3) is largest in regions of the parameter space in which the first mutant is not infinitesimally unlikely to produce a second mutant (and f_k and f_1 would be infinitesimally small), but rather in parameter regions in which this first mutant has a non-negligible chance to produce a second mutant. The absolute effect $(f_k - f_1)$ therefore peaks around fitness-neutral first mutations. While the next comment (below) says that our empirical investigations more closely resemble comparisons of relative effects and not absolute effects, we would expect that the observations in our data come preferentially from multi-step adaptations with large absolute effect since the absolute effect is maximal when both f_k and f_1 are relatively high.

In summary, we believe Figure 2, while having exaggerated parameters for very defensible reasons, is not a misleading illustration of the general phenomenon or of its applicability in biological settings, as effect sizes remain large when moving to biologically realistic parameter ranges. To clarify this issue, we have largely rewritten the relevant paragraphs in the results section and have added two additional figures (Figures SF3 and SF4) as well as a section in the SI with detailed discussion (SI2).

(A2) More generally, parameter values affect the magnitude of the effect. The authors note, for example, that the relative effect decreases with mutation rate. They suggest that the absolute effect, which increases, is more important, but the relative effect seems more relevant and is what is assessed empirically.

Thank you for this comment. As noted in the replies to the above comments, we have now included extensive investigations of how sensitive effect sizes are to different parameter choices. We also apologize for insufficiently clearly communicating how the quantities in Figure 4 relate to the findings of our theoretical models.

The challenge in relating our results to single-timepoint sequencing data is that we only observe the mutations that a tumor has acquired, but we do not directly observe the mutation rate histories that brought about these mutations. As an alternative readout, we therefore consider (through rough proxies: TSGs and APOBEC signatures) the amount of 2-step adaptations per acquired/retained mutation. While we unfortunately cannot control for the average mutation rate in a sample, we motivate using this “TSG-deactivation score” by the hypothesis that for any given mutation rate, we expect a positive relationship between the amount of temporal clustering and the amount of 2-step adaptations per acquired/retained mutation. This hypothesis follows directly from our theoretical model where it formally translates to the statement that for a fixed μ , is increasing in μ .

However, while both quantities f_k/f_1 or $(f_k - f_1)$ from our theoretical model relate to this hypothesis – both are increasing in k –, neither of them maps directly onto the formulation of our empirical hypothesis.

We have now rewritten the relevant passages of the manuscript to more clearly convey our motivation for constructing our TSG deactivation score in this form (P. 4-6).

(A3) Routes to inactivation of both copies of a TSG that are not accelerated by punctuation will dilute any effects of punctuation. An example is a single somatic mutation followed by loss of heterozygosity. Such mechanisms are not included in the

theoretical analysis nor assessed empirically. If, for example, 90% of double inactivations were the result of such mechanisms with a constant mutation rate, a factor of two effect of punctuated mutagenesis would increase the overall rate by only 10%. Consideration of the rate of apparent inactivation of just one TSG copy and of deletion of both copies would shed some light on the importance of this consideration.

This is a very good point, thank you. In our empirical analyses, the main motivation was to investigate whether we would observe patterns that are qualitatively consistent with our theoretical predictions, i.e. whether we would find positive associations between valley-crossing and temporal clustering. Our aim in the empirical analyses was not to provide a quantitative estimate of how strongly temporally clustered mutation processes affect mutation accumulation in human cancers. We hence restricted attention to only one mutation process which is well characterized to be temporally clustered (APOBEC mutagenesis) and to only one category of (epi)genomic changes (SNPs, in which APOBEC signatures are well characterized). Of course, such an analysis ignores that other mutation processes (e.g. LOH, copy number changes, methylation in promoter regions, etc.) may interact with the mechanisms that we consider in deactivating Tumor suppressor genes.

We have now updated the text to include further discussion of this limitation and further elaboration to convey that our empirical analyses are not intended as a complete quantification of the effect of temporal clustering on mutagenesis in-vivo (P. 10,11).

Several factors besides the effects of punctuated mutation might explain or contribute to the empirical observations:

(B1) High APOBEC3 activity can select for inactivation of TSGs (references in Butler and Banday 2023, PMID 36978147). This selective force is another plausible explanation for the empirical observations.

Thank you for making this point. We agree that increased APOBEC3 activity, or any other similar perturbation, can change the fitness effect that any further changes/perturbations to the cell would bring about. Our empirical analyses therefore rely on the assumption that there are no major confounding structural differences in selection pressures between tumors with different levels of APOBEC signature contributions. We have expanded our discussion section to elaborate on this potential limitation (P. 10-11).

While the hypothesis that APOBEC3 activity selects for inactivation of TSGs has been suggested, there remain other explanations. Either way, the ways in which selective pressures have been suggested to change would not interfere relevantly with the effects we describe. The paper cited in the comment argues that “high APOBEC3 activity may generate a selective pressure favoring” TSG mutations as “APOBEC creates a high [mutation] burden, so cells with impaired DNA damage response (DDR) due to tumor suppressor mutations are more likely to avert apoptosis and continue proliferating”. To motivate this reasoning, in the same passage, the authors cite a high prevalence of TP53 mutations across several cancer types with “high burden of APOBEC3-induced mutations”, but also note that “this trend could arise from higher APOBEC3 expression in p53-mutated tumors since p53 may suppress APOBEC3B transcription via p21 and DREAM proteins”.

Translated to our theoretical framework, this reasoning builds on the idea that APOBEC3 activity increases the selective advantage of mutants with inactivation of both copies of a TSG. In contrast, the mechanism we describe acts by altering the chances of mutants with only one TSG allele inactivated to inactivate the second allele before going extinct. If homozygous inactivation of TSGs generally conveys relatively strong fitness advantages, lineages with homozygous inactivation would already be unlikely to go extinct. Further increasing the fitness advantage of such lineages would thus manifest mostly in a quicker spread of these lineages, rather than in changes in the chance that these lineages survive. In

turn, such a change would have limited effect on the “rate” at which such 2-step adaptations occur, but would mostly affect the speed at which they fixate. It would be interesting to investigate these effects empirically by quantifying the speed of proliferation and chance of going extinct for lineages that newly acquired inactivating mutations in TSGs.

Beyond this explicit mention of selection pressures, the cited paper also discusses high occurrences of mutations in TSGs in relation to APOBEC. These enrichments, however, are not uniquely explained by an APOBEC-driven change in selection pressures. Indeed, our analyses would also predict such enrichments.

(B2) Without punctuation, the rate of multistep adaptation is expected to rise more than linearly with mutation rate. Thus, if APOBEC signatures are correlated with a high mutation rate due to the action of APOBEC, this alone could explain the correlation with TSG inactivation.

Thank you for making this point. Indeed, an identifying assumption that we make is that average mutation rates are balanced between samples with a higher vs lower APOBEC signature contribution. We cannot cleanly test this assumption, as we only observe aggregate mutation counts but not mutation rates. However, the fact that we observe an enrichment for APOBEC-associated mutations among the set of TSG-inactivating mutations (see Figure 4F) would be consistent with APOBEC-mutations driving the correlations in Fig 4D, rather than just average mutation rates. We have now added a paragraph to our manuscript to discuss these points (P. 10-11).

(B3) The nature of mutations caused by APOBEC might explain the results. Notably, one of the two APOBEC mutation signatures, SBS13, is particularly likely to produce nonsense mutations. The authors count both nonsense and missense mutations, but nonsense mutations are more likely to inactivate the gene, and hence to be selected.

Thank you for making this point. We have included it in our discussion of potential confounders/limitations in the revised manuscript (P. 10-11).

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

Specific questions/comments/suggestions:

(1) For the theoretical investigation, the authors use the Wright-Fisher model with specific parameters for the decrease/increase in the fitness (0.5,1.5). This model is not so relevant to cancer, because it assumes a constant population size, while in cancer, the population is dynamic (increasing, if the tumor grows). Although I see they mention relevance to the branching process (in SI), I think the branching process should be bold in the main text and the Wright-Fisher in SI (or even dropped).

Thank you for this comment. We agree that too little attention had been given to the branching process in the original version of our manuscript. While the Wright-Fisher process is computationally efficient to simulate and thus lends itself to clean simulations for illustrative examples, it did lead us to put undue emphasis on populations of constant size.

The added Figures SF2 and SF3 now focus on branching processes, and we have substantially expanded our discussion of how dynamics differ as a function of the population-size trajectory (constant vs growing; SI2, P. 4,9,10). Generally, we do believe that it is appropriate to consider both regimes. If tumors evolve from being confined within their site of origin to progressively invading adjacent tissues and organ compartments, they traverse different regions of the birth-death ratio parameter space. Moreover, the timing of transitions between phases of more or less constrained growth is likely closely tied to adaptation dynamics, since

breaching barriers to expansion requires adapting to novel environments and selection pressures.

We hope that the revised version of the manuscript conveys these points more clearly, and thank you for alerting us to this imbalance in the original version of our manuscript.

(2) The parameters 0.5 (decrease in fitness) and 1.5 (increase in fitness) seem exaggerated (the typical values for the selective advantage are usually much lower (by an order of magnitude). The same goes for the mutation rate. The authors chose values of the order 0.001, while in cancer (and generally) it is much lower than that (10^{-5} - 10^{-6}). I think that generally, the authors should present a more systematic analysis of the sensitivity of the results to these parameters.

Thank you very much for this very important comment. We have made this a major focus in our revisions (see our reply to the editor's comments). As suggested, we have now added further analyses to explore more biologically relevant parameter regimes. Reviewer 2 has made a similar remark, and to avoid redundancies, we point for a more detailed response to our response to that comment (A1).

(3) In Figure 3, the authors explore the sensitivity to μ (mutation rate) and k (temporal clustering) and find a non-monotonic behavior (Figure 3C). However, this behavior is not well explained. I think some more explanations are required here.

Thank you for pointing this out. We had initially relegated the more detailed explanations to the SI2 (which in the revised manuscript became SI4), but are happy to provide more elaboration in the main text, and have done so now (P. 5).

For μ , the non-monotonicity reflects the exploration-exploitation tradeoff that this section is dedicated to: very small values (little exploration) prevent the population from finding fitness peaks. In contrast, once a fitness peak is reached, excessively large values (little exploitation) scatter the population away from this peak to points of lower fitness.

For k , the most relevant dynamic is that at high k , the population becomes unable to find close-by fitness improvements (1-step adaptations) if it is not in a burst. As k increases, this delay in adaptation (until a burst occurs) eventually comes to outweigh the benefits of high k (better ability to undergo multi-step adaptations). Additionally, if $k \cdot \mu$ becomes very large, clonal interference eventually leads to diminishing exploration-returns when k is increased further (Fig 5C), as the per-cell likelihood of finding a specific fitness peak eventually saturates and increasing only causes multiple cells to find the same peak, rather than one cell finding this peak and its lineage fixating in the population.

(4) In Figure 5, where the authors show the accumulation of the first (red; deleterious mutation) and second (blue; advantageous mutation), it seems that the fraction of deleterious mutations is much lower than that of advantageous mutations. This is opposite to the case of cancer, where most of the mutations are 'passengers', (slightly) deleterious or neutral mutations. Can the author explain this discrepancy and generally the relation of their parametrization to deleterious vs. advantageous mutations?

Thank you for this comment. In general, we have focused attention in our paper on sequences of mutations that bring about a fitness increase. We call those sequences 'adaptations' and categorize these as one-step or multi-step, depending on whether or not they contain intermediate states with a fitness disadvantage.

In our modelling, we do not consider mutations that are simply deleterious and are not a necessary part of a multi-step adaptation sequence. The motivation for this abstraction is, firstly, to focus on adaptation dynamics, and secondly, that in certain limits (small μ and large constant population sizes), lineages with only deleterious mutations have a probability

close to one of going extinct, so that any emerging deleterious mutant would likely be 'washed out' of the population before a new mutation emerges.

However, whether the dynamics of how neutral or deleterious passenger mutations are acquired also vary relevantly with the extent of temporal clustering is a valid and interesting question that would warrant its own study. The types of theoretical arguments for such an investigation would be very similar to the ones we use in our paper.

(5) The theoretical investigation assumes a multi/2-step adaptation scenario where the first mutation is deleterious and the second is advantageous. I think this should be generalized and further explored. For example, what happens when there are multiple mutations that are slightly deleterious (as probably is the case in cancer) and only much later mutations confer a selective advantage? How stable is the "valley crossing" if more deleterious mutations occur after the 2 steps?

This is also an important point and relates in part to the previous comment (4). For discussion of interactions with deleterious mutations, please see the reply to comment (4).

Regarding generalizations of this valley-crossing scenario, note that any sequence of mutations that increases fitness can be decomposed into sequences of either one-step or multi-step adaptations, as defined in the paper. Therefore, if all intermediate states before the final selectively advantageous state have a selective disadvantage making the lineages of such cells likely to go extinct, then our derivations in S1 apply, and the relative effect of temporal clustering becomes $\frac{f_k}{f_1} = k^n$ where n is the number of intermediate states. If, conversely, any of the intermediate states already had a selective advantage, then our model would consider the subsequence until this first mutation with a selective advantage as its individual (one-step or multi-step) "adaptation".

The second question, "How stable is the "valley crossing" if more deleterious mutations occur after the 2 steps?", touches on a different property of the population dynamics, namely on how the fate of a mutant lineage depends on how this lineage emerged. In our paper, we compare different levels of temporal clustering for a *fixed* average mutation rate. This choice implies that, if we assume that the mutant that emerges from a valley-crossing does not go extinct, then the number of deleterious mutations expected to occur in this lineage, once emerged, will not depend on the extent of temporal clustering. However, if in-burst mutation rates increased the expected burden of early acquired deleterious mutations sufficiently much to affect the probability that the lineage with a multi-step adaptation goes extinct before the burst ends, then there may indeed be an interaction between effects of deleterious passengers and temporal clustering. We would, however, expect effects on this probability of early extinction to be relatively minor, since such a lineage with a selective advantage would quickly grow to large cell-numbers implying that it would require a large number of co-occurring and sufficiently deleterious mutations across these cells for the lineage to go extinct.

(6) For the empirical analysis of TCGA cohorts, the authors focus on the contribution of APOBEC mutations (via signature analysis) to temporal mutagenesis. They find only a few cancer types (Figure 4D) that follow their prediction (in Figure 4C) of a correlation between TSG deactivation and temporal mutations in bursts. I think two main points should be addressed:

Thank you for this comment. We will respond in detail to the corresponding points below, but would like to note here that while we find this correlation "in only a few cancer types", we also show that only few cancer types have relevant proportions of mutations caused by APOBEC, and it is precisely in these cancer types that we find a correlation. We have clarified this aspect in the revised version of the manuscript (P.7).

(i) APOBEC is not the only cause for temporal mutagenesis. For example, elevated ROS and hypoxia are also potential contributors - it might therefore be important to extend the signature analysis (to include more possible sources for temporal mutagenesis). Potentially, such an extension may show that more cancer types follow the author's prediction.

Thank you for this interesting suggestion. We have now included analogous analyses for contributions of signature SBS18 which is associated with ROS mutagenesis, and for the joint contribution of signatures SBS17a, SBS17b, SBS18 and SBS36, which all have been shown (some in a more context-dependent manner) to be associated with ROS mutagenesis. When doing so, we do not find a clear trend. However, we also do not find these signatures to account for substantial proportions of the acquired mutations, meaning that ROS mutagenesis likely also does not account for much of the variation in how temporally clustered the mutation rate trajectories of different tumors are. We have incorporated these results and their discussion in the manuscript (SI5 and Fig S8).

(ii) The TSG deactivation score used by the authors only counts the number of mutations and does not consider if the 2 mutations are biallelic, which is highly important in this case. There are ways to investigate the specific allele of mutations in TCGA data (for example, see Ciani et al. Cell Sys 2022 PMID: 34731645). Given the focus on TSG of this study, I think it is important to account for this in the analysis.

Thank you for making this point. We did initially consider inferring allele-specific mutation status, but decided against it as this would have shrunk our dataset substantially, thus potentially introducing unwanted biases. Determining whether two mutations lie on the same or on different alleles requires either (1) observing sequencing reads that either cover the loci of both mutations, or (2) tracing whether (sets of) other SNPs on the same gene co-occur exclusively with one of the two considered mutations. These requirements lead to a substantial filtering of the observed mutations. Moreover, this filtering would be especially strong for tumors with a small overall mutation burden, as these would have fewer co-occurring SNPs to leverage in this inference. We would have hence preferentially filtered out TSG-deactivating mutations in tumors with low mutation burden. We have modified the text to address this point (P.14).

(7) To continue point 4. I wonder why some known cancer types with high APOBEC signatures (e.g., lung, mentioned in the introduction) do not appear in the results of Figure 4. Can the author explain why it is missed?

We do provide complete results for all categories in Supplementary Figure 3. To not overwhelm the figure in the main text, we only show the four categories with the highest average APOBEC signature contribution, beyond those four, average APOBEC signature contributions quickly drop. Lung-related categories do not feature in these top four (Lung squamous cell carcinoma are fifth and Lung adenocarcinoma are eighth in this ordering).

Minors:

(1) It is worth mentioning the relevance to resistance to treatment (see <https://www.nature.com/articles/s41588-025-02187-1> .

Thank you for this suggestion. We have included a mention of the relation to this paper in the discussion section (P. 11).

(2) Some of the figures' resolution should be improved - specifically, Figures 4, S1, and S5, which are not clear/readable.

Thank you for pointing this out. This was the result of conversion to a word document. We will provide tif files in the revisions to have better resolution.

(3) Regarding Figure 3e,f. How come that moving from $K=1$ to $K=I$ doesn't show any changes in fitness - it looks as if in both cases the value fluctuates around comparable mean fitness? Is that the case?

While fitness differences between simulations with different k manifest robustly over long time-horizons (see Fig 3C with results over generations), there are various sources of substantial stochasticity that make the fitness values in these short-term plots (Fig3D-F) imperfect illustrations of how long-term average fitness behaves. For instance, fitness landscapes are drawn randomly which introduces variability in how high and how close-by different fitness peaks are. Similarly, there is substantial randomness since both the type (direction on the 2-D fitness landscape) and the timing of mutation are stochastic.

The short-term plots in Fig3D-F are intended to showcase representative dynamics of transitions between points on the genotype space with different fitness values following a redrawing of the landscape – but not necessarily to provide a comparison between the height of the attained (local) fitness-maxima.

(4) Figures 4c,d - correlation should be Spearman, not Pearson (it's not a linear relationship).

Thank you for this comment. As a robustness check, we have generated the same figures using Spearman and not Pearson correlations and find results that are qualitatively consistent with the initially shown results. Indeed, using Spearman correlations, all four cancer types from Fig 4D have significant correlations.

(5) Typo for E) "...in samples of the cancer types in (C) were caused by APOBEC" - it should be D (not C) I guess.

Thank you for catching this. We fixed the typo.

(6) Figure 5 - the mutation rate is too high (0.001), sensitivity to that? Also the fitness change is exaggerated (0.5, 1.5), and the division of mutations to 100 and 100 (200 in total) loci is not clear.

Thank you for making this point. In this simulation setting it is unfortunately computationally prohibitively expensive to perform simulations at biologically realistic mutation rates. Therefore, we have scaled up the mutation rate while scaling down the population size. Moreover, the choice of model here is not meant to resemble a biologically realistic dynamic, but rather to create a stylized setting to be able to consider the interplay between clonal interference and facilitated valley-crossing in isolation. The key result from this figure is the separation of time scales at which low or high temporal clustering maximizes adaptability.

However, known parameter dependencies in these models allow us to reason about how tuning individual parameters of this stylized model would affect the relative importance of effects of clonal interference. This relative importance is largest when mutants are likely to co-occur on different competing clones in a population. The likelihood of such co-occurrences decreases substantially if decreasing the mutation rate to biologically realistic values. However, this likelihood also sensitively depends on the time that it takes a clone with a one-step adaptation to spread through the population. Smaller fitness advantages, as well as larger population sizes, slow down this process of taking over the population, which increases the likelihood of clonal interference. We now discuss these points in our revised manuscript (P. 8).

1. In the results text (last section) "Performing simulations for 2-step adaptations, we found that fixation rates are non-monotone in k . While at low k increasing k leads to a steep increase in the fixation rate, this trend eventually levels off and becomes negative, with further increases in k leading to a decrease in the fixation rate". Where are the results of this? It should be bold and apparent.

Thank you for alerting us that this is unclear. The relevant figure reference is indeed Fig 5C as in the preceding passage in the manuscript. However, we noticed that due to the presence of the steadily decreasing black line for 1-step adaptations, it is not easy to see that also the blue line is downward sloping. We have added a further reference to Fig 5C, and have adapted the grid spacing in the background of that figure-panel to make this trend more easily visible.

(8) Although not inconceivable, conclusions regarding resistance in the discussion are overstated. If you want to make this statement, you need to show that in resistant tumors, the temporal mutagenesis is responsible for progression vs. non-resistant/sensitive cases (is that the case), otherwise this should be toned down.

Thank you for pointing this out. We have tempered these conclusions in the revised version of the manuscript (P. 11).

Reviewer #2 (Recommendations for the authors):

(1) It might be useful to look specifically at X-linked TSGs. On the authors' interpretation, their relative inactivation rates should not be correlated with APOBEC signatures in males (but should be in females), though the size of the dataset may preclude any definite conclusions.

Thank you for this suggestion. Indeed, the size of the dataset unfortunately makes such analyses infeasible. Moreover, it is not clear whether X-linked TSGs might have structurally different fitness dynamics than TSGs on other chromosomes. However, this is an interesting suggestion worth following up on as more synergistic pairs confined to the X-chromosome are getting identified.

(2) Might there be value in distinguishing tumors that carry mutations expected to increase APOBEC expression from those that do not? Among several reasons, an APOBEC signature due to such a mutation and an APOBEC signature due to abortive viral infection may differ with respect to the degree of punctuation.

This is also an interesting suggestion for future investigations, but for which we unfortunately do not have sufficient information to build a meaningful analysis. In particular, it is unclear to what extent the degree and manifestation of episodicity/punctuation varies between these different mechanisms. Burst duration and intensity, as well as out-of-burst baseline rates of APOBEC mutagenesis likely differ in ways that are yet insufficiently characterized, which would make any result of analyses like these in Fig 4 hard to interpret.

(3) Also, in that paragraph, is "proportional to" used loosely to mean "an increasing function of"?

Thank you for this comment. We are not quite sure which paragraph is meant, but we use the term "proportional" in a literal sense at every point it is mentioned in the paper.

For the occurrences of the term on pages 3, 10 and 11, the word is used in reference to probabilities of reproduction (division in the branching process, or 'being drawn to populate a spot in the next generation' in the WF process) being "proportional" to fitness. These probabilities are constructed by dividing each individual cell's fitness by the total fitness

summed across all cells in the population. As the population acquires fitness-enhancing mutations, the resulting proportionality constant ($1/\text{total_fitness}$) changes, so that the mapping from ‘fitness’ to probability of reproduction in the next reproduction event changes over time. Nevertheless, this mapping always remains fitness-proportional.

On page 4, the term is used as follows: “the absolute rates f_k and f_1 are proportional to μ^{n+1} ”. Here, proportionality in the literal sense follows from the equations on page 20, when setting $\tilde{\mu} = N \mu$, so that the second factor becomes μ^{n+1} . We have included a clarifying sentence to address this in the derivations (SI1).

(4) It could be mentioned in the main text that the time between bursts (d) must not be too short in order for the effect to be substantial. I would think that the relevant timescale depends on how deleterious the initial mutation is.

Thank you for making this interesting and very relevant point. We have included a section (SI3) and Figure (Fig S4) in the supplement to investigate the dependence on d. In short, we find that effects are weaker for small inter-burst intervals. The sensitivity to the burst size is highest for inter-burst intervals that are sufficiently small so that the lineage of the first mutant has relevant probability of surviving long enough to experience multiple burst phases.

(5) Why not report that relative rate for Figure 2E as for 2D, as the former would seem to be more relevant to TSGs? And why was it assumed that the first inactivation is deleterious in the simulations in Figure 4 if the goal is to model TSGs?

Thank you for noting this. For how we revised the paper to better connect Figures 2 and 4, please see our comment (A1) above. In general, neither 2E nor 2D should serve as quantitative predictions for what effect size we should expect in real world data, but are rather curated illustrations of the general phenomenon that we describe: we chose high mutation rates and exaggerated fitness effects so that dynamics become visually tractable in small simulation examples.

For figure 4, assuming that the first inactivation is deleterious achieves that the branching process for the mutant lineage becomes subcritical, which keeps the simulation example simple and illustrative. For more comprehensive motivation of the approach in 4D, and especially the discussion of how fitness effects of different magnitudes may or may not be subject to the effects we describe depending on whether the population is in a phase of constant or growing population size, we refer the reader to our added section SI2, and the added discussion on pages 6 and 10.

(6) Figure 2, D and E. I'm not sure why heatmaps with height one were provided rather than simple plots over time. It is difficult, for example, to determine from a heatmap whether the increase is linear or the relative rates with and without punctuation.

Thank you for this comment. These are not heatmaps with height one, but rather for every column of pixels, different segments of that column correspond to different clones within that population. This approach is intended to convey the difference in dynamics between the results in Fig 2 and the analogous results for a branching process in Fig S1. In Fig 2, valley-crossings happen sequentially, with subsequent fixations of adapted mutants. In Fig S1, with a growing population size, multiple clones with different numbers of adaptations coexist. We have now adapted the caption of Fig 2 to clarify this point.

(7) Page 3: "High mutation rates are known to limit the rate of 1-step adaptations due to clonal interference." This is a bit misleading, as it makes it sound like increasing the mutation rate decreases the rate of one-step adaptations.

Thank you for alerting us to this poor phrasing. We have changed it in the revised version of the manuscript (P. 3).

(8) Page 4: "*proportional to μ^{n+1}* " Is "*proportional*" being used loosely for "*an increasing function of*"?

It is meant in the literal mathematical sense (see response to comment (3))

(9) Page 5, near bottom: "*at least two mutations across the population*". In the same genome?

We counted mutations irrespective of whether they emerged in the same genome, to remain analogous to the TCGA analyses for which we also do not have single cell-resolved information.

(10) Page 6: "*missense or nonsense mutation*". What about indels? If these are not affected by APOBEC, omitting them will exaggerate the effect of punctuation.

Thank you for pointing out that this focus on single nucleotide substitutions conveys an exaggerated image of the importance of this effect of APOBEC-driven mutagenesis. There are of course several other classes of (epi)genomic alterations (e.g. chromatin modifications, methylation changes, copy number changes) that we do not consider in this part of our analysis. APOBEC mutagenesis serves as an example of a temporally clustered mutation process, which we investigate in its domain of action.

We have added further discussion (P. 10-11) to convey that our empirical results merely constitute an investigation of whether empirical patterns are consistent with our hypothesis, but that the narrow focus on only SNVs, only TSGs, and only APOBEC mutagenesis does not allow for a general quantitative statement about the in-vivo relevance of the phenomena we describe.

(11) Page 6: "*normalized by the total number of single nucleotide substitutions*." It is difficult to know how to normalize correctly, but I might think that the square of the number of substitutions would be more appropriate. Perhaps the total numbers are close enough that it matters little.

Thank you for noting this. In the revised manuscript we have now expanded this passage in the text to more clearly convey our motivations for why we normalize by the total number of single nucleotide substitutions. While the likelihood for crossing a fitness valley with 2 mutations is indeed proportional to the square of the mutation rate, we do not directly observe mutation rates from our data. Rather, we observe the number of acquired single nucleotide substitutions for every tumor sample, but since tumors in our data differ in the time since initiation and therefore differ in the numbers of divisions their cells have undergone before being sequenced, we cannot directly infer mutation rates. One way to phrase our main result about valley-crossing is that temporally clustered mutation processes have an increased rate of successful valley-crossings per attempted valley crossing. Our TSG deactivation score is constructed to reflect this idea. The number of TSGs serves as a proxy for successful valley-crossings and the total mutation burden serves as a proxy for attempted valley-crossings.

To convey these points more clearly, we have rewritten the first paragraph in the Section "Proxies for valley crossing and for temporal clustering found in patient data" (P.6)

(12) Perhaps embed links to the COSMIC web pages for SBS2 and SBS13 in the text.

Thank you for this suggestion. We have embedded the links at the first mention of SBS2 and SBS13 in the text.

<https://doi.org/10.7554/eLife.108058.2.sa0>