

In search of nonlipogenic ABCA1 inducers (NLAI): precision coregulator TR-FRET identifies diverse signatures for LXR ligands

Megan S Laham, Martha Ackerman-Berrier, Fahmida Alam, Sarah Turner, Ganga Reddy Velma, Christopher Penton, Soumya Reddy Musku, Manan Rana, Senthil Kumar, Anandhan Annadurai, Maha Ibrahim Sulaiman, Nina Ma, Gregory RJ Thatcher 

Department of Chemistry & Biochemistry, Colleges of Science & Medicine, University of Arizona, Tucson, United States • Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States • Arizona Center for Drug Discovery, University of Arizona, Tucson, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

The study highlights development of a multiplex coregulator TR-FRET (CRT) assay that detects ligands with theoretical full agonist, partial agonist, antagonist, and inverse agonist signatures within the same chemical series. The findings are **valuable** and will have theoretical and practical implications in the subfield, with respect to guiding the design of non-lipogenic liver X receptor (LXR) agonists. The strength of the evidence is **solid**, whereby the methods, data, and analyses broadly support the claims with only minor weaknesses that can be dealt with through improvements in the data analysis and the discussion. This study will be of interest to experts working in the areas of pharmacology, medicinal chemistry, and drug discovery in Alzheimer's diseases and dementias.

<https://doi.org/10.7554/eLife.109146.1.sa2>

Abstract

APOE4, the major genetic risk factor for Alzheimer's disease (AD), and ABCA1, required for lipidation of APOE are gene products of the liver X receptor (LXR) receptor. LXR agonists have been validated in animal models as therapeutics for AD, atherosclerosis, and many other diseases. Clinical progress has been thwarted by unwanted hepatic lipogenesis. Structurally diverse LXR ligands were profiled in coregulator TR-FRET (CRT) assays analyzing ligand-induced coactivator recruitment, coactivator selectivity, corepressor dissociation, and LXR isoform selectivity. A multiplex CRT assay was developed to measure synchronous ligand-induced displacement of corepressor by coactivator. Potency for coactivator recruitment to LXR β correlated with induction of ABCA1 in human astrocytoma cells. Correlation with lipogenic activation of sterol response element (SRE) in hepatocarcinoma cells, was more complex. CRT response was diverse revealing ligands with theoretical full agonist, partial

agonist, antagonist, and inverse agonist signatures within the same chemical series, suggesting the scope for precision CRT to guide nonlipogenic LXR agonist design.

Introduction

Definitive diagnosis of Alzheimer's disease (AD) requires post-mortem hallmark neuropathology of amyloid- β (A β) plaques and neurofibrillary tangles of intraneuronal tau protein, in combination with characteristic in-life cognitive decline. Alzheimer's disease and related dementia (ADRD) defines a broader patient population, taking account of observations that: (i) hallmark pathology is not necessary and sufficient for progression to dementia; and (ii) at least half of AD patients manifest multiple forms of neuropathology at death. The original description, by Alois Alzheimer, of AD pathology in the brain of Auguste Deter, in addition to A β and tau, has been interpreted to include a third neuropathology: lipid granule accumulation in glial cells.¹ Although this pathology, associated with lipid droplets, has been relatively forgotten, several modern theories of ADRD pathogenesis, such as the "small particle HDL (high density lipoprotein) hypothesis"² propose a central role for cholesterol mobilization and lipoprotein transport. The 2020 Lancet Commission on Dementia stated that reducing the risk from diseases such as Type-2 diabetes (T2D) and obesity would have a significant impact on healthy aging with cognition intact.³ Subsequently, the 2024 Lancet Commission introduced LDL (low-density lipoprotein) cholesterol as the highest-contributing modifiable risk factor for dementia.^{4–5}

Lipid droplets contain esterified cholesterol, phospholipids, triglycerides and other debris from phagocytosis of neurons and are found in astrocytes and microglia as a normal component of lipid homeostasis.⁶ The processing of dying cells, a primary task of microglia, relies upon cholesterol metabolism and transport, which when microglia are over-whelmed requires recruitment of astrocytes to assist in phagocytosis, leading to astrogliosis.^{7–8} Under physiological conditions, lipid droplets undergo lipophagy via mitochondrial cytochrome P450 (CYP)-mediated oxidation of cholesterol to 24-hydroxycholesterol (24HC) and 27-hydroxycholesterol (27HC), metabolites that cross the blood-brain barrier (BBB). Normal cholesterol metabolism and transport utilizes peripheral reverse cholesterol transport (RCT) for clearance to the liver, where cholesterol and metabolites are converted to bile acids. However, aberrant accumulation of lipid droplets in microglia and astrocytes leads to reduced lipophagy, accumulation of cholesterol and reduced production of 27HC. The consequence is elevated cholesterol levels in all cellular membranes, including the plasma membrane where lipid raft malfunction leads to disruption of synapses and multiple signaling pathways. Lipophagy and clearance of excess cholesterol via RCT requires efflux to lipoproteins, which is orchestrated by the nuclear hormone receptor (NR) liver-X-receptor (LXR) and its gene products, including *APOE* (apolipoprotein-E) and *ABCA1* (ATP-binding cassette-A1).⁹ *APOE4* is the major genetic risk factor for ADRD and is associated with aberrant lipid accumulation, notably in microglia.^{10–11}

ABCA1 is expressed ubiquitously throughout the human body, with levels in the periphery highest in liver hepatocytes.¹² The 2,261 amino acid human *ABCA1* protein is an integral membrane protein consuming ATP to transport cholesterol, phospholipids, and other lipids to apolipoprotein carriers that bind to *ABCA1*.^{13–15} *ABCA1* associates with cholesterol-rich membrane domains in the plasma membrane and intracellular organelle membranes, including endosomes and lysosomes.¹⁶ *ABCA1* functions, in part, to protect cells from excessive and potentially cytotoxic accumulation of free cholesterol. *ABCA1* transports lipids to specialized apolipoproteins, APOA1, APOA2, APOA4, APOC1-3, and APOE, to assemble HDL particles.¹⁷ *ABCA1* is uniquely important in early loading of apolipoproteins in the minimally lipidated state,¹⁸ in contrast to related ABC family members (e.g. ABCG1/G4) and cholesterol transporters (e.g. SR-B1) that interact with apolipoproteins in their more lipidated states.^{16, 19} Owing to the direct influence of *ABCA1* on cellular cholesterol homeostasis, increased *ABCA1* activity has many indirect physiological

functions. Taken together, the evidence supports ABCA1 as a promising target for therapeutics that might broadly attenuate pathophysiologic processes in the periphery and in the central nervous system (CNS). In the brain, ABCA1 is found in neurons, astrocytes, and microglia.²⁰

ABCA1 associates with cholesterol-rich membrane domains, where it protects cells from excessive, and potentially cytotoxic, cholesterol and lipid accumulation.^{13–16} LDL particles contribute to vascular cholesterol deposition, inflammation, and atherogenesis to drive atherosclerosis.^{21–23} Conversely, HDL particles are anti-atherogenic. HDL facilitates removal of cholesterol from vascular walls and peripheral tissues to the liver for catabolism and excretion, via RCT. HDL, thereby prevents formation of cytotoxic oxidized lipid species and reduces inflammation and atherogenic lesion formation.²⁴ ABCA1 is essential for formation of nascent small HDL particles^{14, 19, 25}, and with ABCG1, promotes formation of mature HDL particles.²⁶

Reduced plasma cholesterol efflux capacity, which is directly tied to ABCA1 and ABCG1 expression, synergizes with other components of metabolic syndrome to promote atherogenesis.²⁷ Oxidation of LDL and other pathological changes that occur during atherogenesis, may down-regulate ABCA1 expression.^{28–29} Tangier disease, characterized by *ABCA1* loss-of-function mutations, leads to premature atherosclerosis;³⁰ furthermore, *ABCA1* variants that increase AD risk have also been associated with T2D and cardiovascular disease (CVD) risk.^{31–33} Peripheral atheroprotection via ABCA1 induction promotes cerebrovascular health and preserves BBB integrity and function.^{34–35} Therefore, the beneficial role of ABCA1 in CNS disorders may derive from central actions and peripheral effects.^{36–37}

ABCA1 and other genes involved in cholesterol metabolism, mobilization, and signaling are gene products of LXR. Therapeutic approaches to enhance ABCA1 have been reviewed.^{38–40} LXR is the primary NR target for pharmacologic induction of ABCA1 expression. LXR agonists have been studied in many disease models, with the most intensive efforts targeting a therapeutic strategy to treat atherosclerosis and hypercholesterolemia.^{41–45}

The two NR isoforms, LXRα (NR1H3) and LXRβ (NR1H2) mediate gene transcription via transcriptional complexes that bind to specific or shared NR response elements on DNA. LXRα is expressed in liver, small intestine, and adipose, and LXRβ is ubiquitously expressed. LXR forms heterodimers with retinoid X receptor (RXR). Transcriptional complexes are formed by recruitment of coregulator proteins that provide scaffolds for binding of epigenetic tools for DNA and histone writing, erasing, and reading. Since coregulators are shared between NRs, competition for coregulators is one of the many elements allowing crosstalk between different NRs. The canonical mechanism of lig- and activation is the binding of an agonist to the NR dimer, in this case the LXR:RXR heterodimer, leading to coactivator recruitment to the transcriptional complex. In an alternate mechanism, transcriptional repression is maintained by corepressor binding to the unliganded heterodimer (**Fig. 1A**), with ligand binding leading to activation/derepression (**Fig. 1B**). The endogenous ligands for LXR are oxysterols (including 22(*R*)-hydroxycholesterol, 24HC, and 27HC)^{9, 46–47} that activate LXR via canonical or alternate mechanisms.

LXR controls several transcriptional programs one of which, mediated by the master hepatic regulator sterol-response element binding protein 1c (SREBP1c) leads to unwanted lipogenesis, triglyceride (TG) synthesis, and elevated LDL (**Fig. 1C**). Adverse effects associated with elevated levels of TG, include steatohepatitis, and hepatomegaly.⁷ The challenge in developing LXR agonists as therapeutics for T2D, CVD, and ADRD has been to avoid the adverse activation of the lipogenic axis (**Fig. 1C**). The dominant approach has been to develop agonists selective for LXRβ over LXRα.⁸ The rationale for developing LXRβ-selective agonists is based on the high expression of LXRα in the liver; however, despite considerable efforts by pharmaceutical companies, selective LXRβ-selective agonists have yet to be successfully translated to the clinic.^{48–49}




Fig. 1.

LXR activation mediates beneficial effects and unwanted lipogenesis.

LXR transcriptional complexes are RXR heterodimers that can be activated by a canonical mechanism of coregulator recruitment or de-repression of (A) a constitutively repressed complex by (B) corepressor displacement. (C) ABCA1 gene transcription leads to cholesterol efflux and mobilization, APOE lipidation and potentially A β clearance, in addition to other mechanisms of potential benefit in ADRD; whereas, SREBF1 transcription initiates a lipogenic program in the liver leading to unwanted LDL formation and TG elevation.

A recent paper from Astra Zeneca researchers (Belorusova et al. ⁵⁰) specifically addressed the challenge in discovery of LXR ligands as nonlipogenic ABCA1 inducers to treat atherosclerosis. Mouse intestinal ABCA1 and plasma TG were used as binary ligand descriptors that were correlated with LXR readouts, most notably H/D-exchange mass spectrometry (HDX-MS) measurements that inferred ligand-induced conformational changes in the LXR ligand binding domain (LBD).⁵⁰ Coregulator TR-FRET (CRT) measurements of coregulator binding to the LBD have been widely used in the NR field.⁵¹ A detailed and comprehensive CRT study of LXR ligands, both clinical and late preclinical, has not been reported.⁵² CRT data has been reported for individual ligands; for example, to claim that LXR agonist IMB-808 was nonlipogenic because of selective recruitment of specific coactivators.^{53–54} Since repression of lipogenesis in the liver requires corepressor binding to LXR α , corepressors must be included in CRT studies.

We profiled diverse LXR ligands by CRT, including those studied in clinical trials: (i) to assess NR promiscuity by measuring coactivator recruitment to 9 steroid NRs; (ii) to measure LXR α and LXR β selectivity by measuring SRC1 coactivator recruitment; (iii) to measure differential coactivator recruitment; (iv) to delineate the effect of putative antagonists, partial agonists, and inverse agonists; and (v) to define corepressor binding in the presence and absence of a coactivator using multiplexed precision CRT (pCRT).

CRT data was correlated with data from reporter assays in astrocytoma and hepatocarcinoma cells, measuring induction of ABCA1 and SREBP1c, respectively. The results from CRT analysis reveal a surprisingly diverse set of responses that would support diverse phenotypes. NRs are viewed as druggable targets given that the efficacy of 16% of approved drugs is mediated by NRs.⁵⁵ The clinical success of selective estrogen receptor modulators (SERMs) is based upon tissueselective transcriptional activity that is not predictable *a priori* but is thought to be governed by coregulator recruitment.⁵⁶ More precise understanding of ligand-specific coactivator and corepressor binding, will provide a predictive framework for nonlipogenic LXR modulators and reinvestigate the therapeutic targeting of NRs by small molecules.


Results and Discussion

Selection of benchmark LXR ligands

First reported in 2000⁵⁷, T0901317 (T0) is commonly used as a benchmark or control LXR agonist. Cross-reactivity of LXR agonists with other NRs has been reported.⁵⁸ T0 has been claimed to be an agonist for farnesoid X receptor (FXR) and pregnane X receptor (PXR).^{59–61} Although, in our CRT assays potent PXR agonism was observed without evidence for FXR binding. The cross-reactivity of T0 with PXR could confound its use as a control LXR agonist in cellular systems. However, T0 is unique in showing no selectivity between LXR isoforms, which makes it ideal for normalization of CRT data. T0 is the most widely studied LXR agonist, often as a control compound or LXR chemical probe, with dozens of papers reporting effects in preclinical animal models of disease, including AD.^{62–65} T0 is reliably lipogenic in mouse, hamster, and other models, causing elevated TGs and hepatic steatosis.

A different chemotype was identified using a cell-based ABCA1 reporter system, leading to GSK3987, an LXR agonist that in cell cultures showed concentration-dependent increases in ABCA1, SREBP1c, and cholesterol efflux and concentration-dependent decreases in IL-6.⁶⁶ There is a lack of *in vivo* data, which may explain why this compound has not been used as a chemical probe.

GW3965, first reported in 2002, is widely used as an orally bioavailable selective LXR agonist in preclinical animal models.^{67–68} Although the earliest report on GW3965 indicated PXR agonist activity, no binding to PXR is seen in CRT (*vide infra*). In C57Bl/6 mice, the effects of T0 and



Scheme 1

Chemical structures and compound codes of literature LXR ligands with biphenyl sulfone ligands shown in blue.

GW3965 were compared after 3 days treatment, measuring plasma lipids and gene induction in the liver and small intestine.⁶⁸ Responses were highly dose-dependent; however, T0 caused increases in liver weight and TGs, whereas even at 100mg/kg/day, GW3965 had no effect on these liver markers. T0 induced SREBP1c and greatly induced fatty acid synthase (FAS) (38-fold) while GW3965 induced minimal increases in SREBP1c and FAS expression in the liver. GW3965 administration resulted in ABCA1 expression increased 8-fold in the small intestine and 7-fold in peripheral macrophages. Recent dramatic evidence of the benefits of ABCA1 induction by GW3965 was reported in a tauopathy mouse model relevant to AD.⁶⁹ The beneficial effects of low dose GW3965 were replicated by genetic upregulation of *ABCA1*.

The LXR agonist, RGX104 (Abequolixron or SB742881), described as a first-in-class immunotherapy is being studied in cancer clinical trials. RGX104 is the methyl derivative of GW3965 and was compared to GW3965 in hamsters.⁷⁰ Hamsters, unlike mice, express cholesteryl ester transfer protein (CETP) and are argued to have a more humanlike lipid metabolism. Both GW3965 and RGX104 caused a dose-dependent increase in plasma TG, LDL-C, and very-low-density-lipoprotein cholesterol (VLDL-C). Despite structural similarity, RGX104 significantly increased liver triglycerides but GW3965 did not. Similarly, hepatic gene transcription was weaker for GW3965, although comparable in intestines and peritoneal macrophages. In cynomolgus monkeys, GW3965 increased LDL and HDL at 7 days treatment only, with no effect on TG; whereas, RGX104 gave a robust time-dependent increase in TG and LDL. Without overinterpretation, this study hints at possible tissue-selective actions differentiating GW3965 from RGX104: tissue selectivity is well-known for modulators of other NRs.

Cholesterol mobilization was compared in HepG2 cells for RGX104, GW3965 and T0.⁷¹ In these cell cultures, all LXR agonists gave similar induction of ABCA1, ABCG1, FAS, and SREBP1c, which was argued to result in serum-dependent cholesterol efflux with induction of compensatory cholesterol synthesis at a level still leading to cellular cholesterol depletion. Interestingly, steroidal agonists, 22(R)-hydroxycholesterol, 24(S),25-epoxycholesterol, and DMHCA (N,N-dimethyl-3 β -hydroxy-cholenamide) did not induce the same response.

The synthetic steroidal LXR agonist, DMHCA, was reported to increase LXR α , ABCA1 and SREBP1c genes in HepG2 and THP-1 cells to a lesser extent than T0 but to decrease SREBP1c in the J774 macrophage cell line and not significantly to elevate FAS in THP-1 cells.⁷² In THP-1 cells, DMHCA attenuated SREBP1c gene transcription elicited by GW3965 co-treatment, implying partial agonist activity. C57BL/6 mice treated for 6 or 7 days with T0, GW3965, or DMHCA showed increased hepatic gene expression of ABCA1, ABCG1, SREBP1c, FAS, and acetyl-CoA carboxylase (ACC) for all agonists. In mice, T0 and GW3965 were reported to increase plasma TG and VLDL-C. DMHCA is metabolically labile with poor oral bioavailability⁷²; nevertheless, low dose oral delivery has been studied in the ob/ob mouse in the context of diabetic retinopathy and bone marrow pathology.⁷³

Like DMHCA, LXR-623 (WAY-252623) is referred to as an LXR partial agonist.⁷⁴ Comparison of LXR-623 with GW3965 in LDLR^{-/-} mice confirmed the expected anti-atherosclerotic efficacy of both LXR agonists.⁷⁵ In hamsters, LXR-623 did not cause a dose-dependent increase in plasma TG. In primates a significant increase in liver TG was observed but only at high dose (50 mg/kg/day p.o.) and returned to baseline at day 7 of treatment. Tissues from primates showed elevation of *ABCA1* and *ABCG1* in duodenum and suppression of hepatic genes including *SREBF1*, *FAS*, *CYP7A1*, and *LDLR*. The authors concluded that unfavorable lipogenesis was not a class effect of LXR agonists. In the only report on the LXR-623 clinical trial, no data on lipogenesis is reported. Detailed PK/PD correlations were made with ABCA1 transcripts in blood samples.⁷⁶ No serious adverse events were reported; however, all 6 participants on the highest 300 mg dose reported neurologic or psychiatric adverse effects of unknown cause.

In contrast to LXR-623, the clinical trial of XL041 (BMS85297) reported on both activation of ABCA1 and lipogenic genes. As seen for LXR-623, treatment with XL041 increased ABCA1 expression in trial subjects.^{48–49} XL041 demonstrated good pharmacokinetics, safety, and lipid profiles in mice and cynomolgus monkeys.⁴⁹ However, in clinical trials, after multi-day dosing, elevated plasma and liver lipids and neutropenia were observed in healthy and statin-treated subjects. Notably co-administration of statins caused incomplete attenuation of lipid elevation.^{49, 77} The peripheral side effects observed with XL041, included neutropenia and elevated TG and LDL levels, both of which were reversible. No serious adverse effects were reported.

Published data from Merck's LXR program is limited. *Compound 9* (MK9) was studied in 12-month-old Tg2576 familial Alzheimer's disease (FAD) mice dosed subcutaneously for 3 weeks at 50 mg/kg/day.⁷⁸ Significant increases in ABCA1 and apoE protein levels in the brain were not accompanied by significant changes in plasma or liver TG, in contrast to mice treated with T0. In monkeys administered MK9 (20 mg/kg qd p.o.) for 2 weeks, in cerebrospinal fluid (CSF) a threefold increase in APOE and A β was observed: CSF A β is often associated with enhanced clearance from the brain. These changes were accompanied by no change in liver fat content relative to control, in contrast to T0.

AZ876, classified by Belorusova et al. as lipogenic in mice,⁵⁰ showed dose-dependent elevation of plasma TG accompanied by a 91% reduction in atherosclerotic lesion area.⁷⁹ Rescue of cardiac hypertrophy in mice using low dose AZ876 was not accompanied by increases in plasma TG.⁸⁰

CL2-57 was the result of early optimization of a hit discovered from cell-based reporter screening for activation of ABCA1-luciferase in astrocytoma cells, counterscreening using a SREBP1c-luciferase reporter in HepG2 cells.⁸¹ In an obesogenic high-fat diet (HFD) mouse model, CL2-57 corrected T2D-associated readouts, attenuated diet-induced increases in plasma and hepatic TG levels and ameliorated other HFD-associated metabolomic changes.⁸² Further optimization guided by phenotypic screening resulted in CL3-3, a more potent LXR agonist and ABCA1 inducer. CL3-3 was tested in E3/4FAD mice that express one copy each of the ϵ -3 and ϵ -4 alleles of human APOE, mimicking the majority of AD patients. These mice also express five mutant amyloid precursor protein (APP) and PSEN1 human transgenes linked to FAD. Treatment with CL3-3 increased ABCA1 expression, enhanced APOE lipidation and reversed multiple AD phenotypes, without increasing TG. This study is the first in a human APOE-expressing model with hallmark amyloid- β pathology ALX101 (Rovazolac), SR9238, GSK2033, XL652, and BE1218 are examples of the many reported LXR ligands in the biphenyl-3-methylsulfone class, which also includes XL041. Reported in a 2013 patent publication, ALX101 is listed as the subject of a 2018 clinical trial in moderate atopic dermatitis. XL652, an LXR agonist studied in the program that led to XL041, is described as an LXR β -selective partial agonist.^{83–84} In mice XL652 administration caused increased ABCA1 and ABCG1 induction with elevated plasma triglycerides only at the highest dose (30 mg/kg qd), accompanied by induction of hepatic SREBP1c but not FAS.

The structurally related sulfones, BE1218, SR9238, SR9243, and GSK2033 have all been described as LXR inverse agonists. SR9238 was claimed amongst the first synthetic LXR inverse agonists to be discovered. SR9243 is systemically available; whereas, the metabolically labile SR9238, owing to the metabolism of the ester linker, is reported to show liver selectivity for suppression of lipogenic LXR gene products.^{85–87} In obese mice on HFD, plasma TG was suppressed and in a mouse model of non-alcoholic steatohepatitis (NASH), SR9238 treatment significantly reduced hepatic inflammation and fibrosis. In cell-based models, the activity of GSK2033 was consistent with LXR inverse agonism; however, in a NASH mouse model, the expression of lipogenic genes, including FAS and SREBP1c, was *increased* rather than suppressed and no beneficial effects on hepatic steatosis were observed. Subsequently, GSK2033 was reported to activate PXR, FXR, and RXR, negating utility as an in vivo chemical probe.

IMB-808 is structurally distinct from most other LXR ligands. In cell-based assays, it was claimed to activate LXR with modest selectivity for the α -isoform and to be nonlipogenic in HepG2 cells, which is the rationale for inclusion of this compound in our study.⁸⁸ The selection of other LXR ligands, introduced above, provides a diverse array of structures, selectivity and both lipogenic and nonlipogenic properties. Although most compounds discussed have demonstrated beneficial responses in preclinical models of human disease, the lack of a predictive algorithm for ligand design to avoid or circumvent unwanted lipogenesis has prevented progress in the clinic.

Agonists, antagonists and NR coregulator TR-FRET (CRT)

The ligand binding pocket of NRs can be considered as both orthosteric and allosteric, because it binds endogenous agonists and induces a conformational change that induces recruitment of coregulator proteins to helix-12 (H12) of the NR and hence regulates the transcriptional function of the NR complex at its cognate DNA response element. The labels applied to individual NR ligands (full agonist, partial agonist, antagonist, inverse agonist) can be ambiguous leading to misuse. Owing to the tissue and cell-specific output resulting from ligand binding, individual genes may be regulated in different directions.

In the case of estrogen receptor (ER) ligands, the term “pure antagonist” has been coined and is usually, but not universally, used to describe a ligand that competitively displaces estradiol and induces a conformational change incompatible with transcriptional activation.^{89–90} In the case of agonists (ligands that induce a response in the same direction as the endogenous agonist), the term “biased agonist” is useful for NR ligands and is related directly to tissue selectivity.⁹¹ Study of coregulator binding to the NR-LBD can potentially simplify the terminology used for ligand-dependent responses, which we will revisit below. However, first we used CRT to study cross-reactivity of LXR ligand binding to related NRs.

CRT analysis of NR cross-reactivity

Class II NRs include: the NR1 subfamily, LXR, farnesoid X receptor (FXR), peroxisome proliferator activated receptor (PPAR), pregnane X receptor (PXR), retinoic acid receptor (RAR), constitutive androstane receptor (CAR); and the NR2 subfamily retinoid X receptor (RXR). Heterodimer formation of RXR with NR1 family members is a key step in formation of transcriptional complexes. Most LXR DNA binding sites are shared with RXR; however, only a fraction of RXR binding sites are shared with LXR and genome-wide binding profiles are quite different across cell and tissue types.⁹²

Owing both to the permissive nature of the heterodimer complexes within the sterol NR family and to structural similarities between endogenous ligands and ligand binding pockets, LXR ligands may activate other NR complexes. Cross-reactivity of LXR agonists has been reported:⁵⁸ T0 is claimed to be an FXR agonist;⁵⁹ T0 and GW3965 are reported to be PXR agonists,^{60–61} and GSK2033 binds to FXR, PXR, CAR, RXR, and other NR LBDs in a cell-based assay.⁹³ Therefore, all LXR ligands were examined in a Lanthascreen CRT assay measuring ligand-induced stabilization of the NR:coactivator complex and NR-specific ligand binding. All CRT screens of NR selectivity were performed with a terbium donor exciting a fluorescein-tagged coactivator as a measure of ligand binding and recruitment. Benchmarking used a known agonist for each NR.

NR-CRT assays were developed for RXR α , RXR β , RAR α , RAR β , FXR, PXR, and PPAR δ , in addition to LXR α and LXR β (Table S1). Assays for RXR α and RXR β were optimized using the D22 coactivator peptide and the RXR ligands SR11237 and bexarotene as positive controls. Both benchmark RXR ligands were RXR α -selective full agonists with very similar potency and no response was observed for any of the tested LXR ligands (Fig. S1A).

FXR is critical for maintaining bile acid homeostasis that when disrupted leads to excessive accumulation of cholesterol and its actions are interleaved with those of LXR. FXR represses the expression of SREBP1c.⁹⁴ LXR promotes fatty acid and triglyceride synthesis and storage, whereas FXR activation leads to decreased liver and plasma triglyceride levels.⁹⁵ The CRT assay optimized for FXR used SRC2-2 as coactivator and obeticholic acid (OCA) as benchmark agonist, revealing no agonist activity by LXR ligands above baseline, despite description of T0 as a weak FXR agonist (Fig. S1B).^{59–60}

PPAR isoforms have been reported to modulate ABCA1 expression levels.^{96–98} PPARγ activity is closely associated with glucose metabolism, and PPARα and PPARδ activity is associated with fatty acid metabolism. PPARα and PPARγ agonists have been shown to activate ABCA1, improve TG levels, and have shown beneficial effects in FAD mouse models.⁹⁹ PPARδ-selective agonists have also been reported to increase ABCA1,¹⁰⁰ with examples in AD clinical trials.¹⁰¹ The CRT assay for PPARδ used a C33 peptide coactivator and GW501516 as positive control, with no activity observed for any LXR ligands tested (Fig. S1C).

Of the three isoforms, RARβ is the most widely expressed across tissues: brain, intestine, liver, kidney, etc.^{102–104} The RARα CRT was optimized with D22 coactivator peptide and RARβ with SRC2-2 coactivator peptide. TTNPB was used as the benchmark agonist for both RAR isoforms tested, possessing very similar and high potency for both isoforms. Maximal response was seen at 1 μM of TTNPB, a concentration at which no LXR ligand registered any response (Fig. S1D).

PXR is a xenobiotic nuclear receptor, responsible for upregulating metabolic enzymes for detoxification of exogenous and endogenous compounds and is highly expressed in the liver, intestine, and kidneys.¹⁰⁵ The PXR LBD is larger than other sterol NRs and the hydrophobic ligand binding pocket reflects its function in accommodating a wide range of xenobiotics.¹⁰⁶ Drugs that bind and activate PXR bear a risk of drug-drug interactions (DDIs).¹⁰⁷ The bisphosphonate, SR12813, was used as a benchmark PXR agonist; however, T0 proved to be a full agonist and more potent (EC_{50} = 300 nM) than the benchmark (Fig. S1E).

Activity and isoform selectivity of LXR ligands

The CRT selectivity assay incorporates GST-tagged LXR-LBD, anti-GST-terbium, biotin-SRC-1, and streptavidin-XL665. Notably, the 25 amino acid SRC-1 peptide is the only coactivator tested for LXR binding that has the fluorophore remote from the coactivator peptide. Binding of an agonist to LXR induces recruitment of coactivator SRC-1 (steroid receptor coactivator 1; also known as nuclear receptor coactivator 1, NCOA1). When excited at 340 nm the terbium will emit at 620 nm exciting XL665 and emitting a signal reflecting coregulator binding to LXR. The assay measures the EC_{50} for coactivator recruitment, a measure of ligand binding affinity, and the strength of signal reflects maximal efficacy. For sterol receptors, endogenous ligands, including 24HC, generally induce efficacy lower than synthetic ligands; thus, for LXR binding, T0 was used as a suitable control (Fig. 2).

Perhaps surprisingly, considering the description of multiple LXR ligands as partial agonists, most agonists studied gave maximal response at the same level as T0, behaving as full agonists. The clear exception is the endogenous partial agonist, 24HC; although, both the synthetic steroid DMHCA and CL3-3 showed partial agonist activity at LXRα. Both SR9238 and GSK2033 antagonized the agonism of 24HC at LXRβ without effect on LXRα.

In contrast to efficacy, potency and selectivity were observed to vary widely. As expected, T0 was nonselective as were AZ876, DMHCA, and indeed 24HC. GSK3987 was weakly selective for LXRβ; whereas, most agonists displayed 10–100 fold selectivity for LXRβ over LXRα, with MK9 and BE1218 being the most selective LXRβ agonists. BE1218 was also one of the most potent LXRβ agonists; with the most potent being AZ876 (6 nM) and XL041 (10 nM). The most potent LXRα agonists were the pan-LXR agonists AZ876 (6 nM) and T0 (80 nM).




Fig. 2.

LXR isoform selectivity from CRT.

Concentration-response for recruitment of SRC-1 to LXR α (dotted lines) and LXR β (solid line), determined by CRT assay, showing mean and SD from triplicate experiments, normalized to T0 (100%) as a full LXR agonist (shown in turquoise for reference). GSK2033 (R) and SR9238 (S) titrations were run in the presence of 24HC (3 μ M) and neither ligand stabilized the LXR:SRC1 complex alone (T,U).

The ligand binding pockets of the two LXR isoforms differ by only one amino acid (LXR α -Val263 and LXR β -Ile277). Interestingly, mutation of these residues to alanine (V263A and I277A) does not ablate isoform selectivity in reporter assays.¹⁰⁸ Conformational cooperativity of LBD residues beyond these two amino acids leads to different conformations of Leu274 and Ala275 that generally favor ligand binding to LXR β . Therefore, most reported ligands, and all ligands examined in the current work, are LXR β -selective or non-selective.

ABCA1 induction correlated with LXR β CRT binding

Astrocytoma CCF-STTG1 cells, stably transfected with an ABCA1-promoter linked to a luciferase response element provide a model for ABCA1 induction in astrocytes.⁸¹ In contrast to the SRC1 CRT binding data, showing little evidence for partial agonist activity, concentration-response curves in cell-based reporter assays showed apparent partial agonist and antagonist activity (**Fig. 3**). The biphenyl sulfones, BE1218 and XL041, showed partial agonist response. The antagonist activity of GSK2033 and SR9238 in LXR β :SRC1 CRT was reflected by inverse agonist activity in cells. Apparent inverse agonist activity in cell-based assays can result from true inverse agonist recruitment of corepressors or from antagonist activity towards endogenous sterol LXR agonists. Although we use serum stripped media, antagonism of endogenous LXR agonists cannot be ruled out. As introduced above, in the literature, BE1218, SR9238, SR9243, and GSK2033 have been described as LXR inverse agonists; however, BE1218 acted as a full agonist in recruitment of SRC1 and a partial agonist of cellular ABCA1 induction.

The correlation of potency for cellular ABCA1 induction versus stabilization of the LXR β :SRC1 coactivator complex was excellent with $R^2 = 0.95$ (**Fig. 4B**). The correlations of a) ABCA1 induction with SRC1 recruitment to LXR α (**Fig. S3**) and b) SRC1 recruitment to LXR β versus LXR α (**Fig. 4A**) were relatively poor. To the best of our knowledge, and surprisingly, this is the first such quantitative ABCA1/LXR β correlation across a varied structural array of LXR agonists and supports the targeting of LXR β as a mechanism for induction of ABCA1 and related genes in cholesterol mobilization. The corollary is that LXR α holds the key to avoiding lipogenesis.

Lipogenic SRE activation correlated with LXR α CRT binding

Our phenotypic high throughput screen (HTS) approach to identify nonlipogenic ABCA1 Inducers (NLAI) originally used counterscreening with a SREBP1c-luciferase reporter (linked to the *SREBF1* gene promoter) in human hepatocarcinoma HepG2 cells.⁸¹ Unwanted lipogenesis is mediated by LXR agonists via direct transcriptional activation and indirectly by the effects of SREBP1c.^{57, 109} SREBP1c enhances transcription of genes required for synthesis of fatty acids and TG and as the dominant isoform in liver is associated most closely with lipogenesis and TG elevation.¹⁰⁹ SREBP1c is inactive and ER-resident, requiring transport by SREBP cleavage-activating protein (SCAP) to the Golgi membrane and proteolysis by S1P and S2P to release the mature SREBP1c transcription factor (**Fig. 1C**). Binding of SREBP1c to the sterol response element (SRE) induces transcription of lipogenic genes associated with elevated TG. To better model the lipogenic axis in hepatocytes, we use HepG2 cells stably transfected with a luciferase reporter downstream of a SRE promoter (SRE-luc), activation of which requires SREBP1c maturation.

As observed with induction of ABCA1, concentration-response curves for SRE activation by LXR ligands showed apparent partial and full agonist responses (**Fig. 5**). Three of the biphenyl sulfones, XL041, XL652, and SR9238, gave anomalous activation of SRE at concentrations above 1 μ M (**Fig. S4**). At concentrations ≤ 1 μ M, XL652 and XL041 activated SRE with low efficacy (20–25% of T0) and SR9238 lowered SRE activation below baseline. Owing to the anomalous response in the reporter assay, we measured lipogenic genes, *SREBF1*, *FAS*, and *SCD1* by RT-PCR in HepG2 cells in response to treatment with the biphenyl sulfones (**Figs 5G–I**). The PCR data was internally consistent, with GSK2033 and SR9238 reducing lipogenic gene transcription below baseline,

Fig. 3.

ABCA1-luc reporter in CCF cells.

Concentration-response for LXR ligands in astrocytoma cells measured after 24 hr incubation, normalized to T0 response (100%) showing mean and SD from triplicate experiments (see **Fig. S2** for biological replicates).





Fig. 4.

Correlating potency in cell-free and cell-based assays.

A) Coactivator stabilization by LXR ligands correlated for α and β isoforms showing the general bias towards LXR β selectivity (above the line); T0 and AZ876 lie on the 1:1 correlation line. **B)** Potency for ABCA1 induction in cells shows an excellent correlation for cell-free LXR β binding and coactivator stabilization. **C)** The potency correlation between SRE activation and cell-free LXR α binding is weaker but is still observed. **D)** The correlation between potency for ABCA1 induction versus SRE activation in cell-based assays is weak. Data show mean from at least triplicate observations. (See **Fig. S3** for other correlations).






Fig. 5.

Lipogenic response of HepG2 cells to treatment with LXR ligands.

(A-D) Concentration-response curves for LXR ligands normalized to maximal response for T0 ($EC_{50} = 15$ nM) showing a range of potency (AZ876 $EC_{50} = 4.2$ nM to CL2-57 2.8 μ M) and maximal efficacy (BE-1218 80% to GSK3987 150%). See [Fig. S2](#) for biological replicates. (E-F) Three biphenyl sulfones gave anomalous activation of SRE at higher concentrations ([Fig. S4](#)): XL041 gave a partial agonist response at submicromolar concentrations ($EC_{50} = 5.7$ nM $E_{max} = 22\%$), while SR9238 and GSK2033 had a neutral or antagonist response. (G-I) The transcriptional response to the five biphenyl sulfones was studied by RT-PCR under the same conditions as for SRE-luc measurements replicating the lipogenic induction observed in the reporter assays; both GSK2033 and SR9238 reduced response below baseline. Both DMHCA and 24HC were cytotoxic at higher concentrations. Data show mean and standard deviation for triplicate measurements. See [Fig. S4](#) for full concentration-response curves.

Simplistically, activation of the lipogenic axis (SRE activation) might be expected to correlate with LXR α :SRC1 stabilization; however, this correlation was weaker than that observed between ABCA1 induction and LXR β :SRC1 stabilization (**Fig. 4C**). The correlation between SRE activation and ABCA1 induction was weak (**Fig. 4D**). The correlation analyses used potency measurements both for SRC1 recruitment and for activation of luciferase reporters, which does not incorporate a contribution from the different efficacy seen for partial agonists. On the basis of potency alone: (i) cellular ABCA1 induction can be dissociated from induction of lipogenic genes; and (ii) ABCA1 induction can be predicted by LXR β recruitment of SRC1 measured by CRT. Data analysis to this point does not provide a framework for prediction of lipogenic potency or efficacy. Further, the lipogenic response in cell cultures is more complex and diverse than ABCA1 induction, which correlates with LXR β agonism (**Fig. 4B**).

Coactivator selectivity

The varied physiological functions of a particular NR are controlled by recruitment of specific coregulators.^{110–111} Coactivator families include the p160 family (SRC-1 (NCoA-1), SRC-2 (GRIP1, TIF2, or NCoA-2), SRC-3 (p/CIP, RAC3, ACTR, AIB1, or TRAM-1)), PPAR γ coactivator 1 (PGC-1 α , PGC-1 β), and TRAP/DRIP complex. Thyroid hormone receptor-associated proteins (TRAPs) were first studied in the context of the thyroid hormone receptor.¹¹² A highly conserved LXXLL motif in the p160 family of coactivators is necessary for association of coactivators to the ligand-bound receptor.¹¹³ Two subunits of the TRAP complex, TRAP220 and TRAP100, contain LXXLL domains.

Selective recruitment of coactivators to LXR may provide a rationale for putative ligand selectivity for nonlipogenic gene induction. The lack of recruitment to LXR β of D22 and TRAP220 was proposed to underlie nonlipogenic actions of IMB-808. Kim et al. explored the mechanism of LXR-mediated transcription of metabolic genes with a focus on coactivator complexes that lead to the modulation of specific target gene expression.⁵⁴ Two coactivators were identified as proteins that interact with LXR α in a direct and ligand-dependent manner: TRAP220 and TRAP80. Examination of activation by T0 and GW3965 indicated that TRAP220 stimulates promoters for SREBP1c and ABCA1, whereas TRAP80 was selectively lipogenic. Silencing TRAP80 in mice followed by treatment with GW3965 or T0 was reported to prevent hepatic steatosis and reduce hepatic TG levels; whereas, RCT-related genes and anti-inflammatory effects of the LXR ligands were unaffected. This observation provides a potential mechanism for selective repression of SREBP1c-mediated lipogenesis caused by LXR α .

CRT analysis of coactivator recruitment to LXR of PGC1 α , SRC2-2, D22, and TRAP220 demonstrated that potency for LXR agonists was most highly correlated between SRC2-2/D22 and PGC1 α /TRAP220 pairs (**Fig. S5**). Therefore, for clarity, only D22 and TRAP220 recruitment are shown for ligands other than T0 (**Fig. 6**). Concentration-response curves for LXR agonists were normalized to maximal response for T0. Across the four coactivators studied, T0 was nonselective both for LXR isoform and for coactivator identity; whereas, most other agonists displayed both full and partial agonist profiles in contrast to our observations on SRC1 recruitment. Thus, across the four coactivators studied, most ligands showed LXR β selectivity based on maximal efficacy, notably MK9, CL3-3, and LXR-623. Regarding LXR isoform selectivity, AZ876, does not show isoform selectivity for SRC1 recruitment but selectively recruits SRC2-2, D22, PGC1 α and TRAP220 to LXR α . DMHCA is nonselective across all coactivators; whereas, all other LXR agonists are selective for LXR β across all coactivators studied. GSK2033 and SR9238 are unable to induce an LXR conformation compatible with recruitment of any coactivator studied. Several ligands induced very minimal recruitment of coactivator TRAP220 to LXR α , including XL041, XL652, BE1218, CL2-57, and CL3-3. Most ligands were selective for D22 over TRAP220 recruitment, notably XL041 and XL652.




Fig. 6.

Coactivator recruitment to LXR induced by LXR ligands.

CRT data was normalized to T0 maximal response (100%) for all four CoA studied: LXR α (dashed lines) LXR β (solid lines). For clarity, only TRAP220 and D22 data are shown. As noted in the text, most ligands were β -selective or nonselective, with the exception of AZ876 that selectively stabilizes CoA binding to LXR α . GSK2033 and SR9238 were studied in the absence of agonist. Data shown mean and SD from at least triplicate measurements. (See [Fig. S6](#) [all four CoA studied](#)).

Corepressor stabilization and displacement

Support for the de-repression and canonical mechanisms of LXR activation comes from studies on selective knockout or dual knockout of LXR isoforms in mice and cells. As argued below, stabilization of corepressor binding to LXRA may bias towards nonlipogenic LXR agonist activity

Liver-specific knockout of LXRA attenuated the effect of T0 treatment on elevation of lipogenic genes and plasma TG.¹¹⁴ Although T0-induced elevation of liver ABCA1 was also attenuated by LXRA^{-/-}, these observations support the pursuit of LXRβ-selective agonists to avoid adverse effects in the liver. In LXRA^{-/-} mouse livers, the basal expression of some genes (*ABCA1* and *SREBF1*) was unchanged; whereas, others (e.g. *FAS*) were significantly decreased and *LXRβ* was greatly increased.¹³ The elevation of *ABCA1*, *SREBF1*, and *FAS* induced by LXR agonists was attenuated in LXRA^{-/-} relative to wild-type (WT). In the livers of dual LXRA^{-/-}/LXRβ^{-/-} mice, liver lipogenesis and TG levels were lower with normal or high-cholesterol diets.¹¹⁵ In another study, increases in hepatic LXR target genes induced by T0 and GW3965 were attenuated in LXRA^{-/-} versus WT mice; however, higher doses of LXR agonists significantly elevated *ABCA1* and *SREBF1* in both mice.¹¹⁶ The observation that agonist-induced *ABCA1* expression was amplified in the intestines was interpreted to reflect the tissue selectivity of LXR agonists and desirability of selecting against LXRA agonism in the liver.

Transcriptomic analysis of LXR isoform knockout adds complexity. In contrast to ubiquitous LXRβ expression, LXRA is localized in specific cells and tissues: e.g. liver and macrophages. Immortalized macrophages, LXR-null, or expressing either LXRβ or LXRA, were studied using GW3965 and/or GSK2033 as pharmacologic probes.¹¹⁷ Three distinct transcriptional mechanisms were proposed, one of which being pharmacologically nonresponsive and repressed in LXR-null macrophages can be assumed less relevant to LXR ligand design. In the classical or canonical mechanism (Mode II), transcription is weakly active or inactive in LXR-null and is active in the presence of an LXR agonist that induces a conformational change leading to coactivator binding. In the de-repression mechanism (Mode I), LXR-null is active, basal activity is repressed and the LXR agonist induces a conformational change that displaces the corepressor in favor of coactivator binding. *ABCA1* and *SREBF1* are used as example of genes controlled by Mode I and Mode II mechanisms, respectively.

A similar transcriptomic approach to study of LXR-null mouse livers emphasized that knockout of LXR has significant effects on other nuclear receptors and transcription factors, notably PPARα.¹¹⁸ Perhaps surprisingly, the majority of genes differentially expressed in WT versus LXR-null liver were insensitive to treatment with T0 and GW3965. Genes downregulated in LXR-null and induced by agonists in WT livers included lipogenic genes *SREBF1*, *SCD1*, *FAS*, and *LPCAT3*. Although genes mediating cholesterol metabolism and efflux (e.g. *ABCA1*, *APOA1*) are induced by LXR agonists, these were generally unchanged in WT and LXR-null livers. Thus, in the liver, induction of *ABCA1* and *SREBF1* may be dominated by canonical and de-repression mechanisms, respectively.

Arguing that stabilization of corepressor binding to LXRA biases towards nonlipogenic activity, we used CRT to measure ligand-dependent, LXR:corepressor interactions. Nuclear receptor corepressor (NCoR) and silencing mediator of retinoic acid and thyroid hormone receptor (SMRT) are large proteins (ca. 270 kDa) that share about 40% sequence identity. SMRT and NCoR are largely disordered platform proteins that act as a scaffold upon which the enzymatic machinery of the repression complex is built.¹¹⁹ In a transcriptionally repressed complex, a corepressor typically binds to the LBD.

The interaction motifs of coactivator and corepressor proteins compete for LBD binding. Binding of the coactivator interaction motif LXXLL is dependent on hydrophobic amino-acids in helix 12.¹²⁰ The corepressor interaction motif ($\phi x x \phi$ where ϕ is a hydrophobic amino-acid and x is any amino acid) uses additional flanking sequences, but contains a similar amphipathic core to the coactivator motif.^{121–123}

NCOR2 and SMRT2 were observed to bind to LXR isoforms in the absence of ligand, with CRT analysis showing the expected displacement of corepressor (loss of signal) by LXR agonists (**Fig. 7**). For some ligands, ligand stabilization of the corepressor-bound complex (gain of signal) was observed (**Fig. 7**, **S7**). Titration with LXR agonists, GW3965, its methyl-derivative (RGX-104), and T0 caused displacement of NCOR2 from both LXR isoforms; whereas LXR-623 and DMHCA had little or no effect (**Fig. 7A,B**). A similar pattern of ligand response was seen for binding of SMRT2 to LXR (**Fig. 7C,D**). Across isoforms and corepressors, several ligands showed little or no ability to displace NCOR2 and SMRT2 corepressors from LXR: CL2-57, CL3-3, LXR-623, and the antagonist GSK2033 (**Figs 7**, **S7**).

A subset of ligands elicited strong agonist responses (potency and efficacy for corepressor displacement) that were not strongly dependent on LXR isoform nor the identity of the corepressor: T0, RGX-104, GSK3987, and perhaps surprisingly, SR9238 that often displays the properties of an inverse agonist. AZ876 showed similar activity, with selectivity for corepressor displacement from LXR α (**Fig. 7**), the same selectivity seen for coactivator recruitment (**Fig. 6**). MK9 and ALX101 elicited similar responses to the “strong agonists” in displacement of corepressors from LXR β ; however, these ligands were less effective or ineffective in displacement of corepressors from LXR α . BE1218 was less efficacious in general and was ineffective in corepressor displacement from LXR α . Thus, several LXR ligands only stabilize conformations of LXR α that allow corepressor binding.

A strong agonist is expected to potently destabilize corepressor binding to LXR. In this context, SR9238 has strong agonist properties (**Fig. S7Q**), which is perhaps surprising, since it can act as an inverse agonist. In contrast to corepressor displacement, SR9238 was unable to recruit coactivators to LXR α and very weakly supported coactivator recruitment to LXR β . Thus, SR9238 displays unique interactions with LXR, strongly displacing corepressors and with little effect on coactivator recruitment. SR9238 cannot stabilize a conformation of LXR that allows corepressor binding, nor one that stabilizes coactivator binding. The conformation stabilized by SR9238 also cannot be that of the aporeceptor, because the unliganded LXR apo-receptor binds corepressors.

Based on CRT analysis of coregulator binding, SR9238 would be predicted to act as an antagonist; therefore, it is useful to compare with GSK2033, universally described as an antagonist. GSK2033 stabilizes a conformation of LXR that is incompatible with coactivator binding and equally GSK2033 does not stabilize a conformation that disrupts corepressor binding to LXR α (**Fig. S7P**). GSK2033 does selectively displace SMRT2 from LXR β . Two interesting conclusions result from this CRT analysis: 1) GSK2033 is able selectively to displace corepressors from LXR β ; and 2) GSK2033 is expected to act as a pure antagonist of LXR α -mediated transcription. In contrast, strong corepressor destabilization by SR9238 resulting in de-repression of LXR, is expected to activate LXR-mediated transcription. CRT data clearly differentiates SR9238 from the antagonist GSK2033.

Several ligands were unable to displace corepressors from LXR and XL041 stabilized LXR complexes with corepressor bound, as shown by the concentration-dependent increase in the TR-FRET signal (**Fig. S7K**). Similar behavior was seen for the structurally related XL652 in stabilization of SMRT2 corepressor binding to LXR α ; although, XL652 did not perturb SMRT2 binding to LXR β and weakly destabilized the NCOR2 complex with both LXR isoforms (**Fig. S7L**). XL041 binding induced LXR α conformations that strongly stabilized the corepressor-bound




Fig. 7.

Corepressor binding to LXR isoforms: ligand dependence.

Concentration-response from CRT measurements of: NCOR2 binding to LXR α (dashed lines) (A); NCOR2 binding to LXR β (solid lines) (B); SMRT2 binding to LXR α (dashed lines) (C); SMRT2 binding to LXR β (solid lines) (D). Data shown mean and SD from at least triplicate measurements. (See Fig. S7 [for alternative layout](#)).

complexes. The signal associated with corepressor binding to LXR β was not increased to the same degree; furthermore, the signal associated with SMRT2 binding was increased to a higher maximal response than that for NCOR2 binding.

Corepressor binding to LXR gave a much more diverse set of ligand-dependent responses in CRT to those observed for coactivator binding, primarily, because unliganded LXR binds corepressors but not coactivators. Of the observations that stand out from the CRT corepressor binding analysis, the diverse response from the biphenyl sulfone ligands is perhaps the most surprising, with a spectrum of corepressor binding/stabilization to displacement/destabilization: XL041 > XL652 > GSK2033 > BE1218 > SR9238.

Simplistically, if expulsion of the corepressor from the LXR α complex is correlated with hepatic LXR activation and lipogenesis, AZ876, GSK3987, T0, RGX-104, and SR9238 would be predicted to be lipogenic. This mechanism cannot be dominant in HepG2 cells, because SR9238 acts as an inverse agonist at < 3 μ M concentration. Therefore, corepressor binding alone cannot be considered in the absence of coactivator recruitment.

Synchronous coregulator recruitment by multiplex, precision CRT

We established a multiplexed, precision CRT (pCRT) assay using a one donor/two-acceptor system allowing for simultaneous measurement of corepressor dissociation and coactivator recruitment. Terbium has multiple emission wavelengths therefore it is an ideal fluorescence donor. Utilizing two spectrally different acceptor fluorophores such as XL665 and fluorescein allows measurement of two signals in the same assay as the fluorophore acceptor emissions do not overlap. pCRT utilized as corepressor, NCoR2-fluorescein, and as coactivator, the same construct, biotin-SRC1, used to determine LXR isoform selectivity (**Fig. 2**). The pCRT assay was optimized for a total volume of 12 μ L incubating LXR (4 nM), anti-GST-Tb (0.4 nM), SA-XL665 (9.44 nM), SRC1-btn (200 nM) and NCoR2-FL (200 nM), measuring TR-FRET emission ratios at 520/490 nm and 665/615 nm. Coregulators were in excess and concentrations of assay components were kept constant while ligand concentration-response was examined.

It is clear from the pCRT curves that the varied response of corepressor NCoR2 to ligand binding seen in CRT (**Fig. 7**) is “normalized” in the presence of coactivator SRC1 (**Fig. 8**). For every ligand studied by pCRT, ligand binding induced a conformation of LXR compatible with coactivator recruitment and concomitant corepressor displacement. The synchronous exchange of corepressor by coactivator leads to pCRT binding curves in which the potency for ligand-induced binding of SRC1 binding is almost identical to that for ligand-induced displacement of NCOR2. For example: (1) LXR-623 does not displace NCOR2 from LXR α in the absence of SRC1; (2) DMHCA does not displace NCOR2 from LXR β in the absence of SRC1; and (3) although GW3965 does displace NCOR2 from both LXR isoforms, the concentration-response for GW3965 shows a left-shift in the NCOR2 displacement curve of 100-fold (**Fig. 8**). In contrast to NCOR2 response, the SRC1 binding curves show only small shifts in the presence of NCOR2 from pCRT analysis (**Fig. 8**). For LXR β , the correlation between potency for SRC1 recruitment in monoplex and multiplexed CRT is good (data not shown).

Titration of the biphenyl sulfone ligands, XL041 and XL652, in the absence of SRC1 induced binding of NCOR2 to LXR α (for both ligands) and to LXR β (XL041 only) (**Fig. 7**). From pCRT analysis, the presence of SRC1 causes ligand-induced displacement of NCOR2 from LXR. For LXR β , in the presence of SRC1, the corepressor is fully displaced; whereas, the maximal signal for NCOR2-bound LXR α , is significantly higher than zero in the presence of saturating XL041 (**Fig. 8**) indicative of an equilibrium for the repressed XL041:NCOR2:LXR α and activated XL041:NCOR2:LXR α complexes.




Fig. 8.

Ligand-induced conformational change to LXR causing corepressor (NCOR2) replacement by coactivator (SRC1) measured by pCRT.

Concentration-response from CRT for SRC1 binding relative to apo-LXR (black dashed line) and NCOR2 binding relative to apo-LXR (magenta dashed line). Concentration-response from pCRT for SRC1 binding relative to apo-LXR (black solid line) and NCOR2 binding relative to apo-LXR (magenta solid line) with titration of ligand into solution of LXR, NCOR2, and SRC1. Data show mean of triplicate measurements.

CRT analysis of coactivator binding has often been used to determine LXR ligand potency, efficacy, and isoform selectivity. CRT analysis of corepressors is more rarely studied. Concentration-dependence of four LXR ligands (T0, GW3965, LN6500 and 22(R)-hydroxycholesterol) was studied by CRT for coactivator NCoA3 and separately for NCOR1.⁵² Weak recruitment of NCOR1 to LXR α was reported for GW3965 (> 1 μ M). This study was supported by cellular two-hybrid reporter assays in HepG2 and THP1 cells. T0 and GW3965 were classed as full and partial agonists, respectively. In our study, GW3965 and its methylated derivative, RGX-104, behaved as full agonists in CRT, with the exception of coactivator (PGC1 α , SRC2, D22, TRAP220) recruitment to LXR α (**Figs 6**, **6S**). Both agonists displaced NCOR2 and SMRT2 from LXR. In contrast, two biphenyl sulfone LXR agonists (XL041, XL265) strongly stabilized the corepressorbound conformation.

Correlation of CRT signatures with phenotype

With the knowledge that corepressors bind to the apo-receptor; whereas coactivator binding requires a ligand-induced conformational change, we can categorize all ligands using their CRT profiles: ligands induce 3 conformations with CoR (destabilizing D, stabilizing S, neutral N) and only 2 conformations with CoA (stabilizing S, neutral N) (Table 1). We categorized ligands based upon these CRT characteristics using labels compatible with CRT profiles: full agonist (FA), weak agonist (WA), partial agonist (PA), inverse agonist (IA), soft antagonist (SA), and hard antagonist (HA). The ligand category is not always identical for LXR α vs. LXR β . Similarly, the phenotype associated with each classification is isoform dependent. Recruitment of CoA to LXR β by FA, WA, and PA will activate transcription; whereas, lack of recruitment by SA, HA, and IA will antagonize transcription (Table 1). Stabilization of CoA and/or destabilization of CoR bound to LXR α is a derepression mechanism leading to activation/transcription for FA, WA, and IA; whereas partial activation is expected for PA.

To determine which LXR ligands fit with these theoretical CRT signatures, concentration-response curves were interpolated to measure normalized response at 1 μ M ligand concentration and plotted as a heat map (**Figs 9A**, **S8**). Selecting response at 1 μ M ligand is pharmacologically relevant and takes account of potency and efficacy. As bench-mark ligand, the response to T0 is uniform, without LXR isoform selectivity and with strong induction of ABCA1 and activation of SRE in cell cultures. Across the other 13 ligands with agonist activity, all are LXR β -selective, with the exception of AZ876 that is LXR α -selective (**Fig. S8**).

The CRT signatures of the three agonist and three antagonist classifications were modelled in the same heat map using signatures defined in Table 1 and the measured responses to the real LXR ligands in our study. There are several ligands with weak agonist (WA) signatures and GSK3987 fits a full agonist (FA) profile. XL041 has the signature of a partial agonist (PA), which matches its cellular phenotype. GSK2033 and SR9238 match the CRT signatures of a hard antagonist (HA) and inverse agonist (IA), respectively. We hypothesized that the signature of an NLAI, which by definition induces ABCA1 but does not activate SRE, should reflect a CRT profile of LXR β activation without activating LXR α : this leads to a theoretical NLAI profile, shown in **Fig. 9A**, which is not an exact match for any theoretical or experimental LXR ligand.

To give these signatures and labels a more thermodynamic basis, we can take learnings from CRT and pCRT measurements: (i) CoA-bound LXR is of relatively high energy, since it is not observed in absence of ligand; (ii) apo-LXR exists in equilibrium with CoR bound LXR, since CoR-bound LXR is observed in absence of ligand; (iii) The ligand-bound CoA:LXR complex is lowest energy, because CoA binding leads to full or partial CoR displacement. The relative stabilization and relative energy associated with each complex therefore follows (**Fig. 9B**). Thus, an inverse agonist (IA) CRT signature requires the IA to stabilize apo-LXR and a partial agonist (PA) stabilizes both coregulator-bound complexes.




Fig. 9.

CRT signatures and correlation with stabilization and structure of LXR:coregulator complexes.

(Table 1) Categorization of LXR ligands as agonists and antagonists conforming to CRT and pCRT signatures. **(A)** Heat map of responses (measured at 1 μ M) in cell-free and cell-based assays to LXR ligands together with theoretical response to agonist and antagonist ligand signatures. **(B)** Relative energy perturbations of apo and coregulator-bound LXR caused by each class of agonist and antagonist identified and classified in Table 1. **(C)** Co-crystal structure of LXR α :SRC1-2 co-crystal with GW3965 (PDB 3IPQ) showing key helices. **(D)** Superposition of LXR α :SRC1-2:GW3965 co-crystal with structure of LXR α :SMRT2 simulated using AlphaFold, showing the significant structural perturbation of H11 and H12 on ligand binding (red shades) compared to the ligand-free LXR α (blue shades).

Implications for discovery of nonlipogenic LXR ligands

As outlined in the Introduction, Belorusova et al., carried out multi-component analysis on an in-house chemical series of 9 LXR ligands, T0, GW3965, LXR-623, XL041, and AZ876, claiming identification of an “escape route from the adverse lipogenic effects of LXR ligands”.⁵⁰ Although the study was limited by binary definitions of *in vivo* lipogenicity, the in-depth HDX-MS analysis led to inferences on ligand-induced conformational change that might be associated with lipogenesis. Not unexpectedly, given the complexity of the problem, simple principles for NLAI design were not revealed. It was concluded that selection of ligands with “modest potencies” in cell-based transcriptional assays would lead to reduced lipogenesis.

Correlation analysis driven by HDX-MS data inferred the importance of helix-12 (H12) in activating a lipogenic transcriptional program. H12 is the key structural element of Activation Function-2 (AF-2), often described as a ligand-dependent switch and providing a binding surface for coactivator recruitment in the agonist-induced conformation (**Fig. 9C**). There are no published crystal structures of corepressor-bound LXR α , with most PDB entries being liganded coactivator-bound LXR β . To emphasize the conformational change in LXR α induced by agonist binding, we modelled SMRT2-bound apo-LXR α using AlphaFold and superimposed the corepressed apo-structure with a SRC1-bound agonist co-crystal structure (PDB 3IPQ **Fig. 9D**). This illustration shows the CoR, in the absence of ligand, bound to a surface that in the presence of ligand is occupied by H12. A ligand-induced conformation that stabilizes CoR binding to LXR cannot result from direct displacement of H12 by ligand.

Helix-3 (H3) forms part of the coactivator binding surface and the ligand-binding pocket. H3 contributes residues that interact directly with ligands. Helix 4 (H4) interacts with H3 and other helices to form the hydrophobic core, stabilize coactivator-bound conformations, and indirectly to influence receptor dimerization through helix-9 (H9). In LXR, helix-5 (H5) interacts directly with ligands and supports a more flexible ligand-binding pocket than in many other NRs. HDX-MS analysis of liganded LXR α structures correlated stabilization of H3 and H5 with selective ABCA1 induction. It was speculated that ligand interactions with H3 and H5 could induce a conformation leading to corepressor dissociation, which could lead to selectivity owing to the greater flexibility of H3 in LXR α . Conversely, ligands that stabilized H12, itself, were correlated with lipogenesis. Similarly, with PPAR γ , direct stabilization of H12 is associated with full agonism; whereas, partial agonists do not directly stabilize H12, which leads to biased agonism.¹²⁴

As in our study, Belorusova et al noted that the properties of XL014, were differentiated other LXR ligands. The HDX-MS analysis showed XL014 to act as an antagonist: i.e. XL014 did not induce an agonist conformation of H12. XL041 is generally described as an LXR β -selective partial agonist.¹²⁵ Our CRT analysis of coactivator binding (**Figs 6**, **S6**) reflects LXR β -selective partial agonist activity, with the exception of SRC1 CRT data. In cell-based phenotypic assays, XL041 is a potent partial agonist with selectivity (at concentration $\leq 1 \mu\text{M}$) for induction of ABCA1 over SRE activation (**Figs 3**, **4**). Taken together with the XL041-induced stabilization of corepressor-bound LXR (**Fig. 7**) the overall CRT signature of XL041 uniquely predicts partial agonist activity (**Fig. 9**).

XL041 stabilizes corepressor binding to LXR α (**Fig. 7**); although, in the presence of coactivator, SRC1, the corepressor is dissociated (**Fig. 8**). In our study, the biphenyl sulfone class of ligands yielded diverse responses in cell-free and cell-based assays, suggesting that this chemotype can be manipulated to enhance isoform and coregulator selectivity. The CRT profile of XL652, a close homologue of XL041, includes corepressor stabilization for LXR α (**Fig. 7**) and a response in HepG2 cells that shows little or no activation of SRE at submicromolar concentrations and more than 100-fold weaker potency for ABCA1 induction compared to XL041 (**Figs 3**, **5**). The phenotype of ALX101 is a full agonist towards ABCA1 induction and a partial agonist of SRE

activation; whereas, BE1218 has a partial agonist phenotype in both cell-based assays. The partial agonist phenotype is compatible with a relative inability to destabilize the corepressor-bound LXR α complex (Fig. S7 [↗](#)).

GSK2033 and SR9238 are very close structural congeners of BE1218, yet the response in phenotypic assays is quite different, with BE1218 a partial agonist, SR9238 displaying an inverse agonist and GSK2033 a mixed antagonist/inverse agonist phenotype (Figs 3 [↗](#), 5 [↗](#)). In CRT, both GSK2033 and SR9238 displace SRC1 and 24HC from an activated LXR β complex (Fig. 2 [↗](#)) but GSK2033 more closely resembles BE1218 in its weak effects on corepressor-bound LXR α (Fig. S7 [↗](#)). There is scope for further exploration of this chemotype using CRT/pCRT with a focus on interactions with the LXR α corepressor complex.

Limitations of the present study

Measurement of absolute affinity for coregulator binding to LXR by TR-FRET or other methods is complementary to the CRT approach used in this study, which measures relative potency and efficacy for ligand-induced coregulator binding. Absolute measurements require saturating ligand concentrations, which is problematic for weak LXR ligands, which by their nature are hydrophobic with limited aqueous solubility. Elegant studies measuring PPAR/coregulator binding free energy by fluorescence polarization have incorporated RXR heterodimers, full length NRs, and oligonucleotides.^{91 [↗](#), 124 [↗](#)} These studies demonstrate that key aspects of ligand-induced coregulator binding are replicated by the reductionist system of NR-LBD with short coregulator peptides as used in the present work.

Conclusions

Partial agonists are often described as a desirable therapeutic modality with anticipated efficacy and improved safety; although, the rationale for this improved therapeutic window is often intuitive rather than quantitative. The traditional medicinal chemistry approach to optimization of small molecule therapeutics is to increase potency and selectivity against an off-target: the approach in LXR agonist design has invariably been LXR β versus LXR α isoform selectivity. A nominated drug candidate will also maximize target exposure. XL041 ticks all the boxes as a potent, LXR β -selective ligand with partial agonist response in many *in vitro* assays. In human volunteers, XL041 caused a non-linear, dose-dependent increase in blood ABCA1 and plasma TG and LDL-C, with maximal response reached at approximately 2.5 mg/kg p.o. At this dose, the drug concentration in plasma on Day 14 was $C_{\max} \approx 30$ nM. Levels of TG and LDL-C returned to baseline after drug withdrawal; however, with $T_{1/2} \approx 13$ hours and the high potency of XL041, target exposure would be expected to be persistent. Limiting target exposure and potency may be required to attenuate activation of lipogenesis even by partial agonists. The clinical observations on the biphenyl sulfone XL041 are compatible with *in vitro* data reported by others and in this present study.

Preclinical animal studies have shown that enhancing ABCA1 activity with an LXR agonist could be an effective therapeutic strategy in a wide range of human conditions, including diabetes, cardiovascular disease, kidney failure, retinal degeneration, and neurodegenerative diseases, including AD.^{38 [↗](#), 126 [↗](#)–128 [↗](#)} In preclinical proof-of-concept studies T0, RGX104 and GW3965 continue to be used; for example, GW3965 in a tauopathy model of pathologic glial cell activation associated with AD.^{69 [↗](#)} We have reported beneficial effects of CL3-3 in FAD mice expressing human *APOE4*, the major genetic risk factor for AD.^{129 [↗](#)} We observe an excellent correlation between cell-free stabilization of CoA:LXR β complexes versus induction of ABCA1 in cells, supportive of the therapeutic focus on LXR β selectivity. Hepatic lipogenesis as a side effect, caused by LXR α activation in the liver, cannot be tolerated in many chronic diseases. Hepatic lipogenesis, observed in the XL041 clinical trial, is not tolerable in a population prone to atherosclerosis, the

disease target of XL041. The diverse responses of LXR ligands reflected in CoR:LXR α stabilization and SRE activation in HepG2 cells suggests that fine tuning of ligands with respect to LXR α binding is a route to attenuation of unwanted lipogenesis.

Experimental Section

Cell Culture

CCF-STTG1 and HepG2 cell lines were purchased from ATCC and cultured per the supplier's instructions. All cultures were maintained at 37°C in a humidified environment containing 5% CO₂. CCF-STTG1 cultures were grown in RPMI-1640 media (ATCC) and HepG2 in Eagles Modified Essential Medium (EMEM, ATCC 30-2003) supplemented with 10% fetal bovine serum (FBS, Gibco A52567-01) and 1% Penicillin/Streptomycin (P/S, Gibco 15140-122).

Luciferase reporter assays

ABCA1-luc assay

Low passage CCF-STTG1 cells (20,000 cells/well) were seeded in 96-well plates in media containing charcoalstripped FBS and incubated at 37 °C and 5% CO₂ for 24 hours prior to a 24-hour treatment. All compounds were dissolved in DMSO, and final DMSO concentrations never exceeded 1%. After a 24-hour incubation period, add a volume of ONE-GLO™EX Reagent that is equal to the volume of culture medium in each well and incubate for 3 to 5 minutes on the shaker. The luminescent activity of compounds (X to Y μ M, 10-fold dilution) was examined using the ONE-GLO™EX assay (Promega). The data was normalized to DMSO. The assays were performed in duplicates for each concentration, and the EC₅₀ values were determined from non-linear regression analysis of the dose-response curve generated in GraphPad Prism 9.

SRE-luc assay

HepG2 cells were cotransfected with SRE (LDLR)-NlucP SuperPiggyBac Transposon and a separate Transposase plasmid (System Biosciences). Cells were expanded for multiple passages followed by selection with puromycin (1 μ g/ml). Stable cells constitutively express GFP allowed for FACS enrichment and/or cell cloning. Initially, stable pools were compared to low expressing and high expressing enriched pools. The optimal pool was selected based on assay performance (Z') and was grown and maintained as above, with the addition of 1.0 μ g/mL of puromycin biweekly.

Cells were split into flasks containing CS medium (EMEM supplemented with 10% charcoal stripped FBS (Gibco A52567-01) and 1% P/S and grown at 37°C /5% CO₂ for 48 hours before plating. Cells were plated in 384-well, white clear tissue culture plates (Greiner 781098) and allowed to grow at 37°C /5% CO₂ for 24 hours before medium was aspirated and compound dilutions prepared in CS medium were added. Following a 24-hour incubation, the medium was aspirated, and cells were lysed/luminescence quantified using the Nano-Glo Luciferase Assay System (Promega N1130). Luminescence was measured on a CLARIOstar® Plus multi-mode plate reader (BMG Labtech, Ortenberg, Germany). Experimental data were normalized to controls and fit with a four parameter Hill equation using Graphpad Prism software (Boston, MA) to determine EC₅₀ values.

qPCR. HepG2 cells were seeded in 12-well plates and treated with compounds for 24 hours. Following treatment, cells were washed once with PBS and then lysed in 350 μ L of RLT buffer (Qiagen). Lysates were immediately frozen at -80 °C until RNA extraction. Total RNA was extracted using RNeasy Plus Mini columns (Qiagen) according to the manufacturer's instructions. RNA concentration and purity were assessed using the A260/A280 ratio measured on a Synergy Neo2 plate reader (BioTek). For cDNA synthesis, 2 μ g of total RNA was reverse transcribed using

SuperScript III reverse transcriptase (Invitrogen), oligo(dT)12–18 primers, and dNTP mix, following the manufacturer's protocol. Quantitative PCR was performed using TaqMan Gene Expression Master Mix (Invitrogen) with 2 μ L of cDNA and gene-specific TaqMan probes in a 20 μ L reaction volume per well. The following TaqMan assays were used: ABCA1: Hs01059101_m1, SREBF1: Hs01088691_m1, SCD: Hs01682761_m1, FASN: Hs01005622_m1, and HPRT1 (endogenous control): Hs02800695_m1. qPCR was performed on a StepOne Real-Time PCR System (Applied Biosystems), and gene expression was quantified using the $\Delta\Delta$ Ct method.

Coactivator recruitment TR-FRET/HTRF (CRT)

LXR α/β . To determine compound EC50 values, 10mM compound stocks were prepared in DMSO. An initial 8-concentration, 3-fold dilution series in DMSO was performed in 96-well v-bottom, polypropylene plates at 100x the final desired concentration. Intermediate dilution plates were prepared by diluting the 100x DMSO stocks 25-fold in assay buffer (50mM Tris, pH 7.4, 50mM KCl, 0.05% Triton-X 100, 10% glycerol, 0.1% BSA-(added fresh), 5mM DTT (added fresh)), creating a 4x stock. The following assay components were prepared individually at a 4x final concentration in assay buffer: (1) LXR α/β - GST tag (Thermo Fisher Scientific PV4657/PV PV4660); (2) a mixture of anti-GST-Tb (CisBio/Perkin Elmer 61GSTTLA, lot 14A) and SA-XL665 (CisBio/Perkin Elmer 610SAXLB); (3) SRC1-btn (AnaSpec AS62152/Fisher Scientific NC1565694). A 3uL aliquot of the 4x lig- and was then added to each well of the 384-well small volume non-binding black assay plate (Greiner 784900) followed by a 3uL aliquot of each reaction component for a total volume of 12uL. Control wells contained all components minus ligands, replaced by positive control T0 (Enzo 270-309-M010/VWR 89147-616) or DMSO. Incubated at 1hr at room temperature before measurement of TR-FRET. Final concentrations of assay components LXR α/β (4nM), anti-GST-Tb (0.4nM), SA-XL665(9.44nM), and SRC1-btn (100nM). TR-FRET measurements were performed on a CLARIOstar® Plus multi-mode plate reader (BMG Labtech, Ortenberg, Germany), excitation 337nm (laser), emission1 615nm and emission2 665nm. The signal is expressed as a ratio of TR-FRET measurements at 665 nm/615nm $\times$ 1000. Experimental data were normalized to controls and fit with a four parameter Hill equation using Graphpad Prism software (Boston, MA) to determine EC50s. Full details for other NR CRT assays are included in **Supporting Information**.

LXR (coregulator selectivity)

For all assay, ligands were first prepared at 100x of desired final concentration in DMSO. Ligands were serially diluted in a 3x manner in 96-well polypropylene plate. The ligands were then diluted 4x final concentration in assay buffer. The following assay components were prepared individually at a 4x final concentration in assay buffer: (1) LXR α/β -GST tag (Thermo Fisher Scientific PV4657/PV PV4660); (2) a mixture of anti-GST-Tb (Cis-Bio/Perkin Elmer 61GSTTLA, lot 14A) (3) coregulator-FL. A 3uL aliquot of the 4x ligand was then added to each well of the 384-well small volume non-binding black assay plate (Greiner 784900) followed by a 3uL aliquot of each reaction component for a total volume of 12uL. Control wells contained all components minus ligands, replaced by positive control T0 (Enzo 270-309-M010/VWR 89147-616) or DMSO. Incubated at 1hr at room temperature before measurement of TR-FRET. Final concentrations of assay components LXR α/β (3nM), anti-GST-Tb (0.4nM), and coregulator-FL (200nM). TR-FRET measurements were performed on a CLARIOstar® Plus multi-mode plate reader (BMG Labtech, Ortenberg, Germany), excitation 337 nm (laser), emission1 520nm and emission2 490 nm. The signal is expressed as a ratio of TR-FRET measurements at 520nm/490nm $\times$ 1000. Experimental data were normalized to controls and fit with a four parameter Hill equation using Graphpad Prism software (Boston, MA) to determine EC50s.

Multiplex LXR pCRT one donor/two-acceptor fluorophore

For all assays, ligands were first prepared at 100x of desired final concentration in DMSO. Ligands were serially diluted in a 3x manner in 96-well polypropylene plate. The ligands were then diluted to a 4x final concentration in assay buffer. The following assay components were prepared

individually at a 4x final concentration in assay buffer: (1) LXR α /β; (2) a mixture of anti-GST-Tb and SA-XL665; (3) a mixture of SRC1-btn and NCoR2-FL. A 3uL aliquot of the 4x ligand was then added to each well of the 384-well black assay plate followed by a 3uL aliquot of each reaction component for a total volume of 12uL. Incubated at 1hr at room temperature before measurement of TR-FRET. Final concentrations of assay components LXR α /β (4 nM), anti-GST-Tb (0.4 nM), SA-XL665 (9.44 nM), SRC1-btn (200 nM) and NCoR2-FL (200 nM). TR-FRET measurements were performed on a CLARI-Ostar® Plus multi-mode plate reader (BMG Labtech, Ortenberg, Germany). Two different measurements were taken. The first measurement-excitation 337nm (laser), emission1 520nm and emission2 490nm. The signal is expressed as a ratio of TR-FRET measurements at 520 nm/490nm $\times$ 1000. The second measurement-excitation 337nm (laser), emission1 615nm and emission2 665nm. The signal is expressed as a ratio of TR-FRET measurements at 665 nm/615nm $\times$ 1000. Experimental data were normalized to controls and fit with a four parameter Hill equation using Graphpad Prism software (Boston, MA) to determine EC₅₀ values.

Modelling. Structural Alignment and Model Generation Using PyMOL and AlphaFold

The SMRT2 peptide sequence (CPSSHSLTERHKILHRLQEGSPS) and FASTA sequence of LXR α (PDB 3IPQ) were submitted together to the AlphaFold server using the “Sequences” tool with the prediction mode set to “Protein” to yield a predicted model of the LXR α -SMRT2 complex. To evaluate this model, we aligned it with the LXR α :SRC1-2 co-crystal with GW3965.

Chemical synthesis

Detailed methods are provided in Supporting Information for CL2-57, CL3-3 and Merck Compound 9, including characterization and purity.

Additional information

Author contributions

GRJT conceived the project. MSL designed and coordinated the experimental approach. In vitro studies were conducted by MSL, MAB, ST, FA, MR, SRM, AA, SK, MS, CP and NM. Chemistry was conducted by GRV. The manuscript was written by GRJT and MSL with input from all authors.

Notes

GRJT is an inventor on patents owned by the University of Illinois.

Funding Sources

This work was supported by U01AG076450. MSL was supported by NIH T32 GM008804.

Abbreviations

- (AD): Alzheimer’s disease
- (ADRD): Alzheimer’s disease and related dementia
- (Aβ): amyloid-β
- (APOA1): Apolipoprotein A1
- (APOE): Apolipoprotein E
- (ABC): ATP-binding cassette transporter
- (ACC): acetyl-CoA carboxylase
- (AF-2): activation function-2
- (APP): amyloid precursor protein

- (BBB): blood-brain barrier
- (CAR): constitutive androstane receptor
- (CETP): cholesteryl ester transfer protein
- (CNS): central nervous system
- (CRT): coregulator time-resolved fluorescence energy transfer
- (pCRT): precision CRT
- (CSF): cerebrospinal fluid
- (CYP): cytochrome p450
- (CVD): cardiovascular disease
- (DDIs): drugdrug interactions
- (DMHCA): N,N-dimethyl-3 β -hydroxycholeamide
- (EMEM): eagles modified essential medium
- (ER): estrogen receptor
- (FA): full agonist
- (FAS): fatty-acid synthase
- (FAD): familial AD
- (FXR): farnesoid X receptor
- (FBS): fetal bovine serum
- (HA): hard antagonist
- (H3): helix-3
- (H5): Helix-5
- (H): helix-9
- (H12): helix-12
- (HDL): high-density lipoprotein
- (HFD): high-fat diet
- (HTS): high-throughput screen
- (HDX-MS): H/D-exchange mass spectrometry
- (IA): inverse agonist
- (KO): knockout
- (LDL): low-density lipoprotein
- (LXR): liver X receptor
- (MK9): Merck Compound 9
- (NASH): non-alcoholic steatohepatitis
- (NCOA1): nuclear receptor coactivator 1
- (NCoR): nuclear receptor corepressor
- (NHR): nuclear hormone receptor
- (NLAI): nonlipogenic ABCA1 inducer
- (OCA): obeticholic acid
- (PA): partial agonist
- (PPAR): peroxisome proliferator activated receptor
- (PGC-1): PPAR γ coactivator 1
- (PXR): pregnane X receptor
- (P/S): penicillin/streptomycin
- (RCT): reverse cholesterol transport
- (RAR): retinoic acid receptor
- (RXR): retinoid X receptor
- (SA): soft antagonist
- (SERMs): selective estrogen receptor modulators
- (SMRT): silencing mediator of retinoic acid and thyroid hormone receptor
- (SRC-1): steroid receptor coactivator 1
- (SRE): sterol response element
- (SREBP1c): sterolresponse element binding protein 1c
- (SCAP): SREBP cleavage-activating protein
- (SRE-luc): SRE promoter luciferase reporter assay

- (TG): triglyceride
- (TR-FRET): time-resolved fluorescence energy transfer
- (TRAPs): thyroid hormone receptor-associated proteins
- (T0): T0901317
- (T2D): type-2 diabetes
- (WA): weak agonist
- (WT): wild-type
- (VLDL-C): very low-density lipoprotein cholesterol
- (24OHC): 24-hydroxycholesterol
- (24HC): 24(S)-hydroxycholesterol
- (27OHC): and 27-hydroxycholesterol.

Additional files

Supplementary file 1. Tables S1 and S2 showing NR Panel assay components and CRT peptide sequences. Fig S1 showing NR Panel responses. Fig S2 showing control reproducibility in cellular experiments. Fig S3 showing additional correlation plots of cellular data and LXR isoform selectivity. Fig S4 showing full dose response curves for HepG2 data. Fig S5 showing coactivator correlation plots. Fig S6 showing all coactivator responses for ligands. Fig S7 corepressor responses separated by individual ligands. Fig S8 heat map showing coregulator efficacy across ligands. Experimental details for NR Panel experiments and chemical synthesis. NMR spectra for synthesized compounds. [↗](#)

References

1. Foley P. (2010) **Lipids in Alzheimer's disease: A century-old story** *Biochim Biophys Acta* **1801**:750–3 [Google Scholar](#)
2. Martinez A. E., Weissberger G., Kuklenyik Z., He X., Meuret C., Parekh T., Rees J. C., Parks B. A., Gardner M. S., King S. M., Collier T. S., Harrington M. G., Sweeney M. D., Wang X., Zlokovic B. V., Joe E., Nation D. A., Schneider L. S., Chui H. C., Barr J. R., Han S. D., Krauss R. M., Yassine H. N. (2023) **The small HDL particle hypothesis of Alzheimer's disease** *Alzheimers Dement* **19**:391–404 [Google Scholar](#)
3. Livingston G., Huntley J., Sommerlad A., Ames D., Ballard C., Banerjee S., Brayne C., Burns A., Cohen-Mansfield J., Cooper C., Costafreda S. G., Dias A., Fox N., Gitlin L. N., Howard R., Kales H. C., Kivimaki M., Larson E. B., Ogunniyi A., Orgeta V., Ritchie K., Rockwood K., Sampson E. L., Samus Q., Schneider L. S., Selbaek G., Teri L., Mukadam N. (2020) **Dementia prevention, intervention, and care: 2020 report of the Lancet Commission** *Lancet* **396**:413–446 [Google Scholar](#)
4. Livingston G., Huntley J., Liu K. Y., Costafreda S. G., Selbaek G., Alladi S., Ames D., Banerjee S., Burns A., Brayne C., Fox N. C., Ferri C. P., Gitlin L. N., Howard R., Kales H. C., Kivimaki M., Larson E. B., Nakasujja N., Rockwood K., Samus Q., Shirai K., Singh-Manoux A., Schneider L. S., Walsh S., Yao Y., Sommerlad A., Mukadam N. (2024) **Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission** *Lancet* **404**:572–628 [Google Scholar](#)
5. Alzheimer A., Stelzmann R. A., Schnitzlein H. N., Murtagh F. R. (1995) **An English translation of Alzheimer's 1907 paper, "Über eine eigenartige Erkrankung der Hirnrinde"** *Clin Anat* **8**:429–31 [Google Scholar](#)

6. Ralhan I., Chang C. L., Lippincott-Schwartz J., Ioannou M. S. (2021) **Lipid droplets in the nervous system** *J Cell Biol* **220** [Google Scholar](#)
7. Tcw J., Qian L., Pipalia N. H., Chao M. J., Liang S. A., Shi Y., Jain B. R., Bertelsen S. E., Kapoor M., Marcora E., Sikora E., Andrews E. J., Martini A. C., Karch C. M., Head E., Holtzman D. M., Zhang B., Wang M., Maxfield F. R., Poon W. W., Goate A. M. (2022) **Cholesterol and matrisome pathways dysregulated in astrocytes and microglia** *Cell* **185**:2213–2233 [Google Scholar](#)
8. Smolic T., Zorec R., Vardjan N. (2021) **Pathophysiology of Lipid Droplets in Neuroglia** *Antioxidants (Basel)* **11** [Google Scholar](#)
9. Abildayeva K., Jansen P. J., Hirsch-Reinshagen V., Bloks V. W., Bakker A. H., Ramaekers F. C., de Vente J., Groen A. K., Wellington C. L., Kuipers F., Mulder M. (2006) **24(S)-hydroxycholesterol participates in a liver X receptor-controlled pathway in astrocytes that regulates apolipoprotein E-mediated cholesterol efflux** *J Biol Chem* **281**:12799–808 [Google Scholar](#)
10. Victor M. B., Leary N., Luna X., Meharena H. S., Scannail A. N., Bozzelli P. L., Samaan G., Murdock M. H., von Maydell D., Effenberger A. H., Cerit O., Wen H. L., Liu L., Welch G., Bonner M., Tsai L. H. (2022) **Lipid accumulation induced by APOE4 impairs microglial surveillance of neuronal-network activity** *Cell Stem Cell* **29**:1197–1212 [Google Scholar](#)
11. Haney M. S., Palovics R., Munson C. N., Long C., Johansson P. K., Yip O., Dong W., Rawat E., West E., Schlachetzki J. C. M., Tsai A., Guldner I. H., Lamichhane B. S., Smith A., Schaum N., Calcuttawala K., Shin A., Wang Y. H., Wang C., Koutsodendris N., Serrano G. E., Beach T. G., Reiman E. M., Glass C. K., Abu-Remaileh M., Enejder A., Huang Y., Wyss-Coray T. (2024) **APOE4/4 is linked to damaging lipid droplets in Alzheimer’s disease microglia** *Nature* **628**:154–161 [Google Scholar](#)
12. Wellington C. L., Walker E. K., Suarez A., Kwok A., Bissada N., Singaraja R., Yang Y. Z., Zhang L. H., James E., Wilson J. E., Francone O., McManus B. M., Hayden M. R. (2002) **ABCA1 mRNA and protein distribution patterns predict multiple different roles and levels of regulation** *Lab Invest* **82**:273–83 [Google Scholar](#)
13. Fitzgerald M. L., Mendez A. J., Moore K. J., Andersson L. P., Panjeton H. A., Freeman M. W. (2001) **ATP-binding cassette transporter A1 contains an NH2-terminal signal anchor sequence that translocates the protein’s first hydrophilic domain to the exoplasmic space** *J Biol Chem* **276**:15137–45 [Google Scholar](#)
14. Oram J. F. (2003) **HDL apolipoproteins and ABCA1: partners in the removal of excess cellular cholesterol** *Arterioscler Thromb Vasc Biol* **23**:720–7 [Google Scholar](#)
15. Dean M., Hamon Y., Chimini G. (2001) **The human ATP-binding cassette (ABC) transporter superfamily** *J Lipid Res* **42**:1007–17 [Google Scholar](#)
16. Oram J. F., Heinecke J. W. (2005) **ATP-binding cassette transporter A1: a cell cholesterol exporter that protects against cardiovascular disease** *Physiol Rev* **85**:1343–72 [Google Scholar](#)
17. Remaley A. T., Stonik J. A., Demosky S. J., Neufeld E. B., Bocharov A. V., Vishnyakova T. G., Eggeman T. L., Patterson A. P., Duverger N. J., Santamarina-Fojo S., Brewer H. B. (2001) **Apolipoprotein specificity for lipid efflux by the human ABCA1 transporter** *Biochem Biophys Res Commun* **280**:818–23 [Google Scholar](#)

18. Francis G. A., Knopp R. H., Oram J. F. (1995) **Defective removal of cellular cholesterol and phospholipids by apolipoprotein A-I in Tangier Disease** *J Clin Invest* **96**:78–87 [Google Scholar](#)
19. Frambach S., de Haas R., Smeitink J. A. M., Rongen G. A., Russel F. G. M., Schirris T. J. J. (2020) **Brothers in Arms: ABCA1- and ABCG1-Mediated Cholesterol Efflux as Promising Targets in Cardiovascular Disease Treatment** *Pharmacological reviews* **72**:152–190 [Google Scholar](#)
20. Kim W. S., Guillemin G. J., Glaros E. N., Lim C. K., Garner B. (2006) **Quantitation of ATP-binding cassette subfamily-A transporter gene expression in primary human brain cells** *NeuroReport* **17**:891–6 [Google Scholar](#)
21. Feingold K. R. (2000) **Introduction to Lipids and Lipoproteins** In: Feingold K. R., Anawalt B., Boyce A., Chrousos G., de Herder W. W., Dungan K., Grossman A., Hershman J. M., Hofland J., Kaltsas G., Koch C., Kopp P., Korbonits M., McLachlan R., Morley J. E., New M., Purnell J., Singer F., Stratakis C. A., Trencle D. L., Wilson D. P., editors. *Endotext* [Google Scholar](#)
22. Goldstein J. L., Brown M. S. (2009) **The LDL receptor** *Arterioscler Thromb Vasc Biol* **29**:431–8 [Google Scholar](#)
23. Schnitzler J. G., Hoogeveen R. M., Ali L., Prange K. H. M., Waissi F., van Weeghel M., Bachmann J. C., Versloot M., Borrelli M. J., Yeang C., De Kleijn D. P. V., Houtkooper R. H., Koschinsky M. L., de Winther M. P. J., Groen A. K., Witztum J. L., Tsimikas S., Stroes E. S. G., Kroon J. (2020) **Atherogenic Lipoprotein(a) Increases Vascular Glycolysis, Thereby Facilitating Inflammation and Leukocyte Extravasation** *Circ Res* **126**:1346–1359 [Google Scholar](#)
24. Rye K. A., Bursill C. A., Lambert G., Tabet F., Barter P. J. (2009) **The metabolism and anti-atherogenic properties of HDL** *J Lipid Res* **50**:S195–200 [Google Scholar](#)
25. Segrest J. P., Jones M. K., De Loof H., Brouillette C. G., Venkatachalapathi Y. V., Anantharamaiah G. M. (1992) **The amphipathic helix in the exchangeable apolipoproteins: a review of secondary structure and function** *J Lipid Res* **33**:141–66 [Google Scholar](#)
26. Tall A. R., Yvan-Charvet L. (2015) **Cholesterol, inflammation and innate immunity** *Nat Rev Immunol* **15**:104–16 [Google Scholar](#)
27. Gall J., Frisdal E., Bittar R., Le Goff W., Bruckert E., Lesnik P., Guerin M., Giral P. (2016) **Association of Cholesterol Efflux Capacity With Clinical Features of Metabolic Syndrome: Relevance to Atherosclerosis** *J Am Heart Assoc* **5** [Google Scholar](#)
28. Zhu Y., Liao H., Xie X., Yuan Y., Lee T. S., Wang N., Wang X., Shyy J. Y., Stemerman M. B. (2005) **Oxidized LDL downregulates ATP-binding cassette transporter-1 in human vascular endothelial cells via inhibiting liver X receptor (LXR)** *Cardiovasc Res* **68**:425–32 [Google Scholar](#)
29. Shao B., Tang C., Sinha A., Mayer P. S., Davenport G. D., Brot N., Oda M. N., Zhao X. Q., Heinecke J. W. (2014) **Humans with atherosclerosis have impaired ABCA1 cholesterol efflux and enhanced high-density lipoprotein oxidation by myeloperoxidase** *Circ Res* **114**:1733–42 [Google Scholar](#)
30. Schaefer E. J., Santos R. D., Asztalos B. F. (2010) **Marked HDL deficiency and premature coronary heart disease** *Curr Opin Lipidol* **21**:289–97 [Google Scholar](#)
31. Villarreal-Molina M. T., Flores-Dorantes M. T., Arellano-Campos O., Villalobos-Comparan M., Rodriguez-Cruz M., Miliar-Garcia A., Huertas-Vazquez A., Menjivar M., Romero-Hidalgo S.,

- Wacher N. H., Tusie-Luna M. T., Cruz M., Aguilar-Salinas C. A., Canizales-Quinteros S., Metabolic Study, G (2008) **Association of the ATP-binding cassette transporter A1 R230C variant with early-onset type 2 diabetes in a Mexican population** *Diabetes* **57**:509–13 [Google Scholar](#)
32. Doosti M., Najafi M., Reza J. Z., Nikzamir A. (2010) **The role of ATP-binding-cassette-transporter-A1 (ABCA1) gene polymorphism on coronary artery disease risk** *Transl Res* **155**:185–90 [Google Scholar](#)
33. Jung D., Cao S., Liu M., Park S. (2018) **A Meta-Analysis of the Associations Between the ATP-Binding Cassette Transporter ABCA1 R219K (rs2230806) Polymorphism and the Risk of Type 2 Diabetes in Asians** *Horm Metab Res* **50**:308–316 [Google Scholar](#)
34. Tai L. M., Thomas R., Marottoli F. M., Koster K. P., Kanekiyo T., Morris A. W., Bu G. (2016) **The role of APOE in cerebrovascular dysfunction** *Acta Neuropathol* **131**:709–23 [Google Scholar](#)
35. Bowman G. L., Kaye J. A., Quinn J. F. (2012) **Dyslipidemia and blood-brain barrier integrity in Alzheimer's disease** *Curr Gerontol Geriatr Res* **2012**:184042 [Google Scholar](#)
36. Koldamova R., Fitz N. F., Lefterov I. (2014) **ATP-binding cassette transporter A1: from metabolism to neurodegeneration** *Neurobiol Dis* **72**:13–21 [Google Scholar](#)
37. Tang Q., Wang F., Yang J., Peng H., Li Y., Li B., Wang S. (2019) **Revealing a Novel Landscape of the Association Between Blood Lipid Levels and Alzheimer's Disease: A Meta-Analysis of a Case-Control Study** *Front Aging Neurosci* **11**:370 [Google Scholar](#)
38. Lewandowski C. T., Laham M. S., Thatcher G. R. J. (2022) **Remembering your A, B, C's: Alzheimer's disease and ABCA1** *Acta Pharm Sin B* **12**:995–1018 [Google Scholar](#)
39. Karch C. M., Goate A. M. (2015) **Alzheimer's disease risk genes and mechanisms of disease pathogenesis** *Biol Psychiatry* **77**:43–51 [Google Scholar](#)
40. Arenas F., Garcia-Ruiz C., Fernandez-Checa J. C. (2017) **Intracellular Cholesterol Trafficking and Impact in Neurodegeneration** *Front Mol Neurosci* **10**:382 [Google Scholar](#)
41. Joseph S. B., McKilligin E., Pei L., Watson M. A., Collins A. R., Laffitte B. A., Chen M., Noh G., Goodman J., Hagger G. N., Tran J., Tippin T. K., Wang X., Lusi A. J., Hsueh W. A., Law R. E., Collins J. L., Willson T. M., Tontonoz P. (2002) **Synthetic LXR ligand inhibits the development of atherosclerosis in mice** *Proc Natl Acad Sci U S A* **99**:7604–9 [Google Scholar](#)
42. van der Hoorn J., Linden D., Lindahl U., Bekkers M., Voskuilen M., Nilsson R., Oscarsson J., Lindstedt E., Princen H. (2011) **Low dose of the liver X receptor agonist, AZ876, reduces atherosclerosis in APOE*3Leiden mice without affecting liver or plasma triglyceride levels** *Br J Pharmacol* **162**:1553–63 [Google Scholar](#)
43. Srivastava R. A. (2011) **Evaluation of anti-atherosclerotic activities of PPAR-alpha, PPAR-gamma, and LXR agonists in hyperlipidemic atherosclerosis-susceptible F(1)B hamsters** *Atherosclerosis* **214**:86–93 [Google Scholar](#)
44. Fessler M. B. (2018) **The challenges and promise of targeting the Liver X Receptors for treatment of inflammatory disease** *Pharmacology & therapeutics* **181**:1–12 [Google Scholar](#)
45. Loren J., Huang Z., Laffitte B. A., Molteni V. (2013) **Liver X receptor modulators: a review of recently patented compounds (2009 - 2012)** *Expert Opin Ther Pat* **23**:1317–35 [Google Scholar](#)

46. Fu X., Menke J. G., Chen Y., Zhou G., MacNaul K. L., Wright S. D., Sparrow C. P., Lund E. G. (2001) **27-hydroxycholesterol is an endogenous ligand for liver X receptor in cholesterol-loaded cells** *J Biol Chem* **276**:38378–87 [Google Scholar](#)
47. Matsuda A., Nagao K., Matsuo M., Kioka N., Ueda K. (2013) **24(S)-hydroxycholesterol is actively eliminated from neuronal cells by ABCA1** *J Neurochem* **126**:93–101 [Google Scholar](#)
48. Katz A., Udata C., Ott E., Hickey L., Burczynski M. E., Burghart P., Vesterqvist O., Meng X. (2009) **Safety, pharmacokinetics, and pharmacodynamics of single doses of LXR-623, a novel liver X-receptor agonist, in healthy participants** *J Clin Pharmacol* **49**:643–9 [Google Scholar](#)
49. Kirchgessner T. G., Sleph P., Ostrowski J., Lupisella J., Ryan C. S., Liu X., Fernando G., Grimm D., Shipkova P., Zhang R., Garcia R., Zhu J., He A., Malone H., Martin R., Behnia K., Wang Z., Barrett Y. C., Garmise R. J., Yuan L., Zhang J., Gandhi M. D., Wastall P., Li T., Du S., Salvador L., Mohan R., Cantor G. H., Kick E., Lee J., Frost R. J. (2016) **Beneficial and Adverse Effects of an LXR Agonist on Human Lipid and Lipoprotein Metabolism and Circulating Neutrophils** *Cell metabolism* **24**:223–33 [Google Scholar](#)
50. Belorusova A. Y., Evertsson E., Hovdal D., Sandmark J., Bratt E., Maxvall I., Schulman I. G., Akerblad P., Lindstedt E. L. (2019) **Structural analysis identifies an escape route from the adverse lipogenic effects of liver X receptor ligands** *Commun Biol* **2**:431 [Google Scholar](#)
51. Lin W., Chen T. (2018) **Using TR-FRET to Investigate Protein-Protein Interactions: A Case Study of PXR-Coregulator Interaction** *Adv Protein Chem Struct Biol* **110**:31–63 [Google Scholar](#)
52. Albers M., Blume B., Schlueter T., Wright M. B., Kober I., Kremoser C., Deuschle U., Koegl M. (2006) **A novel principle for partial agonism of liver X receptor ligands. Competitive recruitment of activators and repressors** *J Biol Chem* **281**:4920–30 [Google Scholar](#)
53. Li N., Wang X., Xu Y., Lin Y., Zhu N., Liu P., Lu D., Si S. (2017) **Identification of a novel liver X receptor agonist that regulates the expression of key cholesterol homeostasis genes with distinct pharmacological characteristics** *Molecular Pharmacology* [Google Scholar](#)
54. Kim G. H., Oh G. S., Yoon J., Lee G. G., Lee K. U., Kim S. W. (2015) **Hepatic TRAP80 selectively regulates lipogenic activity of liver X receptor** *J Clin Invest* **125**:183–93 [Google Scholar](#)
55. Santos R., Ursu O., Gaulton A., Bento A. P., Donadi R. S., Bologa C. G., Karlsson A., Al-Lazikani B., Hersey A., Oprea T. I., Overington J. P. (2017) **A comprehensive map of molecular drug targets** *Nat Rev Drug Discov* **16**:19–34 [Google Scholar](#)
56. Smith C. L., O'Malley B. W. (2004) **Coregulator function: a key to understanding tissue specificity of selective receptor modulators** *Endocr Rev* **25**:45–71 [Google Scholar](#)
57. Schultz J. R., Tu H., Luk A., Repa J. J., Medina J. C., Li L., Schwendner S., Wang S., Thoolen M., Mangelsdorf D. J., Lustig K. D., Shan B. (2000) **Role of LXRs in control of lipogenesis** *Genes Dev* **14**:2831–8 [Google Scholar](#)
58. Toporova L., Grimaldi M., Boulahtouf A., Balaguer P. (2020) **Assessing the Selectivity of FXR, LXRs, CAR, and RORgamma Pharmaceutical Ligands With Reporter Cell Lines** *Front Pharmacol* **11**:1122 [Google Scholar](#)
59. Houck K. A., Borchert K. M., Hepler C. D., Thomas J. S., Bramlett K. S., Michael L. F., Burris T. P. (2004) **T0901317 is a dual LXR/FXR agonist** *Mol Genet Metab* **83**:184–7 [Google Scholar](#)

60. Mitro N., Vargas L., Romeo R., Koder A., Saez E. (2007) **T0901317 is a potent PXR ligand: implications for the biology ascribed to LXR** *FEBS Lett* **581**:1721–6 [Google Scholar](#)
61. Collins J. L., Fivush A. M., Watson M. A., Galardi C. M., Lewis M. C., Moore L. B., Parks D. J., Wilson J. G., Tippin T. K., Binz J. G., Plunket K. D., Morgan D. G., Beaudet E. J., Whitney K. D., Klierer S. A., Willson T. M. (2002) **Identification of a nonsteroidal liver X receptor agonist through parallel array synthesis of tertiary amines** *J Med Chem* **45**:1963–6 [Google Scholar](#)
62. Chisholm J. W., Hong J., Mills S. A., Lawn R. M. (2003) **The LXR ligand T0901317 induces severe lipogenesis in the db/db diabetic mouse** *J Lipid Res* **44**:2039–48 [Google Scholar](#)
63. Koldamova R. P., Lefterov I. M., Staufenbiel M., Wolfe D., Huang S., Glorioso J. C., Walter M., Roth M. G., Lazo J. S. (2005) **The liver X receptor ligand T0901317 decreases amyloid beta production in vitro and in a mouse model of Alzheimer's disease** *J Biol Chem* **280**:4079–88 [Google Scholar](#)
64. Baranowski M., Zabielski P., Blachnio-Zabielska A. U., Harasim E., Chabowski A., Gorski J. (2014) **Insulin-sensitizing effect of LXR agonist T0901317 in high-fat fed rats is associated with restored muscle GLUT4 expression and insulin-stimulated AS160 phosphorylation** *Cell Physiol Biochem* **33**:1047–57 [Google Scholar](#)
65. Cui W., Sun Y., Wang Z., Xu C., Xu L., Wang F., Chen Z., Peng Y., Li R. (2011) **Activation of liver X receptor decreases BACE1 expression and activity by reducing membrane cholesterol levels** *Neurochem Res* **36**:1910–21 [Google Scholar](#)
66. Jaye M. C., Krawiec J. A., Campobasso N., Smallwood A., Qiu C., Lu Q., Kerrigan J. J., De Los Frailes Alvaro M., Laffitte B., Liu W. S., Marino J. P., Meyer C. R., Nichols J. A., Parks D. J., Perez P., Sarov-Blat L., Seepersaud S. D., Steplewski K. M., Thompson S. K., Wang P., Watson M. A., Webb C. L., Haigh D., Caravella J. A., Macphee C. H., Willson T. M., Collins J. L. (2005) **Discovery of substituted maleimides as liver X receptor agonists and determination of a ligand-bound crystal structure** *J Med Chem* **48**:5419–22 [Google Scholar](#)
67. Collins J. L., Fivush A. M., Watson M. A., Galardi C. M., Lewis M. C., Moore L. B., Parks D. J., Wilson J. G., Tippin T. K., Binz J. G., Plunket K. D., Morgan D. G., Beaudet E. J., Whitney K. D., Klierer S. A., Willson T. M. (2002) **Identification of a Nonsteroidal Liver X Receptor Agonist through Parallel Array Synthesis of Tertiary Amines** *Journal of Medicinal Chemistry* **45**:1963–1966 [Google Scholar](#)
68. Miao B., Zondlo S., Gibbs S., Cromley D., Hosagrahara V. P., Kirchgessner T. G., Billheimer J., Mukherjee R. (2004) **Raising HDL cholesterol without inducing hepatic steatosis and hypertriglyceridemia by a selective LXR modulator** *J Lipid Res* **45**:1410–7 [Google Scholar](#)
69. Litvinchuk A., Suh J. H., Guo J. L., Lin K., Davis S. S., Bien-Ly N., Tycksen E., Tabor G. T., Remolina Serrano J., Manis M., Bao X., Lee C., Bosch M., Perez E. J., Yuede C. M., Cashikar A. G., Ulrich J. D., Di Paolo G., Holtzman D. M. (2024) **Amelioration of Tau and ApoE4-linked glial lipid accumulation and neurodegeneration with an LXR agonist** *Neuron* **112**:384–403 [Google Scholar](#)
70. Groot P. H., Pearce N. J., Yates J. W., Stocker C., Sauermelch C., Doe C. P., Willette R. N., Olzinski A., Peters T., d'Epagnier D., Morasco K. O., Krawiec J. A., Webb C. L., Aravindhan K., Jucker B., Burgert M., Ma C., Marino J. P., Collins J. L., Macphee C. H., Thompson S. K., Jaye M. (2005) **Synthetic LXR agonists increase LDL in CETP species** *J Lipid Res* **46**:2182–91 [Google Scholar](#)

71. Aravindhan K., Webb C. L., Jaye M., Ghosh A., Willette R. N., DiNardo N. J., Jucker B. M. (2006) **Assessing the effects of LXR agonists on cellular cholesterol handling: a stable isotope tracer study** *J Lipid Res* **47**:1250–60 [Google Scholar](#)
72. Quinet E. M., Savio D. A., Halpern A. R., Chen L., Miller C. P., Nambi P. (2004) **Gene-selective modulation by a synthetic oxysterol ligand of the liver X receptor** *J Lipid Res* **45**:1929–42 [Google Scholar](#)
73. Vieira C. P., Fortmann S. D., Hossain M., Longhini A. L., Hammer S. S., Asare-Bediako B., Crossman D. K., Sielski M. S., Adu-Agyeiwaah Y., Dupont M., Floyd J. L., Li Calzi S., Lydic T., Welner R. S., Blanchard G. J., Busik J. V., Grant M. B. (2020) **Selective LXR agonist DMHCA corrects retinal and bone marrow dysfunction in type 2 diabetes** *JCI Insight* **5** [Google Scholar](#)
74. Wrobel J., Steffan R., Bowen S. M., Magolda R., Matelan E., Unwalla R., Basso M., Clerin V., Gardell S. J., Nambi P., Quinet E., Reminick J. I., Vlasuk G. P., Wang S., Feingold I., Huselton C., Bonn T., Farnegardh M., Hansson T., Nilsson A. G., Wilhelmsson A., Zamaratski E., Evans M. J. (2008) **Indazole-based liver X receptor (LXR) modulators with maintained atherosclerotic lesion reduction activity but diminished stimulation of hepatic triglyceride synthesis** *J Med Chem* **51**:7161–8 [Google Scholar](#)
75. Quinet E. M., Basso M. D., Halpern A. R., Yates D. W., Steffan R. J., Clerin V., Resmini C., Keith J. C., Berrodin T. J., Feingold I., Zhong W., Hartman H. B., Evans M. J., Gardell S. J., DiBlasio-Smith E., Mounts W. M., LaVallie E. R., Wrobel J., Nambi P., Vlasuk G. P. (2009) **LXR ligand lowers LDL cholesterol in primates, is lipid neutral in hamster, and reduces atherosclerosis in mouse** *J Lipid Res* **50**:2358–70 [Google Scholar](#)
76. Katz A., Udata C., Ott E., Hickey L., Burczynski M. E., Burghart P., Vesterqvist O., Meng X. (2009) **Safety, Pharmacokinetics, and Pharmacodynamics of Single Doses of LXR-623, a Novel Liver X-Receptor Agonist, in Healthy Participants** *The Journal of Clinical Pharmacology* **49**:643–649 [Google Scholar](#)
77. Kick E. K., Busch B. B., Martin R., Stevens W. C., Bollu V., Xie Y., Boren B. C., Nyman M. C., Nanao M. H., Nguyen L., Plonowski A., Schulman I. G., Yan G., Zhang H., Hou X., Valente M. N., Narayanan R., Behnia K., Rodrigues A. D., Brock B., Smalley J., Cantor G. H., Lupisella J., Sleph P., Grimm D., Ostrowski J., Wexler R. R., Kirchgessner T., Mohan R. (2016) **Discovery of Highly Potent Liver X Receptor beta Agonists** *ACS Med Chem Lett* **7**:1207–1212 [Google Scholar](#)
78. Stachel S. J., Zerbinatti C., Rudd M. T., Cosden M., Suon S., Nanda K. K., Wessner K., DiMuzio J., Maxwell J., Wu Z., Uslaner J. M., Michener M. S., Szczerba P., Brnardic E., Rada V., Kim Y., Meissner R., Wuelfing P., Yuan Y., Ballard J., Holahan M., Klein D. J., Lu J., Fradera X., Parthasarathy G., Uebele V. N., Chen Z., Li Y., Li J., Cooke A. J., Bennett D. J., Bilodeau M. T., Renger J. (2016) **Identification and in Vivo Evaluation of Liver X Receptor β -Selective Agonists for the Potential Treatment of Alzheimer's Disease** *Journal of Medicinal Chemistry* **59**:3489–3498 [Google Scholar](#)
79. van der Hoorn J., Lindén D., Lindahl U., Bekkers M., Voskuilen M., Nilsson R., Oscarsson J., Lindstedt E., Princen H. (2011) **Low dose of the liver X receptor agonist, AZ876, reduces atherosclerosis in APOE*3Leiden mice without affecting liver or plasma triglyceride levels** *Br J Pharmacol* **162**:1553–1563 [Google Scholar](#)
80. Cannon M. V., Yu H., Candido W. M., Dokter M. M., Lindstedt E.-L., Silljé H. H. W., van Gilst W. H., de Boer R. A. (2015) **The liver X receptor agonist AZ876 protects against pathological**

cardiac hypertrophy and fibrosis without lipogenic side effects *European Journal of Heart Failure* **17**:273–282 [Google Scholar](#)

81. Ben Aissa M., Lewandowski C. T., Ratia K. M., Lee S. H., Layden B. T., LaDu M. J., Thatcher G. R. J. (2021) **Discovery of Nonlipogenic ABCA1 Inducing Compounds with Potential in Alzheimer's Disease and Type 2 Diabetes** *ACS Pharmacol Transl Sci* **4**:143–154 [Google Scholar](#)
82. Lewandowski C. T., Khan M. W., BenAissa M., Dubrovskiy O., Ackerman-Berrier M., LaDu M. J., Layden B. T., Thatcher G. R. J. (2021) **Metabolomic analysis of a selective ABCA1 inducer in obesogenic challenge provides a rationale for therapeutic development** *EBioMedicine* **66**:103287 [Google Scholar](#)
83. Kick E., Martin R., Xie Y., Flatt B., Schweiger E., Wang T.-L., Busch B., Nyman M., Gu X.-H., Yan G., Wagner B., Nanao M., Nguyen L., Stout T., Plonowski A., Schulman I., Ostrowski J., Kirchgessner T., Wexler R., Mohan R. (2015) **Liver X Receptor (LXR) partial agonists: Biaryl pyrazoles and imidazoles displaying a preference for LXR β** *Bioorganic & Medicinal Chemistry Letters* **25**:372–377 [Google Scholar](#)
84. Kirchgessner T. G., Martin R., Sleph P., Grimm D., Liu X., Lupisella J., Smalley J., Narayanan R., Xie Y., Ostrowski J., Cantor G. H., Mohan R., Kick E. (2015) **Pharmacological characterization of a novel liver X receptor agonist with partial LXR α activity and a favorable window in nonhuman primates** *J Pharmacol Exp Ther* **352**:305–14 [Google Scholar](#)
85. Griffett K., Solt L. A., El-Gendy B. E.-D. M., Kamenecka T. M., Burris T. P. (2013) **A Liver-Selective LXR Inverse Agonist That Suppresses Hepatic Steatosis** *ACS Chemical Biology* **8**:559–567 [Google Scholar](#)
86. Griffett K., Welch R. D., Flaveny C. A., Kolar G. R., Neuschwander-Tetri B. A., Burris T. P. (2015) **The LXR inverse agonist SR9238 suppresses fibrosis in a model of non-alcoholic steatohepatitis** *Mol Metab* **4**:353–7 [Google Scholar](#)
87. Elgendy B., Griffett K., Hegazy L., Di Fruscia P., Sample K., Schoepke E., Kamenecka T. M., Burris T. P. (2022) **Synthesis and structure activity relationship of the first class of LXR inverse agonists** *Bioorg Chem* **119**:105540 [Google Scholar](#)
88. Li N., Wang X., Xu Y., Lin Y., Zhu N., Liu P., Lu D., Si S. (2017) **Identification of a Novel Liver X Receptor Agonist that Regulates the Expression of Key Cholesterol Homeostasis Genes with Distinct Pharmacological Characteristics** *Mol Pharmacol* **91**:264–276 [Google Scholar](#)
89. Wardell S. E., Marks J. R., McDonnell D. P. (2011) **The turnover of estrogen receptor alpha by the selective estrogen receptor degrader (SERD) fulvestrant is a saturable process that is not required for antagonist efficacy** *Biochem Pharmacol* **82**:122–30 [Google Scholar](#)
90. Wakeling A. E., Bowler J. (1987) **Steroidal pure antioestrogens** *J Endocrinol* **112**:R7–10 [Google Scholar](#)
91. Rayl M. L., Nemetchek M. D., Voss A. H., Hughes T. S. (2024) **Agonists of the Nuclear Receptor PPARgamma Can Produce Biased Signaling** *Mol Pharmacol* **106**:309–318 [Google Scholar](#)
92. Boergesen M., Pedersen T. A., Gross B., van Heeringen S. J., Hagenbeek D., Bindesboll C., Caron S., Lalloyer F., Steffensen K. R., Nebb H. I., Gustafsson J. A., Stunnenberg H. G., Staels B., Mandrup S. (2012) **Genome-wide profiling of liver X receptor, retinoid X receptor, and peroxisome proliferator-activated receptor alpha in mouse liver reveals extensive sharing of binding sites** *Mol Cell Biol* **32**:852–67 [Google Scholar](#)

93. Griffett K., Burris T. P. (2016) **Promiscuous activity of the LXR antagonist GSK2033 in a mouse model of fatty liver disease** *Biochem Biophys Res Commun* **479**:424–428 [Google Scholar](#)
94. Watanabe M., Houten S. M., Wang L., Moschetta A., Mangelsdorf D. J., Heyman R. A., Moore D. D., Auwerx J. (2004) **Bile acids lower triglyceride levels via a pathway involving FXR, SHP, and SREBP-1c** *J Clin Invest* **113**:1408–18 [Google Scholar](#)
95. Kalaany N. Y., Mangelsdorf D. J. (2006) **LXRS and FXR: the yin and yang of cholesterol and fat metabolism** *Annual review of physiology* **68**:159–91 [Google Scholar](#)
96. Chinetti G., Lestavel S., Bocher V., Remaley A. T., Neve B., Torra I. P., Teissier E., Minnich A., Jaye M., Duverger N., Brewer H. B., Fruchart J. C., Clavey V., Staels B. (2001) **PPAR-alpha and PPAR-gamma activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway** *Nat Med* **7**:53–8 [Google Scholar](#)
97. Chawla A., Boisvert W. A., Lee C. H., Laffitte B. A., Barak Y., Joseph S. B., Liao D., Nagy L., Edwards P. A., Curtiss L. K., Evans R. M., Tontonoz P. (2001) **A PPAR gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis** *Mol Cell* **7**:161–71 [Google Scholar](#)
98. Wang X., Luo J., Li N., Liu L., Han X., Liu C., Zuo X., Jiang X., Li Y., Xu Y., Si S. (2018) **E3317 promotes cholesterol efflux in macrophage cells via enhancing ABCA1 expression** *Biochem Biophys Res Commun* **504**:68–74 [Google Scholar](#)
99. Chandra S., Pahan K. (2019) **Gemfibrozil, a Lipid-Lowering Drug, Lowers Amyloid Plaque Pathology and Enhances Memory in a Mouse Model of Alzheimer's Disease via Peroxisome Proliferator-Activated Receptor alpha** *J Alzheimers Dis Rep* **3**:149–168 [Google Scholar](#)
100. Oliver W. R., Shenk J. L., Snaith M. R., Russell C. S., Plunket K. D., Bodkin N. L., Lewis M. C., Winegar D. A., Sznaidman M. L., Lambert M. H., Xu H. E., Sternbach D. D., Kliewer S. A., Hansen B. C., Willson T. M. (2001) **A selective peroxisome proliferator-activated receptor delta agonist promotes reverse cholesterol transport** *Proc Natl Acad Sci U S A* **98**:5306–11 [Google Scholar](#)
101. Chamberlain S., Gabriel H., Strittmatter W., Didsbury J. (2020) **An Exploratory Phase IIa Study of the PPAR delta/gamma Agonist T3D-959 Assessing Metabolic and Cognitive Function in Subjects with Mild to Moderate Alzheimer's Disease** *J Alzheimers Dis* **73**:1085–1103 [Google Scholar](#)
102. Huang P., Chandra V., Rastinejad F. (2014) **Retinoic Acid Actions through Mammalian Nuclear Receptors** *Chemical Reviews* **114**:233–254 [Google Scholar](#)
103. Dollé P. (2009) **Developmental expression of retinoic acid receptors (RARs)** *Nucl Recept Signal* **7**:e006 [Google Scholar](#)
104. Giguère V., Yang N., Segui P., Evans R. M. (1988) **Identification of a new class of steroid hormone receptors** *Nature* **331**:91–94 [Google Scholar](#)
105. Zhang H., LeCulyse E., Liu L., Hu M., Matoney L., Zhu W., Yan B. (1999) **Rat Pregnane X Receptor: Molecular Cloning, Tissue Distribution, and Xenobiotic Regulation** *Archives of Biochemistry and Biophysics* **368**:14–22 [Google Scholar](#)

106. Watkins R. E., Wisely G. B., Moore L. B., Collins J. L., Lambert M. H., Williams S. P., Willson T. M., Kliewer S. A., Redinbo M. R. (2001) **The human nuclear xenobiotic receptor PXR: structural determinants of directed promiscuity** *Science* **292**:2329–33 [Google Scholar](#)
107. Fuhr U. (2000) **Induction of drug metabolising enzymes: pharmacokinetic and toxicological consequences in humans** *Clin Pharmacokinet* **38**:493–504 [Google Scholar](#)
108. Ding P., Chen Z., Chen H., Zhang Z., Liu Z., Yan X., Zhou H., Gu Q., Li C., Xu J. (2019) **Structurally Selective Mechanism of Liver X Receptor Ligand: In Silico and In Vitro Studies** *J Chem Inf Model* **59**:3277–3290 [Google Scholar](#)
109. Brown M. S., Goldstein J. L. (1997) **The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor** *Cell* **89**:331–40 [Google Scholar](#)
110. Wu D. Y., Ou C. Y., Chodankar R., Siegmund K. D., Stallcup M. R. (2014) **Distinct, genome-wide, gene-specific selectivity patterns of four glucocorticoid receptor coregulators** *Nucl Recept Signal* **12**:e002 [Google Scholar](#)
111. Stallcup M. R., Poulard C. (2020) **Gene-Specific Actions of Transcriptional Coregulators Facilitate Physiological Plasticity: Evidence for a Physiological Coregulator Code** *Trends Biochem Sci* **45**:497–510 [Google Scholar](#)
112. Fondell J. D., Ge H., Roeder R. G. (1996) **Ligand induction of a transcriptionally active thyroid hormone receptor coactivator complex** *Proc Natl Acad Sci U S A* **93**:8329–33 [Google Scholar](#)
113. Aranda A., Pascual A. (2001) **Nuclear hormone receptors and gene expression** *Physiol Rev* **81**:1269–304 [Google Scholar](#)
114. Zhang Y., Breevoort S. R., Angdisen J., Fu M., Schmidt D. R., Holmstrom S. R., Kliewer S. A., Mangelsdorf D. J., Schulman I. G. (2012) **Liver LXR α expression is crucial for whole body cholesterol homeostasis and reverse cholesterol transport in mice** *J Clin Invest* **122**:1688–99 [Google Scholar](#)
115. Korach-André M., Archer A., Gabbi C., Barros R. P., Pedrelli M., Steffensen K. R., Pettersson A. T., Laurencikienė J., Parini P., Gustafsson J. (2011) **Liver X receptors regulate de novo lipogenesis in a tissue-specific manner in C57BL/6 female mice** *Am J Physiol Endocrinol Metab* **301**:E210–22 [Google Scholar](#)
116. Quinet E. M., Savio D. A., Halpern A. R., Chen L., Schuster G. U., Gustafsson J. A., Basso M. D., Nambi P. (2006) **Liver X receptor (LXR)-beta regulation in LXRalpha-deficient mice: implications for therapeutic targeting** *Mol Pharmacol* **70**:1340–9 [Google Scholar](#)
117. Ramon-Vazquez A., de la Rosa J. V., Tabraue C., Lopez F., Diaz-Chico B. N., Bosca L., Tontonoz P., Alemany S., Castrillo A. (2019) **Common and Differential Transcriptional Actions of Nuclear Receptors Liver X Receptors alpha and beta in Macrophages** *Mol Cell Biol* **39** [Google Scholar](#)
118. Bideyan L., Fan W., Kaczor-Urbanowicz K. E., Priest C., Casero D., Tontonoz P. (2022) **Integrative analysis reveals multiple modes of LXR transcriptional regulation in liver** *Proc Natl Acad Sci U S A* **119** [Google Scholar](#)
119. Watson P. J., Fairall L., Schwabe J. W. (2012) **Nuclear hormone receptor co-repressors: structure and function** *Mol Cell Endocrinol* **348**:440–9 [Google Scholar](#)

120. Heery D. M., Kalkhoven E., Hoare S., Parker M. G. (1997) **A signature motif in transcriptional co-activators mediates binding to nuclear receptors** *Nature* **387**:733–736 [Google Scholar](#)
121. Nagy L., Kao H. Y., Love J. D., Li C., Banayo E., Gooch J. T., Krishna V., Chatterjee K., Evans R. M., Schwabe J. W. (1999) **Mechanism of corepressor binding and release from nuclear hormone receptors** *Genes Dev* **13**:3209–16 [Google Scholar](#)
122. Hörlein A. J., Näär A. M., Heinzel T., Torchia J., Gloss B., Kurokawa R., Ryan A., Kamei Y., Söderström M., Glass C. K., Rosenfeld M. G. (1995) **Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor** *Nature* **377**:397–404 [Google Scholar](#)
123. Hu X., Lazar M. A. (1999) **The CoRNR motif controls the recruitment of corepressors by nuclear hormone receptors** *Nature* **402**:93–96 [Google Scholar](#)
124. Nemetchek M. D., Chrisman I. M., Rayl M. L., Voss A. H., Hughes T. S. (2022) **A structural mechanism of nuclear receptor biased agonism** *Proc Natl Acad Sci U S A* **119**:e2215333119 [Google Scholar](#)
125. Kick E., Martin R., Xie Y., Flatt B., Schweiger E., Wang T. L., Busch B., Nyman M., Gu X. H., Yan G., Wagner B., Nanao M., Nguyen L., Stout T., Plonowski A., Schulman I., Ostrowski J., Kirchgessner T., Wexler R., Mohan R. (2015) **Liver X receptor (LXR) partial agonists: biaryl pyrazoles and imidazoles displaying a preference for LXRBeta** *Bioorg Med Chem Lett* **25**:372–7 [Google Scholar](#)
126. Wright M. B., Varona Santos J., Kemmer C., Maugeais C., Carralot J. P., Roeveer S., Molina J., Ducasa G. M., Mitrofanova A., Sloan A., Ahmad A., Pedigo C., Ge M., Pressly J., Barisoni L., Mendez A., Sgrignani J., Cavalli A., Merscher S., Prunotto M., Fornoni A. (2021) **Compounds targeting OSBPL7 increase ABCA1-dependent cholesterol efflux preserving kidney function in two models of kidney disease** *Nature communications* **12**:4662 [Google Scholar](#)
127. Mirza Z., Al-Saedi D. A., Saddeek S., Almowallad S., AlMassabi R. F., Huwait E. (2023) **Atheroprotective Effect of Fucoidan in THP-1 Macrophages by Potential Upregulation of ABCA1** *Biomedicines* **11** [Google Scholar](#)
128. Wang Y., Liu Y., Wang Y., Wu Y., Chen Z., Wang F., Wan X., Wang F., Sun X. (2023) **Macrophage Sult2b1 promotes pathological neovascularization in age-related macular degeneration** *Life Sci Alliance* **6** [Google Scholar](#)
129. Velma G. R., Laham M. S., Lewandowski C., Valencia-Olvera A. C., Balu D., Moore A., Ackerman-Berrier M., Rychetsky P., Penton C., Musku S. R., Annadurai A., Sulaiman M. I., Ma N., Thatcher G.R.J. (2024) **Nonlipogenic ABCA1 Inducers (NLAI) for Alzheimer’s Disease Validated in a Mouse Model Expressing Human APOE3/APOE4** *J Med Chem* **67**:15061–15079 [Google Scholar](#)

Author information

Megan S Laham

Department of Chemistry & Biochemistry, Colleges of Science & Medicine, University of Arizona, Tucson, United States
 ORCID iD: [0000-0002-9803-3323](https://orcid.org/0000-0002-9803-3323)

Martha Ackerman-Berrier

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Fahmida Alam

Department of Chemistry & Biochemistry, Colleges of Science & Medicine, University of Arizona, Tucson, United States

Sarah Turner

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Ganga Reddy Velma

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Christopher Penton

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States, Arizona Center for Drug Discovery, University of Arizona, Tucson, United States

Soumya Reddy Musku

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Manan Rana

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Senthil Kumar

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Anandhan Annadurai

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Maha Ibrahim Sulaiman

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Nina Ma

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States

Gregory RJ Thatcher

Department of Pharmacology & Toxicology, R Ken Coit College of Pharmacy, University of Arizona, Tucson, United States, Department of Chemistry & Biochemistry, Colleges of Science & Medicine, University of Arizona, Tucson, United States
ORCID iD: [0000-0002-7757-1739](https://orcid.org/0000-0002-7757-1739)

For correspondence: grjthatcher@arizona.edu

Editors

Reviewing Editor

Warren Andrew Andayi

Murang'a University of Technology, Murang'a, Kenya

Senior Editor

Amy Andreotti

Iowa State University, Ames, United States of America

Reviewer #1 (Public review):

Summary:

This important study functionally profiled ligands targeting the LXR nuclear receptors using biochemical assays in order to classify ligands according to pharmacological functions. Overall, the evidence is solid, but nuances in the reconstituted biochemical assays and cellular studies and terminology of ligand pharmacology limit the potential impact of the study. This work will be of interest to scientists interested in nuclear receptor pharmacology.

Strengths:

- (1) The authors rigorously tested their ligand set in CRTs for several nuclear receptors that could display ligand-dependent cross-talk with LXR cellular signaling and found that all compounds display LXR selectivity when used at $\sim 1 \mu\text{M}$.
- (2) The authors tested the ligand set for selectivity against two LXR isoforms (alpha and beta). Most compounds were found to be LXRbeta-specific.
- (3) The authors performed extensive LXR CRTs, performed correlation analysis to cellular transcription and gene expression, and classification profiling using heatmap analysis-seeking to use relatively easy-to-collect biochemical assays with purified ligand-binding domain (LBD) protein to explain the complex activity of full-length LXR-mediated transcription.

Weaknesses:

- (1) The descriptions of some observations lack detail, which limits understanding of some key concepts.
- (2) The presence of endogenous NR ligands within cells may confound the correlation of ligand activity of cellular assays to biochemical assay data.
- (3) The normalization of biochemical assay data could confound the classification of graded activity ligands.
- (4) The presence of >1 coregulator peptide in the biplex ($n=2$ peptides) CRT (pCRT) format will bias the LBD conformation towards the peptide-bound form with the highest binding affinity, which will impact potency and interpretation of TR-FRET data.
- (5) Correlation graphical plots lack sufficient statistical testing.
- (6) Some of the proposed ligand pharmacology nomenclature is not clear and deviates from classifications used currently in the field (e.g., hard and soft antagonist; weak vs. partial agonist, definition of an inverse agonist that is not the opposite function to an agonist).

Reviewer #2 (Public review):**Summary:**

In this manuscript by Laham and co-workers, the authors profiled structurally diverse LXR ligands via a coregulator TR-FRET (CRT) assay for their ability to recruit coactivators and kick off corepressors, while identifying coregulator preference and LXR isoform selectivity.

The relative ligand potencies measured via CRT for the two LXR isoforms were correlated with ABCA1 induction or lipogenic activation of SRE, depending on cellular contexts (i.e, astrocytoma or hepatocarcinoma cells). While these correlations are interesting, there is some leeway to improve the quantitative presentation of these correlations. Finally, the CRT signatures were correlated with the structural stabilization of the LXR: coregulator complexes. In aggregate, this study curated a set of LXR ligands with disparate agonism signatures that may guide the design of future nonlipogenic LXR agonists with potential therapeutic applications for cardiovascular disease, Alzheimer's, and type 2 diabetes, without inducing mechanisms that promote fat/lipid production.

Strengths:

This study has many strengths, from curating an excellent LXR compound set to the thoughtful design of the CRT and cellular assays. The design of a multiplexed precision CRT (pCRT) assay that detects corepressor displacement as a function of ligand-induced coactivator recruitment is quite impressive, as it allows measurement of ligand potencies to displace corepressors in the presence of coactivators, which cannot be achieved in a regular CRT assay that looks at coactivator recruitment and corepressor dissociation in separate experiments.

Weaknesses:

I did not identify any major weaknesses.

<https://doi.org/10.7554/eLife.109146.1.sa0>