

Interplay between cohesin and TORC1 links chromosome segregation and gene expression to environmental changes

Reviewed Preprint

v1 • November 18, 2025

Not revised

Dorian Besson, Sabine Vaur, Stéphanie Vazquez, Sylvie Tournier, Yannick Gachet, Adrien Birot, Stéphane Claverol, Adèle Marston, Anastasios Damdimopoulos, Karl Ekwall, Jean-Paul Javerzat 

CNRS, Université de Bordeaux - Institut de Biochimie et Génétique Cellulaires, UMR5095 Bordeaux, France • MCD, Centre de Biologie Intégrative, Université de Toulouse, CNRS, UPS, Toulouse, France • Univ. Bordeaux, Bordeaux Proteome, Bordeaux, France • Centre for Cell Biology, Institute of Cell Biology, University of Edinburgh, Edinburgh, United Kingdom • Department of Medicine, Karolinska Institutet, Stockholm, Sweden

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study describes a new link between nutrient signaling and chromosome regulation, providing **compelling** evidence that reduced activity in the central nutrient-sensing pathway governed by TORC1 improves chromosome stability and alters gene expression in *S. pombe* through effects on cohesin. While the biological importance of this newly described circuit is not yet fully known, and some data would benefit from further clarification, the overall body of evidence supports the main conclusions.

<https://doi.org/10.7554/eLife.108275.1.sa2>

Abstract

Cohesin is a DNA tethering complex essential for chromosome structure and function. In fission yeast, defects in the cohesin loader Mis4 result in chromosome segregation defects and dysregulated expression of genes near chromosome ends. A genetic screen for suppressors of the thermosensitive growth defect of *mis4-G1487D* identified several hypomorphic mutants of the Target of Rapamycin Complex 1 (TORC1), a conserved kinase that integrates cellular signals to regulate growth and metabolism through substrate-specific phosphorylation. Here, we demonstrate that the TORC1 pathway modulates cohesin functions in chromosome segregation and gene expression. In the context of compromised cohesin loading, the incidence of chromosome segregation defects was modulated by the growth medium in a TORC1-dependent manner. Pharmacological or genetic down-regulation of TORC1 activity restored cohesin binding to its chromosomal sites and improved mitotic chromosome segregation. Notably, reduced TORC1 activity also increased cohesin binding and chromosome transmission fidelity in wild-type cells. These results suggest that environmental cues influence chromosome stability via TORC1. Biochemically, TORC1 co-purified with cohesin and reduced TORC1 activity correlated with decreased phosphorylation

of specific residues on Mis4 and cohesin. Mutations in cohesin that mimic the non-phosphorylated state mirrored the effects of TORC1 downregulation, showing that TORC1 is part of the network that controls cohesin phosphorylation to modulate its functions. Finally, we show that the functional interaction between TORC1 and Mis4 extends to the regulation of stress-responsive genes. Our findings reveal a TORC1-cohesin link that may facilitate cellular adaptation to environmental changes. Given that TORC1 inhibitors and calorie restriction extend lifespan in diverse species, this connection raises the intriguing possibility that cohesin-mediated changes in chromosome structure contribute to these effects.

Introduction

The cohesin complex is a pivotal player in genome structure and function. Its activity is critical for a variety of biological processes, including sister chromatid cohesion, nuclear division, DNA replication and repair, and gene expression. Cohesin is an ATP-powered molecular machine that is capable of capturing DNA. Intra-chromosomal DNA capture folds the interphase genome into loops, which contributes to gene regulation, particularly during development and cell fate decisions. Conversely, DNA capture in trans provides cohesion between sister chromatids, which is essential for chromosome segregation and repair (Zheng & Xie, 2019 [DOI](#); Davidson & Peters, 2021 [DOI](#); Oldenkamp & Rowland, 2022 [DOI](#)). The core cohesin complex consists of two Structural Maintenance of Chromosome proteins, Smc1 and Smc3 (Psm1 and Psm3 in the fission yeast *Schizosaccharomyces pombe*) whose ATPase heads are bridged by a kleisin subunit (Rad21/Scc1) to which a fourth subunit (hSTAG1-2, spPsc3, scScc3) binds. DNA capture by cohesin requires the so-called loading complex NIPBL/MAU2 (spMis4/Ssl3, scScc2/Scc4) which binds to cohesin and DNA. Inactivation of the cohesin loading machinery before S phase results in the failure of sister-chromatid cohesion establishment and aberrant chromosome segregation during the ensuing mitosis (Furuya *et al*, 1998 [DOI](#); Ciosk *et al*, 2000 [DOI](#); Bernard *et al*, 2006 [DOI](#)). Similarly, DNA loop extrusion is dependent on the NIPBL-Cohesin holocomplex (Davidson *et al*, 2019 [DOI](#)). Chromatin loops are dynamic structures that form and dissolve on a time-scale of minutes. In human, the DNA binding factor CTCF appears to delineate loop domains, often referred to as topologically associated domains (TADs). Dynamically extruding cohesin complexes are thought to facilitate enhancer-promoter scanning *in cis*. TAD boundaries (insulation) appear regulated as the strengths of CTCF-anchored loop domains are enhanced when mouse embryonic stem cells exit pluripotency. Consistently, mutations in the genes encoding CTCF and cohesin components are linked to human diseases and developmental abnormalities (Zheng & Xie, 2019 [DOI](#); Davidson & Peters, 2021 [DOI](#); Oldenkamp & Rowland, 2022 [DOI](#)). The remodelling of genome architecture appears particularly important for appropriate gene expression during cell fate decisions, suggesting that signalling pathways should convey cellular cues to cohesin. In this context, a genetic screen in mammalian cells identified a set of kinases that alter chromosome folding when inactivated (Park *et al*, 2023 [DOI](#)).

Here we reveal a connection between fission yeast cohesin, its loader Mis4 and the Target of Rapamycin Complex 1 (TORC1). TORC1 is a master regulator of cell growth and metabolism, highly conserved in eukaryotes. The kinase activity of TORC1 is stimulated by a variety of intra- and extracellular signals, including nutrients, growth factors, hormones and cellular energy levels. Once activated, TORC1 promotes cell growth and metabolism through the phosphorylation of multiple effectors (González & Hall, 2017 [DOI](#); Otsubo *et al*, 2017 [DOI](#)). In mammalian species, growth factors and cellular energy stimulate the activity of mTORC1 through the Rheb GTPase, and the inhibition of the tuberous sclerosis complex (TSC), which functions as a GTPase-activating protein for Rheb. In response to amino acid availability, mTORC1 is activated via RAG GTPases in a TSC-independent manner. Under nutrient-rich conditions, TORC1 promotes anabolic processes, such as protein, nucleotide, and lipid synthesis, while inhibiting catabolic processes, such as autophagy.

In fission yeast, Tor2 provides the catalytic activity of TORC1, while Mip1, the Raptor equivalent, participates in substrate recognition (Morozumi *et al*, 2021 [↗](#)). The complex contains three additional subunits, the mLST8 orthologue Wat1/Pop3, Toc1 and Tco89 (Hayashi *et al*, 2007 [↗](#)). TORC1 plays a crucial role in switching between cell proliferation and differentiation by sensing nitrogen source. When deprived of nitrogen, fission yeast cells arrest in G1, mate and undergo meiosis. In the absence of a mating partner, G1 cells enter a G0, quiescent state. Like its mammalian counterpart, the Rheb GTPase Rbh1 is an essential activator of TORC1. Upon nitrogen deprivation, TORC1 activity is restrained by the Gtr1-Gtr2 GTPases, the TSC complex, and the Gcn2 kinase (van Slegtenhorst *et al*, 2004 [↗](#); Chia *et al*, 2017 [↗](#); Fukuda & Shiozaki, 2018 [↗](#); Fukuda *et al*, 2021 [↗](#)). In conditions of abundant nutrients, TORC1 activity is high but nevertheless attenuated by the Gtr1-Gtr2 heterodimer, which is analogous to the mammalian RAG GTPases (Chia *et al*, 2017 [↗](#); Fukuda and Shiozaki, 2018 [↗](#)). TORC1 is essential for cell growth, and loss of the TORC1 activity results in cell cycle arrest in G1. Rapamycin only partially inhibits TORC1 activity in *S. pombe* and does not inhibit growth (Otsubo *et al*, 2017 [↗](#)).

The link between cohesin and TORC1 emerged from a genetic screen for mutants able to suppress the thermosensitive growth (Ts) defect of *mis4-G1487D*, which is defective for cohesin loading and chromosome segregation at the restrictive temperature (Biro *et al*, 2020 [↗](#)). In principle the screen had the potential to identify regulators of Mis4, with the rationale that loss of a negative regulator should upregulate residual Mis4^{G1487D} activity and restore growth at the restrictive temperature. Alongside the cyclin-dependent kinase (CDK) Pef1, a known negative regulator of Mis4 (Biro *et al*, 2020 [↗](#)), several mutants of TORC1 were recovered.

In this study, we explore the connection between cohesin and TORC1. Mis4 plays a crucial role in loading cohesin onto chromosomes and establishing sister chromatid cohesion during the S phase of the cell cycle. Cohesin loading is reduced in the *mis4-G1487D* mutant at the non-permissive temperature, resulting in defective sister-chromatid cohesion and chromosome segregation defects during mitosis. The downregulation of TORC1 enhanced cohesin binding to Cohesin-Associated Regions (CARs) and reduced chromosome segregation defects. Likewise, the incidence of chromosome segregation defects was modulated by the growth medium in a TORC1-dependent manner. These observations suggested that the TORC1 complex may control cohesin dynamics. Therefore, we investigated the impact of reducing TORC1 activity on wild-type cells. Indeed, the binding of cohesin to CAR increased. Furthermore, this was accompanied by an increase in the fidelity of chromosome transmission during vegetative growth. These findings raise the intriguing possibility that environmental cues could influence the robustness of chromosome segregation processes.

Affinity purification experiments showed that TORC1 and cohesin co-purified from protein extracts. The phosphorylation level of specific residues on cohesin and its loader was reduced in TORC1 mutants. Phospho-mutants recapitulated most of the effects observed when TORC1 was downregulated, arguing that TORC1 signalling contributes to the phosphorylation status of cohesin and thereby to its regulation.

Finally, we asked whether the TORC1-cohesin link may extend to the regulation of gene expression. Previous work from our group had shown that Mis4 plays a role in controlling gene expression in the sub-telomeric regions of chromosomes (Dheur *et al*, 2011 [↗](#)). Since TORC1 is involved in environmental adaptation, we performed transcriptome profiling of Mis4 and TORC1 mutants under different culture conditions. The analysis confirmed that Mis4 affects the expression of a small number of genes with a localisation bias away from centromeres and close to telomeres. Interestingly, the identity and number of affected genes differed widely between experimental conditions, suggesting a defective adaptive response. Combining all experiments, 338 genes were differentially expressed in *mis4-G1487D* compared to wild type. Remarkably,

almost all of these genes were also regulated by TORC1. In many cases, the genes were deregulated in the same way and without additive effect, suggesting that Mis4 and TORC1 act in the same pathway.

Collectively, the data presented indicate that TORC1 influences the association of cohesin complexes with chromosomes, regulating their functions in chromosome segregation during mitosis and in the transcriptional response to environmental changes.

Results

A genetic screen for suppressors of *mis4-G1487D* identified components of TORC1

The G1487D substitution in the cohesin loader Mis4 renders the strain thermosensitive for growth (Ts, **Figure 1A**). At the restrictive temperature the amount of cohesin associated with chromosomes is reduced and cells exhibit defects in chromosome segregation during mitosis (Briot *et al*, 2020). To identify putative regulators of Mis4, we previously carried out a genetic screen for suppressors of the Ts phenotype of *mis4-G1487D* (Briot *et al*, 2020). One of the suppressors identified was *pef1*, a CDK whose activity restrains cohesin binding to Cohesin Associated Regions (CARs). Besides *pef1*, the genetic screen identified three other genes: *tor2* (1 allele), *mip1* (5 alleles) and *caa1* (1 allele). The suppressive effect of the *caa1* mutant was modest (**Figure 1—figure supplement 1**). On the other hand, *tor2* and *mip1* mutants clearly suppressed the Ts phenotype of *mis4-G1487D*, at a level similar to that conferred by the deletion of the *pef1* gene (**Figure 1A** and **Figure 1—figure supplement 1**).

The *caa1* gene encodes an aspartate aminotransferase that is required for full activation of the TORC1 complex (Reidman *et al*, 2019). Tor2 and Mip1 are components of TORC1 (Hayashi *et al*, 2007), **Figure 1B**). Both genes are essential for cell survival. The *tor2* allele recovered in our genetic screen bears a substitution in the kinase domain which affected colony growth suggesting that TORC1 activity is largely affected. The five Mip1/Raptor mutants did not show this phenotype, the growth being similar to that of a wild strain. Still they all show reduced phosphorylation of the S6-kinase Psk1 (**Figure 1C**), a well-known TORC1 substrate (Nakashima *et al*, 2012). Likewise, the *mip1-Y533A* mutant which is deficient for Psk1 binding and phosphorylation (Morozumi *et al*, 2021), suppressed the Ts phenotype of *mis4-G1487D* (**Figure 1C**). In a similar manner, the pharmacological downregulation of TORC1 by rapamycin resulted in enhanced growth of *mis4-G1487D* at the restrictive temperature (**Figure 1D**). Other thermosensitive mutants of the cohesin pathway (*eso1-H17* and *rad21-K1*) were not rescued ruling out a general suppressing effect. Hence, the suppression of *mis4-G1487D* may stem from reduced TORC1 kinase activity. Conversely, the genetic upregulation of TORC1 exacerbated the Ts phenotype of *mis4-G1487D*. TORC1 activity upon nitrogen deprivation is restrained by the Gtr1-Gtr2 GTPases, the TSC complex, and the Gcn2 kinase. Among these, the Gtr1-Gtr2 GTPases dampen TORC1 even in nutrient replete conditions (**Figure 1E**, (van Slegtenhorst *et al*, 2004; Chia *et al*, 2017; Fukuda & Shiozaki, 2018; Fukuda *et al*, 2021)). The deletion of *gcn2* or *tsc2* was essentially neutral in terms of *mis4-G1487D* suppression, although the suppression by *mip1-R401G* was reduced in a *tsc2*-deleted background. Deletion of *iml1*, *gtr1* or the *gtr1-Q61L* allele (mimicking the GTP-bound inactive state of the Gtr1 GTPase, (Chia *et al*, 2017)) exacerbated the Ts phenotype of *mis4-G1487D* while mimicking the active GDP-bound form by using the *gtr1-16N* allele, (Chia *et al*, 2017) was neutral. The deletion of *tor1*, encoding the catalytic subunit of the related complex TORC2 exacerbated the Ts phenotype of the *mis4* mutant (**Figure 1—figure supplement 1**). As TORC1 kinase activity is upregulated when *tor1* is deleted (Ikai *et al*, 2011), this observation is in line with the notion that *mis4-G1487D* is sensitive to hyperactive TORC1.




Figure 1.

Genetic evidence linking TORC1 to cohesin.

(A) Model structure of Mis4. The G1487D substitution confers a thermosensitive growth (Ts) phenotype. The cell growth assay shows the suppression of the Ts phenotype by the indicated mutations. **(B)** Composition of fission yeast TORC complexes (adapted from Hayashi et al, 2007). **(C-D)** Down-regulation of TORC1 suppresses the Ts growth defect of *mis4-G1487D*. **(C)** Model structure of Mip1 highlighting amino-acid substitutions in the suppressor mutants (RNC: Raptor N-terminal CASPase-like domain, ARM: Armadillo-type fold, WD40: WD-40 repeats). All *mip1* mutant alleles from the genetic screen showed reduced phosphorylation of the S6-kinase Psk1, a known TORC1 substrate. Conversely, *mip1-Y533A* which is deficient for Psk1 binding and phosphorylation, suppressed the Ts phenotype of *mis4-G1487D*. **(D)** The TORC1 inhibitor rapamycin rescued *mis4-G1487D* but not *eso1* and *rad21* mutants. **(E)** Genetic upregulation of TORC1 exacerbates the Ts phenotype of *mis4-G1487D*. The deletion of *gcn2* or *tsc2* was essentially neutral, although the suppression by *mip1-R401G* was reduced in a *tsc2* deleted background. Deletion of *gtr1*, *iml1* or *gtr1-Q61L* mimicking the GTP-bound, inactive state of GATOR exacerbated the Ts phenotype of *mis4-G1487D* while mimicking the active GDP-bound form (*gtr1-S16N*) was neutral. **(F)** *psm3^{K105N-K106N}* confers a Ts phenotype when combined with the deletion of *pph3*, encoding the catalytic subunit of PP4, a phenotype efficiently rescued by *mip1-R401G*.

The above genetic analyses show that the Ts phenotype of the *mis4* mutant is dampened when TORC1 activity is down-regulated and exacerbated when TORC1 is up-regulated. Because *mip1* alleles barely affected growth on their own and were excellent *mis4* suppressors we focused our analyses on one of them, *mip1-R401G*. We found that *mip1-R401G* acted as suppressor in another genetic context of impaired cohesin loading. The acetyl-mimicking forms of Psm3 have been demonstrated to impede the loading of cohesin (Murayama & Uhlmann, 2015 [↗](#); Hu *et al*, 2015 [↗](#); Birot *et al*, 2020 [↗](#)). This phenotype is exacerbated by the deletion of *pph3*, encoding the catalytic subunit of Protein Phosphatase 4 and the double mutant strain is Ts for growth (Birot *et al*, 2020 [↗](#)). The *mip1-R401G* mutant efficiently suppressed the Ts phenotype of *psm3-K105NK106N pph3Δ* (Figure 1F [↗](#)), indicating that the suppression is not restricted to *mis4-G1487D* but appears generally related to cohesin loading.

The occurrence of chromosome segregation defects in *mis4-G1487D* is modulated by TORC1 and growth conditions

When *mis4-G1487D* cells were shifted to the restrictive temperature cell proliferation was reduced and viability dropped ~25 fold after 24 hours when compared to wild-type. The *mip1-R401G* mutation efficiently suppressed the proliferation and viability defects (Figure 2A [↗](#)). A hallmark of cohesin mutants is the occurrence of abnormal mitoses in which the spindle apparatus attempts to segregate un-cohered sister chromatids. A single kinetochore is captured by microtubules emanating from opposite spindle poles (merotelic attachment) and moves back and forth on the anaphase spindle as a result of opposite forces (Pidoux *et al*, 2000 [↗](#); Gregan *et al*, 2011 [↗](#)). Live cell analysis confirmed the occurrence of merotelic kinetochores in the *mis4* mutant (Figure 2—figure supplement 1 [↗](#)). In fixed cells, a merotelically-attached chromatid appears as DAPI-stained material lagging on the anaphase spindle. Anaphases with lagging DNA were frequent for *mis4-G1487D* after one doubling of the cell population at the restrictive temperature and their occurrence was significantly reduced by *mip1-R401G* (Figure 2B [↗](#)). Rapamycin exposure during a single cell cycle also reduced the frequency of anaphases with lagging DNA, although the effect was weaker (Figure 2C [↗](#)). This is consistent with the notion that the down-regulation of TORC1 reduced chromosome segregation defects of the *mis4* mutant. The suppressor effect of the *mip1-R401G* mutation was not restricted to the *mis4* mutant. The frequency of anaphases with lagging DNA was efficiently reduced in the *psm3K105N-K106N pph3Δ* background (Figure 2D [↗](#)). As for *mis4-G1487D*, cohesin loading is impaired in this genetic setup (Birot *et al*, 2020 [↗](#)) but *Mis4* is wild-type. This shows that the suppression is not allele-specific, but may instead stem from an upregulation of the cohesin loading machinery.

Merotelic kinetochores oppose spindle forces and reduce the rate of spindle elongation during anaphase. As the number of merotelic kinetochores increases, the spindle elongation rate slows down and may even become negative resulting in spindle shrinkage (Courtheoux *et al*, 2009 [↗](#)). The rate of spindle elongation was strongly affected in *mis4-G1487D* and spindle shrinkage events were frequent, indicative of a high level of merotelic attachment. Spindle shrinkage occurred in ~53% of anaphases in *mis4-G1487D* whereas it was reduced to ~21% in the *mip1-R401G* background (Figure 2E [↗](#) and Figure 2—figure supplement 2 [↗](#)). Particularly striking was the observation of spindle shrinkage followed by asymmetric spindle displacement resulting in the generation of anucleate daughter cells after cytokinesis (Figure 2F [↗](#)). Nearly 80% of cells showed this phenotype in *mis4-G1487D* for the late time points whereas the frequency was reduced to ~20% in the presence of *mip1-R401G*. Altogether, these analyses indicate that the *mis4* mutant experienced numerous merotelic events during anaphase at the restrictive temperature, leading to asymmetric spindle shrinkage and the production of anucleate cells. The suppression of these phenotypes by *mip1-R401G* is consistent with improved sister-chromatid cohesion, leading to a lower incidence of merotelic attachments and chromosome segregation defects. At centromeres, defective cohesion results in increased distance between sister-kinetochores, a phenotype that can be evidenced by microscopy. When early mitotic cells are abruptly cooled on ice, microtubules disassemble leaving unattached kinetochore pairs (Gachet *et al*, 2008 [↗](#)). In some instances a kinetochore pair drifts




Figure 2.

The incidence of chromosome segregation defects in *mis4-G1487D* is modulated by TORC1 and the growth medium.

(A) Exponentially growing cells at 25°C were shifted at 36.5°C for 24 hours. The growth curve shows that *mip1-R401G* suppressed the *mis4-G1487D* temperature growth defect. Cell survival after the 24 hours temperature shift was addressed by plating serial 5-fold cell dilutions at permissive temperature for *mis4-G1487D* (4×10^4 cells in the first row). Cell survival was reduced ~25 fold in *mis4-G1487D* and restored to wild-type levels when combined with *mip1-R401G*. (B) Exponentially growing cells at 25°C were shifted at 36°C for one doubling. DNA was stained with DAPI (red, pseudo-colour) and tubulin detected by indirect immunofluorescence (green). Lagging DNA (white arrow) was defined as DAPI-stained material on the anaphase spindle (length > 5 μm). (C) Rapamycin reduces the incidence of anaphases with lagging DNA. Rapamycin (RAPA, 200 ng/ml) or solvent alone (DMSO) was added to cycling cells at 25°C and the cultures shifted to 36°C for one doubling of the cell population. Samples were treated as in B. (D) The *mip1* mutant reduced the frequency of anaphases with lagging DNA in the *psm3^{K105-K106N} pph3Δ* background. Cells were treated as in B. (E) Spindle shrinkage events (shrinkage of more than 0.5 μm) were detected by live analysis of cells undergoing anaphase at 36.5°C (Fig. 2 supplement 2). (F) Anaphase cells with displaced spindle. Cells were cultured at 36.5°C, fixed and stained with Calcofluor to visualize septa (blue). Shown are mean and SD from three independent experiments (n > 65 cells per strain and per experiment). (G) Measurement of inter-kinetochore distance. Cells were shifted to 36.5°C for 5 hours and subsequently chilled at 4°C for 30 minutes. Cells were fixed, DNA stained with DAPI and the distance between Mis6 signals was measured. The images on the top show scattered kinetochore pairs in the *mis4-G1487D* mutant. Bottom, graph showing the distribution of inter-kinetochore distances (wt, n = 46; *mis4-G1487D*, n = 48; *mis4-G1487D mip1-R401G*, n = 48). (H) Top, cell growth assay showing that the Ts growth defect of *mis4-G1487D* is more severe in YES+A medium than in EMM2. Bottom, the frequency of anaphases with lagging DNA is higher in YES+A than in EMM2. The inactive form of the TORC1 inhibitor Gtr1 (Gtr1-Q61L) increased the frequency of abnormal anaphases while the active form was neutral (Gtr1-S16N). Cells were treated as in B. (*) (**) (***), P-values from two-sided Fisher's exact test.

away from the other chromosomes, allowing the measurement of inter-kinetochore distance. The inter-kinetochore distance increased in the *mis4* mutant compared to the wild type. Importantly, the defect was reduced in the *mip1-R401G* background (**Figure 2G** [↗](#)). The suppression was nevertheless partial, suggesting sister-centromere cohesion was not fully restored. Using a GFP-tagged *ade6* locus on the right arm of chromosome III, we found that sister-chromatid cohesion remained defective at that site (**Figure 2—figure supplement 3** [↗](#)). The level of Psm3 acetylation, a marker of sister-chromatid cohesion establishment, was low in the *mis4* mutant and was not restored by *mip1-R401G* or rapamycin. These data suggest that sister chromatid cohesion remains defective overall, but has improved sufficiently, particularly at the centromeres, to enable more effective chromosome segregation during mitosis.

Physiologically TORC1 acts as a rheostat, its activity rising with nutrient availability. YES is a rich undefined medium containing numerous sources of nitrogen, whereas ammonia is the sole source of nitrogen in EMM2. The Ts phenotype of *mis4-G1487D* was stronger and the frequency of lagging DNA was higher in the rich, complete medium YES than in the synthetic minimal medium EMM2 (**Figure 2H** [↗](#)). The artificial increase of TORC1 activity by *gtr1-Q61L* resulted in an elevated rate of anaphases with lagging DNA in EMM2 medium (**Figure 2H** [↗](#)). Conversely, the use of glutamate, a nitrogen source slightly poorer than ammonia (Davie et al, 2015 [↗](#)) reduced chromosome segregation defects (**Figure 2—figure supplement 4** [↗](#)). Altogether, these analyses show that the incidence of chromosome segregation defects of *mis4-G1487D* is modulated by growth conditions in a TORC1-dependent manner.

Reduced TORC1 activity increased Mis4 and cohesin binding to Cohesin Associated Regions (CARs)

The cohesin loading complex performs its essential function during G1/S (Furuya et al, 1998 [↗](#); Ciosk et al, 2000 [↗](#); Bernard et al, 2006 [↗](#)). The down-regulation of TORC1 may rescue *mis4-G1487D* by up-regulating its residual cohesin loading activity. To address this issue, cells were arrested at the end of the G1 phase by the use of the *cdc10-129* Ts mutation and cohesin binding to chromosomes monitored by Rad21 ChIP-qPCR at known CARs (**Figure 3A** [↗](#)). Cycling *S. pombe* cells are essentially in the G2 phase of the cell cycle (Carlson et al, 1999 [↗](#)). When shifted at 36.5°C, *cdc10-129* cells progress through G2, M and arrest in G1. As reported previously, Rad21 binding to CARs was reduced in a *mis4-G1487D* background (Feytout et al, 2011 [↗](#); Vaur et al, 2012 [↗](#); Birot et al, 2020 [↗](#)). The *mip1-R401G* mutation efficiently restored cohesin binding to near wild-type levels (**Figure 3B** [↗](#)). A similar effect was observed in the *psm3^{K105N-K106N} pph3Δ* background (**Figure 3—figure supplement 1** [↗](#)) indicating that *mip1-R401G* restores cohesin loading in these two distinct genetic contexts. Importantly, a similar stimulating effect was observed in cells with a wild-type cohesin loading machinery. In an otherwise wild-type background, Mis4 and Rad21 binding to CARs were increased in *mip1-R401G* cells at most chromosomal sites examined (**Figure 3C-D** [↗](#)). Rapamycin treatment during a single cell cycle provoked a similar effect although less pronounced (**Figure 3—figure supplement 1** [↗](#)). We conclude that TORC1 down-regulation increases cohesin abundance at CARs, presumably by stimulating cohesin loading.

TORC1 components co-purify with cohesin and its loader

To see whether TORC1 and cohesin may interact we affinity-purified Mis4 and Rad21 complexes and analysed associated proteins by label-free mass spectrometry (**Figure 4A** [↗](#)). To avoid cell cycle effects, extracts were prepared from G1 (*cdc10-129*) arrested cells. All TORC1 components co-purified with Mis4 and Rad21 and conversely, additional co-immunoprecipitation experiments showed that cohesin co-purified with TORC1 (**Figure 4B-C** [↗](#)).




Figure 3.

Increased cohesin binding to Cohesin Associated Regions in the raptor mutant *mip1-R401G*.

(A) Chromosomal sites examined were Cohesin Associated Regions along the arms (438, 1806, 2898 and 3323) and centromere of chromosome 2, the non-transcribed spacer region (NTS) of the rDNA gene cluster on chromosome 3 and the chromosome 1 right telomere. Within the centromere, the central core (cc2) which is the site of kinetochore assembly, the imr and dg repeats that flank the central core on either side and at tRNA rich domains that delineate the centromere. (B) The *mip1-R401G* mutation restores cohesin binding in *mis4-G1487D* cells. Cohesin binding to chromatin was monitored by Rad21-FLAG ChIP in G1 cells (*cdc10-129* arrest) at 36.5°C. Error bars = SD from 4 ChIPs, except for *mis4-G1487D* (3 ChIPs). *** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$ by two-tailed, unpaired t-test with 95% confidence interval. (C-D) The *mip1-R401G* mutation increased Mis4 and Rad21 binding in *mis4⁺* cells. ChIP assays were made with *cdc10-129* arrested cells. (C) Mean ratios $\pm$ SD were calculated from 2 independent experiments and 4 technical replicates per experiment. (D) Mean ratios $\pm$ SD were calculated from 4 independent experiments and 4 technical replicates per experiment. *** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, by two-tailed, one sample t-test with 95% confidence interval.




Figure 4.

TORC1 components co-purify with Rad21 and Mis4.

(A) Affinity purifications (triplicate samples) were made from G1-arrested cells (*cdc10-129* arrest) and proteins analysed by label-free mass spectrometry. (B-C) Reciprocal co-immunoprecipitation experiments. Western blots were probed with the indicated antibodies. (B) Rad21-PK IP (*cdc10-129* arrested cells). (C) Tor2-FLAG IP (cycling cells).

TORC1 signalling affects the phosphorylation levels of Mis4-S183 and Psm1-S1022

To see whether TORC1 activity may affect cohesin or Mis4 phosphorylation status, Mis4 and Rad21 complexes were purified from *mip1-R401G* and *mip1*⁺ cells and triplicate samples analysed by label-free mass spectrometry. As before, protein extracts were prepared from G1-arrested cells to avoid cell cycle induced changes. A serine residue (S183) within the N terminal domain of Mis4 was less phosphorylated in *mip1-R401G* compared to wild-type. Likewise, Psm1-S1022 phosphorylation level was decreased (**Figure 5A** [↗](#)). Antibodies were raised against the phosphorylated forms (**Figure 5—figure supplement 1-2** [↗](#)) and this confirmed that the levels of Mis4-S183p and Psm1-S1022p were reduced in *mip1-R401G* cells (**Figure 5BC** [↗](#)). A similar reduction was also apparent in the unrelated mutant *mip1-Y533A* (**Figure 5—figure supplement 1** [↗](#)). Rapamycin exposure during one complete cell cycle led to a reduction of Psm1-S1022 phosphorylation (**Figure 5D** [↗](#)). Mis4-S183 behaved differently as its phosphorylation was increased by rapamycin (**Figure 5E** [↗](#)). Altogether, these data show that the phosphorylation levels Mis4-S183 and Psm1-S1022 are responsive to alterations in TORC1-signalling.

Both phosphorylation sites on Psm1 and Mis4 adhere to the CDK consensus (S-T/P) but were not described as direct targets of Cdc2, the main fission yeast CDK (Swaffer *et al*, 2018 [↗](#)). Label-free mass spectrometry indicated that the phosphorylation of Mis4-S183 was not dependent on the CDK Pef1 either (**Figure 5—figure supplement 2** [↗](#)). However, Psm1-S1022 phosphorylation was strongly reduced in *pef1* deleted cells (**Figure 5—figure supplement 2** [↗](#)). Pef1 was reported as a positive regulator of TORC1 (Matsuda *et al*, 2020 [↗](#)), suggesting Pef1 may promote the phosphorylation of Psm1 through TORC1 activation. Indeed, the phosphorylation level of the TORC1 substrate Psk1 was reduced in *pef1* deleted cells and the double mutant *pef1Δ mip1-R401G* grew very poorly (**Figure 5—figure supplement 2** [↗](#)), consistent with further reduced TORC1 activity. The phosphorylation of Psm1 is therefore possibly dependent on the activation of TORC1 by Pef1. Conversely, Pef1 activity does not seem to depend on TORC1 as the phosphorylation status of Rad21-T262, which is a Pef1 substrate (Biot *et al*, 2020 [↗](#)), was not affected by *mip1-R401G* (**Figure 5—figure supplement 2** [↗](#)).

TORC1 may regulate cohesin phosphorylation directly, or via the activation of downstream kinases. In fission yeast there are three known AGC kinases phosphorylated and thereby presumably activated by TORC1: Psk1, Sck1 and Sck2 (Nakashima *et al*, 2012 [↗](#); Chica *et al*, 2016 [↗](#)). The ribosomal S6 kinase Psk1 was introduced earlier. The closest homologs of Sck1-2 are SGK1-3 (Serum/glucocorticoid regulated kinase 1, 2 and 3) in human and SCH9 in budding yeast. AGC kinase genes are dispensable for growth in fission yeast, either alone or in combination, which rendered possible a straightforward analysis.

Deletion of the *psk1* gene was essentially neutral for *mis4-G1487D* (**Figure 5—figure supplement 3** [↗](#)). The deletion of *sck1* had a weak suppressor effect whereas *sck2* deletion suppressed the growth defect to a level similar to *mip1-R401G*. The double deletion of *sck1* and *sck2* had the same effect as the single deletion of *sck2*. Using Rad21-FLAG ChIP-qPCR on G1-arrested cells we determined the relative enrichment of cohesin localised at CARs in the mutant strains versus wild-type (**Figure 5—figure supplement 3** [↗](#)). A small decrease was observed for *sck1Δ* at several CARs, such as 438, tRNA-L, cc2, imr2-L and 1806. The *sck2Δ* single mutant behaved quite similar to wild-type, at least in G1-arrested cells. In contrast, the *sck1Δ sck2Δ* double mutant showed an increase relative to wild-type at almost all loci examined and at a level similar to *mip1-R401G*. This indicates a synthetic effect of the simultaneous deletion of *sck1* and *sck2* on cohesin association at CARs in G1-arrested cells. The frequency of chromosome segregation defects of *mis4-G1487D* was markedly reduced in a *sck2*-deleted background and further decreased by the additional deletion of *sck1* (**Figure 5—figure supplement 3** [↗](#)).




Figure 5.

TORC1 signalling affects the phosphorylation levels of Psm1-S1022 and Mis4-S183.

(A) Mis4-GFP and Cohesin (Rad21-PK) were affinity-purified from G1-arrested cells (*cdc10-129* arrest). Triplicate samples were analyzed by label-free mass spectrometry. Mis4-S183 and Psm1-S1022 phosphorylation levels were reduced in the *mip1-R401G* background. (B) Cohesin was purified by Rad21-FLAG immunoprecipitation from *cdc10-129* arrested cells and probed with anti-Psm1-S1022p and anti-Psm1 antibodies. Mean ratios $\pm$ SD from three experiments. (C) Mis4-GFP was affinity-purified from *cdc10-129* arrested cells and probed with anti-Mis4-S183p and anti-GFP antibodies. Mean ratios $\pm$ SD from three experiments. (D-E) Effect of rapamycin on the phosphorylation status of Psm1-S1022 and Mis4-S183. Rapamycin (RAPA, 200ng/ml) or solvent alone (DMSO) was added to cycling cells at 25°C and the cultures shifted to 36°C for one doubling of the cell population. (C) Mis4-GFP was affinity purified and probed with anti-Mis4-S183p and anti-GFP antibodies. Mean ratios $\pm$ SD from two experiments. (D) Cohesin was purified by Rad21-FLAG immunoprecipitation and probed with anti-Psm1-S1022p and anti-Psm1 antibodies. Mean ratios $\pm$ SD from two experiments

The phosphorylation level of Psm1-S1022 was similar to wild-type in the *sck1Δ* and *sck2Δ* single mutants. The *sck1Δ sck2Δ* double mutant showed a 33% decrease in phosphorylation (**Figure 5—figure supplement 3** [↗](#)). These results suggest that *sck1* and *sck2* are redundant for the regulation of Psm1-S1022 phosphorylation. The phosphorylation level of Mis4-S183 was increased 9-fold in *sck1Δ* compared to wild-type. Conversely, Mis4-S183p was reduced in the *sck2Δ* strain (3.7 fold decrease relative to wild-type) and the double mutant showed a 5-fold higher signal than the wild-type (**Figure 5—figure supplement 3** [↗](#)). As Mis4-S183 is phosphorylated in the double mutant this indicates that Mis4-S183 is phosphorylated by another, unknown kinase, at least in this genetic context. In this scenario, the unknown kinase would be positively regulated by Sck2 and negatively by Sck1.

The interplay between TORC1, Pef1 and other kinases downstream of TORC1 appears complex and remains to be elucidated. Nevertheless, the above data show that TORC1 signalling affects the phosphorylation level of Mis4 and Psm1, the abundance of cohesin at CARs and the suppression of *mis4-G1487D* phenotypes.

Mis4-S183 and Psm1-S1022 are relevant targets for the regulation of cohesin by TORC1

The *mip1-R401G* mutant is a strong suppressor of *mis4-G1487D* phenotypes and the phosphorylation levels of Mis4-S183 and Psm1-S1022 are reduced. To address the causality of the phosphorylation status, we generated mutants mimicking the non-phosphorylated (S to A) or phosphorylated state (S to E/D). Non-phosphorylatable *mis4* and *psm1* mutants rescued *mis4-G1487D* growth and chromosome segregation defects while phospho-mimetics had the opposite effect (**Figure 6A-D** [↗](#)). Cohesin binding to CARs as assayed by ChIP tend to increase for the mutants mimicking the non-phosphorylated state and to decrease with the phospho-mimicking forms (**Figure 6E** [↗](#)). Likewise, non-phosphorylatable *mis4* and *psm1* mutants efficiently rescued growth and chromosome segregation defects in the *psm3^{K105N-K106N} pph3Δ* background and ChIP analyses revealed that cohesin binding to CARs was clearly stimulated (**Figure 6—figure supplement 1** [↗](#)). These data show that the phenotypes of *mip1-R401G* are largely recapitulated by mutants mimicking the non-phosphorylated state, arguing that Mis4 and Psm1 are relevant downstream effectors for cohesin regulation by the TOR signalling pathway.

Rapamycin increased the fidelity of chromosome transmission of wild-type cells in a pathway involving Psm1 phosphorylation

The data we have obtained so far show that reducing TORC1 activity reduced the severity of chromosome segregation defects in genetic contexts where cohesin loading is compromised. We therefore wondered whether the alteration of TORC1 activity could affect the fidelity of chromosome transmission in the wild type. The loss frequency of whole, natural chromosomes can be estimated by measuring the breakdown of diploids (Allshire *et al.*, 1995 [↗](#)). Remarkably, diploids were more stable when grown in the presence of rapamycin when compared to solvent alone (**Figure 7A** [↗](#)). We also measured the loss frequency of CM3112, a circular, non-essential minichromosome derived from the centromere region of chromosome III (Matsumoto *et al.*, 1990 [↗](#)). In the wild type, it is lost at a rate of around 1 per 100 divisions. Again, rapamycin led to a substantial decrease in the incidence of CM3112 loss (**Figure 7B** [↗](#)). The frequency of CM3112 loss was reduced in the *mip1-R401G* mutant when compared to wild-type. Furthermore, the combination of rapamycin and *mip1-R401G* did not result in an additive effect, consistent with rapamycin and *mip1-R401G* acting in the same pathway. In a similar manner, *psm1-S1022A* showed minimal responsiveness to rapamycin, while *psm1-S1022D* exhibited a discernible response. The involvement of Mis4-S183 phosphorylation was less clear. The *mis4-S183A* mutant did show a reduced response to rapamycin (**Figure 7—figure supplement 1** [↗](#)). However, the phosphomimetic allele *mis4-S183E* did not restore the response to rapamycin, and the double




Figure 6.

Mimicking the non phosphorylated state of Psm1-S1022 and Mis4-S183 alleviates *mis4-G1487D* phenotypes.

(A) Cell growth assays. (B) Anaphases with lagging DNA. Exponentially growing cells at 25°C were shifted at 36°C for 3.5hrs (one doubling). DNA was stained with DAPI and tubulin detected by indirect immunofluorescence. Lagging DNA was defined as DAPI-stained material on the anaphase spindle (length>5μm). P-values from two-sided Fisher's exact test. (C) Anaphase cells with displaced spindle. Cells were cultured at 36°C, fixed and stained with Calcofluor to visualize septa. Mean and SD from three independent experiments (n>70 cells per strain and per experiment). P-value from two-sided Fisher's exact test. (D) Inter-kinetochore distance. Cells were treated as in **Figure 2G** (n=51 for each mutant). Data from **Figure 2G** are shown for comparison (dotted lines). (E) Rad21 ChIP. Exponentially growing cells at 25°C were shifted at 36°C for 3.5hrs (one doubling). Mean and SD from 6 ratios and 2 independent experiments. **P ≤ 0.01, *P ≤ 0.05, by two-tailed, one sample t-test with 95% confidence interval.

mutants did not reveal a clear genetic interaction, which complicated the interpretation. Taken together, these data indicate that rapamycin increases the stability of chromosomes through a pathway involving Mip1 and the phosphorylation of the cohesin subunit Psm1.

Mis4 and TORC1 control a common set of genes involved in the response to environmental changes

In a previous study, gene expression profiling of *mis4-G1487D* identified a small number of upregulated and downregulated genes within subtelomeric domains (Dheur *et al.*, 2011 [↗](#)). The genetic link between TORC1 and cohesin prompted us to ask how the two pathways may interact in terms of gene expression. As TORC1 adapts cells to changes in their environment, we looked at the transcriptome by RNA-sequencing of wild-type, *mis4-G1487D*, *mip1-R401G* and the double mutant in various experimental conditions (**Figure 8A** [↗](#)). These included cycling cells in the rich complete medium YES at 25°C (V25) and upon a temperature shift to 36.5°C for one doubling (V36.5) or arrested in G1 (*cdc10-129*) at this temperature (G1_36.5). We also looked upon nitrogen starvation by shifting actively cycling cells from the synthetic medium EMM2 to EMM2 deprived of a nitrogen source (EMM2-N). In this situation, TORC1 activity is restrained, cells arrest in G1 and either mate to enter the reproductive cycle (meiosis) or enter a G0, quiescent state. As no mating partner was available in our experiments, cells homogeneously arrested in G1 after nitrogen deprivation (**Figure 8—figure supplement 1** [↗](#)). RNA-sequencing was performed 24 hours after the shift to EMM2-N (T0-N). One half of the remaining culture was shifted to 36.5°C for 4 days (T4D36.5) while the other half was kept at 25°C for 4 days (T4D25). The analysis of the results is shown in **Figure 8** [↗](#), and the complete data set is given in Supplementary file 2.

As in the previous study (Dheur *et al.*, 2011 [↗](#)), a small number of genes were found to be differentially expressed in the *mis4-G1487D* mutant compared to the wild type (**Figure 8A** [↗](#)). In actively growing cells in rich medium 12 genes were affected, 50 after one cell doubling at 36.5°C and 27 in G1-arrested cells at 36.5°C. The number of differentially expressed genes increased upon nitrogen deprivation (T0-N, 118 genes, T4D25, 125 genes). A similar number of genes were affected in *mis4-G1487D* when nitrogen depleted cells were shifted to 36.5°C for 4 days (T4D36.5, 76 genes). A much larger number of genes were differentially expressed in *mip1-R401G* versus wild-type (**Figure 8A** [↗](#)). As illustrated by the Venn diagrams, a significant proportion of *mis4*-affected genes were also affected in the *mip1* mutant.

Most *mis4*-affected genes were specific to a particular experimental condition (**Figure 8B** [↗](#)). For example, of the genes that are up-regulated in the *mis4-G1487D* mutant compared to the wild-type, 184 are specific to a single experimental condition, 24 are up-regulated in two experiments, two are up-regulated in three experiments, and two are up-regulated in four experiments. A similar trend is seen for down-regulated genes. Thus, most of the *Mis4*-sensitive genes are only sensitive under specific culture conditions. These observations suggest that *mis4-G1487D* affects the transcriptional response to environment changes.

In aggregate, 338 genes were differentially expressed in *mis4-G1487D* relative to wild-type across all experiments (**Figure 8B** [↗](#)). The analysis of gene lists with AnGeLi (Bitton *et al.*, 2015 [↗](#)) indicated that *mis4*-regulated genes showed a localization bias towards chromosome ends (**Figure 8C** [↗](#) and **Figure 8—figure supplement 2** [↗](#)). The response to nitrogen starvation and differentiation (meiosis), as well as enrichment for “stress genes”, were observed, consistent with a defective adaptive response. Genes encoding membrane transporters were significantly enriched, suggesting a role in cell homeostasis. A strong overlap was apparent with a published set of genes whose expression was upregulated in a *tor2* conditional mutant at the restrictive temperature (**Figure 8—figure supplement 2** [↗](#), (Wei *et al.*, 2021 [↗](#))). A significant overlap was also seen with genes up-regulated in G0 (Zahedi *et al.*, 2023 [↗](#)), suggesting *Mis4* may participate in the proper establishment and/or maintenance of the quiescent state (**Figure 8—figure**




Figure 7.

Rapamycin reduces the frequency of chromosome loss.

(A) The rate of chromosome loss was estimated by measuring the breakdown of diploids after growth in medium containing rapamycin (RAPA, 200 ng/ml) or solvent alone (DMSO). Data are mean $\pm$ SD, $n=2$ per group, $**P < 0.01$, unpaired t-test. **(B)** Loss rate of the CM3112 minichromosome. The loss rate was determined in the presence of rapamycin (RAPA, 200 ng/ml) or solvent alone (DMSO). Data are mean $\pm$ SEM, $n=4$ per group ($n=10$ for the wild-type), $***P < 0.001$, repeated measures two-way ANOVA followed by Bonferroni post-hoc test.




Figure 8.

mis4-G1487D affects gene response to environmental changes.

(A) RNA-sequencing (biological triplicates) was performed on cells growing in rich medium at 25°C (V25), after one cell doubling at 36.5°C (V36.5) and on G1-arrested cells (*cdc10-129*) at 36.5°C (G1_36.5). For nitrogen starvation, cells grown in EMM2 were deprived from nitrogen for 24H at 25°C (T0-N). One half of the culture was shifted to 36.5°C for 4 days (T4D36.5) while the other half was left at 25°C for 4 days (T4D25). The Venn diagrams show the overlap number of differentially expressed genes (FDR<0.05) for each experimental condition. P-value for under- or over-enrichment were calculated based on the cumulative distribution function of the hypergeometric distribution. (B) Differentially expressed genes in *mis4-G1487D* across all experiments. There were 408 hits involving 338 genes. The lower graphs show that most of the affected genes are specific to a single experimental condition. (C) Gene features and GO-term analysis for the 338 *mis4*-regulated genes. (D) Overlap between genes differentially expressed in *mis4-G1487D* and *mip1-R401G* versus wild-type in all experimental conditions. (E) Among the 338 genes differentially expressed in *mis4*, 100 show a similar (+/- 10%) fold change in the *mis4*, *mip1* and *mis4 mip1* mutant strains. See also **Figure 8—figure supplement 6**.

supplement 2 [↗](#)). Cell survival remains high in G0 in the *mis4* and *mip1* mutants (**Figure 8—figure supplement 3-4** [↗](#)) but upon refeeding a delay in G0 exit was noticeable (**Figure 8—figure supplement 5** [↗](#)). This phenotype was quite strong for *mip1-R401G* and more subtle for the *mis4* mutant.

Most (300/338) genes affected by *mis4-G1487D* were also affected by *mip1-R401G* (**Figure 8D** [↗](#)). If we go into more detail, we can identify a large group of genes (100/300) for which the quantitative variations were similar, moving in the same direction, with little or no additive effect in the double mutant (**Figure 8E** [↗](#) and **Figure 8—figure supplement 6** [↗](#)). For this particular set of genes the epistatic relationship suggests that Mis4 and TORC1 may act in the same pathway. In other instances, the changes in gene expression relative to the wild-type were in opposite directions in *mis4* and *mip1* mutants (64 occurrences, 62 genes, **Figure 8—figure supplement 7** [↗](#)). Overall, these data show that Mis4 and TORC1 functionally interact for the regulation of genes involved in the response to environmental changes.

Discussion

The data presented in this work provide evidence for a regulation of cohesin by the TOR signalling pathway. This is, to the best of our knowledge, the first report to demonstrate such a relationship. Previously, a link had been established between the two pathways, but in the opposite direction. mTORC1 is indeed down-regulated in cells from patients with Roberts syndrome, caused by mutational alteration of the ESCO2 cohesin acetyl-transferase (Xu *et al*, 2013 [↗](#)). The down-regulation of the cohesin loader NIPBL in breast cancer cells induced cell cycle arrest, apoptosis and autophagy through the caspase 3 and mTOR signalling pathways (Zhou *et al*, 2017 [↗](#)). The first hint for a regulation of cohesin by the TOR pathway came from the observation in fission yeast that the growth of a conditional mutant of *mis4* was enhanced by rapamycin (Sajiki *et al*, 2018 [↗](#)), although the underlying molecular mechanisms remained unclear. In mammalian species, a fraction of mTOR controls nuclear processes, notably transcription (Zhao *et al*, 2024 [↗](#)). Likewise, fission yeast TORC1 is involved in the regulation of facultative heterochromatin (Wei *et al*, 2021 [↗](#); Hirai *et al*, 2023 [↗](#)) and transcriptional processes (Larabee & Weisman, 2020 [↗](#)). The present work establishes, for the first time, a close link between cohesin and the TOR pathway, suggesting the exciting possibility that extracellular signals could remodel the functional architecture of chromosomes during differentiation or adaptation processes.

Mis4 and Psm1 as downstream targets of TORC1

We present evidence showing that cohesin and TORC1 components co-purify from protein extracts and Mis4-S183 and Psm1-S1022 are less phosphorylated in *mip1* mutants. Non-phosphorylatable mutants recapitulated most of the effects of TORC1 down-regulation while mutants mimicking the phosphorylated state produced phenotypes similar to TORC1 hyper activation. Whether cohesin and Mis4 are substrates of TORC1 or other kinases regulated by TORC1 is currently unknown. Meanwhile, our work provides the first evidence that cohesin is a downstream effector of TORC1. It is worth mentioning that TORC1 components were found to co-purify with the budding yeast cohesin loader (Mattingly *et al*, 2022 [↗](#)), suggesting that the link between cohesin and TORC1 may extend to other species.

Both phosphorylation events on Mis4 and Psm1 have been reported in the literature but their biological functions are unknown (Kettenbach *et al*, 2015 [↗](#); Swaffer *et al*, 2018 [↗](#); Tay *et al*, 2019 [↗](#); Halova *et al*, 2021 [↗](#)). Mis4-S183 is located within an unstructured, flexible region connecting the Ssl3 interaction domain and the hook domain that provides Mis4 catalytic activity. It was proposed that phosphorylation at CDK sites within the unstructured linker may affect the flexibility of the complex and modulate its activity (Chao *et al*, 2015 [↗](#)). Mis4-S183 and the CDK consensus are conserved in hNIPBL and the residue was found phosphorylated in a number of

phosphoproteomic studies (Ruse *et al.*, 2008 [↗](#); Dephoure *et al.*, 2008 [↗](#); Pan *et al.*, 2009 [↗](#); Mayya *et al.*, 2009 [↗](#); Christensen *et al.*, 2010 [↗](#); Rigbolt *et al.*, 2011 [↗](#); Beli *et al.*, 2012 [↗](#)). Cryo-EM structures of cohesin bound to its loader and DNA revealed that the SMC coiled coils are folded around their elbow (Collier *et al.*, 2020 [↗](#); Shi *et al.*, 2020 [↗](#)). The S1022 residue of Psm1 lies close to the “joint” region below the ATPase heads that the “hinge” domain contacts when cohesin is in its folded conformation. Whether the phosphorylation of Psm1-S1022 affects the outcome of DNA transactions remain to be investigated.

TORC1 activity restrains cohesin binding to CARs

Reducing TORC1 activity by the *mip1-R401G* mutation enhanced cohesin binding to CARs. Remarkably, a similar effect was observed within a single cell cycle of rapamycin treatment. Increased cohesin binding at CARs occurred in two genetic contexts of sensitized cohesin loading and with a wild-type cohesin loading machinery. The most straightforward interpretation is that TORC1 restrains cohesin loading onto chromosomes. One distinct possibility would be that TORC1 down-regulation promotes cohesin accumulation at CARs without *de novo* loading. It was proposed that CARs correspond to the borders of yeast TAD-like structures (Costantino *et al.*, 2020 [↗](#)). Increased abundance of cohesin at CARs may result from the accumulation of cohesin with loop extrusion activity at TAD boundaries. Considering that loop extrusion requires the NIPBL-Cohesin holocomplex (Davidson *et al.*, 2019 [↗](#)), it is interesting to note that Mis4 binding to CARs was similarly enhanced. TORC1 may also impinge on cohesin sliding by transcription-related processes (Lengronne *et al.*, 2004 [↗](#); Schmidt *et al.*, 2009 [↗](#)) and / or influence cohesin accumulation through the phosphorylation of chromatin-bound proteins. Future work will aim at clarifying these issues.

TORC1 and chromosome segregation

Cells with an impaired cohesin loading machinery experience chromosome segregation defects during mitosis. We have shown here that the intensity of the defect is dependent on TORC1 activity. The most prominent defect is the merotelic attachment of kinetochores which, when left uncorrected, lead to aberrant chromosome segregation. Although the down-regulation of TORC1 reduced the frequency of aberrant anaphases, sister-chromatid cohesion remained largely impaired, even though cohesin binding to CARs in the *mis4* mutant was restored to near wild-type levels in cells arrested at the G1/S boundary. The increase in chromosome bound cohesin may have enhanced cohesion establishment during S-phase. Indeed the inter-kinetochore distance was reduced but sister-chromatid cohesion remained overall defective at a distal, chromosome arm locus. Consistently, cohesin acetylation was not improved by TORC1 down-regulation. The acetylation of Psm3, mediated by Eso1 in fission yeast, counteracts the cohesin releasing activity of Wapl and is thought to stabilize cohesin binding and cohesion (Feytout *et al.*, 2011 [↗](#); Kagami *et al.*, 2011 [↗](#); Vaur *et al.*, 2012 [↗](#); Birot *et al.*, 2017 [↗](#)). However, deletion of the *wpl1* gene does not rescue the *mis4-G1487D* mutant and *S. pombe* cells can survive with non-detectable acetylation or in a context in which Psm3 cannot be acetylated (Feytout *et al.*, 2011 [↗](#); Birot *et al.*, 2020 [↗](#)). The primary defect of *mis4-G1487D* may therefore stem from inability to generate sister-chromatid cohesion rather than its maintenance. *In vitro* assays have suggested that sister-chromatid capture by cohesin is a two-step process catalysed by Mis4 (Murayama *et al.*, 2018 [↗](#)). The down-regulation of TORC1 may increase the residual loading activity of Mis4 but second DNA capture may remain largely deficient, resulting in poor sister-chromatid cohesion. Increased cohesin binding at peri-centromeres may improve chromosome segregation through other routes. Cohesin organizes the peri-centromere regions in budding yeast (Lawrimore *et al.*, 2018 [↗](#); Paldi *et al.*, 2020 [↗](#)). Local depletion of cohesin at centromeres in fission yeast lead to a high incidence of merotelic attachments (Bernard *et al.*, 2001 [↗](#)). A recent study revealed that vertebrate kinetochores are bipartite structures and highlighted a role for cohesin in bridging to the two parts into a functional unit (Sacristan *et al.*, 2024 [↗](#)). Lastly, enhanced cohesin binding to centromeres may facilitate the

correction of merotelic attachments by the Aurora B kinase. The localization of Aurora B relies on histone H3-T3 phosphorylation by Haspin whose localization to centromeres depends largely on the cohesin subunit Pds5 in fission yeast (Yamagishi *et al*, 2010 [↗](#); Goto *et al*, 2017 [↗](#)).

That chromosome segregation defects increased with TORC1 activity suggests the intriguing possibility that the fidelity of chromosome segregation could be modulated by environmental cues. Chromosome segregation defects in the *mis4* mutant were most intense in the complete rich medium YES than in the synthetic medium EMM2. The use of glutamate, a nitrogen source slightly poorer than ammonia (Davie *et al*, 2015 [↗](#)) further reduced chromosome segregation defects. Furthermore, Rapamycin reduced the frequency of chromosome loss in wild-type cells in a pathway involving the phosphorylation state of Psm1-S1022. It is striking that rapamycin increased the stability of chromosomes during vegetative growth. The idea that nutritional inputs affect the stability of chromosomes could make sense in ecological terms. Microbial species in the wild compete with each other. When nutrients are abundant, it may be preferable to divide in order to colonise the niche rapidly, even at the expense of a few cells. Conversely, when nutrients become scarce, survival of the species may rely more heavily on the fitness of individual cells.

Mis4 and TORC1 control the gene response to environmental changes

In a previous study we reported that *mis4-G1487D* affects the expression of genes located in subtelomeric domains (Dheur *et al*, 2011 [↗](#)). Here we extend the study by challenging cells with various culture conditions. Several conclusions can be drawn. Very few genes were affected when cells were actively cycling in rich media, when TORC1 activity is high. The gene list was much longer upon nitrogen deprivation, when TORC1 activity is repressed, and the list of genes evolved with time spent in G0. In aggregate 338 genes were affected in *mis4-G1487D* when compared with wild-type with a bias towards chromosome ends and far from centromeres. Lastly, most genes misregulated in *mis4-G1487D* were misregulated in the *mip1-R401G* mutant suggesting Mis4 and Mip1 act in a same pathway. Paradoxically, the *mis4* and *mip1* mutants remained viable during G0 although a delay was noticeable upon re-entry into the cell cycle at elevated temperature. A defect in quiescence exit was previously reported when the *mis4* gene is affected and the phenotype appears strongly dependent on the allele considered (Sajiki *et al*, 2009 [↗](#); Suma *et al*, 2024 [↗](#)). It must be kept in mind that *mis4* mutants were originally screened for a defect during vegetative growth. It would be interesting to screen specifically for *mis4* alleles affecting cell survival in quiescence.

Given the role of cohesin and its loader in chromosome folding, it is tempting to speculate that TORC1 might direct cohesin to establish a chromosome architecture conducive to transcriptional responses to environmental changes. If increased binding of Mis4 and cohesin at CARs does reflect extruding cohesin complexes that have reached TAD boundaries, chromosomes might be more compacted when TORC1 is down regulated. Conversely, high TORC1 would promote the opposite. Even if this hypothesis turned out to be correct, it does not explain why regulated genes show a bias towards chromosome ends. Chromosomes in fission yeast adopt a Rabl configuration in interphase (Mizuguchi *et al*, 2015 [↗](#)). Centromeres cluster at the SPBs while telomeres cluster in a few patches at the opposing hemisphere near the nuclear periphery. These constraints generate specific chromosomal regions with distinct molecular environments. Chromosome conformation capture and the mapping of DNA binding sites of inner nuclear membrane proteins have provided evidence of functional sub-nuclear environments that correlate with gene expression activity (Steglich *et al*, 2012 [↗](#); Grand *et al*, 2014 [↗](#)). The studies suggest more internal locations of actively expressed genes, and nuclear peripheral localization of poorly expressed genes. Chromosome regions containing gene clusters that are up-regulated upon nitrogen starvation re-position from the nuclear periphery to the interior (Alfredsson-Timmins *et al*, 2009 [↗](#)). In this regard, Mis4-regulated genes tend to be lowly expressed and enriched for nitrogen responsive genes. Modifying chromosome compaction may alter the Rabl configuration or generate forces that modify the

properties of the inner nuclear envelope, leading to misregulated gene expression. Indeed, loss of cohesin causes an increase in inter-chromosomal contacts and a reduction in chromosomal territoriality (Mizuguchi *et al*, 2014 [↗](#)).

Cohesin and TORC1 have been implicated with a wide range of human pathologies, including cancer (Yoon, 2020 [↗](#); Waldman, 2020 [↗](#)). Rapamycin has attracted much attention for its putative ability to mimic caloric restriction, resulting in improved health and lifespan (Bjedov & Partridge, 2011 [↗](#)). Our study suggests the intriguing possibility that some effects are mediated by the ability of cohesin to remodel chromosomes in response to various extracellular signals, including nutrients.

Materials and methods

Strains, media and genetic techniques

General fission yeast methods, reagents and media are described in (Moreno *et al*, 1991 [↗](#)). All strains are listed in Supplementary file 1. Experiments were carried out using YES medium unless otherwise stated. For cell growth assays serial dilutions of cells were spotted on YES medium and the plates incubated at the indicated temperatures for 3 days unless otherwise stated. EMM-GLU is EMM2 in which NH_4Cl was replaced with 20mM glutamate. Rapamycin (Calbiochem, 4.57mg/ml stock in DMSO) was added to the culture medium to 200ng/ml. For nitrogen deprivation, cycling cells in EMM2 ($\sim 5 \cdot 10^6$ - 10^7 cells/ml) were washed three times in EMM2-N and incubated in EMM2-N at the same density. Cell cycle arrest in YES medium was achieved by shifting cycling *cdc10-129* cells ($\sim 10^7$ cells/ml) at 36°C for 3.5 hours. Hydroxyurea (HU) was used to arrest cells in early S-phase. HU (12mM) was added to cycling cells in YES ($\sim 10^7$ /ml) and shifted to 36.5°C for 3.5 hours. All cell cycle arrests were checked by flow cytometry analysis of DNA content. The genetic screen for *mis4* suppressors was described in (Biro *et al*, 2020 [↗](#)). The suppressors fell into four linkage groups. To identify the mutated loci, genomic DNA was extracted from one mutant from each group and from the wild-type *S. pombe* reference strain and co-hybridized to tiling arrays as described (Biro *et al*, 2020 [↗](#)). For the *mip1* group (5 alleles) the mutated site in each strain was identified by PCR followed by DNA sequencing. Phospho-mutants were generated using the SpEDIT method (Torres-Garcia *et al*, 2020 [↗](#)). The introduction of the desired changes was confirmed by PCR followed by DNA sequencing. Gene mapping, sequence and annotations were from Pombase (Rutherford *et al*, 2024 [↗](#)). Chromosome loss assays were as described (Allshire *et al*, 1995 [↗](#)) using the calculation method of Kipling and Kearsey (Kipling & Kearsey, 1990 [↗](#)). Cells containing the CM3112 minichromosome are Ade⁺ and form white colonies on YES medium. Loss of the minichromosome generates Ade⁻ cells that form red colonies. White colonies grown on YES medium (5-10) were picked, resuspended in 1 ml of YES+Adenine medium, and the cell density was measured. Cells were diluted to 10^4 /ml in YES+Adenine and incubated at 25°C for 3 days. Cells (500-1000) were plated on YES plates at day 0 and day 3 to estimate the proportion of cells containing the minichromosome. The rate of chromosome loss per division was calculated using the formula: loss rate = $1 - (F/I)^{1/N}$, where F is the final percentage of cells bearing the minichromosome, I is the initial percentage of cells carrying the minichromosome, and N is the number of generations between I and F. The same method was applied to measure the breakdown of diploid strains but with two successive rounds of growth in liquid YES+Adenine medium before the final plating.

Cytological techniques

DNA content was measured by flow cytometry with an Accuri C6 Flow cytometer after Sytox Green staining of ethanol-fixed cells (Knutsen *et al*, 2011 [↗](#)). Data were presented using the FlowJo software. Indirect immunofluorescence was done as described (Biro *et al*, 2020 [↗](#)). Cells were imaged using a Leica DMRXA microscope and a 63X objective. Measurements were made using MetaMorph software.

Live cell analysis was performed in an imaging chamber (CoverWell PCI-2.5, Grace Bio-Labs, Bend, OR) filled with 1 ml of 1% agarose in minimal medium and sealed with a 22 × 22-mm glass coverslip. Time-lapse images of Z stacks (maximum five stacks of 0.5 μm steps, to avoid photobleaching) were taken at 60 sec intervals. Images were acquired with a CCD Retiga R6 camera (QImaging) fitted to a DM6B upright microscope with a x63 objective, using MetaMorph as a software. Intensity adjustments were made using the MetaMorph, Image J, and Adobe Photoshop packages (Adobe Systems France, Paris, France). To determine the percentage of mispositioned spindles, cells were fixed in 3.7% formaldehyde for 6 min at room temperature, washed twice in PBS, and observed in the presence of DAPI/calcofluor.

The position of the SPBs, kinetochores and centromeres were determined by visualization of the Cdc11–GFP, Ndc80–GFP and Mis6–RFP signals. Maximum intensity projections were prepared for each time point, with the images from each channel being combined into a single RGB image. These images were cropped around the cell of interest, and optional contrast enhancement was performed in MetaMorph, Image J or Photoshop where necessary. The cropped images were exported as 8-bit RGB-stacked TIFF files, with each frame corresponding to one image of the time-lapse series. For all channels, custom peak detection was performed. The successive positions of the SPBs and kinetochores/centromeres were determined.

For the measurement of inter-kinetochore distance, cells were grown overnight at 25°C, then shifted to 36.5°C for 5 hours and subsequently chilled at 4°C for 30 minutes. At this stage, as previously shown (Gachet *et al*, 2008 [DOI](#)), metaphase spindles collapsed, microtubules were absent, and sister kinetochores were frequently misaligned relative to the two spindle pole bodies (SPBs). After the cold shock, cells were fixed in 3.7% formaldehyde for 6 minutes at room temperature, washed twice in PBS, stained with DAPI and the distance between Mis6–RFP signals was measured.

Antibodies, protein extracts, immunoprecipitation, western blotting, phosphatase treatment, chromatin immunoprecipitation (ChIP)

Rabbit polyclonal antibodies against Rad21, Psm1, Psm3, Psm3-K106Ac, Rad21-T262 have been described previously (Feytout *et al*, 2011 [DOI](#); Dheur *et al*, 2011 [DOI](#); Birot *et al*, 2020 [DOI](#)). The mouse monoclonal anti-tubulin antibody TAT1 is from (Woods *et al*, 1989 [DOI](#)). Anti-Psm1-S1022p and anti-Mis4-S183p antibodies were raised by Biotem (Apprieu, France). Rabbits were immunized with the KLH-coupled peptides. Sera were immune-depleted by affinity with the non-phosphorylated form of the peptide and antibodies were affinity purified against the phosphorylated peptide. Other antibodies were of commercial source. Rabbit polyclonal anti-GFP A11122 (Molecular Probes), mouse monoclonal anti-GFP (Roche), anti-PK antibodies (monoclonal mouse anti V5 tag, AbD serotec), mouse monoclonal anti-FLAG (Sigma) and mouse monoclonal anti-myc (Santa-Cruz). For the detection of Psk1 (Thr-415) phosphorylation, protein extracts were prepared by the TCA method and western blots probed with the anti-phospho-p70 S6K antibody (cat. no. 9206, Cell Signaling Technology) as described (Morozumi *et al*, 2021 [DOI](#)). Protein extracts, immunoprecipitation (IP), cell fractionation and western blotting were as described (Birot *et al*, 2020 [DOI](#)). For quantitative western blot analyses, signals were captured with the ChemiDoc MP Imaging System and quantified using the Image Lab software. On beads phosphatase treatment of Mis4–GFP was done as described (Birot *et al*, 2017 [DOI](#)). Chromatin Immunoprecipitation (ChIP) was as described (Birot *et al*, 2020 [DOI](#)) using anti-FLAG, anti-GFP (A11122) and anti-Rad21 antibodies.

RNA sequencing

Total RNA from biological triplicate samples was subjected to quality control with Agilent Tapestation according to the manufacturer's instructions. To construct libraries suitable for Illumina sequencing, rRNA was depleted using the Ribominus kit (Thermo Fisher) starting with 2000 ng total RNA and then followed by the Illumina stranded ligation sample preparation

protocol starting with 100 ng rRNA depleted RNA. The protocol includes fragmentation, denaturation of RNA, cDNA synthesis, ligation of adapters, and amplification of indexed libraries. The yield and quality of the amplified libraries were analyzed using Qubit (Thermo Fisher) and the quality of the libraries was checked by using the Agilent Tapestation. The indexed cDNA libraries were normalized and combined, and the pools were sequenced on the Illumina Nextseq 2000 machine using a P3 100 cycle sequencing run, producing a sequencing length of 58 base paired end reads with dual index.

RNA-Seq Data processing and analysis

Bcl files were converted and demultiplexed to fastq using the bcl2fastq v2.20.0.422 program. STAR 2.7.10a (Dobin *et al*, 2013 [DOI](#)) was used to index the *S. pombe* reference genome (ASM294v2) and align the resulting fastq files. Mapped reads were then counted in annotated exons using featureCounts v1.5.1 (Liao *et al*, 2014 [DOI](#)). The gene annotations (Schizosaccharomyces_pombe.ASM294v2.35.gff3) and reference genome were obtained from Ensembl Fungi. The count table from featureCounts was imported into R/Bioconductor and differential gene expression was performed using the EdgeR (Robinson *et al*, 2010 [DOI](#)) package and its general linear models pipeline. For the gene expression analysis genes with no or very low expression were filtered out using the filterByExpr function and subsequently normalized using TMM normalization. Genes with an FDR adjusted p value < 0.05 were termed significantly regulated. Venn diagrams were generated using jvenn (Bardou *et al*, 2014 [DOI](#)). The p-values for under- or over-enrichment were calculated based on the cumulative distribution function (CDF) of the hypergeometric distribution. The total number of genes was set to 6403, which corresponds to the total number of genes that were detected across all our experiments.

Mass spectrometry

Sample preparation and protein digestion. Protein samples were solubilized in Laemmli buffer and proteins were deposited onto SDS-PAGE gel. After colloidal blue staining, each lane was cut out from the gel and was subsequently cut in 1 mm x 1 mm gel pieces. Gel pieces were destained in 25 mM ammonium bicarbonate 50% ACN, rinsed twice in ultrapure water and shrunk in ACN for 10 min. After ACN removal, gel pieces were dried at room temperature, covered with the trypsin solution (10 ng/μl in 50 mM NH₄HCO₃), rehydrated at 4°C for 10 min, and finally incubated overnight at 37°C. Spots were then incubated for 15 min in 50 mM NH₄HCO₃ at room temperature with rotary shaking. The supernatant was collected, and an H₂O/ACN/HCOOH (47.5:47.5:5) extraction solution was added onto gel slices for 15 min. The extraction step was repeated twice. Supernatants were pooled and dried in a vacuum centrifuge. Digests were finally solubilized in 0.1% HCOOH.

nLC-MS/MS analysis and Label-Free Quantitative Data Analysis

Peptide mixture was analyzed on a Ultimate 3000 nanoLC system (Dionex, Amsterdam, The Netherlands) coupled to a Electrospray Orbitrap Fusion™ Lumos™ Tribrid™ Mass Spectrometer (Thermo Fisher Scientific, San Jose, CA). Ten microliters of peptide digests were loaded onto a 300-μm-inner diameter x 5-mm C18 PepMap™ trap column (LC Packings) at a flow rate of 10 μL/min. The peptides were eluted from the trap column onto an analytical 75-mm id x 50-cm C18 Pep-Map column (LC Packings) with a 5–27.5% linear gradient of solvent B in 105 min (solvent A was 0.1% formic acid and solvent B was 0.1% formic acid in 80% ACN) followed by a 10 min gradient from 27.5% to 40% solvent B. The separation flow rate was set at 300 nL/min. The mass spectrometer operated in positive ion mode at a 2-kV needle voltage. Data were acquired using Xcalibur software in a data-dependent mode. MS scans (m/z 375–1500) were recorded in the Orbitrap at a resolution of R = 120 000 (@ m/z 200) and an AGC target of 4 x 10⁵ ions collected within 50 ms. Dynamic exclusion was set to 60 s and top speed fragmentation in HCD mode was performed over a 3 s cycle. MS/MS scans were collected in the Orbitrap with a resolution of 30,000 and maximum fill time of 54 ms. Only +2 to +7 charged ions were selected for fragmentation. Other settings were

as follows: no sheath nor auxiliary gas flow, heated capillary temperature, 275 °C; normalized HCD collision energy of 30%, isolation width of 1.6 m/z, AGC target of 5×10^4 and normalized AGC target of 100%. Monoisotopic precursor selection (MIPS) was set to Peptide and an intensity threshold was set to 2.5×10^4 .

Database search and results processing

Data were searched by SEQUEST and BYONIC through Proteome Discoverer 2.5 (Thermo Fisher Scientific Inc.) against an *Schizosaccharomyces pombe* uniprot database (5,098 entries in v2021-01). Spectra from peptides higher than 5000 Da or lower than 350 Da were rejected. Precursor Detector node was included. Search parameters were as follows: mass accuracy of the monoisotopic peptide precursor and peptide fragments was set to 10 ppm and 0.02 Da respectively. Only b- and y-ions were considered for mass calculation. Sequest HT was used as the search algorithm : Oxidation of methionines (+16 Da), methionine loss (-131 Da), methionine loss with acetylation (-89 Da), protein N-terminal acetylation (+42 Da) and phosphorylation of Serine, Threonine and Tyrosine (+80 Da) were considered as variable modifications while carbamidomethylation of cysteines (+57 Da) was considered as fixed modification. Two missed trypsin cleavages were allowed. Peptide validation was performed using Percolator algorithm (Kall *et al*, 2007 [\[4\]](#)) and only “high confidence” peptides were retained corresponding to a 1% False Positive Rate at peptide level. Peaks were detected and integrated using the Minora algorithm embedded in Proteome Discoverer. Normalization was performed based on total peptide amount. Protein ratios were calculated as the median of all possible pairwise peptide ratios. A t-test was calculated based on background population of peptides or proteins. Quantitative data were considered at peptide level.

Figure supplements

Figure 1—figure supplement 1.

(A) Cell growth assay showing the suppression of the Ts phenotype of *mis4-G1487D* by the indicated mutations. Plates were incubated for 6 days at the indicated temperatures. The *caa1* mutant is slow growing and 6 days were required to see a weak suppressor effect. (B) The deletion of *tor1*, encoding the catalytic subunit of TORC2, shows a negative genetic interaction with *mis4-G1487D*.





Figure 2—figure supplement 1.

Live analysis of *mis4-G1487D* undergoing anaphase at 36.5°C.

A merotelic kinetochore is evidenced by the stretching of the outer kinetochore protein Ndc80 (line scan time 28 to 36 min).






Figure 2—figure supplement 2.

Live analysis of anaphase cells revealed spindle shrinkage events.

(A) Example of a *mis4-G1487D* cell showing spindle shrinkage during anaphase. (B) As shown in [Figure 2E](#) the proportion of mitotic cells with the occurrence of spindle shrinkage ($>0.5\mu\text{m}$; red curves) is reduced in the *mip1-R401G* background (wt, $n=30$; *mis4-G1487D*, $n=34$; *mis4-G1487D mip1-R401G*, $n=34$).

Figure 2—figure supplement 3.

(A) Sister-chromatid cohesion assay at the *ade6* locus.

Exponentially growing cells at 25°C were shifted at 36°C for one doubling. Fixed cells were treated for detection of GFP dots and tubulin by indirect immunofluorescence. The sample image on the left shows *mis4-G1487D* cells. DNA appears in blue, tubulin in red and the GFP-marked *ade6* locus in green. Separated GFP dots (>0.5µm) were scored in G2 cells as judged by the presence of the interphase array of microtubules and a single nucleus. P-value from two-sided Fisher's exact test. (B-C) Psm3-K106 acetylation levels. (B) Exponentially growing cells at 25°C were shifted at 36.5°C. HU (12mM) was added at the time of the shift and cells incubated for 3.5 hours. The level of Psm3-K106 acetylation was determined by western blotting. Cell cycle arrest in early S-phase was checked by DNA-content analysis (Right). (C) Rapamycin (200ng/ml) or solvent alone (DMSO) was added to exponentially growing cells at 25°C and the cultures shifted to 36.5°C for one doubling of the cell population. Protein extracts were analyzed by western blotting using antibodies detecting phosphorylated Psk1 and the cohesin core subunits Rad21, Psm1 and Psm3. The acetylated form of Psm3 was detected with anti Psm3-K106ac antibodies.




Figure 2—figure supplement 4.

The frequency of anaphases with lagging DNA is dependent on the culture medium.

Exponentially growing *mis4-G1487D* cells at 25°C in the indicated media were shifted at 36.5°C for one doubling of the cell population. DNA was stained with DAPI and tubulin detected by indirect immunofluorescence. Lagging DNA was defined as DAPI-stained material on the anaphase spindle (length>5µm). EMM-GLU is EMM2 in which NH₄Cl was replaced by 20 mM glutamate. (*) P-values from two-sided Fisher's exact test.




Figure 3—figure supplement 1.

(A) The *mip1-R401G* mutation increases cohesin binding to Cohesin Associated Regions in *pph3Δ psm3^{K105N-K106N}* cells. Exponentially growing cells at 25°C were shifted to 36°C for one doubling of the cell population. Cohesin binding to chromatin was monitored by ChIP using anti-Rad21 antibodies. Mean and SD from 6 ChIPs. **(B)** Effect of rapamycin treatment on cohesin binding. Exponentially growing cells at 25°C were shifted to 36°C for one doubling of the cell population. Rapamycin (200ng/ml) or solvent alone (dmsO) was added at the time of the temperature shift. Cohesin binding to chromatin was monitored by ChIP using anti-Rad21 antibodies. Mean and SD from 4 ChIPs. ***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05, by two-tailed, one sample t-test with 95% confidence interval.





Figure 5—figure supplement 1.

(A) Phosphatase assay showing the specificity of anti-Mis4-S183p antibodies. Mis4-GFP was affinity-purified from *cdc10-*arrested cells. Phosphatase treatment was performed in the presence or absence of phosphatase inhibitors and proteins analyzed by western blotting with the indicated antibodies. **(B)** Mis4-GFP immunoprecipitations made from cycling cells. Western-blot were probed with anti-Mis4-S183p and anti-GFP antibodies. **(C)** Rad21-FLAG immunoprecipitations made from G1 cells (*cdc10-129* arrest). Western blots were probed with anti-Psm1-S1022p and anti-Psm1 antibodies.






Figure 5—figure supplement 2.

(A) The phosphorylation of Mis4-S183 is not dependent on Pef1. Mis4-GFP was affinity-purified from *pef1⁺* and *pef1Δ* cells (*cdc10-129* arrest) and triplicate samples analyzed by label-free mass spectrometry. (B-C) The phosphorylation of Psm1-S1022 is dependent on Pef1. (B) Rad21-PK was affinity-purified (triplicate samples) from *pef1⁺* and *pef1Δ* cells (*cdc10-129* arrest) and proteins were analyzed by label-free mass spectrometry (C) Rad21-FLAG was affinity-purified from *pef1⁺* and *pef1Δ* cells (*cdc10-129* arrest) and proteins analyzed by western blotting using the indicated antibodies. Consistent with the mass spectrometry analyses, the anti-Psm1-S1022p antibodies did react with Psm1 of wild-type but not from *pef1Δ* cells. (D) Psk1 phosphorylation is reduced in *pef1Δ*. TCA extracts from cycling cells were probed by western blotting with the indicated antibodies. (E) Cell growth assay showing the negative genetic interaction between *pef1Δ* and *mip1-R401G*.




Figure 5—figure supplement 3.

(A) Cell growth assay showing the genetic interaction between *mis4-G1487D* and null mutants of AGC kinase genes. (B) Cohesin binding to chromatin was monitored by Rad21-FLAG ChIP in G1 cells (*cdc10-129* arrest). Mean ratios and SD were calculated from 2 independent experiments and 4 technical replicates per experiment. *** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, by two-tailed, one sample t-test with 95% confidence interval. (C) Exponentially growing cells at 25°C were shifted at 36.5°C for one doubling of the cell population and fixed. DNA was stained with DAPI and tubulin detected by indirect immunofluorescence. Lagging DNA was defined as DAPI-stained material lagging on the anaphase spindle (length > 5 μm). P-values from two-sided Fisher's exact tests. (D-E) Psm1-S1022 and Mis4-S183 phosphorylation levels in AGC kinase mutants in *cdc10-129*-arrested cells. Rad21-FLAG (D) and Mis4-GFP (E) immunoprecipitates were analyzed by western blotting using the indicated antibodies and the ratios of signal intensities were calculated.




Figure 6—figure supplement 1.

Mimicking the non phosphorylated state of Psm1-S1022 and Mis4-S183 suppressed $pph3\Delta$ $psm3$ -K105K106N phenotypes.

(A) Cell growth assay showing the suppression of the Ts phenotype. (B) Exponentially growing cells at 25°C were shifted at 36°C for one doubling. DNA was stained with DAPI and tubulin detected by indirect immunofluorescence. Lagging DNA was defined as DAPI-stained material on the anaphase spindle (length>5μm). P-values from two-sided Fisher's exact tests. (C) Exponentially growing cells at 25°C were shifted to 36°C for one doubling of the cell population. Cohesin binding to chromatin was monitored by ChIP using anti-Rad21 antibodies. Mean and SD from 6 ChIPs. *** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, by two-tailed, one sample t-test with 95% confidence interval.

Figure 7—figure supplement 1.

CM3112 loss rate was determined in the presence of rapamycin (RAPA, 200 ng/ml) or solvent alone (DMSO).

Data are mean $\pm$ SEM, n=4 per group (n=10 for the wild-type), ***P<0.001, *P<0.05 repeated measures two-way ANOVA followed by Bonferroni post-hoc test.




Figure 8—figure supplement 1.

Response to nitrogen starvation and refeeding.

Exponentially growing cells in EMM2 medium at 25°C were washed three times in EMM2-N and incubated in EMM2-N for 16 hours at 25°C (time 0). DNA content analysis showed that all strains arrested with a 1C DNA content. G1-arrested cells were then released into the cell cycle in rich medium at 36.5°C. DNA content analysis showed that all strains progressed through S phase with similar kinetics.




Figure 8—figure supplement 2.

(A) Browser view of *mis4*-regulated genes showing the distribution bias towards the ends of chromosomes 1 and 2. (B-C) Overlap between the 338 differentially expressed genes in *mis4-G1487D* versus wild-type with genes upregulated in *tor2ts6* (Wei et al., 2021 [DOI](#)) and genes upregulated during G0 (Zahedi et al., 2023 [DOI](#)).




Figure 8—figure supplement 3.

Cell survival during quiescence.

Exponentially growing cells in EMM2 medium at 25°C were washed three times in EMM2-N and incubated in EMM2-N for 24 hours at 25°C (time 0). Half of the culture was transferred to 36.5°C while the other half remained at 25°C. Cell viability was monitored over time by methylene blue staining (sample image on the right).




Figure 8—figure supplement 4.

Quiescence exit at 25°C.

Exponentially growing cells in EMM2 medium at 25°C were washed three times in EMM2-N and incubated in EMM2-N for 24 hours at 25°C. Half of the culture was transferred to 36.5°C while the other half remained at 25°C. After 24 hours, cells were plated on YES medium (25°C) to assess cell's ability to exit G0 and form colonies.





Figure 8—figure supplement 5.

Quiescence exit at 36.5°C.

Exponentially growing cells in EMM2 medium at 25°C were washed three times in EMM2-N and incubated in EMM2-N for 24 hours at 25°C. Half of the culture was transferred to 36.5°C while the other half remained at 25°C. After 3 days, nitrogen (NH_4Cl 5g/l) was added (time 0). Exit from G0 was monitored by flow cytometry analysis of DNA content. At 25°C (G0 25°C => exit 25°C) the onset of DNA replication occurred ~7hrs after nitrogen addition for all strains. At 36.5°C (G0 36.5°C => exit 36.5°C) DNA replication was slow or asynchronous even for the wild-type. The numbers above the peaks indicate the fraction of cells with 1C and 2C DNA content for the 0 and 12h time points. The *mis4* mutant behaved mostly as wild-type although a larger fraction of cells did not replicate 12 hours after nitrogen addition. For the two strains harbouring *mip1-R401G*, a large fraction of cells did not replicate during the duration of the experiment.






Figure 8-figure supplement 6.

Among the 338 genes differentially expressed in mis4-G1487D, 100 (107 occurrences, 100 genes) show a similar fold change (+/-10%) in mis4-G1487D, mip1-R401G and the double mutant.

Graphs show the log2 fold change over wild-type for the 3 mutant strains for each gene_experiment.




Figure 8—figure supplement 7.

Genes differentially expressed in mis4-G1487D showing a change in the opposite direction in the mip1 mutant.

(A) mis4-G1487D log2 fold change over wild-type >0 and mip1-R401G log2 fold change over wild-type <0 ; 43 occurrences, 41 genes. (B) mis4-G1487D log2 fold change over wild-type <0 and mip1-R401G log2 fold change over wild-type >0 ; 21 occurrences, 21 genes. Graphs show the log2 fold change over wild-type for the 3 mutant strains for each gene_experiment.

Acknowledgements

We would like to thank Tomoyuki Fukuda, Yuichi Morozumi, Janni Petersen, Kazuhiro Shiozaki, Ronit Weisman, Mitsuhiro Yanagida and the National BioResource Project for providing *S. pombe* strains and advices. Keith Gull for the gift of anti-tubulin antibodies. This work was supported by the Centre National de la Recherche Scientifique, l'Université de Bordeaux, la Région Aquitaine, la Fondation ARC pour la recherche contre le cancer (ARCPJA2021060003810) and La Ligue contre le Cancer. Dorian Besson was supported by the International Doctorate Program – IdEx University of Bordeaux and a fellowship from the Fondation ARC. Adele Marston is funded through a Wellcome Investigator Award [220780] and the Wellcome Discovery Research Platform for Hidden Cell Biology [226791]. Work in the Karl Ekwall laboratory was financed by the Swedish research council (VR-MH) and the Swedish Cancer Society (CF). We also would like to thank BEA, the core facility for Bioinformatics and Expression Analysis, which is supported by the board of research at the Karolinska Institutet and the research committee at the Karolinska hospital.

Additional files

Supplementary file 1 [↗](#)

Supplementary file 2 [↗](#)

References

1. Alfredsson-Timmins J, Kristell C, Henningson F, Lyckman S, Bjerling P (2009) **Reorganization of chromatin is an early response to nitrogen starvation in *Schizosaccharomyces pombe*** *Chromosoma* **118**:99–112 [Google Scholar](#)
2. Allshire RC, Nimmo ER, Ekwall K, Javerzat JP, Cranston G (1995) **Mutations derepressing silent centromeric domains in fission yeast disrupt chromosome segregation** *Genes Dev* **9**:218–233 [Google Scholar](#)
3. Bardou P, Mariette J, Escudié F, Djemiel C, Klopp C (2014) **jvenn: an interactive Venn diagram viewer** *BMC Bioinformatics* **15**:293 [Google Scholar](#)
4. Beli P, Lukashchuk N, Wagner SA, Weinert BT, Olsen JV, Baskcomb L, Mann M, Jackson SP, Choudhary C (2012) **Proteomic Investigations Reveal a Role for RNA Processing Factor THRAP3 in the DNA Damage Response** *Molecular Cell* **46**:212–225 [Google Scholar](#)
5. Bernard P, Drogat J, Maure J-F, Dheur S, Vaur S, Genier S, Javerzat J-P (2006) **A screen for cohesion mutants uncovers Ssl3, the fission yeast counterpart of the cohesin loading factor Scc4** *Curr Biol* **16**:875–881 [Google Scholar](#)
6. Bernard P, Maure JF, Partridge JF, Genier S, Javerzat JP, Allshire RC (2001) **Requirement of heterochromatin for cohesion at centromeres** *Science* **294**:2539–2542 [Google Scholar](#)
7. Birot A, Eguienta K, Vazquez S, Claverol S, Bonneu M, Ekwall K, Javerzat J-P, Vaur S (2017) **A second Wpl1 anti-cohesion pathway requires dephosphorylation of fission yeast kleisin Rad21 by PP4** *EMBO J* **36**:1364–1378 [Google Scholar](#)

8. Birot A, Tormos-Pérez M, Vaur S, Feytout A, Jaegy J, Alonso Gil D, Vazquez S, Ekwall K, Javerzat J-P (2020) **The CDK Pef1 and protein phosphatase 4 oppose each other for regulating cohesin binding to fission yeast chromosomes** *eLife* :9 [Google Scholar](#)
9. Bitton DA, Schubert F, Dey S, Okoniewski M, Smith GC, Khadayate S, Pancaldi V, Wood V, Bähler J (2015) **AnGeLi: A Tool for the Analysis of Gene Lists from Fission Yeast** *Front Genet* **6**:330 [Google Scholar](#)
10. Bjedov I, Partridge L (2011) **A longer and healthier life with TOR down-regulation: genetics and drugs** *Biochemical Society Transactions* **39**:460–465 [Google Scholar](#)
11. Carlson CR, Grallert B, Stokke T, Boye E (1999) **Regulation of the start of DNA replication in *Schizosaccharomyces pombe*** *J Cell Sci* **112**:939–46 [Google Scholar](#)
12. Chao WC, Murayama Y, Munoz S, Costa A, Uhlmann F, Singleton MR (2015) **Structural Studies Reveal the Functional Modularity of the Scc2-Scc4 Cohesin Loader** *Cell Rep* **12**:719–25 [Google Scholar](#)
13. Chia KH, Fukuda T, Sofyantor F, Matsuda T, Amai T, Shiozaki K (2017) **Ragulator and GATOR1 complexes promote fission yeast growth by attenuating TOR complex 1 through Rag GTPases** *eLife* :6 [Google Scholar](#)
14. Chica N, Rozalén AE, Pérez-Hidalgo L, Rubio A, Novak B, Moreno S (2016) **Nutritional Control of Cell Size by the Greatwall-Endosulfine-PP2A-B55 Pathway** *Curr Biol* **26**:319–330 [Google Scholar](#)
15. Christensen GL, Kelstrup CD, Lyngsø C, Sarwar U, Bøgebo R, Sheikh SP, Gammeltoft S, Olsen JV, Hansen JL (2010) **Quantitative phosphoproteomics dissection of seven-transmembrane receptor signaling using full and biased agonists** *Mol Cell Proteomics* **9**:1540–1553 [Google Scholar](#)
16. Ciosk R, Shirayama M, Shevchenko A, Tanaka T, Toth A, Nasmyth K (2000) **Cohesin's binding to chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins** *Mol Cell* **5**:243–254 [Google Scholar](#)
17. Collier JE, Lee BG, Roig MB, Yatskevich S, Petela NJ, Metson J, Voulgaris M, Gonzalez Llamazares A, Lowe J, Nasmyth KA (2020) **Transport of DNA within cohesin involves clamping on top of engaged heads by Scc2 and entrapment within the ring by Scc3** *eLife* :9 [Google Scholar](#)
18. Costantino L, Hsieh TS, Lamothe R, Darzacq X, Koshland D (2020) **Cohesin residency determines chromatin loop patterns** *eLife* :9 [Google Scholar](#)
19. Courtheoux T, Gay G, Gachet Y, Tournier S (2009) **Ase1/Prc1-dependent spindle elongation corrects merotelically during anaphase in fission yeast** *J Cell Biol* **187**:399–412 [Google Scholar](#)
20. Davidson IF, Bauer B, Goetz D, Tang W, Wutz G, Peters JM (2019) **DNA loop extrusion by human cohesin** *Science* [Google Scholar](#)
21. Davidson IF, Peters J-M (2021) **Genome folding through loop extrusion by SMC complexes** *Nat Rev Mol Cell Biol* **22**:445–464 [Google Scholar](#)
22. Davie E, Forte GMA, Petersen J (2015) **Nitrogen regulates AMPK to control TORC1 signaling** *Curr Biol* **25**:445–454 [Google Scholar](#)

23. Dephoure N, Zhou C, Villén J, Beausoleil SA, Bakalarski CE, Elledge SJ, Gygi SP (2008) **A quantitative atlas of mitotic phosphorylation** *Proc Natl Acad Sci U S A* **105**:10762–10767 [Google Scholar](#)
24. Dheur S, Saupe SJ, Genier S, Vazquez S, Javerzat J-P (2011) **Role for cohesin in the formation of a heterochromatic domain at fission yeast subtelomeres** *Mol Cell Biol* **31**:1088–1097 [Google Scholar](#)
25. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) **STAR: ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21 [Google Scholar](#)
26. Feytout A, Vaur S, Genier S, Vazquez S, Javerzat J-P (2011) **Psm3 acetylation on conserved lysine residues is dispensable for viability in fission yeast but contributes to Eso1-mediated sister chromatid cohesion by antagonizing Wpl1** *Mol Cell Biol* **31**:1771–1786 [Google Scholar](#)
27. Fukuda T, Shiozaki K (2018) **The Rag GTPase-Ragulator complex attenuates TOR complex 1 signaling in fission yeast** *Autophagy* **14**:1105–1106 [Google Scholar](#)
28. Fukuda T, Sofyantor F, Tai YT, Chia KH, Matsuda T, Murase T, Morozumi Y, Tatebe H, Kanki T, Shiozaki K (2021) **Tripartite suppression of fission yeast TORC1 signaling by the GATOR1-Sea3 complex, the TSC complex, and Gcn2 kinase** *eLife* **10** [Google Scholar](#)
29. Furuya K, Takahashi K, Yanagida M (1998) **Faithful anaphase is ensured by Mis4, a sister chromatid cohesion molecule required in S phase and not destroyed in G1 phase** *Genes Dev* **12**:3408–3418 [Google Scholar](#)
30. Gachet Y, Reyes C, Courthéoux T, Goldstone S, Gay G, Serrurier C, Tournier S (2008) **Sister kinetochore recapture in fission yeast occurs by two distinct mechanisms, both requiring Dam1 and Klp2** *Mol Biol Cell* **19**:1646–1662 [Google Scholar](#)
31. González A, Hall MN (2017) **Nutrient sensing and TOR signaling in yeast and mammals** *EMBO J* **36**:397–408 [Google Scholar](#)
32. Goto Y, Yamagishi Y, Shintomi-Kawamura M, Abe M, Tanno Y, Watanabe Y (2017) **Pds5 Regulates Sister-Chromatid Cohesion and Chromosome Bi-orientation through a Conserved Protein Interaction Module** *Current Biology* **27**:1005–1012 [Google Scholar](#)
33. Grand RS, Pichugina T, Gehlen LR, Jones MB, Tsai P, Allison JR, Martienssen R, O’Sullivan JM (2014) **Chromosome conformation maps in fission yeast reveal cell cycle dependent sub nuclear structure** *Nucleic Acids Research* **42**:12585–12599 [Google Scholar](#)
34. Gregan J, Polakova S, Zhang L, Tolic-Norrellykke IM, Cimini D (2011) **Merotelic kinetochore attachment: causes and effects** *Trends Cell Biol* **21**:374–81 [Google Scholar](#)
35. Halova L, Cogley D, Franz-Wachtel M, Wang T, Morrison KR, Krug K, Nalpas N, Mačák B, Hagan IM, Humphrey SJ, et al (2021) **A TOR (target of rapamycin) and nutritional phosphoproteome of fission yeast reveals novel targets in networks conserved in humans** *Open Biol* **11**:200405 [Google Scholar](#)
36. Hayashi T, Hatanaka M, Nagao K, Nakaseko Y, Kanoh J, Kokubu A, Ebe M, Yanagida M (2007) **Rapamycin sensitivity of the Schizosaccharomyces pombe tor2 mutant and organization of two highly phosphorylated TOR complexes by specific and common subunits** *Genes Cells* **12**:1357–70 [Google Scholar](#)

37. Hirai H, Sen Y, Tamura M, Ohta K (2023) **TOR inactivation triggers heterochromatin formation in rDNA during glucose starvation** *Cell Reports* :42 [Google Scholar](#)
38. Hu B, Petela N, Kurze A, Chan KL, Chapard C, Nasmyth K (2015) **Biological chromodynamics: a general method for measuring protein occupancy across the genome by calibrating ChIP-seq** *Nucleic Acids Res* **43**:e132 [Google Scholar](#)
39. Ikai N, Nakazawa N, Hayashi T, Yanagida M (2011) **The reverse, but coordinated, roles of Tor2 (TORC1) and Tor1 (TORC2) kinases for growth, cell cycle and separase-mediated mitosis in *Schizosaccharomyces pombe*** *Open Biology* **1**:110007 [Google Scholar](#)
40. Kagami A, Sakuno T, Yamagishi Y, Ishiguro T, Tsukahara T, Shirahige K, Tanaka K, Watanabe Y (2011) **Acetylation regulates monopolar attachment at multiple levels during meiosis I in fission yeast** *EMBO Rep* **12**:1189–95 [Google Scholar](#)
41. Kall L, Canterbury JD, Weston J, Noble WS, MacCoss MJ (2007) **Semi-supervised learning for peptide identification from shotgun proteomics datasets** *Nature methods* **4**:923–5 [Google Scholar](#)
42. Kettenbach AN, Deng L, Wu Y, Baldissard S, Adamo ME, Gerber SA, Moseley JB (2015) **Quantitative phosphoproteomics reveals pathways for coordination of cell growth and division by the conserved fission yeast kinase pom1** *Mol Cell Proteomics* **14**:1275–87 [Google Scholar](#)
43. Kipling D, Kearsley SE (1990) **Reversion of autonomously replicating sequence mutations in *Saccharomyces cerevisiae*: creation of a eucaryotic replication origin within procaryotic vector DNA** *Mol Cell Biol* **10**:265–272 [Google Scholar](#)
44. Knutsen JH, Rein ID, Rothe C, Stokke T, Grallert B, Boye E (2011) **Cell-cycle analysis of fission yeast cells by flow cytometry** *PLoS One* **6**:e17175 [Google Scholar](#)
45. Larabee RN, Weisman R (2020) **Nuclear Functions of TOR: Impact on Transcription and the Epigenome** *Genes* :11 [Google Scholar](#)
46. Lawrimore J, Doshi A, Friedman B, Yeh E, Bloom K (2018) **Geometric partitioning of cohesin and condensin is a consequence of chromatin loops** *MBoC* **29**:2737–2750 [Google Scholar](#)
47. Lengronne A, Katou Y, Mori S, Yokobayashi S, Kelly GP, Itoh T, Watanabe Y, Shirahige K, Uhlmann F (2004) **Cohesin relocation from sites of chromosomal loading to places of convergent transcription** *Nature* **430**:573–8 [Google Scholar](#)
48. Liao Y, Smyth GK, Shi W (2014) **featureCounts: an efficient general purpose program for assigning sequence reads to genomic features** *Bioinformatics* **30**:923–930 [Google Scholar](#)
49. Matsuda S, Kikkawa U, Uda H, Nakashima A (2020) ***S. pombe* Pef1/CDK5 regulates sexual differentiation through control of the TORC1 pathway and autophagy** *J Cell Sci* [Google Scholar](#)
50. Matsumoto T, Murakami S, Niwa O, Yanagida M (1990) **Construction and characterization of centric and acentric linear minichromosomes in fission yeast** *Curr Genet* **18**:323–330 [Google Scholar](#)
51. Mattingly M, Seidel C, Muñoz S, Hao Y, Zhang Y, Wen Z, Florens L, Uhlmann F, Gerton JL (2022) **Mediator recruits the cohesin loader Scc2 to RNA Pol II-transcribed genes and promotes**

sister chromatid cohesion *Curr Biol* **32**:2884–2896 [Google Scholar](#)

52. Mayya V, Lundgren DH, Hwang S-I, Rezaul K, Wu L, Eng JK, Rodionov V, Han DK (2009) **Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions** *Sci Signal* **2**:ra46 [Google Scholar](#)
53. Mizuguchi T, Barrowman J, Grewal SIS (2015) **Chromosome domain architecture and dynamic organization of the fission yeast genome** *FEBS Lett* **589**:2975–2986 [Google Scholar](#)
54. Mizuguchi T, Fudenberg G, Mehta S, Belton JM, Taneja N, Folco HD, FitzGerald P, Dekker J, Mirny L, Barrowman J, et al (2014) **Cohesin-dependent globules and heterochromatin shape 3D genome architecture in *S. pombe*** *Nature* **516**:432–435 [Google Scholar](#)
55. Moreno S, Klar A, Nurse P (1991) **Molecular genetic analysis of fission yeast *Schizosaccharomyces pombe*** *Methods Enzymol* **194**:795–823 [Google Scholar](#)
56. Morozumi Y, Hishinuma A, Furusawa S, Sofyantoro F, Tatebe H, Shiozaki K (2021) **Fission yeast TOR complex 1 phosphorylates Psk1 through an evolutionarily conserved interaction mediated by the TOS motif** *J Cell Sci* :134 [Google Scholar](#)
57. Murayama Y, Samora CP, Kurokawa Y, Iwasaki H, Uhlmann F (2018) **Establishment of DNA-DNA Interactions by the Cohesin Ring** *Cell* **172**:465–477 [Google Scholar](#)
58. Murayama Y, Uhlmann F (2015) **DNA Entry into and Exit out of the Cohesin Ring by an Interlocking Gate Mechanism** *Cell* **163**:1628–40 [Google Scholar](#)
59. Nakashima A, Otsubo Y, Yamashita A, Sato T, Yamamoto M, Tamanoi F (2012) **Psk1, an AGC kinase family member in fission yeast, is directly phosphorylated and controlled by TORC1 and functions as S6 kinase** *J Cell Sci* **125**:5840–5849 [Google Scholar](#)
60. Oldenkamp R, Rowland BD (2022) **A walk through the SMC cycle: From catching DNAs to shaping the genome** *Mol Cell* **82**:1616–1630 [Google Scholar](#)
61. Otsubo Y, Nakashima A, Yamamoto M, Yamashita A (2017) **TORC1-Dependent Phosphorylation Targets in Fission Yeast** *Biomolecules* **7**:50 [Google Scholar](#)
62. Paldi F, Alver B, Robertson D, Schalbetter SA, Kerr A, Kelly DA, Baxter J, Neale MJ, Marston AL (2020) **Convergent genes shape budding yeast pericentromeres** *Nature* **582**:119–123 [Google Scholar](#)
63. Pan C, Olsen JV, Daub H, Mann M (2009) **Global Effects of Kinase Inhibitors on Signaling Networks Revealed by Quantitative Phosphoproteomics*** *Molecular & Cellular Proteomics* **8**:2796–2808 [Google Scholar](#)
64. Park DS, Nguyen SC, Isenhardt R, Shah PP, Kim W, Barnett RJ, Chandra A, Luppino JM, Harke J, Wai M, et al (2023) **High-throughput Oligopaint screen identifies druggable 3D genome regulators** *Nature* **620**:209–217 [Google Scholar](#)
65. Pidoux AL, Uzawa S, Perry PE, Cande WZ, Allshire RC (2000) **Live analysis of lagging chromosomes during anaphase and their effect on spindle elongation rate in fission yeast** *J Cell Sci* **113**:4177–4191 [Google Scholar](#)
66. Reidman S, Cohen A, Kupiec M, Weisman R (2019) **The cytosolic form of aspartate aminotransferase is required for full activation of TOR complex 1 in fission yeast** *J Biol*

Chem **294**:18244–18255 [Google Scholar](#)

67. Rigbolt KTG, Prokhorova TA, Akimov V, Henningsen J, Johansen PT, Kratchmarova I, Kassem M, Mann M, Olsen JV, Blagoev B (2011) **System-wide temporal characterization of the proteome and phosphoproteome of human embryonic stem cell differentiation** *Sci Signal* **4**:rs3 [Google Scholar](#)
68. Robinson MD, McCarthy DJ, Smyth GK (2010) **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data** *Bioinformatics* **26**:139–140 [Google Scholar](#)
69. Ruse CI, McClatchy DB, Lu B, Cociorva D, Motoyama A, Park SK, Yates JR (2008) **Motif-specific sampling of phosphoproteomes** *J Proteome Res* **7**:2140–2150 [Google Scholar](#)
70. Rutherford KM, Lera-Ramírez M, Wood V (2024) **PomBase: a Global Core Biodata Resource-growth, collaboration, and sustainability** *Genetics* **227**:iyae007 [Google Scholar](#)
71. Sacristan C, Samejima K, Ruiz LA, Deb M, Lambers MLA, Buckle A, Brackley CA, Robertson D, Hori T, Webb S, et al (2024) **Vertebrate centromeres in mitosis are functionally bipartite structures stabilized by cohesin** *Cell* :0 [Google Scholar](#)
72. Sajiki K, Hatanaka M, Nakamura T, Takeda K, Shimanuki M, Yoshida T, Hanyu Y, Hayashi T, Nakaseko Y, Yanagida M (2009) **Genetic control of cellular quiescence in *S. pombe*** *J Cell Sci* **122**:1418–29 [Google Scholar](#)
73. Sajiki K, Tahara Y, Villar-Briones A, Pluskal T, Teruya T, Mori A, Hatanaka M, Ebe M, Nakamura T, Aoki K, et al (2018) **Genetic defects in SAPK signalling, chromatin regulation, vesicle transport and CoA-related lipid metabolism are rescued by rapamycin in fission yeast** *Open Biol* **8**:170261 [Google Scholar](#)
74. Schmidt CK, Brookes N, Uhlmann F (2009) **Conserved features of cohesin binding along fission yeast chromosomes** *Genome Biol* **10**:R52 [Google Scholar](#)
75. Shi Z, Gao H, Bai XC, Yu H (2020) **Cryo-EM structure of the human cohesin-NIPBL-DNA complex** *Science* **368**:1454–1459 [Google Scholar](#)
76. van Slegtenhorst M, Carr E, Stoyanova R, Kruger WD, Henske EP (2004) **Tsc1+ and tsc2+ regulate arginine uptake and metabolism in *Schizosaccharomyces pombe*** *J Biol Chem* **279**:12706–12713 [Google Scholar](#)
77. Steglich B, Fillion GJ, van Steensel B, Ekwall K (2012) **The inner nuclear membrane proteins Man1 and Ima1 link to two different types of chromatin at the nuclear periphery in *S. pombe*** *Nucleus* **3**:77–87 [Google Scholar](#)
78. Suma M, Arakawa O, Tahara Y, Sajiki K, Saitoh S, Yanagida M (2024) **In quiescent G0 phase, *Schizosaccharomyces pombe* Mis4 ensures full nuclear separation during the subsequent M phase** *bioRxiv* <https://doi.org/10.1101/2024.03.29.587322> | [Google Scholar](#)
79. Swaffer MP, Jones AW, Flynn HR, Snijders AP, Nurse P (2018) **Quantitative Phosphoproteomics Reveals the Signaling Dynamics of Cell-Cycle Kinases in the Fission Yeast *Schizosaccharomyces pombe*** *Cell Rep* **24**:503–514 [Google Scholar](#)
80. Tay YD, Leda M, Spanos C, Rappsilber J, Goryachev AB, Sawin KE (2019) **Fission Yeast NDR/LATS Kinase Orb6 Regulates Exocytosis via Phosphorylation of the Exocyst Complex**

Cell Rep **26**:1654–1667 [Google Scholar](#)

81. Torres-Garcia S, Di Pompeo L, Eivers L, Gaborieau B, White SA, Pidoux AL, Kanigowska P, Yaseen I, Cai Y, Allshire RC (2020) **SpEDIT: A fast and efficient CRISPR/Cas9 method for fission yeast** *Wellcome open research* **5**:274 [Google Scholar](#)
82. Vaur S, Feytout A, Vazquez S, Javerzat J-P (2012) **Pds5 promotes cohesin acetylation and stable cohesin-chromosome interaction** *EMBO Rep* **13**:645–652 [Google Scholar](#)
83. Waldman T (2020) **Emerging themes in cohesin cancer biology** *Nature reviews Cancer* **20**:504–515 [Google Scholar](#)
84. Wei Y, Lee NN, Pan L, Dhakshnamoorthy J, Sun LL, Zofall M, Wheeler D, Grewal SIS (2021) **TOR targets an RNA processing network to regulate facultative heterochromatin, developmental gene expression and cell proliferation** *Nat Cell Biol* **23**:243–256 [Google Scholar](#)
85. Woods A, Sherwin T, Sasse R, MacRae TH, Baines AJ, Gull K (1989) **Definition of individual components within the cytoskeleton of Trypanosoma brucei by a library of monoclonal antibodies** *J Cell Sci* **93**:491–500 [Google Scholar](#)
86. Xu B, Lee KK, Zhang L, Gerton JL (2013) **Stimulation of mTORC1 with L-leucine rescues defects associated with Roberts syndrome** *PLoS Genet* **9**:e1003857 [Google Scholar](#)
87. Yamagishi Y, Honda T, Tanno Y, Watanabe Y (2010) **Two Histone Marks Establish the Inner Centromere and Chromosome Bi-Orientation** *Science* **330**:239–243 [Google Scholar](#)
88. Yoon MS (2020) **Nanotechnology-Based Targeting of mTOR Signaling in Cancer** *International journal of nanomedicine* **15**:5767–5781 [Google Scholar](#)
89. Zahedi Y, Zeng S, Ekwall K (2023) **An essential role for the Ino80 chromatin remodeling complex in regulation of gene expression during cellular quiescence** *Chromosome Res* **31**:14 [Google Scholar](#)
90. Zhao T, Fan J, Abu-Zaid A, Burley SK, Zheng XFS (2024) **Nuclear mTOR Signaling Orchestrates Transcriptional Programs Underlying Cellular Growth and Metabolism** *Cells* **13**:781 [Google Scholar](#)
91. Zheng H, Xie W (2019) **The role of 3D genome organization in development and cell differentiation** *Nature reviews Molecular cell biology* **20**:535–550 [Google Scholar](#)
92. Zhou H, Zheng L, Lu K, Gao Y, Guo L, Xu W, Wang X (2017) **Downregulation of Cohesin Loading Factor Nipped-B-Like Protein (NIPBL) Induces Cell Cycle Arrest, Apoptosis, and Autophagy of Breast Cancer Cell Lines** *Med Sci Monit* **23**:4817–4825 [Google Scholar](#)

Author information

Dorian Besson*

CNRS, Université de Bordeaux - Institut de Biochimie et Génétique Cellulaires, UMR5095
Bordeaux, France

*Equal contribution

Sabine Vaur*

CNRS, Université de Bordeaux - Institut de Biochimie et Génétique Cellulaires, UMR5095
Bordeaux, France

*Equal contribution

Stéphanie Vazquez

CNRS, Université de Bordeaux - Institut de Biochimie et Génétique Cellulaires, UMR5095
Bordeaux, France

Sylvie Tournier

MCD, Centre de Biologie Intégrative, Université de Toulouse, CNRS, UPS, Toulouse, France
ORCID iD: [0000-0002-7869-6891](https://orcid.org/0000-0002-7869-6891)

Yannick Gachet

MCD, Centre de Biologie Intégrative, Université de Toulouse, CNRS, UPS, Toulouse, France
ORCID iD: [0000-0001-7548-7974](https://orcid.org/0000-0001-7548-7974)

Adrien Birot

CNRS, Université de Bordeaux - Institut de Biochimie et Génétique Cellulaires, UMR5095
Bordeaux, France
ORCID iD: [0000-0002-8299-8518](https://orcid.org/0000-0002-8299-8518)

Stéphane Claverol

Univ. Bordeaux, Bordeaux Proteome, Bordeaux, France

Adèle Marston

Centre for Cell Biology, Institute of Cell Biology, University of Edinburgh, Edinburgh, United Kingdom
ORCID iD: [0000-0002-3596-9407](https://orcid.org/0000-0002-3596-9407)

Anastasios Damdimopoulos

Department of Medicine, Karolinska Institutet, Stockholm, Sweden
ORCID iD: [0000-0003-4425-1189](https://orcid.org/0000-0003-4425-1189)

Karl Ekwall

Department of Medicine, Karolinska Institutet, Stockholm, Sweden
ORCID iD: [0000-0002-3029-4041](https://orcid.org/0000-0002-3029-4041)

Jean-Paul Javerzat

CNRS, Université de Bordeaux - Institut de Biochimie et Génétique Cellulaires, UMR5095
Bordeaux, France
ORCID iD: [0000-0002-9671-6753](https://orcid.org/0000-0002-9671-6753)

For correspondence: JPaul.Javerzat@ibgc.cnrs.fr

Editors

Reviewing Editor

Megan King

Yale School of Medicine, New Haven, United States of America

Senior Editor

Lori Sussel

University of Colorado Anschutz Medical Campus, Aurora, United States of America

Reviewer #1 (Public review):

Summary:

In this study, Besson et al. investigate how environmental nutrient signals regulate chromosome biology through the TORC1 signaling pathway in *Schizosaccharomyces pombe*. Specifically, the authors explore the impact of TORC1 on cohesin function - a protein complex essential for chromosome segregation and transcriptional regulation. Through a combination of genetic screens, biochemical analysis, phospho-proteomics, and transcriptional profiling, they uncover a functional and physical interaction between TORC1 and cohesin. The data suggest that reduced TORC1 activity enhances cohesin binding to chromosomes and improves chromosome segregation, with implications for stress-responsive gene expression, especially in subtelomeric regions.

Strengths:

This work presents a compelling link between nutrient sensing and chromosome regulation. The major strength of the study lies in its comprehensive and multi-disciplinary approach. The authors integrate genetic suppression screens, live-cell imaging, chromatin immunoprecipitation, co-immunoprecipitation, and mass spectrometry to uncover the functional connection between TORC1 signaling and cohesin. The use of phospho-mutant alleles of cohesin subunits and their loader provides mechanistic insight into the regulatory role of phosphorylation. The addition of transcriptomic analysis further strengthens the biological relevance of the findings and places them in a broader physiological context. Altogether, the dataset convincingly supports the authors' main conclusions and opens up new avenues of investigation.

Weaknesses:

While the study is strong overall, a few limitations are worth noting. The consistency of cohesin phosphorylation changes under different TORC1-inhibiting conditions (e.g., genetic mutants vs. rapamycin treatment) is unclear and could benefit from further clarification. The phosphorylation sites identified on cohesin subunits do not match known AGC kinase consensus motifs, raising the possibility that the modifications are indirect. The study relies heavily on one TORC1 mutant allele (*mip1-R401G*), and additional alleles could strengthen the generality of the findings. Furthermore, while the results suggest that nutrient availability influences cohesin function, this is not directly tested by comparing growth or cohesin dynamics under defined nutrient conditions.

<https://doi.org/10.7554/eLife.108275.1.sa1>

Reviewer #2 (Public review):

Summary:

In this study, the authors follow up on a previous suppressor screen of a temperature-sensitive allele of *mis4* (*mis4*-G1487D), the cohesin loading factor in *S. pombe*, and identify additional suppressor alleles tied to the *S. pombe* TORC1 complex. Their analysis suggests that these suppressor mutations attenuate TORC1 activity, while enhanced TORC1 activity is deleterious in this context. Suppression of TORC1 activity also ameliorates chromosome segregation and spindle defects observed in the *mis4*-G1487D strain, although some more subtle effects are not reconstituted. The authors provide evidence that this genetic suppression is also tied to the reconstitution of cohesin loading. Moreover, disrupting TORC1 also enhances *Mis4*/cohesin association with chromatin (likely reflecting enhanced loading) in WT cells, while rapamycin treatment can enhance the robustness of chromosome transmission. These effects likely arise directly through TORC1 or its downstream effector kinases, as TORC1 co-purifies with *Mis4* and *Rad21*; these factors are also phosphorylated in a TORC1-dependent fashion. Disrupting *Sck2*, a kinase downstream of TORC1, also suppresses the *mis4*-G1487D allele while simultaneous disruption of *Sck1* and *Sck2* enhances cohesin association with chromatin, albeit with differing effects on phosphorylation of *Mis4* and *Psm1/Scm1*. Phosphomutants of *Mis4* and *Psm1* that mimic observed phosphorylation states identified by mass spectrometry that are TORC1-dependent also suppressed phenotypes observed in the *mis4*-G1487D background. Last, the authors provide evidence that the *mis4*-G1487D background and TORC1 mutant backgrounds display an overlap in the dysregulation of genes that respond to environmental conditions, particularly in genes tied to meiosis or other "stress".

Overall, the authors provide compelling evidence from genetics, biochemistry, and cell biology to support a previously unknown mechanism by which nutrient sensing regulates cohesin loading with implications for the stress response. The technical approaches are generally sound, well-controlled, and comprehensive.

Specific Points:

- (1) While the authors favor the model that the enhanced cohesin loading upon diminished TORC1 activity helps cells to survive harsh environmental conditions, as starvation of *S. pombe* also drives commitment to meiosis, it seems as plausible that enhanced cohesin loading is related to preparing the chromosomes to mate.
- (2) Related to Point 1, the lab of Sophie Martin previously published that phosphorylation of *Mis4* characterizes a cluster of phosphotargets during starvation/meiotic induction (PMID: 39705284). This work should be cited, and the authors should interrogate how their observations do or do not relate to these prior observations (are these the same phosphosites?).
- (3) It would be useful for the authors to combine their experimental data sets to interrogate whether there is a relationship between the regions where gene expression is altered in the *mis4*-G1487D strain and changes in the loading of cohesin in their ChIP experiments.
- (4) Given that the genes that are affected are predominantly sub-telomeric while most genes are not affected in the *mis4*-G1487D strain, one possibility that the authors may wish to consider is that the regions that become dysregulated are tied to heterochromatic regions where *Swi6/HP1* has been implicated in cohesin loading.
- (5) It would be helpful to show individual data points from replicates in the bar graphs - it is not always clear what comprises the data sets, and superplots would be of great help.

<https://doi.org/10.7554/eLife.108275.1.sa0>