

# Salmonella exploits host- and bacterial-derived $\beta$ -alanine for replication inside host macrophages

Shuai Ma, Bin Yang, Yuyang Sun, Xinyue Wang, Houliang Guo, Ruiying Liu, Ting Ye, Chenbo Kang, Jingnan Chen, Lingyan Jiang 

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

Reviewed Preprint

v3 • May 15, 2025

Revised by authors

Reviewed Preprint

v2 • April 16, 2025

Reviewed Preprint

v1 • November 21, 2024

## eLife Assessment

The authors use a multidisciplinary approach to provide a link between Beta-alanine and *S. Typhimurium* (STM) infection and virulence. This **valuable** work shows how Beta-alanine synthesis mediates zinc homeostasis regulation, possibly contributing to virulence. The work is **convincing** as it adds to the existing knowledge of metabolic flexibility displayed by STM during infection.

<https://doi.org/10.7554/eLife.103714.3.sa3>

## Abstract

*Salmonella* is a major foodborne pathogen that can effectively replicate inside host macrophages to establish life-threatening systemic infections. *Salmonella* must utilize diverse nutrients for growth in nutrient-poor macrophages, but which nutrients are required for intracellular *Salmonella* growth is largely unknown. Here, we found that either acquisition from the host or *de novo* synthesis of a nonprotein amino acid,  $\beta$ -alanine, is critical for *Salmonella* replication inside macrophages. The concentration of  $\beta$ -alanine is decreased in *Salmonella*-infected macrophages, while the addition of exogenous  $\beta$ -alanine enhances *Salmonella* replication in macrophages, suggesting that *Salmonella* can uptake host-derived  $\beta$ -alanine for intracellular growth. Moreover, the expression of *panD*, the rate-limiting gene required for  $\beta$ -alanine synthesis in *Salmonella*, is upregulated when *Salmonella* enters macrophages. Mutation of *panD* impaired *Salmonella* replication in macrophages and colonization in the mouse liver and spleen, indicating that *de novo* synthesis of  $\beta$ -alanine is essential for intracellular *Salmonella* growth and systemic infection. Additionally, we revealed that  $\beta$ -alanine influences *Salmonella* intracellular replication and *in vivo* virulence partially by increasing expression of the zinc transporter genes *znuABC*, which in turn facilitates the uptake of the essential micronutrient zinc by *Salmonella*. Taken together, these findings highlight the important role of  $\beta$ -alanine in the intracellular replication and virulence of *Salmonella*, and *panD* is a promising target for controlling systemic *Salmonella* infection.

## Introduction

*Salmonella* is a major foodborne pathogen worldwide that can cause self-limiting gastroenteritis or life-threatening systemic disease in a wide range of animals (Fàbrega & Vila, 2013 [↗](#); Ohl & Miller, 2001 [↗](#)). *Salmonella* infection remains a significant global public health concern. An estimated 93.8 million cases of gastroenteritis and 27 million cases of systemic diseases caused by *Salmonella* species occur annually worldwide, with 355,000 deaths (Kim et al., 2019 [↗](#); Majowicz et al., 2010 [↗](#)). The ability to survive and replicate in host macrophages is a key determinant for *Salmonella* to induce systemic infection (Fields, Swanson, Haidaris, & Heffron, 1986 [↗](#); LaRock, Chaudhary, & Miller, 2015 [↗](#); Leung & Finlay, 1991 [↗](#)). After internalization by macrophages, *Salmonella* delivers a set of more than 30 effector proteins to the macrophage cytoplasm, mainly through a type III secretion system (T3SS) encoded by *Salmonella* pathogenicity island-2 (SPI-2) (Pillay et al., 2023 [↗](#)). SPI-2 effectors manipulate diverse cellular processes to promote the formation of a membrane-bound compartment, termed the *Salmonella*-containing vacuole (SCV), a niche where *Salmonella* resides and grows (Castanheira & García-Del Portillo, 2017 [↗](#); Rosenberg, Riquelme, Prince, & Avraham, 2022 [↗](#); Steeb et al., 2013 [↗](#)). SCV protects *Salmonella* from contact with antimicrobial agents in macrophages (Figueira & Holden, 2012 [↗](#); Li et al., 2023 [↗](#)). Moreover, SPI-2 effectors induce the formation of specific tubular membrane compartments that extend from the SCV, known as *Salmonella*-induced filaments (SIFs). These filaments allow *Salmonella* to access various types of endocytosed nutrients, thereby facilitating efficient replication within macrophages (Brumell, Tang, Mills, & Finlay, 2001 [↗](#); Liss et al., 2017 [↗](#); Rajashekar, Liebl, Seitz, & Hensel, 2008 [↗](#)).

As the SCV of macrophages is a nutrient-poor environment (Fields et al., 1986 [↗](#); Kehl, Noster, & Hensel, 2020 [↗](#)), to effectively replicate in the SCV, *Salmonella* needs to acquire a wide range of host nutrients or host-derived metabolites and synthesize metabolites *de novo* that cannot be sufficiently accessed from the host (Dandekar et al., 2014 [↗](#); Röder, Felgner, & Hensel, 2021 [↗](#); Tuli & Sharma, 2019 [↗](#)). Nutrients/metabolites are used by intracellular *Salmonella* either as carbon sources to generate energy or for the synthesis of fatty acids and proteins (Dandekar et al., 2014 [↗](#); Steeb et al., 2013 [↗](#)). Moreover, several metabolites were found to be employed by *Salmonella* as environmental cues to induce the expression of virulence genes (Jiang et al., 2021 [↗](#); X. Wang et al., 2023 [↗](#)). In recent years, an increasing number of studies have focused on the intracellular nutrition of *Salmonella* (Bumann & Schothorst, 2017 [↗](#); Liss et al., 2017 [↗](#); Röder et al., 2021 [↗](#)); however, the nutrients that are required for *Salmonella* replication in macrophages remain largely unknown.

$\beta$ -Alanine, also known as 3-aminopropionic acid (3-AP), is the only naturally occurring  $\beta$ -type amino acid and is found in all living organisms (Song, Zhang, Wang, Xu, & Wei, 2023 [↗](#)).  $\beta$ -Alanine can be synthesized *de novo* by bacteria, fungi, and plants, whereas animals need to obtain it from food or generate it via the catabolism of cytosine and uracil (Gojković, Sandrini, & Piskur, 2001 [↗](#); L. Wang, Mao, Wang, Ma, & Chen, 2021 [↗](#)). In bacteria,  $\beta$ -alanine is synthesized via the decarboxylation of L-aspartate, a reaction catalyzed by L-aspartate decarboxylase (PanD) (Begley, Kinsland, & Strauss, 2001 [↗](#); Schmitzberger, Smith, Abell, & Blundell, 2003 [↗](#); West, Traut, Shanley, & O'Donovan, 1985 [↗](#)). The *panD* gene is conserved among most bacteria (Nozaki, Webb, & Niki, 2012 [↗](#)). Although  $\beta$ -alanine is a nonprotein amino acid that is not incorporated into proteins, it has important physiological functions in the metabolism of organisms (L. Wang et al., 2021 [↗](#); Yuan et al., 2022 [↗](#)). First,  $\beta$ -alanine forms a part of pantothenate (vitamin B5), which is the key precursor for the biosynthesis of coenzyme A (CoA) (Webb, Smith, & Abell, 2004 [↗](#); White, Gunyuzlu, & Toyn, 2001 [↗](#)). CoA is an essential cofactor involved in many metabolic pathways, including the synthesis and degradation of fatty acids, pyruvate oxidation through the tricarboxylic acid (TCA) cycle, and the production of secondary metabolites (Davaapil, Tsuchiya, & Gout, 2014 [↗](#); Gout, 2019 [↗](#); Sibon & Strauss, 2016 [↗](#); Theodoulou, Sibon, Jackowski, & Gout,

2014 [\[1\]](#)). Second,  $\beta$ -alanine is a limiting precursor of carnosine, a nonenzymatic free radical scavenger and a natural antioxidant, with anti-inflammatory and neuroprotective effects in animals (Boldyrev, Aldini, & Derave, 2013 [\[2\]](#); Hoffman, Varanoske, & Stout, 2018 [\[3\]](#)). In the past 15 years,  $\beta$ -alanine has become one of the most commonly used sports supplements worldwide (Bellinger, 2014 [\[4\]](#); Hoffman et al., 2018 [\[5\]](#); Huerta Ojeda, Tapia Cerda, Poblete Salvatierra, Barahona-Fuentes, & Jorquera Aguilera, 2020 [\[6\]](#)). Although both *Salmonella* and host cells are capable of producing  $\beta$ -alanine, whether  $\beta$ -alanine contributes to the pathogenicity and intracellular growth of *Salmonella* remains unknown.


In this work, using targeted metabolic profiling, *in vitro* and *in vivo* infection assays, and many other molecular techniques, we demonstrated that the utilization of  $\beta$ -alanine is essential for *Salmonella* replication in host macrophages and virulence in mice. *Salmonella* acquires  $\beta$ -alanine both via the uptake of  $\beta$ -alanine from host macrophages and the *de novo* synthesis of  $\beta$ -alanine. Further investigation revealed the molecular mechanism underlying the contribution of  $\beta$ -alanine to *Salmonella* intracellular replication and pathogenicity, wherein  $\beta$ -alanine promotes the expression of zinc transporter genes to facilitate the uptake of the essential micronutrient zinc by intracellular *Salmonella*, therefore promoting *Salmonella* replication in macrophages and subsequent systemic infection. Taken together, these findings demonstrate a correlation between *Salmonella*  $\beta$ -alanine utilization and zinc uptake during intracellular infection and provide new insights into the intracellular nutrition of *Salmonella*. The rate-limiting gene (*panD*) in the  $\beta$ -alanine synthesis pathway of *Salmonella* might be a future target for the prevention and treatment of this pathogen.

## Results

### Host-derived $\beta$ -alanine promotes *Salmonella* replication inside macrophages

To explore changes in the levels of different amino acids inside macrophages upon *Salmonella* infection, we performed targeted metabolomics analysis of mouse RAW264.7 macrophages that were mock-infected or infected with wild-type *Salmonella* (*Salmonella enterica* serovar Typhimurium ATCC 14028s, STM) for 8 h using liquid chromatography-tandem mass spectrometry (LC[MS/MS]) (Figure 1A [\[1\]](#)). Principal component analysis (PCA) demonstrated a clear separation between the mock- and *Salmonella*-infected groups (Figure 1B [\[2\]](#)). A total of 26 free amino acids were analyzed, and 8 showed significant differences in abundance between the two groups (VIP (Variable Importance in the Projection) > 1 and  $P < 0.05$ ; FC (fold change) > 1.5 or < 0.667) (Figure 1C, D [\[3\]](#)). Compared with those in the mock-infected group, the concentrations of 3 amino acids (L-hydroxyproline, L-citrulline and L-cysteine) were upregulated (Figure 1C [\[4\]](#)), and 5 amino acids (L-asparagine, L-serine, L-aspartate,  $\beta$ -alanine and  $\gamma$ -aminobutyric acid) were downregulated (Figure 1D [\[5\]](#)) in the *Salmonella*-infected group. Consistent with previous findings, intracellular serine concentrations were downregulated due to the reprogramming of macrophage glucose metabolism during *Salmonella* infection (Jiang et al., 2021 [\[6\]](#)). *Salmonella* can use host-derived aspartate and asparagine for growth in macrophages (Popp et al., 2015 [\[7\]](#)); therefore, the decrease in intracellular aspartate and asparagine upon *Salmonella* infection is likely due to their utilization by bacteria. Interestingly,  $\beta$ -alanine concentrations were also downregulated in the *Salmonella*-infected group (Figure 1D [\[8\]](#)), suggesting that intracellular *Salmonella* may use host-derived  $\beta$ -alanine for growth.

To investigate whether host-derived  $\beta$ -alanine can promote intracellular *Salmonella* replication, we added an additional 0.5, 1, 2, 4 mM  $\beta$ -alanine (Schneider, Krämer, & Burkovski, 2004 [\[9\]](#)) to the culture medium (RPMI) of RAW264.7 cells and then infected them with *Salmonella* to test the influence of  $\beta$ -alanine addition on the ability of *Salmonella* to replicate in macrophages. The results showed that the replication of *Salmonella* in RAW264.7 cells significantly ( $P < 0.001$ )



**Figure 1.**

### Host-derived β-alanine promotes *Salmonella* replication inside macrophages.

(A) Schematic workflow for targeted metabolomics investigation of mock- and *Salmonella*-infected (STM) mouse RAW264.7 macrophages. Picture materials were used from bioicons (<https://bioicons.com/>). (B) Principal component analysis (PCA) score plots of metabolic profiles in the mock- and *Salmonella*-infected (STM) groups (n = 4 biologically independent samples). (C) The concentrations of upregulated amino acids in the mock- and *Salmonella*-infected groups (n = 4 biologically independent samples). (D) The concentrations of downregulated amino acids in the mock- and *Salmonella*-infected groups (n = 4 biologically independent samples). (E) Fold intracellular replication (20 h vs. 2 h) of *Salmonella* WT in RAW264.7 cells in the presence of 0.5, 1, 2, 4 mM β-alanine. Data are presented as the mean ± SD, n = 3 independent experiments. (F) Growth curves of *Salmonella* WT and the *argT* mutant (*ΔargT*) in N-minimal (left) and M9 minimal (right) medium supplemented with β-alanine (1 mM) as the sole carbon source. Data are presented as mean ± SD, n = 4 independent experiments. Statistical significance was assessed using two-sided Student's *t*-test (C, D) and one-way ANOVA (E).

increased with the addition of 1, 2, or 4 mM  $\beta$ -alanine (**Figure 1E**). Furthermore,  $\beta$ -alanine enhanced *Salmonella* intracellular replication in a dose-dependent manner (**Figure 1E**). The results suggest that host-derived  $\beta$ -alanine facilitates *Salmonella* replication inside macrophages. We then investigated whether  $\beta$ -alanine-mediated *Salmonella* growth promotion is due to the changes in antimicrobial activity of the macrophages. We observed that the addition of 1 mM  $\beta$ -alanine did not influence the ROS (reactive oxygen species) and RNS (reactive nitrogen species) levels in *Salmonella*-infected RAW264.7 cells (**Figure 1-figure Supplement 1**). Flow cytometry analysis indicated that the addition of 1 mM  $\beta$ -alanine did not affect the percentage of pro-inflammatory M1 macrophages (CD86+) and anti-inflammatory M2 macrophages (CD163+) during *Salmonella* infection (**Figure 1-figure Supplement 2**), implying that the addition of  $\beta$ -alanine to macrophages does not change their immune response. Combining these results, we can further infer that *Salmonella* use host-derived  $\beta$ -alanine for intracellular growth.


Direct validation of *Salmonella* using host-derived  $\beta$ -alanine for intracellular growth requires a mutant that has a defect in  $\beta$ -alanine uptake. *Escherichia coli* uptakes  $\beta$ -alanine via the transporter protein CycA (Schneider et al., 2004). However, the *Salmonella*  $\Delta$ cycA mutant was able to use  $\beta$ -alanine as the sole carbon source for growth in minimal medium (**Figure 1-figure Supplement 3**, **Figure 1-figure Supplement 4**), indicating that CycA is not a transporter for  $\beta$ -alanine in *Salmonella*. Consistent with these results, mutation of cycA did not influence the replication of *Salmonella* in RAW264.7 cells (**Figure 1-figure Supplement 5**) or colonization in mouse systemic tissues (liver and spleen; **Figure 1-figure Supplement 6**). In *E. coli*, GabP transports  $\gamma$ -aminobutyric acid (GABA), a structural analogue of  $\beta$ -alanine, and may also transport  $\beta$ -alanine (Pavić et al., 2021). Nevertheless, *Salmonella*  $\Delta$ gabP mutant displayed no growth defect in minimal medium with  $\beta$ -alanine as the sole carbon source (**Figure 1-figure Supplement 7**, **Figure 1-figure Supplement 8**), indicating that GabP is not involved in  $\beta$ -alanine uptake in *Salmonella*. Strikingly, the  $\Delta$ argT mutant—defective in arginine uptake—showed markedly decreased growth in the minimal medium with  $\beta$ -alanine as the sole carbon source (**Figure 1F**), suggesting that ArgT also transports  $\beta$ -alanine in *Salmonella*. It has been reported that ArgT is essential for *Salmonella* replication within macrophages and full virulence *in vivo* (Das et al., 2010). Given that ArgT is involved in both arginine and  $\beta$ -alanine uptake (as verified in this study), whether the attenuated virulence of the  $\Delta$ argT mutant is due to a deficiency in  $\beta$ -alanine or arginine requires further investigation.

## De novo $\beta$ -alanine synthesis is critical for *Salmonella* replication inside macrophages

*Salmonella* can *de novo* synthesize  $\beta$ -alanine via the decarboxylation of L-aspartate, which is catalyzed by L-aspartate decarboxylase (PanD) (**Figure 2A**) and is reportedly the rate-limiting step of  $\beta$ -alanine generation (Begley et al., 2001; Schmitzberger et al., 2003; West et al., 1985). To further assess the role of  $\beta$ -alanine in *Salmonella* intracellular replication, we analyzed the expression level of the *Salmonella* panD gene in macrophages and the impact of panD mutation on the ability of *Salmonella* to replicate in macrophages. Quantitative real-time PCR (qRT[PCR]) assays revealed that the expression level of panD was significantly ( $P < 0.01$ ) greater in RAW264.7 cells than in RPMI-1640 medium (**Figure 2B**). Increased expression of panD was also observed in N-minimal medium, a widely used medium that mimics the conditions inside macrophages, as revealed by qRT[PCR] and bioluminescent reporter assays (**Figure 2C, D**). These results demonstrate that panD expression is enhanced during *Salmonella* growth inside macrophages, suggesting a relationship between panD expression and intracellular *Salmonella* growth.

We then constructed the panD mutant strain  $\Delta$ panD and compared the replication ability of the  $\Delta$ panD strain and the *Salmonella* Typhimurium 14028s wild type (WT) strain in RAW264.7 cells. Gentamicin protection assays showed that the replication of  $\Delta$ panD in RAW264.7 cells decreased 2.4-fold at 20 h postinfection compared with that of the WT strain ( $P < 0.001$ ), while





**Figure 2.**

### **De novo β-alanine synthesis is critical for *Salmonella* replication inside macrophages.**


(A) Scheme of β-alanine and the downstream CoA biosynthesis pathway in *Salmonella*. (B) qRT[PCR analysis of the expression of the *Salmonella panD* gene in RAW264.7 cells (8 h postinfection) and RPMI-1640 medium. (C) qRT[PCR analysis of the expression of the *Salmonella panD* gene in N-minimal medium and LB medium. (D) Expression of the *panD*-lux transcriptional fusion in N-minimal medium and LB medium. Luminescence values were normalized to 10<sup>5</sup> bacterial CFUs. (E) Fold intracellular replication (20 h vs. 2 h) of *Salmonella* Typhimurium 14028s wild type (WT), the *panD* mutant (Δ*panD*) and the complemented strain (*cpanD*) in RAW264.7 cells. (F) Number of intracellular *Salmonella* WT, Δ*panD*, and *cpanD* strains per RAW264.7 cell at 2 and 20 h postinfection. The number of intracellular bacteria per infected cell was estimated in random fields, n = 80 cells per group from 3 independent experiments. (G) Representative immunofluorescence images of *Salmonella* WT, Δ*panD*, and *cpanD* in RAW264.7 cells at 20 h postinfection (green, *Salmonella*; blue, nuclei; scale bars, 50 μm). Images are representative of three independent experiments. (H) Replication of *Salmonella* WT and Δ*panD* in RAW264.7 cells in the presence or absence of 1 mM β-alanine. The data are presented as the mean ± SD, n = 3 (B–E, H) independent experiments. Statistical significance was assessed using two-sided Student's *t*-test (B–D) or one-way ANOVA (E, F, H). ns, not Significant.

complementation of  $\Delta panD$  with the *panD* gene restored the replication ability of the mutant strain in RAW264.7 cells (**Figure 2E**). Immunofluorescence analysis revealed that the number of  $\Delta panD$  in each infected RAW264.7 cell was comparable to that of the WT strain at the initial infection stage (2 h), but at 20 h postinfection, the number of  $\Delta panD$  in each infected RAW264.7 cell was significantly ( $P < 0.0001$ ) lower than that of the WT strain (**Figure 2F, G**). These results indicate that *panD* contributes to *Salmonella* replication in macrophages. The growth rates of  $\Delta panD$  in LB medium and RPMI medium resembled those of the WT (**Figure 2-figure Supplement 1**, **Figure 2-figure Supplement 2**), indicating that the impaired intracellular replication ability of the mutant was not due to a growth defect. Moreover, the replication defect of  $\Delta panD$  in RAW264.7 cells was relieved by the addition of 1 mM  $\beta$ -alanine to the RPMI medium (**Figure 2H**). Furthermore, we examined the role of  $\beta$ -alanine synthesis in the intracellular replication of *Salmonella* within another typical serovar, *Salmonella enterica* serovar Typhi (S. Typhi), a serovar specific to humans and the causative agent of typhoid fever (de Jong, Parry, van der Poll, & Wiersinga, 2012). We found that the replication of S. Typhi  $\Delta panD$  in human THP-1 macrophages was reduced by 2.6-fold compared to the S. Typhi Ty2 WT strain ( $P < 0.01$ ) (**Figure 2-figure Supplement 3**), suggesting that *panD* also facilitates S. Typhi replication within human macrophages.

These data collectively suggest that  $\beta$ -alanine synthesis is critical for *Salmonella* replication inside macrophages.

## De novo $\beta$ -alanine synthesis is critical for systemic *Salmonella* infection in mice

As replication in macrophages is a key determinant of systemic *Salmonella* infection, we reasoned that  $\beta$ -alanine synthesis also influences *Salmonella* systemic infection *in vivo*. To determine whether  $\beta$ -alanine influences systemic *Salmonella* infection, we conducted mouse infection assays using intraperitoneal (i.p.) injection. This method allows *Salmonella* to disseminate directly to systemic sites via the lymphatic and bloodstream system, bypassing the need for intestinal invasion (Pandeya et al., 2023; Silva et al., 2016). BALB/c mice were infected by i.p. injection of 5,000 CFU of WT,  $\Delta panD$ , or the complemented strain *cpanD*. The survival rate, body weight, bacterial burden in the liver and spleen, and liver histopathological alterations of the infected mice were measured (**Figure 3A**). The WT-infected mice exhibited high lethality and marked loss of body weight within 5 days, and all mice died within 9 days of infection (**Figure 3B, C**). In contrast, the  $\Delta panD$ -infected mice displayed significantly improved survival rates and body weights, and no mice died within the 10-day surveillance period (**Figure 3B, C**). Consistent with these results, the bacterial burden in the liver and spleen of the  $\Delta panD$ -infected mice was significantly decreased, and the body weight was significantly increased compared with that of the WT-infected mice on day 3 postinfection (**Figure 3D**). Complementation of  $\Delta panD$  with *panD* significantly decreased the survival rate and body weight of infected mice but significantly increased the bacterial burden in the liver and spleen of the infected mice (**Figure 3B-D**). Through immunofluorescence staining, we examined the bacterial count in liver macrophages of mice infected with WT,  $\Delta panD$ , and complemented strain. The results showed that the bacterial count in each macrophage from  $\Delta panD$ -infected mice was significantly ( $P < 0.0001$ ) lower than that in WT-infected mice, on day 5 post-infection. Complementation of  $\Delta panD$  with *panD* restored the bacterial count in each macrophage to WT level (**Figure 3E**). Furthermore, H&E staining revealed increased aggregation of inflammatory cells and pyknosis in the livers of the WT-infected mice on day 5 postinfection, while these histopathological alterations were obviously reduced in the livers of  $\Delta panD$ -infected mice (**Figure 3F**). Taken together, these results reveal that  $\beta$ -alanine synthesis is critical for systemic *Salmonella* infection in mice.



**Figure 3.**

### ***De novo* β-alanine synthesis is critical for systemic *Salmonella* infection in mice.**

(A) Schematic illustration of the mouse infection assays. Picture materials were used from bioicons (<https://bioicons.com/>). (B, C) Survival curves (B) and body weight dynamics (C) of mice infected i.p. with *Salmonella* WT,  $\Delta panD$ , or *cpanD*.  $n = 10$  mice per group. (D) Liver and spleen bacterial burdens and body weights of mice infected with *Salmonella* WT,  $\Delta panD$ , or *cpanD* on day 3 postinfection.  $n = 7$  mice per group. (E) Representative immunofluorescence images and intracellular bacterial counts of *Salmonella* WT,  $\Delta panD$ , and *cpanD* in mouse liver at 5 days post-infection (green, *Salmonella*; blue, nuclei; scale bars, 50  $\mu$ m). Images are representative of three independent experiments. The number of intracellular bacteria per infected cell was estimated in random fields, with  $n = 80$  cells per group from 3 independent experiments. (F) Representative H&E-stained liver sections from mice that were left uninfected or infected with *Salmonella* WT,  $\Delta panD$ , or *cpanD* on day 5 postinfection. Arrows indicate severe inflammatory cell infiltration in the mouse liver. Images are representative of three independent experiments. The data are presented as the mean  $\pm$  SD (B-E). Statistical significance was assessed using the log-rank Mantel-Cox test (B), two-sided Student's *t*-test (C), or one-way ANOVA (D, E).




## β-Alanine is involved in the regulation of several metabolic pathways in *Salmonella*

To explore the mechanism(s) associated with β-alanine-mediated promotion of *Salmonella* replication in macrophages and *in vivo* virulence, we performed RNA sequencing (RNA-seq) to reveal the differences in gene transcripts between *Salmonella* WT and  $\Delta panD$  strains grown in N-minimal medium. PCA plot of the global transcriptomic profiles clearly demonstrated separation between the WT and  $\Delta panD$  strains (**Figure 4A** [↗](#)). Remarkable transcriptional changes were observed due to the mutation of *panD*. Compared with those in the WT strain, 1379 genes were differentially expressed in the  $\Delta panD$  strain, with 561 upregulated genes and 618 downregulated genes (fold change  $\geq 2$  and  $P$  value  $< 0.05$ ; **Figure 4B** [↗](#)). Gene Ontology (GO) enrichment analysis revealed that the differentially expressed genes (DEGs) were mainly involved in the metabolism and biosynthesis of several amino acids (including arginine, leucine, histidine, and branched amino acids), carboxylic acid metabolism, small molecule biosynthesis, and aerobic respiration (**Figure 4C** [↗](#)). Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis also revealed a high frequency of terms related to metabolism, including amino acid metabolism, lipid metabolism, carbohydrate metabolism, energy metabolism, and nucleotide metabolism (**Figure 4D** [↗](#)). These data collectively indicate that β-alanine is involved in the regulation of a series of metabolic pathways in *Salmonella*.

Further analysis of the downregulated DEGs (activated by *PanD*) revealed that mutation of *panD* decreased the expression of genes involved in 7 pathways that are associated with the virulence of *Salmonella* or other bacterial pathogens, including methionine metabolism, fatty acid β-oxidation, histidine biosynthesis, and the transport of zinc, galactose, potassium, and polyamine (**Figure 4E** [↗](#)). Zinc and potassium uptake are associated with the virulence of *Salmonella* (zinc acquisition promotes *Salmonella* Typhimurium virulence in mice, and potassium acquisition promotes *Salmonella* Enteritidis virulence in chickens) (Ammendola et al., 2007 [↗](#); Battistoni, Ammendola, Chiancone, & Ilari, 2017 [↗](#); Ilari et al., 2016 [↗](#); Liu et al., 2013 [↗](#)), while the remaining 5 pathways are involved in the pathogenicity of other pathogens (Basavanna et al., 2013 [↗](#); de Paiva et al., 2016 [↗](#); Feldman et al., 2016 [↗](#); Lauriano et al., 2004 [↗](#); Martínez-Gutián et al., 2019 [↗](#)). In addition, the expression of the LysR-type transcriptional regulator *LeuO*, which activates the expression of the *leuABCD* leucine synthesis operon and numerous virulence genes in *Salmonella* Typhimurium (Dillon et al., 2012 [↗](#); Guadarrama, Villaseñor, & Calva, 2014 [↗](#); Hernández-Lucas et al., 2008 [↗](#)), was also downregulated in the  $\Delta panD$  strain (**Figure 4E** [↗](#)). In line with the decreased expression of *leuO*, the expression of *leuABCD* was downregulated in the  $\Delta panD$  strain (**Figure 4E** [↗](#)).

We selected 16 downregulated DEGs (including the regulatory gene *leuO* and genes from the above 7 pathways) for qRT[PCR analysis. The results showed that the expression of all 16 genes significantly ( $P < 0.05$ ) decreased in the  $\Delta panD$  mutant compared with the WT strain (**Figure 4F** [↗](#)), and complementation of  $\Delta panD$  with *panD* restored the gene expression to the WT level (**Figure 4F** [↗](#)), thus confirming the positive regulation of these pathways and *LeuO* by β-alanine.

*Salmonella* pathogenesis largely depends on virulence genes encoded by *Salmonella* pathogenicity islands (SPIs), with SPI-1 to SPI-5 being well-characterized for their involvement in *Salmonella* virulence (Han et al., 2024 [↗](#)). SPI-2 gene expression is essential for *Salmonella* replication in macrophages and systemic infection (Deng et al., 2017 [↗](#)), yet the expression of SPI-2 genes was unaffected by *panD* mutation (**Figure 4-figure Supplement 1** [↗](#)). Moreover, the gene expression of four other virulence-associated SPIs, SPI-1, SPI-3, SPI-4, and SPI-5, is also unaffected by *panD* mutation (**Figure 4-figure Supplement 2** [↗](#), **Figure 4-figure Supplement 3** [↗](#)).



**Figure 4.**

### **$\beta$ -Alanine is involved in the regulation of several metabolic pathways in *Salmonella*.**

(A) Principal component analysis (PCA) score plots of transcriptomic profiles of *Salmonella* WT and  $\Delta panD$  (n = 3 biologically independent samples). (B) Volcano plot of the differentially expressed genes (DEGs) in *Salmonella* WT versus  $\Delta panD$ . The upper right section (red dots) indicates the upregulated DEGs, and the upper left section (green dots) indicates the downregulated DEGs. (C) Gene Ontology (GO) enrichment analysis of DEGs. Bubble chart showing the top 20 enriched GO terms. (D) Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis of DEGs. (E) Expression of the downregulated pathways (activated by PanD) is shown in the Z score-transformed heatmap, with red representing higher abundance and blue representing lower abundance. (F) qRT-PCR analysis of the mRNA levels of 16 selected downregulated DEGs in *Salmonella* WT,  $\Delta panD$ , and  $cpanD$ . The data are presented as the mean  $\pm$  SD, n = 3 independent experiments. Statistical significance was assessed using two-way ANOVA.

Taken together, these data suggest that  $\beta$ -alanine might promote *Salmonella* intracellular replication and virulence by activating virulence-associated pathway(s) or activating the virulence-associated regulator LeuO, rather than by activating the expression of virulence genes encoded within pathogenicity islands.

## **$\beta$ -Alanine promotes *Salmonella* virulence *in vivo* partially by increasing the expression of zinc transporter genes**

Next, we inactivated the 7 downregulated pathways mentioned above, as well as the regulatory gene *leuO* in *Salmonella*, to uncover the mechanism(s) by which  $\beta$ -alanine promotes *Salmonella* virulence *in vivo*. Mouse infection assays revealed that mutations in *fadAB*, *metR*, *hisABCDGHL*, *kdpABC*, *mglABC*, and *potFGHI*, which are associated with fatty acid  $\beta$ -oxidation, methionine metabolism, histidine biosynthesis, potassium uptake, galactose uptake, and polyamine uptake, respectively, did not influence *Salmonella* colonization in the mouse liver or spleen (**Figure 5A, B**) or the body weight of infected mice (**Figure 5C**). Interestingly, although LeuO has been reported to be associated with the regulation of a diverse set of virulence factors (Dillon et al., 2012; Guadarrama et al., 2014), mutation of the regulatory gene *leuO* did not influence *Salmonella* colonization in the mouse liver or spleen (**Figure 5A, B**) or the body weight of infected mice (**Figure 5C**). In contrast, mutation of the zinc transporter gene *znuA* significantly decreased *Salmonella* colonization in the mouse liver and spleen (**Figure 5D**, left and middle panels); this result is consistent with previous studies (Ammendola et al., 2007; Battistoni et al., 2017; Ilari et al., 2016). Accordingly, the body weight of  $\Delta znuA$ -infected mice was significantly ( $P < 0.001$ ) greater than that of the WT-infected mice (**Figure 5D**, right panel). These results indicate that  $\beta$ -alanine might promote *Salmonella* virulence *in vivo* by promoting zinc uptake.

To test this hypothesis, we initially evaluated the zinc content in the livers of WT- and  $\Delta panD$ -infected mice, on day 3 post-infection. We observed that the zinc concentration in the macrophages of  $\Delta panD$ -infected mouse livers was 3.2-fold higher than in those of WT-infected mice ( $P < 0.0001$ ; **Figure 5E**), suggesting that the *panD* gene and  $\beta$ -alanine are crucial for *Salmonella* to obtain zinc from host cells.


Next, we constructed a double mutant,  $\Delta panD\Delta znuA$ , and compared colonization of the mouse liver and spleen of the double mutant to that of the single mutant,  $\Delta znuA$ . The results showed that colonization of the liver and spleen of infected mice by  $\Delta panD\Delta znuA$  was significantly lower than that of infected mice colonized by  $\Delta znuA$  (**Figure 5D**, left and middle panels). In agreement with these results, the body weight of  $\Delta panD\Delta znuA$ -infected mice was greater than that of  $\Delta znuA$ -infected mice (**Figure 5D**, right panel), suggesting that the contribution of *panD* to the virulence of *Salmonella* is partially dependent on *znuA*.

Collectively, these data indicate that  $\beta$ -alanine promotes *in vivo* virulence of *Salmonella* partially by increasing the expression of zinc transporter genes.

## **$\beta$ -Alanine promotes *Salmonella* replication within macrophages partially by increasing the expression of zinc transporter genes**

To determine whether  $\beta$ -alanine influences *Salmonella* intracellular replication by acting on zinc transporters, the zinc content in RAW 264.7 macrophages infected with WT and  $\Delta panD$  was also examined. We observed that the zinc concentration in  $\Delta panD$ -infected RAW 264.7 cells increased by 1.8-fold compared to WT-infected cells ( $P < 0.0001$ ; **Figure 6A**), further indicating the *panD* gene and  $\beta$ -alanine are crucial for *Salmonella* to absorb zinc from macrophages.


We then analyzed the ability of  $\Delta znuA$  and  $\Delta panD\Delta znuA$  to replicate in RAW264.7 macrophages via gentamicin protection assays. The results showed that the replication of  $\Delta panD\Delta znuA$  in RAW264.7 cells was significantly reduced compared with that of the single mutant  $\Delta znuA$  (**Figure 6B**),



**Figure 5.**

**$\beta$ -Alanine promotes *Salmonella* virulence in vivo partially by increasing the expression of zinc transporter genes.**

(A, B, C) Liver (A) and spleen (B) bacterial burdens and body weight (C) of mice infected with *Salmonella* WT,  $\Delta fadAB$ ,  $\Delta metR$ ,  $\Delta hisABCDFGHL$ ,  $\Delta kdpABC$ ,  $\Delta mglABC$ ,  $\Delta potFGHI$ , or  $\Delta leuO$  on day 3 postinfection.  $n = 5$  mice per group. (D) Liver and spleen bacterial burdens and body weights of mice infected with *Salmonella* WT,  $\Delta panD$ ,  $\Delta znuA$  or  $\Delta panD\Delta znuA$  on day 3 postinfection.  $n = 5$  mice per group. (E) The zinc levels in the livers of mice infection with either *Salmonella* WT or  $\Delta panD$  for 3 days,  $n = 5$  mice per group. The data are presented as the mean  $\pm$  SD (A-E). Statistical significance was assessed using one-way ANOVA (A-D), two-sided Student's *t*-test (E). ns, not Significant.



**Figure 6.**

**$\beta$ -Alanine promotes *Salmonella* replication within macrophages partially by increasing the expression of zinc transporter genes.**

(A) The zinc levels in RAW264.7 cells after infection with *Salmonella* WT or  $\Delta$ panD for 8 h. (B) Replication of *Salmonella* WT,  $\Delta$ panD,  $\Delta$ znuA and  $\Delta$ panD $\Delta$ znuA in RAW264.7 cells. (C) Replication of *Salmonella* WT and  $\Delta$ panD in RAW264.7 cells in the presence or absence of 100  $\mu$ M ZnSO<sub>4</sub>. (D) Replication of *Salmonella* WT and  $\Delta$ znuA in RAW264.7 cells in the presence or absence of 1 mM  $\beta$ -alanine. The data are presented as the mean  $\pm$  SD, n = 3 independent experiments (A–D). Statistical significance was assessed using two-sided Student's *t*-test (A), one-way ANOVA (B–D). ns, not Significant. (E) Schematic model of  $\beta$ -alanine-mediated *Salmonella* replication inside macrophages. In macrophages, *Salmonella* acquires  $\beta$ -alanine both via the uptake of  $\beta$ -alanine from host macrophages and the *de novo* synthesis of  $\beta$ -alanine.  $\beta$ -alanine promotes the expression of zinc transporter genes *ZnuABC*, which facilitate the uptake of zinc by intracellular *Salmonella*, therefore promote *Salmonella* replication in macrophages and subsequent systemic infection.



implying that the contribution of *panD* to the intracellular replication of *Salmonella* is partially dependent on *znuA*. The addition of 100  $\mu$ M zinc to RPMI medium increased the replication of  $\Delta$ *panD* in RAW264.7 cells (**Figure 6C**), while the addition of 1 mM  $\beta$ -alanine to RPMI medium did not increase the replication of  $\Delta$ *znuA* (**Figure 6D**), suggesting that the impaired replication due to the decrease in  $\beta$ -alanine can be relieved by zinc supplementation.

Taken together, these data indicate that  $\beta$ -alanine promotes *Salmonella* replication within macrophages by increasing the expression of zinc transporter genes.

## Discussion

Replication within host macrophages is a crucial step for *Salmonella* to cause life-threatening systemic infection in the host (Bomjan, Zhang, & Zhou, 2019; Lathrop et al., 2015), while the crosstalk between *Salmonella* and macrophages at the metabolic interface is critical for intracellular *Salmonella* replication (Dandekar et al., 2014; Herrero-Fresno & Olsen, 2018; Lynch & Lesser, 2021; Rosenberg et al., 2021; Thompson, Fulde, & Tedin, 2018). Emerging evidence suggests that several metabolites affect the replication of *Salmonella* in macrophages. The promotion of intracellular replication by metabolites is possibly achieved in three ways: i) metabolites are utilized by *Salmonella* as nutrients for intracellular growth (Bowden, Rowley, Hinton, & Thompson, 2009; Eisenreich, Dandekar, Heesemann, & Goebel, 2010; J. Wang et al., 2021); ii) *Salmonella* senses metabolites as environmental cues to activate the expression of virulence genes (Jiang et al., 2021; X. Wang et al., 2023); and iii) metabolites can regulate the immune responses of macrophages (Michelucci et al., 2013; Peace & O'Neill, 2022; Yang & Cong, 2021). In this study, we demonstrated that *Salmonella* promotes its replication inside macrophages by utilizing both host- and bacterial-derived  $\beta$ -alanine (**Figure 6E**). We showed that  $\beta$ -alanine promotes *Salmonella* intracellular replication and systemic infection partially by increasing the expression of zinc transporter genes and therefore the uptake of zinc by intracellular *Salmonella* (**Figure 6E**). Therefore, this work identified another metabolite that can influence the replication of *Salmonella* in macrophages and illustrated the mechanism by which  $\beta$ -alanine promotes intracellular *Salmonella* replication.

We observed that *Salmonella*-infected macrophages contained lower  $\beta$ -alanine levels than mock-infected macrophages, while  $\beta$ -alanine supplementation in the cell medium increased the replication of *Salmonella* in macrophages, revealing that *Salmonella* uptakes host-derived  $\beta$ -alanine to promote intracellular replication. In addition, a deficiency in the biosynthesis of  $\beta$ -alanine (via mutation of the rate-limiting gene *panD*) reduced *Salmonella* replication in macrophages and systemic infection in mice, suggesting that *Salmonella* also utilizes bacterial-derived  $\beta$ -alanine to promote intracellular replication and pathogenicity. It is known that bacteria are quite stringent with their energy resources (Bergkessel, 2020; Bosdriesz, Molenaar, Teusink, & Bruggeman, 2015), while the results of this work indicate that either acquisition from the host or *de novo* synthesis of  $\beta$ -alanine is critical for *Salmonella* replication inside macrophages. We speculate that *Salmonella* relies on a large amount of  $\beta$ -alanine to efficiently replicate in macrophages, thereby highlighting the importance of  $\beta$ -alanine for *Salmonella* intracellular growth. Nevertheless, unlike the closely related species *E. coli*, which takes up  $\beta$ -alanine via the transporter protein CycA (Schneider et al., 2004), *Salmonella* does not use CycA to uptake  $\beta$ -alanine. GabP, the GABA transporter, is potentially involved in the uptake of  $\beta$ -alanine in *E. coli* (Pavić et al., 2021). However, *Salmonella* does not utilize GabP to uptake  $\beta$ -alanine either. Ultimately, we revealed that ArgT, the transporter of arginine, is involved in the transport of  $\beta$ -alanine in *Salmonella*. ArgT has been reported to be essential for *Salmonella* replication within macrophages and for full virulence *in vivo* (Das et al., 2010). However, the attenuated virulence of the  $\Delta$ *argT* mutant is due to a deficiency in  $\beta$ -alanine or arginine requires further investigation.

Several amino acids, including lysine, proline, arginine, aspartate and asparagine, have previously been reported to be associated with the pathogenicity of *Salmonella* (Popp et al., 2015 [↗](#); Steeb et al., 2013 [↗](#)). These amino acids are involved in the synthesis of proteins in *Salmonella*. In contrast,  $\beta$ -alanine is not incorporated into proteins but can participate in the regulation of bacterial activity through the synthesis of pantothenate and CoA (Webb et al., 2004 [↗](#); White et al., 2001 [↗](#); Yuan et al., 2022 [↗](#)). Accordingly, our transcriptome data showed that a deficiency in  $\beta$ -alanine biosynthesis affected the expression of *Salmonella* genes involved in a series of important metabolic pathways. Importantly, although  $\beta$ -alanine does not influence the gene expression of SPIs, it activates methionine metabolism; fatty acid  $\beta$ -oxidation; histidine biosynthesis; and the transport of zinc, galactose, potassium, and polyamine, which have been previously known to be associated with the virulence of *Salmonella* and other bacterial pathogens. Further analysis revealed that  $\beta$ -alanine promotes *Salmonella* intracellular replication and systemic infection partially by promoting the uptake of zinc. As for the other virulence-related metabolic pathways activated by  $\beta$ -alanine, methionine metabolism, fatty acid  $\beta$ -oxidation, and histidine biosynthesis contribute to the virulence of *Streptococcus pneumoniae* (Basavanna et al., 2013 [↗](#)), *Yersinia pestis* (Feldman et al., 2016 [↗](#)), and *Acinetobacter baumannii* (Martínez-Gutián et al., 2019 [↗](#)), respectively; the uptake of galactose and polyamine influences the pathogenicity of *Francisella tularensis* (Lauriano et al., 2004 [↗](#)) and the avian pathogenic *Escherichia coli* (de Paiva et al., 2016 [↗](#)), respectively; and the uptake of potassium is associated with the virulence of *Salmonella* Enteritidis in chickens (Liu et al., 2013 [↗](#)). However, blocking these pathways did not influence the systemic *Salmonella* Typhimurium infection in mice, implying that different bacterial pathogens adopt different virulence strategies to establish infection. Determining other mechanism(s) by which  $\beta$ -alanine promotes the intracellular replication and systemic infection of *Salmonella* require further investigation.

We observed that  $\beta$ -alanine also activates the expression of the LysR-type transcriptional regulator LeuO, which is known to regulate the expression of a wide variety of *Salmonella* genes that impact the stress response and virulence (Dillon et al., 2012 [↗](#); Guadarrama et al., 2014 [↗](#); Hernández-Lucas et al., 2008 [↗](#)). Typically, LeuO activates the synthesis of the quiescent porins OmpS1 and OmpS2, which are required for *Salmonella* virulence in mice (De la Cruz et al., 2007 [↗](#); Fernández-Mora, Puente, & Calva, 2004 [↗](#); Rodríguez-Morales et al., 2006 [↗](#)). Consistent with the positive regulation of OmpS1 and OmpS2 by LeuO, lack of *leuO* in *Salmonella* also attenuated virulence in a mouse model (Rodríguez-Morales et al., 2006 [↗](#)). However, the attenuated phenotypes of the *leuO* mutant in mice were not evident after i.p. injection relative to oral infection, as a previous report showed that the competitive index for the *leuO* mutant indicated approximately 1,000-fold reduced colonization in mouse systemic tissues after oral infection but much less reduced colonization after i.p. injection (less than 10-fold) (Rodríguez-Morales et al., 2006 [↗](#)). These results imply that LeuO might be predominantly associated with the invasion and intestinal infection of *Salmonella* but is weakly implicated in *Salmonella* intracellular replication and systemic infection. Therefore, it is not surprising that lack of *leuO* did not significantly affect *Salmonella* colonization in the systemic tissues of mice after i.p. injection, as revealed by our results.

Zinc is an essential micronutrient for all living organisms and is used as a cofactor for various enzymes and proteins (Bock, Müller, & Blankenfeldt, 2016 [↗](#); Ilari et al., 2016 [↗](#); Tan, Bramlett, & Lindahl, 2004 [↗](#)). In bacteria, zinc-binding proteins account for approximately 5% of the bacterial proteome and play crucial roles in bacterial metabolism and virulence (Andreini, Banci, Bertini, & Rosato, 2006 [↗](#)). Knockout of the zinc transporter ZnuABC reduces the virulence of *Salmonella*, *Campylobacter jejuni*, *Haemophilus ducreyi*, *Moraxella*, and urinary tract pathogenic *Escherichia coli* (UPEC) in the host (Ilari et al., 2016 [↗](#)). Moreover, zinc is also utilized by *Salmonella* to subvert the antimicrobial host defense of macrophages by inhibiting NF- $\kappa$ B activation and impairing NF- $\kappa$ B-dependent bacterial clearance (Jennings, Esposito, Rittinger, & Thurston, 2018 [↗](#); Wu et al., 2017 [↗](#)). Therefore, the efficient acquisition of zinc may be crucial for *Salmonella*'s survival and replication within macrophages, where zinc availability is limited (Ammendola et al., 2007 [↗](#); Ilari et al., 2016 [↗](#)). It has been reported that *Salmonella* utilizes the high-affinity ZnuABC zinc

transporter to optimize zinc availability in host cells (Ammendola et al., 2007 [↗](#)). In this study, we found that  $\beta$ -alanine can increase the expression of the zinc transporter genes *znuABC*, which could represent an additional mechanism for the efficient zinc uptake of *Salmonella* in macrophages. The results demonstrate a correlation between *Salmonella*  $\beta$ -alanine utilization and zinc uptake during intracellular infection and provide evidence that  $\beta$ -alanine can influence the macrophage immune response by acting on zinc uptake.

Overall, our findings suggest a model in which *Salmonella* exploits host- and bacterial-derived  $\beta$ -alanine to efficiently replicate in host macrophages and cause systemic disease. We propose that *Salmonella* requires a large amount of  $\beta$ -alanine during intracellular infection. The utilization of  $\beta$ -alanine promotes *Salmonella* uptake of the essential micronutrient zinc, which was previously shown to be required for the metabolic needs of intracellular *Salmonella* and to subvert the antimicrobial defense of macrophages by *Salmonella*. These observations provide new insight into *Salmonella* pathogenesis and the crosstalk between *Salmonella* and macrophages during intracellular infection. Considering that the *panD* gene is present in the genomes of all *Salmonella* strains and that mutation of *panD* markedly reduced *Salmonella* replication ability in macrophages, as well as virulence in the mouse model, this gene may be used as a potential target to control systemic *Salmonella* infection.

## Methods

### Ethics statement

Six-week-old female BALB/c mice were obtained from Beijing Vital River Laboratory Animal Technology (Beijing, China). Mice were housed in barrier facilities under specific pathogen-free conditions with a 12 h light/dark cycle at a temperature of  $24 \pm 2^\circ\text{C}$  and a relative humidity of  $50 \pm 5\%$ . Mice were fed a standard mouse chow diet, and they consumed food and water *ad libitum* throughout the experiment. All animal experiments were conducted in accordance with the policies of the Institutional Animal Care Committee of Nankai University (Tianjin, China) and performed under protocol no. 2021-SYDWLL-000029.

### Cell culture

The RAW264.7 mouse macrophage-like cell line (ATCC TIB-71) was obtained from the Shanghai Institute of Biochemistry and Cell Biology of the Chinese Academy of Sciences (Shanghai, China). Cells were cultured in RPMI-1640 medium (Gibco #11879020) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco #10100147) at  $37^\circ\text{C}$  with 5%  $\text{CO}_2$ . Cells were seeded in 24-well tissue culture plates at  $1 \times 10^5$  cells per well 24 h before infection.

### Bacterial strains, plasmids, and growth conditions

The bacterial strains and plasmids used in this study are listed in **Supplementary Table 1** [↗](#). The *Salmonella enterica* serovar Typhimurium (STM) strain ATCC 14028s was used as the WT strain throughout this study and for the construction of the mutants. Mutant strains were generated using the  $\lambda$  Red recombination system with the plasmid pSIM17 (Jiang et al., 2017 [↗](#)). To construct the complemented strain of  $\Delta\text{panD}$ , the amplified DNA fragments of the *panD* ORF and its upstream promoter were digested and inserted into the BamHI–EcoRI site of the low-copy-number plasmid pBR322. To generate the *panD-lux* reporter fusion, the amplification products of the *panD* promoter region were digested and cloned into the XhoI–BamHI site of the plasmid pMS402, which carries a promoter-less *luxCDABE* reporter gene cluster (Liang, Li, Dong, Surette, & Duan, 2008 [↗](#)). The sequences of primers used for the construction of the strains are listed in **Supplementary Table 2** [↗](#). All the strains were verified by PCR amplification and sequencing.

Bacterial strains were conventionally grown overnight in Luria–Bertani (LB) medium (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl) or in N-minimal medium (10  $\mu$ M MgCl<sub>2</sub>, 110  $\mu$ M KH<sub>2</sub>PO<sub>4</sub>, 7.5 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.5 mM K<sub>2</sub>SO<sub>4</sub>, 5 mM KCl, 38 mM glycerol, and 0.1% [w/v] casamino acids) supplemented with appropriate antibiotics at 37 °C with shaking at 180 rpm or on LB agar plates. All antibiotics were used at their standard concentrations (chloramphenicol, 25  $\mu$ g/mL; kanamycin, 50  $\mu$ g/mL; ampicillin, 100  $\mu$ g/mL; gentamicin, 10 or 100  $\mu$ g/mL) unless otherwise mentioned.

## Growth curve

Bacterial strains were conventionally grown overnight in LB medium. The next day, they were subcultured (1:100) in new LB medium and RPMI-1640 medium or subcultured in N-minimal medium supplemented with glycerol or  $\beta$ -alanine as the sole carbon source at 37 °C with shaking at 180 rpm. To measure the growth of bacteria, 200  $\mu$ L of the bacterial cultures were transferred to the microplate wells. The absorbance (OD<sub>600</sub>) of the bacteria was measured every half hour for 12 h with a Spark multimode microplate reader (Tecan).

## Bioluminescent reporter assays

STM WT carrying the *panD-luxCDABE* fusion plasmid was conventionally grown overnight in LB medium, and the next day, the cells were subcultured (1:100) in new LB medium or N-minimal medium for 8 h. The luminescence of the cultured bacteria (200  $\mu$ L) was measured with a Spark multimode microplate reader (Tecan). Moreover, the cultured bacteria (100  $\mu$ L) were serially diluted and plated on LB agar plates to estimate bacterial CFUs. Bacterial CFUs were used to normalize luminescence values.

## Salmonella infection of macrophages

Bacterial strains were conventionally grown overnight in LB medium to the late stationary phase, and the next day, the bacteria were diluted to  $2 \times 10^6$  CFUs/mL and opsonized in RPMI-1640 medium supplemented with 10% FBS for 15 min. The macrophage monolayers were infected with the opsonized bacteria culture (0.5 mL/well, multiplicity of infection (MOI) = 10) and centrifuged at 800 $\times$ g for 5 min to synchronize infection. The infected cells were incubated for 30 min at 37 °C in 5% CO<sub>2</sub> and then washed three times with 1 $\times$  PBS. Fresh RPMI-1640 medium containing 100  $\mu$ g/mL gentamicin was added to the infected cells to kill extracellular bacteria. After 1 h, fresh RPMI-1640 medium containing 10  $\mu$ g/mL gentamicin was added to the infected cells for the remainder of the experiment. To assess the intracellular growth of *Salmonella*, the infected cells were lysed with 1% Triton X-100 at 2 hpi and 20 hpi, and the abundance of the intracellular bacteria CFUs was estimated on LB agar plates. The relative fold replication of intracellular bacterial strains was denoted as the CFUs recovered at 20 hpi relative to those at 2 hpi. The relative fold change in replication was normalized to the number of RAW264.7 cells. When indicated, 1 mM  $\beta$ -alanine or 100  $\mu$ M ZnSO<sub>4</sub> were added after 1 h of gentamicin treatment.

## Targeted metabolomics analysis of amino acids in macrophages

Targeted metabolomics was conducted as previously described (Jiang et al., 2021 [DOI](#)), with the combined metabolites of infected cells and intracellular bacteria were extracted for analysis. The metabolites have been confirmed to be dominated by the host metabolites (Jiang et al., 2021 [DOI](#)). RAW264.7 cells were mock-infected or infected with STM WT for 8 h. The cells were harvested and washed with precooled PBS solution to remove the medium. Cellular metabolites were extracted using ice-cold extraction solvent (40:40:20 vol/vol/vol acetonitrile:methanol:water, 0.1 M formic acid), incubated at –20 °C for 20 min, and then centrifuged for 10 min at 12,000 $\times$ g and 4 °C to obtain the supernatant. Subsequently, the supernatant was transferred to an LC–MS vial and analyzed using ultrahigh-performance liquid chromatography (Acquity; Waters, Milford, MA, USA) coupled with mass spectrometry (Q Exactive Hybrid Quadrupole-Orbitrap; Thermo Fisher Scientific,

Waltham, MA, USA). Metabolites were separated with a Luna NH<sub>2</sub> column (2 mm × 100 mm, 3 μm particle size; Phenomenex). Mobile phase A was 20 mM ammonium acetate (pH 9.0), and mobile phase B was acetonitrile containing 0.1% formic acid. The flow rate was 0.4 ml/min.

Xcalibur 4.0 software (Thermo Fisher) was used for data acquisition and processing. Metabolite identification was achieved by high-resolution mass and retention time matching to authentic standards. The absolute quantification of amino acids was performed using the standard curve method, and the values were normalized to the cell number. Four biological replicates of each sample were analyzed.

## Immunofluorescence staining

RAW264.7 cells were infected with STM WT or  $\Delta$ *panD* and the complemented strain *cpaND* as described above. After 2 and 20 h of cultivation, the infected cells were fixed for 15 min in 4% paraformaldehyde, washed in PBS, and permeabilized with 0.1% Triton X-100 in PBS for 15 min. The fixed samples were blocked in 5% bovine serum albumin for 30 min, followed by staining with a FITC-conjugated anti-*Salmonella* antibody (1:100 dilution, Abcam #ab20320) for 1 h at room temperature in the dark. The nuclei were then stained with DAPI (Invitrogen #D21490) for 2 min. A confocal laser scanning microscope (Zeiss LSM800) and ZEN 2.3 software (blue edition) were used to acquire and analyze the cell images (Objective lense: 40×; The number of intracellular bacteria per infected cell was estimated in random fields by Fiji-ImageJ).

## RNA isolation

RNA was extracted from *Salmonella* strains cultured in N-minimal medium or LB medium. To investigate the expression of the *Salmonella panD* gene inside macrophages, we obtained RNA from intracellular bacteria in RAW264.7 cells at 8 h postinfection and from bacteria in RPMI medium. RNA was extracted using an EASYspinPlus bacterial RNA rapid extraction kit (Aidlab #RN0802) according to the manufacturer's protocol. RNA quantity and purity were determined using a NanoDrop 2000 spectrophotometer (NanoDrop Technologies). RNA samples were stored at -80 °C before use.

## Quantitative real-time PCR (qRT-PCR)

According to the manufacturer's protocols, qRT-PCR was performed using 2× RealStar Power SYBR qPCR Mix (Genstar #A304) in a QuantStudio 5 Real Time PCR system (Applied Biosystems). cDNA was synthesized using a StarScript III RT Kit (Genstar #A232). Each sample was subjected to qRT-PCR in triplicate. The expression level of the 16S *rRNA* gene was used to normalize that of the target genes. We estimated the expression of each target gene using the 2<sup>-ΔΔCt</sup> method.

## RNA sequencing and analyses

The STM WT and  $\Delta$ *panD* strains were conventionally grown overnight in LB medium, subcultured (1:100) in N-minimal medium for 8 h and then collected by centrifugation for RNA extraction. Sequencing libraries were generated using the NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs) according to the manufacturer's instructions, and sequencing was conducted using the Illumina HiSeq 2000 platform at Shanghai Majorbio Bio-Pharm Technology Co., Ltd. (Shanghai, China). The sequencing data were deposited in the NCBI Sequence Read Archive under accession number (SRA, PRJNA1124637). The clean reads were mapped to the STM ATCC 14028 reference genome (CP001363 and CP001362) by using the short-sequence alignment software Bowtie 2. Gene expression was evaluated using the fragments per kilobase of transcript per million mapped reads (FPKM) method. DEGs in the *panD* mutant relative to the WT were determined using the R statistical package software EdgeR. The thresholds for statistically significant differences were set to a fold change  $\geq 2$  or  $\leq 0.5$  and a false discovery rate (FDR)  $\leq 0.05$ . P values were adjusted using the Benjamini-Hochberg procedure for controlling the FDR. Enrichment analysis of DEGs was conducted using GO and KEGG enrichment analyses.



## Mouse infection

*Salmonella* strains were conventionally cultured overnight in LB medium, and the next day, they were subcultured (1:100) in new LB medium and grown at 37 °C with shaking at 200 rpm to stationary phase ( $OD_{600} \sim 2$ ). The bacteria were diluted to  $5 \times 10^4$  CFUs/mL in 0.9% NaCl. Groups of BALB/c mice were infected i.p. with 0.1 mL of the NaCl suspension. For survival assays, we recorded and monitored the mortality and body weight of the infected mice daily for 10 days. To analyze the bacterial burden of the mouse liver and spleen, we weighed the infected mice first on day 3 postinfection and then harvested the liver and spleen. The liver and spleen of infected mice were homogenized in ice-cold PBS, serially diluted, and plated on LB plates containing the appropriate antibiotics to determine bacterial CFUs. H&E staining of the mouse liver was performed to investigate the histopathological alterations in the liver of infected mice. To evaluate the histopathological alterations in the mouse liver, we harvested the liver of the infected mice on day 5 postinfection. The mouse liver was washed with 0.9% NaCl, fixed with 10% neutral formalin for 48 h and subsequently processed for routine paraffin embedding. Paraffin-embedded tissues were sectioned at a thickness of 5  $\mu$ m and then stained with hematoxylin (Sangon Biotech #E607318) and eosin (Sangon Biotech #E607318) for histopathological examination. The stained sections were then examined by light microscopy (Leica DM2500 LED).

## Flow cytometry

RAW264.7 cells were infected with *Salmonella* WT for 8 h, in the absence or presence of 1 mM  $\beta$ -alanine, which was added to the infected cells at 1 h post-infection. The infected cells were fixed for 15 min in 4% paraformaldehyde, washed in PBS, and permeabilized with 0.1% Triton X-100 in PBS for 15 min. The fixed samples were blocked in 5% bovine serum albumin for 30 min, followed by staining with an anti-CD86 antibody (Abcam, ab119857), an anti-CD163 antibody (Abcam, ab182422) for 30 min, and a goat anti-rat IgG H&L (Alexa Fluor® 488) (Abcam, ab150165), a donkey anti-rabbit IgG H&L (Alexa Fluor® 647) (Abcam, ab150075) for 30 min in the dark. The infected cells were analyzed using a BD FACSaria Flow Cytometer (BD Biosciences).

## Assessment of intracellular ROS/ RNS levels

RAW264.7 cells were infected with *Salmonella* WT for 8 hours, either in the absence or presence of 1 mM  $\beta$ -alanine. The  $\beta$ -alanine was added to the infected cells at 1 h post-infection. Following the infection, the cells were washed three times with PBS, and the levels of ROS and RNS were measured using a ROS fluorescence probe (BBoxiProbe® O06, Bestbio, BB-46051) and an RNS fluorescence probe (BBoxiProbe® O52, Bestbio, BB-470567), respectively, in accordance with the manufacturer's protocol. The ROS/ RNS levels were normalized to the cell count.

## Assessment of zinc levels in mouse liver and RAW 264.7 macrophages

To assess the zinc level in mouse liver, mice were i.p. infected with approximately 5000 CFUs of *Salmonella* WT or  $\Delta$ *panD* for 3 days, with three mice in each injection group. Following infection, mouse livers were collected and then dissociated into single cells using the gentleMACS/Mouse Liver Dissociation Kit (Miltenyi Biotec, 130-105-807) in accordance with the manufacturer's protocol. The zinc concentration in mouse liver was determined using a zinc fluorescence probe (Zinquin ethyl ester, MKBio, MX4516) following the manufacturer's instructions and normalized by the cell count.

To assess the zinc level in RAW 264.7 macrophages, RAW 264.7 cells were infected with *Salmonella* WT or  $\Delta$ *panD* for 8 hours. The infected cells were washed three times with PBS, and zinc concentration was measured using a zinc fluorescence probe (Zinquin ethyl ester, MKBio,

MX4516) following the manufacturer's protocol. The zinc concentration was normalized by the cell count.

### Statistical analysis

The data are presented as the mean  $\pm$  SD. All in vitro experiments were conducted in duplicate and repeated at least three times ( $n \geq 3$ ). Mouse assays were performed twice, with at least two mouse ( $n \geq 2$ ) in each injection group, and the combined data from the two experiments was used for statistical analysis. Statistical analyses were performed using GraphPad InStat software (version GraphPad Prism 9.5.1, San Diego, CA, USA) with two-sided Student's *t* tests, one-way ANOVA, two-way ANOVA, log-rank Mantel–Cox tests, or Mann[Whitney U tests according to the test requirements (as stated in the figure legends). A *P* value  $< 0.05$  indicated a statistically significant difference. ns represents no statistical significance.

### Data availability statement

The RNA-seq data generated in this study have been deposited in the NCBI Sequence Read Archive (SRA) database under the accession number PRJNA1124637. All other data associated with this study are available in the main text and supplementary materials. Source data are provided in Supplementary Data 1.

### Figure supplements

Figure 1-figure supplement 1.

The levels of ROS and RNS in RAW264.7 cells after infection with *Salmonella* WT for 8 hours, in the presence or absence of 1 mM  $\beta$ -alanine.

Data are presented as mean  $\pm$  SD, n = 3 independent experiments. Statistical significance was assessed using two-sided Student's *t*-test. ns, not Significant.





Figure 1-figure supplement 2.

Flow cytometry analysis was conducted to determine the percentage of pro-inflammatory M1 macrophages (CD86+) and anti-inflammatory M2 macrophages (CD163+).


RAW264.7 cells were infected with *Salmonella* WT for 8 hours, in the presence or absence of 1 mM  $\beta$ -alanine. Subsequently, the infected cells were collected for flow cytometry analysis. Representative dot plots and quantification of M1 (CD86+) and M2 (CD163+) macrophages are displayed in the left and right panels, respectively. Data are presented as mean  $\pm$  SD, n = 3 independent experiments. Statistical significance was assessed using two-way ANOVA. ns, not Significant.



### Figure 1-figure supplement 3.

Growth curves of *Salmonella* WT and the *cycA* mutant ( $\Delta cycA$ ) in N-minimal medium supplemented with  $\beta$ -alanine (1 mM) as the sole carbon source.

Data are presented as mean  $\pm$  SD, n = 3 independent experiments.



### Figure 1-figure supplement 4.

Growth curves of *Salmonella* WT and the *cycA* mutant ( $\Delta cycA$ ) in M9 minimal medium supplemented with  $\beta$ -alanine (1 mM) as the sole carbon source.

Data are presented as mean  $\pm$  SD, n = 3 independent experiments.




Figure 1-figure supplement 5.

**Fold intracellular replication (20 h vs. 2 h) of *Salmonella* wild-type and  $\Delta cycA$  in RAW264.7 cells.**

Data are presented as mean  $\pm$  SD, n = 4 independent experiments. Statistical significance was assessed using two-way ANOVA. ns, not Significant.




Figure 1-figure supplement 6.

**Liver and spleen bacterial burdens, and body weight of mice infected with *Salmonella* wild-type (WT) and  $\Delta cycA$  mutant ( $\Delta cycA$ ), at day 3 post-infection.**

n = 5 mice per group. Statistical significance was assessed using Mann-Whitney U test. ns, not Significant.






Figure 1-figure supplement 7.

Growth curves of *Salmonella* WT and the *gabP* mutant ( $\Delta gabP$ ) in N-minimal medium supplemented with  $\beta$ -alanine (1 mM) as the sole carbon source.

Data are presented as mean  $\pm$  SD, n = 3 independent experiments.




Figure 1-figure supplement 8.

Growth curves of *Salmonella* WT and the *gabP* mutant ( $\Delta gabP$ ) in M9 minimal medium supplemented with  $\beta$ -alanine (1 mM) as the sole carbon source.

Data are presented as mean  $\pm$  SD, n = 3 independent experiments.




Figure 2-figure supplement 1.

Growth curves of *Salmonella* WT, *panD* mutant ( $\Delta panD$ ) and the complemented strain (*cpanD*) in LB medium.

Data are presented as mean  $\pm$  SD, n = 4 independent experiments




Figure 2-figure supplement 2.

Growth curves of *Salmonella* WT, *panD* mutant ( $\Delta panD$ ) and the complemented strain (*cpanD*) in RPMI-1640 medium (B).

Data are presented as mean  $\pm$  SD, n = 4 independent experiments.




Figure 2-figure supplement 3.

Fold intracellular replication (20 h vs. 2 h) of *Salmonella enterica* serovar Typhi Ty2 wild type (WT), the *panD* mutant ( $\Delta panD$ ) in human THP-1 monocyte like cell line (ATCC TIB-22).

Data are presented as mean  $\pm$  SD, n = 4 independent experiments. Statistical significance was assessed using two-sided Student's *t*-test. ns, not Significant.




Figure 4-figure supplement 1.

Expression of SPI-2 is shown in the Z-score transformed heatmap, with orange representing higher and blue representing lower abundance.




Figure 4-figure supplement 2.

Expression of SPI-1 is shown in the Z-score transformed heatmap, with orange representing higher and blue representing lower abundance.





Figure 4-figure supplement 3.

Expression of SPI-3, SPI-4, and SPI-5 genes is shown in the Z-score transformed heatmap, with orange representing higher and blue representing lower abundance.



| Plasmid or strain                    | Genotype or description                                                                                      | Source                |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------|-----------------------|
| Plasmid                              |                                                                                                              |                       |
| pSim17                               | temperature-induced $\lambda$ -Red recombinase system; $\text{Bla}^{\text{S}}$                               | Laboratory collection |
| pKD3                                 | template plasmid containing the Cm cassette for $\lambda$ -Red recombination; $\text{Cm}^{\text{R}}$         | Laboratory collection |
| pKD4                                 | template plasmid containing the Km cassette for $\lambda$ -Red recombination; $\text{Km}^{\text{R}}$         | Laboratory collection |
| pBR322                               | low-copy-number expression vector; $\text{Ap}^{\text{R}}$                                                    | Laboratory collection |
| pMS402                               | Containing a promoterless <i>luxCDABE</i> reporter gene cluster; $\text{Km}^{\text{R}}$                      | Laboratory collection |
| pBR322- <i>panD</i>                  | pBR322 carrying the STM <i>panD</i> gene; $\text{Ap}^{\text{R}}$                                             | This study            |
| pMS402- <i>panD</i> strain           | pMS402 carrying the STM <i>panD</i> promoter; $\text{Km}^{\text{R}}$                                         | This study            |
| wild-type (WT)                       | wild-type <i>Salmonella enterica</i> serovar Typhimurium strain                                              | Laboratory collection |
| STM                                  | ATCC 14028s                                                                                                  | Laboratory collection |
| $\Delta\text{panD}$                  | WT strain <i>panD</i> ::Cm; $\text{Cm}^{\text{R}}$                                                           | This study            |
| <i>cpanD</i>                         | $\Delta\text{panD}$ containing plasmid pBR322- <i>panD</i> ; $\text{Cm}^{\text{R}}$ , $\text{Ap}^{\text{R}}$ | This study            |
| $\Delta\text{cycA}$                  | WT strain <i>cycA</i> ::Cm; $\text{Cm}^{\text{R}}$                                                           | This study            |
| $\Delta\text{fadAB}$                 | WT strain <i>fadAB</i> ::Cm; $\text{Cm}^{\text{R}}$                                                          | This study            |
| $\Delta\text{metR}$                  | WT strain <i>metR</i> ::Cm; $\text{Cm}^{\text{R}}$                                                           | This study            |
| $\Delta\text{leuABCD}$               | WT strain <i>leuABCD</i> ::Cm; $\text{Cm}^{\text{R}}$                                                        | This study            |
| $\Delta\text{leuO}$                  | WT strain <i>leuO</i> ::Km; $\text{Km}^{\text{R}}$                                                           | This study            |
| $\Delta\text{hisABCD}^{\text{FGHL}}$ | WT strain <i>hisABCD</i> <sup>FGHL</sup> ::Cm; $\text{Cm}^{\text{R}}$                                        | This study            |
| $\Delta\text{kdpABC}$                | WT strain <i>kdpABC</i> ::Cm; $\text{Cm}^{\text{R}}$                                                         | This study            |
| $\Delta\text{mglABC}$                | WT strain <i>mglABC</i> ::Cm; $\text{Cm}^{\text{R}}$                                                         | This study            |
| $\Delta\text{potFGHI}$               | WT strain <i>potFGHI</i> ::Cm; $\text{Cm}^{\text{R}}$                                                        | This study            |
| $\Delta\text{znuA}$                  | WT strain <i>znuA</i> ::Cm; $\text{Cm}^{\text{R}}$                                                           | This study            |
| $\Delta\text{argT}$                  | WT strain <i>argT</i> ::Km; $\text{Km}^{\text{R}}$                                                           | This study            |
| $\Delta\text{gabP}$                  | WT strain <i>gabP</i> ::Cm; $\text{Cm}^{\text{R}}$                                                           | This study            |

## Supplementary Table 1

### Strains and plasmids used in this study

| Targets                                                                | Primer Sequences (5'-3') |                                                                                                                                                                                   |
|------------------------------------------------------------------------|--------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primers used for the construction and identification of mutant strains |                          |                                                                                                                                                                                   |
| <i>ΔcycA</i>                                                           | construction             | F: GTATCATAGACCAAAGGCCGTAGAGCCCGCACAAACACAG<br>ACAGGTACAGGAAGAAAAACGTGTAGGCTGGAGCTGCTTC<br>R: CCTCCAAATCAACGTTACGCTGTAAGCCCGGTAAGCGCC<br>AGCGCCACCGGGCAAAACAACATATGAATATCCTCCTTAG |
|                                                                        | identification           | F: TGTGAGCTAATTCGCATTATCAAAG<br>R: ACGGCATTAATGAGATGATTGATGA                                                                                                                      |
| <i>ΔpanD</i>                                                           | construction             | F: CGGATTCGCTGGAAACCATGTCGCGGCTGATCAGCAACC<br>AGCCGCAGGGATAAGGACTAGTGTAGGCTGGAGCTGCTT<br>R: TTTCCAAAGCAGGCCACAAGCGCCTGCTTAGCCAAGGTA<br>AACGACAGGGTAAAGAAGTTCATATGAATATCCTCCTTAG   |
|                                                                        | identification           | F: CGGTTATAACGCGACCGCGATCAAA<br>R:GCTGCTGGAGCTGACGGAAACCAGC                                                                                                                       |
| <i>ΔfadAB</i>                                                          | construction             | F: GGATTCGCTCAGTTGCCGCTGCGCTGCAATGCGAGTTATT<br>TAGGGGATATTATCTTTGAGTGTAGGCTGGAGCTGCTTC<br>R: AATACACACTTCGCTTCATCTGGTACGACCAGATCACTTT<br>GTGGATTGAGGAGCTGACCATATGAATATCCTCCTTAG   |
|                                                                        | identification           | F: AAACGGCAACTAAACTGTTTCCCGT<br>R: AGTGATTCCATTTTTACCTTCTG                                                                                                                        |
| <i>ΔmetR</i>                                                           | construction             | F: TAAAGCTTGGCCAGTGGGGCTGCTGATGCCAGATGGCAA<br>GATTAAACCAGCAGTCCTGCGGTGTAGGCTGGAGCTGCTTC<br>R: GAAAGTCCTTCACTTCGCCATGAACAAATTGCGCTTGAG<br>GAATATACAGTACCTTTACACATATGAATATCCTCCTTAG |
|                                                                        | identification           | F: CCATTGCAGCGTGGTTTCCGGCAGC<br>R: TGCAAGCCGAACTATTAGAGTTTT                                                                                                                       |
| <i>ΔleuABCD</i>                                                        | construction             | F: GAATATAATGGCAACCGGGTGATATTGCATAACCTGTAG<br>GCCCGGTAAGGCGTCAAGGGGTGTAGGCTGGAGCTGCTTC<br>R: CCGTTGCGCGGGTTTTTTATGCCTGACGCAAGGCGCCCC<br>TGGAGACAAGGACCACATCCATATGAATATCCTCCTTAG   |
|                                                                        | identification           | F: TTTTTATTTCGGCTGTTTCTGGAT<br>R: GTTGACATTAAACGGCATATCCAGT                                                                                                                       |
| <i>ΔleuO</i>                                                           | construction             | F: CCAGAAAAAGGGAGTTAAGCGTGACAGTGGAGTTAAATGTG<br>TAGGCTGGAGCTGCTTC<br>R: ATAAACCAGAAATTGTTTCTGATTTATCTGCCCGGTTCATATG<br>AATATCCTCCTTAG                                             |
|                                                                        | identification           | F: CATTATGAATCGCAATGGTGTGAC<br>R: CTCCGTCTGAATCACACCTGGT                                                                                                                          |

## Supplementary Table 2

### Primers involved in this study



| Targets                                                                | Primer Sequences (5'-3') |                                                                                                                                                                                  |
|------------------------------------------------------------------------|--------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Primers used for the construction and identification of mutant strains |                          |                                                                                                                                                                                  |
| <i>ΔhisABCD<sub>FGH</sub></i><br><i>L</i>                              | construction             | F: GGCCTAAAAGTGGTTTAGGTTAAAAGGTATCAAATGAATA<br>AGCATTTCATCGGAATTTTTGTGTAGGCTGGAGCTGCTTC<br>R: GATGGCTGGCATCAGGCCATCGGTTTTTTCCAGTCCAGC<br>TCGCGGCGTGTGTGCTCTGCATATGAATATCCTCCTTAG |
|                                                                        | identification           | F: CCTAACCAACCTAAACCGACAATTG<br>R: AAACGCTGTTTCGTGCGTGAGAAGA                                                                                                                     |
| <i>ΔkdpABC</i>                                                         | construction             | F: GTGGGGCGGGGGCGTTTGTTCACAAGCGATCCGGATC<br>AGGACGCAAGGGTTCGTCGTGTGTAGGCTGGAGCTGCTTC<br>R:CTGGTTTTCTTACTTTTAGGTTATCTGGTCTATGCCCTGATTAAT<br>GCGGAGGCGTTCGCATATGAATATCCTCCTTAG     |
|                                                                        | identification           | F: CAGAATGGTCAGCCCGTCAAG<br>R: ATACTTTTTTACACTCCGCC                                                                                                                              |
| <i>ΔmgIABC</i>                                                         | construction             | F: CCGGCGCGCAGTTTCGCTGCGCCGGTTATCTGATAAGC<br>AAGGCAATAGGTCTGGATAAGTGTAGGCTGGAGCTGCTTC<br>R: GTGCTGAACAGCCGGGCATTTTTTACGCTATACCCCTACA<br>TAATAAAACCGAGCTACCCATATGAATATCCTCCTTAG   |
|                                                                        | identification           | F: AAAAAGAGCACGCGTTAACACTGTA<br>R: AGGCTCTAAATACGCTTCGGC                                                                                                                         |
| <i>ΔpotFGHI</i>                                                        | construction             | F: ATGTTTTAAACCACGCCTAATGGGTTTCATTTGTTAACGGAT<br>TTCAGAAGGAAAGCGATGGTGTAGGCTGGAGCTGCTTC<br>R: GAAAATGATGCCGCGACCCGCGCGCAACGCATTGCCAT<br>AGTGGAAGATTTTAGTGGCTCATATGAATATCCTCCTTAG |
|                                                                        | identification           | F: AAAGGCGTTTTTTAATCTGGGCTAT<br>R: TATTATAATCATCAGCACATCGAG                                                                                                                      |
| <i>ΔznuA</i>                                                           | construction             | F: AGATTATTAATGCCAGGGCGACAGAGCGGGCTATCTGTT<br>GCACGTATTCACTTCCTCGGTGTAGGCTGGAGCTGCTTC<br>R: AAGTGATAGAATGTTATAATATCACATTCACACATTCATT<br>ACGATGATTAGTCGCATTCATATGAATATCCTCCTTAG   |
|                                                                        | identification           | F: AGGGAACGAATCTCGCTTTTCTC<br>R: ATTCAAGCGACAGCTCAGAGAGGAC                                                                                                                       |
| <i>ΔargT</i>                                                           | construction             | F: CGTGATAGTTCCCGAGCGCGCGCGCTTATCCCTTCCCGTGTA<br>GGCTGGAGCTGCTTC<br>R: GTAAAACATAAGAAAATGACGCCACTTGAGGGGTATGTCATAT<br>GAATATCCTCCTTAG                                            |
|                                                                        | identification           | F: CAAAGCATTCACCTTGTGAGCCGG<br>R: CATACCTGCTATCTTCAACATCAGG                                                                                                                      |

**Supplementary Table 2** (continued)

| Targets                                                                | Primer Sequences (5'-3')                           |                                                               |
|------------------------------------------------------------------------|----------------------------------------------------|---------------------------------------------------------------|
| Primers used for the construction and identification of mutant strains |                                                    |                                                               |
| <i>ΔgabP</i>                                                           | construction                                       | F: CCGGTAAACGAGCGCAATAACAATAAAGAGGTTTTAGGGTGTAGGCTGGAGCTGCTTC |
|                                                                        |                                                    | R: TTACTCCTGAGAAGCTTCCCGGCTGACGCCGGAATAACATATGAATATCCTCCTTAG  |
|                                                                        | identification                                     | F: CTTTGATGAGGCGAAACAAGC                                      |
|                                                                        |                                                    | R: AATGTCATTTTAAAGCCAGCGAT                                    |
| Primers used for the construction of complementary strains             |                                                    |                                                               |
| pBR322- <i>panD</i>                                                    | F: CCCTTTCGTCTTCAAGAATTCGCTGCTGGAGCTGACGGAAC       | R: TGCCTCCGGCGTAGAGGATCCTCAGGCAACCTGTACCGGAAT                 |
| pBR322                                                                 | F: GACACGGAAATGTTGAATACTCATA                       | R: CAAGGAATGGTGCATGCAAG                                       |
| Primers used for luciferase reporter system                            |                                                    |                                                               |
| pMS402- <i>panD</i>                                                    | F: CGAGGCCCTTTCGTCTTCACCTCGAGTCAAGGAAATCCTGGTCAATG | R: ATTTTGGCGCGCAACTAGAGGATCCCGGTAATCGTGAGCTGC                 |
| pMS402                                                                 | F: CTGTCCTCTTGATCAGATCTT                           | R: TTATTCCATTACAATCAAT                                        |
| Primers used for RT-qPCR analysis                                      |                                                    |                                                               |
| <i>16s</i>                                                             | F:GAAAGCGTGGGAGCAAAC                               | R:ACATGCTCCACCGCTTG TG                                        |
| <i>panD</i>                                                            | F:AAGGTTCTGTGCCATTGAC                              | R:ACCGTTCACCGAGATGATTC                                        |
| <i>fadH</i>                                                            | F:GAAACCCACATGCCGATAAC                             | R:AAACTGACCGCCAATTTACAC                                       |
| <i>fadB</i>                                                            | F:GTTGCTGTCTGGAAGAAGG                              | R: TACTGCTGCGCCATATCAAG                                       |
| <i>fadI</i>                                                            | F:TCTGACGCTTTTGTGACATGC                            | R:TTCATGCAACGTTTGGGTTA                                        |
| <i>fadD</i>                                                            | F:AAGATCTGGCTTTCCTGCAA                             | R:CCCAGCTCGATAAAAAGCAG                                        |
| <i>fadE</i>                                                            | F:AATTGGTCGTCGTCACTTC                              | R: GCGTCTACCGACAAACATT                                        |
| <i>metR</i>                                                            | F:TTTGGTGATGACGTCGGATA                             | R:CGGGTAAATCAACAGCGTTT                                        |
| <i>hisG</i>                                                            | F:GTGCGTGATGATGACATTCC                             | R:GGAGGTGCGGATATGAGGTA                                        |
| <i>hisF</i>                                                            | F:TGTATTGTCGTCGGGATTGA                             | R:TTTCAACTGCGTCAGATCG                                         |
| <i>leuD</i>                                                            | F:CCGGAATTCGTGTTGAACTT                             | R:GTAGAAGATGTCGGCGAAGC                                        |
| <i>leuO</i>                                                            | F:AGATATGGGCAAAACACAGC                             | R: CCTGACGGA                                                  |

**Supplementary Table 2** (continued)

| Targets                           | Primer Sequences (5'-3')                          |
|-----------------------------------|---------------------------------------------------|
| Primers used for RT-qPCR analysis |                                                   |
| <i>mgIA</i>                       | F:AGCGATTAACCACGGTTTTG<br>R:ATTGGGTGTCGCTTTTCATC  |
| <i>mgIB</i>                       | F:AGAATCCGGTGTGATTCAGG<br>R:CAGCCAGGCATCCATCTTAT  |
| <i>kdpA</i>                       | F:ACTAATCTGGCGCAGATGCT<br>R:ACCCCTCCATGTTGACGCTAC |
| <i>kdpB</i>                       | F:TCGGTAAGCAGATGCTGATG<br>R:CCAGCGGGATCAGAAAAATA  |
| <i>potG</i>                       | F:ATTACGAGCATCCGACGAC<br>R:AGAGGCATCTGCATCGACTT   |
| <i>potF</i>                       | F:TCTGATGGTCGTCTCTGTCTG<br>R:GCCTTCCAGCACTTCGTTAG |
| <i>znuA</i>                       | F:TAAACCGCTTGGGTTTCATC<br>R:AACGGTTTTACATCGGCAAG  |

**Supplementary Table 2** (continued)

## Acknowledgements

This work was supported by the National Natural Science Foundation of China (NSFC) Program (grant no. 32170110, 32270191), the Natural Science Foundation of Tianjin (grant no. 22JCYBJC01060), and the Fundamental Research Funds for the Central Universities, Nankai University (grant no. 63241588, 63243161).

## Additional information

### Author contributions

LJ designed the research; SM, YS, XW, HG, RL, TY, CK, and JC performed the research; BY provided technical support and insights; SM and BY analyzed the data; and LJ and SM wrote the manuscript.

## Additional files

**Figure source data.** [🔗](#)

## References

- Ammendola S., Pasquali P., Pistoia C., Petrucci P., Petrarca P., Rotilio G., Battistoni A (2007) **High-affinity Zn<sup>2+</sup> uptake system ZnuABC is required for bacterial zinc homeostasis in intracellular environments and contributes to the virulence of *Salmonella enterica*** *Infect Immun* **75**:5867–5876 <https://doi.org/10.1128/iai.00559-07>
- Andreini C., Banci L., Bertini I., Rosato A (2006) **Zinc through the three domains of life** *J Proteome Res* **5**:3173–3178 <https://doi.org/10.1021/pr0603699>
- Basavanna S., Chimalapati S., Maqbool A., Rubbo B., Yuste J., Wilson R. J., ..., Brown J. S. (2013) **The effects of methionine acquisition and synthesis on *Streptococcus pneumoniae* growth and virulence** *PLoS One* **8**:e49638 <https://doi.org/10.1371/journal.pone.0049638>
- Battistoni A., Ammendola S., Chiancone E., Ilari A (2017) **A novel antimicrobial approach based on the inhibition of zinc uptake in *Salmonella enterica*** *Future Med Chem* **9**:899–910 <https://doi.org/10.4155/fmc-2017-0042>
- Begley T. P., Kinsland C., Strauss E (2001) **The biosynthesis of coenzyme A in bacteria** *Vitam Horm* **61**:157–171 [https://doi.org/10.1016/s0083-6729\(01\)61005-7](https://doi.org/10.1016/s0083-6729(01)61005-7)
- Bellinger P. M (2014) **β-Alanine supplementation for athletic performance: an update** *J Strength Cond Res* **28**:1751–1770 <https://doi.org/10.1519/jsc.0000000000000327>
- Bergkessel M (2020) **Regulation of protein biosynthetic activity during growth arrest** *Curr Opin Microbiol* **57**:62–69 <https://doi.org/10.1016/j.mib.2020.07.010>
- Bock T., Müller R., Blankenfeldt W (2016) **Crystal structure of AibC, a reductase involved in alternative de novo isovaleryl coenzyme A biosynthesis in *Myxococcus xanthus*** *Acta Crystallogr F Struct Biol Commun* **72**:652–658 <https://doi.org/10.1107/s2053230x16011146>
- Boldyrev A. A., Aldini G., Derave W (2013) **Physiology and pathophysiology of carnosine** *Physiol Rev* **93**:1803–1845 <https://doi.org/10.1152/physrev.00039.2012>
- Bomjan R., Zhang M., Zhou D (2019) **YshB promotes intracellular replication and is required for *salmonella* virulence** *J Bacteriol* **201**:e00314–19 <https://doi.org/10.1128/jb.00314-19>
- Bosdriesz E., Molenaar D., Teusink B., Bruggeman F. J (2015) **How fast-growing bacteria robustly tune their ribosome concentration to approximate growth-rate maximization** *Febs j* **282**:2029–2044 <https://doi.org/10.1111/febs.13258>
- Bowden S. D., Rowley G., Hinton J. C., Thompson A (2009) **Glucose and glycolysis are required for the successful infection of macrophages and mice by *Salmonella enterica* serovar Typhimurium** *Infect Immun* **77**:3117–3126 <https://doi.org/10.1128/iai.00093-09>
- Brumell J. H., Tang P., Mills S. D., Finlay B. B (2001) **Characterization of *Salmonella*-induced filaments (Sifs) reveals a delayed interaction between *Salmonella*-containing vacuoles and late endocytic compartments** *Traffic* **2**:643–653 <https://doi.org/10.1034/j.1600-0854.2001.20907.x>

- Bumann D., Schothorst J (2017) **Intracellular Salmonella metabolism** *Cell Microbiol* **19**:10.1111/cmi.12766 <https://doi.org/10.1111/cmi.12766>
- Castanheira S., García-Del Portillo F (2017) **Salmonella populations inside host cells** *Front Cell Infect Microbiol* **7**:432 <https://doi.org/10.3389/fcimb.2017.00432>
- Dandekar T., Fieselmann A., Fischer E., Popp J., Hensel M., Noster J (2014) **Salmonella-how a metabolic generalist adopts an intracellular lifestyle during infection** *Front Cell Infect Microbiol* **4**:191 <https://doi.org/10.3389/fcimb.2014.00191>
- Das P., Lahiri A., Lahiri A., Sen M., Iyer N., Kapoor N., ..., Chakravortty D. (2010) **Cationic amino acid transporters and Salmonella Typhimurium ArgT collectively regulate arginine availability towards intracellular Salmonella growth** *PLoS One* **5**:e15466 <https://doi.org/10.1371/journal.pone.0015466>
- Davaapil H., Tsuchiya Y., Gout I (2014) **Signalling functions of coenzyme A and its derivatives in mammalian cells** *Biochem Soc Trans* **42**:1056–1062 <https://doi.org/10.1042/bst20140146>
- de Jong H. K., Parry C. M., van der Poll T., Wiersinga W. J. (2012) **Host-pathogen interaction in invasive Salmonellosis** *PLoS Pathog* **8**:e1002933 <https://doi.org/10.1371/journal.ppat.1002933>
- De la Cruz M. A., Fernández-Mora M., Guadarrama C., Flores-Valdez M. A., Bustamante V. H., Vázquez A., Calva E. (2007) **LeuO antagonizes H-NS and StpA-dependent repression in Salmonella enterica ompS1** *Mol Microbiol* **66**:727–743 <https://doi.org/10.1111/j.1365-2958.2007.05958.x>
- de Paiva J. B., da Silva L. P., Casas M. R., Conceição R. A., Nakazato G., de Pace F., ..., da Silveira W. D. (2016) **In vivo influence of in vitro up-regulated genes in the virulence of an APEC strain associated with swollen head syndrome** *Avian Pathol* **45**:94–105 <https://doi.org/10.1080/03079457.2015.1125995>
- Deng W., Marshall N. C., Rowland J. L., McCoy J. M., Worrall L. J., Santos A. S., ..., Finlay B. B. (2017) **Assembly, structure, function and regulation of type III secretion systems** *Nat Rev Microbiol* **15**:323–337 <https://doi.org/10.1038/nrmicro.2017.20>
- Dillon S. C., Espinosa E., Hokamp K., Ussery D. W., Casadesús J., Dorman C. J (2012) **LeuO is a global regulator of gene expression in Salmonella enterica serovar Typhimurium** *Mol Microbiol* **85**:1072–1089 <https://doi.org/10.1111/j.1365-2958.2012.08162.x>
- Eisenreich W., Dandekar T., Heesemann J., Goebel W (2010) **Carbon metabolism of intracellular bacterial pathogens and possible links to virulence** *Nat Rev Microbiol* **8**:401–412 <https://doi.org/10.1038/nrmicro2351>
- Fàbrega A., Vila J (2013) **Salmonella enterica serovar Typhimurium skills to succeed in the host: virulence and regulation** *Clin Microbiol Rev* **26**:308–341 <https://doi.org/10.1128/cmr.00066-12>
- Feldman M., Harbeck M., Keller M., Spyrou M. A., Rott A., Trautmann B., ..., Krause J. (2016) **A High-Coverage Yersinia pestis Genome from a Sixth-Century Justinianic Plague Victim** *Mol Biol Evol* **33**:2911–2923 <https://doi.org/10.1093/molbev/msw170>
- Fernández-Mora M., Puente J. L., Calva E (2004) **OmpR and LeuO positively regulate the Salmonella enterica serovar Typhi ompS2 porin gene** *J Bacteriol* **186**:2909–2920 <https://doi.org/10.1128/JB.186.10.2909-2920.2004>



[.org/10.1128/jb.186.10.2909-2920.2004](https://doi.org/10.1128/jb.186.10.2909-2920.2004)

Fields P. I., Swanson R. V., Haidaris C. G., Heffron F (1986) **Mutants of *Salmonella* Typhimurium that cannot survive within the macrophage are avirulent** *Proc Natl Acad Sci U S A* **83**:5189–5193 <https://doi.org/10.1073/pnas.83.14.5189>

Figueira R., Holden D. W (2012) **Functions of the *Salmonella* pathogenicity island 2 (SPI-2) type III secretion system effectors** *Microbiology (Reading)* **158**:1147–1161 <https://doi.org/10.1099/mic.0.058115-0>

Gojković Z., Sandrini M. P., Piskur J (2001) **Eukaryotic beta-alanine synthases are functionally related but have a high degree of structural diversity** *Genetics* **158**:999–1011 <https://doi.org/10.1093/genetics/158.3.999>

Gout I (2019) **Coenzyme A: a protective thiol in bacterial antioxidant defence** *Biochem Soc Trans* **47**:469–476 <https://doi.org/10.1042/bst20180415>

Guadarrama C., Villaseñor T., Calva E (2014) **The subtleties and contrasts of the *LeuO* regulator in *Salmonella* Typhi: implications in the immune response** *Front Immunol* **5**:581 <https://doi.org/10.3389/fimmu.2014.00581>

Han J., Aljahdali N., Zhao S., Tang H., Harbottle H., Hoffmann M., ..., Foley S. L. (2024) **Infection biology of *Salmonella enterica*** *EcoSal Plus* **12**:eesp00012023 <https://doi.org/10.1128/ecosalplus.esp-0001-2023>

Hernández-Lucas I., Gallego-Hernández A. L., Encarnación S., Fernández-Mora M., Martínez-Batallar A. G., Salgado H., ..., Calva E (2008) **The *LysR*-type transcriptional regulator *LeuO* controls expression of several genes in *Salmonella enterica* serovar Typhi** *J Bacteriol* **190**:1658–1670 <https://doi.org/10.1128/jb.01649-07>

Herrero-Fresno A., Olsen J. E (2018) ***Salmonella* Typhimurium metabolism affects virulence in the host - A mini-review** *Food Microbiol* **71**:98–110 <https://doi.org/10.1016/j.fm.2017.04.016>

Hoffman J. R., Varanoske A., Stout J. R (2018) **Effects of  $\beta$ -Alanine supplementation on carnosine elevation and physiological performance** *Adv Food Nutr Res* **84**:183–206 <https://doi.org/10.1016/bs.afnr.2017.12.003>

Huerta Ojeda Á., Tapia Cerda C., Poblete Salvatierra M. F., Barahona-Fuentes G., Jorquera Aguilera C. (2020) **Effects of beta-Alanine supplementation on physical performance in aerobic-anaerobic transition zones: a systematic review and meta-analysis** *Nutrients* **12**:2490 <https://doi.org/10.3390/nu12092490>

Ilari A., Pescatori L., Di Santo R., Battistoni A., Ammendola S., Falconi M., ..., Chiancone E. (2016) ***Salmonella enterica* serovar Typhimurium growth is inhibited by the concomitant binding of Zn(II) and a pyrrolyl-hydroxamate to ZnuA, the soluble component of the ZnuABC transporter** *Biochim Biophys Acta* **1860**:534–541 <https://doi.org/10.1016/j.bbagen.2015.12.006>

Jennings E., Esposito D., Rittinger K., Thurston T. L. M (2018) **Structure-function analyses of the bacterial zinc metalloprotease effector protein GtgA uncover key residues required for deactivating NF- $\kappa$ B** *J Biol Chem* **293**:15316–15329 <https://doi.org/10.1074/jbc.RA118.004255>

- Jiang L., Feng L., Yang B., Zhang W., Wang P., Jiang X., Wang L. (2017) **Signal transduction pathway mediated by the novel regulator *LoiA* for low oxygen tension induced *Salmonella* Typhimurium invasion** *PLoS Pathog* **13**:e1006429 <https://doi.org/10.1371/journal.ppat.1006429>
- Jiang L., Wang P., Song X., Zhang H., Ma S., Wang J., ..., Wang L. (2021) ***Salmonella* Typhimurium reprograms macrophage metabolism via T3SS effector *SopE2* to promote intracellular replication and virulence** *Nat Commun* **12**:879 <https://doi.org/10.1038/s41467-021-21186-4>
- Kehl A., Noster J., Hensel M. (2020) **Eat in or Take out? Metabolism of Intracellular *Salmonella enterica*** *Trends Microbiol* **28**:644–654 <https://doi.org/10.1016/j.tim.2020.03.005>
- Kim S., Lee K. S., Pak G. D., Excler J. L., Sahastrabuddhe S., Marks F., ..., Mogasale V. (2019) **Spatial and temporal patterns of typhoid and paratyphoid fever outbreaks: a worldwide review, 1990–2018** *Clin Infect Dis* **69**:S499–s509 <https://doi.org/10.1093/cid/ciz705>
- LaRock D. L., Chaudhary A., Miller S. I. (2015) ***Salmonellae* interactions with host processes** *Nat Rev Microbiol* **13**:191–205 <https://doi.org/10.1038/nrmicro3420>
- Lathrop S. K., Binder K. A., Starr T., Cooper K. G., Chong A., Carmody A. B., Steele-Mortimer O. (2015) **Replication of *Salmonella enterica* serovar Typhimurium in human monocyte-derived macrophages** *Infect Immun* **83**:2661–2671 <https://doi.org/10.1128/iai.00033-15>
- Lauriano C. M., Barker J. R., Yoon S. S., Nano F. E., Arulanandam B. P., Hassett D. J., Klose K. E. (2004) ***MglA* regulates transcription of virulence factors necessary for *Francisella tularensis* intraamoebae and intramacrophage survival** *Proc Natl Acad Sci U S A* **101**:4246–4249 <https://doi.org/10.1073/pnas.0307690101>
- Leung K. Y., Finlay B. B. (1991) **Intracellular replication is essential for the virulence of *Salmonella* Typhimurium** *Proc Natl Acad Sci U S A* **88**:11470–11474 <https://doi.org/10.1073/pnas.88.24.11470>
- Li W., Ren Q., Ni T., Zhao Y., Sang Z., Luo R., ..., Li S. (2023) **Strategies adopted by *Salmonella* to survive in host: a review** *Arch Microbiol* **205**:362 <https://doi.org/10.1007/s00203-023-03702-w>
- Liang H., Li L., Dong Z., Surette M. G., Duan K. (2008) **The *YebC* family protein PA0964 negatively regulates the *Pseudomonas aeruginosa* quinolone signal system and pyocyanin production** *J Bacteriol* **190**:6217–6227 <https://doi.org/10.1128/jb.00428-08>
- Liss V., Swart A. L., Kehl A., Hermanns N., Zhang Y., Chikkaballi D., ..., Hensel M. (2017) ***Salmonella enterica* Remodels the Host Cell Endosomal System for Efficient Intravacuolar Nutrition** *Cell Host Microbe* **21**:390–402 <https://doi.org/10.1016/j.chom.2017.02.005>
- Liu Y., Ho K. K., Su J., Gong H., Chang A. C., Lu S. (2013) **Potassium transport of *Salmonella* is important for type III secretion and pathogenesis** *Microbiology (Reading)* **159**:1705–1719 <https://doi.org/10.1099/mic.0.068700-0>
- Lynch J. P., Lesser C. F. (2021) **A host metabolite promotes *Salmonella* survival** *Science* **371**:344–345 <https://doi.org/10.1126/science.abf8414>
- Majowicz S. E., Musto J., Scallan E., Angulo F. J., Kirk M., O’Brien S. J., ..., Hoekstra R. M. (2010) **The global burden of nontyphoidal *Salmonella* gastroenteritis** *Clin Infect Dis* **50**:882–

889 <https://doi.org/10.1086/650733>

Martínez-Gutián M., Vázquez-Ucha J. C., Álvarez-Fraga L., Conde-Pérez K., Lasarte-Monterrubio C., Vallejo J. A., ..., Beceiro A. (2019) **Involvement of HisF in the persistence of *Acinetobacter baumannii* during a pneumonia infection** *Front Cell Infect Microbiol* **9**:310 <https://doi.org/10.3389/fcimb.2019.00310>

Michelucci A., Cordes T., Ghelfi J., Pailot A., Reiling N., Goldmann O., ..., Hiller K. (2013) **Immune-responsive gene 1 protein links metabolism to immunity by catalyzing itaconic acid production** *Proc Natl Acad Sci U S A* **110**:7820–7825 <https://doi.org/10.1073/pnas.1218599110>

Nozaki S., Webb M. E., Niki H (2012) **An activator for pyruvoyl-dependent l-aspartate  $\alpha$ -decarboxylase is conserved in a small group of the  $\gamma$ -proteobacteria including *Escherichia coli*** *Microbiologyopen* **1**:298–310 <https://doi.org/10.1002/mbo3.34>

Ohl M. E., Miller S. I (2001) **Salmonella: a model for bacterial pathogenesis** *Annu Rev Med* **52**:259–274 <https://doi.org/10.1146/annurev.med.52.1.259>

Pandeya A., Zhang Y., Cui J., Yang L., Li J., Zhang G., ..., Wei Y. (2023) **Inflammasome activation and pyroptosis mediate coagulopathy and inflammation in *Salmonella* systemic infection** *Microbiol Res* **275**:127460 <https://doi.org/10.1016/j.micres.2023.127460>

Pavić A., Ji Y., Serafini A., Garza-Garcia A., McPhillie M. J., Holmes A. O. M., ..., Postis V. L. G. (2021) **Functional Characterization of the  $\gamma$ -Aminobutyric Acid Transporter from *Mycobacterium smegmatis* MC(2) 155 Reveals Sodium-Driven GABA Transport** *J Bacteriol* **203**:e00642–20 <https://doi.org/10.1128/jb.00642-20>

Peace C. G., O'Neill L. A (2022) **The role of itaconate in host defense and inflammation** *J Clin Invest* **132**:e148548 <https://doi.org/10.1172/jci148548>

Pillay T. D., Hettiarachchi S. U., Gan J., Diaz-Del-Olmo I., Yu X. J., Muench J. H., ..., Pearson J. S. (2023) **Speaking the host language: how *Salmonella* effector proteins manipulate the host** *Microbiology (Reading)* **169**:001342 <https://doi.org/10.1099/mic.0.001342>

Popp J., Noster J., Busch K., Kehl A., Zur Hellen G., Hensel M (2015) **Role of host cell-derived amino acids in nutrition of intracellular *Salmonella enterica*** *Infect Immun* **83**:4466–4475 <https://doi.org/10.1128/iai.00624-15>

Rajashekar R., Liebl D., Seitz A., Hensel M (2008) **Dynamic remodeling of the endosomal system during formation of *Salmonella*-induced filaments by intracellular *Salmonella enterica*** *Traffic* **9**:2100–2116 <https://doi.org/10.1111/j.1600-0854.2008.00821.x>

Röder J., Felgner P., Hensel M (2021) **Comprehensive Single Cell Analyses of the Nutritional Environment of Intracellular *Salmonella enterica*** *Front Cell Infect Microbiol* **11**:624650 <https://doi.org/10.3389/fcimb.2021.624650>

Rodríguez-Morales O., Fernández-Mora M., Hernández-Lucas I., Vázquez A., Puente J. L., Calva E (2006) ***Salmonella enterica* serovar Typhimurium ompS1 and ompS2 mutants are attenuated for virulence in mice** *Infect Immun* **74**:1398–1402 <https://doi.org/10.1128/iai.74.2.1398-1402.2006>

Rosenberg G., Riquelme S., Prince A., Avraham R (2022) **Immunometabolic crosstalk during bacterial infection** *Nat Microbiol* **7**:497–507 <https://doi.org/10.1038/s41564-022-01080-5>

- Rosenberg G., Yehezkel D., Hoffman D., Mattioli C. C., Fremder M., Ben-Arosh H., ..., Avraham R. (2021) **Host succinate is an activation signal for Salmonella virulence during intracellular infection** *Science* **371**:400–405 <https://doi.org/10.1126/science.aba8026>
- Schmitzberger F., Smith A. G., Abell C., Blundell T. L (2003) **Comparative analysis of the Escherichia coli ketopantoate hydroxymethyltransferase crystal structure confirms that it is a member of the (betaalpha)<sub>8</sub> phosphoenolpyruvate/pyruvate superfamily** *J Bacteriol* **185**:4163–4171 <https://doi.org/10.1128/jb.185.14.4163-4171.2003>
- Schneider F., Krämer R., Burkovski A (2004) **Identification and characterization of the main beta-alanine uptake system in Escherichia coli** *Appl Microbiol Biotechnol* **65**:576–582 <https://doi.org/10.1007/s00253-004-1636-0>
- Sibon O. C., Strauss E (2016) **Coenzyme A: to make it or uptake it?** *Nat Rev Mol Cell Biol* **17**:605–606 <https://doi.org/10.1038/nrm.2016.110>
- Silva A. F., Matos M. P., Ralph M. T., Silva D. L., de Alencar N. M., Ramos M. V., Lima-Filho J. V. (2016) **Comparison of immunomodulatory properties of mannose-binding lectins from Canavalia brasiliensis and Cratylia argentea in a mice model of Salmonella infection** *Int Immunopharmacol* **31**:233–238 <https://doi.org/10.1016/j.intimp.2015.12.036>
- Song P., Zhang X., Wang S., Xu W., Wei F (2023) **Advances in the synthesis of β-alanine** *Front Bioeng Biotechnol* **11**:1283129 <https://doi.org/10.3389/fbioe.2023.1283129>
- Steeb B., Claudi B., Burton N. A., Tienz P., Schmidt A., Farhan H., Bumann D (2013) **Parallel exploitation of diverse host nutrients enhances Salmonella virulence** *PLoS Pathog* **9**:e1003301 <https://doi.org/10.1371/journal.ppat.1003301>
- Tan X., Bramlett M. R., Lindahl P. A (2004) **Effect of Zn on acetyl coenzyme a synthase: evidence for a conformational change in the alpha subunit during catalysis** *J Am Chem Soc* **126**:5954–5955 <https://doi.org/10.1021/ja039600z>
- Theodoulou F. L., Sibon O. C., Jackowski S., Gout I (2014) **Coenzyme A and its derivatives: renaissance of a textbook classic** *Biochem Soc Trans* **42**:1025–1032 <https://doi.org/10.1042/bst20140176>
- Thompson A., Fulde M., Tedin K (2018) **The metabolic pathways utilized by Salmonella Typhimurium during infection of host cells** *Environ Microbiol Rep* **10**:140–154 <https://doi.org/10.1111/1758-2229.12628>
- Tuli A., Sharma M (2019) **How to do business with lysosomes: Salmonella leads the way** *Curr Opin Microbiol* **47**:1–7 <https://doi.org/10.1016/j.mib.2018.10.003>
- Wang J., Ma S., Li W., Wang X., Huang D., Jiang L., Feng L (2021) **Salmonella enterica serovar Typhi induces host metabolic reprogramming to increase glucose availability for intracellular replication** *Int J Mol Sci* **22**:10003 <https://doi.org/10.3390/ijms221810003>
- Wang L., Mao Y., Wang Z., Ma H., Chen T (2021) **Advances in biotechnological production of β-alanine** *World J Microbiol Biotechnol* **37**:79 <https://doi.org/10.1007/s11274-021-03042-1>
- Wang X., Yang B., Ma S., Yan X., Ma S., Sun H., ..., Jiang L. (2023) **Lactate promotes Salmonella intracellular replication and systemic infection via driving macrophage M2 polarization** *Microbiol Spectr* **11**:e0225323 <https://doi.org/10.1128/spectrum.02253-23>

- Webb M. E., Smith A. G., Abell C (2004) **Biosynthesis of pantothenate** *Nat Prod Rep* **21**:695–721 <https://doi.org/10.1039/b316419p>
- West T. P., Traut T. W., Shanley M. S., O'Donovan G. A (1985) **A *Salmonella typhimurium* strain defective in uracil catabolism and beta-alanine synthesis** *J Gen Microbiol* **131**:1083–1090 <https://doi.org/10.1099/00221287-131-5-1083>
- White W. H., Gunyuzlu P. L., Toyn J. H (2001) **Saccharomyces cerevisiae is capable of de novo pantothenic acid biosynthesis involving a novel pathway of beta-alanine production from spermine** *J Biol Chem* **276**:10794–10800 <https://doi.org/10.1074/jbc.M009804200>
- Wu A., Tymoszyk P., Haschka D., Heeke S., Dichtl S., Petzer V., ..., Weiss G. (2017) **Salmonella utilizes zinc to subvert antimicrobial host defense of macrophages via modulation of NF- $\kappa$ B signaling** *Infect Immun* **85**:e00418–17 <https://doi.org/10.1128/iai.00418-17>
- Yang W., Cong Y (2021) **Gut microbiota-derived metabolites in the regulation of host immune responses and immune-related inflammatory diseases** *Cell Mol Immunol* **18**:866–877 <https://doi.org/10.1038/s41423-021-00661-4>
- Yuan S. F., Nair P. H., Borbon D., Coleman S. M., Fan P. H., Lin W. L., Alper H. S (2022) **Metabolic engineering of *E. coli* for  $\beta$ -alanine production using a multi-biosensor enabled approach** *Metab Eng* **74**:24–35 <https://doi.org/10.1016/j.ymben.2022.08.012>

## Author information

### Shuai Ma<sup>†</sup>

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

<sup>†</sup>Contributed equally to this work.

### Bin Yang<sup>†</sup>

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

<sup>†</sup>Contributed equally to this work.

### Yuyang Sun

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

### Xinyue Wang

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

### Houliang Guo

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

**Ruiying Liu**

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

**Ting Ye**

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

**Chenbo Kang**

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

**Jingnan Chen**

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

**Lingyan Jiang**

National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, TEDA Institute of Biological Sciences and Biotechnology, Nankai University, Tianjin, China

ORCID iD: [0000-0001-8580-606X](https://orcid.org/0000-0001-8580-606X)

**For correspondence:** [jianglingyan@nankai.edu.cn](mailto:jianglingyan@nankai.edu.cn)

**Editors**

Reviewing Editor

**Amit Singh**

Indian Institute of Science, Bangalore, India

Senior Editor

**Dominique Soldati-Favre**

University of Geneva, Geneva, Switzerland

**Reviewer #1 (Public review):**

Summary:

Ma & Yang et al. report a new investigation aimed at elucidating one of the key nutrients S. Typhimurium (STM) utilizes with the nutrient-poor intracellular niche within macrophage, focusing on the amino acid beta-alanine. From these data, the authors report that beta-alanine plays important roles in mediating STM infection and virulence. The authors employ a multidisciplinary approach that includes some mouse studies, and ultimately propose a mechanism by which panD, involved in B-Ala synthesis, mediates regulation of zinc homeostasis in Salmonella.

Strengths and weaknesses:

The results and model are adequately supported by the authors' data. Further work will need to be performed to learn whether the Zn<sup>2+</sup> functions as proposed in their mechanism. By performing a small set of confirmatory experiments in S. Typhi, the authors provide some evidence of relevance to human infections.

Impact:

This work adds to the body of literature on the metabolic flexibility of *Salmonella* during infection that enable pathogenesis.

<https://doi.org/10.7554/eLife.103714.3.sa2>

**Reviewer #3 (Public review):**

*Salmonella* is interesting due to its life within a compact compartment, which we call SCV or *Salmonella* containing vacuole in the field of *Salmonella*. SCV is a tight-fitting vacuole where the acquisition of nutrients is a key factor by *Salmonella*. The authors among many nutrients, focused on beta-alanine. It is also known that *Salmonella* requires beta-alanine from many other studies. The authors have done in vitro RAW macrophage infection assays and In vivo mouse infection assays to see the life of *Salmonella* in the presence of beta-alanine. They concluded by comprehending that beta-alanine modulates the expression of many genes including zinc transporters which is required for pathogenesis.

[Editors' note: The authors have appropriately addressed the previous reviewers' concerns.]

<https://doi.org/10.7554/eLife.103714.3.sa1>

**Author response:**

The following is the authors' response to the previous reviews

**Reviewer #1 (Public review):**

*Summary:*

*Ma & Yang et al. report a new investigation aimed at elucidating one of the key nutrients S. Typhimurium (STM) utilizes with the nutrient-poor intracellular niche within macrophage, focusing on the amino acid beta-alanine. From these data, the authors report that beta-alanine plays important roles in mediating STM infection and virulence. The authors employ a multidisciplinary approach that includes some mouse studies, and ultimately propose a mechanism by which panD, involved in B-Ala synthesis, mediates regulation of zinc homeostasis in Salmonella.*

*Strengths and weaknesses:*

*The results and model are adequately supported by the authors' data. Further work will need to be performed to learn whether the Zn<sup>2+</sup> functions as proposed in their mechanism. By performing a small set of confirmatory experiments in S. Typhi, the authors provide some evidence of relevance to human infections.*

*Impact:*

*This work adds to the body of literature on the metabolic flexibility of Salmonella during infection that enable pathogenesis.*

**Reviewer #1 (Recommendations for the authors):**

*No further suggestions. The authors have adequately addressed my prior concerns through new data and revisions to the text.*

Thank you for considering this work. We appreciate your efforts in aiding us to improve our manuscript.



**Reviewer #3 (Public review):**

**Summary:**

*Salmonella* is interesting due to its life within a compact compartment, which we call SCV or *Salmonella* containing vacuole in the field of *Salmonella*. SCV is a tight-fitting vacuole where the acquisition of nutrients is a key factor by *Salmonella*. The authors among many nutrients, focussed on beta-alanine. It is also known that *Salmonella* requires beta-alanine from many other studies. The authors have done in vitro RAW macrophage infection assays and In vivo mouse infection assays to see the life of *Salmonella* in the presence of beta-alanine. They concluded by comprehending that beta-alanine modulates the expression of many genes including zinc transporters which is required for pathogenesis.

**Strengths:**


Made a couple of knockouts in *Salmonella* and did transcriptomic to understand the global gene expression pattern

**Weaknesses:**

(1) Transport of Beta-alanine to SCV is not yet elucidated. Is it possible to determine whether the Zn transporter is involved in B-alanine transport?

Thank you for the comment. Following your suggestion, we investigated the growth of *Salmonella* WT and the  $\Delta znuA$  mutant cultured in N-minimal and M9 minimal medium, with  $\beta$ -alanine as the sole carbon source. We observed no significant difference in growth kinetics between the  $\Delta znuA$  mutant and WT strain under either culture condition (please refer to Author response image 1). The results indicate that ZnuA is not involved in  $\beta$ -alanine transport in *Salmonella*.

**Author response image 1.**



(2) Beta-alanine can also be shuttled to form carnosine along with histidine. If beta-alanine is channelled to make more carnosine, then the virulence phenotypes may be very different.

Our study reveals that  $\beta$ -alanine availability, whether obtained from the host or synthesized *de novo* via the *panD*-dependent pathway, is important for *Salmonella* pathogenesis. We have shown that  $\beta$ -alanine influences *Salmonella* intracellular replication and *in vivo* virulence partly by enhancing the expression of the zinc transporter genes.

Although  $\beta$ -alanine can also be shuttled to form carnosine along with histidine in animals, the *Salmonella* genome lacks canonical carnosine synthase (CARNS) orthologs that catalyze the condensation of  $\beta$ -alanine and histidine into carnosine. Therefore, we believe that the carnosine biosynthetic pathway does not influence the virulence phenotypes of *Salmonella*.

*(3) Some amino acid transporters can be knocked out to see if beta-alanine uptake is perturbed. Like ArgT transport Arginine, and its mutation perturbs the uptake of beta-alanine. What is the beta-alanine concentration in the SCV? SCVS can be purified at different time points, and the Beta-alanine concentration can be measured*

Thank you for the comment. As suggested, we have investigated the role of other amino acid transporters in the uptake of  $\beta$ -alanine. In *E. coli*, GabP transports  $\gamma$ -aminobutyric acid (GABA), a structural analogue of  $\beta$ -alanine, and may also transport  $\beta$ -alanine (J Bacteriol. 2021, 203(4):e00642-20). Nevertheless, *Salmonella*  $\Delta$ *gabP* mutant displayed no growth defect in minimal medium with  $\beta$ -alanine as the sole carbon source (Figure 1\_figure Supplement 7, Figure 1\_figure Supplement 8), indicating that GabP is not involved in  $\beta$ -alanine uptake in *Salmonella*. Strikingly, the  $\Delta$ *argT* mutant—defective in arginine uptake—showed markedly decreased growth in the minimal medium with  $\beta$ -alanine as the sole carbon source (Figure 1F), suggesting that ArgT also transports  $\beta$ -alanine in *Salmonella*. We have added the results in the revised manuscript (lines 167-179).

It has been reported that ArgT is essential for *Salmonella* replication within macrophages and full virulence in vivo (PloS one. 2010, 5(12):e15466). Given that ArgT is involved in both arginine and  $\beta$ -alanine uptake (as verified in this study), whether the attenuated virulence of the  $\Delta$ *argT* mutant is due to a deficiency in  $\beta$ -alanine or arginine requires further investigation. We have also included a discussion on this issue (lines 409-415).

In this work, to avoid delays and alterations in metabolite concentrations during the isolation of bacterial contents from macrophages, we directly assessed the combined metabolite concentrations within infected cells and *Salmonella*. It has been previously verified that these metabolites are primarily of host origin (Nat Commun. 2021, 12(1):879.). We noted a decrease in  $\beta$ -alanine levels in macrophages infected with *Salmonella*. The process of separating SCV is intricate and encompasses dissociation and sonication (Nat Commun. 2018, 9(1):2091). These steps may potentially result in alterations of metabolite concentrations during the separation procedure. Therefore, we did not measure the  $\beta$ -alanine concentration in the SCV.

**Reviewer #3 (Recommendations for the authors):**

*The Authors have done meticulous experiments to address the questions asked by the reviewers. My one question of beta-alanine transport inside the SCV remains undone, though the authors have tried.*

*Was Zinc transporter mutant checked? It is possible that the Zn transporter can take up Beta-alanine.*

Thank you for the comment. Following your suggestion, we investigated the growth of *Salmonella* WT and the  $\Delta$ *znuA* mutant cultured in N-minimal and M9 minimal medium, with  $\beta$ -alanine as the sole carbon source. We observed no significant difference in growth kinetics between the  $\Delta$ *znuA* mutant and WT strain under either culture condition (please refer to Author response image 1). The results indicate that ZnuA is not involved in  $\beta$ -alanine transport in *Salmonella*.

Additionally, we have investigated the role of other amino acid transporters in the uptake of  $\beta$ -alanine and have ultimately identified that ArgT, the arginine transporter, is involved in the uptake of  $\beta$ -alanine in *Salmonella* (please refer to our previous response).

<https://doi.org/10.7554/eLife.103714.3.sa0>