

The evolution of gene expression in seasonal environments

Reviewed Preprint

v2 • October 21, 2025

Revised by authors

Reviewed Preprint

v1 • August 13, 2025

Shuichi N Kudo, Yuka Ikezaki, Junko Kusumi, Hideki Hirakawa, Sachiko Isobe, Akiko Satake 

Graduate School of Systems Life Sciences, Kyushu University, Fukuoka, Japan • Department of Biology, Faculty of Science, Kyushu University, Fukuoka, Japan • Department of Environmental Changes, Faculty of Social and Cultural Studies, Kyushu University, Fukuoka, Japan • Kazusa DNA Research Institute, Kisarazu, Japan • Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka, Japan • Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

The authors collected time-course RNA-seq data from four tree species in natural environments and analyzed seasonal patterns of gene expression. This **fundamental** study substantially advances our understanding of how seasonal environments shape gene expression. The evolutionary effects of seasonal environments on gene expression are rarely studied at this scale and the dataset is extensive. The evidence supporting the conclusions is **compelling**, with caveats and limitations clearly described. The work will be of broad interest to colleagues studying evolution and gene expression.

<https://doi.org/10.7554/eLife.107309.2.sa2>

Abstract

The biological activities of organisms are closely linked to seasonality. Phenology, the temporal orchestration of biological activities, is governed by gene expression, yet the evolutionary dynamics underlying seasonal gene expression remain unclear. To investigate these dynamics, we compared genome-wide expression dynamics (molecular phenology) in four dominant evergreen Fagaceae species in Asia (*Quercus glauca*, *Q. acuta*, *Lithocarpus edulis*, and *L. glaber*), using leaf and bud tissues over two seasonal cycles. We assembled high-quality reference genomes, identifying 11749 single-copy orthologous genes. Seasonal transcriptomic profiling of these orthologous genes revealed highly conserved gene expression across species in winter when temperatures fall below ~10°C. Rhythmic gene expression with significant periodic oscillations was more prevalent in buds (51.9%) than in leaves (40.6%), with most rhythmic genes (78.4–92.0%) exhibiting annual periodicity, while a smaller fraction (1.2–11.9%) followed half-annual cycles. The seasonal peaks of rhythmic genes were highly synchronized across species in winter but diverged during the growing season, reflecting species-specific timing of leaf flushing and flowering. These findings suggest that the four species share a common molecular calendar in winter, which constrains the evolution of gene expression under seasonal environments.

Introduction

Fundamental biological processes, such as growth, mortality, and reproduction, rarely remain constant over time; instead, they exhibit pronounced variation in response to seasonal environmental changes (Fretwell, 1972 [↗](#)). This temporal organization of a series of biological activities, known as phenology, plays a critical role in the evolution of life histories in response to seasonal fluctuating environments (Forrest and Miller-Rushing, 2010 [↗](#); Morellato et al., 2013 [↗](#); Piao et al., 2019 [↗](#); Schwartz, 2013 [↗](#)). Phenology is regulated by the coordinated expression of genes that enable organisms to exhibit essential functions at the appropriate times (Kudoh, 2016 [↗](#); Satake et al., 2024a [↗](#), 2022 [↗](#)). However, gene expression dynamics underpinning seasonal responses in organisms remain not fully understood.

To explore the evolution of gene expression, transcriptome analyses are widely employed to compare transcript profiles across organs, sexes, species, and developmental stages within a given environment. Such studies have been conducted in mammals (Brawand et al., 2011 [↗](#); Cardoso-Moreira et al., 2019 [↗](#); Naqvi et al., 2019 [↗](#)), *Drosophila* (Lemos et al., 2005 [↗](#); Nuzhdin et al., 2004 [↗](#)), and Tanzanian cichlids (El Taher et al., 2021 [↗](#)), providing key insights into the evolutionary dynamics of gene expression and its central role in shaping phenotypic diversity. These studies have revealed that gene expression profiles tend to be more conserved across species within the same tissue, whereas the evolutionary rate of gene expression varies among organs. In mammals, for instance, nervous tissue exhibits a slower rate of gene expression change compared to the testis (Brawand et al., 2011 [↗](#); Cardoso-Moreira et al., 2019 [↗](#); Naqvi et al., 2019 [↗](#)), suggesting the presence of tissue-specific functional constraints that influence the evolution of gene expression.

In addition to tissue-dependent constraints, seasonal environmental changes may play a crucial role in shaping the evolution of gene expression, particularly by influencing the temporal coordination of gene expression under natural conditions. Gene expression responses to certain seasons may be more resistant to change, whereas in other seasons, divergence in gene expression timing across species may occur, contributing to temporal niche differentiation. Despite its potential importance, the influence of seasonal environmental fluctuations on the gene expression evolution remains poorly understood, largely because most transcriptomic analyses have been conducted under tightly controlled laboratory conditions.

To investigate the evolution of gene expression in seasonal environments, we performed a molecular phenology analysis and phenological observation of leaf flushing and flowering in Fagaceae tree species, which are widely distributed across the Northern Hemisphere, with its center of diversity in subtropical Southeast Asia (Hipp et al., 2020 [↗](#); Kremer et al., 2012 [↗](#); Manos et al., 2001 [↗](#); Zhou et al., 2022 [↗](#)). Molecular phenology refers to transcriptome analysis conducted under natural fluctuating conditions (Komoto et al., 2024 [↗](#); Kudoh, 2016 [↗](#); Nagano et al., 2019 [↗](#); Satake et al., 2023 [↗](#)) and provides a valuable framework for studying gene–environment interactions and their role in orchestrating the temporal biological functions essential for adaptation to environmental fluctuations (Satake et al., 2022 [↗](#)). In this study, we selected four evergreen Fagaceae species—two *Quercus* species (*Q. glauca* and *Q. acuta*) and two *Lithocarpus* species (*L. edulis* and *L. glaber*)—that co-occur in the same locality in Japan. *Q. glauca*, *L. edulis* and *L. glaber* inhabit nearly identical environment, while *Q. acuta* is restricted to the high-altitude regions with slightly colder environment compared to the other three species. Leaf flushing and flowering phenology vary across species, with *Q. glauca* and *Q. acuta* flowering in early spring, almost simultaneously with leaf flushing, *L. edulis* flowering in late spring, and *L. glaber* flowering in autumn, approximately one to two months after leaf flushing. This setting enabled us to compare seasonal gene expression profiles across species with diverse phenology, both within and between genera, under nearly identical environmental fluctuations. By assessing

and comparing seasonal transcriptomic dynamics across Fagaceae species, our study provides new insights into the evolution of gene expression and adaptation to seasonal natural environments.

Results

Genome assemblies of *Q. glauca* and *L. edulis*

To quantify transcript abundance of non-model Fagaceae species, we first constructed reference genomes for *Q. glauca*, as a representative of genus *Quercus*, and *L. edulis*, as a representative of genus *Lithocarpus* (Fig. 1A). To obtain high-quality chromosome-level reference genomes, we conducted a hybrid genome assembly for the HiFi reads and the high-throughput chromosome conformation capture (Hi-C) reads (Table S1). The resultant scaffolds were 867.7 and 835.7 Mb and consisted of 797 and 750 scaffolds including 12 chromosome-level scaffolds with contig N50 of 71.7 Mb and 68.8 Mb for *Q. glauca* and *L. edulis*, respectively (Table S1). Approximately 94.8% and 97.2% of the final assemblies were assigned to 12 pseudochromosomes of *Q. glauca* and *L. edulis*, respectively (Fig. 1B). The number of chromosome-scale scaffolds were identical to the known numbers in other *Quercus* (Zoldos et al., 1999) and *Lithocarpus* species (Liu et al., 2023). The genomes were estimated to contain 39758 and 34059 protein-coding genes (33947 and 30246 genes were annotated within 12 chromosomes), covering 98.3% and 95.0% of complete Benchmarking Universal Single Copy Orthologs (BUSCO) genes (eudicots_odb10) for *Q. glauca* and *L. edulis*, respectively. The genome sizes of *Q. glauca* and *L. edulis* were estimated to be 891.2 Mb and 868.7 Mb, respectively, using KmerGenie (Chikhi and Medvedev, 2014) and GenomeScope2 (Ranallo-Benavidez et al., 2020).

Overall, the abundance and distribution of protein-coding genes, GC content, and repeat sequences (Fig. 1C, D; Table S1) and syntenic structure (Fig. 1B) were similar between *Q. glauca* and *L. edulis*. This pattern of synteny conservation appears to be broadly shared across Fagaceae, except for the genus *Fagus*, which shows a distinct genome structure (Ikezaki et al., 2025). The homology between different chromosomes within the genome suggests that small-scale segmental duplications likely occurred, as previously proposed (Chen et al., 2014). Using OrthoFinder2 (Emms and Kelly, 2019), we clustered genes from the two newly assembled genomes into orthologous groups. We identified 13427 single-copy orthologs, enabling pairwise comparisons between species (Fig. 1E).

Seasonal gene expression dynamics

To characterize the evolution of seasonal gene expressions, we generated a total of 483 transcriptome profiles every four weeks over two years using leaf and bud tissues from *Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber* during 2021–2023. Reads from the two *Quercus* species were mapped to the newly constructed *Q. glauca* reference genome, achieving a mean mapping rate of $92.6 \pm 2.2\%$ and $88.6 \pm 2.3\%$ for *Q. glauca* and *Q. acuta*, respectively. Similarly, reads from the two *Lithocarpus* species were mapped to the newly constructed *L. edulis* reference genome, with a mean mapping rate of $89.3 \pm 2.7\%$ and $84.3 \pm 5.4\%$ for *L. edulis* and *L. glaber*, respectively. The number of reads uniquely mapped on the reference genome was 19.9 million reads per sample, on average. We used single-copy orthologous genes for further analyses because these genes exhibited significantly higher average expression levels than multi-copy genes (Steel-Dwass test, $p < 0.05$; Fig. 1F, G). Among the single copy orthologous genes, we excluded genes with low expression levels (mean of Reads Per Kilobase (RPK) across samples for each species < 1) and normalized the transcript count data using the GeTMM method (Smid et al., 2018). This process resulted in a high-quality expression dataset for 11749 genes (Fig. S1A). For further analyses, we used the mean expression levels across three biological replicates (three individuals per species),




Figure 1.

Synteny and distribution of genomic features of *Q. glauca* and *L. edulis* genomes.

(A) A phylogenetic tree of five Fagaceae species from three genera, *Quercus*, *Lithocarpus*, and *Fagus*. Bootstrap values for all branches are 100. (B) Synteny relationship between the *Q. glauca* and *L. edulis* genomes. The collinear blocks within the genomes were displayed by lines. Black lines show reverse alignment and gray lines represent regular. Numbers correspond to the 12 chromosomes. (C–D) Circos plots of the *Q. glauca* (C) and *L. edulis* genomes (D). Lanes depict circular representation of chromosome length (Mbp) (a), gene density (b), GC density (c), and repeat (d). Lines in the inner circle represent links between syntenic paralogs. (E) Orthogroup classification based on the orthogroup size difference between the *Q. glauca* and *L. edulis* genomes. The bar plot illustrates the number of genes in gene families categorized into 1:1 orthologs, N:N (many-to-many), 1:N (one-to-many), N:1 (many-to-one), and species-specific unique genes. (F–G) Mean gene expression levels in each copy number variation category for *Q. glauca* (F) and *L. edulis* (G). Different letters denote statistically significant differences ($p < 0.05$, Steel-Dwass test).

except for *L. edulis*, where only one individual was available. We confirmed that expression patterns of individual genes were highly consistent across all the three biological replicates (Fig. S2) in *Q. glauca* (Pearson's correlation $r = 0.947$), *Q. acuta* ($r = 0.948$) and *L. glaber* ($r = 0.948$).

To gain an initial overview of gene expression dynamics in a seasonal environment, we assessed the relative magnitude of seasonal variation in expression levels for each gene by calculating the standard deviation divided by the mean for each gene (σ/μ) and compared it to the overall variation across all genes in the genome (σ/μ) (Fig. S1B). A prior study using 3025 yeast RNA-seq experiments reported that the dynamic range of individual genes across experiments was considerably smaller than the overall genomic variation under laboratory conditions (Zrimec et al., 2021). Consistently, our results demonstrate that despite substantial seasonal fluctuations in environmental conditions (Fig. S3), individual gene expression remains relatively conserved. On average, the magnitude of genome-wide variation was 15.8 times greater than the seasonal variation observed for individual genes (Fig. S1B).

To assess the similarity of gene expression profiles across tissues, species, and seasons, we performed a hierarchical clustering of seasonal transcriptome profiles from leaf and bud tissues across four species and identified five distinct clusters (cluster L1, L2, B1–B3) according to the Elbow method (Fig. 2A; Fig. S4A; Dataset S1). Transcriptional profiles first grouped into leaves (cluster L1 and L2) and buds (cluster B1–B3), irrespective of seasons and species (Fig. 2A), except that newly flushed leaf samples were grouped closely with buds (Fig. 2A; Fig. S5). Within the leaf cluster, transcriptome profiles were first clustered by genus and then by species in *Lithocarpus* (cluster L1; Fig. 2A). In contrast, samples from *Quercus* species formed a mixture of two species (cluster L2; Fig. 2A). Notably, the winter samples of *Q. acuta*, collected during periods of low mean temperatures, were distinctly separated from those collected in other seasons (cluster L2; Fig. 2A). Within the bud cluster, transcriptome profiles were predominantly grouped by season—winter (cluster B3; November to February, with the exception of a March sample from *L. edulis*; Dataset S1) versus the other growing seasons (cluster B1 and B2; March to November; Dataset S1)—and subsequently organized by genus and species. This suggests that the winter season exerts a stronger influence on gene expression in buds than phylogenetic relationships. We confirmed the robustness of our findings to differences in sampling periods between *Quercus* and *Lithocarpus* species by reanalyzing expression data limited to common sampling periods (Fig. S6). Even when using only the data from this shared period, the clustering patterns were consistently reproduced. To assess whether the higher-elevation site of *Q. acuta* introduced confounding environmental effects, we reanalyzed the data after excluding this species. Hierarchical clustering still revealed that winter bud samples formed a distinct cluster regardless of species identity (Fig. S7), consistent with our original findings.

To assess the biological relevance of the clusters identified, we tested whether the temperatures and photoperiod at the time of sample collection differed significantly among the clusters. Within the bud cluster, transcriptional profiles in the winter cluster (cluster B3) were associated with significantly lower temperatures—approximately below 10 °C—compared to those in other seasons (Fig. 2B; Fig. S8A). In contrast, no such temperature-related distinction was evident within the leaf cluster (Fig. 2B). Additionally, photoperiods were markedly shorter in the winter cluster in buds (Fig. S8B). These findings suggest that the buds of the four species exhibit a conserved transcriptional response to environmental cues, namely cold temperatures and short photoperiods, characteristic of the winter season. To further investigate the relationships among all samples, we conducted a principal component analysis (PCA). The first principal component (PC1) clearly separated samples by tissue type, while the second (PC2) and the third principal component (PC3) gradually differentiated winter samples from those collected during other seasons (Fig. S9).




Figure 2.

Seasonal gene expression dynamics.

(A) Hierarchical clustering of 213 time points encompassing two tissues (leaf and bud) and four species (*Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*), collected over two-year. Newly flushed leaf samples are indicated as asterisks. (B) Distribution of daily mean temperature on each sampling date across clusters, from L1 (top) to B3 (bottom). The Mann-Whitney U test was used to compare daily mean temperature between cluster L1 and L2 ($p = 0.136$). The Steel-Dwass test was applied for multiple comparisons across clusters B1–B3 ($p < 0.05$). Different letters denote statistically significant differences ($p < 0.05$). (C–D) Hierarchical clustering of 11749 genes based on the similarity of gene expression dynamics. Daily mean temperature, flowering and leaf flushing phenology (proportion) are shown in the upper three panels of the heatmaps.

To investigate seasonal expression dynamics of each gene, we conducted hierarchical clustering on the transcriptional dynamics of 11749 genes and identified five distinct gene sets characterizing different transcriptional modes (gene set I–V; **Fig. 2C, D** [Fig. 2C, D](#); Fig. S4B; Dataset S2). Gene set I exhibited predominant expression in buds, with seasonal peaks during summer, whereas gene set II was primarily expressed in leaves, highlighting clear tissue-specificity (**Fig. 2C, D** [Fig. 2C, D](#)). Gene sets III and V demonstrated genus-specific expression patterns, with higher expression levels observed in *Lithocarpus* and *Quercus*, respectively (**Fig. 2C, D** [Fig. 2C, D](#)). In contrast, gene set IV was expressed during winter across both leaf and bud tissues (**Fig. 2C, D** [Fig. 2C, D](#)).

Detection of rhythmic genes

To identify genes revealing seasonal expression dynamics and determine the period and seasonal peaks in gene expression, we applied the RAIN algorithm (Thaben and Westermark, 2014 [Thaben and Westermark, 2014](#)). This approach enabled us to detect genes with rhythmic expression (hereafter rhythmic genes) and analyze their periods and phases (peak months). Our analysis revealed that 44.6–58.8% of genes in bud tissues and 33.9–45.9% in leaf tissues exhibited significant rhythmic expression across species (Fig. S10A; Dataset S3). Notably, bud tissues demonstrated a higher proportion of rhythmic genes, with 51.9% across four species on average, compared to 40.6% in leaf tissues (Fig. S10A).

Among the rhythmic genes, the majority (78.4–92.0%) exhibited annual periodicity, with a period of 12 ± 1 month observed across species and tissues (**Fig. 3A, B** [Fig. 3A, B](#), Dataset S3). Additionally, hundreds of genes (1.2–11.9%) displayed half-annual periodicity (6 ± 1 month) with these genes being most frequently observed in the bud samples of *L. edulis* (**Fig. 3A, B** [Fig. 3A, B](#)). The fraction of genes exhibiting other periodicities was relatively small (5.7–14.1%; **Fig. 3A, B** [Fig. 3A, B](#)). Based on the observed period distributions, we classified genes with periods within the range of 8–16 months as exhibiting annual periodicity (96.0%) and those with periods shorter than 8 months as having half-annual periodicity (4.0%), accounting for detection noise (Fig. S10A). Rhythmic genes with longer periods (>17 months) were also identified, although their periodicity may be less reliable due to the limited number of time points. Many genes with annual periodicity were shared across all species, including 1280 genes in leaves and 1894 genes in buds (Fig. S10B). In contrast, most genes exhibiting half-annual periodicity were species-specific, with no genes shared among all the species (Fig. S10C).

Peak month distribution of rhythmic genes and intra-genus and inter-genera comparison

To determine the seasonal timing of gene expression peaks, we focused on genes with annual or half-annual periodicity and visualized the density and direction of these peaks throughout the year using a circular histogram (**Fig. 3C, D** [Fig. 3C, D](#); Dataset S3). On average, 34.9% and 41.3% of annual rhythmic genes peaked in winter (December–February) in leaf and bud samples, respectively. Additionally, another peak corresponding to or occurring slightly after leaf flushing and flowering events was observed, with its timing varying by species: April in *Q. glauca* (15.6%), July in *Q. acuta* (15.3%), June in *L. edulis* (16.0%), September in *L. glaber* (18.1%), for leaf tissues (**Fig. 3C** [Fig. 3C](#)). Similar peaks corresponding to leaf flushing or flowering were also detected in bud samples, occurring approximately one to two months earlier than in leaf samples (**Fig. 3D** [Fig. 3D](#)). For genes with half-annual periodicity, the majority ($63.2 \pm 4.8\%$) exhibited expression peaks both in spring (March–May) and autumn (September–November) in both tissues (Fig. S10D; Dataset S3), with the exception of *L. glaber*, an autumn flowering species. Interestingly, *WRKY22*, a transcription factor regulating ethylene biosynthesis and senescence (Zhu et al., 2024 [Zhu et al., 2024](#)), exhibited half-annual periodicity in *L. glaber* (Fig. S10E). In addition, *SNF1-RELATED PROTEIN KINASE REGULATORY SUBUNIT GAMMA 1 (KINy1)*, a subunit of SNF1-related kinase (*SnRK*) involved in stress response such as starvation (Emanuelle et al., 2016 [Emanuelle et al., 2016](#)), showed two distinct peaks in *Q. glauca* and *L. edulis*, occurring in both summer (July) and winter (January) (Fig. S10E). This result suggests that both summer and winter pose stressful conditions for these species.




Figure 3.

Distribution of period and peak month in rhythmic gene expression.

(A–B) Distribution of period length (month) among the genes with significant rhythmicity calculated by RAIN for leaf (A) and bud (B) samples of each species. (C–D) Distribution of seasonal peaks among the genes with annual periodicity. The angle of each bar represents the peak month, while the bar height indicates the number of genes. Blue arrows indicate the month with the highest number of genes peaking during winter (December–February), while red arrows indicate the month with the highest number of genes peaking during the growing season (March–November). Purple and green squares around each rose plot denote the observed phenology of flowering and leaf flushing (proportion), respectively.

To compare the peak month of gene expression with annual periodicity across species, we quantified the number of genes within each peak combination for both intra-genus (**Fig. 4A**) and inter-genera species pairs in buds (**Fig. 4B**) and leaves (Fig. S11A, B). During winter, a large number of genes exhibited highly synchronized expression peaks across species, with high pairwise Pearson's correlation coefficients observed (**Fig. 4A, B**). In contrast, the peak timing of gene expression from spring to autumn varied across species, aligning with differences in leaf flushing and flowering times (**Fig. 4A, B**). Some genes exhibited pronounced divergence, showing negative Pearson's correlation coefficients (< -0.3) between species. Notably, the number of such genes was larger in inter-genera comparisons (11–77 genes; **Fig. 4B**) than in intra-genus comparisons (4–12 genes; **Fig. 4A**). A similar pattern was observed in leaf samples (Fig. S11A, B). When comparing seasonal expression peaks between tissues, the majority of genes were synchronously activated in winter in both tissues (Fig. S11C). During spring to summer, slight variations in seasonal expression peaks were observed among tissues (Fig. S11C).

We quantified the degree of divergence in seasonal gene expression patterns across species by calculating the peak month differences between two species, where one species exhibits an expression peak in the given month. We defined the molecular phenology divergence index (D) for each month by quantifying the proportion of genes with peak month differences greater than 2 months between two species, relative to the total number of genes with expression peaks in that month. From spring to autumn, the mean of D over all species pairs in buds ranged from 0.47 to 0.66, but it declined sharply during winter, reaching a minimum of 0.12 in January in buds (Nemenyi test, $p < 0.05$; **Fig. 4C**). A similar trend was observed in leaf samples, with another trough in summer (Fig. S11D). This seasonal trend remained consistent even after excluding *Q. acuta*, which was sampled from a slightly cooler environment than the other three species, from the analysis (Fig. S12), indicating that the observed patterns are not driven by environmental differences associated with elevation. The values of D were not consistently smaller in inter-genera pairs compared to intra-genus pairs. These findings suggest that there is a seasonal constraint in gene expression in which the degree of molecular phenology divergence is not constant but varies significantly across seasons, with the least divergence occurring in winter.

Phylogenetic constraints in the evolution of seasonal gene expression

We next investigated whether phylogenetic relationship influences the evolution of seasonal gene expression by calculating pairwise Pearson's correlations of seasonal expression for each gene across all species pairs (Dataset S4). The median of Pearson's correlation coefficients across all genes were significantly higher between species within the same genus compared to those between species from different genera in both leaves (Steel-Dwass test, $p < 0.05$; Fig. S13A; Fig. S14) and buds (Steel-Dwass test, $p < 0.05$; **Fig. 5A**; Fig. S14). The intra-genus comparison in genus *Lithocarpus* exhibited the highest correlation in their seasonal gene expression, despite their notable differences in flowering phenology, with *L. edulis* flowering in spring and *L. glaber* in autumn. In contrast, the intra-genus comparison in genus *Quercus* resulted in a relatively low correlation in their gene expression regardless of relatively similar phenology in leaf flushing and flowering. This difference may be attributed to environmental factors, as *Q. acuta* is the only species found in the high-altitude study site with colder climates compared to the other three species.

Rhythmic genes exhibited more conserved expression patterns than arrhythmic genes in both tissues (**Fig. 5A**; Fig. S13A). In particular, genes with seasonal expression peaks in winter showed high correlation across species in both tissues (**Fig. 5B**; Fig. S13B). The proportion of rhythmic genes with annual periodicity declined as the mean correlation coefficient across species decreased, with a more rapid decline observed for genes peaking in winter compared to those with peak expression during the growing seasons (**Fig. 5B**; Fig. S13B). Additionally, the mean absolute difference in seasonal gene expression dynamics was significantly smaller between




Figure 4.

Comparison of peak months in seasonal gene expression across species in buds.

(A) Intra-genus and (B) inter-genera comparisons of peak months in seasonal gene expression. The size and color of the circles represent the number of genes and the mean of Pearson's correlation, respectively. Genes with negative correlations lower than -0.3 are highlighted in blue, while the month of leaf flushing is highlighted in green. (C) Plot of molecular phenology divergence index D across months. Triangles and circles indicate the values of D calculated from intra-genus and inter-genera comparison, respectively. The line and shaded envelope indicate the mean and standard deviation (s.d.). Different letters denote statistically significant differences (Nemenyi test, $p < 0.05$).




Figure 5.

Phylogenetic constraints in the evolution of seasonal gene expression and the relationship between seasonal gene expression divergence and sequence evolution.

(A) Distribution of pairwise Pearson's correlation coefficients for gene expression across species (Le: *L. edulis*, Lg: *L. glaber*, Qg: *Q. glauca*, Qa: *Q. acuta*; $n = 11749$). Colors indicate rhythmicity categories. Black bars show median correlation values, and different letters indicate significant differences (Nemenyi test, $p < 0.05$). (B) Heatmap of pairwise correlation coefficients for each gene, with mean gene counts in four seasonal expression categories: (1) annual rhythmic genes peaking in the growing season (March–November), (2) annual rhythmic genes peaking in winter (December–February), (3) half-annual rhythmic genes, and (4) arrhythmic genes. Genes are ordered by mean correlation across species pairs. (C–D) Conserved (C) and diverged (D) gene expression patterns, based on the top and bottom 5% of pairwise correlations. Solid lines and error bars indicate mean $\pm$ SD. Numbers in parentheses show peak month differences between *Q. glauca* and *L. edulis*. Peak month difference for *BFT* was not calculated due to arrhythmic expression in *Q. glauca*. (E) Relationship between evolutionary rate (dN/dS) and peak month difference ($\Delta\phi$) for genes with annual periodicity in *Q. glauca* and *L. edulis* ($n = 951$). Black lines and transparent bands represent regression lines and 95% confidence intervals. The regression coefficient (β) and corresponding p -value are shown in each panel.

species within the same genus than between species from different genera in both tissues (Fig. S15A, B). The relationship between the mean absolute difference and the proportion of rhythmic genes was less clear compared to that of the mean correlation coefficient (Fig. S15C, D). These findings suggest that, in addition to seasonal constraints, the overall dynamics of seasonal gene expression are also shaped by phylogenetic constraints.

Functional annotation of genes with conserved or divergent expression across species

To functionally annotate genes with conserved or divergent expression across species, we extracted genes with the top and bottom 5% based on the distribution of mean correlation (Fig. 5B; Fig. S13B) and conducted Gene Ontology (GO) enrichment analysis. GO analysis revealed that the top 5% of genes in leaves were associated with photosynthesis, while those in buds were linked to cell fate determination (Fig. S16A; Dataset S5). The gene with the highest correlation in leaves was *STRESS RESPONSE SUPPRESSOR1* (*STRS1*; mean of pairwise correlation = 0.91; Fig. 5C), a DEAD-box RNA helicase previously reported to enhance tolerance to abiotic stresses such as cold (Kant et al., 2007). In buds, the most highly correlated gene was the histone deacetylase *HDA19* (mean of pairwise correlation = 0.93; Fig. 5C), which plays a key role in environmental stress responses through histone modification and chromatin remodeling (Chen and Wu, 2010). These findings suggest that genes associated with photosynthesis during the growing season and stress responses during winter exhibit highly conserved seasonal expression patterns across species.

The bottom 5% of genes were predominantly associated with DNA replication and cell division in leaves (Fig. S16B), aligning with the divergent leaf phenology observed (Fig. 3B). In contrast, GO terms related to immune response were weakly enriched in buds (Fig. S16B). Notably, we identified MADS-box genes linked to floral transition among the bottom 5% of genes: *BROTHER OF FT AND TFL1* (*BFT*) in leaves (mean of pairwise correlation = -0.094) and *FRUITFULL* (*FUL*) in buds (mean of pairwise correlation = -0.069). *BFT*, a floral repressor (Yoo et al., 2010), was expressed in autumn in the spring-flowering species *L. edulis* and in spring in the autumn-flowering species *L. glaber* leaves (Fig. 5D). Meanwhile, *FRUITFULL* (*FUL*), a positive regulator of floral transition (Smaczniak et al., 2012), was expressed in spring in *L. edulis* and in autumn in *L. glaber* (Fig. 5D), mirroring the timing of flowering. In *Q. glauca* and *Q. acuta*, *FUL* was expressed in winter (Fig. 5D), nearly out of phase with their expressions in *Lithocarpus* species. These results suggest that genes involved in reproductive processes exhibit species-specific divergence in expression patterns in response to seasonal environmental changes.

Relating seasonal gene expression divergence to sequence divergence

If divergence of seasonal timing of gene expression is related to selective constraint, we would expect a correlation between sequence divergence patterns and the divergence in seasonal expression peaks. To explore the relationship between the divergence of seasonal gene expression and protein sequence evolution, we calculated synonymous (dS) and non-synonymous substitution rates (dN) and their ratio (dN/dS) for each gene using coding sequences from newly assembled genomes of *Q. glauca* and *L. edulis*. We first confirmed that the highly expressed genes evolve more slowly by identifying a significant negative correlation between evolutionary rate (dN/dS) and mean of the gene expression ($r = -0.21$, $p = 1.8 \times 10^{-118}$ for dN/dS; Fig. S17A; Dataset S6). This finding aligns with previous studies in yeasts, mammals, and plants (Drummond et al., 2005; Gaut et al., 2011; Liao and Zhang, 2006; Yang and Gaut, 2011). By targeting the genes that exhibited annual periodicity across both species and tissues, we examined the relationship between these evolutionary measures and peak month differences in seasonal gene expression between species ($\Delta\phi$) in leaves and bud tissues using linear regression. Assessment of the significance of the regression coefficient using a *t*-test revealed that genes with smaller peak

month differences between species exhibited lower values of dN/dS in buds ($p = 2.6 \times 10^{-2}$) and marginally lower values of dN/dS in leaves ($p = 5.7 \times 10^{-2}$). The lower values were also found in dN in both leaves ($p = 8.1 \times 10^{-3}$) and buds ($p = 7.8 \times 10^{-3}$), whereas no significant trend was observed for dS in both tissues (**Fig. 5E**; Fig. S17B, C; Dataset S6). We further investigated whether the association between the divergence in seasonal expression peaks and sequence divergence patterns varies between genes expressed in winter and in the growing season. We classified genes that exhibited annual periodicity across both species and tissues into four groups based on their expression peak timing (winter or growing seasons) and tissue specificity (leaves or buds) and compared the extent of protein sequence divergence across these groups. We found no significant differences in dN/dS (Kruskal Wallis test, $p = 2.8 \times 10^{-1}$; Fig. S17D; Dataset S6) between genes expressed in winter in both tissues and those in growing seasons. The same result was observed even when focusing only on genes with conserved peaks (peak month difference < 2), with no significant differences in dN/dS between these gene sets (Kruskal Wallis test, $p = 3.5 \times 10^{-1}$; Fig. S17E; Dataset S6). These results suggest that coding-sequence variation is unlikely to be the primary driver of seasonal gene expression patterns.

Discussion

A central goal in biology is to unravel the molecular basis of phenotypic evolution, with gene expression serving as a fundamental driver of this process. In this study, we investigated the impact of seasonal environmental fluctuations on the evolution of gene expression and its relationship with phenological traits in four dominant *Fagaceae* tree species in forest ecosystems. We found that gene expression profiles were more resistant to change during winter compared to the growing season, indicating the presence of a seasonal constraint on the evolution of gene expression. This constraint becomes particularly pronounced in buds during winter when temperatures fall below approximately 10°C (**Fig. 2B**). This temperature threshold aligns with previous findings, where winter-specific transcript profiles were reported at temperatures below 8.7–11.3°C in cherry trees (Miyawaki - Kuwakado et al., 2024), 11.3°C in ever-green *Fagaceae* (Satake et al., 2023), and 10.5°C in perennial *Arabidopsis* (Aikawa et al., 2010). These results suggest the presence of a conserved mechanism for gene expression regulation under low-temperature environments across different species. The greater conservation of gene expression patterns in buds compared to leaves across species may be attributed to their functional role protecting essential meristematic tissues, including the shoot apex and leaf and flower primordia. Additionally, buds likely play a crucial role in maintaining common physiological functions necessary for the resumption of growth in spring. This functional consistency may impose evolutionary constraints on gene expression, contributing to its conservation across species.

It is intriguing to compare the seasonal constraints identified in this study with developmental constraints in gene expression evolution, where evolutionary conservation and variation across developmental stages during the life cycle have been observed, such as in the early conservation model and the developmental hourglass model (Fig. S18A) (Irie and Kuratani, 2011; Lotharukpong et al., 2024; Quint et al., 2012). Previous studies have demonstrated that certain developmental stages exhibit remarkably similar gene expression patterns (Drost et al., 2016; Kalinka et al., 2010; Levin et al., 2016). Our findings highlight that such highly similar gene expression patterns are observed during the winter season (Fig. S18B), suggesting that a common physiological state is essential for enduring harsh winter conditions. This seasonal constraint on gene expression evolution may have the potential to limit temporal niche partitioning at the phenotypic level and slow species divergence rates in seasonal environments (Cleland and Wolkovich, 2024; Kubo and Iwasa, 1996). Although completely disentangling the effects of developmental stages and seasonal changes on gene expression is challenging when using molecular phenology data collected from natural environments, the observation that species with different phenology—i.e., different developmental timing in seasonal environments—exhibit convergent gene expression patterns during winter underscores the profound influence of winter

environmental conditions on the evolution of gene expression. Future research is needed to elucidate the mechanisms driving the establishment of such conserved gene expression patterns, including the dynamic reorganization of chromatin architecture in response to cold (Fischl et al., 2020 [DOI](#); Nishio et al., 2020 [DOI](#); Zhang et al., 2023 [DOI](#)).

We found that the evolution of coding-sequences was only weakly correlated with the divergence of seasonal peaks in gene expression across species. To understand the evolutionary mechanism underlying the seasonal constraint on gene expression that results in similar expression patterns during winter, an important future direction is to explore the evolution of non-coding sequences that modulate gene expression levels in response to seasonal environmental changes. Regulatory variation has been recognized as a major driver of evolutionary innovation (Prud'homme et al., 2007 [DOI](#); True and Haag, 2001 [DOI](#)). Recent deep learning studies utilizing genome and transcriptome data from yeast and humans have highlighted the significance of co-evolution between coding and non-coding sequences in shaping gene expression levels (Zrimec et al., 2021 [DOI](#), 2020 [DOI](#)). Understanding how seasonal environments drive the co-evolution of coding and regulatory sequences, as well as the divergence of molecular phenology and its phenological outputs, will require expanding analyses to a broader range of species. Such efforts will facilitate the development of predictive models for evolutionary processes influenced by seasonal dynamics. Comparison of molecular phenology across wide climatic zones, including both tropical and temperate zones, will also be effective to identify key genes for the phenological response to ongoing climate change.

Materials and Methods

Sample collection for genome sequencing

To construct genomes of *Q. glauca* and *L. edulis*, we collected approximately 5g of fresh leaves from one individual in spring, respectively. We shipped these samples to Dovetail Genomics, LLC (Scotts Valley, CA, USA) where Dovetail Staff performed DNA extraction, library preparation, sequencing, and assembly steps.

Genome sequencing and assembly

DNA extraction was performed using the QIAamp DNA Mini Kit (Qiagen, Germantown, MD, USA) using the manufacturer's recommended protocol. Mean fragment length of the extracted DNA was 60 kbp for *Q. glauca* and 65 kbp for *L. edulis*. DNA samples were quantified using Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, CA, USA). The library preparation, sequencing and scaffolding were carried out by Dovetail Genomics (California, USA) according to their standard genome assembly pipeline (<https://dovetailgenomics.com/> [DOI](#)). The PacBio SMRTbell library (~20kb) for PacBio Sequel was constructed using SMRTbell Express Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA) using the manufacturer's recommended protocol. The library was bound to polymerase using the Sequel II Binding Kit 2.0 (PacBio) and loaded onto PacBio Sequel II. Sequencing was performed on PacBio Sequel II 8M SMRT cells. PacBio CCS reads were used as an input to Hifiasm1 v0.15.4-r347 with default parameters. For *Q. glauca*, we also generated sequence data using PacBio Revio. BLASTn results of the contigs obtained by Hifiasm assembly against the NCBI's NT database were used as input for blobtools2 v1.1.1 and contigs identified as possible contamination were removed from the assembly. Finally, purge_dups3 v1.2.5 was used to remove haplotigs and contig overlaps.

Assembly scaffolding with HiRise

A proximity ligation library was generated by the Dovetail's Omni-C technique, followed by sequencing on an Illumina HiSeqX platform. Chromatin was fixed in place in the nucleus with formaldehyde before extraction (<https://dovetailgenomics.com/wp-content/uploads/2021/09/Omni-C->

Tech-Note.pdf [↗](#)). Fixed chromatin was digested with DNase I, and the fragmented chromatin ends were repaired and biotinylated to adapters followed by proximity ligation. Crosslinks were then reversed, the DNA purified, and the biotin subsequently removed. The DNA library was prepared and sequenced to produce 2×150 bp paired-end reads (around 30× coverage). The Omni-C technology uses a sequence-independent endonuclease which provides even, unbiased genome coverage. The input *de novo* assembly and Omni-C library reads were used as input data for HiRise, a software pipeline designed specifically for using proximity ligation data to scaffold genome assemblies (Putnam et al., 2016 [↗](#)). The Omni-C library sequences were aligned to the contigs assembled by Hifiasm using bwa (Li and Durbin, 2009 [↗](#)). The separations of Omni-C reads from paired-end mapped within the contigs were analyzed by HiRise to produce a likelihood model for genomic distance between paired-end reads, and the model was used to identify and break putative misjoins, to score prospective joins, and make joins above a threshold.

Sample collection for RNA sequencing

To monitor the genome-wide gene expression profiles, we collected leaf, bud, and flower samples from four Fagaceae species, *Q. glauca*, *Q. acuta*, *L. edulis* and *L. glaber*, every four weeks for two years (October 2019 to October 2021 for the *Lithocarpus* species and April 2021 to April 2023 for the *Quercus* species) across Fukuoka and Saga prefectures in Kyushu, Japan. We targeted three individuals, and each sample was collected from three branches per individual, except for *L. edulis*. Given the consistent expression patterns observed among individuals in a previous study (Satake et al., 2023 [↗](#)), we focused on only one individual for *L. edulis*. Study sites for *Q. glauca* is the biodiversity reserve on the Ito campus of Kyushu University (33°35' N, 130°12' E, 20 to 57 m a.s.l.). *L. edulis* and *L. glaber* were studied at the Imajuku Field Activity Center (33°33' N, 130°16' E, 84 to 111 m a.s.l.). The Imajuku and Ito Campus study sites, located only 7.3 km apart, experience nearly identical temperature conditions. *Q. acuta* was monitored in Sefuri mountains (33°26' N, 130°22' E, 970 to 974 m a.s.l.), a higher-altitude, cooler site. Environmental data from Ito Campus and the Imajuku Field Activity Center were obtained from the Japan Meteorological Agency, while data from Sefuri Mountain were collected using a logger installed at the sampling site. Additionally, to assess the potential impact of different sampling periods between the *Lithocarpus* and *Quercus* species on the gene expression analysis, we conducted an additional year of sampling for *L. edulis* and *L. glaber*, collecting leaf and bud samples from November 2021 to November 2022, and used the samples collected within the same period for all the four species (May 2021 to November 2022).

Samples for RNA-seq were taken from the sun-exposed crown (approximately 2–5 m from the ground) using long pruning shears from 11:30 to 12:30 h. For each sample, 0.2–0.4 g of tissue was preserved in a 2 ml microtube containing 1.5 ml of RNA-stabilizing reagent (RNAlater; Thermo Fisher Scientific, Waltham, MA, USA) immediately after harvesting. The samples were transferred to the laboratory within 2 hr after sampling, stored at 4°C overnight and then stored at –80°C until RNA extraction. During transport to the laboratory, the samples were kept in a cooler box with ice to maintain a low temperature.

RNA extraction and RNA-seq analysis

RNA was extracted independently from bud and leaf samples of each tree. Total RNA of the leaf sample was extracted in accordance with the method described in a previous study (Miyazaki et al., 2014 [↗](#)). Total RNA of the bud sample was extracted using the PureLink™ Plant RNA Reagent (Thermo Fisher Scientific, Waltham, MA, USA) and subsequently processed with DNase using the TURBO DNA-free kit™ (Thermo Fisher Scientific) according to their protocols and purified using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to its RNA cleanup protocol. RNA integrity was examined using the Agilent RNA 6000 Nano kit on a 2100 Bioanalyzer (Agilent Technologies), and the RNA yield was determined using a NanoDrop ND-2000 spectrophotometer (Thermo Fisher Scientific). Five to six micrograms of RNA extracted from each sample were sent to

Hangzhou Veritas Genetics Medical Institute Co., Ltd., where a cDNA library was prepared with a NEBNext Ultra II RNA Library Prep Kit for Illumina, and 150 base paired-end transcriptome sequencing of each sample was conducted using an Illumina NovaSeq6000 sequencer (Illumina).

The quantification of gene expression followed a three-step process. First, quality filtering was performed using the fastp v0.23.2 with default options: phred quality ≥ 15 , the maximum number of N base in one read = 5 (Chen et al., 2018 [↗](#)). Second, RNA-seq reads were mapped to the reference genome using STAR v2.7.10b (Dobin et al., 2013 [↗](#)). Third, RNA-seq reads were quantified using RSEM v1.3.1 (Li and Dewey, 2011 [↗](#)). We excluded the genes with low expression level using the threshold, the value of mean Reads Per Kilobase (RPK) < 1 for each species. Then, we further normalized the expression level using the value of Gene length corrected Trimmed Mean of M-values (GeTMM) (Smid et al., 2018 [↗](#)) using an R package “edgeR” v3.42.4. For the downstream analysis we calculated the mean expression levels of GeTMM of each gene across multiple individuals and transformed them to $\log_2(\text{GeTMM}+1)$. To confirm that the effect of the difference in sampling period is negligible, we added the RNA-Seq data of *L. edulis* and *L. glauca* collected from December 2021 to December 2022. We extracted the same genes in the existing data that curated based on the mean RPK values across two- year sampling periods. Then we further normalized the expression level using GeTMM.

Genome annotation

The RNA-seq reads of *Q. glauca* (8 samples obtained by a previous study (20) collected from May to December 2017; GEO’s accession number: GSE211384) and *L. edulis* (20 samples covering leaf, bud and flower tissues collected in January, April, July, and November 2021) were respectively mapped against the scaffolds of *Q. glauca* and *L. edulis* obtained by HiRise with HISAT2 v2.1.0 (Kim et al., 2019 [↗](#)). The bam files generated by the mappings and amino acid sequences of the 9 species obtained from NCBI (*Castanea mollissima*: GCA_000763605.2 (NCBI’s accession number), *Q. lobata*: GCF_001633185.1, *Q. robur*: GCF_932294415.1, *Q. suber*: GCF_002906115.1, *Carpinus fangiana*: GCA_006937295.1, *Carya illinoensis*: GCF_018687715.1, *Juglans macrocarpa* $\times$ *J. regia*: GCF_004785595.1, *Juglans regia*: GCF_001411555.2, *Morella rubra*: GCA_003952965.2) were used for ab initio gene prediction by Braker2 v2.1.6 (Brůna et al., 2021 [↗](#)), and the genes with highest-score were selected to exclude splicing variants. In parallel, the gene models were respectively constructed in *Q. glauca* and *L. edulis* by GeMoMa v1.7.1 (Keilwagen et al., 2019 [↗](#)) with RNA-seq reads and amino acid sequences used for the gene prediction in Braker2, and the genes with highest-score were selected to exclude splicing variants. The duplicated genes between Braker2 and GeMoMa were excluded by GFFcompare v0.11.2 (Pertea and Pertea, 2020 [↗](#)). The remaining genes were searched against UniProtKB (Bateman et al., 2025) by DIAMOND v2.0.5 (Buchfink et al., 2014 [↗](#)) in ‘more-sensitive’ option with E-value $\leq 10^{-20}$ and identity $\geq 25\%$. Next, gene expression analysis was performed by Salmon v1.4.0 (Patro et al., 2017 [↗](#)), and the genes with TPM values > 0.1 were considered as expressed. The orthologous genes were searched against EggNOG (Hernández-Plaza et al., 2023 [↗](#)) using eggNOG-mapper v2.1.8 (Cantalapiedra et al., 2021) with E-value $\leq 10^{-10}$. The genes were searched against protein sequences of *Castanea mollissima* (Accession: GCA_000763605.2), *Quercus lobata* (GCF_001633185.2), and *Arabidopsis thaliana* (Araport11 (Krishnakumar et al., 2015 [↗](#))) by BLASTp v2.13.0 (Camacho et al., 2009 [↗](#)) with E-value $\leq 10^{-20}$. According to these similarity searches, the genes whose product names contain keywords related to transposable elements (TEs) were categorized into TE. The genes having similarity hits and being expressed by Salmon were classified into high-confidence (HC) genes, while remaining genes were categorized into low-confidence (LC) genes. The genes categorized into HC were used for further analyses.

Genome synteny analysis

To investigate the syntenic relationship between *Q. glauca* and *L. edulis*, we first performed an all-vs-all BLASTp search of the protein sequences, using a cut-off E-value of 10^{-5} . Collinearity blocks within and between the genomes were then identified using MScanX (Wang et al., 2012 [↗](#)) based

on the results of the BLASTp searches. Finally, the collinearity blocks were visualized using SynVisio (<https://synvisio.github.io>).

Illumina short-read sequencing and library preparation

For genome size estimation and phylogenetic reconstruction, we also generated short-read sequencing of *Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*. DNA extraction was performed using a modified version of the method described previously (Satake et al., 2024b). The DNA samples were sent to MacroGen Inc. (Republic of Korea) for sequencing on the Illumina HiSeqX platform (Illumina, San Diego, CA, USA). DNA was sheared to around 500 bp fragments in size using dsDNA fragmentase (New England BioLabs, Ipswich, MA, USA). Library preparation was performed using the NEBNext Ultra II DNA Library Prep Kit (New England BioLabs) according to the manufacturer's protocol, and the libraries were individually indexed with the NEBNext Multiplex Oligos for Illumina (New England BioLabs) by PCR. The quality and quantity of each amplified library were analyzed using the Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA), the High Sensitivity DNA kit (Agilent Technologies), and the NEBNext Library Quant Kit for Illumina (New England BioLabs).

Genome size estimation

The genome size of *Q. glauca* was estimated by GenomeScope2 v2.0 (Ranallo-Benavidez et al., 2020) using *k*-mer frequency plot obtained by kmc v3.2.4 (Kokot et al., 2017) with *k*-mer size 61 against the HiFi reads. We also used pair-end short reads for genome size estimation of *Q. glauca* and confirmed that resulted estimates were not different. For *L. edulis*, paired-end short reads from four individuals were merged and quality-trimmed using fastp v0.23.2 (Chen et al., 2018). The genome size was then estimated using KmerGenie (Chikhi and Medvedev, 2014) with a *k*-mer size of 55 applied to the trimmed reads.

Phylogenetic reconstruction

We constructed a phylogenetic tree using an alignment-free method (Fan et al., 2015) based on short-read data from *Q. acuta* and *L. glaber* and genome assembly data from *Q. glauca* and *L. edulis*. Additionally, genomic data for *Fagus sylvatica* was obtained from beechgenome.net (Mishra et al., 2022) and included as an outgroup. Due to the large volume of short-read data for *Q. acuta* and *L. glaber* (18 Gb each), we randomly subsampled half of the reads. Bootstrap values were estimated through 100 resampling of the *k*-mer table.

Identification of orthologs

The prediction of orthogroups was based on a BLASTp all-against-all comparison of the protein sequences (E-value < 10^{-5}) of 5 Fagaceae species: *L. edulis*, *Q. glauca*, *Q. robur*, *Fagus sylvatica* and *Juglans regia*. Clustering was conducted with OrthoFinder2, using version 2.2.6 for BLAST data generation and version 2.5.5 for orthogroup determination (Emms and Kelly, 2019). The genomic data of *Q. robur* and *J. regia* were downloaded from NCBI (dhQueRobu3.1 and Walnut 2.0), while *F. sylvatica* data were sourced from beechgenome.net (Mishra et al., 2022). Since the *F. sylvatica* dataset contained numerous predicted transposable elements (TEs), we re-annotated genes using three databases—EggNOG5 (EggNOG-mapper), Pfam35.0 (HMMER v3.3.2), and UniRef90_2023_0628 (DIAMOND v2.1.8.162)—and excluded genes identified as TEs in at least one database. In total, we identified 26692 orthogroups and focused on single-copy orthogroups after filtering out genes mapped outside the 12 chromosomes. This resulted in a final dataset of 11749 genes for further analysis (Data S2).

Hierarchical clustering and principal component analysis (PCA)

To assess the similarity of the genome-wide transcriptional profiles across orthologous genes and 483 samples collected in different seasons, we performed hierarchical clustering using the monthly time series data of 11749 orthologous genes from October 2019 to April 2023. For each orthologous gene, there were 26 or 27 time points, with three individuals each for *Q. glauca*, *Q. acuta*, *L. glaber* and one individual for *L. edulis*. We calculated the mean expression levels of each orthologous gene across three individuals in each species and subsequently normalized the values by adjusting the mean to zero and the standard deviation to one. We performed hierarchical clustering using the Ward method and the Euclidian distance using the pheatmap function in R (v 1.0.12). To determine the number of the clusters, we calculated Within-Cluster-Sum of Squared Error (WSS) and determined the number of clusters as an elbow point of it (Fig. S4). To assess the seasonal expression dynamics of 11749 orthologous genes, we also performed PCA of the gene expression profiles from all samples using the prcomp function in R (v 4.3.1).

Identification of genes with rhythmic expression

To identify genes with rhythmic expression and determine the period and peak month of gene expression, we applied the RAIN algorithm (Thaben and Westermark, 2014 [\[1\]](#)) using the rain function in R (v 1.34.0). We set the following settings: `deltat = 1`, `period = 12`, `period.delta = 11`, `peak.border = (0,1)`, `nr.series = 1`, `method = "independent"`, `na.rm = T`, `adjp.method = "ABH"`. Based on the results of the RAIN analysis, we identified genes with significant periodicity using a Benjamini-Hochberg (BH) adjusted *q*-value threshold of < 0.01 . Since the period length and peak month in the RAIN output are expressed in terms of the number of time points, we converted them into months using the sampling span (four weeks) and sampling dates. Subsequently, we categorized the rhythmic genes into three categories based on the period length *T* (in months): “half-annual” ($0 \leq T < 8$), “annual” ($8 \leq T \leq 16$), and “long” ($T > 16$).

Calculation of Pearson’s correlation coefficient across species

To quantify the similarity of seasonal gene expression pattern across species, we calculated pairwise Pearson’s correlations of seasonal expression for each gene across all the species pairs. For species within the same genus, the monitoring period was identical, allowing direct comparison. On the other hand, the monitoring periods differed between *Quercus* and *Lithocarpus* species: *Q. glauca* and *Q. acuta* were monitored from May 2021 to April 2023, while *L. edulis* and *L. glaber* were monitored from October 2019 to October 2021 (Fig. S3). To ensure comparable time series data for inter-genera comparisons, we extracted the time series from *Q. glauca* and *Q. acuta* spanning May 19, 2021, to November 2, 2022, and from *L. edulis* and *L. glaber* spanning May 13, 2020, to October 20, 2021, for correlation analysis. The mean absolute differences for each gene across all the species pairs were also calculated in a similar manner.

Molecular phenology divergence index (*D*)

To quantify the seasonal difference of peak month divergence across species, we defined the molecular phenology divergence index (*D*) for each month *t* and species pair A-B as the proportion of genes that peak in month *t* in the reference species A but exhibit a peak month difference ($\Delta\phi$) greater than two months in species B using following formula: $D := \# \text{ Genes with peak month } t \text{ in species A and } t \pm \Delta\phi (\geq 2) \text{ in species B} / \# \text{ Genes with peak month } t \text{ in species A}$

Gene Ontology (GO) enrichment analysis

First, we assigned GO terms for each gene in *Lithocarpus* and *Quercus* genomes based on the annotation data of *Arabidopsis thaliana* orthologs downloaded from PLAZA Database (<http://bioinformatics.psb.ugent.be/plaza/> [\[2\]](#)). Then, GO enrichment analysis was conducted using the GOstats (Falcon and Gentleman, 2007 [\[3\]](#)) function in R (v 2.66.0) with the following settings:

ontology = BP, pvalueCutoff = 0.05, conditional = TRUE, and testDirection = over. The universe gene set was defined as genes associated with GO terms common to both of the genomes. Genes without annotated GO terms were removed from the universe set, and those with differing GO term annotations between the two genomes were also excluded. The BH adjusted q -values were calculated from the p -values generated by GStats to account for multiple testing.

Phenology measurement: leaf unfolding and flowering

To compare seasonal changes in genome-wide gene expression profiles with phenology, we conducted observations of leaf unfolding and flowering phenology on the three terminal branches of each of the three individuals used for gene expression monitoring. Leaf unfolding was defined as the stage where both the entire leaf blade and the leaf stalk were visible (Fig. S5). Flowering was defined as the stage where either male or female flowers were visible (Fig. S5). We subsequently plotted the proportion of branches exhibiting leaf unfolding and flowering. The data are provided in Dataset S2.

Estimation of protein-coding sequence divergence

Pairwise distances for non-synonymous (dN) and synonymous (dS) substitutions, as well as the dN/dS ratio, were estimated for each orthologous gene pair between *L. edulis* and *Q. glauca* using codeml (PAML 4.0)(Yang, 2007 [link](#)). The analysis was performed with the following settings: seqtype = 1, CodonFreq = 2, Runmode = - 2, and the transition/transversion ratio estimated from the data. To ensure reliability, genes exhibiting signs of saturated divergence or underestimated dS values were excluded, as codeml produces robust results only within a moderate range of sequence divergence. Specifically, we removed 43 orthologs with dN/dS $\geq$ 99 and 483 genes with dS > 10, dS < 0.001, or dN < 0.0003. The thresholds of dN and dS were determined by examining the relationship between dN and dS and excluding outliers that deviated from the main distribution. After filtering, the final dataset comprised 11223 orthologous genes.

Linear regression analyses were performed to examine the relationship between dN, dS, dN/dS and peak month differences ($\Delta\phi$) in gene expression between *Q. glauca* and *L. edulis*, as well as mean expression levels. These analyses were conducted using the lm function in R software (v 4.3.1). Specifically, we modeled the evolutionary rates (dN, dS, and dN/dS) as the response variable and the peak month difference ($\Delta\phi$) or the mean levels in gene expression as the predictor variable. To account for the skewed distributions of evolutionary rates, logarithm transformations with base 2 were applied.

Data deposition and code availability

All sequencing data generated in this study have been deposited in the DNA Data Bank of Japan (DDBJ) and are available under the BioProject accession number PRJDB20498. RNA-seq reads used for expression analyses are available under the DDBJ Sequence Read Archive (DRA) run accession numbers DRR659185- DRR659374 and DRR664882-DRR665873, while DNA sequences used for genome assembly, genome size estimation and phylogenetic reconstruction are available under DRR659899-DRR659929. The assembled genomes are also available from DDBJ: the *Q. glauca* genome under accession numbers BAAHOF010000001-BAAHOF010000797, and the *L. edulis* genome under BAAHOE010000001-BAAHOE010000750. All codes used in the analysis are available at https://github.com/ku-biomath/Evolution_of_gene_expression_in_seasonal_environment [link](#).

Acknowledgements

We thank Atsuko Miyawaki-Kuwakado for her valuable technical advice on RNA-seq analysis. We also appreciate the technical support provided by Kayoko Ohta. Additionally, we thank Koharu Yamaguchi, Yuta Aoyagi, and Ryosuke Imai for their help with sample collection.

Additional information

Funding

This work was supported by JSPS KAKENHI Grant Numbers JP23H04965 and JP23H04966.

Author Contributions

S.N.K. and A.S. designed research; S.N.K., Y.I., J.K., H.H., S.I., and A.S. performed research; S.N.K., Y.I., J.K., H.H., S.I., and A.S. analyzed data; S.N.K. and A.S. wrote the paper with inputs from all the authors.

Funding

Japan Society for the Promotion of Science (JP23H04965)

Japan Society for the Promotion of Science (JP23H04966)

Additional files

Dataset S1. Summary of clustering 213 gene expression profiles. [↗](#)

Dataset S2. Hierarchical clustering of 11749 gene expression dynamics. [↗](#)

Dataset S3. Identification of genes with rhythmic expression. [↗](#)

Dataset S4. Comparison of correlation and absolute difference among the species pairs. [↗](#)

Dataset S5. List of genes with conserved and diverged gene expression across species. [↗](#)

Dataset S6. Statistical test of evolutionary rates. [↗](#)

Supplementary figures




Figure S1.

Dynamic range of gene expression in seasonal environments.

(A) Dynamic range of seasonal gene expression in leaves and buds for *Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*. Each dot represents an expression level ($\log_2(\text{GeTMM}+1)$) on each sampling date, while black lines indicate the mean expression level of each gene over the entire sampling period. The bars on the left side of the panels distinguish between rhythmic (blue) and arrhythmic (gray) genes. (B) Density plots showing the distribution of the ratio of standard deviation divided by the mean for each gene (σ_i/μ_i) and the overall variation across all genes in the genome (σ_g/μ_g) in leaves and buds for *Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*.




Figure S2.

Comparison of gene expression across individuals.

Scatter plots showing the expression levels of each gene measured from different individuals for *L. glaber* (A), *Q. glauca* (B), and *Q. acuta* (C). The Pearson's correlation coefficient for each pair of individuals is shown in the corresponding panel.

Figure S3.

Seasonal fluctuations in environmental factors at the three study sites.

Daily mean temperature and photoperiods at the three study sites (Ito campus, Kyushu university; Imajuku Field Activity Center; Mt. Sefuri in Kyushu, Japan) during the monitoring periods. The solid lines indicate the daily mean temperature, and the shaded area indicates the photoperiod. The study sites are located within a relatively small region, with distances between them ranging from approximately 10 to 20 km. The elevations vary, with Ito Campus (20–57 m a.s.l.) and Imajuku Field Activity Center (84–111 m a.s.l.) at lower altitudes, while Mt. Sefuri is at a higher elevation (970–974 m a.s.l.). Environmental data from Ito Campus and the Imajuku Field Activity Center were obtained from the Japan Meteorological Agency, while data from Sefuri Mountain were collected using a logger installed at the sampling site since July 15, 2021.





Figure S4.

Determining the optimal number of clusters in hierarchical clustering.

(A) Relationship between the number of clusters k and total Within-Cluster-Sum of Squared Error (WSS) on the clustering of 213 samples. We adopted the optimal $k = 5$ because it was the elbow point of the curve. (B) Relationship between the number of clusters k and total WSS on the clustering of 11749 genes. The elbow point was $k = 4$ or 5 and we selected the number of cluster $k = 5$. If $k = 4$ is selected, gene set IV and V (Fig. 2D, E in main text) are merged.






Figure S5.

Phenological observations of leaf flushing, flowering (male and female flower) and winter buds of the four target species.




Figure S6.

Seasonal gene expression dynamics observed within the same sampling period (from May 2021 to October 2022).

(A) Hierarchical clustering of 160 time points encompassing two tissues (leaf and bud) and four species (*Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*). Newly flushed leaf samples are indicated as asterisks. (C-D) Hierarchical clustering of 11749 genes based on the similarity of gene expression dynamics. Daily mean temperature is shown in the top of the heatmaps.




Figure S7.

Seasonal gene expression dynamics under the identical environmental condition.

(A) Hierarchical clustering of 161 time points from two tissues (leaf and bud) across three species (*Q. glauca*, *L. edulis*, and *L. glaber*) excluding *Q. acuta* sampled at higher elevation. Newly flushed leaf samples are indicated as asterisks. (C-D) Hierarchical clustering of 11749 genes based on the similarity of their gene expression dynamics. Daily mean temperature is shown in the top of the heatmaps.




Figure S8.

Distribution of environmental factors across clusters.

Distribution of mean temperature for the 7 days prior to the sampling (A) and photoperiod for each sampling date (B) across clusters identified through hierarchical clustering in [Fig. 2A](#). The Mann-Whitney U test was used to compare climatic factors between leaf clusters (L1 and L2), while the Steel-Dwass test was applied for multiple comparisons among bud clusters (B1, B2, and B3). Newly flushed leaves were included within cluster B1. Different letters denote statistically significant differences ($p < 0.05$).




Figure S9.

3D plot of PC1–PC3 resulting from the PCA of 11749 orthologous genes in four Fagaceae species.

Leaf samples are represented in a color gradient from green to blue, while bud samples range from orange to yellow. The color gradients indicate the mean temperature on the respective sampling dates for bud and leaf samples. The percentages in parentheses on each axis denote the variance explained by each principal component. The numbers within the plot indicate sampling months.




Figure S10.

Number of rhythmic genes across species and distribution of peak months across half-annual genes.

(A) Number of genes classified into annual (period range: 8-16 months), half-annual (< 8 month), and arrhythmic periodicity. (B-C) Number of common rhythmic genes shared among species within each species combination, categorized by annual (B) and half-annual (C) periodicity. Black dots beneath each bar indicate the species included in each combination. (D) Distribution of seasonal peaks across genes with half-annual periodicity. (E) Seasonal expression patterns of *WRKY22* in leaves and *KIN1* in buds. For each gene, the species exhibiting a half-annual periodicity is highlighted with a thick, dark line, while the expression patterns in the other species are shown with thin, light lines.




Figure S11.

Comparison of peak months in seasonal gene expression across species in leaves.

(A) Intra-genus and (B) inter-genera comparison of peak months in seasonal gene expression. (C) Comparison of peak months between leaves and buds. The size and color of the circles represent the number of genes and the mean of Pearson's correlation, respectively. Genes with negative correlations lower than -0.3 are highlighted in blue, while the month of leaf flushing is highlighted in green. (D) Plot of molecular phenology divergence index D across months. Triangles and circles indicate the D values calculated from intra- and inter- genera comparison, respectively. The line and shaded envelope indicate the mean and standard deviation (s.d.). Different letters denote statistically significant differences (Nemenyi test, $p < 0.05$).




Figure S12.

Seasonal difference of molecular phenology divergence index D calculated for species under the identical environmental condition.

Plot of molecular phenology divergence index D across months in buds (A) and leaves (B) calculated from the gene expression in *Q. glauca*, *L. edulis*, and *L. glaber*. Triangles and circles indicate the values of D calculated from intra-genus and inter-genera comparison, respectively. The line and shaded envelope indicate the mean and standard deviation (s.d.). Different letters denote statistically significant differences (Nemenyi test, $p < 0.05$).




Figure S13

Phylogenetic constraints in the evolution of seasonal gene expression in leaves.

(A) Distribution of pairwise Pearson's correlation coefficients for gene expression across species (*Le*: *L. edulis*, *Lg*: *L. glaber*, *Qg*: *Q. glauca*, *Qa*: *Q. acuta*; $n = 11749$). Colors indicate three categories based on the presence or absence of rhythmicity in gene expression. The median values of Pearson's correlation across all the genes were depicted as black bars. Different letters denote significantly different values (Nemenyi test, $p < 0.05$). (B) Heatmap of pairwise Pearson's correlation coefficients for each gene, along with the mean count of genes in four categories of seasonal gene expression: (1) rhythmic genes with annual periodicity peaking in the growing season (March–November), (2) rhythmic genes with annual periodicity peaking in winter (December–February), (3) rhythmic genes with half-annual periodicity, and (4) arrhythmic genes. The mean count was calculated for each block of 100 genes. Genes were ordered based on the mean Pearson's correlation of their expression across all species pairs.




Figure S14

Distribution of pairwise Pearson's correlation coefficients for gene expression across species under the identical environmental condition.

Density plot of pairwise Pearson's correlation coefficients in buds (A) and leaves (B), calculated from the gene expression in *Q. glauca*, *L. edulis*, and *L. glaber* inhabiting nearly identical environments. Colors indicate rhythmicity categories. Black bars show median correlation values, and different letters indicate significant differences (Nemenyi test, $p < 0.05$).





Figure S15.

Pairwise mean absolute difference of gene expression dynamics.


(A– B) Distribution of mean absolute difference of gene expression dynamics ($\text{Log}_2(\text{GeTMM}+1)$) in leaf (A) and bud samples (B) across species (Le: *L. edulis*, Lg: *L. glaber*, Qg: *Q. glauca*, Qa: *Q. acuta*; $n = 11749$). Colors indicate three categories based on the presence or absence of rhythmicity in gene expression. The median values across all the genes were depicted as black bars. Different letters denote significantly different values (Nemenyi test, $p < 0.05$). (C–D) Heatmap of the mean absolute difference for each gene, along with the mean count of genes in four categories of seasonal gene expression: (1) rhythmic genes with annual periodicity peaking in the growing season (March–November), (2) rhythmic genes with annual periodicity peaking in winter (December–February), (3) rhythmic genes with half-annual periodicity, and (4) arrhythmic genes. The mean count was calculated for each block of 100 genes. Genes were ordered based on the mean absolute difference of their expression across all the species pairs.

A Genes with conserved expression

Leaf (589 genes)




Bud (590 genes)



B Genes with diverged expression

Leaf (587 genes)



Bud (590 genes)




Figure S16.

Gene Ontology (GO) enrichment analysis.

(A) Top 5 GO terms (ranked by the raw p-values) for the genes with conserved expression patterns across species (a mean correlation in gene expression ranking in the top 5%) in leaves and buds. (B) Top 5 GO terms for the genes with diverged expression patterns across species (a mean correlation in gene expression ranking in the bottom 5%) in leaves and buds. Genes used for the analysis are listed in Dataset S5.




Figure S17.

Relationship between sequence divergence and gene expression divergence.

(A) Relationships between evolutionary measures (dN/dS, dN, dS), and mean expression levels of genes except for the outliers ($n = 11223$). (B–C) Relationships between the evolutionary measures (dN, dS) and peak differences ($\Delta\phi$) between *Q. glauca* and *L. edulis* in leaves (B) and buds (C) using the same gene set as in **Fig. 5E** ($n = 951$). The regression slope and corresponding p-value are shown in each panel. Black lines and transparent bands represent regression lines and 95% confidence intervals. (D–E) Comparison of the evolutionary measures (dN/dS, dN, dS) among genes with conserved annual periodicity across four categories ($n = 951$). Genes were classified into four categories: those expressed in winter in both tissues, those expressed in winter only in buds, those expressed in winter only in leaves, and others. (E) shows a subset of (D), including only genes with conserved seasonal peaks (peak month difference < 2).




Figure S18.

Developmental constraints and seasonal constraints in the evolution of gene expression.

(A) The hourglass model of developmental constraints, illustrating similar gene expression during the mid-embryonic stage (40–45). (B) Seasonal constraints identified in this study. Similar gene expression patterns were observed during the winter season in both leaves and buds, as well as during the summer in leaves. The vertical axis represents time (developmental stages or seasons), while the horizontal axis represents the divergence of gene expression patterns.

	<i>Q. glauca</i>	<i>L. edulis</i>
Sequencing		
Raw bases of PacBio (Gb)	32.6	273.6 (subreads)
Raw bases of Hi-C (Gb)	24.8	50.1
Estimated genome size (Mb)	891.2	868.7
Assembly		
Final assembly size (Mb)	867.7	835.7
Number of chromosomes	12	12
Number of contigs	797	750
N50 of contigs (Mb)	71.7	68.8
GC percent	35.68	35.74
Annotation		
Number of predicted genes	39,758	34,059
Mean gene length (bp)	3,889.70	4,099.28
Mean CDS length (bp)	1,221.90	1,164.30
Annotated to EggNOG	37,408 (94%)	31,668 (93%)

Table S1.

The statistics for genome assemblies of two new sequenced species

Note

This reviewed preprint has been updated to correct level headings.

References

1. Aikawa S, Kobayashi MJ, Satake A, Shimizu KK, Kudoh H (2010) **Robust control of the seasonal expression of the Arabidopsis FLC gene in a fluctuating environment** *PNAS* **107**:11632–11637 <https://doi.org/10.1073/pnas.0914293107> | Google Scholar
2. Bateman A (2025) **UniProt: the Universal Protein Knowledgebase in 2025** *Nucleic Acids Research* **53**:D609–D617 <https://doi.org/10.1093/nar/gkae1010> | Google Scholar
3. Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti A, Aximu-Petri A, Kircher M, Albert FW, Zeller U, Khaitovich P, Grützner F, Bergmann S, Nielsen R, Pääbo S, Kaessmann H (2011) **The evolution of gene expression levels in mammalian organs** *Nature* **478**:343–348 <https://doi.org/10.1038/nature10532> | Google Scholar
4. Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M (2021) **BRAKER2: Automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database** *NAR Genomics and Bioinformatics* **3**:1–11 <https://doi.org/10.1093/nargab/lqaa108> | Google Scholar
5. Buchfink B, Xie C, Huson DH (2014) **Fast and sensitive protein alignment using DIAMOND** *Nature Methods* <https://doi.org/10.1038/nmeth.3176> | Google Scholar
6. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) **BLAST+: Architecture and applications** *BMC Bioinformatics* **10** <https://doi.org/10.1186/1471-2105-10-421> | Google Scholar
7. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J (2021) **eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale** *Molecular Biology and Evolution* **38**:5825–5829 <https://doi.org/10.1093/molbev/msab293> | Google Scholar
8. Cardoso-Moreira M, Halbert J, Vallotton D, Velten B, Chen C, Shao Y, Liechti A, Ascensão K, Rummel C, Ovchinnikova S, Mazin P V., Xenarios I, Harshman K, Mort M, Cooper DN, Sandi C, Soares MJ, Ferreira PG, Afonso S, Carneiro M, Turner JMA, VandeBerg JL, Fallahshahroudi A, Jensen P, Behr R, Lisgo S, Lindsay S, Khaitovich P, Huber W, Baker J, Anders S, Zhang YE, Kaessmann H (2019) **Gene expression across mammalian organ development** *Nature* **571**:505–509 <https://doi.org/10.1038/s41586-019-1338-5> | Google Scholar
9. Chen LT, Wu K (2010) **Role of histone deacetylases HDA6 and HDA19 in ABA and abiotic stress response** *Plant Signaling & Behavior* **5**:1318–1320 <https://doi.org/10.4161/psb.5.10.13168> | Google Scholar
10. Chen S, Zhou Y, Chen Y, Gu J (2018) **. fastp: an ultra-fast all-in-one FASTQ preprocessor** *Bioinformatics* **34**:i884–i890 <https://doi.org/10.1093/bioinformatics/bty560> | Google Scholar

11. Chen SC, Cannon CH, Kua CS, Liu JJ, Galbraith DW (2014) **Genome size variation in the Fagaceae and its implications for trees** *Tree Genetics and Genomes* **10**:977–988 <https://doi.org/10.1007/s11295-014-0736-y> | Google Scholar
12. Chikhi R, Medvedev P (2014) **Informed and automated k-mer size selection for genome assembly** *Bioinformatics* **30**:31–37 <https://doi.org/10.1093/bioinformatics/btt310> | Google Scholar
13. Cleland EE, Wolkovich EM (2024) **Effects of phenology on plant community assembly and structure. Annual Review of Ecology Evolution, and Systematics** **55**:471–492 <https://doi.org/10.1146/annurev-ecolsys-102722-011653> | Google Scholar
14. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) **STAR: Ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21 <https://doi.org/10.1093/bioinformatics/bts635> | Google Scholar
15. Drost HG, Bellstädt J, Ó'Maoiléidigh DS, Silva AT, Gabel A, Weinholdt C, Ryan PT, Dekkers BJW, Bentsink L, Hilhorst HWM, Ligterink W, Wellmer F, Grosse I, Quint M (2016) **Post-embryonic hourglass patterns mark ontogenetic transitions in plant development** *Mol Biol Evol* **33**:1158–1163 <https://doi.org/10.1093/molbev/msw039> | Google Scholar
16. Drummond DA, Bloom JD, Adami C, Wilke CO, Arnold FH (2005) **Why highly expressed proteins evolve slowly** *PNAS* **102**:14338–14343 <https://doi.org/10.1073/pnas.0504070102> | Google Scholar
17. Taher A El, Böhne A, Boileau N, Ronco F, Indermaur A, Widmer L, Salzburger W (2021) **Gene expression dynamics during rapid organismal diversification in African cichlid fishes** *Nature Ecology and Evolution* **5**:243–250 <https://doi.org/10.1038/s41559-020-01354-3> | Google Scholar
18. Emanuelle S, Doblin MS, Stapleton DI, Bacic A, Gooley PR (2016) **Molecular insights into the enigmatic metabolic regulator *SnRK1*** **1**:341–353 <https://doi.org/10.1016/j.tplants.2015.11.001> | Google Scholar
19. Emms DM, Kelly S (2019) **OrthoFinder: Phylogenetic orthology inference for comparative genomics** *Genome Biology* **20** <https://doi.org/10.1186/s13059-019-1832-y> | Google Scholar
20. Falcon S, Gentleman R (2007) **Using GOstats to test gene lists for GO term association** *Bioinformatics* **23**:257–258 <https://doi.org/10.1093/bioinformatics/btl567> | Google Scholar
21. Fan H, Ives AR, Surget-Groba Y, Cannon CH (2015) **An assembly and alignment- free method of phylogeny reconstruction from next-generation sequencing data** *BMC Genomics* **16** <https://doi.org/10.1186/s12864-015-1647-5> | Google Scholar
22. Fischl H, McManus D, Oldenkamp R, Schermelleh L, Mellor J, Jagannath A, Furger A (2020) **Cold-induced chromatin compaction and nuclear retention of clock mRNAs resets the circadian rhythm** *EMBO Journal* **39** <https://doi.org/10.15252/embj.2020105604> | Google Scholar
23. Forrest J, Miller-Rushing AJ (2010) **Toward a synthetic understanding of the role of phenology in ecology and evolution** *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**:3101–3112 <https://doi.org/10.1098/rstb.2010.0145> | Google Scholar

24. Fretwell SD (1972) **Populations in a seasonal environment** *Monographs in Population Biology* **5**:1–217 [Google Scholar](#)
25. Gaut B, Yang L, Takuno S, Eguiarte LE (2011) **The patterns and causes of variation in plant nucleotide substitution rates** *Annual Review of Ecology, Evolution, and Systematics* **42**:245–266 <https://doi.org/10.1146/annurev-ecolsys-102710-145119> | [Google Scholar](#)
26. Hernández-Plaza A, Szklarczyk D, Botas J, Cantalapiedra CP, Giner-Lamia J, Mende DR, Kirsch R, Rattei T, Letunic I, Jensen LJ, Bork P, von Mering C, Huerta-Cepas J. (2023) **eggNOG 6.0: enabling comparative genomics across 12 535 organisms** *Nucleic Acids Research* **51**:D389–D394 <https://doi.org/10.1093/nar/gkac1022> | [Google Scholar](#)
27. Hipp AL, Manos PS, Hahn M, Avishai M, Bodénès C, Cavender-Bares J, Crowl AA, Deng M, Denk T, Fitz-Gibbon S, Gailing O, González-Elizondo MS, González- Rodríguez A, Grimm GW, Jiang XL, Kremer A, Lesur I, McVay JD, Plomion C, Rodríguez-Correa H, Schulze ED, Simeone MC, Sork VL, Valencia-Avalos S (2020) **Genomic landscape of the global oak phylogeny** *New Phytologist* **226**:1198–1212 <https://doi.org/10.1111/nph.16162> | [Google Scholar](#)
28. Ikezaki Y, Kudo SN, Nakata T, Koita S, Munakata R, Yazaki K, Torimaru T, Tomaru N, Isobe S, Hirakawa H, Kusumi J, Satake A (2025) **Molecular evolution of terpene synthase underlying the diversification of isoprene emission in Fagaceae** *bioRxiv* <https://doi.org/10.1101/2025.07.31.667835> | [Google Scholar](#)
29. Irie N, Kuratani S (2011) **Comparative transcriptome analysis reveals vertebrate phylotypic period during organogenesis** *Nature Communications* **2** <https://doi.org/10.1038/ncomms1248> | [Google Scholar](#)
30. Kalinka AT, Varga KM, Gerrard DT, Preibisch S, Corcoran DL, Jarrells J, Ohler U, Bergman CM, Tomancak P (2010) **Gene expression divergence recapitulates the developmental hourglass model** *Nature* **468**:811–816 <https://doi.org/10.1038/nature09634> | [Google Scholar](#)
31. Kant P, Kant S, Gordon M, Shaked R, Barak S (2007) **Stress Response Suppressor1 and Stress Response Suppressor2, two Dead-box RNA helicases that attenuate Arabidopsis responses to multiple abiotic stresses** *Plant Physiology* **145**:814–830 <https://doi.org/10.1104/pp.107.099895> | [Google Scholar](#)
32. Keilwagen J, Hartung F, Grau J (2019) **GeMoMa: Homology-based gene prediction utilizing intron position conservation and RNA-seq data** *Methods in Molecular Biology Humana Press Inc* :161–177 https://doi.org/10.1007/978-1-4939-9173-0_9 | [Google Scholar](#)
33. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) **Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype** *Nature Biotechnology* **37**:907–915 <https://doi.org/10.1038/s41587-019-0201-4> | [Google Scholar](#)
34. Kokot M, Dlugosz M, Deorowicz S (2017) **KMC 3: counting and manipulating k-mer statistics** *Bioinformatics* **33**:2759–2761 <https://doi.org/10.1093/bioinformatics/btx304> | [Google Scholar](#)
35. Komoto H, Nagahama A, Miyawaki-Kuwakado A, Hata Y, Kyojuka J, Kajita Y, Toyama H, Satake A (2024) **The transcriptional changes underlying the flowering phenology shift of Arabidopsis halleri in response to climate warming** *Plant Cell & Environment* **47**:174–186 <https://doi.org/10.1111/pce.14716> | [Google Scholar](#)
36. Kremer A, Abbott AG, Carlson JE, Manos PS, Plomion C, Sisco P, Staton ME, Ueno S, Vendramin GG (2012) **Genomics of Fagaceae** *Tree Genetics & Genomes* **8**:583–610 <https://doi.org/10.1007>

37. Krishnakumar V, Hanlon MR, Contrino S, Ferlanti ES, Karamycheva S, Kim M, Rosen BD, Cheng CY, Moreira W, Mock SA, Stubbs J, Sullivan JM, Krampis K, Miller JR, Micklem G, Vaughn M, Town CD (2015) **Araport: The Arabidopsis Information Portal** *Nucleic Acids Research* **43**:D1003–D1009 <https://doi.org/10.1093/nar/gku1200> | [Google Scholar](#)
38. Kubo T, Iwasa Y (1996) **Phenological pattern of tree regeneration in a model for forest species diversity** *Theoretical Population Biology* **49**:90–117 <https://doi.org/10.1006/tpbi.1996.0004> | [Google Scholar](#)
39. Kudoh H (2016) **Molecular phenology in plants: in natura systems biology for the comprehensive understanding of seasonal responses under natural environments** *New Phytologist* **210**:399–412 <https://doi.org/10.1111/nph.13733> | [Google Scholar](#)
40. Lemos B, Bettencourt BR, Meiklejohn CD, Hartl DL (2005) **Evolution of proteins and gene expression levels are coupled in Drosophila and are independently associated with mRNA abundance, protein length, and number of protein- protein interactions** *Molecular Biology and Evolution* **22**:1345–1354 <https://doi.org/10.1093/molbev/msi122> | [Google Scholar](#)
41. Levin M, Anavy L, Cole AG, Winter E, Mostov N, Khair S, Senderovich N, Kovalev E, Silver DH, Feder M, Fernandez-Valverde SL, Nakanishi N, Simmons D, Simakov O, Larsson T, Liu SY, Jerafi-Vider A, Yaniv K, Ryan JF, Martindale MQ, Rink JC, Arendt D, Degnan SM, Degnan BM, Hashimshony T, Yanai I (2016) **The mid-developmental transition and the evolution of animal body plans** *Nature* **531**:637–641 <https://doi.org/10.1038/nature16994> | [Google Scholar](#)
42. Li B, Dewey CN (2011) **RSEM: Accurate transcript quantification from RNA-Seq data with or without a reference genome** *BMC Bioinformatics* **12** <https://doi.org/10.1186/1471-2105-12-323> | [Google Scholar](#)
43. Li H, Durbin R (2009) **Fast and accurate short read alignment with Burrows- Wheeler transform** *Bioinformatics* **25**:1754–1760 <https://doi.org/10.1093/bioinformatics/btp324> | [Google Scholar](#)
44. Liao BY, Zhang J (2006) **Low rates of expression profile divergence in highly expressed genes and tissue-specific genes during mammalian evolution** *Molecular Biology and Evolution* **23**:1119–1128 <https://doi.org/10.1093/molbev/msj119> | [Google Scholar](#)
45. Liu H, Zhang R, Zhou BF, Shen Z, Chen XY, Gao J, Wang B (2023) **Chromosome- scale genome assembly of sweet tea (Lithocarpus polystachyus Rehder)** *Scientific Data* **10** <https://doi.org/10.1038/s41597-023-02791-y> | [Google Scholar](#)
46. Lotharukpong JS, Zheng M, Luthringer R, Liesner D, Drost H-G, Coelho SM (2024) **A transcriptomic hourglass in brown algae** *Nature* **635**:129–135 <https://doi.org/10.1038/s41586-024-08059-8> | [Google Scholar](#)
47. Manos PS, Zhou Z, Cannon CH (2001) **Systematics of Fagaceae: phylogenetic tests of reproductive trait evolution** *International Journal of Plant Sciences* **162**:1361–1379 <https://doi.org/10.1086/322949> | [Google Scholar](#)
48. Mishra B, Ulaszewski B, Meger J, Aury JM, Bodénès C, Lesur-Kupin I, Pfenninger M, Da Silva C, Gupta DK, Guichoux E, Heer K, Lalanne C, Labadie K, Opgenoorth L, Ploch S, Le Provost G, Salse J, Scotti I, Wötzel S, Plomion C, Burczyk J, Thines M. (2022) **A chromosome-level genome assembly of the European Beech (Fagus sylvatica) reveals anomalies for organelle DNA**

integration, repeat content and distribution of SNPs *Frontiers in Genetics* **12** <https://doi.org/10.3389/fgene.2021.691058> | Google Scholar

49. Miyawaki-Kuwakado A, Han Q, Kitamura K, Satake A (2024) **Impacts of climate change on the transcriptional dynamics and timing of bud dormancy release in Yoshino - cherry tree** *Plants, People, Planet* :1505–1521 <https://doi.org/10.1002/ppp3.10548> | Google Scholar
50. Miyazaki Y, Maruyama Y, Chiba Y, Kobayashi MJ, Joseph B, Shimizu KK, Mochida K, Hiura T, Kon H, Satake A (2014) **Nitrogen as a key regulator of flowering in *Fagus crenata*: Understanding the physiological mechanism of masting by gene expression analysis** *Ecology Letters* **17**:1299–1309 <https://doi.org/10.1111/ele.12338> | Google Scholar
51. Morellato LPC, Camargo MGG, Gressler E (2013) **A review of plant phenology in South and Central America** In: *Phenology: An Integrative Environmental Science* Springer | Google Scholar
52. Nagano AJ, Kawagoe T, Sugisaka J, Honjo MN, Iwayama K, Kudoh H (2019) **Annual transcriptome dynamics in natural environments reveals plant seasonal adaptation** *Nature Plants* **5**:74–83 <https://doi.org/10.1038/s41477-018-0338-z> | Google Scholar
53. Naqvi S, Godfrey AK, Hughes JF, Goodheart ML, Mitchell RN, Page DC (2019) **Conservation, acquisition, and functional impact of sex-biased gene expression in mammals** *Science* **1979** <https://doi.org/10.1126/science.aaw7317> | Google Scholar
54. Nishio H, Nagano AJ, Ito T, Suzuki Y, Kudoh H (2020) **Seasonal plasticity and diel stability of H3K27me3 in natural fluctuating environments** *Nature Plants* **6**:1091–1097 <https://doi.org/10.1038/s41477-020-00757-1> | Google Scholar
55. Nuzhdin S V., Wayne ML, Harmon KL, McIntyre LM (2004) **Common pattern of evolution of gene expression level and protein sequence in *Drosophila*** *Molecular Biology and Evolution* **21**:1308–1317 <https://doi.org/10.1093/molbev/msh128> | Google Scholar
56. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C (2017) **Salmon provides fast and bias-aware quantification of transcript expression** *Nature Methods* **14**:417–419 <https://doi.org/10.1038/nmeth.4197> | Google Scholar
57. Pertea M, Pertea G (2020) **GFF Utilities: GffRead and GffCompare** *F1000Res* <https://doi.org/10.12688/f1000research.23297.1> | Google Scholar
58. Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, Liu L, Lian X, Shen M, Zhu X (2019) **Plant phenology and global climate change: Current progresses and challenges** *Global Change Biology* **25**:1922–1940 <https://doi.org/10.1111/gcb.14619> | Google Scholar
59. Prud'homme B, Gompel N, Carroll SB (2007) **Emerging principles of regulatory evolution** *PNAS* **104**:8605–8612 <https://doi.org/10.1073/pnas.0700488104> | Google Scholar
60. Putnam NH, O'Connell BL, Stites JC, Rice BJ, Blanchette M, Calef R, Troll CJ, Fields A, Hartley PD, Sugnet CW, Haussler D, Rokhsar DS, Green RE (2016) **Chromosome-scale shotgun assembly using an in vitro method for long-range linkage** *Genome Research* **26**:342–350 <https://doi.org/10.1101/gr.193474.115> | Google Scholar
61. Quint M, Drost HG, Gabel A, Ullrich KK, Bönn M, Grosse I (2012) **A transcriptomic hourglass in plant embryogenesis** *Nature* **490**:98–101 <https://doi.org/10.1038/nature11394> | Google Scholar

62. Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020) **GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes** *Nature Communications* **11** <https://doi.org/10.1038/s41467-020-14998-3> | [Google Scholar](#)
63. Satake A, Hagiwara T, Nagano AJ, Yamaguchi N, Sekimoto K, Shiojiri K, Sudo K (2024a) **Plant molecular phenology and climate feedbacks mediated by BVOCs** *Annual Review of Plant Biology* **75**:605–627 <https://doi.org/10.1146/annurev-arplant-060223-032108> | [Google Scholar](#)
64. Satake A, Imai R, Fujino T, Tomimoto S, Ohta K, Na'iem M, Indrioko S, Widiyatno W, Purnomo S, Morales AM, Nizhynska V, Tani N, Suyama Y, Sasaki E, Kasahara M (2024b) **Somatic mutation rates scale with time not growth rate in long-lived tropical trees** *eLife* **12** <https://doi.org/10.7554/eLife.88456> | [Google Scholar](#)
65. Satake A, Nagahama A, Sasaki E (2022) **A cross - scale approach to unravel the molecular basis of plant phenology in temperate and tropical climates** *New Phytologist* **233**:2340–2353 <https://doi.org/10.1111/nph.17897> | [Google Scholar](#)
66. Satake A, Ohta K, Takeda-Kamiya N, Toyooka K, Kusumi J (2023) **Seasonal gene expression signatures of delayed fertilization in Fagaceae** *Molecular Ecology* **32**:4801–4813 <https://doi.org/10.1111/mec.17079> | [Google Scholar](#)
67. Schwartz MD (2013) **Phenology: An Integrative Environmental Science** Springer Dordrecht [Google Scholar](#)
68. Smaczniak C, Immink RGH, Muiño JM, Blanvillain R, Busscher M, Busscher- Lange J, Dinh QD, Liu S, Westphal AH, Boeren S, Parcy F, Xu L, Carles CC, Angenent GC, Kaufmann K (2012) **Characterization of MADS-domain transcription factor complexes in Arabidopsis flower development** *PNAS* **109**:1560–1565 <https://doi.org/10.1073/pnas.1112871109> | [Google Scholar](#)
69. Smid M, Coebergh van den Braak RRJ, van de Werken HJG, van Riet J, van Galen A, de Weerd V, van der Vlugt-Daane M, Bril SI, Lalmahomed ZS, Kloosterman WP, Wilting SM, Foekens JA, IJzermans JNM, Martens JWM, Sieuwerts AM (2018) **Gene length corrected trimmed mean of M-values (GeTMM) processing of RNA-seq data performs similarly in intersample analyses while improving intrasample comparisons** *BMC Bioinformatics* **19** <https://doi.org/10.1186/s12859-018-2246-7> | [Google Scholar](#)
70. Thaben PF, Westermark PO (2014) **Detecting rhythms in time series with rain** *Journal of Biological Rhythms* **29**:391–400 <https://doi.org/10.1177/0748730414553029> | [Google Scholar](#)
71. True JR, Haag ES (2001) **Developmental system drift and flexibility in evolutionary trajectories** *Evolution & Development* **3**:109–119 <https://doi.org/10.1046/j.1525-142x.2001.003002109.x> | [Google Scholar](#)
72. Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee TH, Jin H, Marler B, Guo H, Kissinger JC, Paterson AH (2012) **MCScanX: A toolkit for detection and evolutionary analysis of gene synteny and collinearity** *Nucleic Acids Research* **40** <https://doi.org/10.1093/nar/gkr1293> | [Google Scholar](#)
73. Yang L, Gaut BS (2011) **Factors that contribute to variation in evolutionary rate among arabidopsis genes** *Molecular Biology and Evolution* **28**:2359–2369 <https://doi.org/10.1093/molbev/msr058> | [Google Scholar](#)
74. Yang Z (2007) **PAML 4: Phylogenetic analysis by maximum likelihood** *Molecular Biology and Evolution* **24**:1586–1591 <https://doi.org/10.1093/molbev/msm088> | [Google Scholar](#)

75. Yoo SJ, Chung KS, Jung SH, Yoo SY, Lee JS, Ahn JH (2010) **BROTHER of FT and TFL1 (BFT) has TFL1-like activity and functions redundantly with TFL1 in inflorescence meristem development in Arabidopsis** *Plant Journal* **63**:241–253 <https://doi.org/10.1111/j.1365-313X.2010.04234.x> | Google Scholar
76. Zhang X, Yu G, Dai Y, Zhang H, Wang K, Han J (2023) **High-resolution Hi-C maps highlight multiscale chromatin architecture reorganization during cold stress in Brachypodium distachyon** *BMC Plant Biology* **23** <https://doi.org/10.1186/s12870-023-04269-w> | Google Scholar
77. Zhou BF, Yuan S, Crowl AA, Liang YY, Shi Y, Chen XY, An QQ, Kang M, Manos PS, Wang B (2022) **Phylogenomic analyses highlight innovation and introgression in the continental radiations of Fagaceae across the Northern Hemisphere** *Nature Communications* **13** <https://doi.org/10.1038/s41467-022-28917-1> | Google Scholar
78. Zhu GQ, Qu L, Xue HW (2024) **Casein kinase 1 AELs promote senescence by enhancing ethylene biosynthesis through phosphorylating WRKY22 transcription factor** *New Phytologist* **244**:116–130 <https://doi.org/10.1111/nph.19785> | Google Scholar
79. Zoldos V, Papes D, Cerbah M, Panaud O, Besendorfer V, Siljak-Yakovlev S (1999) **Molecular-cytogenetic studies of ribosomal genes and heterochromatin reveal conserved genome organization among 11 Quercus species** *Theoretical and Applied Genetics* **99**:969–977 <https://doi.org/10.1007/s001220051404> | Google Scholar
80. Zrimec J, Börlin CS, Buric F, Muhammad AS, Chen R, Siewers V, Verendel V, Nielsen J, Töpel M, Zelezniak A (2020) **Deep learning suggests that gene expression is encoded in all parts of a co-evolving interacting gene regulatory structure** *Nature Communications* **11** <https://doi.org/10.1038/s41467-020-19921-4> | Google Scholar
81. Zrimec J, Buric F, Kokina M, Garcia V, Zelezniak A (2021) **Learning the Regulatory Code of Gene Expression** *Frontiers in Molecular Biosciences* **8** <https://doi.org/10.3389/fmolb.2021.673363> | Google Scholar

Author information

Shuichi N Kudo

Graduate School of Systems Life Sciences, Kyushu University, Fukuoka, Japan
ORCID iD: [0009-0003-9385-1794](https://orcid.org/0009-0003-9385-1794)

Yuka Ikezaki

Department of Biology, Faculty of Science, Kyushu University, Fukuoka, Japan

Junko Kusumi

Department of Environmental Changes, Faculty of Social and Cultural Studies, Kyushu University, Fukuoka, Japan
ORCID iD: [0000-0002-1693-7071](https://orcid.org/0000-0002-1693-7071)

Hideki Hirakawa

Kazusa DNA Research Institute, Kisarazu, Japan, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka, Japan
ORCID iD: [0000-0002-7128-7390](https://orcid.org/0000-0002-7128-7390)

Sachiko Isobe

Kazusa DNA Research Institute, Kisarazu, Japan, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan

ORCID iD: [0000-0002-9555-5054](https://orcid.org/0000-0002-9555-5054)

Akiko Satake

Department of Biology, Faculty of Science, Kyushu University, Fukuoka, Japan

ORCID iD: [0000-0002-0831-8617](https://orcid.org/0000-0002-0831-8617)

For correspondence: satake.akiko.269@m.kyushu-u.ac.jp

Editors

Reviewing Editor

Vincent Lynch

University at Buffalo, State University of New York, Buffalo, United States of America

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #2 (Public review):

This study investigates how seasonal environments shape the evolution of gene expression by analyzing two-year time-series transcriptomes from leaves and buds of four Fagaceae tree species. The revised manuscript incorporates additional data and analyses that directly address earlier concerns about sampling design and environmental variation, thereby strengthening the robustness of the conclusions.

The major strengths of this work are the scale and quality of the dataset, the integration of genome assemblies with time-series transcriptomics, and the careful analyses showing that winter bud expression is strongly conserved across species. The additional samples and re-analyses demonstrate convincingly that these results are not artifacts of sampling period or site differences. The study also links gene expression dynamics to phenological observations and frames its findings in relation to broader evolutionary concepts such as phenological synchrony and the developmental hourglass model.

Remaining limitations include the absence of direct mechanistic analyses of cis-regulatory and chromatin-level processes, the relatively coarse resolution of phenological trait measurements, and the weak association between seasonal expression divergence and sequence divergence. Importantly, these limitations are now explicitly acknowledged in the revised Discussion and framed as directions for future research.

Overall, the authors have substantially achieved their aims. This revised version represents a robust and convincing contribution that provides valuable data resources and conceptual insights into how seasonal environments constrain and shape gene expression. It will be of interest not only to evolutionary biologists and plant scientists, but also to researchers considering the broader role of environmental cycles in gene regulatory evolution.

<https://doi.org/10.7554/eLife.107309.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews**Reviewer #1 (Public review):***Summary:*

The authors performed genome assemblies for two Fagaceae species and collected transcriptome data from four natural tree species every month over two years. They identified seasonal gene expression patterns and further analyzed species-specific differences.

Strengths:

The study of gene expression patterns in natural environments, as opposed to controlled chambers, is gaining increasing attention. The authors collected RNA-seq data monthly for two years from four tree species and analyzed seasonal expression patterns. The data are novel. The authors could revise the manuscript to emphasize seasonal expression patterns in three species (with one additional species having more limited data). Furthermore, the chromosome-scale genome assemblies for the two Fagaceae species represent valuable resources, although the authors did not cite existing assemblies from closely related species.

Thank you for your careful assessment of our manuscript.

Weaknesses:

Comment; The study design has a fundamental flaw regarding the evaluation of genetic or evolutionary effects. As a basic principle in biology, phenotypes, including gene expression levels, are influenced by genetics, environmental factors, and their interaction. This principle is well-established in quantitative genetics.

In this study, the four species were sampled from three different sites (see Materials and Methods, lines 543-546), and additionally, two species were sampled from 2019-2021, while the other two were sampled from 2021-2023 (see Figure S2). This critical detail should be clearly described in the Results and Materials and Methods. Due to these variations in sampling sites and periods, environmental conditions are not uniform across species.

Even in studies conducted in natural environments, there are ways to design experiments that allow genetic effects to be evaluated. For example, by studying co-occurring species, or through transplant experiments, or in common gardens. To illustrate the issue, imagine an experiment where clones of a single species were sampled from three sites and two time periods, similar to the current design. RNA-seq analysis would likely detect differences that could qualitatively resemble those reported in this manuscript.

One example is in line 197, where genus-specific expression patterns are mentioned. While it may be true that the authors' conclusions (e.g., winter synchronization, phylogenetic constraints) reflect real biological trends, these conclusions are also predictable even without empirical data, and the current dataset does not provide quantitative support.

If the authors can present a valid method to disentangle genetic and environmental effects from their dataset, that would significantly strengthen the manuscript. However, I

do not believe the current study design is suitable for this purpose.

Unless these issues are addressed, the use of the term "evolution" is inappropriate in this context. The title should be revised, and the result sections starting from "Peak months distribution..." should be either removed or fundamentally revised. The entire Discussion section, which is based on evolutionary interpretation, should be deleted in its current form.

*If the authors still wish to explore genetic or evolutionary analyses, the pair of *L. edulis* and *L. glaber*, which were sampled at the same site and over the same period, might be used to analyze "seasonal gene expression divergence in relation to sequence divergence." Nevertheless, the manuscript would benefit from focusing on seasonal expression patterns without framing the study in evolutionary terms.*

We sincerely thank the reviewer for the detailed and thoughtful comments. We fully recognize the importance of carefully distinguishing genetic and environmental contributions in transcriptomic studies, particularly when addressing evolutionary questions. The reviewer identified two major concerns regarding our study design: (1) the use of different monitoring periods across species, and (2) the use of samples collected from different study sites. We addressed both concerns with additional analyses using 112 new samples and now present new evidence that supports the robustness of our conclusions.

(1) Monitoring period variation does not bias our conclusions

To address concerns about the differing monitoring periods, we added new RNA-seq data (42 samples each for bud and leaf samples for *L. glaber* and 14 samples each for bud and leaf samples for *L. edulis*) collected from November 2021 to November 2022, enabling direct comparison across species within a consistent timeframe. Hierarchical clustering of this expanded dataset (Fig. S6) yielded results consistent with our original findings: winter-collected samples cluster together regardless of species identity. This strongly supports our conclusion that the seasonal synchrony observed in winter is not an artifact of the monitoring period and demonstrates the robustness of our conclusions across datasets.

(2) Site variation is limited and does not confound our findings

Although the study included three sites, two of them (Imajuku and Ito Campus) are only 7.3 km apart, share nearly identical temperature profiles (see Fig. S2), and are located at the edge of similar evergreen broadleaf forests. Only *Q. acuta* was sampled from a higher-altitude, cooler site. To assess whether the higher elevation site of *Q. acuta* introduced confounding environmental effects, we reanalyzed the data after excluding this species. Hierarchical clustering still revealed that winter bud samples formed a distinct cluster regardless of species identity (Fig. S7), consistent with our original finding.

Furthermore, we recalculated the molecular phenology divergence index D (Fig. 4C) and the interspecific Pearson's correlation coefficients (Fig. 5A) without including *Q. acuta*. These analyses produced results that were similar to those obtained from the full dataset (Fig. S12; Fig. S14), indicating that the observed patterns are not driven by environmental differences associated with elevation.

(3) Justification for our approach in natural systems

We agree with the reviewer that experimental approaches such as common gardens, reciprocal transplants, and the use of co-occurring species are valuable for disentangling genetic and environmental effects. In fact, we have previously implemented such designs in studies using the perennial herb *Arabidopsis halleri* (Komoto et al., 2022, <https://doi.org/10.1111/pce.14716>) and clonal *Someiyoshino* cherry trees (Miyawaki-Kuwakado et al., 2024, <https://doi.org/10.1002/ppp3.10548>) to examine environmental effects on gene expression. However, extending these approaches to long-lived tree species in diverse natural ecosystems poses significant logistical and biological challenges. In this study, we addressed this

limitation by including three co-occurring species at the same site, which allowed us to evaluate interspecific differences under comparable environmental conditions. Importantly, even when we limited our analyses to these co-occurring species, the results remained consistent, indicating that the observed variation in transcriptomic profiles cannot be attributed to environmental factors alone and likely reflects underlying genetic influences.

Accordingly, we added four new figures (Fig. S6, Fig. S7, Fig. S12 and Fig. S14) and revised the manuscript to clarify the limitations and strengths of our design, to tone down the evolutionary claims where appropriate, and to more explicitly define the scope of our conclusions in light of the data. We hope that these efforts sufficiently address the reviewer's concerns and strengthen the manuscript.

To better support the seasonal expression analysis, the early RNA-seq analysis sections should be strengthened. There is little discussion of biological replicate variation or variation among branches of the same individual. These could be important factors to analyze. In line 137, the mapping rate for two species is mentioned, but the rates for each species should be clearly reported. One RNA-seq dataset is based on a species different from the reference genome, so a lower mapping rate is expected. While this likely does not hinder downstream analysis, quantification is important.

We thank the reviewer 1 for the helpful comment. To evaluate the variation among biological replicates, we compared the expression level of each gene across different individuals. We observed high correlation between each pair of individuals (*Q. glauca* ($n=3$): an average correlation coefficient $r = 0.947$; *Q. acuta* ($n=3$): $r = 0.948$; *L. glaber* ($n=3$): $r = 0.948$). This result suggests that the seasonal gene expression pattern is highly synchronized across individuals within the same species. We mentioned this point in the Result section in the revised manuscript. We also calculated the mean mapping rates for each species. As the reviewer expected, the mapping rate was slightly lower in *Q. acuta* ($88.6 \pm 2.3\%$) and *L. glaber* ($84.3 \pm 5.4\%$), whose RNA-Seq data were mapped to reference genomes of related but different species, compared to that in *Q. glauca* ($92.6 \pm 2.2\%$) and *L. edulis* ($89.3 \pm 2.7\%$). However, we minimized the impact of these differences on downstream analysis. These details have been included in the revised main text.

In Figures 2A and 2B, clustering is used to support several points discussed in the Results section (e.g., lines 175-177). However, clustering is primarily a visualization method or a hypothesis-generating tool; it cannot serve as a statistical test. Stronger conclusions would require further statistical testing.

We thank the reviewer for the helpful comment. As noted, we acknowledge that hierarchical clustering (Fig. 2A) is primarily a visualization and hypothesis-generating method. To assess the biological relevance of the clusters identified, we conducted a Mann-Whitney U test or the Steel-Dwass test to evaluate whether the environmental temperatures at the time of sample collection differed significantly among the clusters. This analysis (Fig. 2B) revealed statistically significant differences in temperature in the cluster B3 ($p < 0.01$), indicating that the gene expression clusters are associated with seasonal thermal variation. These results support the interpretation that the clusters reflect coordinated transcriptional responses to environmental temperature. We revised the Results section to clarify this point.

The quality of the genome assemblies appears adequate, but related assemblies should be cited and discussed. Several assemblies of Fagaceae species already exist, including Quercus mongolica (Ai et al., Mol Ecol Res, 2022), Q. gilva (Front Plant Sci, 2022), and Fagus sylvatica (GigaScience, 2018), among others. Is there any novelty here? Can you compare your results with these existing assemblies?

We agree that genome assemblies of Fagaceae species are becoming increasingly available. However, our study does not aim to emphasize the novelty of the genome assemblies per se. Rather, with the increasing availability of chromosome-level genomes, we regard genome assembly as a necessary foundation for more advanced analyses. The main objective of our study is to investigate how each gene is expressed in response to seasonal environmental changes, and to link genome information with seasonal transcriptomic dynamics. To address the reviewer's comment in line with this objective, we added a discussion on the syntenic structure of eight genome assemblies spanning four genera within the Fagaceae, including a species from the genus *Fagus* (Ikezaki et al. 2025, <https://doi.org/10.1101/2025.07.31.667835>). This addition helps to position our work more clearly within the context of existing genomic resources.

Most importantly, Figure 1B-D shows synteny between the two genera but also indicates homology between different chromosomes. Does this suggest paleopolyploidy or another novel feature? These chromosome connections should be interpreted in the main text-even if they could be methodological artifacts.

A previous study on genome size variation in Fagaceae suggested that, given the consistent ploidy level across the family, genome expansion likely occurred through relatively small segmental duplications rather than whole-genome duplications. Because Figure 1B-D supports this view, we cited the following reference in the revised version of the manuscript. Chen et al. (2014) <https://doi.org/10.1007/s11295-014-0736-y>

In both the Results and Materials and Methods sections, descriptions of genome and RNA-seq data are unclear. In line 128, a paragraph on genome assembly suddenly introduces expression levels. RNA-seq data should be described before this. Similarly, in line 238, the sentence "we assembled high-quality reference genomes" seems disconnected from the surrounding discussion of expression studies. In line 632, Illumina short-read DNA sequencing is mentioned, but it's unclear how these data were used.

We relocated the explanation regarding the expression levels of single-copy and multi-copy genes to the section titled "Seasonal gene expression dynamics." Additionally, we clarified in the Materials and Methods section that short-read sequencing data were used for both genome size estimation and phylogenetic reconstruction.

Reviewer #2 (Public review):

Summary:

This study explores how gene expression evolves in response to seasonal environments, using four evergreen Fagaceae species growing in similar habitats in Japan. By combining chromosome-scale genome assemblies with a two-year RNA-seq time series in leaves and buds, the authors identify seasonal rhythms in gene expression and examine both conserved and divergent patterns. A central result is that winter bud expression is highly conserved across species, likely due to shared physiological demands under cold conditions. One of the intriguing implications of this study is that seasonal cycles might play a role similar to ontogenetic stages in animals. The authors touch on this by comparing their findings to the developmental hourglass model, and indeed, the recurrence of phenological states such as winter dormancy may act as a cyclic form of developmental canalization, shaping expression evolution in a way analogous to embryogenesis in animals.

Strengths:

- (1) *The evolutionary effects of seasonal environments on gene expression are rarely studied at this scale. This paper fills that gap.*
- (2) *The dataset is extensive, covering two years, two tissues, and four tree species, and is well suited to the questions being asked.*
- (3) *Transcriptome clustering across species (Figure 2) shows strong grouping by season and tissue rather than species, suggesting that the authors effectively controlled for technical confounders such as batch effects and mapping bias.*
- (4) *The idea that winter imposes a shared constraint on gene expression, especially in buds, is well argued and supported by the data.*
- (5) *The discussion links the findings to known concepts like phenological synchrony and the developmental hourglass model, which helps frame the results.*

We are grateful for the reviewer for the detailed and thoughtful review of our manuscript.

Weaknesses:

- (1) *While the hierarchical clustering shown in Figure 2A largely supports separation by tissue type and season, one issue worth noting is that some leaf samples appear to cluster closely with bud samples. The authors do not comment on this pattern, which raises questions about possible biological overlap between tissues during certain seasonal transitions or technical artifacts such as sample contamination. Clarifying this point would improve confidence in the interpretation of tissue-specific seasonal expression patterns.*

Leaf samples clustered into the bud are newly flushed leaves collected in April for *Q. glauca*, May for *Q. acuta*, May and June for *L. edulis*, and August and September for *L. glaber*. To clarify this point, we highlighted these newly flushed leaf samples as asterisk in the revised figure (Fig. 2A).

- (2) *While the study provides compelling evidence of conserved and divergent seasonal gene expression, it does not directly examine the role of cis-regulatory elements or chromatin-level regulatory architecture. Including regulatory genomic or epigenomic data would considerably strengthen the mechanistic understanding of expression divergence.*

We thank the reviewer for this insightful comment. As noted in the Discussion section, we hypothesize that such genome-wide seasonal expression patterns—and their divergence across species—are likely mediated by cis-regulatory elements and chromatin-level mechanisms. While a direct investigation of regulatory architecture was beyond the scope of the present study, we fully agree that incorporating regulatory genomic and epigenomic data would significantly deepen the mechanistic understanding of expression divergence. In this regard, we are currently working to identify putative cis-regulatory elements in non-coding regions and are collecting epigenetic data from the same tree species using ChIP-seq. We believe the current study provide a foundation for these future investigations into the regulatory basis of seasonal transcriptome variation. We made a minor revision to the Discussion to note that an important future direction is to investigate the evolution of non-coding sequences that regulate gene expression in response to seasonal environmental changes.

- (3) *The manuscript includes a thoughtful analysis of flowering-related genes and seasonal GO enrichment (e.g., Figure 3C-D), providing an initial link between gene*

expression timing and phenological functions. However, the analysis remains largely gene-centric, and the study does not incorporate direct measurements of phenological traits (e.g., flowering or bud break dates). As a result, the connection between molecular divergence and phenotypic variation, while suggestive, remains indirect.

We would like to note that phenological traits have been observed in the field on a monthly basis throughout the sampling period and the phenological data were plotted together with molecular phenology (e.g. Fig. 2A, C; Fig. 3C, D). Although the temporal resolution is limited, these observations captured species-specific differences in key phenological events such as leaf flushing and flowering times. We revised the manuscript to clarify this point.

*(4) Although species were sampled from similar habitats, one species (*Q. acuta*) was collected at a higher elevation, and factors such as microclimate or local photoperiod conditions could influence expression patterns. These potential confounding variables are not fully accounted for, and their effects should be more thoroughly discussed or controlled in future analyses.*

We fully agree with the reviewer that local environmental conditions, including microclimate and photoperiod differences, could potentially influence gene expression patterns. To assess whether the higher elevation site of *Q. acuta* introduced confounding environmental effects, we reanalyzed the data after excluding this species. Hierarchical clustering still revealed that winter bud samples formed a distinct cluster regardless of species identity (Fig. S7), consistent with our original finding.

Furthermore, we recalculated the molecular phenology divergence index D (Fig. 4C) and the interspecific Pearson's correlation coefficients (Fig. 5A) without including *Q. acuta*. These analyses produced results that were qualitatively similar to those obtained from the full dataset (Fig. S12; Fig. S14), indicating that the observed patterns are not driven by environmental differences associated with elevation.

We believe these additional analyses help to decouple the effects of environment and genetics, and support our conclusion that both seasonal synchrony and phylogenetic constraints play key roles in shaping transcriptome dynamics. We added four new figures (Fig. S6, Fig. S7, Fig. S12 and Fig. S14) and revised the text accordingly to clarify this point and to acknowledge the potential impact of site-specific environmental variation.

(5) Statistical and Interpretive Concerns Regarding $\Delta\phi$ and dN/dS Correlation (Figures 5E and 5F):

a) Statistical Inappropriateness: $\Delta\phi$ is a discrete ordinal variable (likely 1-11), making it unsuitable for Pearson correlation, which assumes continuous, normally distributed variables. This undermines the statistical validity of the analysis.

We thank the reviewer for the insightful comment. We would like to clarify that the analysis presented in Figures 5E and 5F was based on linear regression, not Pearson's correlation. Although $\Delta\phi$ is a discrete variable, it takes values from 0 to 6 in 0.5 increments, resulting in 13 levels. We treated it as a quasi-continuous variable for the purposes of linear regression analysis. This approach is commonly adopted in practice when a discrete variable has sufficient resolution and ordering to approximate continuity. To enhance clarity, we revised the manuscript to explicitly state that linear regression was used, and we now reported the regression coefficient and associated *p*-value to support the interpretation of the observed trend.

b) Biological Interpretability: Even with the substantial statistical power afforded by genome-wide analysis, the observed correlations are extremely weak. This suggests that

the relationship, if any, between temporal divergence in expression and protein-coding evolution is negligible.

Taken together, these issues weaken the case for any biologically meaningful association between $\Delta\phi$ and dN/dS . I recommend either omitting these panels or clearly reframing them as exploratory and statistically limited observations.

We agree with the reviewer's comment. While we retained the original panels, we reframed our interpretation to emphasize that, despite statistical significance, the observed correlation is very weak—suggesting that coding region variation is unlikely to be the primary driver of seasonal gene expression patterns. Accordingly, we revised the “Relating seasonal gene expression divergence to sequence divergence” section in the Results, as well as the relevant part of the Discussion.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

Sentences around lines 250-251 are incomplete and need revision.

We thank the reviewer for pointing this out. We revised the sentences in the subsection “Peak month distribution of rhythmic genes and intra-genus and inter-genera comparison” in the Results section to ensure clarity and completeness. In addition, to improve the interpretability of the peak month distribution, we added arrows to indicate the major peaks in the circular histograms shown in Fig. 3C and 3D.

Reviewer #2 (Recommendations for the authors):

(1) In Figure 1E-G, the term Copy number or Copy number variation could be misleading, as it is commonly associated with inter-individual gene copy number variation in a population. Since the analysis here refers to orthology relationships rather than population-level variation, a more precise term, such as orthogroup classification, may be preferable.

We thank the reviewer for this helpful suggestion. We agree that the term “copy number” could be misleading in this context. Accordingly, we updated the labeling in Fig. 1 to reflect the more precise term “orthogroup classification.”

(2) In Figure 3A, the x-axis label Period (month) may be misleading, as it could be mistaken for calendar months rather than referring to the periodicity of gene expression cycles. A more explicit label, such as Expression periodicity (months), might improve clarity for the reader.

We thank the reviewer for this valuable suggestion. In the original version of Fig. 3A, we used the label “Period (month),” which could indeed be misinterpreted as referring to calendar months. To clarify that this axis represents the length of gene expression cycles, we revised the label to “Period length (months).” This change also aligns with the terminology used throughout the manuscript, where “Period” refers specifically to cycle length, and “Periodicity” denotes the presence or absence of rhythmic expression.

Other minor revisions

We also made minor revisions for the reference list and the grant number details, and included the accession numbers for all DNA and RNA sequence data deposited in the DNA Data Bank of Japan (DDBJ) in the Data deposition and code availability section, in addition to the BioProject ID.

<https://doi.org/10.7554/eLife.107309.2.sa0>