

Occupancy of the HbYX hydrophobic pocket is sufficient to induce gate opening in the archaeal 20S proteasomes

Janelle JY Chuah, Madalena R Daugherty, David M Smith 

Department of Biochemistry and Molecular Medicine, West Virginia University School of Medicine, Morgantown, United States • Department of Neuroscience, Rockefeller Neuroscience Institute, West Virginia University, Morgantown, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **useful** manuscript describes cryo-EM structures of archaeal proteasomes that reveal insights into how occupancy of binding pockets on the 20S protease regulates proteasome gating. The evidence supporting these claims is **convincing**, although inclusion of more quantitative comparisons would help strengthen the conclusions. This work will be of special interest to researchers interested in proteasome structure and regulation.

<https://doi.org/10.7554/eLife.106611.1.sa2>

Abstract

Enhancing proteasome function has been a long-standing but challenging target of interest for the potential treatment of neurodegenerative diseases, emphasizing the importance of understanding proteasome activation mechanisms. Most proteasome activator complexes use the C-terminal HbYX (hydrophobic-tyrosine-almost any residue) motif to bind and trigger gate-opening in the 20S proteasome. This study defines a critical molecular interaction in the HbYX mechanism that triggers gate opening. We focus on the Hb site interaction and find it plays a surprisingly central and crucial role in driving the allosteric conformational changes that induce gate opening in the archaeal 20S. We examined the cryo-EM structure of two mutant archaeal proteasomes, α V24Y T20S and α V24F T20S. These two mutants were engineered to place a bulky aromatic residue in the HbYX hydrophobic pocket; both mutants are highly active, though their mechanisms of activation are undefined. Collectively, our findings indicate that the interaction between the Hb group of the HbYX motif and its corresponding hydrophobic pocket is sufficient to induce gate opening in a mechanistically similar way to the HbYX motif. The activation mechanism studied here involves the expansion of the hydrophobic binding site, allosterically altering the state of the IT switch thus triggering gate-opening. Furthermore, we show that the canonical α K66 residue, previously understood to be critical for proteasome activator binding, also plays a key role in

stabilizing the open gate, irrespective of activator binding. This study differentiates between the residues in the HbYX motif that support binding interactions (“YX”) versus those that allosterically contribute to gate opening (“Hb”). The insights reported here will guide future drug development efforts, particularly in designing small molecule proteasome activators, by targeting the identified hydrophobic pocket.

Introduction

The proteasome is a molecular machine responsible for degrading misfolded, damaged, and unneeded proteins that are targeted via multiple mechanisms including the conventional pathway of ubiquitination(1–6). Proteasomal degradation regulates major cellular processes and functions, such as synaptic plasticity and memory formation(7–11). Impairment of proteasome function has correlated with areas of the brain affected by neurodegenerative diseases, supporting its implication in these diseases(12–17). Furthermore, increasing proteasome activity has been shown to induce protective or rescue effects in models of neurodegenerative diseases(18–23), strengthening the rationale to develop small molecule proteasome activators. The objective of this study is to provide a clear mechanistic framework of how proteasome activation occurs to inform drug discovery efforts.

The eukaryotic proteasome, also referred to as the 20S or the Core Particle (CP), consists of four stacked heteroheptameric (homoheptameric in archaea) rings arranged as $\alpha - \beta - \beta - \alpha$ (24). Three of the seven β subunits carry protease sites with distinct protease activities: caspase-like (β 1), trypsin-like (β 2), and chymotrypsin-like (β 5). The α subunits have an N-termini that extends toward the central pore, forming a gate that regulates substrate entry. The gate is primarily formed by the N-termini of α 2, 3, and 4 with α 3’s N-termini (the longest) acting as the lynchpin(24). The interactions that stabilize the open conformation have been reported in crystal and cryo-EM structures(24–31). A YDR motif, found on the α N-termini, interacts with the neighboring α subunit’s N-terminal tail, to stabilize the gate in the open state. Most recently, we discovered that the open and closed states of the α N-termini conformation are stabilized by two alternating conformations of the IT switch(31). Two α subunit residues, I12 and T13, in the archaeal *Thermoplasma acidophilum* proteasome (T20S), switch positions to occupy the pocket adjacent to Helix 0 and stabilize the open and closed gate conformation respectively. This IT switch is functionally conserved in the eukaryotic 20S(31). Basal kinetics of the archaeal proteasome gate, quantified using NMR(32), indicates that the gate fluctuates between open and close on a time scale of seconds; however, the gate primarily exists in the closed state of isolated 20S.

The 20S proteasome is known to associate with regulatory complexes, collectively referred to as Proteasome Activators (PAs)(33). PA700 or 19S, which associates with the 20S to form the 26S, has been identified as a critical component of the Ubiquitin-Proteasome Pathway. PA700 recognizes ubiquitinated substrates and unfolds them to be degraded by the 20S. Like many PAs, PA700 induces 20S gate-opening via the HbYX-dependent mechanism, as does the archaeal homolog of PA700 called PAN(33, 34). Three of the six C-terminal tails of PA700’s ATPases carry a HbYX motif (hydrophobic-tyrosine-almost any residue), whereby the “X” residue must be the C-terminal residue. These HbYX motifs interact with the inter-subunit pockets in the 20S α ring and allosterically induce gate-opening. The precise molecular mechanism of how the motif’s binding in inter-subunit pockets induces gate opening is not known, though high-resolution structures of the 26S(26–29, 35) and the 20S interacting with a di-peptide HbYX-mimetic(31) have generated plausible models. Structures of the 26S reveal that the C-terminal tails bind in a particular order during activation, though the correlation between the number of C-terminal tails/HbYX-bound to induce gate opening is controversial. A study indicated that partial gate opening occurs when Rpt2, 3, and 5, which have HbYX motifs, dock in their corresponding α inter-subunit pockets and full gate opening occurs when Rpt6, which does not have a HbYX motif, additionally binds(26). Another study indicated that the binding of Rpt2, 3, and 5 did not open

the gate, but the additional binding of Rpt6 causes full gate opening(28 [↗](#)). Interestingly, multiple studies suggest that Rpt1, which carries the penultimate tyrosine, but not the Hb group (partial-HbYX), must also bind in combination with Rpt2, 3, 5, and 6 for gate opening(27 [↗](#), 29 [↗](#), 35 [↗](#)). Collectively, these structures suggest that each inter-subunit pocket contributes to gate opening to varying extents, which correlates with the fact that each α subunit's N-terminal tails differentially participate in the closed state, though the minimum occupancy of inter-subunit pockets required for gate opening is unresolved.

While some PAs are HbYX-dependent like PA700, the 11S family, such as PA26 or PA28 activate the proteasome via a HbYX-independent mechanism. Despite the divergence between mechanisms, they converge on the critical molecular interactions and conformational changes (e.g. the IT switch) that stabilize the open gate state(31 [↗](#)). For example, both HbYX-dependent and HbYX-independent PAs have been shown to interact with α K66 side chain and mutating α K66 to alanine impedes PAs association and activation, validated by velocity sedimentation assay and proteasomal peptidase activity(30 [↗](#), 34 [↗](#)). Both mechanisms also involve a displacement of α Pro17(30 [↗](#), 31 [↗](#)) and switching of the IT switch(31 [↗](#)). Therefore, though the cause of the conformational changes mentioned may differ between HbYX-dependent and HbYX-independent mechanisms, they share in some of the PA-20S interactions that support the open conformational state of the gate.

Our early studies showed that the HbYX motif could function even on short 6-7 residue peptides(34 [↗](#)); recently we showed that HbYX-dependent gate opening could be reduced to a carboxybenzyl (CBZ)-blocked YA dipeptide called ZYA—a minimal HbYX mimetic. Our ZYA-T20S structure(31 [↗](#)) indicated the minimum molecular interactions necessary for optimal HbYX-dependent gate-opening by a small molecule. In our 1.9Å resolution structure, ZYA was observed to interact with residues in the α subunit (G19, K66, L81, etc.) in a similar orientation and fashion as the native HbYX motif of various PA complexes. Our cryo-EM data indicated that α V24 (and α A154) hydrophobically interacts with the carboxybenzyl of ZYA and the hydrophobic side chain of the PAs HbYX motif. Interestingly, our previous study also indicated that a single residue mutation on α V24 (i.e., α V24Y, α V24F) increased proteasome activity, though the mechanism of activation by this mutation was not determined. To assess the significance of these mutations in activating the proteasome via the HbYX-dependent pathway, we determined the cryo-EM structure of α V24Y T20S and α V24F T20S. The results of this study indicate that occupancy of a hydrophobic pocket by a bulky hydrophobic group such as α V24Y within the α inter-subunit pocket is sufficient to trigger 20S gate opening via the HbYX-dependent mechanism, demonstrating the surprising importance of the HbYX motif's single Hb group for allosteric activation of the 20S. The findings reported here are expected to translate to the human proteasome as ZYA is a robust activator of the mammalian 20S and the HbYX-dependent mechanism is conserved; thereby, uncovering a highly specific pharmaceutical target for proteasome activation.

Results

Cryo-EM map of α V24Y T20S shows HbYX-dependent-like conformational changes and an open gate state

To elucidate how the mutation, α V24Y, induce gate-opening in the archaeal proteasome (α V24Y T20S), we resolved its structure using cryo-EM (Fig. 1 [↗](#); Fig. S1 [↗](#) & S2 [↗](#)). The *Thermoplasma acidophilum* 20S (T20S) has D7 symmetry, which was applied during reconstruction, and the final map resolved to 2.38Å (EMD-44914). Comparing the α V24Y T20S electron density map against our wild-type T20S (WT T20S) map (EMD-28878) revealed conformational changes similar to those previously noted in our ZYA-T20S model(31 [↗](#)) (PDB:8F7K) (Fig. 1 [↗](#)). Alignment of the maps indicate the α V24Y mutation induced minimal conformational changes in the β rings, while the α subunits exhibited rotational changes that closely mimicked HbYX-dependent conformational

changes, such as those induced by ZYA binding(31, 36) (Fig. 1A & B). Comparison of the maps show the presence of the expected larger density of the mutant tyrosine side chain in the hydrophobic pocket of the α V24Y T20S map with proximity to the docking location of the carboxybenzyl group (Z) of ZYA (Fig. 1C, D, & E).

Our prior biochemical analysis showed that the α V24Y mutation caused substantial acceleration of peptide degradation, suggesting that this mutation might stimulate gate-opening(31); therefore, we sought to determine the conformational changes in the structure of α V24Y T20S. We observed significant densities corresponding to the YDR motif in the open state, designating an open gate conformation (Fig. 1F, G, & H). Additionally, the densities corresponding to the “IT switch” were in similar conformation to previously known open gate proteasome cryo-EM structures: ZYA-T20S (Fig. 1I, J, & K) and L81Y-T20S(31). It is also apparent in the map of α V24Y T20S that there is a lack of densities in the central cavity of the α ring that corresponding to the closed state of the α N-termini, definitively confirming that V24Y T20S’s gate is open (Fig. 1A & B). Collectively, the comparison of maps demonstrates that V24Y T20S is in an open state with global conformational changes in the α -subunits that mirror those of a HbYX-bound proteasome (Fig. 1).

α V24Y T20S shows HbYX dependent conformational changes that are greater in magnitude than those seen in the ZYA-T20S structure

To gain a more detailed understanding of the conformational changes induced by α V24Y, we generated an atomic model using the α V24Y T20S map. To evaluate the global conformational changes in the α subunits, we aligned the α V24Y T20S and the ZYA-T20S to the WT T20S model via their β subunits and overlayed all three models (Fig. 2). The α subunits in α V24Y T20S rotate roughly as a rigid body on a pivot around Helix 2 (Fig. 2A, B, C, & 3A), using WT T20S as a reference. This rotation around Helix 2 is also true for the ZYA-T20S structure and was noted previously(31), but the rotation in α V24Y T20S is greater than in ZYA-T20S (see arrows in Fig. 2A). The helical bundle composed of Helices 4, 5, and 6 (see asterisks in Fig. 2A) move the most with some residues in these helices moving 3–4Å in the α V24Y T20S model relative to WT, which is about twice the distance moved in the ZYA-T20S model(31). In addition, α P17 in the α V24Y moves about 1.6Å and Helix 0 moves ~1.4Å compared to WT (Fig. 2C), which is also about twice the distance these residues move in the ZYA-T20S structure. Taken together, the conformational changes induced by α V24Y are highly similar to those induce by binding of the HbYX mimetic ZYA, but they are of greater magnitude in these structures. The observed greater change is likely attributable to the α V24Y mutation affecting the entire proteasome population within that structure. In contrast, ZYA is a low-affinity ligand, anticipated to occupy only a subset of the binding sites in its structure’s population. Since cryo-EM analysis averages all structures, the reduced population with conformational changes in the ZYA-bound structure represents an average of closed and open states and thus, apparent conformational changes are smaller in the ZYA structure relative to the α V24Y mutant.

α K66 has long been known to be critical for binding to the C-terminal carboxyl of PA’s(30, 31, 34). We previously observed that α K66 was repositioned by the binding of ZYA’s carboxyl(31). Since the inter-subunit pocket in the α V24Y T20S has no ligand bound, we expected α K66 to have a similar conformation to WT, but we were surprised to find that K66 was similarly repositioned in the α V24Y T20S mutant, as it was in ZYA-T20S, except to a slightly greater extent (Fig. 2D). This suggests α K66, by itself, may play mechanistic roles in gate opening, in addition to its role in PAs binding. Based on the comparison of the WT, ZYA bound and α V24Y T20S structures, we conclude that the α V24Y mutation by itself induces HbYX-dependent conformational changes in the 20S proteasome.




Fig. 1

Overlay of cryo EM maps shows α V24Y T20S has similar conformational changes as ZYA bound T20S and is in the gate-open state

A Overlay of wild-type (WT) T20S (yellow) and α V24Y T20S (red) electron density maps, showing the α and β rings. Whole maps were aligned to one another.

B Overlay of α V24Y T20S (red) and ZYA-T20S (blue) electron density maps, showing the α and β rings. V24Y T20S shows similar but larger conformational changes than does ZYA-T20S.

C Close up of the intersubunit pocket in the WT T20S (yellow) electron density map, highlighted in **A** in the larger neon green box outline.

D, E Same as **C**, but for α V24Y T20S (red) and ZYA-T20S (blue). Arrows point to the density corresponding to α V24Y and Z of ZYA, respectively.

F Close up of the WT T20S (yellow) electron density map, corresponding to the YDR motif, the missing density for tyrosine is highlighted by the neon green dotted circle, which is expected in the closed state. The relative position of the close-up is indicated by the smaller neon blue box outline in **A**.

G, H Same as **F**, but for α V24Y T20S (red) and ZYA-T20S (blue). Neon green dotted circle highlight the YDR tyrosine side chain density, which is expected to be in this position in the open gate conformation.

I Close up of the WT T20S (yellow) electron density map, corresponding to the IT Switch, highlighted by the neon green dotted circle.

J, K Same as **I** but for α V24Y T20S (red) and ZYA-T20S (blue). Changes in the conformation of the IT switch indicates switching from the closed to the open state.




Fig. 2

Atomic model of αV24Y T20S compared to WT and ZYA-T20S shows it has undergone HbYX-dependent conformational changes associated with 20S gate-opening

A Atomic model overlay of WT T20S (yellow), αV24Y T20S (red), and ZYA-T20S (blue) with bound ZYA (gold), aligned by the β rings to show conformational changes in the α subunits. Top half of sideview of 20S is shown. Conformational changes in the α-ring are clearly visible (black arrows). Helix 4, 5, and 6 are labeled with black asterisks outlined in white.

B Top view of **A**. Conformational changes noted by (black arrows).

C Closeup view of one of the α intersubunit pockets as in **B**. αPro17 is shown with a black arrow. Helix 0 is labeled with a black asterisk outlined in white.

D Closeup view of αK66 which is labeled and shown in sticks.

Models in panel b through d are aligned to show intersubunit conformational changes (i.e., aligned by β rings)

Comparison of V24Y T20S to the WT T20S reveals specific intra-subunit conformational changes induced by α V24Y mutation

It is remarkable that a single point mutation in the T20S can mimic a 3-residue motif, let alone the activation capacity of a large complex like PAN or PA26. While the α ring is an allosteric system with each neighboring α subunit affecting the next (e.g., a rigid body rotation of one α subunits is expected to affect its neighbor and so on), it's also likely that HbYX binding or the α V24Y mutation could cause conformational changes in a single α subunit as well, which could trigger allosteric transitions to the open state. Thus, we sought to identify the intra-subunit (within a single subunit) conformational changes that could lead to larger inter-subunit (between subunits) conformational changes (Fig. 3A & B) that perpetuate around the entire α -ring and induce gate-opening (Fig. 2B). The substitution of α V24 for tyrosine presents a much larger and aromatic side chain in the hydrophobic pocket. This tyrosine could impose intra-subunit conformational changes on the surrounding secondary structures. To measure these changes, we aligned a single α subunit from the WT T20S to α V24Y T20S. We found that the distance between α L21 and α A154, residues that surround the hydrophobic pocket, indicate that the pocket is widened by $\sim 1.3\text{\AA}$ due to the α V24Y mutation (Fig. 3C vs. D). Another measurement of this pocket width between T124 and L21 also widens by $\sim 1\text{\AA}$ (Fig. 4G). However, the distance between α V24 or α V24Y to α L81 is consistent (Fig. 3C vs. D), suggesting that the hydrophobic pocket has not enlarged in all directions.

We hypothesize that the widening of the hydrophobic pocket that α V24Y occupies is the causal point of origin triggering the inter-subunit conformational changes that we observed in Fig 2. The HbYX hydrophobic pocket occupied by α V24Y side chain is adjacent to another hydrophobic pocket found on the opposite side of Helix 0 (on the 20S central channel), where the side chain of α I12 (an IT Switch residue) occupies in the open state (Fig. 3E & F). The α V24Y T20S model coincides with the α V24Y T20S map, showing that the IT switch is in the open state (Fig. 1), where the α I12 side chain is oriented to occupy the IT switch hydrophobic pocket adjacent to Helix 0 (Fig. 3E vs. F). Due to the adjacent proximity of both hydrophobic pockets (on either side of and below Helix 0), a conformational change in the HbYX hydrophobic pocket would also affect the IT switch's hydrophobic pocket since these pockets share common space. It is highly plausible that the change in the HbYX hydrophobic pocket, resulting from the larger side chain of tyrosine (compared to valine), could facilitate the switch between α T13 and α I12 in the IT switch hydrophobic pocket (Fig. 3E vs. F). The α V24Y mutation increases the hydrophobicity and size of the pocket, likely enhancing the binding affinity for the larger hydrophobic side chain of isoleucine. In conclusion, it appears that occupancy of the HbYX hydrophobic pocket with an aromatic ring of tyrosine allosterically affects the IT switch hydrophobic pocket, as intra-subunit conformational changes allow the IT switch to reconfigure to an open state.

α Proline17 (P17) displacement is one of the first identified inter-subunit conformational changes associated with triggering gate opening, specifically by the 11S family members(30, 33, 37). Interestingly, when the IT switch switches to an open configuration, it generates "slack" between the IT switch and P17(31) (Fig. S4A & S4B). In α V24Y T20S, this "slack" appears to allow for an increase in the intra-subunit distance between Helix 0 and α P17 (Fig. S4B). In addition, we observe an intra-subunit displacement (not rotation based) of P17 by $1.5\text{\AA}$ (Fig. 3G) consistent with the additional slack from the switching of the IT switch. In addition, intra-subunit analysis shows α P17 is displaced by $1\text{\AA}$ away from its neighbor's Helix 0 (Fig. S4C); however, this distance between P17 and the neighboring Helix 0 does not change globally (Fig. S4A & S4B) due to the rigid body rotation of the α subunits which maintain packing between the P17 loop and the neighboring Helix 0. In short, the conformational change of the IT switch to the open state appears to provide flexibility for the Pro17 loop to absorb, or perhaps allow, the rigid body




Fig. 3

α V24Y mutation induces intrasubunit conformational changes in the HbYX hydrophobic pocket, IT switch pocket, and P17 loop

A Overlay of WT T20S (yellow) and α V24Y T20S (red) α subunit models, aligned by the β rings.

B Same as **A** except the models were aligned according to individual α subunit.

C Close up of the intersubunit pocket in α V24Y T20S (red), showing distances between the α -carbon selected residues (labeled and shown in sticks) to demonstrate conformational changes when compared to the WT T20S (yellow) model in **D**. Models are aligned by individual α subunit.

D Same as **C** except it is the WT T20S (yellow) model.

E Close up on the residues of the IT Switch (sticks) and α V24Y (sticks), showing the distance between the α -carbon of α V24Y and α A154 (sticks) to demonstrate conformational changes in the open state, when compared to the WT T20S (yellow) model in **F**. Models are aligned by individual α subunit.

F Same as **E** except it is the WT T20S (yellow) model.

G Close up of Helix 0 and selected surrounding residues (shown in sticks) in an overlay of WT T20S (yellow) and α V24Y T20S (red) α subunits, aligned by individual α subunit. Distances are measured between α -carbons and colored according to the model it corresponds to.

H Close up of α V24Y (shown in sticks) in the α V24Y T20S model, demonstrating its side chain's polar interaction (yellow dotted line) with α D152 (shown in sticks).




Fig. 4

α V24Y mutation and ZYA binding both induce similar intrasubunit conformational changes

- A** Same as **3A** except for α V24Y T20S (red) and ZYA-T20S (blue). Models are aligned by the β ring.
- B** Same as **3B** except for α V24Y T20S (red) and ZYA-T20S (blue). Models are aligned by individual α subunit.
- C** Same as **B** except showing a closeup of the Helix 4, 5, and 6 bundle.
- D** α V24Y T20S and ZYA-T20S are aligned by β subunits and α V24Y side chain and carboxybenzyl group (Z) of ZYA are both shown in space-fill model to appreciate the level of spatial overlap between the α V24Y tyrosine ring (red) and the Z ring (gold) of ZYA, demonstrating they each occupy the same HbYX hydrophobic pocket in the T20S.
- E** Close up of the α V24Y T20S (red) α intersubunit pocket showing measurements between the α V24Y tyrosine and labeled α carbons (shown in sticks). Comparison against **F** indirectly demonstrates the position of α V24Y's side chain relative to the carboxybenzyl group of ZYA in the ZYA-T20S model.
- F** Measurements of distances between ZYA's carboxybenzyl group (gold) and the α -carbon of surrounding residues (shown in sticks) in the ZYA-T20S model (blue).
- G** The HbYX hydrophobic pocket measurement: α T124 to α L21 (main chain shown in sticks) distance in WT T20S (yellow), α V24Y T20S (red), and ZYA-T20S (blue).
- H** Same as **3E & F**, except with alignment of ZYA-T20S (blue) and α V24Y T20S (red) α subunit models. Distances noted are colored according to the models they correspond to representing intrasubunit differences.

rotation associated with HbYX induced gate-opening. Moreover, our structure suggests the widening of the hydrophobic pocket between α L21 and α A154 may also contribute to the shift in α P17, as α L21 is connected on the other side of the α P17 loop (**Fig. 3G** [↗](#)).

We observed that the side chain of α E25 is involved in an intrahelical hydrogen(H)-bond with α R28 in the WT structure (**Fig. S5A** [↗](#) & **S5B** [↗](#)), which is displaced by the α V24Y mutation and instead H-bonds with α R20 (**Fig. S5C** [↗](#) & **S5D** [↗](#)). The displacement slightly increases the distance between α E25 and α R20 (Fig. SB vs. **S5D** [↗](#)) further providing “slack” for the α P17 movement. We previously showed the mutation of α E25 to alanine, activated gate opening and prevented PAs from being able to induce gate opening in the T20S(**31** [↗](#)). Therefore, biochemical evidence also supports an important role for this α E25 rearrangement in destabilizing the closed state and perhaps stabilizing the open state.

We also considered other mechanisms by which the α V24Y mutation might influence gate opening such as its side chain’s hydroxyl polar interactions. We find that α Y24 side chain interacted with the side chain of α D152 (**Fig. 3H** [↗](#)); however, no significant conformational changes were observed along the β -sheet that α D152 is positioned on. Thus, it is not apparent how the polar interaction between α Y24 and α D152 might be associated with gate opening, other than the fact that the H-bond may help orient this tyrosine rotamer.

Comparison of activated T20S forms: α V24Y T20S and ZYA-T20S

The cryo-EM structure presented here (**Fig. 1** [↗](#) & **2** [↗](#)) and our previously published biochemical data(**31** [↗](#)) clearly demonstrate the open state of the α V24Y-T20S mutant. The substitution of α V24 for tyrosine was engineered to mimic how the “Hb” group of the HbYX motif binds to its hydrophobic pocket, but does it? As previously mentioned, the global conformational changes in the α V24Y 20S were similar to a HbYX-bound 20S (i.e., minimal conformational changes in the β subunits, rigid body rotation of α subunits pivoting on Helix 2) (**Fig. 1B** [↗](#) & **2** [↗](#)), suggesting that the mechanism of gate-opening by α V24Y mutation has mechanistic overlap with the HbYX mechanism (e.g. ZYA-T20S). Therefore, we sought to identify, at the molecular level, how these two open T20S models compare and identify any intra-subunit conformational changes that can be specifically attributed to the occupancy of the hydrophobic pocket by the Hb component of the HbYX motif, without influence from the Y and X residues of the HbYX motif. The α V24Y T20S model allows for analysis of the isolated effects of Hb occupancy in the hydrophobic pocket.

Comparison of α V24Y T20S against ZYA-T20S reveals that the intra-subunit conformational changes induced by α V24Y were highly similar, as there were few differences between the α subunits when aligned to one another (**Fig. 4A**, **B** [↗](#) & **C** [↗](#)). Comparison of the hydrophobic pocket between the two open models, ZYA-T20S and α V24Y T20S, shows that the Z group of ZYA and the α V24Y tyrosine both clearly bind to the HbYX hydrophobic pocket and overlap in space, though they bind in different orientations (**Fig. 4D** [↗](#)). In addition, the measurements of α V24Y’s side chain or Z of ZYA to surrounding residues within each individual subunit provide the most relevant visual approximation and overlap comparison of the “Z” and “ α V24Y” aromatic group positions in the HbYX hydrophobic pocket (**Fig. 4E** [↗](#) & **F** [↗](#)). For comparison, the distance between α V24 or α V24Y to α L81 marginally differs by ~ 0.3 Å while the distance between α L21 and α L81 is comparable in both models (**Fig. S6A** [↗](#) & **S6B** [↗](#)). The measurement of the hydrophobic pocket between α L21 and α T124 is widened in the α V24Y T20S (~ 0.7 Å; **Fig. 4G** [↗](#)). This pocket measurement also enlarges in the ZYA T20S structure compared to WT, though to a lesser extent (**Fig. 4G** [↗](#)), consistent with the observed extent of changes in these models (**Fig. 2** [↗](#)). However, the largest difference in this comparison between ZYA bound and α V24Y is in the measurement between α L21 and α A154, as the V24Y tyrosine is better positioned to influence this specific measurement. (**Fig. 4H** [↗](#)). To validate this important conformational change in the maps raw data in a statistically significant way we used CCP-EM to generate confidence maps of WT and V24Y structures at an FDR 0.01. These confidence maps clear show conformational differences in A154 in the V24Y structure (**Fig. S3B** [↗](#)). Lastly, the placement of α I12’s side chain in the IT switch

hydrophobic pocket between Helix 0 and Helix 2 protrudes $\sim 0.8\text{\AA}$ deeper in the pocket in $\alpha V24Y$ T20S compared to ZYA-T20S (Fig. 4H), consistent with the larger hydrophobic pocket under Helix 0 in the mutant.

Collectively, evidence suggests that ZYA's mechanism as an activator is due to its hydrophobic group occupying the space between HbYX hydrophobic pocket, similar to $\alpha V24Y$, but ZYA's mechanism also relies on others parts of the dipeptide as modifications to tyrosine affected the small molecule's ability to activate (38). In combination with the evidence shown here, we conclude that the hydrophobic group of ZYA is likely the primarily feature allosterically affecting gate-opening but the insertion of this Hb group into the hydrophobic pocket also relies on interactions of the Y and X residues with the inter-subunit pocket to provide sufficient binding affinity. As for $\alpha V24Y$ T20S, the hydrophobic component (tyrosine's side chain) is covalently held in place in the hydrophobic pocket and thus does not require the "YX" interactions and therefore, sufficiently supports stable gate opening through a similar mechanism to ZYA.

$\alpha K66$ supports gate opening through a previously unknown mechanism

Many studies in the past have highlighted the importance of $\alpha K66$, found in the back loop of the inter-subunit pocket (Fig. 2D & 5A), and our $\alpha V24Y$ T20S structure shows that $\alpha K66$ is reorganized similar to when ZYA binds. These observations inspired us to elucidate how intra-subunit conformational changes in $\alpha K66$ may be affected by $\alpha V24Y$. $\alpha K66$ is well known to form hydrogen or salt bonds with the C-termini of PAs and is required for PA binding to the 20S (30), but how is $\alpha K66$ affected when there is no bound PA? The back loop of the inter-subunit pocket is naturally a more flexible loop but in $\alpha V24Y$ T20S, this loop does have significant intra-subunit conformational changes (Fig. 5A), with this loop being displaced by $\sim 3\text{\AA}$ (inter-subunit measurements are similar). Additionally, we noticed $\alpha K66$ moved $\sim 1\text{\AA}$ towards $\alpha T78$ and its side chain is oriented to be in closer proximity to hydrogen bond with both $\alpha T78$ and $\alpha E211$ (Fig. 5A & B). Such interactions are not observed in the WT T20S (Fig. 5C) but are seen in the ZYA-T20S (Fig. 5D). $\alpha K66A$ mutants do not exhibit activation by HbYX-dependent or HbYX-independent activators leading prior studies to conclude that $\alpha K66$ was important for binding of proteasome activators (30, 34). However, the repositioning of $\alpha K66$ in the $\alpha V24Y$ T20S structure, which does not have a proteasome activator bound, indicates that $\alpha K66$ may also be involved in stabilizing the open state on its own. To test this hypothesis, we generated a double mutant proteasome, $\alpha V24Y$ - $\alpha K66A$ T20S. We isolated this mutant and measured fluorogenic peptide hydrolysis activity (LFP). We found that $\alpha V24Y$ - $\alpha K66A$ T20S has similar activity to WT and is not hyperactive like $\alpha V24Y$ T20S (Fig. 5E). Thus, $\alpha K66$ is essential for $\alpha V24Y$ induced gate-opening. These enzymatic results support the conclusion that $\alpha K66$ does not only interact with proteasome activators to support gate opening but is also critically involved in stabilizing the open state of the α subunit gate by forming new interactions with $\alpha T78$ and $\alpha E211$ in the open state.

$\alpha V24Y$ mutation affected the hydrophilicity of the inter-subunit pocket

Our previous study of ZYA-T20S suggested that the HbYX mechanism of activation involved the rearrangement of water molecules, so we sought to determine if the same applies in $\alpha V24Y$ T20S. The mutation essentially introduces a larger hydrophobic side chain to the inter-subunit pocket and therefore, we expect rearrangement of water molecules. Our map and model suggest that the $\alpha V24Y$ mutation either excluded or rearranged water molecules found in the WT T20S inter-subunit pocket (Fig. 7A vs. B). Additionally, none of the water molecules placement in $\alpha V24Y$ T20S coincided with the exact placement and interactions as in ZYA-T20S (Fig. 6C), suggesting that waters involved in ZYA's binding to the inter-subunit pocket may not be relevant in $\alpha V24Y$ T20S. This is unsurprising given that ZYA is a molecule that occupies a good portion of the inter-




Fig. 5

αK66 plays a key role in stabilizing the open state of the T20S, even in the absence of proteasome activator binding

A Overlay of WT T20S (yellow) and αV24Y T20S (red), aligned by individual α subunits, showing a close-up of the αK66 (shown in sticks) back loop. Distance measured demonstrates the extent of intrasubunit conformational changes in the “back loop” and K66 α carbon.

B Close up of αV24Y T20S (red) model, showing the polar interactions (yellow dotted lines) between αK66 and surrounding residues (shown in sticks). Distances are measured between the exact atoms involved in the polar interactions.

C Same as **B** except it is the WT T20S (yellow) model.

D Same as **B** except it is the ZYA-T20S (blue) model and ZYA (gold).

E WT T20S and mutants were measured for activity based on LFP degradation rate (rfu/min), normalized to the WT T20S as a control. Data (means) are representative of three or more independent experiments, each performed in triplicate. Error bars represent ± standard deviation.

subunit pocket, relative to the small change caused by the α V24Y mutation. Interestingly the inter-subunit pockets of α V24Y appears to be more dehydrated relative to the WT, but structures with equivalent resolutions would be necessary to confirm. The results collectively suggest that while water interactions in the inter-subunit pockets change substantially in the open versus closed states (at these resolutions), we do not observe specific waters shared between these two open 20S models that would suggest waters being critically involved in stabilization of the open gate by these two different activation mechanisms (Fig. 6 [↗](#)).

α V24F T20S is not activated similarly to α V24Y T20S

Similar to α V24Y T20S, the α V24F T20S mutant showed increased basal activity compared to WT T20S biochemically([31](#) [↗](#)). Therefore, we sought to elucidate whether the mechanism of activation would be similar if the substitution was to a phenylalanine as opposed to a tyrosine. Comparison of the α V24F T20S map (Fig. 7A [↗](#), Fig. S7 [↗](#) & S8 [↗](#)) to α V24Y T20S and WT T20S maps and models reveal a distinct lack of densities in the α subunits when α V24 is mutated to phenylalanine (Fig. 7B [↗](#) & C [↗](#)). The lack of densities is most apparent from the N-termini and through Helix 0, though the map was also lacking densities in other selected regions of the α subunit. Globally analyzing the α V24F T20S map suggests that the α V24F mutation affected the proper folding of secondary structures surrounding the α V24F mutation (e.g. on Helix 0), but interestingly, this lack of Helix 0 folding did not affect the assembly of the proteasome itself (Fig. 7B [↗](#) & C [↗](#)). The difference between maps indicates that the β subunits are minimally affected in the assembly of the proteasome and that the lack of densities affected the α subunit throughout, though they were most apparent around the gating regions (Fig. 7D [↗](#) & E [↗](#)). The lack of densities in α V24F T20S map suggests that substrates could enter the mutant proteasome more easily through the larger opening and therefore, biochemically exhibited kinetics comparable to or even exceeding the activated proteasome([31](#) [↗](#)).

Discussion

Proteasome impairment and reduced proteostasis are common observations in clinical cases and animal models of neurodegenerative diseases such as Alzheimer's and Parkinson's, which has inspired the search for small molecule proteasome activators for years. Yet, there is a lack of robust small molecule proteasome activators to date, in part due to a lack of understanding in proteasome activation mechanisms. The α inter-subunit pockets have long been known to be the binding site of PA complexes but the exact molecular interactions between the inter-subunit pockets and the PAs that cause gate-opening have been elusive, preventing efficacious design of proteasome-activating small molecules. While no FDA-approved *in vivo* proteasome-activating small molecule exists as a tool for proof of concept, our previous work([18](#) [↗](#)) to express constitutively active 20S proteasomes in *C. elegans* indicated that the mutant proteasomes supported an increased resistance to proteotoxicities and increased lifespan, supporting the rationale for this study. Here, we present a single specific binding site within the inter-subunit pocket that when occupied, is sufficient to induce gate opening and therefore, a plausible and well-defined pharmaceutical target for proteasome activators.

To identify the fundamental molecular interaction(s) required for gate-opening, we focused on the archaeal proteasome as it is homoheptameric, which is particularly an advantage when desiring a high resolution cryo-EM structure to map detailed interactions. Moreover, the archaeal proteasome still carries the conserved structures involved in gating, and both human and archaeal 20S can be opened by the same HbYX peptides (e.g. ZYA), implying that the findings of this study could be applied to the human system with some caveats discussed below. We recently demonstrated that the α V24Y mutant archaeal proteasomes exhibited upregulated activity([31](#) [↗](#)). Our α V24Y T20S structure here reveals inter-subunit conformational changes essentially identical




Fig. 6

The intersubunit pocket of the α V24Y T20S is less populated with waters, relative to ZYA T20S and WT T20S

A Intersubunit pocket of WT T20S model (yellow) showing waters (spheres) and their interactions (yellow dotted lines) with the residues (shown in sticks, listed in table) in the intersubunit pocket. Spheres are colored individually and match the colors on the table (right) to identify residues it interacts with.

B, C Same as **A** except α V24Y T20S model (red) and ZYA-T20S model (blue).




Fig. 7

A large portion of the N-terminal domain of the α V24F T20S mutant is unstructured

A Angled view of α V24F T20S 2.2Å electron density map (green), showing the α and β rings.

B Overlay of α V24F T20S map with α V24Y T20S map (red; transparent surface; left) and WT T20S map (yellow; transparent surface; right).

C Same as **B** except α V24Y T20S model (red, cartoon, left) and WT T20S model (yellow; cartoon; right).

E Difference map generated by subtracting α V24F T20S map from α V24Y T20S map, using ChimeraX.

F Same as **E** except WT T20S map instead of α V24Y T20S map.

to a HbYX-bound archaeal proteasome (ZYA-T20S)([31](#)), demonstrating that it is in an open state, allowing us to further elucidate how this single residue mutation opens the gate, and by extension, better understand the mechanism of gate opening by the HbYX motif itself.

Collectively, our findings in α V24Y T20S focus on a mechanism of gate-opening originating at the occupancy of a hydrophobic pocket between α L21 and α T124. The α V24Y mutation places an aromatic side chain in between α L21 and α T124 (as well α L21 and α A154), similar to the position of the benzyl group (Z) of ZYA in the ZYA-T20S structure. The occupancy of this hydrophobic pocket by α V24Y expands the adjacent pocket where α I12 of the IT Switch is previously found to bind in open gate proteasome structures. α L21, displaced by α V24Y's larger side chain compared to α V24 (**Fig. 3C**), is connected to the α P17 loop, which has been commonly shown to be displaced in open gate proteasome structures. The results here also suggest that this mechanism applies to the ZYA-T20S (**Fig. 4G**), though it may not be the only mechanism supporting ZYA's gate-opening as ZYA is a dipeptide as opposed to a single residue mutation (α V24Y) that occupies the hydrophobic pocket. For example, the YA residues bridge the subunit from α K66 to the α P17 loop, which also likely stabilizes the open state([31](#)). The interaction of ZYA's carboxy group with α K66 may also help stabilize the open-state conformation of α K66 (**Fig. 5B**), which we showed is essential for gate-opening here (**Fig. 5E**). The conformational changes between the two models are consistent, although α V24Y T20S exhibits larger conformational changes likely due cryo-EM averaging effects in the lower populated ligand-bound states (ZYA) compared to the V24Y mutant. ZYA as a dipeptide has the ability to and perhaps, even rely on interacting with other parts of the inter-subunit pocket to supports its binding and orient the Z group in the hydrophobic pocket. Results from both α V24Y T20S and ZYA-T20S both suggest that occupying the hydrophobic pocket is a key to gate opening in the archaeal proteasome, and potentially the mammalian proteasome if placed in the right pocket(s).

The two key residues that “sandwiches” α V24Y's side chain and forms the hydrophobic pocket, α L21 and α A154, are conserved in the human 20S α inter-subunit pockets (**Fig. S9**), indicating that a similar mechanism of occupying this hydrophobic pocket is used for gate-opening in the human system. Five out of seven H20S α subunits carry leucine in a similar position to α L21 in the T20S. The remaining two subunits carry an isoleucine (α 6: PSMA1) and valine (α 7: PSMA3), which are hydrophobic like leucine. Additionally, while α A154 is only conserved as alanine in α 1: PSMA6, the rest of the subunits carry a methionine in this position, which is also hydrophobic. Previous analysis([31](#)) also indicated that the H20S has analogs of the IT switch in selected α subunits, which suggests that targeting this hydrophobic pocket at selected α subunits with the conserved IT switch would have a greater effect on gate-opening than others. Any application of this mechanism in the human system is still complicated and should also consider the prior prediction that multiple inter-subunit pockets likely need to be occupied by the HbYX motifs for gate-opening([38](#)). The conservation of activation motif, residues, and conformational changes between archaeal and human proteasomes for gate opening demonstrates potential for the findings in this study to translate to activating the human 20S.

Interestingly, our α V24F T20S structure indicates that the position of this residue in the protein chain plays a role in α subunit folding. Our α V24F T20S cryo-EM map reveals that the α subunits are partially assembled, with the majority of regions proximal to the gating residues lacking densities, suggesting disorder. Yet, the β subunits appear functional, both structurally and biochemically([31](#)), which suggests that the proteasome does not require perfectly formed α subunits for complete proteasome assembly. We notice that the hydroxyl group of α V24Y's side chain is involved in polar interactions; however, we do not presume this to be involved in α subunit assembly as valine lacks the side chain to interact similarly. Rather, we speculate that α V24F may introduce too large of a hydrophobic group that, without a hydroxyl to interact with α D152 (**Fig. 3H**) prevents proper folding interactions.

Collectively, our analysis indicates that the HbYX hydrophobic pocket expands due to α V24Y's more bulky side chain, which likely sets off a chain of events starting with the IT switch, which are modeled and schematized in **Figure 8**. The wider and now more hydrophobic IT switch pocket likely promotes IT switch to switch to the open state, providing flexibility for the α P17 loop intra-subunit motions, or "slack". In addition, the widening of the hydrophobic pocket and rearrangement of the α E25 H-bonding pattern also appears to allow more flexibility for α P17 loop motions. Combined, these intra-subunit conformational changes collectively elongate the α P17 loop, increasing its flexibility. This enhanced flexibility is crucial because the α P17 loop interfaces with Helix 0 of neighboring α subunits. The loop's increased plasticity accommodates rigid-body rotations against the adjacent α subunits, which would otherwise be constrained. As a result, these rotational movements can propagate allosterically around the proteasome α ring. Though this causal chain of events is supported by these structures, the timing and order of these events cannot be elucidated with these static structures. The structures here suggest a plausible molecular mechanism of how a single residue mutation in the HbYX hydrophobic pocket can induce 20S gate opening (**Fig. 8**).

Our findings in this study also highlight and revise an understanding of the α K66's role in proteasomal gate-opening. Previous findings have consistently alluded to its critical role for the association of HbYX-dependent and HbYX-independent PAs, as α K66 was shown to interact with and orient the C-terminal tails of PAs for gate-opening (30, 31, 35, 37, 39). To our surprise, the α V24Y T20S structure demonstrates that α K66's side chain conformation changes, even though it does not interact with a PA or HbYX mimetic such as ZYA. Our double mutant α V24Y&K66A T20S's basal activity supports the necessity of the α K66 conformational change observed in the α V24Y model and indicates the involvement of α K66 in stabilizing the open-gate conformation through a previously unreported mechanism. The conformational change in α K66 is supported by changes in H-bonding interactions between α K66 and surrounding residues, which are absent in the closed structure (WT T20S). These findings therefore demonstrate that α K66's conformational changes support other conformational changes such as the conformation of the back loop, which it is connected to, and has been shown to be involved in HbYX-dependent gate opening (31). Essentially, α K66 is deduced to contribute more than interacting with activators, and also appears to be necessary to stabilize the open state of the 20S.

Our prior study showed the possible roles waters could play in proteasome activation and gating regulation. Our ZYA-T20S structure indicated that the hydroxyl of ZYA's tyrosine side chain interacted with a water molecule that further hydrogen bonded with α L21 and α E25. To elucidate the relevance and importance of this water interaction, we mutated α E25 (31). α E25A proteasome mutants exhibited an upregulated basal activity that was not further activated by PAN or PA26, suggesting it plays an important role in gating. To elucidate whether a similar mechanism occurs in α V24Y T20S open gate conformation, we sought to identify where waters might be found within the inter-subunit pocket. Broadly, we observed significantly fewer waters in the inter-subunit pockets of α V24Y T20S than we did in the WT T20S and ZYA-T20S. We note that α V24Y T20S's resolution is not as high as ZYA-T20S but is reasonable for predicting water coordinates. Because no common water was found in the two different open states, the necessity and role of specific water molecule interactions in gate-opening could not be elucidated from these structures.

Ultimately, the findings described indicate that the occupancy of the hydrophobic pocket within the α inter-subunit pocket is sufficient to induce gate-opening in archaeal proteasomes. The application of this finding in the mammalian system relies on further identification of the critical inter-subunit pockets that need to be occupied by the HbYX motif in that system to stabilize gate-opening. Our prior studies involving ZYA have indicated a minimum of two HbYX motifs binding for gate-opening in the mammalian system (38). However, the mammalian 20S has 7 different pockets that can be occupied in different ways by 6 different Rpt1-6 C-termini, so the specific inter-subunit pockets that need C-termini/HbYX motifs bound to activate gate opening has not been worked out. All cryo-EM studies to date agree that occupancy of five Rpt C-termini is sufficient for




Fig. 8

Model for induction of gate opening by binding of the Hb group binding to the HbYX hydrophobic site in the 20S intersubunit pockets

Binding of the Hb group of the HbYX motif, or tyrosine of $\alpha V24Y$, is expected to initiate an allosteric chain of events that leads to 20S gate opening. Hb group binding causes:

- 1) Increase the size of the Hb pocket
- 2) Affects IT switch Hb pocket promoting IT switching to open state
- 3) IT switching lengthens P17 loop (increases loop slack)
- 4) E25 H-bond rearrangement further lengthens P17 loop
- 5) Combine effects on P17 loop slack promotes rigid body rotations against the neighboring Helix 0.
- 6) Rigid body rotations effect back loop (not shown), which repositions K66 to further stabilize open state.

gate opening, but which of those are required for gate opening is not known. Further progress on the HbYX mechanism based on the results thus far would be to identify the specific inter-subunit pockets that need to be occupied by the HbYX motif to induce gate-opening in the mammalian proteasomes.

The work presented here is conducted with the intentions of understanding proteasome activation mechanisms and developing the molecular framework for restoring or activating proteasome activity to combat the impairment consistently observed in neurodegenerative diseases, which is barred by our limited understanding of proteasome regulation. Here, we advance the field's understanding of proteasomal gate-opening by extending beyond the vague idea that HbYX motif binding induces gate opening, to pinpointing a sole molecular interaction that significantly and sufficiently opens the proteasome's gate. Our findings inform proteasome activating drug discovery efforts to focus on small molecules that occupies the HbYX hydrophobic pocket.

Materials and methods

Proteasome purifications

T. acidophilum wild-type 20S, α V24Y 20S, or α V24F 20S proteasomes were similarly purified as described(40), using the 8XHis tags on the C-terminus of the β subunits. All T20S mutants were generated by overlapping PCR site-directed mutagenesis.

Proteasome activity assays

Fluorogenic substrate peptide, LFP (Mca-AKVYPYPME-Dpa(Dnp)-amide), were synthesized by from EZBiolabs. LFP was dissolved in DMSO and incubated with proteasomes at 3 μ M final concentration. The final concentration of DMSO in activity assays was 1%. Protein concentrations were determined by Bradford assay (Thermo Scientific). 0.25 μ g archaeal proteasomes and LFP peptides were added to 25 μ L Tris-based reaction buffer at 45°C. Assays were performed for 30mins to an hour and analyzed using BioTek Gen5 Data Analysis software. Activity was measured as relative fluorescence units/minute (rfu/min), generating a curve which was used to calculate the initial velocity, according to the slope of the curves. "Fold Activation" was calculated by dividing the average initial velocity of mutant proteasomes against the average initial velocity of wild type proteasomes.

Cryo-EM Sample Preparation and Data Collection

Copper Quantifoil R 1.2/1.3 300 mesh (EMS) grids were cleaned using a PELCO easiGlow Glow Discharge cleaning system. A volume of 3 μ L of 0.75mg/mL α V24Y-T20S or α V24F T20S (suspended in 50mM Tris pH 7.4, 150mM NaCl) sample was placed onto a grid, and then flash frozen in liquid ethane using a manual plunge freeze apparatus. Data collection was done using a Titan Krios transmission electron microscope (Thermo Fisher) operated at 300kW and a magnification of x81,000, which resulted in 0.503Å/px. Images were collected using a Falcon III EC direct electron detector camera equipped with a K3/GIF operating in counting and super resolution modes. Electron dose per pixel of 50 e-/Å² was saved as 40 frame movies within a target defocus range of -2.5 to -1.25. All the data was collected using cryoSPARC software (Structura Biotechnology Inc) (41).

Cryo-EM Single Particle Analysis

Cryo-EM images of the α V24Y T20S and α V24F T20S proteasome were analyzed using cryoSPARC. Schematic for cryo-EM single-particle data processing available in supplement. α V24Y T20S: From 3663 movies collected, we picked 396542 particles after two rounds of 2D Classification to generate

an Ab-initio model, which was used for homogeneous refinement (using D7 symmetry). Particle processing scheme and validations are shown in **Fig. S1** & **S2**.

α V24F T20S: From 4920 movies collected, we picked 542722 after two rounds of 2D Classification to generate an Ab-initio model, which was used for homogenous refinement (using D7 symmetry). Particle processing scheme and validations are shown in **Fig. S6** & **S7**.

All representations of the T20S proteasome complex were created using PyMOL 2.5.4 and UCSF ChimeraX v1.7.1(42, 43).

Confidence map generation

Small conformational changes (e.g. $\sim 1\text{\AA}$) with larger allosteric effects altering 20S gating and enzymatic activity are common features in the 20S proteasome, which has been published extensively. However, to more rigorously demonstrate that these changes are statistically significant at the map data level, we used CPP-EM to generate confidence maps of our WT and V24Y structures (**Fig. S3B**). Confidence maps statistically interpret cryo-EM densities by testing each voxel against background noise and controlling the false discovery rate (FDR). We thresholded our maps of WT and V24Y at an FDR of 0.01 to highlight regions with statistically significant density. We then overlaid these maps in Chimera to show they clearly show conformational changes even at an FDR of 0.01, especially at the specific regions the reviewer was concerned about.

Atomic model building

The atomic models were built using PDB: 8F7K as a template, rigid body fitting into the electron density map using PHENIX 1.19.2-4158(44–46). The docked models were subjected to a cycle of morphing and simulated annealing, five real-space refinement macrocycles with atomic displacement parameters, secondary structure restraints, and local grid searched in PHENIX. Consequently, the models were refined by oscillating between manual real-space refinement in WinCoot(47) 0.9.8.1 EL and real-space refinement in PHENIX (five macrocycles, without morphing and simulated annealing). Waters were added to models using PHENIX Douse. The final model of α V24Y T20S was deposited at wwPDB: 9BUZ.

Statistical analysis

Data were analyzed in Graph Pad or excel using an unpaired Student's t-test (Prism). For all statistical analyses, a value of $p < 0.05$ was considered significant.

Data and materials availability

Cryo-EM maps are deposited in the Electron Microscopy Data Bank (EMDB) under accession codes, EMD-44914 (α V24F T20S) and coordinates are available from the RCSB Protein Data Bank under accession codes, 9BUZ (α V24Y T20S), 8F7K (ZYA-T20S), 8F6A (WT T20S). The authors declare that data supporting the findings of this study are available within the paper and its supplementary information files and are available from the corresponding author upon request.

Acknowledgements

We thank the members of the Smith Lab for the helpful and valuable discussions, especially Giovanni Howells for his meticulous review of this manuscript. We also thank Thomas (Tom) C. Terwilliger, PhD at the Los Alamos National Laboratory for his gracious guidance and assistance in




Fig. S1

α V24Y T20S Processing Scheme

A Cryo-EM workflow for alphaV24Y T20S. All steps were performed in Cryosparc.

B Final 2D class averages after 2 rounds of 2D Classification shown with a mixture of top and side views.

C Homogenous refinement of class of particles that showed the open gate conformation.




Fig. S2

αV24Y T20S Validation

A Micrograph representative showing αV24Y T20S. Scale bar represents 5nm.

B. Standard FSC-0.143 graph showing corrected 2.38Å resolution.

C View of particle angle distribution showing a mixture of side and top view particles.

D Particles from selected ab-initio model were refined with homogenous refinement and D7 symmetry applied. D7 symmetry was confirmed as appropriate by homogenous refinement with C1 symmetry; the resulting C1 map was copied then flipped 180° around the 2-fold axis then rotated 180° about the 7-fold axis. The flipped and rotated map was essentially identical to the original, confirming D7 symmetry of the complex. Final 2.38Å D7 map was colored in Chimera to show range of resolution of 3D reconstructed map.




Fig. S3

α V24Y T20S differ from WT T20S

A Close up of α V24Y T20S sharpened map and model, demonstrating quality of map and model fit in the shown region.

B. Overlay of CCP-EM generated confidence maps for α V24Y T20S (yellow) and WT T20S (cyan), indicating statistically significant conformational changes. Maps were set at a threshold of 0.01 FDR. View is similar to **Figure 4H** showing clear conformational changes in A154 relative to helix 0. Arrows point out the distance differences between the same two residues in the WT or the V24Y maps. Gaussian smoothing was applied to both maps.





Fig. S4

Additional measurements in the intersubunit pocket in V24Y T20S

Complementary to **Figure 3**. **A** Measurement between P17 and neighboring Helix 0 for β ring-aligned WT and α V24Y T20S structures representing absence of intersubunit conformational differences in this distance.

B Measurement between P17 and its own Helix 0 (E25) show intrasubunit changes to this distance.

C Close up of intrasubunit conformational change in the IT switch and α P17 loop providing flexibility to allow for intersubunit rigid body rotation. Distance shown are intrasubunit conformational change in the α V24Y T20S model and do not represent the global conformational changes, as the P17 packing against the neighboring Helix 0 is similar in open and closed states (SFig. 3a)



Sig. S5

V24Y mutation and ZYA binding both alter E25's H-bonding partner

E25 hydrogen bonding is affected by hydrophobic group occupancy in the HbYX hydrophobic pocket as it is positioned at the top of this pocket.

A E25 is H-bonded to R28 in the WT T20S.

B Distance between E25 and R20 in the WT T20S.

C E25 H-bonds with R20 in the V24Y T20S.

D Distance between E25 and R20 in the V24Y T20S

E E25 H-bonds with R20 in ZYA-T20S structure.

F Distance between E25 and R20 in the ZYA-T20S.




Fig. S6

Comparison of HbYX hydrophobic pocket in V24Y T20S and ZYA-T20S structures

A Distances between the α -carbon of selected residues (shown in sticks) to be compared against **B** to demonstrate the similarities and minimal level of intrasubunit conformational changes between these structures. **B** Same as **A** except the ZYA-T20S model (blue) was used.




Fig. S7

α V24F T20S Processing Scheme

A Cryo-EM workflow for α V24F T20S. All steps were performed in Cryosparc.

B Final 2D class averages after 2 rounds of 2D Classification shown with a mixture of top and side views.

C Homogenous refinement of class of particles that showed the open gate conformation.




Fig. S8

αV24Y T20S Validation

A Micrograph representative showing αV24F T20S. Scale bar represents 5nm.

B Standard FSC-0.143 graph showing corrected 2.20Å resolution.

C View of particle angle distribution showing a mixture of side and top view particles.

D Particles from selected ab-initio model were refined with homogenous refinement and D7 symmetry applied (D7 symmetry was appropriately confirmed as discussed in [SFig. 2d](#)). Final 2.20Å D7 map was colored in Chimera to show range of resolution of 3D reconstructed map.




Fig. S9

αV24Y T20S Validation

A Alignment showing sequences in the first 30 positions of the α subunits of T20S and H20S (listed according to gene names). Alignment was performed using ClustalΩ. Conservation and figure was generated using Jalview 2.11.2.5. Alignments are colored using the Clustal color scheme and color intensity indicates conservation (darkest: highly conserved). The bar graph in yellow indicates the conservation score, accounting for size, charge, hydrophobicity, and polarity. Red arrow points to αL21 and the corresponding residues in the alignment.

B Same as **A** except the alignment shows sequences in the positions 145 to 176 of the α subunits of T20S and H20S (listed according to gene names). The red arrow points to αA154 and the corresponding residues in the alignment.

using Phenix. Transmission electron micrographs were recorded at the University of Virginia Molecular Electron Microscopy Core facility (RRID:SCR_019031), which is supported in part by the School of Medicine and built with NIH grant G20-RR31199. In addition, the Titan Krios (S10-RR025067), Falcon II/3EC direct detector (S10-OD018149), and K3/GIF (U24-GM116790) were purchased in part or in full using the designated NIH grants. Molecular graphics and analyses performed with UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy, and Infectious Diseases([43](#) [↗](#)).

Additional information

Funding

This work was supported by National Institute of Health grant R01AG064188 (DMS).

Author contributions

JJYC and **MRD** prepared WT T20S, α V24Y T20S, and α V24F T20S for cryo-EM and activity assays. **JJYC** and **DMS** analyzed cryo-EM data, reconstructed the α V24Y T20S and α V24F T20S structures. **JJYC** and **MRD** performed enzymatic assays of the archaeal proteasomes. **JJYC** and **DMS** generated PyMOL figures. Results were analyzed and interpreted by **JJYC** and **DMS**. Manuscript was prepared by **JJYC**, **MRD**, and **DMS**. All authors reviewed the results and approved the final version of this manuscript.

References

1. Cohen-Kaplan V., Livneh I., Avni N., Cohen-Rosenzweig C., Ciechanover A. (2016) **The ubiquitin-proteasome system and autophagy: Coordinated and independent activities** *Int. J. Biochem. Cell Biol* **79**:403–418 [Google Scholar](#)
2. Bialek W., Collawn J. F., Bartoszewski R. (2023) **Ubiquitin-Dependent and Independent Proteasomal Degradation in Host-Pathogen Interactions** *Molecules* **28**:6740 [Google Scholar](#)
3. Demasi M., da Cunha F. M. (2018) **The physiological role of the free 20S proteasome in protein degradation: A critical review** *Biochim. Biophys. Acta BBA - Gen. Subj* **1862**:2948–2954 [Google Scholar](#)
4. Khaminets A., Behl C., Dikic I. (2016) **Ubiquitin-Dependent And Independent Signals In Selective Autophagy** *Trends Cell Biol* **26**:6–16 [Google Scholar](#)
5. Makaros Y., Raiff A., Timms R. T., Wagh A. R., Gueta M. I., Bekturova A., Guez-Haddad J., Brodsky S., Opatowsky Y., Glickman M. H., Elledge S. J., Koren I. (2023) **Ubiquitin-independent proteasomal degradation driven by C-degron pathways** *Mol. Cell* **83**:1921–1935 [Google Scholar](#)
6. Tsuchiya H., Endo A., Saeki Y. (2020) **Multi-Step Ubiquitin Decoding Mechanism for Proteasomal Degradation** *Pharm. Basel Switz* **13** [Google Scholar](#)
7. Hegde A. N. (2017) **Proteolysis, synaptic plasticity and memory** *Neurobiol. Learn. Mem* **138**:98–110 [Google Scholar](#)
8. Park H., Kaang B.-K. (2019) **Balanced actions of protein synthesis and degradation in memory formation** *Learn. Mem* **26**:299–306 [Google Scholar](#)
9. Patrick M. B., Omar N., Werner C. T., Mitra S., Jarome T. J. (2023) **The ubiquitin-proteasome system and learning-dependent synaptic plasticity – A 10 year update** *Neurosci. Biobehav. Rev* **152** [Google Scholar](#)
10. Lip P. Z. Y., Demasi M., Bonatto D. (2017) **The role of the ubiquitin proteasome system in the memory process** *Neurochem. Int* **102**:57–65 [Google Scholar](#)
11. Jarome T. J., Devulapalli R. K. (2018) **The Ubiquitin-Proteasome System and Memory: Moving Beyond Protein Degradation** *The Neuroscientist* **24**:639–651 [Google Scholar](#)
12. Keller J. N., Hanni K. B., Markesbery W. R. (2000) **Impaired Proteasome Function in Alzheimer's Disease** *J. Neurochem* **75**:436–439 [Google Scholar](#)
13. Ciechanover A., Brundin P. (2003) **The ubiquitin proteasome system in neurodegenerative diseases: sometimes the chicken, sometimes the egg** *Neuron* **40**:427–446 [Google Scholar](#)

14. McNaught K. S., Olanow C. W., Halliwell B., Isacson O., Jenner P. (2001) **Failure of the ubiquitin-proteasome system in Parkinson's disease** *Nat. Rev. Neurosci* **2**:589–594 [Google Scholar](#)
15. Ortega Z., Díaz-Hernández M., Lucas J. J. (2007) **Is the ubiquitin-proteasome system impaired in Huntington's disease?** *Cell. Mol. Life Sci. CMLS* **64**:2245–2257 [Google Scholar](#)
16. McKinnon C., Tabrizi S. J. (2014) **The ubiquitin-proteasome system in neurodegeneration** *Antioxid. Redox Signal* **21**:2302–2321 [Google Scholar](#)
17. McNaught K. S. P., Björklund L. M., Belizaire R., Isacson O., Jenner P., Olanow C. W. (2002) **Proteasome inhibition causes nigral degeneration with inclusion bodies in rats** *Neuroreport* **13**:1437–1441 [Google Scholar](#)
18. Anderson R. T., Bradley T. A., Smith D. M. (2022) **Hyperactivation of the proteasome in *Caenorhabditis elegans* protects against proteotoxic stress and extends lifespan** *J. Biol. Chem* **298** [Google Scholar](#)
19. Njomen E., Tepe J. J. (2019) **Proteasome Activation as a New Therapeutic Approach To Target Proteotoxic Disorders** *J. Med. Chem* **62**:6469–6481 [Google Scholar](#)
20. VerPlank J. J. S., Lokireddy S., Zhao J., Goldberg A. L. (2019) **26S Proteasomes are rapidly activated by diverse hormones and physiological states that raise cAMP and cause Rpn6 phosphorylation** *Proc. Natl. Acad. Sci* **116**:4228–4237 [Google Scholar](#)
21. VerPlank J. J. S., Tyrkalska S. D., Fleming A., Rubinsztein D. C., Goldberg A. L. (2020) **cGMP via PKG activates 26S proteasomes and enhances degradation of proteins, including ones that cause neurodegenerative diseases** *Proc. Natl. Acad. Sci* **117**:14220–14230 [Google Scholar](#)
22. Myeku N., Wang H., Figueiredo-Pereira M. E. (2012) **cAMP stimulates the ubiquitin/proteasome pathway in rat spinal cord neurons** *Neurosci. Lett* **527**:126–131 [Google Scholar](#)
23. Wang X., Yen J., Kaiser P., Huang L. (2010) **Regulation of the 26S Proteasome Complex During Oxidative Stress** *Sci. Signal* **3**:ra88–ra88 [Google Scholar](#)
24. Groll M., Bajorek M., Köhler A., Moroder L., Rubin D. M., Huber R., Glickman M. H., Finley D. (2000) **A gated channel into the proteasome core particle** *Nat. Struct. Biol* **7**:1062–1067 [Google Scholar](#)
25. Rabl J., Smith D. M., Yu Y., Chang S.-C., Goldberg A. L., Cheng Y. (2008) **Mechanism of gate opening in the 20S proteasome by the proteasomal ATPases** *Mol. Cell* **30**:360–368 [Google Scholar](#)
26. Ding Z., Xu C., Sahu I., Wang Y., Fu Z., Huang M., Wong C. C. L., Glickman M. H., Cong Y. (2019) **Structural Snapshots of 26S Proteasome Reveal Tetraubiquitin-Induced Conformations** *Mol. Cell* **73**:1150–1161 [Google Scholar](#)
27. Dong Y., Zhang S., Wu Z., Li X., Wang W. L., Zhu Y., Stoilova-McPhie S., Lu Y., Finley D., Mao Y. (2019) **Cryo-EM structures and dynamics of substrate-engaged human 26S proteasome** *Nature* **565**:49–55 [Google Scholar](#)
28. Wehmer M., Rudack T., Beck F., Aufderheide A., Pfeifer G., Plitzko J. M., Förster F., Schulten K., Baumeister W., Sakata E. (2017) **Structural insights into the functional cycle of the ATPase**

module of the 26S proteasome *Proc. Natl. Acad. Sci. U. S. A* **114**:1305–1310 [Google Scholar](#)

29. Eisele M. R., Reed R. G., Rudack T., Schweitzer A., Beck F., Nagy I., Pfeifer G., Plitzko J. M., Baumeister W., Tomko R. J., Sakata E. (2018) **Expanded Coverage of the 26S Proteasome Conformational Landscape Reveals Mechanisms of Peptidase Gating** *Cell Rep* **24**:1301–1315 [Google Scholar](#)
30. Förster A., Masters E. I., Whitby F. G., Robinson H., Hill C. P. (2005) **The 1.9 Å structure of a proteasome-11S activator complex and implications for proteasome-PAN/PA700 interactions** *Mol. Cell* **18**:589–599 [Google Scholar](#)
31. Chuah J. J. Y., Rexroad M. S., Smith D. M. (2023) **High resolution structures define divergent and convergent mechanisms of archaeal proteasome activation** *Commun Biol* **6**:1–16 [Google Scholar](#)
32. Religa T. L., Sprangers R., Kay L. E. (2010) **Dynamic regulation of archaeal proteasome gate opening as studied by TROSY NMR** *Science* **328**:98–102 [Google Scholar](#)
33. Stadtmueller B. M., Hill C. P. (2011) **Proteasome activators** *Mol. Cell* **41**:8–19 [Google Scholar](#)
34. Smith D. M., Chang S.-C., Park S., Finley D., Cheng Y., Goldberg A. (2007) **Docking of the Proteasomal ATPases' C-termini in the 20S Proteasomes alpha Ring Opens the Gate for Substrate Entry** *Mol. Cell* **27**:731–744 [Google Scholar](#)
35. de la Peña A. H., Goodall E. A., Gates S. N., Lander G. C., Martin A. (2018) **Substrate-engaged 26S proteasome structures reveal mechanisms for ATP-hydrolysis-driven translocation** *Science* **362** [Google Scholar](#)
36. Yu Y., Smith D. M., Kim H. M., Rodriguez V., Goldberg A. L., Cheng Y. (2010) **Interactions of PAN's C-termini with archaeal 20S proteasome and implications for the eukaryotic proteasome-ATPase interactions** *EMBO J* **29**:692–702 [Google Scholar](#)
37. Hill C. P., Masters E. I., Whitby F. G. (2002) **The 11S regulators of 20S proteasome activity** *Curr. Top. Microbiol. Immunol* **268**:73–89 [Google Scholar](#)
38. Chuah J. J., Thibaudeau T. A., Smith D. M. (2023) **Minimal mechanistic component of HbYX-dependent proteasome activation that reverses impairment by neurodegenerative-associated oligomers** *Commun. Biol* [Google Scholar](#)
39. Dong Y., Zhang S., Wu Z., Li X., Wang W. L., Zhu Y., Stoilova-McPhie S., Lu Y., Finley D., Mao Y. (2019) **Cryo-EM structures and dynamics of substrate-engaged human 26S proteasome** *Nature* **565**:49–55 [Google Scholar](#)
40. Dorn I. T., Eschrich R., Seemüller E., Guckenberger R., Tampé R. (1999) **High-resolution AFM-imaging and mechanistic analysis of the 20 S proteasome** *J. Mol. Biol* **288**:1027–1036 [Google Scholar](#)
41. Punjani A., Rubinstein J. L., Fleet D. J., Brubaker M. A. (2017) **cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination** *Nat. Methods* **14**:290–296 [Google Scholar](#)

42. Goddard T. D., Huang C. C., Meng E. C., Pettersen E. F., Couch G. S., Morris J. H., Ferrin T. E. (2018) **UCSF ChimeraX: Meeting modern challenges in visualization and analysis** *Protein Sci. Publ. Protein Soc* **27**:14–25 [Google Scholar](#)
43. Pettersen E. F., Goddard T. D., Huang C. C., Meng E. C., Couch G. S., Croll T. I., Morris J. H., Ferrin T. E. (2021) **UCSF ChimeraX: Structure visualization for researchers, educators, and developers** *Protein Sci. Publ. Protein Soc* **30**:70–82 [Google Scholar](#)
44. Liebschner D., Afonine P. V., Baker M. L., Bunkóczi G., Chen V. B., Croll T. I., Hintze B., Hung L.-W., Jain S., McCoy A. J., Moriarty N. W., Oeffner R. D., Poon B. K., Prisant M. G., Read R. J., Richardson J. S., Richardson D. C., Sammito M. D., Sobolev O. V., Stockwell D. H., Terwilliger T. C., Urzhumtsev A. G., Videau L. L., Williams C. J., Adams P. D. (2019) **Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix** *Acta Crystallogr. Sect. Struct. Biol* **75**:861–877 [Google Scholar](#)
45. Adams P. D., Afonine P. V., Bunkóczi G., Chen V. B., Davis I. W., Echols N., Headd J. J., Hung L.-W., Kapral G. J., Grosse-Kunstleve R. W., McCoy A. J., Moriarty N. W., Oeffner R., Read R. J., Richardson D. C., Richardson J. S., Terwilliger T. C., Zwart P. H. (2010) **PHENIX: a comprehensive Python-based system for macromolecular structure solution** *Acta Crystallogr. D Biol. Crystallogr* **66**:213–221 [Google Scholar](#)
46. Afonine P. V., Poon B. K., Read R. J., Sobolev O. V., Terwilliger T. C., Urzhumtsev A., Adams P. D. (2018) **Real-space refinement in PHENIX for cryo-EM and crystallography** *Acta Crystallogr. Sect. Struct. Biol* **74**:531–544 [Google Scholar](#)
47. Emsley P., Lohkamp B., Scott W. G., Cowtan K. (2010) **Features and development of Coot** *Acta Crystallogr. D Biol. Crystallogr* **66**:486–501 [Google Scholar](#)

Author information

Janelle JY Chuah

Department of Biochemistry and Molecular Medicine, West Virginia University School of Medicine, Morgantown, United States
ORCID iD: [0000-0002-1139-694X](#)

Madalena R Daugherty

Department of Biochemistry and Molecular Medicine, West Virginia University School of Medicine, Morgantown, United States

David M Smith

Department of Biochemistry and Molecular Medicine, West Virginia University School of Medicine, Morgantown, United States, Department of Neuroscience, Rockefeller Neuroscience Institute, West Virginia University, Morgantown, United States
ORCID iD: [0000-0002-1502-676X](#)

For correspondence: dmsmith@hsc.wvu.edu

Editors

Reviewing Editor

Luke Wiseman

Scripps Research Institute, La Jolla, United States of America

Senior Editor

Volker Dötsch

Goethe University Frankfurt, Frankfurt am Main, Germany

Reviewer #1 (Public review):

Summary:

In this manuscript, Chua, Daugherty, and Smith analyze a new set of archaeal 20S proteasomes obtained by cryo-EM that illustrate how the occupancy of the HbYX binding pocket induces gate opening. They do so primarily through a V24Y mutation in the α -subunit. These results are supported by a limited set of mutations in K66 in the α subunit, bringing new emphasis to this unit.

Strengths:

The new structure's analysis is comprehensive, occupying the entire manuscript. As such, the scope of this manuscript is very narrow, but the strength of the data is solid, and they offer an interesting and important new piece to the gate-opening literature.

Weaknesses:

Major Concerns

- (1) This manuscript rests on one new cryo-EM structure, leading to a single (albeit convincing) experiment demonstrating the importance of occupying the pocket and moving K66. Could a corresponding bulky mutation at K66 not activate the 20S proteasome?
- (2) To emphasize the importance of this work, the authors highlight the importance of gate-opening to human 20S proteasomes. However, the key distinctions between these proteasomes are not given sufficient weight.
 - (a) As the authors note, the six distinct Rpt C-termini can occupy seven different pickets. However, how these differences would impact activation is not thoroughly discussed.
 - (b) With those other sites, the relative importance of various pockets, such as the one controlling the α 3 N-terminus, should be discussed more thoroughly as a potential critical difference.
 - (c) These differences can lead to eukaryote 20S gates shifting between closed and open and having a partially opened state. This becomes relevant if the goal is to lead to an activated 20S. It would have been interesting to have archaea 20S with a mix of WT and V24Y α -subunits. However, one might imagine the subclassification problem would be challenging and require an extraordinary number of particles.
 - (d) Furthermore, the conservation of the amino acids around the binding pocket was not addressed. This seems particularly important in the relative contribution of a residue analogous to K66 or V24.

<https://doi.org/10.7554/eLife.106611.1.sa1>

Reviewer #2 (Public review):**Summary:**

The manuscript by Chuah et al. reports the experimental results that suggest the occupancy of the HbYX pockets suffices for proteasome gate opening. The authors conducted cryo-EM reconstructions of two mutant archaeal proteasomes. The work is technically sound and may be of special interest in the field of structural biology of the proteasomes.

Strengths:

Overall, the work incrementally deepens our understanding of the proteasome activation and expands the structural foundation for therapeutic intervention of proteasome function. The evidence presented appears to be well aligned with the existing literature, which adds confidence in the presentation.

Weaknesses:

The paper may benefit from some minor revision by making improvements on the figures and necessary quantitative comparative studies.

<https://doi.org/10.7554/eLife.106611.1.sa0>