

Reviewed Preprint

v1 • September 1, 2025

Not revised

Reviewed Preprint

v2 • December 5, 2025

Revised by authors

For correspondence:

lmahadev@g.harvard.edu

Competing interests: No competing interests declared

Funding: See [page 21](#)

Reviewing editor: Saad Jbabdi, University of Oxford, United Kingdom

© 2025, Choi et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Biophysical basis for brain folding and misfolding patterns in ferrets and humans

Gary PT Choi¹, Chunzi Liu², Sifan Yin², Gabrielle Séjourné^{2,3}, Richard S Smith⁴, Christopher A Walsh^{5,6}, L Mahadevan^{2,7,8} ✉

¹Department of Mathematics, The Chinese University of Hong Kong, Hong Kong, China • ²School of Engineering and Applied Sciences, Harvard University, Cambridge, United States • ³Department of Cell Biology, Duke University, Durham, United States • ⁴Department of Pharmacology, Feinberg School of Medicine, Northwestern University, Chicago, United States • ⁵Division of Genetics and Genomics, Manton Center for Orphan Disease, and Howard Hughes Medical Institute, Boston, United States • ⁶Boston Children's Hospital, Boston, United States • ⁷Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States • ⁸Department of Physics, Harvard University, Cambridge, United States

eLife Assessment

This **important** study characterises the morphogenesis of cortical folding in the ferret and human cerebral cortex using complementary physical and computational modelling. Notably, these approaches are applied to charting, in the ferret model, known abnormalities of cortical folding in humans. The study finds **convincing** evidence that variation in cortical thickness and expansion account for deviations in morphology, and supports these findings using cutting-edge approaches from both physical gel models and numerical simulations. The study will be of broad interest to the field of developmental neuroscience.

<https://doi.org/10.7554/eLife.107141.2.sa3>

Abstract

A mechanistic understanding of neurodevelopment requires us to follow the multiscale processes that connect molecular genetic processes to macroscopic cerebral cortical formations and thence to neurological function. Using magnetic resonance imaging of the brain of the ferret, a model organism for studying cortical morphogenesis, we create *in vitro* physical gel models and *in silico* numerical simulations of normal brain gyrification. Using observations of genetically manipulated animal models, we identify cerebral cortical thickness and cortical expansion rate as the primary drivers of dysmorphogenesis and demonstrate that *in silico* models allow us to examine the causes of aberrations in morphology and developmental processes at various stages of cortical ontogenesis. Finally, we explain analogous cortical malformations in human brains, with comparisons with human phenotypes induced by the same genetic defects, providing a unified perspective on brain morphogenesis that is driven proximally by genetic causes and affected mechanically via variations in the geometry of the brain and differential growth of the cortex.

Understanding the growth and form of normal and abnormal cortical convolutions (gyri and sulci) is important for the study of human neurodevelopmental diseases [Lui et al. \(2011\)](#) [Geschwind and Rakic \(2013\)](#) [Molnár et al. \(2019\)](#) [Del-Valle-Anton and Borrell \(2022\)](#) [Akula et al. \(2023a\)](#). During early brain development, the cortical plate expands tangentially relative to the underlying white matter [Welker \(1990\)](#). This pattern of growth is the central cause of gyrification; indeed tangential cortical expansion creates compressive forces on the faster-growing outer layer of the cortex and tensile forces on the attached slower-growing inner layer, and the relative-growth induced forces cause cortical folding as suggested more than a century

ago His (1868 [↗](#)) and first quantitatively elucidated nearly fifty years ago Richman et al. (1975 [↗](#)). At a molecular and cellular level, neurogenesis, neuronal migration, and neuronal differentiation all contribute to the tangential growth of the developing cortex via processes such as an increase in either the number or size of cells Fietz et al. (2010) [↗](#); Hansen et al. (2010) [↗](#); Borrell (2018) [↗](#); Van Essen (2022 [↗](#)). Recent models that take these facts into account attempt to explain gyrification in terms of a simple mechanical instability, termed sulcification Hohlfeld and Mahadevan (2011 [↗](#)), that when iterated with variations Tallinen et al. (2013 [↗](#)) shows that tangential expansion of the gray matter constrained by the white matter can explain a range of different morphologies seen in the brains of different organisms Tallinen et al. (2014) [↗](#); Kroenke and Bayly (2018 [↗](#)). Furthermore, when deployed over developmental time to simulate normal human cortical convolution, the results can capture a substantial range of features seen in normal human fetal brain morphogenesis Tallinen et al. (2016 [↗](#)). However, these and other similar studies Toro and Burnod (2005) [↗](#); Toro et al. (2008) [↗](#); Herculano-Houzel (2009) [↗](#); Azevedo et al. (2009) [↗](#); Hutton et al. (2009) [↗](#); Nie et al. (2010) [↗](#); Bayly et al. (2013) [↗](#); Budday et al. (2014) [↗](#); Garcia et al. (2021) [↗](#); Heuer et al. (2023) [↗](#); Pang et al. (2023) [↗](#); Schwartz et al. (2023 [↗](#)) do not allow us to understand malformations of cortical development (MCD), neurodevelopmental disorders that result from disrupted human cerebral cortex formation during embryonic brain development Desikan and Barkovich (2016 [↗](#)), owing to our inability to probe the development of the human fetal brain *in utero*. In addition, MCDs are difficult to relate to the physical properties of cerebral cortical folding because of the uncertainties in defining the effects of specific human genetic abnormalities on specific cortical features such as thickness and surface area.

An alternative strategy is to turn to model organisms to study the developmental trajectory of MCDs. Since commonly used animal models such as the mouse and rat have lissencephalic cortices, the ferret, a gyrencephalic non-primate, has been favored as an experimentally tractable laboratory organism that demonstrates cortical folding patterns that are roughly similar to that observed in the human Neal et al. (2007) [↗](#); Fietz et al. (2010) [↗](#); Sawada and Watanabe (2012) [↗](#); Johnson et al. (2018) [↗](#); Gilardi and Kalebic (2021 [↗](#)). Furthermore, since the process of cortical folding in the ferret is almost exclusively postnatal, with the progressive development of cortical gyri and sulci from postnatal day 0 (P0) to adolescence (Fig. 1 [↗](#)), it is more easily observable. Finally, the ability to perform region-specific genetic manipulation of the ferret brain through *in utero* electroporation Masuda et al. (2015) [↗](#); Tabata and Nakajima (2001 [↗](#)) makes the ferret an ideal system for modeling normal and abnormal neurodevelopmental processes.

Inspired by our previous studies using physical experiments with swelling gels and computational models of brain growth Tallinen et al. (2014) [↗](#); Tallinen and Biggins (2015) [↗](#); Tallinen et al. (2016 [↗](#)), we model the folding of a normal ferret brain using a physical gel model and a computational model based on the principle of constrained cortical expansion and compare the simulation results with the real brain development using various geometric morphometric approaches. We then use the computational and physical models to reproduce defective developmental processes of the ferret brain and show that they are consistent with biological experiments that manipulate different molecular drivers of neurogenesis, neuronal migration and cell growth in the cortex that underlie its relative thickness and expansion rate. Taken together, our studies provide a mesoscopic approach to brain morphogenesis that combines computational *in silico* and physical gel *in vitro* models with morphological and molecular analysis of ferret cortical disease models, and shed light on analogous MCDs in human brains.

Physical gel model

Inspired by the observation that soft physical gels swell and fold superficially when immersed in solvents, we constructed a physical simulacrum of ferret brain folding following our previous protocols Tallinen et al. (2014 [↗](#), 2016). Specifically, we produced two-layer PDMS gel models of the ferret brain at various ages based on surfaces reconstructed from MR Images (see SI Section S1 for details). We then immersed the two-layer gel brain model in n-hexane, which led to folding patterns by solvent-driven swelling of the outer layers (Fig. 2a [↗](#)).




Figure 1. Time course of ferret brain morphogenesis.

a, Whole brain samples from ferrets of various ages show progressive development of cortical gyri and sulci. **b**, Ferret brains show an increase in complexity of sulcal pattern and in sulcal depth throughout development. The rhinal sulcus (rs), cruciate sulcus (crs), coronal sulcus (cns), suprasylvian sulcus (sss), pseudosylvian sulcus (pss), lateral sulcus (ls), and ansate sulcus (as) are labelled. Schematic by G. Séjourné.




Figure 2. Physical gel model of ferret brain morphogenesis.

a, Schematic of the gel experiment. We first produced a two-layer gel model of a ferret brain from MRI scans as previously described Tallinen et al. (2016). We then immersed the gel model in n-hexane for 1.5 hours, which induced the outer layer to swell by absorbing the solvent over time, resulting in the development of cortical gyri and sulci. **b**, The swelling experiment for the P8 ferret model, in which it can be observed that the swelling of the cortical layer produces sulcal patterns and characteristics comparable to the real ferret brain. **c**, The swelling experiment for the P16 gel model. Scale bar = 1cm. Notation guide: cruciate sulcus (crs), coronal sulcus (cns), suprasylvian sulcus (sss), rhinal sulcus (rs), pseudosylvian sulcus (pss), lateral sulcus (ls), ansate sulcus (as).

Fig. 2b [↗](#) shows the experimental results for a P8 gel brain; it swells nonuniformly and folds progressively from an initial state that has invaginations corresponding to the cruciate sulcus (crs), the coronal sulcus (cns), and the suprasylvian sulcus (sss). The post-swelling state shows the development of sulci corresponding in location and self-contacting nature to the crs, cns, sss, and the formation of rhinal sulcus (rs), the pseudosylvian sulcus (pss), and lateral sulcus (ls), and ansate sulcus (as) observed in real ferrets aged P21 and older. Fig. 2b [↗](#) shows another swelling experiment with a starting shape corresponding to the P16 gel brain, from which we observe a similar progression in the folding patterns. We see that our minimal physical model can capture the qualitative aspects of the folding transitions in the ferret brain (see also SI Section S1 and Video S1–S2).

Computational model

To complement our physical experiments with quantitative simulations of ferret brain development, we followed the approach in Tallinen et al. (2014 [↗](#), 2016) and considered a neo-Hookean material model for the brain cortex consisting of a layer of gray matter on top of a deep layer of white matter with volumetric strain energy density

$$W = \frac{\mu}{2} [Tr(\mathbf{F}\mathbf{F}^T)J^{-2/3} - 3] + \frac{K}{2} (J - 1)^2, \quad (1)$$

where $\mathbf{F}$ is the deformation gradient, $J = \det(\mathbf{F})$, μ is the shear modulus and K is the bulk modulus. We assume that $K = 5\mu$ for a modestly compressible material. Computer simulations were then performed on tetrahedral meshes of ferret brains to model the gyrification (see SI Section S2 for more details).

We considered both simulations that modeled the changes in brain morphology from P0 to P32 as one continuous process (Fig. 3b [↗](#), see also SI Movie S3) and stepwise simulations that considered the growth process in stages, i.e. from P0 to P4, from P4 to P8, from P8 to P16 and from P16 to P32 (Fig. 3b [↗](#), see also SI Movie S4). In both sets of simulations, the emergence of cortical folding can be observed. In the continuous simulation approach, we observed the appearance of multiple minor folds since the continuous simulations only depend on the P0 initial brain, so that the effect of minor features in the P0 brain on the brain growth may accumulate over time. By contrast, in the stepwise simulation approach which focuses on multiple shorter growth periods, and thus reduces the accumulation of shape variations over time. Comparing the P16 results of stepwise numerical simulation, the gel experiment and the P16 real brain, we observe that the folding patterns are visually very similar (Fig. 4b [↗](#)). For a more quantitative comparison, we applied a method based on aligning landmarks using spherical mapping termed FLASH (Fast Landmark Aligned Spherical Harmonic Parameterization) Choi et al. (2015) [↗](#) to parameterize the simulated P16 ferret brain and the P16 brain surface generated from the MRI scans onto the unit sphere using landmark-aligned optimized conformal mappings, with the coronal sulcus (cns), suprasylvian sulcus (sss), presylvian sulcus (prs), and pseudosylvian sulcus (pss) on both the left and right hemispheres used as landmarks (see Fig. 4b [↗](#), and SI Section S3 for more details). We then assessed the geometric similarity of the two brain surfaces on the spherical domain in terms of their shape index Koenderink and van Doorn (1992 [↗](#)), which is a surface measure defined based on the surface mean curvature and Gaussian curvature. The similarity of the two shape indices suggests that the folding pattern produced by our simulation is close to the actual folding pattern (see SI Section S3 for additional analyses and Yin et al. (2025 [↗](#)) for more details of the morphometric method). We further utilized spherical harmonic-based representations for comparing the real and simulated ferret brains at different maximum orders, which also show that they have consistent geometric similarities (see SI Section S3).

Physical gel and computational models for misfolding ferret brains

The effectiveness of our differential growth-based model in quantifying the normative development of normal ferret brains begs the question of whether we can use the same framework to study malformations of cortical development (MCD). Here, we first consider simulating ferret cortical misfolding using both modified physical gel and computational models.




Figure 3. Numerical model of ferret brain morphogenesis.

a, Continuous growth simulation from P0 to adolescence. The P0 brain tetrahedral mesh was used as the input for the numerical simulation. **b**, Stepwise growth simulation from P0 to P4, P4 to P8, P8 to P16, and P16 to P32. For different growth intervals, we use different brain tetrahedral meshes as the input for the numerical simulation. Different views of the input P0 brain and the simulated P16 brain are provided.




Figure 4. Comparison of cortical folding in real and simulated ferret brain models.

a, The top row shows the increase in complexity of sulcal pattern and in sulcal depth of ferret brains from P8 to P16. The middle row shows a numerical model of a P8 brain and its deformed state mimicking progression to P16. The bottom row shows a physical gel model of P8 and its post-swelling state mimicking progression to P16 (scale bar = 1cm). The P8 initial states have invaginations corresponding to the cruciate sulcus (crs), coronal sulcus (cns) and suprasylvian sulcus (sss), and both the numerical deformed state and the physical post-swelling state have sulci corresponding in location and self-contacting nature to the crs, cns, sss, lateral sulcus (ls), and ansate sulcus (as) observed in P16 real ferrets. **b**, The real P16 brain reconstructed from MRI scans, the simulated P16 brain, and their respective landmark-aligned spherical mappings obtained by the FLASH algorithm Choi et al. (2015), each color-coded with the shape index Koenderink and van Doorn (1992) of the brain.

In Fig. 5b [↗](#), we performed a modified numerical experiment on the P8 brain with the cortical thickness reduced to 1/4 of the original thickness globally. We also performed a modified gel brain experiment by surfacecoating 1 layer of PDMS gel onto the core layer instead of 4 layers as in the original gel model in Fig. 2 [↗](#), equivalent to reducing the cortical layer thickness to 1/4 of the original one. In both the numerical and gel experimental results, it can be observed that small, tightly packed folds are formed. In Fig. 5b [↗](#), we performed another modified numerical experiment with the cortical thickness doubled globally. We also performed a modified gel brain experiment by surface-coating 8 layers of PDMS gel onto the core layer instead of 4 layers as in the original gel model to double the cortical layer thickness. In both the numerical and gel experimental results, it can be observed that the number of small folds is significantly reduced. In both cases, the numerical and gel results show a good qualitative match.

Neurology of ferret and human cortical malformations

After performing the modified physical gel and computational experiments for ferret brain misfoldings, we proceed to consider the effect of human genetic variants on brain gyrification in ferrets in prior studies (see Table 1 [↗](#)). For instance, *SCN3A* encodes a sodium channel and specific missense mutations of it are associated with the MCD polymicrogyria (PMG) Smith et al. (2018 [↗](#)), while genetic manipulations of *ASPM* have also been shown to produce severe microcephaly in human and ferret brains Johnson et al. (2018 [↗](#)), while disrupting *TMEM161B* leads to cortical malformations in humans and ferrets Akula et al. (2023b) [↗](#). In these and other examples, various genetic causes lead to variations in the cortical thickness ratio h/R and/or the tangential growth ratio g in space and time, which we know to be critical geometric parameters that change the physical nature of the sulcification instability driving cortical folding.

In Fig. 6b [↗](#), we show a control human brain MRI (top), a wild-type P16 ferret brain MRI (middle), and the stepwise numerical simulation result under the normal parameter setup (bottom). In Fig. 6b [↗](#), (top) we show the MCD polymicrogyria (PMG) phenotype in humans associated with over-expression of a mutated *SCN3A* gene Smith et al. (2018 [↗](#)). In Fig. 6b [↗](#) (middle), we show that the same mutation in ferrets leads to an increased number of tightly packed folds in the perisylvian region Smith et al. (2018 [↗](#)). To model this cortical malformation, we performed a modified numerical simulation with the cortical thickness reduced to 1/4 of the original thickness at a localized zone around the perisylvian region of the P8 brain model. From the modified numerical simulation result (Fig. 6b [↗](#), bottom), we can see that our model can qualitatively capture perisylvian PMG (see also SI Section S4).

In Fig. 6b [↗](#) (top) and Fig. 6b [↗](#) (middle) Johnson et al. (2018 [↗](#)) we show that *ASPM* mutants produce severe microcephaly in human and ferret brains, because the cortical surface area is reduced while there is no significant change in the cortical thickness. To reproduce this malformation, we consider a modified computational experiment with a reduction of the growth rate to 1/4 of the original rate. In Fig. 6b [↗](#) (bottom), we show the results of numerical simulations that lead to less prominent folding when compared to the normal brain, consistent with observations in human and ferret brains.

Finally, in Fig. 6b [↗](#) (top) and Fig. 6b [↗](#) (middle) Akula et al. (2023b) [↗](#), we show that disrupting *TMEM161B* also leads to cortical malformations in human and ferret brains, with shallower sulci. Modifying the morphogenetic simulations with a reduction of the growth rate to 3/4 of the original rate and an increase of the cortical thickness to 1.5 times the original thickness leads to results shown in Fig. 6b [↗](#) (bottom) that match the experimentally observed malformation patterns qualitatively.

In SI Section S4, we present additional computational experiments to demonstrate the effect of different combinations of the growth rate and cortical thickness parameters on the cortical malformation results.

Figure 5. Numerical and gel experiments on the P8 ferret brain with globally modified brain gyrification.

a, The numerical and gel experimental results with a global reduction of the cortical thickness to 1/4 of the original thickness.
b, The numerical and gel experimental results with a global increase in the cortical thickness to twice the original thickness.
 Scale bar = 1cm.




Table 1. Human genetic variants modeled in ferret developmental brain phenotypes.

Gene	Change in geometry	Relevant brain developmental disorder
<i>SCN3A</i> Smith et al. (2018)	Excessive number of tightly packed folds	Polymicrogyria
<i>ASPM</i> Johnson et al. (2018)	Reduced brain size and cortical surface area	Microcephaly
<i>Cdk5</i> Shinmyo et al. (2017)	Reduced depth of the sulcus	Lissencephaly
<i>ARHGAP11B</i> Kalebic et al. (2018)	Expansion in both the radial and tangential dimensions	Megalencephaly
<i>TMEM161B</i> Akula et al. (2023b)	Reduced gyrus size and sulcal depth	Lissencephaly




Figure 6. Modeling malformations of cortical development (MCD) using our model.

a, Control. **b**, *SCN3A*. **c**, *ASPM*. **d**, *TMEM161B*. For each example, we show human (top) and ferret (middle) brain MRIs (images adapted from Smith et al. (2018) [DOI](#); Johnson et al. (2018) [DOI](#); Akula et al. (2023b) [DOI](#)). We then perform a modified numerical brain simulation on the P8 model with different tangential growth rate and cortical thickness parameters, including (**a**) the original growth rate and cortical thickness, (**b**) a reduction of the cortical thickness at a localized zone, (**c**) a reduction of the growth rate globally, and (**d**) a reduction of the growth rate and an increase of the cortical thickness globally. All numerical simulation results (bottom) qualitatively capture the cortical malformations.

© 2018 Elsevier Inc. The ferret (middle) brain MRI image in panel a is reprinted from Smith et al. (2018) [DOI](#) with permission. It is not covered by the CC-BY 4.0 licence and further reproduction of this image would need permission from the copyright holder.

Discussion

Understanding the growth and form of the cerebral cortex is a fundamental question in neurobiology, and the experimentally accessible progressive postnatal development of the ferret brain makes it an ideal system for analysis. Here, we have used a combination of physical and computational models based on differential growth to show how ferret brain morphologies arise. One may also use the physical and computational models to compare the ferret brain folding patterns with those in macaques and humans, as in our companion study Yin et al. (2025 [DOI](#)).

By modifying the scaled cortical layer thickness and the tangential growth profile in our model, we have qualitatively reproduced various cortical malformations and shown how developmental mechanisms lead to morphological manifestations with potential functional implications. As several diseases, such as ion channels Smith and Walsh (2020 [DOI](#)); Smith et al. (2021 [DOI](#)), converge on brain malformations, future studies to validate across disease pathways could leverage these results. All together, our study elucidates the normal and abnormal folding in the ferret brain as a function of its genetic antecedents that lead to changes in the geometry of the cortex and thence to different physical folding patterns with functional consequences. A computational and physical-gel brain study informed by detailed MRI of ferret and human fetal brains allows us to move towards a synthesis of the genetic, physical and morphological basis for cortical malformations. Natural next steps include accounting for varying spatio-temporal expansion rates of the cortex to capture the quantitative differences in the development of fetal folding patterns in both ferrets and humans, and understanding the functional consequences as a result of impaired connectivity due to misfolding.

Methods

Physical gel model for ferret brain folding

Beginning with T2-weighted motion-corrected anatomical MR images of ferret brains of various ages Toro et al. (2018 [DOI](#)), digital maps of the surfaces of native (pre-swollen) brain were recreated. Then, we followed our prior experimental approach Tallinen et al. (2014 [DOI](#), 2016 [DOI](#)) and produced twolayer PDMS gel models of the ferret brain at various ages based on the reconstructed brain surfaces. Specifically, we first generated a negative rubber mold with Ecoflex 00-30 from a 3D-printed brain plastic model and then the core gel with SYLGARD 184 at a 1:45 crosslinker:base ratio. To mimic the cortical layer, we surface-coated 4 layers of PDMS gel at a 1:35 crosslinker:base ratio onto the core layer. Finally, tangential cortical growth was mimicked by immersing the two-layer gel brain model in n-hexane for 1.5 hours, which resulted in solvent-driven swelling of the outer layers, leading to folding patterns. See SI Section S1 for details.

Computational model for ferret brain folding

Three geometrical parameters of the 3D brain models control its morphogenesis: the average brain size R (determined for example by its volume), the average cortical thickness T and the average tangential expansion ratio of the cortex relative to the white matter, g^2 . To characterize brain development in the ferret, we followed the empirical scaling laws for gray-matter volume to thickness described in Tallinen et al. (2014 [DOI](#)) and set $R/T \approx 10$ with the tangential expansion ratio $g \approx 1.9$, along with an indicator function $\theta(y) = (1 + e^{10(y/T - 1)})^{-1}$, with y the distance from surface in a material reference frame used to distinguish between the cortical gray matter layer (with $\theta = 1$) from the deeper white matter (with $\theta = 0$).

Using MRIs of ferret brains, we created a computational model of the initial brain size and shape, which was discretized using tetrahedral meshes with over one million tetrahedral elements using Netgen NGSolve Team (2019 [DOI](#)). Using a finite element method implemented using a discretized version of the energy of the system (1), we minimized the energy by quasistatic equilibration using an explicit solver Tallinen et al. (2016 [DOI](#)), while growth was applied incrementally using the form described earlier by expanding the tetrahedral elements with inversion handling Stomakhin et al. (2012 [DOI](#)) and a nodal pressure formulation Bonet and Burton (1998 [DOI](#)). Self-avoidance of the

surface was handled using the penalty-based vertex-triangle contact processing [Ericson \(2004\)](#). We also enforce the condition that there is no growth in the central part as well as in the bottom part of the brain to better simulate the development of ferret brains. See SI Section S2 for more details.

Malformations of cortical development

In [Fig. 6](#), we showed various human and ferret brain MRIs with malformations of cortical development. The *SCN3A* MRI images were adapted from [Smith et al. \(2018\)](#) (human: 15y; ferret: P33). The *ASPM* MRI images were adapted from [Johnson et al. \(2018\)](#) (human: 4y 10 mo; ferret: P16). The *TMEM161B* MRI images were adapted from [Akula et al. \(2023b\)](#) (human: 2y; ferret: P21).

The numerical simulation results for cortical malformations were obtained using the computational model described above, with the cortical thickness and growth rate parameters modified either locally or globally. See SI Section S4 for more details.

Data availability

Requests for the ferret brain data should be made to [Toro et al. \(2018\)](#). All other data are included in the article and/or supplementary material.

Supplementary material

S1 Physical gel model for ferret brain folding

S1.1 Surface reconstruction

Surfaces of initial (pre-swelling) brain states were reconstructed from T2-weighted motion-corrected anatomical MR images of ferret brains at various ages [1, 2] by R. Toro and K. Heuer. The reconstruction can be achieved by first computing masks of the pial and inner cortical surfaces from MRI data using the Nilearn Python module to generate and threshold image histograms [3], and then utilizing the Brainbox web application [4] to manually correct region of interest (ROI) selections and convert masks from Nifti (.nii.gz) files to three-dimensional triangular meshes, with conservative corrections to the output meshes including smoothing and inversion of inward-pointing normal vectors applied, with further mesh corrections performed using Meshlab.

S1.2 Swelling experiment

In main text [Fig. 2](#), we presented the swelling experimental results for an input P8 gel model and an input P16 gel model. In [Fig. S1](#), we present the swelling experiment for the P4 ferret brain with the same experimental setup, where the 2-layer PDMS model was immersed in n-hexane for 1.5 hours. It can be observed that while the sulcal pattern is not particularly prominent in the input P4 gel model, complex folding patterns can be observed in the resulting swollen gel.

We remark that comparing the P4 swelling result with the P8 and P16 gel brain experiments in main text [Fig. 2](#), one can see that the folding patterns are notably different. A possible reason is that all gel models were immersed in n-hexane for swelling with the same duration (1.5 hours) regardless of their size and shape, which might result in the variation in the final gyrification patterns.

S2 Computational model for ferret brain folding

S2.1 Formulation

For the numerical simulation of ferret brain development, we follow the approach in [5, 6] and consider a material consisting of a layer of gray matter on top of a deep layer of white matter. The material is assumed to be neo-Hookean with volumetric strain energy density

$$W = \frac{\mu}{2} [\text{Tr}(\mathbf{F}\mathbf{F}^T)J^{-2/3} - 3] + \frac{K}{2}(J - 1)^2, \quad (\text{S1})$$

where $\mathbf{F}$ is the deformation gradient, $J = \det(\mathbf{F})$, μ is the shear modulus and K is the bulk modulus. We assume that $K = 5\mu$ for a modestly compressible material.

Three geometrical parameters of the 3D brain models are the brain size R , the cortical thickness T and the tangential expansion g^2 . For ferret brain development, we follow the empirical scaling law for gray-matter volume to thickness and set $R/T \approx 10$ with the tangential expansion ratio $g \approx 1.9$. An indicator function

$$\theta(y) = \frac{1}{1 + e^{10(y/T-1)}} \quad (\text{S2})$$




Figure S1. Swelling experiment for an input P4 gel brain. Here, the two-layer PDMS model of a P4 ferret brain was immersed in n-hexane for 1.5 hours. Analogous to the gel brain experiments presented in the main text, gyrfication patterns can be observed in the swollen gel brain. Scale bar = 1cm.

is applied for distinguishing between the cortical layer (with $\theta = 1$) and white matter zone (with $\theta = 0$). Here, y is the distance from surface in material coordinates.

S2.2 Numerical experiments

Tetrahedral meshes were generated from the MRI-based reconstructed surfaces using the Netgen software [7], with each of them consisting of over 1 million tetrahedral elements. The numerical simulation procedure was implemented in C++. The energy of the system was minimized by quasistatic equilibration using an explicit scheme. Growth was applied by expanding the tetrahedral elements with inversion handling [8] and nodal pressure formulation [9]. We further set the innermost part and the bottom part of the brain tetrahedral meshes to be non-growing regions to better simulate the development of ferret brains. See [5, 6] for more details of the computation setup.




Figure S2. Stepwise growth simulation of ferret brain. The P0 brain and the simulated brains from P0 to P4, from P4 to P8, from P8 to P16, and from P16 to P32 are shown. For each brain, six different views are provided (not to scale).

We simulated the stepwise growth of the ferret brain from P0 to P4, from P4 to P8, from P8 to P16, and from P16 to P32. Fig. S2 shows different views of the simulation results.




Figure S3. Mesh independence for the numerical simulation. The continuous growth simulations from P0 to adolescence using P0 brain tetrahedral meshes with 250,000 tets (first row), 1 million tets (second row), 2 million tets (third row), and 3 million tets (fourth row). It can be observed that the resulting folding patterns are highly similar.

S2.3 Mesh independence

One may ask about the effect of mesh size on the numerical simulation. Fig. S3 shows the continuous simulation results starting from P0 with different mesh resolutions. It can be observed that the results are similar, which shows the mesh independence of the numerical simulation.

S3 Morphometric analysis

S3.1 Analyzing the brain surfaces using surface parameterization

The FLASH (Fast Landmark-Aligned Spherical Harmonic Parameterization) method [10] was used for quantifying the similarity of the numerically simulated brain and the MRI-based brain. Denote the two brain surfaces to be compared by $\mathcal{M}$ and $\mathcal{N}$. The FLASH method produces two bijective spherical mappings $f: \mathcal{M} \rightarrow \mathbb{S}^2$ and $g: \mathcal{N} \rightarrow \mathbb{S}^2$, so that both surfaces are mapped to the unit sphere with prescribed landmark pairs on them optimally aligned. Here, g is a conformal map from $\mathcal{N}$ to the unit sphere $\mathbb{S}^2$, and f is a landmark-constrained optimized conformal map from $\mathcal{M}$ to $\mathbb{S}^2$. More specifically, f achieves a balance between the landmark mismatch error and the conformality distortion, with the following combined energy minimized:

$$E(f) = \int_{\mathcal{M}} \|\nabla f\|^2 + \eta \sum_{i=1}^t \|f(p_i) - g(q_i)\|^2, \quad (\text{S3})$$

where $\{p_i, q_i\}$ are corresponding landmark curves on $\mathcal{M}$ and $\mathcal{N}$ respectively, and η is a balancing parameter. The coronal sulcus (cns), suprasylvian sulcus (sss), presylvian sulcus (prs), and pseudosylvian sulcus (pss) on both the left and right hemispheres of the ferret brain are used as the landmark curves to ensure that the two spherical parameterizations are consistent. We set $\eta = 10$ to achieve an accurate alignment of the two spherical parameterizations without inducing a large conformal distortion.

After obtaining the spherical parameterizations with the landmarks optimally aligned, we can compare the folding patterns of the two brain surfaces by evaluating the similarity of their shape indices on $\mathbb{S}^2$. The shape index is a dimensionless surface measure defined based on the surface mean curvature and the surface Gaussian curvature [11]. Specifically, the mean curvature H and Gaussian curvature K of any surface $\mathcal{S}$ are given by

$$H(v) = \frac{1}{2}(\kappa_1(v) + \kappa_2(v)) \quad (\text{S4})$$

and

$$K(v) = \kappa_1(v)\kappa_2(v), \quad (\text{S5})$$

where v is a point on $\mathcal{S}$, and $\kappa_1(v)$, $\kappa_2(v)$ are the principal curvatures at v . The shape index $I(v)$ [11] at each point v is then given by

$$I(v) = \frac{2}{\pi} \arctan \left(\frac{H(v)}{\sqrt{(H(v))^2 - K(v)}} \right). \quad (\text{S6})$$

It is easy to see that $I(v) \in [-1, 1]$ for any v . Intuitively, for brain cortical surfaces, the sulci are with a smaller value of I and the gyri are with a larger value of I .

Now, with the aid of the spherical parameterizations, we can easily evaluate the similarity of the shape index distributions of two brain surfaces. Specifically, the similarity s of the shape index distributions $I_{\mathcal{M}}, I_{\mathcal{N}}$ of the two surfaces $\mathcal{M}, \mathcal{N}$ is defined as

$$s(\mathcal{M}, \mathcal{N}) = 1 - \frac{1}{2m} \sum_{k=1}^m |I_{\mathcal{M}}(v_k) - I_{\mathcal{N}}((g^{-1} \circ f)(v_k))|, \quad (\text{S7})$$

where $\{v_k\}_{k=1}^m$ are the vertices of $\mathcal{M}$. In particular, since $I_{\mathcal{M}}, I_{\mathcal{N}} \in [-1, 1]$, we have

$$0 \leq |I_{\mathcal{M}}(v_k) - I_{\mathcal{N}}((g^{-1} \circ f)(v_k))| \leq 2 \quad (\text{S8})$$

for all k . Consequently, we have $s \in [0, 1]$, and $s = 1$ if the two brain surfaces are identical. As discussed in the main text, our result of $s \approx 0.83$ shows that the real and simulated brain surfaces are highly similar.

Besides analyzing the entire brain surfaces, one can also quantify the shape difference for each half-brain surface using the mapping and shape similarity quantification method and different norms as described in [12]:

$$s(\mathcal{M}, \mathcal{N}) = 1 - \frac{1}{2m^{1/p}} \|I_{\mathcal{M}}(v_k) - I_{\mathcal{N}}((g^{-1} \circ f)(v_k))\|_p, \quad (\text{S9})$$

where $\|\cdot\|_p$ is the p -norm. As shown by the quantification results in Table S1, the real and simulated half-brain surfaces are highly similar. These additional results again demonstrate the effectiveness of our model for simulating ferret brain folding.

S3.2 Spherical harmonics representations

With the spherical parameterizations, we further computed the spherical harmonics representation of the two surfaces with different maximum order N used. Denote (θ, ϕ) as the spherical coordinates of $\mathbb{S}^2$, where $\theta \in [0, \pi]$ is the elevation angle and $\phi \in [-\pi, \pi]$ is the azimuth angle. The spherical harmonics functions $Y_n^m : [0, \pi] \times [-\pi, \pi] \rightarrow \mathbb{R}$ of order n and degree m are given as follows [13]:

$$Y_n^m(\theta, \phi) = \begin{cases} (-1)^{|m|} \sqrt{2} K_n^m \sin(|m|\phi) P_n^{|m|}(\cos \theta) & : m < 0, \\ (-1)^{|m|} \sqrt{2} K_n^m \cos(m\phi) P_n^m(\cos \theta) & : m > 0, \\ K_n^0 P_n^0(\cos \theta) & : m = 0, \end{cases} \quad (\text{S10})$$

where

$$K_n^m = \sqrt{\frac{(2n+1)(n-|m|)!}{4\pi(n+|m|)!}}, \quad (\text{S11})$$

Surface	Similarity (1-norm)	Similarity (2-norm)	Similarity (∞ -norm)
Real P8 and Simulated P8 (left)	0.8040	0.7432	1.0000
Real P8 and Simulated P8 (right)	0.7893	0.7260	1.0000
Real P16 and Simulated P16 (left)	0.8247	0.7632	1.0000
Real P16 and Simulated P16 (right)	0.7980	0.7339	1.0000

Table S1. Similarity between the real and simulated ferret half-brain surfaces.

and

$$P_n^m(x) = \frac{(-1)^m}{2^n n!} (1-x^2)^{m/2} \frac{d^{n+m}}{dx^{n+m}} (x^2-1)^n. \quad (\text{S12})$$

With the maximum order N prescribed, we can consider all Y_n^m with $n = 0, 1, \dots, N$ and $m = -n, -n+1, \dots, 0, \dots, n-1, n$. We can then approximate any given surface as a linear combination of this set of SH functions:

$$\begin{cases} x(\theta, \phi) \approx \sum_{n=0}^N \sum_{m=-n}^n a_n^m Y_n^m(\theta, \phi), \\ y(\theta, \phi) \approx \sum_{n=0}^N \sum_{m=-n}^n b_n^m Y_n^m(\theta, \phi), \\ z(\theta, \phi) \approx \sum_{n=0}^N \sum_{m=-n}^n c_n^m Y_n^m(\theta, \phi), \end{cases} \quad (\text{S13})$$

where the coefficients a_n^m, b_n^m, c_n^m can be computed from the spherical parameterization (see [14] for details). As the maximum order N increases, more geometric details of the surface can be captured in the spherical harmonics representation.

From Fig. S4, it can be observed that for different maximum order N , the spherical harmonics representations of MRI-based brain and the numerically simulated brain match very well. More specifically, when $N = 5$ is used, both representations give an overall smooth brain shape without folding. When $N = 15$ is used, both of them start to exhibit some folding at consistent locations. When $N = 25$ is used, both of them show highly similar sulci and gyri patterns. This again indicates that the two brain surfaces are geometrically similar.




Figure S4. The spherical harmonics representations of the MRI-based brain and the numerically simulated brain with different maximum order N used.

S4 Cortical malformations

In the main text, we considered various types of cortical malformations and discussed how our proposed model can be modified to capture their features qualitatively. In this section, additional analyses are provided.

In Fig. S5 [\[link\]](#), we provide a comparison between human brain surface MRI reconstructions and our ferret brain simulations for the MCD polymicrogyria (PMG). Here, the human brain surface reconstruction images (SCN3A case and control age-matched subject) are adapted from [15], where MRI stacks were processed to extract the surfaces of the gray/white matter and gray matter/cerebrospinal fluid boundaries, and to define the cortical surface using FreeSurfer software. It can be observed that using our computational model with a modification in the cortical thickness, we obtain ferret brain simulation results with a good qualitative match.

In Fig. S6 [\[link\]](#), we further performed numerical simulations on the P8 ferret brain with different modifications in the tangential growth rate and cortical thickness at a localized region (dotted ellipse). Specifically, we first considered increasing the tangential growth rate at the localized region to 1.5 times or 2 times the original rate. In the numerical simulation results, more prominent folds can be observed at the corresponding region. Also, we considered reducing the cortical thickness to 1/2 or 1/4 of the original thickness at the localized region. In the simulation results, we can see that new folds are created at the localized region. Finally, we considered modifying both the tangential growth rate and the cortical thickness. In this case, a higher level of abnormal gyrification can be observed locally in the simulation results. Altogether, the modified brain experiments show that different cortical malformations can be effectively simulated by modifying the tangential growth rate and cortical thickness.




Figure S5. Comparison between human brain surface MRI reconstructions and our ferret brain simulations for the MCD polymicrogyria (PMG). **a.** Surface MRI reconstructions of human MRIs performed using FreeSurfer software of control (top) and age-matched affected individual with a gain-of-function *SCN3A* variant resulting in polymicrogyria. Red box outlines the PMG of perisylvian and surrounding areas, where the normal gyri and sulci form microgyri/sulci, pushing together to make an appearance of “Moroccan leather”. **b.** The numerical growth simulations on the P8 ferret brain without modification in the cortical thickness (top) and with a reduced cortical thickness to 1/4 of the original thickness, i.e., 0.25T (bottom). The differences are highlighted by the red boxes.

© 2018 Elsevier Inc. Images in panel a are reprinted from [Smith et al. \(2018\)](#) with permission. They are not covered by the CC-BY 4.0 licence and further reproduction of this panel would need permission from the copyright holder.




Figure S6. Numerical simulations with locally modified brain growth. We performed numerical simulations on the P8 brain with different modifications in the tangential growth rate g and the cortical layer thickness T at a localized region (dotted ellipse). The results without modification (normal), with an increased growth rate to 1.5 times the original growth rate (1.5g) or 2 times the original growth rate (2g), with a reduced cortical thickness to 1/2 of the original thickness (0.5T) or 1/4 of the original thickness (0.25T), and with a modification in both the growth rate and the cortical thickness (1.5g and 0.5T, 2g and 0.25T) are presented.

Acknowledgements

We thank Roberto Toro and Katja Heuer for the ferret brain data, and Jun Young Chung and James Weaver for their help with preliminary experiments. This work is supported in part by the Harvard Quantitative Biology Initiative and the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard, award no. 1764269 (to G.P.T.C. and L.M.), the Simons Foundation and the Henri Seydoux Fund (to L.M.). G.P.T.C. is also supported by the CUHK Faculty of Science Direct Grant for Research (Project Code 4053650). C.A.W. is supported by the NINDS through R01NS032457 and R37NS035129, grant 62587 from the John Templeton Foundation (the opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation), and by the Allen Discovery Center for Human Brain Evolution, a Paul G. Allen Frontiers Group advised program of the Paul G. Allen Family Foundation. C.A.W. is an Investigator of the Howard Hughes Medical Institute. R.S.S. was supported by R00NS112604 and R01NS140046.

Additional information

Code Availability Statement

Computer codes for numerical simulations and morphometric analyses are available on GitHub at <https://github.com/garyptchoi/ferret-brain-morphogenesis> .

Funding

Funder	Grant reference number	Author
Harvard Quantitative Biology Initiative and the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard	1764269	L Mahadevan
Simons Foundation (SF)		L Mahadevan
Henri Seydoux Fund		L Mahadevan
CUHK Faculty of Science Direct Grant for Research	4053650	Gary Choi
NINDS	R01NS032457	Christopher A Walsh
NINDS	R37NS035129	Christopher A Walsh
John Templeton Foundation (JTF)	62587	Christopher A Walsh
AI The Frontiers Group Allen Discovery Center (The Allen Discovery Centers)		Christopher A Walsh
NINDS	R00NS112604	Richard S Smith
NINDS	R01NS140046	Richard S Smith

Author ORCID iDs

Gary PT Choi: <http://orcid.org/0000-0001-5407-9111>

Chunzi Liu: <http://orcid.org/0000-0002-3469-3904>

Sifan Yin: <http://orcid.org/0000-0002-0296-3981>

Gabrielle Séjourné: <http://orcid.org/0000-0002-4633-2700>

Richard S Smith: <http://orcid.org/0000-0003-1937-6529>

Christopher A Walsh: <http://orcid.org/0000-0002-0156-2238>

L Mahadevan: <http://orcid.org/0000-0002-5114-0519>

Additional files

Video S1.  Time-lapse video of the swelling experiment for the P8 gel brain, where the 2-layer PDMS model was immersed in n-hexane for 1.5 hours.

Video S2.  Time-lapse video of the swelling experiment for a P16 gel brain, where the 2-layer PDMS model was immersed in n-hexane for 1.5 hours.

Video S3.  Continuous numerical simulation of the ferret brain folding from P0 to P32.

Video S4. [Stepwise numerical simulations of the ferret brain folding from P0 to P4, from P4 to P8, from P8 to P16, and from P16 to P32.](#)

References

1. **Akula SK**, Exposito-Alonso D, Walsh CA (2023a) Shaping the brain: The emergence of cortical structure and folding. *Dev Cell* **58**:2836-2849
2. **Akula SK**, Marciano JH, Lim Y, Exposito-Alonso D, Hylton NK, Hwang GH, Neil JE, Dominado N, Bunton-Stasyshyn RK, Song JHT, *et al.* (2023b) TMEM161B regulates cerebral cortical gyration, Sonic Hedgehog signaling, and ciliary structure in the developing central nervous system. *Proc Natl Acad Sci* **120**:e2209964120
3. **Azevedo FAC**, Carvalho LRB, Grinberg LT, Farfel JM, Ferretti REL, Leite REP, Filho WJ, Lent R, Herculano-Houzel S. (2009) Equal numbers of neuronal and nonneuronal cells make the human brain an isometrically scaled-up primate brain. *J Comp Neurol* **513**:532-541
4. **Bayly P**, Okamoto R, Xu G, Shi Y, Taber L. (2013) A cortical folding model incorporating stress-dependent growth explains gyral wavelengths and stress patterns in the developing brain. *Phys Biol* **10**:016005
5. **Bonet J**, Burton A. (1998) A simple average nodal pressure tetrahedral element for incompressible and nearly incompressible dynamic explicit applications. *Comm Numer Meth Eng* **14**:437-449
6. **Borrell V**. (2018) How cells fold the cerebral cortex. *J Neurosci* **38**:776-783
7. **Budday S**, Raybaud C, Kuhl E. (2014) A mechanical model predicts morphological abnormalities in the developing human brain. *Sci Rep* **4**:5644
8. **Choi PT**, Lam KC, Lui LM (2015) FLASH: Fast landmark aligned spherical harmonic parameterization for genus-0 closed brain surfaces. *SIAM J Imaging Sci* **8**:67-94
9. **Del-Valle-Anton L**, Borrell V. (2022) Folding brains: from development to disease modeling. *Physiol Rev* **102**:511-550
10. **Desikan RS**, Barkovich AJ (2016) Malformations of cortical development. *Ann Neurol* **80**:797-810
11. **Ericson C**. (2004) *Real-time collision detection* CRC Press.
12. **Fietz SA**, Kelava I, Vogt J, Wilsch-Bräuninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R, *et al.* (2010) OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci* **13**:690-699
13. **Garcia KE**, Wang X, Kroenke CD (2021) A model of tension-induced fiber growth predicts white matter organization during brain folding. *Nat Commun* **12**:6681
14. **Geschwind DH**, Rakic P. (2013) Cortical evolution: judge the brain by its cover. *Neuron* **80**:633-647
15. **Gilardi C**, Kalebic N. (2021) The ferret as a model system for neocortex development and evolution. *Front Cell Dev Biol* **9**:661759
16. **Hansen DV**, Lui JH, Parker PRL, Kriegstein AR (2010) Neurogenic radial glia in the outer subventricular zone of human neocortex. *Nature* **464**:554-561
17. **Herculano-Houzel S**. (2009) The human brain in numbers: a linearly scaled-up primate brain. *Front Hum Neurosci* **3**:857
18. **Heuer K**, Traut N, de Sousa AA, Valk SL, Clavel J, Toro R. (2023) Diversity and evolution of cerebellar folding in mammals. *eLife* **12**:e85907 <https://doi.org/10.7554/eLife.85907>
19. **His W**. (1868) *Untersuchungen über die erste Anlage des Wirbelthierleibes: die erste Entwicklung des Hühnchens im Ei* **1** FCW Vogel.
20. **Hohlfeld E**, Mahadevan L. (2011) Unfolding the sulcus. *Phys Rev Lett* **106**:105702
21. **Hutton C**, Draganski B, Ashburner J, Weiskopf N. (2009) A comparison between voxel-based cortical thickness and voxel-based morphometry in normal aging. *Neuroimage* **48**:371-380

22. Johnson MB, Sun X, Kodani A, Borges-Monroy R, Girsakis KM, Ryu SC, Wang PP, Patel K, Gonzalez DM, Woo YM, *et al.* (2018) Aspm knockout ferret reveals an evolutionary mechanism governing cerebral cortical size. *Nature* **556**:370-375
23. Kalebic N, Gilardi C, Albert M, Namba T, Long KR, Kostic M, Langen B, Huttner WB (2018) Human-specific ARHGAP11B induces hallmarks of neocortical expansion in developing ferret neocortex. *eLife* **7**:e41241 <https://doi.org/10.7554/eLife.41241>
24. Koenderink JJ, van Doorn AJ (1992) Surface shape and curvature scales. *Image Vis Comput* **10**:557-564
25. Kroenke CD, Bayly PV (2018) How forces fold the cerebral cortex. *J Neurosci* **38**:767-775
26. Lui JH, Hansen DV, Kriegstein AR (2011) Development and evolution of the human neocortex. *Cell* **146**:18-36
27. Masuda K, Toda T, Shinmyo Y, Ebisu H, Hoshiba Y, Wakimoto M, Ichikawa Y, Kawasaki H. (2015) Pathophysiological analyses of cortical malformation using gyrencephalic mammals. *Sci Rep* **5**:1-15
28. Molnár Z, Clowry GJ, estan N, Alzu'bi A, Bakken T, Hevner RF, Hüppi PS, Kostović I, Rakic P, Anton E, *et al.* (2019) New insights into the development of the human cerebral cortex. *J Anat* **235**:432-451
29. Neal J, Takahashi M, Silva M, Tiao G, Walsh CA, Sheen VL (2007) Insights into the gyrification of developing ferret brain by magnetic resonance imaging. *J Anat* **210**:66-77
30. NGSolve Team (2019) NetGen. <https://ngsolve.org/>
31. Nie J, Guo L, Li G, Faraco C, Miller LS, Liu T. (2010) A computational model of cerebral cortex folding. *J Theor Biol* **264**:467-478
32. Pang JC, Aquino KM, Oldehinkel M, Robinson PA, Fulcher BD, Breakspear M, Fornito A. (2023) Geometric constraints on human brain function. *Nature* **618**:566-574
33. Richman DP, Stewart RM, Hutchinson JW, Caviness VS (1975) Mechanical model of brain convolutional development. *Science* **189**:18-21
34. Sawada K, Watanabe M. (2012) Development of cerebral sulci and gyri in ferrets (*Mustela putorius*). *Congenit Anom* **52**:168-175
35. Schwartz E, Nenning KH, Heuer K, Jeffery N, Bertrand OC, Toro R, Kasprian G, Prayer D, Langs G. (2023) Evolution of cortical geometry and its link to function, behaviour and ecology. *Nat Commun* **14**:2252
36. Shinmyo Y, Terashita Y, Duong TAD, Horiike T, Kawasumi M, Hosomichi K, Tajima A, Kawasaki H. (2017) Folding of the cerebral cortex requires Cdk5 in upper-layer neurons in gyrencephalic mammals. *Cell Rep* **20**:2131-2143
37. Smith RS, Florio M, Akula SK, Neil JE, Wang Y, Hill RS, Goldman M, Mullally CD, Reed N, Bello-Espinosa L, *et al.* (2021) Early role for a Na⁺, K⁺-ATPase (ATP1A3) in brain development. *Proc Natl Acad Sci* **118**:e2023333118
38. Smith RS, Kenny CJ, Ganesh V, Jang A, Borges-Monroy R, Partlow JN, Hill RS, Shin T, Chen AY, Doan RN, *et al.* (2018) Sodium channel SCN3A (NaV1. 3) regulation of human cerebral cortical folding and oral motor development. *Neuron* **99**:905-913
39. Smith RS, Walsh CA (2020) Ion channel functions in early brain development. *Trends Neurosci* **43**:103-114
40. Stomakhin A, Howes R, Schroeder C, Teran JM (2012) Energetically consistent invertible elasticity. In: Proceedings of the 11th ACM SIGGRAPH/Eurographics conference on Computer Animation. pp. 25-32
41. Tabata H, Nakajima K. (2001) Efficient in utero gene transfer system to the developing mouse brain using electroporation: visualization of neuronal migration in the developing cortex. *Neuroscience* **103**:865-872
42. Tallinen T, Biggins JS (2015) Mechanics of invagination and folding: Hybridized instabilities when one soft tissue grows on another. *Phys Rev E* **92**:022720

43. Tallinen T, Biggins JS, Mahadevan L. (2013) Surface sulci in squeezed soft solids. *Phys Rev Lett* **110**:024302
44. Tallinen T, Chung JY, Biggins JS, Mahadevan L. (2014) Gyrification from constrained cortical expansion. *Proc Natl Acad Sci* **111**:12667-12672
45. Tallinen T, Chung JY, Rousseau F, Girard N, Lefèvre J, Mahadevan L. (2016) On the growth and form of cortical convolutions. *Nat Phys* **12**:588-593
46. Toro R, Burnod Y. (2005) A morphogenetic model for the development of cortical convolutions. *Cereb Cortex* **15**:1900-1913
47. Toro R, Perron M, Pike B, Richer L, Veillette S, Pausova Z, Paus T. (2008) Brain size and folding of the human cerebral cortex. *Cereb Cortex* **18**:2352-2357
48. Toro R, Tiesinga P, Delzescaux T, Evans A, et al. (2018) FIIND: Ferret Interactive Integrated Neuro Development Atlas. <https://neuroanatomy.github.io/fiind/>
49. Van Essen DC (2022) Elsevier. Biomechanical models and mechanisms of cellular morphogenesis and cerebral cortical expansion and folding. *Semin Cell Dev Biol* **140**:90-104
50. Welker W. (1990) Why does the cortex fissure and fold: a review of determinants of gyri and sulci. In: Jones EG, Peters A (Eds). *Cerebral Cortex* New York, NY: Plenum Press. pp. 3-136
51. Yin S, Liu C, Choi GPT, Jung Y, Toro R, Heuer K, Mahadevan L. (2025) Morphogenesis and morphometry of brain folding patterns across species. *eLife* **14**:RP107138 <https://doi.org/10.7554/eLife.107138>
1. Toro R., Bakker R., Delzescaux T., Evans A., Tiesinga P. (2018) FIIND: Ferret interactive integrated neurodevelopment atlas. *Res. Ideas Outcomes* **4**:e25312
2. (2018) FIIND: Ferret interactive integrated neuro development atlas. <https://neuroanatomy.github.io/fiind/>
3. Abraham A., Pedregosa F., Eickenberg M., Gervais P., Mueller A., Kossaifi J., Gramfort A., Thirion B., Varoquaux G. (2014) Machine learning for neuroimaging with scikit-learn. *Front. Neuroinform* **8**:14
4. Toro R. (2017) "BrainBox github repository.". <https://github.com/r03ert0/BrainBox>
5. Tallinen T., Chung J. Y., Biggins J. S., Mahadevan L. (2014) Gyrification from constrained cortical expansion. *Proc. Natl. Acad. Sci* **111**, no. 35, pp. 12667–12672
6. Tallinen T., Chung J. Y., Rousseau F., Girard N., Mahadevan L. (2016) On the growth and form of cortical convolutions. *Nat. Phys* **12**:588-593
7. (2019) "NetGen.". <https://ngsolve.org/>
8. Stomakhin A., Howes R., Schroeder C., Teran J. M. (2012) Energetically consistent invertible elasticity. In: Proceedings of the 11th ACM SIGGRAPH/Eurographics Conference on Computer Animation. pp. 25-32
9. Bonet J., Burton A. (1998) A simple average nodal pressure tetrahedral element for incompressible and nearly incompressible dynamic explicit applications. *Comm. Numer. Meth. Eng* **14**:437-449
10. Choi P. T., Lam K. C., Lui L. M. (2015) FLASH: Fast landmark aligned spherical harmonic parameterization for genus-0 closed brain surfaces. *SIAM J. Imaging Sci* **8**:67-94
11. Koenderink J. J., van Doorn A. J. (1992) Surface shape and curvature scales. *Image Vis. Comput* **10**:557-564
12. Yin S., Liu C., Choi G. P. T., Jung Y., Toro R., Heuer K., Mahadevan L. (2025) Morphogenesis and morphometry of brain folding patterns across species. *eLife* **14**:RP107138 <https://doi.org/10.7554/eLife.107138>
13. Müller C. (2006) *Spherical harmonics* **17** Springer.
14. Brechbühler C., Gerig G., Kußbier O. (1995) Parametrization of closed surfaces for 3-D shape description. *Comput. Vis. Image Underst* **61**:154-170

15. Smith R. S., Kenny C. J., Ganesh V., Jang A., Borges-Monroy R., Partlow J. N., Hill R. S., Shin T., Chen A. Y., Doan R. N., *et al.* (2018) Sodium channel SCN3A (Nav1. 3) regulation of human cerebral cortical folding and oral motor development. *Neuron* **99**:905-913

Peer reviews

Reviewer #1 (Public review):

The manuscript by Choi and colleagues investigates the impact of variation in cortical geometry and growth on cortical surface morphology. Specifically, the study uses physical gel models and computational models to evaluate the impact of varying specific features/parameters of the cortical surface. The study makes use of this approach to address the topic of malformations of cortical development and finds that cortical thickness and cortical expansion rate are the drivers of differences in morphogenesis.

The study is composed of two main sections. First, the authors validate numerical simulation and gel model approaches against real cortical postnatal development in the ferret. Next, the study turns to modelling malformations in cortical development using modified tangential growth rate and cortical thickness parameters in numerical simulations. The findings investigate three genetically linked cortical malformations observed in the human brain to demonstrate the impact of the two physical parameters on folding in the ferret brain.

This is a tightly presented study that demonstrates a key insight into cortical morphogenesis and the impact of deviations from normal development. The dual physical and computational modeling approach offers the potential for unique insights into mechanisms driving malformations. This study establishes a strong foundation for further work directly probing the development of cortical folding in the ferret brain.

<https://doi.org/10.7554/eLife.107141.2.sa2>

Reviewer #2 (Public review):

Summary:

Based on MRI data of the ferret (a gyrencephalic non-primate animal, in whom folding happens postnatally), the authors create in vitro physical gel models and in silico numerical simulations of typical cortical gyrification. They then use genetic manipulations of animal models to demonstrate that cortical thickness and expansion rate are primary drivers of atypical morphogenesis. These observations are then used to explain cortical malformations in humans.

Strengths:

The paper is very interesting and original, and combines physical gel experiments, numerical simulations, as well as observations in MCD. The figures are informative, and the results appear to have good overall face validity.

Comment on the revised version from the Reviewing Editor:

The reviewers are happy with the authors replies and the eLife Assessment has been amended accordingly.

<https://doi.org/10.7554/eLife.107141.2.sa1>

Reviewer #1 (Public review):

The manuscript by Choi and colleagues investigates the impact of variation in cortical geometry and growth on cortical surface morphology. Specifically, the study uses physical gel models and computational models to evaluate the impact of varying specific features/parameters of the cortical surface. The study makes use of this approach to address the topic of malformations of cortical development and finds that cortical thickness and cortical expansion rate are the drivers of differences in morphogenesis.

The study is composed of two main sections. First, the authors validate numerical simulation and gel model approaches against real cortical postnatal development in the ferret. Next, the study turns to modelling malformations in cortical development using modified tangential growth rate and cortical thickness parameters in numerical simulations. The findings investigate three genetically linked cortical malformations observed in the human brain to demonstrate the impact of the two physical parameters on folding in the ferret brain.

This is a tightly presented study that demonstrates a key insight into cortical morphogenesis and the impact of deviations from normal development. The dual physical and computational modeling approach offers the potential for unique insights into mechanisms driving malformations. This study establishes a strong foundation for further work directly probing the development of cortical folding in the ferret brain. One weakness of the current study is that the interpretation of the results in the context of human cortical development is at present indirect, as the modelling results are solely derived from the ferret. However, these modelling approaches demonstrate proof of concept for investigating related alterations more directly in future work through similar approaches to models of the human cerebral cortex.

We thank the reviewer for the very positive comments. While the current gel and organismal experiments focus on the ferret only, we want to emphasize that our analysis does consider previous observations of human brains and morphologies therein (Tallinen et al., Proc. Natl. Acad. Sci. 2014; Tallinen et al., Nat. Phys. 2016), which we compare and explain. This allows us to analyze the implications of our study broadly to understand the explanations of cortical malformations in humans using the ferret to motivate our study. Further analysis of normal human brain growth using computational and physical gel models can be found in our companion paper (Yin et al., 2025), now also published to eLife: S. Yin, C. Liu, G. P. T. Choi, Y. Jung, K. Heuer, R. Toro, L. Mahadevan, Morphogenesis and morphometry of brain folding patterns across species. eLife, 14, RP107138, 2025. doi:10.7554/eLife.107138

In future work, we plan to obtain malformed human cortical surface data, which would allow us to further investigate related alterations more directly. We have added a remark on this in the revised manuscript (please see page 8–9).

Reviewer 2 (Public review):
Summary:

Based on MRI data of the ferret (a gyrencephalic non-primate animal, in whom folding happens postnatally), the authors create in vitro physical gel models and in silico numerical simulations of typical cortical gyrification. They then use genetic manipulations of animal models to demonstrate that cortical thickness and expansion rate are primary drivers of atypical morphogenesis. These observations are then used to explain cortical malformations in humans.

Strengths:

The paper is very interesting and original, and combines physical gel experiments, numerical simulations, as well as observations in MCD. The figures are informative, and the results appear to have good overall face validity.

We thank the reviewer for the very positive comments.

Weaknesses:

On the other hand, I perceived some lack of quantitative analyses in the different experiments, and currently, there seems to be rather a visual/qualitative interpretation of the different processes and their similarities/differences. Ideally, the authors also quantify local/pointwise surface expansion in the physical and simulation experiments, to more directly compare these processes. Time courses of eg, cortical curvature changes, could also be plotted and compared for those experiments. I had a similar impression about the comparisons between simulation results and human MRI data. Again, face validity appears high, but the comparison appeared mainly qualitative.

We thank the reviewer for the comments. Besides the visual and qualitative comparisons between the models, we would like to point out that we have included the quantification of the shape difference between the real and simulated ferret brain models via spherical parameterization and the curvature-based shape index as detailed in main text Fig. 4 and SI Section 3. We have also utilized spherical harmonics representations for the comparison between the real and simulated ferret brains at different maximum order N . In our revision, we have included more calculations for the comparison between the real and simulated ferret brains at more time points in the SI (please see SI page 6). As for the comparison between the malformation simulation results and human MRI data in the current work, since the human MRI data are two-dimensional while our computational models are three-dimensional, we focus on the qualitative comparison between them. In future work, we plan to obtain malformed human cortical surface data, from which we can then perform the parameterization-based and curvature-based shape analysis for a more quantitative assessment.

I felt that MCDs could have been better contextualized in the introduction.

We thank the reviewer for the comment. In our revision, we have revised the description of MCDs in the introduction (please see page 2).

Reviewer #1 (Recommendations for the authors):

The study is beautifully presented and offers an excellent complement to the work presented by Yin et al. In its current form, the malformation portion of the study appears predominantly reliant on the numerical simulations rather than the gel model. It might be helpful, therefore, to further incorporate the results presented in Figure S5 into the main text, as this seems to be a clear application of the physical gel model to modelling malformations. Any additional use of the gel models in the malformation portion of the study would help to further justify the necessity and complementarity of the dual methodological approaches.

We thank the reviewer for the suggestion. We have moved Fig. S5 and the associated description to the main text in the revised manuscript (please see the newly added Figure 5 on page 6 and the description on page 5–7). In particular, we have included a new section on the physical gel and computational models for ferret cortical malformations right before the section on the neurology of ferret and human cortical malformations.

One additional consideration is that the analyses in the current study focus entirely on the ferret cortex. Given the emphasis in the title on the human brain, it may be

worthwhile to either consider adding additional modelling of the human cortex or to consider modifying the title to more accurately align with the focus of the methods/results.

We thank the reviewer for the suggestion. While the current gel and organismal experiments focus on the ferret only, we want to emphasize that our analysis does consider previous observations of human brains and morphologies therein (Tallinen et al., Proc. Natl. Acad. Sci. 2014; Tallinen et al., Nat. Phys. 2016), which we compare and explain. This allows us to analyze the implications of our study broadly to understand the explanations of cortical malformations in humans using the ferret to motivate our study. Therefore, we think that the title of the paper seems reasonable. To further highlight the connection between the ferret brain simulations and human brain growth, we have included an additional comparison between human brain surface reconstructions adapted from a prior study and the ferret simulation results in the SI (please see SI Section S4 and SI Fig. S5 on page 9–10).

Two additional minor points:

Table S1 seems sufficiently critical to the motivation for the study and organization of the results section to justify inclusion in the main text. Of course, I would leave any such minor changes to the discretion of the authors.

We thank the reviewer for the suggestion. We have moved Table S1 and the associated description to the main text in the revised manuscript (please see Table 1 on page 7).

Page 7, Column 1: “macacques” → “macaques”.

We thank the reviewer for pointing out the typo. We have fixed it in the revised manuscript (please see page 8).

Reviewer #2 (Recommendations for the authors):

The methods lack details on the human MRI data and patients.

We thank the reviewer for the comment. Note that the human MRI data and patients were from prior works (Smith et al., Neuron 2018; Johnson et al., Nature 2018; Akula et al., Proc. Natl. Acad. Sci. 2023) and were used for the discussion on cortical malformations in Fig. 6. In the revision, we have included a new subsection in the Methods section and provided more details and references of the MRI data and patients (please see page 9–10).

<https://doi.org/10.7554/eLife.107141.2.sa0>