

A split-GAL4 driver line resource for *Drosophila* neuron types

Reviewed Preprint

v2 • January 6, 2025

Revised by authors

Reviewed Preprint

v1 • July 30, 2024

Geoffrey W Meissner , Allison Vannan, Jennifer Jeter, Kari Close, Gina M DePasquale, Zachary Dorman, Kaitlyn Forster, Jaye Anne Beringer, Theresa V Gibney, Joanna H Hausenfluck, Yisheng He, Kristin Henderson, Lauren Johnson, Rebecca M Johnston, Gudrun Ihrke, Nirmala Iyer, Rachel Lazarus, Kelley Lee, Hsing-Hsi Li, Hua-Peng Liaw, Brian Melton, Scott Miller, Reeham Motaher, Alexandra Novak, Omotara Ogundeyi, Alyson Petruncio, Jacquelyn Price, Sophia Protopapas, Susana Tae, Jennifer Taylor, Rebecca Vorimo, Brianna Yarbrough, Kevin Xiankun Zeng, Christopher T Zugates, Heather Dionne, Claire Angstadt, Kelly Ashley, Amanda Cavallaro, Tam Dang, Guillermo A Gonzalez III, Karen L Hibbard, Cuizhen Huang, Jui-Chun Kao, Todd Laverty, Monti Mercer, Brenda Perez, Scarlett Pitts, Danielle Ruiz, Viruthika Vallanadu, Grace Zhiyu Zheng, Cristian Goina, Hideo Otsuna, Konrad Rokicki, Robert R Svirskas, Han SJ Cheong, Michael-John Dolan, Erica Ehrhardt, Kai Feng, Basel El Galfi, Jens Goldammer, Stephen J Huston, Nan Hu, Masayoshi Ito, Claire McKellar, Ryo Minegishi, Shigehiro Namiki, Aljoscha Nern, Catherine E Schretter, Gabriella R Sterne, Lalanti Venkatasubramanian, Kaiyu Wang, Tanya Wolff, Ming Wu, Reed George, Oz Malkesman, Yoshinori Aso , Gwyneth M Card , Barry J Dickson , Wyatt Korff , Kei Ito , James W Truman , Marta Zlatic , Gerald M Rubin , FlyLight Project Team

Institute of Zoology, University of Cologne, Cologne, Germany • Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, United States • Queensland Brain Institute, University of Queensland, Brisbane, Australia • Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, United States • Department of Cell & Molecular Biology, University of California, Berkeley, Berkeley, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **valuable** study presents a resource for researchers using *Drosophila* to study neural circuits, in the form of a collection of split-Gal4 lines with an online search engine, which will facilitate the mapping of neuronal circuits. The evidence is **convincing** to demonstrate the utility of these new tools, and of the search engine, for understanding expression patterns in adults and larvae, and differences between the sexes. These resources will be of broad interest to *Drosophila* researchers in the field of neurobiology.

<https://doi.org/10.7554/eLife.98405.2.sa3>

Abstract

Techniques that enable precise manipulations of subsets of neurons in the fly central nervous system have greatly facilitated our understanding of the neural basis of behavior. Split-GAL4 driver lines allow specific targeting of cell types in *Drosophila melanogaster* and other species. We describe here a collection of 3060 lines targeting a range of cell types in the adult *Drosophila* central nervous system and 1373 lines characterized in third-instar larvae. These tools enable functional, transcriptomic, and proteomic studies based on precise anatomical targeting. NeuronBridge and other search tools relate light microscopy images of these split-

GAL4 lines to connectomes reconstructed from electron microscopy images. The collections are the result of screening over 77,000 split hemidriver combinations. Previously published and new lines are included, all validated for driver expression and curated for optimal cell type specificity across diverse cell types. In addition to images and fly stocks for these well-characterized lines, we make available 300,000 new 3D images of other split-GAL4 lines.

Introduction

The ability to manipulate small subsets of neurons is critical to many of the experimental approaches used to study neuronal circuits. In *Drosophila*, researchers have generated genetic lines that express an exogenous transcription factor, primarily GAL4, in a subset of neurons (Griffith 2012 [↗](#); Venken et al., 2011 [↗](#)). The GAL4 protein then drives expression of indicator or effector genes carried in a separate UAS transgenic construct (Fischer et al., 1988 [↗](#); Brand & Perrimon, 1993 [↗](#)). This modular approach has proven to be very powerful but depends on generating collections of lines with reproducible GAL4 expression limited to different, specific subsets of cells. In the 1990s, so-called enhancer trap lines were the method of choice (Bellen et al., 1989 [↗](#)). In this method, a GAL4 gene that lacks its own upstream control elements is inserted as part of a transposable element into different genome locations where its expression might come under the control of nearby endogenous regulatory elements. However, the resultant patterns were generally broad, with expression in hundreds of cells, limiting their use for manipulating specific neuronal cell types (Manseau et al., 1997 [↗](#); Yoshihara & Ito, 2000 [↗](#); Ito et al., 2003 [↗](#)).

The FlyLight Project Team (<https://www.janelia.org/project-team/flylight> [↗](#)) was started to address this limitation with the overall goal of generating a large collection of GAL4 lines that each drove expression in a distinct small subset of neurons—ideally individual cell types. Many labs studying the upstream regulatory elements of individual genes in the 1980s and 1990s had observed that short segments of DNA located upstream of protein coding regions or in introns, when assayed for enhancer function, frequently drove expression in small, reproducible subsets of cells (see Levine and Tjian 2003 [↗](#)). Pfeiffer et al. (2008) [↗](#) developed an efficient strategy for scaling up such assays of individual DNA fragments and showed that a high percentage of 2-3 kb genomic fragments, when cloned upstream of a core promoter driving GAL4 expression, produced distinct patterns of expression. Importantly, these patterns were much sparser than those observed with enhancer traps (Pfeiffer et al., 2008 [↗](#)). The approach also took advantage of newly developed methods for site specific integration of transgenes into the genome (Groth et al. 2004 [↗](#)), which facilitated the ability to compare constructs by placing them in the same genomic context. FlyLight was established in 2009 to scale up this approach. In 2012, the project reported the expression patterns produced by 6,650 different genomic segments in the adult brain and ventral nerve cord (Jenett et al., 2012 [↗](#)) and later in the larval central nervous system (CNS; Li et al., 2014 [↗](#)). In the adult central brain, we estimated that this ‘Generation 1’ (Gen1) collection contained 3,850 lines in which the number of labeled central-brain neurons was in the range of 20 to 5,000. These GAL4 driver lines and a collection of similarly constructed LexA lines have been widely used by hundreds of laboratories. However, we concluded that less than one percent of our lines had expression in only a single cell type, highlighting the need for a better approach to generating cell-type-specific driver lines.

To gain more specific expression the project turned to intersectional methods. These methods require two different enhancers to be active in a cell to observe expression of a functional GAL4 transcriptional activator in that cell. We adopted the split-GAL4 approach that was developed by Luan et al. (2008) and subsequently optimized by Pfeiffer et al. (2010) [↗](#) in which an enhancer drives either the activation domain (AD) or the DNA binding domain (DBD) of GAL4 (or optimized alternatives) in separate proteins. When present in the same cell the proteins carrying the AD and DBD domains, each inactive in isolation, dimerize to form a functional GAL4 transcription factor.

The work of Jenett et al. (2012) [described](#) the expression patterns of thousands of enhancers. Using these data, anatomical experts could identify Gen1 enhancers that express in the cell type of interest and cross flies that express the AD or DBD half of GAL4 under the control of two of the same enhancers and observe the resultant intersected expression pattern. Large collections of such AD or DBD genetic drivers, which we refer to as hemidriver lines, were generated at Janelia (Dionne et al., 2017) and at the IMP in Vienna (Tirian et al., 2017). In ~5-10% of such crosses, the cell type of interest was still observed, but now as part of a much sparser expression pattern than displayed by either of the initial enhancers. In about 1-2% of these genetic intersections, expression appeared to be limited to a single cell type.

At Janelia, early efforts were directed at targeting neuronal populations in the optic lobes (Tuthill et al., 2013 [described](#); Nern et al., 2015 [described](#); Wu et al., 2016 [described](#)) and mushroom bodies (Aso et al., 2014 [described](#)). We were encouraged by the fact that we were able to generate lines specific for the majority of cell types in these populations. With the involvement of additional collaborating groups, the project was extended to several other CNS regions (**Table 1** [described](#)). In our initial studies we relied on expert human annotators performing extensive visual surveys of expression data to identify candidate enhancers to intersect. More recently, two advances have greatly facilitated this process. First, we have developed computational approaches (Otsuna et al., 2018 [described](#); Hirsh et al., 2020; Mais et al., 2021 [described](#); Meissner et al., 2023 [described](#)) to search databases of neuronal morphologies generated by stochastic labeling (Nern et al., 2015 [described](#)) of several thousand of the Jenett et al. (2012) [described](#) and Tirian et al. (2017) GAL4 lines. Second, electron microscopy (EM) datasets (Scheffer et al., 2020 [described](#); Cheong et al., 2023 [described](#); Marin et al., 2023 [described](#); Takemura et al., 2023 [described](#); Nern et al., 2024 [described](#); Schlegel et al., 2024 [described](#)) have provided comprehensive cell type inventories for many brain regions.

In this report, we summarize the results obtained over the past decade and present a collection of the best cell-type-specific split-GAL4 lines that were identified.

Results

Cell-type-specific split-GAL4 line collection

We describe here a collection of 3060 split-GAL4 lines targeting cell types across the adult *Drosophila* CNS. All lines were created in collaborations with the FlyLight Project Team at Janelia Research Campus from 2013 to 2023, based on examination of over 77,000 split combinations. 1644 lines were published previously and are drawn together here as part of the larger collection. **Table 1** [described](#) summarizes prior publications. The remaining 1416 lines in the collection are described in other in preparation manuscripts or newly reported here as shown in Figure 1 Supplemental File 1, which specifies proper citations and other line-specific information.

To confirm and consistently document the expression patterns of included lines, all were rescreened by crossing to a UAS-CsChrimson-mVenus reporter inserted in at a specific genomic location (attP18) (**Figure 1** [described](#); Klapoetke et al., 2014). At least one male and one female CNS were dissected, antibody labeled, and imaged per line. Expression patterns were validated by manual qualitative comparison to prior data, where available, and scored for specificity and consistency (see below and Methods). Male/female differences were observed in 109 of these 3060 lines, confirming previously reported sexual dimorphisms or suggesting potential areas for further study (**Figure 1F** [described](#); Figure 1 Supplemental File 1).

Previously published and newer lines were evaluated as a whole, taking advantage of new screening data to highlight existing lines or identify new lines that best label cell types. The lines were selected for inclusion based on several factors:

Publication DOI	First author(s)	Year	Split-GAL4 lines	Anatomical region/cell types for lines
10.1016/j.neuron.2013.05.024	Tuthill, Nern	2013	22	Lamina (optic lobe)
10.1016/j.neuron.2014.05.017	Feng, Palfreyman	2014	1	SAG ascending neurons
10.7554/eLife.04577	Aso	2014	63	Mushroom body
10.7554/eLife.16135	Aso	2016	2	Mushroom body
10.7554/eLife.21022	Wu, Nern	2016	56	Lobula columnar neurons (optic lobe)
10.1016/j.neuron.2017.03.010	Strother	2017	9	T4 neurons & inputs (optic lobe)
10.1016/j.neuron.2017.05.036	von Reyn	2017	1	Lobula columnar neuron LC4 (optic lobe)
10.1038/nature24626	Klapoetke	2017	3	LPLC2 neurons & inputs (optic lobe)
10.7554/eLife.24394	Takemura	2017	1	CT1 neurons (optic lobe)
10.1002/cne.24512	Wolff	2018	46	Central complex
10.7554/eLife.34272	Namiki	2018	137	Descending neurons
10.1016/j.cub.2019.01.009	Jovanic	2019	2	Larval anemotaxis & adult descending neuron
10.7554/eLife.43079	Dolan	2019	2	Lateral horn
10.1016/j.cub.2020.07.083	Wang, Wang	2020	5	Descending neurons DNp13
10.1016/j.neuron.2020.08.006	Turner-Evans	2020	2	Central complex
10.1038/s41467-020-19936-x	Feng	2020	80	Leg motor MDN targets
10.1038/s41586-020-2055-9	Wang, Wang	2020	9	Descending neurons oviDN
10.1371/journal.pone.0236495	Bogovic	2020	1	Brain
10.7554/eLife.50901	Davis, Nern	2020	40	Optic lobe
10.7554/eLife.57685	Morimoto	2020	10	Central brain targets of lobula LC6 neurons
10.7554/eLife.58942	Schretter	2020	12	Central brain pC1d aIPg
10.1038/s41467-023-43566-8	Longden	2021	1	L1 neurons (optic lobe)
10.1038/s41586-020-2972-7	Wang, Wang	2021	7	Descending neurons vpoDN vpoEN
10.1101/2021.07.23.453511	Mais	2021	1	Central brain pC1e
10.7554/eLife.66039	Hulse, Haberkern, Franconville, Turner-Evans	2021	1	Central complex
10.7554/eLife.71679	Sterne	2021	67	Subesophageal zone
10.7554/eLife.71858	Kind, Longden, Nern, Zhao	2021	3	R7 & R8 photoreceptor targets (optic lobe)
10.1016/j.cub.2022.01.008	Namiki, Ros	2022	9	Descending neurons flight
10.1016/j.cub.2022.06.019	Baker	2022	28	Central brain AMMC, WED, AVLP, PVLP
10.1016/j.neuron.2022.02.013	Klapoetke	2022	2	Lobula LC18 & LC25 neurons (optic lobe)
10.1101/2022.12.14.520178	Zhao	2022	1	H2 neurons (optic lobe)
10.1038/s41586-023-06271-6	Vijayan	2023	2	Descending neurons oviDN
10.1101/2023.05.31.542897	Ehrhardt, Whitehead	2023	164	Dorsal VNC
10.1101/2023.06.07.543976	Cheong, Eichler, Stuermer	2023	11	VNC premotor
10.1101/2023.06.21.546024	Isaacson	2023	6	Lobula plate LPC & LLPC (optic lobe)
10.1101/2023.10.16.562634	Zhao	2023	2	MeLp2 & LPi4b neurons (optic lobe)
10.1101/2023.11.29.569241	Garner, Kind	2023	5	Medulla to AOTU (MeTu) neurons (optic lobe)
10.25378/janelia.23726103	Minegishi	2023	52	Ascending neurons
10.1016/j.cub.2024.01.015	Lillis	2024	22	VNC song generation
10.1016/j.cub.2024.01.071	Cheong, Boone, Bennett	2024	1	Ascending neurons flight
10.1038/s41586-024-07222-5	Gorko	2024	3	Neck motor neurons
10.1101/2024.03.15.585289	Schretter	2024	3	Aggression circuit
10.1101/2024.04.16.589741	Nern	2024	320	Optic lobe
10.1101/2024.10.21.619448	Wolff	2024	240	Central complex
10.7554/eLife.90523	Rubin	2024	21	Mushroom body output neurons
10.7554/eLife.94168	Shuai	2024	168	Mushroom body
Dionne et al., in prep	Dionne		81	Accessory medulla (optic lobe) and clock
Rubin et al. in prep	Rubin		50	Anterior optic tubercle
This paper			1285	
Total			3060	

Table 1.

Publications reporting split-GAL4 lines from the adult cell-type-specific collection.

Publications are listed by year. The number of lines from the collection in each publication is listed, along with the CNS regions and/or cell types most commonly labeled. Many of these publications describe additional lines that were not included in the collection described here.




Figure 1.

Example cell-type-specific lines.

(A) Split-GAL4 line SS52577 is expressed in P-FNv neurons arborizing in the protocerebral bridge, fan-shaped body, and nodulus (Wolff & Rubin, 2018 [DOI](#)).

(B) Split-GAL4 line MB043C is expressed primarily in PAM- α 1 dopaminergic neurons that mediate reinforcement signals of nutritional value to induce stable olfactory memory for driving wind-directed locomotion and higher-order learning (Aso et al., 2014 [DOI](#); Ichinose et al., 2015 [DOI](#); Aso & Rubin, 2016 [DOI](#); Aso et al., 2023 [DOI](#); Yamada et al., 2023 [DOI](#)).

(C) Split-GAL4 line SS40265 is expressed in members of the 8B(t1) cluster of cholinergic neurons that connect the lower tectulum neuropil of the prothorax with the gnathal neuropil and the ventral most border of the vest neuropil of the brain ventral complex.

(D) Split-GAL4 line SS60203 is expressed in ascending neurons likely innervating the wing neuropil.

(E) Split-GAL4 line SS47938 is expressed in LBL40, mediating backwards walking (Feng et al., 2020 [DOI](#); same sample used in Fig 5b, CC-BY license).

(F) Split-GAL4 line SS36564 is expressed in female-specific aIPg neurons (F1) and not observed in males (F2; Schretter et al., 2020 [DOI](#)).

(G) MCFO of split-GAL4 line SS72207 with specific expression in DNg34, a cell type described in Namiki et al., 2018 [DOI](#).

Scale bars, 50 μ m. See Figure 1 Supplemental file 1 for more line information and Figure 1 Supplemental file 2 for images of all cell-type-specific lines.

1. Diversity: Where cell type information was available, especially from comparison to EM volumes (Scheffer et al., 2020 [↗](#); Cheong et al., 2023 [↗](#); Marin et al., 2023 [↗](#); Takemura et al., 2023 [↗](#); Nern et al., 2024 [↗](#)), we generally limited each cell type to the two best split-GAL4 lines, so as to cover a wider range of cell types.
2. Specificity: 1767 lines were scored as highest quality, well suited to activation-based behavioral studies, with strong and consistent labeling of a single identified cell type and minimal detected off-target expression (**Figure 2A** [↗](#); Figure 1 Supplemental File 1). 1258 lines showed spatially segregated off-target expression that doesn't interfere with neuron visualization for anatomy or physiology (**Figure 2B-C** [↗](#); Figure 1 Supplemental File 1). A control line with minimal detected expression was also included (SS01062; Namiki et al., 2018 [↗](#)).
3. Consistency: 34 lines were very specific but showed weaker or less consistent expression (**Figure 2D** [↗](#); **Figure 2 — figure supplement 1** [↗](#); Figure 1 Supplemental File 1). These lines reveal anatomy but may be challenging to use for manipulations without examination of expression in each individual fly.
4. Regions of interest: Lines were generated in collaboration with Janelia labs and collaborators studying particular CNS regions or classes of neurons. While this collection includes many high-quality lines across the CNS, most efforts were directed to regions of interest within the CNS (see below and **Table 1** [↗](#)).

We examined the distribution of the cell type collection across the male and female CNS. To visualize the distribution, we aligned each image to a unisex template and segmented neuron patterns from rescreening images into a heat map (**Figure 3** [↗](#)). The cell type lines show neuronal expression across most of the CNS, with 96% of voxels having expression from at least 5 lines, 90% with 10 or more lines, and 66% with 20 or more lines (Figure 3 Supplemental File 1). Prominently labeled areas include the fan-shaped body, lobula, and superior medial protocerebrum in the brain, and the T1 medial ventral association center and intermediate tectulum in the ventral nerve cord (VNC). The antennal lobe is very rarely labeled. Other relatively rarely labeled areas include the anterior ventrolateral protocerebrum and prow in the brain and the ovoid (accessory mesothoracic neuropil) in the VNC.

We compared the line distribution between female and male images (**Figure 3C-D** [↗](#)). We observed more female split-GAL4 expression in the epaulette and abdominal ganglion, and more male expression in the antennal mechanosensory and motor center and several regions of the mushroom body (Figure 3 Supplemental File 1). Two example sexually dimorphic split-GAL4 lines from the cell type collection illustrate regions of increased split-GAL4 expression in females (**Figures 3E-H** [↗](#)). The lines were identified based on NeuronBridge searches of the 'female minus male' image (Clements et al., 2021 [↗](#); Clements et al., 2024 [↗](#)). Examination of individual male images suggests that (in contrast to the female) the subtraction approach highlighted male artifacts of the reporter system: either background incompletely segmented away from neurons (e.g. in the metathoracic ganglion) or repetitive neuron signal unrelated to other line-specific expression ('sparse-T' neuron and projections into the AMMC; Meissner, et al., 2023 [↗](#)).

The collection of lines described above has been deposited with Bloomington Drosophila Stock Center (Bloomington, IN) for availability until at least August 31, 2026. Images and line metadata are available at <https://splitgal4.janelia.org> [↗](#). Anatomical searching for comparison to other light microscopy (LM) and EM data is available at <https://neuronbridge.janelia.org> [↗](#).

Raw image data collection and analysis workflow

In addition to the line collection described above, we describe a split-GAL4 image data resource. It consists of all good quality FlyLight split-GAL4 images and associated metadata generated in the project's first nine years of split-GAL4 characterization (Figure 1 Supplemental File 4). It includes

Figure 2.

Examples of line quality levels.

The 3060 cell type lines were scored for expression.

(A) Quality level 1 (1767 lines): Split-GAL4 line OL0015B (Wu, Nern, et al., 2016) is specifically and strongly expressed in a single cell type. Occasional weak expression may be seen in other cells.

(B) Quality level 2 (1232 lines): Split-GAL4 line SS59643 (Wolff, et al., 2024) has expression in two cell types. Occasional weak expression may be seen in other cells.

(C) Quality level 3 (26 lines): Split-GAL4 line SS59643 (Wolff, et al., 2024) has expression in three or more cell types.

(D) Quality level 4 (34 lines): Split-GAL4 line SS61022 has specific expression but weak or variable labeling efficiency. See Figure 2 Supplement 1 for examples of variable expression.

(E) Quality level 5: IS36417 is an Initial Split combination not selected for stabilization. Groups of neurons are visible, but the cell type of interest was not labeled with sufficient specificity for further work. Such lines were only included in the raw image collection.

Scale bars, 50 μ m.





Figure 2 — figure supplement 1.

Example expression variability in Quality level 4 lines. Five samples from split-GAL4 line SS61022 show variable expression. (A-C) Male flies. (D-E) Female flies. Scale bars, 50 μ m.






Figure 3.

Spatial distribution of cell type lines.

(A-D) Images of one male and one female sample from 3029 cell type rescreening lines were aligned to JRC2018 Unisex (Bogovic et al., 2020), segmented from background (see Methods), binarized, overlaid, and maximum intensity projected, such that brightness indicates the number of lines with expression. All images were scaled uniformly to a maximum brightness equal to 206 lines on Fiji's "royal" LUT (scale inset in D). This saturated a small portion of the male antennal mechanosensory and motor center (AMMC) that reached a peak value of 260 lines per voxel, for the purpose of better visualizing the rest of the CNS.

(A) Female CNS.

(B) Male CNS.

(C) Female image stack minus male, then maximum intensity projected.

(D) Male image stack minus female, then maximum intensity projected.

(E-F) Split-GAL4 line SS56987 (Schretter, et al., 2020) is sex-specifically expressed in pC1d neurons that largely lie within the region of the female central brain highlighted in (C). Male and female images have different brightness scales.

(G-H) Split-GAL4 line SS35230 (Shuai, et al., 2024) is sex-specifically expressed in abdominal ganglion neurons that largely lie within the region of the female VNC highlighted in (C). Male and female images have different brightness scales. All scale bars, 50 μm .

additional data for the lines described above, together with data for many additional split-Gal4 combinations. Many of these represent lower quality lines that label multiple nearby cell types (**Figure 2D** and see below), but the image collection also includes additional high quality lines that were not chosen for stabilization or are currently not maintained as stable lines (for example, because of similarity to other lines in the collection).

FlyLight's analysis workflow consisted of multiple related image-generating pipelines (**Figure 4A-B**). Typically, we screened an "Initial Split" (IS) cross temporarily combining the two candidate split hemidriviers and reporter before building ("stabilizing") a genetically stable "Stable Split" (SS) line. IS and SS data with the same 5-6 digit code reflect the same combination of split hemidriviers. Some split-GAL4 lines use region-specific nomenclatures, with names prefixed by "MB" (mushroom body), "OL" (optic lobe), or "LH" (lateral horn), and in most cases label neurons within or nearby those regions. We examined about 77,000 IS crosses and after quick visual evaluation chose to image a fly CNS sample from about half of them. 36,033 such IS samples (individual flies) are included in this image collection (**Table 2**). About 10% of IS crosses were made into SS lines and imaged again, with 8,273 such lines included.

Further documentation of the full SS pattern at higher quality with varying levels of pre-synaptic labeling was generated by the Polarity pipeline (Aso et al., 2014; Sterne et al., 2021), followed by full 20x imaging and selected regions of interest imaging at 63x, with 37,409 such samples from 7,039 lines included. Single neuron stochastic labeling by MultiColor FlpOut (MCFO; Nern, et al., 2015) revealed single neuron morphology and any diversity latent within the full SS pattern, with 54,807 such samples from 7,679 lines included in the image collection.

In total the raw image collection consists of 46,653 IS/SS combinations, 129,665 samples (flies), 612,124 3D image stacks, and 4,371,364 secondary (processed) image outputs, together 192 TB in size. The IS data may be particularly valuable for work on understudied CNS regions, as it contains biological intersectional results (as opposed to computational predictions) that may have specific expression outside regions focused on in prior publications (**Table 1**).

Due to the size of the raw image collection, it has not undergone the same level of validation as our other image collections, and caution is recommended in interpreting the data therein. We believe it is nonetheless of overall good quality. The images are available at <https://flylight-raw.janelia.org> and are gradually being made searchable via NeuronBridge due to the size of the dataset. At the time of this version of record, the 36,000 IS images have been added, with the remaining data planned to be added by the end of 2024. For a further guide to interpreting the image data, see Figure 4 Supplemental file 1.

Larval split-GAL4 line collection

We include 982 split-GAL4 lines, 31 Generation 1 GAL4 lines, and 353 Generation 1 LexA lines selected for specific expression in the third-instar larval CNS (see Figure 1 Supplemental file 1). Seven more split-GAL4 lines were selected based on both larval and adult expression.

We selected larval lines based on the following criteria: 1) The projection pattern had to be sparse enough for individual neurons in a line to be identifiable; 2) The line had to be useful for imaging of neural activity or patch-clamp recording; and 3) the line had to be useful for establishing necessity or sufficiency of those neurons in behavioral paradigms. We thus selected lines with no more than 2 neurons per brain hemisphere or per subesophageal zone (SEZ) or VNC hemisegment. When there are more neurons in an expression pattern, overlap in projections makes it hard to uniquely identify these neurons. Also, interpreting neural manipulation results in behavioral experiments becomes much more challenging.

Figure 4.

Split-GAL4 workflow and FlyLight data release statistics




Table 2.

Summary of FlyLight adult fly line and image releases

Includes descriptions of [Jenett et al., 2012](#); [Dionne et al., 2018](#); [Tirian & Dickson, 2017](#); publications in [Table 1](#); and stock distribution by Bloomington Drosophila Stock Center and Janelia Fly Facility. Section with italic text is specific to publications from [Table 1](#) and this publication. Image counts are unique between categories, whereas line counts overlap extensively between categories. “Image tiles” are considered as unique 3D regions, with each MCFO tile captured using two LSM image stacks. Stock shipments count each shipment of each stock separately. Raw and processed images are available at <https://www.janelia.org/gal4-gen1>, <https://gen1mcfo.janelia.org>, <https://splitgal4.janelia.org>, and many can be searched based on anatomy at <https://neuronbridge.janelia.org>.

	Gen1 GAL4	Gen1 LexA	Gen1 MCFO	AD/ DBD	IS Screen	SS Screen	SS MCFO	SS Polarity	Total
Lines	9,389	1,500	5,155	8,882	35,708	8,986	7,823	7,050	67,562
Samples imaged	9,389	1,500	74,363	NA	36,182	16,974	55,702	37,526	231,636
20x image tiles	18,891	2,966	29,780	NA	72,346	33,924	66,484	61,787	286,178
40x image tiles	0	0	111,563	NA	0	0	0	0	111,563
63x image tiles	0	0	22,372	NA	0	0	77,039	49,117	148,528
Samples with 63x images	0	0	8,489	NA	0	0	31,849	17,619	57,957
Stocks currently available	2,871	1,781	NA	8,358	NA	3,013	NA	NA	16,023
Janelia & Bloomington stock shipments as of September 2024	167,082	29,227	NA	69,546	NA	40,067	NA	NA	305,922

If lines had sparse expression in the brain or SEZ but also had expression in the VNC (sparse or not), we nevertheless considered them useful for studying the brain or SEZ neurons because 1) the brain neurons could be identified; 2) the line could be used for imaging or patch-clamp recording in the brain; and 3) the split-GAL4 line could be cleaned up for behavioral studies, using teashirt-killer-zipper to eliminate VNC expression (Dolan et al., 2017 [DOI](#)). Similarly, lines with specific VNC expression but with undesired additional expression in the brain or SEZ were still useful for imaging and behavioral studies in the VNC, because a loss of phenotype following teashirt-killer-zipper could establish causal involvement of VNC neurons. For LexA lines, we included a few with 3 or more neurons per hemisphere or per hemisegment because there are fewer LexA lines overall and they could still be used for functional connectivity studies.

Images of all of the selected lines are available at Virtual Fly Brain (VFB; <https://raw.larval.flylight.virtualflybrain.org/> [DOI](#)) and will be further integrated into the main VFB website (<https://virtualflybrain.org/> [DOI](#)). We anticipate these lines will be useful tools in conjunction with electron microscopy volumes of the larval nervous system.

Discussion

This collection of cell-type-specific split-GAL4 lines and associated image resource is a toolkit for studies of many *Drosophila* neurons. The adult cell-type-specific line collection covers many identified cell types, and the images are being made instantly searchable for LM and EM comparisons at NeuronBridge, enabling selection of existing or new combinations of genetic tools for specific targeting based on anatomy. In total, the Janelia FlyLight Project Team has contributed 540,000 3D images of 230,000 GAL4, LexA, and split-GAL4 fly CNS samples. As of this publication, Janelia and Bloomington *Drosophila* Stock Center have distributed FlyLight stocks 300,000 times to thousands of groups in over 50 countries. We have also released standardized protocols for fly dissection and immunolabeling at the FlyLight protocols website <https://www.janelia.org/project-team/flylight/protocols> [DOI](#).

Several caveats should be kept in mind when using these tools and other transgenic systems. Many split-GAL4 lines still have some off-target expression (see **Figure 2** [DOI](#)). It is always good practice to validate results using multiple lines labeling the same neurons. In most cases, the off-target expression will not be present in multiple lines, supporting assigning a phenotype to the shared neurons. The less precise a set of lines, the more lines and other supporting evidence should be used. Different effectors and genomic insertion sites also vary in strength and specificity of expression and should be validated for each application (Pfeiffer et al., 2010 [DOI](#); Aso et al., 2014 [DOI](#)). For example, MCFO images can have brighter single neurons than in full split-GAL4 patterns. UAS-CsChrimson-mVenus allows for a direct correlation between labeling and manipulation that isn't available for every effector. See Figure 4 Supplemental File 1 for other examples of differences between effectors used in this study. If multiple transgenes are inserted into the same genomic locus, they can interact directly in unanticipated ways (transvection; Mellert & Truman, 2012 [DOI](#)). Fly age should also be controlled, as expression can vary over fly development (Harris et al., 2015 [DOI](#)), and effectors may take time to mature after expression (Baird et al., 2000 [DOI](#)).

Cell type consensus naturally evolves as new observational modalities are developed (e.g. connectomics and RNA sequencing) and comparisons are made across existing data, such as between EM volumes. Not all lines in this collection have cell type information, and some types will likely change. Different cell types also vary widely in their number of constituent neurons, and within a type there is often some variation between hemispheres and individuals. Thus, for any cell types with more than one neuron we cannot claim to have fully labeled every neuron of a type.

Despite extensive efforts, we have not developed specific lines for every cell type in the fly CNS. Moreover, the peripheral nervous system and the rest of the body were beyond the scope of this effort. Our hemidriviers likely do not effectively label some CNS cell types (discounting extremely broad expression), as they are based on Generation 1 collections that together may only label about half of all biological enhancers. As a rule of thumb, we estimate being able to use these tools to create precise split-GAL4 lines for one third of cell types, imprecise lines for another third, and no usable line for the remaining third of cell types. This hit rate nonetheless is often enough to characterize key circuit components. Continued iteration with the existing toolset is expected to lead to diminishing returns on improved CNS coverage.

An ideal toolkit with complete coverage of cell types with specific lines would require additional development. Capturing a full range of enhancers requires alternatives to our model of short genetic fragments inserted into a small number of genomic locations. Predicting enhancer combinations based on transcriptomic data and capturing them at their native location holds promise (e.g., Pavlou et al., 2016 [DOI](#); Chen et al., 2023 [DOI](#)), though high-quality transcriptomics data with cell type resolution are not yet available for much of the CNS. Insertions throughout the genome bring their own challenges of unpredictable side-effects, however, and specific targeting of every cell type remains a challenge. Further restriction by killer zipper, GAL80, Flp, or other intersectional techniques will likely remain an area of development (Pavlou et al., 2016 [DOI](#); Dolan et al., 2017 [DOI](#); Ewen-Campen et al., 2023 [DOI](#)).

The Janelia FlyLight Project Team has achieved its goal of developing tools to study the *Drosophila* nervous system. The team and collaborators have delivered GAL4/LexA lines labeling many neurons, single-neuron images enabling correlation with EM and the prediction of split-GAL4 lines, this collection of cell-type-specific split-GAL4 lines, an image resource of many more split combinations, and a series of focused studies of the nervous system.

Materials and methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (<i>Drosophila melanogaster</i>)	MCFO-1; hsPESTOP::attP3_3stop1_X_0036; (w, pBPhsFlp2::PEST in attP3;; pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFP-HA in VK00005, pJFRC240-10XUAS-FRT>STOP>FRT-myr::smGFP-V5-THS-10XUAS-FRT>STOP>FRT-myr::smGFPFLAG in su(Hw)attP1/TM3,Sb)	Nern et al., 2015	RRID:BDSC_64085 (Janelia stock 1117734)	
Genetic reagent (<i>Drosophila melanogaster</i>)	MCFO-2; pBPhsFLP_PEST_HAV5_FLAG_OLLAS_X3_0095; (w, pBPhsFlp2::PEST in attP3;; pJFRC210-10XUAS-FRT>STOP>FRT-myr::smGFP-OLLAS in attP2, pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFP-HA in VK00005, pJFRC240-10XUAS-FRT>STOP>FRT-myr::smGFP-V5-THS10XUAS-FRT>STOP>FRT-myr::smGFPFLAG in su(Hw)attP1/TM2)	Nern et al., 2015	RRID:BDSC_64086 (Janelia stock 3022015)	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (Drosophila melanogaster)	MCFO-4; 57C10wt_attp8_3stop1; (w, R57C10-Flp2 in su(Hw)attP8;; pJFRC201-10XUASFRT >STOP > FRT-myr::smGFP-HA in VK00005,pJFRC240-10XUAS- FRT>STOP > FRT-myr::smGFP-V5-THS-10XUASFRT >STOP > FRT-myr::smGFP-FLAG in su(Hw)attP1)	Nern et al., 2015	RRID:BDSC_64088 (Janelia stock 1116898)	
Genetic reagent (Drosophila melanogaster)	MCFO-5; 57C10PEST_attp8_3stop1; (w, R57C10-Flp2::PEST in su(Hw)attP8;; pJFRC201- 10XUAS-FRT>STOP > FRT-myr::smGFPHA in VK00005, pJFRC240-10XUAS-FRT>STOP > FRTmyr::smGFP- V5-THS-10XUAS-FRT>STOP > FRTmyr::smGFP- FLAG in su(Hw)attP1/TM2)	Nern et al., 2015	RRID:BDSC_64089 (Janelia stock 1116876)	
Genetic reagent (Drosophila melanogaster)	MCFO-6; 57C10L_attp8_4stop1; (w, R57C10-FlpL in su(Hw)attP8;; pJFRC210-10XUASFRT >STOP > FRT- myr::smGFP-OLLAS in attP2, pJFRC201-10XUAS- FRT>STOP > FRT-myr::smGFP-HA in VK00005, pJFRC240-10XUAS-FRT>STOP > FRT-myr::smGFP- V5-THS10XUAS-FRT>STOP > FRT-myr::smGFPFLAG in su(Hw)attP1/TM2)	Nern et al., 2015	RRID:BDSC_64090 (Janelia stock 1116894)	
Genetic reagent (Drosophila melanogaster)	MCFO-7; 57C10PEST_attp18_4stop1; (w, R57C10- Flp2::PEST in attP18;; pJFRC210-10XUASFRT >STOP > FRT-myr::smGFP-OLLAS in attP2, pJFRC201- 10XUAS-FRT>STOP > FRT-myr::smGFPHA in VK00005, pJFRC240-10XUAS-FRT>STOP > FRTmyr::smGFP-V5-THS-10XUAS-FRT>STOP > FRTmyr::smGFP-FLAG in su(Hw)attP1/TM2)	Nern et al., 2015	RRID:BDSC_64091 (Janelia stock 1116875)	
Genetic reagent (Drosophila melanogaster)	MCFO-3 derivative; 57C10L_brp_SNAP_MCFO_X23_0117; (w, R57C10-FlpL in su(Hw)attP8; brp::Snap / CyO; pJFRC201-10XUAS-FRT>STOP > FRT-myr::smGFPHA in VK00005,pJFRC240-10XUAS- FRT>STOP > FRTmyr::smGFP-V5-THS-10XUAS- FRT>STOP > FRTmyr::smGFP-FLAG in su(Hw)attP1/ TM6B)	Nern et al., 2015; Kohl et al., 2014	RRID:BDSC_64087 (Janelia stock 3023700)	
Genetic reagent (Drosophila melanogaster)	57C10PEST_brp_SNAP_MCFO_X23_0099; (w, R57C10- Flp2::PEST in attP18; brp::Snap / CyO; pJFRC201-10XUASFRT >STOP > FRT-myr::smGFP- HA in VK00005,pJFRC240- 10XUAS-FRT>STOP > FRT-myr::smGFP-V5-THS-10XUASFRT >STOP > FRT- myr::smGFP-FLAG in su(Hw)attP1/TM6B)	Nern et al., 2015	(Janelia stock 3023701)	
Genetic reagent (Drosophila melanogaster)	MCFO-1 derivative; pBPhsFlp2_PEST_brp_SNAP_MCFO_0128; (w, pBPhsFlp2::PEST in attP3; brp::Snap / CyO; pJFRC201- 10XUAS-FRT>STOP > FRT-myr::smGFPHA in VK00005,pJFRC240-10XUAS- FRT>STOP > FRTmyr::smGFP-V5-THS-10XUAS- FRT>STOP > FRTmyr::smGFP-FLAG in su(Hw)attP1/ TM6B)	Nern et al., 2015; Kohl et al., 2014	RRID:BDSC_64085 (Janelia stock 3023951)	
Genetic reagent (Drosophila melanogaster)	pJFRC2-10XUAS-IVS-mCD8::GFP	Pfeiffer et al., 2010	RRID:BDSC_32185 (Janelia stock 1115125)	
Genetic reagent (Drosophila melanogaster)	UAS_Chrimson_Venus_X_0070; (20XUAS-CsChrimson-mVenus trafficked in attP18)	Klapoetke et al., 2014	RRID:BDSC_55134 (Janelia stock 1150416)	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (Drosophila melanogaster)	UAS_CsChrimson_Venus_X_0107; (UAS-Syt-HA, 20XUAS-CsChrimson-mVenus trafficked in attP18)	Klapoetke et al., 2014; Robinson et al., 2002	(Janelia stock 3028408)	
Genetic reagent (Drosophila melanogaster)	hsPESTOPT_attP3_3stop1_X_0036; (MCFO-1; (w, pBPhsFlp2::PEST in attP3;; pJFRC201-10XUASFRT>STOP>FRT-myr::smGFP-HA in VK00005,pJFRC240-10XUAS-FRT>STOP>FRT-myr::smGFP-V5-THS-10XUASFRT>STOP>FRT-myr::smGFPFLAG in su(Hw)attP1/TM3,Sb))	Nern et al., 2015	RRID: BDSC_64085 (Janelia stock 1117734)	
Genetic reagent (Drosophila melanogaster)	57C10L_brp_SNAP_MCFO_X23_0117; (MCFO-3 derivative; (w, R57C10-FlpL in su(Hw)attP8; brp::Snap / CyO; pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFPHA in VK00005,pJFRC240-10XUAS-FRT>STOP>FRTmyr::smGFP-V5-THS-10XUAS-FRT>STOP>FRTmyr::smGFP-FLAG in su(Hw)attP1/TM6B))	Kohl et al., 2014; Nern et al., 2015	(Janelia stock 3023700)	
Genetic reagent (Drosophila melanogaster)	57C10PEST_attp8_3stop1; (MCFO-5; (w, R57C10-Flp2::PEST in su(Hw)attP8;; pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFPHA in VK00005, pJFRC240-10XUAS-FRT>STOP>FRTmyr::smGFP-V5-THS-10XUAS-FRT>STOP>FRTmyr::smGFP-FLAG in su(Hw)attP1/TM2))	Nern et al., 2015	RRID: BDSC_64089 (Janelia stock 1116876)	
Genetic reagent (Drosophila melanogaster)	57C10L_attp8_4stop1; (MCFO-6; (w, R57C10-FlpL in su(Hw)attP8;; pJFRC210-10XUASFRT>STOP>FRT-myr::smGFP-OLLAS in attP2, pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFP-HA in VK00005, pJFRC240-10XUAS-FRT>STOP>FRT-myr::smGFP-V5-THS10XUAS-FRT>STOP>FRT-myr::smGFPFLAG in su(Hw)attP1/TM2))	Nern et al., 2015	RRID: BDSC_64090 (Janelia stock 1116894)	
Genetic reagent (Drosophila melanogaster)	57C10PEST_attp18_4stop1; (MCFO-7; (w, R57C10-Flp2::PEST in attP18;; pJFRC210-10XUASFRT>STOP>FRT-myr::smGFP-OLLAS in attP2, pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFPHA in VK00005, pJFRC240-10XUAS-FRT>STOP>FRTmyr::smGFP-V5-THS-10XUAS-FRT>STOP>FRTmyr::smGFP-FLAG in su(Hw)attP1/TM2))	Nern et al., 2015	RRID: BDSC_64091 (Janelia stock 1116875)	
Genetic reagent (Drosophila melanogaster)	LexAop_IVS-myr_3_0008; (pJFRC19-13XLexAop2-IVS-myr::GFP in attP2)	Pfeiffer et al., 2012	RRID: BDSC_32209 (Janelia stock 1116736)	
Genetic reagent (Drosophila melanogaster)	UAS_IVS-mCD8_3_0007; (pJFRC2-10XUAS-IVS-mCD8::GFP)	Pfeiffer et al., 2010	RRID: BDSC_32185 (Janelia stock 1115125)	
Genetic reagent (Drosophila melanogaster)	HAV5_X_0083; (pJFRC200-10XUAS-IVS-myr::smGFP-HA (attP18), pJFRC216-13XLexAop2-IVS-myr::smGFP-V5 (su(Hw)attP8; Dr e /TM6B))	Nern et al., 2015	(Janelia stock 3020172)	
Genetic reagent (Drosophila melanogaster)	IVS-myr-FLAG_Syt-HA_3_0055; (w;;5XUAS-IVS-myr::smFLAG in VK00005, pJFRC51-3XUAS-IVS-Syt::smHA in su(Hw)attP1)	Aso et al., 2014	(Janelia stock 3001064)	
Genetic reagent (Drosophila melanogaster)	UAS_IVS_myr_3_0009; (pJFRC12-10XUAS-IVS-myr::GFP in attP2)	Pfeiffer et al., 2010	(Janelia stock 1115116)	

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (Drosophila melanogaster)	TLN-V5_myr-FLAG_Syt-HA_23_0037; (w; 3XpJFRC-TLN-smV5 in su(Hw)attP5; 5XUAS-IVS-myr::smFLAG in VK00005_pJFRC51-3XUAS-IVS-Syt::smHA in su(Hw)attP1 /CyO::TM6b)	Nern et al., 2015	(Janelia stock 2600002)	
Genetic reagent (Drosophila melanogaster)	TLN-V5_myr-FLAG_Syt-HA_23_0038; (w; 3XpJFRC-TLN-smV5 in su(Hw)attP5; 5XUAS-IVS-myr::smFLAG in VK00005_pJFRC51-3XUAS-IVS-Syt::smHA in su(Hw)attP2 /CyO::TM6b)	Nern et al., 2015	(Janelia stock 2600003)	
Genetic reagent (Drosophila melanogaster)	UAS_IVS_mCD8_3_0045; (pJFRC2-10XUAS-IVS-mCD8::GFP in VK00005)	Pfeiffer et al., 2010	(Janelia stock 1115339)	
Genetic reagent (Drosophila melanogaster)	IVS_myr_GFP_X_0072; (pJFRC200-10XUAS-IVS-myr::smGFP-HA in attP18)	Nern et al., 2015	RRID: BDSC_62145 (Janelia stock 1116624)	
Genetic reagent (Drosophila melanogaster)	UAS_IVS_mCD8_3_0073; (pJFRC7-20XUAS-IVS-mCD8::GFP in attP2)	Pfeiffer et al., 2010	RRID: BDSC_32194 (Janelia stock 1115387)	
Genetic reagent (Drosophila melanogaster)	LexAop2_Chrimson_Venus_X_0082; (13XLexAop2-CsChrimson-mVenus trafficked in attP18)	Klapoetke et al., 2014	(Janelia stock 1150410)	
Genetic reagent (Drosophila melanogaster)	57C10L_suHwattP8_HAV5_FLAG_0098; (R57C10-FlpL in su(Hw)attP8;; pJFRC201-10XUAS-FRT>STOP>FRT-myr::smGFP-HA in VK00005, pJFRC240-10XUAS-FRT>STOP>FRT-myr::smGFP-V5-THS-10XUAS-FRT>STOP>FRT-myr::smGFP-FLAG in su(Hw)attP1 /TM2)	Nern et al., 2015	RRID: BDSC_64087 (Janelia stock 3022016)	
Antibody	Anti-Brp mouse monoclonal nc82	Developmental Studies Hybridoma Bank (DSHB)	RRID: AB_2314866	1:30
Antibody	Anti-HA rabbit monoclonal C29F4	Cell Signaling Technologies: 3724S	RRID: AB_1549585	1:300
Antibody	Anti-FLAG rat monoclonal DYKDDDDK Epitope Tag Antibody	Novus Biologicals: NBP1-06712	RRID: AB_1625981	1:200
Antibody	DyLight 550 conjugated anti-V5 mouse monoclonal	AbD Serotec: MCA1360D550GA	RRID: AB_2687576	1:500
Antibody	Anti-rat IgG (H&L) Goat Polyclonal Antibody ATTO 647N Conjugated	Rockland: 612-156-120	RRID: AB_10893386	1:300
Antibody	Alexa Fluor 594 AffiniPure Donkey Polyclonal Anti-Rabbit IgG (H+L)	Jackson ImmunoResearch Labs: 711-585-152	RRID: AB_2340621	1:500
Antibody	Anti-Green Fluorescent Protein (GFP) Rabbit Polyclonal Antibody, Unconjugated	Thermo Fisher Scientific: A-11122	RRID: AB_221569	1:1000
Antibody	Goat Polyclonal anti-Rabbit IgG (H+L) Highly Cross- Adsorbed Antibody, Alexa Fluor 488	Thermo Fisher Scientific: A-11034	RRID: AB_2576217	1:800
Antibody	Goat Polyclonal anti-Mouse IgG (H+L) Highly Cross- Adsorbed Antibody, Alexa Fluor 568	Thermo Fisher Scientific: A-11031	RRID: AB_144696	1:800
Antibody	Anti-HA High Affinity; Rat monoclonal antibody (clone 3F10)	Roche: 11867423001	RRID: AB_390918	1:100
Antibody	Cy2-AffiniPure Goat Anti-Mouse IgG (H+L) (min X Hu,Bov,Hrs,Rb,Rat Sr Prot)	Jackson ImmunoResearch Labs: 115-225-166	RRID: AB_2338746	1:600

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Antibody	Cy3-AffiniPure Goat Anti-Rabbit IgG (H+L) (min X Hu,Ms,Rat Sr Prot)	Jackson ImmunoResearch Labs: 111-165-144	RRID: AB_2338006	1:1000
Software, algorithm	Janelia Workstation	Rokicki et al., 2019; https://github.com/JaneliaSciComp/workstation ; Howard Hughes Medical Institute, 2023	RRID: SCR_014302	
Software, algorithm	NeuronBridge codebase	Clements et al., 2022; Clements et al., 2021 https://doi.org/10.25378/janelia.12159378.v2		
Software, algorithm	Fiji	https://fiji.sc	SCR_0022852	
Software, algorithm	Affinity Designer	https://affinity.serif.com/designer/	RRID: SCR_016952	
Other	FlyLight protocols for dissection, immunohistochemistry, and mounting	https://www.janelia.org/project-team/flylight/protocols	Protocol	

Fly driver stocks

Included split-GAL4 driver stocks and references are listed in Figure 1 Supplemental File 1. See **Table 2** [for](#) counts of driver stocks available from Bloomington Drosophila Stock Center.

Fly reporter stocks

Rescreening of the cell type collection made use of a 20XUAS-CsChrimson-mVenus in attP18 reporter (Klapoetke et al, 2014). Additional reporters are listed in the Key Resources Table and Figure 1 Supplemental File 4.

Fly crosses and dissection

Flies were raised on standard cornmeal molasses food typically between 21 and 25 C and 50% humidity. Crosses with hs-Flp MCFO were at 21-22 C, except for a 10-60 minute 37 C heat shock. Other crosses were typically at 25 C for most of development and 22 C in preparation for dissection. Flies were typically dissected at 1-5 days old for the cell type collection rescreening, 1-8 days old for other non-MCFO crosses and 3-8 days old for MCFO images reported in this study. Dissection and fixation were carried out as previously described (Aso et al., 2014 [for](#); Nern et al., 2015 [for](#)). Protocols are available at <https://www.janelia.org/project-team/flylight/protocols> [for](#).

Immunohistochemistry, imaging, and image processing

Immunohistochemistry, imaging, and image processing were carried out as previously described (Aso et al., 2014 [for](#); Nern et al., 2015 [for](#); Sterne et al., 2021 [for](#); Meissner et al., 2023 [for](#)). See Key Resources table for antibodies. Additional details of image processing, including source code for processing scripts, are available at <https://data.janelia.org/pipeline> [for](#). See Figure 4 Supplemental Figure 1 for a user guide to interpreting the image data.

Cell type collection rescreening

If inconsistencies were observed in expression between rescreened images and older image data for a line, the cross was repeated. If still inconsistent, the other copy of the stock (all are kept in duplicate) was examined. If only one of the two stock copies gave the expected expression pattern,

it was used to replace the other copy. Repeated problems with expression pattern, except for stochasticity within a cell type, were grounds for removal from the collection. If sex differences were observed, crosses were also repeated. Lines reported with sex differences showed the same difference at least twice. We could not eliminate the possibility of stock issues appearing as sex differences by coincidentally appearing different between sexes and consistent within them.

Line quality levels

Cell type lines were qualitatively scored first by visual examination of averaged color depth Maximum Intensity Projection (MIP) CNS images (a version of Figure 1 Supplemental File 2 with less image compression; [Otsuna et al., 2018](#)), and then by a second observer examining individual images. Scores were applied as described in **Figure 2**. As images were taken from a variety of reporters, an effort was made to discount background specific to the reporters, especially 5XUAS-IVS-myr::smFLAG in VK00005.

Line distribution analysis

3,029 cell type split-GAL4 lines were analyzed, each including images of one male and one female fly sample from rescreening. 6,058 brains and 6,058 VNCs were included. Images were aligned to the JRC2018 Unisex template ([Bogovic et al., 2020](#)) and segmented using Direction-Selective Local Thresholding (DSLTL; [Kawase et al., 2015](#)). DSLTL weight value, neuron volume percentage, MIP ratio against the aligned template, and MIP-weight ratio were adjusted to refine segmentation. DSLTL connected fragmented neurons with up to a 5 μ m gap, enhancing neuronal continuity. Noise was eliminated and objects were separated into individual neurons using a connecting component algorithm. The algorithm distinguishes individual neurons by recognizing connected voxel clusters as distinct entities based on DSLTL 3D segmentation, followed by removal of voxels below 45/4095 (12 bit) brightness and objects smaller than 30 voxels.

The segmentation produced 19,313 non-overlapping brain objects and 18,860 VNC objects, consisting of neurons, trachea, debris, and antibody background. Most non-neuronal objects and some very dim neurons were eliminated using color depth MIP-based 2D shape filters and machine learning filters, followed by visual inspection, yielding 10,778 brain and 9,310 VNC objects. We observed 74 objects containing tissue background with neurons. The background was manually 3D removed using VVDViewer. Neuron masks were combined into a single volume heat map with a newly developed Fiji plugin “Seg_volume_combine_to_heatmap.jar” (<https://github.com/JaneliaSciComp/3D-fiber-auto-segmentation/tree/main>). Voxel brightness indicates the number of objects (at most one per line) within each voxel.

Acknowledgements

This work is part of the FlyLight Project Team at Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA. Author order includes the following alphabetical groups: FlyLight Project Team, Janelia Fly Facility, Janelia Scientific Computing Shared Resource, and contributing laboratories. We thank Kristin Scott for visitor project mentorship and the generation of lines labeling the subesophageal ganglion. We thank Arnim Jenett and Teri Ngo for early contributions to split-GAL4 screening. We thank Yichun Shuai for annotations of lines generated with the Aso lab. We thank Project Technical Resources for management coordination and staff support. We thank Melanie Radcliff and Crystal Sullivan Di Pietro for administrative support. We thank Mark Bolstad, Tom Dolafi, Leslie L Foster, Sean Murphy, Donald J Olbris, Todd Safford, Eric Trautman, and Yang Yu for their work on software infrastructure. We thank Olivia DeThomasis, Raj Nair, Hina Naz, Jessalyn Pugh, Brian Rebhorn, Rose Rogall, and Rodney Simmons for preparation of fly food. We thank David Shepherd and Sam Whitehead for their help with analyzing VNC interneuron types. We thank Stephan Saalfeld for imaging discussions. Stocks obtained from the

Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study. We thank them, especially Annette Parks, Cale Whitworth, and Sam Zheng, for the maintenance and distribution of stocks from the *Janelia* collection. We thank Rob Court, Alex McLachlan, and David Osumi-Sutherland for their work on Virtual Fly Brain and coordination on image distribution. We thank Brian Crooks and Zbigniew Iwinski of Zeiss for microscopy assistance. This work was supported in part by the *Janelia* Visiting Scientist Program. This article is subject to HHMI's Open Access to Publications policy. HHMI lab heads and project team leads have previously granted a nonexclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their research articles. Pursuant to those licenses, the author-accepted manuscript of this article can be made freely available under a CC BY 4.0 license immediately upon publication.

Author contributions

Geoffrey W Meissner, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing—original draft, Writing—review & editing, Visualization, Supervision, Project administration; Allison Vannan, Validation, Investigation, Data curation, Project administration; Jennifer Jeter, Validation, Investigation, Visualization, Project administration; Kari Close, Investigation; Gina M DePasquale, Investigation; Zachary Dorman, Validation, Investigation, Project administration; Kaitlyn Forster, Investigation; Jaye Anne Beringer, Investigation; Theresa V Gibney, Investigation; Joanna H Hausenfluck, Investigation; Yisheng He, Investigation; Kristin Henderson, Investigation, Writing—review & editing; Lauren Johnson, Investigation; Rebecca M Johnston, Methodology, Validation, Investigation, Project administration; Gudrun Ihrke, Writing—review & editing, Supervision, Project administration; Nirmala Iyer, Investigation; Rachel Lazarus, Validation, Investigation, Project administration; Kelley Lee, Validation, Investigation, Project administration; Hsing-Hsi Li, Validation, Investigation, Project administration; Hua-Peng Liaw, Investigation; Brian Melton, Validation, Investigation; Scott Miller, Investigation; Reeham Motaher, Investigation; Alexandra Novak, Investigation; Omotara Ogundeyi, Investigation; Alyson Petruncio, Investigation; Jacquelyn Price, Validation, Investigation, Project administration; Sophia Protopapas, Investigation; Susana Tae, Validation, Investigation, Project administration; Jennifer Taylor, Investigation; Rebecca Vorimo, Validation, Investigation, Project administration; Brianna Yarbrough, Investigation; Kevin Xiankun Zeng, Validation, Investigation; Christopher T Zugates, Investigation, Supervision, Project administration; Heather Dionne, Investigation; Claire Angstadt, Investigation; Kelly Ashley, Investigation; Amanda Cavallaro, Validation, Investigation, Supervision, Project administration; Tam Dang, Investigation; Guillermo A Gonzalez III, Investigation; Karen L Hibbard, Investigation; Cuizhen Huang, Investigation; Jui-Chun Kao, Validation, Investigation; Todd Laverty, Supervision, Project administration; Monti Mercer, Investigation; Brenda Perez, Investigation; Scarlett Pitts, Validation, Investigation; Danielle Ruiz, Investigation; Viruthika Vallanadu, Investigation; Grace Zhiyu Zheng, Investigation; Cristian Goina, Software; Hideo Otsuna, Methodology, Software, Formal analysis, Data curation, Visualization; Konrad Rokicki, Software, Writing—review & editing, Visualization, Supervision; Robert R Svirskas, Software, Visualization; Han SJ Cheong, Investigation, Data curation; Michael-John Dolan, Investigation, Data curation; Erica Ehrhardt, Investigation, Data curation; Kai Feng, Investigation, Data curation; Basel El Galfi, Investigation, Data curation; Jens Goldammer, Investigation, Data curation, Writing—review & editing; Stephen J Huston, Investigation, Data curation; Nan Hu, Investigation, Data curation; Masayoshi Ito, Investigation, Data curation; Claire McKellar, Investigation, Data curation; Ryo Minegishi, Investigation, Data curation; Shigehiro Namiki, Investigation, Data curation; Aljoscha Nern, Conceptualization, Methodology, Investigation, Data curation, Writing—review & editing; Catherine E Schretter, Investigation, Data curation; Gabriella R Sterne, Investigation, Data curation, Writing—review & editing; Lalanti Venkatasubramanian, Investigation, Data curation; Kaiyu Wang, Investigation, Data curation; Tanya Wolff, Investigation, Data curation, Writing—review & editing; Ming Wu, Investigation, Data curation; Reed George, Supervision; Oz Malkesman, Supervision, Project administration; Yoshinori Aso, Conceptualization, Methodology, Investigation, Data curation, Writing—review & editing.

Supervision; Gwyneth M Card, Conceptualization, Supervision; Barry J Dickson, Conceptualization, Supervision; Wyatt Korff, Conceptualization, Supervision, Funding acquisition; Kei Ito, Conceptualization, Writing—review & editing, Supervision; James W Truman, Conceptualization, Supervision; Marta Zlatic, Conceptualization, Writing—review & editing, Supervision; Gerald M Rubin, Conceptualization, Methodology, Investigation, Writing—original draft, Writing—review & editing, Supervision, Project administration, Funding acquisition; FlyLight Project Team, Investigation, Validation

Data availability

The footprint of this image resource (~200 TB) exceeds our known current practical limits on standard public data repositories. Thus, we have made all the primary adult data (and a variety of processed outputs) used in this study freely available under a CC BY 4.0 license at <https://doi.org/10.25378/janelia.21266625.v1> and through the publicly accessible websites <https://splitgal4.janelia.org> and <https://flylight-raw.janelia.org>. Many images are being made searchable with the same permissions on the user-friendly NeuronBridge website <https://neuronbridge.janelia.org>. NeuronBridge code is available at [Clements et al., 2021](#) and the application and implementation are discussed further in [Clements et al., 2024](#). Larval images are available at <https://raw.larval.flylight.virtualflybrain.org/>. All other data generated or analyzed during this study are included in the manuscript and supporting files.

References

- Aso Y *et al.* (2014) **The neuronal architecture of the mushroom body provides a logic for associative learning** *eLife* **3** <https://doi.org/10.7554/eLife.04577>
- Aso Y, Rubin GM (2016) **Dopaminergic neurons write and update memories with cell-type-specific rules** *Elife* **5** <https://doi.org/10.7554/eLife.16135>
- Aso Y, Yamada D, Bushey D, Hibbard KL, Sammons M, Otsuna H, Shuai Y, Hige T (2023) **Neural circuit mechanisms for transforming learned olfactory valences into wind-oriented movement** *Elife* **12** <https://doi.org/10.7554/eLife.85756>
- Baird GS, Zacharias DA, Tsien RY (2000) **Biochemistry, mutagenesis, and oligomerization of DsRed, a red fluorescent protein from coral** *Proc Natl Acad Sci U S A* **97** <https://doi.org/10.1073/pnas.97.22.11984>
- Baker CA, McKellar C, Pang R, Nern A, Dorkenwald S, Pacheco DA, Eckstein N, Funke J, Dickson BJ, Murthy M (2022) **Neural network organization for courtship-song feature detection in *Drosophila*** *Curr Biol* **32** <https://doi.org/10.1016/j.cub.2022.06.019>
- Bellen HJ, O’Kane CJ, Wilson C, Grossniklaus U, Pearson RK, Gehring WJ (1989) **P-element-mediated enhancer detection: a versatile method to study development in *Drosophila*** *Genes Dev* **3**:1288–1300
- Bogovic JA *et al.* (2020) **An unbiased template of the *Drosophila* brain and ventral nerve cord** *PLoS One* **15** <https://doi.org/10.1371/journal.pone.0236495>
- Brand AH, Perrimon N (1993) **Targeted gene expression as a means of altering cell fates and generating dominant phenotypes** *Development* **118**:401–415
- Cheong HSJ *et al.* (2023) **Transforming descending input into behavior: The organization of premotor circuits in the *Drosophila* Male Adult Nerve Cord connectome** *bioRxiv* <https://doi.org/10.1101/2023.06.07.543976>
- Cheong HSJ *et al.* (2024) **Organization of an ascending circuit that conveys flight motor state in *Drosophila*** *Current Biology* <https://doi.org/10.1016/j.cub.2024.01.071>
- Chen C, Aymanns F, Minegishi R, Matsuda VDV, Talabot N, Günel S, Dickson BJ, Ramdya P (2023) **Ascending neurons convey behavioral state to integrative sensory and action selection brain regions** *Nat Neurosci* **26** <https://doi.org/10.1038/s41593-023-01281-z>
- Chen YD, Chen Y, Rajesh R, Shoji N, Jacy M, Lacin H, Erclik T, Desplan C (2023) **Using single-cell RNA sequencing to generate cell-type-specific split-GAL4 reagents throughout development** *bioRxiv* <https://doi.org/10.1101/2023.02.03.527019>
- Clements J, Goina C, Otsuna H, Kazimiers A, Kawase T, Olbris DJ, Svirskas R, Rokicki K (2021) **NeuronBridge Codebase** *figshare* <https://doi.org/10.25378/janelia.12159378.v2>

- Clements J, Goina C, Hubbard PM, Kawase T, Olbris DJ, Otsuna H, Svirkas R, Rokicki K (2024) **NeuronBridge: an intuitive web application for neuronal morphology search across large data sets** *BMC Bioinformatics* **25** <https://doi.org/10.1186/s12859-024-05732-7>
- Davis FP, Nern A, Picard S, Reiser MB, Rubin GM, Eddy SR, Henry GL (2020) **A genetic, genomic, and computational resource for exploring neural circuit function** *Elife* **9** <https://doi.org/10.7554/eLife.50901>
- Dionne H, Hibbard KL, Cavallaro A, Kao JC, Rubin GM (2018) **Genetic reagents for making Split-GAL4 lines in Drosophila** *Genetics* **209**:31–35 <https://doi.org/10.1534/genetics.118.300682>
- Dolan M, Luan H, Shropshire WC, Sutcliffe B, Cocanougher B, Scott RL, Frechter S, Zlatić M, Jefferis GSXE, White BH (2017) **Facilitating neuron-specific genetic manipulations in Drosophila melanogaster using a split GAL4 repressor** *Genetics* **206** <https://doi.org/10.1534/genetics.116.199687>
- Dolan M *et al.* (2019) **Neurogenetic dissection of the Drosophila lateral horn reveals major outputs, diverse behavioural functions, and interactions with the mushroom body** *Elife* **8** <https://doi.org/10.7554/eLife.43079>
- Ehrhardt E *et al.* (2023) **Single-cell type analysis of wing premotor circuits in the ventral nerve cord of Drosophila melanogaster** *bioRxiv* <https://doi.org/10.1101/2023.05.31.542897>
- Ewen-Campen B, Luan H, Xu J, Singh R, Joshi N, Thakkar T, Berger B, White BH, Perrimon N (2023) **. split-intein Gal4 provides intersectional genetic labeling that is repressible by Gal80** *Proc Natl Acad Sci U S A* **120** <https://doi.org/10.1073/pnas.2304730120>
- Feng K, Palfreyman MT, Häsemeyer M, Talsma A, Dickson BJ (2014) **Ascending SAG neurons control sexual receptivity of Drosophila females** *Neuron* **83** <https://doi.org/10.1016/j.neuron.2014.05.017>
- Feng K, Sen R, Minegishi R, Dübberdt M, Bockemühl T, Büschges A, Dickson BJ (2020) **Distributed control of motor circuits for backward walking in Drosophila** *Nat Commun* **11** <https://doi.org/10.1038/s41467-020-19936-x>
- Fischer JA, Giniger E., Maniatis T., Ptashne M. (1988) **GAL4 activates transcription in Drosophila** *Nature* **332**:853–856
- Garner D, Kind E, Nern A, Houghton L, Zhao A, Sancer G, Rubin GM, Wernet MF, Kim SS (2023) **Connectomic reconstruction predicts the functional organization of visual inputs to the navigation center of the Drosophila brain** *bioRxiv* <https://doi.org/10.1101/2023.11.29.569241>
- Gorko B *et al.* (2024) **Motor neurons generate pose-targeted movements via proprioceptive sculpting** *Nature* <https://doi.org/10.1038/s41586-024-07222-5>
- Griffith LC (2012) **Identifying behavioral circuits in Drosophila melanogaster: moving targets in a flying insect** *Curr. Opin. Neurobiol* **22**:609–614
- Groth AC, Fish M, Nusse R, Calos MP (2004) **Construction of transgenic Drosophila by using the site-specific integrase from phage phiC31** *Genetics* **166**:1775–1782

Guo F, Chen X, Rosbash M (2017) **Temporal calcium profiling of specific circadian neurons in freely moving flies** *Proc Natl Acad Sci U S A* **114** <https://doi.org/10.1073/pnas.1706608114>

Harris RM, Pfeiffer BD, Rubin GM, Truman JW (2015) **Neuron hemilineages provide the functional ground plan for the Drosophila ventral nervous system** *Elife* **4** <https://doi.org/10.7554/eLife.04493>

Hulse BK *et al.* (2021) **A connectome of the Drosophila central complex reveals network motifs suitable for flexible navigation and context-dependent action selection** *Elife* **10** <https://doi.org/10.7554/eLife.66039>

Ichinose T, Aso Y, Yamagata N, Abe A, Rubin GM, Tanimoto H (2015) **Reward signal in a recurrent circuit drives appetitive long-term memory formation** *Elife* **4** <https://doi.org/10.7554/eLife.10719>

Isaacson MD *et al.* (2023) **Small-field visual projection neurons detect translational optic flow and support walking control** *bioRxiv* <https://doi.org/10.1101/2023.06.21.546024>

Ito K, Okada R, Tanaka NK, Awasaki T (2003) **Cautionary observations on preparing and interpreting brain images using molecular biology-based staining techniques** *Microsc. Res. Tech* **62**:170–186

Jenett A *et al.* (2012) **A GAL4-driver line resource for Drosophila neurobiology** *Cell Reports* **2**:991–1001 <https://doi.org/10.1016/j.celrep.2012.09.011>

Jovanic T, Winding M, Cardona A, Truman JW, Gershow M, Zlatić M (2019) **Neural substrates of Drosophila larval anemotaxis** *Curr Biol* **29** <https://doi.org/10.1016/j.cub.2019.01.009>

Kawase T, Sugano SS, Shimada T, Hara-Nishimura I (2015) **A direction-selective local-thresholding method, DSLT, in combination with a dye-based method for automated three-dimensional segmentation of cells and airspaces in developing leaves** *Plant J* **81** <https://doi.org/10.1111/tpj.12738>

Kind E *et al.* (2021) **Synaptic targets of photoreceptors specialized to detect color and skylight polarization in Drosophila** *Elife* **10** <https://doi.org/10.7554/eLife.71858>

Klapoetke NC, Nern A, Peek MY, Rogers EM, Breads P, Rubin GM, Reiser MB, Card GM (2017) **Ultra-selective looming detection from radial motion opponency** *Nature* **551** <https://doi.org/10.1038/nature24626>

Klapoetke NC, Nern A, Rogers EM, Rubin GM, Reiser MB, Card GM (2022) **A functionally ordered visual feature map in the Drosophila brain** *Neuron* **110** <https://doi.org/10.1016/j.neuron.2022.02.013>

Levine M, Tjian R (2003) **Transcription regulation and animal diversity** *Nature* **424**:147–51 <https://doi.org/10.1038/nature01763>

Li HH, Kroll JR, Lennox SM, Ogundeyi O, Jeter J, Depasquale G, Truman JW (2014) **A GAL4 driver resource for developmental and behavioral studies on the larval CNS of Drosophila** *Cell Rep* **8**:897–908 <https://doi.org/10.1016/j.celrep.2014.06.065>

Liang X, Holy TE, Taghert PH (2017) **A series of suppressive signals within the Drosophila circadian neural circuit generates sequential daily outputs** *Neuron* **94** <https://doi.org/10.1016/j.neuron.2017.05.007>

- Lillvis JL, Wang K, Shiozaki HM, Xu M, Stern DL, Dickson BJ (2024) **Nested neural circuits generate distinct acoustic signals during *Drosophila* courtship** *Curr Biol* **34** <https://doi.org/10.1016/j.cub.2024.01.015>
- Longden KD, Rogers EM, Nern A, Dionne H, Reiser MB (2023) **Different spectral sensitivities of ON- and OFF-motion pathways enhance the detection of approaching color objects in *Drosophila*** *Nat Commun* **14** <https://doi.org/10.1038/s41467-023-43566-8>
- Luan H, Peabody NC, Vinson CR, White BH (2006) **Refined spatial manipulation of neuronal function by combinatorial restriction of transgene expression** *Neuron* **52**:425–436 <https://doi.org/10.1016/j.neuron.2006.08.028>
- Mais L *et al.* (2021) **PatchPerPixMatch for automated 3d search of neuronal morphologies in light microscopy** *bioRxiv* <https://doi.org/10.1101/2021.07.23.453511>
- Manseau L *et al.* (1997) **GAL4 enhancer traps expressed in the embryo, larval brain, imaginal discs, and ovary of *Drosophila*** *Developmental Dynamics* **209**:310–322
- Marin EC *et al.* (2023) **Systematic annotation of a complete adult male *Drosophila* nerve cord connectome reveals principles of functional organisation** *bioRxiv* <https://doi.org/10.1101/2023.06.05.543407>
- Meissner GW *et al.* (2023) **A searchable image resource of *Drosophila* GAL4 driver expression patterns with single neuron resolution** *Elife* **12** <https://doi.org/10.7554/eLife.80660>
- Mellert DJ, Truman JW (2012) **Transvection is common throughout the *Drosophila* genome** *Genetics* **191** <https://doi.org/10.1534/genetics.112.140475>
- Minegishi R, Dickson BJ, Team FP (2023) **Ascending Neurons 2023 split-GAL4 lines** *figshare* <https://doi.org/10.25378/janelia.23726103.v1>
- Morimoto MM, Nern A, Zhao A, Rogers EM, Wong AM, Isaacson MD, Bock DD, Rubin GM, Reiser MB (2020) **Spatial readout of visual looming in the central brain of *Drosophila*** *Elife* **9** <https://doi.org/10.7554/eLife.57685>
- Namiki S, Dickinson MH, Wong AM, Korff W, Card GM (2018) **The functional organization of descending sensory-motor pathways in *Drosophila*** *Elife* **7** <https://doi.org/10.7554/eLife.34272>
- Namiki S, Ros IG, Morrow C, Rowell WJ, Card GM, Korff W, Dickinson MH (2022) **A population of descending neurons that regulates the flight motor of *Drosophila*** *Curr Biol* **32** <https://doi.org/10.1016/j.cub.2022.01.008>
- Nern A, Pfeiffer BD, Rubin GM (2015) **Optimized tools for multicolor stochastic labeling reveal diverse stereotyped cell arrangements in the fly visual system** *Proc Natl Acad Sci U S A* **112**:E2967–76 <https://doi.org/10.1073/pnas.1506763112>
- Nern A *et al.* (2024) **Connectome-driven neural inventory of a complete visual system** *bioRxiv* <https://doi.org/10.1101/2024.04.16.589741>
- Otsuna H, Ito M, Kawase T (2018) **Color depth MIP mask search: a new tool to expedite Split-GAL4 creation** *bioRxiv* <https://doi.org/10.1101/318006>

Pavlou HJ, Lin AC, Neville MC, Nojima T, Diao F, Chen BE, White BH, Goodwin SF (2016) **Neural circuitry coordinating male copulation** *Elife* 5 <https://doi.org/10.7554/eLife.20713>

Pfeiffer BD *et al.* (2008) **Tools for neuroanatomy and neurogenetics in Drosophila** *PNAS* 105:9715–9720 <https://doi.org/10.1073/pnas.0803697105>

Pfeiffer BD, Ngo TT, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM (2010) **Refinement of tools for targeted gene expression in Drosophila** *Genetics* 186:735–755 <https://doi.org/10.1534/genetics.110.119917>

Pfeiffer BD, Truman JW, Rubin GM (2012) **Using translational enhancers to increase transgene expression in Drosophila** *Proc Natl Acad Sci U S A* 109

Robinson IM, Ranjan R, Schwarz TL (2002) **Synaptotagmins I and IV promote transmitter release independently of Ca(2+) binding in the C(2)A domain** *Nature* 418 <https://doi.org/10.1038/nature00915>

Rubin GM, Aso Y (2024) **New genetic tools for mushroom body output neurons in Drosophila** *Elife* 12 <https://doi.org/10.7554/eLife.90523>

Scheffer LK *et al.* (2020) **A connectome and analysis of the adult Drosophila central brain** *Elife* 9 <https://doi.org/10.7554/eLife.57443>

Schindelin J *et al.* (2012) **Fiji: an open-source platform for biological-image analysis** *Nat Methods* 9 <https://doi.org/10.1038/nmeth.2019>

Schlegel P *et al.* (2024) **Whole-brain annotation and multi-connectome cell typing of Drosophila** *Nature* 634 <https://doi.org/10.1038/s41586-024-07686-5>

Schlichting M, Díaz MM, Xin J, Rosbash M (2019) **Neuron-specific knockouts indicate the importance of network communication to Drosophila rhythmicity** *Elife* 8 <https://doi.org/10.7554/eLife.48301>

Schretter CE *et al.* (2020) **Cell types and neuronal circuitry underlying female aggression in Drosophila** *Elife* 9 <https://doi.org/10.7554/eLife.58942>

Schretter CE *et al.* (2024) **Social state gates vision using three circuit mechanisms in Drosophila** *bioRxiv* <https://doi.org/10.1101/2024.03.15.585289>

Shuai Y *et al.* (2024) **Driver lines for studying associative learning in Drosophila** *Elife* 13 <https://doi.org/10.7554/elife.94168.1>

Sterne GR, Otsuna H, Dickson BJ, Scott K (2021) **Classification and genetic targeting of cell types in the primary taste and premotor center of the adult Drosophila brain** *Elife* 10 <https://doi.org/10.7554/eLife.71679>

Strother JA, Wu S, Wong AM, Nern A, Rogers EM, Le JQ, Rubin GM, Reiser MB (2017) **The Emergence of Directional Selectivity in the Visual Motion Pathway of Drosophila** *Neuron* 94 <https://doi.org/10.1016/j.neuron.2017.03.010>

Takemura S, Nern A, Chklovskii DB, Scheffer LK, Rubin GM, Meinertzhagen IA (2017) **The comprehensive connectome of a neural substrate for 'ON' motion detection in Drosophila** *Elife* 6 <https://doi.org/10.7554/eLife.24394>

- Takemura S *et al.* (2023) **A Connectome of the Male Drosophila Ventral Nerve Cord** *bioRxiv* <https://doi.org/10.1101/2023.06.05.543757>
- Tirian L, Dickson BJ (2017) **The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the Drosophila nervous system** *bioRxiv* <https://doi.org/10.1101/198648>
- Turner-Evans DB *et al.* (2020) **The neuroanatomical ultrastructure and function of a biological ring attractor** *Neuron* **108** <https://doi.org/10.1016/j.neuron.2020.08.006>
- Tuthill JC, Nern A, Holtz SL, Rubin GM, Reiser MB (2013) **Contributions of the 12 neuron classes in the fly lamina to motion vision** *Neuron* **79**:128–40 <https://doi.org/10.1016/j.neuron.2013.05.024>
- von Reyn CR, Nern A, Williamson WR, Breads P, Wu M, Namiki S, Card GM (2017) **Feature integration drives probabilistic behavior in the Drosophila escape response** *Neuron* **94** <https://doi.org/10.1016/j.neuron.2017.05.036>
- Wang F, Wang K, Forknall N, Patrick C, Yang T, Parekh R, Bock D, Dickson BJ (2020) **Neural circuitry linking mating and egg laying in Drosophila females** *Nature* **579** <https://doi.org/10.1038/s41586-020-2055-9>
- Wang F, Wang K, Forknall N, Parekh R, Dickson BJ (2020) **Circuit and behavioral mechanisms of sexual rejection by Drosophila females** *Curr Biol* **30** <https://doi.org/10.1016/j.cub.2020.07.083>
- Wang K, Wang F, Forknall N, Yang T, Patrick C, Parekh R, Dickson BJ (2021) **Neural circuit mechanisms of sexual receptivity in Drosophila females** *Nature* **589** <https://doi.org/10.1038/s41586-020-2972-7>
- Wolff T, Rubin GM (2018) **Neuroarchitecture of the Drosophila central complex: A catalog of nodulus and asymmetrical body neurons and a revision of the protocerebral bridge catalog** *J Comp Neurol* **526** <https://doi.org/10.1002/cne.24512>
- Wolff T, Eddison M, Chen N, Nern A, Sundaramurthi P, Sitaraman D, Rubin GM (2024) **Cell type-specific driver lines targeting the Drosophila central complex and their use to investigate neuropeptide expression and sleep regulation** *bioRxiv* <https://doi.org/10.1101/2024.10.21.619448>
- Wu M, Nern A, Williamson WR, Morimoto MM, Reiser MB, Card GM, Rubin GM (2016) **Visual projection neurons in the Drosophila lobula link feature detection to distinct behavioral programs** *Elife* **5** <https://doi.org/10.7554/eLife.21022>
- Venken K.J., Simpson J.H., Bellen H.J. (2011) **Genetic manipulation of genes and cells in the nervous system of the fruit fly** *Neuron* **72**:202–230
- Vijayan V, Wang F, Wang K, Chakravorty A, Adachi A, Akhlaghpour H, Dickson BJ, Maimon G (2023) **A rise-to-threshold process for a relative-value decision** *Nature* **619** <https://doi.org/10.1038/s41586-023-06271-6>
- Yamada D, Bushey D, Li F, Hibbard KL, Sammons M, Funke J, Litwin-Kumar A, Hige T, Aso Y (2023) **Hierarchical architecture of dopaminergic circuits enables second-order conditioning in Drosophila** *Elife* **12** <https://doi.org/10.7554/eLife.79042>

Yoshihara M, Ito K (2000) **Improved GAL4 screening kit for large-scale generation of enhancer-trap strains** *Drosophila Information Service* **83**:199–202

Zhao A *et al.* (2022) **Eye structure shapes neuron function in *Drosophila* motion vision** *bioRxiv* <https://doi.org/10.1101/2022.12.14.520178>

Zhao A *et al.* (2023) **A comprehensive neuroanatomical survey of the *Drosophila* Lobula Plate Tangential Neurons with predictions for their optic flow sensitivity** *bioRxiv* <https://doi.org/10.1101/2023.10.16.562634>

Zwart MF, Pulver SR, Truman JW, Fushiki A, Fetter RD, Cardona A, Landgraf M (2016) **Selective inhibition mediates the sequential recruitment of motor pools** *Neuron* **91** <https://doi.org/10.1016/j.neuron.2016.06.031>

Author information

Geoffrey W Meissner

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0003-0369-9788](https://orcid.org/0000-0003-0369-9788)

For correspondence: rubing@janelia.hhmi.org

Allison Vannan

Institute of Zoology, University of Cologne, Cologne, Germany

Jennifer Jeter

Institute of Zoology, University of Cologne, Cologne, Germany

Kari Close

Institute of Zoology, University of Cologne, Cologne, Germany

Gina M DePasquale

Institute of Zoology, University of Cologne, Cologne, Germany

Zachary Dorman

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0001-9933-7217](https://orcid.org/0000-0001-9933-7217)

Kaitlyn Forster

Institute of Zoology, University of Cologne, Cologne, Germany

Jaye Anne Beringer

Institute of Zoology, University of Cologne, Cologne, Germany

Theresa V Gibney

Institute of Zoology, University of Cologne, Cologne, Germany

Joanna H Hausenfluck

Institute of Zoology, University of Cologne, Cologne, Germany

Yisheng He

Institute of Zoology, University of Cologne, Cologne, Germany

Kristin Henderson

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0002-9265-7709](https://orcid.org/0000-0002-9265-7709)

Lauren Johnson

Institute of Zoology, University of Cologne, Cologne, Germany

Rebecca M Johnston

Institute of Zoology, University of Cologne, Cologne, Germany

Gudrun Ihrke

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0003-4604-735X](https://orcid.org/0000-0003-4604-735X)

Nirmala Iyer

Institute of Zoology, University of Cologne, Cologne, Germany

Rachel Lazarus

Institute of Zoology, University of Cologne, Cologne, Germany

Kelley Lee

Institute of Zoology, University of Cologne, Cologne, Germany

Hsing-Hsi Li

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0009-0001-9883-8201](https://orcid.org/0009-0001-9883-8201)

Hua-Peng Liaw

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0009-0003-4721-3556](https://orcid.org/0009-0003-4721-3556)

Brian Melton

Institute of Zoology, University of Cologne, Cologne, Germany

Scott Miller

Institute of Zoology, University of Cologne, Cologne, Germany

Reeham Motaher

Institute of Zoology, University of Cologne, Cologne, Germany

Alexandra Novak

Institute of Zoology, University of Cologne, Cologne, Germany

Omotara Ogundeyi

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0003-2770-0026](https://orcid.org/0000-0003-2770-0026)

Alyson Petruncio

Institute of Zoology, University of Cologne, Cologne, Germany

Jacquelyn Price

Institute of Zoology, University of Cologne, Cologne, Germany

Sophia Protopapas

Institute of Zoology, University of Cologne, Cologne, Germany

Susana Tae

Institute of Zoology, University of Cologne, Cologne, Germany

Jennifer Taylor

Institute of Zoology, University of Cologne, Cologne, Germany

Rebecca Vorimo

Institute of Zoology, University of Cologne, Cologne, Germany

Brianna Yarbrough

Institute of Zoology, University of Cologne, Cologne, Germany

Kevin Xiankun Zeng

Institute of Zoology, University of Cologne, Cologne, Germany

Christopher T Zugates

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0003-1882-3665](https://orcid.org/0000-0003-1882-3665)

Heather Dionne

Institute of Zoology, University of Cologne, Cologne, Germany

Claire Angstadt

Institute of Zoology, University of Cologne, Cologne, Germany

Kelly Ashley

Institute of Zoology, University of Cologne, Cologne, Germany

Amanda Cavallaro

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0002-0650-4651](https://orcid.org/0000-0002-0650-4651)

Tam Dang

Institute of Zoology, University of Cologne, Cologne, Germany

Guillermo A Gonzalez III

Institute of Zoology, University of Cologne, Cologne, Germany

Karen L Hibbard

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0002-2001-6099](https://orcid.org/0000-0002-2001-6099)

Cuizhen Huang

Institute of Zoology, University of Cologne, Cologne, Germany

Jui-Chun Kao

Institute of Zoology, University of Cologne, Cologne, Germany

Todd Lavery

Institute of Zoology, University of Cologne, Cologne, Germany

Monti Mercer

Institute of Zoology, University of Cologne, Cologne, Germany

Brenda Perez

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0002-2420-402X](https://orcid.org/0000-0002-2420-402X)

Scarlett Pitts

Institute of Zoology, University of Cologne, Cologne, Germany

Danielle Ruiz

Institute of Zoology, University of Cologne, Cologne, Germany

Viruthika Vallanadu

Institute of Zoology, University of Cologne, Cologne, Germany

Grace Zhiyu Zheng

Institute of Zoology, University of Cologne, Cologne, Germany

Cristian Goina

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0003-2835-7602](https://orcid.org/0000-0003-2835-7602)

Hideo Otsuna

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0002-2107-8881](https://orcid.org/0000-0002-2107-8881)

Konrad Rokicki

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0002-2799-9833](https://orcid.org/0000-0002-2799-9833)

Robert R Svirskas

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0001-8374-6008](https://orcid.org/0000-0001-8374-6008)

Han SJ Cheong

Institute of Zoology, University of Cologne, Cologne, Germany

ORCID iD: [0000-0001-5233-3438](https://orcid.org/0000-0001-5233-3438)

Michael-John Dolan

Institute of Zoology, University of Cologne, Cologne, Germany

Erica Ehrhardt

Institute of Zoology, University of Cologne, Cologne, Germany, Janelia Research Campus,
Howard Hughes Medical Institute, Ashburn, United States
ORCID iD: [0000-0002-9252-1414](https://orcid.org/0000-0002-9252-1414)

Kai Feng

Queensland Brain Institute, University of Queensland, Brisbane, Australia, Janelia Research
Campus, Howard Hughes Medical Institute, Ashburn, United States
ORCID iD: [0000-0002-5506-6889](https://orcid.org/0000-0002-5506-6889)

Basel El Galfi

Institute of Zoology, University of Cologne, Cologne, Germany

Jens Goldammer

Institute of Zoology, University of Cologne, Cologne, Germany, Janelia Research Campus,
Howard Hughes Medical Institute, Ashburn, United States
ORCID iD: [0000-0002-5623-8339](https://orcid.org/0000-0002-5623-8339)

Stephen J Huston

Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York,
United States, Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, United
States
ORCID iD: [0000-0002-0834-3257](https://orcid.org/0000-0002-0834-3257)

Nan Hu

Institute of Zoology, University of Cologne, Cologne, Germany

Masayoshi Ito

Institute of Zoology, University of Cologne, Cologne, Germany

Claire McKellar

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-3580-7336](https://orcid.org/0000-0002-3580-7336)

Ryo Minegishi

Queensland Brain Institute, University of Queensland, Brisbane, Australia, Janelia Research
Campus, Howard Hughes Medical Institute, Ashburn, United States
ORCID iD: [0000-0002-2895-9438](https://orcid.org/0000-0002-2895-9438)

Shigehiro Namiki

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0003-1559-799X](https://orcid.org/0000-0003-1559-799X)

Aljoscha Nern

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-3822-489X](https://orcid.org/0000-0002-3822-489X)

Catherine E Schretter

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-3957-6838](https://orcid.org/0000-0002-3957-6838)

Gabriella R Sterne

Department of Cell & Molecular Biology, University of California, Berkeley, Berkeley, United States, Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, United States
ORCID iD: [0000-0002-7221-648X](https://orcid.org/0000-0002-7221-648X)

Lalanti Venkatasubramanian

Institute of Zoology, University of Cologne, Cologne, Germany

Kaiyu Wang

Institute of Zoology, University of Cologne, Cologne, Germany

Tanya Wolff

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-8681-1749](https://orcid.org/0000-0002-8681-1749)

Ming Wu

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-2193-8271](https://orcid.org/0000-0002-2193-8271)

Reed George

Institute of Zoology, University of Cologne, Cologne, Germany

Oz Malkesman

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0003-2219-7476](https://orcid.org/0000-0003-2219-7476)

Yoshinori Aso

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-2939-1688](https://orcid.org/0000-0002-2939-1688)

For correspondence: asoy@janelia.hhmi.org

Gwyneth M Card

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0002-7679-3639](https://orcid.org/0000-0002-7679-3639)

For correspondence: gc3017@columbia.edu

Barry J Dickson

Queensland Brain Institute, University of Queensland, Brisbane, Australia
ORCID iD: [0000-0003-0715-892X](https://orcid.org/0000-0003-0715-892X)

For correspondence: b.dickson@uq.edu.au

Wyatt Korff

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0001-8396-1533](https://orcid.org/0000-0001-8396-1533)

For correspondence: korffw@janelia.hhmi.org

Kei Ito

Institute of Zoology, University of Cologne, Cologne, Germany, Janelia Research Campus,
Howard Hughes Medical Institute, Ashburn, United States
ORCID iD: [0000-0002-7274-5533](https://orcid.org/0000-0002-7274-5533)

For correspondence: k.ito@uni-koeln.de

James W Truman

Institute of Zoology, University of Cologne, Cologne, Germany

For correspondence: jwt@uw.edu

Marta Zlatic

Institute of Zoology, University of Cologne, Cologne, Germany

For correspondence: mz209@cam.ac.uk

Gerald M Rubin

Institute of Zoology, University of Cologne, Cologne, Germany
ORCID iD: [0000-0001-8762-8703](https://orcid.org/0000-0001-8762-8703)

For correspondence: meissnerg@janelia.hhmi.org

FlyLight Project Team

Institute of Zoology, University of Cologne, Cologne, Germany

Editors

Reviewing Editor

Tania Reis

University of Colorado Anschutz Medical Campus, Aurora, United States of America

Senior Editor

Claude Desplan

New York University, New York, United States of America

Reviewer #1 (Public review):

Summary:

Meissner et al describe an update on the collection of split-GAL4 lines generated by a consortium led by Janelia Research Campus. This follows the same experimental pipeline described before and presents as a significant increment to the present collection. This will strengthen the usefulness and relevance of "splits" as a standard tool for labs that already use this tool and attract more labs and researchers to use it.

Strengths:

This manuscript presents a solid step to establish Split-GAL4 lines as a relevant tool in the powerful *Drosophila* toolkit. Not only the raw number of available lines contribute to the relevance of this tool in the "technical landscape" of genetic tools, but additional features of this effort contribute to the successful adoption. These include:

- (1) A description of expression patterns in the adult and larvae, expanding the "audience" for these tools
 - (2) A classification of line combination according to quality levels, which provides a relevant criterion while deciding to use a particular set of "splits".
 - (3) Discrimination between male and female expression patterns, providing hints regarding the potential role of these gender-specific circuits.
 - (4) The search engine seems to be user-friendly, facilitating the retrieval of useful information.
 - (5) An acknowledgement of the caveats and challenges that splits (like any other genetic tool) can carry.
- Overall, the authors employed a pipeline that maximizes the potential of the Split-GAL4 collection to the scientific community.

Weaknesses:

My concerns were resolved regarding the existence of caveats while using these tools that researchers should be aware of, particularly those using them for the first time.

<https://doi.org/10.7554/eLife.98405.2.sa2>

Reviewer #2 (Public review):

Summary:

This manuscript describes the creation and curation of a collection of genetic driver lines that specifically label small numbers of neurons, often just a single to handful of cell types, in the central nervous system of the fruit fly, *Drosophila melanogaster*. The authors screened over 77,000 split hemidriver combinations to yield a collection of 3060 lines targeting a range of cell types in the adult *Drosophila* central nervous system and 1373 lines characterized in third-instar larvae. These genetic driver lines have already contributed to several important publications and will no doubt continue to do so. It is a truly valuable resource that represents the cooperation of several labs throughout the *Drosophila* community.

Strengths:

The authors have thoughtfully curated and documented the lines that they have created, so that they may be maximally useful to the greater community. This documentation includes confocal images of neurons labeled by each driver line and when possible, a list of cell types labeled by the genetic driver line and their identity in an EM connectome dataset. The authors have also made available some information from the other lines they created and tested but deemed not specific or strong enough to be included as part of the collection. This additional resource will be a valuable aid for those seeking to label cell types that may not be included in the main collection.

The added revisions help to clarify important points relating to the creation of the lines, which lines were included as part of this specific collection, and caveats to be mindful of when using any of the described lines. These revisions will increase the manuscript's utility to users who may be less familiar with this resource.

Weaknesses:

The major weakness, which is also in some ways a strength, is the stringent requirement that lines that be included be highly specific across the CNS. As a result, the lines that are part of this specific collection are sparse and specific but also limited in which cell types they cover. Doubtless there are many missing cell types.

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public Review):

Summary:

Meissner et al describe an update on the collection of split-GAL4 lines generated by a consortium led by Janelia Research Campus. This follows the same experimental pipeline described before and presents as a significant increment to the present collection. This will strengthen the usefulness and relevance of "splits" as a standard tool for labs that already use this tool and attract more labs and researchers to use it.

Strengths:

This manuscript presents a solid step to establish Split-GAL4 lines as a relevant tool in the powerful Drosophila toolkit. Not only does the raw number of available lines contribute to the relevance of this tool in the "technical landscape" of genetic tools, but additional features of this effort contribute to the successful adoption. These include:

- (1) A description of expression patterns in the adult and larvae, expanding the "audience" for these tools*
- (2) A classification of line combination according to quality levels, which provides a relevant criterion while deciding to use a particular set of "splits".*
- (3) Discrimination between male and female expression patterns, providing hints regarding the potential role of these gender-specific circuits.*
- (4) The search engine seems to be user-friendly, facilitating the retrieval of useful information.*

Overall, the authors employed a pipeline that maximizes the potential of the Split-GAL4 collection to the scientific community.

Weaknesses:

The following aspects apply:

The use of split-GAL4 lines has improved tremendously the genetic toolkit of Drosophila and this manuscript is another step forward in establishing this tool in the genetic repertoire that laboratories use. Thus, this would be a perfect opportunity for the authors to review the current status of this tool, addressing its caveats and how to effectively implement it into the experimental pipeline.

- (1) While the authors do bring up a series of relevant caveats that the community should be aware of while using split-GAL4 lines, the authors should take the opportunity to address some of the genetic issues that frequently arise while using the described genetic tools. This is particularly important for laboratories that lack the experience using split-GAL4 lines and wish to use them. Some of these issues are covertly brought up, but not entirely clarified.*

First, why do the authors (wisely) rescreen the lines using UAS-CsChrimson-mVenus? One reason is that using another transgene (such as UAS-GFP) and/or another genomic locus can drive a different expression pattern or intensities. Although this is discussed, this should be made more explicit and the readers should be aware of this.

Second, it would be important to include a discussion regarding the potential of hemidriver lines to suffer from transvection effects whenever there is a genetic element in the same locus. These are serious issues that prevent a more reliable use of split-GAL4 lines that, once again, should be discussed.

We added additional explanatory text to the discussion.

(2) The authors simply mention that the goal of the manuscript is to "summarize the results obtained over the past decade.". A better explanation would be welcomed in order to understand the need of a dedicated manuscript to announce the availability of a new batch of lines when previous publications already described the Split-GAL4 lines. At the extreme, one might question why we need a manuscript for this when a simple footnote on Janelia's website would suffice.

We added an additional mention of the cell type split-GAL4 collection at the relevant section and added more emphasis on the curation process adding value to the final selections. We feel that the manuscript is useful to document the methods used for the contained analysis and datasets and gives a starting point to the reader to go through the many split-GAL4 publications and images.

Reviewer #2 (Public Review):

*Summary: This manuscript describes the creation and curation of a collection of genetic driver lines that specifically label small numbers of neurons, often just a single to handful of cell types, in the central nervous system of the fruit fly, *Drosophila melanogaster*. The authors screened over 77,000 split hemidriver combinations to yield a collection of 3060 lines targeting a range of cell types in the adult *Drosophila* central nervous system and 1373 lines characterized in third-instar larvae. These genetic driver lines have already contributed to several important publications and will no doubt continue to do so. It is a truly valuable resource that represents the cooperation of several labs throughout the *Drosophila* community.*

Strengths:

The authors have thoughtfully curated and documented the lines that they have created, so that they may be maximally useful to the greater community. This documentation includes confocal images of neurons labeled by each driver line and when possible, a list of cell types labeled by the genetic driver line and their identity in an EM connectome dataset. The authors have also made available some information from the other lines they created and tested but deemed not specific or strong enough to be included as part of the collection. This additional resource will be a valuable aid for those seeking to label cell types that may not be included in the main collection.

Weaknesses:

None, this is a valuable set of tools that took many years of effort by several labs. This collection will continue to facilitate important science for years to come.

We thank the reviewer for their positive feedback.

Reviewer #3 (Public Review):

Summary:

The manuscript by Meissner et al. describes a collection of 3060 Drosophila lines that can be used to genetically target very small numbers of brain cells. The collection is the product of over a decade of work by the FlyLight Project Team at the Janelia Research Campus and their collaborators. This painstaking work has used the intersectional split-Gal4 method to combine pairs of so-called hemidriviers into driver lines capable of highly refined expression, often targeting single cell types. Roughly one-third of the lines have been described and characterized in previous publications and others will be described in manuscripts still in preparation. They are brought together here with many new lines to form one high-quality collection of lines with exceptional selectivity of expression. As detailed in the manuscript, all of the lines described have been made publicly available accompanied by an online database of images and metadata that allow researchers to identify lines containing neurons of interest to them. Collectively, the lines include neurons in most regions of both the adult and larval nervous systems, and the imaging database is intended to eventually permit anatomical searching that can match cell types targeted by the lines to those identified at the EM level in emerging connectomes. In addition, the manuscript introduces a second, freely accessible database of raw imaging data for many lower quality, but still potentially useful, split-Gal4 driver lines made by the FlyLight Project Team.

Strengths:

Both the stock collection and the image databases are substantial and important resources that will be of obvious interest to neuroscientists conducting research in Drosophila. Although many researchers will already be aware of the basic resources generated at Janelia, the comprehensive description provided in this manuscript represents a useful summary of past and recent accomplishments of the FlyLight Team and their collaborators and will be very valuable to newcomers in the field. In addition, the new lines being made available and the effort to collect all lines that have been generated that have highly specific expression patterns is very useful to all.

Weaknesses:

The collection of lines presented here is obviously somewhat redundant in including lines from previously published collections. Potentially confusing is the fact that previously published split-Gal4 collections have also touted lines with highly selective expression, but only a fraction of those lines have been chosen for inclusion in the present manuscript. For example, the collection of Shuai et al. (2023) describes some 800 new lines, many with specificity for neurons with connectivity to the mushroom body, but only 168 of these lines were selected for inclusion here. This is presumably because of the more stringent criteria applied in selecting the lines described in this manuscript, but it would be useful to spell this out and explain what makes this collection different from those previously published (and those forthcoming).

We added more description of how this collection is focused on the best cell-type-specific lines across the CNS. An important requirement for inclusion was this degree of specificity across the CNS, while many prior publications had a greater emphasis on lines with a narrower focus of specificity.

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

Luckily for us, genetics is for the most part an exact science. However, there's still some "voodoo" in a lot of genetic combinations that the authors should disclose and be as clear as possible in the manuscript. This allows for the potential users to gauge expectations and devise a priori alternative plans.

We attempted to comprehensively cover the caveats inherent in our genetic targeting approach.

Minor points:

(1) The authors mention that fly age should be controlled as expression can vary. Is there any reference to support this claim?

We added a reference describing driver expression changes over development.

(2) There should be a citation for "Flies were typically 1-5 days old at dissection for the cell type collection rescreening, 1-8 days old for other non-MCFO crosses and 3-8 days old for MCFO".

We clarified that these descriptions were of our experimental preparations, not describing other citable work.

Reviewer #3 (Recommendations For The Authors):

General Points:

Overall, the manuscript is very clear, but there are a couple of points where more explicit information would be useful. One of these is with respect to the issue of selectivity of targeting. The cell type specificity of lines is often referred to, but cell types can range from single pairs of neurons to hundreds of indistinguishable neurons with similar morphology and function. It would be useful if the authors explained whether their use of the term "cell type" distinguishes cell type from cell number. It would also be useful if lines that target many neurons of a single cell type were identified.

We added further discussion of cell types vs. cell numbers. Our labeling strategy was not optimized for counting cell numbers labeled by each line. We believe EM studies are best positioned to comprehensively evaluate the number of cells making up each type.

The second point relates to vagueness about the intended schedule for providing resources that will match (or allow matching of) neurons to the connectome. For example, on pp. 5-6 it is stated that: "In the future all of the neurons in these lines will be uniquely identified and linked to neurons reconstructed in the electron microscopy volume of the larva" but no timeline is provided. Similarly, for the adult neurons it is stated on p. 4 that: "Anatomical searching for comparison to other light microscopy (LM) and EM data is being made available." A more explicit statement about what resources are and are not yet available, a timeline for full availability, and an indication of how many lines currently have been matched to EM data would be helpful.

During the review and revision period we have made progress on processing the images in the collection. We updated the text with the current status and anticipated timeline for completion.

Specific Points:

p. 4 "Although the lines used for these comparisons are not a random sample, the areas of greatest difference are in the vicinity of previously described sexual dimorphisms..." In the vicinity of is a very vague statement of localization. A couple of examples of what is meant here would be useful.

We added example images to Figure 3.

p. 5 "...may have specific expression outside our regions of interest." It's not clear what "our regions of interest" refers to here. Please clarify.

We clarified that we were referring to the regions studied in the publications listed in Table 1.

p. 5 "...lines that were sparse in VNC but dirty in the brain or SEZ..." A more quantitative descriptor than "dirty" would be helpful.

We unfortunately did not quantify the extent of undesired brain/SEZ expression, but attempted to clarify the statement.

p. 6 "...the images are being made instantly searchable for LM and EM comparisons at NeuronBridge..." Here again it is hard to know what is meant by "being made instantly searchable." How many have been made searchable and what is the bottleneck in making the rest searchable?

We updated the text as described above. The bottleneck has been available processing capacity for the hundreds of thousands of included images.

Figure 1 Supplemental File 2: The movie is beautiful, but it seems more useful as art than as a reference. Perhaps converting it to a pdf of searchable images for each line would make it more useful.

We replaced the movie with a searchable PDF.

Fig. 2(B) legend: "Other lines may have more than two types." It is not clear what "other lines" are being referred to.

As part of making the quality evaluation more robust, we scored lines for the clear presence of three or more cell types. We updated the text accordingly.

Fig. 2(C): Presumably the image shown is an example of variability in expression rather than weakness, but it is hard to know without a point of comparison. Perhaps show the expression patterns of other samples? Or describe briefly in the legend what other samples looked like?

We added Figure 2 - figure supplement 1 with examples of variable expression in a split-GAL4 line.

<https://doi.org/10.7554/eLife.98405.2.sa0>