

## Reviewed Preprint

v1 • November 18, 2025

Not revised

## Reviewed Preprint

v2 • March 17, 2026

Revised by authors

### ✉ For correspondence:

[jj.phillips@exeter.ac.uk](mailto:jj.phillips@exeter.ac.uk)

[d.n.woolfson@bristol.ac.uk](mailto:d.n.woolfson@bristol.ac.uk)

[mark.dodding@bristol.ac.uk](mailto:mark.dodding@bristol.ac.uk)

\* These authors contributed equally.

### Competing interests:

No competing interests declared

### Funding:

See [page 30](#)

### Reviewing editor:

Julien Roche,  
Iowa State University, United States

© 2025, Shukla et al. This article is

distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted

use and redistribution provided that

the original author and source are

credited.

# Kinesin-1 conformational dynamics are controlled by a cargo-sensitive TPR switch

Shivam Shukla<sup>1,2,\*</sup>, Jessica A Cross<sup>1,2,3,\*</sup>, Monika Kish<sup>4,5,\*</sup>, Sathish KN Yadav<sup>1</sup>, Johannes F Weijman<sup>1</sup>, Laura O'Regan<sup>1</sup>, Judith Mantell<sup>1</sup>, Ufuk Borucu<sup>6</sup>, Xiyue Leng<sup>2</sup>, Christiane Schaffitzel<sup>1</sup>, Jonathan J Phillips<sup>4,5</sup> ✉, Derek N Woolfson<sup>1,2,7</sup> ✉, Mark P Dodding<sup>1</sup> ✉

<sup>1</sup>School of Biochemistry, University of Bristol, Bristol, United Kingdom • <sup>2</sup>School of Chemistry, University of Bristol, Bristol, United Kingdom • <sup>3</sup>School of Engineering Mathematics and Technology, University of Bristol, Bristol, United Kingdom • <sup>4</sup>Living Systems Institute, University of Exeter, Exeter, United Kingdom • <sup>5</sup>Department of Biosciences, University of Exeter, Exeter, United Kingdom • <sup>6</sup>GW4 Facility for High-Resolution Electron Cryo-Microscopy, University of Bristol, Bristol, United Kingdom • <sup>7</sup>Max Planck-Bristol Centre for Minimal Biology, University of Bristol, Bristol, United Kingdom

## eLife Assessment

The revised manuscript by Shukla et al. provides **important** mechanistic insights into kinesin-1 autoinhibition and cargo-mediated activation. Through a **convincing** integration of protein engineering, computational modeling, biophysical assays, HDX-MS, and electron microscopy, the study delineates how cargo binding induces an allosteric transition that propagates along the coiled-coil stalk to the motor domains, enhancing MAP7 engagement. The revisions substantially improve clarity, figure annotation, and methodological transparency, leaving the remaining limitations, primarily those inherent to conformational heterogeneity and resolution, appropriately acknowledged. Overall, the updated manuscript presents a coherent mechanism for kinesin-1 activation that will be of broad interest to the motor protein, structural biology, and cell biology communities.

<https://doi.org/10.7554/eLife.109462.2.sa4>

## Abstract

Kinesin-1 is a dynamic heterotetrameric assembly of two heavy and two light chains (KHC and KLC) that mediates microtubule-based intracellular transport of many different cargoes. The complex adopts a compact, autoinhibited state that is activated by cargo-adaptor proteins containing specific short linear peptide motifs (SLiMs). These motifs interact with the tetratricopeptide repeat (TPR) domains of the KLCs. The mechanism coupling SLiM recognition to activation-associated conformational changes in the complex is unknown. Here we combine protein design, computational modelling, biophysical analysis, and electron microscopy to examine the structural and mechanistic consequences of SLiM binding to the KLC-TPR domain within the complete heterotetrameric holoenzyme. We show that coiled coil 1 (CC1) of the KHC docks KLC TPR domains in the autoinhibited complex, forming the 'shoulder' feature observed in electron microscopy. Disrupting this interaction or binding an activating SLiM dislocates the TPR shoulder, freeing the motor domains and promoting transition between its closed, inactive, and open states. Opening the kinesin-1 complex facilitates binding to the microtubule-associated kinesin-1 cofactor, MAP7. Therefore, cargo-mediated dislocation of the TPR shoulder serves as a key initial step in kinesin-1 activation, allosterically linking cargo binding to motor dynamics.


## Introduction

Intracellular transport relies on the conformational dynamics of cytoskeletal motor proteins. One such process involves ATPase-driven binding and movement along cytoskeletal tracks. Another, less-understood process, involves shape changes in motor complexes, which regulate movement in response to inputs like cargo or regulatory protein binding, ensuring precise transport control (1–4). Conformational transitions between a compact inhibited state and an open active state are essential for the regulation of kinesin-1, a ubiquitous and prototypic family of microtubule motors involved in transporting proteins, ribonucleoproteins, vesicles, and organelles in cells (1, 2, 5–8). Kinesin-1 is also hijacked by pathogens during infection, and its dysregulation is linked to neurological disorders (9, 10).

Heterotetrameric kinesin-1 consists of two kinesin heavy chains (KHCs) and two kinesin light chains (KLCs). In mammals, KHCs are encoded by three paralogs—KIF5A, KIF5B, and KIF5C—while KLCs are encoded by four paralogs (KLC1–KLC4) (11–14). The KHCs have ATPase motor domains at the amino terminus, followed by coiled-coil domains (CC0–CC4) that mediate KHC dimerization, CC0 forms the neck between the amino-terminal motor domains and the coiled-coil stalk which comprises CC1 – CC4. The carboxy-terminus contains an unstructured domain with the autoinhibitory 'IAK' motif, which regulates ATPase activity in the compact state, but is not essential for its formation (15–23). The KLCs, which also help to regulate kinesin-1 activity, have a coiled-coil domain that binds to CC3 of the KHCs, followed by an unstructured linker to a tetratricopeptide repeat (TPR) domain critical for cargo binding (2, 24–26) (Fig. 1A). In several KLC isoforms, the TPR domain is followed by an intrinsically disordered domain, which includes a membrane-binding amphipathic helix that can stabilise interactions with membranous cargoes (27–29).

Kinesin-1 activity is regulated by its conformational state (6–8, 24, 30, 31). Recent studies using negative stain electron microscopy (NS-EM), AlphaFold2 modelling, small-angle X-ray scattering (SAXS) and cross-linking mass spectrometry have revealed the architecture of the compact, autoinhibited state, named the “lambda particle” (22, 32, 33). Key features of this particle include the ATPase motor domains, forming the “head,” and a flexion point in the coiled-coil scaffold between CC2 and CC3—the “elbow”—that enables the autoinhibited conformation to form (22, 32, 33). This flexion is functionally significant, as the complex can be activated or stabilized in its inhibited state through targeted mutations or binding of *de novo* designed peptide-based activators (32, 34). NS-EM and SAXS also reveal a “shoulder” near the motor domains. Domain-deletion experiments indicate that the shoulder is formed by at least one KLC TPR domain (32). Cross-linking mass spectrometry, NS-EM, and AlphaFold2 modelling suggest that the motor domains invert relative to CC1, binding back upon it and interact with CC4 in a hierarchical inhibition mechanism (22, 33) and propose many possible contacts between the KLC-TPR domains with the motor domains and coiled-coil body (22). Key features of the inhibited complex appear to be conserved across the closely related mammalian KHC paralogues (22, 32, 33).

Understanding the position and function of TPR domains in the inhibited state is crucial for uncovering how kinesin-1 is activated, as these domains provide binding sites for activating cargo adaptors with W-acidic SLiMs (2, 35, 36). W-acidic motifs bind the concave groove of the TPR superhelix, increase its curvature, and displace an autoregulatory interaction with the CC-TPR linker region to modulate KLC conformational state (24). This molecular switch is also the target for the kinesin-1 activating small-molecule kinesore (37). Adaptors carrying W-acidic motifs include the lysosomal adaptor SKIP (35, 38, 39), the nuclear envelope adaptors Nesprin-2/4 (40, 41), and the amyloid-precursor protein (APP) vesicle adaptor calsynenin-1 (42–44), and others (2). Recently, we have designed a small-peptide ligand, KinTag, for the TPR binding site. This combines features of natural micromolar affinity W-acidic and related Y-acidic motifs, to give high-affinity (low nanomolar) binding through simultaneous occupation of the W-, Y- and F- pockets on the TPR receptor (26, 45, 46). Notably, KinTag drives changes in TPR conformation like the natural ligands. Therefore, KinTag provides a useful tool to probe the mechanistic consequences of stable motor-cargo interactions, which in the natural system, are also enabled by other co-operative protein-



**Figure 1. Computational modelling and single particle EM analysis identify a TPR docking site (TDS) on CC1**

(A) Schematic representation of heterotetramer kinesin-1 in an open conformation. (B) AF3 model in 3 orientations alongside a schematic showing KHC(KIF5C) coiled-coil assembly with KLCs. KHCs are shown in blue and cyan, KLCs in orange and purple. The complete AF3 output, including models for the full tetramer, and sequences used are described in [Figure S1](#). Schematic shows predicted domain positioning, omitting linker sequences and separating the KHC coiled-coils for clarity (C) Representative NS electron micrograph (from 3 independent experiments) showing ElbowLock complexes (scale bar is 50 nm) with selected 2D classes from NS-EM (scale bar is 26 nm). (D) Reference-free 2D class averages showing two orientations of complete heterotetrameric kinesin-1 complexes, compared to low-pass filtered back-projection of the AF3 model (scale bar is 22 nm). (E) Structure of the nanobody-TPR complex highlighting binding on exterior of TPRs 4 and 5 (pdb:6fuz). Cargo JIP1 Y-acidic peptide was removed for clarity.(F) Representative NS electron micrograph showing ElbowLock-nanobody complexes (scale bar is 50 nm) with selected 2D classes from NS-EM (scale bar is 26 nm).

protein and protein-lipid interactions (27, 47). Importantly, when fused to an integral membrane protein, KinTag, or its parental W-acidic sequence, is sufficient to drive kinesin-organelle recruitment and axonal transport (36, 42, 48, 49). Thus, structural and biophysical analysis of isolated TPR domains in vitro and functional analysis of organelle transport in cells demonstrates key role for SLiM-TPR binding in motor activation and has defined some important components of the pathway. However, the mechanism that couples SLiM cargo recognition to activation-linked conformational change in the complete kinesin-1 holoenzyme remains unknown.

To define this mechanism, we use protein design and engineering approaches to overcome intrinsic conformational dynamics and low-affinity motor-cargo interactions to stabilize intermediates along the kinesin-1 inhibition-activation pathway, enabling their biochemical characterization and visualization. We show that the KLC TPR domains dock onto CC1 in the autoinhibited complex to form the shoulder. Disrupting this interaction or binding an activating SLiM dislocates the shoulder. This makes the motor domains more accessible and dynamic as judged by hydrogen-deuterium exchange measurements in the open, active, and cargo-bound states compared with the closed, inactive form. Opening the complex also promotes interaction with the microtubule-associated cofactor, MAP7. Therefore, cargo-mediated dislocation of the TPR shoulder serves as a key step in kinesin-1 activation, allosterically linking cargo recognition to conformational dynamics.

## Results

### Coiled coil 1 (CC1) provides a TPR docking site (TDS) in the autoinhibited kinesin-1 complex

To begin, we sought to clarify the position of the TPR domains in the lambda particle by modelling their potential for associations using AlphaFold3 (AF3). AF3-generated assemblies of the heterotetrameric complexes of the KHC coiled coils (KIF5C) and KLC (KLC1A), that included the TPR domains (KIF5C/KLC1A) yielded models that were invariably folded-over at the elbow, with a global coiled-coil architecture consistent with our previous work (32, 34) (Fig. 1B, Fig. S1A). Interestingly, the two TPR domains were confidently predicted to bind with C2 symmetry on CC1 ‘end on’ via the first repeats (TPR1) at a position concurrent with the shoulder observed in NS-EM. This TPR conformation closely resembles their known mode of binding to the coiled-coil leucine zipper motif of the cargo adaptor JIP3 (50). The concave cargo binding surface on the TPR domain that recognises SLiM adaptors was typically occupied by the linker region that connects the TPR to the KHC-KLC coiled-coil interface (24). Models of the complete heterotetramer, including the motor domains were similar. The motor domains were inverted around Hinge 1 enabling interactions between them and the first part of CC1 and the end of CC4 with the neck coiled coil (CC0) projecting away from the motors, in line with recent reports (22, 33) (Fig. S1B,C,D).

To test these predictions experimentally, we examined purified kinesin-1 complexes using negative stain electron microscopy (NS-EM) followed by 2D-classification of single particles (Fig. S2, S3, S4). We focused on the ElbowLock background. This KIF5C construct contains a short, 5 amino acid, deletion that restricts flexibility around the elbow and helps maintain particles in their lambda conformation, providing homogenous samples, and facilitating subsequent analysis (34). Proteins were also stabilised using the amine-to-amine crosslinker BS3 that was important for achieving reproducibly high-quality samples for imaging. As observed previously for closed wild-type particles, the TPR ‘shoulder’ feature was readily apparent, and typically appeared as a single globular density proximal to the two motor domains (22, 32) (Fig. 1C).

To visualise finer structural details, we turned to single-particle cryoEM analysis of frozen-hydrated samples. We were unable to obtain optimal samples suitable for determining the complete structure. Nonetheless, we obtained reference-free 2D class averages that appeared to show full-length ‘side’ views of the complex with clear definition of the elbow, hinge 2, and KHC-KLC (coiled-coil) interface features (Fig. 1D). The motor domains were poorly resolved in these

classes, suggesting that the head assembly is somewhat flexible relative to the coiled coil/TPR body. A comparison to low-pass filtered back-projections from the AF3 model (without motor domains) revealed density at a position concurrent with the docked TPR domains (Fig. 1D [↗](#)).


However, TPR domain stoichiometry at the shoulder remained unclear. We reasoned that if the AF3 models were correct then the latter TPR repeats (4–6) would be positioned away from the interface with the CC1. The KLC TPR domains are small and known to be conformationally dynamic, and so this might make it difficult to resolve them in 2D class averages (24, 35, 45). To test this hypothesis, we assembled complexes with a nanobody to bind and stabilise TPRs 4 and 5 and add additional mass (13 kDa per TPR) (36, 45) (Fig. 1E [↗](#), Fig. S2 [↗](#)). This resulted in NS-EM 2D classes with density extending either side of the coiled-coil scaffold that were fully consistent with the AF3 tetramer models and binding of both TPR domains on CC1 (Fig. 1E, F [↗](#)). There was no evidence of additional density elsewhere in the complex. Thus, AF3 predicts a model for TPR domain association with the coiled-coil scaffold that is supported by electron microscopy experiments. Because we have used a stabilisation approach, we do not exclude the possibility that there may be differences in TPR stability, dynamics and occupancy at the two potential CC1 sites in the absence of stabilisation. i.e. only one TPR may be typically bound at one time, and/or they may have potential for exchange. Hereafter, the predicted TPR docking site on CC1 is called the TDS.

## Removal of the TDS dislocates the kinesin-1 shoulder

To test the role of the TDS in positioning the KLC TPRs directly, we removed it while preserving the overall length and coiled-coil structure of CC1. To do this, four heptad repeats of CC1 of KIF5C, flanking and incorporating TDS, were replaced with a *de novo*-designed homodimeric coiled coil, CC-Di (51). The aim was to retain the dimeric coiled-coil structure and length of CC1 while eliminating side chains that promote its interface with the TPR ( $\Delta$ TDS) (Fig. 2A [↗](#)). Supporting this, models of  $\Delta$ TDS complexes using AF3 showed the expected seamless insertion of CCDi into CC1, with displacement of the TPR domains to a variety of different positions, in 5 models, all with high position error with respect to KHC (Fig. S5 [↗](#)). We used size-exclusion chromatography (SEC) to further validate our design. ElbowLock proteins elute as a single lambda peak and this elution profile was unchanged by the  $\Delta$ TDS modification, indicating that complex assembly or folding is not disrupted (Fig. 2B [↗](#)). ElbowLock- $\Delta$ TDS complexes were then analyzed by NS-EM and 2D classification (Fig. 2 C,D [↗](#)) and quantitatively compared to the parental ElbowLock (Fig. 2E [↗](#)). The shoulder was lost in the  $\Delta$ TDS background, suggesting that TPR domains had undocked from the core of the complex.

## The TDS binds directly to the KLC TPR domain

To determine whether the TPR binds TDS directly, we measured the interaction between fluorescently labelled synthetic peptides corresponding to the TDS region and the TPR domain directly using fluorescence polarisation (FP) assays. Short peptides from the TDS region were unfolded in the solution, which we attributed to an imperfect hydrophobic repeat pattern, lacking leucine residues typical of the cores of dimeric coiled coils. Therefore, we made 3 mutations of core residues (A479, V486, K500) to leucine and templated folding using 7 flanking residues on either side of the sequence from CC-Di (Fig. S6A [↗](#)). Circular dichroism (CD) spectroscopy and analytical ultracentrifugation (AUC) confirmed that the resulting peptide (Eng-TDS) formed a helical dimer (Fig. S6B, C, D, E [↗](#)). Subsequent FP assays showed that isolated KLC1-TPR interacted with Eng-TDS with low affinity (approx.  $K_d$  of 25  $\mu$ M), but not with CC-Di alone. (Fig. 2F [↗](#)). We obtained similar results for KLC2-TPR with tighter binding to Eng-TDS (Fig. S6F [↗](#)) ( $K_d$  = 2.7  $\mu$ M), suggesting that this interaction is conserved between KLC paralogues. We note that these binding measurements are for an inter-molecular process but in the context of the holoenzyme the interaction is effectively intra-molecular because the TPR domain is tethered to the coiled-coil scaffold. Therefore, the effective local concentration is substantially higher, resulting in a high equilibrium fraction of bound complexes, consistent with our EM experiments. Prior work shows that constructs with deletions of the first helix of TPR1 of KLC2 result in unfolding of TPR1 (35, 52). Therefore, to test a key role for the TPR1 interface, we deleted the first helix of TPR1. This



**Figure 2.** The CC1 TDS is required for the formation of the kinesin-1 shoulder.

(A) Schematic showing strategy to delete the TDS in CC1 and replace it with an equivalent length of a *de novo* designed homodimeric coiled coil, CC-Di. (B) SEC traces showing the elution profile of wild type, ElbowLock, and ElbowLock-ΔTDS proteins. Constructs in the ElbowLock background elute exclusively in the peak associated with closed lambda particles. Representative of 3 independent experiments. (C) Representative NS electron micrographs (from 3 independent experiments) showing ElbowLock and ElbowLock-ΔTDS complexes. Scale bar is 50 nm (D) Selected 2D classes from NS-EM highlighting the presence of the shoulder in ElbowLock but not ElbowLock-ΔTDS complexes. The scale bar is 26 nm. Full sets of classes are provided in Fig. S4 (E) Quantification of number of particles in classes with and without a prominent shoulder of 3 independent experiments. \*\*\* indicates  $p < 0.001$  (F) Fluorescence polarisation binding assays show fluorescently labelled TDS and SKIP (W-acidic motif, positive control) peptides, but not the control peptide CC-Di, bind to isolated TPR domains. Error bars show S.E.M. from 3 replicates.

effectively abrogated any detectable binding (Fig. S6G). Together, this modelling, electron microscopy analysis and biophysical experiments clearly demonstrate that the TDS provides a TPR domain docking site on the lambda particle.

## Cargo adaptor binding dislocates the TPR shoulder

To understand the consequences of cargo binding, we incorporated an activating W-/Y-acidic cargo adaptor ligand into the complex (KinTag) via fusion on a flexible linker at the carboxy-terminus of KLC; i.e., an intramolecular mimic of stable cargo binding, following a similar strategy used to solve the structure by X-ray crystallography (36) (Fig. 3A). In a wild-type background, this resulted in a shift in the SEC elution profile from the two peaks corresponding to open and closed conformations to a single peak at an intermediate position (Fig. 3B), which negative stain EM analysis revealed consisted of a mixture of open and closed particles, but with no discrete intermediate state (Fig. S7). From this, we conclude that the main effect of SLiM-TPR interaction is to increase the rate of exchange between open and closed conformations such that they are no longer resolved by SEC. Supporting this, additional incorporation of the closed-stabilising ElbowLock modification shifted the equilibrium to a position corresponding to the closed peak, demonstrating that the cargo-induced SEC shift can be suppressed by restricting conformational flexibility at the elbow, and therefore results from opening the complex (Fig. 3B).


Analysis of the resulting ElbowLock-KinTag particles by NS-EM and 2D classification resulted in clear loss of the TPR shoulder (Fig. 3C-E), showing that the intramolecular adaptor interaction promotes TPR dissociation from the closed complex. To substantiate this further, we generated 3D models of ElbowLock, ElbowLock-ΔTDS and ElbowLock-KinTag complexes from the NS-EM data (Fig. S8). For ElbowLock complexes, this resulted in classes with and without a prominent shoulder, in agreement with 2D classification. For ElbowLock-ΔTDS and ElbowLock-KinTag complexes, no prominent shoulder containing classes were observed. Moreover, these modifications appeared to result in a slight loosening of the whole coiled-coil assembly (Fig. 3F); note the indentations between the densities of the two main coiled-coil regions, which we interpret as the start of a separation of these two domains that is restricted by the ElbowLock mutation. Consistent with a KinTag-induced inhibition of TDS-TPR association, inclusion of KinTag on the KLC-TPR carboxy-terminus effectively abrogated any detectable binding to Eng-TDS, or its direct competitor, the W-acidic motif of SKIP, in FP assays (Fig. S6H). Together, these data support a model where adaptor-TPR interaction dislocates the TPR shoulder from the complex (Fig. 3G).

## Cargo-mediated dislocation of the shoulder enhances motor domain accessibility

To determine how the local structural changes from adaptor binding and shoulder dislocation affected the dynamics of kinesin-1 complexes in solution, as directly and least invasively as possible, and without the risk of cross-linker artefacts, we recorded millisecond time-resolved hydrogen/deuterium-exchange mass spectrometry (HDX-MS) time courses, resolved at the peptide level (Fig. S9, S10, Table S1) (53–57). Sequence coverage was good (overall 88%) with the exception of KHC-CC0 (neck coil) and the acidic-linker region that connects the KLC coiled-coil to the TPR domains where coverage was more limited.


To begin, we established baseline parameters by comparing the wild-type and DeltaElbow complexes. The latter stabilizes the fully extended conformation by promoting helical readthrough at the elbow and thus forces TPR undocking from CC1 (32, 34). The data for wild type were subtracted from those of the DeltaElbow to generate difference plots (Fig. S11A, S12A), which were then mapped onto the X-ray crystal structures of the motor and TPR domains (Fig. 4) (52, 58).

First, as a general observation for the KHC chain, the largest differences in HDX were observed outside of its coiled-coil (CC) stalk. This is consistent with hydrogen-bonded α-helical structure and the known stabilities of coiled-coil assemblies in general that is protective against hydrogen-



**Figure 3. Adaptor-binding to KLC-TPR dislocates the kinesin-1 shoulder.**

(A) Comparison of X-ray crystal structures of ligand-free (grey, pdb: 3nf1) and KinTag (W/Y acidic)-ligand bound (purple, pdb: 6swu) TPR domains highlighting the ligand-binding site and ligand-induced change in TPR curvature. (B) SEC traces showing elution profile of wild type, ElbowLock and ElbowLock-KinTag proteins. (C) Representative NS electron micrographs (from 3 independent experiments) showing ElbowLock and ElbowLock-KinTag complexes. Scale bar = 50 nm. (D) 2D classes from NS-EM highlighting the loss of the shoulder in ElbowLock-KinTag complexes. The scale bar is 26 nm. (E) Quantification of the percentage of particles in classes with and without a prominent shoulder from 3 independent experiments. \*\* indicates  $p < 0.001$  and \*\*\* indicates  $p < 0.001$  (F) 3D-reconstructions from the ElbowLock, ElbowLock- $\Delta$ TDS and ElbowLock-KinTag datasets show dislocation of the shoulder either upon deletion of the TDS or incorporation of KinTag. (G) Model showing dislocation of the shoulder induced upon KinTag binding.



**Figure 4. Adaptor-binding to the KLC TPR domain promotes motor accessibility.**

Summaries of comparative HDX-MS analysis of wild type, DeltaElbow, ElbowLock and ElbowLock-Kintag complexes. At each time point, the rates of exchange were subtracted as indicated (X minus Y) and filtered and filtered using a hybrid significance test (global significance and Welch's t-test), then transposed onto X-ray crystal structures of the motor domains (pdb: 3kin) and TPR domain (pdb: 3nf1). Grey indicates non-significant differences or limited peptide coverage. (A) Comparison of deltaElbow (X) against wild type (Y). (B) ElbowLock (X) against wild type (Y). (C) ElbowLock-Kintag (X) against ElbowLock (Y). Relative protection (blue) indicates less solvent exposure or increased H-bonding for X, while deprotection (red) indicates more solvent exposure or H-bond loss for X.

exchange and may partially mask changes in solvent accessibility (59). Nonetheless, there were several small but statistically significant differences in labelling between the two constructs in the KHC CCs consistent with the opening of the DeltaElbow complex. These included deprotection/increased accessibility at the proposed motor-CC1 interface (22, 33), the TDS, and the KHC-KLC interface in CC3, running into the amino terminus of CC4 (Fig. S11A). Interestingly, for the KLC chain in the open state, we observed more deprotection/increased accessibility in its CC and even protection at its amino terminus suggesting larger-scale activation associated structural changes in this component of the kinesin complex (Fig. S12A).

Second and strikingly, the DeltaElbow modification significantly increased HDX rates of the motor domains (Fig. 4A). This is consistent with their liberation from the folded-back and locked conformation of the lambda particle (22). The increase in HDX was not tightly localised. Rather, the motor domain as a whole was significantly deprotected. This is consistent with enhanced solvent accessibility and increased local subdomain dynamics throughout the motors in response to opening of the elbow. In addition, opening the complex resulted in significant deprotection/increased solvent accessibility of TPR1. This is consistent with its role in docking the TPR onto the CC assembly and its forced undocking by helical readthrough in the DeltaElbow variant. We also observed additional deprotection of TPR6, extending through to the C-terminal sequence that follows the TPR, indicating possible occlusion of these features in the inhibited state.

Next, we compared the ElbowLock (stabilised closed) and wild-type complexes. Opposite to DeltaElbow, both the motor domains and the TPR1 sites were more protected/less accessible in ElbowLock variant (Fig. 4B, S11B, S12B). There was also protection of the C-terminal tail of KHC. Thus, shifting the open-closed equilibrium to either of its extremes enables visualization of the accessibility of key interfaces in the complex that maintain autoinhibition, and support our findings emerging from AF3 modelling, negative-stain EM, and biophysical analyses.

Finally, we compared the ElbowLock-KinTag and ElbowLock complexes to examine the consequences of adaptor binding in the closed state (Fig. 4C, S11C, S12C). As expected, inclusion of KinTag significantly decreased HDX in, and therefore stabilized, the central TPR repeats at its structurally characterised binding site (36). This was juxtaposed with the striking deprotection of TPR1, which is consistent with ligand-induced TPR domain undocking from the lambda particle at this epitope (Fig. 3). The motor domains were also deprotected. Changes on the KHC CC were again much lower in magnitude, but we did note statistically significant deprotection at the predicted motor-CC1 docking site and C-terminal tail, suggesting that TPR-adaptor binding triggers release of the motors from their folded back, locked conformation.

Together, these data provide direct in-solution measurements revealing both predicted and unanticipated changes in solvent accessibility associated with the opening and closing of the complex that demonstrate allosteric coupling between adaptor-induced dislocation of the TPR shoulder and increased motor domain accessibility.

## Opening the kinesin-1 complex enhances its association with MAP7

Thus far, our findings support a model where cargo-binding destabilises the compact conformation and so, may unmask protein-protein interaction interfaces that facilitate subsequent steps in the activation pathway. The microtubule associated protein MAP7 is an essential kinesin-1 cofactor that promotes binding to microtubules and activation. Its binding site has been biochemically mapped to a short region of CC1 that overlaps the TDS (60–67). Indeed, AlphaFold3 predictions suggest that although the TPRs and MAP7 may bind to CC1 via distinct mechanisms (near perpendicular TPR helix-turn-helix motif vs parallel single MAP7 alpha helix), their binding sites directly overlap, and as such, they may compete for KHC binding (Fig. 5A). One prediction of this model is that opening of the complex at the Elbow to force TPR undocking would enhance binding to MAP7. To test this, we immunoprecipitated GFP-tagged MAP7 from cells expressing wildtype, DeltaElbow and ElbowLock kinesin-1 complexes. This resulted in a roughly 5-fold increase in kinesin-1 associated with MAP7 in the DeltaElbow background, compared to wildtype


or ElbowLock constructs (Fig 5B,C). We also noted slightly elevated expression of ElbowLock complexes and slightly lower expression of DeltaElbow complexes, suggesting that opening/closing of the complex could impact on kinesin-1 turnover. Similarly, in immunofluorescence experiments, the DeltaElbow modification resulted in a striking increase in kinesin-1 associated with MAP7-positive microtubules and pronounced microtubule bundling (Fig 5D). Together, these data reveal a coupling between kinesin-1 conformational state and its capacity to associate with the microtubule associated co-factor MAP7.

## Discussion

Over the past two decades, studies have identified a key role for SLiM-containing adaptor proteins in recruiting and activating kinesin-1 by binding to the KLC-TPR domains (2, 26, 35, 36, 39, 40, 42–45). Recent work has revealed the low-resolution architecture of the autoinhibited lambda state, including the TPR shoulder resting on the coiled-coil body (22, 32, 33). Increasing evidence suggests that a balanced equilibrium between inhibited and active states ensures responsiveness to stimuli (34, 40, 68). The challenge is to understand how adaptor binding to the TPR drives the transition from the closed, inhibited state to the open, active state; intrinsic conformational dynamics and unitarily low-affinity, cooperative, motor-cargo interactions obscure key intermediates in this pathway. To address this here, we have used *de novo* protein design and protein engineering to stabilise these intermediates alongside biophysical, biochemical, and electron microscopy measurements of the complete heterotetrametric holoenzyme. Our results lead us to propose a model where adaptor binding dislocates the TPR shoulder, destabilizing the autoinhibited state, enhancing motor domain accessibility and promoting the transition from the inactive closed state to the open forms (Fig. 6).


Integration of the present findings with prior work suggests a unified model for the role of TPR binding cargo adaptors in motor activation. We reveal that TPR1, the first TPR repeat, forms a key interface for TPR domain-CC association. TPR1 also binds the adaptor protein JIP3 via its coiled-coil leucine zipper, resembling TDS interaction predictions. However, a detailed evolutionary analysis in the same study suggests that TPR1 plays a distinct role beyond cargo binding (50). Moreover, a comparison of high-resolution X-ray crystal structures of KLC-TPR domains, with and without SLiM cargo, demonstrates structural plasticity in TPR1, which may enable ligand binding to modulate TPR1 orientation alongside larger scale conformational changes resulting from an increase in TPR domain curvature in response to SLiM binding (24, 35, 45, 69). We propose that SLiM ligand binding to the TPR groove drives conformational changes that reduces TPR1 affinity for the TDS to dislocate the shoulder; JIP3 may compete for the same site on TPR1 to the same effect. Our previous work has shown that SLiM-TPR interaction gates access to important cargo and microtubule binding sites on CC4, that is predicted to be closely juxtaposed to CC1 in models of the inhibited state (24, 37, 47). Our computational predictions do not indicate a direct interaction between KLC TPR and CC4 although they are predicted to be in proximity. Whilst we do not exclude this possibility of a direct interaction, we favour a model where removal of steric blocking effects from the bulky TPRs and docked motors on CC1, as well as opening of the coiled-coil assembly enables subsequent steps such as cargo binding to additional sites on CC4.

The TDS is located within biochemically defined binding site for the essential kinesin-1 cofactor MAP7 (60–67). Recent studies show clear cooperativity between MAP7 and W-acidic SLiM adaptors in motor activation (40, 70). This work provides a compelling explanation for this co-operativity: shoulder dislocation would be predicted to destabilise autoinhibition and expose the MAP7 binding site, facilitating subsequent activation steps such as enhanced microtubule recruitment and transition to and/or maintenance of the fully extended state (Fig. 6). Consistent with this, our models suggest that CC1 binding to MAP7 or KLC TPR would be mutually exclusive if the symmetrical binding sites on CC1 were both occupied (2xMAP7 or 2xKLC-TPR). However, it is not clear at this point whether this is the case. In the absence of nanobody-mediated TPR stabilisation, negative stain EM hints at an asymmetric TPR conformation and we remain open to the possibility



**Figure 5. Opening the kinesin-1 complex promotes its association with MAP7.**

(A) Overlay of two AlphaFold3 models of a tetrameric CC1/TPR assembly and a tetrameric CC1/MAP7-kinesin binding domain assembly, aligned on CC1. For clarity, only one CC1 model is shown. (B) GFP-MAP7 immunoprecipitation experiments showing enhanced association of KHC and KLC in the DeltaElbow background. Cells were transfected with the indicated constructs (GFP/GFP-MAP7, HA-KLC2, HA-KIF5C (WT, ElbowLock, DeltaElbow)). Complexes were immunoprecipitated using GFP-TRAP beads, and input and bound samples were analysed using western blotting with anti-GFP and anti-HA antibodies. (C) Quantification of b. from 4 independent experiments. Error bars show S.E.M. \*\*\*= $p < 0.001$  using 1-way ANOVA with Tukey's multiple comparison test. (D) Immunofluorescence images of HeLa cells transfected with GFP-MAP7, HA-KHC, and KLC2-Halo (TMR-labelled) showing enhance association of kinesin-1 with GFP-MAP7 positive microtubules in the DeltaElbow background. Images are representative of 3 independent experiments. Scale bar is 20  $\mu$ m.



**Figure 6. Model for cargo-mediated initiation of kinesin-1 activation.**

In the autoinhibited state, a TPR domain(s) are docked at the TDS forming the shoulder. i. Binding to the cargo adaptor dislocates the TPR shoulder and promotes motor domain accessibility, and initiates separation of the coiled-coil domains (ii). This allosteric coupling between cargo and motor would facilitate recruitment to microtubules (supported by MAP7) and subsequent transport (iii).

that only one TPR is bound at one time and that they may exchange. This would, in principle, leave one CC1 binding site free to interact with MAP7 and one TPR domain free to make first contact with cargo.

It will be important to understand this dynamic interplay and to sequence these events. A cargo initiated opening up of the complex is consistent a model for MAP7 function where it facilitates tethered diffusion to bypass obstacles on the microtubule and enhances kinesin-1 processivity (61, 62), and activation without cargo-recognition would seem counter-productive. However, recently, MAP7 binding to KHC was shown to promote KLC binding to W-acidic and Y-acidic cargo (70). Here, we have used a high-affinity ligand to mimic stable cargo attachment, but natural motifs typically bind with low affinity and act cooperatively with additional protein-protein and protein-lipid interactions (27, 35, 39, 47, 71). Therefore, one might also consider a two-way switch model where MAP7 binding also facilitates shoulder dislocation to modulate cargo binding properties of the KLC TPR domains. In either case, this points to a critical interplay between regulator and cargo, converging on CC1, establishing a new target for intervention in kinesin-1 mediated transport processes, and design of new reagents to target this site will likely be critical to dissect its dynamic functions.

In summary, our findings demonstrate that kinesin-1 autoinhibition and initiation of activation by SLiM-cargo pivots on the association and dislocation of its TPR shoulder from the coiled-coil scaffold. One important implication of these findings is that kinesin-1 is unlikely to be stably inhibited when bound to cargo but instead may be primed for further regulatory inputs that control its activity. This may be important for the coordination of bidirectional transport in multi-motor/adaptor systems.

## Materials and Methods

### AlphaFold3 modelling

All of the models shown in the manuscript were predicted using the AlphaFold3 server (72). The algorithm was asked to model a heterotetramer composed of KIF5C (NP\_001101200.1) and KLC1A (NP\_032476.2) with varying input sequences described in the figure legends. Predicted aligned error statistics are as output by the software with manual annotations to highlight important features. For the models presented in Fig. 5 (3), the algorithm was asked to model a heterotetramer of KIF5C CC1 (S410-G558) with the kinesin binding helix of MAP7 (NP\_001185564.1, P467-K610), or KLC1 TPR domain (NP\_032476.2, G205-K495). Similar models and predicted binding sites were obtained for heterotrimers (2xCC1, 1xMAP7/TPR). Output models were visualised/aligned and prepared for publication using in UCSF Chimera (73).

### Constructs for protein expression

A 2-plasmid system to express heterotetrameric kinesin-1 in *E. coli* cells has been described previously (32, 34) and constructs used in this study were derived from these. Wild-type kinesin heavy chain was KIF5C expressed in the pMW bacterial expression vector with an N-terminal 6X His tag. KLC1 was expressed pET28a with the His tag removed by site-directed mutagenesis. ElbowLock was originally designed by deleting five amino acids from the elbow loop (E685-L690) (34), and ElbowLock-ΔTDS was prepared by additionally replacing E475-S501 in CC1 with 4-heptads of CC-Di through subcloning of a synthetic gene fragment. KLC1-KinTag construct was prepared by fusing KinTag peptide via a (Thr-Gly-Ser)<sub>10</sub>-Gly flexible linker at C-terminal of KLC1 (36). The TPR domain of KLC1 (A211-A495) and KLC2 (A196-K480, delta helix 1 P219-K480) were cloned in the bacterial expression vector pET28His-Thrombin and have been described previously (24, 35). Anti-TPR nanobody (sequence from pdb: 6fv0)(45) was codon optimised and synthesised and cloned into NcoI and XhoI site of pet22b with N-terminal pelB sequence and C-terminal 6X His tag, by Genscript Inc. The identity of all plasmids was confirmed by DNA sequencing.

## Protein expression and purification

All kinesin-1 constructs were expressed in *E. coli* BL21(DE3) cells as a heterotetramer adapting the two-plasmid system as described previously (32, 34). Briefly, expression plasmids encoding individual KIF5C constructs and KLC1 constructs were transformed into *E. coli* BL21(DE3). Positive clones were selected based on a double antibiotic selection marker. Cells were cultured in LB medium (miller) supplemented with ampicillin (50 µg/ml) and kanamycin (50 µg/ml) in a shaking incubator under conditions of 37°C/180 rpm. When the optical density (OD 600) measured at 600 nm reached 0.6, the temperature was reduced to 18°C. Protein expression was induced by addition of 0.2 µM IPTG. Cultures were incubated overnight with shaking at 18°C. Cells were harvested by centrifugation at 5000 g for 15 min at 4°C. Cells were resuspended in Buffer A (40 mM Hepes (pH 7.4), 500 mM NaCl, 40 mM imidazole, 5% glycerol, and 5 mM β-mercaptoethanol), supplemented with protease inhibitors. Lysis was carried out by sonication (70% amplitude, 5s on/15 s off) for 7.5 min on ice. Lysates were centrifuged at 40,000g for 45 min at 4°C to obtain supernatant and filtered through a 0.45 µm membrane. The supernatant was loaded onto a His-Trap (1ml) column equilibrated with Buffer A. The column was washed with at least 50 CV of Buffer A with 60 mM Imidazole. Bound proteins were eluted using a gradient of Buffer A with increasing concentrations up to 500 mM of Imidazole. Eluted protein fractions were analysed on SDS-PAGE, then concentrated using a 30,000-Da MWCO ultrafiltration device up to 500 µl and immediately loaded on a size exclusion chromatography (SEC) column. SEC was carried out using a Superose 6 10/300 column pre-equilibrated with 20 mM Hepes (pH 7.4), 150 mM NaCl, 1 mM MgCl<sub>2</sub>, 0.1 mM ADP, and 0.5 mM TCEP. Eluted protein fractions were analysed using SDS-PAGE. Appropriate fractions were aliquoted and stored at -80°C freezer and used further for downstream applications.

TPR constructs were expressed in *E. coli* BL21(DE3) and affinity purified in buffer (25 mM Hepes pH 7.4, 500 mM NaCl, 5 mM β-mercaptoethanol) over a gradient of 20 mM to 500 mM imidazole on a His-Trap (1ml) column (24, 35). SEC was carried out using a HiLoad 16/600 Superdex 75 prep grade column (GE Healthcare) equilibrated with 25 mM HEPES (pH 7.4), 500 mM NaCl, 5 mM β-mercaptoethanol. Eluted fractions were analysed on SDS-PAGE.

Nanobody was expressed in *E. coli* BL21(DE3) and isolated from periplasmic space in 0.2 M Tris pH 8, 0.5 M EDTA, 0.5 M Sucrose and purified on His-Trap (1ml) column. SEC was carried out using Superose 6 10/300 column pre-equilibrated with PBS. Eluted fractions were concentrated and mixed in 10-fold excess with SEC purified ElbowLock (5.56 µM: 66 µM). Eluted complexes were analysed on SDS-PAGE.

## Negative stain electron microscopy

Protein aliquots (50 µl) were thawed from -80°C storage and cross-linked with 0.6 µl of 50 mM BS3 (final concentration 0.59 mM) at room temperature for 30 minutes. After cross-linking, the protein sample was placed on ice, and serial dilutions were prepared to achieve optimal single-particle distribution on grids. Formvar/carbon-coated 300-mesh copper grids were glow-discharged for 30 seconds at 20 mA. A 5 µl aliquot of the protein sample was applied to the grid and incubated for 1 minute before blotting with filter paper. The grid was subsequently stained by sequentially picking and blotting into three 5 µl drops of 3% uranyl acetate. The first drop served as a quick rinse, the second was left on the grid for 2 minutes, and the third also served as a quick rinse. The prepared grids were air-dried and subsequently imaged on a Talos L120C transmission electron microscope operated at 120 kV, equipped with a 4k × 4k Ceta CMOS camera. Micrographs were acquired at a magnification of ×57,000, corresponding to a pixel size of 2.48 Å/pixel. Datasets of approximately 500 micrographs were acquired using EPU automated collection software for all samples. The total electron dose was 52 e<sup>-</sup>/Å<sup>2</sup>, and micrographs were recorded with defocus values ranging from -1.5 µm to -2.5 µm.

## Negative stain electron microscopy image processing

Micrographs were analysed using CryoSPARC (74). After importing the micrographs, CTF correction was performed using the Patch CTF tool. Approximately 1,000 particles were manually picked and subjected to 2D classification to generate templates for automated particle picking. Using these templates, particles were autopicked, extracted (63.4 nm box size), and fourier-cropped to a pixel size of 4.96 Å/pixel. Extracted particles were 2D classified into 200 classes. A cleanup process was conducted by discarding particles that did not resemble kinesins through iterative rounds of curation and 2D classification. The number of particles in classes showing full ‘side views’ of the complex were counted and distinguished as particles with or without a prominent shoulder. The data quantified in this way was statistically analysed using 2-way ANOVA (using Sidak’s multiple comparison test). For figure presentation only, and to allow for intuitive comparison to AlphaFold models, final particle sets were reclassified with the CryoSPARC ‘align filament classes vertically’ option ticked. This resulted in no obvious morphological differences, but produced classes aligned on long axis of the complex. For 3D reconstruction, cleaned particle sets were used to generate four ab-initio models in CryoSPARC. A heterogeneous refinement of ab-initio models was performed, followed by homogeneous refinement, with alignment resolution restricted to 20 Å. Resulting models were low-pass filtered to 40 Å resolution and visualized using ChimeraX (73).

## Cryo-electron microscopy and image processing

Purified cross-linked ElbowLock complexes (5 µl at 3 mg/ml) were applied to glow-discharged Quantifoil R1.2/1.3 200-mesh Cu grids. The grids were blotted for 2 seconds at 4 °C and 100% humidity, then flash-frozen in 37% ethane/propane mix kept at approximately at –195°C using an FEI Vitrobot Mark IV (Thermo Fisher). Images were acquired on a Talos Artica microscope (FEI) operating at 200 kV with a Gatan K2 Summit direct detector. Automatic image collection was performed using EPU software (Thermo Fisher). A total of 3943 micrographs were captured with a total dose of 58 e/Å<sup>2</sup>, dose-fractionated into 64 movie frames at a defocus ranging from –1 µm to –2.5 µm. Images were recorded at 130,000× nominal magnification, resulting in a pixel size of 1.05 Å per pixel. Data processing was carried out in CryoSPARC (74). Motion correction and CTF estimation were performed using Patch Motion Correction and Patch CTF, respectively. Particles (focussing on full length side views) were manually picked from selected micrographs to create templates for automated particle picking. The autopicked particles were extracted and binned to 4.2 Å/pixel with a box size of 128 pixels / 53.76 nm. Extracted particles underwent multiple rounds of 2D classification to produce the class averages shown in Fig. 1. For comparison to the AF3 model, simulated density was generated using the molmap command in ChimeraX (73) filtering to 15 Å, and projections were generated/selected automatically using the Reference Based Auto Selected 2D function in CryoSPARC.

## Hydrogen-deuterium exchange mass spectrometry

Hydrogen-deuterium exchange (HDX) was conducted using “ms2min”, a fully automated, millisecond-resolution HDX labelling online quench-flow device (Applied Photophysics Ltd), which was connected to a Waters HDX manager. In the labelling experiments, 14 µL (10 µM) of DeltaElbow, wildtype, ElbowLock and ElbowLock-KinTag was introduced into the labelling mixer separately. The HDX reaction was initiated by diluting the sample 20-fold with a labelling buffer at 20°C. This buffer consisted of 20 mM HEPES, 150 mM NaCl, 1 mM MgCl<sub>2</sub>, 0.1 mM ADP, 0.5 mM TCEP in D<sub>2</sub>O, adjusted to a pH of 7.40 at 20°C. Samples were labelled at seven different time points (0.3, 0.5, 1, 3, 10, 30, 300) in triplicate. The HDX reaction was immediately quenched by mixing 1:1 with quench buffer (8 M urea, pH 2.55, at 0°C), and the sample was digested online using a Waters Enzymate pepsin column. The resulting peptides were trapped on a 2.1 × 5 mm VanGuard ACQUITY BEH C18 column (Waters) for 3 minutes at a flow rate of 155 µL/min and then separated on a 1 × 100 mm ACQUITY BEH C18 column (1.7 µm particle size) using a 7-minute linear gradient of 5–40% acetonitrile with 0.1% formic acid. Peptides were analysed using a Synapt G2-Si mass spectrometer (Waters) in HDMSE mode across a mass range of 50–2000 m/z. Instrument settings

were as follows: capillary voltage of 3.0 kV, cone voltage of 50 V, trap collision energy of 4 V, traveling wave ion mobility with a speed of 475 m/s, wave amplitude of 36.5 V, and nitrogen pressure of 2.75 mbar. Low-energy scans applied a transfer collision energy of 4 V, while high-energy scans utilized four distinct collision energy ramps between 15 and 55 V.

ProteinLynx Global Server (PLGS 2.5.1, Waters) was employed to analyse MSE reference data and to identify all detectable peptic peptides. The raw data files were processed, and isotopic distributions assigned using DynamX 3.0 (Waters, USA). The mean values and standard deviations (SD) of these replicates were calculated to assess the significance of changes, with a global significance threshold applied and a T-test (75). The peptide level data were also flattened per amino acid (76, 77). Custom scripts developed in Python 3.4.0 (Python Software Foundation) were utilized for post-processing analysis, and visualizations were generated using PyMOL (Schrödinger, Inc.).

## Peptide synthesis

Standard Fmoc solid-phase peptide synthesis was performed on a 0.1 mM scale using CEM (Buckingham, UK) Liberty Blue automated peptide synthesis apparatus with inline UV monitoring. Activation was achieved with DIC/Oxyma. Fmoc deprotection was performed with 20% v/v morpholine/DMF. Double couplings were used for  $\beta$ -branched residues and the subsequent amino acid. Peptides were synthesised from C to N terminus as the C-terminal amide on Rink amide resin. For fluorescently labelled peptides TAMRA (0.1 mM, 2 eq.), HATU (0.095 mM, 1.9 eq.) and DIPEA (0.225 mM, 4.5 eq) in DMF (3 mL) were added to DMF washed peptide resin (0.05 mM) with agitation for 3 hours. Resin was washed with 20% piperidine in DMF (5 mL) for 2 x 30 minutes to remove any excess dye. All manipulations were carried out under foil to exclude light. Peptides were cleaved from the solid support by addition of TFA (9.5 mL), TIPS (0.25 mL) and water (0.25 mL) for 3 hours with shaking at rt. The cleavage solution was reduced to approximately 1 mL under a flow of nitrogen. Crude peptide was precipitated upon addition of ice-cold diethyl ether (40 mL) and recovered via centrifugation. The resulting precipitant was dissolved in 1:1 acetonitrile and water ( $\approx$  15 mL) and lyophilised to yield crude peptide as a solid.

## Peptide purification

Peptides were purified by reverse phase HPLC on a Phenomenex (Macclesfield, UK) Luna C18 stationary phase column (150 x 10 mm, 5  $\mu$ m particle size, 100 Å pore size) using a preparative JASCO HPLC system. A linear gradient of 20-80% acetonitrile and water was applied over 30 minutes. Chromatograms were monitored at wavelengths of 220 and 280 nm. The identities of the peptides were confirmed using MALDI-TOF mass spectrometer using a Bruker ultrafleXtreme II instrument in reflector mode. Peptides were spotted on a groundsteel target plate using dihydroxybenzoic acid as the matrix. Peptide purities were determined using a JASCO analytical HPLC system, fitted with a reverse-phase Kinetex® C18 analytical column (Phenomenex, 5  $\mu$ m particle size, 100 Å pore size, 100 x 4.6 mm). Fractions containing pure peptide were pooled and lyophilised. Peptides were dissolved in buffer (25 mM HEPES pH 7.4 buffer with 150mM NaCl, 5mM BME) and their concentrations determined by UV-Vis on a ThermoScientific (Hemel Hemstead, UK) Nanodrop 2000 spectrophotometer using measurement of UV absorbance at 555 nm ( $\epsilon_{555}$ (TAMRA) = 85 000 mol<sup>-1</sup> cm<sup>-1</sup>).

## Circular-dichroism spectroscopy

CD data were collected on a JASCO J-810 or J-815 spectropolarimeter fitted with a Peltier temperature controller (Jasco UK). Peptide samples were dissolved at 100  $\mu$ M concentration in PBS (8.2 mM sodium phosphate, 1.8 mM potassium phosphate, 137 mM sodium chloride, 2.7 mM potassium chloride at pH 7.4). CD spectra were recorded in 1-mm path length quartz cuvettes at 20 °C. The instruments were set with a scan rate of 100 nm min<sup>-1</sup>, a 1-nm interval, a 1-nm bandwidth and a 1-s response time and scans are an average of 8 scans recorded for the same sample. The

spectra were converted from ellipticities (deg) to mean residue ellipticities (MRE;  $\text{deg.cm}^2.\text{dmol}^{-1}.\text{res}^{-1}$ ) by normalising for concentration of peptide bonds and the cell path length using the equation:

$$MRE(\text{deg.cm}^2.\text{dmol}^{-1}.\text{res}^{-1}) = \frac{\theta \times 100}{c \times l \times b}$$

Where the variable  $\theta$  is the measured difference in absorbed circularly polarised light in millidegrees,  $c$  is the millimolar concentration of the specimen,  $l$  is the path-length of the cuvette in centimetres and  $b$  is the number of amide bonds in the polypeptide, for which the N-terminal acetyl bond was included but not the C-terminal amide.

## Sedimentation-equilibrium analytical ultracentrifugation

Analytical ultracentrifugation (AUC) sedimentation-equilibrium experiments were conducted at 20 °C in a Beckman Optima XL-I analytical ultracentrifuge using an An-60 Ti rotor (Beckmann Coulter). Solutions were made up in PBS at 100  $\mu\text{M}$  total peptide concentration. The experiments were run in two-channel centrepiece. The samples were centrifuged at speeds in the range 44–60 krpm and scans at each recorded speed were duplicated. Data were fitted to single, ideal species models using SEDFIT (v15.2b)/SEDPHAT, comprising a minimum of four speeds. 95% confidence limits were obtained via Monte Carlo analysis of the obtained fits.


## Fluorescence polarisation

TAMRA conjugated peptides were diluted to 150 nM and incubated with increasing concentrations of KLC-1/2 TPR protein (typically 0 – 25  $\mu\text{M}$ ) in assay buffer (25mM HEPES pH 7.4, 5 mM  $\beta$ -mercaptoethanol and 150mM NaCl). Measurements were performed on a CLARIOstar (BMG Labtech) microplate reader at room temperature.  $\Delta\text{FP}$  values at each concentration were calculated by subtraction of measurements made without TPR protein. Binding constants ( $K_d$ ) were determined using a one-site specific binding model using GraphPad Prism software. Data were normalized to the calculated  $B_{\text{max}}$ . Where binding was lost or undetectable (e.g. delta Helix1, KinTag-fusion), data were normalised to the  $B_{\text{max}}$  of the parental protein (e.g. full length KLC1/2 TPR).


## Cell Culture, Immunoprecipitation and Fluorescence Imaging

HeLa cells were grown and maintained in high-glucose Dulbecco's modified Eagle's medium (Sigma-Aldrich) supplemented with 10% fetal bovine serum (Sigma-Aldrich) and 1% penicillin/streptomycin (Gibco) at 37°C with 5% CO<sub>2</sub>. For immunoprecipitation experiments, cells were transfected with plasmids encoding GFP-MAP7 (Addgene plasmid 46076)(64) with HA-KLC2 and HA-KIF5C (47). Cell lysis and GFP-TRAP immunoprecipitation experiments were performed as described in (24). For fluorescence imaging, cells were transfected with HA-KIF5C, KLC2-Halo(24), and GFP-MAP7, and labelled with Halo-TMR ligand as previously described (24), before fixation and staining with Mouse anti-HA (HA-7) from Sigma-Aldrich, and Alexa 647-conjugated anti-mouse (Thermo Fisher Scientific, and imaging on a confocal microscope.

# Supplementary Figures




**Figure S1. AlphaFold3 models of KHC-KLC complexes.** (A) Models of kinesin-1 heterotetramer coiled-coil domains (KIF5C CC1-CC4; aa 410-917) with two KLCs (KLC1A CC-TPR; aa 1-480). (B) Models of complete full-length (KIF5C/KLC1A) kinesin-1 heterotetramers, including motor domains and C-terminal KLC sequences. (C) Representative predicted aligned error plot highlighting confidence in intra and inter domain interactions in a full length heterotetrameric model. (D) Representative model showing pLDDT. Plots are typical of other models.




**Figure S2. Representative samples of protein complexes used in this study.**

Coomassie-stained SDS-PAGE gels show samples of proteins after size-exclusion chromatography, with and without BS3 crosslinker (top panel). Coomassie-stained SDS-PAGE gel showing purification of ElbowLock-Nanobody complex from size-exclusion chromatography (bottom panel).




**Figure S3. Representative NS-EM processing workflow.**

Schematic shows typical workflow for 2D classification steps to process NS-EM. Example shown is for ElbowLock complexes.




**Figure S4. Representative sets of 2D classes for all complexes.**

The scale bar is 26 nm.



**Figure S5. AlphaFold3 models of  $\Delta$ TDS kinesin-1 complexes.**

(A) 5 models showing  $\Delta$ TDS complexes (CCDi replaces the TDS sequence showing in green). In all models, the TPR domains are displaced from their original CC1-docked position (compare with Figure 1 [S1](#)) and occupy a variety of different positions with very low confidence (B) Representative position error plot (from model 1). Box highlights low confidence in TPR positioning.




**Figure S6.** Eng-TDS design, characterisation and interaction with KLC.


(A) Schematic showing design of Eng-TDS construct with flanking CCDi sequence and leucine mutations in the hydrophobic core. (B) CD analysis of Eng-TDS indicates alpha helical structure. (C) AUC analysis of Eng-TDS indicates assembly of a dimeric coiled-coil. (D, E) Chromatogram showing the purification of Eng-TDS, accompanied by mass spectrometry analysis confirming a molecular mass of 4895 Da. (F) FP assay showing binding of fluorescently labelled peptides from Eng-TDS and SKIP W-acidic motif to KLC2. (G) FP assay showing no detectable binding of Eng-TDS to KLC2 lacking the first helix of TPR1. (H) Fluorescence polarisation binding assays show no detectable binding of fluorescently labelled TDS and SKIP (W-acidic motif) peptides to isolated KLC1-TPR-KinTag fusion proteins.

**Figure S7.** Representative NS-EM electron micrograph showing wild type-KinTag particles with selected open and close particles expanded on right.


The scale bar is 50 nm.




**Figure S8.** EM processing workflow for 3D reconstructions.




**Figure S9.** Schematic representation of the HDX experiment.




**Figure S10.** Coverage map of mass spectral assignment of peptides in 'bottom-up' HDX-MS experiments for kinesin-1 HC (A) and LC (B).





**Figure S11. Difference plots showing total HDX across timepoints comparing heavy chain of DeltaElbow and Elbowlock to wild type and ElbowLock-KinTag to ElbowLock.**

The difference in the total observed deuterium uptake at 0.3, 0.5, 1, 3, 10, 30, and 300 seconds was summed and compared between DeltaElbow and wild type (A) and ElbowLock and wild type (B), and ElbowLock-KinTag to ElbowLock (C). In each case, the data for the later were subtracted from the former. Therefore, relative protection in a region of the former results in a more negative value (blue bars extending downward), while deprotection, such as from an exposed domain interface, results in a more positive value (red bars extending upward). Each vertical bar represents a distinct peptide, and the horizontal axis corresponds to amino acid number from N to C. Significant differences in deuterium labelling per peptide and time point, derived from a T-test, are marked with pink asterisks.



**Figure S12.** Difference plots showing total HDX across all timepoints comparing light chain of DeltaElbow and Elbowlock to wild type and ElbowLock-KinTag to ElbowLock.

The difference in the total observed deuterium uptake at 0.3, 0.5, 1, 3, 10, 30, and 300 seconds was summed and compared between DeltaElbow and wild type (A) and ElbowLock and wildtype (B), and ElbowLock-KinTag to ElbowLock (C). In each case, the data for the later were subtracted from the former. Therefore, relative protection in a region of the former results in a more negative value (blue bars extending downward), while deprotection, such as from an exposed domain interface, results in a more positive value (red bars extending upward). Each vertical bar represents a distinct peptide, and the horizontal axis corresponds to amino acid number from N to C. Significant differences in deuterium labelling per peptide and time point, derived from a T-test, are marked with pink asterisks.

|                                                  |                                                                                                               |
|--------------------------------------------------|---------------------------------------------------------------------------------------------------------------|
| Data Set                                         | Wild type; DeltaElbow; ElbowLock;<br>ElbowLock-KinTag                                                         |
| HDX reaction details                             | 20mM Hepes, 150mM NaCl, 1mM MgCl <sub>2</sub> ,<br>0.1mM ADP, 0.5mM TCEP, pD <sub>read</sub> = 7.00,<br>23 °C |
| HDX time course (s)                              | 0.3, 0.5, 1, 3, 10, 30, 300                                                                                   |
| Back-exchange (mean / IQR)                       |                                                                                                               |
| # of Peptides                                    | 318                                                                                                           |
| Sequence coverage                                | 88%                                                                                                           |
| Average peptide length / Redundancy              | 14 / 3.55                                                                                                     |
| Replicates                                       | 3 (technical)                                                                                                 |
| Repeatability                                    | 0.08 (average standard deviation)                                                                             |
| Significant differences in HDX (delta HDX > X D) | 0.286 D (99% CI)                                                                                              |

**Table S1. HDX summary table**

## Data availability

Data required to evaluate this manuscript is provided within the preprint

## Acknowledgements

This work was funded by BBSRC grants BB/W005581/1 and BB/Z517276/1. JAC was supported by an EPSRC-funded Doctoral Prize Fellowship. MPD acknowledges support from a Lister Institute of Preventative Medicine Fellowship. CS acknowledges funding by a Wellcome Trust Investigator award (210701/Z/18/Z). JJP and MK acknowledge funding from UKRI Future Leaders Fellowship (MR/T02223X/1). We thank the University of Bristol School of Chemistry Mass Spectrometry Facility for access to the EPSRC-funded Bruker Ultraflex MALDI-TOF instrument (EP/K03927X/1), the BBSRC-funded BrisSynBio centre for access to peptide synthesis and the plate reader (BB/L01386X/1) and the Wolfson Bioimaging Facility for access to this Talos L120C microscope supported by BBSRC equipment grant (BB/X019799/1). We are grateful for the assistance and access to equipment at the GW4 Facility for High-Resolution Electron Cryo-Microscopy, funded by the Wellcome Trust (202904/Z/16/Z and 206181/Z/17/Z). We are grateful to Alex Walker and Bram Mylemans for helpful discussions and Ferdos Abid Ali for comments on the manuscript.

## Additional information

### Author Contributions

SS purified kinesin-1 complexes, performed SEC, and NS-EM experiments and analysis. JAC designed and engineered the  $\Delta$ TDS KIF5C construct, designed, synthesised and characterised peptides for FP binding assays, purified TPR proteins and performed and analysed FP experiments. MK, SS and JJP designed, carried out and analysed HDX-MS experiments. SS, SKNY, CS and MPD analysed cryoEM data. JW designed and engineered the KLC1-KinTag fusion protein and other constructs. UB and JM supported NS and cryo-EM sample preparation and data acquisition. XL performed AUC assays. MPD, DNW and SS wrote the paper with input from all the authors.

### Funding

| Funder                                                               | Grant reference number | Author                           |
|----------------------------------------------------------------------|------------------------|----------------------------------|
| UKRI   Biotechnology and Biological Sciences Research Council (AFRC) | BB/Z517276/1           | Mark P Dodding<br>Dek N Woolfson |
| UKRI   Biotechnology and Biological Sciences Research Council (AFRC) | BB/W005581/1           | Mark P Dodding<br>Dek N Woolfson |

### Author ORCID iDs

**Shivam Shukla:** <https://orcid.org/0000-0003-2856-5081>  
**Jessica A Cross:** <https://orcid.org/0000-0001-7725-3821>  
**Monika Kish:** <https://orcid.org/0000-0002-3661-8641>  
**Sathish KN Yadav:** <https://orcid.org/0000-0003-0779-0941>  
**Johannes F Weijman:** <https://orcid.org/0000-0002-2082-5777>  
**Laura O'Regan:** <https://orcid.org/0000-0002-6101-6300>  
**Judith Mantell:** <https://orcid.org/0000-0002-9061-8933>  
**Ufuk Borucu:** <https://orcid.org/0000-0001-6746-5409>  
**Xiyue Leng:** <https://orcid.org/0000-0002-4441-2352>  
**Christiane Schaffitzel:** <https://orcid.org/0000-0002-1516-9760>  
**Jonathan J Phillips:** <https://orcid.org/0000-0002-5361-9582>  
**Derek N Woolfson:** <https://orcid.org/0000-0002-0394-3202>  
**Mark P Dodding:** <https://orcid.org/0000-0001-8091-6534>

## References

1. Yildiz A. (2024) Mechanism and regulation of kinesin motors. *Nat Rev Mol Cell Biol*
2. Cross J. A., Dodding M. P. (2019) Motor-cargo adaptors at the organelle-cytoskeleton interface. *Curr Opin Cell Biol* **59**:16-23
3. Cross R. A. (2016) Review: Mechanochemistry of the kinesin-1 ATPase. *Biopolymers* **105**:476-482
4. Reck-Peterson S. L., Redwine W. B., Vale R. D., Carter A. P. (2018) The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol* **19**:382-398
5. Vale R. D., Reese T. S., Sheetz M. P. (1985) Identification of a novel force-generating protein, kinesin, involved in microtubule-based motility. *Cell* **42**:39-50
6. Hackney D. D., Levitt J. D., Suhan J. (1992) Kinesin undergoes a 9 S to 6 S conformational transition. *J Biol Chem* **267**:8696-8701
7. Hisanaga S., et al. (1989) The molecular structure of adrenal medulla kinesin. *Cell Motil Cytoskeleton* **12**:264-272
8. Hirokawa N., et al. (1989) Submolecular domains of bovine brain kinesin identified by electron microscopy and monoclonal antibody decoration. *Cell* **56**:867-878
9. Dodding M. P., Way M. (2011) Coupling viruses to dynein and kinesin-1. *Embo j* **30**:3527-3539
10. Sleigh J. N., Rossor A. M., Fellows A. D., Tosolini A. P., Schiavo G. (2019) Axonal transport and neurological disease. *Nat Rev Neurol* **15**:691-703
11. Xia C., Rahman A., Yang Z., Goldstein L. S. (1998) Chromosomal localization reveals three kinesin heavy chain genes in mouse. *Genomics* **52**:209-213
12. Rahman A., Friedman D. S., Goldstein L. S. (1998) Two kinesin light chain genes in mice. Identification and characterization of the encoded proteins. *J Biol Chem* **273**:15395-15403
13. Gauger A. K., Goldstein L. S. (1993) The Drosophila kinesin light chain. Primary structure and interaction with kinesin heavy chain. *J Biol Chem* **268**:13657-13666
14. Fan J., Amos L. A. (1994) Kinesin Light Chain Isoforms in Caenorhabditis elegans. *Journal of Molecular Biology* **240**:507-512
15. Hackney D. D. (2007) Jump-starting kinesin. *J Cell Biol* **176**:7-9
16. Kaan H. Y., Hackney D. D., Kozielski F. (2011) The structure of the kinesin-1 motor-tail complex reveals the mechanism of autoinhibition. *Science* **333**:883-885
17. Dietrich K. A., et al. (2008) The kinesin-1 motor protein is regulated by a direct interaction of its head and tail. *Proc Natl Acad Sci U S A* **105**:8938-8943
18. Hackney D. D., Baek N., Snyder A. C. (2009) Half-site inhibition of dimeric kinesin head domains by monomeric tail domains. *Biochemistry* **48**:3448-3456
19. Friedman D. S., Vale R. D. (1999) Single-molecule analysis of kinesin motility reveals regulation by the cargo-binding tail domain. *Nat Cell Biol* **1**:293-297
20. Coy D. L., Hancock W. O., Wagenbach M., Howard J. (1999) Kinesin's tail domain is an inhibitory regulator of the motor domain. *Nature Cell Biology* **1**:288-292
21. Atherton J., et al. (2024) Microtubule association induces a Mg-free apo-like ADP pre-release conformation in kinesin-1 that is unaffected by its autoinhibitory tail. *bioRxiv*
22. Tan Z., et al. (2023) Autoinhibited kinesin-1 adopts a hierarchical folding pattern. *eLife* **12**:RP86776 <https://doi.org/10.7554/eLife.86776>
23. Yang J. T., Laymon R. A., Goldstein L. S. (1989) A three-domain structure of kinesin heavy chain revealed by DNA sequence and microtubule binding analyses. *Cell* **56**:879-889
24. Yip Y. Y., et al. (2016) The light chains of kinesin-1 are autoinhibited. *Proc Natl Acad Sci U S A* **113**:2418-2423

25. Verhey K. J., et al. (1998) Light chain-dependent regulation of Kinesin's interaction with microtubules. *J Cell Biol* **143**:1053-1066
26. Verhey K. J., et al. (2001) Cargo of kinesin identified as JIP scaffolding proteins and associated signaling molecules. *J Cell Biol* **152**:959-970
27. Antón Z., et al. (2021) Molecular mechanism for kinesin-1 direct membrane recognition. *Science Advances*
28. Cyr J. L., Pfister K. K., Bloom G. S., Slaughter C. A., Brady S. T. (1991) Molecular genetics of kinesin light chains: generation of isoforms by alternative splicing. *Proc Natl Acad Sci U S A* **88**:10114-10118
29. Woźniak M. J., Allan V. J. (2006) Cargo selection by specific kinesin light chain 1 isoforms. *The EMBO Journal* **25**:5457-5468-5468
30. Cai D., Hoppe A. D., Swanson J. A., Verhey K. J. (2007) Kinesin-1 structural organization and conformational changes revealed by FRET stoichiometry in live cells. *J Cell Biol* **176**:51-63
31. Stock M. F., et al. (1999) Formation of the Compact Conformer of Kinesin Requires a COOH-terminal Heavy Chain Domain and Inhibits Microtubule-stimulated ATPase Activity\*. *Journal of Biological Chemistry* **274**:14617-14623
32. Weijman J. F., et al. (2022) Molecular architecture of the autoinhibited kinesin-1 lambda particle. *Science Advances* **8**:eabp9660
33. Carrington G., et al. (2024) A multiscale approach reveals the molecular architecture of the autoinhibited kinesin KIF5A. *J Biol Chem* **300**:105713
34. Cross J. A., et al. (2024) A de novo designed coiled coil-based switch regulates the microtubule motor kinesin-1. *Nat Chem Biol* **20**:916-923
35. Pernigo S., Lamprecht A., Steiner R. A., Dodding M. P. (2013) Structural basis for kinesin-1: cargo recognition. *Science* **340**:356-359
36. Cross J. A., Chegkazi M. S., Steiner R. A., Woolfson D. N., Dodding M. P. (2021) Fragment-linking peptide design yields a high-affinity ligand for microtubule-based transport. *Cell Chem Biol*
37. Randall T. S., et al. (2017) A small-molecule activator of kinesin-1 drives remodeling of the microtubule network. *Proc Natl Acad Sci U S A* **114**:13738-13743
38. Rosa-Ferreira C., Munro S. (2011) Arl8 and SKIP act together to link lysosomes to kinesin-1. *Dev Cell* **21**:1171-1178
39. Dodding M. P., Mitter R., Humphries A. C., Way M. (2011) A kinesin-1 binding motif in vaccinia virus that is widespread throughout the human genome. *EMBO J* **30**:4523-4538
40. Chiba K., Ori-McKenney K. M., Niwa S., McKenney R. J. (2022) Synergistic autoinhibition and activation mechanisms control kinesin-1 motor activity. *Cell Rep* **39**:111016
41. Wilson M. H., Holzbaur E. L. (2015) Nesprins anchor kinesin-1 motors to the nucleus to drive nuclear distribution in muscle cells. *Development* **142**:218-228
42. Kawano T., et al. (2012) A small peptide sequence is sufficient for initiating kinesin-1 activation through part of TPR region of KLC1. *Traffic* **13**:834-848
43. Araki Y., et al. (2007) The novel cargo Alcadein induces vesicle association of kinesin-1 motor components and activates axonal transport. *Embo j* **26**:1475-1486
44. Konecna A., et al. (2006) Calsyntenin-1 Docks Vesicular Cargo to Kinesin-1. *Mol Biol Cell* **17**:3651-3663
45. Pernigo S., et al. (2018) Structural basis for isoform-specific kinesin-1 recognition of Y-acidic cargo adaptors. *eLife* **7**
46. Nguyen T. Q., et al. (2018) Characterization of the binding mode of JNK-interacting protein 1 (JIP1) to kinesin-light chain 1 (KLC1). *J Biol Chem* **293**:13946-13960
47. Sanger A., et al. (2017) SKIP controls lysosome positioning using a composite kinesin-1 heavy and light chain-binding domain. *J Cell Sci* **130**:1637-1651

48. Pu J., et al. (2015) BORC, a multisubunit complex that regulates lysosome positioning. *Dev Cell* **33**:176-188
49. Farias G. G., Guardia C. M., Britt D. J., Guo X., Bonifacio J. S. (2015) Sorting of Dendritic and Axonal Vesicles at the Pre-axonal Exclusion Zone. *Cell Rep* **13**:1221-1232
50. Cockburn J. J. B., et al. (2018) Insights into Kinesin-1 Activation from the Crystal Structure of KLC2 Bound to JIP3. *Structure* **26**:1486-1498.e1486
51. Fletcher J. M., et al. (2012) A basis set of de novo coiled-coil peptide oligomers for rational protein design and synthetic biology. *ACS Synth Biol* **1**:240-250
52. Zhu H., et al. (2012) Crystal Structures of the Tetratricopeptide Repeat Domains of Kinesin Light Chains: Insight into Cargo Recognition Mechanisms. *PLOS One* **7**:e33943
53. Kish M., et al. (2023) Allosteric Regulation of Glycogen Phosphorylase by Order/Disorder Transition of the 250' and 280s Loops. *Biochemistry* **62**:1360-1368
54. Guo Z., et al. (2022) Engineering and exploiting synthetic allostery of NanoLuc luciferase. *Nature Communications* **13**:789
55. Bai Y., Milne J. S., Mayne L., Englander S. W. (1993) Primary structure effects on peptide group hydrogen exchange. *Proteins: Structure, Function, and Bioinformatics* **17**:75-86
56. Garcia Natalie K., Guttman M., Ebner Jamie L., Lee Kelly K. (2015) Dynamic Changes during Acid-Induced Activation of Influenza Hemagglutinin. *Structure* **23**:665-676
57. Eggertson M. J., Fadgen K., Engen J. R., Wales T. E., Matthiesen R. (2020) *Mass Spectrometry Data Analysis in Proteomics* New York, NY: Springer New York. pp. 407-435
58. Kozielski F., et al. (1997) The Crystal Structure of Dimeric Kinesin and Implications for Microtubule-Dependent Motility. *Cell* **91**:985-994
59. Woolfson D. N. (2023) Understanding a protein fold: The physics, chemistry, and biology of  $\alpha$ -helical coiled coils. *J Biol Chem* **299**:104579
60. Barlan K., Lu W., Gelfand V. I. (2013) The microtubule-binding protein ensconsin is an essential cofactor of kinesin-1. *Curr Biol* **23**:317-322
61. Ferro L. S., et al. (2022) Structural and functional insight into regulation of kinesin-1 by microtubule-associated protein MAP7. *Science* **375**:326-331
62. Hooikaas P. J., et al. (2019) MAP7 family proteins regulate kinesin-1 recruitment and activation. *J Cell Biol* **218**:1298-1318
63. Chaudhary A. R., et al. (2019) MAP7 regulates organelle transport by recruiting kinesin-1 to microtubules. *J Biol Chem* **294**:10160-10171
64. Metzger T., et al. (2012) MAP and kinesin-dependent nuclear positioning is required for skeletal muscle function. *Nature* **484**:120-124
65. Tymanskyj S. R., Yang B. H., Verhey K. J., Ma L. (2018) MAP7 regulates axon morphogenesis by recruiting kinesin-1 to microtubules and modulating organelle transport. *eLife* **7**:e36374 <https://doi.org/10.7554/eLife.36374>
66. Berisha A.-M., et al. (2025) Spatial activation of Kinesin-1 by Ensconsin shapes microtubule networks via ncMTOCs recruitment. *bioRxiv*
67. Monroy B. Y., et al. (2018) Competition between microtubule-associated proteins directs motor transport. *Nature Communications* **9**:1487
68. Smith E. R., Turner E. D., Abdelhamid M. A. S., Craggs T. D., Twelvetrees A. E. (2024) Kinesin-1 is highly flexible and adopts an open conformation in the absence of cargo. *bioRxiv*
69. Nguyen T. Q., et al. (2017) Structural plasticity of the N-terminal capping helix of the TPR domain of kinesin light chain. *PLoS One* **12**:e0186354
70. Sahabandu N., et al. (2025) Active microtubule-actin cross-talk mediated by a nesprin-2G-kinesin complex. *Sci Adv* **11**:eadq4726

71. Blasius T. L., Cai D., Jih G. T., Toret C. P., Verhey K. J. (2007) Two binding partners cooperate to activate the molecular motor Kinesin-1. *J Cell Biol* **176**:11-17
72. Abramson J., et al. (2024) Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**:493-500
73. Pettersen E. F., et al. (2004) UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**:1605-1612
74. Punjani A., Fleet D. (2022) 3D Flexible Refinement: Structure and Motion of Flexible Proteins from Cryo-EM. *Microscopy and Microanalysis* **28**:1218-1218
75. Hageman T. S., Weis D. D. (2019) Reliable Identification of Significant Differences in Differential Hydrogen Exchange-Mass Spectrometry Measurements Using a Hybrid Significance Testing Approach. *Analytical Chemistry* **91**:8008-8016
76. Seetaloo N., Kish M., Phillips J. J. (2022) HDflex: Software for Flexible High Structural Resolution of Hydrogen/Deuterium-Exchange Mass Spectrometry Data. *Analytical Chemistry* **94**:4557-4564
77. Keppel T. R., Weis D. D. (2015) Mapping Residual Structure in Intrinsically Disordered Proteins at Residue Resolution Using Millisecond Hydrogen/Deuterium Exchange and Residue Averaging. *Journal of the American Society for Mass Spectrometry* **26**:547-554

## Peer reviews

### Reviewer #1 (Public review):

The authors aim to interrogate the sets of intramolecular interactions that cause kinesin-1 hetero-tetramer autoinhibition and the mechanism by which cargo interactions via the light chain tetratricopeptide repeat domains can initiate motor activation. The molecular mechanisms of kinesin regulation remain a key question with respect to intracellular transport and this study adds important perspectives to our understanding. It has implications for the accuracy and efficiency of motor transport by different motor families, for example the direction of cargos in one or other direction on microtubules.

The authors focus on the response of inactivated kinesin-1 to peptides found in cargos and the cascade of conformational changes that are induced. They also test the effects of the known activator of kinesin-1 - MAP7 - in the context of their model. The study benefits from multiple complementary, albeit relatively low-resolution, methods - structural prediction using AlphaFold3, 2D and 3D analysis of (mainly negative stain) TEM images of several engineered kinesin constructs, biophysical characterisation of the complexes, peptide design, hydrogen/deuterium-exchange mass spectrometry and simple cell-based imaging. Each set of experiments is carefully designed and the intrinsic limitations of each method are offset by other approaches, such that the assembled data convincingly supports the authors' regulatory model of kinesin activation.

This study benefits from prior work by the authors on this system and the tools and constructs they previously accrued, as well as from other recent contributions to the field. This work will be of broad interest to cell and structural biologists, especially those seeking to tackle small and flexible macromolecular complexes, as well as biophysicists and those interested in protein engineering.

<https://doi.org/10.7554/eLife.109462.2.sa3>

### Reviewer #2 (Public review):

Summary:

In this paper, Shukla, Cross, Kish, and colleagues investigate how binding of a cargo-adaptor mimic (KinTag) to the TPR domains of the kinesin-1 light chain, or disruption of the TPR docking site (TDS) on the kinesin-1 heavy chain, triggers release of the TPR domains from the holoenzyme. This dislocation provides a plausible mechanism for transition out of the autoinhibited lambda-particle toward the open and active conformation of kinesin-1. Using a combination of negative-stain electron microscopy, AlphaFold modeling, biochemical assays, hydrogen-deuterium exchange mass spectrometry (HDX-MS) and other methods, the authors show how TPR undocking propagates conformational changes through the coiled-coil stalk to the motor domains, increasing their mobility, and enhances interactions with the microtubule-bound cofactor MAP7. Together, they propose a model in which the TDS on CC1 of the heavy chain forms a "shoulder" in the compact, autoinhibited state. Cargo-adaptor binding, mimicked here by KinTag, dislodges this shoulder, liberating the motor domains and promoting MAP7 association, driving kinesin-1 activation.

#### Strengths:

Throughout the study, the authors use clever construct design - e.g. delta-Elbow, ElbowLock, CC-Di and the high-affinity KinTag - to test specific mechanisms by directly perturbing structural contacts or effecting interactions. The proposed mechanism of releasing autoinhibition via adaptor-induced TPR undocking is also interrogated with a number of complementary techniques that converge on a convincing model for activation that can be further tested in future studies.

#### Weaknesses:

These reflect limits of what the current data can establish rather than flaws in execution. It remains to be tested if the open state of kinesin-1 initiated by TPR undocking is indeed an active state of kinesin-1 capable of processive movement and/or cargo transport. It also remains to be determined what the mechanism of motor domain undocking from the autoinhibited conformation is. But this important study provides the groundwork for testing these open questions.

#### Comments on revisions:

My original minor concerns have been addressed in the revision.

<https://doi.org/10.7554/eLife.109462.2.sa2>

### Reviewer #3 (Public review):

#### Summary:

The manuscript by Shukla and colleagues presents a comprehensive study that addresses a central question in kinesin-1 regulation-how cargo binding to the kinesin light chain (KLC) tetratricopeptide repeat (TPR) domains triggers activation of full-length kinesin-1 (KHC). The authors combine AlphaFold3 modeling, biophysical analysis (fluorescence polarization, hydrogen-deuterium exchange), and electron microscopy to derive a mechanistic model in which the KLC-TPR domains dock onto coiled-coil 1 (CC1) of the KHC to form the "TPR shoulder," stabilizing the autoinhibited ( $\lambda$ -particle) conformation. Binding of a W/Y-acidic cargo motif (KinTag) or deletion of the CC1 docking site (TDS) dislocates this shoulder, liberating the motor domains and enhancing accessibility to cofactors such as MAP7. The results link cargo recognition to allosteric structural transitions and present a unified model of kinesin-1 activation. I recommend acceptance of the manuscript subject to the following additions:

#### Strengths:

- (1) The study addresses a fundamental and long-standing question in kinesin-1 regulation using a multidisciplinary approach that combines structural modeling, quantitative biophysics, and electron microscopy.
- (2) The mechanistic model linking cargo-induced dislocation of the TPR shoulder to activation of the motor complex is well supported by both structural and biochemical evidence.
- (3) The authors employ elegant protein-engineering strategies (e.g., ElbowLock and  $\Delta$ TDS constructs) that enable direct testing of model predictions, providing clear mechanistic insight rather than purely correlative data.
- (4) The data are internally consistent and align well with previous studies on kinesin-1 regulation and MAP7-mediated activation, strengthening the overall conclusion.

#### Weaknesses:

- (1) While the EM and HDX-MS analyses are informative, the conformational heterogeneity of the complex limits structural resolution, making some aspects of the model (e.g., stoichiometry or symmetry of TPR docking) indirect rather than directly visualized.
- (2) The dynamics of KLC-TPR docking and undocking remain incompletely defined; it is unclear whether both TPR domains engage CC1 simultaneously or in an alternating fashion.
- (3) The interplay between cargo adaptors and MAP7 is discussed but not experimentally explored, leaving open questions about the sequence and exclusivity of their interactions with CC1.

#### Comments on revisions:

The authors have addressed my comments satisfactorily.

<https://doi.org/10.7554/eLife.109462.2.sa1>

### Author response:

The following is the authors' response to the original reviews

#### **eLife Assessment**

*The manuscript by Shukla et al. provides important mechanistic insights into kinesin-1 autoinhibition and cargo-mediated activation. Using a convincing combination of protein engineering, computational modeling, biophysical assays, HDX-MS, and electron microscopy, the authors reveal how cargo binding induces an allosteric transition that propagates to the motor domains and enhances MAP7 binding. Despite limitations arising from conformational heterogeneity and structural resolution, the study presents a unified mechanism for kinesin-1 activation that will be of broad interest to the motor protein, structural biology, and cell biology communities.*

We are grateful for the time and effort from the reviewers and editors in providing fair and constructive comments that have helped to improve the manuscript. Our point-by-point response is provided below.

#### **Public Reviews:**

##### **Reviewer #1 (Public review):**

*Summary:*

*The authors aim to interrogate the sets of intramolecular interactions that cause kinesin-1 hetero-tetramer autoinhibition and the mechanism by which cargo interactions via the light chain tetratricopeptide repeat domains can initiate motor activation. The molecular mechanisms of kinesin regulation remain an important question with respect to intracellular transport. It has implications for the accuracy and efficiency of motor transport by different motor families, for example, the direction of cargos towards one or other microtubules.*

**Strengths:**

*The authors focus on the response of inactivated kinesin-1 to peptides found in cargos and the cascade of conformational changes that occur. They also test the effects of the known activator of kinesin-1 - MAP7 - in the context of their model. The study benefits from multiple complementary methods - structural prediction using AlphaFold3, 2D and 3D analysis of (mainly negative stain) TEM images of several engineered kinesin constructs, biophysical characterisation of the complexes, peptide design, hydrogen/deuterium-exchange mass spectrometry, and simple cell-based imaging. Each set of experiments is thoughtfully designed, and the intrinsic limitations of each method are offset by other approaches such that the assembled data convincingly support the authors' conclusions. This study benefits from prior work by the authors on this system and the tools and constructs they previously accrued, as well as from other recent contributions to the field.*

**Weaknesses:**

*It is not always straightforward to follow the design logic of a particular set of experiments, with the result that the internal consistency of the data appears unconvincing in places.*

*For example, i) the Figure 1 AlphaFold3 models do not include motor domains whereas the nearly all of the rest of the data involve constructs with the motor domains;*

We appreciate the reviewer's comment regarding the absence of the motor domains in the AlphaFold3 models shown in Figure 1. These domains were intentionally excluded to improve visual clarity and to better highlight the interaction between the TPR domains and CC1 in the inhibited kinesin-1 conformation. We felt that this simplified presentation in the main figure helps readers focus on the key mechanistic advance introduced in this work at the outset of the paper. For completeness, we have provided full-length kinesin-1 AlphaFold3 models that include the motor domains in the Supplementary Information (Fig. S1), and they are described in detail in the main text. In addition, we have added a note to the Figure 1 legend to explicitly direct readers to these full-length models.

*ii) the kinesin constructs are chemically cross-linked prior to TEM sample preparation - this is clear in the Methods but should be included in the Results text, together with some discussion of how this might influence consistency with other methods where crosslinking was not used.*

Thank you. Chemical crosslinking is typically important for obtaining high-quality negative-stain TEM grids of kinesin-1 complexes and has been employed in all prior EM studies by our group and others. While this was described in the Methods, we agree that it should also be stated explicitly in the Results. Accordingly, we have added a sentence to the Results section noting that the proteins were stabilized using the amine-to-amine crosslinker BS3 ("Proteins were also stabilised using the amine-to-amine crosslinker BS3 that was important for achieving reproducibly high-quality samples for imaging.").

Please see point below for acknowledgement of risks of using crosslinker.

*Can those cross-links themselves be used to probe the intramolecular interactions in the molecular populations by mass spec?*

We had considered this, however, cross-linking mass spectrometry (XL-MS) has been applied extensively to essentially identical kinesin-1 complexes by Tan et al. (eLife 2023). That work provided important insights into the overall architecture of the complex, including the new head-CC1 interactions. However, as fully acknowledged by the authors, significant ambiguity remained with respect to the positioning of the TPR domains, with many cross-links that could not be straightforwardly rationalized in a single model. These unresolved aspects provided part of the motivation for the present study, as highlighted in the Introduction.

We believe that this ambiguity likely reflects an underlying conformational equilibrium of the kinesin-1 complex (e.g. opening/closing transitions) and/or dynamic docking and undocking of the TPR domains, and lysine-rich features of the TPR domains (most notably the loops that connect the TPR alpha helices) which may make them prone to lock in non-native states, which limits the interpretability of static cross-linking data in this system. In this context therefore, we feel that XL-MS has already been thoroughly explored for kinesin-1 and that its practical limitations in resolving these TPR interactions have been reached.

This consideration was a primary motivation for pursuing cross-linker-free, solution-based approaches, particularly HDX-MS, which we argue provide the most relevant new insights into the assembly and conformational dynamics of the complex. To make this rationale clearer, we have added an explicit note in the HDX-MS section emphasizing that this is a cross-linker-free method. The added text reads:

“To determine how the local structural changes from adaptor binding and shoulder dislocation affected the dynamics of kinesin-1 complexes in solution, as directly and least invasively as possible, and without the risk of cross-linker artefacts.”

*In general, the information content of some of the figure panels can also be improved with more annotations (e.g. angular relationship between views in Figure 1B, approximate interpretations of the various blobs in Fig 3F, and more thought given to what the reader should extract from the representative micrographs in several figures - inclusion of the raw data is welcome but extraction and magnification of exemplar particles (as is done more effectively in Fig S5) could convey more useful information elsewhere.*

We appreciate these suggestions. We have modified the figures throughout the manuscript in line with the reviewer's points. Raw data is now provided at higher magnification throughout so the reader can better distinguish individual particles, angular relationships have been added and further annotations provided on 2D class averages. We do not want the reader to draw too many conclusions from images of single closed particles (with the exception of open vs closed in Fig S7) as these require averaging and 2D classification to obtain meaningful insights, and so we have not added zoom panels in these cases. Figure 3F has been annotated as requested.

#### **Reviewer #2 (Public review):**

##### *Summary:*

*In this paper, Shukla, Cross, Kish, and colleagues investigate how binding of a cargo-adaptor mimic (KinTag) to the TPR domains of the kinesin-1 light chain, or disruption of the TPR docking site (TDS) on the kinesin-1 heavy chain, triggers release of the TPR domains from the holoenzyme. This dislocation provides a plausible mechanism for transition out of the autoinhibited lambda-particle toward the open and active conformation of kinesin-1. Using a combination of negative-stain electron microscopy,*

*AlphaFold modeling, biochemical assays, hydrogen-deuterium exchange mass spectrometry (HDX-MS), and other methods, the authors show how TPR undocking propagates conformational changes through the coiled-coil stalk to the motor domains, increasing their mobility and enhancing interactions with the microtubule-bound cofactor MAP7. Together, they propose a model in which the TDS on CC1 of the heavy chain forms a "shoulder" in the compact, autoinhibited state. Cargo-adaptor binding, mimicked here by KinTag, dislodges this shoulder, liberating the motor domains and promoting MAP7 association, driving kinesin-1 activation.*

**Strengths:**

*Throughout the study, the authors use a clever construct design - e.g., delta-Elbow, ElbowLock, CC-Di, and the high-affinity KinTag - to test specific mechanisms by directly perturbing structural contacts or affecting interactions. The proposed mechanism of releasing autoinhibition via adaptor-induced TPR undocking is also interrogated with a number of complementary techniques that converge on a convincing model for activation that can be further tested in future studies. The paper is well-written and easy to follow, though some more attention to figure labels and legends would improve the manuscript (detailed in recommendations for the authors).*

**Weaknesses:**

*These reflect limits of what the current data can establish rather than flaws in execution. It remains to be tested if the open state of kinesin-1 initiated by TPR undocking is indeed an active state of kinesin-1 capable of processive movement and/or cargo transport. It also remains to be determined what the mechanism of motor domain undocking from the autoinhibited conformation is, and perhaps this could have been explored more here. The authors have shown by HDX-MS that the motor domains become more mobile on KinTag binding, but perhaps molecular dynamics would also be useful for modelling how that might occur.*

We are grateful for the reviewer's comments. We agree that the weaknesses the reviewer has outlined define the limitations of the study and establish important priorities for future work, that includes molecular dynamics simulations. An important prerequisite for the latter is a starting model that one has confidence in. We think that our study and earlier work now provide a good experimentally supported foundation for using AF3 generated assemblies for this purpose, by ourselves and others.

**Reviewer #3 (Public review):**

**Summary:**

*The manuscript by Shukla and colleagues presents a comprehensive study that addresses a central question in kinesin-1 regulation - how cargo binding to the kinesin light chain (KLC) tetratricopeptide repeat (TPR) domains triggers activation of full-length kinesin-1 (KHC). The authors combine AlphaFold3 modeling, biophysical analysis (fluorescence polarization, hydrogen-deuterium exchange), and electron microscopy to derive a mechanistic model in which the KLC-TPR domains dock onto coiled-coil 1 (CC1) of the KHC to form the "TPR shoulder," stabilizing the autoinhibited ( $\lambda$ -particle) conformation. Binding of a W/Y-acidic cargo motif (KinTag) or deletion of the CC1 docking site (TDS) dislocates this shoulder, liberating the motor domains and enhancing accessibility to cofactors such as MAP7. The results link cargo recognition to allosteric structural transitions and present a unified model of kinesin-1 activation.*

**Strengths:**

*(1) The study addresses a fundamental and long-standing question in kinesin-1 regulation using a multidisciplinary approach that combines structural modeling, quantitative biophysics, and electron microscopy.*

*(2) The mechanistic model linking cargo-induced dislocation of the TPR shoulder to activation of the motor complex is well supported by both structural and biochemical evidence.*

*(3) The authors employ elegant protein-engineering strategies (e.g., ElbowLock and  $\Delta$ TDS constructs) that enable direct testing of model predictions, providing clear mechanistic insight rather than purely correlative data.*

*(4) The data are internally consistent and align well with previous studies on kinesin-1 regulation and MAP7-mediated activation, strengthening the overall conclusion.*

**Weaknesses:**

*(1) While the EM and HDX-MS analyses are informative, the conformational heterogeneity of the complex limits structural resolution, making some aspects of the model (e.g., stoichiometry or symmetry of TPR docking) indirect rather than directly visualized.*

We agree with the reviewers point. Conformational heterogeneity is a significant challenge, and the model has been developed from multiple complementary approaches. A higher resolution cryoEM study remains a priority, but is challenging because of the size, shape and flexibility of the particle, but we hope that some the approaches used here (e.g. nanobody TPR stabilisation, ElbowLock) will provide a path to achieve this.

*(2) The dynamics of KLC-TPR docking and undocking remain incompletely defined; it is unclear whether both TPR domains engage CC1 simultaneously or in an alternating fashion.*

We agree that this is a limitation. We strongly suspect that the TPR domains dynamic and are working to overcome experimental challenges to resolve this important outstanding question. We have expanded the discussion section to better highlight this important priority.

*(3) The interplay between cargo adaptors and MAP7 is discussed but not experimentally explored, leaving open questions about the sequence and exclusivity of their interactions with CC1.*

We agree that this is a limitation but will be an important priority for future studies.

**Recommendations for the authors:**

**Reviewer #1 (Recommendations for the authors):**

*There are a number of places where the text could be more precise or clear, or the figures could be designed to be more informative:*

*(1) The word "unitarily" is used in several places, and I don't know what it means in this context.*

We have changed the phrasing throughout the manuscript to this term. We were attempting to contrast with presumed cooperative multivalent interactions in the context of the kinesin-1 tetramer but agree that this choice of word doesn't quite achieve that.

*(2) On page 5 the phrase "We focused on the ElbowLock background" is introduced and needs to be explained more clearly.*

Thank you. We have amended the text to read “This KIF5C construct contains a short 5 amino acid deletion that restricts flexibility around the elbow and helps maintain particles in their lambda conformation, providing homogenous samples, and facilitating subsequent analysis (34).”

(3) On page 6, the phrase “To improve the resolution of our images, we turned to single-particle cryoEM analysis” is imprecise - what do the authors mean by the resolution of the images? Cryo-EM data does not always guarantee a higher resolution structure, but it offers the possibility of visualising finer structural features. This is probably what is meant here, but needs to be stated more precisely.

We have amended the text to ‘visualise finer structural details’ as suggested.

(4) Page 7 - “suggesting that TPR domains had loosely dissociated from the core” - I don't think the evidence points to dissociation of KLCs from the complex, but the phrase “loosely dissociated” implies this - would benefit from rephrasing.

We have changed this to ‘undocked’ for consistency with other descriptions in the manuscript.

(5) Was the effect of the CC-Di insertion ( $\Delta$ TDS) detectable by AlphaFold prediction? It would be interesting to include this, partly for completeness and partly because a slightly imperfect and maybe a more dynamic coiled-coil in this region of the molecule may be important in supporting the conformational changes required for activation.

Thank you for this suggestion. Modelling of  $\Delta$ TDS complex indeed shows displacement of the TPR domains. In the standard 5 output models, the TPR domains now occupy a variety of different positions, all with essentially zero confidence (high position error). Consistent with biochemical data, the CCDi insertion is modelled with no overall disruption to the architecture or length of CC1 as expected. We think that this is a valuable addition to the study and have included it as a new supplementary figure (Fig S5), with main text reading.

.... “Supporting this, models of  $\Delta$ TDS complexes using AF3 showed the expected seamless insertion of CCDi into CC1, with displacement of the TPR domains to a variety of different positions, in 5 models, all with high position error with respect to KHC (Fig S5).”

(6) Figure S1 has two sections designated (C) in the legend.

Corrected

(7) Figure S3 - given the resolution and level of interpretation of the 3D reconstructions, it is not relevant to include an FSC curve, but other standard information, such as angular distribution and any evidence of variability from 3D classifications (and how many particles per 3D class) should be included for all structures.

Thank you, a complete workflow for all complexes has now been provided in Figure S8 with the information requested. In each case there were typically two ‘good’ classes. For ElbowLock, this included one without a prominent shoulder, consistent with 2D classification and quantification. We assume this may reflect a docking/undocking equilibrium. For the  $\Delta$ TDS and KinTag particles, neither class showed the shoulder feature. The main text has been modified to reflect this and reads “For ElbowLock complexes, this resulted in classes with and without a prominent shoulder, in agreement with 2D classification. For ElbowLock- $\Delta$ TDS and ElbowLock-KinTag complexes, no prominent shoulder containing classes were observed.”

**Reviewer #2 (Recommendations for the authors):**

Overall, the figures would benefit from more labels for clarity, some examples and suggestions below:

(1) Figure 1A - Connect motors to the rest of the structure e.g., wiggly lines.

Corrected.

(2) Figure 1B - Add arrows and angles to indicate different views of the model.

Corrected.

(3) Figure 1B - Label TPR1-6 (e.g., inset zoom in).

Corrected.

(4) Figure 2D and 3D - Label the lack of a shoulder in all averages (perhaps with an arrow instead of a circle to not obscure density), include an example average which shows prominent shoulder density.

Corrected. Full sets of classes showing shoulder like features for deltaTDS and KinTag complexes are now shown in Figure S4.

(5) Figure 3D: Label motor domains and elbow as in other figures.

Corrected.

(6) Methods: Include more information on how EM classes were compared to AF projections (e.g., Figure 1D). Was this done visually or computationally? Likewise, more information is needed on how classes were judged to have prominent/weak shoulder density (Figure 2D). In the figure legend, there is a statement that "Full sets of classes are provided in Fig. S4" but this is absent in the supplement.

Thank you. This information has been added to the methods.

"For comparison to the AF3 model, simulated density was generated using the molmap command in ChimeraX (73) filtering to 15 Å, and projections were generated/selected automatically using the Reference Based Auto Selected 2D function in CryoSPARC".

Full sets of classes are now provided in Figure S4.

(7) Figure 1-3 - Raw micrographs are a very useful inclusion but would benefit from being a more zoomed-in view (e.g., Figure S5 scale). Particularly useful for 3C, where the mixture of open and closed would be good to see.

Higher zoom micrographs have been provided throughout.

(8) Figure 5D: Panels too small to see the result, suggest making full width and moving E below.

Thank you. We have expanded the panel and moved the model to a new Figure 6.

(9) Figure S1: PAE plot convincing, but pLDDT colour models needed.

A representative model coloured for pLDDT has been added to Figure S1. Most of the structure sits within the light blue confident range ( $90 > \text{pLDDT} > 70$ ) with the exception of the disordered regions and neck coil.

(10) Figure 5B: Reason for the variable inputs?

The reviewer raises an interesting point. The slightly reduced expression of deltaElbow and slightly increased expression of ElbowLock is a consistent feature of these experiments. We note that this effect is in the ‘opposite direction’ to the impact on binding to MAP7 and so does not affect our conclusions from the experiment. However, we wonder whether opening and closing of the complex may impact on turnover of kinesin proteins, which could have implications for their normal homeostasis and possible degradation after transport in polarised cells. We are considering how to explore this going forwards. We have added a note to the results section to highlight this interesting observation to the reader.

“We also noted slightly elevated expression of ElbowLock complexes and slightly lower expression of DeltaElbow complexes, suggesting that opening/closing of the complex could impact on kinesin-1 turnover”

(11) *Figure legend 5B: Insufficient detail, the end result is stated, but the three separate gels are not described.*

Legend has been expanded.

(12) *Figure 3F: Currently somewhat problematic. It is unclear if the models are in the same view, and so comparison is difficult. Figure 1C (bottom right) shows class averages with a clear, separate CC density, so the relatively featureless model in this region is puzzling. A statement on how the three model views are related to each other, if aligned with each other, would be useful.*

We appreciate the reviewers point. Models were aligned in Chimera, using the fit in map command. Because of the limited features of the models presumably due to flexibility, achieving a good alignment for all three models was challenging, but we think that showing the 180-degree rotations is probably about the best we can achieve here.

(13) *The following statement is too strong: "Nonetheless, we obtained reference-free 2D class averages that appeared to show full-length 'side' views of the complex with clear definition of the elbow, hinge 2, and KHC-KLC (coiled-coil) interface features which enabled us to identify CC1 confidently (Fig. 1D)". Given that the negative-stain EM data were collected primarily to validate the AlphaFold model, the assignment of CC1 should be described as consistent with rather than confidently identified from the class averages. The resolution of the EM data does not independently support such an assignment, and the wording needs to be softened.*

We appreciate the reviewer’s point, we have softened the wording as suggested. The paragraph now reads.

“To visualise finer structural details, we turned to single-particle cryoEM analysis of frozen-hydrated samples. We were unable to obtain optimal samples suitable for determining the complete structure. Nonetheless, we obtained reference-free 2D class averages that appeared to show full-length ‘side’ views of the complex with clear definition of the elbow, hinge 2, and KHC-KLC (coiled-coil) interface features (Fig. 1D). The motor domains were poorly resolved in these classes, suggesting that the head assembly is somewhat flexible relative to the coiled coil/TPR body. A comparison to low-pass filtered back-projections from the AF3 model (without motor domains) revealed density at a position concurrent with the docked TPR domains (Fig. 1D).”

(14) *There is a typo in the figure legend of Figure 3 - (E) and (F) should be (F) and (G).*

Corrected

**Reviewer #3 (Recommendations for the authors):**

*I recommend the following additions:*

*(1) Figure 1 labeling - In panel A, please label the "linker domain" and the "KLC subunits" explicitly to help orient the reader. In panel B, please mark the "TPR shoulder" corresponding to the docked TPR domains on CC1; this will help the reader connect parts B and C.*

Thank you, we have modified Figure 1A with this additional information.

*(2) The TPR docking site (TDS) is a central structural element, and its sequence boundaries are provided in the Methods. It would help to visualize this directly in Figure 2A or in an inset.*

We hope that the reviewer agrees that the zoomed in model in Figure 5A (alongside MAP7) provides a sufficiently detailed view of the structural interface to highlight the orientation of TPR1 with respect to CC1. The side chain contacts in the model are very plausible and confidently predicted (and can be straightforwardly reproduced in AF3 using the sequence information provided in the methods), but as our study has not explored this interaction at the single residue level, we would prefer not to imply this to the reader at this stage.

*(3) The authors' model of cargo-induced TPR dislocation is convincing. However, the Discussion could benefit from a clarification on whether both KLC-TPR domains are expected to be bound simultaneously or if a dynamic exchange occurs, as the EM data suggest potential asymmetry.*

Thank you, please see point 5 below where we have modified the discussion to reflect the reviewer's thoughtful comments.

*(4) The HDX-MS analysis is comprehensive, but the authors may want to briefly comment on the coverage of low-signal regions (especially within CC2-CC3) to enhance clarity.*

We have added an additional supplementary figure (S10) showing sequence coverage. Overall, this is 88% but with some lower coverage around KHC-CC0 (neck) and the acidic linker that connects the KLC coiled-coil to the TPR. We have added a note to the main text to reflect this.

"Sequence coverage was high (overall 88%) with the exception of KHC-CC0 (neck coil) and the acidic-linker region that connects the KLC coiled-coil to the TPR domains where coverage was lower"

*(5) In the Discussion, the proposed interplay between MAP7 and cargo adaptors is intriguing, especially considering the results from Anna Akhmanova's lab showing that MAP7 activates kinesin-1 processivity. Do the authors suggest that competition for CC1 is mutually exclusive or sequential? The answer has mechanistic implications.*

We have been considering questions for some time, and the short answer is that we don't fully understand the dynamics yet. However, we appreciate the reviewer's prompt to clarify our thinking on this. We have attempted to do this in a revised discussion section where we more explicitly outline these outstanding questions.

<https://doi.org/10.7554/eLife.109462.2.sa0>