

Molecular and Mechanical Signatures Contributing to Epidermal Differentiation and Barrier Formation

Reviewed Preprint

v1 • October 28, 2024

Not revised

Alexandra Prado-Mantilla, Wenxiu Ning, Terry Lechler 

Department of Dermatology, Duke University Medical Center, Durham, United States • Department of Cell Biology, Duke University Medical Center, Durham, United States

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

This **important** paper by Lechler and colleagues describes the transcriptomic signature and fate of intermediate cells (ICs), a transient and poorly defined embryonic cell type in the skin. The paper **convincingly** shows through lineage tracing that ICs are granular and not spinous cell precursors, and through ectopic expression in vivo, that cell contractility, a mechanical feature of ICs, lies upstream of differentiation.

<https://doi.org/10.7554/eLife.100961.1.sa4>

Abstract

Formation of the skin barrier requires rapid proliferation coupled with differentiation and stratification of the embryonic epidermis. Basal progenitors give rise to progeny throughout development – first to intermediate cells, a transient proliferative suprabasal cell population, and later to spinous cells. Neither the function nor the differentiation trajectory of intermediate cells has been documented. We generated transcriptomes of intermediate and spinous cells and identified specific markers that distinguish these two populations. Further, we found that intermediate cells express a subset of genes in common with granular cells of the epidermis – the terminal living cell type that helps establish the barrier. Lineage tracing revealed that most intermediate cells directly transition to granular cells without expressing markers specific to spinous cells, thus revealing a distinct lineage pathway leading to granular fate. In addition to their transcriptional similarities, intermediate and granular cells both had hallmarks of increased actomyosin contractility. We found that rather than simply lying downstream of cell fate pathways, contractility was sufficient to suppress spinous fate and promote granular gene expression. Together, these data establish the molecular and mechanical characteristics of the developing epidermis that allow this tissue to rapidly develop barrier activity.

Introduction

Development of many tissues is characterized by rapid proliferation coupled with morphogenesis and differentiation. This process often uses specialized and transient cell types that are not found in homeostatic tissue, necessitating distinct pathways for differentiation (Singh and Tiwari, 2023). During the embryonic development of the epidermis, an initial single layer of progenitor cells gives rise to a multi-layered and differentiated tissue that acts as a chemical and mechanical barrier at birth (Moreci and Lechler, 2020; Sumigray and Lechler, 2015). This process begins at embryonic day (E) 14.5 in the mouse back skin when basal progenitor cells start to give rise to a suprabasal cell layer of intermediate cells (Damen et al., 2021; Lechler and Fuchs, 2005; Smart, 1970). Subsequently, basal cells divide to generate spinous cells, which are post-mitotic. These cells then further differentiate into granular cells, the last living cell types in the epidermis which are integral to forming the epidermal barrier. Intermediate cells are a transient cell type that expresses differentiation markers like Keratin 1 and 10 but remain mitotically active (Weiss and Zelickson, 1975). At this time point, the proliferation rate of intermediate cells is comparable to basal cells, suggesting that they are a significant contributor to tissue expansion (Damen et al., 2021; Smart, 1970). These cells lack basement membrane attachment, and thus, they proliferate in a tissue environment that is normally quiescent in the adult. Currently, we do not understand if they play functional roles outside of proliferation, their cell fate trajectory, or how they compare to later generated spinous cells at a molecular level. Further, we lack specific markers for intermediate cells despite recent scRNA-Seq data that has characterized the transcriptomes of these cells (Jacob et al., 2023). Generation of intermediate cells is transient (E14-15.5), and by E16.5 basal cells give rise to post-mitotic spinous cells (**Fig 1A**). It is not known whether this switch in progeny type is driven by intrinsic changes in basal progenitors, changes in signals from the surrounding periderm, and/or by systemic signals (circulating factors or amniotic fluid (Huebner et al., 2012)).

In addition to chemical cues, mechanical information is assessed by cells and can instruct proliferation and differentiation decisions (Vining and Mooney, 2017). This has been most clearly demonstrated by substrates of differential stiffness eliciting altered differentiation pathways in mesenchymal stem cells (Engler et al., 2006). Whether intra-tissue mechanical information is also instructive in differentiation has not been as thoroughly addressed. That said, in many tissues, differentiation leads to changes in contractility and or stiffness. For example, in the epidermis, there is increased contractility in the granular cell layer, and this contractility is important for the formation of tight junctions in this cell layer (Miroshnikova et al., 2018; Rubsam et al., 2017; Sumigray et al., 2012). While this flow of information from transcriptome to contractility has largely been assumed to be unidirectional, some works have demonstrated bidirectional interactions (Le et al., 2016; Meyer-ter-Vehn et al., 2006). When physical linkages between the cytoskeleton and nucleus were removed, there was a premature differentiation of keratinocytes and this was hypothesized to involve alterations in the transcription of the epidermal differentiation complex (EDC), a genetic locus that undergoes a change in localization in response to differentiation (Carley et al., 2021). Further, increased contractility in differentiated cells of the epidermis results in a non-cell autonomous effect on proliferation of their progenitors (Ning et al., 2021). However, we lack clear evidence that contractility can directly affect differentiation in the epidermis.

Here, we used bulk sequencing to characterize the transcriptomes of intermediate and spinous cells, as well as their progenitors. We defined markers that are specific for intermediate and spinous cells and showed marked transcriptomic differences in the basal cell progenitors that give rise to these distinct cell types. Further, our data demonstrate that intermediate cells express many genes associated with granular cells – a cell type that emerges days later and is responsible for secreting lipids that constitute part of the epidermal barrier. Rather than moving through a




Figure 1.

Transcriptomic analysis of embryonic epidermal cell populations.

(A) Diagram depicting the stratification process and different cell populations (basal and K10+ suprabasal layers) existing in the epidermis from E14.5 to E16.5.

(B) Images of EdU staining (white) at E14.5, E15.5 and E16.5. Immunofluorescence of suprabasal cells labeled with K10 in red and the basement membrane, with β4-integrin in green. Scale bars: 20 μm.

(C) Quantification of the percentage of EdU+ cells in the basal and the first two layers of suprabasal cells at E14.5, E15.5 and E16.5. n= 3 embryos/time point. Data are presented as mean ± standard deviation (SD). ns: not significant, *: p < 0.05, **: p < 0.01, ***: p < 0.001, ordinary two-way ANOVA, Tukey's multiple comparisons test.

(D) Schematic showing basal and suprabasal cell populations collected for RNA sequencing: K14-RFP;K10-rtTA;TRE-H2B-GFP pregnant dams were all fed with doxycycline from E12.5, and then sacrificed at either E14.5 or E16.5. The E14.5 intermediate cells (K10-GFP⁺;K14-RFP⁻), E16.5 spinous cells (K10-GFP⁺;K14-RFP⁺), and basal cells at E14.5 and E16.5 (K10-GFP⁺;K14-RFP⁺) cells were collected separately and sent for RNA sequencing. Granular cells were excluded from E16.5 samples.

(E) PCA score plot of the first two principal components (PC1: 33.5% variance and PC2: 21.1% variance) for gene expression levels from samples of cell populations indicated in (D) (n=3 embryos/cell population).

spinous state, as has long been assumed (Koster and Roop, 2005 [↗](#)), intermediate cells appear to directly transition to granular cells. Intermediate cells also show similarity to granular cells in having increased apparent actomyosin contractility. Using genetic models to induce contractility in spinous cells, we demonstrate that contractility is sufficient to induce a granular-like state. Together this work reveals the lineage pathways and the influences of contractility on epidermal barrier formation.

Results

Transcriptomic analysis of embryonic epidermal differentiation

During early epidermal stratification at E14.5, intermediate cells (ICs) are the first layer of suprabasal cells that express the differentiation marker keratin 10 (K10) and proliferate at a rate similar to basal cells (Fig. 1A–C [↗](#)) (Damen et al., 2021 [↗](#); Smart, 1970 [↗](#)). The proliferation rate of suprabasal cells decreases over the next 48 hours, and these cells are largely postmitotic by E16.5. To further validate these suprabasal cell dynamics during epidermal stratification, we performed live imaging of back skin explants from K10-rtTA;TRE-H2B-GFP embryos, which express H2B-GFP in K10-positive suprabasal cells. Consistent with data from fixed embryos, we observed numerous suprabasal mitotic events in explants collected from E14.5 embryos, while suprabasal cells in explants from E16.5 embryos were mitotically inactive (Fig. 1, Supplement 1 [↗](#)). All divisions visualized were planar to the epithelium.

To understand the differences between suprabasal cells at E14.5 and E16.5, we explored the transcriptomic changes occurring between these two time points. We used K14-RFP;K10-rtTA;TRE-H2B-GFP mice and FACS to purify H2B-GFP⁺ suprabasal cells and RFP⁺ basal cells at E14.5 and E16.5 and performed bulk RNA sequencing of each of these cell populations (Fig. 1D [↗](#)). Given that granular cells also form part of the suprabasal compartment at E16.5, we excluded the more superficial cells that bind *Ulex europaeus* agglutinin I lectin from E16.5 samples in order to uniquely compare suprabasal cells that lie immediately above the basal layer (ICs and spinous cells) (Brabec et al., 1980 [↗](#)).

Principal component analysis (PCA) of the sequenced cell populations revealed that biological replicates grouped together and that samples were segregated by cell type (basal vs suprabasal) in the first component, and by age in the second component (E14.5 vs. E16.5). Notably, ICs at E14.5 and spinous cells at E16.5 were not clustered together, indicating that they are distinct populations, as are their corresponding basal cells (Fig. 1E [↗](#)). Though not addressed further here, the developmental transition in suprabasal cells from E14.5 to E16.5 may reflect the very different states of basal cells at these time points.

To identify specific molecular markers for ICs and spinous cells, we filtered the genes that were uniquely enriched in ICs (at E14.5 compared to both E16.5 spinous cells and E14.5 basal cells) and in spinous cells (compared to both E14.5 ICs and E16.5 basal cells). The IC gene signature consisted of 610 genes and included *Scara5*, a scavenger receptor, *St8sia6*, which encodes a sialyltransferase enzyme, and *Tgm1* (transglutaminase 1), an enzyme involved in covalent cross-linking of proteins (Fig. 2A [↗](#)). To validate these candidates, we performed RNAScope and detected mRNA of *Scara5* and *St8sia6* in ICs but not in suprabasal cells at E16.5 (Fig. 2B–E [↗](#)). It is worthwhile to note that *Scara5* expression was specific to ICs, while *St8sia6* mRNA was also present in granular cells at E16.5 (Fig. 2C [↗](#)). This was also reflected in the presence of Loricrin and *St8sia6* protein in the granular layers at E16.5 by antibody staining; however, we did not detect their fluorescence signal in ICs at E14.5 (Fig. 2, Supplement 1 [↗](#)). This suggests that some granular genes are transcriptionally enriched in ICs but not yet appreciably expressed at the protein level. On the contrary, we found *Tgm1* protein expression in intermediate cells, as described in more detail below (Fig. 4C [↗](#)).




Figure 2.

Identifying markers for intermediate and spinous cells.

(A) The core gene signature of intermediate cells was obtained from the list of genes only upregulated in ICs (log2 fold > 1, $p < 0.05$; as compared with both basal cells at E14.5 and spinous cells at E16.5). Total genes in the IC signature were 610. From this list, Scara5 and St8sia6 were two of the most highly upregulated genes compared to spinous cells at E16.5.

(B and C) RNAscope of Scara5 (B) and St8sia6 (C) in WT embryos at E14.5 and E16.5. Basement membrane is indicated as a dotted line. Scale bars: 20 μ m.

(D and E) Quantification of the percentage of Scara5+ (D) and St8sia6+ (E) basal and suprabasal cells at E14.5 and E16.5. $n = 3$ embryos/time point. Data are presented as the mean $\pm$ SD. ****: $p < 0.0001$, ordinary two-way ANOVA, Sidak's multiple comparisons test.

(F) The core gene signature of spinous cells was obtained from the list of genes only upregulated in spinous cells (log2 fold > 1, $p < 0.05$ compared with both basal cells at E16.5 and intermediate cells at E14.5). Total genes in the spinous cell signature were 163.

(G and H) Immunofluorescence staining of MafB (G) and Cox1 (H) in WT embryos at E14.5 and E16.5. Basement membrane is indicated as a dotted line. Scale bars: 20 μ m.

(I) Quantification of the percentage of MafB+ basal and suprabasal cells at E14.5 and E16.5. $n = 3$ embryos/time point. Data are presented as the mean $\pm$ SD. ****: $p < 0.0001$, ordinary two-way ANOVA, Sidak's multiple comparisons test.

(J) Quantification of Cox1 puncta/standardized area in intermediate cells at E14.5 was compared to the first two layers of spinous cells at E16.5. $n = 3$ embryos/time point. Data are presented as the mean $\pm$ SD. *: $p < 0.05$, two-tailed unpaired t-test.

There were 163 genes whose transcripts were specifically upregulated in spinous cells at E16.5 (**Fig 2F**). Among this list, the transcription factor MafB was a prominent candidate since it has been implicated in epidermal differentiation in cultured human epidermal cells (Lopez-Pajares et al., 2015), and it is expressed in suprabasal cells *in vivo* during development (Miyai et al., 2016). Consistent with our transcriptional data, immunofluorescence staining of MafB at E16.5 revealed that it was present in the first suprabasal cell layers in the epidermis but was excluded from granular cells (**Fig 2G,I**), and it was undetectable in ICs at E14.5. Another highly upregulated gene in this list, *Ptgs1*, which encodes the enzyme Cox1, was also enriched in spinous cells but not in ICs at the protein level (**Fig 2H,J**). Therefore, MafB and Cox1 are specific markers for spinous cells in the embryonic epidermis.

MafB inhibits proliferation of intermediate cells

To test the functional role of Maf transcription factors in determining spinous cell fate, we prematurely induced MafB expression in ICs. For this, we collected K10-rtTA;TRE-MafB-HA (here after called K10-MafB) embryos that were collected at E14.5, when suprabasal cells are ICs (**Fig 3, Supplement 1A**). In these embryos, MafB-HA was expressed in about 30% of all suprabasal cells (**Fig 3, Supplement 1B**). Examination of EdU incorporation revealed that HA⁺ cells had a significantly lower rate of incorporation than ICs in controls, demonstrating that MafB expression is sufficient to decrease cell proliferation (**Fig 3A,B**). Notably, however, we found that there was increased proliferation of surrounding MafB-negative cells in mutant embryos (**Fig 3A,B**); while the overall suprabasal proliferation rate was not significantly different in mutants vs controls (**Fig 3, Supplement 1C**). This suggests a potential homeostatic mechanism for maintaining total proliferation during the epidermal stratification process. To determine whether MafB expression was sufficient to repress IC-specific gene expression, we examined IC markers, including Tgm1 and Scara5. While we found a decrease in the levels of Tgm1 protein, Scara5 mRNA levels were unchanged (**Fig 3C-F**). These data demonstrate that MafB may repress parts of the IC signature but does not globally repress all markers. Given that it altered the mitotic status of ICs into a spinous-like state, we also examined whether MafB expression was sufficient for induction of spinous markers. However, we found that the spinous marker Cox1 was not induced by MafB (**Fig 3G,H**). Therefore, MafB is sufficient for some aspects of spinous fate (like loss of proliferation), but not for the expression of all spinous marker genes.

Intermediate cells are granular cell precursors

Using the IC-specific gene signature we identified above, we turned to Gene Ontology (GO) analysis to determine biological pathways characteristic of these cells. Categories that were uniquely enriched in ICs at E14.5 included pathways involved in lipid metabolic processes and establishment of the skin barrier (**Fig 4A**). These pathways are hallmarks of granular cells, which produce and secrete lipids that contribute to the formation of the watertight cornified layer that acts as the epidermal barrier. These findings were unexpected since granular cells have not formed yet at E14.5. Comparing the IC gene signature to a granular gene signature that we generated from published data ((Matsui et al., 2021) GSE168011) revealed a striking overlap with over 1/3 of the genes in the IC signature also being part of the granular cell signature (**Fig 4B**). GO analysis of these shared genes revealed pathways such as keratinocyte differentiation, endomembrane system (reminiscent of lamellar bodies) and lipid processes (**Fig 4, Supplement 1**). Notably, both Grhl3 and Hopx, two transcription factors that control late epidermal differentiation were upregulated in both of these cell types (Chalmers et al., 2006; Chen et al., 2019; Obarzanek-Fojt et al., 2011; Ting et al., 2005a; Ting et al., 2005b; Yu et al., 2006).

To validate the expression of canonical granular markers in ICs, we performed immunofluorescence staining of Tgm1, an enzyme that performs protein cross-linking of structural proteins for cornified envelope formation, and Abca12, which plays a role in




Figure 3

Mis-expression of MafB in intermediate cells is sufficient to reduce proliferation but not to induce spinous fate.

(A) Image of EdU staining (magenta) and MafB-expressing ICs immunolabeled with HA (green) in K10-MafB and control mice at E14.5, doxycycline fed since E12. All scale bars in this panel: 20 μ m. Dotted lines represent the basement membrane.

(B) Percentage of dividing suprabasal cells (EdU+) in control, and HA+ and HA-cells in K10-MafB mice at E14.5 (n= 3 embryos/genotype). Data are presented as the mean $\pm$ SD. *: p < 0.05, **: p < 0.01, ***: p < 0.001, ordinary one-way ANOVA, Tukey's multiple comparisons test.

(C) Immunofluorescence staining of Tgm1 in K10-MafB vs control mice at E14.5.

(D) Quantification of Tgm1 average fluorescence in suprabasal cells of control vs K10-MafB at E14.5 (n= 3 embryos/genotype). Data are presented as the mean $\pm$ SD. *: p < 0.05, two-tailed unpaired t-test.

(E) RNAScope of Scara5 in red and immunostaining with MafB in white.

(F) Quantification of the average number of Scara5 RNAScope puncta per cell in suprabasal layers of control, and MafB+ and MafB-cells of K10-MafB mice at E14.5 (n= 3 embryos/genotype). Data are presented as the mean. ns: not significant, repeated-measures one-way ANOVA, Tukey's multiple comparisons test.

(G) Immunofluorescence staining of Cox1 in K10-MafB vs control mice at E14.5.

(H) Quantification of number of Cox1 puncta per standardized area of suprabasal layer in K10-MafB vs control mice at E14.5. n= 3 embryos/genotype. Data are presented as the mean. ns: not significant, two-tailed unpaired t-test.




Figure 4.

Intermediate cells are transcriptionally similar to granular cells and express granular markers.

(A) Gene Ontology (GO) term enrichment analysis of genes in the intermediate cell signature.

(B) Venn diagram illustrating the number of genes in the intermediate cell signature compared with the granular signature (log2 fold > 1, p < 0.05) for each cell population. A total of 209 genes were commonly enriched in ICs and granular cells, which represents 34% of the total IC signature.

(C and E) Immunofluorescence staining of Tgm1 (C) and Abca12 (E) in green and K14 (red) to mark basal cells in WT embryos at E14.5 vs E16.5. Scale bars: 20 μ m.

(D) Quantification of Tgm1 average fluorescence intensity at suprabasal cells at E14.5 (IC) vs the first two layers of suprabasal cells at E16.5 (spinous cells). n = 3 embryos/time point. Bar represents the mean. *: p < 0.05, two-tailed paired t-test.

(F) Quantification of Abca12 average fluorescence intensity at suprabasal cells at E14.5 (ICs) vs the first two layers of suprabasal cells at E16.5 (spinous cells). n = 3 embryos/time point. Data are presented as the mean $\pm$ SD. *: p < 0.05, two-tailed unpaired t-test.

transporting lipids into lamellar bodies and is disrupted in Harlequin Ichthyosis (Akiyama, 2010; Thomas et al., 2006). Both proteins were already expressed at the protein level in ICs, but not in spinous cells (Fig 4C-F).

Together, these data suggested that in addition to promoting rapid amplification of keratinocytes, intermediate cells may serve as progenitors for the first granular cells that form during development. To test this hypothesis, we performed a pulse-chase experiment in K10-rtTA;TRE-H2B-GFP embryos by injecting a low dose of doxycycline to induce the expression of H2B-GFP in K10-positive cells at E14.5 (i.e. only ICs) (Fig 5, Supplement 1A). We followed the H2B-GFP+ cells over time, and we found that at E15.5, these cells were already more superficial, separated from basal cells by a layer of H2B-GFP negative cells (Fig 5A). Based on their expression of MafB, these cells are the first spinous cells produced (Fig 5A). Strikingly, about 95% of H2B-GFP+ cells did not stain for MafB (Fig 5A'). Thus, rather than converting to a spinous cell fate, they are displaced upwards. When we chased these cells until E16.5, they were located in the upper most layers of the epidermis (Fig 5B), and they co-stained with loricrin, a granular cell marker (Fig 5, Supplement 1A). This indicates that most ICs transition into granular cells without going through a MafB+ state. When labeled cells were chased until E18.5, GFP signal was clearly present in the cornified layer, demonstrating that the traced ICs had terminally differentiated before birth (Fig 5C). The above data suggest that the first spinous cells arise from basal cells and not from differentiation of ICs. To test this, we generated K14-rtTA;TRE-H2B-GFP mice and pulsed them with low levels of doxycycline at E14.5. With these conditions, we specifically labeled basal cells at E14.5 and followed their progeny. In embryos chased to E15.5, we observed that 80% of basal cells and 70% of all MafB+ cells were H2B-GFP+, demonstrating that most spinous cells arise from basal divisions (Fig 5D,D'). Altogether, these data demonstrate that most ICs are direct precursors for granular cells and do not pass through a MafB+ spinous intermediate.

Contractility status changes through epidermal development and is sufficient to drive aspects of granular cell fate

Further analysis of our RNA-Seq dataset suggested that intermediate cells may be mechanically as well as molecularly distinct from spinous cells. We examined the list of genes that were upregulated in ICs versus spinous cells, regardless of their expression levels in basal cells (Fig 6A). GO term analysis revealed that regulation of cell morphogenesis and the actin cytoskeleton were among the most highly upregulated pathways in ICs (Fig 6A). Heat map analysis of the contractome gene set (Zaidel-Bar et al., 2015), demonstrated clear differences in suprabasal cell expression at E14.5 and E16.5 (Fig 6B). To validate these findings, we stained for markers that indicate contractility status. We observed that F-actin levels were higher in basal and suprabasal cells at E14.5, as well as in granular cells at E16.5. However, they were low in spinous cells at this later time point (Fig 6C-C'). Myosin IIA, a major contributor to actomyosin contractility in the epidermis (Miroshnikova et al., 2018; Sumigray et al., 2012), was also higher in ICs than spinous cells (Fig 6D-D'). Similarly, α 18, an antibody that recognizes an epitope of α -catenin that is exposed when adherens junctions are under tension (Yonemura et al., 2010), was also higher in basal and suprabasal cells at E14.5 vs E16.5 (Fig 6E-E'). Furthermore, active nuclear YAP, a mechanoresponsive transcriptional co-activator, was present in ICs and granular cells at E16.5, but not in spinous cells (Fig 6, Supplement 1). Similarly, the percentage of basal cells at E14.5 with active YAP was significantly higher than basal cells at E16.5. Apart from its role in mechanotransduction, the active form of YAP is an important driver of proliferation in the basal layers of the epidermis (Aragona et al., 2020; Schlegelmilch et al., 2011; Zhang et al., 2011a). To test a potential role of YAP in IC proliferation, we used K10-rtTA;TRE-YAP^{S112A}-GFP (hereafter called K10-YAP^{CA}) mice to induce an active form of YAP, marked by H2B-GFP, in suprabasal cells at E16.5. Active YAP was not sufficient to induce proliferation in spinous cells; however, it resulted in a non-cell autonomous increase in proliferation of basal cells (Fig 6, Supplement 2).




Figure 5.

Intermediate cells do not differentiate into spinous cells.

(A, B and C) Immunofluorescence staining of spinous cells with MafB in red and labeled cells with H2B-GFP from K10-rtTA;TRE-H2B-GFP mice that were injected with a low dose of doxycycline at E14.5 to label only ICs and sacrificed at E15.5 (A), E16.5 (B) or E18.5 (C).

Basement membrane is indicated as a dotted line. Scale bars: 20 μ m.

(A' and B') Percentage of MafB+ and MafB-suprabasal cells expressing H2B-GFP at E15.5 (n = 3 embryos) (A) and E16.5 (n = 4 embryos) (B). Data are presented as the mean $\pm$ SD. ****: $p < 0.0001$, two-tailed unpaired t-test.

(D) Immunofluorescence staining of spinous cells with MafB in red and labeled cells with H2B-GFP in K14-rtTA;TRE-H2B-GFP mice that were injected with a low dose of doxycycline at E14.5 and sacrificed at E15.5. Scale bar: 20 μ m.

(D') Percentage of basal cells and MafB+ suprabasal cells expressing H2B-GFP. Paired samples are quantifications from the same mouse (n = 3 K14-rtTA;TRE-H2B-GFP embryos).




Figure 6

Intermediate cells show upregulation of contractility markers as compared to spinous cells.

(A) GO term enrichment analysis of genes upregulated in intermediate cells vs spinous cells at E16.5, as shown in the above diagram, regardless of their expression in basal cells ($\log_2$ fold > 1, $p < 0.05$) reveals upregulation of biological processes related to cytoskeleton organization.

(B) RNA-Seq heatmap depicting differential expression of contractome genes in intermediate and spinous cells at E16.5. Gene expression by FPKM was $\log_2$ -transformed.

(C, D and E) Immunofluorescence staining of Phalloidin (C), MyosinIIA (D) and α 18 (E) in green at E14.5 vs E16.5. Basement membrane is indicated as a dotted line. Scale bars: 20 μ m.

(C', D' and E') Fluorescence intensity of suprabasal-suprabasal cell boundaries in the first two layers of suprabasal cells. Line scans were performed in WT embryos at E14.5 (between intermediate cells) and E16.5 (between spinous cells). Line scan graphs show centered measurements across six cell-cell boundaries at each time point. Data are presented as the mean $\pm$ SD (shown as colored shadows in green at E14.5 and pink at E16.5). Scale bars: 20 μ m.

(C'', D'' and E'') Quantification of average fluorescence intensity from the line scan maximum values at suprabasal-suprabasal cell boundaries in WT E14.5 and E16.5. For bar plots, bars represent the mean, joined dots are paired samples from each time point. $n = 3$ embryos/time point, average maximum values of at least 15 line scans at suprabasal cell boundaries in each embryo were calculated. *: $p < 0.05$, two-tailed paired t-test.

The apparent changes in contractility through epidermal development raised the question of whether the shared contractility between ICs and granular cells lies downstream of differentiation, or if it contributes to their similar gene expression. To test the effects of heightened contractility in the suprabasal cells, we turned to mouse models we previously developed that allow doxycycline-induced actomyosin contractility: K10-rtTA; TRE-Spstin and K10-rtTA, TRE-Arhgef11^{CA}, hereafter referred as K10-Spstin and K10-Arhgef11^{CA}, respectively (Hinnant et al., 2024 [↗](#); Muroyama and Lechler, 2017 [↗](#); Muroyama et al., 2018 [↗](#); Ning et al., 2021 [↗](#)). In both cases this expression is confined to suprabasal cells. K10-Spstin expresses the active form of the microtubule severing protein Spstin which increases contractility (Ning et al., 2021 [↗](#)); and K10-Arhgef11^{CA} induces expression of a constitutively active Rho-GEF that activates RhoA GTPase, a key positive regulator of actomyosin contractility (Ning et al., 2021 [↗](#)).

We started by examining the transcriptomic changes that are induced by contractility (from K10-Spstin vs controls at E16.5, log2 fold change ≥ 1 ; GEO: GSE158786) and compared it with the intermediate and granular cell signatures. Remarkably, we found that 47% of contractility-induced genes overlapped with the granular gene signature (**Fig 7, Supplement 1A** [↗](#))—demonstrating a major change in gene expression and differentiation state by intracellular contractility (hypergeometric test representation factor 3.4, $p=2 \times 10^{-101}$). GO analysis revealed that the common signature was highly enriched in keratinocyte differentiation, cornified envelope, lamellar bodies and lipid processes – all hallmarks of granular cells (**Fig 7A** [↗](#)). When comparing with the IC signature, there was still significant overlap – 80 of the 610 genes in the IC signature (13%) were also found in the contractile signature, though less than that found in the granular cells (**Fig 7, Supplement 1B** [↗](#)). These included the late epidermal differentiation transcriptional regulators Grhl3 and Hopx. The most enriched terms for the overlapping genes between ICs and the contractile gene signature were lipid processing and transport (**Fig 7B** [↗](#)). In addition, mainly lipid metabolic genes were upregulated in the shared ICs, granular cell, and contractility signatures (**Fig 7, Supplement 1C** [↗](#)), suggesting that these were the pathways commonly upregulated in IC and granular cells as a consequence of contractility. Notably, many EDC genes were enriched in the contractile and granular signatures (30/61), but many fewer in the IC signature (5/61). This demonstrates specific differentiation pathways are enriched in ICs, while the contractile gene signature is much more similar to the full granular gene expression program.

In contrast to an upregulation of granular gene markers in hypercontractile skin, we noted that about 1/3 of all the spinous signature genes were downregulated in this mutant (55/163) genes, demonstrating a partial suppression of the spinous cell fate.

We next examined the effects of increased contractility through analysis of specific markers for spinous cells we identified earlier in this study. We used both models of increased contractility in suprabasal cells: K10-Spstin and K10-Arhgef11^{CA}, where expression of the transgene for both is detectable by HA staining. In both mouse models, suprabasal cells with increased contractility (at E16.5 and later) did not express the spinous marker MafB (**Fig 7C,E** [↗](#)), even when they were positioned in the first layers on top of basal cells, where they should clearly be spinous cells. In contrast, these cells precociously expressed the granular marker Tgm1 (**Fig 7D,F** [↗](#)). Similar results were obtained in back and glabrous skin of K10-Arhgef11^{CA} adult mice (**Fig 7, Supplement 2** [↗](#)), and at E17.5, when suprabasal contractility was induced after initial spinous cell specification (**Fig 7, Supplement 3** [↗](#)). These latter results suggest that the effects of increased suprabasal contractility are the same regardless of the time of its induction.

Analysis of the IC marker Scara5 revealed that increasing suprabasal contractility did not alter Scara5 mRNA levels in HA+ cells compared to controls; suggesting that contractility induces a granular-like rather than intermediate-like cell type in mutant mice (**Fig 7G** [↗](#)).




Figure 7.

Increased contractility in spinous cells is sufficient to induce many granular genes.

(A and B) GO term analysis of the list of genes commonly upregulated in suprabasal K10-Spастin (vs control spinous cells at E16.5, log 2-fold > 1, $p < 0.05$) and granular cell signature (A) or intermediate cell signature (B).

(C and D) Immunofluorescence staining of MafB (C) or Tgm1 (D) in red, and suprabasal cells expressing Spастin marked by HA (green) in K10-Spастin and controls. Images of E16.5 embryos, doxycycline fed starting at E14.5. Insets to the left of all images in this panel show close-ups of HA+ cells in suprabasal layers adjacent to basal cells. Scale bars: 20 μ m.

(E and F) Immunofluorescence staining of MafB (F) or Tgm1 (H) in red, and suprabasal cells expressing Arhgef11^{CA} marked by HA (green) in K10-Arhgef11 and controls. Images of E17.5 embryos, doxycycline fed from E10.5. Scale bars: 20 μ m.

(C' and E') Percentage of MafB+ suprabasal cells in controls, and in HA+ and HA- cells in K10-Spастin at E16.5 (C') or K10-Arhgef11^{CA} at E17.5 (E'). Total number of mice analyzed was three embryos per genotype. Data are presented as the mean $\pm$ SD. *: $p < 0.05$, ****: $p < 0.0001$, repeated-measures one-way ANOVA, Tukey's multiple comparisons test. For all plots in this panel, cells located in the first two layers of suprabasal cells immediately above the basal layer were considered for quantification.

(D' and F') Quantification of average fluorescence intensity of Tgm1 in suprabasal cells of control and HA+ cells in K10-Spастin at E16.5 (D') or K10-Arhgef11 at E17.5 (F'). Total number of mice analyzed was three embryos per genotype. *: $p < 0.05$, two-tailed paired t-test.

(G) RNAscope of Scar5 in red and immunostaining with HA in green in control and K10-Arhgef11. Images of E16.5 embryos, doxycycline fed starting at E14.5. Scale bars: 20 μ m. (G') Violin plot of number of Scar5 puncta per suprabasal cell in control, and HA+ cells and neighbor HA-cells in K10-Spастin at E16.5. $n = 224$ HA+ cells and 160 WT cells from 2 embryos per genotype. ns: not significant, two-tailed unpaired t-test.

YAP concentrates within the nucleus in suprabasal epidermal cells in both of our mouse models that increase contractility (Ning et al., 2021 [↗](#)), and YAP is also localized to the nucleus in granular cells (Fig 6, Supplement 1 [↗](#)). To test the role of YAP in inducing a granular-like state, we again used the mouse model K10-YAP^{CA}. We observed that inducing active YAP did not induce Tgm1 in spinous cells and did not suppress the expression of MafB, the spinous cell marker (Fig 7, Supplement 4 [↗](#)). These data suggest that YAP activation lies downstream of contractility but is not sufficient to induce the effects observed by increased contractility in spinous cells. In addition to YAP activation, stretching of keratinocytes results in epigenetic changes, including an increase in H3K27me3 levels (Le et al., 2016 [↗](#)). However, inducing contractility did not cause a significant change in nuclear levels of this epigenetic mark (Fig 7, Supplement 5 [↗](#)).

Discussion

Embryonic epidermal cells coordinate their growth and differentiation to allow rapid stratification and barrier formation. Our data here characterize intermediate cells in epidermal development, revealing novel trajectories of cell differentiation. Further, our data directly demonstrate that actomyosin contractility in the developing epidermis can play an instructive role in differentiation – promoting granular-like gene expression and suppressing spinous markers.

Bulk RNA-Seq analysis has allowed us to characterize embryonic epidermal cells in greater depth. This revealed that basal cells undergo significant changes between E14.5 and E16.5, consistent with the fact that they generate distinct cell progeny at these time points (intermediate vs spinous cells). In addition, like suprabasal cells, basal cells at these time points also show differences in mechanical properties. Whether this is in response to changes in their physical environment or is mediated by internal timers remains unknown. That said, understanding specific markers and transcriptional regulators of these maturing basal cells may allow us to manipulate these transitions to maintain or bypass distinct stages of epidermal development.

Analysis of the intermediate cell transcriptome revealed an unexpected expression of granular genes in these transient cells. This gene expression program was especially enriched for lipid-modifying enzymes that play a key role in barrier formation. Lipids are synthesized in granular cells and stored in lamellar bodies, before secreting them to the intercellular spaces (Prado-Mantilla and Lechler, 2023 [↗](#); Vietri Rudan and Watt, 2021 [↗](#)). Our data suggest that this process may begin early in intermediate cells. In contrast, few EDC genes were upregulated in intermediate cells, suggesting that only parts of the granular program are expressed early in these cells, and many of the structural components of the eventual barrier are expressed only later.

Further, our short-term lineage tracing suggests that most intermediate cells transition into granular cells without going through a MafB+ (a spinous cell marker) stage. This reveals two distinct differentiation pathways leading to the granular cell state, one through intermediate cells during development, and one through spinous cells post-natally. This raises questions about whether each route will require distinct regulatory and transcriptional pathways.

Finally, our data revealed mechanical similarities between intermediate and granular cells, in addition to their similarities at the transcriptomic level. Actomyosin regulators were transcriptionally upregulated in both of these cell types as compared to spinous cells, and we were able to validate these findings in intact skin. Importantly, we found that contractility was sufficient to induce granular-like gene expression and to repress some spinous cell markers. This reveals a rather direct effect of contractility lying upstream, as well as presumably downstream, of differentiation pathways. It also explains the observation that increased contractility results in premature formation of the epidermal barrier (Muroyama and Lechler, 2017 [↗](#); Ning et al., 2021 [↗](#)). It has been appreciated for some time that granular cells have increased actomyosin contractility, which is important for the function and placement of tight junctions specifically in

this layer of the epidermis (Itoh et al., 2012 [DOI](#); Prado-Mantilla and Lechler, 2023 [DOI](#); Rubsam et al., 2017 [DOI](#); Sumigray et al., 2012 [DOI](#)). There are many intriguing possibilities for how contractility may affect cell fates, such as mechanosensitive transcriptional regulators that mediate this effect. We tested a possible role for YAP1, a transcriptional regulator whose activation is induced by contractility in the epidermis. We found that active YAP1 was insufficient to drive these gene expression changes. Future work is required to determine the mechanism of mechanosensitive differentiation in the epidermis. The epidermis increases contractility in response to externally applied forces that result in stretch, and in response to perturbation of cell adhesions (Aragona et al., 2020 [DOI](#); Sumigray et al., 2014 [DOI](#)). It will be important to test whether these stimuli alter differentiation pathways through their effects on contractility.

Of note, we found that contractility had a strong effect on the EDC, whose positioning within the nucleus changes with differentiation (Gdula et al., 2013 [DOI](#); Mardaryev et al., 2014 [DOI](#); Williams et al., 2002 [DOI](#)). This suggests that aspects of epigenetics and chromatin organization may be downstream of contractility, as shown by external forces directly stretching chromatin (Tajik et al., 2016 [DOI](#)) and extrinsic topographical cues constricting the nucleus to induce chromatin reprogramming (Wang et al., 2023 [DOI](#)). Notably, we found that there is a decrease in mRNA levels of Sun1 and Nesprin1/3, parts of a complex that tether the actin cytoskeleton in the cytoplasm to chromatin in the nucleus when contractility is increased, in hypercontractile cells. Loss of Sun1/2 also leads to precocious differentiation in the epidermis (Carley et al., 2021 [DOI](#)) through effects on integrin-mediated mechanical integration, though whether regulation of these connections mediates the contractility effect seen in suprabasal cells will require further investigation.

Material and methods

Mice

All animal work was approved by Duke University's Institutional Animal Care and Use Committee and performed in accordance with their committee guidelines. Mice were genotyped by PCR, and both male and female mice were analyzed in this study. All mice were maintained in a barrier facility with 12-hour light/dark cycles. Mouse strains used in this study include: K14-RFP (Zhang et al., 2011b [DOI](#)), K10-rtTA and TRE-Spstin (Muroyama and Lechler, 2017 [DOI](#)), TRE-Arhgef11^{CA} (Ning et al., 2021 [DOI](#)), CD1 (Charles River, strain code: 022). Other mice were from the Jackson Laboratories, and their stock numbers are as follows: TRE-H2B-GFP (005104), TRE-YAP1^{S112A}-GFP (031279, (Gao et al., 2013 [DOI](#))).

Generation of the TRE-MafB-HA mouse line

MafB with an HA tag on the C-terminus followed by a stop codon was synthesized by GenScript and ligated into the pTRE2 vector. To verify the proper doxycycline-dependent expression of the TRE-MafB-HA cassette, the vector was co-transfected with a K14-rtTA plasmid into cultured keratinocytes and placed in doxycycline-containing media. TRE-MafB-HA plasmid was linearized, purified, and used by the Duke Transgenic Core to generate transgenic TRE-MafB-HA mice via pronuclear injection.

Skin explant preparation

Back skin of embryonic mouse skin was dissected, rinsed, and gently unfolded in sterile PBS, then placed on a 2% agarose pad diluted in a 1:1 sterile water:media mixture (10% FBS in DMEM, doxycycline (2 µg/ml, to induce H2B-GFP expression) and 1:100 Penicillin/Streptomycin). Note that the dermis of the skin explant faces the agarose pad. The skin explants on agarose were gently placed into a 6 cm dish containing 1 mL 10% FBS in DMEM (Gibco, 11965). The explants were cultured at 37°C and 7.5% CO₂ for at least 2 hours for recovery before live imaging.

Sample preparation for RNA-Seq

K10-rtTA;TRE-H2B-GFP;K14-RFP pregnant dams were all fed with doxycycline chow at E12.5, and then sacrificed at either E14.5 or E16.5. Embryos were checked under a dissecting microscope for GFP and RFP expression before use, and their tails were taken to confirm genotypes. E16.5 back skins were cut off and treated with 1.4 U/ml dispase II (Roche, 4942078001) in HBSS at room temperature for 1 hour, then peeled off the epidermis from the dermis. The epidermis of E14.5 back skin was not peeled off due to its low thickness. Both E16.5 epidermis and E14.5 skin samples were digested in 1:1 trypsin (GIBCO, 25200-056) with versene (GIBCO, 15040-066) at 37°C and rotated for 20 minutes. Samples were then mixed 1:1 with FACS buffer (HBSS with 2.5% FBS and 10 µg/ml DNAaseI) and centrifuged at low speed. The cell pellet was diluted into FACS buffer with propidium iodide solution (Sigma, P4864, to exclude dead cells), and filtered using sterile Celltrics 30 µm filters (Sysmex, 04-004-2326). The lectin UEA1 (Ulex europaeus agglutinin I) binds to carbohydrates in cell surfaces present in uppermost keratinocytes. To exclude them, we used biotinylated UEA1 (Vector Laboratories, B-1065) and detected them with APC-streptavidin (BioLegend, 405207). The E14.5 intermediate cells (K10-GFP⁺;K14-RFP⁺;APC-UEA1⁻), E16.5 spinous cells (K10-GFP⁺;K14-RFP⁺;APC-UEA1⁻), and basal cells at E14.5 and E16.5 (K10-GFP⁺;K14-RFP⁺) cells were FACS sorted separately. RNA was extracted using a QIAGEN RNAeasy Mini kit (QIAGEN, 74104) following the manufacturer's protocols, with DNA digested using RNase-Free DNase (QIAGEN, 79254). Biological replicates, three independent RNA samples from each cell population, were collected and sent for sequencing and analysis by Novogene.

RNA-Seq analysis

Differential expression analysis between two time points or cell types (three biological replicates per group) was performed using the DESeq2 R package. The gene signature of a cell population was obtained by the overlapping genes that were enriched compared to the suprabasal cells at a different time point and enriched compared to the basal cells at its corresponding time point. A significance threshold of adj p-value < 0.05, log2 fold change ≥ 1 and FPKM ≥ 1 was used for defining marker genes of each cell population.

GO term analysis for differentially expressed genes and gene signatures was performed using the Gene Ontology enrichment analysis (Ashburner et al., 2000 [DOI](#); Gene Ontology et al., 2023 [DOI](#); Thomas et al., 2022 [DOI](#)).

FPKM of differentially expressed genes in the contractome list were log2-transformed and used for heatmaps. Heatmaps were generated using the online resource Heatmapper (Babicki et al., 2016 [DOI](#)), using the average linkage clustering method and Pearson's distance measurement methods.

PCA plot of all cell population data was generated using the web tool Clustvis: <https://biit.cs.ut.ee/clustvis/> (Metsalu and Vilo, 2015 [DOI](#)).

Granular cell signature was determined from genes significantly upregulated in ETA sheets vs epidermal sheets (p-value < 0.05, log2 fold change ≥ 1) obtained from the previously published dataset: GSE168011 (Matsui et al., 2021 [DOI](#)).

Immunofluorescence

Fresh tissue was embedded in OCT (Sakura), frozen, and sectioned at 10 µm using a cryostat. Sections were fixed with 4% paraformaldehyde (PFA) in PBS for 8 minutes at room temperature or ice-cold acetone (for Tgm1 and St8sia6 staining) for 2 minutes, washed with PBS containing 0.2% Triton (PBST) for 5 minutes, then blocked with blocking buffer (3% bovine serum albumin with 5% normal goat serum (Gibco, 16210064), and 5% normal donkey serum (Sigma-Aldrich, D9663) in

PBST) for 15 minutes. Sections were incubated with primary antibodies diluted in blocking buffer for 1h at room temperature (α 18 antibody was incubated for 15 minutes), then three times washed with PBST, and incubated in secondary antibodies and stains, such as Hoechst 34580 or Phalloidin, for 15 minutes at room temperature. After three washes with PBST, sections were finally mounted in the anti-fade buffer (90% glycerol in PBS plus 2.5 mg/ml p-Phenylenediamine (Thermo Fisher, 417481000)) and sealed using clear nail polish in the borders.

Primary antibodies used in this study: rat anti-HA (Sigma-Aldrich, 11867423001), chicken anti-keratin 5/14 (generated in the Lechler lab), rabbit anti-keratin 10 (Covance, 905401), guinea pig anti-keratin 10 (Progen, GP-K10), rat anti- β 4 integrin (BD Biosciences, 553745), rabbit anti-MyosinIIA (Biolegend, PRB-440P), rat anti- α 18 (gift from Akira Nagafuchi, Kumamoto University), rabbit anti-YAP/TAZ (Cell Signaling Technology, 8418S), rabbit anti-Tgm1(Proteintech,12912-3-AP), rabbit anti-MafB (Novus, NBP1-81342), rabbit anti-St8sia6 (Sigma, HPA011635), rabbit anti-Abca12 (gift from Dr. Wong, University of Michigan). F-actin was stained with phalloidin-488 (Invitrogen, A12379).

EdU labeling

Pregnant dams were intraperitoneally injected with 10 mg/kg of EdU and sacrificed after 1 hour for tissue dissection. Back skins of embryos were collected, and tails were taken for genotyping. Tissue sections were fixed with 4% PFA and stained with primary and secondary antibodies, then EdU was detected following Click-iT EdU (Thermo Fisher, C10337) protocol.

Lineage tracing/pulse-chase

K10-rtTA;TRE-H2B-GFP pregnant dams were intraperitoneally injected with a low dose of doxycycline (1mg/kg) at E14.5 to label only intermediate cells. Dams were sacrificed, and their embryos were collected after 1 day (E15.5), 2 days (E16.5), or 4 days (E18.5) for tissue dissection. Their tails were kept for genotyping and K10-rtTA;TRE-H2B-GFP embryos were sectioned and analyzed.

RNA-Scope

RNA-Scope was performed using the Multiplex Fluorescent v1 and v2 systems (ACD, 323100) followed by antibody co-staining. Back skins from mouse embryos were freshly frozen in OCT and sectioned at 10 μ m. Tissue sections were fixed for 1 hour with 4% PFA at 4°C. After fixation, standard RNA-Scope protocols were used according to manufacturer's instructions. The following probes were used: St8sia6 (ACD, 887831-C1) and Scara5 (ACD, 522301-C1). TSA Vivid Fluorophore 570 (ACD, 323272) was used to develop probe signal. Then, antibody staining was performed to quantify the probe signal in suprabasal cells, using guinea pig anti-keratin 10 antibody, and in basal cells, using chicken anti-keratin 5/14. HA⁺ cells were marked with rat anti-HA or rabbit anti-MafB (for K10-MafB samples). Coverslips were mounted using Prolong Gold (Invitrogen, P10144).

Imaging

For live imaging of K10⁺ cell division, prepared skin explants were placed upside down in a Lumox dish 35 (Sarstedt, 94.6077.331) with the epidermal side facing toward the membrane. Samples were imaged at 15 minute-intervals overnight using the MetaMorph software on an Andor XD revolution spinning disc confocal microscope at 37°C and 5% CO₂ using a 20x/0.5 UplanFL N dry objective.

For immunofluorescence staining, tissue sections were imaged on a Zeiss AxioImager Z1 microscope with Apotome.2 attachment, Plan-APOCHROMAT 20X/0.8 objective, Plan-NEOFLUAR 40X/1.3 oil objective, or Plan-NEOFLUAR 63X/1.4 oil objective, AxioCam 506 mono camera, and acquired using Zen software (Zeiss). When making intensity measurement comparisons, all images within one experiment were taken with identical exposure times.

Image quantification and statistics

All image quantifications were done using FIJI software. Quantifications of fluorescence intensity of cortical F-actin, Myosin IIA and α 18 were measured by drawing 10-pixel wide lines across cell-cell boundaries. Maximum values from plot profiles were aligned to yield the final line scan plots. The mean of the maximum values of each line scan was calculated per mouse and compared between conditions or time points to determine statistical significance.

EdU, Scara5, St8sia6, YAP and MafB measurements were determined by calculating the percentage of the number of positive cells in the total number of suprabasal cells in the first two suprabasal layers adjacent to basal cells. Quantifications are from at least three fields per mouse.

Tgm1 average fluorescence intensity was calculated by measuring the mean value of suprabasal cells above basal cells. This area was defined by drawing 50-pixel width lines in the suprabasal area, along the labeled basal layer, using the Freehand line tool in FIJI.

All statistical analyses were performed using GraphPad Prism 10 software. Data shown in bar plots was presented as mean $\pm$ standard deviation (SD), and significance was determined using two-tailed Student's t-test, or, for multiple comparisons, one-way or two-way ANOVA, followed by Tukey's or Sidak tests. Further details in statistical analysis were specified in figure legends. Data were determined to be statistically significant when p value < 0.05 . Asterisks denote statistical significance (ns = not significant, *: p < 0.05 , **: p < 0.01 , ***: p < 0.001 , ****: p < 0.0001).

Data and software availability

Raw data files for the RNA sequencing analysis have been deposited in the NCBI Gene Expression Omnibus under accession number GEO: TBD.

Acknowledgements

We thank Julie Underwood for expert care of our mice and lab management as well as members of the Lechler Lab for comments on the manuscript. Maggie Bara provided initial characterization of the TRE-MafB line, David Kirsch provided mouse lines, and Sunny Wong provided Abca12 antibody.




Figure 1-Supplement 1

Live imaging of mitotic suprabasal cells during early epidermal stratification.

(A) Diagram depicting alleles used for live imaging of suprabasal cells, and treatment timeline: Pregnant dams were doxycycline fed 48 hrs before embryos were collected and their tissue harvested for explant imaging.

(B) Snapshots from live imaging of back skin explants from E14.5, E15.5 and E16.5 K10-rtTA;TRE-H2B-GFP embryos, which express H2B-GFP in suprabasal cells. Images are from the first plane of H2B-GFP⁺ cells, corresponding to the suprabasal layer immediately on top of basal cells from E14.5, E15.5 and E16.5 explants. Time interval between snapshots: 15 min. Scale bars: 20 μ m.

(C) Percentage of mitotic events counted during 2 hrs of live imaging. Only suprabasal cells from layers on top of basal cells were quantified. $n=5$ different fields for E14.5 and E15.5, and $n=4$ different fields for E16.5 from at least 2 embryos for each time point were analyzed. ****: $p < 0.001$, ordinary one-way ANOVA, Tukey's multiple comparisons test.

Figure 2-Supplement 1

Intermediate cells do not express some granular markers at the protein level.

(A and B) Immunofluorescence staining of Loricrin(A) or St8sia6 (B) in green labeling granular cells at E16.5 but not in intermediate cells at E14.5. Basement membrane is marked with β 4-integrin in red. Scale bars: 20 μ m.




Figure 3-Supplement 1

The overall rate of suprabasal cell proliferation in K10-MafB is not significantly different from controls.

(A) Diagram depicting alleles in K10-MafB used to induce MafB expression in intermediate cells. Pregnant dams were fed with doxycycline chow from E12.5 and sacrificed at E14.5.

(B) Percentage of suprabasal cells expressing MafB (HA+) in K10-MafB mice at E14.5.

(C) Percentage of dividing cells among all suprabasal cells at E14.5 in K10-MafB vs controls is not significantly different. n= 3 embryos/genotype. Data are presented as the mean. ns: not significant, two-tailed unpaired t-test.




Figure 4-Supplement 1

Intermediate and granular cells shared pathways related to lipid biosynthetic processes and endomembrane systems.

GO analysis from commonly enriched genes in intermediate and granular cell signatures.

Commonly upregulated pathways in IC and granular cells





Figure 5-Supplement 1

Intermediate cells differentiate into granular cells

(A) Image showing H2B-GFP expression in intermediate cells in a E14.5 K10-rtTA;TRE-H2B-GFP mouse fed with doxycycline from E12.5. Basement membrane is indicated as a dotted line. Scale bars: 20 μm .

(B) Immunofluorescence staining of Loricrin in red colocalizing with H2B-GFP labeled cells in K10-rtTA;TRE-H2B-GFP mice injected with a low dose of doxycycline at E14.5 and sacrificed at E16.5. Scale bars: 20 μm .






Figure 6-Supplement 1

Like granular cells, more intermediate cells have nuclear YAP than spinous cells.

(A) Immunofluorescence staining of YAP/TAZ at E14.5 and E16.5. Scale bars: 20 μ m.

(B) Percentage of nuclear active YAP in basal cells and in the first two suprabasal layers at E14.5 and E16.5. Active YAP was also quantified in granular cells at E16.5. *: $p < 0.05$, multiple unpaired t-test. $n = 3$ embryos/time point.




Figure 6-Supplement 2

Inducing constitutively active YAP in spinous cells does not increase their proliferation.

(A) Image of EdU staining in magenta and immunofluorescence of Loricrin in red from control and K10-YAP^{CA} embryos at E16.5. Cells expressing constitutively active nuclear YAP are labeled with H2B-GFP in green. Mice were fed with doxycycline from E13.5. Scale bars: 20 μ m.

(B) Percentage of dividing suprabasal cells in control vs K10-YAP^{CA}, $n = 3$ embryos/genotype. For all plots in this panel, cells located in the first two layers of suprabasal cells immediately above the basal layer were considered for quantification. *: $p < 0.05$, ns: not significant, two-tailed unpaired t-test.




Figure 7-Supplement 1

K10-Spастin cells are more granular-like than intermediate-like, and lipid metabolic processes are commonly upregulated among K10-Spастin, IC and granular cells.

(A and B) Size proportional-Venn Diagrams show that K10-Spастin cells are more granular-like (A: 46.6% similarity) than intermediate-like (B: 11.7% similarity). Green: genes enriched in K10-Spастin vs WT embryos at E16.5; blue: genes in intermediate cell signature, pink: genes in granular signature.

(C) GO term analysis of genes commonly upregulated in ICs, granular cells and K10-Spастin (total: 50 genes) are lipid metabolic processes.




Figure 7-Supplement 2

Increased contractility in spinous cells of adult epidermis induces some granular markers in adult epidermis.

(A and B) Immunofluorescence staining of MafB in back skin (A) or glabrous skin (B) in red, and suprabasal cells expressing Arhgef11^{CA} marked by HA (green) in adult K10-Arhgef11 and controls. Mice were fed with doxycycline since adulthood. Insets show close-ups of HA+ cells in suprabasal layers adjacent to basal cells. Scale bars: 20 μ m.

(C and D) Immunofluorescence staining of Tgm1 in back skin (C) or glabrous skin (D) in red, and suprabasal cells expressing Arhgef11^{CA} marked by HA (green) in adult K10-Arhgef11 and controls. Insets show close-ups of HA+ cells in suprabasal layers adjacent to basal cells. Mice were fed with doxycycline since adulthood. Scale bars: 20 μ m.

Figure 7-Supplement 3

Increased contractility in spinous cells accelerates acquisition of granular markers.

(A and B) Immunofluorescence staining of MafB (A) or Tgm1 (B) in red, and suprabasal cells expressing Arhgef11^{CA} marked by HA (green) in adult K10-Arhgef11 and controls. Images from E17.5 embryos, fed with doxycycline starting at E15.5. Insets show close-ups of HA+ cells in suprabasal layers adjacent to basal cells. Scale bars: 20





Figure 7-Supplement 4

Inducing constitutively active YAP in spinous cells does not induce granular markers

(A and C) Immunofluorescence of MafB (A) or Tgm1 (C) in red, active YAP in green (YAP-H2B-GFP) in control, and K10-YAP^{CA} at E16.5. Scale bars: 20 μm.

(B) Average MafB fluorescence intensity per cell in control and K10-YAP^{CA}. n=113 suprabasal cells from 3 control embryos, and 103 from 3 K10-YAP^{CA} embryos. ns: not significant, two-tailed unpaired t-test.

(D) Average fluorescence intensity of suprabasal cells, area above basal cells in control and K10-YAP^{CA} embryos at E16.5 were measured. n= 3 embryos/genotype. ns: not significant, two-tailed unpaired t-test.






Figure 7-Supplement 5

Increased contractility does not alter H3K27me3 levels.

Quantifications of nuclear H3K27me3 levels in control spinous and K10-Arhgef-expressing spinous cells.

References

- Akiyama M (2010) **ABCA12 mutations and autosomal recessive congenital ichthyosis: a review of genotype/phenotype correlations and of pathogenetic concepts** *Human mutation* **31**:1090–1096
- Aragona M. *et al.* (2020) **Mechanisms of stretch-mediated skin expansion at single-cell resolution** *Nature* **584**:268–273
- Ashburner M. *et al.* (2000) **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium** *Nature genetics* **25**:25–29
- Babicki S., Arndt D., Marcu A., Liang Y., Grant J.R., Maciejewski A., Wishart D.S. (2016) **Heatmapper: web-enabled heat mapping for all** *Nucleic Acids Res* **44**:W147–153
- Brabec R.K., Peters B.P., Bernstein I.A., Gray R.H., Goldstein I.J. (1980) **Differential lectin binding to cellular membranes in the epidermis of the newborn rat** *Proceedings of the National Academy of Sciences of the United States of America* **77**:477–479
- Carley E., Stewart R.M., Ziemann A., Jalilian I., King D.E., Zubek A., Lin S., Horsley V., King M.C. (2021) **The LINC complex transmits integrin-dependent tension to the nuclear lamina and represses epidermal differentiation** *Elife* **10**
- Chalmers A.D., Lachani K., Shin Y., Sherwood V., Cho K.W., Papalopulu N. (2006) **Grainyhead-like 3, a transcription factor identified in a microarray screen, promotes the specification of the superficial layer of the embryonic epidermis** *Mechanisms of development* **123**:702–718
- Chen S.Y., Ishii M.A., Cheng B., Otten A.B.C., Sun B.K. (2019) **HOPX Is a ZNF750 Target that Promotes Late Epidermal Differentiation** *The Journal of investigative dermatology* **139**:2039–2042
- Damen M., Wirtz L., Soroka E., Khatif H., Kukat C., Simons B.D., Bazzi H. (2021) **High proliferation and delamination during skin epidermal stratification** *Nat Commun* **12**
- Engler A.J., Sen S., Sweeney H.L., Discher D.E. (2006) **Matrix elasticity directs stem cell lineage specification** *Cell* **126**:677–689
- Gao T., Zhou D., Yang C., Singh T., Penzo-Mendez A., Maddipati R., Tzatsos A., Bardeesy N., Avruch J., Stanger B.Z. (2013) **Hippo signaling regulates differentiation and maintenance in the exocrine pancreas** *Gastroenterology* **144**:1543–1553
- Gdula M.R., Poterlowicz K., Mardaryev A.N., Sharov A.A., Peng Y., Fessing M.Y., Botchkarev V.A. (2013) **Remodeling of three-dimensional organization of the nucleus during terminal keratinocyte differentiation in the epidermis** *The Journal of investigative dermatology* **133**:2191–2201
- Gene Ontology C. *et al.* (2023) **The Gene Ontology knowledgebase in 2023** *Genetics* **224**
- Hinnant T., Ning W., Lechler T. (2024) **Compartment specific responses to contractility in the small intestinal epithelium** *PLoS Genet* **20**

- Huebner A.J., Dai D., Morasso M., Schmidt E.E., Schafer M., Werner S., Roop D.R. (2012) **Amniotic fluid activates the nrf2/keap1 pathway to repair an epidermal barrier defect in utero** *Developmental cell* **23**:1238–1246
- Itoh M., Tsukita S., Yamazaki Y., Sugimoto H. (2012) **Rho GTP exchange factor ARHGEF11 regulates the integrity of epithelial junctions by connecting ZO-1 and RhoA-myosin II signaling** *Proceedings of the National Academy of Sciences of the United States of America* **109**:9905–9910
- Jacob T. *et al.* (2023) **Molecular and spatial landmarks of early mouse skin development** *Developmental cell* **58**:2140–2162
- Koster M.I., Roop D.R. (2005) **Asymmetric cell division in skin development: a new look at an old observation** *Developmental cell* **9**:444–446
- Le H.Q. *et al.* (2016) **Mechanical regulation of transcription controls Polycomb-mediated gene silencing during lineage commitment** *Nature cell biology* **18**:864–875
- Lechler T., Fuchs E. (2005) **Asymmetric cell divisions promote stratification and differentiation of mammalian skin** *Nature* **437**:275–280
- Lopez-Pajares V. *et al.* (2015) **A LncRNA-MAF:MAFB transcription factor network regulates epidermal differentiation** *Developmental cell* **32**:693–706
- Mardaryev A.N. *et al.* (2014) **p63 and Brg1 control developmentally regulated higher-order chromatin remodelling at the epidermal differentiation complex locus in epidermal progenitor cells** *Development (Cambridge, England)* **141**:101–111
- Matsui T. *et al.* (2021) **A unique mode of keratinocyte death requires intracellular acidification** *Proceedings of the National Academy of Sciences of the United States of America* **118**
- Metsalu T., Vilo J. (2015) **ClustVis: a web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap** *Nucleic Acids Res* **43**:W566–570
- Meyer-ter-Vehn T., Sieprath S., Katzenberger B., Gebhardt S., Grehn F., Schlunck G. (2006) **Contractility as a prerequisite for TGF-beta-induced myofibroblast transdifferentiation in human tenon fibroblasts** *Invest Ophthalmol Vis Sci* **47**:4895–4904
- Miroshnikova Y.A. *et al.* (2018) **Adhesion forces and cortical tension couple cell proliferation and differentiation to drive epidermal stratification** *Nature cell biology* **20**:69–80
- Miyai M., Hamada M., Moriguchi T., Hiruma J., Kamitani-Kawamoto A., Watanabe H., Hara-Chikuma M., Takahashi K., Takahashi S., Kataoka K. (2016) **Transcription Factor MafB Coordinates Epidermal Keratinocyte Differentiation** *The Journal of investigative dermatology* **136**:1848–1857
- Moreci R.S., Lechler T. (2020) **Epidermal structure and differentiation** *Curr Biol* **30**:R144–R149
- Muroyama A., Lechler T. (2017) **A transgenic toolkit for visualizing and perturbing microtubules reveals unexpected functions in the epidermis** *Elife* **6**
- Muroyama A., Terwilliger M., Dong B., Suh H., Lechler T. (2018) **Genetically induced microtubule disruption in the mouse intestine impairs intracellular organization and transport** *Molecular biology of the cell* **29**:1533–1541

- Ning W., Muroyama A., Li H., Lechler T. (2021) **Differentiated Daughter Cells Regulate Stem Cell Proliferation and Fate through Intra-tissue Tension** *Cell Stem Cell* **28**:436–452
- Obarzanek-Fojt M., Favre B., Kypriotou M., Ryser S., Huber M., Hohl D. (2011) **Homeodomain-only protein HOP is a novel modulator of late differentiation in keratinocytes** *Eur J Cell Biol* **90**:279–290
- Prado-Mantilla A., Lechler T. (2023) **Polarity in skin development and cancer** *Curr Top Dev Biol* **154**:317–336
- Rubsam M. *et al.* (2017) **E-cadherin integrates mechanotransduction and EGFR signaling to control junctional tissue polarization and tight junction positioning** *Nat Commun* **8**
- Schlegelmilch K. *et al.* (2011) **Yap1 acts downstream of alpha-catenin to control epidermal proliferation** *Cell* **144**:782–795
- Singh A., Tiwari V.K. (2023) **Transcriptional networks of transient cell states during human prefrontal cortex development** *Front Mol Neurosci* **16**
- Smart I.H. (1970) **Variation in the plane of cell cleavage during the process of stratification in the mouse epidermis** *The British journal of dermatology* **82**:276–282
- Sumigra K., Zhou K., Lechler T. (2014) **Cell-cell adhesions and cell contractility are upregulated upon desmosome disruption** *PLoS One* **9**
- Sumigra K.D., Foote H.P., Lechler T. (2012) **Non-centrosomal microtubules and type II myosins potentiate epidermal cell adhesion and barrier formation** *The Journal of cell biology* **199**
- Sumigra K.D., Lechler T. (2015) **Cell adhesion in epidermal development and barrier formation** *Curr Top Dev Biol* **112**:383–414
- Tajik A., Zhang Y., Wei F., Sun J., Jia Q., Zhou W., Singh R., Khanna N., Belmont A.S., Wang N. (2016) **Transcription upregulation via force-induced direct stretching of chromatin** *Nat Mater* **15**:1287–1296
- Thomas A.C. *et al.* (2006) **ABCA12 is the major harlequin ichthyosis gene** *The Journal of investigative dermatology* **126**:2408–2413
- Thomas P.D., Ebert D., Muruganujan A., Mushayahama T., Albou L.P., Mi H. (2022) **PANTHER: Making genome-scale phylogenetics accessible to all** *Protein Sci* **31**:8–22
- Ting S.B. *et al.* (2005) **A homolog of Drosophila grainy head is essential for epidermal integrity in mice.** *Science (New York N.Y)* **308**:411–413
- Ting S.B., Caddy J., Wilanowski T., Auden A., Cunningham J.M., Elias P.M., Holleran W.M., Jane S.M. (2005) **The epidermis of grhl3-null mice displays altered lipid processing and cellular hyperproliferation** *Organogenesis* **2**:33–35
- Vietri Rudan M., Watt F.M. (2021) **Mammalian Epidermis: A Compendium of Lipid Functionality** *Front Physiol* **12**
- Vining K.H., Mooney D.J. (2017) **Mechanical forces direct stem cell behaviour in development and regeneration** *Nat Rev Mol Cell Biol* **18**:728–742

Wang X. *et al.* (2023) **Chromatin reprogramming and bone regeneration in vitro and in vivo via the microtopography-induced constriction of cell nuclei** *Nat Biomed Eng* **7**:1514–1529

Weiss L.W., Zelikson A.S. (1975) **Embryology of the epidermis: ultrastructural aspects. II. Period of differentiation in the mouse with mammalian comparisons** *Acta dermatovenereologica* **55**:321–329

Williams R.R., Broad S., Sheer D., Ragoussis J. (2002) **Subchromosomal positioning of the epidermal differentiation complex (EDC) in keratinocyte and lymphoblast interphase nuclei** *Experimental cell research* **272**:163–175

Yonemura S., Wada Y., Watanabe T., Nagafuchi A., Shibata M. (2010) . **alpha-Catenin as a tension transducer that induces adherens junction development** *Nature cell biology* **12**:533–542

Yu Z. *et al.* (2006) **The Grainyhead-like epithelial transactivator Get-1/Grhl3 regulates epidermal terminal differentiation and interacts functionally with LMO4** *Dev Biol* **299**:122–136

Zaidel-Bar R., Zhenhuan G., Luxenburg C. (2015) **The contractome--a systems view of actomyosin contractility in non-muscle cells** *Journal of cell science* **128**:2209–2217

Zhang H., Pasolli H.A., Fuchs E. (2011) **Yes-associated protein (YAP) transcriptional coactivator functions in balancing growth and differentiation in skin** *Proceedings of the National Academy of Sciences of the United States of America* **108**:2270–2275

Zhang L., Stokes N., Polak L., Fuchs E. (2011) **Specific microRNAs are preferentially expressed by skin stem cells to balance self-renewal and early lineage commitment** *Cell Stem Cell* **8**:294–308

Editors

Reviewing Editor

Elaine Fuchs

Howard Hughes Medical Institute, The Rockefeller University, New York, United States of America

Senior Editor

Utpal Banerjee

University of California, Los Angeles, Los Angeles, United States of America

Reviewer #1 (Public review):

Summary:

The authors address a fundamental question for cell and tissue biology using the skin epidermis as a paradigm and ask how stratifying self-renewing epithelia induce differentiation and upward migration in basal dividing progenitor cells to generate suprabasal barrier-forming cells that are essential for a functional barrier formed by such an epithelium. The authors show for the first time that an increase in intracellular actomyosin contractility, a hallmark of barrier-forming keratinocytes, is sufficient to trigger terminal differentiation. Hence the data provide in vivo evidence of the more general interdependency

of cell mechanics and differentiation. The data appear to be of high quality and the evidences are strengthened through a combination of different genetic mouse models, RNA sequencing, and immunofluorescence analysis.

To generate and maintain the multilayered, barrier-forming epidermis, keratinocytes of the basal stem cell layer differentiate and move suprabasally accompanied by stepwise changes not only in gene expression but also in cell morphology, mechanics, and cell position. Whether any of these changes is instructive for differentiation itself and whether consecutive changes in differentiation are required remains unclear. Also, there are few comprehensive data sets on the exact changes in gene expression between different states of keratinocyte differentiation. In this study, through genetic fluorescence labeling of cell states at different developmental time points the authors were able to analyze gene expression of basal stem cells and suprabasal differentiated cells at two different stages of maturation: E14 (embryonic day 14) when the epidermis comprises mostly two functional compartments (basal stem cells and suprabasal so-called intermediate cells) and E16 when the epidermis comprise three (living) compartments where the spinous layer separates basal stem cells from the barrier-forming granular layer, as is the case in adult epidermis. Using RNA bulk sequencing, the authors developed useful new markers for suprabasal stages of differentiation like *MafB* and *Cox1*. The transcription factor *MafB* was then shown to inhibit suprabasal proliferation in a *MafB* transgenic model.

The data indicate that early in development at E14 the suprabasal intermediate cells resemble in terms of RNA expression, the barrier-forming granular layer at E16, suggesting that keratinocytes can undergo either stepwise (E16) or more direct (E14) terminal differentiation.

Previous studies by several groups found an increased actomyosin contractility in the barrier-forming granular layer and showed that this increase in tension is important for epidermal barrier formation and function. However, it was not clear whether contractility itself serves as an instructive signal for differentiation. To address this question, the authors use a previously published model to induce premature hypercontractility in the spinous layer by using spastin overexpression (K10-Spastin) to disrupt microtubules (MT) thereby indirectly inducing actomyosin contractility. A second model activates myosin contractility more directly through overexpression of a constitutively active RhoA GEF (K10-Arhgef11CA). Both models induce late differentiation of suprabasal keratinocytes regardless of the suprabasal position in either spinous or granular layer indicating that increased contractility is key to induce late differentiation of granular cells. A potential weakness of the K10-spastin model is the disruption of MT as the primary effect which secondarily causes hypercontractility. However, their previous publications provided some evidence that the effect on differentiation is driven by the increase in contractility (Ning et al. cell stem cell 2021). Moreover, the data are confirmed by the second model directly activating myosin through RhoA. These previous publications already indicated a role for contractility in differentiation but were focused on early differentiation. The data in this manuscript focus on the regulation of late differentiation in barrier-forming cells. These important data help to unravel the interdependencies of cell position, mechanical state, and differentiation in the epidermis, suggesting that an increase in cellular contractility in most apical positions within the epidermis can induce terminal differentiation. Importantly the authors show that despite contractility-induced nuclear localization of the mechanoresponsive transcription factor YAP in the barrier-forming granular layer, YAP nuclear localization is not sufficient to drive premature differentiation when forced to the nucleus in the spinous layer.

Overall, this is a well-written manuscript and a comprehensive dataset. Only the RNA sequencing result should be presented more transparently providing the full lists of regulated genes instead of presenting just the GO analysis and selected target genes so that this analysis can serve as a useful repository. The authors themselves have profited from and used

published datasets of gene expression of the granular cells. Moreover, some of the previous data should be better discussed though. The authors state that forced suprabasal contractility in their mouse models induces the expression of some genes of the epidermal differentiation complex (EDC). However, in their previous publication, the authors showed that major classical EDC genes are actually not regulated like filaggrin and loricrin (Muroyama and Lechler eLife 2017). This should be discussed better and necessitates including the full list of regulated genes to show what exactly is regulated.

<https://doi.org/10.7554/eLife.100961.1.sa3>

Reviewer #2 (Public review):

Summary:

The manuscript from Prado-Mantilla and co-workers addresses mechanisms of embryonic epidermis development, focusing on the intermediate layer cells, a transient population of suprabasal cells that contributes to the expansion of the epidermis through proliferation. Using bulk-RNA they show that these cells are transcriptionally distinct from the suprabasal spinous cells and identify specific marker genes for these populations. They then use transgenesis to demonstrate that one of these selected spinous layer-specific markers, the transcription factor MafB is capable of suppressing proliferation in the intermediate layers, providing a potential explanation for the shift of suprabasal cells into a non-proliferative state during development. Further, lineage tracing experiments show that the intermediate cells become granular cells without a spinous layer intermediate. Finally, the authors show that the intermediate layer cells express higher levels of contractility-related genes than spinous layers and overexpression of cytoskeletal regulators accelerates the differentiation of spinous layer cells into granular cells.

Overall the manuscript presents a number of interesting observations on the developmental stage-specific identities of suprabasal cells and their differentiation trajectories and points to a potential role of contractility in promoting differentiation of suprabasal cells into granular cells. The precise mechanisms by which MafB suppresses proliferation, how the intermediate cells bypass the spinous layer stage to differentiate into granular cells, and how contractility feeds into these mechanisms remain open. Interestingly, while the mechanosensitive transcription factor YAP appears differentially active in the two states, it is shown to be downstream rather than upstream of the observed differences in mechanics.

Strengths:

The authors use a nice combination of RNA sequencing, imaging, lineage tracing, and transgenesis to address the suprabasal to granular layer transition. The imaging is convincing and the biological effects appear robust. The manuscript is clearly written and logical to follow.

Weaknesses:

While the data overall supports the authors' claims, there are a few minor weaknesses that pertain to the aspect of the role of contractility. The choice of spastin overexpression to modulate contractility is not ideal as spastin has multiple roles in regulating microtubule dynamics and membrane transport which could also be potential mechanisms explaining some of the phenotypes. Use of Arghap11 overexpression mitigates this effect to some extent but overall it would have been more convincing to manipulate myosin activity directly. It would also be important to show that these manipulations increase the levels of F-actin and myosin II as shown for the intermediate layer. It would also be logical to address if further

increasing contractility in the intermediate layer would enhance the differentiation of these cells.

The gene expression analyses are relatively superficial and rely heavily on GO term analyses which are of course informative but do not give the reader a good sense of what kind of genes and transcriptional programs are regulated. It would be useful to show volcano plots or heatmaps of actual gene expression changes as well as to perform additional analyses of for example gene set enrichment and/or transcription factor enrichment analyses to better describe the transcriptional programs

Claims of changes in cell division/proliferation changes are made exclusively by quantifying EdU incorporation. It would be useful to more directly look at mitosis. At minimum Y-axis labels should be changed from "% Dividing cells" to % EdU+ cells to more accurately represent findings

Despite these minor weaknesses the manuscript is overall of high quality, sheds new light on the fundamental mechanisms of epidermal stratification during embryogenesis, and will likely be of interest to the skin research community.

<https://doi.org/10.7554/eLife.100961.1.sa2>

Reviewer #3 (Public review):

Summary:

This is an interesting paper by Lechler and colleagues describing the transcriptomic signature and fate of intermediate cells (ICs), a transient and poorly defined embryonic cell type in the skin. ICs are the first suprabasal cells in the stratifying skin and unlike later-developing suprabasal cells, ICs continue to divide. Using bulk RNA seq to compare ICs to spinous and granular transcriptomes, the authors find that IC-specific gene signatures include hallmarks of granular cells, such as genes involved in lipid metabolism and skin barrier function that are not expressed in spinous cells. ICs were assumed to differentiate into spinous cells, but lineage tracing convincingly shows ICs differentiate directly into granular cells without passing through a spinous intermediate. Rather, basal cells give rise to the first spinous cells. They further show that transcripts associated with contractility are also shared signatures of ICs and granular cells, and overexpression of two contractility inducers (Spastin and ArhGEF-CA) can induce granular and repress spinous gene expression. This contractility-induced granular gene expression does not appear to be mediated by the mechanosensitive transcription factor, Yap. The paper also identifies new markers that distinguish IC and spinous layers and shows the spinous signature gene, MafB, is sufficient to repress proliferation when prematurely expressed in ICs.

Strengths:

Overall this is a well-executed study, and the data are clearly presented and the findings convincing. It provides an important contribution to the skin field by characterizing the features and fate of ICs, a much-understudied cell type, at high levels of spatial and transcriptomic detail. The conclusions challenge the assumption that ICs are spinous precursors through compelling lineage tracing data. The demonstration that differentiation can be induced by cell contractility is an intriguing finding and adds a growing list of examples where cell mechanics influence gene expression and differentiation.

Weaknesses:

A weakness of the study is an over-reliance on overexpression and sufficiency experiments to test the contributions of MafB, Yap, and contractility in differentiation. The inclusion of loss-

of-function approaches would enable one to determine if, for example, contractility is required for the transition of ICs to granular fate, and whether MafB is required for spinous fate. Second, whether the induction of contractility-associated genes is accompanied by measurable changes in the physical properties or mechanics of the IC and granular layers is not directly shown. The inclusion of physical measurements would bolster the conclusion that mechanics lies upstream of differentiation.

Finally, whether the expression of granular-associated genes in ICs provides them with some sort of barrier function in the embryo is not addressed, so the role of ICs in epidermal development remains unclear. Although not essential to support the conclusions of this study, insights into the function of this transient cell layer would strengthen the overall impact.

<https://doi.org/10.7554/eLife.100961.1.sa1>

Author response:

Public Reviews:

Reviewer #1 (Public review):

Summary:

The authors address a fundamental question for cell and tissue biology using the skin epidermis as a paradigm and ask how stratifying self-renewing epithelia induce diCere differentiation and upward migration in basal dividing progenitor cells to generate suprabasal barrier-forming cells that are essential for a functional barrier formed by such an epithelium. The authors show for the first time that an increase in intracellular actomyosin contractility, a hallmark of barrier-forming keratinocytes, is sufficient to trigger terminal diCere differentiation. Hence the data provide in vivo evidence of the more general interdependency of cell mechanics and diCere differentiation. The data appear to be of high quality and the evidences are strengthened through a combination of diCere genetic mouse models, RNA sequencing, and immunofluorescence analysis.

To generate and maintain the multilayered, barrier-forming epidermis, keratinocytes of the basal stem cell layer diCere differentiate and move suprabasally accompanied by stepwise changes not only in gene expression but also in cell morphology, mechanics, and cell position. Whether any of these changes is instructive for diCere differentiation itself and whether consecutive changes in diCere differentiation are required remains unclear. Also, there are few comprehensive data sets on the exact changes in gene expression between diCere states of keratinocyte diCere differentiation. In this study, through genetic fluorescence labeling of cell states at diCere developmental time points the authors were able to analyze gene expression of basal stem cells and suprabasal diCere differentiated cells at two diCere stages of maturation: E14 (embryonic day 14) when the epidermis comprises mostly two functional compartments (basal stem cells and suprabasal so-called intermediate cells) and E16 when the epidermis comprise three (living) compartments where the spinous layer separates basal stem cells from the barrier-forming granular layer, as is the case in adult epidermis. Using RNA bulk sequencing, the authors developed useful new markers for suprabasal stages of diCere differentiation like MafB and Cox1. The transcription factor MafB was then shown to inhibit suprabasal proliferation in a MafB transgenic model.

The data indicate that early in development at E14 the suprabasal intermediate cells resemble in terms of RNA expression, the barrier-forming granular layer at E16, suggesting that keratinocytes can undergo either stepwise (E16) or more direct (E14) terminal diCere differentiation.

Previous studies by several groups found an increased actomyosin contractility in the barrier-forming granular layer and showed that this increase in tension is important for epidermal barrier formation and function. However, it was not clear whether contractility itself serves as an instructive signal for diCere differentiation. To address this question, the authors use a previously published model to induce premature hypercontractility in the spinous layer by using spastin overexpression (K10-Spstin) to disrupt microtubules (MT) thereby indirectly inducing actomyosin contractility. A second model activates myosin contractility more directly through overexpression of a constitutively active RhoA GEF (K10Arhgef11CA). Both models induce late diCere differentiation of suprabasal keratinocytes regardless of the suprabasal position in either spinous or granular layer indicating that increased contractility is key to induce late diCere differentiation of granular cells. A potential weakness of the K10-spastin model is the disruption of MT as the primary eCect which secondarily causes hypercontractility. However, their previous publications provided some evidence that the eCect on diCere differentiation is driven by the increase in contractility (Ning et al. cell stem cell 2021). Moreover, the data are confirmed by the second model directly activating myosin through RhoA. These previous publications already indicated a role for contractility in diCere differentiation but were focused on early diCere differentiation. The data in this manuscript focus on the regulation of late diCere differentiation in barrier-forming cells. These important data help to unravel the interdependencies of cell position, mechanical state, and diCere differentiation in the epidermis, suggesting that an increase in cellular contractility in most apical positions within the epidermis can induce terminal diCere differentiation. Importantly the authors show that despite contractility-induced nuclear localization of the mechanoresponsive transcription factor YAP in the barrier-forming granular layer, YAP nuclear localization is not sufficient to drive premature diCere differentiation when forced to the nucleus in the spinous layer.

Overall, this is a well-written manuscript and a comprehensive dataset. Only the RNA sequencing result should be presented more transparently providing the full lists of regulated genes instead of presenting just the GO analysis and selected target genes so that this analysis can serve as a useful repository. The authors themselves have profited from and used published datasets of gene expression of the granular cells. Moreover, some of the previous data should be better discussed though. The authors state that forced suprabasal contractility in their mouse models induces the expression of some genes of the epidermal diCere differentiation complex (EDC). However, in their previous publication, the authors showed that major classical EDC genes are actually not regulated like filaggrin and loricrin (Muroyama and Lechler eLife 2017). This should be discussed better and necessitates including the full list of regulated genes to show what exactly is regulated.

We thank all the reviewers for their suggestions and comments.

Thank you especially for the reminder to include gene lists. We had an excel document with all this data but neglected to upload it with the initial manuscript decision. This includes all the gene signatures for the diCere cell compartments across development. We will also include a page that lists all EDC genes and whether they were up-regulated in intermediate cells and cells in which contractility was induced. Further, we note that all the RNA-Seq datasets are available for use on GEO.

In our previous publication, we indeed included images showing a lack of change in loricrin and filaggrin in the embryos where spastin was expressed in the diCere differentiated epidermis. Consistent with this, there is no change in Lor mRNA levels by RNA-Seq, (it is one of the rare EDC genes that is unchanged). In contrast, Flg mRNA was up in the RNASeq, though we didn't see a dramatic change in protein levels. We have not further pursued whether this reflects

translational regulation. That said, our data clearly show that other genes associated with granular fate were increased in the contractile skin.

Reviewer #2 (Public review):

Summary:

The manuscript from Prado-Mantilla and co-workers addresses mechanisms of embryonic epidermis development, focusing on the intermediate layer cells, a transient population of suprabasal cells that contributes to the expansion of the epidermis through proliferation. Using bulk-RNA they show that these cells are transcriptionally distinct from the suprabasal spinous cells and identify specific marker genes for these populations. They then use transgenesis to demonstrate that one of these selected spinous layer-specific markers, the transcription factor MafB is capable of suppressing proliferation in the intermediate layers, providing a potential explanation for the shift of suprabasal cells into a non-proliferative state during development. Further, lineage tracing experiments show that the intermediate cells become granular cells without a spinous layer intermediate. Finally, the authors show that the intermediate layer cells express higher levels of contractility-related genes than spinous layers and overexpression of cytoskeletal regulators accelerates the diCereentiation of spinous layer cells into granular cells.

Overall the manuscript presents a number of interesting observations on the developmental stage-specific identities of suprabasal cells and their diCereentiation trajectories and points to a potential role of contractility in promoting diCereentiation of suprabasal cells into granular cells. The precise mechanisms by which MafB suppresses proliferation, how the intermediate cells bypass the spinous layer stage to diCereentiate into granular cells, and how contractility feeds into these mechanisms remain open. Interestingly, while the mechanosensitive transcription factor YAP appears deferentially active in the two states, it is shown to be downstream rather than upstream of the observed diCereences in mechanics.

Strengths:

The authors use a nice combination of RNA sequencing, imaging, lineage tracing, and transgenesis to address the suprabasal to granular layer transition. The imaging is convincing and the biological eCects appear robust. The manuscript is clearly written and logical to follow.

Weaknesses:

While the data overall supports the authors' claims, there are a few minor weaknesses that pertain to the aspect of the role of contractility, The choice of spastin overexpression to modulate contractility is not ideal as spastin has multiple roles in regulating microtubule dynamics and membrane transport which could also be potential mechanisms explaining some of the phenotypes. Use of Arghap11 overexpression mitigates this eCect to some extent but overall it would have been more convincing to manipulate myosin activity directly. It would also be important to show that these manipulations increase the levels of F-actin and myosin II as shown for the intermediate layer. It would also be logical to address if further increasing contractility in the intermediate layer would enhance the diCereentiation of these cells.

We agree with the reviewer that the development of additional tools to precisely control myosin activity will be of great use to the field. That said, our series of publications has clearly demonstrated that ablating microtubules results in increased contractility and that this phenocopies the eCects of Arhgef11 induced contractility (Ning et al, Cell Stem Cell 2021).

Further, we showed that these phenotypes were rescued by myosin inhibition with blebbistatin. Our prior publications also showed a clear increase in junctional acto-myosin through expression of either spastin or Arhgef11, as well as increased staining for the tension sensitive epitope of alpha-catenin (alpha-18) (also in Ning et al, 2021). We are not aware of tools that allow direct manipulation of myosin activity that currently exist in mouse models.

The gene expression analyses are relatively superficial and rely heavily on GO term analyses which are of course informative but do not give the reader a good sense of what kind of genes and transcriptional programs are regulated. It would be useful to show volcano plots or heatmaps of actual gene expression changes as well as to perform additional analyses of for example gene set enrichment and/or transcription factor enrichment analyses to better describe the transcriptional programs

We will include an excel document that lists all the gene signatures. Additionally, all of our data are deposited in GEO for others to perform their own analyses.

Claims of changes in cell division/proliferation changes are made exclusively by quantifying EdU incorporation. It would be useful to more directly look at mitosis. At minimum Y-axis labels should be changed from "% Dividing cells" to % EdU+ cells to more accurately represent findings

We will change the axis label to precisely match our analysis.

Despite these minor weaknesses the manuscript is overall of high quality, sheds new light on the fundamental mechanisms of epidermal stratification during embryogenesis, and will likely be of interest to the skin research community.

Reviewer #3 (Public review):

Summary:

This is an interesting paper by Lechler and colleagues describing the transcriptomic signature and fate of intermediate cells (ICs), a transient and poorly defined embryonic cell type in the skin. ICs are the first suprabasal cells in the stratifying skin and unlike later-developing suprabasal cells, ICs continue to divide. Using bulk RNA seq to compare ICs to spinous and granular transcriptomes, the authors find that IC-specific gene signatures include hallmarks of granular cells, such as genes involved in lipid metabolism and skin barrier function that are not expressed in spinous cells. ICs were assumed to differentiate into spinous cells, but lineage tracing convincingly shows ICs differentiate directly into granular cells without passing through a spinous intermediate. Rather, basal cells give rise to the first spinous cells. They further show that transcripts associated with contractility are also shared signatures of ICs and granular cells, and overexpression of two contractility inducers (Spastin and ArhGEF-CA) can induce granular and repress spinous gene expression. This contractility-induced granular gene expression does not appear to be mediated by the mechanosensitive transcription factor, Yap. The paper also identifies new markers that distinguish IC and spinous layers and shows the spinous signature gene, MafB, is sufficient to repress proliferation when prematurely expressed in ICs.

Strengths:

Overall this is a well-executed study, and the data are clearly presented and the findings convincing. It provides an important contribution to the skin field by characterizing the features and fate of ICs, a much-understudied cell type, at high levels of spatial and transcriptomic detail. The conclusions challenge the assumption that ICs are spinous precursors through compelling lineage tracing data. The demonstration that

diCerentiation can be induced by cell contractility is an intriguing finding and adds a growing list of examples where cell mechanics influence gene expression and diCerentiation.

Weaknesses:

A weakness of the study is an over-reliance on overexpression and suCiciency experiments to test the contributions of MafB, Yap, and contractility in diCerentiation. The inclusion of loss-of-function approaches would enable one to determine if, for example, contractility is required for the transition of ICs to granular fate, and whether MafB is required for spinous fate. Second, whether the induction of contractility-associated genes is accompanied by measurable changes in the physical properties or mechanics of the IC and granular layers is not directly shown. The inclusion of physical measurements would bolster the conclusion that mechanics lies upstream of diCerentiation.

We agree that loss of function studies would be useful. For MafB, these have been performed in cultured human keratinocytes, where loss of MafB and its ortholog cMaf results in a phenotype consistent with loss of spinous diCerentiation (Lopes-Pajares, Dev Cell 2015). Due to the complex genetics involved, generating these double mutant mice is beyond the scope of this study. Loss of function studies of myosin are also complicated by genetic redundancy of the non-muscle type II myosin genes, as well as the role for these myosins in actin cross linking in addition to contractility. In addition, we have found that these myosins are quite stable in the embryonic intestine, with loss of protein delayed by several days from the induction of recombination. Therefore, elimination of myosins by embryonic day e14.5 with our current drivers is not likely possible. Thus, generation of inducible inhibitors of contractility is a valuable future goal.

A number of recent papers have used AFM of skin sections to probe tissue rigidity. We have not attempted these studies and are unclear about the spatial resolution and whether, in the very thin epidermis at these stages we could spatially resolve diCerenes. That said, we previously assessed the macro-contractility of tissues in which myosin activity was induced and demonstrated that there was a significant increase in this over a tissue-wide scale (Ning et al, Cell Stem Cell, 2021).

Finally, whether the expression of granular-associated genes in ICs provides them with some sort of barrier function in the embryo is not addressed, so the role of ICs in epidermal development remains unclear. Although not essential to support the conclusions of this study, insights into the function of this transient cell layer would strengthen the overall impact.

By traditional dye penetration assays, there is no epidermal barrier at the time that intermediate cells exist. One interpretation of the data is that cells are beginning to express mRNAs (and in some cases, proteins) so that they are able to rapidly generate a barrier as they become granular cells. We have attempted experiments to ablate intermediate cells with DTA expression - this resulted in ineCicient and delayed cell death and thus did not yield strong conclusions. Our findings that transcriptional regulators of granular diCerentiation (such as Grhl3 and Hopx) are also present in intermediate cells, should allow future analysis of the eCects of their ablation on the earliest stages of granular diCerentiation from intermediate cells.

<https://doi.org/10.7554/eLife.100961.1.sa0>