

Axonal injury signaling is restrained by a spared synaptic branch

Reviewed Preprint

v1 • December 27, 2024

Not revised

Laura J Smithson, Juliana Zang, Lucas Junginger, Thomas J Waller, Lauren Reilly-Jankowiak, Sophia Khan, Ye Li, Dawen Cai, Catherine A Collins ✉

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, USA •

Department of Neurosciences, Case Western Reserve University, Cleveland, USA • Department of Cell and

Developmental Biology, University of Michigan, Ann Arbor, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study leverages the power of *Drosophila* genetics and sparsely-labeled neurons to propose a new model for neuronal injury signaling. The authors present **convincing** evidence to support that the somatic response to axonal injury is suppressed if the injury is not complete, suggesting the presence of an integration of axonal injury-related signaling. While the underlying mechanism of this fascinating observation is unknown, the phenomenon itself will be of broad significance in the field.

<https://doi.org/10.7554/eLife.104896.1.sa4>

Abstract

The intrinsic ability of injured neurons to degenerate and regenerate their axons facilitates nervous system repair, however this ability is not engaged in all neurons and injury locations. Here we investigate the regulation of a conserved axonal injury response pathway with respect to the location of damage in branched motoneuron axons in *Drosophila* larvae. The dileucine zipper kinase DLK, (also known as MAP3K12 in mammals and Wallenda (Wnd) in *Drosophila*), is a key regulator of diverse responses to axonal injury. In three different populations of motoneurons, we observed the same striking result that Wnd/DLK signaling becomes activated only in response to injuries that remove all synaptic terminals. Injuries that spare even a small part of a synaptic terminal fail to activate Wnd/DLK signaling, despite the presence of extensive axonal degeneration. The regulation of injury-induced Wnd/DLK signaling occurs independently of its previously known regulator, the Hiw/PHR ubiquitin ligase. We propose that Wnd/DLK signaling regulation is linked to the trafficking of a synapse-to-nucleus axonal cargo and that this mechanism enables neurons to respond to impairments in synaptic connectivity.

Introduction

Repair of nervous system damage requires an ability of neurons to regenerate axons and synaptic connections. However this ability is not universally induced following injury, most notably following spinal cord injury (SCI). While many studies have identified extrinsic factors that influence or inhibit axonal regeneration, several studies have noted that the location of damage can influence the intrinsic ability of the neuron to mount a transcriptional response to the damage (Fernandes et al., 1999 [↗](#); Lorenzana et al., 2015 [↗](#); Mason et al., 2003 [↗](#); Wang et al., 2024 [↗](#)). Some studies have noted an effect of distance from the cell body for long axons in the spinal cord (Fernandes et al., 1999 [↗](#); Mason et al., 2003 [↗](#); Wang et al., 2024 [↗](#)). Other studies have noted that the location of injury with respect to an axonal branching point also strongly influences the response (Lorenzana et al., 2015 [↗](#); Wu et al., 2007 [↗](#)). Even in a strongly inhibitory environment to regeneration, dorsal column sensory axons show robust axonal growth when injured proximal to their bifurcation in the spinal cord (Lorenzana et al., 2015 [↗](#)).

Here we investigate the regulation of a conserved axonal injury response pathway with respect to the location of axonal injury. The dileucine zipper kinase DLK, (also known as MAP3K12 in mammals and Wallenda (Wnd) in *Drosophila*), is a key regulator of diverse responses to axonal injury. These include an essential role in the ability of damaged neurons to initiate axonal regeneration in worm and fly models (Chen et al., 2011 [↗](#); Hammarlund et al., 2009 [↗](#); Stone et al., 2014 [↗](#); X. Xiong et al., 2010 [↗](#); Yan et al., 2009 [↗](#)), synaptic repair and recovery following CCR5 inhibition in a stroke model (Joy et al., 2019 [↗](#)), and enhanced regeneration and mechanical allodynia following PNS nerve damage in mice (Hu et al., 2019 [↗](#); Wlaschin et al., 2018 [↗](#)).

Dichotomously, DLK is also required for the death of retinal ganglion cells following optic nerve damage (Watkins et al., 2013 [↗](#); Welsbie et al., 2017 [↗](#), 2013 [↗](#)). In mammalian as well as in fly neurons, this kinase associates with vesicles that are physically transported in axons (Holland et al., 2016 [↗](#); Xiong et al., 2010 [↗](#)), while downstream nuclear signaling requires functional axonal transport machinery (Xiong et al., 2010 [↗](#)). DLK is therefore considered to function as a ‘sensor’ of axonal damage, whose activation can confer responses of repair or death, depending upon the cellular context (Asghari Adib et al., 2018 [↗](#)).

While the responses gated by DLK are impactful for neurons and their circuits, the mechanism(s) that lead to DLK signaling activation are still poorly understood. A number of observations have documented DLK signaling activation in neurons that are not mechanically damaged but have experienced some form of cellular stress. These include the presence of cytoskeletal mutations (Bounoutas et al., 2011 [↗](#); Chen et al., 2014 [↗](#); Kurup et al., 2015 [↗](#); Valakh et al., 2013 [↗](#)) and the presence of chemotherapy agents (Bhattacharya et al., 2012 [↗](#); DeVault et al., 2024 [↗](#); Valakh et al., 2015 [↗](#)) known to impair axonal cytoskeleton integrity and transport. DLK activation is also responsible for phenotypes associated with mutations in the unc-104/KIF1A kinesin (Li et al., 2017 [↗](#)), a major carrier of synaptic vesicle precursors in axons (Guedes-Dias and Holzbaur, 2019 [↗](#); Petzoldt, 2023 [↗](#)). Inhibition of DLK is protective in mouse models of ALS and Alzheimer’s Disease (Le Pichon et al., 2017 [↗](#); Patel et al., 2017 [↗](#)). These observations have fostered growing interest in DLK as a potential therapeutic target, and in understanding the mechanisms that control DLK signaling activation in neurons.

Here we test the hypothesis that DLK/Wnd signaling is tuned to the synaptic connectivity of a neuron. A shared feature of nerve injuries and stressors that disrupt axonal cytoskeleton and transport is a loss in downstream connections. A conserved regulator of DLK/Wnd, the E3-ubiquitin ligase PAM/Highwire/Rpm-1/Phr1 (Collins et al., 2006 [↗](#); Huntwork-Rodriguez et al., 2013 [↗](#); Nakata et al., 2005 [↗](#)), is hypothesized to function at synaptic terminals (Abrams et al., 2008 [↗](#); Opperman and Grill, 2014 [↗](#); Xiong et al., 2012 [↗](#); Zhen et al., 2000 [↗](#)). This led us to ask

whether interactions at an intact synaptic terminal are responsible for restraining Wnd signaling in uninjured neurons. We probed this hypothesis through injuries to branched motoneuron axons in *Drosophila* larvae, which allowed us to compare injuries that leave spared synaptic terminals to injuries that lead to complete denervation. In three different populations of motoneurons, we observed the same striking result that Wnd signaling becomes activated only in response to injuries that remove all synaptic terminals. Injuries that spared even a small part of a synaptic terminal did not activate Wnd signaling, despite the presence of extensive axonal degeneration. Surprisingly, removal of all synapses led to additive induction of Wnd signaling in *hiw* mutants.

These observations suggest the existence of a mechanism that restrains Wnd signaling at synaptic terminals independently of the Hiw ubiquitin ligase.

Results

The presence of a spared synaptic branch restrains Wnd-mediated injury signaling in SNc motoneurons

To determine whether synaptic connections influence injury signaling by Wnd/Dlk, we established methods to injure single synaptic branches of defined larval motoneurons. The m12 (5053A)-Gal4 driver line (Ritzenthaler et al., 2000) that we have used in previous nerve injury studies (Xiong et al., 2010; Xiong and Collins, 2012), drives expression in two single motoneurons that project closely fasciculated axons to innervate body wall muscles 26, 27 and 29 (Figure 1A). This pattern was previously attributed to a motoneuron (MN) named MNSNc, which was noted to have ‘two cell bodies’ (Kim et al., 2009). We used the Bitbow2 (Li et al., 2021) multi-colored cell labeling approach to resolve the two neurons, and observed an invariable pattern that one MNSNc neuron innervates muscles 26 and 29, while the other innervates muscle 27. (Figure 1B shows two examples: the neuron that innervates muscle 27 (middle images) expresses a set of colors that distinguish it from the terminals on muscles 26 and 29). We therefore refer to these two neurons labeled by the m12Gal4 driver as MNSNc-26/29 and MNSNc-27 (Figures 1A, B, Supplemental Figures 1A). MNSNc-26/29 (cartooned in blue in Figure 1A) has three collateral branches: two branches innervate muscle 26, and a single branch innervates muscle 29. MNSNc-27 (cartooned in red in Figure 1A) has two collateral branches that innervate muscle 27: these two branches most often remain together and occasionally bifurcate separately onto muscle 27 (Supplemental Figure 1A). This innervation is stereotyped across segments and animals. As previously noted (Kim et al., 2009), the paired cell bodies of the MNSNc neurons are found on the lateral sides of the abdominal region of the ventral nerve cord (VNC). Bitbow2 expression revealed that the MNSNc-27 cell somas are positioned more ventrally compared to the cell somas of MNSNc-26/29 in the VNC (Supplemental Figure 1B).

We used a pulsed dye laser to carry out axotomies of MNSNc motoneuron axons at different locations in intact immobilized larvae (described in methods and (Smithson et al., 2024a)). Successful injuries were confirmed by the degeneration of distal stumps within 24h post-injury. To assess the activation of Wnd signaling, we probed for the induction of *puckered* expression from the *puc-lacZ* enhancer trap (Martín-Blanco et al., 1998). Previous studies from this reporter line have shown that expression of nuclear localized lacZ is strongly induced following nerve injury, requiring Wnd, JNK and the Fos transcription factor (Xiong et al., 2010). We confirmed this is the case for MNSNc neurons; axotomy of the m12-Gal4 labeled neurons upstream of the synaptic branches led to a four fold induction of *puc-lacZ* expression in either neuron 24h after the injury; this was abolished in neurons that co-express double stranded RNAi targeting *wnd* (Figure 1C). Similar results were observed for nerve crush injuries (Supplemental Figure 1C).




Figure 1

A spared synaptic branch restrains Wnd-dependent injury signaling in SNc motoneurons.

A) Schematic representation of the two SNc motoneurons innervating muscles 26 and 29 (MNSNc-26/29, red) and muscle 27 (MNSNc-27, blue), which are labeled by expression of the m12-Gal4 driver.

B) Example images of NMJ terminals from m12-Gal4/+; UAS-BitBow2 (Li et al., 2021) /+ third instar larvae, used to define the connectivity shown in A. The neuron that innervates muscle 27 (MNSNc-27) expresses a distinct set of colors from the Bitbow2 (Li et al., 2021) reporter than the neuron that innervates muscles 26 and 29 (MNSNc-26/29).

C) Example NMJs (top row) and cell bodies (bottom row) of MNSNc neurons (from m12-Gal4, UAS-mCD8GFP/*puc-lacZ* larvae) following injury at a location upstream of all of the synaptic branches, indicated by the arrow. The left column shows uninjured wild type animals, the middle column shows wild type animals 24h post injury. The far right column is from an animal that co-expresses UAS-*wnd*-RNAi. LacZ expression (red channel) in MNSNc nuclei indicates the induction of Wnd-dependent signaling in the injured neurons.

D) Example MNSNc neurons injured at different locations. The top row shows *lucida* drawings of the NMJ terminals in the middle row, while the bottom row shows *puc-lacZ* expression (red), in GFP(gray)-expressing cell bodies (from m12-Gal4, UAS-mCD8GFP/*puc-lacZ* larvae).

E) Quantification of *puc-lacZ* intensity measurements in MNSNc-26/29 (blue) and MNSNc-27 (red) following injuries that remove all synaptic branches versus injuries that leave a spared synaptic branch. A one-way ANOVA with Tukey test for multiple comparisons was performed for each neuron. **** $p < 0.0001$; ** $p = 0.0011$; comparisons not denoted were not significant.

In contrast to axotomies that removed all synaptic branches, laser injury to single collateral branches of MNSNc-26/29, including branches innervating either muscle 26 or 29 and resulting in spared synaptic branches, fail to induce *puc-lacZ* expression (**Figures 1D, E**). In addition, laser ablation of the anterior branch of MNSNc-27 also failed to activate Wnd signaling (**Figures 1D, E**). We carried out similar experiments in aCC motoneurons, which can be labeled with the Dpr4- Gal4 driver) (Pérez-Moreno and O’Kane, 2019). Laser axotomies that removed all of the synaptic branches resulted in four-fold increase in *puc-LacZ* levels, while injuries that left spared synaptic boutons failed to induce *puc-lacZ* expression. (**Supplemental Figure 1D,E**).

Despite the small distance from the disconnected muscle, none of the injured MNSNc synaptic branches were able to re-innervate the muscle. We think this is due to an absence of axon growth-promoting cues, since MNSNc axons did show robust but misdirected axonal growth into the segmental nerve SNa following injuries in locations upstream of the synaptic branches (data not shown). However we did notice differences in the trafficking of proteins to injured proximal stumps. An example of this is shown for ectopically expressed kinase-dead Wnd transgenic protein, GFP- Wnd-KD in **Supplemental Figure 1F, G**. (We were only able to track kinase-inactive Wnd since overexpression of Wnd causes dramatic morphological defects to neurons (Collins et al., 2006; Feoktistov and Herman, 2016; Xiong et al., 2010)). GFP-Wnd-KD was strongly induced and accumulated at the proximal stump following injuries that removed all synaptic branches, but was barely detectable following injuries that left spared synaptic branches (**Supplemental Figure 1F, G**). Collectively, these observations suggest that the presence of spared synaptic branches affect the subsequent events that occur in the injured axon. These include the stability and/or trafficking of Wnd protein in injured axons and the activation of Wnd-regulated signaling in the neuron soma.

Restraint of Wnd-dependent injury signaling in bifurcated axons of type II VUM motoneurons

A more extreme example of axonal branching is illustrated by the ventral unpaired median (VUM) neurons, which project symmetric bifurcated axons through separate nerves to innervate multiple body wall muscles on both left and right sides of the larva (**Figure 2A**, **Supplemental Figure 2** and (Koon et al., 2011; Monastirioti et al., 1995; Vömel and Wegener, 2008)). VUM neurons can be specifically labeled based on their expression of tyrosine decarboxylase 2 using the Tdc2-Gal4 driver. Each abdominal segment has 3 VUM neurons, each of which sends a single axon dorsally which then bifurcates in the midline VNC (**Supplemental Figure 2**, and (Vömel and Wegener, 2008)). The two bifurcations then project through separate nerves to symmetrically innervate both left and right halves of the larval body. Through nerve crush injuries to only one ventral side of the animal, we were able to injure one bifurcation while leaving the other bifurcation intact. Successful injuries were determined based on the degeneration of the distal axon and synaptic terminals at 24 hours post-crush (**Figure 2B**). Whether both bifurcations, a single bifurcation or neither bifurcation was injured was scored for each VUM neuron by tracing the Tdc2-Gal4, UAS- mCD8GFP labeled axons from the nerve to the cell body.

Similarly to other motoneurons(Xiong et al., 2010), *puc-lacZ* expression is barely detectable in uninjured VUM motoneurons and, compared to uninjured VUM neurons, is induced almost 3-fold at 24h following complete/full nerve crush injuries that damage both bifurcations and remove all synaptic connections (**Figure 2C,D**). This induction is abolished in VUM neurons that co-express *wnd-RNAi* (**Figure 2C,D**). Note that *wnd-RNAi* is only expressed in the VUM neurons and does not affect the other MNs which do not express Tdc2-Gal4. In contrast to full nerve crushes, injuries to nerves on a single side of the animal that the other bifurcation intact (‘half crush’ injuries) failed to induce *puc-lacZ* expression in VUM neurons (**Figure 2C,D**). Most MNs make ipsilateral and not bilateral projections; in ‘half-crush’ injuries, *puc-lacZ* is induced in most non-VUM motoneurons on the side of the crush, but is not induced in VUMs. These combined observations




Figure 2

Restraint of Wnd-mediated injury signaling by spared branch in bifurcated neurons.

A) Cartoon of Ventral Unpaired Motoneurons (VUM), which have bifurcated axons that symmetrically innervate body wall muscles on both the left and right sides of the animal. Nerve crush to either left or right side of the animal can axotomize a single bifurcation while leaving the other bifurcated axon intact.

B) Example images of VUM axons (visualized in Tdc2-Gal4, UAS-mCD8-GFP larvae) in segmental nerves on the uninjured and injured sides following nerve crush to a single side.

C) Example images of *puc-lacZ* expression in the VNC (ventral nerve cord) of larvae following nerve crush to a single side (half crush) versus crush to all the segmental nerves (full crush). *puc-lacZ* expression (red) is induced in VUM neurons (white) only after full crush. In contrast, other motoneurons, which innervate a single side, are induced by both half and full crush injuries. Co-expression of UAS-*wnd*-RNAi in VUM neurons cell autonomously inhibits *puc-lacZ* induction.

D) Quantification of *puc-lacZ* intensity measurements in VUM neurons. A one-way ANOVA with Tukey test for multiple comparisons was performed. **** $p < 0.0001$. Scale bars = 20 μ m.

suggest that in *Drosophila* motoneurons, Wnd signaling is not tuned to detect axonal damage *per se*, but is instead uniquely tuned to detect a complete loss of innervation, which occurs following some injuries but not others.

Restraint of Wnd signaling at spared branches does not require synaptic transmission

Since the presence of intact synaptic boutons restrains Wnd signaling activation, we considered whether cellular events associated with evoked or spontaneous synaptic transmission are associated with this mechanism. Summarized in **Supplementary Table 1** [\[link\]](#), we tested a total of 22 genetic manipulations expected to inhibit synaptic transmission, but none led to a change in *puc-lacZ* expression. These include electrical silencing of SNC motoneurons by Gal4/UAS mediated expression of the *Drosophila* open rectifier K⁺ channel (dORK) (Nitabach et al., 2002 [\[link\]](#)), silencing of transmission using temperature-sensitive mutations in dynamin (Kitamoto, 2001 [\[link\]](#)), and light-induced silencing of neurons expressing *Guillardia theta* anion channelrhodopsin 1 (gtACR1) (Mohammad et al., 2017 [\[link\]](#)) (**Supplementary Table 1** [\[link\]](#)). Consistent with these negative results, we noted that previous studies have described many genetic manipulations that perturb evoked and/or spontaneous synaptic transmission (Choi et al., 2014 [\[link\]](#); Daniels et al., 2006 [\[link\]](#); DiAntonio and Schwarz, 1994 [\[link\]](#); Han et al., 2022 [\[link\]](#)) do not yield synaptic phenotypes (of synaptic overgrowth or decreased VGlut expression levels) associated with Wnd activation (Collins et al., 2006 [\[link\]](#); Li et al., 2017 [\[link\]](#)).

Restraint of Wnd-mediated axon injury signaling is independent of the Highwire ubiquitin ligase

We then asked whether a known upstream regulator of Wnd signaling, the Pam/Hiw/Rpm-1 (PHR) ubiquitin ligase, functions to restrain Wnd at synaptic branches. PHR is a large protein with multiple evolutionarily conserved domains, including a RING-finger domain, which regulates DLK/Wnd in invertebrate (*C. elegans* and *Drosophila*) (Collins et al., 2006 [\[link\]](#); Nakata et al., 2005 [\[link\]](#)) and vertebrate (Huntwork-Rodriguez et al., 2013 [\[link\]](#)) model organisms. PHR is an attractive candidate since it localizes to synaptic terminals (Abrams et al., 2008 [\[link\]](#); Opperman and Grill, 2014 [\[link\]](#); Zhen et al., 2000 [\[link\]](#)) and loss of PHR function leads to increased levels of Wnd/DLK at synapses (Collins et al., 2006 [\[link\]](#); Nakata et al., 2005 [\[link\]](#)). This hypothesis predicts that removal of all synaptic branches would be equivalent to a genetic loss in PHR function. We tested this in null mutants for the *Drosophila* ortholog of PHR, Highwire (Hiw). The *hiw*^{ΔN} mutation deletes the entire N-terminal half of the *hiw* gene and abolishes expression of the Hiw protein (Wu et al., 2005 [\[link\]](#)). Since *hiw*^{ΔN} animals are viable, we were able to carry out injury assays in neurons that completely lack Hiw function.

Consistent with previously reported phenotypes for *hiw* in other motoneuron types (Collins et al., 2006 [\[link\]](#); Wan et al., 2000 [\[link\]](#); Wu et al., 2005 [\[link\]](#)), uninjured MNSNc neurons in male *hiw*^{ΔN} mutants have an increased number of axon collateral and terminal branches at muscles 26, 29 and 27 NMJs (**Figure 3A** [\[link\]](#) and **Supplemental Figure 3A** [\[link\]](#)). Also consistent with previous observations, uninjured neurons show elevated expression of *puc-lacZ* in *hiw*^{ΔN} mutants compared to control animals (**Figure 3C** [\[link\]](#) and **Supplemental Figure 3A** [\[link\]](#)). Strikingly, injuries that removed all synaptic terminals led to an even further elevation of *puc-lacZ* expression in *hiw*^{ΔN} mutant neurons. This was the case for laser axotomies that removed all synaptic branches from either MNSNc-26/29 and/or MNSNc-27 neurons (**Figure 3A** [\[link\]](#), right column, and **Figure 3B** [\[link\]](#)), and also for VUM neurons following nerve crush injuries (**Figure 3C** [\[link\]](#)). Injuries to a single synaptic branch (on muscle 29) of MNSNc-26/29 in *hiw*^{ΔN} mutants had a similar level of *puc-lacZ* expression as uninjured *hiw*^{ΔN} neurons (**Figures 3A, B** [\[link\]](#)). Collectively, these observations suggest that the presence of a spared synapse is capable of restraining Wnd signaling independently of Hiw's function.




Figure 3

Presence of spared synaptic branch restrains Wnd signaling independently of Hiw.

A) Laser axotomy is carried out to MNSNc neurons at a location that completely removes the synaptic terminal of MNSNc-27 (red neuron). The injury also leads to loss of the MNSNc-26/29 (blue) terminal on muscle 29 but not 26 (hence leaves a spared synaptic branch). The final column shows an axotomy that fully removes the terminals for both MNSNc neurons. These injuries were repeated in control animals versus the background of a *hiw* null mutant, *hiw^{ΔN}*.

B) Quantification of *puc-lacZ* expression for individual MNSNc neurons after full versus spared axotomies, compared to uninjured neurons. Basal *puc-lacZ* expression is already elevated in uninjured *hiw^{ΔN}* neurons compared to control. This can be further elevated in axotomies that remove all synapses, but not in axotomies that leave spared branches.

C) Quantification of *puc-lacZ* in VUM neurons (labeled by Tdc-2-Gal4; UAS-mCD8-GFP) 24h and 48h following full nerve crush in control versus *hiw^{ΔN}* mutants. A two-way ANOVA with Tukey test for multiple comparisons was performed. **** $p < 0.0001$; *** $p < 0.001$; ** $p < 0.01$.

Discussion

Axonal branches and spared synaptic connections influence the ability of injured axons to regenerate

Since the foundational observations of Ramon y Cajal, we have known that the ability of axons to regrow following damage varies strongly according to the location of damage ([Ramón y Cajal, 1928](#)). Most widely considered are differences in extrinsic factors that influence the ability of axons to grow following PNS injuries, and the impediments to axonal regeneration in the CNS that inhibit repair following spinal cord injuries ([Huebner and Strittmatter, 2009](#)). A less well studied but important intrinsic determinant of regeneration ability is the location of the injury with respect to axonal branches. For *C. elegans* PLM mechanosensory neurons, robust regeneration occurs following injuries that remove both the synaptic and sensory branches, but not following injuries that leave a synaptic branch intact ([Wu et al., 2007](#)). In the mammalian spinal cord, an elegant study following responses to laser-induced microsurgery to ascending and descending central projections of sensory neurons observed that remarkable regeneration occurred following injuries proximal to the branching point (which led to removal of all branches) but not distal (which left one branch intact) ([Lorenzana et al., 2015](#)). Studies in mice have also implicated synaptic proteins alpha2-delta, Munc13 and RIM in restraining regenerative ability of axons ([Hilton et al., 2022](#); [Tedeschi et al., 2016](#)). These observations are consistent with the possibility that spared afferent synaptic connections that remain following injuries distal to the branch point inhibit the regeneration ability of centrally projecting axons in the spinal cord.

It is noteworthy that many of the axons that project over great distances in the spinal cord have at least one synaptic branch. Neurons that project through the cortical spinal tract (CST), whose poor regeneration ability following spinal cord injury is most widely studied, form synaptic branches in the red nucleus, brainstem, and throughout the spinal cord ([Raineteau and Schwab, 2001](#)). A recent study has profiled the responses of CST neurons at different injury locations ([Wang et al., 2024](#)). Strikingly, regeneration associated genes (RAGs) and phosphorylated cJun, a marker of DLK signaling activation, are not induced in CST neurons following spinal cord injury, but are robustly induced following intracortical injuries close to CST cell bodies ([Mason et al., 2003](#); [Wang et al., 2024](#)). Since the latter may be the only form of injury that removes all synaptic branches from the CST neurons, we propose that restraint of DLK signaling activation by spared synaptic branches could be a prominent feature of the poor intrinsic regeneration capacity of neurons following spinal cord injury.

Restraint of Wnd/DLK signaling at synaptic terminals

Using genetic manipulations that inhibit or perturb synaptic transmission and/or neuronal excitability, we did not detect a requirement for synaptic transmission in the restraint of Wnd by spared synaptic branches. The PHR ubiquitin ligase, known as Hiw in *Drosophila*, was a logical candidate to regulate Wnd at synapses, since studies in multiple model organisms have shown that loss of this enzyme leads to increased levels of Wnd/DLK in axons ([Collins et al., 2006](#); [Huntwork-Rodriguez et al., 2013](#); [Nakata et al., 2005](#)). The *puc-lacZ* reporter implies that Wnd signaling is elevated in *hiw* mutants. However the restraint conferred by spared synaptic branches is still active in the absence of Hiw, since Wnd signaling can be further elevated by axotomy of all synapses in *hiw* null mutants (**Figure 3**). In a previous study we noted that the nerve crush injury led to a rapid down-regulation of an ectopically expressed Hiw-GFP transgene, and speculated that impairment of restraint by Hiw leads to activation of Wnd signaling in injured axons ([Xiong et al., 2010](#)). Our current data do not rule out a role for Hiw, but suggest the existence of additional mechanisms that restrain Wnd signaling at intact synapses. This has also

been suggested from developmental studies of photoreceptor growth cone termination, in which Hiw-independent downregulation of Wnd protein occurs concomitantly with the development of presynaptic boutons (Feoktistov and Herman, 2016).

We speculate that the regulation of Wnd is linked to the trafficking of organelles between the synaptic terminal and cell body, akin to neurotrophin signaling, which relies on retrograde trafficking of signaling endosomes in axons (Cosker et al., 2008). Consistent with this idea, DLK signaling becomes activated following nerve growth factor withdrawal from distal axons (Ghosh et al., 2011; Larhammar et al., 2017). Previous studies of Wnd signaling have documented its dependence on retrograde axonal transport machinery (Xiong et al., 2010). Moreover, mutations that disrupt axonal cytoskeleton and the unc-104/kif1A kinesin also lead to Wnd/DLK signaling activation (Li et al., 2017; Valakh et al., 2015, 2013). Both Wnd and its homologue DLK in mice show regulated association with organelle membranes via palmitoylation, and disruption of its palmitoylation abolishes DLK's signaling ability (Holland et al., 2016; Kim et al., 2024; Niu et al., 2022). Palmitoylation and depalmitoylation are dynamically regulated in axons (Ramzan et al., 2023; Zhang et al., 2024), hence comprise an attractive mechanism for mediating restraint of DLK at synaptic branches. Future delineation of the organelle(s) that Wnd/DLK associates with may provide important clues to its mechanism of regulation.

Our observations that Wnd signaling could be restrained by an intact axonal bifurcation suggests that at least one level of regulation could occur in the cell body. Consistent with this idea, a recent study has shown that Wnd signaling can be ectopically activated in the cell body when its transport to distal synapses is impaired in *rab11* mutants (Kim et al., 2024). Given the many factors that have been documented thus far that regulate DLK/Wnd protein or signaling (Asghari Adib et al., 2018), we think that this kinase must be tightly regulated both at synapses and cell bodies (Figure 4). Regulation at both locations gives the neuron a way to monitor the state of its entire axon and restrict signaling activation to scenarios where all efferent connections of the axon are disrupted.

Materials and methods

Drosophila stocks/Genetics

All fly crosses were raised at 25°C in a 12h light/dark cycle on standard sucrose and yeast food. Both male and female larvae were used unless otherwise stated. The following strains were used in this study: BL labeled flies were obtained from Bloomington Drosophila Stock Center (Indiana University), v labeled flies were obtained from the Vienna Drosophila Resource Center (VDRC, Vienna Biocenter Core Facilities). UAS-mCD8- GFP (Lee and Luo, 1999) (RRID:BDSC_5137), m12-gal4 (P(Gal4)^{5053A}) (Ritzenthaler et al., 2000) (RRID:BDSC_2702), BG380-gal4 (C80-Gal4) (Budnik et al., 1996; Sanyal, 2009), puc-lacZ^{E69} (Martín-Blanco et al., 1998), *hiw*^{ΔN} (Wu et al., 2005) (RRID:BDSC_51637), UAS-Bitbow2 (Li et al., 2021), Tdc2-Gal4 (RRID:BDSC_9313), UAS-*wnd* RNAi (RRID:BDSC_35369), UAS-LexA RNAi (RRID:BDSC_67947). Additional stocks tested are listed in Supplementary Table 1.

Immunohistochemistry

Wandering second and third instar larvae were dissected in ice-cold 1x PBS, fixed in 4% paraformaldehyde (16% diluted in 1X PBS, Electron microscopy Labs) for 20 mins at room temperature and washed thrice in 1X PBS. Tissues were blocked for a minimum 30 mins in 5% normal goat serum (NGS) in 0.25% triton X 100 in 1X PBS (PBST). Primary antibodies were incubated overnight in 5% NGS in 0.25% PBST at room temperature unless otherwise stated. The following antibodies and concentrations were used: mouse anti-lacZ (1:100, DSHB Cat# 40-1a, RRID:AB_528100), rabbit anti-DsRed (1:1000, Takara Bio Cat# 632496, RRID:AB_10013483), A488 rabbit anti-GFP (1:1000, for 2 hours at room temperature, Molecular Probes Cat# A-21311,




Figure 4

Potential mechanisms for regulation of Wnd signaling from synaptic terminals.

In green, Wnd signaling may be regulated in the cell body downstream of a retrogradely transported signal. (For example, neurotrophin signaling). In blue, Wnd signaling activation is restrained locally at synaptic terminals, perhaps by regulating the levels or activation of Wnd itself. Activated Wnd or a downstream signaling factor is then retrogradely transported to the cell body. Previous observations that inhibition of retrograde transport blocks the induction of Wnd signaling following axonal injury favors the latter (blue) possibility. However the restraint conferred by a physically separate bifurcation suggests that an inhibitor of Wnd signaling activation can be retrogradely transported (green). We speculate that both mechanisms act as dual checkpoints to restrain Wnd signaling activation in the context of healthy circuits.

RRID:AB_221477). Tissues were washed thrice in 1X PBS and incubated at room temperature for 2 hours in the appropriate secondary antibodies (1:1000 in 5% NGS in 0.25% PBST): AlexFluor 568 goat anti-mouse (ThermoFisher, A11004, RRID:AB_2534072), AlexFluor 488 goat anti-mouse, (ThermoFisher A32723, RRID:AB_2633275), AlexFluor 568 goat anti-rabbit, ThermoFisher A-11011, RRID:AB_143157). Tissues were washed thrice in 1x PBS and mounted onto superfrost plus slides (Fisher Scientific) and coverslipped with ProLong Diamond Antifade media (ThermoFisher Scientific, P36970). All experimental and control larval groups were processed (dissected, fixed, stained and images captured) together using identical confocal settings.

Nerve crush assays

Peripheral nerve crushes in wandering 2nd instar larvae were performed as previously described (Waller et al., 2024 [↗](#); Xiong et al., 2010 [↗](#)). Briefly, with the ventral side facing upwards, anesthetized larvae (taken from 1X PBS and placed on CO₂ pad for ~3-5 mins) had a small region of the cuticle (around abdominal segment 2) with underlying segmental nerves gently pinched with a pair of Dumont #5 fine forceps (Roboz Surgical RS4978). After injury, larvae were placed in small dishes containing standard fly food and kept at 25°C (12h light/dark) for 24h. Injuries were confirmed by the presence of posterior tail paralysis. For the new spared nerve crush assay, we performed injuries on wandering 2nd instar larvae with labeled type II octopaminergic neurons driven by Tdc2-gal4. The Tdc2-gal4 drives expression in 3 ventrally located midline neurons that sends one axon to the left side and one axon to the right side of the larvae. Similarly to complete nerve crushes, anesthetized larvae (ventral side up) were gently pinched with fine forceps on the left side (around abdominal segment 2), injuring nerves on one side while sparing nerves on the other side. Animals were placed in small dishes containing standard fly food and kept at 25°C (12h light/dark) for 24-72 hours.

Axon/synaptic branch laser injury

As previously described (Ghannad-Rezaie et al., 2012 [↗](#); Mishra et al., 2014 [↗](#); Smithson et al., 2024b [↗](#); Waller et al., 2024 [↗](#)), single wandering 2nd instar larva were taken from a dish containing 1X PBS and gently placed onto a kim wipe to remove excess PBS and then dipped into halocarbon oil 700 (H8898, Sigma). Each larva was then placed onto a glass coverslip dorsal side up (for inverted confocal) with an anterior to posterior (left to right) orientation. A PDMS microfluidic chip (<https://www.ukosmos.com/> [↗](#)) was placed on top of the larvae applying light force. A tight seal was created by applying gentle suction from a 30-ml syringe plunger. This vacuum suction immobilizes the intact larvae. The coverslip containing the microfluidic chip and larvae was mounted onto an Improvion spinning disk confocal system (PerkinElmer) connected to a Micropoint Laser Illumination and Ablation System (Andor Technology). The method for laser-induced microsurgery is described in (Smithson et al., 2024a [↗](#)). Prior to injury, the laser strength and region of interest were calibrated and optimized. Individual axonal branches (membrane GFP labeled) were identified and laser ablated. Confirmation of injury was demonstrated by a small absence of membrane GFP and degeneration of the proximal axon was detected as early as 8 hours after laser injury.


Imaging, Quantification and Analysis

Both experimental and control larvae were imaged on an Improvion spinning disk confocal system (PerkinElmer) and quantified using identical confocal settings. To quantify *puc*-lacZ levels, the mean intensity of lacZ immunolabeling was measured in specific injury confirmed neurons. Injured m12-gal4 and Tdc2-gal4 axons were traced back to identify corresponding cell bodies in the lateral and medial ventral nerve cord. *Puc*-lacZ contains a NLS sequence fused to lacZ, resulting in nuclear expression of *puc*. The mean lacZ intensity was measured per neuron and normalized to uninjured control neurons. All imaging, image contrast/color adjustments, and

quantification were conducted using Volocity software 6.2 (Improvision, Perkin Elmer). Data were analyzed and statistical tests were performed using Prism (GraphPad) and presented as mean $\pm$ SEM.

Acknowledgements

We would like to thank Mariana Jimenez, Luis Rivera, and Sarah Cooke for help with tissue processing, Eric Robertson and Chris Jasinski for technical assistance. We thank Heather Broihier, Dion Dickman, Dhananjay Yellajoshyula and Jerry Silver for helpful discussions and comments on the manuscript. We thank Yuanquan Song (University of Pennsylvania) for sharing piezo flies, Yujia (Henry) Hu and Bing Ye (University of Michigan) for gtACR1 flies and technical advice. Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537) and Vienna Drosophila Resource Center (VDRC, www.vdrc.at [↗](#) (Dietzl et al., 2007 [↗](#)) were used in this study. This research was funded by the National Institutes for Health (NS069844 to CAC) and by the Canadian Institutes for Health Research (CIHR Fellowship to LJS).



Supplemental Figure 1

Further characterization of branched injury assays to SNc and aCC motoneurons.

A) cartoon denoting nomenclature of MNSNc-26/29 (red) and MNSNc-27 (blue) branches.

B) Example images of nerve terminals and cell bodies of m12-Gal4; UAS-Bitbow2 labeled MNSNc neurons, used to confirm the anatomy.

C) Similarly to laser axotomy in [Figure 1C](#), nerve crush injury (24h) induces *puc-lacZ* expression in both MNSNc neurons, but not in MNSNc neurons that co-express *wnd*-RNAi.


D) Synaptic terminals (top row) and cell bodies (bottom row) of aCC motoneurons on muscle 1, labeled in Dpr4-Gal4, UAS-mCD8-GFP larvae. *puc-lacZ* expression (red) is induced following injuries that result in loss of all synaptic boutons but not following injuries to one branch that leave the other branch intact.

E) Quantification of *puc-lacZ* intensities in aCC neurons. A one-way ANOVA with Tukey test for multiple comparisons was performed. **** $p < 0.0001$.

F-G) Full (F) but not partial (G) removal of synaptic branches induces stability and trafficking ectopically expressed kinase-dead GFP-Wnd-KD (in UAS-GFP-Wnd-KD; m12-Gal4, UAS-mCD8-RFP animals). Synaptic branches from muscle 27 (F), or muscle 29 (G) were axotomized by laser surgery and imaged following 24h.

F) GFP-Wnd-KD protein accumulates at the proximal tip of axons that have lost all synaptic boutons.

G) GFP-Wnd-KD is barely detectable in axons following injuries that leave spared synaptic branches. Not shown, GFP-Wnd-KD levels in G are similar in uninjured MNSNc axons.



Supplemental Figure 2

Anatomy and laser surgery of Tdc2 bifurcated neurons.

A-E) Views of Tdc2-Gal4, UAS-mCD8-GFP expressing VUM neurons to illustrate their anatomy.


A) Individual confocal plane that shows 3 cell bodies in individual segments, which lie in the middle of the nerve cord. **B)** Side view and **C)** top view that show the locations of the bifurcations.

D) Cartoon of the 3 neurons from one segment, showing their bifurcations to symmetrical sides of the animal.


E) composite and *camera lucida* views of the NMJ terminals for the 3 neurons on one abdominal hemisegment. The 3 Tdc2 neurons each form stereotyped branches to innervate a unique group of muscles.

F) Example results from laser axotomies to individual Tdc2/VUM neurons (24h after injury) on either one side or both sides of the animal. Surgeries were carried out at the indicated locations which lie upstream of the final synaptic branches (at the transition zone between the segmental nerve and abdominal muscles). The stereotyped anatomy allows for identification of each VUM neuron (labeled 1, 2 and 3). Injured branches are marked with an asterisk while spared are marked with squares. For each neuron, only injuries to both bifurcations allowed for induction of *puc-lacZ*. Scale bars = 20 μm.

A



B



Supplemental Figure 3

Confirmation that elevated *puc-lacZ* expression in *hiw* mutants requires Wnd.

A) Top row shows *puc-lacZ* expression (red) together with a marker for nuclei (Draq5, gray). Control (lexA)-RNAi or wnd-RNAi UAS lines are driven by the BG380Gal4 driver in the background of control versus *hiw*^{ΔN}. Bottom row shows example NMJs in these genotypes. Scale bars = 20 μm.

B) Quantification of *puc-lacZ* expression in A. A one-way ANOVA with Tukey test for multiple comparisons was performed. **** *p* < 0.0001.

Name ^[LS1]	Stock ID	Target/Function	ref(s)	Driver (Gal4)	observation
UAS lines					
UAS- <i>shibire</i> -ts	RRID:BDSC_44222	dominant temperature sensitive mutation in dynamin - confers rapid inhibition of synaptic vesicle endocytosis and exocytosis.	(Kitamoto, 2001)	m12	temperature shift to 32C for 24h. (nSyb-Gal4, UAS-Shi-ts flies are paralyzed when shifted to 32C for 20 minutes). No difference between shifted (32C) and unshifted (25C) for uninjured, single branch injured, and complete axotomy. (shift was done immediately after laser surgery).
UAS-dORK-delta-C	RRID:BDSC_8928	constitutively open K+ rectifying channel. Confers electrical silencing.	(Nitabach et al., 2002)	m12	No difference from control (CS) flies, both uninjured and full axotomy.
UAS-dORK-delta-NC (non-conducting)	RRID:BDSC_6587	Non-conducting version of UAS-ORK1 (control)	(Nitabach et al., 2002)	m12	No difference from control (CS) flies, both uninjured and full axotomy.
UAS-gtACR1	RRID:BDSC_92983	light-inducible Cl- channel	(Mohammad et al., 2017)	m12	Kept in dark until 24h before dissection, when larvae were shifted to 530 nm light (for 24h). No difference between vehicle (ethanol) and ATR (1mM) control. As a control for the method, BG380Gal4; UAS-gtACR1 (ATR) flies showed rapid paralysis with shifted 530 nm light, but animals raised with vehicle food did not.
UAS-Sh-DN	RRID:BDSC_40974	mutated Shaker K+ channel which attenuates membrane excitability	(Mosca et al., 2005)	m12	No difference from control (CS) flies, uninjured

Supplementary Table 1

Summary of negative results from genetic manipulations that impair synaptic transmission and/or signaling at synapses.

The listed UAS lines, UAS-RNAi lines and genetic mutations were tested for their ability to alter the expression of *puc-lacZ* in motoneurons. Outcomes and experimental details are noted.

UAS- <i>tkv</i> -CA	RRID:BDSC_36537	consitutively activated type I BMP receptor (thickveins)		BG380	causes elevation of Gbb/BMP signaling (measured by pSMAD), but does not alter the induction of <i>puc-lacZ</i> in either injured (24h nerve crush) or uninjured animals.
endogenous mutations					
<i>paralytic (para)</i> -ts	RRID:BDSC_26701	nit of voltage-gated sodi	(Pittendrigh et al., 1997)		tested 6h at 37C, and 24h at 29. No difference between shifted and unshifted for uninjured animals. Also no difference between shifted and unshifted for full axotomy. (shift was done immediately after nerve crush).
<i>Dm-piezo</i> KO		peizo mechanosensitive Ca2+ channel	(Kim et al., 2012)		
RNAi lines					
<i>cacophony (cac)</i>	vdrc 101478	alpha1 subunit of voltage-gated calcium channel		m12 with dicer2	no difference from UAS-control (luciferase) RNAi
				BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>straightjacket (stj)</i>	RRID:BDSC_25807	alpha 2 delta subunit of voltage-gated calcium channel		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
				D42	no difference from UAS-control (luciferase) RNAi
<i>paralytic (para)</i>	RRID:BDSC_31471	nit of voltage-gated sodi		BG380	lethal
<i>paralytic (para)</i>	RRID:BDSC_31676	nit of voltage-gated sodi		BG380, D42	no difference from UAS-control (luciferase) RNAi
<i>shaker (sh)</i>	RRID:BDSC_31680	shaker (sh) voltage gated K+ channel		BG380, Dicer 2	no difference from UAS-control (luciferase) RNAi
<i>shaker cognate (shaw)</i>	RRID:BDSC_28346	voltage gated K channel Kv3.1		BG380, Dicer 2	no difference from UAS-control (luciferase) RNAi
	RRID:BDSC_27306			BG380, Dicer2	no difference from UAS-control (luciferase) RNAi

Supplementary Table 1 (continued)

<i>synaptosomal-associated protein (SNAP25)</i>	RRID:BDSC_34377	t-SNARE, regulation of neurotransmitter release		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
				D42	no difference from UAS-control (luciferase) RNAi
<i>synaptobrevin (nSyb)</i>	RRID:BDSC_31983	v-SNARE, role in synaptic vesicle exocytosis		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
				m12	no difference from UAS-control (luciferase) RNAi
<i>syntaxin 4 (syx4)</i>	RRID:BDSC_44054	tSNARE, role in synaptic vesicle exocytosis		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>syntaxin 1A (syx1A)</i>	RRID:BDSC_25811	tSNARE, role in synaptic vesicle exocytosis		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>synaptotagmin 4 (syt4)</i>	RRID:BDSC_39016	calcium dependent exocytosis of secretory vesicles.		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>vesicular glutamate transporter (vGlut)</i>	v104324	vesicular glutamate transporters		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	RRID:BDSC_40927			D42	no difference from UAS-control (luciferase) RNAi
				BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>peizo</i>	vdrc 2796	peizo mechanosensitive Ca2+ channel			no difference from UAS-control (luciferase) RNAi
<i>no mechanoreceptor potential C (nompC)</i>	vdrc 105579	pore-forming subunit of mechanotransduction channel		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>shortwing (sw)</i>	vdrc 101559	intermediate chain subunit of dynein		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	vdrc 48334			BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>neurexin 1 (nrx-1)</i>	vdrc 36326	presynaptic protein, cell-cell interactions, exocytosis, synaptic signaling		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	RRID:BDSC_27502			BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
				D42	no difference from UAS-control (luciferase) RNAi

Supplementary Table 1 (continued)

Supplementary Table 1 (continued)

	RRID:BDSC_32408		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
			D42	no difference from UAS-control (luciferase) RNAi
<i>neuroligin 1 (nlg1)</i>	vdrc 42616	postsynaptic cell surface protein, cell adhesion, excitatory synapses	BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>defective proboscis extension response 4 (dpr4)</i>	vdrc 28519	cell surface receptor for DIPs, adhesion protein, synapse organization	BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	vdrc 102905		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	vdrc 28518		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>fasciculin II (fasII)</i>	RRID:BDSC_34084	cell adhesion molecule, axonal pathfinding/guidance	BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	vdrc 8393		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
	vdrc 36350		BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>comatose/NSF1 (comt)</i>	RRID:BDSC_27890	N-ethylmaleimide-sensitive factor 1, role disassembly of SNARE complexes	D42	no difference from UAS-control (luciferase) RNAi
<i>semaphorin 2b (sema2b)</i>	RRID:BDSC_28932	secreted semaphorin, involed in embryonic axonal guidance	BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>plexin B (plexB)</i>	RRID:BDSC_57813	Acts as a receptor for Sema-2a and seems to suppress axon branching	BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>neuroglian (Nrg)</i>	RRID:BDSC_37496	L1-type cell adhesion molecule	BG380, Dicer2	no difference from UAS-control (luciferase) RNAi
<i>wishful thinking (wit)</i>	RRID:BDSC_25949	BMP type II receptor with known functions synaptic development and maintenance at the Drosophila NMJ	BG380, Dicer2	no effect on <i>puc-lacZ</i> induction in uninjured animals. <i>puc-lacZ</i> induction 24 post peripheral nerve crush is similar to wild type animals.
	vdrc 103808		BG380, Dicer2	"
<i>thickveins (tkv)</i>	RRID:BDSC_41904	BMP type I receptor with known functions synaptic development and maintenance at the Drosophila NMJ	BG380, Dicer2	no effect on <i>puc-lacZ</i> induction in uninjured animals. <i>puc-lacZ</i> induction 24 post peripheral nerve crush is similar to wild type animals.

References

- Abrams B, Grill B, Huang X, Jin Y (2008) **Cellular and molecular determinants targeting the *Caenorhabditis elegans* PHR protein RPM-1 to perisynaptic regions** *Dev Dyn* **237**:630–639
- Asghari Adib E, Smithson LJ, Collins CA (2018) **An axonal stress response pathway: degenerative and regenerative signaling by DLK** *Curr Opin Neurobiol* **53**:110–119
- Bhattacharya MRC, Gerdts J, Naylor SA, Royse EX, Ebstein SY, Sasaki Y, Milbrandt J, DiAntonio A (2012) **A model of toxic neuropathy in *Drosophila* reveals a role for MORN4 in promoting axonal degeneration** *J Neurosci* **32**:5054–5061
- Bounoutas A, Kratz J, Emtage L, Ma C, Nguyen KC, Chalfie M (2011) **Microtubule depolymerization in *Caenorhabditis elegans* touch receptor neurons reduces gene expression through a p38 MAPK pathway** *Proc Natl Acad Sci U S A* **108**:3982–3987
- Budnik V, Koh YH, Guan B, Hartmann B, Hough C, Woods D, Gorczyca M (1996) **Regulation of synapse structure and function by the *Drosophila* tumor suppressor gene *dlg*** *Neuron* **17**:627–640
- Chen C-H, Lee A, Liao C-P, Liu Y-W, Pan C-L (2014) **RHGF-1/PDZ-RhoGEF and retrograde DLK-1 signaling drive neuronal remodeling on microtubule disassembly** *Proc Natl Acad Sci U S A* **111**:16568–16573
- Chen L, Wang Z, Ghosh-Roy A, Hubert T, Yan D, O'Rourke S, Bowerman B, Wu Z, Jin Y, Chisholm AD (2011) **Axon regeneration pathways identified by systematic genetic screening in *C. elegans*** *Neuron* **71**:1043–1057
- Choi BJ, Imlach WL, Jiao W, Wolfram V, Wu Y, Grbic M, Cela C, Baines RA, Nitabach MN, McCabe BD (2014) **Miniature neurotransmission regulates *Drosophila* synaptic structural maturation** *Neuron* **82**:618–634
- Collins CA, Wairkar YP, Johnson SL, DiAntonio A (2006) **Highwire restrains synaptic growth by attenuating a MAP kinase signal** *Neuron* **51**:57–69
- Cosker KE, Courchesne SL, Segal RA (2008) **Action in the axon: generation and transport of signaling endosomes** *Curr Opin Neurobiol* **18**:270–275
- Daniels RW, Collins CA, Chen K, Gelfand MV, Featherstone DE, DiAntonio A (2006) **A single vesicular glutamate transporter is sufficient to fill a synaptic vesicle** *Neuron* **49**:11–16
- DeVault L, Mateusiak C, Palucki J, Brent M, Milbrandt J, DiAntonio A (2024) **The response of Dual-leucine zipper kinase (DLK) to nocodazole: Evidence for a homeostatic cytoskeletal repair mechanism** *PLoS One* **19**
- DiAntonio A, Schwarz TL (1994) **The effect on synaptic physiology of synaptotagmin mutations in *Drosophila*** *Neuron* **12**:909–920
- Dietzl G *et al.* (2007) **A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*** *Nature* **448**:151–156

- Feoktistov AI, Herman TG (2016) **Wallenda/DLK protein levels are temporally downregulated by Tramtrack69 to allow R7 growth cones to become stationary boutons** *Development* **143**:2983–2993
- Fernandes KJ, Fan DP, Tsui BJ, Cassar SL, Tetzlaff W (1999) **Influence of the axotomy to cell body distance in rat rubrospinal and spinal motoneurons: differential regulation of GAP-43, tubulins, and neurofilament-M** *J Comp Neurol* **414**:495–510
- Ghannad-Rezaie M, Wang X, Mishra B, Collins C, Chronis N (2012) **Microfluidic chips for in vivo imaging of cellular responses to neural injury in Drosophila larvae** *PLoS One* **7**
- Ghosh AS, Wang B, Pozniak CD, Chen M, Watts RJ, Lewcock JW (2011) **DLK induces developmental neuronal degeneration via selective regulation of proapoptotic JNK activity** *J Cell Biol* **194**:751–764
- Guedes-Dias P, Holzbaur ELF (2019) **Axonal transport: Driving synaptic function** *Science* **366** <https://doi.org/10.1126/science.aaw9997>
- Hammarlund M, Nix P, Hauth L, Jorgensen EM, Bastiani M (2009) **Axon regeneration requires a conserved MAP kinase pathway** *Science* **323**:802–806
- Han Y, Chien C, Goel P, He K, Pinales C, Buser C, Dickman D (2022) **Botulinum neurotoxin accurately separates tonic vs. phasic transmission and reveals heterosynaptic plasticity rules in Drosophila** *Elife* **11** <https://doi.org/10.7554/eLife.77924>
- Hilton BJ *et al.* (2022) **An active vesicle priming machinery suppresses axon regeneration upon adult CNS injury** *Neuron* **110**:51–69
- Holland SM, Collura KM, Ketschek A, Noma K, Ferguson TA, Jin Y, Gallo G, Thomas GM (2016) **Palmitoylation controls DLK localization, interactions and activity to ensure effective axonal injury signaling** *Proc Natl Acad Sci U S A* **113**:763–768
- Huebner EA, Strittmatter SM (2009) **Axon regeneration in the peripheral and central nervous systems** *Results Probl Cell Differ* **48**:339–351
- Huntwork-Rodriguez S, Wang B, Watkins T, Ghosh AS, Pozniak CD, Bustos D, Newton K, Kirkpatrick DS, Lewcock JW (2013) **JNK-mediated phosphorylation of DLK suppresses its ubiquitination to promote neuronal apoptosis** *J Cell Biol* **202**:747–763
- Hu Z, Deng N, Liu K, Zeng W (2019) **DLK mediates the neuronal intrinsic immune response and regulates glial reaction and neuropathic pain** *Exp Neurol* **322**
- Joy MT *et al.* (2019) **CCR5 Is a Therapeutic Target for Recovery after Stroke and Traumatic Brain Injury** *Cell* **176**:1143–1157
- Kim MD, Wen Y, Jan Y-N (2009) **Patterning and organization of motor neuron dendrites in the Drosophila larva** *Dev Biol* **336**:213–221
- Kim SE, Coste B, Chadha A, Cook B, Patapoutian A (2012) **The role of Drosophila Piezo in mechanical nociception** *Nature* **483**:209–212
- Kim S, Quagraine Y, Singh M, Kim JH (2024) **Rab11 suppresses neuronal stress signaling by localizing Dual leucine zipper kinase to axon terminals for protein turnover** *bioRxiv* <https://doi.org/10.1101/2023.04.18.537392>

- Kitamoto T (2001) **Conditional modification of behavior in *Drosophila* by targeted expression of a temperature-sensitive shibire allele in defined neurons** *J Neurobiol* **47**:81–92
- Koon AC, Ashley J, Barria R, DasGupta S, Brain R, Waddell S, Alkema MJ, Budnik V (2011) **Autoregulatory and paracrine control of synaptic and behavioral plasticity by octopaminergic signaling** *Nat Neurosci* **14**:190–199
- Kurup N, Yan D, Goncharov A, Jin Y (2015) **Dynamic microtubules drive circuit rewiring in the absence of neurite remodeling** *Curr Biol* **25**:1594–1605
- Larhammar M, Huntwork-Rodriguez S, Rudhard Y, Sengupta-Ghosh A, Lewcock JW (2017) **The Ste20 Family Kinases MAP4K4, MINK1, and TNIK Converge to Regulate Stress-Induced JNK Signaling in Neurons** *J Neurosci* **37**:11074–11084
- Lee T, Luo L (1999) **Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis** *Neuron* **22**:451–461
- Le Pichon CE *et al.* (2017) **Loss of dual leucine zipper kinase signaling is protective in animal models of neurodegenerative disease** *Sci Transl Med* **9** <https://doi.org/10.1126/scitranslmed.aag0394>
- Li J *et al.* (2017) **Restraint of presynaptic protein levels by Wnd/DLK signaling mediates synaptic defects associated with the kinesin-3 motor Unc-104** *Elife* **6** <https://doi.org/10.7554/eLife.24271>
- Li Y *et al.* (2021) **Bitbow Enables Highly Efficient Neuronal Lineage Tracing and Morphology Reconstruction in Single *Drosophila* Brains** *Front Neural Circuits* **15**
- Lorenzana AO, Lee JK, Mui M, Chang A, Zheng B (2015) **A surviving intact branch stabilizes remaining axon architecture after injury as revealed by in vivo imaging in the mouse spinal cord** *Neuron* **86**:947–954
- Martín-Blanco E, Gampel A, Ring J, Virdee K, Kirov N, Tolkovsky AM, Martínez-Arias A (1998) . **puckered encodes a phosphatase that mediates a feedback loop regulating JNK activity during dorsal closure in *Drosophila*** *Genes Dev* **12**:557–570
- Mason MRJ, Lieberman AR, Anderson PN (2003) **Corticospinal neurons up-regulate a range of growth-associated genes following intracortical, but not spinal, axotomy** *Eur J Neurosci* **18**:789–802
- Mishra B, Ghannad-Rezaie M, Li J, Wang X, Hao Y, Ye B, Chronis N, Collins CA (2014) **Using microfluidics chips for live imaging and study of injury responses in *Drosophila* larvae** *J Vis Exp* **e 50998**
- Mohammad F, Stewart JC, Ott S, Chlebikova K, Chua JY, Koh T-W, Ho J, Claridge-Chang A (2017) **Optogenetic inhibition of behavior with anion channelrhodopsins** *Nat Methods* **14**:271–274
- Monastirioti M, Gorczyca M, Rapus J, Eckert M, White K, Budnik V (1995) **Octopamine immunoreactivity in the fruit fly *Drosophila melanogaster*** *J Comp Neurol* **356**:275–287
- Mosca TJ, Carrillo RA, White BH, Keshishian H (2005) **Dissection of synaptic excitability phenotypes by using a dominant-negative Shaker K⁺ channel subunit** *Proc Natl Acad Sci U S A* **102**:3477–3482

- Nakata K, Abrams B, Grill B, Goncharov A, Huang X, Chisholm AD, Jin Y (2005) **Regulation of a DLK-1 and p38 MAP kinase pathway by the ubiquitin ligase RPM-1 is required for presynaptic development** *Cell* **120**:407–420
- Nitabach MN, Blau J, Holmes TC (2002) **Electrical silencing of *Drosophila* pacemaker neurons stops the free-running circadian clock** *Cell* **109**:485–495
- Niu J, Holland SM, Ketschek A, Collura KM, Hesketh NL, Hayashi T, Gallo G, Thomas GM (2022) **Palmitoylation couples the kinases DLK and JNK3 to facilitate prodegenerative axon- to-soma signaling** *Sci Signal* **15**
- Opperman KJ, Grill B (2014) **RPM-1 is localized to distinct subcellular compartments and regulates axon length in GABAergic motor neurons** *Neural Dev* **9**
- Patel S *et al.* (2017) **Selective Inhibitors of Dual Leucine Zipper Kinase (DLK, MAP3K12) with Activity in a Model of Alzheimer’s Disease** *J Med Chem* **60**:8083–8102
- Pérez-Moreno JJ, O’Kane CJ (2019) **GAL4 Drivers Specific for Type Ib and Type Is Motor Neurons in *Drosophila*** *G3* **9**:453–462
- Petzoldt AG (2023) **Presynaptic Precursor Vesicles-Cargo, Biogenesis, and Kinesin-Based Transport across Species** *Cells* **12** <https://doi.org/10.3390/cells12182248>
- Pittendrigh B, Reenan R, ffrench-Constant RH, Ganetzky B (1997) **Point mutations in the *Drosophila* sodium channel gene para associated with resistance to DDT and pyrethroid insecticides** *Mol Gen Genet* **256**:602–610
- Raineteau O, Schwab ME (2001) **Plasticity of motor systems after incomplete spinal cord injury** *Nat Rev Neurosci* **2**:263–273
- Ramón y Cajal S, May Raoul M. (1928) **Degeneration & Regeneration of the Nervous System** London: Oxford University Press
- Ramzan F, Abrar F, Mishra GG, Liao LMQ, Martin DDO (2023) **Lost in traffic: consequences of altered palmitoylation in neurodegeneration** *Front Physiol* **14**
- Ritzenthaler S, Suzuki E, Chiba A (2000) **Postsynaptic filopodia in muscle cells interact with innervating motoneuron axons** *Nat Neurosci* **3**:1012–1017
- Sanyal S (2009) **Genomic mapping and expression patterns of C380, OK6 and D42 enhancer trap lines in the larval nervous system of *Drosophila*** *Gene Expr Patterns* **9**:371–380
- Smithson LJ, Waller TJ, Collins CA (2024) **Laser Microsurgery in *Drosophila* Larvae Using the MicroPoint Ablation System** *Cold Spring Harb Protoc* <https://doi.org/10.1101/pdb.prot108171>
- Smithson LJ, Waller TJ, Collins CA (2024) **Immobilizing Second-Instar *Drosophila* Larvae for Imaging and Surgery Using the Larva Chip** *Cold Spring Harb Protoc* <https://doi.org/10.1101/pdb.prot108170>
- Stone MC, Albertson RM, Chen L, Rolls MM (2014) **Dendrite injury triggers DLK-independent regeneration** *Cell Rep* **6**:247–253

- Tedeschi A, Dupraz S, Laskowski CJ, Xue J, Ulas T, Beyer M, Schultze JL, Bradke F (2016) **The Calcium Channel Subunit Alpha2delta2 Suppresses Axon Regeneration in the Adult CNS** *Neuron* **92**:419–434
- Valakh V, Frey E, Babetto E, Walker LJ, DiAntonio A (2015) **Cytoskeletal disruption activates the DLK/JNK pathway, which promotes axonal regeneration and mimics a preconditioning injury** *Neurobiol Dis* **77**:13–25
- Valakh V, Walker LJ, Skeath JB, DiAntonio A (2013) **Loss of the spectraplakins short stop activates the DLK injury response pathway in Drosophila** *J Neurosci* **33**:17863–17873
- Vömel M, Wegener C (2008) **Neuroarchitecture of aminergic systems in the larval ventral ganglion of Drosophila melanogaster** *PLoS One* **3**
- Waller TJ, Smithson LJ, Collins CA (2024) **Peripheral Nerve Crush in Drosophila Larvae** *Cold Spring Harb Protoc* <https://doi.org/10.1101/pdb.prot108169>
- Wang Z, Kumaran M, Batsel E, Testor-Cabrera S, Beine Z, Ribelles AA, Tsoulfas P, Venkatesh I, Blackmore MG (2024) **Injury distance limits the transcriptional response to spinal injury** *bioRxiv* <https://doi.org/10.1101/2024.05.27.596075>
- Wan HI, DiAntonio A, Fetter RD, Bergstrom K, Strauss R, Goodman CS (2000) **Highwire regulates synaptic growth in Drosophila** *Neuron* **26**:313–329
- Watkins TA, Wang B, Huntwork-Rodriguez S, Yang J, Jiang Z, Eastham-Anderson J, Modrusan Z, Kaminker JS, Tessier-Lavigne M, Lewcock JW (2013) **DLK initiates a transcriptional program that couples apoptotic and regenerative responses to axonal injury** *Proc Natl Acad Sci U S A* **110**:4039–4044
- Welsbie DS *et al.* (2017) **Enhanced Functional Genomic Screening Identifies Novel Mediators of Dual Leucine Zipper Kinase-Dependent Injury Signaling in Neurons** *Neuron* **94**:1142–1154
- Welsbie DS *et al.* (2013) **Functional genomic screening identifies dual leucine zipper kinase as a key mediator of retinal ganglion cell death** *Proc Natl Acad Sci U S A* **110**:4045–4050
- Wlaschin JJ, Gluski JM, Nguyen E, Silberberg H, Thompson JH, Chesler AT, Le Pichon CE (2018) **Dual leucine zipper kinase is required for mechanical allodynia and microgliosis after nerve injury** *Elife* **7** <https://doi.org/10.7554/eLife.33910>
- Wu C, Wairkar YP, Collins CA, DiAntonio A (2005) **Highwire function at the Drosophila neuromuscular junction: spatial, structural, and temporal requirements** *J Neurosci* **25**:9557–9566
- Wu Z, Ghosh-Roy A, Yanik MF, Zhang JZ, Jin Y, Chisholm AD (2007) **Caenorhabditis elegans neuronal regeneration is influenced by life stage, ephrin signaling, and synaptic branching** *Proc Natl Acad Sci U S A* **104**:15132–15137
- Xiong X, Collins CA (2012) **A conditioning lesion protects axons from degeneration via the Wallenda/DLK MAP kinase signaling cascade** *J Neurosci* **32**:610–615
- Xiong X, Hao Y, Sun K, Li J, Li X, Mishra B, Soppina P, Wu C, Hume RI, Collins CA (2012) **The Highwire ubiquitin ligase promotes axonal degeneration by tuning levels of Nmnat protein** *PLoS Biol* **10**

Xiong X, Wang X, Ewanek R, Bhat P, Diantonio A, Collins CA (2010) **Protein turnover of the Wallenda/DLK kinase regulates a retrograde response to axonal injury** *J Cell Biol* **191**:211–223

Xiong X, Wang X, Ewanek R, Bhat P, Diantonio A, Collins CA (2010) **Protein turnover of the Wallenda/DLK kinase regulates a retrograde response to axonal injury** *J Cell Biol* **191**:211–223

Yan D, Wu Z, Chisholm AD, Jin Y (2009) **The DLK-1 kinase promotes mRNA stability and local translation in *C. elegans* synapses and axon regeneration** *Cell* **138**:1005–1018

Zhang X, Jeong H, Niu J, Holland SM, Rotanz BN, Gordon J, Einarson MB, Childers WE, Thomas GM (2024) **Novel inhibitors of acute, axonal DLK palmitoylation are neuroprotective and avoid the deleterious side effects of cell-wide DLK inhibition** *bioRxiv* <https://doi.org/10.1101/2024.04.19.590310>

Zhen M, Huang X, Bamber B, Jin Y (2000) **Regulation of presynaptic terminal organization by *C. elegans* RPM-1, a putative guanine nucleotide exchanger with a RING-H2 finger domain** *Neuron* **26**:331–343

Author information

Laura J Smithson

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, USA

Juliana Zang

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, USA

ORCID iD: [0000-0002-5738-8355](https://orcid.org/0000-0002-5738-8355)

Lucas Junginger

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, USA

Thomas J Waller

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, USA

Lauren Reilly-Jankowiak

Department of Neurosciences, Case Western Reserve University, Cleveland, USA

Sophia Khan

Department of Neurosciences, Case Western Reserve University, Cleveland, USA

Ye Li

Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, USA
ORCID iD: [0000-0002-8647-384X](https://orcid.org/0000-0002-8647-384X)

Dawen Cai

Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, USA
ORCID iD: [0000-0003-4471-2061](https://orcid.org/0000-0003-4471-2061)

Catherine A Collins

Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, USA, Department of Neurosciences, Case Western Reserve University, Cleveland, USA
ORCID iD: [0000-0002-1608-6692](https://orcid.org/0000-0002-1608-6692)

For correspondence: catherinecollins2@case.edu

Editors

Reviewing Editor

Dion Dickman

University of Southern California, Los Angeles, United States of America

Senior Editor

Claude Desplan

New York University, New York, United States of America

Reviewer #1 (Public review):

This manuscript presents an interesting exploration of the potential activation mechanisms of DLK following axonal injury. While the experiments are beautifully conducted and the data are solid, I feel that there is insufficient evidence to fully support the conclusions made by the authors.

In this manuscript, the authors exclusively use the puc-lacZ reporter to determine the activation of DLK. This reporter has been shown to be induced when DLK is activated. However, there is insufficient evidence to confirm that the absence of reporter activation necessarily indicates that DLK is inactive. As with many MAP kinase pathways, the DLK pathway can be locally or globally activated in neurons, and the level of DLK activation may depend on the strength of the stimulation. This reporter might only reflect strong DLK activation and may not be turned on if DLK is weakly activated. The results presented in this manuscript support this interpretation. Strong stimulation, such as axotomy of all synaptic branches, caused robust DLK activation, as indicated by puc-lacZ expression. In contrast, weak stimulation, such as axotomy of some synaptic branches, resulted in weaker DLK activation, which did not induce the puc-lacZ reporter. This suggests that the strength of DLK activation depends on the severity of the injury rather than the presence of intact synapses. Given that this is a central conclusion of the study, it may be worthwhile to confirm this further. Alternatively, the authors may consider refining their conclusion to better align with the evidence presented.

As noted by the authors, DLK has been implicated in both axon regeneration and degeneration. Following axotomy, DLK activation can lead to the degeneration of distal axons, where synapses are located. This raises an important question: how is DLK activated in distal axons? The authors might consider discussing the significance of this "synapse connection-dependent" DLK activation in the broader context of DLK function and activation mechanisms.

<https://doi.org/10.7554/eLife.104896.1.sa3>

Reviewer #2 (Public review):**Summary:**

The authors study a panel of sparsely labeled neuronal lines in *Drosophila* that each form multiple synapses. Critically, each axonal branch can be injured without affecting the others, allowing the authors to differentiate between injuries that affect all axonal branches versus those that do not, creating spared branches. Axonal injuries are known to cause Wnd (mammalian DLK)-dependent retrograde signals to the cell body, culminating in a transcriptional response. This work identifies a fascinating new phenomenon that this injury response is not all-or-none. If even a single branch remains uninjured, the injury signal is not activated in the cell body. The authors rule out that this could be due to changes in the abundance of Wnd (perhaps if incrementally activated at each injured branch) by Wnd, Hiw's known negative regulator. Thus there is both a yet-undiscovered mechanism to regulate Wnd signaling, and more broadly a mechanism by which the neuron can integrate the degree of injury it has sustained. It will now be important to tease apart the mechanism(s) of this fascinating phenomenon. But even absent a clear mechanism, this is a new biology that will inform the interpretation of injury signaling studies across species.

Strengths:

- (1) A conceptually beautiful series of experiments that reveal a fascinating new phenomenon is described, with clear implications (as the authors discuss in their Discussion) for injury signaling in mammals.
- (2) Suggests a new mode of Wnd regulation, independent of Hiw.

Weaknesses:

- (1) The use of a somatic transcriptional reporter for Wnd activity is powerful, however, the reporter indicates whether the transcriptional response was activated, not whether the injury signal was received. It remains possible that Wnd is still activated in the case of a spared branch, but that this activation is either local within the axons (impossible to determine in the absence of a local reporter) or that the retrograde signal was indeed generated but it was somehow insufficient to activate transcription when it entered the cell body. This is more of a mechanistic detail and should not detract from the overall importance of the study
- (2) That the protective effect of a spared branch is independent of Hiw, the known negative regulator of Wnd, is fascinating. But this leaves open a key question: what is the signal?

<https://doi.org/10.7554/eLife.104896.1.sa2>

Reviewer #3 (Public review):**Summary:**

This manuscript seeks to understand how nerve injury-induced signaling to the nucleus is influenced, and it establishes a new location where these principles can be studied. By identifying and mapping specific bifurcated neuronal innervations in the *Drosophila* larvae, and using laser axotomy to localize the injury, the authors find that sparing a branch of a complex muscular innervation is enough to impair Wallenda-puc (analogous to DLK-JNK-cJun) signaling that is known to promote regeneration. It is only when all connections to the target are disconnected that cJun-transcriptional activation occurs.

Overall, this is a thorough and well-performed investigation of the mechanism of spared-branch influence on axon injury signaling. The findings on control of wnd are important because this is a very widely used injury signaling pathway across species and injury models. The authors present detailed and carefully executed experiments to support their conclusions. Their effort to identify the control mechanism is admirable and will be of aid to the field as they continue to try to understand how to promote better regeneration of axons.

Strengths:

The paper does a very comprehensive job of investigating this phenomenon at multiple locations and through both pinpoint laser injury as well as larger crush models. They identify a non-hiw based restraint mechanism of the wnd-puc signaling axis that presumably originates from the spared terminal. They also present a large list of tests they performed to identify the actual restraint mechanism from the spared branch, which has ruled out many of the most likely explanations. This is an extremely important set of information to report, to guide future investigators in this and other model organisms on mechanisms by which regeneration signaling is controlled (or not).

Weaknesses:

The weakest data presented by this manuscript is the study of the actual amounts of Wallenda protein in the axon. The authors argue that increased Wnd protein is being anterogradely delivered from the soma, but no support for this is given. Whether this change is due to transcription/translation, protein stability, transport, or other means is not investigated in this work. However, because this point is not central to the arguments in the paper, it is only a minor critique.

As far as the scope of impact: because the conclusions of the paper are focused on a single (albeit well-validated) reporter in different types of motor neurons, it is hard to determine whether the mechanism of spared branch inhibition of regeneration requires wnd-puc (DLK/cJun) signaling in all contexts (for example, sensory axons or interneurons). Is the nerve-muscle connection the rule or the exception in terms of regeneration program activation?

Because changes in puc-lacZ intensity are the major readout, it would be helpful to better explain the significance of the amount of puc-lacZ in the nucleus with respect to the activation of regeneration. Is it known that scaling up the amount of puc-lacZ transcription scales functional responses (regeneration or others)? The alternative would be that only a small amount of puc-lacZ is sufficient to efficiently induce relevant pathways (threshold response).

<https://doi.org/10.7554/eLife.104896.1.sa1>

Author response:

Reviewer #1 (Public review):

This manuscript presents an interesting exploration of the potential activation mechanisms of DLK following axonal injury. While the experiments are beautifully conducted and the data are solid, I feel that there is insufficient evidence to fully support the conclusions made by the authors.

In this manuscript, the authors exclusively use the puc-lacZ reporter to determine the activation of DLK. This reporter has been shown to be induced when DLK is activated. However, there is insufficient evidence to confirm that the absence of reporter activation

necessarily indicates that DLK is inactive. As with many MAP kinase pathways, the DLK pathway can be locally or globally activated in neurons, and the level of DLK activation may depend on the strength of the stimulation. This reporter might only reflect strong DLK activation and may not be turned on if DLK is weakly activated. The results presented in this manuscript support this interpretation. Strong stimulation, such as axotomy of all synaptic branches, caused robust DLK activation, as indicated by puc-lacZ expression. In contrast, weak stimulation, such as axotomy of some synaptic branches, resulted in weaker DLK activation, which did not induce the puc-lacZ reporter. This suggests that the strength of DLK activation depends on the severity of the injury rather than the presence of intact synapses. Given that this is a central conclusion of the study, it may be worthwhile to confirm this further. Alternatively, the authors may consider refining their conclusion to better align with the evidence presented.

We wish to further clarify a striking aspect of puc-lacZ induction following injury: it is bimodal. It is either induced (in various injuries that remove all synaptic boutons), or not induced, including in injuries that spared only 1-2 remaining boutons. This was particularly evident for injuries that spared the NMJ on muscle 29, which is comprised of only a few boutons. In some instances, only a single bouton was evident on muscle 29. While our injuries varied enormously in the number of branches and boutons that were lost, we did not see a comparable variability in puc-lacZ induction. In the revision we will include additional images to better demonstrate this observation.

The reviewer (and others) fairly point out that our current study focuses on puc-lacZ as a reporter of Wnd signaling in the cell body. We consider this to be a downstream integration of events in axons that are more challenging to detect. It is striking that this integration appears strongly sensitized to the presence of spared synaptic boutons. Examination of Wnd's activation in axons and synapses is a goal for our future work.

As noted by the authors, DLK has been implicated in both axon regeneration and degeneration. Following axotomy, DLK activation can lead to the degeneration of distal axons, where synapses are located. This raises an important question: how is DLK activated in distal axons? The authors might consider discussing the significance of this "synapse connection-dependent" DLK activation in the broader context of DLK function and activation mechanisms.

While it has been noted that inhibition of DLK can mildly delay Wallerian degeneration (Miller et al., 2009), this does not appear to be the case for retinal ganglion cell axons following optic nerve crush (Fernandes et al., 2014). It is also not the case for *Drosophila* motoneurons and NMJ terminals following peripheral nerve injury (Xiong et al., 2012; Xiong and Collins, 2012). Instead, overexpression of Wnd or activation of Wnd by a conditioning injury leads to an opposite phenotype - an increase in resiliency to Wallerian degeneration for axons that have been previously injured (Xiong et al., 2012; Xiong and Collins, 2012). The downstream outcome of Wnd activation is highly dependent on the context; it may be an integration of the outcomes of local Wnd/DLK activation in axons with downstream consequences of nuclear/cell body signaling. The current study suggests some rules for the cell body signaling, however, how Wnd is regulated at synapses and why it promotes degeneration in some circumstances but not others are important future questions.

For the reviewer's suggestion, it is interesting to consider DLK's potential contributions to the loss of NMJ synapses in a mouse model of ALS (Le Pichon et al., 2017; Wlaschin et al., 2023). Our findings suggest that the synaptic terminal is an important locus of DLK regulation, while dysfunction of NMJ terminals is an important feature of the 'dying back' hypothesis of disease etiology (Dadon-Nachum et al., 2011; Verma et al., 2022). We propose that the regulation of DLK at synaptic terminals is an important area for future study, and may reveal how DLK might be modulated to curtail disease progression. Of note, DLK inhibitors are in clinical

trials (Katz et al., 2022; Le et al., 2023; Siu et al., 2018), but at least some have been paused due to safety concerns (Katz et al., 2022). Further understanding of the mechanisms that regulate DLK are needed to understand whether and how DLK and its downstream signaling can be tuned for therapeutic benefit.

Reviewer #2 (Public review):

Summary:

The authors study a panel of sparsely labeled neuronal lines in Drosophila that each form multiple synapses. Critically, each axonal branch can be injured without affecting the others, allowing the authors to differentiate between injuries that affect all axonal branches versus those that do not, creating spared branches. Axonal injuries are known to cause Wnd (mammalian DLK)-dependent retrograde signals to the cell body, culminating in a transcriptional response. This work identifies a fascinating new phenomenon that this injury response is not all-or-none. If even a single branch remains uninjured, the injury signal is not activated in the cell body. The authors rule out that this could be due to changes in the abundance of Wnd (perhaps if incrementally activated at each injured branch) by Wnd, Hiw's known negative regulator. Thus there is both a yet-undiscovered mechanism to regulate Wnd signaling, and more broadly a mechanism by which the neuron can integrate the degree of injury it has sustained. It will now be important to tease apart the mechanism(s) of this fascinating phenomenon. But even absent a clear mechanism, this is a new biology that will inform the interpretation of injury signaling studies across species.

Strengths:

(1) A conceptually beautiful series of experiments that reveal a fascinating new phenomenon is described, with clear implications (as the authors discuss in their Discussion) for injury signaling in mammals.

(2) Suggests a new mode of Wnd regulation, independent of Hiw.

Weaknesses:

(1) The use of a somatic transcriptional reporter for Wnd activity is powerful, however, the reporter indicates whether the transcriptional response was activated, not whether the injury signal was received. It remains possible that Wnd is still activated in the case of a spared branch, but that this activation is either local within the axons (impossible to determine in the absence of a local reporter) or that the retrograde signal was indeed generated but it was somehow insufficient to activate transcription when it entered the cell body. This is more of a mechanistic detail and should not detract from the overall importance of the study

We agree. The puc-lacZ reporter tells us about signaling in the cell body, but whether and how Wnd is regulated in axons and synaptic branches, which we think occurs upstream of the cell body response, remains to be addressed in future studies.

(2) That the protective effect of a spared branch is independent of Hiw, the known negative regulator of Wnd, is fascinating. But this leaves open a key question: what is the signal?

This is indeed an important future question, and would still be a question even if Hiw were part of the protective mechanism by the spared synaptic branch. Our current hypothesis (outlined in Figure 4) is that regulation of Wnd is tied to the retrograde trafficking of a signaling organelle in axons. The Hiw-independent regulation complements other

observations in the literature that multiple pathways regulate Wnd/DLK (Collins et al., 2006; Feoktistov and Herman, 2016; Klinedinst et al., 2013; Li et al., 2017; Russo and DiAntonio, 2019; Valakh et al., 2013). It is logical for this critical stress response pathway to have multiple modes of regulation that may act in parallel to tune and restrain its activation.

Reviewer #3 (Public review):

Summary:

This manuscript seeks to understand how nerve injury-induced signaling to the nucleus is influenced, and it establishes a new location where these principles can be studied. By identifying and mapping specific bifurcated neuronal innervations in the Drosophila larvae, and using laser axotomy to localize the injury, the authors find that sparing a branch of a complex muscular innervation is enough to impair Wallenda-puc (analogous to DLK-JNK-cJun) signaling that is known to promote regeneration. It is only when all connections to the target are disconnected that cJun-transcriptional activation occurs.

Overall, this is a thorough and well-performed investigation of the mechanism of spared-branch influence on axon injury signaling. The findings on control of wnd are important because this is a very widely used injury signaling pathway across species and injury models. The authors present detailed and carefully executed experiments to support their conclusions. Their effort to identify the control mechanism is admirable and will be of aid to the field as they continue to try to understand how to promote better regeneration of axons.

Strengths:

The paper does a very comprehensive job of investigating this phenomenon at multiple locations and through both pinpoint laser injury as well as larger crush models. They identify a non-hiw based restraint mechanism of the wnd-puc signaling axis that presumably originates from the spared terminal. They also present a large list of tests they performed to identify the actual restraint mechanism from the spared branch, which has ruled out many of the most likely explanations. This is an extremely important set of information to report, to guide future investigators in this and other model organisms on mechanisms by which regeneration signaling is controlled (or not).

Weaknesses:

The weakest data presented by this manuscript is the study of the actual amounts of Wallenda protein in the axon. The authors argue that increased Wnd protein is being anterogradely delivered from the soma, but no support for this is given. Whether this change is due to transcription/translation, protein stability, transport, or other means is not investigated in this work. However, because this point is not central to the arguments in the paper, it is only a minor critique.

We agree and are glad that the reviewer considers this a minor critique; this is an area for future study. In Supplemental Figure 1 we present differences in the levels of an ectopically expressed GFP-Wnd-kinase-dead transgene, which is strikingly increased in axons that have received a full but not partial axotomy. We suspect this accumulation occurs downstream of the cell body response because of the timing. We observed the accumulations after 24 hours (Figure S1F) but not at early (1-4 hour) time points following axotomy (data not shown). Further study of the local regulation of Wnd protein and its kinase activity in axons is an important future direction.

As far as the scope of impact: because the conclusions of the paper are focused on a single (albeit well-validated) reporter in different types of motor neurons, it is hard to

determine whether the mechanism of spared branch inhibition of regeneration requires wnd-puc (DLK/cjun) signaling in all contexts (for example, sensory axons or interneurons). Is the nerve-muscle connection the rule or the exception in terms of regeneration program activation?

DLK signaling is strongly activated in DRG sensory neurons following peripheral nerve injury (Shin et al., 2012), despite the fact that sensory neurons have bifurcated axons and their projections in the dorsal spinal cord are not directly damaged by injuries to the peripheral nerve. Therefore it is unlikely that protection by a spared synapse is a universal rule for all neuron types. However the molecular mechanisms that underlie this regulation may indeed be shared across different types of neurons but utilized in different ways. For instance, nerve growth factor withdrawal can lead to activation of DLK (Ghosh et al., 2011), however neurotrophins and their receptors are regulated and implemented differently in different cell types. We suspect that the restraint of Wnd signaling by the spared synaptic branch shares a common underlying mechanism with the restraint of DLK signaling by neurotrophin signaling. Further elucidation of the molecular mechanism is an important next step towards addressing this question.

Because changes in puc-lacZ intensity are the major readout, it would be helpful to better explain the significance of the amount of puc-lacZ in the nucleus with respect to the activation of regeneration. Is it known that scaling up the amount of puc-lacZ transcription scales functional responses (regeneration or others)? The alternative would be that only a small amount of puc-lacZ is sufficient to efficiently induce relevant pathways (threshold response).

While induction of puc-lacZ expression correlates with Wnd-mediated phenotypes, including sprouting of injured axons (Xiong et al., 2010), protection from Wallerian degeneration (Xiong et al., 2012; Xiong and Collins, 2012) and synaptic overgrowth (Collins et al., 2006), we have not observed any correlation between the degree of puc-lacZ induction (eg modest, medium or high) and the phenotypic outcomes (sprouting, overgrowth, etc). Rather, there appears to be a striking all-or-none difference in whether puc-lacZ is induced or not induced. There may indeed be a threshold that can be restrained through multiple mechanisms. We posit in figure 4 that restraint may take place in the cell body, where it can be influenced by the spared bifurcation.

References Cited:

- Collins CA, Wairkar YP, Johnson SL, DiAntonio A. 2006. Highwire restrains synaptic growth by attenuating a MAP kinase signal. *Neuron* 51:57–69.
- Dadon-Nachum M, Melamed E, Offen D. 2011. The “dying-back” phenomenon of motor neurons in ALS. *J Mol Neurosci* 43:470–477.
- Feoktistov AI, Herman TG. 2016. Wallenda/DLK protein levels are temporally downregulated by Tramtrack69 to allow R7 growth cones to become stationary boutons. *Development* 143:2983–2993.
- Fernandes KA, Harder JM, John SW, Shrager P, Libby RT. 2014. DLK-dependent signaling is important for somal but not axonal degeneration of retinal ganglion cells following axonal injury. *Neurobiol Dis* 69:108–116.
- Ghosh AS, Wang B, Pozniak CD, Chen M, Watts RJ, Lewcock JW. 2011. DLK induces developmental neuronal degeneration via selective regulation of proapoptotic JNK activity. *J Cell Biol* 194:751–764.

- Hao Y, Frey E, Yoon C, Wong H, Nestorovski D, Holzman LB, Giger RJ, DiAntonio A, Collins C. 2016. An evolutionarily conserved mechanism for cAMP elicited axonal regeneration involves direct activation of the dual leucine zipper kinase DLK. *Elife* 5. doi:10.7554/eLife.14048
- Huntwork-Rodriguez S, Wang B, Watkins T, Ghosh AS, Pozniak CD, Bustos D, Newton K, Kirkpatrick DS, Lewcock JW. 2013. JNK-mediated phosphorylation of DLK suppresses its ubiquitination to promote neuronal apoptosis. *J Cell Biol* 202:747–763.
- Katz JS, Rothstein JD, Cudkowicz ME, Genge A, Oskarsson B, Hains AB, Chen C, Galanter J, Burgess BL, Cho W, Kerchner GA, Yeh FL, Ghosh AS, Cheeti S, Brooks L, Honigberg L, Couch JA, Rothenberg ME, Brunstein F, Sharma KR, van den Berg L, Berry JD, Glass JD. 2022. A Phase 1 study of GDC-0134, a dual leucine zipper kinase inhibitor, in ALS. *Ann Clin Transl Neurol* 9:50–66.
- Klinedinst S, Wang X, Xiong X, Haenfler JM, Collins CA. 2013. Independent pathways downstream of the Wnd/DLK MAPKKK regulate synaptic structure, axonal transport, and injury signaling. *J Neurosci* 33:12764–12778.
- Le K, Soth MJ, Cross JB, Liu G, Ray WJ, Ma J, Goodwani SG, Acton PJ, Buggia-Prevot V, Akkermans O, Barker J, Conner ML, Jiang Y, Liu Z, McEwan P, Warner-Schmidt J, Xu A, Zebisch M, Heijnen CJ, Abrahams B, Jones P. 2023. Discovery of IACS-52825, a potent and selective DLK inhibitor for treatment of chemotherapy-induced peripheral neuropathy. *J Med Chem* 66:9954–9971.
- Le Pichon CE, Meilandt WJ, Dominguez S, Solanoy H, Lin H, Ngu H, Gogineni A, Sengupta Ghosh A, Jiang Z, Lee S-H, Maloney J, Gandham VD, Pozniak CD, Wang B, Lee S, Siu M, Patel S, Modrusan Z, Liu X, Rudhard Y, Baca M, Gustafson A, Kaminker J, Carano RAD, Huang EJ, Foreman O, Weimer R, Scarce-Levie K, Lewcock JW. 2017. Loss of dual leucine zipper kinase signaling is protective in animal models of neurodegenerative disease. *Sci Transl Med* 9. doi:10.1126/scitranslmed.aag0394
- Li J, Zhang YV, Asghari Adib E, Stanchev DT, Xiong X, Klinedinst S, Soppina P, Jahn TR, Hume RI, Rasse TM, Collins CA. 2017. Restraint of presynaptic protein levels by Wnd/DLK signaling mediates synaptic defects associated with the kinesin-3 motor Unc-104. *Elife* 6. doi:10.7554/eLife.24271
- Miller BR, Press C, Daniels RW, Sasaki Y, Milbrandt J, DiAntonio A. 2009. A dual leucine kinase-dependent axon self-destruction program promotes Wallerian degeneration. *Nat Neurosci* 12:387–389.
- Nihalani D, Merritt S, Holzman LB. 2000. Identification of structural and functional domains in mixed lineage kinase dual leucine zipper-bearing kinase required for complex formation and stress-activated protein kinase activation. *J Biol Chem* 275:7273–7279.
- Russo A, DiAntonio A. 2019. Wnd/DLK is a critical target of FMRP responsible for neurodevelopmental and behavior defects in the *Drosophila* model of fragile X syndrome. *Cell Rep* 28:2581–2593.e5.
- Shin JE, Cho Y, Beirowski B, Milbrandt J, Cavalli V, DiAntonio A. 2012. Dual leucine zipper kinase is required for retrograde injury signaling and axonal regeneration. *Neuron* 74:1015–1022.
- Siu M, Sengupta Ghosh A, Lewcock JW. 2018. Dual Leucine Zipper Kinase Inhibitors for the Treatment of Neurodegeneration. *J Med Chem* 61:8078–8087.
- Valakh V, Walker LJ, Skeath JB, DiAntonio A. 2013. Loss of the spectraplakins short stop activates the DLK injury response pathway in *Drosophila*. *J Neurosci* 33:17863–17873.

Verma S, Khurana S, Vats A, Sahu B, Ganguly NK, Chakraborti P, Gourie-Devi M, Taneja V. 2022. Neuromuscular junction dysfunction in amyotrophic lateral sclerosis. *Mol Neurobiol* 59:1502–1527.

Wlaschin JJ, Donahue C, Gluski J, Osborne JF, Ramos LM, Silberberg H, Le Pichon CE. 2023. Promoting regeneration while blocking cell death preserves motor neuron function in a model of ALS. *Brain* 146:2016–2028.

Xiong X, Collins CA. 2012. A conditioning lesion protects axons from degeneration via the Wallenda/DLK MAP kinase signaling cascade. *J Neurosci* 32:610–615.

Xiong X, Hao Y, Sun K, Li J, Li X, Mishra B, Soppina P, Wu C, Hume RI, Collins CA. 2012. The Highwire ubiquitin ligase promotes axonal degeneration by tuning levels of Nmnat protein. *PLoS Biol* 10:e1001440.

Xiong X, Wang X, Ewanek R, Bhat P, Diantonio A, Collins CA. 2010. Protein turnover of the Wallenda/DLK kinase regulates a retrograde response to axonal injury. *J Cell Biol* 191:211–223.

<https://doi.org/10.7554/eLife.104896.1.sa0>