

Erythrocytosis-inducing PHD2 mutations implicate biological role for N-terminal prolyl-hydroxylation in HIF1 α oxygen-dependent degradation domain

Reviewed Preprint

v1 • June 27, 2025

Not revised

Cassandra C Taber, Wenguang He, Geneviève MC Gasmi-Seabrook, Mia Hubert, Fraser G Ferens, Mitsuhiro Ikura, Jeffery E Lee, Michael Ohh 

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada • Department of Biochemistry, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada • Princess Margaret Cancer Centre, University Health Network, Toronto, Canada • Department of Medical Biophysics, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

In this **valuable** study, Taber et al used a battery of biophysical and structural approaches to characterize the impact of erythrocytosis-related mutations in prolyl hydroxylase domain protein 2 (PHD2). The authors show that PHD2 mutant proteins are destabilized, thus supporting the tenet that dysregulation of PHD2/hypoxia induced factor (HIF) axis underpins erythrocytosis, while providing **incomplete** evidence that N-terminal ODD prolyl hydroxylation of HIF is indispensable for these phenotypes. Notwithstanding that this study was found to be of broad interest for a variety of fields focusing on oxygen sensing in homeostasis and pathological states, resolving inconsistencies in the biophysical analysis (e.g., NMR, SEC, and BLI/MST) was thought to be warranted to further corroborate the proposed model.

<https://doi.org/10.7554/eLife.107121.1.sa4>

Abstract

Mutations in *EGLN1*, the gene encoding for hypoxia-inducible factor (HIF) prolyl-4-hydroxylase 2 (PHD2), cause erythrocytosis and in rare cases the development of neuroendocrine tumors. In the presence of oxygen, PHD2 hydroxylates one or both conserved prolines in the oxygen-dependent degradation domain (ODD) of HIF α subunits, sufficiently marking HIF α for binding and ubiquitylation via the von Hippel-Lindau (VHL) tumor suppressor protein-containing E3 ubiquitin ligase and subsequent degradation by the 26S proteasome. However, prolyl-hydroxylation in the C-terminal ODD appears to be the predominant and sufficient event in triggering the oxygen-dependent destruction of HIF α , rendering the biological significance of N-terminal ODD proline unclear. Here, we examined 7 disease-associated *EGLN1* mutations scattered across the catalytic core and show

definitively that all PHD2 mutants have a structural and/or catalytic activity defect as measured by time-resolved nuclear magnetic resonance. Notably, we identified one of the PHD2 mutants, P317R, to retain comparably wild-type capacity to hydroxylate the predominant proline in the C-terminal ODD but had uniquely compromised ability to hydroxylate the N-terminal ODD proline. These findings support the notion that deregulation of HIF ultimately underlies PHD2-driven erythrocytosis and challenge the currently held uncertainty that the N-terminal ODD prolyl-hydroxylation event is dispensable in normal hypoxic signaling pathway.

Introduction

About half of blood volume is made up of erythrocytes (red blood cells) whose role is to deliver oxygen from the lungs to the tissues in the body¹. Low concentrations of red blood cells result in insufficient oxygen delivery to tissues while high concentrations can induce a propensity towards thrombotic events rendering red blood cell mass regulation crucial. Erythropoietin (EPO) is a hormone produced primarily in the kidneys that stimulates erythropoiesis^{2,3}. Through binding to EPO receptors on erythroid progenitors, EPO can activate a signaling cascade that induces differentiation into erythrocytes⁴. EPO production is regulated in response to physiological oxygen concentrations through hypoxia response elements (HRE). HREs are enhancers that transcription factors called hypoxia-inducible factors (HIFs) bind to and subsequently upregulate the expression of numerous hypoxic response genes including *EPO*⁵. Disrupted EPO regulation can be pathogenic and lead to an autonomous overproduction of erythrocytes causing erythrocytosis⁶. Due to EPO's inherent link to the oxygen response, mutations in the metazoan oxygen-sensing pathway have frequently been found to underlie erythrocytosis.

The oxygen-sensing pathway has been conserved in all animals and is essential for the maintenance of cellular activities in low oxygen settings⁷. Under hypoxia, oxygen-labile HIF1-3 α subunits complex with its constitutively expressed HIF β subunits to translocate to the nucleus as active heterodimeric transcription factors⁸. Following translocation, HIF binds to HREs across the genome to trigger the transactivation of numerous genes such as *GLUT1*, *EPO* and *VEGF* to promote anaerobic metabolism, erythropoiesis and angiogenesis, respectively, to induce adaptation to low oxygen availability⁹. In normal oxygen conditions, HIF prolyl hydroxylases (PHDs) hydroxylate HIF α subunit at one or both of the two conserved proline sites located in the oxygen-dependent degradation domain (ODD), which initiates its ubiquitylation via the von Hippel-Lindau tumor suppressor protein (pVHL)-containing E3 ubiquitin ligase (Elongin BC/Cullin 2-Rbx1/pVHL) and subsequent destruction via the 26S proteasome. Notably, only one proline is required to be hydroxylated to trigger its rapid destruction and whether HIF α is singly or doubly hydroxylated does not appear to change the rate of its ubiquitylation or proteasome-mediated destruction¹⁰.

Moreover, the C-terminally located proline within ODD or CODD (e.g., P564 on HIF1 α) has been shown to be the predominant residue that becomes hydroxylated prior to the modification of proline in the N-terminal ODD or NODD (e.g., P402 on HIF1 α)^{10,12}. The reason for this preferential prolyl-hydroxylation is incompletely understood, and the biological significance or the mechanistic purpose of the N-terminal proline remains unclear.

Clinically observed mutations have been identified in the major components of the oxygen-sensing pathway including *VHL* (pVHL)^{13,14}, *EPAS1* (HIF2 α)^{15–17}, *EGLN1* (PHD2)^{18,19}, and *EGLN2* (PHD1)²⁰. These have been associated with the development of diseases with overlapping phenotypes with erythrocytosis being the most common. Collectively, these diseases are referred to as pseudohypoxic diseases as they involve an inappropriate hypoxic response. Erythrocytosis caused by mutations in *EGLN1* is the most recently discovered pseudohypoxic

disease. Since its first reported case in 2006²¹, 152 cases with 96 distinct mutations have been reported. *EGLN1* mutation-driven erythrocytosis is referred to as erythrocytosis, familial, 3 (ECYT3) by the Online Mendelian Inheritance in Man (OMIM). However, three individuals with distinct *EGLN1* mutations (W51X, A228S and H374R) also developed recurrent pheochromocytoma and paraganglioma (PPGL)^{20,22,23}, which are neuroendocrine tumors that develop from chromaffin cells either within (pheochromocytoma) or outside (paraganglioma) the adrenal gland²⁴.

Of three isozymes of PHD, PHD2 is thought to be the most critical with high expression seen in a variety of cell types²⁵ and the only PHD able to effectively target both prolines in HIF1 α and 2 α ODDs^{11,26,27}. PHD2 belongs to the 2-oxoglutarate-dependent-dioxygenase superfamily²⁸ in which enzymes contain a double-stranded β -helix fold with a highly conserved HXE/D...H motif that coordinates Fe²⁺ at its catalytic site²⁹. In PHD2, the iron coordinating residues are His313, Asp315 and His374³⁰. Erythrocytosis-causing mutations at (H374R) and near these conserved catalytic residues have been shown to be defective at regulating HIF α in a cellular context^{18,21,23}, which appears to be a common mechanism observed in the other pseudohypoxic diseases. For example, numerous studies have revealed that pVHL mutants have varied abilities to recognize and regulate HIF α leading to the development of PPGL, erythrocytosis and other cancers such as clear-cell renal cell carcinoma and hemangioblastoma^{31–33}. Recently, we showed that phenotypic severity of Pacak-Zhuang syndrome, caused by *EPAS1* mutations, is correlated to the ability of HIF2 α mutants to escape degradation mediated by pVHL and PHD2^{15,34}.

Together, these observations suggest that disproportionately high levels of HIF α are responsible for pseudohypoxic disease phenotypes such as erythrocytosis and PPGL. In keeping with this notion, disease-causing mutants of PHD2 are likely defective in catalyzing HIF α hydroxylation to some extent, resulting in less HIF α degradation and correspondingly increased hypoxia response signaling. However, the precise mechanism(s) underlying such defect in PHD2 function is unclear. Recently, we devised a time-resolved nuclear magnetic resonance (NMR) assay to directly measure the rate of hydroxylation on the two conserved proline residues simultaneously in HIF1 α ODD¹². Here, we examined 7 disease-associated PHD2 mutants via biophysical analyses and show that all PHD2 mutants have structural and/or catalytic activity defects. While all mutants had the anticipated defect in binding and hydroxylating the predominant CODD proline, we identified PHD2 P317R mutant that retained comparably wild-type level of hydroxylating CODD proline (i.e., HIF1 α P564) but failed to hydroxylate NODD proline (i.e., HIF1 α P402) as measured by NMR. Unexpectedly, PHD2 P317R was capable of binding to both NODD and CODD peptides with similar affinity. These results provide a biophysical mechanism that, at least in part, underlies PHD2-driven erythrocytosis, and serendipitously reveal a direct biochemical evidence to suggest that NODD prolyl-hydroxylation is necessary for normal physiological response to hypoxia.

Results

Mutations across the PHD2 enzyme induce erythrocytosis

We compiled and analyzed all reported cases of PHD2-driven erythrocytosis as of January 2025. 96 distinct variants have been reported, including two missense variants commonly found in Tibetan populations that are hypothesized to be adaptations to high altitude conditions (D4E and C127S)^{35,36}. Consistent with erythrocytosis in the general population, most patients with PHD2-driven erythrocytosis were male (~70%). About half of the case reports provided information on family history of PHD2-driven erythrocytosis, of which ~75% were reported to be familial while ~25% were *de novo* mutations; a similar rate of *de novo* mutations is observed with VHL disease. PHD2-driven erythrocytosis is typically classified as secondary erythrocytosis, which is defined as being caused by an EPO-inducing mechanism downstream of the primary defect,

ultimately resulting in normal to high EPO levels that drive erythropoiesis. This classification is aligned with the suspected mechanism of PHD2-driven disease in which mutated PHD2 dysregulates HIF leading to increased EPO production. Curiously however, EPO was reported to be largely normal in patients with PHD2-driven erythrocytosis with few exceptions (**Supplementary Data**), which is consistent with the loss of PHD2 in mice leading to EPO hypersensitivity in erythroid progenitors, enabling a ‘normal’ amount of EPO to generate a pathogenic response³⁷.

All cases of PHD2-driven erythrocytosis were reported as germline heterozygous mutations of PHD2, suggesting haploinsufficiency of PHD2 is enough to induce erythrocytosis. This is consistent with previous reports that HIF α regulation is dose dependent on PHD2 knockdown via siRNA²⁵. As stated previously, in three exceptional cases, individuals with PHD2 mutations developed PPGL (W51X, A228S and H374R). Biopsies of each tumor revealed a loss of heterozygosity for PHD2, implying a complete loss of wild-type PHD2 in chromaffin cells is necessary to induce tumorigenesis.

The majority of the reported mutations were missense (~74%) (**Figure 1A**). Interestingly, mutations that should theoretically severely disrupt PHD2 structure (nonsense, frameshift and deletions) caused the same phenotype (erythrocytosis) as less disruptive missense mutations, suggesting an unclear or lack of correlation between mutation type and phenotypic severity. Thus, a potential phenotypic mechanism of PHD2-driven disease can be hypothesized where erythrocytosis will develop upon deleterious mutation in one PHD2 allele while PPGL will be induced upon a loss of heterozygosity in chromaffin cells.

Mutations are distributed across all exons of PHD2 with some clustering across the catalytic core (**Figure 1B**). When mapped onto the three-dimensional crystal structure of the PHD2 catalytic core, which we and others have previously solved^{34,38,39}, these mutations do not seem to display a specific pattern (**Figure 1C**). Thus, we selected 7 PHD2 mutants with distinct residue changes in the catalytic core of PHD2 for study.

Some PHD2 mutants are less stable and defective in regulating HRE-driven transcription

We asked whether the enzymatic activity of PHD2 mutants can be inferred from downstream HIF activity using a dual luciferase reporter assay under control of HRE. Notably, we first generated CRISPR/Cas9-mediated PHD2 knockout (-/-) HEK293A cells to minimize the impact of endogenous PHD2 on the prolyl-hydroxylation of HIF α . As expected, wild-type (WT) ectopic PHD2 markedly reduced the HRE-luciferase activity driven by ectopic HIF1 α compared to cells without ectopic PHD2 expression (**Figure 2A**). Two disease-associated PHD2 mutants, P317R and H374R, displayed significantly reduced ability to knockdown luciferase activity (~2-fold and ~2.5-fold, respectively) compared to PHD2 WT. Notably, the other mutants showed comparable activity to PHD2 WT in down-regulating HRE-driven luciferase signal under these *in cellulo* experimental conditions (**Figure 2A**).

We next performed cycloheximide (CHX) chase assay to determine the protein stability of PHD2 mutants. We observed that mutants G206C, W334R, F366L, R371H, and H374R showed noticeable instability by 24 hours (**Figure 2B and C**). These results suggest that while most PHD mutants showed diminished protein stability in comparison to PHD2 WT, the stability of PHD2 mutants cannot alone explain the impact of these disease-associated mutations on HIF1 α -driven HRE-luciferase activity in cells.




Figure 1.

Analysis of disease causing PHD2 mutants.

(A) Distribution of mutation type calculated from clinical case reports. (B) Linear map of mutant location and frequency on PHD2. The zinc finger is comprised of residues 21-58 while the catalytic core ranges from 181-426. Each dot represents a clinical report of a disease-causing mutation at the given residue. (C) Structure of PHD2 HIF1αCDD complex (PDB:5L9B) with PHD2 mutant locations highlighted (yellow). PHD2 (grey) and HIF1αCDD (red) are depicted as ribbons. PHD2 mutants selected for analysis are highlighted in green and labelled. The structure was turned 260° on the y-axis to highlight all mutant locations.




Figure 2.

Methods in HEK293A cells detect defects in some PHD2 mutants but not all.

(A) Dual luciferase reporter assays were performed to measure HIF1 α transcriptional activity in the presence of PHD2 mutants in PHD2 $-/-$ HEK293A cells. Individual data points are plotted. Loading accuracy was evaluated via immunoblotting for FLAG-tagged PHD2 and vinculin. (B) PHD2 mutant stability was measured via cycloheximide chase assay. HEK293A cells were transfected with wild type or mutant PHD2 constructs, with the amount of transfected adjusted to ensure equal expression at 0 hours. After 24 hours, the transfected cells were treated with CHX to halt protein production and monitor the stability of PHD2. Cells were harvested at various time points up to 24 hours and lysates were immunoblotted to measure PHD2 levels. (C) FLAG immunoblot density was quantified at each time point and normalized with vinculin density to yield a relative density. For A and C, bars represent mean values, and standard error is represented by error bars. * indicates $P < 0.0332$, ** indicates $P < 0.0021$, *** indicates $P < 0.0002$, and **** indicates $P < 0.0001$ (two-tailed t-test).

PHD2 mutants are thermally unstable and some are prone to aggregation

We asked if the disease-causing mutations compromised other biophysical characteristics of PHD2. We purified catalytic cores of PHD2 harboring the selected mutations for the subsequent biophysical studies. Size exclusion chromatography (SEC) was performed as a polishing step and chromatograms were acquired with monomeric PHD2 eluting from the column around 17 mL (**Figure 3A**). Mutants were eluted from the column at different volumes ranging from 16.9 mL (G206C) to 17.9 mL (P317R), while WT PHD2 was eluted consistently at 17 mL. These results suggest that mutations may affect the hydrodynamic size of PHD2, hinting to mutation-induced structural alteration. Moreover, PHD2 G206C, W334R and H374R aggregated severely in SEC and throughout the purification process and could not be purified.

We next subjected the remaining four mutants (A228S, P317R, F366L, and R371H) in comparison to WT PHD2 to circular dichroism (CD) to determine structural abnormalities. The CD spectra of the mutant PHD2s did not display appreciable differences from WT PHD2 (**Figure 3B**), indicating that these PHD2 mutants' overall structure is mostly unaffected.

To further probe the stability of PHD2 mutants, molar ellipticity was measured at 220 nm from 25 to 95 °C to determine melting temperature (T_m). Alpha helices produce a strong minimum at 220 nm so this wavelength was selected to measure thermal protein unfolding⁴⁰. Melting curves were created by setting the peak value of each spectrum at 25 °C to 1.0. The data were then fitted with a sigmoidal curve to produce melting curves and calculate T_m 's (**Figure 3C** and **Table 1**). WT PHD2's T_m , estimated as the temperature at which half of the proteins are unfolded, was determined to be 52.3°C. All four mutants had lower T_m 's than WT, suggesting a degree of thermal instability. Although it is reasonable to assume that instability would result in less PHD2, it is unclear if the unstable enzyme is active prior to degradation and whether the remaining portion of mutant PHD2s would be sufficient to bind and hydroxylate HIF α .

PHD2 mutants have weaker binding to HIF α than WT PHD2

Assuming intact catalytic function, the strength of PHD2 binding affinity to HIF α should translate to the extent of prolyl-hydroxylation of HIF α via PHD2. We performed microscale thermophoresis (MST), which utilizes the motion of molecules in a temperature gradient to determine dissociation constants (K_d), to measure the binding deficiencies, if any, between PHD2 mutants and HIF α . In addition to being highly sensitive, MST is also advantageous for measuring protein-ligand binding in an untethered setting. We showed previously that this technique was effective at distinguishing between disease classes of HIF2 α mutants in Pacak-Zhuang syndrome by measuring their binding affinities to WT PHD2³⁴. We performed a similar protocol with PHD2 mutants to identify potential binding defects with HIF1 α and 2 α peptides of the C-terminal oxygen-dependent degradation domain (C-ODD) that contains single target proline, which is thought to be predominant over the N-terminal oxygen-dependent degradation domain (N-ODD)¹⁰. PHD2 WT bound tightly to HIF1 α C-ODD and HIF2 α C-ODD peptides (7.5 and 15 μ M, respectively). Notably, all but one tested PHD2 mutants (A228S, F366L and R371H) displayed mild binding defects, up to approximately 2-fold, to HIF1 α C-ODD in comparison to WT PHD2 while P317R showed a severe binding defect with high K_d value (320 μ M) (**Figure 4A** and **Table 2**). Similar pattern of binding deficiencies was noted for PHD2 mutants to HIF2 α C-ODD (**Figure 4B** and **Table 2**), suggesting no significant preferential binding defects to either HIF α paralog.

Figure 3.

PHD2 mutants display instability through aggregation and decreased thermostability.

(A) Chromatograms of PHD2 catalytic cores were acquired via SEC on a Superdex200 column. Curves have been normalized according to their area under curves. Dashed lines indicate mutants that were not purified. (B) CD was performed on PHD2 mutants to predict secondary structure variations. The far-UV spectra (190-260 nm) of the purified catalytic cores was measured and converted to molar ellipticity. (C) Molar ellipticity of PHD2 mutants was monitored at 220 nm from 25 to 95 °C to evaluate thermal stability. (D) Melting curves were generated from the CD melt. Molar ellipticity values were normalized and transformed into a fraction of folded protein and fitted with a sigmoidal curve.




Table 1.

PHD2 mutant melting temperatures calculated via CD.

	T_m (°C)	R^2
WT	51.9	0.871
A228S	48.3	0.707
P317R	46.2	0.748
F366L	43.8	0.878
R371H	45.1	0.843




Figure 4.

PHD2 mutants have minor binding defects to HIF1α peptides.

Microscale thermophoresis was performed on fluorescently labelled PHD2 and HIF1α 555-574 CDD (A) or HIF2α 522-542 CDD (B) peptides. PHD2 P317R displayed a severe binding defect whereas the other three mutants had minor binding defects. It is suspected that amine reactive fluorescent labelling induced a binding defect on PHD2 P317R.

PHD2	K _d (μM)	
	HIF1α	HIF2α
WT	7.5 ± 0.52	15 ± 1.1
A228S	9.3 ± 0.57	22 ± 1.6
R371H	13 ± 1.6	39 ± 4.4
F366L	16 ± 1.8	43 ± 4.7
P317R	320 ± 20	580 ± 38

Table 2.

MST determined dissociation constants between PHD2 mutants and HIFα peptides.

NMR reveals major defect in PHD2 P317R-mediated hydroxylation of NODD proline

Previously, we developed an assay utilizing NMR to directly measure PHD2 hydroxylation of the critical two prolines located within the intrinsically disordered ODD of HIF1 α ¹². Under the catalysis of PHD2, movement of resonances corresponding to the target prolines as well as adjacent residues are observed, reflecting local conformational changes due to the post-translational modification of HIF1 α ODD. The time-dependent shift in peak intensity from unhydroxylated to hydroxylated species can be interpreted as a function of the reaction progress. Notably, the enzymatic kinetics of the hydroxylation of both target prolines (P402 in NODD, P564 in CODD) can be measured simultaneously over time. Since binding experiments with PHD2 mutants did not implicate marked preferential defects for either HIF1 α or HIF2 α , these assays were performed with HIF1 α ODD of which more than 95% of amino acids have been assigned to their corresponding 3D NMR backbone cross-peak¹².

We selected PHD2 A228S for NMR study to determine if this mutant, which showed negligible to minute defect in the above studies from HRE-driven luciferase assay to MST based binding experiments, exhibited any appreciable enzymatic defect in hydroxylating P564 or P402. We included as controls PHD2 WT and P317R that showed major defect in binding HIF1 α CODD peptide and transcriptional activation of HRE-luciferase. As shown previously¹², PHD2 WT showed robust and rapid hydroxylation of P564 in CODD and a markedly slower hydroxylation of P402 in NODD (**Figure 5A**). PHD2 A228S mutant effectively hydroxylated P564 faster than P402 with negligible differences in the enzymatic rate to PHD2 WT (**Figure 5A**). Unexpectedly, P317R displayed a near wild-type hydroxylation of P564 but failed to hydroxylate P402 (**Figure 5A**).

Superimposition of ^1H - ^{15}N HSQC spectra showed that cross-peak of P564 shifts with PHD2 WT, A228S, and P317R mutant, but P402 only shifts when incubated with PHD2 WT and A228S, not PHD2 P317R (**Figure 5B**). These results are inconsistent with observations made via MST that showed P317R having a severe binding defect for HIF1 α CODD peptide. Furthermore, it is of particular interest to investigate whether the lack of activity of PHD2 P317R on NODD is due to a loss of binding interaction.

PHD2 P317R binds effectively to both CODD and NODD via BLI

We performed bio-layer interferometry (BLI) using immobilized biotinylated HIF1 α CODD and NODD peptides with purified PHD2 WT and P317R, which showed P317R binding to CODD peptide only slightly weaker than PHD2 WT (**Table 3**). The discrepancy between the MST and BLI results was likely due to the amine-linked fluorescent labelling of PHD2 P317R for MST where the fluorescent label bound to the newly available amine residue on the mutant arginine, thereby disrupting the binding to HIF1 α CODD peptide. The removal of this label allowed for proper, unimpeded interaction to occur as observed with BLI. These results are consistent with the near-WT, efficient hydroxylation of HIF1 α CODD by PHD2 P317R as revealed by NMR (See **Figure 5C**).

The binding affinity between PHD2 P317R and HIF1 α NODD was also measured to examine whether the catalytic defect observed via NMR would manifest as a binding defect. However, PHD2 P317R bound to NODD similarly to PHD2 WT suggesting that its failure to hydroxylate P402 is unlikely attributable to binding.




Figure 5.

PHD2 P317R does not hydroxylate HIF1 α ODD, while PHD2 A228S has very minor enzymatic defects.

An assay measuring hydroxylation of HIF1 α ODD (394-574) by PHD2 via NMR was performed. (A) Resonance shifting was monitored in real time to compare hydroxylation rates of A228S (blue), P317R (red) and WT (black) PHD2. PHD2 A228S displayed minorly impaired hydroxylation of both ODDs. PHD2 P317R displayed no activity on the P402 (NODD), while retaining near WT activity on P564 (CODD). (B) HSQC spectra display the resonance shifting pattern of HIF1 α ODD upon prolyl hydroxylation catalyzed by PHD2 (WT, P317R, A228S) over the course of 20.2 h. A403 and I566 were used to monitor hydroxylation of P402 and P564 respectively. Spectra recorded at 0 h is shown in black while spectra recorded at the endpoint (20.2 h) is shown in red.

	K_d (μ M)	k_a ($\times 10^4$) (1/Ms)	k_d ($\times 10^{-3}$) (1/s)
PHD2 WT + HIF1 α CODD	2.1 $\pm$ 0.8	5.2 $\pm$ 1.4	101 $\pm$ 20
PHD2 P317R + HIF1 α CODD	6.0 $\pm$ 0.7	0.21 $\pm$ 0.11	12.3 $\pm$ 5
PHD2 WT + HIF1 α NODD	2.0 $\pm$ 0.8	0.26 $\pm$ 0.093	4.7 $\pm$ 0.2
PHD2 P317R + HIF1 α NODD	2.0 $\pm$ 1.0	0.17 $\pm$ 0.049	3.0 $\pm$ 0.7

Table 3.

BLI determined binding constants between PHD2 WT and P317R and HIF α 1 NODD and CODD peptides.

Discussion

Mutations occurring in the core components of the hypoxia sensing pathway have been shown to cause various phenotypes ranging from erythrocytosis and neuroendocrine tumors to hemangioblastoma and renal cell carcinoma. Collectively, these diseases are referred to as pseudohypoxic diseases due to the inappropriate hypoxic response induced by the mutations. It is presumed that the abnormal response is the reason for overlapping phenotypes caused by mutations in different proteins of the same pathway (pVHL, HIF2 α and PHD2). Specifically, we have proposed that the common phenotypes of pseudohypoxic diseases are caused by the gain-of-function ability of HIF α , in particular HIF2 α , to escape degradation in the presence of the disease-causing mutants³¹. It has been shown that VHL disease-causing mutants have varying ability to target HIF1/2 α for ubiquitin-mediated destruction^{32,41} and Pacak-Zhuang syndrome-causing HIF2 α mutants are able to escape recognition to a varying degree from both pVHL and PHD2^{15,34}. It follows then that in the case of PHD2-driven erythrocytosis, PHD2 mutants would have a hindered ability to promote proper oxygen-dependent hydroxylation of HIF1 α and 2 α .

Complete deletion of PHD2 in mice causes embryonic lethality due to venous congestion and cardiomyopathy, theorized to be directly related to high blood volume induced upon PHD2 loss⁴². Conditional inactivation of PHD2 whether globally⁴³ or in renal erythropoietin-producing cells⁴⁴ has been observed to cause erythrocytosis in mice. Moreover, while inactivation of PHD2 alone was sufficient to induce erythrocytosis, inactivating PHD1 or PHD3 alone did not result in erythrocytosis. However, when PHD1 or PHD3 were inactivated in addition to PHD2, erythrocytosis phenotype worsened, suggesting a supporting role for the PHD paralogs⁴⁵.

Here, we examined 7 disease-causing PHD2 mutations across the catalytic core, the most frequently mutated region within PHD2. G206C, W334R and H374R exhibited gross protein stability defects and were prone to severe aggregation, which precluded their purification necessary for further biophysical analyses. One of these mutants, H374R, is located at the catalytic site integral to the function of PHD2, which combined with the stability defect likely abrogated its ability to promote the oxygen-dependent hydroxylation of ectopic HIF1 α , leading to increased HRE-luciferase transcriptional activity in *PHD2*^{-/-} HEK293A cells. Previous work has shown mutating analogous His iron-coordinating residues to a charged residue (Arg or Glu) in the prolyl-4-hydroxylase responsible for hydroxylating prolines of collagen abolishes enzymatic activity⁴⁶. Given the high structural conservation in the Fe/2OG-oxygenase enzyme family²⁹, it can be predicted that a comparable mutation on PHD2 would also abolish activity. However, PHD2 G206C and W334R mutants showed similar capacity as the PHD2 WT to inhibit the transcriptional activity of ectopic HIF1 α . This may have been due to the nature of transfected PHD2 in which mutant PHD2s that are prone to aggregation and instability are being translated throughout the luciferase assay, which may generate a small pool of monomeric PHDs that is sufficient to inhibit the transcriptional activity of HIF1 α comparable to PHD2 WT.

The remaining four mutants (A228S, P317R, F366L, and R371H) not prone to aggregation displayed an overall structure similar to PHD2 WT, as determined by CD, and with the exception of PHD2 P317R (discussed later), were able to effectively suppress the ectopic HIF1 α transcriptional activity. These mutants also exhibited slight thermal instability with mildly weaker binding affinity to HIF1 α CODD peptide, which may be sufficient to cause the disease phenotype *in vivo*. PHD2 A228S exhibited the most minute of defects amongst all mutants tested and concordantly showed marginal reduction in enzymatic activity as measured via time-resolved NMR. Notably, despite

these defects, albeit subtle, it is evident that the HRE-luciferase reporter assay is in general not reliable to reveal the disease-causing mutations on the functional output of PHD2 mutants, including those prone to aggregation (G206C and W334R).

We recently developed a time-resolved NMR approach to directly and simultaneously measure PHD2-mediated hydroxylation of P564 within CODD and P402 within NODD in the context of full-length HIF1 α ODD. We showed that P564 is the predominant oxygen-dependent hydroxylation site over P402. Considering that pVHL only requires one hydroxylated proline for binding and triggering rapid ubiquitin-mediated proteasomal degradation of HIF1 α ^{10,12,47}, the biological utility of P402 was thought to be inconsequential in normal hypoxic response. Here, our results revealed a previously unappreciated intricacy in the enzymatic selectivity of PHD2 P317R between CODD and NODD, challenging the current notion of NODD insignificance. NMR results showed that the disease-causing PHD2 P317R retained near-WT activity on CODD P564 but had negligible activity on NODD P402. Furthermore, despite effective activity on P564, ectopic PHD2 P317R generated abnormally high HRE-driven luciferase reporter signal, suggesting the negative impact of defective P402 hydroxylation on an otherwise rapid oxygen-dependent degradation of HIF1 α . This is congruous with Schofield's earlier work that showed, using an indirect metabolic method with individual peptides, no activity on NODD by PHD2 P317R while effectively hydroxylating CODD³⁸. Given the pathogenic nature of P317R mutation, it appears likely that this selective defect on HIF1 α NODD contributes to the development of erythrocytosis and therefore, the involvement of NODD in normal oxygen-sensing pathway is not insignificant.

The molecular basis for PHD2 P317R's selective defect for NODD is currently unknown. Intriguingly, BLI results showed that PHD2 P317R binds to both CODD and NODD peptides comparable to PHD2 WT. Substrates of Fe/2OG-oxygenases typically bind the enzymes non-covalently with a prime separation distance of 4-5 Å between the target atom and the catalytic iron⁴⁸. One possibility is that the Pro to Arg substitution at position 317 may increase the distance between HIF1 α P402 and the active site within the catalytic core without compromising the interaction between the substrate and enzyme. However, such mutation did not markedly impact the efficient turnover of P564, suggesting that the varied residues between NODD and CODD may influence the substrate selectivity of PHD2.

PHD2-driven erythrocytosis is a relatively newly discovered genetic disease with the first reported case in 2006. Since then, there has been a large increase in reports owing to the addition of *EGLN1* mutant screening in recent years for cases of idiopathic erythrocytosis³². As further awareness of the disease grows, screening will increase, leading to the likely identification of more disease-associated mutations. This study, therefore, will add to the groundwork knowledge of how mutant PHD2 induces disease in the context of the hypoxic response.

Materials and methods

Plasmids, antibodies, and peptides

Previously described plasmids were used as follows: pET-46-HIS₆-PHD2¹⁵, pGL3-VEGFA-HRE³⁴, pCDF-Ek/LIC-StrepII-HIS₆-HIF1 α ODD¹², pcDNA3-HA-HIF1 α ¹⁰, pRL-SV40 was procured from Promega. pcDNA3-3XFLAG-EGLN1 was generated by amplifying the EGLN1 gene from HA-EglN1-pcDNA3 (a gift from William Kaelin; Addgene plasmid # 18963) using primers 5' – GCG GCG GGA TCC ATG GCC AAT GAC AGC G – 3' and 5' – GCG GCG TCT AGA CTA GAA GAC GTC TTT ACC GAC – 3'. The product was digested with BamHI/XbaI and subcloned into pcDNA3 vector containing an N-terminal 3XFLAG tag. pX330-U6-Chimeric_CBh-hSpCas9 was gifted to the lab from Feng Zhang (Addgene plasmid #42230). PHD2 mutants of pET-46-HIS₆-PHD2 and pcDNA3-3XFLAG-PHD2 were generated through QuikChange site directed mutagenesis (Agilent). Mutant sequences were confirmed via DNA sequencing. The following antibodies were used: α -HA (C29F4) from Cell

Signaling, α -FLAG M2 (F1804) from Sigma-Aldrich, and α -PHD2/EGLN1 (D31E11) from Cell Signaling. Primary antibodies were diluted in TBS-T (20 mM Tris pH 7.6, 137 mM NaCl, 0.05% (v/v) Tween 20) with 0.02% (w/v) sodium azide for immunoblotting. Peptides were N-terminally acetylated for microscale thermophoresis (MST) and biotinylated for bio-layer interferometry (BLI). All peptides were amidated at the C-terminus. For microscale thermophoresis, peptides were reconstituted in 50mM Tris (tris(hydroxymethyl)aminomethane) pH 8.0 buffer. For bio-layer interferometry, peptides were reconstituted in 100% (v/v) DMSO.

PHD2 $-/-$ HEK293A Cells

Wild-type (WT) human epithelial kidney (HEK293A) cells were subjected to the clustered regularly interspaces short palindromic repeats (CRISPR) method to generate homozygous PHD2 knockout ($-/-$) cells. The CRISPOR design tool was used to generate CRISPR guide RNAs (gRNA) targeting exon 1 in *EGLN1*⁴⁹. Forward and reverse gRNA oligonucleotides were phosphorylated and annealed before ligation into *BbsI*-digested pX330-U6-Chimeric_CBh-hSpCas9. WT HEK293A cells were transfected with pX330-EGLN1 gRNA using Lipofectamine 2000 according to manufacturer instructions. Single cell colonies were seeded into 96-well plates from the previously transfected cells at a density of 1.25 cells/mL. 200 μ l of cells were seeded into each well resulting in a density of $\sim$ 0.25 cells/well. Single cell clones were screened for PHD2 knockout via immunoblotting for PHD2.

Dual Luciferase Assay

A dual luciferase assay was performed according to a modified previously reported protocol³⁴. PHD2 $-/-$ HEK293A cells were grown and maintained in Dulbecco's Modified Eagle Medium (DMEM, Wisent) with 10% (v/v) Fetal Bovine Serum (FBS, Wisent) and 100 IU/mL penicillin and streptomycin in a humidified incubator at 37°C and 5% CO₂. Transfection complexes consisted of 2 μ g total DNA: 0.4 μ g pcDNA3-HA, 0.9 μ g pGL3-VEGF α -HRE, 0.1 μ g pRL-SV40, 0.4 μ g pcDNA3-HA-HIF1 α , and 0.2 μ g pcDNA3-FLAG3X-PHD2 (WT or mutant). Complexes were incubated for 15 minutes at room temperature with 8 μ g polyethylenimine pH 7.2 (Polysciences) in 400 μ l of OptiMEMTM Reduced Serum Media (Gibco). Following incubation, transfection complexes were added to 5 x 10⁵ PHD2 $-/-$ HEK293A cells suspended in 400 μ l OptiMEMTM and incubated for 5 minutes. Cells were transferred to six well plates and incubated at 37°C, 5% CO₂ for 24 hours. Cells were lysed and processed according to the Promega Dual-Luciferase Reporter Assay System (#E1960) instructions. Firefly luciferase and Renilla luciferase activity were measured on a Varioskan Lux microplate reader (Thermo Fisher Scientific). Luciferase activity was normalized by dividing Firefly RLU values by the constitutively expressed, control Renilla RLU values. Normalized RLU values were transformed by setting the RLU value of the WT PHD2 sample to 1 and dividing the other sample values by the same transformation factor. A two-tailed t-test was performed to determine statistically significant differences between samples. A p-value below 0.0332 was considered significant. Protein levels of PHD2 were determined by immunoblotting for FLAG3X-PHD2. Vinculin was probed as a loading control.

Cycloheximide Chase Assay

HEK293A cells were cultured as described above. Transfection complexes consisting of 1.5 μ g total DNA were incubated with 6 μ g Polyethylenimine pH 7.2 in 400 μ l of OptiMEMTM for 15 min at room temperature. Transfection complexes consisted of either 0.5 μ g (WT, A228S, P317R, F366L, R371H) or 1.5 μ g (G206C, W334R, H374R) of pcDNA3-FLAG3X-PHD2. Additional pcDNA3-FLAG3X-PHD2 was transfected for less stable mutants to ensure equal amounts of PHD2 expression. The total amount of DNA was brought to 1.5 μ g with the addition of 1 μ g of pcDNA3 if necessary.

Transfection complexes were added to 1.5 x 10⁵ cells suspended in 400 μ l OptiMEMTM and incubated for 5 minutes. Cells were plated in 60 mm plates with 5 plates for each mutant and 2 for WT. Cells were incubated for 24 hours at 37°C and 5% CO₂. Following incubation, media was

replaced to remove Polyethyleneimine and fresh media with 10 µg/ml cycloheximide (CHX) was added. Cells were incubated in CHX supplemented media for 0, 3, 6, 12 and 24 hours. WT samples were incubated only for 0 and 24 hours. Cells were harvested through manual scraping and lysed via sonication in 250 µl of cold EBC lysis buffer (50 mM Tris, 120 mM NaCl, 0.5% (v/v) NP-40) supplemented with 1X protease inhibitor cocktail (BioShop PIC002.1). Protein levels were determined via Bradford assay and equal protein amounts of lysate were denatured by boiling for 5 minutes with 3X sample buffer (62.5 mM Tris, 10% (v/v) glycerol, 2% (w/v) SDS, 0.01% (w/v) bromophenol blue). Samples were then resolved on a 10% acrylamide SDS-PAGE gel for 90 minutes at 120 V. Proteins were transferred to polyvinylidene fluoride (PVDF) membrane via wet electrophoretic transfer for 70 minutes at 110 V. Membranes were blocked for 60 minutes at room temperature in 5% (w/v) skim milk in TBS-T. Membranes were then incubated with α-FLAG-M2 (1:1500) and α-vinculin (1:1500) at 4°C overnight. Following three washes with TBS-T, membranes were incubated with HRP-conjugated secondary antibodies (1:10,000 in 5% (v/v) skim milk in TBS-T) for 60 minutes at room temperature. Proteins were detected using a chemiluminescence solution and imaged with a ChemiDoc™ MP Imaging System (BioRad). Immunoblot band density was quantified using BioRad Image Lab densitometry software. FLAG3X-PHD2 levels were normalized according to corresponding vinculin levels. Each 0-hour sample was set to 1 and each other time point was divided by this value to reflect relative protein levels. A two-tailed t-test was performed to identify significant differences between 0 hour and 24-hour timepoints. A p-value below 0.0332 was considered significant.

PHD2 (181-426) Expression and Purification

BL21 (DE3) *E. coli* cells (Novagen, #69450) were transformed with pET-46-HIS₆-PHD2 (wild type or mutant). 1 liter Luria Broth (LB) cell cultures were grown at 37°C until OD₆₀₀ reached ~0.8. Cultures were then induced with a final concentration of 1 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG) for 16 hours at 16°C. Cultures were harvested via centrifugation and stored at -80°C for future use. His₆-PHD2 (181-426) was purified according to a previously described protocol⁵⁰. Bacterial pellets were resuspended in lysis buffer (50 mM Tris-HCl pH 7.9, 500 mM NaCl, 5 mM imidazole) supplemented with 1X SigmaFast protease inhibitor cocktail (Sigma-Aldrich). Cells were lysed with 3 passes through a cell disruptor at 30 kpsi. Cell lysate was cleared via centrifugation at 30,000 g for 45 minutes. Cleared lysates were applied to a 2 mL Ni-NTA resin (ThermoFisher). The resin was washed with 10 mL of lysis buffer followed by 2 separate 30 mL washes with Wash buffer (50 mM Tris-HCl pH 7.9, 500 mM NaCl, 30 mM imidazole). His₆-PHD2 was eluted from the resin with 10 mL of elution buffer (50 mM Tris-HCl pH 7.9, 500 mM NaCl, 1 M imidazole). The eluate was dialyzed against 2 liters of dialysis buffer (50 mM Tris-HCl pH 7.5) using a 10 kDa cut-off dialysis membrane. After 4 hours, the dialysis buffer was replaced with 2 fresh liters and 1.5 units of thrombin was added per mg of His₆-PHD2. His₆-PHD2 was dialyzed overnight with thrombin at 4°C. The dialyzed PHD2 was applied to Ni-NTA resin equilibrated with dialysis buffer to separate thrombin-cleaved protein from uncleaved protein and His₆ tags. The resin was washed with dialysis buffer supplemented with 5 mM imidazole. The flowthrough and wash were combined and concentrated to ~600 µl using an Amicon 10 kDa cut-off centrifugal concentrator (Millipore). The concentrated PHD2 was cleared at 13,000 rpm for 10 minutes then applied to a Superdex™ 200 Increase 10/300 (Cytiva) column equilibrated with dialysis buffer or labeling buffer (50 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid) pH 7.5, 150 mM NaCl). Elution was monitored via UV absorbance and samples with PHD2 were collected. SDS-PAGE gels were run to confirm purity. PHD2 proteins were concentrated to > 2 mg/ml. The final concentration was measured via UV absorbance at 280 nm ($\epsilon_{0.1\%} = 1.34$). Unlabeled protein was flash frozen and stored at -80°C for use in circular dichroism and nuclear magnetic resonance.

PHD2 proteins used for microscale thermophoresis (MST) were labelled with Alexa Fluor-647 N-hydroxysuccinimide ester dye (Thermo Scientific) following purification according to a previously reported protocol³⁴. PHD2 proteins were diluted to 1 mg/ml and Alexa Fluor-647 dye was added to 500 µl of protein in a 4:1 molar ratio (dye: PHD2). Following incubation at room temperature in the dark for 1-hour, excess dye was removed using a PD-10 Sephadex™ G-25 M desalting column

(Cytiva) equilibrated with dialysis buffer. Labelled protein subjected to a 10-fold dilution in dialysis buffer and was then concentrated using a 10 kDa cut-off Amicon centrifugal concentrator to remove excess dye. Alexa Fluor-647 labelled PHD2 was then brought to 10 mM in dialysis buffer, flash frozen and stored at -80°C for use in MST.

HIF1 α ODD (394-574) Expression and Purification

DNA corresponding to the sequence of human HIF-1 α ODD (residue 394-574) with a C-terminal Strep Tag II (WSHPQFEK) was cloned into a pCDF Ek/LIC vector (EMD Millipore). Thrombin cleavage sites were included after the N-terminal-His₆ tag and before the C-terminal Strep Tag II. HIF-1 α ODD was expressed in Rosetta-2 (DE3) cells. To produce ^{15}N -labeled proteins, cells were grown in 2X M9 minimal media, supplemented with a final concentration of 1 g/L $^{15}\text{NH}_4\text{Cl}$ (ACP chemicals) and 4 g/L D-glucose (Bioshop). Expression and purification of HIF-1 α ODD was previously described¹². Briefly, cleared lysate was applied to Ni-NTA resin (ThermoFisher) and eluted with 300 mM imidazole. The elution was then incubated with StrepTactin XT 4Flow resin (IBA Lifesciences) overnight. Protein was eluted from the StrepTactin resin with 50 mM biotin. This process was repeated with the flowthrough from the StrepTactin beads to capture remaining unbound protein. Protein was concentrated and run on a Bio-Rad SEC650 10/300 column. Fractions containing HIF-1 α ODD were then run on a HiResQ 5/50 (Cytiva) anion exchange column. After tag cleavage with thrombin, purified HIF-1 α ODD was concentrated to 0.9 mM and stored at -80°C for future use.

Circular Dichroism

Purified PHD2 proteins were buffer exchanged on PD-10 SephadexTM G-25 M desalting columns (Cytiva) for 5 mM $\text{KH}_2\text{PO}_4/\text{K}_2\text{HPO}_4$ pH 7.4 buffer. Proteins were then diluted to 0.1 mg/ml. Far-UV wavelength scans were collected from 190-260 nm on a Jasco-J-1500 CD spectrophotometer with a 1-mm pathlength quartz cuvette (Helma). CD values were averaged from 20 accumulations, baseline buffer subtracted and converted into mean residue ellipticity (degrees $\text{cm}^2 \text{dmol}^{-1}$). Spectra were smoothed using Jasco Spectra Analysis software. Wavelength scans were run in triplicate and the average mean residue ellipticity was plotted with other PHD2 CD curves.

Melting curves were generated by monitoring CD values for each PHD2 protein from $25-95^{\circ}\text{C}$ at 220 nm with 1°C steps. Values were baseline buffer subtracted and converted to mean residue ellipticity. Triplicate non-transformed data was plotted. Curves were transformed by setting the average peak value between $25-37^{\circ}\text{C}$ to 1.0 and normalizing other points according to the same normalization factor. Curves were fitted with a sigmoidal curve on Prism. Corresponding R^2 values are recorded in **Table 1**. Melting temperature (T_m) for each protein were determined from the EC50 of each sigmoidal curve. Curves were plotted using GraphPad Prism version 10.2.3.

Microscale Thermophoresis

Microscale Thermophoresis (MST) binding tests and analysis were performed according to a previously recorded protocol³⁴. A Monolith NT.115 (Nanotemper Technologies) was used at room temperature to record MST measurements. 2X stock solutions of Alexa Fluor-647-labeled PHD2 with necessary hydroxylation reagents was made (50 nM Alexa Fluor-647-PHD2, 50 mM Tris-HCl pH 7.5, 1 mM N-oxalylglycine (NOG), 10 μM ferrous sulfate, 5 mg/ml BSA). ~ 1 mg lyophilized HIF1 α ₅₅₅₋₅₇₄ CDD and HIF2 α ₅₂₂₋₅₄₂ CDD peptides were dissolved in 200 μl of 50 mM Tris HCl pH 8.0 and their concentrations were measured via UV absorbance with the following extinction coefficient: 1490 $\text{M}^{-1}\text{cm}^{-1}$. Starting with the stock HIF α peptide solutions, two-fold serial dilutions were performed to generate 16 different 15 μl HIF α concentrated solutions. 15 μl of 2X Alexa-647-PHD2 solution was then added to each HIF α peptide dilution resulting in a final Alexa-647-PHD2 concentration of 25 nM. The solutions were incubated at room temperature for 10 minutes then loaded into standard MST capillaries (Nanotemper Technologies). MST measurements were recorded using 20% fluorescence excitation LED power and 40% (medium)

infrared (IR) laser power. Measurements were recorded for 5 seconds prior to IR laser activation, 20 seconds during IR activation and 3 seconds following IR deactivation. The experimental data was analyzed and fit with a 1:1 K_d binding model using M.O. affinity analysis software (Nanotemper Technologies). Curves were plotted on GraphPad Prism (version 10.2.3).

Bio-layer Interferometry

The binding affinities of HIF1 α peptides and WT or P317R PHD2 were measured by bio-layer interferometry (BLI) using a single channel BLItz instrument (Sartorius). Purified PHD2 was concentrated to 0.45 mM and diluted into BLI Kinetic Buffer (20 mM Tris-HCl pH 7.4, 100 mM NaCl, 1 mM L-ascorbic acid (Sigma), 100 μ M ferrous chloride tetrahydrate (Sigma), 1 mM N-oxalylglycine (Sigma), 0.1% (w/v) Bovine Serum Albumin and 0.1% (v/v) Tween-20). Following reconstitution to a final concentration of 2 mg/ml in DMSO, biotinylated HIF1 α peptides were diluted 200-fold in the BLI Kinetic Buffer. HIF1 α peptides were immobilized on a streptavidin-coated biosensor (Sartorius) for 120 seconds before measuring association to WT PHD2 or P317R PHD2 in a concentration series over 120 seconds. Subsequently, the biosensor was immersed in the BLI Kinetic Buffer for 120 seconds to measure dissociation of the analyte. The sensorgrams were referenced and step corrected, and the binding affinities and kinetic rates were calculated based on global fit of the data to a 1:1 binding model using the BLItz Pro v.1.3.0.5 software. All binding measurements were performed in technical triplicates.

Real-time Prolyl-hydroxylation Kinetic Measurement

Real-time prolyl hydroxylation of HIF1 α ODD was measured according to a previously reported protocol based on Nuclear Magnetic Resonance (NMR)¹². NMR experiments were performed at the UHN Nuclear Magnetic Resonance Core Facility on a Bruker 800 MHz US2 equipped with a NEO console and a 5 mm triple-resonance 13 C optimized TXO cryoprobe, Z-gradient. NMR samples were prepared in 50 mM Tris-HCl pH 7.5, 100 mM NaCl, and 0.5 mM Tris(2-carboxyethyl) phosphine hydrochloride (TCEP-HCl) (NMR buffer) supplemented with 5% D₂O. Prior to the experiments, NMR buffer was gassed with 99.6% oxygen (Messer) for 40 minutes and then used to dilute 0.9 mM of HIF1 α ODD to a final concentration of 0.18 μ M. Samples were transferred to a 5-mm NMR tube pre-purged with 99.6% oxygen for 10 minutes. The real-time hydroxylation experiments were conducted at 25°C using the ^1H - ^{15}N HSQC pulse sequence with sodium 2,2-dimethyl-2-silapentane-5-sulfonate (DSS) as reference. For each experiment, a ^1H - ^{15}N HSQC spectrum of HIF1 α ODD was acquired prior to initiating the hydroxylation reaction with the addition of purified WT or mutant PHD2 at a molar ratio of 1:20 versus HIF1 α ODD with 5 mM α -ketoglutaric acid disodium salt dihydrate (Sigma), 2 mM L-ascorbic acid (Sigma), and 0.1 mM ferrous chloride tetrahydrate. Multiple ^1H - ^{15}N HSQC spectra with acquisition time of 602 seconds were acquired over the course of 20.18 hours. After 6 consecutive acquisitions of ^1H - ^{15}N HSQC spectra, the subsequent spectra were acquired with interscan delays of 2400 seconds. Kinetic measurements with WT, A228S, and P317R PHD2 were performed in technical duplicates.

Real-time hydroxylation of P402 and P564 of HIF1 α was measured by monitoring resonance shifting of A403 and I566¹², respectively, in all ^1H - ^{15}N HSQC spectra. Both resonances shifted as reflections of changes in the local chemical environments upon prolyl-hydroxylation. Reaction progress was reported as the peak intensity of the shifted resonance over the sum of peak intensities for both shifted and unshifted resonances. All spectra were processed with the NMRPipe software⁵¹ and analyzed using the CcpNmr Version-3 software⁵². Kinetic curves were plotted using GraphPad Prism (version 10.2.3).

Supplementary Information

Supplementary Table 1.


Sequences of the HIF1 α and HIF2 α peptides used in the study.
Peptides used for MST and BLI were N-terminally biotinylated.


Peptide	Sequence
HIF1 α ₅₅₅₋₅₇₄ CODD	DLDLEMLAPYIPMDDDFQL
HIF2 α ₅₂₂₋₅₄₂ CODD	ELDLETLAPYIPMDGEDFQL
HIF1 α ₃₉₅₋₄₁₁ NODD	DALTLLAPAAGDTIISLDF

Supplemental Figure 1.

Unnormalized Superdex 200 Increase Chromatogram of wild type and mutant His₆-PHD2 (181-426).

Monomeric PHD2 elutes around 16-17 mL. Peaks before are aggregates and contaminants.






Supplemental Figure 2.

Biolayer interferometry was performed to confirm the binding defect of PHD2 P317R observed via MST.

Measurements with HIF1αCDD and HIF1αNODD peptides indicated that PHD2 P317R did not display a severe binding defect compared to PHD2 WT. While a minor defect was observed with HIF1αCDD and PHD2 P317R, no binding defect was observed with HIF1αNODD. Binding measurements were performed in technical triplicates.



Supplemental Figure 3.

Purification of PHD2 181-426. His₆-PHD2 (181-426) was expressed in BL21 (DE3) *E. coli* cells and purified using a Ni-NTA agarose and size exclusion column (SEC).

10 μ l samples were taken throughout the purification and analyzed via Coomassie stained SDS-PAGE gel. The expected final size of His₆-PHD2 (181-426) is 27.76 kDa. (1) BLUelf prestained protein ladder, 15 μ l (2) Lysate from BL21 (DE3) (3) Ni-NTA agarose flowthrough (4) Ni-NTA agarose 5mM imidazole wash (5) Ni-NTA agarose 30 mM imidazole wash 1 (6) Ni-NTA agarose 30 mM imidazole wash 2 (7) PHD2 elution from Ni-NTA agarose (8) Thrombin cleavage to remove His₆ tag (9) Reverse Ni-NTA agarose flowthrough (10) Reverse Ni-NTA agarose wash (11) Reverse Ni-NTA agarose elution (12) Pooled SEC fractions containing PHD2 (13) BLUelf prestained protein ladder, 5 μ l.

Acknowledgements

We thank the members of the Lee and Ohh labs for their critical comments and helpful discussions. This study was supported by funds from the Canadian Institutes of Health Research (PJT-191811 to M.O. and MOP-133694 to J.E.L.). The UHN NMR Core Facility was supported by the Princess Margaret Cancer Foundation. Infrastructure for biophysics and NMR was funded by the Canada Foundation for Innovation (CFI).

Additional information

Author contributions

C.C.T. designed and performed the experiments, interpreted the data, and wrote the manuscript. W. H. performed the Bio-layer interferometry. W.H. and G.M.C.G.-S. performed the real time Prolyl-hydroxylation assay. M.H. assisted in performing the CHX chase assays. C.C.T., F.G.F. and M.O. conceptualized the project. G.M.C.G.-S., M.I., and J.E.L. contributed new reagents and analytical tools. M.O. interpreted the data, wrote and edited the manuscript. All authors reviewed and edited the manuscript.

Funding

Canadian Institutes of Health Research (PJT-191811)

Canadian Institutes of Health Research (MOP-133694)

Additional files

Supplementary Data [↗](#)

References

- 1 Lee F. S., Percy M. J. (2011) **The HIF pathway and erythrocytosis** *Annu Rev Pathol* **6**:165–192 <https://doi.org/10.1146/annurev-pathol-011110-130321> | Google Scholar
- 2 Weidemann A., et al. (2009) **The glial cell response is an essential component of hypoxia-induced erythropoiesis in mice** *J Clin Invest* **119**:3373–3383 <https://doi.org/10.1172/JCI39378> | Google Scholar
- 3 Richmond T. D., Chohan M., Barber D. L (2005) **Turning cells red: signal transduction mediated by erythropoietin** *Trends Cell Biol* **15**:146–155 <https://doi.org/10.1016/j.tcb.2005.01.007> | Google Scholar
- 4 Bhoopalan S. V., Huang L. J., Weiss M. J (2020) **Erythropoietin regulation of red blood cell production: from bench to bedside and back** *F1000Res* **9** <https://doi.org/10.12688/f1000research.26648.1> | Google Scholar
- 5 Semenza G. L., Wang G. L (1992) **A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation** *Mol Cell Biol* **12**:5447–5454 <https://doi.org/10.1128/mcb.12.12.5447-5454.1992> | Google Scholar
- 6 Mithoowani S., Laureano M., Crowther M. A., Hillis C. M. (2020) **Investigation and management of erythrocytosis** *Cmaj* **192**:E913–E918 <https://doi.org/10.1503/cmaj.191587> | Google Scholar
- 7 Lippl K., et al. (2018) **Born to sense: biophysical analyses of the oxygen sensing prolyl hydroxylase from the simplest animal *Trichoplax adhaerens*** *Hypoxia (Auckl)* **6**:57–71 <https://doi.org/10.2147/HP.S174655> | Google Scholar
- 8 Dengler V. L., Galbraith M., Espinosa J. M (2014) **Transcriptional regulation by hypoxia inducible factors** *Crit Rev Biochem Mol Biol* **49**:1–15 <https://doi.org/10.3109/10409238.2013.838205> | Google Scholar
- 9 Schodel J., et al. (2011) **High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq** *Blood* **117**:e207–217 <https://doi.org/10.1182/blood-2010-10-314427> | Google Scholar
- 10 He W., Batty-Stuart S., Lee J. E., Ohh M (2021) **HIF-1α Hydroxyprolines Modulate Oxygen-Dependent Protein Stability Via Single VHL Interface With Comparable Effect on Ubiquitination Rate** *J Mol Biol* **433** <https://doi.org/10.1016/j.jmb.2021.167244> | Google Scholar
- 11 Chan D. A., Sutphin P. D., Yen S. E., Giaccia A. J (2005) **Coordinate regulation of the oxygen-dependent degradation domains of hypoxia-inducible factor 1 α** *Mol Cell Biol* **25**:6415–6426 <https://doi.org/10.1128/MCB.25.15.6415-6426.2005> | Google Scholar
- 12 He W., Gasmi-Seabrook G. M. C., Ikura M., Lee J. E., Ohh M (2024) **Time-resolved NMR detection of prolyl-hydroxylation in intrinsically disordered region of HIF-1α** *Proc Natl Acad Sci U S A* **121**:e2408104121 <https://doi.org/10.1073/pnas.2408104121> | Google Scholar

- 13 Clifford S. C., Maher E. R (2001) **Von Hippel-Lindau disease: clinical and molecular perspectives** *Adv Cancer Res* **82**:85–105 [https://doi.org/10.1016/s0065-230x\(01\)82003-0](https://doi.org/10.1016/s0065-230x(01)82003-0) | [Google Scholar](#)
- 14 Neumann H. P., Wiestler O. D (1991) **Clustering of features and genetics of von Hippel-Lindau syndrome** *Lancet* **338** [https://doi.org/10.1016/0140-6736\(91\)90401-a](https://doi.org/10.1016/0140-6736(91)90401-a) | [Google Scholar](#)
- 15 Tarade D., Robinson C. M., Lee J. E., Ohh M (2018) **HIF-2alpha-pVHL complex reveals broad genotype-phenotype correlations in HIF-2alpha-driven disease** *Nat Commun* **9**:3359 <https://doi.org/10.1038/s41467-018-05554-1> | [Google Scholar](#)
- 16 Pacak K., et al. (2013) **New syndrome of paraganglioma and somatostatinoma associated with polycythemia** *J Clin Oncol* **31**:1690–1698 <https://doi.org/10.1200/JCO.2012.47.1912> | [Google Scholar](#)
- 17 Pacak K., et al. (2014) **Ocular manifestations of hypoxia-inducible factor-2alpha paraganglioma-somatostatinoma-polycythemia syndrome** *Ophthalmology* **121**:2291–2293 <https://doi.org/10.1016/j.ophtha.2014.06.019> | [Google Scholar](#)
- 18 Delamare M., et al. (2023) **Characterization of genetic variants in the EGLN1/PHD2 gene identified in a European collection of patients with erythrocytosis** *Haematologica* **108**:3068–3085 <https://doi.org/10.3324/haematol.2023.282913> | [Google Scholar](#)
- 19 Gardie B., et al. (2014) **The role of PHD2 mutations in the pathogenesis of erythrocytosis** *Hypoxia (Auckl)* **2**:71–90 <https://doi.org/10.2147/HP.S54455> | [Google Scholar](#)
- 20 Yang C., et al. (2015) **Germ-line PHD1 and PHD2 mutations detected in patients with pheochromocytoma/paraganglioma-polycythemia** *J Mol Med (Berl)* **93**:93–104 <https://doi.org/10.1007/s00109-014-1205-7> | [Google Scholar](#)
- 21 Percy M. J., et al. (2006) **A family with erythrocytosis establishes a role for prolyl hydroxylase domain protein 2 in oxygen homeostasis** *Proc Natl Acad Sci U S A* **103**:654–659 <https://doi.org/10.1073/pnas.0508423103> | [Google Scholar](#)
- 22 Provenzano A., et al. (2022) **Novel Germline PHD2 Variant in a Metastatic Pheochromocytoma and Chronic Myeloid Leukemia, but in the Absence of Polycythemia** *Medicina (Kaunas)* **58** <https://doi.org/10.3390/medicina58081113> | [Google Scholar](#)
- 23 Ladroue C., et al. (2008) **PHD2 mutation and congenital erythrocytosis with paraganglioma** *N Engl J Med* **359**:2685–2692 <https://doi.org/10.1056/NEJMoa0806277> | [Google Scholar](#)
- 24 Lenders J. W., Eisenhofer G., Mannelli M., Pacak K. (2005) **Phaeochromocytoma** *Lancet* **366**:665–675 [https://doi.org/10.1016/S0140-6736\(05\)67139-5](https://doi.org/10.1016/S0140-6736(05)67139-5) | [Google Scholar](#)
- 25 Berra E., et al. (2003) **HIF prolyl-hydroxylase 2 is the key oxygen sensor setting low steady-state levels of HIF-1alpha in normoxia** *EMBO J* **22**:4082–4090 <https://doi.org/10.1093/emboj/cdg392> | [Google Scholar](#)
- 26 Appelhoff R. J., et al. (2004) **Differential function of the prolyl hydroxylases PHD1, PHD2, and PHD3 in the regulation of hypoxia-inducible factor** *J Biol Chem* **279**:38458–38465 <https://doi.org/10.1074/jbc.M406026200> | [Google Scholar](#)
- 27 Flashman E., et al. (2008) **Kinetic rationale for selectivity toward N- and C-terminal oxygen-dependent degradation domain substrates mediated by a loop region of hypoxia-**

- inducible factor prolyl hydroxylases** *J Biol Chem* **283**:3808–3815 <https://doi.org/10.1074/jbc.M707411200> | Google Scholar
- 28 Schofield C. J., Ratcliffe P. J (2004) **Oxygen sensing by HIF hydroxylases** *Nat Rev Mol Cell Biol* **5**:343–354 <https://doi.org/10.1038/nrm1366> | Google Scholar
- 29 Clifton I. J., et al. (2006) **Structural studies on 2-oxoglutarate oxygenases and related double-stranded beta-helix fold proteins** *J Inorg Biochem* **100**:644–669 <https://doi.org/10.1016/j.jinorgbio.2006.01.024> | Google Scholar
- 30 Figg W. D., et al. (2023) **Structural basis for binding of the renal carcinoma target hypoxia-inducible factor 2alpha to prolyl hydroxylase domain 2** *Proteins* **91**:1510–1524 <https://doi.org/10.1002/prot.26541> | Google Scholar
- 31 Ohh M., Taber C. C., Ferens F. G., Tarade D (2022) **Hypoxia-inducible factor underlies von Hippel-Lindau disease stigmata** *eLife* **11** <https://doi.org/10.7554/eLife.80774> | Google Scholar
- 32 Rechsteiner M. P., et al. (2011) **VHL gene mutations and their effects on hypoxia inducible factor HIFalpha: identification of potential driver and passenger mutations** *Cancer Res* **71**:5500–5511 <https://doi.org/10.1158/0008-5472.CAN-11-0757> | Google Scholar
- 33 Maher E. R., Kaelin W. G. (1997) **von Hippel-Lindau disease** *Medicine (Baltimore)* **76**:381–391 <https://doi.org/10.1097/00005792-199711000-00001> | Google Scholar
- 34 Ferens F. G., et al. (2024) **Deficiency in PHD2-mediated hydroxylation of HIF2alpha underlies Pacak-Zhuang syndrome** *Commun Biol* **7** <https://doi.org/10.1038/s42003-024-05904-4> | Google Scholar
- 35 Song D., et al. (2014) **Defective Tibetan PHD2 binding to p23 links high altitude adaption to altered oxygen sensing** *J Biol Chem* **289**:14656–14665 <https://doi.org/10.1074/jbc.M113.541227> | Google Scholar
- 36 Xiang K., et al. (2013) **Identification of a Tibetan-specific mutation in the hypoxic gene EGLN1 and its contribution to high-altitude adaptation** *Mol Biol Evol* **30**:1889–1898 <https://doi.org/10.1093/molbev/mst090> | Google Scholar
- 37 Arsenault P. R., et al. (2013) **A knock-in mouse model of human PHD2 gene-associated erythrocytosis establishes a haploinsufficiency mechanism** *J Biol Chem* **288**:33571–33584 <https://doi.org/10.1074/jbc.M113.482364> | Google Scholar
- 38 Chowdhury R., et al. (2016) **Structural basis for oxygen degradation domain selectivity of the HIF prolyl hydroxylases** *Nat Commun* **7** <https://doi.org/10.1038/ncomms12673> | Google Scholar
- 39 McDonough M. A., et al. (2006) **Cellular oxygen sensing: Crystal structure of hypoxia-inducible factor prolyl hydroxylase (PHD2)** *Proc Natl Acad Sci U S A* **103**:9814–9819 <https://doi.org/10.1073/pnas.0601283103> | Google Scholar
- 40 Greenfield N. J (2006) **Using circular dichroism spectra to estimate protein secondary structure** *Nat Protoc* **1**:2876–2890 <https://doi.org/10.1038/nprot.2006.202> | Google Scholar
- 41 Knauth K., Cartwright E., Freund S., Bycroft M., Buchberger A (2009) **VHL mutations linked to type 2C von Hippel-Lindau disease cause extensive structural perturbations in pVHL** *J Biol*

Chem **284**:10514–10522 <https://doi.org/10.1074/jbc.M809056200> | Google Scholar

- 42 Minamishima Y. A., et al. (2008) **Somatic inactivation of the PHD2 prolyl hydroxylase causes polycythemia and congestive heart failure** *Blood* **111**:3236–3244 <https://doi.org/10.1182/blood-2007-10-117812> | Google Scholar
- 43 Li X., Sutherland S., Takeda K., Fong G. H., Lee F. S (2010) **Integrity of the prolyl hydroxylase domain protein 2:erythropoietin pathway in aging mice** *Blood Cells Mol Dis* **45**:9–19 <https://doi.org/10.1016/j.bcmd.2010.03.003> | Google Scholar
- 44 Franke K., et al. (2013) **HIF-1alpha is a protective factor in conditional PHD2-deficient mice suffering from severe HIF-2alpha-induced excessive erythropoiesis** *Blood* **121**:1436–1445 <https://doi.org/10.1182/blood-2012-08-449181> | Google Scholar
- 45 Takeda K., et al. (2008) **Regulation of adult erythropoiesis by prolyl hydroxylase domain proteins** *Blood* **111**:3229–3235 <https://doi.org/10.1182/blood-2007-09-114561> | Google Scholar
- 46 Myllyharju J., Kivirikko K. I (1997) **Characterization of the iron- and 2-oxoglutarate-binding sites of human prolyl 4-hydroxylase** *EMBO J* **16**:1173–1180 <https://doi.org/10.1093/emboj/16.6.1173> | Google Scholar
- 47 Tian Y. M., et al. (2011) **Differential sensitivity of hypoxia inducible factor hydroxylation sites to hypoxia and hydroxylase inhibitors** *J Biol Chem* **286**:13041–13051 <https://doi.org/10.1074/jbc.M110.211110> | Google Scholar
- 48 Aik W., McDonough M. A., Thalhammer A., Chowdhury R., Schofield C. J (2012) **Role of the jelly-roll fold in substrate binding by 2-oxoglutarate oxygenases** *Curr Opin Struct Biol* **22**:691–700 <https://doi.org/10.1016/j.sbi.2012.10.001> | Google Scholar
- 49 Concordet J. P., Haeussler M (2018) **CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens** *Nucleic Acids Res* **46**:W242–W245 <https://doi.org/10.1093/nar/gky354> | Google Scholar
- 50 Hewitson K. S., Schofield C. J., Ratcliffe P. J (2007) **Hypoxia-inducible factor prolyl-hydroxylase: purification and assays of PHD2** *Methods Enzymol* **435**:25–42 [https://doi.org/10.1016/S0076-6879\(07\)35002-7](https://doi.org/10.1016/S0076-6879(07)35002-7) | Google Scholar
- 51 Delaglio F., et al. (1995) **NMRPipe: a multidimensional spectral processing system based on UNIX pipes** *J Biomol NMR* **6**:277–293 <https://doi.org/10.1007/BF00197809> | Google Scholar
- 52 Skinner S. P., et al. (2016) **CcpNmr AnalysisAssign: a flexible platform for integrated NMR analysis** *J Biomol NMR* **66**:111–124 <https://doi.org/10.1007/s10858-016-0060-y> | Google Scholar

Author information

Cassandra C Taber

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

Wenguang He

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada, Department of Biochemistry, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

Geneviève MC Gasmi-Seabrook

Princess Margaret Cancer Centre, University Health Network, Toronto, Canada

Mia Hubert

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

Fraser G Ferens

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

Mitsuhiko Ikura

Princess Margaret Cancer Centre, University Health Network, Toronto, Canada, Department of Medical Biophysics, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

Jeffery E Lee

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

Michael Ohh

Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada, Department of Biochemistry, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada

ORCID iD: [0000-0001-7600-4751](https://orcid.org/0000-0001-7600-4751)

For correspondence: michael.ohh@utoronto.ca

Editors

Reviewing Editor

Ivan Topisirovic

McGill University, Montreal, Canada

Senior Editor

Lynne-Marie Postovit

Queens University, Kingston, Canada

Reviewer #1 (Public review):

Summary:

Taber et al report the biochemical characterization of 7 mutations in PHD2 that induce erythrocytosis. Their goal is to provide a mechanism for how these mutations cause the disease. PHD2 hydroxylates HIF1a in the presence of oxygen at two distinct proline residues (P564 and P402) in the "oxygen degradation domain" (ODD). This leads to the ubiquitylation of HIF1a by the VHL E3 ligase and its subsequent degradation. Multiple mutations have been reported in the EGLN1 gene (coding for PHD2), which are associated with pseudohypoxic

diseases that include erythrocytosis. Furthermore, 3 mutations in PHD2 also cause pheochromocytoma and paraganglioma (PPGL), a neuroendocrine tumour. These mutations likely cause elevated levels of HIF1a, but their mechanisms are unclear. Here, the authors analyze mutations from 152 case reports and map them on the crystal structure. They then focus on 7 mutations, which they clone in a plasmid and transfect into PHD2-KO to monitor HIF1a transcriptional activity via a luciferase assay. All mutants show impaired activation. Some mutants also impaired stability in pulse chase turnover assays (except A228S, P317R, and F366L). In vitro purified PHD2 mutants display a minor loss in thermal stability and some propensity to aggregate. Using MST technology, they show that P317R is strongly impaired in binding to HIF1a and HIF2a, whereas other mutants are only slightly affected. Using NMR, they show that the PHD2 P317R mutation greatly reduces hydroxylation of P402 (HIF1a NODD), as well as P562 (HIF1a CODD), but to a lesser extent. Finally, BLI shows that the P317R mutation reduces affinity for CODD by 3-fold, but not NODD.

Strengths:

- (1) Simple, easy-to-follow manuscript. Generally well-written.
- (2) Disease-relevant mutations are studied in PHD2 that provide insights into its mechanism of action.
- (3) Good, well-researched background section.

Weaknesses:

- (1) Poor use of existing structural data on the complexes of PHD2 with HIF1a peptides and various metals and substrates. A quick survey of the impact of these mutations (as well as analysis by Chowdhury et al, 2016) on the structure and interactions between PHD2 peptides of HIF1a shows that the P317R mutation interferes with peptide binding. By contrast, F366L will affect the hydrophobic core, and A228S is on the surface, and it's not obvious how it would interfere with the stability of the protein.
- (2) To determine aggregation and monodispersity of the PHD2 mutants using size-exclusion chromatography (SEC), equal quantities of the protein must be loaded on the column. This is not what was done. As an aside, the colors used for the SEC are very similar and nearly indistinguishable.
- (3) The interpretation of some mutants remains incomplete. For A228S, what is the explanation for its reduced activity? It is not substantially less stable than WT and does not seem to affect peptide hydroxylation.
- (4) The interpretation of the NMR prolyl hydroxylation is tainted by the high concentrations used here. First of all, there is a likely a typo in the method section; the final concentration of ODD is likely 0.18 mM, and not 0.18 uM (PNAS paper by the same group in 2024 reports using a final concentration of 230 uM). Here, I will assume the concentration is 180 uM. Flashman et al (JBC 2008) showed that the affinity of the NODD site (P402; around 10 uM) for PHD2 is 10-fold weaker than CODD (P564, around 1 uM). This likely explains the much faster kinetics of hydroxylation towards the latter. Now, using the MST data, let's say the P317R mutation reduces the affinity by 40-fold; the affinity becomes 400 uM for NODD (above the protein concentration) and 40 uM for CODD (below the protein concentration). Thus, CODD would still be hydroxylated by the P317R mutant, but not NODD.
- (5) The discrepancy between the MST and BLI results does not make sense, especially regarding the P317R mutant. Based on the crystal structures of PHD2 in complex with the ODD peptides, the P317R mutation should have a major impact on the affinity, which is what is reported by MST. This suggests that the MST is more likely to be valid than BLI, and the

latter is subject to some kind of artefact. Furthermore, the BLI results are inconsistent with previous results showing that PHD2 has a 10-fold lower affinity for NODD compared to CODD.

(6) Overall, the study provides some insights into mutants inducing erythrocytosis, but the impact is limited. Most insights are provided on the P317R mutant, but this mutant had already been characterized by Chowdhury et al (2016). Some mutants affect the stability of the protein in cells, but then no mechanism is provided for A228S or F366L, which have stabilities similar to WT, yet have impaired HIF1a activation.

<https://doi.org/10.7554/eLife.107121.1.sa3>

Reviewer #2 (Public review):

Summary:

Mutations in the prolyl hydroxylase, PHD2, cause erythrocytosis and, in some cases, can result in tumorigenesis. Taber and colleagues test the structural and functional consequences of seven patient-derived missense mutations in PHD2 using cell-based reporter and stability assays, and multiple biophysical assays, and find that most mutations are destabilizing. Interestingly, they discover a PHD2 mutant that can hydroxylate the C-terminal ODD, but not the N-terminal ODD, which suggests the importance of N-terminal ODD for biology. A major strength of the manuscript is the multidisciplinary approach used by the authors to characterize the functional and structural consequences of the mutations. However, the manuscript had several major weaknesses, such as an incomplete description of how the NMR was performed, a justification for using neighboring residues as a surrogate for looking at prolyl hydroxylation directly, or a reference to the clinical case studies describing the phenotypes of patient mutations. Additionally, the experimental descriptions for several experiments are missing descriptions of controls or validation, which limits their strength in supporting the claims of the authors.

Strengths:

- (1) This manuscript is well-written and clear.
- (2) The authors use multiple assays to look at the effects of several disease-associated mutations, which support the claims.
- (3) The identification of P317R as a mutant that loses activity specifically against NODD, which could be a useful tool for further studies in cells.

Weaknesses:

Major:

- (1) The source data for the patient mutations (Figure 1) in PHD2 is not referenced, and it's not clear where this data came from or if it's publicly available. There is no section describing this in the methods.
- (2) The NMR hydroxylation assay.

A. The description of these experiments is really confusing. The authors have published a recent paper describing a method using ¹³C-NMR to directly detect prolyl-hydroxylation over time, and they refer to this manuscript multiple times as the method used for the studies under review. However, it appears the current study is using ¹⁵N-HSQC-based experiments to track the CSP of neighboring residues to the target prolines, so not the target prolines themselves. The authors should make this clear in the text, especially on page 9, 5th line, where they describe proline cross-peaks and refer to the ¹⁵N-HSQC data in Figure 5B.

B. The authors are using neighboring residues as reporters for proline hydroxylation, without validating this approach. How well do CSPs of A403 and I566 track with proline hydroxylation? Have the authors confirmed this using their ¹³C-NMR data or mass spec?

C. Peak intensities. In some cases, the peak intensities of the end point residue look weaker than the peak intensities of the starting residue (5B, PHD2 WT I566, 6 ct lines vs. 4 ct lines). Is this because of sample dilution (i.e., should happen globally)? Can the authors comment on this?

(3) Data validating the CRISPR KO HEK293A cells is missing.

(4) The interpretation of the SEC data for the PHD2 mutants is a little problematic. Subtle alterations in the elution profiles may hint at different hydrodynamic radii, but as the samples were not loaded at equal concentrations or volumes, these data seem more anecdotal, rather than definitive. Repeating this multiple times, using matched samples, followed by comparison with standards loaded under identical buffer conditions, would significantly strengthen the conclusions one could make from the data.

Minor:

(1) Justification for picking the seven residues is not clearly articulated. The authors say they picked 7 mutants with "distinct residue changes", but no further rationale is provided.

(2) A major finding of the paper is that a disease-associated mutation, P317R, can differentially affect HIF1 prolyhydroxylation, however, additional follow-up studies have not been performed to test this in cells or to validate the mutant in another method. Is it the position of the proline within the catalytic core, or the identity of the mutation that accounts for the selectivity?

<https://doi.org/10.7554/eLife.107121.1.sa2>

Reviewer #3 (Public review):

Summary:

This is an interesting and clinically relevant in vitro study by Taber et al., exploring how mutations in PHD2 contribute to erythrocytosis and/or neuroendocrine tumors. PHD2 regulates HIFα degradation through prolyl-hydroxylation, a key step in the cellular oxygen-sensing pathway.

Using a time-resolved NMR-based assay, the authors systematically analyze seven patient-derived PHD2 mutants and demonstrate that all exhibit structural and/or catalytic defects. Strikingly, the P317R variant retains normal activity toward the C-terminal proline but fails to hydroxylate the N-terminal site. This provides the first direct evidence that N-terminal prolyl-hydroxylation is not dispensable, as previously thought.

The findings offer valuable mechanistic insight into PHD2-driven effects and refine our understanding of HIF regulation in hypoxia-related diseases.

Strengths:

The manuscript has several notable strengths. By applying a novel time-resolved NMR approach, the authors directly assess hydroxylation at both HIF1α ODD sites, offering a clear functional readout. This method allows them to identify the P317R variant as uniquely defective in NODD hydroxylation, despite retaining normal activity toward CODD, thereby challenging the long-held view that the N-terminal proline is biologically dispensable. The work significantly advances our understanding of PHD2 function and its role in oxygen

sensing, and might help in the future interpretation and clinical management of associated erythrocytosis.

Weaknesses:

There is a lack of in vivo/ex vivo validation. This is actually required to confirm whether the observed defects in hydroxylation-especially the selective NODD impairment in P317R-are sufficient to drive disease phenotypes such as erythrocytosis.

The reliance on HRE-luciferase reporter assays may not reliably reflect the PHD2 function and highlights a limitation in the assessment of downstream hypoxic signaling.

The study clearly documents the selective defect of the P317R mutant, but the structural basis for this selectivity is not addressed through high-resolution structural analysis (e.g., cryo-EM).

Given the proposed central role of HIF2 α in erythrocytosis, direct assessment of HIF2 α hydroxylation by the mutants would have strengthened the conclusions.

<https://doi.org/10.7554/eLife.107121.1.sa1>

Author response:

Public Reviews:

Reviewer #1 (Public review):

Summary:

Taber et al report the biochemical characterization of 7 mutations in PHD2 that induce erythrocytosis.

Their goal is to provide a mechanism for how these mutations cause the disease. PHD2 hydroxylates HIF1 α in the presence of oxygen at two distinct proline residues (P564 and P402) in the "oxygen degradation domain" (ODD). This leads to the ubiquitylation of HIF1 α by the VHL E3 ligase and its subsequent degradation. Multiple mutations have been reported in the EGLN1 gene (coding for PHD2), which are associated with pseudohypoxic diseases that include erythrocytosis. Furthermore, 3 mutations in PHD2 also cause pheochromocytoma and paraganglioma (PPGL), a neuroendocrine tumour. These mutations likely cause elevated levels of HIF1 α , but their mechanisms are unclear. Here, the authors analyze mutations from 152 case reports and map them on the crystal structure. They then focus on 7 mutations, which they clone in a plasmid and transfect into PHD2-KO to monitor HIF1 α transcriptional activity via a luciferase assay. All mutants show impaired activation. Some mutants also impaired stability in pulse chase turnover assays (except A228S, P317R, and F366L). In vitro purified PHD2 mutants display a minor loss in thermal stability and some propensity to aggregate. Using MST technology, they show that P317R is strongly impaired in binding to HIF1 α and HIF2 α , whereas other mutants are only slightly affected. Using NMR, they show that the PHD2 P317R mutation greatly reduces hydroxylation of P402 (HIF1 α NODD), as well as P562 (HIF1 α CODD), but to a lesser extent. Finally, BLI shows that the P317R mutation reduces affinity for CODD by 3-fold, but not NODD.

Strengths:

(1) Simple, easy-to-follow manuscript. Generally well-written.

(2) Disease-relevant mutations are studied in PHD2 that provide insights into its mechanism of action.

(3) Good, well-researched background section.

Weaknesses:

(1) Poor use of existing structural data on the complexes of PHD2 with HIF1 α peptides and various metals and substrates. A quick survey of the impact of these mutations (as well as analysis by Chowdhury et al, 2016) on the structure and interactions between PHD2 peptides of HIF1 α shows that the P317R mutation interferes with peptide binding. By contrast, F366L will affect the hydrophobic core, and A228S is on the surface, and it's not obvious how it would interfere with the stability of the protein.

Thank you for the comment. We will further analyze the mutations on the available PHD2 crystal structures in complex with HIF α to discern how these substitution mutations may impact PHD2 structure and function.

(2) To determine aggregation and monodispersity of the PHD2 mutants using size-exclusion chromatography (SEC), equal quantities of the protein must be loaded on the column. This is not what was done. As an aside, the colors used for the SEC are very similar and nearly indistinguishable.

Agreed. We will perform additional experiment as suggested by the reviewer to further assess aggregation and hydrodynamic size. The colors used in the graph will be changed for a clearer differentiation between samples.

(3) The interpretation of some mutants remains incomplete. For A228S, what is the explanation for its reduced activity? It is not substantially less stable than WT and does not seem to affect peptide hydroxylation.

We agree with the reviewer that the causal mechanism for some of the tested disease-causing mutants remain unclear. The negative findings also raise the notion, perhaps considered controversial, that there may be other substrates of PHD2 that are impacted by certain mutations, which contribute to disease pathogenesis. We will expand our discussion accordingly.

(4) The interpretation of the NMR prolyl hydroxylation is tainted by the high concentrations used here. First of all, there is a likely a typo in the method section; the final concentration of ODD is likely 0.18 mM, and not 0.18 μ M (PNAS paper by the same group in 2024 reports using a final concentration of 230 μ M). Here, I will assume the concentration is 180 μ M. Flashman et al (JBC 2008) showed that the affinity of the NODD site (P402; around 10 μ M) for PHD2 is 10-fold weaker than CODD (P564, around 1 μ M). This likely explains the much faster kinetics of hydroxylation towards the latter. Now, using the MST data, let's say the P317R mutation reduces the affinity by 40-fold; the affinity becomes 400 μ M for NODD (above the protein concentration) and 40 μ M for CODD (below the protein concentration). Thus, CODD would still be hydroxylated by the P317R mutant, but not NODD.

The HIF1 α concentration was indeed an oversight, which will be corrected to 0.18 mM. The study by Flashman et al.[1] showing PHD2 having a lower affinity to the NODD than CODD likely contributes to the differential hydroxylation rates via PHD2 WT. We showed here via MST that PHD2 P317R had K_d of 320 ± 20 μ M for HIF1 α CODD, which should have led to a severe enzymatic defect, even at the high concentrations used for NMR (180 μ M). However, we observed only a subtle reduction in hydroxylation efficiency in comparison to PHD2 WT. Thus, we performed another binding method using BLI that showed a mild binding defect on CODD by PHD2 P317R, consistent with NMR data. The perplexing result is the WT-like binding

to the NODD by PHD2 P317R, which appears inconsistent with the severe defect in NODD hydroxylation via PHD2 P317R as measured via NMR. These results suggest that there are supporting residues within the PHD2/NODD interface that help maintain binding to NODD but compromise the efficiency of NODD hydroxylation upon PHD2 P317R mutation. We will perform additional binding experiments to further interrogate and validate the binding affinity of PHD2 P317R to NODD and CODD.

(5) The discrepancy between the MST and BLI results does not make sense, especially regarding the P317R mutant. Based on the crystal structures of PHD2 in complex with the ODD peptides, the P317R mutation should have a major impact on the affinity, which is what is reported by MST. This suggests that the MST is more likely to be valid than BLI, and the latter is subject to some kind of artefact. Furthermore, the BLI results are inconsistent with previous results showing that PHD2 has a 10-fold lower affinity for NODD compared to CODD.

The reviewer's structural prediction that P317R mutation should cause a major binding defect, while agreeable with our MST data, is incongruent with our NMR and the data from Chowdhury et al.[2] that showed efficient hydroxylation of CODD via PHD2 P317R. Moreover, we have attempted to model NODD and CODD on apo PHD2 P317R structure and found that the mutation had no major impact on CODD while the mutated residue could clash with NODD, causing a shifting of peptide positioning on the protein. However, these modeling predictions, like any *in silico* projections, would need experimental validation. As mentioned in our preceding response, we also performed BLI, which showed that PHD2 P317R had a minor binding defect for CODD, consistent with the NMR results and findings by Chowdhury et al[2]. NODD binding was also measured with BLI as purified NODD peptides were not amenable for soluble-based MST assay, which showed similar K_d 's for PHD2 WT and P317R. Considering the absence of NODD hydroxylation via PHD2 P317R as measured by NMR and modeling on apo PHD2 P317R, we posit that P317R causes deviation of NODD from its original orientation that may not affect binding due to the other interactions from the surrounding elements but unfortunately disallows NODD from turnover. Further study would be required to validate such notion, which we feel is beyond the scope of this manuscript. However, we will perform additional binding experiments to further interrogate PHD2 P317R binding to NODD.

(6) Overall, the study provides some insights into mutants inducing erythrocytosis, but the impact is limited. Most insights are provided on the P317R mutant, but this mutant had already been characterized by Chowdhury et al (2016). Some mutants affect the stability of the protein in cells, but then no mechanism is provided for A228S or F366L, which have stabilities similar to WT, yet have impaired HIF1a activation.

We thank the reviewer for raising these and other limitations. We will expand on the shortcomings of the present study but would like to underscore that the current work using the recently described NMR assay along with other biophysical analyses suggests a previously under-appreciated role of NODD hydroxylation in the normal oxygen-sensing pathway.

Reviewer #2 (Public review):

Summary:

Mutations in the prolyl hydroxylase, PHD2, cause erythrocytosis and, in some cases, can result in tumorigenesis. Taber and colleagues test the structural and functional consequences of seven patient-derived missense mutations in PHD2 using cell-based reporter and stability assays, and multiple biophysical assays, and find that most mutations are destabilizing. Interestingly, they discover a PHD2 mutant that can hydroxylate the C-terminal ODD, but not the N-terminal ODD, which suggests the

importance of N-terminal ODD for biology. A major strength of the manuscript is the multidisciplinary approach used by the authors to characterize the functional and structural consequences of the mutations. However, the manuscript had several major weaknesses, such as an incomplete description of how the NMR was performed, a justification for using neighboring residues as a surrogate for looking at prolyl hydroxylation directly, or a reference to the clinical case studies describing the phenotypes of patient mutations. Additionally, the experimental descriptions for several experiments are missing descriptions of controls or validation, which limits their strength in supporting the claims of the authors.

Strengths:

- (1) *This manuscript is well-written and clear.*
- (2) *The authors use multiple assays to look at the effects of several disease-associated mutations, which support the claims.*
- (3) *The identification of P317R as a mutant that loses activity specifically against NODD, which could be a useful tool for further studies in cells.*

Weaknesses:

Major:

- (1) *The source data for the patient mutations (Figure 1) in PHD2 is not referenced, and it's not clear where this data came from or if it's publicly available. There is no section describing this in the methods.*

Clinical and patient information on disease-causing PHD2 mutants was compiled from various case reports and summarized in an excel sheet found in the Supplementary Information. The case reports are cited in this excel file. A reference to the supplementary data will be added to the Figure 1 legend and in the introduction.

- (2) *The NMR hydroxylation assay.*

A. The description of these experiments is really confusing. The authors have published a recent paper describing a method using ¹³C-NMR to directly detect prolyl-hydroxylation over time, and they refer to this manuscript multiple times as the method used for the studies under review. However, it appears the current study is using ¹⁵N-HSQC-based experiments to track the CSP of neighboring residues to the target prolines, so not the target prolines themselves. The authors should make this clear in the text, especially on page 9, 5th line, where they describe proline cross-peaks and refer to the ¹⁵N-HSQC data in Figure 5B.

As the reviewer mentioned, the assay that we developed directly measures the target proline residues. This assay is ideal when mutations near the prolines are studied, such as A403, Y565 (He et al[3]). In this previous work, we observed that the shifting of the target proline cross-peaks due to change in electronegativity on the pyrrolidine ring of proline in turn impacted the neighboring residues[3], which meant that the neighboring residues can be used as reporter residues for certain purposes. In this study, we focused on investigating the mutations on PHD2 while leaving the sequence of the HIF-1α unchanged by using solely ¹⁵N-HSQC-based experiments without the need for double-labeled samples. Nonetheless, we thank the reviewer for pointing out the confusion in the text and we will correct and clarify our description of this assay.

B. The authors are using neighboring residues as reporters for proline hydroxylation, without validating this approach. How well do CSPs of A403 and I566 track with proline

hydroxylation? Have the authors confirmed this using their 13C-NMR data or mass spec?

For previous studies, we performed intercalated 15N-HSQC and 13C-CON experiments for the kinetic measurements of wild-type HIF-1 α and mutants. We observed that the shifting pattern of A403 and I566 in the 15N-HSQC spectra aligned well with the ones of P402 and P564, respectively, in the 13C-CON spectra. Representative data will be added to Supplemental Data.

C. Peak intensities. In some cases, the peak intensities of the end point residue look weaker than the peak intensities of the starting residue (5B, PHD2 WT I566, 6 ct lines vs. 4 ct lines). Is this because of sample dilution (i.e., should happen globally)? Can the authors comment on this?

This is an astute observation by the reviewer. We checked and confirmed that for all kinetic datasets, the peak intensities of the end point residue are always slightly lower than the ones of the starting. This includes the cases for PHD2 A228S and P317R in 5B, although not as obvious as the one of PHD2 WT. We agree with the reviewer that the sample dilution is a factor as a total volume of 16 microliters of reaction components was added to the solution to trigger the reaction after the first spectrum was acquired. It is also likely that rate of prolyl hydroxylation becomes extremely slow with only a low amount of substrate available in the system. Therefore, the reaction would not be 100% complete which was detected by the sensitive NMR experimentation.

(3) Data validating the CRISPR KO HEK293A cells is missing.

We thank the reviewer for noting this oversight. Western blots validating PHD2 KO in HEK293A cells will be added to the Supplementary Data file.

(4) The interpretation of the SEC data for the PHD2 mutants is a little problematic. Subtle alterations in the elution profiles may hint at different hydrodynamic radii, but as the samples were not loaded at equal concentrations or volumes, these data seem more anecdotal, rather than definitive. Repeating this multiple times, using matched samples, followed by comparison with standards loaded under identical buffer conditions, would significantly strengthen the conclusions one could make from the data.

Agreed. We will perform additional experiments as suggested with equal volume and concentration of each PHD2 construct loaded onto the SEC column for better assessment of aggregation.

Minor:

(1) Justification for picking the seven residues is not clearly articulated. The authors say they picked 7 mutants with "distinct residue changes", but no further rationale is provided.

Additional justification for the selection of the mutants will be added to the 'Mutations across the PHD2 enzyme induce erythrocytosis' section. Briefly, some mutants were chosen based on their frequency in the clinical data and their presence in potential mutational hot spots. Various mutations were noted at W334 and R371, while F366L was identified in multiple individuals. Additionally, 9 cases of PHD2-driven disease were reported to be caused from mutations located between residues 200 to 210 while 13 cases were reported between residues 369-379, so G206C and R371H were chosen to represent potential hot spots. To examine a potential genotype-phenotype relationship, two of the mutants responsible for neuroendocrine tumor development, A228S and H374R, were also selected. Finally, mutations

located close or on catalytic core residues (P317R, R371H, and H374R) were chosen to test for suspected defects.

(2) A major finding of the paper is that a disease-associated mutation, P317R, can differentially affect HIF1 prolyhydroxylation, however, additional follow-up studies have not been performed to test this in cells or to validate the mutant in another method. Is it the position of the proline within the catalytic core, or the identity of the mutation that accounts for the selectivity?

This is the very question that we are currently addressing but as a part of a follow-up study. Indeed, one thought is that the preferential defect observed could be the result of the loss of proline, an exceptionally rigid amino acid that makes contact with the backbone twice, or the addition of a specific amino acid, namely arginine, a flexible amino acid with an added charge at this site. Although beyond the scope of this manuscript, we will investigate whether such and other characteristics in this region of PHD2/HIF1 α interface contribute to the differential hydroxylation.

Reviewer #3 (Public review):

Summary:

This is an interesting and clinically relevant in vitro study by Taber et al., exploring how mutations in PHD2 contribute to erythrocytosis and/or neuroendocrine tumors. PHD2 regulates HIF α degradation through prolyl-hydroxylation, a key step in the cellular oxygen-sensing pathway.

Using a time-resolved NMR-based assay, the authors systematically analyze seven patient-derived PHD2 mutants and demonstrate that all exhibit structural and/or catalytic defects. Strikingly, the P317R variant retains normal activity toward the C-terminal proline but fails to hydroxylate the N-terminal site. This provides the first direct evidence that N-terminal prolyl-hydroxylation is not dispensable, as previously thought.

The findings offer valuable mechanistic insight into PHD2-driven effects and refine our understanding of HIF regulation in hypoxia-related diseases.

Strengths:

The manuscript has several notable strengths. By applying a novel time-resolved NMR approach, the authors directly assess hydroxylation at both HIF1 α ODD sites, offering a clear functional readout. This method allows them to identify the P317R variant as uniquely defective in NODD hydroxylation, despite retaining normal activity toward CODD, thereby challenging the long-held view that the N-terminal proline is biologically dispensable. The work significantly advances our understanding of PHD2 function and its role in oxygen sensing, and might help in the future interpretation and clinical management of associated erythrocytosis.

Weaknesses:

(1) There is a lack of in vivo/ex vivo validation. This is actually required to confirm whether the observed defects in hydroxylation-especially the selective NODD impairment in P317R-are sufficient to drive disease phenotypes such as erythrocytosis.

We thank the reviewer for this comment, and while we agree with this statement, the objective of this study *per se* was to elucidate the structural and/or functional defect caused by the various disease-associated mutations on PHD2. The subsequent study would be to validate whether the identified defects, in particular the selective NODD impairment, would

lead to erythrocytosis in vivo. However, we feel that such study would be beyond the scope of this manuscript.

(2) The reliance on HRE-luciferase reporter assays may not reliably reflect the PHD2 function and highlights a limitation in the assessment of downstream hypoxic signaling.

Agreed. All experimental assays and systems have limitations. The HRE-luciferase assay used in the present manuscript also has limitations such as the continuous expression of exogenous PHD2 mutants driven via CMV promoter. Thus, we performed several additional biophysical methodologies to interrogate the disease-causing PHD2 mutants. The limitations of the luciferase assay will be expanded in the revised manuscript.

(3) The study clearly documents the selective defect of the P317R mutant, but the structural basis for this selectivity is not addressed through high-resolution structural analysis (e.g., cryo-EM).

We thank the reviewer for the comment. While solving the structure of PHD2 P317R in complex with HIF α substrate is beyond the scope for this study, a structure of PHD2 P317R in complex with a clinically used inhibitor has been solved (PDB:5LAT). In analyzing this structure and that of PHD2 WT in complex with NODD, Chowdhury et al[2] stated that P317 makes hydrophobic contacts with LXXLAP motif on HIF α and R317 is predicted to interact differently with this motif. While this analysis does not directly elucidate the reason for the preferential NODD defect, it supports the possibility that P317R substitution may be more detrimental for enzymatic activity on NODD than CODD. We will discuss this notion in the revised manuscript.

(4) Given the proposed central role of HIF2 α in erythrocytosis, direct assessment of HIF2 α hydroxylation by the mutants would have strengthened the conclusions.

We thank the reviewer for this comment, but we feel that such study would be beyond the scope of the present study. We observed that the PHD2 binding patterns to HIF1 α and HIF2 α were similar, and we have previously assigned >95% of the amino acids in HIF1 α ODD for NMR study[3]. Thus, we first focused on the elucidation of possible defects on disease-associated PHD2 mutants using HIF1 α as the substrate with the supposition that an identified deregulation on HIF1 α could be extended to HIF2 α paralog.

However, we agree with the reviewer that future studies should examine the impact of PHD2 mutants directly on HIF2 α .

References:

- (1) Flashman, E. et al. Kinetic rationale for selectivity toward N- and C-terminal oxygen-dependent degradation domain substrates mediated by a loop region of hypoxia-inducible factor prolyl hydroxylases. *J Biol Chem* 283, 3808-3815 (2008).
- (2) Chowdhury, R. et al. Structural basis for oxygen degradation domain selectivity of the HIF prolyl hydroxylases. *Nat Commun* 7, 12673 (2016).
- (3) He, W., Gasmi-Seabrook, G.M.C., Ikura, M., Lee, J.E. & Ohh, M. Time-resolved NMR detection of prolyl-hydroxylation in intrinsically disordered region of HIF-1 α . *Proc Natl Acad Sci U S A* 121, e2408104121 (2024).

<https://doi.org/10.7554/eLife.107121.1.sa0>