

Dynamics of compartment-specific proteomic landscapes of hepatotoxic and cholestatic models of liver fibrosis

Reviewed Preprint

v1 • November 13, 2024

Not revised

Marketa Jirouskova , Karel Harant, Pavel Cejnar, Srikant Ojha, Katerina Korelova, Lenka Sarnova, Eva Sticova, Christoph Mayr, Herbert Schiller, Martin Gregor 

Laboratory of Integrative Biology, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic • Laboratory of Mass Spectrometry, BIOCEV, Faculty of Science, Charles University in Prague, Prague, Czech Republic • Department of Mathematics, Informatics and Cybernetics, University of Chemistry and Technology, Prague, Czech Republic • Department of Animal Physiology, Faculty of Science, Charles University, Prague, Czech Republic • Clinical and Transplant Pathology Centre, Institute for Clinical and Experimental Medicine, Prague, Czech Republic • Department of Pathology, The Third Faculty of Medicine, Charles University and University Hospital Kralovske Vinohrady, Prague, Czech Republic • Comprehensive Pneumology Center (CPC)/Institute of Lung Health and Immunity (LHI), Helmholtz Munich; Member of the German Center for Lung Research (DZL), Munich, Germany • Institute of Experimental Pneumology, LMU University Hospital, Ludwig-Maximilians University, Munich, Germany

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study suggests that the composition of the extracellular matrix in a mouse model of liver fibrosis changes depending on the cause of liver fibrosis. The data could be used as a foundation for future antifibrotic therapies. The strength of evidence is **solid** with respect to the use of animal models and proteomic analysis. The study provides a helpful inventory of proteins up or down-regulated, but functional analyses are limited and translational data are lacking.

<https://doi.org/10.7554/eLife.98023.1.sa2>

Abstract

Accumulation of extracellular matrix (ECM) in liver fibrosis is associated with changes in protein abundance and composition depending upon etiology of the underlying liver disease. Current efforts to unravel etiology-specific mechanisms and pharmacological targets rely on several models of experimental fibrosis. Here, we characterize and compare dynamics of hepatic proteome remodeling during fibrosis development and spontaneous healing in experimental models of hepatotoxic (CCl₄ intoxication) and cholestatic (3,5-diethoxycarbonyl-1,4-dihydrocollidine (DDC) feeding) injury. Using detergent-based tissue extraction and mass spectrometry, we identified compartment-specific changes in the liver proteome with detailed attention to ECM composition and changes in protein solubility. Our analysis yielded unique time-resolved CCl₄ and DDC signatures underscoring restricted healing with higher carcinogenic potential upon cholestasis. Correlation of protein abundance profiles with fibrous deposits revealed extracellular chaperone clusterin with

implicated role in fibrosis resolution. Dynamics of clusterin expression was validated in the context of human liver fibrosis. Atomic force microscopy of fibrotic livers complemented proteomics with profiles of disease-associated changes in local liver tissue mechanics. This study determined compartment-specific proteomic landscapes of liver fibrosis and delineated etiology-specific ECM components, providing thus a foundation for future antifibrotic therapies.

Introduction

Liver fibrosis and subsequent cirrhosis, two leading causes of liver disease-related deaths worldwide,¹ develop as a result of chronic liver injury of multiple etiologies.² Fibrosis is characterized by excessive accumulation of extracellular matrix (ECM) proteins forming fibrous scar tissue. In the liver, such pathological matrix is deposited by activated hepatic stellate cells (HSCs) and/or portal fibroblasts (PFs) in response to the inflammatory reaction.^{2,3} Recent multi-omics studies have framed fibrosis as a dynamic multicellular process associated with the remodeling of gene expression landscapes⁴ and profound changes in liver protein abundance as well as composition.^{5–7}

Hepatocellular injury alters parenchymal mechanical properties,⁸ thus further facilitating activation of resident HSCs and PFs and their differentiation into myofibroblasts depositing collagen-rich ECM.⁹ Accumulation of abnormal ECM further promotes ECM-depositing myofibroblasts and fosters progressive whole-organ stiffening as a result of ongoing fibrogenesis.¹⁰ Thus, ECM-associated changes are increasingly perceived as causative, rather than consequential, and multiple efforts have focused on the identification of extracellular niche components that drive the pathogenesis of fibrosis.¹¹

The ECM is a complex network of hundreds of proteins acting as a three-dimensional scaffold, supporting cells mechanically and functioning as a reservoir for secreted factors (e.g., growth factors and cytokines).¹² This so-called “matrisome” consists of “core matrisome” (the structural components of the ECM; i.e., collagens, glycoproteins, and proteoglycans) and “matrix-associated matrisome” (the ECM-interacting proteins).¹³ Transitions of extracellular factors between the insoluble matrisome and the soluble pools modulate their signaling capabilities and bioavailability.¹¹ As fibrous ECM assemblies determine also tissue mechanics, matrisome constituents provide matrix-embedded cells with spatially distinct biochemical and biomechanical context.

In terms of etiology, chronic liver injury is inflicted either by hepatotoxic or cholestatic insult.² In recent decades, several corresponding animal models of liver fibrosis have been established.¹⁴ Despite several translational limitations,¹⁵ these experimental models mimic fundamental aspects of human liver fibrosis.¹⁴ For instance, iterative carbon tetrachloride (CCl₄) intoxication causes hepatocellular damage, HSC activation, and development of pericentral liver fibrosis that evolves into severe bridging fibrosis.¹⁴ In contrast, 3,5- diethoxycarbonyl-1,4- dihydrocollidine (DDC) feeding induces obstructive cholestasis, characterized by expansion of PFs and consecutive periportal fibrosis with typical portal–portal septa.¹⁶ As both human and animal studies have shown that liver fibrosis can be ameliorated by either targeting progression or promoting resolution,¹⁵ experimental models have become instrumental for identifying factors and mechanisms central to blocking fibrosis progression and promoting the reversal of advanced fibrosis.

In this comparative study, we employed CCl₄- and DDC-based mouse models to describe proteomic changes during liver injury, fibrosis development, and repair while focusing on matrisome remodeling and associated alterations in tissue biomechanics. Our in-depth analysis of MS data

obtained from total liver lysates and ECM-enriched insoluble fractions defines compartment- and time-resolved proteomic signatures in toxin- versus cholestasis-induced fibrosis and healing and delineates disease-specific matrisome.

Materials and methods

Animals and liver injury models

C57BL/6J male mice 8–10 weeks old were used for the analysis. All experiments were performed in accordance with an animal protocol approved by the Animal Care Committee of The Institute of Molecular Genetics and according to the EU Directive 2010/63/EU for animal experiments. To induce cholestatic liver injury, mice were either injected with 1 μ l/g CCl₄¹⁴ or fed a diet supplemented with 0.1% DDC.¹⁶

Mass spectrometry sample preparation

A sample of ca 1 mm³ excised from the middle section of left lateral liver lobe was cryo-homogenized using TissueLyser II (QIAGEN, Germantown, MD, USA), resuspended, and sonicated. An aliquot of the sample was precipitated and labeled “Total”. The remaining solution was centrifuged at 16,000 $\times$ g for 20 minutes to separate the supernatant containing soluble (S)-fraction from the pellet, which itself contained insoluble ECM- enriched (E)-fraction.

Nano-liquid chromatography mass spectrometry analysis

Nano reversed-phase columns were used for liquid chromatography MS analysis. Eluting peptide cations were analyzed on a Thermo Orbitrap Fusion collision cell, linear ion trap mass analyzer (Q-OT-qIT, Thermo Scientific). Tandem MS was performed by isolation at 1.5 Th with the quadrupole, higher-energy collisional dissociation fragmentation, and rapid scan MS analysis in the ion trap. Peptides were identified and quantified by MaxQuant label-free quantification software (1.6.7 version) using *Mus musculus* UniProt protein database (UniProtKB version July 2020).

Human samples

The use of completely anonymized archived liver tissue samples for research purposes has been approved by the Ethical Committee of the Institute of Experimental and Clinical Medicine and Thomayer University Hospital, Prague, Czech Republic. Written informed consent was obtained from all patients enrolled in the study. All research was conducted in accordance with both the Declarations of Helsinki and Istanbul. Processing of human liver sections for immunofluorescence is detailed in the Supporting Information.

Atomic force microscopy

Atomic force microscopy (AFM) was performed on liver sections 30 μ m thick as previously described.¹⁷ Using the force mapping method, we measured 30 $\times$ 100 μ m² (10 $\times$ 36 pixels) defined regions precisely located by polarized microscopy. Seven areas (two sections per mouse) were chosen from three different mice for each time point (control, T1, T2, and T4).

Statistics

All graphs and statistical tests indicated in graphs were performed using GraphPad Prism (GraphPad Software). All results are presented as mean $\pm$ SEM unless indicated otherwise. In the boxplots, the box represents 25th and 75th percentiles with the median indicated; whiskers reach to the last data point. Statistical tests used are specified in the figure legends. Statistical significance was determined at levels *, $p < 0.05$, **, $p < 0.01$, and ***, $p < 0.001$.

For further details regarding the materials and methods, please refer to the Supporting Information.

Results

Proteomic analysis outlines the gradual fibrosis development and partial healing in both CCl₄- and DDC- induced liver injury

To compare time-resolved, compartment-specific proteomes of CCl₄- and DDC-induced models of liver fibrosis, total liver lysates (Total) together with two protein fractions obtained by one-step detergent extraction (ECM-enriched insoluble fraction (E-fraction) and soluble fraction (S-fraction)) were prepared for MS analysis (**Figure 1A** [↗](#)). The gradual fibrogenesis was assessed at two time points (T1 and T2) corresponding to mild fibrosis with partially bridging septa (T1, 3 weeks for CCl₄ and 2 weeks for DDC treatment) and advanced fibrosis (T2, 6 weeks for CCl₄ and 4 weeks for DDC). To characterize the dynamics of spontaneous fibrosis resolution, we allowed the mice 5 (T3) or 10 days (T4) to recover. Both models developed typical progressive liver damage with fibrotic scarring followed by partial healing upon insult withdrawal as demonstrated by plasma levels of liver injury markers and the extent of collagen-rich deposits in liver sections (Figure S1).

Using the two models, we quantified 4,737 proteins approximately evenly distributed across all time points and fractions (**Figure 1B,C** [↗](#), Supporting Results). Principal component analysis (PCA) revealed clear temporal separation of CCl₄- and DDC-derived samples in both Total and E-fraction (**Figure 1E** [↗](#)). Unsupervised hierarchical cluster analysis of total lysate proteins together with annotation enrichment of the observed clusters further demonstrated substantial differences in the dynamic regulation of protein expression between the two models (Figure S2). This was evidenced by differential Total MS protein intensity temporal profiles of matrisome-enriched cluster constituents, such as vitronectin, galectin 8, fibrillin 1, or a putative portal myofibroblast marker collagen α -1(XV) chain (**Figure 1F** [↗](#)). Moreover, distinct galectin 8 and vitronectin expression profiles obtained for CCl₄ and DDC E-fractions revealed complex changes in protein association with the insoluble proteome. Taken together, our MS data allow for time-resolved identification of proteins differentially expressed during fibrosis progression and resolution in etiologically distinct models of liver fibrosis. Moreover, this approach enables analysis of complex changes in the solubility of the identified proteins and their transition between liver compartments.

Time-resolved analysis of Total proteomes shows limited healing and tumorigenic potential of the DDC model

To capture the proteome differences between toxin- vs. cholestasis-induced fibrogenesis, we determined proteins significantly regulated during fibrogenesis separately in CCl₄ and DDC Total samples (Benjamini-Hochberg false discovery rate (B.H. FDR) < 5%; see Supporting Materials and Methods). In total, 702 proteins were found to be shared by the two models, while 514 (CCl₄) and 1,074 (DDC) proteins were identified as unique for the respective model (**Figure 2A** [↗](#)).

Using Ingenuity Pathway Analysis (IPA), we predicted upstream transcriptional regulators and growth factors together with corresponding downstream biological signaling pathways differentially associated with identified protein groups (**Figures 2B** [↗](#) and S3A). Abundance changes in known targets indicated the activity of key regulators involved in fibrogenesis (TGF β 1 and angiotensinogen), hypoxia response (HIF1A), and inhibition of factors mediating hepatocyte function (HNF4 α and HNF1 α) or involved in HSC inactivation (PPARGC1A) in both fibrotic models (Figure S3A). Interestingly, the pro-proliferative mTOR pathway showed signaling attenuation in




FIGURE 1.

Hepatotoxic and cholestatic injury generate distinct time-resolved and compartment-specific protein signatures. **(A)** Schematic overview of the experimental setup. Six animals were used in each model at each time point. **(B)** Numbers of quantified proteins at the indicated time points and experimental conditions; $n = 6$. **(C)** Box plots show the distribution of protein sequence coverage (coverage of possible tryptic peptides per protein in %) for the indicated matrisome categories (as defined by Naba et al.¹³) and all detected proteins in experimental conditions indicated. **(D)** Principal component analysis (PCA) of Total proteome separates time-dependent fibrogenesis and healing in CCl₄- (closed symbols) and DDC- (open symbols) induced fibrosis. The first two components of data variability of 3,624 proteins identified in Total liver fractions in CCl₄ and 3,521 proteins in DDC are shown; $n = 6$. **(E)** PCA shows the separation of Total (closed symbols) and E-fraction (open symbols) proteomes in time; the first two components of data variability are shown; $n = 6$. **(F)** Line plots show time-dependent changes in mass spectrometry intensities in Total (solid line) and E-fraction (broken line) proteomes for indicated selected proteins in CCl₄ and DDC models; $n = 4-6$.




FIGURE 2.

Time-resolved analysis of Total proteomes shows limited healing in matrisome-enriched protein clusters in the cholestatic model. **(A)** Venn diagram shows relative proportion of 1,276 and 1,836 proteins differentially expressed (*t*-test; Benjamini-Hochberg FDR < 5%) in Total CCl₄ and DDC proteomes. **(B)** Hierarchical cluster analysis of the activity score of the upstream regulators at the indicated time points predicted by Ingenuity Pathway Analysis (IPA) from unique CCl₄ and DDC protein signatures shown in A. **(C,D)** Hierarchical clustering of mean z-scored mass spectrometry (MS) intensities of proteins of Total CCl₄ (C) or DDC (D) proteomes; *n* = 6. Profiles of z-scored MS intensities of proteins from matrisome-annotated clusters for CCl₄ (I–III) and DDC (I and II) models are shown. **(E)** Venn diagram compares proteins from matrisome-annotated clusters shown in C and D. UniProt keyword enrichment annotation for each group within the diagram is indicated (Fisher's test, Benjamini-Hochberg FDR < 4%).

the course of CCl₄ treatment but upregulation in the case of DDC treatment, suggesting hyperplastic potential of the DDC model (Figure S3A). Consistently, analysis of the DDC signature revealed inhibition of tumor suppressor gene *Rb1* with simultaneous activation of potential oncogenic regulators KDM5A, MRTFA, MYC, and YAP, thus underscoring increased carcinogenic risk (Figures 2B and S3B). By contrast, the CCl₄ signature indicated activation of EGF and VEGFα with simultaneous downregulation of the HNF4α (Figure 2B). This supports previous reports on the importance of these factors during hepatotoxic injury.^{18–20} While dynamic changes of regulatory networks and signaling pathways correlated well with fibrosis progression in both models, ineffective downregulation of profibrotic signaling in the DDC-induced model suggested a limited capacity of the liver to heal from cholestatic injury (Figures 2B and S3B).

Next, we analyzed CCl₄ and DDC Total proteomes using unsupervised hierarchical clustering and functional annotation term enrichment (Figure 2C,D). This allowed the separation of proteins with similar temporal abundance profiles while simultaneously revealing clusters comprising matrisome-annotated proteins (see Materials and Methods). In the CCl₄ proteome, three matrisome-annotated clusters (Figure 2C; clusters I, II, and III) containing 755 proteins showed a variable degree of time-dependent abundance decline over the course of healing. By contrast, two large matrisome-annotated clusters (Figure 2D; clusters I and II) with 799 constituents showed almost no decline in abundance over the recovery period in the DDC model. Our data reflect the reversibility of CCl₄-induced fibrogenesis and restricted healing with higher carcinogenic risk in the DDC model.

Matrisome linked with cholestasis is enriched in basement membrane (BM) proteins whereas deposits upon hepatotoxic injury contain matrisome-associated proteins

To further dissect disease-specific processes associated with matrisome-enriched clusters (Figure 2C,D), we followed the functional annotation of proteins using a comprehensive annotation matrix compiled from GO terms, UniProtKB keywords, and the MatrisomeDB matrisome database.^{13, 21} This revealed that 303 proteins shared by CCl₄ and DDC Total proteomes were annotated not only with matrisome-related keywords but also as “Cell adhesion”, “Fibrinogen complex”, “Platelet aggregation”, and “Intermediate filaments”, thus indicating engagement in cellular interaction with ECM (Figure 2E). Consistently, top canonical IPA pathways included “Signaling by Rho family GTPases”, “Integrin signaling” and “Actin cytoskeleton signaling” (Figure S3C). As anticipated, 452 proteins uniquely identified in CCl₄ matrisome-enriched clusters were also associated with inflammatory response, a well-described feature of this model.²² A group of 496 proteins exclusive for DDC was found to associate with “Basement membrane” and “Cell substrate junction” reflecting an increased abundance of BM components synthesized during the development of periductal fibrosis (e.g., laminins, α-chains of collagens IV, VI, and XVIII; Figure 2E).

In a group of 60 matrisome proteins significantly changing in both Total CCl₄ and DDC proteomes (Figure 3A) there were mainly core matrisome proteins, such as collagens (e.g., α-chains of collagens I and V) and ECM glycoproteins (e.g., fibronectin, EMILIN1, and vitronectin). In contrast to previous reports,⁷ we found collagen VI, a protein regulating ECM contractility,²³ upregulated during fibrogenesis in both models (Figure S4). In addition, we identified several matrisome-associated proteins (e.g., NGLY1, PLOD3, S100A4, MUG2, SERPINA7, SERPINF1, and P4HA1) not previously reported in liver fibrosis as uniquely induced in the CCl₄ model.²¹ Among these, ECM glycoprotein lactadherin (MFGE8), has been shown to reduce liver fibrosis in mice.²⁴ In contrast, BM matrisome proteins, such as laminins, collagens IV and XVIII, and perlecan (HSPG2) were exclusively upregulated in the DDC proteome (Figures 3A and S4).




FIGURE 3.

Comparison of matrisome proteins differentially expressed between the CCl₄- and DDC-derived Total proteomes with immunofluorescence (IF) localization of the main core matrisome proteins within the injured livers. **(A)** Venn diagram shows a comparison of matrisome proteins differentially enriched in Total CCl₄ and DDC proteomes. Color coding indicates identified matrisome categories. **(B–D)** Representative IF images of fibronectin (B), collagen I (C), and collagen IV (D), all in green in liver sections from untreated controls (Ctrl), CCl₄-, and DDC-treated mice at time points of fibrosis development (T2) and resolution (T4). Bile ducts were visualized with antibodies to keratin 19 (K19; red); nuclei were stained with DAPI (blue). CV, central vein; PV, portal vein. Scale bar = 100 μ m. Line plots show time-dependent change in respective mass spectrometry intensities in Total (solid line) and E-fraction (broken line) proteomes in CCl₄ and DDC models; n = 4–6.

To validate our findings demonstrating distinct features of the fibrotic models used, we performed immunofluorescence (IF) analysis of major core matrisome proteins abundantly upregulated in both CCl₄ and DDC proteomes (**Figure 3B–D**). Collagen I accumulated at the sites of primary injury and, consistent with proteomics, was only partially reduced upon recovery in the DDC model (**Figure 3B**). Collagen I-rich areas were delineated by fibronectin, a protein serving as a scaffold for collagen fibril assembly.²⁵ Although fibronectin abundance in Total CCl₄ and DDC proteomes decreased with healing, IF analysis revealed persisting deposits within capillarized hepatic sinusoids even after a recovery period of 10 days (**Figure 3C**). Consistent with a report on human cirrhotic livers,²⁶ BM-associated collagen IV accumulated around fibrotic septa in both models and its persistent presence indicated incomplete liver healing after DDC withdrawal (**Figure 3D**).

Integrin α v is specifically induced on the membranes of injured hepatocytes within fibrotic scar tissue in the hepatotoxic CCl₄ model

The dynamic nature of fibrosis stems from interplay between injured hepatocytes, immune cells, and hepatic myofibroblasts.³ Using previously published cell type-specific signatures (see Materials and Methods), we identified 10 different cell populations in CCl₄ and DDC samples and compared their abundances over the course of liver fibrosis (**Figures 4A–D** and S5A). A set of 18 proteins quantified from the signature of hepatocytes exhibited faster decline in the hepatotoxic CCl₄ than DDC model (**Figure 4A**), corresponding well to the extensive parenchymal injury evidenced by high alanine and aspartate transaminase levels (Figure S1A).

Rapid increase in HSC and activated PF signatures paralleled the activation and proliferation of ECM-depositing myofibroblasts in both fibrotic models (**Figure 4B**). However, faster upregulation of activated PFs in DDC confirms their predominant role in cholestatic fibrosis. Temporal abundance profiles of Kupffer cell, macrophage, monocyte, and B-cell signature proteins illustrate faster recovery from inflammation with healing in the CCl₄ than in DDC model (**Figures 4C,D** and S5A). These findings, together with matrisome-enriched protein cluster analysis (**Figure 2E**), emphasized the role of the inflammation component in the context of fibrosis and underscored the model-specific involvement of PFs in cholestatic fibrosis.

Cellular interactions with the changing microenvironment are mediated by integrins, a heterogeneous family of cell adhesion receptors. Thus, temporal changes in cell type-specific integrin subsets reflect the alterations in fibrotic injury-induced cell populations with the potential to predict the fibrogenicity of the ongoing disease. Our profiling revealed increased expression of integrin β 1 (Figure S5B), the most abundant collagen receptor, which has been reported (together with integrins α 1 and α 5) to correlate with the stages of human liver fibrosis.²⁷ Given its cell type-specific expression prevalence, this finding likely corresponds to the expansion of HSCs, simultaneously with induced integrin β 1 expression on LSECs and injured hepatocytes.²⁷ A rapid immune response is documented by extensive upregulation of main leukocyte integrin subunits β 2 and α M (**Figures 4E** and S5B). Their differential expression kinetics between the models correlate well with cell type-specific signatures (**Figures 4C,D** and S5A). Unexpectedly, fibronectin receptor integrin α 5 (typically expressed on HSCs, PFs, and LSECs^{28, 29}) was detected in Total DDC proteome only (**Figure 4E**). Inasmuch as a boosted expression of integrin α 5 has been reported in liver tumors,³⁰ this finding further supports the tumorigenic potential of biliary fibrosis.

Strong induction of TGF β -activating integrin α v in both CCl₄ and DDC Total proteomes (**Figure 4E**) reflected its central role in fibrogenesis.³¹ Strikingly, in E-fraction, integrin α v was found to increase only in the CCl₄ model (**Figure 4E**), suggesting its close association with ECM specifically during hepatotoxic injury. Subsequent IF analysis showed localization of antibodies to




FIGURE 4.

Liver cell type dynamics and cell type-specific integrin expression during fibrogenesis and healing. **(A–D)** Box plots show mean z-scored mass spectrometry (MS) intensities of the indicated cell type-specific protein signatures in time. Hepatocytes (A; $n = 18$), HSCs, and activated portal fibroblasts (PFs) (B; $n = 3$ and 4), Kupffer cells (C; $n = 6$), granulocytes, and macrophages (D; $n = 16$ and 33). One-way ANOVA with Bonferroni's post-test; * $p < 0.05$; ** $p < 0.01$; † $p < 0.001$. **(E)** The line plots show the time-dependent change in MS intensities of indicated selected integrins in Total (solid line) and E-fraction (broken line) proteomes in CCl₄ and DDC models; $n = 4–6$. **(F)** Representative immunofluorescence images of liver sections from untreated controls (Ctrl), CCl₄-, and DDC-treated mice at time points of fibrosis development (T2) and resolution (T4) immunolabeled for integrin α V (green), K19 (red), and α SMA (blue). Arrowheads, integrin α V-positive injured hepatocytes; arrows, integrin α V-positive biliary epithelial cells of reactive ductuli. CV, central vein; PV, portal vein. Boxed areas, ×2 images. Scale bar = 50 μ m.

av mostly to periportal hepatocytes and with little signal found in central hepatocytes of control livers (**Figure 4F**). During CCl₄-induced injury, av staining was enhanced at the periphery of pericentral hepatocytes adjacent to the collagen-rich scars, reflecting ongoing liver periportalization with the injury.²² The zonal distribution of av was partially restored with fibrosis regression. In DDC-driven cholestasis, antibodies to av stained strongly reactive biliary epithelial cells (BECs), the main drivers of fibrogenesis in biliary fibrosis (**Figure 4F**). Thus, our analysis reveals stage-specific induction of integrin av on the surface of pericentral hepatocytes that has been unrecognized to date and suggests its potential as a marker of reactive cholangiocytes in cholestasis.

Analysis of solubility profile dynamics of liver proteome reveals extracellular matrix protein-1 among matrisome proteins induced by fibrogenesis

As ECM remodeling during fibrogenesis entails also changes in the solubility of its constituents and associated proteins,¹¹ we next set out to identify proteins that become increasingly insoluble with fibrosis progression. We found 1,273 (CCl₄) and 762 (DDC) differentially expressed proteins, defined by at least 3-fold higher expression in E- than in S-fraction (see Materials and Methods). Hierarchical clustering of their MS intensity ratios clearly reflected changes in protein solubility profiles in the course of disease progression (Figure S6A,B). Proteins grouped into three (CCl₄, 406 proteins) and two (DDC, 63 proteins) ECM-enriched clusters exhibited increasing insolubility during fibrogenesis. Matrisome proteins found in both fibrotic models were mainly collagens and ECM glycoproteins (**Figure 5A**). Interestingly, we identified 93 (CCl₄) and 16 (DDC) proteins not present in control samples but induced by treatment (Figure S6C, Supporting Results). Abundance of most of these proteins in E-fraction decreased with healing in the CCl₄ but not in the DDC model (**Figure 5B**), suggesting model-specific incorporation of matrisome-associated proteins into insoluble ECM.

Among proteins upregulated with disease onset in E-fractions of both fibrotic models was extracellular matrix protein-1 (ECM1; **Figures 5C** and S6C). Subsequent IF analysis confirmed weak ECM1 staining (corresponding to low expression levels) in control livers, which was confined mostly to Kupffer cells and the apical membrane of BECs (**Figures 5E** and S6B). Increased ECM1 abundance during fibrogenesis was paralleled by its increasing localization within the infiltrating inflammatory cells and activated Kupffer cells. In addition, ECM1 heavily decorated apical membranes of BECs forming reactive ductuli in DDC-treated livers, corresponding thus to higher MS intensity of ECM1 during the development of biliary fibrosis. Induction of ECM1 expression in the inflammatory cells at the sites of primary injury in the CCl₄ model and in reactive ductuli in the DDC model strongly indicates its involvement in fibrogenesis and healing. Although a previous study had identified ECM1 expression to be hepatocyte-specific,³² our data suggest that its role might be more complex than anticipated.

Correlation of dynamic changes of hepatotoxic proteome and fibrotic deposits identifies clusterin as a novel protein associated with fibrosis resolution

The CCl₄ model gradually developed typical bridging liver fibrosis¹⁴ followed by fibrosis resolution as documented by liver injury markers (Figure S1A) and collagen deposits (Figure S1B), as well as by results of our MS profiling (**Figure 2**). These dynamic changes allowed us to correlate protein abundance profiles (from both Total and E-fraction proteomes) with disease dynamics (captured as fibrotic area) in individual mice (**Figure 6**). For example, leukocyte-specific integrin β2-interacting protein coronin 1a displayed a significant positive correlation with




FIGURE 5.

Solubility profiling provides in-depth analysis of model-specific matrisome composition. **(A)** Venn diagram shows relative proportion of proteins from E-fraction proteome identified as proteins with increasing insolubility over the course of fibrosis in each model (see Figure S6 and Supporting Materials and Methods). Matrisome proteins are highlighted with color coding to indicate identified matrisome categories. **(B)** Line plots show time-dependent change in mass spectrometry (MS) intensities of indicated selected matrisome proteins uniquely identified in E-fraction (broken line) proteomes in CCl₄ and DDC models; n = 4–6. (Solid line shows MS intensity profile in Total proteome.) **(C)** Representative immunofluorescence images of liver sections from untreated controls (Ctrl), CCl₄-, and DDC-treated mice at indicated time points of fibrosis development (T1 and T2) and resolution (T4) immunolabeled for ECM1 (green) and collagen 1 (red). Nuclei were stained with DAPI (blue). Arrows, ECM1-positive reactive biliary epithelial cells. CV, central vein; PV, portal vein. Boxed areas, ×2 images. Scale bar = 50 μm.

fibrogenesis reflecting the time-course of the immune response while negatively correlating methionine cycle enzyme adenosylhomocysteinase corresponded to hepatocellular death induced by fibrogenesis (**Figure 6B,C**).

In the Total CCl₄ proteome, proteins positively correlating with fibrogenesis included matrisome proteins (e.g., angiotensinogen, fibrinogen α , β , γ -chain, fibrinogen-like protein 1, and fibronectin). The IPA analysis linked these to the inflammation and pathways associated with ECM-cell interactions, such as “Actin cytoskeleton signaling” and “Integrin signaling” (Figure S7A). Proteins with negative correlation were mostly enzymes involved in hepatocyte metabolism, such as alpha-enolase (ENO1), an enzyme shown to participate as plasminogen receptor in ECM degradation.³³ Analogously to Total proteome, proteins of CCl₄ E-fraction positively correlating with fibrosis area were linked to fibrogenesis (Figure S7B). Displaying negative correlation was a group of 41 proteins consisting mainly of enzymes associated with amino acid metabolism, necrosis, and apoptosis (Figure S7A,B). In addition, three core matrisome proteins – decorin, MFAP2, and collagen α -1 (I) – also were identified. As shown for collagen α -1 (I), their MS intensity temporal profiles indicated their upregulation in E-fraction while their overall abundance decreased with healing (**Figure 6D**), likely corresponding to the increased association of these matrisome proteins with degraded insoluble ECM.

Among proteins positively correlating with fibrotic scar development and resolution in both Total and insoluble proteomes was clusterin, a glycoprotein implicated in several diseases (**Figure 6D**).³⁴ As clusterin imparts extracellular chaperone function not previously linked to fibrosis, we next investigated clusterin’s localization by IF microscopy (**Figure 6E**). In untreated livers, clusterin was detected in BECs and partially in hepatocytes. Upon CCl₄ intoxication, IF staining revealed its prominent expression by damaged hepatocytes. Strikingly, clusterin increasingly re-localized along the collagen deposits over the healing period. In DDC- treated livers, clusterin was found mostly in BECs forming reactive ductuli and hepatocytes, with minimal changes upon healing (**Figure 6E**). Moreover, extracellular clusterin deposition in close proximity of collagen fibers was further demonstrated in fibrotic and cirrhotic human livers of various etiologies at different stages of fibrosis (**Figures 6F** and S8). The gradual increase in clusterin expression with the progression of fibrosis against the background of chronic hepatitis C (METAVIR score F1-F4) further emphasized its role in the development of liver fibrotic diseases.

Interface hepatocyte elasticity responds dynamically to the pericentral injury in the course of fibrosis development

Identification of “Rho-A”, “Rho family GTPases”, “Actin cytoskeleton”, and “Integrin/ILK signaling” among top canonical pathways elicited by proteins of Total CCl₄ proteome as well as proteins of matrisome- enriched clusters (Figure S3A,C) suggested that the hepatotoxic injury leads to increased cytoskeletal tension in injured hepatocytes. This was further supported by significant enrichment in intermediate filament proteins (**Figure 2E**) and in myocardin-related transcription factor (MRTF) targets (e.g., ITGA1, THBS1, ACTR2, and MSN; **Figure 7A**), key regulators involved in cell and tissue mechanics, cytoskeletal dynamics, and mechanosensing.³⁵

As variations in ECM composition and content also substantially affect the biomechanics of liver tissue, we decided to examine correlation between our proteomic results and mapping of dynamic changes in the biomechanical properties of CCl₄-treated livers. We used atomic force microscopy (AFM) combined with polarized microscopy¹⁷ to precisely locate and probe the following compartments: 1) regions of collagen-rich scar tissue in close vicinity of the central vein, 2) injured hepatocyte regions next to the collagen scar, and 3) regions of hepatocytes on the interface between injured and not visibly damaged hepatocytes (so-called interface hepatocytes³⁶) (**Figure 7B**). In parallel, we also measured stiffness of corresponding regions in nontreated control liver sections (**Figure 7C**).




FIGURE 6.

Correlation analysis of protein abundance and changes in fibrotic deposits in the hepatotoxic model associates clusterin with fibrosis resolution. **(A)** Schematic illustrates the correlation of protein abundance changes with changes in sirius red-positive (SR⁺) areas of fibrous ECM deposits in CCl₄-treated animals at the indicated time points. **(B)** The regulator of the actin cytoskeleton, coronin 1a (CORO1A), serves as an example of a protein with a positive slope of the correlation fit. The methionine cycle enzyme, adenosylhomocysteinase (AHCY), serves as an example of a protein with a negative slope of the correlation fit. **(C,D)** The scatter plots show the linear regression slope and the Pearson correlation coefficient for proteins of CCl₄ Total (C), and E- fraction (D) proteomes. Statistical significance of the correlation is color-coded as indicated. The line plots show time-dependent change in mass spectrometry intensities of indicated representative proteins with significant correlation in Total (solid line) and E-fraction (broken line) proteomes in CCl₄ and DDC models; n = 4–6. **(E)** Representative immunofluorescence (IF) images of liver sections from untreated controls (Ctrl), CCl₄-, and DDC-treated mice at indicated time points of fibrosis development (T2) and resolution (T3 and T4) immunolabeled for clusterin (green), keratin 19 (K19, red), and collagen 1 (Col1, blue). Nuclei were stained with DAPI (blue). Arrowheads, clusterin staining signal delineating collagen deposits; arrows, clusterin-positive injured hepatocytes; yellow arrows, clusterin-positive biliary epithelial cells. CV, central vein; PV, portal vein. Boxed areas, ×2 images. Scale bar = 50 μm. **(F)** Representative IF images of human liver sections from different stages of chronic liver diseases of various etiologies (biliary-type, steatotic liver disease, and chronic HCV infection) immunolabeled for clusterin (Clu, green) and collagen 1 (Col1, magenta). Nuclei were stained with DAPI (blue). Top row shows increase in clusterin expression along collagen fibrils in biliary-type and metabolic syndrome-related cirrhosis compared to the stage of mild fibrosis. Bottom row documents change in clusterin staining pattern with chronic hepatitis C progression from fibrosis stage F1 to stage F4 (METAVIR grading system: F1, portal fibrosis; F2, periportal fibrosis; F3, bridging septal fibrosis; F4, cirrhosis). Arrowheads, clusterin staining delineating collagen deposits; arrows, clusterin-positive capillarized sinusoids; yellow arrows, clusterin-positive bile canaliculi (stage F1 only). PV, portal vein. Boxed areas, ×4 images. Scale bar = 50 μm.

Our AFM analysis revealed no substantial topological variations across defined compartments in control livers, with median Young's moduli ranging between 1.2 and 1.6 kPa (**Figure 7C**, Table S1). Fibrogenic response triggered progressive tissue stiffening apparent in all analyzed compartments (**Figure 7D–F**, Table S1), with collagen-rich regions the stiffest at maximal fibrosis (~4.4 kPa at T2; **Figure 7D**). Unexpectedly, interface regions exhibited initial softening (T1; median ~0.8 kPa vs. 1.2 kPa in control livers), which was followed by increase in stiffness up to ~2.0 kPa (T2). With healing (T4), all compartments demonstrated clear decrease in stiffness (**Figure 7D–F**), although this was only marginal in the regions corresponding to injured hepatocytes. Significant softening of collagen-rich scar tissue during healing indicated partial resolution of scar-associated ECM while it associated with the incorporation of several core matrisome proteins (e.g., COL1A1 and FBN1) into the scar tissue (**Figures 1F** and **6D**). Although the heterogeneity of AFM-based measurements constituted a limitation on our correlation analysis relative to proteomic data (Figure S9), this analysis provides a coherent framework for better understanding the dynamics of proteomic landscapes in the context of fibrosis-associated changes in the local mechanics of liver tissue.

Discussion

In this study, we comprehensively characterized and compared dynamic proteomic landscapes of liver fibrosis development and repair in two mouse models based either on repetitive CCl₄ intoxication¹⁴ or DDC feeding.¹⁶ These models are widely used to reliably mimic either hepatotoxic (CCl₄) or cholestatic (DDC) liver injuries leading to fibrosis in humans.¹⁴ Previous attempts to grasp the complexity of etiology-specific fibrotic proteomes with a focus on diseased ECM^{6,7} were limited due to origins of the analyzed material (mostly decellularized samples) and yielded only fragmented insight into the time course of the disease. To fill this research gap, we employed detergent-based tissue extraction and analyzed both Total and insoluble fraction proteomes at multiple time points of disease progression and spontaneous healing. This approach allowed us to 1) compare time-resolved, compartment-specific proteomes of CCl₄ and DDC models, 2) define etiology-specific elements of fibrogenesis, and 3) detect even low-abundance matrisome and matrisome-interacting proteins in the ECM-enriched insoluble fraction samples. Further, our detailed AFM-based profiling of CCl₄-treated livers enabled us to relate our proteomic data to disease-associated changes in the local mechanics of liver tissue.

We and others have demonstrated that the nature of liver injury determines the set of components assembled in diseased ECM,⁷ which thus reflects unique features of the injury. For instance, the initial decline in abundance of hepatocyte- and activated BEC-expressed COL18A1³⁷ and its consecutive increase thereafter in the hepatotoxic CCl₄ model corresponds to hepatocyte death followed by expansion of both COL18A1-expressing cell types.³⁸ By contrast, extensive COL18A1 expression in DDC-induced cholestasis was already apparent in the early fibrotic deposits due to a massive ductular reaction. A prominent group of BM-associated proteins uniquely identified in the cholestatic model can be attributed to activated BECs in proliferating bile ducts. Some of them (e.g., COL18A1, COL4A2, COL6A2, and HSPG2) are among the 14-gene signature potently predicting human patient cirrhosis progression and survival.³⁹ Other DDC matrisome constituents (e.g., COL3A1, elastin, and LAMC1) have been recently linked to the deterioration of hepatocyte functions in connection with ECM stiffening.⁴⁰ Notably, elastin is present in liver biopsies from patients with advanced fibrosis and adverse clinical outcome⁴¹ and is associated with the irreversibility of liver fibrosis. Further, the increased incorporation of crosslinking protein LOXL1 and TGFβ1-related protein LTBP4 into DDC-specific ECM underscores its resemblance to human cirrhotic liver ECM.⁴² Hence, our findings indicate that cholestasis-driven ECM deposits contain numerous proteins detected in more advanced stages of liver disease favoring hepatocarcinogenesis with a compromised ability to heal. Moreover, identified cholestasis-induced unique signaling pathways were analogous to those recently identified in a subtype of mouse cancer models mimicking human cholangiocarcinoma-like hepatocellular cancer.⁴³

Our observations are concordant with recently published studies (reviewed in^{2,36}) revealing that in the fibrotic liver resident and nonresident cell types wire into dynamic intercellular hubs with shifting cell populations, reflecting and shaping the disease course in an etiology-specific manner. Many of the feedback mechanisms between the cells and ECM regulating fibrosis are mediated by members of the integrin family. In agreement with previous studies,^{27,44} we observed in both models significant upregulation of all detected integrin subunits during fibrosis progression. Strong induction of integrin α M expression in cholestatic but not hepatotoxic injury documents the role of leukocyte-specific integrin α M β 2 in the modulation of biliary fibrosis,⁴⁴ suggesting that the integrin α M might thus serve as a possible selective target for treatment of cholestatic liver disease. Further, a prominent transition of TGF β 1-activating integrin α V toward insoluble ECM exclusively upon hepatotoxic injury was accompanied by its localization to injured pericentral hepatocytes near collagen-rich scars and α SMA-positive HSCs. This finding, together with a measured increase in the stiffness of injured hepatocytes, implies an active role of integrin α V in the targeted activation of TGF β 1 in the local microenvironment during centrilobular fibrosis. Indeed, integrin α V located on hepatocytes in the vicinity of biliary fibrotic septa has been suggested to indicate hepatocyte biliary transformation.⁴⁵ Here, we propose that compartment-specific induction of α V expression on the surface of scar-associated hepatocytes could be a general mechanism to promote fibrosis progression.

In contrast to the cholestatic DDC model, hepatotoxic injury in the CCl₄ model showed a substantial ability to heal 10 days after challenge withdrawal. Such healing capacity allowed us to correlate the dynamics in CCl₄ proteomes with the changes in fibrotic deposits not only during fibrosis progression but also in resolution. Among proteins correlating with scar tissue formation in both Total and insoluble proteomes, we identified a small molecular chaperone clusterin. Clusterin is believed to be associated with elastin fibers in cholestasis,⁴⁶ and recently it was linked to the attenuation of mouse hepatic fibrosis.⁴⁷ Here, we localized clusterin exclusively to injured, pathologically stiffer hepatocytes. Interestingly, the clusterin promoter comprises binding motifs for mechanosensitive transcription factors such as Fos and AP-1/Jun.^{48,49} This suggests that local clusterin induction can reflect the dynamic changes of microenvironmental mechanical cues. Most intriguingly, we also found clusterin to associate with collagen-rich ECM deposits in the course of mouse fibrosis regression and with pathological human matrix of various etiologies. As clusterin overexpression associates with increased activity of ECM-degrading matrix metalloproteinases,⁵⁰ we hypothesize that clusterin accumulation facilitates the remodeling and resolution of scar tissue. Although the specific molecular function of clusterin in fibrotic tissue repair processes remains to be determined, our results strongly suggest clusterin to be an attractive antifibrotic target.

Liver cell and tissue mechanics play a pivotal role in the processes that initiate and resolve fibrotic injury.³ Initial disruption of mechanical homeostasis prompts cytoskeletal remodeling that alters cell-generated forces and cellular biomechanics. Here, such a shift to a higher stiffness regime is illustrated by prominent changes in cytoskeleton-related signatures (actin and intermediate filaments) and signatures of mechanosensitive transcription regulators (e.g., MRTF) accompanied by a significant stiffening of parenchymal compartments devoid of apparent collagen deposits. This indicates substantial cell-driven changes in the biomechanical properties of tissue microenvironment. An unexpected decrease in the stiffness of interface hepatocytes with the initial fibrotic changes revealed by our AFM analysis suggests that the initial softening of interface hepatocytes upon injury counterbalances the stiffening of the injured hepatocytes and/or fibrous scar tissue regions. As the interface hepatocytes have been shown to undergo a phenotypic shift in response to the injury,³⁶ it will now be interesting to determine whether altered mechanical properties serve as the cue leading to the genetic fetal reprogramming or if this initial softening is due to the expression of fetal markers. Together, our AFM and proteomic data underscore the role of local tissue stiffening in fibrotic response and postulate involvement of tissue softening during resolution not only in the collagen scar tissue but also in regenerating parenchyma.

Acknowledgements

We would like to thank S.M. Meier-Menches (University of Vienna, Austria) and J. Masek (Charles University, Prague, Czech Republic) for critical reading of the manuscript; and D. Hadraba (Institute of Physiology, Prague, Czech Republic) for sharing his expertise. We also acknowledge the Light Microscopy Core Facility, IMG CAS, Prague, Czech Republic for support with the microscopy imaging presented herein.

This work was supported by the Grant Agency of the Czech Republic (18-02699S and 21-21736S); the Grant Agency of the Ministry of Health of the Czech Republic (NU21-04-00100); the Institutional Research Project of the Czech Academy of Sciences (RVO 68378050); the National Institute for Cancer Research (Programme EXCELES, LX22NPO5102) - Funded by the European Union - Next Generation EU; the Grant Agency of Charles University (273723), and MEYS CR projects (LM2023050, LM2018126, LQ1604 NPU II, LO1419, and LM2015040). CIISB, Instruct-CZ Centre of Instruct-ERIC EU consortium, funded by MEYS CR infrastructure project LM2018127 and European Regional Development Fund-Project “UP CIISB” (No. CZ.02.1.01/0.0/0.0/18_046/0015974) financially supported the measurements at the CF Nanobiotechnology, CEITEC MU (AFM measurements). The funding sources were not involved in the study design, data collection and analysis, decision to publish, or preparation of the article.

Presentation: none.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Additional information

Author Contributions

Conceptualization: M.G. and M.J. Acquisition of data: M.J., K.H., S.O., K.K., and L.S. Analysis and interpretation of data: M.J., K.H., P.C., S.O., and M.G. Drafting of the manuscript: M.J. and M.G. Critical revision of the manuscript for important intellectual content: all authors. Funding: M. G. and M.J. Technical and material support: H.S., E.S., and C.M.

Conflict of interest

Nothing to declare.

List of Abbreviations

- AFM: atomic force microscopy
- BECs: biliary epithelial cells
- B.H. FDR: Benjamini- Hochberg false discovery rate
- CCl₄: carbon tetrachloride
- DDC: 3,5-diethoxycarbonyl-1,4-dihydrocollidine

- ECM: extracellular matrix
- ECM1: extracellular matrix protein-1
- GO term: gene ontology
- HSC: hepatic stellate cell
- IPA: Ingenuity Pathway Analysis
- MRTF: myocardin-related transcription factor
- MS: mass spectrometry
- PCA: principal component analysis
- PF: portal fibroblasts.

References

1. Karlsen TH, Sheron N, Zelber-Sagi S, et al. (2022) **The EASL-Lancet Liver Commission: protecting the next generation of Europeans against liver disease complications and premature mortality** *Lancet* **399**:61–116
2. Kisseleva T, Brenner D (2021) **Molecular and cellular mechanisms of liver fibrosis and its regression** *Nat Rev Gastroenterol Hepatol* **18**:151–166
3. Henderson NC, Rieder F, Wynn TA (2020) **Fibrosis: from mechanisms to medicines** *Nature* **587**:555–566
4. Ramachandran P, Dobie R, Wilson-Kanamori JR, et al. (2019) **Resolving the fibrotic niche of human liver cirrhosis at single-cell level** *Nature* **575**:512–518
5. Niu L, Geyer PE, Gupta R, et al. (2022) **Dynamic human liver proteome atlas reveals functional insights into disease pathways** *Mol Syst Biol* **18**
6. Massey VL, Dolin CE, Poole LG, et al. (2017) **The hepatic "matrisome" responds dynamically to injury: Characterization of transitional changes to the extracellular matrix in mice** *Hepatology* **65**:969–982
7. Klaas M, Kangur T, Viil J, et al. (2016) **The alterations in the extracellular matrix composition guide the repair of damaged liver tissue** *Sci Rep* **6**
8. Georges PC, Hui JJ, Gombos Z, et al. (2007) **Increased stiffness of the rat liver precedes matrix deposition: implications for fibrosis** *Am J Physiol Gastrointest Liver Physiol* **293**:G1147–54
9. Olsen AL, Bloomer SA, Chan EP, et al. (2011) **Hepatic stellate cells require a stiff environment for myofibroblastic differentiation** *Am J Physiol Gastrointest Liver Physiol* **301**:G110–8
10. Saneyasu T, Akhtar R, Sakai T (2016) **Molecular Cues Guiding Matrix Stiffness in Liver Fibrosis** *Biomed Res Int* **2016**
11. Schiller HB, Fernandez IE, Burgstaller G, et al. (2015) **Time- and compartment-resolved proteome profiling of the extracellular niche in lung injury and repair** *Mol Syst Biol* **11**
12. Hynes RO (2009) **The extracellular matrix: not just pretty fibrils** *Science* **326**:1216–9
13. Naba A, Clauser KR, Hoersch S, et al. (2012) **The matrisome: in silico definition and in vivo characterization by proteomics of normal and tumor extracellular matrices** *Mol Cell Proteomics* **11**
14. Liu Y, Meyer C, Xu C, et al. (2013) **Animal models of chronic liver diseases** *Am J Physiol Gastrointest Liver Physiol* **304**:G449–68
15. Trautwein C, Friedman SL, Schuppan D, et al. (2015) **Hepatic fibrosis: Concept to treatment** *J Hepatol* **62**:S15–24

16. Fickert P, Stoger U, Fuchsbichler A, et al. (2007) **A new xenobiotic-induced mouse model of sclerosing cholangitis and biliary fibrosis** *Am J Pathol* **171**:525–36
17. Ojha S, Pribyl J, Klimovic S, et al. (2022) **Measurement of Liver Stiffness using Atomic Force Microscopy Coupled with Polarization Microscopy** *J Vis Exp*
18. Komuves LG, Feren A, Jones AL, et al. (2000) **Expression of epidermal growth factor and its receptor in cirrhotic liver disease** *J Histochem Cytochem* **48**:821–30
19. Yang L, Kwon J, Popov Y, et al. (2014) **Vascular endothelial growth factor promotes fibrosis resolution and repair in mice** *Gastroenterology* **146**:1339–50
20. Yue HY, Yin C, Hou JL, et al. (2010) **Hepatocyte nuclear factor 4alpha attenuates hepatic fibrosis in rats** *Gut* **59**:236–46
21. Shao X, Taha IN, Clauser KR, et al. (2020) **MatrisomeDB: the ECM-protein knowledge database** *Nucleic Acids Res* **48**:D1136–D1144
22. Ghallab A, Myllys M, Holland CH, et al. (2019) **Influence of Liver Fibrosis on Lobular Zonation** *Cells* **8**
23. Williams LM, McCann FE, Cabrita MA, et al. (2020) **Identifying collagen VI as a target of fibrotic diseases regulated by CREBBP/EP300** *Proc Natl Acad Sci U S A* **117**:20753–20763
24. An SY, Jang YJ, Lim HJ, et al. (2017) **Milk Fat Globule-EGF Factor 8, Secreted by Mesenchymal Stem Cells, Protects Against Liver Fibrosis in Mice** *Gastroenterology* **152**:1174–1186
25. Sottile J, Shi F, Rublyevska I, et al. (2007) **Fibronectin-dependent collagen I deposition modulates the cell response to fibronectin** *Am J Physiol Cell Physiol* **293**:C1934–46
26. Mak KM, Chen LL, Lee TF (2013) **Codistribution of collagen type IV and laminin in liver fibrosis of elderly cadavers: immunohistochemical marker of perisinusoidal basement membrane formation** *Anat Rec (Hoboken)* **296**:953–64
27. Nejari M, Couvelard A, Mosnier JF, et al. (2001) **Integrin up-regulation in chronic liver disease: relationship with inflammation and fibrosis in chronic hepatitis C** *J Pathol* **195**:473–81
28. Olsen AL, Sackey BK, Marcinkiewicz C, et al. (2012) **Fibronectin extra domain-A promotes hepatic stellate cell motility but not differentiation into myofibroblasts** *Gastroenterology* **142**:928–937
29. Couvelard A, Scoazec JY, Feldmann G (1993) **Expression of cell-cell and cell-matrix adhesion proteins by sinusoidal endothelial cells in the normal and cirrhotic human liver** *Am J Pathol* **143**:738–52
30. Lai KK, Shang S, Lohia N, et al. (2011) **Extracellular matrix dynamics in hepatocarcinogenesis: a comparative proteomics study of PDGFC transgenic and Pten null mouse models** *PLoS Genet* **7**
31. Reed NI, Jo H, Chen C, et al. (2015) **The alphavbeta1 integrin plays a critical in vivo role in tissue fibrosis** *Sci Transl Med* **7**

32. Fan W, Liu T, Chen W, et al. (2019) **ECM1 Prevents Activation of Transforming Growth Factor beta, Hepatic Stellate Cells, and Fibrogenesis in Mice** *Gastroenterology* **157**:1352–1367
33. Didiasova M, Schaefer L, Wygrecka M (2019) **When Place Matters: Shuttling of Enolase-1 Across Cellular Compartments** *Front Cell Dev Biol* **7**
34. Wilson MR, Zoubeidi A (2017) **Clusterin as a therapeutic target** *Expert Opin Ther Targets* **21**:201–213
35. Esnault C, Stewart A, Gualdrini F, et al. (2014) **Rho-actin signaling to the MRTF coactivators dominates the immediate transcriptional response to serum in fibroblasts** *Genes Dev* **28**:943–58
36. Ben-Moshe S, Veg T, Manco R, et al. (2022) **The spatiotemporal program of zonal liver regeneration following acute injury** *Cell Stem Cell* **29**:973–989
37. Schuppan D, Cramer T, Bauer M, et al. (1998) **Hepatocytes as a source of collagen type XVIII endostatin** *Lancet* **352**:879–80
38. Manco R, Clerbaux LA, Verhulst S, et al. (2019) **Reactive cholangiocytes differentiate into proliferative hepatocytes with efficient DNA repair in mice with chronic liver injury** *J Hepatol* **70**:1180–1191
39. Wu Y, Cao Y, Xu K, et al. (2021) **Dynamically remodeled hepatic extracellular matrix predicts prognosis of early-stage cirrhosis** *Cell Death Dis* **12**
40. Acun A, Oganessian R, Uygun K, et al. (2021) **Liver donor age affects hepatocyte function through age- dependent changes in decellularized liver matrix** *Biomaterials* **270**
41. Kendall TJ, Dolman GE, Duff CM, et al. (2018) **Hepatic elastin content is predictive of adverse outcome in advanced fibrotic liver disease** *Histopathology* **73**:90–100
42. Mazza G, Telese A, Al-Akkad W, et al. (2019) **Cirrhotic Human Liver Extracellular Matrix 3D Scaffolds Promote Smad-Dependent TGF-beta1 Epithelial Mesenchymal Transition** *Cells* **9**
43. Tang M, Zhao Y, Zhao J, et al. (2022) **Liver cancer heterogeneity modeled by in situ genome editing of hepatocytes** *Sci Adv* **8**
44. Joshi N, Kopec AK, Ray JL, et al. (2016) **Fibrin deposition following bile duct injury limits fibrosis through an alphaMbeta2-dependent mechanism** *Blood* **127**:2751–62
45. Popov Y, Patsenker E, Stickel F, et al. (2008) **Integrin alphavbeta6 is a marker of the progression of biliary and portal liver fibrosis and a novel target for antifibrotic therapies** *J Hepatol* **48**:453–64
46. Aigelsreiter A, Janig E, Sostaric J, et al. (2009) **Clusterin expression in cholestasis, hepatocellular carcinoma and liver fibrosis** *Histopathology* **54**:561–70
47. Seo HY, Lee SH, Lee JH, et al. (2019) **Clusterin Attenuates Hepatic Fibrosis by Inhibiting Hepatic Stellate Cell Activation and Downregulating the Smad3 Signaling Pathway** *Cells* **8**
48. Jin G, Howe PH (1999) **Transforming growth factor beta regulates clusterin gene expression via modulation of transcription factor c-Fos** *Eur J Biochem* **263**:534–42

49. Miyamoto-Sato E, Fujimori S, Ishizaka M, et al. (2010) **A comprehensive resource of interacting protein regions for refining human transcription factor networks** *PLoS One* **5**
50. Shim YJ, Kang BH, Jeon HS, et al. (2011) **Clusterin induces matrix metalloproteinase-9 expression via ERK1/2 and PI3K/Akt/NF-kappaB pathways in monocytes/macrophages** *J Leukoc Biol* **90**:761–9

Author information

Marketa Jirouskova

Laboratory of Integrative Biology, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic
ORCID iD: [0009-0004-7173-2177](https://orcid.org/0009-0004-7173-2177)

For correspondence: marketa.jirouskova@img.cas.cz

Karel Harant

Laboratory of Mass Spectrometry, BIOCEV, Faculty of Science, Charles University in Prague, Prague, Czech Republic
ORCID iD: [0000-0002-9654-5392](https://orcid.org/0000-0002-9654-5392)

Pavel Cejnar

Department of Mathematics, Informatics and Cybernetics, University of Chemistry and Technology, Prague, Czech Republic

Srikant Ojha

Laboratory of Integrative Biology, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic, Department of Animal Physiology, Faculty of Science, Charles University, Prague, Czech Republic
ORCID iD: [0009-0002-3930-6624](https://orcid.org/0009-0002-3930-6624)

Katerina Korelova

Laboratory of Integrative Biology, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic
ORCID iD: [0009-0008-6439-5474](https://orcid.org/0009-0008-6439-5474)

Lenka Sarnova

Laboratory of Integrative Biology, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic

Eva Sticova

Clinical and Transplant Pathology Centre, Institute for Clinical and Experimental Medicine, Prague, Czech Republic, Department of Pathology, The Third Faculty of Medicine, Charles University and University Hospital Kralovske Vinohrady, Prague, Czech Republic
ORCID iD: [0000-0003-2486-6266](https://orcid.org/0000-0003-2486-6266)

Christoph Mayr

Comprehensive Pneumology Center (CPC)/Institute of Lung Health and Immunity (LHI), Helmholtz Munich; Member of the German Center for Lung Research (DZL), Munich,

Germany

Herbert Schiller

Comprehensive Pneumology Center (CPC)/Institute of Lung Health and Immunity (LHI), Helmholtz Munich; Member of the German Center for Lung Research (DZL), Munich, Germany, Institute of Experimental Pneumology, LMU University Hospital, Ludwig-Maximilians University, Munich, Germany

ORCID iD: [0000-0001-9498-7034](https://orcid.org/0000-0001-9498-7034)

Martin Gregor

Laboratory of Integrative Biology, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic

ORCID iD: [0000-0001-6841-9527](https://orcid.org/0000-0001-6841-9527)

For correspondence: martin.gregor@img.cas.cz

Editors

Reviewing Editor

Pramod Mistry

Yale University, New Haven, United States of America

Senior Editor

Pramod Mistry

Yale University, New Haven, United States of America

Reviewer #1 (Public review):

Summary:

Jirouskova and colleagues in their study have carried out an in-depth proteomic characterization of the dynamics of the liver fibrotic response and the resulting resolution in two distinct models of liver injury: CCl₄-induced model of hepatotoxicity and pericentral/bridging liver fibrosis and the DDC feeding model of obstructive cholestasis and periportal fibrosis. They focussed on both the insoluble extracellular matrix (ECM) components as well as the soluble secreted factors produced by hepatic stellate cells (HSCs) and/or portal fibroblasts (PFs). They identified compartment- and time-resolved proteomic signatures in the two models with disease-specific factors or matrisomes. Their study also identified phenotypic differences between the models such as that while the CCl₄-induced model induced profound hepatotoxicity followed by resolution, the DDC model induced more lasting liver damage and proteomic changes that resembled advanced human liver fibrosis favouring hepatocarcinogenesis.

Overall, this comprehensive and very well-conducted study is rigorous and well-planned. The conclusions are supported by compelling studies and analyses. One caveat is the lack of mechanistic experiments to prove causality, but this can be carried out in follow-up studies.

Strengths:

(1) A major strength of the study is that the experiments are rigorous and very well conducted. For instance, the authors utilized two models of liver fibrosis to study different aspects of the pathology - hepatotoxicity vs cholestasis. In addition, 4 time points for each model were investigated - 2 for fibrosis development and 2 for fibrosis resolution. They have taken 3 components for proteomic analyses - total lysates, insoluble ECM components as well

as the soluble secreted factors. Thus, the authors provide a comprehensive overview of the fibrosis and resolution process in these models.

(2) Another great strength of the study is that the methodology utilized was able to dissect unique pathways relevant to each model as well as common targets. For example, the authors identified known pathways such as mTOR signalling to be differentially regulated in the CCl₄ vs DDC model. mTOR signalling was increased in the DDC model which is associated with hyperproliferation. Thus showing that the approach taken is specific enough to distinguish between the two similar (both induce fibrosis) but distinct mechanisms (hepatotoxicity vs cholestasis) is a strong point of the study.

Weaknesses:

(1) The authors themselves propose in their Introduction that the "ECM-associated changes are increasingly perceived as causative, rather than consequential"; however, they have not conducted mechanistic (gain of function/loss of function) studies either in vitro or in vivo from any of their identified targets to truly prove causality. This remains one of the limitations of this study. Thus, future studies should investigate this point in detail. For instance, it would have been intriguing to dissect if knocking out specific genes involved in one specific model or genes common to both would yield distinct phenotypic outcomes.

(2) The majority of the conclusions are derived primarily from the proteomic analyses. Although well conducted, it would strengthen the study to corroborate some of the major findings by other means such as IHC/IF with the corresponding quantifications and not only representative images.

<https://doi.org/10.7554/eLife.98023.1.sa1>

Reviewer #2 (Public review):

Summary:

The authors suggest that ECM abundance and composition change depending on the aetiology of liver fibrosis. To understand this they have investigated the proteome in two models of animal fibrosis and resolution. They suggest their findings could provide a foundation for future anti-fibrotic therapies.

Strengths:

The animal models used are widely studied models of liver fibrosis from both parenchymal and biliary damage aspects. Both would allow analysis of resolution. The CCl₄ model in particular fully reverts to a 'healthy' liver following cessation of the insult. I am less clear whether/how quickly the ductal plugs clear in DDC models and thus this may not provide the response they are looking for in terms of reversibility. I believe there have been several extensive studies using a transcriptomics approach in assessing genes and cells involved in the CCl₄ model of resolution. Even more multiomic models of general fibrosis progression in many of the mouse models of fibrosis. However, the proteomic approach they have used is robust and they have made some attempts to integrate with cell-type specific signatures from previously published data.

Although there is minimal data, hepatocyte elasticity is a very interesting part of their study. Additional data and focussed attention on the mechanisms underpinning this would be very insightful.

Weaknesses:

As it currently stands, the data, whilst extensive, is primarily focussed on the proteomic data which is fairly descriptive and I am not clear on the additional insight gained in their approach that is not already detailed from the extensive transcriptomic studies. The manuscript overall would benefit from some mechanistic functional insight to provide new additional modes of action relevant to fibrosis progression. Whilst there is some human data presented it is a minimal analysis without quantification that would imply relevance to disease state.

Although studying disease progression in animals is a fundamental aspect of understanding the full physiological response of fibrotic disease, without more human insight makes any analysis difficult to fulfil their suggestion that these targets identified will be of use to treat human disease.

Some of the terminology is incorrect while discussing these models of injury used and care should be taken. For example - both models are toxin-induced and I do not think these data have any support that the DDC model has a higher carcinogenic risk. An investigation into the tumour-induced risk would require significant additional models. These types of statements are incorrect and not supported by this study.

<https://doi.org/10.7554/eLife.98023.1.sa0>