

Structural mechanism of strand exchange by the RAD51 filament

Reviewed Preprint

v2 • July 15, 2025

Revised by authors

Reviewed Preprint

v1 • May 27, 2025

Luay Joudeh, Robert E Appleby, Joseph D Maman, Luca Pellegrini 

Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom • Francis Crick Institute, London, United Kingdom

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **landmark** study describes the structure of the human RAD51 filament with a recombination intermediate called the displacement loop (D-loop). Using cryogenic structural, biochemical, and single-molecule analyses, the authors provide **compelling** evidence on how the RAD51 filament promotes strand exchange between single-stranded and double-stranded DNAs. The findings are highly relevant to the fields of homologous recombination, DNA repair, and genome stability.

<https://doi.org/10.7554/eLife.107114.2.sa4>

Abstract

Homologous Recombination (HR) preserves genomic stability by repairing double-strand DNA breaks and ensuring efficient DNA replication. Central to HR is the strand-exchange reaction taking place within the three-stranded synapsis wherein a RAD51 nucleoprotein filament binds to a donor DNA. Here we present the cryoEM structure of a displacement loop of human RAD51 that captures the synaptic state when the filament has become tightly bound to the donor DNA. The structure elucidates the mechanism of strand exchange by RAD51, including the filament engagement with the donor DNA, the strand invasion and pairing with the complementary sequence of the donor DNA, the capture of the non-complementary strand and the polarity of the strand-exchange reaction. Our findings provide fundamental mechanistic insights into the biochemical reaction of eukaryotic HR.

Introduction

The transfer of genetic information between DNA molecules promoted by HR is essential in the repair of DNA double-strand breaks^{1,2} and to overcome replicative stress during chromosomal DNA synthesis^{3,4}. The molecular apparatus of HR is also responsible for the crossover reactions that generate genetic diversity in meiosis⁵. HR deficiency causes genetic instability that predisposes to cancer^{4,6,7} and its synthetic lethality with other DNA repair pathways is exploited to induce cancer cell death in therapy^{8,9}.

In eukaryotic cells, the reaction of HR is performed by RAD51^{10,11}, a member of a highly-conserved family of ATPases that includes bacterial RecA and archaeal RadA^{12,13}. RAD51 polymerises on the single-stranded (ss) DNA overhangs that form upon long-range resection of DNA ends¹⁴. The RAD51 nucleoprotein filament invades the donor DNA duplex and searches for sequence homology within a three-stranded synaptic filament intermediate^{15,16}. When homology is found, the invading strand of the filament becomes stably paired to the complementary sequence of the donor DNA, yielding a structure known as a displacement loop (D-loop)^{17–19} (**Fig. 1A**), which ultimately progresses to strand-exchange. Although cellular HR is controlled by a complex network of recombination mediators such as BRCA2²⁰, the RAD51 paralogues^{21–23} and FIGL1^{24,25}, RAD51 can catalyse the biochemical reaction of strand-exchange independently.

The RecA/RAD51/RadA protein family has been extensively investigated to elucidate the mechanistic basis for the reaction of strand exchange. Low-resolution electron microscopy (EM) and X-ray crystallography have demonstrated conservation in the helical parameters of the nucleoprotein filament^{26,27}, the structure of the ATPase domain^{28,29} and the protomer self-association into a filament^{30,31}. CryoEM analysis of the RAD51 nucleoprotein filament have revealed further conservation with RecA in the mode of interaction with DNA, such as the observation that DNA stretching results from its separation into B-form nucleotide triplets^{32–35}. Furthermore, single-molecule experiments have determined the kinetic parameters of filament nucleation and growth^{36,37}, have shown that a minimum length of 8 homologous nucleotides is required for formation of stable synaptic joints^{38,39} and that homologous pairing takes place in steps of three nucleotides^{40,41}.

High-resolution information about the RAD51 synaptic state is still missing, limiting our mechanistic understanding of the strand-exchange reaction. In this study, we have carried out the biochemical reconstitution, cryoEM structure and biophysical validation of a D-loop of human RAD51. The structure elucidates the key steps of strand-exchange, including the mechanism of filament engagement with the donor DNA, strand invasion and pairing with the complementary strand, and capture of the exchanged strand. Comparison with the synaptic state of RecA filaments⁴² reveals both conserved and distinct features of the strand-exchange reaction, shedding light on the molecular evolution of the biochemical reaction of HR.

Results

Electron microscopy (EM) analysis of filamentous RAD51 relies normally on helical averaging, which is unsuitable for reconstruction of an aperiodic structure such as a D-loop. We therefore sought to prepare a D-loop sample of human RAD51 that is amenable to single-particle cryoEM. We designed a 32-nucleotide (nt) poly(dT) with a central, unique 10-nt long sequence complementary to a 50-nt donor double-stranded (ds) DNA; the complementary sequence of the donor DNA was incorporated in a 10-nt mismatch bubble to favour D-loop formation (Fig. S1A and Table S1). The length of the complementary sequence was decided based on evidence that RAD51 requires a homologous tract of at least 8 nucleotides for stable capture of the donor DNA³⁹. As RAD51 can bind both ss- and dsDNA, we defined experimentally appropriate stoichiometry and buffer conditions for D-loop reconstitution. Electrophoretic mobility shift assay (EMSA) with fluorescently labelled oligos showed successful D-loop formation (Fig. S1B and Table S1).

We collected an initial cryoEM dataset of 2792 movies of the RAD51 D-loop sample and processed the data by single-particle analysis. 2D classification showed clear evidence of the presence of synaptic particles and yielded a D-loop reconstruction at 3.31 Å resolution (Fig. S2 and Table S2). To improve data homogeneity and resolution, we collected a larger dataset of 8960 movies of a D-loop sample with ss- and dsDNA that had been biotinylated at both ends and capped with monomeric streptavidin (Fig. S1C)⁴³, yielding a structure of the D-loop at 2.64 Å (Fig. S3 and




Figure 1.

CryoEM structure of the RAD51 D-loop.

A Schematic drawing of RAD51 D-loop formation during double-strand DNA break repair by Homologous Recombination. **B** CryoEM map of the RAD51 D-loop, coloured according to local resolution. **C** Drawing of the RAD51 D-loop structure. The nine RAD51 protomers in the filament are shown as cylinder cartoon, coloured alternatively in lavender and pink. The DNA strands of the D-loop are represented as light yellow tubes for the phosphate ribose backbone and the nucleotides are drawn in stick representation with filled rings. **D** D-loop drawing that highlights the trajectories of the DNA strands of the donor DNA and the invading DNA. The complementary and exchanged strands of the donor DNA are coloured light green and orange; the invading strand of the filament is coloured as the exchanged strand to emphasise that they would share sequence homology *in vivo*. The protein filament is drawn as a transparent surface in light blue. **E** Nucleotide sequences and observed base pairing of the D-loop DNA. Sequences are coloured as in panel D. **F** Cartoon drawing of the DNA strands at the 5'-arm junction. The phospho-ribose backbone is shown as narrow tube, the nucleotide bases as filled rings. DNA strands are coloured as in panel D. The Watson-Crick hydrogen bonds between base pairs are drawn as dashed purple lines. **G** Cartoon drawing of the DNA strands at the 3'-arm junction, coloured and annotated as in panel F.

Table S2). The final single-particle reconstruction based on 102,596 particles comprises a nucleoprotein filament of 9 RAD51 protomers and 26 nucleotides of single-stranded DNA, bound to a dsDNA containing the central bubble region flanked by 15- and 16-bp long segments of donor DNA. The cryoEM map ranges in resolution between 2.5 Å and 2.9 Å for the homologous pairing region of the D-loop and 3 Å to 6 Å for the flanking dsDNA arms (**Fig. 1B** [↗](#)).

D-loop structure

In the D-loop structure, RAD51 maintains the right-handed helical arrangement of protomers observed in pre- and post-synaptic filaments^{32,33} [↗](#), with the same helical twist and a slightly compressed pitch relative to the pre-synaptic filament (**Fig. 1C** [↗](#), Fig. S4A and Movie S1). Continuous density for the exchanged strand (es), that becomes excluded upon homologous pairing of the invading strand (is) with the complementary strand (cs) of the donor DNA, connects the arms of the donor DNA (Fig. S4B). The filament-bound esDNA follows a peripheral path along the filament groove that runs parallel to the isDNA (**Fig. 1D** [↗](#)). The conformation of the duplex formed by pairing of the isDNA and csDNA strands resembles closely that of the dsDNA in the post-synaptic filament (Fig. S4C)^{33,34} [↗](#), with sharp bends at the junctions where the donor DNA enters and exits the filament (**Fig. 1C, D** [↗](#)).

The arms of the donor DNA flanking the complementary region project away from the filament in opposite directions and with different tilt angles relative to the filament axis: the dsDNA arm closer to the 5'-end of the filament, hereinafter referred to as the 5'-arm, is almost perpendicular to the filament axis, with its filament-proximal end pointing slightly towards the 3'-end of the filament (3'-tilt), whereas the dsDNA arm closer to the 3'-end of the filament, hereinafter referred to as the 3'-arm, exhibits a more pronounced tilt in the opposite direction, towards the 5'-end of the ssDNA (5'-tilt) (**Fig. 1D** [↗](#) and Fig. S4D). The opposite tilt of the donor arms favours their connection via the esDNA strand in the D-loop structure.

Homologous pairing

A continuous span of 10 base-pair interactions is present between the isDNA and csDNA (**Fig. 1E** [↗](#) and Fig. S5A). Base pairing includes the entirety of the csDNA sequence designed to be homologous to the invading strand, with the exception of G26_{cs} (**Fig. 1E, F** [↗](#) and Fig. S5A): its guanine base adopts a minor-occupancy conformation in the direction of C8 in the isDNA but too distant for Watson-Crick hydrogen bonding, while the predominant conformation is pointing away (Fig. S5B).

Base pairing is fully maintained in the transition between the csDNA : isDNA duplex and the flanking arms of the donor DNA (**Fig. 1F, G** [↗](#)), with the exception of the aforementioned G26_{cs}. Unexpectedly, an additional, mismatched base pair is formed at the junction of the csDNA : isDNA duplex with the 3'-arm of the donor DNA, where T18_{is} becomes base paired with G16_{cs} (**Fig. 1E, G** [↗](#) and Fig. S5A). Thus, the complementary pairing observed in the D-loop structure does not entirely follow our experimental design, with one unrealised G26_{cs} : C8_{is} base pair and one mismatched G16_{cs} : T18_{is} base pair (Fig. S5). This observation indicates that factors additional to hydrogen-bonding potential contribute to determine pairing between the donor DNA and the invading strand of the filament.

Unwinding and strand invasion of donor DNA

Biochemical and structural analyses of pre- and post-synaptic filaments have revealed that RAD51 loops L1 and L2 mediate DNA binding, with prominent roles for amino acids L1 R235 and L2 V273 in partitioning filament DNA into nucleotide triplets^{33,44} [↗](#). However, L2 residues Q272 to P283 remained disordered in these filament structures (**Fig. 2A** [↗](#)).




Figure 2.

Donor DNA unwinding.

A Multiple sequence alignment of RAD51 L2 loop sequences, coloured according to conservation based on the Clustal scheme (*Hsa*: human, *Dre*: zebrafish, *Dme*: fly, *Ath*: arabidopsis, *Sce*: budding yeast, *Spo*: fission yeast). **B** Schematic drawing of the RAD51 D-loop. The N- and C-terminal domains of each RAD51 protomer are shown as spheres of size proportional to mass. RAD51 E, F, G and H are coloured in blue, the other protomers in pale blue. **C** Ribbon drawing of the L2 loop insertion into the donor DNA for RAD51-E, F, G and H. The L2 loops are drawn as thick blue tubes and the rest of the RAD51 protein is represented as transparent surface. The DNA strands are drawn as narrow tubes with the nucleotides shown as sticks; the DNA strands are coloured dark magenta except for the homologous sequences of the invading and complementary strands, coloured in pink. **D** Nucleotide sequences and observed base pairing of the D-loop DNA. Sequences are coloured in dark magenta and pink as in panel C. The approximate position of the L2 loop insertion in the unwound donor DNA is marked with the name of the RAD51 protomer. Vertical dashed lines mark the extent of donor DNA unwinding for each L2 insertion. **E-H** Cartoon drawings of the interface between the L2 loop of RAD51-E (panel E), RAD51-F (panel F), RAD51-G (panel G), RAD51-H (panel H) with the donor DNA. L2 loop amino acids and DNA-strand nucleotides are drawn as sticks and carbon-coloured as in panel C. The nucleotide bases are shown as filled rings and the Watson-Crick hydrogen bonds are drawn as dashed lines.

In our D-loop structure, the L2 sequences of four contiguous RAD51 protomers – E, F, G and H (**Fig. 2B**) – have become ordered and traceable in the density map, folding in random-coil conformation with 3_{10} -helix elements. The structure shows that filamentous RAD51 achieves strand separation by sequential L2-loop insertion of adjacent RAD51 protomers between the strands of the donor DNA (**Fig. 2C, D**). Insertion of the RAD51-H L2 loop at the 5'-arm junction causes a drastic widening of the minor groove, progressing to full separation of the complementary and exchanged strands upon L2-loop insertion by RAD51-G (**Fig. 2C, D, G, H**). In total, donor DNA unwinding by RAD51-H and -G exposes a continuous csDNA stretch of five nucleotides for homologous pairing: A22 - C25 of csDNA base pair with G9 - T12 of isDNA, physically organised as a base-pair triplet of (A22 - A24)_{cs} with (T10 - T12)_{is} and one unstacked C25_{cs} : G9_{is} base pair, while G26_{cs} remains unpaired (**Figs. 1F** and **2C, D, G, H**).

Successive L2-loop insertions extend the unwinding of donor DNA, with each insertion exposing a triplet of csDNA nucleotides for homology pairing with the invading strand (**Fig. 2C**). Thus, L2-loop insertions by RAD51-F and -E present (A19 - C21)_{cs} for base pairing with (G13 - T15)_{is}, and (G16 - T18)_{cs} for base pairing with (A16 - T18)_{is}, respectively (**Fig. 2C-F**). After completion of homologous pairing, the strands of the donor DNA rejoin in double-helical form at the 3'-arm junction (**Figs. 1G** and **2C**).

The L2 loop acts as a physical spacer, directly separating the unwound strands of the donor DNA. Its sequence is dominated by hydrophobic amino acids (**Fig. 2A**) that make extensive contacts with the ribose-phosphate backbone of the esDNA and culminate in stacking of F279's phenylalanine side chain against the last base pair of the 5'-arm duplex (RAD51-H) or the aromatic bases of nucleotides in the exchanged strand (RAD51-G, F, E) (**Fig. 2E-H**). The L2 loop displays a considerable degree of conformational heterogeneity, likely because each loop engages in both common and unique contacts with the DNA. This is best exemplified by the L2 loops of RAD51-E and -H that interact with the duplex arms of the donor DNA in different ways. RAD51-H L2 initiates donor DNA unwinding by buttressing the F279 side chain against G27_{cs} : C15_{is}, the last base pair of the 5'-arm duplex (**Fig. 2H**). Conversely, contact with RAD51-E L2 does not alter the conformation of the 3'-arm, as its F279 side chain stacks against C26 of the exchanged strand (**Fig. 2E**).

The asymmetric binding of RAD51-H and -E L2 loops at the boundaries between the arms of the donor DNA and the three-stranded synaptic DNA suggests that strand invasion and pairing follows a preferred direction. Thus, invasion by filament DNA likely begin at the 3'-end of the homology region in the complementary strand of the donor DNA and extends in the 5'- direction, as incrementally more nucleotides of csDNA are exposed by L2 loop insertions and become available for homologous pairing.

Capture of the exchanged strand

Strand invasion leads to displacement of the exchanged strand in the donor DNA. The structure of the D-loop shows that the esDNA is sequestered in a basic channel running parallel to the ssDNA-binding groove of the RAD51 nucleoprotein filament (**Fig. 3A, B**). One side of the channel is formed by RAD51 residues 303 to 313 (**Fig. 3C**), that fold in a beta hairpin on the inside of the filament groove, whereas the opposite side is formed by the C-terminal half of the L2-loop sequence (**Fig. 3B, D**). In the filament, the beta hairpins of successive protomers align to generate a continuous ssDNA-binding stripe (**Fig. 3D**). The esDNA is bound in the basic channel so that its aromatic bases remain largely exposed to solvent whereas the phosphodiester backbone faces the protein (**Fig. 3B, D**).

Three adjacent RAD51 protomers – F, G and H – engage the exchanged strand in the D-loop (**Fig. 3D**). The majority of contacts comprise charged interactions between four invariant RAD51 residues R303, K304, R306 and K313 and the phosphate groups of the esDNA (**Fig. 3E-G**). In addition to the electrostatic interactions, R306 is part of a GRG motif at the tip of the beta hairpin




Figure 3.

Capture of the exchanged strand.

A Nucleotide sequences and observed base pairing of the D-loop DNA. The esDNA sequence is in yellow, the rest of the D-loop DNA in light brown. The span of esDNA bound by each of RAD51-H, -G and -F is underlined. **B** Drawing of esDNA capture in a basic channel of the filament, aligned with the filament axis. The DNA is shown as full-atom model with filled rings for the ribose and the bases, coloured in yellow for the captured esDNA nucleotides and light brown for the rest. Filament RAD51 is shown in surface representation, coloured according to electrostatic potential from blue (basic) to red (acidic). **C** Multiple sequence alignment of RAD51 esDNA-binding sequences, coloured according to conservation based on the Clustal scheme (*Hsa*: human, *Dre*: zebrafish, *Dme*: fly, *Ath*: arabidopsis, *Sce*: budding yeast, *Spo*: fission yeast). **D** Ribbon drawing of the interface between the filament and the captured esDNA, highlighting the position of the esDNA-binding beta hairpins of RAD51-H, -G and -F in blue. The DNA is drawn as in panel B. The inset shows a schematic cartoon of the RAD51 D-loop, drawn as in [figure 2B](#), to emphasise the position of RAD51-H, -G, -F in the filament. **E-G** Drawings of the interface between the beta-hairpin amino acids of RAD51-H (panel E), RAD51-G (panel F), RAD51-F (panel G) and the esDNA. Beta-hairpin residues are drawn as sticks; amino acids are coloured according to atom type, whereas esDNA nucleotides are coloured as in panel D.

that comes into close contact with the phosphate-ribose backbone of the esDNA and makes stacking interactions with the aromatic bases of nucleotides A19_{es} and T25_{es} in RAD51-H and -G, respectively (Fig. 3E, F). The interaction of the 305-GRG-307 motif with esDNA resembles that of L1-loop 234-GRG-236 with csDNA in the homologous duplex of the D-loop.

Of the three RAD51 protomers F, G and H that bind the esDNA, RAD51-G makes the most extensive set of contacts, spanning five nucleotides, from A21_{es} to T25_{es}, involving all four basic residues R303, K304, R306 and K313 (Fig. 3F). Binding of RAD51-G to the esDNA is further strengthened by phosphate-group contacts of L2-loop K284 with C22_{es} and of R130 with C23_{es} and T24_{es}. RAD51-H contacts four nucleotides, from A17_{es} to A20_{es}, with R303, K304 and 305-GRG-307 (Fig. 3E), whereas RAD51-F makes a single long-range interaction between K313 and T25_{es} (Fig. 3G). In addition, R303 and R306 of RAD51-F engage the phosphate groups of T27, A28 and T29 of the 3'-arm duplex, possibly to guide the 3'-arm in the appropriate direction as it exits the filament groove, towards the RAD51-A NTD (Fig. 3G).

In total, the filament interacts with 9 of the 11 nucleotides of unpaired esDNA via contiguous interactions of the two RAD51 protomers G and H (5 and 4 nt, respectively). This compares with donor DNA opening by L2 loop insertion, which progresses by exposure of 3 nucleotides per RAD51 protomer.

Filament binding of donor DNA arms

The D-loop structure shows that the nucleoprotein filament interacts with the double-stranded arms of the donor DNA via the N-terminal domains of RAD51 protomers A and D (Fig. 4A, B). Although the position of the RAD51-A and -D NTDs relative to the donor DNA is broadly similar, their DNA-binding interfaces differ significantly in the extent and mode of their engagement with the 3'- and 5'-arm duplexes, respectively.

The RAD51-D NTD engages the minor groove of the 5'-arm DNA (Fig. 4C). The side chains of K39, K40 and K64 contact the phosphate groups of 10-GCTC-13 and 34-CG-35 in the upper and lower rungs of the minor groove respectively, while K70 and K73 contribute long-range electrostatic interactions to the binding (Fig. 4C, E). The association of RAD51-D NTD with 5'-arm is closer and more extensive than that of RAD51-A with the 3'-arm, and includes a direct hydrogen bond between the main-chain nitrogen of G65 and the phosphate group of T12 (Fig. 4C, E). In contrast, RAD51-A NTD binds in the major groove of the 3'-arm DNA, using electrostatic interactions mediated by the same set of lysine residues as in RAD51-D (Fig. 4D). However, the RAD51-A association with the 3'-arm involves a more limited set of direct contacts with the phosphate groups of C5 and 31-TTG-33 across the major groove (Fig. 4D, E).

The interaction of the NTD with the arms of the donor DNA highlights a lack of direct contacts with the bases and the predominantly electrostatic nature of the association. Such plasticity in the NTD - DNA interaction might be required for effective D-loop formation, as the direction at which the 3'-arm exits the filament upon D-loop formation is a function of the number of nucleotides involved in homologous pairing.

Structure-based mutagenesis of RAD51 residues important for D-loop formation

The atomic model of the RAD51 D-loop has revealed several new protein - DNA interfaces, including those involved in strand separation, capture of the exchanged strand and anchoring of the flanking DNA arms to the filament. To assess the relative importance and contribution of RAD51 residues involved in D-loop formation, we performed alanine mutagenesis of F279 (strand separation); R303, K304, R306, K313 (esDNA capture); K39, K40, K64, K70 and K73 (arm duplex binding) (Fig. 5A).




Figure 4.

Filament engagement of the donor DNA arms.

A Drawing of the D-loop structure that highlights the interaction of RAD51-A and -D with the 3'- and 5'-arms of the duplex donor DNA, respectively. RAD51-A and -D are drawn as cylinder cartoons in pale blue with their NTDs in blue. The donor DNA arms are shown as molecular surfaces in light brown while the rest of the DNA is drawn as narrow tubes. The filament is shown as a transparent surface. The inset shows a schematic cartoon of the D-loop structure to highlight the position of the RAD51-A and -D protomers. **B** Multiple sequence alignment of RAD51 NTD sequences, coloured according to conservation based on the Clustal scheme (*Hsa*: human, *Dre*: zebrafish, *Dme*: fly, *Ath*: arabidopsis, *Sce*: budding yeast, *Spo*: fission yeast). **C**, **D** The interaction of RAD51-D and -A NTD with the 5'- and 3'-arms, respectively. The protein is drawn as a ribbon, with the side chains as sticks. The hydrogen bond between the main-chain nitrogen of G65 and the phosphate group of T12 is drawn as a dashed line. The DNA is in spacefill representation; the nucleotides contacted by the NTD are coloured yellow, and the rest of the DNA is light brown. **e** Nucleotide sequence of the donor DNA, coloured as in panels C and D.




Figure 5.

Structure-based mutagenesis of the RAD51 D-loop.

A Ribbon drawing of RAD51, highlighting residues targeted for alanine mutagenesis, colour-coded according to their role in D-loop formation (F279 in pink; R303, K304, R306, K313 in magenta; K39, K40, K64, K70, K73 in light blue). **B** Strand-exchange assay showing the change in fluorescence for each RAD51 mutant following injection of the complementary strand after 9 minutes, with traces coloured according to amino acid as in panel A. Error bars denote the standard deviation of three independent replicates. The upper panel shows a diagrammatic representation of the assay, with DNA shown as light or dark brown lines and the fluorophore and quencher represented as in the key. **C** Schematic drawing of the experimental setup for single molecule analysis of RAD51 D-loop formation. **D** Kymograph of a representative D-loop formation event. The upper panel shows a time-lapse fluorescence intensity scan of the λ DNA, with a stable binding event at the expected target site (10.6 kb, indicated by the arrow). The white dash line marks the first 60-second interval that was used in frame acquisition. The lower panel shows a summed projection of the fluorescence signal over the acquisition time. **E** Quantification of RAD51 filament binding events. The left-side histogram displays the binding profile of wild-type RAD51 (0.5 μ M), showing a Gaussian distribution of binding events centered on the expected position of the complementary sequence ($\sim$ 10.6 kb), consistent with specific ssDNA pairing and D-loop formation. The right-side histogram shows the binding profile of the KK39,40AA RAD51 double mutant (2.5 μ M). **F** Mechanism of D-loop formation by RAD51 nucleoprotein filaments.

The designed mutations are distant from the binding site of ssDNA in filamentous RAD51, and therefore are not predicted to reduce the ability of the mutant protein to form nucleoprotein filaments on ssDNA. As expected, EMSA analysis showed that the F279 and NTD mutants retained wild-type ability to form filaments on ssDNA (Fig. S6A). However, RAD51 mutants R303A, K304A, R306A and K313A showed a clear decrease in ssDNA binding (Fig. S6A), suggesting that, in addition to their role in ssDNA capture, they might mediate the initial contact of the incoming ssDNA during pre-synaptic filament nucleation. All RAD51 mutants showed a moderate decrease in affinity for dsDNA, with the exception of K304 (Fig. S6B).

We evaluated the ability of the alanine mutants to perform DNA-strand exchange using an *in vitro* assay that measures the increase in fluorescent signal when a fluorescently-labelled 60-nt oligo that is quenched in duplex form is strand-exchanged with the unlabelled sequence (Fig. 5B).⁴⁵ The results of the strand-exchange assay revealed that all mutants decreased the efficiency of the reaction, but with varying degrees of impairment (Fig. 5B). The L2 loop F279A showed a modest impact, in agreement with earlier findings,⁴⁴ whereas the ssDNA-capture mutants R303A, K304A, R306A and K313A displayed a more pronounced effect. Notably, the NTD mutations showed the largest negative impact on strand exchange: the double lysine-to-alanine substitutions of K39, K40 and K70, K73 reduced strand exchange to a fraction of the yield for the wild-type protein.

To provide further support for the role of the RAD51 NTD in D-loop formation, we developed a single-molecule D-loop assay using fluorescence imaging with optical tweezers.⁴⁶ In the assay, RAD51 filaments were reconstituted on an Alexa 488-labeled 49-nt ssDNA designed to anneal at position 10.6 kb of *I* DNA (Fig. 5C). The filaments interacted with doubly-biotinylated *I* DNA held between streptavidin-coated beads and D-loop formation was monitored by the presence of a fluorescent signal at the expected locus on *I* DNA. Wild-type RAD51 filaments were capable of D-loop formation at the expected site, as assessed by kymograph and binding events analysis (Fig. 5D, E and Fig. S7). In contrast, filaments of the KK39,40AA RAD51 double mutant failed to show measurable binding events; a 5-fold increase in the concentration of the double mutant protein showed non-specific binding and no stable D-loop formation at the expected *I* DNA site (Fig. 5E and Fig. S7).

These findings validate our model for the D-loop structure and highlight the important role of the RAD51 NTD in the strand-exchange reaction, by anchoring the donor dsDNA to the nucleoprotein filament in the correct orientation for the strand invasion and homology search. The biochemical behaviour of our ssDNA-capture site mutants agrees with the impaired D-loop formation of a yeast RAD51 protein bearing alanine mutations at amino acids corresponding to R130, K303 and K313 of human RAD51.⁴⁷

Discussion

The architecture of the RAD51 D-loop structure reveals the mechanism of strand invasion and homology search within a synaptic RAD51 filament. The close engagement of the RAD51-D NTD with the 5'-arm duplex, coupled to minor-groove widening by the RAD51-H L2 loop that initiates DNA unwinding, indicates that synapsis formation starts by interaction of the filament with the 5'-arm of the donor DNA. In comparison, the interaction of the RAD51-A NTD with the 3'-arm is more limited and insertion of the L2 loop of RAD51-E does not remodel the proximal end of the 3'-arm duplex. Furthermore, the first ssDNA : dsDNA base pair formed at the 3'-arm junction is a T : G mismatch that is unlikely to stabilise the initial strand pairing.

Our findings form the basis for a mechanistic model of strand exchange by a RAD51 filament (Fig. 5F). In the model, RAD51-D NTD binding directs the donor duplex towards the filament groove, where L2-loop insertions by RAD51-H and -G initiate strand separation and capture of the

exchanged strand. Such unwinding exposes 5 nucleotides of csDNA, of which 4 are available for pairing with the invading strand of the filament. Initial base pair formation in the presence of homology would drive further donor DNA unwinding by L2-loop insertions of RAD51 -F and -E; each insertion exposes a triplet of csDNA nucleotides, in agreement with experimental evidence that homologous pairing progresses in triplet steps^{39,41}. Concomitant capture of the exchanged strand by RAD51-G after insertion of a third L2 loop by RAD51-F would lead to sequestration of nine esDNA nucleotides, contributing to the formation of a stable synaptic intermediate.

Successful homology pairing drives the engagement of the 3'-arm of the donor dsDNA with the NTD of the RAD51 molecule three protomer positions away towards the 3'-end of the filament, completing D-loop formation. The observed plasticity in the interaction of the NTD with dsDNA would facilitate capture of the 3'-arm DNA emerging at slightly different angles from the filament axis, depending on the exact number of paired nucleotides. Overall, the structural features of the RAD51 D-loop provide a strong indication that strand pairing and exchange begins at the 3'-end of the complementary strand in the donor DNA and progresses with 3'-to-5' polarity (**Fig. 5F**).

The observation of an unrealised base pair involving G26_{cs} at the 3'-end of the unwound csDNA, which was designed to base pair with C8_{is}, indicates that the mechanism of unwinding of the 5'-arm prevents base pairing of the first available csDNA nucleotide, likely due to physical constraints. Further unwinding of the 5'-arm to allow G26_{cs} pairing was not observed, possibly because the presence of three consecutive G : C base pairs in the 5'-arm duplex acted as a GC clamp.

The presence of a T18_{is} : G16_{cs} mismatch as the last base pair at the 5'-end of the paired csDNA is intriguing (Fig. S5B). Of note, T18_{is} and G16_{cs} do not form a 'wobble' base pair⁴⁸, indicating that fulfilment of hydrogen bonding potential is not critical. Rather, mismatch pairing that completes a base-pair triplet might occur regardless of perfect homology, to achieve a full complement of base stacking and protein - DNA interactions with RAD51 L1 and L2 loops. This observation provides a potential mechanism for how RAD51 can tolerate mismatches at certain positions of the homologous sequence during strand exchange.

The observed 3'-to-5' polarity of exchange of the complementary strand by RAD51 is opposite to the 5'-to-3' polarity of bacterial RecA (Fig. S8), that was determined based on cryoEM structures of RecA D-loops⁴². Comparison of RAD51 and RecA D-loop structures reveals that their different polarity follows from the opposite position occupied by their dsDNA-binding domains in the filament: in a vertically-oriented filament with the 5'-end of the ssDNA at the top and the 3'-end at the bottom, the RAD51 NTDs line the bottom of the filament groove whereas the RecA CTDs hang from the top (Fig. S9). The shared mode of minor-groove binding by RAD51-NTD and RecA-CTD and L2 loop insertion into the minor groove half a turn away leads naturally to opposite polarity in the progression of strand invasion and pairing with the complementary strand of the donor DNA. Our mechanistic interpretation of static D-loop structures awaits full reconciliation with earlier efforts to determine strand-exchange polarity for RecA and RAD51 measured under a variety of experimental conditions⁴⁹⁻⁵².

Altogether, the available evidence suggests that the ability to perform DNA-strand exchange by members of the RecA-family of recombinases results from a process of convergent evolution (Fig. S10). Thus, an ancestral RecA-fold ATPase protein might first have evolved to bind ssDNA using the L1 and L2 loops, possibly as a means of protecting the exposed DNA template during DNA replication. Independent acquisition of dsDNA-binding by RecA and RAD51 via incorporation of carboxy- and amino-terminal domains, respectively might have further endowed these proteins with the ability to engage simultaneously ss- and dsDNA and catalyse the exchange of homologous DNA strands.

Materials and methods

RAD51 expression and purification

Human RAD51 was expressed and purified as described⁵³. Briefly, RAD51 was co-expressed with a HisTag - MBP (Maltose Binding Protein) - BRCA2 BRC4 fusion protein in Rosetta BL21(DE3) *E. coli* cells, and purified over successive steps of Ni²⁺ chromatography, Heparin chromatography and size-exclusion chromatography. The purified protein was concentrated and stored in small aliquots at -70°C. Structure-based RAD51 mutations were introduced according to standard protocols and the mutant RAD51 proteins were purified as for the wild-type protein.

DNA substrate preparation

DNA oligos were purchased from Integrated DNA Technologies (IDT). Unmodified or biotinylated oligos were supplied PAGE-purified. Fluorescently-labelled oligos were HPLC-purified. All oligos were resuspended to 100 µM in TE buffer (10 mM Tris pH 7.5, 1 mM EDTA). dsDNA reagents were prepared at a concentration of 10 µM duplex in TE buffer, by mixing equimolar concentrations of the constituent complementary ssDNA oligos and heating at 95°C for 5 minutes before slowly cooling to room temperature.

For the strand-exchange assay, a duplex DNA 60mer in which one strand was 5'-labelled with fluorescein and the complementary strand 3'-labelled with an Iowa Black quencher was prepared by mixing quencher strand to fluorescent strand in a 1.2 : 1 molar ratio to ensure full quenching, and purified on 10% native PAGE.

All DNA sequences can be found in Table S1.

Electrophoretic mobility shift assay (EMSA)

EMSA reactions were prepared in buffer: 25 mM HEPES pH 7.5, 150 mM NaCl, 2 mM DTT, 5 mM CaCl₂ and 2 mM ATP, by mixing fluorescein-labelled ss- or dsDNA 60mer at 250 nM concentration with increasing concentrations of wild-type or mutant RAD51 (0.5, 1.0, 2.0, 4.0, 6.0, 8.0, and 10.0 µM) in a 20 µL volume, and incubated at room temperature for 10 min. 10 µL of 50% sucrose (w/v) was added to each reaction mixture prior to electrophoresis, and samples were resolved on a 0.5 % agarose gel in TB buffer (45 mM Tris, 45 mM boric acid) at 45 V for 2 hours at 4°C.

For D-loop reconstitution experiments, EMSA reactions were prepared in same buffer by mixing 500 nM of either Cy5-labelled ssDNA or doubly-biotinylated ssDNA with increasing concentrations of RAD51 (1.0, 2.5, 5.0, 10.0, 20.0, 30.0 µM), followed by incubation at room temperature for 10 minutes. 500 nM of either Cy3-labelled dsDNA or doubly-biotinylated Cy3-labelled DNA was then added to the reaction and incubated for a further 10 minutes before electrophoresis. In experiments with mono-streptavidin (mSA) capped DNA, mSA was first added to the DNA at 5x molar excess over biotin labels and incubated for 15 minutes at room temperature before adding it to EMSA reactions. SAvPhireTM monomeric streptavidin (mSA) was purchased from Sigma Aldrich and resuspended to a working stock of 4 mg/ml from lyophilised powder according to supplier instructions.

In experiments where unlabelled DNA was used, the gel was stained in a 2x SYBR Gold solution made up in milliQ water (10 min, 25°C) and then destained in milliQ water. Gels were visualised using a Typhoon FLA 9000 fluorescent gel imager, by excitation at 473 nm for fluorescein-labelled DNA or SYBR Gold staining, 532 nm excitation for Cy3-labelled DNA or 635 nm excitation for Cy5-labelled DNA.

DNA Strand Exchange (SE) Assay

Strand exchange assays were performed using a PHERAstar FS instrument (BMG Labtech) equipped with a fluorescence intensity optic module (excitation at 485 nm, emission at 520 nm, 10 nm bandwidth). The gain was calibrated to 80 % of the maximum fluorescence intensity using a 25 nM sample of fluorescein-labelled single-stranded DNA in strand-exchange buffer: 150 mM NaCl, 25 mM HEPES pH 7.5, 2 mM DTT, 2 mM CaCl₂, 2 mM ATP, 0.1 mg/ml BSA.

Reactions were prepared in 50 µL volumes of strand-exchange buffer containing 500 nM RAD51 and 50 nM ssDNA 60mer, corresponding to a 1 : 2 ratio of RAD51 to RAD51-binding sites in the DNA. RAD51 was incubated with ssDNA at 30°C for 9 minutes prior to initiation of the strand-exchange reaction by 50 µL injection of dsDNA labelled with fluorescein at the 5'-end of one strand and with the Iowa Black quencher at the 3'-end of the complementary strand, using the PHERAstar FI Slow Kinetics mode. Fluorescence measurements began 2 seconds after duplex injection, and data were collected at 1-minute intervals for an additional 30 minutes. As a specificity control for strand exchange, the reaction was performed with an oligo(dT) ssDNA 60mer. Blank measurements were performed by injecting dsDNA into strand-exchange buffer lacking both ssDNA and RAD51. All reactions were performed in triplicate.

CryoEM

Sample preparation

5 µM RAD51 was first incubated with 250 nM ssDNA in buffer: 150 mM NaCl, 25 mM HEPES pH 7.5, 2 mM DTT, 2 mM ATP, 5 mM CaCl₂, for 10 minutes to assemble a nucleoprotein filament; 250 nM dsDNA was then added to the sample and incubated for a further 10 minutes. In experiments with mSA-capped DNA, mSA was first added to the DNA at 5x molar excess over biotin labels and incubated for 15 minutes before the addition of other components.

Vitrification protocol

UltrAuFoil R1.2/R1.3 300 mesh gold grids (Quantifoil) were glow-discharged twice for 1 minute using a PELCO easiGlow system (0.38 mBar, 30 mA, negative polarity). 3 µl of sample was applied to the mesh side of each grid before plunge-freezing in liquid ethane using a Vitrobot Mark IV robot (FEI), set to 100 % humidity, 4°C, 2 second blot time and -3 blot force.

Grid screening and data collection

Grids were screened using a Talos Arctica 200 keV transmission electron microscope fitted with a Falcon III direct electron detector (Thermo Fisher). Data were collected using a Titan Krios G3 300 keV transmission electron microscope (Thermo Fisher) fitted with a K3 (Gatan) direct electron detector, using the EPU package (Thermo Fisher). Two datasets were collected: a first dataset with free-end DNA (#1; 2792 movies in super-resolution mode at 0.326 Å/pixel) and a second dataset with mSA-capped ss- and dsDNA (#2, 8960 movies in super-resolution mode at 0.326 Å/pixel and 2xbinned). All data were recorded at the cryoEM facility of the University of Cambridge in the Department of Biochemistry. Data collection parameters are reported in Table S2.

Data processing

Data was processed using CryoSparc 4.4.1⁵⁴[54](#). Micrographs were motion corrected with Patch Motion Correction with output F-crop factor set to ½. The Contrast Transfer Function was estimated using Patch CTF Estimation and exposures were manually curated to remove outliers.

Dataset #1 (Figs. S1, 2). 1477 particles were manually picked from 2711 micrographs and, after 2D classification, 578 particles were selected to generate filament templates for automated picking. Template picking yielded 948,693 particles, which were reduced to 838,403 after outlier removal. 705,678 particles were extracted using a 400-pixel box. Multiple rounds of 2D classification led to the identification of a set of 2522 particles that yielded classes with clear evidence of synaptic filaments and was therefore used for Topaz training. The trained Topaz model picked 66,799 particles, which were manually curated to 64,942 particles; of these, 60,328 were extracted with a 416-pixel box. Ab Initio reconstruction of two 3D volumes followed by Heterogeneous Refinement yielded a 3D shape of a recognisable D-loop structure. The corresponding set of 44,864 particles was subject to 3D classification into 5 classes, three of which presented pronounced D-loop features with strong density for the synaptic dsDNA; these classes were pooled and the combined 29,398 particles refined by Homogenous Refinement to 4.43 Å. The resulting particle set was used for a second round of Topaz training and picking, which yielded 167,005 particles and from which 153,235 particles were extracted with a 416-pixel box. The particles were used directly for Ab Initio reconstruction and Heterogeneous Refinement, generating a D-loop structure from a set of 117,041 particles. These were fractionated into 10 classes by 3D classification, and the particles for four of these classes that presented the best D-loop shapes (50,363 particles) were pooled and subject to Homogenous Refinement to 3.62 Å, followed by Non-Uniform Refinement to yield a final D-loop reconstruction at 3.31 Å.

Dataset #2 (Figs. S1-3). The final 3D reconstruction obtained from dataset #1 processing was low-pass filtered at 8 Å and used to generate templates for particle picking. 2,940,045 particles were picked from 8834 micrographs, 2,680,398 particles were selected after manual curation and 2,245,379 particles extracted with a 208-pixel box. After multiple rounds of 2D classification, classes from a 19,116 particle set showed clear synaptic features and were used for Ab Initio reconstruction and Homogenous Refinement of a 3D volume from 13,626 particles, which presented the expected filament shape and D-loop features such as density attributable to the dsDNA arms of the target DNA. These particles were used for Topaz training and picking of 415,845 particles, of which 413,320 particles were extracted with a 208-pixel box. The extracted particles were fed directly into Ab Initio reconstruction and Heterogeneous Refinement, which yielded a 3D volume of a filament with recognisable D-loop features at lower map contouring for 328,724 extracted particles. 3D classification into 10 classes identified a single class of 43,824 particles of a complete D-loop structure, which refined in Non-Uniform Refinement to 2.96 Å. This particle set was used in a second round of Topaz training and picking that yielded 922,204 particles, of which 900,213 were extracted with a 208-pixel box and used directly in ab Initio reconstruction and Heterogenous Refinement of a filament volume of 711,480 particles with recognisable D-loop features at lower map contouring. 3D classification of this particles set over 10 classes identified one class of 74,483 particles corresponding to a high-quality D-loop structure, which was improved by further classification and refinement by Heterogenous Refinement. The resulting set of 102,679 particles was subject to Non-Uniform refinement to yield a 3D reconstruction at 2.72 Å, followed by Reference-Based Motion Correction of 102,596 particles and another step of Non-Uniform refinement, to yield a final reconstruction at 2.64 Å resolution.

Model building and refinement

An atomic model comprising nine RAD51-ATP protomers, 26 nucleotides of ssDNA and 41 nucleotides for each strand of the 'mismatch bubble' dsDNA was built into the 3D volume of the final dataset #2 reconstruction at 2.64 Å resolution using Coot⁵⁵. Fitting of the model to the map was improved by real-space refinement in Phenix⁵⁶ and model rebuilding in Coot⁵⁵. In the later stages of refinement, map fitting and model stereochemistry were improved using ISOLDE⁵⁷. The register of the DNA sequence in the structure could be unambiguously assigned based on fitting of the nucleoside bases in the homologous dsDNA region, where the map resolution was highest.

Optical-tweezers based single-molecule fluorescence imaging

To reconstitute RAD51 D-loop formation at the single-molecule level, we used a C-Trap® (LUMICKS) instrument that combines optical tweezers and confocal fluorescence microscopy. The experimental setup featured a microfluidic chip with four distinct flow channels separated by laminar flow: one for streptavidin-coated polystyrene beads (Spherotech, 4.43 μm diameter), one for biotinylated λ DNA, and two for buffer exchange or RAD51 nucleoprotein filaments reservoirs.

Doubly-biotinylated λ DNA molecules (48.5 kbp) prepared as previously described³⁷ were tethered between two streptavidin-coated beads in the microfluidic flow cell. Successful capture of a single λ DNA was confirmed by measuring force-extension curves⁵⁸. The optical traps were calibrated to a stiffness of 0.4 pN/nm, enabling stable manipulation of the DNA tether under tension^{59,60}. Fluorescence imaging was performed using a single 488 nm laser for excitation with an illumination intensity of 2 μW . Fluorescence signals were detected on a highly sensitive single-photon counting detector. Confocal fluorescence scans were performed over a region of 300×50 pixels to visualize the DNA tether and the edges of the beads. Pixel width was set to 0.1 μm and each pixel was illuminated for 0.1 ms. During imaging, DNA was held at a constant force of 20 pN, and the flow channels were closed.

RAD51 D-loop formation reactions were examined by incubating 20 μM RAD51 with 1 μM Alexa488-labelled ssDNA 49mer in buffer: 25 mM HEPES pH 7.5, 150 mM NaCl, 2 mM ATP, 2 mM CaCl_2 , 2 mM DTT at room temperature for 20 minutes, 40-fold diluted in the same buffer and injected into the microfluidic flow cell for analysis of the interaction with the λ DNA. The 49 nt-long sequence of the ssDNA is complementary to nucleotides 10,617 to 10,666 of λ DNA, to promote D-loop formation at a single, specific site on the DNA tether. During imaging, the DNA tether tension was maintained at 20 pN, ensuring optimal conditions for detecting RAD51 interactions along the DNA substrate.

Data Extraction and Analysis

Two-dimensional (2D) fluorescence scans (300×50 pixels) of individual λ DNA molecules were recorded for a total acquisition time of 180 seconds at a temporal resolution of one frame every 2 seconds. Given that RAD51 binding induces DNA extension^{61,62}, leading to drift of the D-loop position under the constant applied force regime, only the initial 30 frames were considered for analysis to ensure accurate localization. The stack of frames was summed using the open-source image processing package FIJI (ImageJ), to enhance fluorescence signal detection by summing pixel intensity values across all frames in a stack⁶³. Fluorescence intensity profiles were extracted by applying a linear scan along the DNA contour and bead peripheries, excluding regions near the bead peripheries to minimise fluorescence distortion, enabling precise quantification of binding events.

Due to fluorescence interference and reduced sensitivity near the beads, regions adjacent to each bead were excluded from analysis. The extent of this exclusion region was determined by performing a linear fit of the loss of fluorescence intensity measured at the junction areas versus the physical diameter of the beads and the contour length of λ DNA. Based on this analysis, the exclusion region was estimated to be approximately 0.75 kbp from each end of the DNA molecule.

To automate peak identification, fluorescence intensity profiles were plotted against position along λ -DNA. Using the peak fitting module in Origin Pro (OriginLab), fluorescence intensity peaks were identified and fitted with a 2D-Gaussian model which tracks each peak using a symmetrical bell-shaped curve characterized by amplitude, width (full width at half maximum) and center position (X-centre) to accurately locate and quantify binding events^{64,65}. The X-centre values of the detected peaks were extracted to determine the precise binding positions along λ DNA.

Binding event frequencies were binned at 1 kbp intervals along the DNA molecule length. Similar peak detection and Gaussian fitting were applied to the histogram of binding events, allowing for the identification of the focal binding site.

Data and materials availability

The coordinates and cryoEM map of the RAD51 D-loop structure have been deposited in the Protein Data Bank (accession code [9I62](#)) and Electron Microscopy Data Bank (accession code [EMD-52646](#)).

Acknowledgements

We would like to thank Chen Qian for early work on the project and the Biochemistry cryoEM facility team for assistance with data collection. This work was funded by a Wellcome Trust Investigator award to LP (221892/Z/20/Z) and a UKRI BBSRC PhD studentship award to LP and RA (grant number 2271086).

Additional information

Author contributions

LJ produced wild-type and mutant RAD51 proteins, performed the EMSA and strand-exchange assays, designed and analysed the optical-tweezers single-molecule imaging experiments, collected cryoEM data and supervised REA. REA performed the biochemical reconstitution of the D-loop, collected cryoEM data and performed an initial cryoEM analysis. JDM carried out early work on the RAD51 NTD mutants, supervised and assisted REA and LJ throughout the project. LP conceived and directed the project, solved the 3D structure of the D-loop and wrote the paper with input from all authors.

Funding

Wellcome Trust

<https://doi.org/10.35802/221892>

BBSRC DTP (2271086)

Additional files

Supplementary figures and tables

References

1. Kowalczykowski SC (2015) **An Overview of the Molecular Mechanisms of Recombinational DNA Repair** *Cold Spring Harbor Perspectives in Biology* **7**:a016410 <https://doi.org/10.1101/cshperspect.a016410> | Google Scholar
2. Wright WD, Shah SS, Heyer W-D (2018) **Homologous recombination and the repair of DNA double-strand breaks** *Journal of Biological Chemistry* **293**:10524–10535 <https://doi.org/10.1074/jbc.tm118.000372> | Google Scholar
3. Chakraborty S, Schirmeisen K, Lambert SA (2023) **The multifaceted functions of homologous recombination in dealing with replication-associated DNA damages** *DNA Repair* **129**:103548 <https://doi.org/10.1016/j.dnarep.2023.103548> | Google Scholar
4. Triplett MK, Johnson MJ, Symington LS (2024) **Induction of homologous recombination by site-specific replication stress** *DNA Repair* **142**:103753 <https://doi.org/10.1016/j.dnarep.2024.103753> | Google Scholar
5. Brown MS, Bishop DK (2014) **DNA Strand Exchange and RecA Homologs in Meiosis** *Cold Spring Harbor Perspectives in Biology* **7**:a016659 <https://doi.org/10.1101/cshperspect.a016659> | Google Scholar
6. Prakash R, Zhang Y, Feng W, Jasin M (2015) **Homologous Recombination and Human Health: The Roles of BRCA1, BRCA2, and Associated Proteins** *Cold Spring Harbor Perspectives in Biology* **7**:a016600 <https://doi.org/10.1101/cshperspect.a016600> | Google Scholar
7. Matos-Rodrigues G, Guirouilh-Barbat J, Martini E, Lopez BS (2021) **Homologous recombination, cancer and the ‘RAD51 paradox.’** *NAR Cancer* **3**:zcab016 <https://doi.org/10.1093/narcan/zcab016> | Google Scholar
8. Lord CJ, Ashworth A (2017) **PARP inhibitors: Synthetic lethality in the clinic** *Science* **355**:1152–1158 <https://doi.org/10.1126/science.aam7344> | Google Scholar
9. Dibitetto D, Widmer CA, Rottenberg S. (2024) **PARPi, BRCA, and gaps: controversies and future research** *Trends in Cancer* **10**:857–869 <https://doi.org/10.1016/j.trecan.2024.06.008> | Google Scholar
10. Baumann P, Benson FE, West SC (1996) **Human Rad51 Protein Promotes ATP-Dependent Homologous Pairing and Strand Transfer Reactions In Vitro** *Cell* **87**:757–766 [https://doi.org/10.1016/s0092-8674\(00\)81394-x](https://doi.org/10.1016/s0092-8674(00)81394-x) | Google Scholar
11. Sung P (1994) **Catalysis of ATP-Dependent Homologous DNA Pairing and Strand Exchange by Yeast RAD51 Protein** *Science* **265**:1241–1243 <https://doi.org/10.1126/science.8066464> | Google Scholar
12. Lin Z, Kong H, Nei M, Ma H (2006) **Origins and evolution of the recA/RAD51 gene family: Evidence for ancient gene duplication and endosymbiotic gene transfer** *Proceedings of the*

National Academy of Sciences **103**:10328–10333 <https://doi.org/10.1073/pnas.0604232103> | [Google Scholar](#)

13. Chintapalli SV, et al. (2013) **Reevaluation of the evolutionary events within recA/RAD51 phylogeny** *BMC Genomics* **14**:240 <https://doi.org/10.1186/1471-2164-14-240> | [Google Scholar](#)
14. Cejka P, Symington LS (2021) **DNA End Resection: Mechanism and Control** *Annual Review of Genetics* **55**:285–307 <https://doi.org/10.1146/annurev-genet-071719-020312> | [Google Scholar](#)
15. Renkawitz J, Lademann CA, Jentsch S (2014) **Mechanisms and principles of homology search during recombination** *Nature Reviews Molecular Cell Biology* **15**:369–383 <https://doi.org/10.1038/nrm3805> | [Google Scholar](#)
16. Greene EC (2016) **DNA Sequence Alignment during Homologous Recombination** *Journal of Biological Chemistry* **291**:11572–11580 <https://doi.org/10.1074/jbc.r116.724807> | [Google Scholar](#)
17. Shibata T, DasGupta C, Cunningham RP, Radding CM (1979) **Purified Escherichia coli recA protein catalyzes homologous pairing of superhelical DNA and single-stranded fragments** *Proceedings of the National Academy of Sciences* **76**:1638–1642 <https://doi.org/10.1073/pnas.76.4.1638> | [Google Scholar](#)
18. Jain SK, Cox MM, Inman RB (1995) **Occurrence of three-stranded DNA within a RecA protein filament** *The Journal of biological chemistry* **270**:4943–4949 [Google Scholar](#)
19. Piazza A, et al. (2019) **Dynamic Processing of Displacement Loops during Recombinational DNA Repair** *Molecular Cell* **73**:1255–1266 <https://doi.org/10.1016/j.molcel.2019.01.005> | [Google Scholar](#)
20. Bell JC, et al. (2023) **BRCA2 chaperones RAD51 to single molecules of RPA-coated ssDNA** *Proceedings of the National Academy of Sciences* **120**:e2221971120 <https://doi.org/10.1073/pnas.2221971120> | [Google Scholar](#)
21. Greenhough LA, et al. (2023) **Structure and function of the RAD51B–RAD51C–RAD51D–XRCC2 tumour suppressor** *Nature* **619**:650–657 <https://doi.org/10.1038/s41586-023-06179-1> | [Google Scholar](#)
22. Taylor MRG, et al. (2015) **Rad51 Paralogs Remodel Pre-synaptic Rad51 Filaments to Stimulate Homologous Recombination** *Cell* **162**:271–286 <https://doi.org/10.1016/j.cell.2015.06.015> | [Google Scholar](#)
23. Rawal Y, et al. (2023) **Structural insights into BCDX2 complex function in homologous recombination** *Nature* **619**:640–649 <https://doi.org/10.1038/s41586-023-06219-w> | [Google Scholar](#)
24. Ito M, et al. (2023) **FIGNL1 AAA+ ATPase remodels RAD51 and DMC1 filaments in pre-meiotic DNA replication and meiotic recombination** *Nature Communications* **14**:6857 <https://doi.org/10.1038/s41467-023-42576-w> | [Google Scholar](#)
25. Carver A, et al. (2024) **Molecular basis of FIGNL1 in dissociating RAD51 from DNA and chromatin** *Science* **387**:426–431 <https://doi.org/10.1126/science.adr7920> | [Google Scholar](#)
26. Ogawa T, Yu X, Shinohara A, Egelman EH (1993) **Similarity of the Yeast RAD51 Filament to the Bacterial RecA Filament** *Science* **259**:1896–1899 <https://doi.org/10.1126/science.8456314> | [Google Scholar](#)

27. Yu X, et al. (2001) **Domain structure and dynamics in the helical filaments formed by RecA and Rad51 on DNA** *Proceedings of the National Academy of Sciences* **98**:8419–8424 <https://doi.org/10.1073/pnas.111005398> | Google Scholar
28. Story RM, Weber IT, Steitz TA (1992) **The structure of the E. coli recA protein monomer and polymer** *Nature* **355**:318–325 <https://doi.org/10.1038/355318a0> | Google Scholar
29. Pellegrini L, et al. (2002) **Insights into DNA recombination from the structure of a RAD51–BRCA2 complex** *Nature* **420**:287–293 <https://doi.org/10.1038/nature01230> | Google Scholar
30. Conway AB, et al. (2004) **Crystal structure of a Rad51 filament** *Nature Structural & Molecular Biology* **11**:791–796 <https://doi.org/10.1038/nsmb795> | Google Scholar
31. Wu Y, et al. (2004) **Crystal Structure of Archaeal Recombinase RadA A Snapshot of Its Extended Conformation** *Molecular Cell* **15**:423–435 <https://doi.org/10.1016/j.molcel.2004.07.014> | Google Scholar
32. Short JM, et al. (2016) **High-resolution structure of the presynaptic RAD51 filament on single-stranded DNA by electron cryo-microscopy** *Nucleic acids research* **44**:9017–9030 <https://doi.org/10.1093/nar/gkw783> | Google Scholar
33. Xu J, et al. (2017) **Cryo-EM structures of human RAD51 recombinase filaments during catalysis of DNA-strand exchange** *Nature Structural & Molecular Biology* **24**:40–46 <https://doi.org/10.1038/nsmb.3336> | Google Scholar
34. Appleby R, et al. (2023) **A metal ion-dependent mechanism of RAD51 nucleoprotein filament disassembly** *iScience* **26**:106689 <https://doi.org/10.1016/j.isci.2023.106689> | Google Scholar
35. Chen Z, Yang H, Pavletich NP (2008) **Mechanism of homologous recombination from the RecA–ssDNA/dsDNA structures** *Nature* **453**:489–494 <https://doi.org/10.1038/nature06971> | Google Scholar
36. Galletto R, Amitani I, Baskin RJ, Kowalczykowski SC (2006) **Direct observation of individual RecA filaments assembling on single DNA molecules** *Nature* **443**:875–878 <https://doi.org/10.1038/nature05197> | Google Scholar
37. Candelli A, et al. (2014) **Visualization and quantification of nascent RAD51 filament formation at single-monomer resolution** *Proc Natl Acad Sci U S A* **111**:15090–15095 <https://doi.org/10.1073/pnas.1307824111> | Google Scholar
38. Hsieh P, Camerini-Otero CS, Camerini-Otero RD (1992) **The synopsis event in the homologous pairing of DNAs: RecA recognizes and pairs less than one helical repeat of DNA** *Proceedings of the National Academy of Sciences* **89**:6492–6496 <https://doi.org/10.1073/pnas.89.14.6492> | Google Scholar
39. Qi Z, et al. (2015) **DNA Sequence Alignment by Microhomology Sampling during Homologous Recombination** *Cell* **160**:856–869 <https://doi.org/10.1016/j.cell.2015.01.029> | Google Scholar

40. Ragunathan K, Joo C, Ha T (2011) **Real-Time Observation of Strand Exchange Reaction with High Spatiotemporal Resolution** *Structure* **19**:1064–1073 <https://doi.org/10.1016/j.str.2011.06.009> | Google Scholar
41. Lee JY, et al. (2015) **Base triplet stepping by the Rad51/RecA family of recombinases** *Science* **349**:977–981 <https://doi.org/10.1126/science.aab2666> | Google Scholar
42. Yang H, Zhou C, Dhar A, Pavletich NP (2020) **Mechanism of strand exchange from RecA–DNA synaptic and D-loop structures** *Nature* **586**:801–806 <https://doi.org/10.1038/s41586-020-2820-9> | Google Scholar
43. Appleby R, Joudeh L, Cobbett K, Pellegrini L (2023) **Structural basis for stabilisation of the RAD51 nucleoprotein filament by BRCA2** *Nature Communications* **14**:7003 <https://doi.org/10.1038/s41467-023-42830-1> | Google Scholar
44. Matsuo Y, et al. (2006) **Roles of the human Rad51 L1 and L2 loops in DNA binding** *The FEBS Journal* **273**:3148–3159 <https://doi.org/10.1111/j.1742-4658.2006.05323.x> | Google Scholar
45. Huang F, et al. (2011) **Identification of Specific Inhibitors of Human RAD51 Recombinase Using High-Throughput Screening** *ACS Chemical Biology* **6**:628–635 <https://doi.org/10.1021/cb100428c> | Google Scholar
46. Heller I, et al. (2014) **Optical Tweezers Analysis of DNA–Protein Complexes** *Chemical Reviews* **114**:3087–3119 <https://doi.org/10.1021/cr4003006> | Google Scholar
47. Cloud V, et al. (2012) **Rad51 Is an Accessory Factor for Dmc1-Mediated Joint Molecule Formation During Meiosis** *Science* **337**:1222–1225 <https://doi.org/10.1126/science.1219379> | Google Scholar
48. Hunter WN, et al. (1987) **The structure of guanosine-thymidine mismatches in B-DNA at 2.5-Å resolution** *Journal of Biological Chemistry* **262**:9962–9970 [https://doi.org/10.1016/s0021-9258\(18\)61060-9](https://doi.org/10.1016/s0021-9258(18)61060-9) | Google Scholar
49. Gupta RC, Golub EI, Wold MS, Radding CM (1998) **Polarity of DNA strand exchange promoted by recombination proteins of the RecA family** *Proceedings of the National Academy of Sciences* **95**:9843–9848 <https://doi.org/10.1073/pnas.95.17.9843> | Google Scholar
50. Baumann P, West SC (1997) **The human Rad51 protein: polarity of strand transfer and stimulation by hRP-A** *The EMBO Journal* **16**:5198–5206 <https://doi.org/10.1093/emboj/16.17.5198> | Google Scholar
51. Sung P, Roberson DL (1995) **DNA strand exchange mediated by a RAD51-ssDNA nucleoprotein filament with polarity opposite to that of RecA** *Cell* **82**:453–461 [https://doi.org/10.1016/0092-8674\(95\)90434-4](https://doi.org/10.1016/0092-8674(95)90434-4) | Google Scholar
52. Murayama Y, Kurokawa Y, Mayanagi K, Iwasaki H (2008) **Formation and branch migration of Holliday junctions mediated by eukaryotic recombinases** *Nature* **451**:1018–1021 <https://doi.org/10.1038/nature06609> | Google Scholar
53. Brouwer I, et al. (2018) **Two distinct conformational states define the interaction of human RAD51-ATP with single-stranded DNA** *The EMBO Journal* **37**:EMBJ201798162 <https://doi.org/10.15252/emboj.201798162> | Google Scholar

54. Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA (2017) **cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination** *Nat Methods* **14**:290–296 <https://doi.org/10.1038/nmeth.4169> | Google Scholar
55. Emsley P, Cowtan K (2004) **Coot: Model-building tools for molecular graphics** *Acta Crystallogr D Biol Crystallogr* **60**:2126–2132 | Google Scholar
56. Afonine PV, et al. (2018) **Real-space refinement in PHENIX for cryo-EM and crystallography** *Acta Crystallographica Section D* **74**:531–544 <https://doi.org/10.1107/s2059798318006551> | Google Scholar
57. Croll TI (2018) **ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps** *Acta Crystallographica Section D* **74**:519–530 <https://doi.org/10.1107/S2059798318002425> | Google Scholar
58. Smith SB, Cui Y, Bustamante C (1996) **Overstretching B-DNA: The Elastic Response of Individual Double-Stranded and Single-Stranded DNA Molecules** *Science* **271**:795–799 <https://doi.org/10.1126/science.271.5250.795> | Google Scholar
59. Rouzina I, Bloomfield VA (2001) **Force-Induced Melting of the DNA Double Helix. 2. Effect of Solution Conditions** *Biophysical Journal* **80**:894–900 [https://doi.org/10.1016/s0006-3495\(01\)76068-7](https://doi.org/10.1016/s0006-3495(01)76068-7) | Google Scholar
60. Rouzina I, Bloomfield VA (2001) **Force-Induced Melting of the DNA Double Helix 1. Thermodynamic Analysis** *Biophysical Journal* **80**:882–893 [https://doi.org/10.1016/s0006-3495\(01\)76067-5](https://doi.org/10.1016/s0006-3495(01)76067-5) | Google Scholar
61. Hilario J, Amitani I, Baskin RJ, Kowalczykowski SC (2009) **Direct imaging of human Rad51 nucleoprotein dynamics on individual DNA molecules** *Proceedings of the National Academy of Sciences* **106**:361–368 <https://doi.org/10.1073/pnas.0811965106> | Google Scholar
62. Mameren J van, et al. (2009) **Counting RAD51 proteins disassembling from nucleoprotein filaments under tension** *Nature* **457**:745–748 <https://doi.org/10.1038/nature07581> | Google Scholar
63. Finkelstein IJ, Visnapuu M-L, Greene EC (2010) **Single-molecule imaging reveals mechanisms of protein disruption by a DNA translocase** *Nature* **468**:983–987 <https://doi.org/10.1038/nature09561> | Google Scholar
64. Morin JA, et al. (2022) **Sequence-dependent surface condensation of a pioneer transcription factor on DNA** *Nature Physics* **18**:271–276 <https://doi.org/10.1038/s41567-021-01462-2> | Google Scholar
65. Gorman J, et al. (2007) **Dynamic Basis for One-Dimensional DNA Scanning by the Mismatch Repair Complex Msh2-Msh6** *Molecular Cell* **28**:359–370 <https://doi.org/10.1016/j.molcel.2007.09.008> | Google Scholar

Author information

Luay Joudeh[†]

Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom
ORCID iD: [0000-0001-9338-205X](https://orcid.org/0000-0001-9338-205X)

[†]These authors contributed equally to this work.

Robert E Appleby[†]

Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom, Francis Crick Institute, London, United Kingdom
ORCID iD: [0000-0002-9989-8310](https://orcid.org/0000-0002-9989-8310)

[†]These authors contributed equally to this work.

Joseph D Maman

Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom

Luca Pellegrini

Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom
ORCID iD: [0000-0002-9300-497X](https://orcid.org/0000-0002-9300-497X)

For correspondence: lp212@cam.ac.uk

Editors

Reviewing Editor

Akira Shinohara

Osaka University, Suita/Osaka, Japan

Senior Editor

Adèle Marston

University of Edinburgh, Edinburgh, United Kingdom

Reviewer #1 (Public review):

Summary:

The paper describes the cryoEM structure of RAD51 filament on the recombination intermediate. In the RAD51 filament, the insertion of a DNA-binding loop called the L2 loop stabilizes the separation of the complementary strand for the base-pairing with an incoming ssDNA and the non-complementary strand, which is captured by the second DNA-binding channel called the site II. The molecular structure of the RAD51 filament with a recombination intermediate provides a new insight into the mechanism of homology search and strand exchange between ssDNA and dsDNA.

Strong points:

This is the first human RAD51 filament structure with a recombination intermediate called the D-loop. The work has been done with great care, and the results shown in the paper are compelling based on cryo-EM and biochemical analyses. The paper is really nice and important for researchers in the field of homologous recombination, which gives a new view on the molecular mechanism of RAD51-mediated homology search and strand exchange.

Comments on revisions:

The authors nicely address most of the previous points.

<https://doi.org/10.7554/eLife.107114.2.sa3>

Reviewer #2 (Public review):

Homologous recombination is essential for DNA double-strand break repair, with RAD51-catalyzed strand exchange at its core. This study presents a 2.64 Å resolution cryogenic electron microscopy structure of the RAD51 D-loop complex, achieved through reconstitution of a RAD51 mini-filament. The structure uncovers how specific RAD51 residues drive strand exchange, offering atomic-level insight into the mechanics of eukaryotic HR and DNA repair.

Comments on revisions:

Authors acknowledged:

"We acknowledge that there exists an extensive body of literature that has investigated the polarity of strand exchange by RecA and RAD51 under a variety of experimental conditions, and we have added a brief comment to the text to reflect this, as well as some of the key citations. Undoubtedly, and as we also mention in our reply to the public reviews, further experimental work will be needed for a full reconciliation of the available evidence."

In the revised manuscript, this is reflected in the statement:

"Our mechanistic interpretation of static D-loop structures awaits full reconciliation with earlier efforts to determine strand-exchange polarity for RecA and RAD51 measured under a variety of experimental conditions."

Among the four cited studies, my understanding (as a person who has never studied this subject of polarity) is as follows:

- References 50 (EMBO J. 1997), 51 (Cell. 1995), and 52 (Nature. 2008) suggest that the strand exchange by human RAD51 occurs with a polarity opposite to that of RecA—that is, in the 5' → 3' direction relative to the complementary strand, or 3' → 5' relative to the initiating single-stranded DNA (ssDNA).
- In contrast, reference 49 (PNAS 1998) proposed that 5' → 3' polarity (relative to ssDNA) is conserved across RecA, human RAD51, and yeast RAD51.

Given the substantial structural analysis provided in the current manuscript, it would strengthen the work to include a concise description of these earlier biochemical findings, rather than citing them without context. This would benefit readers who are not familiar with the longstanding studies in the field and allow for a more informed interpretation of how the structural observations may reconcile or contrast with previous work.

<https://doi.org/10.7554/eLife.107114.2.sa2>

Reviewer #3 (Public review):

Summary:

Built on their previous pioneer expertise in studying RAD51 biology, in this paper, the authors aim to capture and investigate the structural mechanism of human RAD51 filament bound with a displacement loop (D-loop), which occurs during the dynamic synaptic state of the homologous recombination (HR) strand-exchange step. As the structures of both pre- and

post-synaptic RAD51 filaments were previously determined, a complex structure of RAD51 filament during strand exchange is one of the key missing pieces of information for a complete understanding of how RAD51 functions in HR pathway. This paper aims to determine the high-resolution cryo-EM structure of RAD51 filament bound with D-loop. Combined with mutagenesis analysis and biophysical assays, the authors aim to investigate the D-loop DNA structure, RAD51 mediated strand separation and polarity, and a working model of RAD51 during HR strand invasion in comparison with RecA.

Strengths:

- (1) The structural work and associated biophysical assays in this paper are solid, elegantly designed and interpreted. These results provide novel insights into RAD51's function in HR.
- (2) The DNA substrate used was well designed, taking into consideration of the nucleotide number requirement of RAD51 for stable capture of donor DNA. This DNA substrate choice lays the foundation for successfully determining the structure of the RAD51 filament on D-loop DNA using single-partial cryo-EM.
- (3) The authors utilised their previous expertise in capping DNA ends using monomeric streptavidin and combined their careful data collection and processing to determine the cryo-EM structure of full-length human RAD51 bound at D-loop in high resolution. This interesting structure forms the core part of this work and allows detailed mapping of DNA-DNA and DNA-protein interaction among RAD51, invading strands, and donor DNA arms (Figures 1, 2, 3, 4). The geometric analysis of D-loop DNA bound with RAD51 and EM density for homologous DNA pairing are also impressive (Figure S5). The previously disordered RAD51's L2-loop is now ordered and traceable in the density map and functions as a physical spacer when bound with D-loop DNA. Interestingly, the authors identified that the side chain position of F279 in the L2_loop of RAD51_H differs from other F279 residues in L2-loops of E, F and G protomers. This asymmetric binding of L2 loops and RAD51_NTD binding with donor DNA arms forms the basis of the proposed working model about the polarity on csDNA during RAD51-mediated strand exchange.
- (4) This work also includes mutagenesis analysis and biophysical experiments, especially EMSA, single-molecule fluorescence imaging using an optical tweezer, and DNA strand exchange assay, which are all suitable methods to study the key residues of RAD51 for strand exchange and D-loop formation (Figure 5).

Weaknesses:

- (1) The proposed model for the 3'-5' polarity of RAD51-mediated strand invasion is based on the structural observations in the cryo-EM structure. This study lacks follow-up biochemical/biophysical experiments to validate the proposed model compared to RecA or developing methods to capture structures of any intermediate states with different polarity models.
- (2) The functional impact of key mutants designed based on structure has not been tested in cells to evaluate how these mutants impact the HR pathway.

The significance of the work for the DNA repair field and beyond:

Homologous recombination (HR) is a key pathway for repairing DNA double-strand breaks and involves multiple steps. RAD51 forms nucleoprotein filaments first with 3' overhang single-strand DNA (ssDNA), followed by a search and exchange with a homology strand. This function serves as the basis of an accurate template-based DNA repair during HR. This research addressed a long-standing challenge of capturing RAD51 bound with the dynamic synaptic DNA and provided the first structural insight into how RAD51 performs this function. The significance of this work extends beyond the discovery biology for the DNA

repair field, into its medical relevance. RAD51 is a potential drug target for inhibiting DNA repair in cancer cells to overcome drug resistance. This work offers a structural understanding of RAD51's function with D-loop and provides new strategies for targeting RAD51 to improve cancer therapies.

<https://doi.org/10.7554/eLife.107114.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

Summary:

The paper describes the cryoEM structure of RAD51 filament on the recombination intermediate. In the RAD51 filament, the insertion of a DNA-binding loop called the L2 loop stabilizes the separation of the complementary strand for the base-pairing with an incoming ssDNA and the non-complementary strand, which is captured by the second DNA-binding channel called the site II. The molecular structure of the RAD51 filament with a recombination intermediate provides a new insight into the mechanism of homology search and strand exchange between ssDNA and dsDNA.

Strengths:

This is the first human RAD51 filament structure with a recombination intermediate called the D-loop. The work has been done with great care, and the results shown in the paper are compelling based on cryo-EM and biochemical analyses. The paper is really nice and important for researchers in the field of homologous recombination, which gives a new view on the molecular mechanism of RAD51-mediated homology search and strand exchange.

Weaknesses:

The authors need more careful text writing. Without page and line numbers, it is hard to give comments.

We would like to thank the reviewer for their kind words of appreciation of our work.

Reviewer #2 (Public review):

Summary:

Homologous recombination (HR) is a critical pathway for repairing double-strand DNA breaks and ensuring genomic stability. At the core of HR is the RAD51-mediated strand-exchange process, in which the RAD51-ssDNA filament binds to homologous double-stranded DNA (dsDNA) to form a characteristic D-loop structure. While decades of biochemical, genetic, and single-molecule studies have elucidated many aspects of this mechanism, the atomic-level details of the strand-exchange process remained unresolved due to a lack of atomic-resolution structure of RAD51 D-loop complex.

In this study, the authors achieved this by reconstituting a RAD51 mini-filament, allowing them to solve the RAD51 D-loop complex at 2.64 Å resolution using a single particle approach. The atomic resolution structure reveals how specific residues of RAD51 facilitate the strand exchange reaction. Ultimately, this work provides unprecedented structural insight into the eukaryotic HR process and deepens the understanding of

RAD51 function at the atomic level, advancing the broader knowledge of DNA repair mechanisms.

Strengths:

The authors overcame the challenge of RAD51's helical symmetry by designing a minifilament system suitable for single-particle cryo-EM, enabling them to resolve the RAD51 D-loop structure at 2.64 Å without imposed symmetry. This high resolution revealed precise roles of key residues, including F279 in Loop 2, which facilitates strand separation, and basic residues on site II that capture the displaced strand. Their findings were supported by mutagenesis, strand exchange assays, and single-molecule analysis, providing strong validation of the structural insights.

Weaknesses:

Despite the detailed structural data, some structure-based mutagenesis data interpretation lacks clarity. Additionally, the proposed 3'-to-5' polarity of strand exchange relies on assumptions from static structural features, such as stronger binding of the 5'-arm-which are not directly supported by other experiments. This makes the directional model compelling but contradicts several well-established biochemical studies that support a 5'-to-3' polarity relative to the complementary strand (e.g., Cell 1995, PMID: 7634335; JBC 1996, PMID: 8910403; Nature 2008, PMID: 18256600).

Overall:

The 2.6 Å resolution cryoEM structure of the RAD51 D-loop complex provides remarkably detailed insights into the residues involved in D-loop formation. The high-quality cryoEM density enables precise placement of each nucleotide, which is essential for interpreting the molecular interactions between RAD51 and DNA. Particularly, the structural analysis highlights specific roles for key domains, such as the N-terminal domain (NTD), in engaging the donor DNA duplex.

This structural interpretation is further substantiated by single-molecule fluorescence experiments using the KK39,40AA NTD mutant. The data clearly show a significant reduction in D-loop formation by the mutant compared to wild-type, supporting the proposed functional role of the NTD observed in the cryoEM model.

However, the strand exchange activity interpretation presented in Figure 5B could benefit from a more rigorous experimental design. The current assay measures an increase in fluorescence intensity, which depends heavily on the formation of RAD51-ssDNA filaments. As shown in Figure S6A, several mutants exhibit reduced ability to form such filaments, which could confound the interpretation of strand exchange efficiency. To address this, the assay should either: (1) normalize for equivalent levels of RAD51-ssDNA filaments across samples, or (2) compare the initial rates of fluorescence increase (i.e., the slope of the reaction curve), rather than endpoint fluorescence, to better isolate the strand exchange activity itself.

We agree with the reviewer that the reduced filament-forming ability of some of the RAD51 mutants complicates a straightforward interpretation of their strand-exchange assay. Interestingly, the RAD51 mutants that appear most impaired are the ssDNA-capture mutants that do not contact the ssDNA in the structure of the pre-synaptic filament. However, the RAD51 NTD mutants, that display the most severe defect in strand-exchange, have a near-WT filament forming ability.

Based on the structural features of the D-loop, the authors propose that strand pairing and exchange initiate at the 3'-end of the complementary strand in the donor DNA and proceed with a 3'-to-5' polarity. This conclusion, drawn from static structural

observations, contrasts with several well-established biochemical studies that support a 5'-to-3' polarity relative to the complementary strand (e.g., Cell 1995, PMID: 7634335; JBC 1996, PMID: 8910403; Nature 2008, PMID: 18256600). While the structural model is compelling and methodologically robust, this discrepancy underscores the need for further experiments.

We would like to thank the reviewer for highlighting the importance of our findings to our understanding of the mechanism of homologous recombination.

The reviewer correctly points out that the polarity of strand exchange by RecA and RAD51 is an extensively researched topic that has been characterised in several authoritative studies. In our paper, we simply describe the mechanistic insights obtained from the structural D-loop models of RAD51 (our work) and RecA (Yang et al, PMID: 33057191). The structures illustrate a very similar mechanism of D-loop formation that proceeds with opposite polarity of strand exchange for RAD51 and RecA. Comparison of the D-loop structures for RecA and RAD51 provides an attractive explanation for the opposite polarity, as caused by the different positions of their dsDNA-binding domains in the filament structure.

We agree with the reviewer that further investigation will be needed for an adequate rationalisation of the available evidence. We will mention the relevant literature in the revised version of the manuscript.

Reviewer #3 (Public review):

Summary:

Built on their previous pioneer expertise in studying RAD51 biology, in this paper, the authors aim to capture and investigate the structural mechanism of human RAD51 filament bound with a displacement loop (D-loop), which occurs during the dynamic synaptic state of the homologous recombination (HR) strand-exchange step. As the structures of both pre- and post-synaptic RAD51 filaments were previously determined, a complex structure of RAD51 filaments during strand exchange is one of the key missing pieces of information for a complete understanding of how RAD51 functions in the HR pathway. This paper aims to determine the high-resolution cryo-EM structure of RAD51 filament bound with the D-loop. Combined with mutagenesis analysis and biophysical assays, the authors aim to investigate the D-loop DNA structure, RAD51-mediated strand separation and polarity, and a working model of RAD51 during HR strand invasion in comparison with RecA.

Strengths:

- (1) The structural work and associated biophysical assays in this paper are solid, elegantly designed, and interpreted. These results provide novel insights into RAD51's function in HR.*
- (2) The DNA substrate used was well designed, taking into consideration the nucleotide number requirement of RAD51 for stable capture of donor DNA. This DNA substrate choice lays the foundation for successfully determining the structure of the RAD51 filament on D-loop DNA using single-particle cryo-EM.*
- (3) The authors utilised their previous expertise in capping DNA ends using monomeric streptavidin and combined their careful data collection and processing to determine the cryo-EM structure of full-length human RAD51 bound at the D-loop in high resolution. This interesting structure forms the core part of this work and allows detailed mapping of DNA-DNA and DNA-protein interaction among RAD51, invading strands, and donor DNA arms (Figures 1, 2, 3, 4). The geometric analysis of D-loop DNA bound with RAD51 and EM density for homologous DNA pairing is also impressive (Figure S5). The*

previously disordered RAD51's L2-loop is now ordered and traceable in the density map and functions as a physical spacer when bound with D-loop DNA. Interestingly, the authors identified that the side chain position of F279 in the L2_loop of RAD51_H differs from other F279 residues in L2-loops of E, F, and G protomers. This asymmetric binding of L2 loops and RAD51_NTD binding with donor DNA arms forms the basis of the proposed working model about the polarity of csDNA during RAD51-mediated strand exchange.

(4) This work also includes mutagenesis analysis and biophysical experiments, especially EMSA, singlemolecule fluorescence imaging using an optical tweezer, and DNA strand exchange assay, which are all suitable methods to study the key residues of RAD51 for strand exchange and D-loop formation (Figure 5).

Weaknesses:

(1) The proposed model for the 3'-5' polarity of RAD51-mediated strand invasion is based on the structural observations in the cryo-EM structure. This study lacks follow-up biochemical/biophysical experiments to validate the proposed model compared to RecA or developing methods to capture structures of any intermediate states with different polarity models.

(2) The functional impact of key mutants designed based on structure has not been tested in cells to evaluate how these mutants impact the HR pathway.

The significance of the work for the DNA repair field and beyond:

Homologous recombination (HR) is a key pathway for repairing DNA double-strand breaks and involves multiple steps. RAD51 forms nucleoprotein filaments first with 3' overhang single-strand DNA (ssDNA), followed by a search and exchange with a homologous strand. This function serves as the basis of an accurate template-based DNA repair during HR. This research addressed a long-standing challenge of capturing RAD51 bound with the dynamic synaptic DNA and provided the first structural insight into how RAD51 performs this function. The significance of this work extends beyond the discovery of biology for the DNA repair field, into its medical relevance. RAD51 is a potential drug target for inhibiting DNA repair in cancer cells to overcome drug resistance. This work offers a structural understanding of RAD51's function with the D-loop and provides new strategies for targeting RAD51 to improve cancer therapies.

We thank the reviewer for their positive comments on the significance of our work. Concerning the proposed polarity of strand exchange based on our structural finding, please see our reply to the previous reviewer; we agree with the reviewer that further experimentation will be needed to reach a settled view on this.

Testing the functional effects of the RAD51 mutants on HR in cells was not an aim of the current work but we agree that it would be a very interesting experiment, which would likely provide further important insights into the mechanism of strand exchange at the core of the HR reaction.

Reviewer #1 (Recommendations for the authors):

Major points:

(1) Structural analysis showed a critical role of F279 in the L2 loop. However, the biochemical study showed that the F279A substitution did not provide a strong defect in the *in vitro* strand exchange, as shown in Figure 5B. Moreover, a previous study by Matsuo et al. *FEBS J*, 2006; ref 43) showed human RAD51-F279A is proficient in the *in vitro* strand exchange. These suggest that human RAD51 F279 is not critical for the

strand exchange. The authors need more discussions of the role of F279 or the L2 for the RAD51-mediated reactions in the Discussion.

In the strand-exchange assay of Figure 5B, the F279A mutant shows the mildest phenotype, in agreement with the findings of Matsuo et al. Accordingly, in the text we describe the F279A mutant as having a “modest impact” on strand-exchange.

We have now added a brief comment to the relevant text, pointing out that the result of the strand exchange assay for F279A are in agreement with the previous findings by Matsuo et al., and adding the reference.

(2) In some parts, the authors cited the newest references rather than the paper describing the original findings. For RAD51 paralogs, why are these three (refs 21,22, 23) selected here? For FIGNL1, why is only one (ref 24) chosen?

The cited publications were chosen to acquaint the reader with the latest structural and mechanistic advances about the function of some of the most important and well-studied recombination mediator proteins. For completeness, we have now added a further reference for FIGNL1 - Ito, Masaru et al, Nat Comm, 2023 – in the Introduction, to provide the reader with an additional pointer to our current knowledge about the mechanism of FIGNL1 in Homologous Recombination.

Minor points:

(1) Page 3, line 1 in the second paragraph, the reaction of "HR": HR should be homology search and strand exchange. HR is used incorrectly throughout the text, please check them. Remove "strandexchange" from ATPases in line 2.

We believe that HR is used correctly in this context, as we refer to the biochemical reactions of HR, which includes the search for homology and strand exchange.

We have removed “strand-exchange” from ATPases in line 2, as requested by the reviewer.

(2) Supplementary Figure 1B, C, "EMSA" experiment: Please indicate an experimental condition in the legend: how ssDNA and dsDNA were mixed with RAD51. In (B), this is not an actual EMSA result, but rather a native gel analysis of reaction products with the D-loop. In (C), was the binding of RAD51 to the pre-formed D-loop examined? Which is correct here? Moreover, why do the authors need streptavidin in this experiment? Please explain why this is necessary for the EMSA assay. Please show where is Cy3 or Cy5 labels on the DNAs should be shown in the schematic drawing.

The conditions for the experiments of Supplementary figure 1B, C are reported in the Methods section.

Panel B shows the mobility shifts of the ssDNA and dsDNA sequences in panel A, so it is appropriate to describe it as an EMSA.

We did not examine the binding of RAD51 to a pre-formed D-loop.

We used streptavidine in the experiment of Supplementary Figure 1C to show that streptavidine binding did not interfere with D-loop reconstitution.

The position of the Cy3, Cy5 labels in the DNAs is reported in Table S1.

(3) Figure S4B, page 6, line 6 from the top, 5'-arm and 3'-arm: please add them to the figure. And also, please explain what 5'-arm and 3'-arm are here in the text, as shown in

lines 3-5 in the second paragraph of the same page.

We thank the reviewer for spotting this slight incongruity. We have removed the reference to 5'- and 3' arms of the donor DNA in the initial description of the D-loop (first paragraph of the "D-loop structure" section, 6 lines from the top), as the nomenclature for the arms of the donor DNA is introduced more appropriately in the following paragraph. Thus, there is no need to re-label Figure S4B; we note that the 5'- and 3'-labels are added to the arms of the donor DNA in Figure S4D.

(4) Page 7, line 4, and Figure 2E, "C24": C24 should be C26 here (Figure 2D shows that position 24 in esDNA is "T").

We thank the reviewer for spotting this typo, that is now corrected in the revised version of Figure 2 and in the text.

(5) Page 8, line 1, K284: It would be nice to show "K284" in Figure 3F.

We have added the side chain of K284 to Figure 3F, as suggested by the reviewer.

(6) Page 8, second paragraph, line 3 from the bottom, "5'-arm" should be "3'-arm" for the binding of RAD51A NTD to ds DNA (Figure 4D).

We thank the reviewer for spotting this typo, that is now corrected in the revised version of the text.

Reviewer #2 (Recommendations for the authors):

I understand that the strand exchange polarity of RAD51 should be opposite to that of RecA. But in the RecA manuscript (Nature 2020), it states (in the extended figure 1) "Because the mini-filament consists of fused RecA protomers, it does not reflect the effects a preferential polarity of RecA polymerization might have on the directionality of strand exchange. Also, our strand exchange reactions do not include the single-stranded DNA binding protein SSB that is involved in strand exchange in vivo and may sequester released DNA strands."

We are aware that the findings by Yang et al, 2020 were obtained with a multi-protomeric RecA chimera and that their construct might not therefore recapitulate a potential effect of RecA polymerisation on the directionality of strand-exchange.

Comparison of the RecA and RAD51 D-loop structures shows that RecA and RAD51 adopt the same asymmetric mechanism of D-loop formation, which begins at one arm of the donor DNA and proceeds with donor unwinding and strand invasion until the second arm is captured, completing D-loop formation. However, the cryoEM structures provide compelling evidence that, after engagement with the donor DNA, RecA and RAD51 proceed to unwind the donor with opposite polarity; the structures provide a clear rationale for this, because of the different position of their dsDNA-binding domains relative to the ATPase domain.

We acknowledge that there exists an extensive body of literature that has investigated the polarity of strand exchange by RecA and RAD51 under a variety of experimental conditions, and we have added a brief comment to the text to reflect this, as well as some of the key citations. Undoubtedly, and as we also mention in our reply to the public reviews, further experimental work will be needed for a full reconciliation of the available evidence.

Reviewer #3 (Recommendations for the authors):

(1) I have a minor comment regarding the DNA shown in the structural figures in this work. The authors have used different colours to differentiate between isDNA, esDNA, and csDNA for easier interpretation. However, these colour codes are inconsistent across Figures 1, 2, 3, and 5. This inconsistency makes it difficult to interpret which strand is which, particularly for readers unfamiliar with D-loops and strand invasion. A consistent colour scheme for the DNA strands would enhance the quality of the structural figures.

We appreciate the reviewer's comment about the colour scheme of the strands in the D-loop. We chose a unique colour scheme for each figure, to help the reader focus on the particular structural features that we wanted to highlight in the figure. So for instance, in figure 1D we chose to highlight the relationship (complementary vs identical) of the donor DNA strands with the the invading strand; in figure 2, the emphasis is on distinguishing the homologously paired dsDNA (pink) from the exchanged strand (magenta), as a consequence of L2 loop binding; etc.

(2) I have another comment regarding the rationale behind naming the RAD51 protomers (A to H) within the structure, which could confuse general readers if not clearly explained. In this paper, the RAD51 protomer is RAD51_A when closest to the 3' end of the isDNA. I assume the authors chose this order because HR generates a 3' ssDNA overhang before strand invasion. It would be beneficial for the introduction and results sections to mention this property of the 3' ssDNA overhang and the reasoning behind this naming strategy. This explanation will help readers understand how it differs from other naming orders used in RecA/RAD51 with ssDNA, where protomer A is closer to the 5' ssDNA.

We thank the reviewer for their insightful comment. We chose to name as chain A the RAD51 protomer nearest to the 3'-end of the isDNA to be consistent with the naming scheme that we use for all our published RAD51 filament structures.

(3) I have highlighted some text within this paper that has contradicting parts for authors to clarify and correct:

"Overall, the structural features of the RAD51 D-loop provide a strong indication that strand pairing and exchange begins at the 3'-end of the complementary strand in the donor DNA and progresses with 3'-to-5' polarity (Fig. 5F)"

"The observed 5'-to-3' polarity of strand-exchange by RAD51 is opposite to the 3'-to-5' polarity of bacterial RecA (Fig. S8), that was determined based on cryoEM structures of RecA D-loops".

We thank the reviewer for alerting us to this inconsistency that has now been corrected in the revised manuscript.

(4) Figure S8 last model: NTD should be CTD in the title; Figure 2B: resolution scale bar needs A unit. We thank the reviewer for spotting this typo that has now been corrected in the revised version of figure S8.

We couldn't find a missing resolution scale bar in Figure 2B; however, we have added a missing resolution bar with A unit to Fig. S3B.

<https://doi.org/10.7554/eLife.107114.2.sa0>