

Reviewed Preprint

v1 • December 11, 2024

Not revised

Reviewed Preprint

v2 • September 24, 2025

Revised by authors

✉ For correspondence:

crf2s@virginia.edu

Competing interests:

No competing interests declared

Funding:

See [page 18](#)

Reviewing editor:

Dolores Shoback,
University of California, San
Francisco, United States

© 2024, Dillard et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Cell type-specific network analysis in Diversity Outbred mice identifies genes potentially responsible for human bone mineral density GWAS associations

Luke J Dillard¹, Gina M Calabrese¹, Larry D Mesner^{1,2}, Charles R Farber^{1,2,3} ✉

¹Center for Public Health Genomics, School of Medicine, University of Virginia, Charlottesville, United States •

²Department of Public Health Sciences, School of Medicine, University of Virginia, Charlottesville, United States •

³Department of Biochemistry and Molecular Genetics, School of Medicine, University of Virginia, Charlottesville, United States

eLife Assessment

This is an **important** study that provides **compelling** data from a diverse set of approaches from single cell transcriptome data and network analysis from genetically diverse mouse cells to identify novel driver genes underlying human GWAS associations. The authors present evidence that network analysis of scRNA-seq data from genetically diverse mouse bone-marrow derived stromal cells can be informative for identifying human BMD GWAS driver genes. Their approach should be broadly used and applicable to other GWAS studies.

<https://doi.org/10.7554/eLife.100832.2.sa3>

Summary

Genome-wide association studies (GWASs) have identified many sources of genetic variation associated with bone mineral density (BMD), a clinical predictor of fracture risk and osteoporosis. Aside from the identification of causal genes, other difficult challenges to informing GWAS include characterizing the roles of predicted causal genes in disease and providing additional functional context, such as the cell type predictions or biological pathways in which causal genes operate. Leveraging single-cell transcriptomics (scRNA-seq) can assist in informing BMD GWAS by linking disease-associated variants to genes and providing a cell type context for which these causal genes drive disease. Here, we use large-scale scRNA-seq data from bone marrow-derived stromal cells cultured under osteogenic conditions (BMSC-OBs) from Diversity Outbred (DO) mice to generate cell type-specific networks and contextualize BMD GWAS-implicated genes. Using trajectories inferred from the scRNA-seq data that map cell state transitions, we identify networks enriched with genes that exhibit the most dynamic changes in expression across trajectories. We discover 21 network driver genes, which are likely to be causal for human BMD GWAS associations that colocalize with expression/splicing quantitative trait loci (eQTL/sQTL). These driver genes, including *Fgfr1l* and *Tpx2*, along with their associated networks, are predicted to be novel regulators of BMD via their roles in the differentiation of mesenchymal lineage cells. In this work, we showcase the use of single-cell transcriptomics from mouse bone-relevant cells to inform human BMD GWAS and prioritize genetic targets with potential causal roles in the development of osteoporosis.

Introduction

Osteoporosis is a complex disease characterized by low bone mineral density (BMD), bone fragility, and an increased risk of fracture¹. BMD, a highly heritable trait, is one of the most important clinical predictors of osteoporotic fracture^{2,3}. Increasing our understanding of the genetic basis of BMD is critical for the development of approaches for the treatment and prevention of osteoporosis. In recent years, genome-wide association studies (GWAS) have made great progress in unraveling BMD genetics by identifying over 1,100 independent associations⁴. Now the challenge lies in pinpointing causal genes, which is necessary for the translation of genetic findings into novel therapies.

A number of approaches exist to identify genes responsible for GWAS associations^{5–8}. Most rely on population-based “-omics” data⁹, which are scarce for human bone, to connect associations to causal genes. Our group has used co-expression networks generated from mouse bone transcriptomic datasets to assist in the identification of genes likely responsible for BMD associations. One significant advantage of this approach is its ability to utilize the network connections of candidate genes to predict how these candidate genes may affect BMD. For example, we generated co-expression networks from bone tissue and primary osteoblasts in mouse genetic reference populations and identified multiple co-expression modules enriched with genes located in BMD associations^{10,11}. We then cross-referenced genes in these modules with those regulated by co-localizing expression quantitative trait loci (eQTLs) from the Gene-Tissue Expression project (GTEx)^{12,13} to identify “high priority” genes. Recently, we expanded our analyses to include directed networks generated via a Bayesian approach using cortical bone RNA-seq data from 192 Diversity Outbred (DO) mice. By combining key driver analysis and GTEx eQTL colocalization data, we identified 19 novel genes, such as *SERTAD4* and *GLT8D2*, which are likely causal for human BMD GWAS associations¹⁴.

To date, our analyses have been reliant on networks generated from heterogeneous bulk transcriptomics (RNA-seq) datasets from mouse bone and primary bone cells. However, leveraging single-cell transcriptomics (scRNA-seq) data would offer the added benefit of resolving the transcriptomic profiles of discrete cell types. Additionally, using scRNA-seq data has the potential to provide context by predicting the specific cell types in which causal genes and their associated networks operate. In recent work, we demonstrated the utility of bone marrow-derived stromal cells cultured under osteogenic condition (BMSC-OB) for the generation of population-scale scRNA-seq data on bone relevant cell types¹⁵. The BMSC-OB model effectively enriches for mesenchymal lineage cells (e.g., mesenchymal progenitors, osteoblasts, osteocyte-like cells) that are highly relevant to the regulation of BMD.

In this work, our goal was to prioritize and contextualize genes implicated by BMD GWAS using an expanded large-scale (N=80) BMSC-OB scRNA-seq dataset on bone cell types. We accomplished this by first generating co-expression and Bayesian networks¹⁴ for each BMSC-OB mesenchymal cell type. We subsequently prioritized networks based on their enrichment for genes exhibiting the most dynamic changes in expression across trajectories inferred from the scRNA-seq data, thus highlighting networks likely associated with the differentiation of BMSC-OBs. We then used these networks linked to osteogenic differentiation to prioritize genes with eQTL and/or splicing quantitative trait loci (sQTL) which colocalize with BMD GWAS associations^{7,16}. In doing so, this analysis provides additional support for a role of these genes in the regulation of BMD and highlights their potential roles in differentiation of cell types essential to skeletal health.

Results

BMSC-OBs derived from DO mice yield diverse cell types that are enriched for mesenchymal lineage cells

We cultured BMSCs from a total of 80 Diversity Outbred (DO) mice, a genetically diverse outbred mouse population^{17,18} (N=75 from the current study and N=5 from¹⁵; N = 49 male and N = 31 females). We cultured BMSCs under osteogenic conditions and subsequently performed scRNA-

seq, as described in¹⁵. After stringent processing and quality control (Materials and Methods), the dataset consisted of 21,831 expressed genes across 139,392 cells. We manually annotated 15 clusters ranging in size from 270 to 27,291 cells and identified cell types of the mesenchymal lineage as well as various other cell types (Fig. 1A¹⁵, Supplementary Table S1).

Based on our prior BMSC-OB scRNA-seq study¹⁵, we expected to identify a large proportion of mesenchymal cells and a smaller fraction of non-mesenchymal cell types. Consistent with this hypothesis, clusters associated with mesenchymal lineages accounted for 74.1% of all cells (Fig. 1A¹⁵). These included mesenchymal progenitor cells (MPCs), late mesenchymal progenitors (LMPs), osteoblast progenitors (OBPs), two mature osteoblast populations (OB1 and OB2), osteocyte-like cells (Ocy), and marrow adipogenic lineage progenitors (MALPs). The non-mesenchymal cell types observed included macrophages, monocytes, granulocytes, T-cells, B-cells, endothelial cells, and osteoclast-like cells (Fig. 1A¹⁵). With regards to the mesenchymal cell types, the only differences in cell clusters relative to our previous report¹⁵ were the presence of MPCs and two mature osteoblast clusters. Upon comparing the two distinct osteoblast clusters, OB1 and OB2 (Fig. 1A¹⁵), both clusters had ubiquitous expression of genes associated with mature osteoblasts (e.g., *Col1a1*, *Bglap*, *Sparc*, and *Ibsp*) (Supplementary Table S1) while many of the “canonical” osteoblast markers were more highly expressed in OB1 compared to OB2 (Supplementary Table S2). Interestingly, MPCs did not have transcriptomic profiles similar to other mesenchymal progenitor cells previously identified by our group or others^{15,19}. All other mesenchymal cell types demonstrated specific expression of canonical marker genes (Fig. 1A, B¹⁵).

We next assessed the variability in cell type frequencies across DO mice by quantifying the proportions of each annotated mesenchymal cell type. All other clusters, which mainly consisted of immune cells of hematopoietic origin, were aggregated into one group (Hem) for each mouse. We observed high variability in the raw proportional abundances of cell types derived from each mouse (Fig. 1C¹⁵, Supplementary Table S3). For example, the proportions of osteoblasts (OB1 and OB2) varied significantly among individual DO mice (Fig. 1D¹⁵). All mice used in the current experiment had been extensively phenotyped for a wide range of bone traits (microCT, histomorphometry, biomechanical bone properties, etc.) as part of a previous genetic analysis of bone strength¹⁴. We correlated cell type frequencies with bone traits, however, none of the cell type proportions were strongly correlated with any bone trait (Supplementary Table S4, S5).

Mesenchymal lineage cells are enriched in BMD heritability

The primary goal of this work was to prioritize and contextualize genes implicated by BMD GWAS. As a first step towards this goal, we sought to determine which cell types were the most relevant to the genetics of BMD. Using the BMD GWAS and the BMSC-OB scRNA-seq data, we performed a CELLECT²⁰ cell type prioritization analysis to identify cell clusters enriched for BMD heritability. We observed that OB1, Ocy, and MALP cell clusters were significantly enriched ($P_{\text{adj}} < 0.05$, red dashed line) for BMD heritability ($P_{\text{adj}} = 0.018, 0.010, 0.006$, respectively) (Fig. 1E¹⁵, Supplementary Tables S6). None of the non-mesenchymal cells identified were significant ($P_{\text{adj}} > 0.05$) (Fig. 1E¹⁵). As a result, all downstream efforts focused solely on using data on mesenchymal cell types to inform GWAS.

Generating mesenchymal cell type-specific Bayesian networks to inform BMD GWAS

We have previously shown that network-based approaches using bulk RNA-seq are powerful tools for the identification of putative causal genes from BMD GWAS data^{10,11,14}. Here, our goal was to apply these same approaches using the BMSC-OB scRNA-seq data to prioritize and contextualize genes we previously identified as having a colocalizing expression quantitative trait locus (eQTL; $N=512$) or splicing QTL (sQTL; $N=732$) in a tissue from the Genotype-Tissue Expression (GTEx) project^{7,13,16}. Genes identified in each study (or both) yielded a list of high priority target genes ($N = 1,037$). While GTEx does not currently contain data for bone tissue, eQTL can be shared across many tissues and may exert their effects in cell types resident to bone²¹. Therefore, utilizing our




Figure 1. Analysis of single cell RNA-seq (scRNA-seq) data for BMSC-OBs derived from 80 Diversity Outbred (DO)

(A) Uniform Manifold Approximation and Projection (UMAP) of 139,392 single cells (BMSC-OBs). Cell numbers and corresponding percentages for the fifteen (15) annotated cell clusters are listed in parenthesis to the right of the annotated cluster name. (B) Dot plot⁶⁵ portraying representative and highly expressed genes for all annotated cell clusters. Dot color indicates the scaled gene expression while the size of the dot corresponds to the percentage of cells of a given cluster that express a given gene. (C) The raw proportional abundances of seven (7) mesenchymal cell clusters and one (1) cluster (Hem) representing the remain cells (i.e., mainly hematopoietic immune cells) across all 80 DO mice. (D) UMAP plots for mesenchymal lineage cell clusters for DO mouse 50 and DO mouse 233. (E) CELLECT (CELL-type Expression-specific integration for Complex Traits) cell type prioritization results displaying the Bonferroni adjusted P-values for the cell clusters. The OB1, Ocy, and MALP cell clusters (red) were significantly enriched ($P_{adj} < 0.05$, red dashed line) for BMD heritability ($P_{adj} = 0.018, 0.010, 0.006$, respectively).

previous work, we hypothesized that generating cell type-specific networks would yield more biologically relevant representations of processes occurring within particular mesenchymal cell types. Additionally, by integrating GWAS, cell type-specific networks, and dynamic gene expression as a function of differentiation, we anticipated we would identify points of intervention in which genetic variation impacts genes involved in the differentiation process.

Our network analysis begins by partitioning genes into groups based on co-expression by applying iterative weighted gene co-expression network analysis (iterativeWGCNA)²² to each mesenchymal cell type (Step 1, Fig. 2 [↗](#)). In total, 535 modules were identified from the BMSC-OB scRNA-seq data, and the number of modules identified for each mesenchymal cell cluster ranged from 26 to 153 with an average of 76 modules per cluster (Supplementary Table S7, S8). We sought to infer causal relationships between genes in each cell type-specific co-expression module and subsequently identify networks involved in processes relevant to BMSC-OB differentiation. To this end, we generated Bayesian networks for each co-expression module, thus enabling us to model directed interactions between co-expressed genes based on conditional independence¹⁴ (Step 2, Fig. 2 [↗](#)).

Identifying putative drivers of mesenchymal cell differentiation

We hypothesized that many genes impacting BMD do so by influencing osteogenic differentiation or possibly bone marrow adipogenic differentiation of key mesenchymal cell types, as suggested by the CELLECT analysis above. Therefore, the next step of our network analysis was to identify cell type-specific Bayesian networks enriched for genes potentially driving mesenchymal differentiation (Step 3, Fig. 2 [↗](#)). To accomplish this, we first performed a pseudotime trajectory analysis to infer paths of differentiation in the mesenchymal lineage cells. We resolved three pseudotime trajectories (two osteogenic, one adipogenic) originating from the MPC cell cluster and ending in either Ocy, OB2, or MALP cell fates (Fig. 3A [↗](#)). It is important to note that given the identification of multiple skeletal stem cells^{23–26}, we do not view these trajectories as lineages, but rather “differentiation paths” (progenitor to mature and/or terminally differentiated cells) that are likely traversed by different subsets of skeletal stem cells.

To identify genes likely impacting BMSC-OB differentiation, we used tradeSeq to identify genes that exhibit dynamic changes in expression along pseudotime²⁷. Prior to performing the tradeSeq analysis, we parsed the pseudotime trajectories into regions that encompass cells associated with each cell type along their respective trajectories (Fig. 3B [↗](#)). We defined multiple cell type boundaries (nine in total) using pseudotime values, which represent points along a trajectory. The tradeSeq analysis was performed for each boundary (Supplementary Table S9). For example, trajectories bifurcate in the LMP cell cluster (Fig. 3A [↗](#)); therefore, cells belonging to the LMP cluster can map to adipogenic and/or osteogenic trajectories depending on their placement along pseudotime. Between a cell type boundary, cells spanning a specific cluster (e.g., LMP) and mapping to a specific trajectory (e.g., osteogenic trajectory) are used as input to analyze gene expression between the start and end points of the cell type boundary (e.g., LMP_to_OBP). We analyzed gene expression within the established cell type boundaries for all trajectories and identified genes that exhibit the most significant differences in expression between the start and end points of the cell type boundaries. The total number of significant trajectory-specific tradeSeq genes ($P_{\text{adj}} < 0.05$) ranged from 87 to 1,697 across the 9 cell type boundaries (Supplementary Table S9, S10–12). The expression of representative marker genes for all cell types as a function of pseudotime were consistent with boundaries defined for each cell type (Fig. 3C [↗](#)).

To provide further support that tradeSeq-identified genes are involved in differentiation, we performed a cell type-specific expression quantitative trait locus (eQTL) analysis for each mesenchymal cell type from the 80 DO mice. We identified 563 genes (eGenes) regulated by a significant *cis*-eQTL in specific cell types of the BMSC-OB scRNA-seq data (Supplementary Table S14). In total, 73 eGenes were also tradeSeq-identified genes in one or more cell type boundaries along their respective trajectories (Supplementary Table S9).

We hypothesized that if tradeSeq genes were responsible for driving mesenchymal differentiation, then the eQTLs that perturb their expression would also impact the proportion of cells at different stages along the cell trajectories. Despite being significantly underpowered for this analysis due to




Figure 2. Overview of the network analysis pipeline

Step 1: For all seven (7) of the mesenchymal lineage cell clusters (MPC, LMP, OBP, OB1, OB2, Ocy, MALP), cell type-specific co-expression modules were generated using iterative Weighted Gene Co-expression Network Analysis (iterativeWGCNA). *Step 2:* Bayesian networks were learned to generate directed networks and model causal interactions between co-expressed genes. *Step 3:* Differentiation Driver Genes (DDGs) were identified by extracting subnetworks (i.e., large 3-step neighborhood) for each gene in each cell type-specific Bayesian network and highlighting those subnetworks that were enriched ($P_{adj} < 0.05$) for trajectory-specific tradeSeq genes for the cell type boundary. *Step 4:* DDGs (and associated networks) were prioritized if the DDG was identified previously as an expression/splicing quantitative trait loci (eQTL/sQTL) that colocated with BMD GWAS associations. Created with *Biorender.com*.




Figure 3. Pseudotime Trajectory Inference analysis and establishment of cell type boundaries for tradeSeq analysis

(A) Three (3) trajectories (two adipogenic, one adipogenic) were inferred from the mesenchymal cell clusters of the BMSC-OB scRNA-seq data using Slingshot. All trajectories originate from the MPC and end in either osteogenic (Ocy, OB2) or adipogenic (MALP) cell fates. (B) For each of the trajectories, cell type boundaries were generated using pseudotime values along the trajectories, which encompass the majority of cells of a cell type mapping to their respective trajectory. (C) Normalized gene expression of select genes associated with each cluster are represented in feature plots (top) and gene plotted as a function of pseudotime (bottom) for all pseudotime trajectories (color corresponds to cell type annotation observed throughout). Vertical lines (red) represent the cell type (pseudotime) boundaries established for each cell type (label). The cell type boundary for OB1 and OB2 are represented as one red line/label for visualization purposes.

our relatively smaller sample size ($N = 80$), we identified two cell type-specific eGenes where the genotype responsible for the *cis*-eQTL effect was also associated with cell type proportions. The first of these genes was Pyruvate Kinase, muscle (*Pkm*), which was identified as a significant global tradeSeq gene ($P_{\text{adj}} = 8.35 \times 10^{-8}$; Supplementary Table S13) associated with the transition from LMPs to OBPs along an osteogenic trajectory (Fig. 4A). Moreover, *Pkm* served as an eGene in the LMP cell cluster (LOD = 9.72; Fig. 4B, Supplementary Table S14). Mice inheriting at least one PWK allele at this locus ($N = 15$) demonstrated lower *Pkm* expression (Fig. 4C) and a notable reduction in mature osteoblasts (OB1) and osteocyte-like cells (Ocy) proportions ($P = 0.030$ and $P = 0.026$, respectively), while LMP proportions were unaffected (Fig. 4D, Supplementary Table S15). Similarly, S100 calcium binding protein A1 (*S100a1*) was an OBP to OB1 transition tradeSeq gene ($P_{\text{adj}} = 0.023$; Fig. 4A, Supplementary Table S13) and an eGene in the OBP cell cluster (LOD = 10.12; Fig. 4B, Supplementary Table S14). Mice inheriting at least one 129 allele at this locus ($N = 30$) had higher *S100a1* expression, while the opposite was observed for mice inheriting NZO alleles ($N = 14$) (Fig. 4C). Additionally, mice inheriting at least one 129 allele showed a significant decrease in LMP proportion and increase in OB1 proportion ($P = 0.008$ and $P = 0.016$, respectively) (Fig. 4D, Supplementary Table S15), while no significant differences were observed in cell type proportions among mice inheriting NZO alleles at this locus (Supplementary Fig. 2, Supplementary Table S15). These data support the role of tradeSeq-identified genes in the differentiation of mesenchymal cell types.

Identification of differentiation driver genes (DDG)

In order to discover BMSC-OB differentiation genes potentially responsible for BMD GWAS associations, the next step of our network analysis leveraged the trajectory-specific tradeSeq genes identified for each cell type boundary (Supplementary Table S10-12) to identify differentiation driver genes (DDGs) (Step 3, Fig. 2). We identified DDGs by extracting subnetworks (i.e., large 3-step neighborhoods; see Methods) for each gene in each cell type-specific Bayesian network and identifying those subnetworks enriched ($P_{\text{adj}} < 0.05$) for trajectory-specific tradeSeq genes for the cell type boundary. The analysis identified 408 significant DDGs (Supplementary Table S16, S17-19). We performed a PANTHER²⁸ Gene Ontology (GO) analysis for the cell type boundaries yielding a sufficient number of DDGs and found that DDGs for the osteogenic cell type boundaries (LMP_to_OBP, OBP_to_OB1, OBP_to_OB2) were enriched for genes associated with the cell cycle (GO:0007049; $N = 23, 18, 23$; $P = 1.12 \times 10^{-6}, 1.29 \times 10^{-13}, 1.0 \times 10^{-14}$, respectively) (Supplementary Table S20-22). The DDGs for the adipogenic cell type boundary (LMP_to_MALP, MALP_to_end) were enriched for genes associated with extracellular matrix organization (GO:0030198; $N = 10$; $P = 1.62 \times 10^{-7}$) and lipid metabolic processes (GO:0006629; $N = 25$; $P = 1.83 \times 10^{-11}$), respectively (Supplementary Table S23-24). Across all 408 DDGs, 49 (12%) were identified in one or more cell type boundaries as genes with a significant alteration ($P < 0.05$) of whole-body BMD when knocked-out/down in mice, as reported by the International Mouse Knockout Consortium (IMPC)²⁹ (Supplementary Table S17-19).

We used our previously generated list of potentially causal BMD GWAS genes ($N=1,037$) to subsequently prioritize the DDGs (Step 4, Fig. 2). Of the 408 DDGs, 21 DDGs in one or more cell type boundaries were genes that have BMD GWAS associations that colocalize with sQTL/eQTL (Table 1). The majority of these DDGs were identified in LMPs along both the osteogenic (LMP_to_OBP) and adipogenic (LMP_to_MALP) trajectories ($N = 10$ and 6, respectively; Supplementary Table S16, S25). The remaining DDGs were identified in OBPs along both osteoblast trajectories (OBP_to_OB1, OBP_to_OB2; $N = 1$ and 3, respectively) and MALPs (MALP_to_end; $N = 6$). Additionally, 3 of the 21 DDGs (*Tet1*, *Tpx2*, *Timp2*) are IMPC genes that exhibit a significant alteration of BMD (Supplementary Table S16, S25).




Figure 4. TradeSeq-identified genes associated with BMSC-OB differentiation exhibit eQTL effects.

(A) *Pkm* was identified as a significant global tradeSeq-identified gene ($P_{\text{adj}} = 8.35 \times 10^{-8}$) for LMP cells along an osteogenic trajectory (LMP_to_OBP) (left). *S100a1* was identified as a significant global tradeSeq-identified gene ($P_{\text{adj}} = 0.023$) for OBP cells along osteogenic trajectory 1 (OBP_to_OB1) (right). (B) Plots indicating the cell type-specific expression quantitative trait loci (eQTL) signal for both *Pkm* and *S100a1*. A negative eQTL effect on *Pkm* expression was observed in LMPs for Diversity Outbred (DO) mice with a PWK haplotype background at the *Pkm* locus (left). A positive eQTL effect on the expression of *S100a1* was observed in OBPs for DO mice with a 129 haplotype background at the *S100a1* locus, while a negative effect was observed for NZO mice (right). (C) The expression of *Pkm* and *S100a1* based on DO mouse (expression values transformed via variance stabilizing transformation (VST), as described in Methods). Genotype abbreviations correspond to DO haplotype background (legend) at the respective gene locus. Mice with at least one PWK allele (genotype abbreviation G) tend to have decreased expression of *Pkm* (left). Mice with at least one 129 allele (genotype abbreviation C) tend to have increased expression of *S100a1*, while NZO mice (genotype abbreviation E) have decreased expression (right). (D) PWK mice had a significant reduction in mature osteoblasts (OB1) and osteocyte-like cells (Ocy) proportions relative to other mice ($P = 0.030$ and $P = 0.026$, respectively; t-test), while LMP proportions were unaffected. Asterisks represent any of the other haplotype backgrounds. 129 mice showed a significant decrease in LMP proportion and increase in OB1 proportion ($P = 0.008$ and $P = 0.016$, respectively; t-test), but OBP proportions were unaffected. No significant effects on cell type proportions were observed in NZO mice (Supplementary Fig. 2).

Trajectory	Cell type boundary	DDG	GTEx Tissue with strongest eQTL colocalization (RCP)	GTEx Tissue with strongest sQTL colocalization (H4PP)
1	LMP to OBP	<i>Tet1</i>	Adipose (Visceral); 0.3191	-
1	LMP to OBP	<i>Tpx2</i>	Testis; 0.2031	-
1	LMP to OBP	<i>Cdk1</i>	-	Pituitary; 0.7795
1	LMP to OBP	<i>Ttyh3</i>	-	Liver; 0.9350
1	LMP to OBP	<i>Olfml3</i>	Artery (Aorta); 0.8048	-
1	LMP to OBP	<i>Izumo4</i>	-	Brain (Hypothalamus); 0.9182
1	LMP to OBP	<i>Sec24d</i>	Nerve (Tibial); 0.2677	-
1	LMP to OBP	<i>Tmem263</i>	Adipose (Subcutaneous); 0.5704	Cultured cells (fibroblasts); 0.9716
1	LMP to OBP	<i>Lmf2</i>	-	Adrenal Gland; 0.8181
1	LMP to OBP	<i>Tln2</i>	Esophagus (Muscularis); 0.9697	-
1	OBP to OB1	<i>Kremen1</i>	Heart (Left Ventricle); 0.8686	-
2	OBP to OB2	<i>Kremen1</i>	Heart (Left Ventricle); 0.8686	-
2	OBP to OB2	<i>Ebf1</i>	-	Testis; 0.8760
2	OBP to OB2	<i>Lrp4</i>	Pancreas; 0.7943	-

Table 1. Prioritized Differentiation Driver Genes (DDGs) that have BMD GWAS associations that colocalize with splicing/expression QTL (eQTL/sQTL) identified in a Genotype-Tissue Expression project (GTEx) tissue.

The tissue with the most significant colocalization (RCP and/or H4PP) is listed for each DDG (26 total, 21 distinct), as determined from Al-Barghouthi *et al.* (2022) and Abood *et al.* (2023) for eQTL and sQTL, respectively^{7,16}. RCP = Regional Colocalization Probability (GWAS and eQTL colocalization). H4PP = H4 Posterior Probability (GWAS and sQTL colocalization).

3	LMP to MALP	<i>Ttyh3</i>	-	Liver; 0.9350
3	LMP to MALP	<i>Fgfr11</i>	Cultured cells (fibroblasts); 0.1611	-
3	LMP to MALP	<i>Ebf1</i>	-	Testis; 0.8760
3	LMP to MALP	<i>Ppp1r12b</i>	-	Nerve (Tibial); 0.8807
3	LMP to MALP	<i>Rhoj</i>	Cultured cells (fibroblasts); 0.352	Breast; 0.7844
3	LMP to MALP	<i>Tln2</i>	Esophagus (Muscularis); 0.9697	-
3	MALP to end	<i>Adh1</i>	-	Esophagus (Gastroesophageal Junction); 0.9999
3	MALP to end	<i>Fgfr11</i>	Cultured cells (fibroblasts); 0.1611	-
3	MALP to end	<i>Adcy5</i>	-	Esophagus (Gastroesophageal Junction); 0.8456
3	MALP to end	<i>Cnn2</i>	-	Spleen; 0.7743
3	MALP to end	<i>Mxra8</i>	-	Pituitary; 0.7545
3	MALP to end	<i>Timp2</i>	-	Testis; 0.9429

Table 1. (continued)

Network analysis predict *Fgfr1* and *Tpx2* as novel regulators of BMD

Here we highlight two DDGs that putatively impact human BMD via their roles in LMP differentiation along either an adipogenic (*Fgfr1*) or osteogenic (*Tpx2*) trajectory, which are genes with potential roles that have been minimally characterized in the context of human BMD. Based on our previous work⁷, *Fgfr1* (fibroblast growth factor receptor-like 1) was identified as a DDG with significant human BMD GWAS associations that also colocalized with eQTL identified in the cultured fibroblast GTEX tissue (RCP = 0.1611, Table 1 [↗](#)). The *Fgfr1* network was enriched for tradeSeq-identified genes (N = 6 genes, $P_{\text{adj}} = 7.5 \times 10^{-3}$) for LMPs along an adipogenic trajectory (Fig. 5A [↗](#)). An increase in the expression of all tradeSeq-identified genes for the *Fgfr1* network was observed (Fig. 5B [↗](#), Supplementary Table S12). Importantly, the expression pattern for the tradeSeq-identified genes were consistent with the cell type boundaries established for LMPs differentiating along the adipogenic trajectory toward the MALP cell state (Fig. 5B [↗](#)). Furthermore, in the surrounding *Fgfr1* network, two genes (*Plpp3* and *Cfap100*) have significant human BMD GWAS associations that also colocalized with sQTL in GTEX tissues, as reported in our previous study¹⁶. In the *Fgfr1* network, many other genes can be associated with adipocyte function (e.g., *Lpl*, *Plpp3*, *Igfbp4*)^{30–32} and the maintenance of cilia (e.g., *Cfap100*, *St5* (*Denn2b*), *Mark1*)^{33–35}.

The other network we identified, the *Tpx2* network, was identified for LMPs along an osteogenic trajectory (Fig. 5C [↗](#)). *Tpx2* (TPX2, microtubule-associated) is a DDG with significant human BMD GWAS associations that also colocalized with eQTL identified in the Testis GTEX tissue (RCP = 0.2031, Table 1 [↗](#)). The network was enriched for tradeSeq-identified genes (N = 9 genes, $P_{\text{adj}} = 5.7 \times 10^{-7}$) for LMPs differentiating along the osteogenic trajectory (Fig. 5C [↗](#)). Furthermore, the expression of the tradeSeq-identified genes for the *Tpx2* network were consistent with the cell type boundaries established for LMPs differentiating along the osteogenic trajectory toward the OBP (osteoblast progenitor) cell state (Fig. 5D [↗](#); Supplementary Table S10). The expression of these genes increase as LMPs differentiate into OBPs and subsequently decrease upon reaching an OBP cell state. Additionally, *Tpx2* exhibited a significant alteration of BMD in both male and female mutant mice (Genotype P-value = 1.03×10^{-3}) from IMPC (Fig. 5E [↗](#)). In regards to the constituents of the *Tpx2* network, additional genes have been tested by the IMPC and result in a significant impact on BMD, such as *Ube2c*, *Top2a*, and *Papss1*. Many other genes in the *Tpx2* network can be associated with cellular division and proliferation, including four of the genes of the kinesin family (*Kif*) motor protein genes³⁶: *Kif4*, *Kif11*, *Kif15*, *Kif23*.

Discussion

BMD GWAS has been successful at identifying thousands of SNPs associated with disease; however, the identification of causal genes and defining their functional role in disease remains challenging. The integration of “-omics” data, particularly transcriptomics, can assist in overcoming this challenge. Leveraging transcriptomics data has proven invaluable to informing GWAS, as demonstrated in studies that use these data to perform eQTL mapping, transcriptome-wide association studies (TWASs), and co-expression/gene regulatory network reconstruction. GWAS associations can colocalize with predicted sources of genetic variation that perturb causal gene function or expression, thus providing a potential mechanism through which associations impact disease. While bulk RNA-seq data has been the foundation of such analyses, scRNA-seq data can provide valuable biological context by predicting the cell type in which causal genes are affected. To inform BMD GWAS, the generation of population-scale transcriptomics data at single-cell resolution in bone-relevant cell types can assist in the discovery of novel gene targets. Here, we perform scRNA-seq on 80 DO mice to generate single-cell transcriptomics data of mesenchymal cell types relevant to bone. Using these data, our goal was to prioritize putative causal genes and provide biological context in which these genes potentially influence disease, at cell type-specific




Figure 5. *Fgfr1* and *Tpx2* are prioritized DDGs and putative drivers of mesenchymal differentiation.

(A) *Fgfr1* was identified as a Differentiation Driver Gene (DDG) of a network for LMPs differentiating along an adipogenic trajectory. The network is enriched ($P_{adj} = 7.5 \times 10^{-3}$) for trajectory-specific tradeSeq-identified genes for the LMP_to_MALP cell type boundary (*Hmt*, *St5*, *Igfbp4*, *Cyp1b1*, *Pdzrn4*, *Mark1*). *Fgfr1* was previously identified as a gene that has BMD GWAS associations that colocalize with an eQTL in the cultured fibroblast GTEx tissue. **(B)** An increase in the expression of tradeSeq-identified genes coincides with the LMP_to_MALP cell type boundary in which they were identified as significant. **(C)** *Tpx2* was identified as a DDG of a network for LMPs differentiating along an osteogenic trajectory. The network is enriched ($P_{adj} = 5.7 \times 10^{-7}$) for tradeSeq-identified genes for the LMP_to_OBP cell type boundary (*Tpx2*, *Top2a*, *Kif4*, *Iqgap3*, *Prc1*, *Kif11*, *Ect2*, *Sgo2a*, *Ube2c*). *Tpx2* is both a tradeSeq gene and previously identified as a gene that has BMD GWAS associations that colocalize with an eQTL in the Testis GTEx tissue. **(D)** An increase in the expression of tradeSeq-identified genes coincides with the LMP_to_OBP cell type boundary in which they were identified as significant. **(E)** Box plot displaying whole-body bone mineral density (BMD) measurements (excluding skull) from the International Mouse Knockout Consortium (IMPC) for *Tpx2* mutant mice, which exhibited a significant increase in BMD (Genotype P-value = 1.03×10^{-3}) in both male and female mice (N = 8 (M) and 8 (F) mutants; N = 2574 (M) and 2633 (F) controls)

resolution. Through our pseudotemporal gene expression and network analyses, we identified 21 networks governed by predicted differentiation driver genes (DDGs) that have corresponding human BMD GWAS associations colocalizing with eQTL/sQTL in a GTEx tissue.

We demonstrate that the BMSC-OB model serves as an effective method to enrich for mesenchymal lineage cells, particularly bone-relevant cells. We characterized cells from 80 mice and identified both osteogenic and adipogenic cells derived from the mesenchymal lineage, such as two populations of osteoblasts (OB1 and OB2), osteocyte-like cells (Ocy), and MALPs. Our trajectory inference analysis identified three distinct trajectories in which mesenchymal progenitor cells give rise to both osteogenic and adipogenic cell types, thus portraying biologically relevant and known paths of differentiation of mesenchymal progenitor cells. Pseudotemporal gene expression was analyzed along each trajectory, in a cell type-specific fashion, to identify genes that were changing the most as a function of pseudotime (tradeSeq-identified genes). Subsequent *cis*-eQTL analysis indicated that the expression of some tradeSeq-identified genes were associated with the relative proportion of cell types. While instances such as these were rare, they illustrate that the potential consequence of genetic variation impacting the expression of tradeSeq-identified genes may impact differentiation and the abundances of certain cell types along a trajectory.

To inform BMD GWAS, we utilized the scRNA-seq data in a network analysis to contextualize causal genes (and their associated networks) by predicting the cell types through which these genes are likely acting. Towards this goal, we generated cell type-specific Bayesian networks from our BMSC-OB scRNA-seq data. Our approach was similar to our previous network analyses where bulk RNA-seq data was leveraged to identify genes with strong evidence of playing central roles in networks^{10,11,14}. In contrast, here we utilized scRNA-seq data to identify DDGs and prioritize networks based on the likelihood that they are involved in the differentiation of mesenchymal lineage cells (based on network connections enriched for tradeSeq-identified genes determined from inferred trajectories). Leveraging our previous work^{7,16}, we prioritized DDGs if they were genes with BMD GWAS associations colocalizing with human eQTL/sQTL in a GTEx tissue. Together, a gene being both a DDG and having BMD GWAS associations that colocalize with eQTL/sQTL is strong support of causality.

We identified 21 DDGs and associated networks, some of which have little to no known prior connection to bone. We contextualize these causal genes and their networks by not only providing cell type predictions in which they likely operate, but also providing information regarding the biological processes they likely affect. For example, the *Tpx2* network was identified in LMPs differentiating along an osteogenic trajectory. *Tpx2* is a microtubule assembly factor that interacts with spindle microtubules during cellular division³⁷. The expression of *Tpx2* and its regulation is associated with osteosarcoma, as well as other cancers³⁸. In our previous study, *Tpx2* was identified as a gene that has BMD GWAS associations that colocalize with eQTL in the Testis GTEx tissue⁷. While GTEx does not maintain bone tissue, eQTL are shared across many tissues²¹; therefore, non-bone eQTL may exert their effects in cell types associated with bone, such as LMPs, and evidence of a human eQTL effect indicates that genetic variation can modulate the expression of *Tpx2*. Additionally, when knocked out by IMPC, *Tpx2* exhibited a significant increase in whole body BMD in mice, thus providing strong support for *Tpx2* influencing the regulation of BMD in humans. In the surrounding gene neighborhood of the *Tpx2* network, other genes can be associated with cellular division as well, such as Topoisomerase 2A (*Top2a*) and the kinesin family (*Kif*) genes^{36,39}. Taken together, these results indicate a potential role of *Tpx2* as a mediator of BMD and genetic variation altering its expression could affect microtubule maintenance during the expansion of osteogenic cell populations.

Additionally, the *Fgfr1* network was identified in LMPs differentiating along an adipogenic trajectory. Fibroblast growth factor receptor-like 1 (*Fgfr1*) is presumed to function as a decoy receptor that interacts with FGF ligands necessary for FGF signaling^{40,41} and *Fgfr1* expression is suggested to play a role in both adipogenic and osteogenic differentiation⁴². Our previous study also identified *Fgfr1*, which has BMD GWAS associations that colocalize with eQTL in the cultured fibroblasts GTEx tissue¹⁴. In the neighborhood of the *Fgfr1* network, *Lpl*, *Plpp3*, *Igfbp4* have well-

established roles in adipocyte function and metabolism^{30–32}; however, other genes can be associated with cilia, such as *Cfap100*, *St5 (Denn2b)*, *Mark1*^{33–35}. Interestingly, the maintenance and remodeling of cilia is essential to the differentiation of mesenchymal stem cells and pre-adipocytes (e.g., MALPs) while mature adipocytes lack cilia⁴³. Moreover, the inactivation of FGF signaling is associated with the length of primary cilia⁴⁴. Thus, genetic variation altering the expression of expression of *Fgfr1* may affect FGF signaling to impact the maintenance of cilia and adipogenic differentiation. Additionally, given the prioritization of MALPs in the CELLECT analysis and the well-established inverse relationship between marrow adiposity and BMD^{45,46}, skewed balance of LMP differentiation toward marrow adipogenic cell fates may affect BMD. In summary, the *Fgfr1* network harbors genes involved in adipogenic function, including cilia maintenance, which may contribute to LMP differentiation along an adipogenic trajectory. Together, these results indicate a potential role of *Fgfr1* as a mediator of BMD via its role in adipogenic differentiation and maintenance of cilia.

Analyses performed here are not without limitations to consider. Our *in vitro* culturing approach and the preparation of single cells for scRNA-seq could be sources of technical variation in our study. Additionally, a pitfall of scRNA-seq is the sparsity of the resulting data, which yields an increased frequency of zero values for the expression of some genes in a proportion of cells, also known as “drop-outs”⁴⁷. While statistical approaches can be employed to impute missing data, the accuracy of such methods and whether or not the resulting improvement in transcriptomic signal recovery is enough to warrant such intervention is contentious^{48,49}. However, this issue may be partially offset given the larger scale of the scRNA-seq performed in this study and the average expression approach performed for network and eQTL analysis. Another limitation of this study is that read alignment of the scRNA-seq data did not account for DO founder genetic variation in RNA transcripts, which could affect read mapping and gene expression measurements. An additional limitation is that the BMSC-OB model does not capture osteoclasts, another cell type associated with bone tissue. Importantly, results from our CELLECT analysis indicate that BMD heritability was not enriched for genes whose expression was more specific to osteoclast-like cells; however, these cells likely represent immature osteoclasts, as mature multinucleated cells would be too large to be captured for sequencing. Lastly, while our study employed 80 DO mice, the issue of statistical power is still a limitation; however, we demonstrate that the BMSC-OB model is amenable to high throughput and the inclusion of hundreds of mice, thus statistical power will be improved in future studies.

In summary, we showcase the use of large-scale scRNA-seq data to inform GWAS by performing a network analysis to contextualize BMD GWAS associations. Through the use of multiple single-cell analyses, we have expanded upon our understanding of the genetics of BMD. Our work exemplifies the power of single-cell transcriptomics from large populations of genetically diverse samples and our network approach for data analysis may guide future studies to consider systems genetics strategies for the discovery of genetic determinants of disease.

Methods

Sample preparation and scRNA-seq

We prepared our samples in the same fashion as performed previously in Al-Barghouthi and colleagues¹⁵. In brief, bone marrow was extracted from the femurs of initially 77 DO mice. BMSCs were grown to confluence after 3 days of incubation in 48-well plates and then underwent *in vitro* osteoblast differentiation for 10 days with osteogenic differentiation media (alpha MEM, 10% FBS, 1% pen/strep, 1% Glutamax, 50 µg/µL ascorbic acid [Sigma, St. Louis, MO, USA], 10 nM B-glycerophosphate [Sigma], 10 nM dexamethasone [Sigma]). After differentiation, single cells were liberated from mineralizing cultures via incubations with 60 mM ethylenediaminetetraacetic acid pH 7.4 (EDTA [Thermo Fisher Scientific], made in DPBS), 8 mg/mL collagenase (Gibco) in HBSS/4 mM CaCl₂ (Fisher), and 0.25% trypsin–EDTA (Gibco). After single-cell isolation, cells from mice were pooled into groups containing cells from four to five mice total and concentrated to 800 cells/µL in PBS supplemented with 0.1% BSA (bovine serum albumin). Pooled single cells were prepared

for sequencing using the 10× Chromium Controller (10× Genomics, Pleasanton, CA, USA) with the Single Cell 3' v2 reagent kit, according to the manufacturer's protocol. Libraries were sequenced on the NextSeq500 (Illumina, San Diego, CA, USA).

scRNA-seq analysis pipeline

The data was subsequently processed using the 10× Genomics Cell Ranger toolkit (version 5.0.0) using the GRCm38 reference genome⁵⁰. Using Seurat⁵¹ (version 4.1.0), a combined Seurat object containing all cells was generated with the inclusion of features detected in at least three cells and cells with at least 200 features detected. We used Souporecell⁵² (version 2.0.0) to deconvolve the genotypes of all mice and to remove doublet cells. Cells were assigned to their associated DO mouse by making a pairwise comparison between allele calls made by the shared variants captured between Souporecell and GigaMUGA genotype arrays generated for all mice in the cohort, as previously performed in Dillard and colleagues¹⁵. Cell derived from two mice (176 and 244) were removed in some analyses due to poor genotyping of their respective Souporecell clusters, thus yielding a total of 75 DO mice from this study and 5 DO mice from our previous study¹⁵ for a total of 80 DO mice. We filtered out cells with more than 6200 reads and less than 400 reads, as well as those cells with more than 10% mitochondrial reads. Further, cells were removed if they expressed greater than 20% *Rpl* and 15% *Rps* reads, which equates to cells approximately exceeding the 98 percentile. After filtering, 139,392 cells remained and the resulting object underwent standard normalization, scaling, and the top 3000 features were modeled from a variance stabilizing transformation (VST) using Seurat. Cell-cycle markers based on Tirosh and colleagues⁵³ were regressed out using the “CellCycleScoring” and scaling functions. For subsequent dimensionality reduction, 15 principal components (PCs) were summarized. Then, a kNN ($k = 20$) graph was created and the Louvain algorithm was used to cluster cells at a resolution of 0.5. Annotation of cell type clusters was performed manually based on differential gene expression analysis using the Seurat “FindAllMarkers” function (Supplementary Table S1).

For subsequent WGCNA and eQTL mapping, transcriptomic profiles for each cell type cluster were generated for each sample using a mean expression approach, as performed similarly by others^{54,55}. For each sample contributing at least 5 cells to a given cluster, unnormalized unique molecular identifier (UMI) counts of gene expression for all cells in the cluster for the sample were averaged and then rounded to the nearest hundredth decimal place. A total of 80, 80, 77, 67, 50, 76, 80 mice contributed enough cells to the MPC, LMP, OBP, OB1, OB2, Ocy, and MALP cell type clusters, respectively. Genes with non-zero expression values in fewer than 15 samples were removed. A total of 11971, 15162, 14857, 13674, 13825, 14136, and 14534 genes remained for the MPC, LMP, OBP, OB1, OB2, Ocy, and MALP clusters, respectively. Samples were normalized by computing CPMs (counts per million) without log transformation for each gene using edgeR⁵⁶ (version 4.0.7), then transformed via VST using DESeq2⁵⁷ (version 1.42.0), and quantile normalized using preprocessCore (version 1.60.2).

Trajectory and tradeSeq Analysis

Trajectory inference analysis was performed using Slingshot⁵⁸ (version 1.8.0) on the mesenchymal lineage cell clusters (seven total) of the BMSC-OB scRNA-seq data. The starting cluster was set as the MPC cluster upon the removal of a small outlier population of cells. Trajectories were inferred using 15 PCs. TradeSeq²⁷ (version 1.4.0) was used to analyze gene expression along the trajectories by fitting a negative binomial generalized additive model (NB-GAM) to each gene using the “fitGAM” function with $nknots = 10$, which was determined by using the “evaluateK” function. Prior to performing the tradeSeq analysis, the scRNA-seq data was downsampled to reduce the size of the dataset to approximately 10,000 cells (sampled at random across all seven clusters).

All cell type boundaries were established to encompass on average 78% of cells of a cell cluster (Supplementary Table S9). To identify genes significantly changing between boundaries in a trajectory-specific fashion, we first performed tradeSeq to compare gene expression within each trajectory (two osteogenic, one adipogenic) to highlight genes with a significant difference in expression between boundaries using the “startVsEndTest” function (Supplementary Table S9, S10-

12). Next, we performed a global test with tradeSeq to compare gene expression between trajectories in order to highlight genes exhibiting a significant difference in expression using the “startVsEndTest” function (Supplementary Table S9, S13). All tests were performed with the $\log_2$ fold change threshold ($\log_2$ fc) = 0.5. For all global and trajectory-specific tests, the P-values associated with each gene were adjusted to control the false discovery rate using the “p.adjust” function from the stats (version 4.2.1) R package and genes were filtered to include those with a $P_{\text{adj}} < 0.05$.

CELLECT Analysis

CELLECT²⁰ (CELL-type Expression-specific integration for Complex Traits) (version 1.1.0) was used to identify likely etiologic cell types underlying complex traits of both the BMSC-OBs scRNA-seq data (Fig. 1E, Supplementary Table S6). CELLECT P-values were adjusted using the Bonferroni correction. CELLECT quantifies the association between the GWAS signal and cell type expression specificity using the S-LDSC genetic prioritization model⁵⁹. Summary statistics from the UK Biobank eBMD and Fracture GWAS (Data Release 2018) and cell type annotations from each scRNA-seq data set were used as input. Cell type expression specificities were estimated using CELLEX²⁰ (CELL-type EXpression-specificity) (version 1.2.1) (Supplementary Table S26).

WGCNA

Cell type-specific mean expression matrices (as obtained above) were used as input to generate signed co-expression network modules (Supplementary Table S7, S8). IterativeWGCNA²² (version 1.1.6) was used from a Singularity container built from a Docker hub image⁶⁰. A soft threshold (power) of 14, which exceeded a R^2 threshold of 0.85 for all cell type clusters, was selected for module construction (Supplementary Fig. 1). Modules were generated using iterativeWGCNA with default parameters for the “blockwiseModules” function, a minimum module size of 20 genes, minCoreKME = 0.7, and minKMEtoStay = 0.5.

Bayesian network learning

Bayesian networks were learned from each of the cell type-specific modules of co-expressed genes with the bnlearn (version 4.8.3). Gene expression matrices containing the genes for each module were used as input to the “mmhc” function which employs the Max-Min Hill Climbing algorithm (MMHC) algorithm⁶¹ to learn the underlying structure of the Bayesian network. From the generated networks, igraph (version 1.6.0) was used to resolve 3-step neighborhoods⁶². Nodes (genes) that were unconnected to a neighborhood or connected to only one neighbor were removed. Neighborhoods were filtered to include those with a size greater than 1 standard deviation from the mean across all neighborhoods generated for the network.

DDGs (differentiation driver genes) are genes that yield large 3-step neighborhoods that are enriched ($P_{\text{adj}} < 0.05$) with tradeSeq-identified genes for a given cell type boundary. We calculated whether each neighborhood contained more tradeSeq-identified genes (for the neighborhoods’ associated cell type boundary) than would be expected by chance using the hypergeometric distribution (“phyper” function) from the stats (version 4.2.1) R package. The arguments were as follows: q : (number of neighbors in a neighborhood that are also tradeSeq-identified genes for a given cell type boundary) – 1; m : total number of tradeSeq-identified genes for a given cell type boundary; n : (total number of identified neighborhoods) – m ; k : neighborhood size (total number of neighbors); lower.tail = false. P-values were adjusted to control the false discovery rate using the “p.adjust” function from the stats (version 4.2.1) R package. These pruning steps resulted in a total of 408 DDGs and associated networks for all cell types (Supplementary Table S16, S17-19).

DO eQTL mapping

Prior to performing the eQTL analysis, DNA was extracted from the tails of the 80 DO mice, using the PureLink Genomic DNA mini kit (Invitrogen) and genotyped using the GigaMUGA array by Neogen Genomics (GeneSeek; Lincoln, NE). Processing and quality control of genotype data, including calculation of genotype/allele probabilities, was performed as previously described in Al-Barghouthi and colleagues¹⁴. Cell type-specific mean expression matrices (as obtained above)

for mesenchymal lineage clusters were used as input for the eQTL mapping, which was performed using a linear mixed model (LMM) via the “scan1” function from the [qtl2](#)⁶³ (version 0.30) R package with allowances for the following covariates: sex, age at sacrifice (in days), weight, length, and DO mouse generation. To identify significant eQTL, we calculated a LOD (logarithm of the odds) threshold; for each cell type cluster, we chose 50 genes at random and then permuted them 1000 times using the “scan1perm” function from [qtl2](#). We established the LOD threshold of 9.68 and 9.49 for the autosomal chromosomes and X chromosome, respectively, by taking the average of the median LOD across each cell type. A total of 563 eQTL exceeded the LOD thresholds and were no more than 1 Mbp from the transcription start site of the associated eGene (Supplementary Table S14).

Cell type proportion analysis

To account for technical sources of variation often retained in scRNA-seq, cell type proportions were transformed using the arcsin (asin) square root transformation from the [speckle](#)⁶⁴ R package (version 0.0.3). Tests of statistical significance were performed using the propeller t-test and ANOVA functions with default parameters. Sex of the mice and the batch each mouse was associated with for sequencing were modeled as covariates. Transformed values were used as input for computing tests of statistical differences of cell type proportions between mice, as well as correlation to phenotypic traits (Supplementary Table S3-S5).

Data Availability Statement

The data that support the findings of this study are openly available in NCBI Gene Expression Omnibus database with accession codes GSE152806 and GSE269583. Processed scRNA-seq data available on Zenodo at <https://zenodo.org/records/15299631> [↗](#).

Acknowledgements

Research reported in this publication was supported in part by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under award numbers R01AR68345, R01AR082880, and R01AR077992 to CRF.

Additional information

Code Availability Statement

Code for analysis is available on GitHub at https://github.com/Farber-Lab/DO80_project [↗](#).

Author Contributions

Luke J Dillard: Writing - review & editing; Visualization; Conceptualization; Methodology; Formal analysis; Data curation; Writing - original draft. Gina M Calabrese: Methodology. Larry D Mesner: Methodology. Charles R Farber: Writing - review & editing; Funding acquisition; Supervision; Conceptualization.

Funding

Funder	Grant reference number
National Institute of Arthritis and Musculoskeletal and Skin Diseases	R01AR68345
National Institute of Arthritis and Musculoskeletal and Skin Diseases	R01AR082880
National Institute of Arthritis and Musculoskeletal and Skin Diseases	R01AR077992

Author ORCID iDs

Luke J Dillard: <http://orcid.org/0000-0001-8293-0479>

Charles R Farber: <http://orcid.org/0000-0002-6748-4711>

Additional files

[Description_of_Supplementary_Tables](#) 

[Supplementary_Figures](#) 

[Supplementary_Tables](#) 

References

1. Lin J. T., Lane J. M. (2004) Osteoporosis: a review. *Clin. Orthop. Relat. Res* 126-134
2. Peacock M., Turner C. H., Econs M. J., Foroud T. (2002) Genetics of osteoporosis. *Endocr. Rev* 23:303-326
3. Johnell O., et al. (2005) Predictive value of BMD for hip and other fractures. *J. Bone Miner. Res* 20:1185-1194
4. Morris J. A., et al. (2019) An atlas of genetic influences on osteoporosis in humans and mice. *Nat. Genet* 51:258-266
5. Cookson W., Liang L., Abecasis G., Moffatt M., Lathrop M (2009) Mapping complex disease traits with global gene expression. *Nat. Rev. Genet* 10:184-194
6. Wen X., Pique-Regi R., Luca F (2017) Integrating molecular QTL data into genome-wide genetic association analysis: Probabilistic assessment of enrichment and colocalization. *PLoS Genet* 13:e1006646
7. Al-Barghouthi B. M., et al. (2022) Transcriptome-wide association study and eQTL colocalization identify potentially causal genes responsible for human bone mineral density GWAS associations. *eLife* 11
8. Li B., Ritchie M. D (2021) From GWAS to Gene: Transcriptome-Wide Association Studies and Other Methods to Functionally Understand GWAS Discoveries. *Front. Genet* 12:713230
9. Akiyama M (2020) Multi-omics study for interpretation of genome-wide association study. *J. Hum. Genet* 66:3-10
10. Calabrese G. M., et al. (2017) Integrating GWAS and Co-expression Network Data Identifies Bone Mineral Density Genes SPTBN1 and MARK3 and an Osteoblast Functional Module. *Cell Syst* 4:46-59.e4
11. Sabik O. L., Calabrese G. M., Taleghani E., Ackert-Bicknell C. L., Farber C. R (2020) Identification of a Core Module for Bone Mineral Density through the Integration of a Co-expression Network and GWAS Data. *Cell Rep* 32:108145
12. GTEx Consortium (2013) The Genotype-Tissue Expression (GTEx) project. *Nat. Genet* 45:580-585
13. GTEx Consortium (2020) The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* 369:1318-1330
14. Al-Barghouthi B. M., et al. (2021) Systems genetics in diversity outbred mice inform BMD GWAS and identify determinants of bone strength. *Nat. Commun* 12:3408
15. Dillard L. J., et al. (2023) Single-Cell Transcriptomics of Bone Marrow Stromal Cells in Diversity Outbred Mice: A Model for Population-Level scRNA-Seq Studies. *J. Bone Miner. Res* 38:1350-1363
16. Abood A., et al. (2023) Long-read proteogenomics to connect disease-associated sQTLs to the protein isoform effectors of disease. *bioRxiv* 2023.03.17.531557
<https://doi.org/10.1101/2023.03.17.531557>
17. Bogue M. A., Churchill G. A., Chesler E. J (2015) Collaborative Cross and Diversity Outbred data resources in the Mouse Phenome Database. *Mamm. Genome* 26:511-520

18. Churchill G. A., Gatti D. M., Munger S. C., Svenson K. L (2012) The Diversity Outbred mouse population. *Mamm. Genome* **23**:713-718
19. Zhong L., et al. (2020) Single cell transcriptomics identifies a unique adipose lineage cell population that regulates bone marrow environment. *eLife* **9**
20. Timshel P. N., Thompson J. J., Pers T. H (2020) Genetic mapping of etiologic brain cell types for obesity. *eLife* **9**
21. GTEx Consortium, et al. (2017) Genetic effects on gene expression across human tissues. *Nature* **550**:204-213
22. Greenfest-Allen E., Cartailier J.-P., Magnuson M. A., Stoeckert C. J. (2017) iterativeWGCNA: iterative refinement to improve module detection from WGCNA co-expression networks. *bioRxiv* 234062 <https://doi.org/10.1101/234062>
23. Chan C. K. F., et al. (2018) Identification of the Human Skeletal Stem Cell. *Cell* **175**:43-56.e21
24. Mizuhashi K., et al. (2018) Resting zone of the growth plate houses a unique class of skeletal stem cells. *Nature* **563**:254-258
25. Debnath S., et al. (2018) Discovery of a periosteal stem cell mediating intramembranous bone formation. *Nature* **562**:133-139
26. Matsushita Y., et al. (2020) A Wnt-mediated transformation of the bone marrow stromal cell identity orchestrates skeletal regeneration. *Nat. Commun* **11**:332
27. Berge Van den, et al. (2020) Trajectory-based differential expression analysis for single-cell sequencing data. *Nat. Commun* **11**:1201
28. Thomas P. D., et al. (2022) PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci* **31**:8-22
29. Groza T., et al. (2023) The International Mouse Phenotyping Consortium: comprehensive knockout phenotyping underpinning the study of human disease. *Nucleic Acids Res* **51**:D1038-D1045
30. Enerbäck S., Ohlsson B. G., Samuelsson L., Bjursell G (1992) Characterization of the human lipoprotein lipase (LPL) promoter: evidence of two cis-regulatory regions, LP-alpha and LP-beta, of importance for the differentiation-linked induction of the LPL gene during adipogenesis. *Mol. Cell. Biol* **12**:4622-4633
31. Federico L., et al. (2018) Lipid phosphate phosphatase 3 regulates adipocyte sphingolipid synthesis, but not developmental adipogenesis or diet-induced obesity in mice. *PLoS One* **13**:e0198063
32. Maridas D. E., DeMambro V. E., Le P. T., Mohan S., Rosen C. J (2017) IGFBP4 Is Required for Adipogenesis and Influences the Distribution of Adipose Depots. *Endocrinology* **158**:3488-3500
33. Sigg M. A., et al. (2017) Evolutionary Proteomics Uncovers Ancient Associations of Cilia with Signaling Pathways. *Dev. Cell* **43**:744-762.e11
34. Kumar R., et al. (2022) A cell-based GEF assay reveals new substrates for DENN domains and a role for DENND2B in primary ciliogenesis. *Sci Adv* **8**:eabk3088
35. Fumoto K., et al. (2019) Mark1 regulates distal airspace expansion through type I pneumocyte flattening in lung development. *J. Cell Sci* **132**
36. Miki H., Setou M., Kaneshiro K., Hirokawa N (2001) All kinesin superfamily protein, KIF, genes in mouse and human. *Proc. Natl. Acad. Sci. U. S. A* **98**:7004-7011
37. Zhang R., Roostalu J., Surrey T., Nogales E (2017) Structural insight into TPX2-stimulated microtubule assembly. *eLife* **6**
38. Zhu D., Xu X., Zhang M., Wang T (2022) TPX2 regulated by miR-29c-3p induces cell proliferation in osteosarcoma via the AKT signaling pathway. *Oncol. Lett* **23**:143
39. Uusküla-Reimand L., Wilson M. D (2022) Untangling the roles of TOP2A and TOP2B in transcription and cancer. *Sci Adv* **8**:eadd4920
40. Trueb B (2011) Biology of FGFR1, the fifth fibroblast growth factor receptor. *Cell. Mol. Life Sci* **68**:951-964

41. Steinberg F., et al. (2010) The FGRL1 receptor is shed from cell membranes, binds fibroblast growth factors (FGFs), and antagonizes FGF signaling in *Xenopus* embryos. *J. Biol. Chem* **285**:2193-2202
42. Kähkönen T. E., et al. (2018) Role of fibroblast growth factor receptors (FGFR) and FGFR like-1 (FGFRL1) in mesenchymal stromal cell differentiation to osteoblasts and adipocytes. *Mol. Cell. Endocrinol* **461**:194-204
43. Hilgendorf K. I (2021) Primary Cilia Are Critical Regulators of White Adipose Tissue Expansion. *Front. Physiol* **12**:769367
44. Neugebauer J. M., Amack J. D., Peterson A. G., Bisgrove B. W., Yost H. J (2009) FGF signalling during embryo development regulates cilia length in diverse epithelia. *Nature* **458**:651-654
45. Fazeli P. K., et al. (2013) Marrow fat and bone--new perspectives. *J. Clin. Endocrinol. Metab* **98**:935-945
46. Veldhuis-Vlug A. G., Rosen C. J (2018) Clinical implications of bone marrow adiposity. *J. Intern. Med* **283**:121-139
47. Haque A., Engel J., Teichmann S. A., Lönnberg T (2017) A practical guide to single-cell RNA-sequencing for biomedical research and clinical applications. *Genome Med* **9**:75
48. Cheng Y., Ma X., Yuan L., Sun Z., Wang P (2023) Evaluating imputation methods for single-cell RNA-seq data. *BMC Bioinformatics* **24**:302
49. Yu X., Abbas-Aghababazadeh F., Chen Y. A., Fridley B. L (2021) Statistical and Bioinformatics Analysis of Data from Bulk and Single-Cell RNA Sequencing Experiments. *Methods Mol. Biol* **2194**:143-175
50. Church D. M., et al. (2009) Lineage-specific biology revealed by a finished genome assembly of the mouse. *PLoS Biol* **7**:e1000112
51. Hao Y., et al. (2021) Integrated analysis of multimodal single-cell data. *Cell* **184**:3573-3587.e29
52. Heaton H., et al. (2020) Souporecell: robust clustering of single-cell RNA-seq data by genotype without reference genotypes. *Nat. Methods* **17**:615-620
53. Tirosh I., et al. (2016) Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352**:189-196
54. Neavin D., et al. (2021) Single cell eQTL analysis identifies cell type-specific genetic control of gene expression in fibroblasts and reprogrammed induced pluripotent stem cells. *Genome Biol* **22**:76
55. van der Wijst M. G. P., et al. (2018) Single-cell RNA sequencing identifies celltype-specific cis-eQTLs and co-expression QTLs. *Nat. Genet* **50**:493-497
56. Robinson M. D., McCarthy D. J., Smyth G. K (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139-140
57. Love M. I., Huber W., Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550
58. Street K., et al. (2018) Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics* **19**:477
59. Finucane H. K., et al. (2015) Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet* **47**:1228-1235
60. Greenfest E., Cartailier J. P., Stoeckert C., Brestelli J. (2018) IterativeWGCNAGitHub. version: 1.1.6 <https://github.com/cstoeckert/iterativeWGCNA>
61. Tsamardinos I., Brown L. E., Aliferis C. F (2006) The max-min hill-climbing Bayesian network structure learning algorithm. *Mach. Learn* **65**:31-78
62. Porter M. D., Smith R. (2010) Network neighborhood analysis. <https://doi.org/10.1109/isi.2010.5484781>
63. Broman K. W., et al. (2019) R/qt2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations. *Genetics* **211**:495-502

64. Phipson B., et al. (2022) propeller: testing for differences in cell type proportions in single cell data. *Bioinformatics* **38**:4720-4726
65. Marsh S., Salmon M., Hoffman P. (2023) Custom Visualizations & Functions for Streamlined Analyses of Single Cell SequencingZenodo. <https://doi.org/10.5281/zenodo.7534950>

Peer reviews

Reviewer #1 (Public review):

In this manuscript, Dillard and colleagues integrate cross-species genomic data with a systems approach to identify potential driver genes underlying human GWAS loci and establish the cell type(s) within which these genes act and potentially drive disease.

Specifically, they utilize a large single cell RNA-seq (scRNA-seq) dataset from an osteogenic cell culture model - bone marrow-derived stromal cells cultured under osteogenic conditions (BMSC-OBs) - from a genetically diverse outbred mouse population called the Diversity Outbred (DO) stock to discover network driver genes that likely underlie human bone mineral density (BMD) GWAS loci. The DO mice segregate over 40M single nucleotide variants, many of which affect gene expression levels, therefore making this an ideal population for systems genetic and co-expression analyses.

The current study builds on previous published work from the same group that used co-expression analysis to identify co-expressed "modules" of genes that were enriched for BMD GWAS associations. In this study, the authors utilized a much larger scRNA-seq dataset from 80 DO BMSC-OBs, inferred co-expression based on Bayesian networks for each identified mesenchymal cell type, focused on networks with dynamic expression trajectories that are most likely driving differentiation of BMSC-OBs, and then prioritized genes ("differentiation driver genes" or DDGs) in these osteogenic differentiation networks that had known expression or splicing QTLs (eQTL/sQTLs) in any GTEx tissue that co-localized with human BMD GWAS loci. The systems analysis is impressive, the experimental methods are described in detail, and the experiments appear to be carefully done. The computational analysis of the single cell data is comprehensive and thorough, and the evidence presented in support of the identified DDGs, including *Tpx2* and *Fgfr11*, is for the most part convincing. Some limitations in the data resources and methods hamper enthusiasm somewhat and are discussed below.

Overall, while this study will no doubt be valuable to the BMD community, the cross-species data integration and analytical framework may be more valuable and generally applicable to the study of other diseases, especially for diseases with robust human GWAS data but for which robust human genomic data in relevant cell types is lacking.

Specific strengths of the study include the large scRNA-seq dataset on BMSC-OBs from 80 DO mice, the clustering analysis to identify specific cell types and sub-types, the comparison of cell type frequencies across the DO mice, and the CELLECT analysis to prioritize cell clusters that are enriched for BMD heritability (Figure 1). The network analysis pipeline outlined in Figure 2 is also a strength, as is the pseudotime trajectory analysis (results in Figure 3).

Potential drawbacks of the authors' approach include their focus on genes that were previously identified as having an eQTL or sQTL in any GTEx tissue. The authors rightly point out that the GTEx database does not contain data for bone tissue, but reason that eQTLs can be shared across many tissues - this assumption is valid for many cis-eQTLs, but it could also exclude many genes as potential DDGs with effects that are specific to bone/osteoblasts. Indeed, the authors show that important BMD driver genes have cell-type specific eQTLs. Another issue concerns potential model overfitting in the iterativeWGCNA analysis of mesenchymal cell type-specific co-expression, which identified an average of 76 co-expression modules per cell cluster (range 26-153). Based on the limited number of genes that

are detected as expressed in a given cell due to sparse per cell read depth (400-6200 reads/cell) and drop outs, it's surprising that as many as 153 co-expression modules could be distinguished within any cell cluster. I would suspect some degree of model overfitting is responsible for these results.

Overall, though, these concerns are minor relative to the many strengths of the study design and results. Indeed, I expect the analytical framework employed by the authors here will be valuable to -- and replicated by -- researchers in other disease areas.

Comments on revisions:

Thank you for addressing my concerns. This is an impressive study and manuscript that you should be proud of.

<https://doi.org/10.7554/eLife.100832.2.sa2>

Reviewer #2 (Public review):

Summary:

In this manuscript, Farber and colleagues have performed single cell RNAseq analysis on bone marrow derived stem cells from DO Mice. By performing network analysis, they look for driver genes that are associated with bone mineral density GWAS associations. They identify two genes as potential candidates to showcase the utility of this approach.

Strengths:

The study is very thorough and the approach is innovative and exciting. The manuscript contains some interesting data relating to how cell differentiation is occurring and the effects of genetics on this process. The section looking for genes with eQTLs that differ across the differentiation trajectory (Figure 4) was particularly exciting.

Weaknesses:

The manuscript is, in parts, hard to read due to the use of acronyms and there are some questions about data analysis that still need to be addressed.

Comments on revisions:

Dillard et al have made several improvements to their manuscript.

(1) We previously asked the authors to determine whether any cell types were enriched for BMD-related traits since the premise of the paper is that 'many genes impacting BMD do so by influencing osteogenic differentiation or ... adipogenic differentiation'. Given the potential for the cell culture method to skew the cell type distribution non-physiologically, it is important to establish which cell types in their assay are most closely associated with BMD traits. The new CELLECT analysis and Figure 1E address this point nicely. However, it would still be nice to see the correlations between these cell types and BMD traits in the mice as this would provide independent evidence to support their physiological importance more broadly.

(2) Shortening the introduction.

(3) Addressing limitations that arise from not accounting for founder genome SNPs when aligning scRNA-seq data.

(4) The main take-away of this paper is, to us, the development of a single cell approach to studying BMD-related traits. It is encouraging that the cells post-culture appear to be representative of those pre-culture (supplemental figure 3).

However, the authors seem to have neglected several comments made by both reviewers. While we share the authors' enthusiasm for the single cell analytical approach, we do not understand their reluctance to perform further statistical tests. We feel that the following comments have still not been addressed:

(1) The manuscript still contains the following:

"To provide further support that tradeSeq-identified genes are involved in differentiation, we performed a cell type-specific expression quantitative trait locus (eQTL) analysis for each mesenchymal cell type from the 80 DO mice. We identified 563 genes (eGenes) regulated by a significant cis-eQTL in specific cell types of the BMSC-OB scRNA-seq data (Supplementary Table S14). In total, 73 eGenes were also tradeSeq-identified genes in one or more cell type boundaries along their respective trajectories (Supplementary Table S9)."

The purpose of this paragraph is to convince readers that the eGenes approach aligns with the tradeSeq approach (and that their approach can therefore be trusted). It is essential that such claims are supported by statistical reasoning. Given that it would be very simple to perform permutation/enrichment analyses to address this point, and both reviewers requested similar analyses, we do not understand the author's reluctance here. Otherwise, this section should be rewritten so that it does not imply that the identification of these genes provides support for their approach.

(2) Given that a central purpose of this manuscript is to establish a systematic workflow for identifying candidate genes, the manuscript could still benefit from more explanation as to why the authors chose to highlight Tpx2 and Fgfr11. Tpx2 does already have a role in bone physiology through the IMPC. The authors should comment on why they did not explore Kremen1, for instance, as this gene seems important for the transition to both OB1 and 2.

A final minor comment is that it would be very helpful if the authors could indicate if the DDGs in Table 1 are also eGenes for the relevant cell type. This is much more meaningful than looking through GTEx.

<https://doi.org/10.7554/eLife.100832.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

In this manuscript, Dillard and colleagues integrate cross-species genomic data with a systems approach to identify potential driver genes underlying human GWAS loci and establish the cell type(s) within which these genes act and potentially drive disease. Specifically, they utilize a large single-cell RNA-seq (scRNA-seq) dataset from an osteogenic cell culture model - bone marrow-derived stromal cells cultured under osteogenic conditions (BMSC-OBs) - from a genetically diverse outbred mouse population called the Diversity Outbred (DO) stock to discover network driver genes that likely underlie human bone mineral density (BMD) GWAS loci. The DO mice segregate over 40M single nucleotide variants, many of which affect gene expression levels, therefore making this an ideal population for systems genetic and co-expression analyses. The current study builds on previously published work from the same group that used co-expression analysis to identify co-expressed "modules" of genes that were enriched for BMD GWAS associations. In this study, the authors utilize a much larger scRNA-seq dataset from 80 DO BMSC-OBs, infer co-expression-based and Bayesian networks for each identified mesenchymal cell type, focused on networks with dynamic expression trajectories that are most likely driving differentiation of BMSC-OBs, and then prioritized genes

*("differentiation driver genes" or DDGs) in these osteogenic differentiation networks that had known expression or splicing QTLs (eQTL/sQTLs) in any GTEx tissue that colocalized with human BMD GWAS loci. The systems analysis is impressive, the experimental methods are described in detail, and the experiments appear to be carefully done. The computational analysis of the single-cell data is comprehensive and thorough, and the evidence presented in support of the identified DDGs, including *Tpx2* and *Fgfr1*, is for the most part convincing. Some limitations in the data resources and methods hamper enthusiasm somewhat and are discussed below. Overall, while this study will no doubt be valuable to the BMD community, the cross-species data integration and analytical framework may be more valuable and generally applicable to the study of other diseases, especially for diseases with robust human GWAS data but for which robust human genomic data in relevant cell types is lacking.*

Specific strengths of the study include the large scRNA-seq dataset on BMSC-OBs from 80 DO mice, the clustering analysis to identify specific cell types and sub-types, the comparison of cell type frequencies across the DO mice, and the CELLECT analysis to prioritize cell clusters that are enriched for BMD heritability (Figure 1). The network analysis pipeline outlined in Figure 2 is also a strength, as is the pseudotime trajectory analysis (results in Figure 3). One weakness involves the focus on genes that were previously identified as having an eQTL or sQTL in any GTEx tissue. The authors rightly point out that the GTEx database does not contain data for bone tissue, but the reason that eQTLs can be shared across many tissues - this assumption is valid for many cis-eQTLs, but it could also exclude many genes as potential DDGs with effects that are specific to bone/osteoblasts. Indeed, the authors show that important BMD driver genes have cell-type-specific eQTLs. Furthermore, the mesenchymal cell type-specific co-expression analysis by iterative WGCNA identified an average of 76 co-expression modules per cell cluster (range 26-153). Based on the limited number of genes that are detected as expressed in a given cell due to sparse per-cell read depth (400-6200 reads/cell) and dropouts, it's hard to believe that as many as 153 co-expression modules could be distinguished within any cell cluster. I would suspect some degree of model overfitting here and would expect that many/most of these identified modules have very few gene members, but the methods list a minimum module size of 20 genes. How do the numbers of modules identified in this study compare to other published scRNA-seq studies that use iterative WGCNA?

In the section "Identification of differentiation driver genes (DDGs)", the authors identified 408 significant DDGs and found that 49 (12%) were reported by the International Mouse Knockout [sic] Consortium (IMPC) as having a significant effect on whole-body BMD when knocked out in mice. Is this enrichment significant? E.g., what is the background percentage of IMPC gene knockouts that show an effect on whole-body BMD? Similarly, they found that 21 of the 408 DDGs were genes that have BMD GWAS associations that colocalize with GTEx eQTLs/sQTLs. Given that there are > 1,000 BMD GWAS associations, is this enrichment (21/408) significant? Recommend performing a hypergeometric test to provide statistical context to the reported overlaps here.

We thank the reviewer for their constructive feedback and thoughtful questions. In regards to the iterativeWGCNA, a larger number of modules is sometimes an outcome of the analysis, as reported in the iterativeWGCNA preprint (Greenfest-Allen et al., 2017). While we did not make a comparison to other works leveraging this tool for scRNA-seq, it has been used broadly across other published studies, such as PMID: 39640571, 40075303, 33677398, 33653874. While model overfitting, as you mention, may be a cause for more modules, our Bayesian network analysis we perform after iterativeWGCNA highlights smaller aspects of coexpression modules, as opposed to focusing on the entirety of any given module.

We did not perform enrichment or statistical tests as our goal was to simply highlight attributes or unique features of these genes for additional context.

Reviewer #2 (Public review):

Summary:

In this manuscript, Farber and colleagues have performed single-cell RNAseq analysis on bone marrow-derived stem cells from DO Mice. By performing network analysis, they look for driver genes that are associated with bone mineral density GWAS associations. They identify two genes as potential candidates to showcase the utility of this approach.

Strengths:

The study is very thorough and the approach is innovative and exciting. The manuscript contains some interesting data relating to how cell differentiation is occurring and the effects of genetics on this process. The section looking for genes with eQTLs that differ across the differentiation trajectory (Figure 4) was particularly exciting.

Weaknesses:

The manuscript is in parts hard to read due to the use of acronyms and there are some questions about data analysis that need to be addressed.

We thank the reviewer for their feedback and shared enthusiasm for our work. We tried to minimize the use of technical acronyms as much as we could without compromising readability. Additionally, we addressed questions regarding aspects of data analysis.

Reviewer #1 (Recommendations for the authors):

(1) For increased transparency and to allow reproducibility, it would be necessary for the scripts used in the analysis to be shared along with the publication of the preprint. Also, where feasible, sharing the processed data in addition to the raw data would allow the community greater access to the results and be highly beneficial.

Thank you for this suggestion. The raw data will be available via GEO accession codes listed in the data availability statement. We will make available scripts for some analyses on our Github (https://github.com/Farber-Lab/DO80_project) and processed scRNA-seq data in a Seurat object (.rds) on Zenodo (<https://zenodo.org/records/15299631>)

(2) Lines 55-76: I think the summary of previous work here is too long. I understand that they would like to cover what has been done previously, but this seems like overkill.

Good suggestion. We have streamlined some of the summary of our previous work.

(3) Did the authors try to map QTL for cell-type proportion differences in their BMSC-OBs? While 80 samples certainly limit mapping power, the data shown in Figs 4C/D suggest that you might identify a large-effect modifier of LMP/OB1 proportions.

We did try to map QTL for cell type proportion differences, but no significant associations were identified.

(4) Methods question: Does the read alignment method used in your analysis account for SNPs/indels that segregate among the DO/CC founder strains? If not, the authors may wish to include this in their discussion of study limitations and speculate on how unmapped reads could affect expression results.

The read alignment method we used does not account for SNPs/indels from the DO founder strains that fall in RNA transcripts captured in the scRNA-seq data. We have included this as a limitation in our discussion (line 422-424).

(5) Much of the discussion reads as an overview of the methods, while a discussion of the results and their context to the existing BMD literature is relatively lacking in comparison.

We have added additional explanation of the results and context to the discussion (line 381-382, 396-407).

(6) Figure 1E and lines 146-149: Adjusted p values should be reported in the figure and accompanying text instead of switching between unadjusted and adjusted p values.

We updated Figure 1e to portray adjusted p-values, listed the adjusted p-values in legend of Figure 1e, and listed them in the main text (line 153-154).

(7) Why do the authors bring the IMPC KO gene list into the analysis so late? This seems like a highly relevant data resource (more so than the GTEx eQTLs/sQTLs) that could have been used much earlier to help identify DDGs.

Given that our scRNA-seq data is also from mice, we did choose to integrate information from the IMPC to highlight supplemental features of genes in networks (i.e., genes that have an experimentally-tested and significant effect on BMD in mice). However, our primary goal was to inform human GWAS and leverage our previous work in which we identified colocalizations between human BMD GWAS and eQTL/sQTL in a human GTEx tissue, which is why this information was used to guide our network analysis.

(8) Does Fgfr1 and/or Tpx2 have a cis-eQTL in your BMSC-OB scRNA-seq dataset?

We did not identify cis-eQTL effects for Fgfr1 and Tpx2.

(9) Figure 4B-C: These eQTLs may be real, but based on the diplotype patterns in Figure 4C, I suspect they are artifacts of low mapping power that are driven by rare genotype classes with one or two samples having outlier expression results. For example, if you look at the results in Fig 4C for S100a1 expression, the genotype classes with the highest/lowest expression have lower sample numbers. In the case of Pkm eQTL showing a PWK-low effect, the PWK genome has many SNPs that differ from the reference genome in the 3' UTR of this gene, and I wonder if reads overlapping these SNPs are not aligning correctly (see point 4 above) and resulting (falsely) in lower expression values for samples with a PWK haplotype.

As mentioned above, our alignment method did not consider DO founder genetic variation that is specifically located in the 3' end of RNA transcripts in the scRNA-seq data. We have included this as a limitation in our discussion (line 422-424).

In future studies, we intend to include larger populations of mice to potentially overcome, as you mention, any artifacts that may be attributable to low statistical power, rare genotype classes, or outlier expression.

Reviewer #2 (Recommendations for the authors):

Major Points

(1) The authors hypothesize "that many genes impacting BMD do so by influencing osteogenic differentiation or possibly bone marrow adipogenic differentiation". However, cell type itself does not correlate with any bone trait. Does this indicate that the hypothesis is not entirely correct, as genes that drive these phenotypes would not be

enriched in one particular cell type? The authors have previously identified "high-priority target genes". So, are there any cell types that are enriched for these target genes? If not, this would indicate that all these genes are more ubiquitously expressed and this is probably why they would have a greater effect on the overall bone traits. Furthermore, are the 73 eGenes (so genes with eQTLs in a particular cell type that change around cell type boundaries) or the DDGs (Table 1) enriched for these high-priority target genes?

The bone traits measured in the DO mice are complex and impacted by many factors, including the differentiation propensity and abundance of certain cell types, both within and outside of bone. Though we did not identify correlations between cell type abundance and the bone traits we measured, we tailored our investigations to focus on cellular differentiation using the scRNA-seq data. However, future studies would need to be performed to investigate any connections between cellular differentiation, cell type abundance, and bone traits.

We did not perform enrichment analyses of either the target genes identified from our other work or eGenes identified here, but instead used the target gene list to center our network analysis and the eGenes to showcase the utility of the DO mouse population.

(2) The readability of the paper could be improved by minimising the use of acronyms and there are several instances of confusing wording throughout the paper. In many cases, this can be solved by re-organising sentences and adding a bit more detail. For example, it was unclear how you arrived at Fgfr1 or Tpx2.

One of the goals of our study was to identify genes that have (to our knowledge) little to no known connection to BMD. We chose to highlight Fgfr1 and Tpx2 because there is minimal literature characterizing these genes in the context of bone, which we speak to in the results (line 296-297). Additionally, we prioritized these genes in our previous work and they were identified in this study by using our network analyses using the scRNA-seq data, which we mention in the results (line 276-279).

(3) Technical aspects of the assay. In Figure 1d you show that the cell populations vary considerably between different DO mice. It would be useful to give some sense of the technical variance of this assay given that the assay involves culturing the cells in an exogenous environment. This could take the form of tests between mice within the same inbred strain, or even between different legs of the same DO mice to show that results are technically very consistent. It might also be prudent to identify that this is a potential limitation of the approach as in vitro culturing has the potential to substantially change the cell populations that are present.

We agree that in vitro culturing, in addition to the preparation of single cells for scRNA-seq, are unavoidable sources of technical variation in this study. However, the total number of cells contributed by each of the 80 DO mice after data processing does not appear to be skewed and the distribution appears normal (see added figures, now included as Supplemental Figure 3). Therefore, technical variation is at least consistent across all samples. Nevertheless, we have mentioned the potential for technical variation artifacts in our study in the discussion (line 414-416).

(4) Need for permutation testing. "We identified 563 genes regulated by a significant eQTL in specific cell types. In total, 73 genes with eQTLs were also tradeSeq-identified genes in one or more cell type boundaries". These types of statements are fine but they need to be backed up with permutation testing to show that this level of enrichment is greater than one would expect by chance.

We did not perform enrichment tests as our only goal was to 1. determine if eQTL could be resolved in the DO mouse population using our scRNA-seq data and 2. predict in what cell type the associated eQTL and associated eGene may have an effect.

(5) The main novelty of the paper seems to be that you have used single-cell RNA seq (given that you appear to have already detailed the candidates at the end). I don't think this makes the paper less interesting, but I think you need to reframe the paper more about the approach, and not the specific results. How you landed on these candidates is also not clear. So the paper might be improved by more robustly establishing the workflow and providing guidelines for how studies like this should be conducted in the future.

We sought to not only devise a rigorous approach to analyze our single cell data, but also showcase the utility of the approach in practice by highlighting targets for future research (i.e., *Fgfr1* and *Tpx2*).

Our goal was to identify novel genes and we landed on these candidate genes (*Fgfr1* and *Tpx2*) because they had substantial data supporting their causality and they have yet to be fully characterized in the context of bone and BMD (line 295-297).

In regards to establishing the workflow, we have included rationale for specific aspects of our approach throughout the paper. For example, Figure 2 itemizes each step of our network analysis and we explain why each step is utilized throughout various parts results (e.g., lines 168-170, 179-181, 191-193, 202-203, 257-260, 276-277).

We have added a statement advocating for large-scale scRNA-seq from genetically diverse samples and network analyses for future studies (line 436-438).

Minor Points

(1) In the summary you use the word "trajectory". Trajectories for what? I assume the transition between cell types, but this is not clear.

We added text to clarify the use of trajectory in the summary (line 34).

(2) This sentence: "By 60 identifying networks enriched for genes implicated in GWAS we predicted putatively causal genes 61 for hundreds of BMD associations based on their membership in enriched modules." is also not clear. Do you mean: we predicted putatively causal genes by identifying clusters of co-expressed genes that were enriched for GWAS genes?" It is not clear how you identify the causal gene in the network. Is this just based on the hub gene?

The aforementioned sentence has since been removed to streamline the introduction, as suggested by Reviewer 1.

In regards to causal gene identification, it is not based on whether it is hub gene. We prioritized a DDG (and their associated networks) if it was a causal gene that we identified in our previous work as having eQTL/sQTL in a GTEx tissue that colocalizes with human BMD GWAS.

(3) Figure 3C. This is good but the labels are quite small. Would be good to make all the font sizes larger.

We have enlarged Figure 3C.

(4) Line 341 in the Discussion should be "pseudotemporal".

We have edited "temporal" to "pseduotemporal".

<https://doi.org/10.7554/eLife.100832.2.sa0>