

A new class of receptors: Lipids regulate mammalian Gs α -stimulated adenylyl cyclase activities via their membrane anchors

Reviewed Preprint

v2 • October 23, 2024

Revised by authors

Reviewed Preprint

v1 • October 1, 2024

Marius Landau, Sherif Elsabbagh, Harald Gross, Adrian Fuchs, Anita CF Schultz, Joachim E Schultz 

Pharmazeutisches Institut der Universität Tübingen, Auf der Morgenstelle 8, Germany • Max-Planck-Institut für Biologie, Max-Planck-Ring 5, 72076 Tübingen, Germany

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

This manuscript describes an **important** study of a new lipid-mediated regulation mechanism of adenylyl cyclases. The biochemical evidence provided is **convincing** and will trigger more research in this mechanism. This manuscript will be of interest to all scientists working on lipid regulation and adenylyl cyclases.

<https://doi.org/10.7554/eLife.101483.2.sa3>

Abstract

The biosynthesis of cAMP by mammalian membrane-bound adenylyl cyclases (mACs) is predominantly regulated by G-protein-coupled-receptors (GPCRs). Up to now the two hexahelical transmembrane domains of mACs were considered to fix the enzyme to membranes. Here we show that the transmembrane domains serve in addition as signal receptors and transmitters of lipid signals that control Gs α -stimulated mAC activities. We identify aliphatic fatty acids and anandamide as receptor ligands of mAC isoforms 1 to 7 and 9. The ligands enhance (mAC isoforms 2, 3, 7, and 9) or attenuate (isoforms 1, 4, 5, and 6) Gs α -stimulated mAC activities *in vitro* and *in vivo*. Substitution of the stimulatory membrane receptor of mAC3 by the inhibitory receptor of mAC5 results in a ligand inhibited mAC5-mAC3 chimera. Thus, we discovered a new class of membrane receptors in which two signaling modalities are at a crossing, direct tonic lipid and indirect phasic GPCR-Gs α signaling regulating the biosynthesis of cAMP.

The structure of the first second messenger, cyclic 3',5'-adenosine monophosphate (cAMP), was reported in 1958 by Sutherland and Rall. It was generated by incubation of a liver extract with the first messengers adrenaline or glucagon (1 ). Since then, cAMP has been demonstrated to be an almost universal second messenger used to translate various extracellular stimuli into a uniform intracellular chemical signal (2 , 3 ). The biosynthetic enzymes for cAMP from ATP, adenylyl cyclases (ACs), have been biochemically investigated in bacteria and eukaryotic cells (4 –6 ). Finally in 1989, the first mammalian AC was sequenced (7 ). The protein displayed two catalytic domains (C1 and C2) which are highly similar in all isoforms. The two hexahelical membrane

anchors (TM1 and TM2) are dissimilar and isoform specifically conserved (8). The latter were proposed to possess a channel or transporter-like function, properties, which were never confirmed (7). The domain architecture of mammalian ACs, TM1-C1-TM2-C2, clearly indicated a pseudoheterodimeric protein composed of two concatenated monomeric bacterial precursor proteins (9). To date, sequencing has identified nine mammalian membrane-delimited AC isoforms (mACs) with identical domain architectures (2). In all nine isoforms the catalytic domains are conserved and share extensive sequence and structural similarities which indicate a similar biosynthetic mechanism (5, 6, 10, 11). On the other hand, the associated two membrane domains differ substantially within each isoform and between all nine mAC isoforms (8, 12, 13).

Bioinformatic studies, however, revealed that the membrane domains, TM1 as well as TM2, are highly conserved in an isoform-specific manner for about 0.5 billion years of evolution (8, 12).

Extensive studies on the regulation of the nine mAC isoforms revealed that the G α subunit of the trimeric G-proteins activate cAMP formation. G α is released intracellularly upon stimulation of G-protein-coupled receptors (GPCR), i.e. the receptor function for mAC regulation was assigned to the diversity of GPCRs which presently are most prominent drug targets. In 1995 Tang and Gilman reported that G α regulation of mammalian mACs does not require the presence of the membrane anchors. A soluble C1-C2 dimer devoid of the membrane domains was fully activated by G α (13). This reinforced the view that the mAC membrane anchors which comprise up to 40 % of the protein were just that and otherwise functionally inert. The theoretical possibility of mACs to be regulated directly, bypassing the GPCRs, was dismissed (14).

We were intrigued by the exceptional evolutionary conservation of the mAC membrane anchors for 0.5 billion years (8, 12). In addition, we identified a cyclase-transducing-element that connects the TM1 and TM2 domains to the attached C1 or C2 catalytic domains. These transducer elements are similarly conserved in a strictly isoform-specific manner (12, 15).

Furthermore, cryo-EM structures of mAC holoenzymes clearly revealed that the two membrane domains, TM1 and TM2, collapse into a tight dodecahedral complex resembling membrane receptors (16–18). Lastly, we created a chimeric model involving the hexahelical quorum-sensing receptor from *Vibrio cholerae* which has a known aliphatic lipid ligand and the mAC 2 isoform. We observed that the ligand directly affected, i.e., attenuated the G α activation of mAC2 (19). As a proof-of concept this demonstrated that the cytosolic catalytic AC dimer serves as a receiver for extracellular signals transmitted through the dyad-related membrane anchor. Accordingly, we proposed a general model of mAC regulation in which the extent of the indirect mAC activation via the GPCR/G α axis is under direct ligand control via the mAC membrane anchor (19).

Here we report the results of a rigorous search for potential ligands of the mAC membrane domains. We identified aliphatic lipids as ligands for mAC isoforms 1 to 7 and 9. Isoform-dependently, the ligands either attenuate or enhance G α -activation *in vitro* and *in vivo* demonstrating a receptor function of the mAC proteins. The receptor properties are transferable as demonstrated by interchanging the membrane anchors between mAC3 and 5. Thus, the results define a new class of membrane receptors and establish a completely new level of regulation of cAMP biosynthesis in mammals in which tonic and phasic signaling processes intersect in a central signaling system, which is the target of frequently used drugs.

Oleic acid enhances Gsa-stimulated mAC3, but not mAC5 activity

In earlier experiments we demonstrated regulation of the mycobacterial AC Rv2212 by lipids (20), the presence of oleic acid in the mycobacterial AC Rv1264 structure (21), and regulation of a chimera consisting of the quorum-sensing receptor from *Vibrio* and the mAC2 catalytic dimer by the aliphatic lipid 3-hydroxytridecan-4-one (22). Therefore, we searched for lipids as ligands. Here, we used bovine lung as a starting material because lipids are important for lung development and function (23, 24). Lipids were extracted from a cleared lung homogenate, acidified to pH 1, with dichloromethane/methanol (2:1). The dried organic phase was chromatographed on silica gel (employing vacuum-liquid-chromatography; Si-VLC) and fractions A to Q were assayed (fractionation scheme in appendix 1). Fraction E enhanced mAC3 activity stimulated by 300 nM Gsa four-fold, whereas mAC5 activity was unaffected (Fig. 1A). The non-maximal concentration of 300 nM Gsa was used because it enabled us to observe stimulatory as well as inhibitory effects.

Fraction E was further separated by RP-HPLC into five subfractions (E1 – E5; appendix 1 Fig. 1). The mAC3 enhancing constituents appeared in fraction E2. It enhanced Gsa-stimulated mAC3 four-fold but had no effect on mAC5 (Fig. 1B). ¹H and ¹³C-NMR spectra of fraction E2 indicated the presence of aliphatic lipids (appendix 1 Fig. 2). Subsequent GC/MS analysis identified palmitic, stearic, oleic and myristic acid in E2 (Fig. 1C). Concentration-response curves were established for these fatty acids with mAC3 and mAC5 stimulated by 300 nM Gsa (Fig. 1D). 20 μM oleic acid enhanced Gsa-stimulated mAC3 activity three-fold (EC₅₀ = 10.4 μM) and 20 μM palmitic acid two-fold (EC₅₀ = 6.4 μM), while stearic or myristic acid had no significant effect. None of these fatty acids affected mAC5 activity (Fig. 1D).

The action of oleic acid on mAC3 was linear for >25 min (appendix 2, Fig. 1). The K_m of mAC3 for ATP (335 μM) was unaffected. V_{max} was increased from 0.62 to 1.23 nmol cAMP/mg/min (appendix 2, Fig. 2). Oleic acid did not affect the activity of a soluble, Gsa-stimulated construct formerly used for generating a C1 and C2 catalytic dimer from mAC1 and 2, ruling out spurious detergent effects (13) (appendix 2, Fig. 3). The effect of oleic acid was further evaluated by Gsa concentration response curves of mAC3 and mAC5 in presence and absence of 20 μM oleic acid (Fig. 2, left and center). For mAC3 the calculated EC₅₀ of Gsa in presence and absence of oleic acid were 549 and 471 nM, respectively (not significant). Over the Gsa concentration range tested with mAC3 the enhancement of cAMP formation by 20 μM oleic was uniformly about 3.4-fold (Fig. 2 right). In the case of mAC5, Gsa stimulation was not enhanced by oleic acid (Fig. 2 center and right).

To explore the ligand space, we tested 18 aliphatic C₁₂ to C₂₀ lipids (Table 1; structures in appendix 2 Fig. 4). At 20 μM, elaidic, *cis*-vaccenic and linoleic acids were efficient enhancers of Gsa-stimulated mAC3 activity. Palmitic, palmitoleic, linolenic, eicosa-pentaenoic acids and oleamide were less efficacious; other compounds were inactive (Fig. 3). Notably, the saturated C₁₈ stearic acid was inactive here and throughout, albeit otherwise variations in chain length, and the number, location, and conformation of double bonds were tolerated to some extent, e.g., *cis*-vaccenic, linoleic and linolenic acids. The relaxed ligand specificity was anticipated as aliphatic fatty acids are highly bendable and bind to a flexible dodecahedral protein dimer embedded in a fluid lipid membrane. The ligand space of mAC3 somewhat resembled the fuzzy and overlapping ligand specificities of the free-fatty-acid receptors 1 and 4 (25–27).




Fig. 1.

Identification of mAC3 activating fatty acids.

(A and B) Effect of 1 μg/assay of fractions E from vacuum-liquid-chromatography (A) and E2 from RP-HPLC (B) on 300 nM Gsα-stimulated mAC isoforms 3 and 5. Activities are shown as % compared to 300 nM Gsα stimulation (100%). n=2, each with two technical replicates. Basal and Gsα activities of mAC3 in (A) were 0.01 and 0.07 and of mAC5 0.06 and 1.32 nmol cAMP•mg⁻¹•min⁻¹, respectively. In (B), basal and Gsα activities of mAC3 were 0.02 and 0.12 and of mAC5 0.09 and 1.1 nmol cAMP•mg⁻¹•min⁻¹, respectively. (C) GC-MS chromatogram of fraction E2. Mass spectra of the fatty acids are shown. Fatty acids' identity was confirmed by comparing with corresponding standards (TMS: Trimethylsilyl). (D) Effect of fatty acids identified by GC-MS on 300 nM Gsα-stimulated mAC3 (left) and mAC5 (right). Basal and Gsα activities of mAC3 were 0.023 ± 0.02 and 0.17 ± 0.03 and of mAC5 0.08 ± 0.02 and 0.44 ± 0.09 nmol cAMP•mg⁻¹•min⁻¹, respectively. n=3-23. EC₅₀ of palmitic and oleic acids for mAC3 were 6.4 and 10.4 μM, respectively. Data represent individual experiments (black dots in A and B) or mean ± SEM (D). One-sample t tests were performed. Significances: **P < 0.01, ***P < 0.001; ****P < 0.0001.




Fig. 2.

Concentration-response curves for Gsα and mAC3 and 5 activities in presence or absence of 20 μM oleic acid.

20 μM oleic acid enhances Gsα stimulation of mAC3 (left) but not of mAC5 (center). **(Left)** EC₅₀ of Gsα in the absence of oleic acid was 549 nM and in the presence of 20 μM oleic acid, it was 471 nM (not significant). mAC3 basal activity was 30 ± 24 pmol cAMP·mg⁻¹·min⁻¹. **(Center)** The EC₅₀ of Gsα in the absence of oleic acid was 245 nM and in the presence of oleic acid, it was 277 nM (not significant). mAC5 basal activity was 84 ± 60 cAMP·mg⁻¹·min⁻¹. n=2-3, each with two technical replicates. **(Right)** Fold stimulation of mAC3 and 5 activities incubated with different concentrations of (Gsα + 20 μM oleic acid) (n = 2-3). Data are mean ± SEM, paired t test for left and center, and one-way ANOVA for right. Significances: *P < 0.05.




Fig. 3.

Effect of 20 μ M lipids on 300 nM Gsa-stimulated mAC3.

Basal and Gsa stimulated activities were 0.02 ± 0.001 and 0.17 ± 0.01 nmol cAMP $\cdot$ mg $^{-1}$ $\cdot$ min $^{-1}$, respectively. EPA: Eicosapentaenoic acid; 9-HSA: 9-hydroxystearic acid. Data are mean $\pm$ SEM, n= 2-23. One-sample t tests, Significances: * P < 0.05, ** P < 0.01, *** P < 0.001; **** P < 0.0001.




Fig. 4.

Oleic acid enhances cAMP formation in mAC3-transfected HEK293 cells.

Effect of oleic acid on HEK293 cells permanently transfected with mACs 3 and 5 stimulated by 2.5 μM isoproterenol (set as 100%). Basal and isoproterenol stimulated cAMP levels of HEK- mAC3 were 0.02 ± 0.006 and 1.35 ± 0.24 and of HEK-mAC5 2.13 ± 0.69 and 2.60 ± 0.88 pmol cAMP/10000 cells, respectively. $n=3-9$, carried out in technical triplicates. Data are mean $\pm$ SEM . One-sample t tests. Significances: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Lauric (Dodecanoic) acid
Myristic (Tetradecanoic) acid
Myristoleic ((9Z)-Tetradecenoic) acid
Palmitic (Hexadecanoic) acid
Palmitoleic ((9Z)-Hexadecenoic) acid
Octadecane
1,18-Octadecanedicarboxylic acid
Stearic (Octadecanoic) acid
9-Hydroxystearic acid
Oleic ((9Z)-Octadecenoic) acid
Oleamide ((9Z)-Octadecenamide)
Methyl oleate
2-Oleoyleglycerol
Triolein
Elaidic ((9E)-Octadecenoic) acid
cis-Vaccenic ((11E)-Octadecenoic) acid
Linoleic ((9Z,12Z)-Octadecadienoic) acid
Linolenic ((9Z,12Z,15Z)-Octadecatrienoic) acid
Arachidonic ((5Z,8Z,11Z,14Z)-eicosatetraenoic) acid
Eicosapentaenoic ((5Z,8Z,11Z,14Z,17Z)-eicosapentaenoic) acid

Table 1.
List of lipids tested against mAC isoforms.

Next, the effect of oleic acid was probed *in vivo* using HEK293 cells permanently transfected with mAC3 (HEK-mAC3) or mAC5 (HEK-mAC5). Intracellular cAMP formation via Gs α was triggered via stimulation of the endogenous β -receptor with 2.5 μ M of the β -agonist isoproterenol, a submaximal concentration (concentration-response curve for isoproterenol and HEK-mAC3 see appendix 2; **Fig. 5** [↗](#)). Addition of oleic acid enhanced cAMP formation in HEK-mAC3 1.5-fold (**Fig. 4** [↗](#)). Stearic acid was inactive. Under identical conditions, cAMP formation in HEK-mAC5 cells was unaffected (**Fig. 4** [↗](#)). The EC₅₀ of oleic acid in HEK293- mAC3 cells was 0.5 μ M, i.e., the potency appeared to be increased compared to respective membrane preparations whereas the efficiency was reduced, possibly reflecting the regulatory interplay within the cell. To exclude experimental artifacts, transfection efficiencies were tested by PCR. mAC3 and mAC5 transfections were similar (appendix 2; **Fig. 6** [↗](#)). Taken together, the results suggest that the enhancement of Gs α -stimulated mAC3 by oleic acid might be due to binding of oleic acid to or into an mAC3 membrane receptor ([12](#) [↗](#), [22](#) [↗](#)).

Oleic acid enhances Gs α -stimulated mAC 2, 7, and 9 activities

Next, we examined other AC isoforms with oleic acid as a ligand. 20 μ M oleic acid significantly enhanced Gs α -stimulated activities of isoforms 2, 7, and 9, mAC1 was slightly attenuated, and isoforms 4, 5, 6, and 8 were unaffected (**Fig. 5A** [↗](#)).

Concentration-response curves were carried out for mACs 2, 7, and 9 (**Fig. 5B** [↗](#)). The EC₅₀ of oleic acid were 8.6, 6.7 and 7.8 μ M, respectively, comparable to that determined for mAC3. Exploration of the ligand space for mACs 2, 7 and 9 with the panel of 18 aliphatic lipids uncovered more active lipids (**Fig. 5C** [↗](#)). In the case of mAC2, 20 μ M *cis*-vaccenic acid doubled cAMP formation (EC₅₀ 10.6 μ M) while other compounds were inactive (**Fig. 5C** and [for](#) [↗](#) additional concentration-response curves see appendix 2 **Fig. 7** [↗](#)). For mAC7 the EC₅₀ of elaidic was 9.7 μ M (concentration-response curve see appendix 2 **Fig. 8** [↗](#)). The range of potential ligands for mAC9 was more comprehensive: 3-4-fold enhancement was observed with 20 μ M palmitoleic, oleic, elaidic, *cis*-vaccenic, and linoleic acid. With 20 μ M myristic, palmitic, palmitoleic, linolenic, and arachidonic acid 1.5- 2 fold enhancements were observed (**Fig. 5C** and [for](#) [↗](#) concentration response curves see appendix 2 **Fig.s 9** [↗](#) and **10** [↗](#)).

Arachidonic acid and anandamide inhibit Gs α -stimulated activities of mAC1, 4, 5 and 6

Testing the panel of lipids at 20 μ M with mAC isoforms 1, 4, 5, 6, and 8 we found that isoforms 1 and 4 were significantly attenuated by arachidonic acid, and somewhat less by palmitoleic acid. Other lipids had no effect (for bar plots and dose response curves see appendix 2 **Fig. 11-13**). Of note, eicosa-pentaenoic acid which resembles arachidonic acid but for an additional *cis*- Δ^{17} double bond had no effect on mAC activities (appendix 2 **Fig. 11, 12**). Concentration-response curves for arachidonic acid with 300 nM Gs α -stimulated mAC1 and 4 yielded IC₅₀ of 23 and 36 μ M, respectively, i.e., about two-fold higher compared to the EC₅₀ of enhancing ligands (**Fig. 6A** [↗](#)).

Next, we examined whether arachidonic acid attenuates mAC1 and 4 in intact HEK 293 cells. Surprisingly, cAMP formation in HEK-mAC1 cells stimulated by 10 μ M isoproterenol was attenuated by arachidonic acid with high potency (IC₅₀ = 250 pM), i.e., with higher potency compared to data with membranes prepared from the same cell line. In contrast, mAC4 activity examined under identical conditions was not attenuated (**Fig. 6B** [↗](#)). Currently, we are unable to rationalize these discrepancies. Possibly, mAC4 has another, more specific lipid ligand which is




Fig. 5.

Fatty acids enhance mAC isoforms 2, 7 and 9 activities.

(A) Effect of 20 μ M oleic acid on 300 nM Gs α -stimulated mAC activities normalized to 100%. Basal and Gs α -stimulated activities for each isoform are in appendix 2 [Table 1](#). n= 2-23. **(B)** Oleic acid activates mACs 2, 7, and 9 stimulated by 300 nM Gs α . n=7-23. **(C)** Fatty acids activating mACs 2, 7, and 9 at 20 μ M. For basal and Gs α -stimulated activities, see appendix 2, Fig.'s 6-9. n= 5-15. Identical colours indicate identical compounds. Data are mean $\pm$ SEM. One-sample *t* tests were performed. Significances: **P* < 0.05, ***P* < 0.01, ****P* < 0.001; *****P* < 0.0001.




Fig. 6.

Arachidonic acid attenuates 300 nM Gsa-stimulated activities of mACs 1 and 4. **(A)** Arachidonic acid attenuates Gsa-stimulated mACs 1 and 4. Basal and Gsa stimulated activities of mAC1 were 0.12 ± 0.01 and 0.42 ± 0.03 and of mAC4 0.02 ± 0.002 and 0.14 ± 0.02 nmol cAMP $\cdot$ mg $^{-1}$ $\cdot$ min $^{-1}$, respectively. IC $_{50}$ of arachidonic acid for mAC1 and mAC4 were 23 and 36 μ M, respectively. $n = 3-9$. **(B)** Effect of arachidonic acid on HEK-mAC1 and HEK-mAC4 cells. Cells were stimulated by 10 μ M isoproterenol (set as 100 %) in the presence of 0.5 mM IBMX (3-isobutyl-1-methylxanthine). Basal and isoproterenol stimulated cAMP levels in HEK-mAC1 were 1.03 ± 0.15 and 1.66 ± 0.28 and in HEK-mAC4 0.20 ± 0.04 and 0.86 ± 0.24 pmol cAMP/10000 cells, respectively. IC $_{50}$ for HEK-mAC1 was 250 pM. $n = 2-11$, each with three replicates. Data are mean $\pm$ SEM. One-sample t tests were performed. Significances: ** $P < 0.01$, *** $P < 0.001$; † $P < 0.0001$. For clarity, not all significances are indicated.




Fig. 7.

Anandamide attenuates 300 nM Gsα-stimulated activities of mACs 1, 4, 5 and 6. **(A)** Effect of anandamide on Gsα-stimulated mAC5 and 6. Basal and Gsα activities of mAC5 were 0.05 ± 0.01 and 0.98 ± 0.12 and of mAC6 0.05 ± 0.01 and 0.78 ± 0.12 nmol cAMP $\cdot$ mg $^{-1}\cdot$ min $^{-1}$, respectively. IC₅₀ of anandamide were 42 and 22 μM, respectively. n= 3-32. **(B)** Anandamide attenuates mAC1 but not mAC4 stimulated by Gsα. Basal and Gsα stimulated activities of mAC1 were 0.12 ± 0.01 and 0.40 ± 0.03 and of mAC4 0.02 ± 0.002 and 0.15 ± 0.02 nmol cAMP $\cdot$ mg $^{-1}\cdot$ min $^{-1}$, respectively. IC₅₀ for mAC1 was 29 μM. n= 3-4, each with two technical replicates. **(C)** Effect of anandamide on 2.5 μM isoproterenol stimulated HEK- mAC5. Basal and isoproterenol stimulated cAMP levels of HEK-mAC5 were 1.8 ± 0.22 and 2.4 ± 0.48 pmol cAMP/10000 cells, respectively. The control bar represents 2.5 μM isoproterenol stimulation alone. n=5-6, each with three technical replicates. IC₅₀ of anandamide was 133 μM. Data are mean $\pm$ SEM. One-sample *t* tests (A-B) and paired *t* test (C) were performed. Significances: ns: not significant $P > 0.05$; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. For clarity, not all significances are indicated.




Fig. 8.

Receptor properties are exchangeable between mAC isoforms. (A, left)

Detection of AC5_(membr)-AC3_(cat) receptor chimera. AC5_(membr)-AC3_(cat) (AC5-3) (36) was expressed in HEK293 cells with an N-terminal tag for labeling with the protein ligase Connectase. The membrane preparation was incubated with fluorophore-conjugated Connectase and separated by SDS-PAGE. A fluorescence scan of the gel detects AC5_(membr)-AC3_(cat) (right), the reagent (fluorophore-conjugated Connectase) is detected when using HEK293 membrane (middle) or a buffer control (left); **(A, right)** Design of the chimeric AC5-3 construct. Numbers are amino acid positions in mAC3 and 5, respectively. **(B)** Gsa concentration response curve of mAC5-3. Basal activity for mAC5-3 was 0.02 pmol cAMP·mg⁻¹·min⁻¹. Error bars denote SEM of n=3, each with two technical replicates. **(C)** Effect of 20 μM oleic acid on 300 nM Gsa-stimulated mACs 3, 5, and 5-3. Basal and Gsa activities of mACs 3, 5, and 5-3 were 0.02 ± 0.003 and 0.11 ± 0.02, 0.05 ± 0.01 and 0.98 ± 0.12, and 0.01 ± 0.004 and 0.2 ± 0.02 nmol cAMP·mg⁻¹·min⁻¹, respectively. n=7-33. **(D)** Effect of 100 μM Anandamide on 300 nM Gsa-stimulated mACs 3, 5, and 5-3. Basal and Gsa activities of mACs 3, 5, and 5-3 were 0.02 ± 0.002 and 0.19 ± 0.02, 0.05 ± 0.01 and 0.98 ± 0.12 and 0.02 ± 0.003 and 0.23 ± 0.04 nmol cAMP·mg⁻¹·min⁻¹, respectively. n=6-9. IC₅₀ for mAC5 and mAC5-3 were 42 and 29 μM, respectively. **(E)** Exchange of TM domains transfers anandamide effect on mAC3. Basal and Gsa stimulated activities of mAC3 were 0.02 ± 0.002 and 0.12 ± 0.02 nmol cAMP·mg⁻¹·min⁻¹, respectively. Basal and Gsa stimulated activities of mAC5 were 0.05 ± 0.005 and 0.98 ± 0.12 nmol cAMP·mg⁻¹·min⁻¹, respectively. Basal and Gsa stimulated activities of mAC5-3 were 0.02 ± 0.002 and 0.22 ± 0.03 nmol cAMP·mg⁻¹·min⁻¹, respectively. Calculated IC₅₀ concentrations of Anandamide for mAC5 and mAC5-3 were 42 and 29 μM, respectively. Data are mean ± SEM. One-sample *t* tests. Significances: ***P* < 0.01; *****P* < 0.0001.




Fig. 9.

mAC5 – mAC3 receptor-transfer analyzed in HEK293 cells.

Effect of anandamide on HEK-mAC5 and HEK-mAC5-3 cells stimulated by 2.5 μM isoproterenol (set as 100 %). Basal and isoproterenol stimulated cAMP levels in HEK-mAC5 were 1.80 ± 0.22 and 2.29 ± 0.39 and in HEK-mAC5-3 (+ 0.5 mM IBMX) 0.17 ± 0.02 and 3.11 ± 0.55 pmol cAMP/10000 cells, respectively. $n = 4-11$. IC_{50} for HEK-mAC5 and HEK-mAC5-3 were 133 and 60 μM, respectively. Anandamide had no effect on basal activity of HEK-mAC5 and stimulated HEK-mAC3 cells in concentrations up to 100 μM (data not shown). Data are mean $\pm$ SEM. One-sample t tests were performed. Significances: * $P < 0.05$; *** $P < 0.001$.




Fig. 10.

Oleic acid concentration-dependently potentiates mAC activity in brain cortical membranes from mouse. Basal and 300 nM Gsα activities were 0.4 ± 0.1 and 2.7 ± 0.7 nmol cAMP•mg⁻¹•min⁻¹, respectively. N= 4-6. One-sample *t* test: **P* < 0.05; ***P* < 0.01 compared to 100% (300 nM Gsα stimulation).

needed in *in vivo*. In general, the enhancing and attenuating effects bolster the hypothesis of specific receptor-ligand interactions and divergent intrinsic activities for different ligands. Of note, it was reported that arachidonic acid at concentrations up to 1 mM inhibits AC activity in brain membrane fractions and that essential fatty acid deficiency effects AC activity in rat heart (28, 29).

At this point we were lacking ligands for mACs 5, 6, and 8 (appendix 2 Fig. 14-16). Possibly, the negative charge of the fatty acid headgroups might impair receptor interactions. A neutral lipid neurotransmitter closely related to arachidonic acid is arachidonylethanolamide (anandamide) (30). Indeed, anandamide attenuated 300 nM Gs α -stimulation of mAC5 and 6 with IC₅₀ of 42 and 23 μ M, respectively, i.e., comparable to the effect of arachidonic acid on mACs 1 and 4, and distinctly less potently than the ligands for mAC 2, 3, 7, and 9 (Fig. 7A).

mACs 5 and 6 may thus represent new targets for anandamide which is part of a widespread neuromodulatory system (31). The concentrations of arachidonic acid and anandamide required may be achieved *in vivo* by local biosynthesis and degradation. An interfacial membrane-embedded phosphodiesterase cleaves the phosphodiester bond of the membrane lipid *N*-arachidonylethanolamine-glycerophosphate releasing anandamide into the extracellular space (30, 32, 33). The lipophilicity and lack of charge should enable it to diffuse readily. Whether the mACs and this biosynthetic phosphodiesterase colocalize or associate with its target mACs is unknown. Degradation of anandamide is by a membrane-bound amidase, generating arachidonic acid and ethanolamine (34). Therefore, we examined whether anandamide at higher concentrations might also affect mAC1 and 4. In fact, anandamide significantly attenuated Gs α -stimulated mAC1, but distinctly not mAC4 (Fig. 7B). The IC₅₀ of anandamide for mAC1 was 29 μ M.

We also tested whether anandamide attenuated cAMP formation *in vivo* using HEK-mAC5 cells primed by 2.5 μ M isoproterenol (Fig. 7C). 100 μ M Anandamide attenuated cAMP formation by only 23 % in HEK293-mAC5 cells, the effect was significant ($P < 0.01$).

At this point, we were unable to identify a ligand for mAC8, presumably another lipid.

Receptor properties are exchangeable between mAC isoforms 3 and 5

To unequivocally validate specific mAC-ligand-receptor interactions and regulation we generated a chimera in which the enhancing membrane domains of mAC3, i.e., mAC3-TM1 and TM2, were substituted by those of mAC5 (design in Fig. 8A, right). The intention was to obtain a chimera, mAC5_(membr)-AC3_(cat), with a loss of receptor function, i.e., no enhancement by oleic acid, and a gain of another receptor function, i.e., attenuation of activity by anandamide. Successful expression and membrane insertion of the chimera in HEK293 cells was demonstrated by specific conjugation to Cy5.5 fluorophore, using the protein ligase connectase (Fig. 8A, left (35)). cAMP synthesis of isolated membranes from these cells was stimulated up to 10-fold by addition of 300 nM Gs α , comparable to membranes with recombinant mAC3 or mAC5 proteins (Fig. 8B). mAC activity in the mAC5_(membr)-AC3_(cat) chimera was not enhanced by oleic acid, i.e., loss of receptor function, but was attenuated by anandamide, i.e., gain of receptor function (Fig. 8C and D). The attenuation was comparable to results obtained with mAC5 membranes, i.e., IC₅₀ was 29 μ M mAC5_(membr)-AC3_(cat) (Fig. 8E) compared to 42 μ M for mAC5. This means that the attenuating receptor property of mAC5 was successfully grafted onto the mAC3-catalytic dimer. We take this to support the hypothesis that the mammalian mAC membrane domains operate as receptors using lipid ligands. The data virtually rule out unspecific lipid effects such as disturbance of membrane

integrity by intercalation and surfactant or detergent effects. In addition, the data demonstrated that the signal most likely originates from the receptor entity and is transmitted through the subsequent linker regions to the catalytic dimer.

The findings were further substantiated *in vivo* using HEK293-mAC5_(membr)-AC3_(cat) cells. cAMP formation primed by 2.5 μ M isoproterenol was attenuated by anandamide in HEK293-mAC5_(membr)-AC3_(cat) cells by 66%, (**Fig. 9**), i.e., a gain of function which remarkably exceeded the anandamide attenuation in HEK293-mAC5 cells of 23 %. In HEK293-mAC5_(membr)-AC3_(cat) cells oleic acid was ineffective, i.e., loss of function (data not shown).

The results bolster the notion that mAC isoforms are receptors with lipids as ligands.

Lastly, we prepared membranes from mouse brain cortex in which predominantly mAC isoforms 2, 3 and 9 are expressed, isoforms with demonstrated enhancement of Gsa stimulation by oleic acid (37). In cortical membranes 20 μ M oleic acid enhanced Gsa stimulated cAMP formation 1.5-fold with an EC₅₀ of 5 μ M, almost identical to the one determined for mAC2, 3, 7 and 9 (**Fig. 10**). This suggests that mACs in brain cortical membranes are similarly affected by fatty acids.

Discussion

In the past, the biology of the two membrane anchors of mACs, highly conserved in an isoform-specific manner, remained unresolved. The theoretical possibility of a receptor function of these large hexahelical anchor domains, each comprising 150-170 amino acids, was considered unlikely (14). Our data are a transformative step toward resolving this issue and introduce lipids as critical participants in regulating cAMP biosynthesis in mammals. The first salient discovery is the identification of the membrane domains of mACs as a new class of receptors for chemically defined ligands which set the level of stimulation by the GPCR/Gsa system. This conclusion is based on (i) the dodecahelical membrane domains of the nine mAC receptors have distinct, conserved isoform-specific sequences for the TM1 and TM2 domains (12); (ii) the receptors have distinct ligand specificities and affinities in the lower micromolar range; (iii) isoform dependently ligands either enhance or attenuate Gsa-stimulated mAC activities; (iv) receptor properties are transferable between isoforms by interchanging membrane domains; (v) isoproterenol-stimulated formation of cAMP *in vivo* is affected by addition of extracellular ligands. (vi) Gsa-stimulated cAMP formation in mouse cortical membranes is enhanced by oleic acid. Therefore, the results establish a new class of receptors, the membrane domains of mACs, with lipids as ligands. The data question the utility of the currently used mAC sub-classification, which groups mAC1, 3, 8, mAC2, 4, 7, mACs 5, 6, and mAC9 together (2). At this point, mAC 1, 4, 5, and 6, which are ligand-attenuated, may be grouped together and a second group may consist of isoforms 2, 3, 7, and 9 which are ligand-enhanced. Our data do not contradict earlier findings concerning regulation of mACs via GPCRs, cellular localization of mAC isoforms or regional cAMP signaling (2). Instead, the data reveal a completely new level of regulation of cAMP biosynthesis in which two independent modalities of signaling, i.e., direct, tonic lipid signaling and indirect phasic signaling via the GPCR/Gsa circuits intersect at the crucial biosynthetic step mediated by the nine mAC isoforms.

The second important finding is the observation that the extent of enhancement of mAC3 activity by 20 μ M oleic acid is uniform up to 1000 nM Gsa (**Fig. 2** right). We suppose that in mAC3 the equilibrium of two differing ground states favors a Gsa unresponsive state and the effector oleic acid concentration-dependently shifts this equilibrium to a Gsa responsive state (19). In contrast, the equilibrium of ground states of mAC5 probably is opposite, i.e. the one accessible to Gsa stimulation predominates and stimulation by Gsa is maximal. In addition, oleic acid has little effect because the mAC5 receptor domain does not bind oleic acid (**Fig. 1D**, right, and **Fig. 2**, center). A ligand for mAC5, e.g., anandamide or arachidonic acid, likely shifts the equilibrium of ground states to a Gsa unresponsive state and inhibits stimulation. The biological balance of

ground states appears to be an intrinsic property which is isoform- specifically imprinted in mACs. Probably, it defines a major element of regulation and enables distinct inhibitory or stimulatory inputs by extracellular lipid ligands. The ground states probably are separated by a low transition energy and are stabilized by receptor occupancy. Hitherto available structures required Gsa and/or forskolin for stabilization and probably did not capture different ground states (14, 16, 17, 38–40). Mechanistically, tonic levels of lipid ligands affect the ground states and thus set the bounds of cAMP formation elicited by phasic GPCR/Gsa-stimulation. As such lipid signaling through the mAC membrane receptors appears to represent a higher level of a systemic regulatory input based on constant monitoring the physiological and nutritional status of an organism.

Lipid signaling is much less characterized than solute signaling (41). Most of the highly functionalized ligands for GPCRs are storable in vesicles and the release, inactivation and removal are strictly controlled. On the other hand, the very nature of lipids, i.e., high flexibility of aliphatic chains, low water solubility, propensity for nonspecific protein binding, membrane permeability and potential effects on membrane fluidity complicate discrimination between extra- and intracellular lipid actions (26). Yet, viewed from an evolutionary perspective, lipids possibly are ideal primordial signaling molecules because for the emergence of the first cell lipids were required to separate an intra- and extracellular space. Conceivably, lipids derived from membrane lipids were used for regulatory purposes early-on. In association with the evolution of bacterial mAC progenitors lipid ligands may have persisted in evolution and regulation by GPCR/Gsa in metazoans was acquired and expanded later.

The concentrations of free fatty acids in serum or interstitial fluid usually are rather low, mostly below the EC₅₀ concentrations determined in this study (27, 42, 43). This raises the question of the origin of lipid ligands under physiological conditions. It is well known that cell membranes are highly dynamic (29, 44). Within limits, cell morphology and lipid composition are in constant flux to accommodate diverse functional requirements.

Remodeling of cell membranes is accomplished by targeted phospholipid biosynthesis and by regulated lipolysis of membrane lipids, e.g., by phospholipase A2, mono- and diacylglycerol lipases or lipoprotein lipases (45). Therefore, a speculative possibility is that lipid ligands are acutely and locally generated directly from membrane lipids (33, 46). Such regulation probably happens at the level of individual cells, cellular networks, complex tissues and whole organs. Additional potential lipid sources available for ligand generation may be, among others, lipids in nutrients (29), exosomes present in blood, serum lipids, chylomicrons, blood triglycerides and lipids originating from the microbiome. On the one hand, the discovery of lipids as ligands for the mAC receptors broadens the basis of regulation of cAMP biosynthesis with potentially wide-ranging consequences in health and disease. On the other hand, the data pose the challenge to identify how the tonic signal is generated and regulated

Materials and methods

Reagents and materials

ATP, creatine kinase, and creatine phosphate were from Merck. Except for lauric acid (Henkel) and 1,18-Octadecanedicarboxylic acid (ThermoFisher Scientific), lipids were from Merck. 10 mM stock solutions were prepared in analytical grade DMSO and kept under nitrogen. For assays the stock solution was appropriately diluted in 20 mM Mops buffer, pH 7.5, suitable to be added to the assays resulting in the desired final concentrations. The final DMSO concentrations in *in vitro* and *in vivo* assays were maximally 1%, a concentration without any biochemical effect as checked in respective control incubations. The constitutively active GsaQ227L mutant protein was expressed and purified as described earlier (47–49).

General Experimental Procedures

For HPLC analysis, a Waters HPLC system (1525 pump, 2996 photodiode array detector, 7725i injector, 200 series PerkinElmer vacuum degasser) was used. Solvents were HPLC or LC-MS grade from Merck-Sigma. One-dimensional ^1H and ^{13}C NMR spectra were recorded on a 400 MHz Bruker AVANCE III NMR spectrometer equipped with a 5 mm broadband SmartProbe and AVANCE III HD Nanobay console. Spectra were recorded in methanol- d_4 and calibrated to the residual solvent signal (δ_{H} 3.31 and δ_{C} 49.15 ppm).

Lung tissue extraction and fractionation

1.24 kg bovine lung was minced in a meat grinder, then mixed and homogenized with 1.2 L 50mM MOPS, pH 7.5, in a Waring blender (4°C) resulting in 2.3 L homogenate. It was centrifuged (30 min, 4°C, 7200×g) resulting in 1.2 L supernatant. The pH was adjusted to 1 using 7% HCl. Equal volumes of $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (2:1) were mixed with the supernatant in a separatory funnel and shaken vigorously. Centrifugation was at 5300×g for 30 min. The lower organic CH_2Cl_2 layer was recovered and the solvent was evaporated yielding 2 g of dried material. It was dissolved in 100 ml petroleum ether and subjected to normal-phase silica gel vacuum liquid chromatography (60 H Supelco). The column was eluted stepwise with solvents of increasing polarity from 90:10 petroleum ether/EtOAc to 100% EtOAc, followed by 100% MeOH. 17 fractions (A-Q) of 300 mL were collected and dried. Fraction E (eluting at 40:60 petroleum ether/EtOAc) was analysed by RP-HPLC using a linear MeOH/ H_2O gradient from 80:20 to 100:0 (0.1% TFA) for 15 min, followed by 100:0 for 30 min (Knauer Eurosphere II C18P 100-5, 250 x 8 mm, 1.2 mL/min flow rate, UV-absorbance monitored at 210 nm) to yield five subfractions; E1-E5. Fraction E2 was analysed by ^1H - and ^{13}C -NMR which indicated the presence of aliphatic lipids and fatty acids (Fig. S2).

GC-MS analysis

Fraction E2 was analysed by GC-MS. Acids were acid trimethylsilylated using *N,O*-bis(trimethylsilyl)trifluoroacetamide + trimethylchlorosilane (BSTFA + TMCS, 99:1 v/v). The mixture was heated for 2 h at 90°C. After cooling and clearing the sample was transferred into a GC vial in 200 μL hexane.

An Agilent Technologies GC system (8890 gas chromatograph and 5977B mass spectrometer equipped with a DB-HP5MS UI column, 30 m x 0.25 mm, film thickness of 0.25 μm) was used. Injection volume was 1 μL . The temperature was kept at 100 C for 5 min, and then increased at 53°C/min to 240 C. The rate was decreased to 3 C/min to reach 305 C. Carrier gas was He_2 (99.9%; 1.2 mL/min). Ionization was with 70 eV and MS spectra were recorded for a mass range m/z 35-800 for 35 min. Compounds were identified by comparing the spectra with those in the NIST library. Individual compound content is given as a relative % of the total peak area.

Plasmid construction and protein expression

hAC sequences were from NCBI were: ADCY1: NM_021116.3; ADCY2: NM_020546.2; ADCY3: NM_004036.4; ADCY4: NM_001198568.2; ADCY5: NM_183357.2; ADCY6: NM_015270.4; ADCY7: NM_001114.4; ADCY8: NM_001115.2; ADCY9: NM_001116.3.

Human mAC genes were from GenScript and fitted with a C-terminal FLAG-tag. The chimera mAC5(TM)_mAC3(cat) had an N-terminal connectase-tag, MPGAFDADPLVVEIAAAGA, followed by AC5(1–402)_AC3(250–631)_AC5(761–1009)_AC3(862–1144). The gene was synthesized by GenScript. HEK293 cells were maintained in DMEM with 10% FBS at 37°C and 5% CO_2 . Transfection with AC plasmids was with PolyJet (SignaGen, Frederick, MD, USA). Permanent cell lines were generated by selection for 7 days with 600 $\mu\text{g}/\text{mL}$ G418 and maintained with 300 $\mu\text{g}/\text{mL}$. For membrane preparation, cells were tyrosinized, collected by centrifugation (3000×g, 5 min) and lysed and homogenized in 20 mM HEPES, pH 7.5, 1 mM EDTA, 2 mM MgCl_2 , 1 mM DTT, one tablet of

cOmplete, EDTA-free (per 50 mL) and 250 mM sucrose by 20 strokes in a potter homogenizer on ice. Debris was removed (5 min at 1000xg, 0°C), membranes were collected at 100,000xg, 60 min at 0°C, suspended and stored at -80°C in 20 mM MOPS, pH 7.5, 0.5 mM EDTA, 2 mM MgCl₂. Membrane preparation from mouse brain cortex was according to (19, 50). Three cerebral cortices were dissected and homogenized in 4.5 ml cold 48 mM Tris- HCl, pH 7.4, 12 mM MgCl₂, and 0.1 mM EGTA with a Polytron hand disperser (Kinematica AG, Switzerland). The homogenate was centrifuged for 15 min at 12000xg at 4°C. The pellet was washed once with 5 mL 1 mM KHCO₃. The final suspension in 2 mL 1 mM KHCO₃ was stored in aliquots at -80 C.

DNA extraction

DNA from 1x10⁶ cells of permanently transfected and non-transfected HEK293 cells was extracted using the High Pure PCR Template Preparation Kit (Roche) according to the manufacturer's instructions. DNA concentrations were determined at 260 nm using a sub- microliter cell (IMPLEN) in a P330 NanoPhotometer (IMPLEN). Elution buffer (Roche) was used for blanks.

Polymerase chain reaction

100 ng of template DNA was mixed with 0.5 µM Forward primer and 0.5 µM Reverse primer. 12.5 µL 2X KAPA2G Fast (HotStart) Genotyping Mix with dye and water was added, total reaction volume 25 µL according to the KAPA2G Fast HotStart Genotyping Mix kit (Roche) protocol. PCR followed the cycling protocol in a Biometra T3000 thermocycler:

Step	Temperature	Duration	Cycles
Initial denaturation	95 C	3 min	1
Denaturation	95 C	15 sec	35
Annealing	60 C	15 sec	
Extension	72 C	30-60 sec*	
Final extension	72 C	2-4 min*	1

*Extension and final extension times were adjusted to the expected amplicon length.

The PCR products were directly loaded on a 1.5% agarose gel. A 1 kb DNA ladder (New England Bio Labs #N3232S) was mixed with Gel Loading Dye Purple 6X (New England Biolabs #B7024S) and water. After running the gel for 15-20 minutes at 90 V in 1X TAE buffer, the gel was stained in an Ethidium bromide bath and left running for another 10-20 minutes. The gels were then evaluated under UV light in a UVP GelStudio PLUS (Analytik Jena) gel imager.

AC isoform	Forward primer (5'-3')	Reverse primer (5'-3')
1	GTCAACAGGTACATCAGCCGCC	AGCCTCCTTCCCAGCTGCTGC
2	AGGAGACTGCTACTACTGTGTATCTGGAC	GGATGCCACGTTGCTCTGGGA
3	TTCATCCTGGTGATGGCAAATGTCGT	GGAGTTGTCCACCACCTGGTG
4	CGGGGATGCCAAGTTCTTCCAGGTCATTG	GCCTAGGGTAGCTGAAGGAGG
5	CCTCATCCTGCGCTGCACCCAGAAGCG	ACTGAGC
6	TCCTGAGCCGTGCCATCGA	ACTGCTGGGGCCCCCATTGAG
7	TCCTCGGCGACTGCTACTACTG	GTTCAGCCCCAGCCCCTGAAA
8	ACTTGCGGAGTGGCGATAAATTGAGA	TGGCAAATCAGATTTGTCTGGTGCC
9	CGCTGTGCTTCCTCCTGGTG	CACACTCTTTGAAACGTTGAGC

Adenylyl cyclase assay

In a volume of 10 μ L, AC activities were measured using 1 mM ATP, 2 mM $MgCl_2$, 3 mM creatine phosphate, 60 μ g/mL creatine kinase, and 50 mM MOPS pH 7.5. The cAMP assay kit from Cisbio (Codolet, France) was used for detection according to the supplier's instructions. For each assay a cAMP standard curve was established. EC_{50} and IC_{50} values were calculated by GraphPad Prism version 8.4.3 for Windows, GraphPad Software, San Diego, California USA, www.graphpad.com.

cAMP accumulation assay

HEK293 cells stably expressing mAC isoforms 3, 5, and mAC5(membr)_mAC3(cat) were plated at 2500-10000 cells/well into 384 well plates. Cells were treated with varying concentrations of lipids and incubated for 10 min at 37°C and 5% CO_2 . 2.5-10 μ M isoproterenol was added to stimulate cAMP formation and cells were incubated for 5 min. HEK293-AC5-3 was assayed in the presence of the phosphodiesterase inhibitor 0.5 mM isobutyl-methyl-xanthine. Addition of Cisbio HTRF detection reagents stopped the reaction and cAMP levels were determined.

Data handling and analysis

Experimental results were evaluated using GraphPad Prism version 8.4.3. Assays were conducted with a minimum of two technical replicates from at least two independent assays, as specified in the figure legends. Average values from duplicate or triplicate experiments were designated as single points and data are expressed as means $\pm$ S.E.M. Graphs were generated by GraphPad and assembled in PowerPoint.

Acknowledgements

We are indebted to Prof. Andrei Lupas, Max-Planck-Institute of Biology, Tübingen, for support, advice and critique. We thank N. Grzegorzek, Organic Chemistry, for GC/MS measurements. Funding was from the Deutsche Forschungsgemeinschaft (Schu275/45-1) and from institutional funds from the Max-Planck-Gesellschaft;

Author contributions

Experimental realization: ML, SE and AS. HPLC, GC/MS, NMR, SE and HG; Connectase assay, AF; Concept, data evaluation and writing JES.

Conflict of interest

The authors declare no conflict of interest with the contents of this article. Data and Materials availability: All data are available in the paper and the two appendices.

Appendix 1: chemical ligand identification


Acidic lipid extract from bovine lung																	
Si-VLC																	
Fraction	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q
Solvent system	100% PE	90:10 PE:EA	80:20 PE:EA	60:40 PE:EA	40:60 PE:EA	20:80 PE:EA	100% EA	25:75 MeOH:EA	100% MeOH	100% MeOH	100% MeOH	100% MeOH	100% MeOH	100% MeOH	100% MeOH	100% MeOH	100% MeOH
weight (mg)	3.1	4.5	459.1	404	285.8	70.1	29.5	21.6	344.6	753.3	95.7	85.3	175.5	136.3	54.7	27.4	17.9
% AC5 activity	116.5	125	100.5	80	73	48.5	84	87.5	102.5	44.5	11	1	30	50	44.5	38	25
% AC3 activity					376	372											

40 mg RP18-HPLC					
Fraction	E-1	E-2	E-3	E-4	E-5
weight (mg)	7.5	5.4	6.8	3.9	0.5
% AC5 activity	129	96	142	157	162
% AC3 activity	389	411	153	208	257

Fractionation Scheme


Si-VLC: Silica-Vacuum Liquid Chromatography, RP-HPLC: Reversed-Phase-High-Performance Liquid Chromatography, EA: Ethyl acetate, PE: petroleum ether, MeOH: Methanol. % AC3 and AC5 activities compared to 300 nM Gsα stimulation (= 100 %; values are means of two experiments carried out in triplicates. Fractions were tested at 1 μg/10 μL assay. Basal AC5 and 300 nM Gsα activities were 0.05 nmol and 2.1 and for mAC3 0.02 and 0.15 nmol cAMP•mg⁻¹•min⁻¹, respectively.

Appendix 2




Appendix 1 - Fig. 1.

RP-HPLC chromatogram of fraction E. UV-absorbance at 210 nm.



Appendix 1 - Fig. 2.


NMR spectra of Fraction E2 in d_4 -MeOH. (Top panel) ^1H -NMR spectrum. (Bottom Panel) ^{13}C -NMR spectrum.



Appendix 2 - Fig. 1.

Time-dependent stimulation of mAC3 by oleic acid.


mAC3 was incubated with 300 nM Gsα ± 20 μM oleic acid at 37°C for the time depicted. Data represent the mean of two independent experiments performed in duplicates. Correlation coefficients were 0.9733 and 0.9864 minus and plus added oleic acid, respectively.



Appendix 2 - Fig. 2.

Hanes-Woolf plot of mAC3 $\pm$ 20 μM oleic acid.

The assay at 37°C, 15 min. K_m of ATP was 335 and 221 μM $\pm$ oleic acid, respectively (not significant). V_{max} $\pm$ oleic acid was 0.62 and 1.23 $\text{nmol cAMP} \cdot \text{mg}^{-1} \cdot \text{min}^{-1}$, respectively. Lineweaver-Burk plots and Eddie- Hofstee plots yielded identical data.



Appendix 2 - Fig. 3.


Oleic acid has no stimulatory effect on the soluble catalytic dimer. **Basal and 300 nM** Gsa activities of the mAC1-C1/mAC2-C2 were 0.02 ± 0.003 and 0.08 ± 0.02 nmol



Appendix 2 - Fig. 4.

Structures of the tested aliphatic lipids listed in


Table 1 [↗](#).



Appendix 2 - Fig. 5.

Effect of isoproterenol on cAMP formation in HEK293 cells transfected with mAC3.


Basal activity was 0.04 ± 0.01 pmol cAMP/10,000. Where applicable, error bars indicate S.E.M. (experiments carried out in triplicates; $n=1-3$). Calculated $\text{EC}_{50} = 1.4 \mu\text{M}$.



Appendix 2 - Fig. 6.


Agarose gels of PCR products from HEK293 cells permanently transfected with mAC1-9.

Expected amplicon lengths were 1667, 2266, 1296, 1642, 1864, 1624, 2270, 1730, and 3614 bp for mAC isoforms 1-9, respectively. As controls, the primers pairs for each isoform were tested with DNA isolated from all other eight cell lines, resulting in no bands (not shown). Further, the untransfected HEK293 cells were tested with the primers specific for each isoform, resulting in no bands (not shown; the primer pairs are listed in the experimental section above).




Appendix 2 - Fig. 7.

Effect of lipids on 300 nM Gsa stimulated mAC2. (Left) Effect of 20 μM lipids on mAC2. **(Right)** Concentration-response curve of *cis*-vaccenic acid. Basal and Gsa activities min, respectively. EC₅₀ of *cis*-vaccenic acid was 10.6 μM. Error bars denote SEM of n=2-7. One-sample *t* test: **P* < 0.05; ****P* < 0.001 compared to 100% (300 nM Gsa stimulation).




Appendix 2 - Fig. 8.

Effect of lipids on 300 nM Gsα stimulated mAC7. (Left) Effect of 20 μM lipids on mAC7. **(Right)** Concentration-response curve for elaidic acid. Basal and Gsα activities were 0.01 ± 0.003 and 0.06 ± 0.01 nmol cAMP•mg min, respectively. EC_{50} was 9.7 μM. Error bars are SEM of n=2-7. One-sample *t* test: **P* < 0.05; ***P* < 0.01 compared to 100% (300 nM Gsα stimulation).




Appendix 2 - Fig. 9.

Effect of lipids on 300 nM Gsa stimulated mAC9. (Left) Effect of 20 μ M lipids on mAC9. **(Right)** Concentration-response curves of elaidic and linoleic acids. EC_{50} for elaidic acid was 5 μ M. Basal and Gsa activities were 0.07 ± 0.005 and 0.95 ± 0.06 nmol cAMP $\cdot$ mg $^{-1}$ min $^{-1}$, respectively. Error bars denote SEM of n=2-15. One-sample *t* test: **P* < 0.05; ***P* < 0.01; ****P* < 0.001; *****P* < 0.0001 compared to 100% (300 nM Gsa stimulation).




Appendix 2 - Fig. 10.

Concentration-response curves of fatty acids activating Gsα stimulated mAC9. Basal and Gsα activities were 0.06 ± 0.005 and 0.92 ± 0.07 nmol cAMP•mgmin, respectively. EC₅₀ for lauric, *cis*-vaccenic, palmitic and arachidonic acids were 13.7, 8.5, 8.6, and 7.5 μM, respectively. Error bars denote SEM of n=3-9. Significances were removed for clarity.




Appendix 2 - Fig. 11.

Effect of 20 μ M lipids on 300 nM G α stimulated mAC1. Basal and G α activities were 0.18 ± 0.02 and 0.46 ± 0.02 nmol cAMP $\cdot$ mg $^{-1}\cdot$ min $^{-1}$, respectively. Error bars denote SEM of n=2-9. One-sample *t* test: **P* < 0.05; ****P* < 0.001 compared to 100% (300 nM G α stimulation).




Appendix 2 - Fig. 12.

Effect of 20 μ M lipids on 300 nM G α stimulated mAC4. Basal and G α activities were 0.03 ± 0.003 and 0.59 ± 0.11 nmol cAMP $\cdot$ mg $^{-1}\cdot$ min $^{-1}$, respectively. Error bars denote SEM of n=2-6. One-sample *t* test: ***P* < 0.01; ****P* < 0.001 compared to 100% (300 nM G α stimulation).




Appendix 2 - Fig. 13.

Palmitoleic acid inhibits mACs 1 and 4 stimulated by 300 nM Gsα. Basal and Gsα stimulated activities of mAC1 were 0.14 ± 0.01 and 0.44 ± 0.02 and of mAC4 were 0.03 ± 0.002 and 0.25 ± 0.02 nmol cAMP•mg⁻¹•min⁻¹. IC₅₀ for mAC1 and 4 were 49 and 20 μM, respectively. Error bars denote SEM of n= 3-6. One-sample *t* test: **P* < 0.05; ***P* < 0.01 ****P* < 0.001; compared to 100% (300 nM Gsα stimulation).




Appendix 2 - Fig. 14.

Effect of 20 μ M lipids on 300 nM G α stimulated mAC5. Basal and G α activities were 0.07 ± 0.01 and 0.46 ± 0.04 nmol cAMP•mg SEM of $n=2-5$, min⁻¹, respectively. Error bars denote



Appendix 2 - Fig. 15.

Effect of 20 μ M lipids on 300 nM G α stimulated mAC6. Basal and G α activities were 0.07 ± 0.01 and 0.50 ± 0.06 nmol cAMP·mg SEM of $n=2-6$, min⁻¹, respectively. Error bars denote



Appendix 2 - Fig. 16.

Effect of 20 μ M lipids on 300 nM Gsa stimulated mAC8. Basal and Gsa activities were 0.19 ± 0.01 and 1.04 ± 0.19 nmol cAMP•mg SEM of $n=2-5 \cdot \text{min}^{-1}$, respectively. Error bars denote

	N	Basal activity	300 nM Gsα activity
AC1	5	0.12 ± 0.02	0.39 ± 0.05
AC2	7	0.22 ± 0.05	1.41 ± 0.21
AC3	23	0.02 ± 0.002	0.17 ± 0.03
AC4	4	0.02 ± 0.007	0.19 ± 0.02
AC5	7	0.05 ± 0.01	0.50 ± 0.11
AC6	3	0.06 ± 0.02	0.29 ± 0.05
AC7	7	0.01 ± 0.001	0.05 ± 0.002
AC8	2	0.08 ± 0.02	0.31 ± 0.06
AC9	15	0.09 ± 0.02	0.97 ± 0.15

Appendix 2 - Table 1.

Basal and Gsα-stimulated activities of mAC isoforms

. Activities are listed as mean ± SEM in nmol cAMP • mg⁻¹ min⁻¹ .

References

1. Sutherland E. W., Rall T. W. (1958) **Fractionation and characterization of a cyclic adenine ribonucleotide formed by tissue particles** *J Biol Chem* **232**:1077–1091
2. Dessauer C. W., et al. (2017) **International Union of Basic and Clinical Pharmacology CI. Structures and Small Molecule Modulators of Mammalian Adenylyl Cyclases.** *Pharmacol Rev* **69**:93–139
3. Ostrom K. F., et al. (2022) **Physiological roles of mammalian transmembrane adenylyl cyclase isoforms** *Physiol Rev* **102**:815–857
4. Khandelwal R. L., Hamilton I. R. (1971) **Purification and properties of adenylyl cyclase from *Streptococcus salivarius*** *J Biol Chem* **246**:3297–3304
5. Linder J. U., Schultz J. E. (2003) **The class III adenylyl cyclases: multi-purpose signalling modules** *Cell Signal* **15**:1081–1089
6. Linder J. U., Schultz J. E. (2008) **Versatility of signal transduction encoded in dimeric adenylyl cyclases** *Current opinion in structural biology* **18**:667–672
7. Krupinski J., et al. (1989) **Adenylyl cyclase amino acid sequence: possible channel- or transporter-like structure** *Science* **244**:1558–1564
8. Bassler J., Schultz J. E., Lupas A. N. (2018) **Adenylate cyclases: Receivers, transducers, and generators of signals** *Cell Signaling* **46**:135–144
9. Guo Y. L., Seebacher T., Kurz U., Linder J. U., Schultz J. E. (2001) **Adenylyl cyclase Rv1625c of *Mycobacterium tuberculosis*: a progenitor of mammalian adenylyl cyclases** *EMBO J* **20**:3667–3675
10. Tesmer J. J., Sunahara R. K., Gilman A. G., Sprang S. R. (1997) **Crystal structure of the catalytic domains of adenylyl cyclase in a complex with G α .GTP γ S** *Science* **278**:1907–1916
11. Sinha S. C., Sprang S. R. (2006) **Structures, mechanism, regulation and evolution of class III nucleotidyl cyclases** *Rev Physiol Biochem Pharmacol* **157**:105–140
12. Schultz J. E. (2022) **The evolutionary conservation of eukaryotic membrane-bound adenylyl cyclase isoforms** *Front Pharmacol* **13**
13. Tang W. J., Gilman A. G. (1995) **Construction of a soluble adenylyl cyclase activated by G α and forskolin** *Science* **268**:1769–1772
14. Schuster D., Khanppnavar B., Kantarci I., Mehta V., Korkhov V. M. (2024) **Structural insights into membrane adenylyl cyclases, initiators of cAMP signaling** *Trends Biochem Sci* **49**:156–168
15. Ziegler M., et al. (2017) **A novel signal transducer element intrinsic to class IIIa and IIIb adenylate cyclases** *FEBS J* **284**:1204–1217

16. Qi C., Sorrentino S., Medalia O., Korkhov V. M. (2019) **The structure of a membrane adenylyl cyclase bound to an activated stimulatory G protein** *Science* **364**:389–394
17. Qi C., et al. (2022) **Structural basis of adenylyl cyclase 9 activation** *Nature communications* **13**
18. Gu C., Sorkin A., Cooper D. M. F. (2001) **Persistent interactions between the two transmembrane clusters dictate the targeting and functional assembly of adenylyl cyclase** *Current biology : CB* **11**:185–190
19. Seth A., Finkbeiner M., Grischin J., Schultz J. E. (2020) **G α stimulation of mammalian adenylate cyclases regulated by their hexahelical membrane anchors** *Cell Signal* **68**
20. Abdel Motaal A., Tews I., Schultz J. E., Linder J. U. (2006) **Fatty acid regulation of adenylyl cyclase Rv2212 from Mycobacterium tuberculosis H37Rv** *Febs J* **273**:4219–4228
21. Findeisen F., et al. (2007) **The structure of the regulatory domain of the adenylyl cyclase Rv1264 from Mycobacterium tuberculosis with bound oleic acid** *J Mol Biol* **369**:1282–1295
22. Beltz S., Bassler J., Schultz J. E. (2016) **Regulation by the quorum sensor from Vibrio indicates a receptor function for the membrane anchors of adenylate cyclases** *Elife* **5**
23. Anggard E., Samuelsson B. (1965) **Biosynthesis of prostaglandins from arachidonic acid in guinea pig lung. Prostaglandins and related factors.** **38** *J Biol Chem* **240**
24. Dautel S. E., et al. (2017) **Lipidomics reveals dramatic lipid compositional changes in the maturing postnatal lung** *Sci Rep* **7**
25. Kimura I., Ichimura A., Ohue-Kitano R., Igarashi M. (2020) **Free Fatty Acid Receptors in Health and Disease** *Physiol Rev* **100**:171–210
26. Samovski D., Jacome-Sosa M., Abumrad N. A. (2023) **Fatty Acid Transport and Signaling: Mechanisms and Physiological Implications** *Annu Rev Physiol* **85**:317–337
27. Grundmann M., Bender E., Schamberger J., Eitner F. (2021) **Pharmacology of Free Fatty Acid Receptors and Their Allosteric Modulators** *Int J Mol Sci* **22**
28. Nakamura J., Okamura N., Usuki S., Bannai S. (2001) **Inhibition of adenylyl cyclase activity in brain membrane fractions by arachidonic acid and related unsaturated fatty acids** *Arch Biochem Biophys* **389**:68–76
29. Alam S. Q., Mannino S. J., Alam B. S., McDonough K. (1995) **Effect of essential fatty acid deficiency on forskolin binding sites, adenylate cyclase and cyclic AMP-dependent protein kinase activity, the levels of G proteins and ventricular function in rat heart** *J Mol Cell Cardiol* **27**:1593–1604
30. Mock E. D., Gagestein B., van der Stelt M. (2023) **Anandamide and other N-acylethanolamines: A class of signaling lipids with therapeutic opportunities** *Prog Lipid Res* **89**
31. Lu H. C., Mackie K. (2016) **An Introduction to the Endogenous Cannabinoid System** *Biological psychiatry* **79**:516–525

32. Simon G. M., Cravatt B. F. (2008) **Anandamide biosynthesis catalyzed by the phosphodiesterase GDE1 and detection of glycerophospho-N-acyl ethanolamine precursors in mouse brain** *J Biol Chem* **283**:9341–9349
33. Liu J., et al. (2006) **A biosynthetic pathway for anandamide** *Proceedings of the National Academy of Sciences of the United States of America* **103**:13345–13350
34. McKinney M. K., Cravatt B. F. (2005) **Structure and function of fatty acid amide hydrolase** *Annu Rev Biochem* **74**:411–432
35. Fuchs A. C. D. (2023) **Specific, sensitive and quantitative protein detection by in-gel fluorescence** *Nature communications* **14**
36. Tews, et al. (2005) **The structure of a pH-sensing mycobacterial adenylyl cyclase holoenzyme** *Science* **308**:1020–1023
37. Sanabra C., Mengod G. (2011) **Neuroanatomical distribution and neurochemical characterization of cells expressing adenylyl cyclase isoforms in mouse and rat brain** *J Chem Neuroanat* **41**:43–54
38. Sunahara R. K., Tesmer J. J., Gilman A. G., Sprang S. R. (1997) **Crystal structure of the adenylyl cyclase activator Gsalpha** *Science* **278**:1943–1947
39. Tesmer J. J., Sprang S. R. (1998) **The structure, catalytic mechanism and regulation of adenylyl cyclase** *Current opinion in structural biology* **8**:713–719
40. Vercellino I., et al. (2017) **Role of the nucleotidyl cyclase helical domain in catalytically active dimer formation** *Proceedings of the National Academy of Sciences of the United States of America* **114**:E9821–E9828
41. Eyster K. M. (2007) **The membrane and lipids as integral participants in signal transduction: lipid signal transduction for the non-lipid biochemist** *Adv Physiol Educ* **31**:5–16
42. Ulven T., Christiansen E. (2015) **Dietary Fatty Acids and Their Potential for Controlling Metabolic Diseases Through Activation of FFA4/GPR120** *Annu Rev Nutr* **35**:239–263
43. Huber A. H., Kleinfeld A. M. (2017) **Unbound free fatty acid profiles in human plasma and the unexpected absence of unbound palmitoleate** *J Lipid Res* **58**:578–585
44. McMahon H. T., Gallop J. L. (2005) **Membrane curvature and mechanisms of dynamic cell membrane remodelling** *Nature* **438**:590–596
45. Brown W. J., Chambers K., Doody A. (2003) **Phospholipase A2 (PLA2) enzymes in membrane trafficking: mediators of membrane shape and function** *Traffic* **4**:214–221
46. Muccioli G. G. (2010) **Endocannabinoid biosynthesis and inactivation, from simple to complex** *Drug Discov Today* **15**:474–483
47. Graziano M. P., Freissmuth M., Gilman A. G. (1989) **Expression of Gs alpha in Escherichia coli. Purification and properties of two forms of the protein** *J Biol Chem* **264**:409–418
48. Graziano M. P., Freissmuth M., Gilman A. G. (1991) **Purification of recombinant Gs alpha** *Methods Enzymol* **195**:192–202

49. Sunahara R. K., Dessauer C. W., Whisnant R. E., Kleuss C., Gilman A. G. (1997) **Interaction of Gsalpha with the cytosolic domains of mammalian adenylyl cyclase** *J Biol Chem* **272**:22265–22271
50. Schultz J. E., Schmidt B. H. (1987) **Treatment of rats with thyrotropin (TSH) reduces the adrenoceptor sensitivity of adenylate cyclase from cerebral cortex** *Neurochem Int* **10**:173–178

Editors

Reviewing Editor

Volker Dötsch

Goethe University Frankfurt, Frankfurt am Main, Germany

Senior Editor

Volker Dötsch

Goethe University Frankfurt, Frankfurt am Main, Germany

Reviewer #1 (Public review):

Summary:

The authors show that the Gas-stimulated activity of human membrane adenylyl cyclases (mAC) can be enhanced or inhibited by certain unsaturated fatty acids (FA) in an isoform-specific fashion. Thus, with IC50s in the 10-20 micromolar range, oleic acid affects 3-fold stimulation of membrane-preparations of mAC isoform 3 (mAC3) but it does not act on mAC5. Enhanced Gas-stimulated activities of isoforms 2, 7, and 9, while mAC1 was slightly attenuated, but isoforms 4, 5, 6, and 8 were unaffected. Certain other unsaturated octadecanoic FAs act similarly. FA effects were not observed in AC catalytic domain constructs in which TM domains are not present. Oleic acid also enhances the AC activity of isoproterenol-stimulated HEK293 cells stably transfected with mAC3, although with lower efficacy but much higher potency. Gas-stimulated mAC1 and 4 cyclase activity were significantly attenuated in the 20-40 micromolar by arachidonic acid, with similar effects in transfected HEK cells, again with higher potency but lower efficacy. While activity mAC5 was not affected by unsaturated FAs, neutral anandamide attenuated Gas-stimulation of mAC5 and 6 by about 50%. In HEK cells, inhibition by anandamide is low in potency and efficacy. To demonstrate isoform specificity, the authors were able to show that membrane preparations of a domain-swapped AC bearing the catalytic domains of mAC3 and the TM regions of mAC5 are unaffected by oleic acid but inhibited by anandamide. To verify in vivo activity, in mouse brain cortical membranes 20 µM oleic acid enhanced Gas-stimulated cAMP formation 1.5-fold with an EC50 in the low micromolar range.

Strengths:

- (1) A convincing demonstration that certain unsaturated FAs are capable of regulating membrane adenylyl cyclases in an isoform-specific manner, and the demonstration that these act at the AC transmembrane domains.
- (2) Confirmation of activity in HEK293 cell models and towards endogenous AC activity in mouse cortical membranes.
- (3) Opens up a new direction of research to investigate the physiological significance of FA regulation of mACs and investigate their mechanisms as tonic or regulated enhancers or inhibitors of catalytic activity.

(4) Suggests a novel scheme for the classification of mAC isoforms.

Comments on revised version:

The issues I raised have largely been addressed. A minor concern relates to the legend for Figure 2C, where, according to the author's rebuttal, the vertical axis is "The ratio would be (Gsa + oleic acid stimulation) / (Gsa stimulation)" Otherwise, my general evaluation of the importance of the manuscript stands as stated in my initial review, namely, that the manuscript presents data and results that add a new dimension to existing paradigms for AC regulation, and will prompt future research into the role of physiological lipids in isoform-specific activation or inhibition of AC in tissues.

<https://doi.org/10.7554/eLife.101483.2.sa2>

Reviewer #3 (Public review):

Summary:

Landau et al. have submitted a manuscript describing for the first time that mammalian adenylyl cyclases can serve as membrane receptors. They have also identified the respective endogenous ligands which act via AC membrane linkers to modify and control Gs-stimulated AC activity either towards enhancement or inhibition of ACs which is family and ligand-specific. Overall, they have used classical assays such as adenylyl cyclase and cAMP accumulation assays combined with molecular cloning and mutagenesis to provide exceptionally strong biochemical evidence for the mechanism of the involved pathway regulation.

Strengths:

The authors have gone the whole long classical way from having a hypothesis that ACs could be receptors to a series of MS studies aimed at ligand identification, to functional studies of how these candidate substances affect the activity of various AC families in intact cells. They have used a large array of techniques with a paper having clear conceptual story and several strong lines of evidence.

Comments on revised version:

In general, the authors have addressed my comments satisfactorily apart from the suggestion to use a lower ISO concentration in their assay or at least to discuss this issue, cite relevant literature etc. Pending this small amendment I would be fine to proceed.

<https://doi.org/10.7554/eLife.101483.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public review):

Summary:

The authors show that the Gas-stimulated activity of human membrane adenylyl cyclases (mAC) can be enhanced or inhibited by certain unsaturated fatty acids (FA) in an isoform-

specific fashion. Thus, with IC₅₀s in the 10-20 micromolar range, oleic acid affects 3-fold stimulation of membrane-preparations of mAC isoform 3 (mAC3) but it does not act on mAC5. Enhanced Gas-stimulated activities of isoforms 2, 7, and 9, while mAC1 was slightly attenuated, but isoforms 4, 5, 6, and 8 were unaffected. Certain other unsaturated octadecanoic FAs act similarly. FA effects were not observed in AC catalytic domain constructs in which TM domains are not present. Oleic acid also enhances the AC activity of isoproterenol-stimulated HEK293 cells stably transfected with mAC3, although with lower efficacy but much higher potency. Gas-stimulated mAC1 and 4 cyclase activity were significantly attenuated in the 20-40 micromolar by arachidonic acid, with similar effects in transfected HEK cells, again with higher potency but lower efficacy. While activity mAC5 was not affected by unsaturated FAs, neutral anandamide attenuated Gas-stimulation of mAC5 and 6 by about 50%. In HEK cells, inhibition by anandamide is low in potency and efficacy. To demonstrate isoform specificity, the authors were able to show that membrane preparations of a domain-swapped AC bearing the catalytic domains of mAC3 and the TM regions of mAC5 are unaffected by oleic acid but inhibited by anandamide. To verify *in vivo* activity, in mouse brain cortical membranes 20 μ M oleic acid enhanced Gas-stimulated cAMP formation 1.5-fold with an EC₅₀ in the low micromolar range.

Strengths:

- (1) A convincing demonstration that certain unsaturated FAs are capable of regulating membrane adenylyl cyclases in an isoform-specific manner, and the demonstration that these act at the AC transmembrane domains.
- (2) Confirmation of activity in HEK293 cell models and towards endogenous AC activity in mouse cortical membranes.
- (3) Opens up a new direction of research to investigate the physiological significance of FA regulation of mACs and investigate their mechanisms as tonic or regulated enhancers or inhibitors of catalytic activity.
- (4) Suggests a novel scheme for the classification of mAC isoforms.

Weaknesses:

- (1) Important methodological details regarding the treatment of mAC membrane preps with fatty acids are missing.

We will address this issue in more detail.

- (2) It is not evident that fatty acid regulators can be considered as "signaling molecules" since it is not clear (at least to this reviewer) how concentrations of free fatty acids in plasma or endocytic membranes are hormonally or otherwise regulated.

Although this question is not the subject of this ms., we will address this question in more detail in the discussion of the revision.

Reviewer #2 (Public review):

Summary:

The authors extend their earlier findings with bacterial adenylyl cyclases to mammalian enzymes. They show that certain aliphatic lipids activate adenylyl cyclases in the absence of stimulatory G proteins and that lipids can modulate activation by G proteins. Adding lipids to cells expressing specific isoforms of adenylyl cyclases could regulate cAMP production, suggesting that adenylyl cyclases could serve as 'receptors'.

Strengths:

This is the first report of lipids regulating mammalian adenylyl cyclases directly. The evidence is based on biochemical assays with purified proteins, or in cells expressing specific isoforms of adenylyl cyclases.

Weaknesses:

It is not clear if the concentrations of lipids used in assays are physiologically relevant. Nor is there evidence to show that the specific lipids that activate or inhibit adenylyl cyclases are present at the concentrations required in cell membranes. Nor is there any evidence to indicate that this method of regulation is seen in cells under relevant stimuli.

Although this question is not the subject of this ms., we will address this question in more detail in the discussion of the revision.

Reviewer #3 (Public review):

Summary:

Landau et al. have submitted a manuscript describing for the first time that mammalian adenylyl cyclases can serve as membrane receptors. They have also identified the respective endogenous ligands which act via AC membrane linkers to modify and control Gs-stimulated AC activity either towards enhancement or inhibition of ACs which is family and ligand-specific. Overall, they have used classical assays such as adenylyl cyclase and cAMP accumulation assays combined with molecular cloning and mutagenesis to provide exceptionally strong biochemical evidence for the mechanism of the involved pathway regulation.

Strengths:

The authors have gone the whole long classical way from having a hypothesis that ACs could be receptors to a series of MS studies aimed at ligand identification, to functional studies of how these candidate substances affect the activity of various AC families in intact cells. They have used a large array of techniques with a paper having clear conceptual story and several strong lines of evidence.

Weaknesses:

(1) At the beginning of the results section, the authors say "We have expected lipids as ligands". It is not quite clear why these could not have been other substances. It is because they were expected to bind in the lipophilic membrane anchors? Various lipophilic and hydrophilic ligands are known for GPCR which also have transmembrane domains. Maybe 1-2 additional sentences could be helpful here.

Will be done as suggested.

(2) In stably transfected HEK cells expressing mAC3 or mAC5, they have used only one dose of isoproterenol (2.5 μ M) for submaximal AC activation. The reference 28 provided here (PMID: 33208818) did not specifically look at Iso and endogenous beta2 adrenergic receptors expressed in HEK cells. As far as I remember from the old pharmacological literature, this concentration is indeed submaximal in receptor binding assays but regarding AC activity and cAMP generation (which happen after signal amplification with a so-called receptor reserve), lower Iso amounts would be submaximal. When we measure cAMP, these are rather 10 to 100 nM but no more than 1 μ M at which concentration response dependencies usually saturate. Have the authors tried lower Iso

concentrations to prestimulate intracellular cAMP formation? I am asking this because, with lower Iso prestimulation, the subsequent stimulatory effects of AC ligands could be even greater.

The best way to address this issue is to establish a concentration-response curve for Iso-stimulated cAMP formation using the permanently transfected cells. We note that in the past isoproterenol concentrations used in biochemical or electrophysiological experiments differed substantially.

(3) The authors refer to HEK cell models as "in vivo". I agree that these are intact cells and an important model to start with. It would be very nice to see the effects of the new ligands in other physiologically relevant types of cells, and how they modulate cAMP production under even more physiological conditions. Probably, this is a topic for follow-up studies.

The last sentence is correct.

Appraisal of whether the authors achieved their aims, and whether the results support their conclusions:

The authors have achieved their aims to a very high degree, their results do nicely support their conclusions. There is only one point (various classical GPCR concentrations, please see above) that would be beneficial to address.

Without any doubt, this is a groundbreaking study that will have profound implications in the field for the next years/decades. Since it is now clear that mammalian adenylyl cyclases are receptors for aliphatic fatty acids and anandamide, this will change our view on the whole signaling pathway and initiate many new studies looking at the biological function and pathophysiological implications of this mechanism. The manuscript is outstanding.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

It is not clear from the methods section how free FAs were applied to membrane preparations or HEK293 cells. Were FAs solubilized in organic solvents, or introduced as micelles?

The requested info is inserted into the M&M section

Could the authors comment on what is known about the concentration of oleic acid and other non-saturated fatty acids in plasma membranes relative to those required to produce allosteric effects on cyclase activity?

This info is now included in the last paragraph of the discussion.

It would be worthwhile to test the effect of FAs on basal (not Gas-stimulated) activity of mACs.

This has been carried with mAC isoforms 2, 3, 7, and 9 in which oleic acid enhances Gsa-stimulated activity. Due to the low levels of basal activities interpretable data were not obtained.

Do triglycerides esterified with oleic acid stimulate mAC3 and other sensitive isoforms?

Experiments were done with triolein and 2-oleoyl-glycerol (the answer is no). The data are presented in Fig. 3 and in the appendix Fig.'s 8, 9, 14; structural formulas in appendix 2 Fig. 4 were updated.

Does the quantity plotted on the vertical axis of Figure 1, right panel represent "Fractional Stimulation by Oleic acid" rather than simply "Fold Stimulation"? Clearly, as shown in the two left-most panels, Gas stimulates both mAC and mAC5. Rather it seems that the ratio (oleic acid stimulation) / (Gas stimulation) remains constant. This observation supports the statement in the discussion that "We suppose that in mAC3 the equilibrium of two differing ground states favors a Gas-unresponsive state and the effector oleic acid concentration-dependently shifts this equilibrium to a Gas-responsive state". It could also be said that the effect of oleic acid is additive, and in constant proportion to that of Gas.

This comment certainly is related to Fig. 2:

The ratio would be (Gsa + oleic acid stimulation) / (Gsa-stimulation), i.e., fractional stimulation by addition of oleic acid is identical to fold stimulation.

We have amended the legend to fig. 2C for clarification.

The last sentence is wrong because oleic acid alone does not stimulate.

It is stated on page 3, 2nd to last line that "The action of oleic acid on mAC3 was instantaneous...". Since the earliest time point is taken at 5 minutes, the claim that the action of the lipid is instantaneous cannot be made. Information about kinetics would be useful to have, since it is possible that the lipid must be released from a micelle and be incorporated into the AC membrane fraction before it is active.

The first point is 3 min.

We deleted the word "instantaneous" and added the correlation coefficients for both conditions in the legend to appendix 2; fig. 1 for clarification.

The data spread in Figure 4 and other figures showing similar data is significant, to the extent that the computed value for EC50 may not be of high precision. Authors should cite the correlation coefficient for the overall fit and uncertainty for the EC50 value (in addition to significances by t-test of individual data points).

This will not add valuable information. Pearsons correlation coefficients are only for linear relationships.

(cf. N.N. Kachouie, W. Deebani (2020) Association Factor for Identifying Linear and Nonlinear Correlations in Noisy Conditions. Entropy 22:440)

The "switch" between relatively low potency and high efficacy in membrane preps to high potency and low efficacy in cells is remarkable. Could this have a methodological basis or is it reflective of the mechanism by which FAs access mACs in membrane preps vs. cell membranes, or perhaps some biochemical transformation of the lipid in cells?

Honestly, we do not know.

The authors should note that there is some precedence for this work:

J Nakamura , N Okamura, S Usuki, S Bannai, Inhibition of adenylyl cyclase activity in brain membrane fractions by arachidonic acid and related unsaturated fatty acids. Arch Biochem Biophys. 2001 May 1;389(1):68-76. doi: 10.1006/abbi.2001.2315.

The effects of FA deficiencies on AC and related activities have been noted:

Alam SQ, Mannino SJ, Alam BS, McDonough K Effect of essential fatty acid deficiency on forskolin binding sites, adenylate cyclase, and cyclic AMP-dependent protein kinase activity, the levels of G proteins and ventricular function in rat heart. J Mol Cell Cardiol. 1995 Aug;27(8):1593-604. doi: 10.1016/s0022-2828(95)90491-3. PMID: 8523422

The latter publications are supportive of, and provide context to, the author's findings.

Both references are mentioned and cited.

Minor points:

The significance of the coloring scheme in Figure 5C bar graph should be stated in the legend.

Done.

In the introduction, it is stated that "The protein displayed two similar catalytic domains (C1 and C2) and two dissimilar hexahelical membrane anchors (TM1 and TM2)". In both cases, the respective domains can be said to be similar in overall fold, but - certainly in the case of the catalytic domains - different in amino acid sequence in functionally important regions of the domain.

Done: Changed wording.

The statement in the introduction that "The domain architecture, TM1-C1-TM2-C2, clearly indicated a pseudoheterodimeric protein composed of two concatenated bacterial precursor proteins" The authors refer to the fact that mammalian enzymes are pseudo heterodimers whereas bacterial type III cyclases are dimers of identical subunits.

Done.

Reviewer #2 (Recommendations for the authors):

The title need not state that a 'new class of receptors' has been identified. There is no direct evidence that the lipids bind to the enzymes, and the affinities can only be surmised from the EC50 graphs. To call a protein a receptor requires evidence to show that the binding is specific by showing that binding can be inhibited by a large excess of 'unlabelled' ligand. This could have been done by procuring labelled lipids for experimental verification.

As is well known, lipids easily bind to proteins. In this study no purified proteins were used. Therefore, binding assays most likely would result in unreliable data.

The paper would have benefitted from showing sequence alignments in the TM domains of the ACs discussed in the paper. Further, a phylogenetic tree of mammalian ACs would also reveal which enzymes from other species may be regulated similarly to those described in the paper. This would be important for researchers who use other model organisms to study cAMP signalling.

Such data are in multiple papers accessible in the literature. Where deemed appropriate we inserted references.

Figures 1A and 1B show data from only two experiments. A third experiment would have been useful in order to show the statistical significance of the data.

At this stage more experiments would not have affected further experimental plans.

Statements made in the text (for example, the last paragraph on page 6) state only the mean value and not the SDs. This would have been important to include even if the data is shown in the appendix. The same is true in the Legend of Figure 2. Why have the authors decided to use SEM and not SDs?

The reason is specified in M&M.

Concentrations of lipids used in biochemical assays are in the micromolar range. This suggests that we have moderate affinity binding, more in the range of an enzyme for a substrate rather than a receptor-ligand interaction.

We happen to disagree. Clearly, the differential activities, enhancing or attenuating Gsa-stimulated mAC activities is most plausibly explained by mAC receptor properties. mACs have enzyme activities using fatty acids as substrates.

The authors add lipids to cells and show changes in cAMP levels in their presence and absence. They also discuss how these extracellular lipids could be produced. Do you think this is necessary in vivo, though? Could the lipids present in membranes naturally act as regulators? Do specific lipid concentrations differ in different cell types, suggesting tissue-specific regulation of these mammalian Acs?

These are things that could be discussed in the manuscript.

The last paragraph of the discussion deals with these questions.

<https://doi.org/10.7554/eLife.101483.2.sa0>