

A scale-invariant log-normal droplet size distribution below the critical concentration for protein phase separation

Reviewed Preprint

v2 • August 6, 2024

Revised by authors

Reviewed Preprint

v1 • February 7, 2024

Tommaso Amico, Samuel Dada, Andrea Lazzari, Michaela Brezinova, Antonio Trovato, Michele Vendruscolo , Monika Fuxreiter , Amos Maritan 

Department of Physics and Astronomy, University of Padova, Italy • Centre for Misfolding Diseases, Department of Chemistry, University of Cambridge, UK • National Institute for Nuclear Physics (INFN), Padova Section, Padova, Italy • Department of Biomedical Sciences, University of Padova, Italy

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Many proteins have been recently shown to undergo a process of phase separation that leads to the formation of biomolecular condensates. Intriguingly, it has been observed that some of these proteins form dense droplets of sizeable dimensions already below the critical concentration, which is the concentration at which phase separation occurs. To understand this phenomenon, which is not readily compatible with classical nucleation theory, we investigated the properties of the droplet size distributions as a function of protein concentration. We found that these distributions can be described by a scale-invariant log-normal function with an average that increases progressively as the concentration approaches the critical concentration from below. The results of this scaling analysis suggest the existence of a universal behaviour independent of the sequences and structures of the proteins undergoing phase separation. While we refrain from proposing a theoretical model here, we suggest that any model of protein phase separation should predict the scaling exponents that we reported here from the fitting of experimental measurements of droplet size distributions. Furthermore, based on these observations, we show that it is possible to use the scale invariance to estimate the critical concentration for protein phase separation.

eLife assessment

In this **valuable** study, the authors analyze droplet size distributions of multiple protein condensates and their fit to a scaling ansatz, highlighting that they exhibit features of first- and second-order phase transitions. The experimental evidence is **solid**, but the interpretation can be improved. The text would benefit from further clarification and connection to the coupled percolation-and-phase-separation model.

<https://doi.org/10.7554/eLife.94214.2.sa3>

Introduction

Many proteins have been shown to undergo a phase separation process into a liquid-like condensed state^{1,5}. This process appears to be of physiological significance since it may lead to the formation of biomolecular condensates^{1,7}. As a consequence, it is closely controlled by the protein homeostasis system^{8,9}, and its dysregulation has been associated with a broad range of human diseases^{10,11}. It is therefore important to understand the fundamental nature of this process^{5,12,13} to provide insights for the identification of ways of modulating it through pharmacological interventions.

To help understand the nature of the transition underlying protein phase separation on the basis of recent experimental observations^{26,27}, here we study the distribution of the size of the droplets below the value of the concentration at which the transition occurs, which here is referred to as the critical concentration, ρ_c . This question appears as a promising starting point to develop new insights since it has been reported that proteins can form droplets of sizeable dimensions already well below the concentration at which phase separation occurs^{26,27}. This behaviour is not predicted by classical nucleation theory²⁸, and not readily consistent with the idea that the protein phase separation process can be described as a first-order phase transition. This is because, in a first-order phase transition, nucleation takes place in a supersaturated system²⁹, while in a subsaturated system particles can still self-assemble, but with a probability that decreases exponentially with the size of the assemblies.

In the present study, we asked whether the experimental data on the droplet size distributions obey scale invariance, a general characteristic of complex systems, which is often used to reveal universal patterns underlying self-assembly. We report the observation that the droplet size distributions of the proteins FUS and α -synuclein follow a scale-invariant log-normal behaviour. These findings are consistent with a universal behaviour resulting from the presence of an increasingly large correlation length, ξ , as the concentration approaches the critical concentration from below. The correlation length is an emergent characteristic, and it is related to the typical spatial range over which density fluctuations are correlated. When ξ is sufficiently large, one can expect scale invariance and finite-size scaling³⁰ to occur within a range of lengths spanning from the molecular size to ξ . This means that physical observables cease to depend explicitly on numerous microscopic details that are peculiar to spatial scales smaller than ξ , thus leading to a universal behaviour characterized by quantities obtained by coarse graining over scales smaller than ξ .

At a first-order phase transition, ξ is finite, and if it is not large enough, the length range discussed above remains too short to observe scale invariance. However, the vicinity of the spinodal line, where nucleation disappears as the dilute phase becomes unstable, to the coexistence curve, where nucleation appears as the dilute phase becomes metastable, might cause a large increase of ξ as the first-order phase transition is approached by increasing the concentration. If it is not preceded by a first-order phase transition, the spinodal line would correspond to a second-order phase transition, resulting in infinite ξ , with scale invariance holding on all length scales larger than the molecular scale. Various alternative scenarios contributing to the emergence of significant correlations will be mentioned in the Discussion section. Nevertheless, our scaling analysis exhibits a high degree of generality, devoid of reliance on specific underlying models.

As an application of the above observations, we address the question of whether scale invariance holds for droplet size distributions near the coexistence curve. As a practical consequence, we use this observation to propose a procedure to overcome the challenge of estimating the critical

concentration, ρ_c , which enters explicitly into our scaling analysis. Such challenge arises from the fact that close to the critical concentration, the timescale required for the equilibration of a system grows together with ξ , thus exceeding the timescale amenable to experimental observation.

Our analysis of experimental data indicates that: (i) scale invariance does indeed hold near the coexistence curve, and (ii) the droplet size distribution is log-normal. Based on the properties of the scale-invariant log-normal distribution of droplet sizes, we investigate a correlation between the moments of the distribution and the distance from ρ_c .

Finally, we note that methods to assess the critical concentration are crucial for understanding the location of proteins in their phase diagram, their proximity to the phase boundary between the native and droplet states, and how pharmacological interventions can modify their phase behaviour. To address this problem, we report how the moments of the distribution can be used as a scale-invariant gauge to estimate the critical concentration. In this way, an accurate estimate of the critical concentration is possible because it is based on measurements carried out away from the critical point, under conditions such that fluctuations are small, and hence experimental errors are smaller than in the proximity of the transition.

Results

Formulation of the scaling ansatz

Empirical evidence indicates that protein self-assembly into liquid-like condensates is characterized by: (i) a phase separation transition at a concentration ρ_c , (ii) a formation of droplets of sizeable dimensions already below ρ_c , and (iii) a droplet size distribution that, after an initial transient, does not change with the experimental observation time, although individual droplets can form, grow, shrink and dissolve. An initial analysis of droplet size distributions observed experimentally led us to ask whether the region near the transition could be described in terms of a scaling theory, as commonly done for critical phenomena (29), as summarized above. We also note that this approach is analogous to analyzing the cluster size distribution in a percolation problem³¹.

In our analysis, we called $P_>(s|\rho)$ the survival distribution function (SDF = 1-CDF, where CDF is the cumulative distribution function) corresponding to the probability to observe a droplet of size greater than s , when the concentration is ρ . The distance from the critical concentration ρ_c is measured in terms of the dimensionless variable $\tilde{\rho} = (\rho - \rho_c)/\rho_c$, which also allows to compare data from different experiments, as explained below. $P_>(s|\rho)$ in general depends separately on s , ρ and many other parameters characterizing the process, including temperature. However, if scale invariance holds in the vicinity of ρ_c , i.e. when $|\tilde{\rho}|$ is small and the correlation length of the system is large enough, we would expect $P_>(s|\rho)$ to depend on ρ and on other details pertaining to the microscopic scales only through the characteristic droplet size, s_c . The characteristic size s_c is defined, apart from a proportionality constant (see below), as the ratio of the second to the first moment of the droplet size distribution. This leads us to formulate the following scaling ansatz³¹.

$$P_>(s|\rho) = s^{-\alpha} f\left(\frac{s}{s_c}\right) \quad (1)$$

where, s_c depends on ρ , and it is expected to diverge at the critical concentration as^{30,31}.

$$s_c = a \left| \frac{\rho - \rho_c}{\rho_c} \right|^{-\phi} = a |\tilde{\rho}|^{-\phi}. \quad (2)$$

where $\alpha \geq$ and $\phi > 0$ are critical exponents, a is a constant and f is the so-called scaling function³⁰.

Thus, the scaling of the SDF is equivalent to saying that, apart from the singular behaviour $s^{-\alpha}$, the remaining s and ρ dependence occurs only through the ratio s/s_c . All extra dependencies are encapsulated in s_c through the constant a , ρ_c and, possibly, on the specific form of the scaling function f . A consequence of scaling and singular behaviour described by Eq. (2), i.e. the divergence of the characteristic size of the droplets, is that details of any specific system may not affect the value of the critical exponents, which are therefore expected to be universal, i.e. independent of specific details of the system.

To determine the exponents α and ϕ , we introduce the moments of $P_{>}(s|\rho)$ for $k>0$

$$\langle s^k \rangle = \int_0^\infty s^k \cdot \left(-\frac{dP_{>}(s|\rho)}{ds} \right) ds = c_k \cdot s_c^{k-\alpha} \quad (3)$$

where the scaling ansatz Eq. (1) has been used in the last step and c_k is given by

$$c_k = k \int_0^\infty x^{k-1-\alpha} f(x) dx$$

which depends on the function f but is independent of $\tilde{\rho}$. From Eq. (3), we deduce that

$$\frac{\langle s^{k+1} \rangle}{\langle s^k \rangle} = \frac{c_{k+1}}{c_k} \cdot s_c = \frac{c_{k+1}}{c_k} \cdot a |\tilde{\rho}|^{-\varphi} \quad (4)$$

If the scaling ansatz of Eq. (1) is correct, by plotting the ratio of moments $\langle s^{k+1} \rangle / \langle s^k \rangle$ for various values of k as a function of $1/|\tilde{\rho}|$ in a double logarithmic scale, we should obtain straight parallel lines with slope ϕ and intercept $\ln(ac_{k+1}/c_k)$.

Scaling behaviour of the droplet size distributions of FUS

We investigated the validity the ansatz in Eq. (2) using experimental data on the RNA-binding protein FUS, which are available for both the untagged and the SNAP-tagged protein²⁷ (Table S1). We calculated the survival distribution function, Eq. (2), and its moments, Eq. (3). We then plotted the moment ratios versus the inverse distance from the critical concentration $1/|\tilde{\rho}|$ in double logarithmic scale (Figure 1). We used the estimates of the critical concentrations, $\rho_c = 5.0 \mu\text{M}$ for FUS and $\rho_c = 5.4 \mu\text{M}$ for SNAP-tagged FUS, obtained below, in a self-consistency check of the validity of the scaling ansatz. We observed that the moment ratios at different distances from the critical concentration fall onto straight lines, as predicted by the scaling ansatz. In addition, the weighted average slope (see Eqs. (15) and (16) below) of the lines for different moment ratios is 0.95 ± 0.05 for untagged FUS, and 0.95 ± 0.05 for SNAP-tagged FUS, which is in good agreement with $\phi=1$ for the exponent in the scaling ansatz in Eq. (1) (Figure 1A,C).

Having determined the exponent ϕ , we can also determine the exponent α using Eq. (3)

$$\langle s^k \rangle = c_k \cdot s_c^{k-\alpha} \propto \left| \frac{\rho - \rho_c}{\rho_c} \right|^{-\varphi \cdot (k-\alpha)} \quad (5)$$

The exponent α is then calculated from

$$m = (k - \alpha)\varphi \rightarrow \alpha = k - \frac{m}{\varphi} \quad (6)$$

where m is the slope of the linear fit of the double logarithmic plot of $\langle s \rangle$ (the first moment, $k=1$) vs $1/|\tilde{\rho}|$. We then plotted the mean droplet size, $\langle s \rangle$, versus the inverse distance from the critical concentration $1/|\tilde{\rho}|$ on a natural logarithm scale, which could be fitted using a line with a slope




Figure 1.

Determination of the critical exponents for FUS of the scaling invariance.

(A,C) Determination of the exponent φ for FUS (A) and SNAP-tagged FUS (C). The ratios of the average moments of the droplet sizes ($\langle s^{k+1} \rangle / \langle s^k \rangle$, at $k=0.5, 1, 1.5, 2$, Eq. (4)) are represented at various distances from the critical concentration ($|\bar{\rho}|$). The exponent φ for each value of k was determined by error weighted linear regressions. The exponent φ and its error were determined as mean and standard deviation of the three independent measurements (Eqs. (15) and (16)). Error bars are shown in inset for graphical clarity. (B,D) Determination of the exponent α for FUS (B) and SNAP-tagged FUS (D). The mean of the droplet size distributions is plotted at various distances from the critical concentration ($|\bar{\rho}|$). The value of the exponent m was determined by error weighted linear regression (Eq. (6)), using $\varphi=1$, where the errors were standard deviations of the three independent measurements (Eq. 16). Error bars, which were obtained as the standard deviation of the three independent measurements are shown in inset for graphical clarity. Error weighted linear regressions are performed in both cases excluding the data point at the lowest concentration $p = 0.125 \mu\text{M}$. The fit corresponding to the scaling ansatz, compatible with $\varphi=1$ and $\alpha=0$, is represented by a dashed gray line with a slope of 1.

$m = 0.99 \pm 0.05$ for untagged FUS, and 0.93 ± 0.07 for SNAP-tagged FUS (**Figure 1B,D**), which is consistent with $m=1$. Using the value of $\phi = 1$, determined based on **Eq. (4)**, we obtain $\alpha = 0$. Taken together, these data support the validity of the scaling ansatz of **Eq. (1)**.

The droplet size distribution of FUS is log-normal

The above analysis suggests that the droplet size distribution may follow a log-normal distribution. The scaling ansatz of **Eq. (1)** for the SDF is equivalent to the following scaling for the probability density distribution

$$P(s|\rho) = -\frac{dP_{>}(s|\rho)}{ds} = s^{-1-\alpha} \cdot F(s/s_c), \quad s_c \propto |\tilde{\rho}|^{-\varphi} \quad (7)$$

where f , the scaling function in **Eq. (1)**, and F are related as follows

$$f(z) = z^{\alpha} \int_z^{\infty} x^{-\alpha-1} F(x) dx \quad (8)$$

The log-normal droplet size distribution $P(s|\rho)$ is

$$P(s|\rho) = s^{-1} \frac{1}{\sigma\sqrt{2\pi}} \exp\left\{-\frac{(\ln(s/s_0))^2}{2\sigma^2}\right\} \quad (9a)$$

With

$$s_0 \equiv s_c e^{-3\sigma^2/2} \quad \text{where} \quad s_c = \frac{\langle s^2 \rangle}{\langle s \rangle} \quad (9b)$$

being the characteristic droplet size as defined above. Consequently, the size survival distribution function is

$$P_{>}(s|\rho) = \frac{1}{2} \cdot \text{erfc}\left(\frac{\ln(s/s_0)}{\sigma \cdot \sqrt{2}}\right) \quad (10)$$

The values of s_0 and σ can be determined as the average and the variance, of $\ln(s/u)$ obtained at each concentration:

$$\begin{aligned} \ln(s_0/u) &\equiv \langle \ln(s/u) \rangle \\ \sigma^2 &\equiv \text{Var}(\ln(s/u)) \equiv \langle \ln^2(s/u) \rangle - \langle \ln(s/u) \rangle^2 \end{aligned} \quad (11)$$

where u is an arbitrary (and irrelevant) constant with the same units as s . In the following, when not stated, it is implicitly assumed that $u=1$ in the same units as s . The droplet sizes follow a log-normal distribution only if the survival distribution functions or, equivalently, the size distribution functions, multiplied by s , collapse when plotted versus **Eq. (9)** or **Eq. (10)** with the values of s_0 and σ of each droplet size distributions obtained at different concentrations (**Eq. (11)**). We determined s_0 and σ values for each distribution (**Figure 2A,B**) and plotted the properly rescaled size distribution functions versus $[\ln(s/s_0)]/\sigma$ (**Figure 2C**). We observed that the size distribution functions collapsed for both FUS and SNAP-tagged FUS (**Figure 2C**). Furthermore, the collapsed curve overlapped with the analytic log-normal distribution we computed with $s_0=1$, $\sigma=1$, the normal distribution in the rescaled variables.

The observed collapse supports the observation that droplet size distributions from different experiments follows a log-normal behaviour.




Figure 2.

Log-normal behavior of FUS and SNAP-tagged FUS size distributions below the critical concentration.

(A, B) Variation of the size distribution with protein concentration: $\ln s_0$ (A) and σ (B). $\ln s_0$ and σ (inset) were computed for FUS (blue) and SNAP-tagged FUS (red) using Eq. (11). Error bars (in inset for graphical clarity) are estimated from three independent measurements. While droplet sizes increase with concentration, the width of distribution does not change considerably. (C) The collapse of the droplet size distribution functions is consistent with a log-normal behaviour. The droplet size distribution functions for both untagged FUS (blue) and SNAP-tagged FUS (red) are plotted after rescaling the sizes by the $\ln s_0$ and σ values, the first and second moment of the logarithm of the droplet size distribution, which are a function of the concentration. The rescaled curves for both the untagged and the tagged protein collapse to the normal distribution (gray dashed), as expected when the non-rescaled droplet sizes follow a log-normal distribution.

Independence of the variance of the distribution from the concentration of FUS

The log-normal behaviour described above is consistent with the scale invariance underlying [Eq. \(1\)](#) if the variance of the log-normal distribution, σ , is independent of ρ or, equivalently, of $\tilde{\rho}$ [32](#). Indeed, comparing [Eq. \(1\)](#) with [Eqs. \(7\)](#), [\(9a,b\)](#) and [\(10\)](#) we obtain that the scaling invariance holds with

$$\begin{aligned} f(x) &= \frac{1}{2} \operatorname{erfc} \left[\frac{\ln(xe^{3\sigma^2/2})}{\sigma\sqrt{2}} \right] \\ F(x) &= \frac{1}{\sigma\sqrt{2\pi}} \exp \left\{ - \left[\frac{\ln(xe^{3\sigma^2/2})}{\sigma\sqrt{2}} \right]^2 \right\} \end{aligned} \quad (12)$$

with $\alpha=0$ and $\phi = 1$ ([Figure 1](#)). Notice that the log-normal distribution implies that $\alpha=0$ whereas the value of ϕ is not determined a priori. Furthermore, the k -th moment of the log-normal distribution is

$$\langle s^k \rangle = s_c^k e^{\sigma^2 k(k-3)/2}, \quad (13)$$

which, compared with the scaling prediction [Eq. \(3\)](#), is also consistent with $\alpha = 0$, appropriate for the log-normal, and

$$c_k = e^{\sigma^2 k(k-3)/2} \quad (14)$$

which is independent of $\tilde{\rho}$ if σ is independent of $\tilde{\rho}$, see [Eq. \(3\)](#).

This prediction is verified in [Figure 2B](#), where the σ^2 values are shown to be nearly uniform at different concentrations, with the exception of the data at the lowest concentration values, i.e. those furthest away from the critical concentration. These results are consistent with the scaling ansatz in the vicinity of $\tilde{\rho} = 0$.

Our analysis does not exclude the possibility that σ might depend on specific experimental conditions, even though the data that we analysed are suggestive of at most a weak dependence. We note that such dependence, even if present, does not invalidate the scaling, as long as the exponents do not depend on the experimental conditions.

Estimation of the critical concentration of FUS using the scale invariance

The fact that the scaling ansatz is satisfied for different set of experiments opens a possibility to estimate the critical concentration. The scaling with $\phi=1$ predicts $(\langle s_k \rangle)^{-1/k}$ vs ρ to be a straight line with a slope depending on k . It is important to note that this is the consequence only of the scaling ansatz and not of the log-normal distribution. The line should intersect the ρ -axis at the critical concentration providing an estimate of ρ_c .

We illustrate this process using the FUS data by plotting $(\langle s_k \rangle)^{-1/k}$ vs ρ . As expected, for different values of k we obtained a straight line fit of the points near ρ_c ([Figure 3](#)). Due to experimental uncertainties, the various lines, one for each value of k , lead to a slightly different estimate of ρ_c . The average of the estimated ρ_c values is $5.0 \pm 0.2 \mu\text{M}$ for FUS and $5.4 \pm 0.4 \mu\text{M}$ for SNAP-tagged FUS ([Figure 3](#)). The different ρ_c values predicted based on the scaling ansatz using different k

values enable the estimation of error of the predicted critical concentration (**Figure 3**). Both estimates are higher than the values of ρ_c originally reported ($\rho_c = 2 \mu\text{M}$ of untagged and $\rho_c = 3 \mu\text{M}$ of tagged FUS²⁷) but consistent with them once taken into account the behaviour of the plateau of the absorbance of a spin-down assay used in that work. We also note that our estimates are compatible with other ones recently reported³³.

We then used both the values of ρ_c to probe the collapse of the size distribution functions (**Figure 4**). We observed that both in cases of untagged and tagged FUS, the SDFs collapsed with our estimated value of ρ_c (**Figure 4**).

These results indicate that the scaling model can be used to estimate the critical concentration based on the distribution of droplet sizes.

Estimation of the critical concentration of α -synuclein using the scale invariance

As α -synuclein droplets (called clusters) were recently reported below the critical concentration³⁴, we aimed at characterising whether such droplets also follow scale invariant size distribution. Using A90C α -synuclein labelled with Alexa Fluor 647, we monitored droplet formation as previously described³⁵. We measured droplet sizes at 5% PEG concentration 10 min after detection of liquid-like condensates using increasing concentrations of α -synuclein (20, 40, 50, 60, 75, 80 and 100 μM) (**Table S2**). Using k values of 0.25, 0.75, 1.25, 1.75, we determined the critical exponents in the scaling ansatz (Eqs. (4) and (6)), obtaining $\phi = 1.3 \pm 0.2$ (**Figure 5A**) and $m=1-\alpha = 0.9 \pm 0.2$ (**Figure 5B**). As control, we also determined the critical exponents using the scaling ansatz in a different way, using **Eq. (19)**, obtaining $\phi = 1.1 \pm 0.2$ and $m=1-\alpha = 0.8 \pm 0.2$, corroborating the validity of the scale invariant model. We then estimated the critical concentration using the two methods obtaining $\rho_c = 137 \pm 10 \mu\text{M}$, using **Eq. (18)** (**Figure 5C**) and $\rho_c = 125 \pm 7 \mu\text{M}$, using **Eq. (19)** (**Figure 5D**). Images of the droplets at the different concentrations are shown in **Figure 6A**. We also observed that the droplet size distributions are stationary below the critical concentration, as expected (**Figure 6B,C**).

Discussion

Although growing experimental evidence indicates the presence of protein condensates both in vitro and in vivo^{1-5,36}, their exact nature, and the mechanisms responsible for their formation are not fully understood. For example, it is unclear whether the droplets observed below the saturation concentration and those detected above result from the same process. Here we aimed at describing phase separation using a scaling ansatz.

With this aim, we analyzed a series of droplet size distributions of FUS and α -synuclein. We found that the droplet size distributions follow a scale-invariant log-normal behaviour. Scale invariance means that the description of the behaviour of a system remains the same regardless of the scale of observation. It has been seen for example in self-similar systems like fractals, which repeat patterns at different length scales³⁷. Scale invariance is commonly observed in physics, biology, and economics, where it helps understand complex systems by identifying consistent patterns and fundamental properties³⁸. The critical exponents of the scaling model are the same for different systems. We also note the generality of the log-normal model for nucleation and grain size growth ranging from crystal seeding^{39,40} to the mass size distributions in organism growth of various sea organisms³².

It is interesting to discuss the links of the scale invariance observed here with the theoretical models that have been proposed to date to explain the phenomenon of protein phase separation. A commonly adopted framework is the Flory-Huggins theory of phase separation^{14,16}. In its




Figure 3

Estimation of the critical concentration of FUS using the scale invariance.

(A,B) Critical concentration of FUS ($5.0 \pm 0.2 \mu\text{M}$) (A) and SNAP-tagged FUS ($5.4 \pm 0.4 \mu\text{M}$) (B). The scaling model predicts that the function of the moments plotted versus the concentration ρ becomes a straight line near the critical concentration p_c and intersects the ρ -axis at p_c , independently of the value of k . Error weighted linear regressions are performed in both cases excluding the data point at the lowest concentration $\rho = 0.125 \mu\text{M}$. The resulting estimate of the critical concentration is shown in green along with the corresponding standard deviation, estimated from three independent measurements.




Figure 4.

Collapse of the droplet size distributions of FUS as predicted by the scale invariance.

If the scaling ansatz of Eq. (2) holds, the standard deviation σ of the log-normal distribution should not depend on the distance from the critical concentration, and a collapse should be achieved by rescaling the size with the distance $|\rho|$ from the critical concentration. **(A,C)** Droplet size distributions derived from the experimental data of untagged FUS at 0.125, 0.25, 0.5, 1.0 and 2.0 μM concentrations (A) and SNAP-tagged FUS at 0.125, 0.25, 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0 μM concentrations (C), and their standard error of the mean from three independent measurements (26). **(B,D)** Collapse of the droplet size distributions rescaled by the estimated critical concentration. The error bars show the standard error of the mean from the three independent measurements (26).




Figure 5.

Estimation of the critical concentration of α-synuclein using the scale invariance.

(A) Determination of the critical exponent ϕ . The ratios of the average moments of the droplet sizes ($\langle s^{k+1} \rangle / \langle s^k \rangle$, at $k=0.25, 0.75, 1.25, 1.75$, Eq. (4)) are represented at various distances $|\rho|$ from the critical concentration. The exponent ϕ and its error for each value of k were determined as a mean and standard deviation of the three independent measurements (Eqs. (15) and (16)). Error bars are shown in inset for clarity. (B) Determination of the critical exponent α . The mean of the droplet size distributions is plotted at various distances $|\rho|$ from the critical concentration. The value of the exponent α was determined by error weighted linear regression (Eq. 6), using $\phi=1$, where the errors were standard deviations of the five independent measurements (Eq. (16)). Error bars, which were obtained as the standard deviation of the five independent measurements are shown in inset for clarity. Error-weighted linear regressions were performed. The fit corresponding to the scaling ansatz, compatible with $\phi=1$ and $\alpha=0$, is represented by a scattered gray line with a slope of 1. (C,D) Determination of the critical concentration for α-synuclein using the scaling ansatz in two different ways, either using Eq. (18) (C), resulting in $\rho_c = 137 \pm 10 \mu\text{M}$, or Eq. 19 (D), resulting in $\rho_c = 125 \pm 7 \mu\text{M}$, which are consistent within errors. The scaling model predicts that the function of the moments plotted versus the concentration ρ becomes a straight line near the critical concentration ρ_c and intersects the ρ -axis at ρ_c , independently of the value of k . Error-weighted linear regressions were performed. The resulting estimates of the critical concentration are shown in green along with the corresponding standard deviation, estimated from four independent measurements. In panel D, ϕ was constrained to 1.0 using Eq. (19).




Figure 6.

The droplet size distribution is stationary below the critical concentration.

(A) Images of α -synuclein droplets at increasing concentrations of α -synuclein. **(B,C)** After an initial transient of 5 min, the droplet size distributions remain approximately stationary below the critical concentration, as shown for the case of 50 μM α -synuclein concentration.

simplest form, this theory describes a first-order phase separation in a system of homopolymers, which can also be adapted to polyampholites¹⁷. The Flory-Huggins theory has been extended to associative polymers, with the aim of modelling the sequence composition of proteins, by Flory and Stockmayer, who described the phase separation in terms of a third-order gelation process^{18,19}. Semenov and Rubinstein modified the Flory-Stockmayer theory, reporting that the gelation process is not a real phase transition, as all the derivatives of the free energy are analytical at the gelation point²⁰. It was also argued that another generalization of the Flory-Stockmayer theory²¹ could describe phase transitions in cytoskeletal networks²². In an older study, γ -crystallin was reported to undergo a second-order phase separation process²⁴, consistently with observations in lysozyme solutions²⁵.

More recently, it has been suggested that the protein phase separation process is coupled with percolation²³. It is notable that percolation models exhibit a continuous transition marked by a diverging correlation length and a scale-invariant droplet size distribution³¹, resembling the general form as presented in Eqs. (1) and (2). However, the critical exponents for 3D percolation at the critical threshold are $\alpha=1.19$ and $\phi=2.21$, and in mean-field ($d>6$) scenarios, they are $\alpha=1.5$ and $\phi=2$, as per existing literature³¹. In contrast, our scaling analysis is consistent with $\alpha=0$ and $\phi=1$. Therefore, a phase separation coupled with percolation model²³ could be consistent with the scaling behavior reported here if this coupling would induce a shift in the universality class from that of standard percolation³¹. This possibility should be further investigated to determine whether this type of models could predict the critical exponents $\alpha=0$ and $\phi=1$.

Scale invariance may appear to be in contrast with the Flory-Huggins theory, which characterizes droplet formation as a consequence of nucleation processes within a metastable state of a supersaturated system. In such cases, however, when the correlation length exceeds the cellular dimensions, the remnants of what should have been a first-order phase transition in an infinite system may effectively exhibit indistinguishable characteristics from a second-order phase transition. We also note that a power law distribution, but above the transition temperature, has recently been reported for nucleoli⁴¹.

Quite generally, the existence of scale invariance for the droplet size distribution, at least under the conditions investigated here, imposes stringent constraints on theoretical models that aim to elucidate protein phase separation. The importance of a scaling analysis lies in its ability to uncover the fundamental aspects of universal phenomena, transcending models confined solely to specific systems for which they were originally designed.

As a practical consequence of the scaling model, we found that the moments of the droplet size distribution can be used to obtain an estimate of the critical concentration for phase separation.

Conclusions and Perspectives

It has been challenging to determine critical concentrations in the study of protein phase separation, particularly in cell systems. As a consequence, it remains still largely unclear when the observed protein droplets are formed at sub-saturation concentrations, and when they represent phase-separated condensates at super-saturation concentrations. To make progress towards understanding this problem, we have analyzed the probability distributions of the droplet sizes at different concentrations. The scale-invariant log-normal behaviour that we have reported offer a tool to determine the critical concentration. This approach should enable the positioning of a system with respect to the phase boundary when experimental data could be obtained about the length (1D), area (2D), or volume (3D) of the condensates. We anticipate that this method will find applications in the study of protein condensates under cellular conditions.

We also note that the scale invariance of the droplet size distribution that we have reported is characteristic of critical phenomena, which tend to be highly sensitive to environmental conditions. The formation and dimensions of protein droplets could therefore be rather tunable. This feature would appear to be favourable for the control of the formation of biomolecular condensates by the protein homeostasis system, and also suggest that protein phase separation could be efficiently modulated pharmacologically.

Methods

Expression and purification of α -synuclein

Human wild-type α -synuclein and A90C cysteine variants were purified from *Escherichia coli* BL21 (DE3)-gold (Agilent Technologies) expressing plasmid pT7-7 encoding for α -synuclein as previously described^{35,42,43}. Following purification in 50 mM trisaminomethane-hydrochloride (Tris-HCL) at pH 7.4, α -synuclein was concentrated using Amicon Ultra-15 centrifugal filter units (Merck Millipore). The protein was subsequently labelled with a 1.5-fold molar excess of C5 maleimide-linked Alexa Fluor 647 (Invitrogen Life Technologies) overnight at 4 °C with constant mixing. The excess dye was removed using an Amicon Ultra-15 centrifugal filter unit and used immediately for phase separation experiments.

Determination of the droplet distribution of α -synuclein

The experimental conditions were determined from a previously described phase boundary³⁵. To induce liquid droplet/condensate formation, wild-type α -synuclein was mixed with an A90C variant labelled with Alexa 647 at a 100:1 molar ratio in 50 mM Tris-HCL and 5% polyethylene glycol 10,000 (PEG) (Thermo Fisher Scientific). The final mixture was pipetted onto a 35-mm glass-bottom dish (P35G-1.5-20-C; MatTek Life Sciences) and immediately imaged on Leica Stellaris Will inverted stage scanning confocal microscope using a 40 $\times$ /1.3 HC PL Apo CS oil objective (Leica Microsystems) at room temperature. The excitation wavelength was 633 nm for all experiments. For liquid droplet size characterisation, images were captured 10 min post-liquid droplet formation. To analyse the possible time dependence of the liquid droplet size distribution, phase separation was induced, and 5 min post onset of phase separation, the glass-bottom dish containing the experiment was sealed and closed to maintain a stable and controlled environment. Subsequently, images were captured at each designated time point. All images were processed and analysed in ImageJ (NIH). Images were analysed by applying a threshold function in ImageJ that excluded the background of the image and identified the liquid droplets as having a circularity of 0.8-1.

Determination of the droplet distributions of FUS

We used previously published data on FUS droplets²⁷, where the fluorescence intensity of the FUS droplets was reported to be proportional to their diameters²⁷. The abundance of untagged FUS droplets formed at 0.125, 0.25, 0.5 and 1.0, 2.0 μ M concentrations were measured by nanoparticle tracking analysis for 51 different droplet sizes with three repetitions for each measure²⁷. Similarly, the abundance of SNAP-tagged FUS droplets at 0.125, 0.25, 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0 μ M concentrations were measured for 64 different sizes with 3 repetitions for each measure²⁷. For each data point, we determined the mean droplet abundance and the standard error of the mean from the three experimental replicates. The survival distribution function was computed as $P_{>x} = N_{>x}/N_{\text{tot}}$, where $N_{>x}$ is the number of droplets above size x , and N_{tot} is the number of droplets. The values of x were chosen according to the original data²⁷.

Determination of the critical exponents for FUS

We computed the k -th moment of each droplet size, determined the average for each value of k and computed the ratios of the subsequent moments (Eq. (4)). We tested several k values from 0.25 to 3 in 0.25 increments, determined the moment ratios in each case. We selected four k values ($k=0.5$, $k=1.0$, $k=1.5$ and $k=2.0$) where the moment ratios at each concentration provided the best linear fit for both FUS and SNAP-tagged FUS. Then we generated a log-log plot of the moments and $\tilde{\rho}$, the distance from the critical concentration (Figure 1). For each value of k , the points were estimated by taking the average of the computation done on the 3 independent measurements. The errors of the data points are estimated as the standard error of the mean. At each k value, we performed a linear regression between the points at each concentration²⁷ determining the slope, resulting in four ϕ_k values, with an associated error from the weighted linear regression. Given the independent observations ϕ_k with variance σ_k^2 , the value of the exponent was obtained as the weighted average

$$\bar{\phi} = \frac{\sum_k \frac{\phi_k}{\sigma_k^2}}{\sum_k \frac{1}{\sigma_k^2}} \quad (15)$$

The error on the exponent ϕ was computed as

$$\text{Var}(\bar{\phi}) = \frac{1}{\sum_k \frac{1}{\sigma_k^2}} \quad (16)$$

The exponent α was determined based on the log-log plot of the mean of the droplet size distribution at various distances from the critical concentration ($|\tilde{\rho}|$) (Figure 1). We performed a linear regression weighted by the inverse of the variance of the data points corresponding, as previously, to the computed mean of the three independent experiments for each concentration (Eq. (6)) using $\phi=1$. The error on the slope from the weighted linear regression is associated with the estimate of the exponent m . The exponent α is then derived using $\alpha = 1 - m$.

At each concentration, from the distribution of $\ln(s)$ we determined s_0 as $\langle \ln(s) \rangle = \ln(s_0)$, and σ as the standard deviation of $\ln(s)$. At each concentration, we plotted the droplet size distribution function versus $\frac{\ln(s/s_0)}{\sigma}$ (Eq. (10)) using the s_0 and σ values determined from the corresponding droplet size distribution at the given concentration. The log-normal behaviour is demonstrated by the resulting collapse (Figure 2). Furthermore, the collapsed curves overlapped with a theoretical log-normal curve computed with $s_0=1$, $\sigma=1$, which is the normal distribution in the rescaled log variables.

Determination of the critical concentration for FUS

The k -th moment of the log-normal distribution (Eq. (13)) and the scaling ansatz with $\alpha=0$ (Eq. (3)) gives

$$(\langle s^k \rangle)^{-1/k} = \frac{1}{s_c} e^{\frac{(3-k)\sigma^2}{2}} \quad (17)$$

where $s_c = a \cdot \left| \frac{\rho - \rho_c}{\rho_c} \right|^{-1}$, a is a proportionality constant independent of k , and σ is the standard deviation of the logarithm of the droplet probability size distribution. Therefore

$$(\langle s^k \rangle)^{-1/k} = \frac{1}{a} e^{\frac{(3-k)\sigma^2}{2}} \left| \frac{\rho - \rho_c}{\rho_c} \right| \quad (18)$$

plotted versus the concentration intersects the x-axis at the value of the critical concentration.

Determination of the critical exponents for α -synuclein

For α -synuclein, the critical exponents were determined in two different ways, the first using the same method as for FUS (Eq. (18) [↗](#)), and the second using

$$\left(\frac{\langle s^k \rangle}{\langle s \rangle}\right)^{-1/(k-1)} = \left(\frac{c_l}{c_k}\right)^{1/(k-1)} \frac{1}{a} \left| \frac{\rho - \rho_c}{\rho_c} \right|^\varphi \quad (19)$$

Determination of the critical concentration of α -synuclein

As for the critical exponents, we determined the critical concentration of α -synuclein using the scaling ansatz in two different ways, either using Eq. (18) [↗](#) or using Eq. (19) [↗](#) with the constrain $\phi=1$. We plotted the moments of the droplet size (Eq. (3) [↗](#)) versus the concentration, using a range of values of k to cover majority of the data. Analogously to what happens in Figure 3 [↗](#), the lines plotted in Figure 5D [↗](#) intercept the x axis on the same point, which corresponds to the critical concentration. We performed a linear regression weighted by the inverse variance, determining the intercept on the x-axis. For each value of k , we calculated the regression errors of both intercept and slopes. We estimated the critical concentration from each fit as the intercept calculated as $\rho_{c,i} = \frac{-q}{m}$, where q and m are the y-axis intercept and the slope retrieved by the fit, respectively, where the subscript i indicates each independent experiment. The estimated ρ_c was obtained as the mean of the different independent values $\rho_{c,i}$. The error on the estimate of ρ_c is obtained as the standard error of the mean of the $\rho_{c,i}$.

Acknowledgements

We acknowledge insightful discussions with Drs. Serena Carra, Jonathan Vinet, Alex Buell and Soumik Ray. This work was supported by AIRC IG 26229 (M.F.).

References

1. Brangwynne C.P., et al. (2009) **Germline P granules are liquid droplets that localize by controlled dissolution/condensation** *Science* **324**:1729–1732
2. Hyman A.A., Weber C.A., Jülicher F. (2014) **Liquid-liquid phase separation in biology** *Annu. Rev. Cell Biol* **30**:39–58
3. Banani S.F., Lee H.O., Hyman A.A., Rosen M.K. (2017) **Biomolecular condensates: Organizers of cellular biochemistry** *Nat. Rev. Mol. Cell Biol* **18**:285–298
4. Alberti S., Hyman A.A. (2021) **Biomolecular condensates at the nexus of cellular stress, protein aggregation disease and ageing** *Nat. Rev. Mol. Cell Biol* **22**:196–213
5. Lyon A.S., Peeples W.B., Rosen M.K. (2021) **A framework for understanding the functions of biomolecular condensates across scales** *Nat. Rev. Mol. Cell Biol* **22**:215–235
6. Buchan J.R., Parker R. (2009) **Eukaryotic stress granules: The ins and outs of translation** *Mol. Cell* **36**:932–941
7. Andersen J.S., et al. (2005) **Nucleolar proteome dynamics** *Nature* **433**:77–83
8. Mateju D., et al. (2017) **An aberrant phase transition of stress granules triggered by misfolded protein and prevented by chaperone function** *EMBO J* **36**:1669–1687
9. Qamar S., et al. (2018) **FUS phase separation is modulated by a molecular chaperone and methylation of arginine cation- π interactions** *Cell* **173**:720–734
10. Mitrea D.M., Mittasch M., Gomes B.F., Klein I.A., Murcko M.A. (2022) **Modulating biomolecular condensates: A novel approach to drug discovery** *Nat. Rev. Drug Discov* **21**:841–862
11. Vendruscolo M., Fuxreiter M. (2022) **Protein condensation diseases: Therapeutic opportunities** *Nat. Comm* **13**
12. Fuxreiter M., Vendruscolo M. (2021) **Generic nature of the condensed states of proteins** *Nat. Cell Biol* **23**:587–594
13. Jawerth L., et al. (2020) **Protein condensates as aging Maxwell fluids** *Science* **370**:1317–1323
14. Flory P.J. (1942) **Thermodynamics of high polymer solutions** *J. Chem. Phys* **10**:51–61
15. Huggins M.L. (1942) **Some properties of solutions of long-chain compounds** *J. Phys. Chem* **46**:151–158
16. Michaels T.C., et al. (2023) **Amyloid formation as a protein phase transition** *Nat. Rev. Phys* **5**:379–397
17. Overbeek J.T., Voorn M.J. (1957) **Phase separation in polyelectrolyte solutions. Theory of complex coacervation** *J. Cell. Comp. Physiol* **49**:7–26

18. Flory P.J. (1941) **Molecular size distribution in three dimensional polymers. I. Gelation** *J. Am. Chem. Soc* **63**:3083–3090
19. Stockmayer W.H. (1943) **Theory of molecular size distribution and gel formation in branched-chain polymers** *J. Chem. Phys* **11**:45–55
20. Semenov A.N., Rubinstein M. (1998) **Thermoreversible gelation in solutions of associative polymers. 1. Statics** *Macromolecules* **31**:1373–1385
21. Tavares J., Teixeira P., Telo da Gama M., Sciortino F. (2010) **Equilibrium self-assembly of colloids with distinct interaction sites: Thermodynamics, percolation, and cluster distribution functions** *J. Chem. Phys* **132**
22. Bueno C., Liman J., Schafer N.P., Cheung M.S., Wolynes P.G. (2022) **A generalized flory-stockmayer kinetic theory of connectivity percolation and rigidity percolation of cytoskeletal networks** *PLoS Comp. Biol* **18**
23. Mittag T., Pappu R.V. (2022) **A conceptual framework for understanding phase separation and addressing open questions and challenges** *Mol. Cell* **82**:2201–2214
24. Thomson J.A., Schurtenberger P., Thurston G.M., Benedek G.B. (1987) **Binary liquid phase separation and critical phenomena in a protein/water solution** *Proc. Natl. Acad. Sci. USA* **84**:7079–7083
25. Ishimoto C., Tanaka T. (1977) **Critical behavior of a binary mixture of protein and salt water** *Phys. Rev. Lett* **39**
26. Stender E.G., et al. (2021) **Capillary flow experiments for thermodynamic and kinetic characterization of protein liquid-liquid phase separation** *Nat. Comm* **12**
27. Kar M., et al. (2022) **Phase-separating RNA-binding proteins form heterogeneous distributions of clusters in subsaturated solutions** *Proc. Natl. Acad. Sci. USA* **119**
28. Li Y., Lubchenko V., Vorontsova M.A., Filobelo L., Vekilov P.G. (2012) **Ostwald-like ripening of the anomalous mesoscopic clusters in protein solutions** *J. Phys. Chem. B* **116**:10657–10664
29. Shimobayashi S.F., Ronceray P., Sanders D.W., Haataja M.P., Brangwynne C.P. (2021) **Nucleation landscape of biomolecular condensates** *Nature* **599**:503–506
30. Brankov J.G., Danchev D.M., Tonchev N.S. (2000) **Theory of critical phenomena in finite-size systems: Scaling and quantum effects**
31. Stauffer D., Aharony A. (2018) **Introduction to percolation theory**
32. Giometto A., Altermatt F., Carrara F., Maritan A., Rinaldo A. (2013) **Scaling body size fluctuations** *Proc. Natl. Acad. Sci. USA* **110**:4646–4650
33. Patel A., et al. (2015) **A liquid-to-solid phase transition of the ALS protein FUS accelerated by disease mutation** *Cell* **162**:1066–1077
34. Ray S., et al. (2023) **Mass photometric detection and quantification of nanoscale α -synuclein phase separation** *Nat. Chem* **15**:1306–1316

35. Dada S.T., et al. (2023) **Spontaneous nucleation and fast aggregate-dependent proliferation of α -synuclein aggregates within liquid condensates at neutral pH** *Proc. Natl. Acad. Sci. USA* **120**
36. Boija A., et al. (2018) **Transcription factors activate genes through the phase-separation capacity of their activation domains** *Cell* **175**:1842–1855
37. Song C., Havlin S., Makse H.A. (2005) **Self-similarity of complex networks** *Nature* **433**:392–395
38. Stanley H. E. (1999) **Scaling, universality, and renormalization: Three pillars of modern critical phenomena** *Rev. Mod. Phys* **71**:S358–S366
39. Binder K., Virnau P. (2016) **Overview: Understanding nucleation phenomena from simulations of lattice gas models** *J. Chem. Phys* **145**
40. Bergmann R.B., Bill A. (2008) **On the origin of logarithmic-normal distributions: An analytical derivation, and its application to nucleation and growth processes** *J. Cryst. Growth* **310**:3135–3138
41. Lee D.S., et al. (2023) **Size distributions of intracellular condensates reflect competition between coalescence and nucleation** *Nat. Phys* **19**:586–596
42. Fusco G., et al. (2014) **Direct observation of the three regions in α -synuclein that determine its membrane-bound behaviour** *Nat. Comm* **5**
43. Hardenberg M.C., et al. (2021) **Observation of an α -synuclein liquid droplet state and its maturation into Lewy body-like assemblies** *J. Mol. Cell Biol* **13**:282–294

Editors

Reviewing Editor

Bin Zhang

Massachusetts Institute of Technology, Cambridge, United States of America

Senior Editor

Qiang Cui

Boston University, Boston, United States of America

Reviewer #1 (Public Review):

The authors analyse droplet size distributions of multiple protein condensates and fit to a scaling ansatz to highlight that they exhibit features of first-order and second-order phase transitions. While the experimental evidence is solid, the text lacks connection and contextualization to the well-understood expectations from the coupling of percolation and phase separation in protein condensates - a phenomenon that is increasingly gaining consensus amongst the community. The evidence supports the percolation+phase separation model rather than being close to a true critical point in the liquid-gas phase space. Overall, the work is useful to the community.

Strengths:

The experimental analysis of distinct protein condensates is very well done and the reported

exponents/scaling framework provides a clear framework to help the community help deconvolve signatures of percolation in condensates.

Weaknesses:

The principal concern this reviewer has is that the reviewers adopt a framing in this paper to present a discovery of second-order features and connections to criticality - however they ignore/miss the connections to percolation (a well-understood second-order transition that is expected to play a major role in protein condensates). I believe this needs to be addressed and the paper suitably revised to help connect with these expectations.

- Protein condensates have been increasingly understood to be described as fluids whose assembly is driven by a connection of density (phase separation, first-order) and connectivity (percolation, second-order) transitions. This has been long known in the polymer community (Flory, Stockmayer, Tanaka, Rubinstein, Semenov and others) and recently repopularized in the condensate community (by Pappu and Mittag, in particular, amongst others). The authors make no connections to any of this frameworks - which actually seem to be the essence of what they are describing.

- Percolation theory, which has been around for more than half-a-century, has clear-cut scaling laws that have essentially similar forms to the ansatz adopted by the authors and the commonalities/differences are not discussed by the authors - this is essential since this provides a physical basis for their ansatz rather than an arbitrary mathematical formulation. In particular, percolation models connect size distribution exponents to factors like dimensionality, valence, etc. and if these connections can be made with this data, that would be very powerful.

- The connections between spinodal decomposition and second-order phase transitions are very confusing. Spinodal decomposition happens when the barriers for first-order phase transitions are zero and systems can phase separate without crossing nucleation barriers. Further, the "criticality" discussed in the paper is confusing since it more likely refers to a percolation threshold and much less likely to a "critical temperature" (T_c - where spinodal and binodals become identical). I would recommend reframing this argument.

It's unlikely, in this reviewer's opinion, that the authors are actually discussing a "first-order" liquid-gas critical point - because saturation concentrations of these proteins can be much higher with temperature and the critical point would thus likely be at much higher concentrations (and ofc temperature). Further the scaling exponents don't fall in that class naturally. However, if the authors disagree, I would appreciate clear quantitative reasons (including through the scaling exponents in that universality class) and be happy to be convinced to change my mind. As provided, the data does not support this model.

<https://doi.org/10.7554/eLife.94214.2.sa2>

Reviewer #2 (Public Review):

In response to the two referee reports, the authors have made substantial improvements. Regarding my previous concerns, the new data provided in Fig.6 for demonstrating that the droplet size distribution is stable over time is particularly valuable.

As to several of my other previous concerns regarding possible change in droplet size distribution over time, etc., the authors responded by stating that their system was below the critical concentration and therefore the possible scenarios pointed out in my previous report were not expected. While there may be a certain degree of validity to their argument, it would be much more helpful to the readers if the authors would bring up my previous

concerns briefly (as readers of the journal will likely have similar concerns) and then address them succinctly within the manuscript.

Apparently, as a key element in the authors' response to the referees, the term "transition concentration" in the originally submitted manuscript is now changed to "critical concentration" (including in the title and abstract). But the two terms do not have identical meaning. A transition concentration is usually recognized as the saturation concentration at which phase separation or some other transition process commences at a given temperature. The transition concentration can be lower than the critical concentration, whereas the critical concentration is associated with the critical temperature, above (or below, depending on the temperature dependence of phase separation) which phase separation is not possible. It will be best if the authors can clarify their usage of transition concentration vs. critical concentration in the version of record of their manuscript.

<https://doi.org/10.7554/eLife.94214.2.sa1>

Author response:

The following is the authors' response to the original reviews

eLife assessment:

In this useful study, the authors analyze droplet size distributions of multiple protein condensates and their fit to a scaling ansatz, highlighting that they exhibit features of first- and second-order phase transitions. The experimental evidence is still incomplete as the measurements were apparently done only at one time point, neglecting the possibility that droplet size distribution can evolve with time. The text would benefit from a connection to and contextualization with the well-understood expectations from the coupling of percolation and phase separation in protein condensates - a phenomenon that is increasingly gaining consensus amongst the community and that emphasizes "liquid-gas" criticality.

We have now carried out new experiments at multiple time points to establish that the droplet size distributions are stationary below the critical concentration. We have also addressed the comments made by the reviewers about the nature of the phase transition.

Our analysis does not depend on a specific hypothesis on the nature of the phase transition, whether it be percolation or a gas-liquid critical transition. The scaling that we observed is an emergent property that is independent from the possible theoretical models used to describe the phase transition. In fact, our scaling analysis indicates that any theoretical model proposed for protein phase separation should predict the critical exponents that we reported.

Reviewer #1:

The authors analyse droplet size distributions of multiple protein condensates and fit to a scaling ansatz to highlight that they exhibit features of first-order and second-order phase transitions. While the experimental evidence is solid, the text lacks connection and contextualization to the well-understood expectations from the coupling of percolation and phase separation in protein condensates - a phenomenon that is increasingly gaining consensus amongst the community. The evidence supports the percolation and phase separation model rather than being close to a true critical point in the liquid-gas phase space. Overall, the work is useful to the community.

We are grateful to the reviewer for these positive comments. We would like to emphasise that our contribution is not to propose a theoretical model, but rather to report a scaling

behaviour in the experimentally measured droplet size distributions. The main implication of our work is that any theoretical model should predict the scaling exponents that we derived from the experimental measurements.

Strengths:

The experimental analysis of distinct protein condensates is very well done and the reported exponents/scaling framework provides a clear framework to help the community deconvolve signatures of percolation in condensates.

Weaknesses:

The principal concern this reviewer has is that the reviewers adopt a framing in this paper to present a discovery of second-order features and connections to criticality - however, they ignore/miss the connections to percolation (a well-understood second-order transition that is expected to play a major role in protein condensates). I believe this needs to be addressed and the paper suitably revised to help connect with these expectations.

The scaling that we found is not characteristic standard percolation, since the exponents that we obtained ($a=0$ and $f=1$) are different from those of percolation ($a=1.19$ and $f=2.21$). This difference indicates that protein phase separation is not in the same universality class of standard percolation. Further studies will be required to understand whether theoretical models based on percolation could predict the observed critical exponents.

- Protein condensates have been increasingly understood to be described as fluids whose assembly is driven by a connection of density (phase separation, first-order) and connectivity (percolation, second-order) transitions. This has been long known in the polymer community (Flory, Stockmayer, Tanaka, Rubinstein, Semenov, and others) and recently repopularized in the condensate community (by Pappu and Mittag, in particular, amongst others). The authors make no connections to any of these frameworks - which actually seem to be the essence of what they are describing.

As mentioned above, our purpose was neither to support an existing theoretical model, nor to propose a new one. Rather, we have reported a scaling behaviour and scaling exponents not noted before. Further studies will be required to establish whether existing theoretical models could account for this scaling behaviour.

- Percolation theory, which has been around for more than half a century, has clear-cut scaling laws that have essentially similar forms to the ansatz adopted by the authors, and the commonalities/differences are not discussed by the authors - this is essential since this provides a physical basis for their ansatz rather than an arbitrary mathematical formulation. In particular, percolation models connect size distribution exponents to factors like dimensionality, valence, etc. and if these connections can be made with this data, that would be very powerful.

The scaling ansatz that we are using is commonly adopted in studies of critical phenomena, and it is not specific to percolation. The scaling exponents depends only on very few attributes like dimensionality, symmetries and if interactions are short or long range. These attributes determine the universality class. As such, scaling does not link with molecular determinants, but can distinguish different classes.

- The connections between spinodal decomposition and second-order phase transitions are very confusing. Spindal decomposition happens when the barriers for first-order phase transitions are zero and systems can phase separate without crossing nucleation

barriers. Further, the "criticality" discussed in the paper is confusing since it more likely refers to a percolation threshold and much less likely to a "critical temperature" (T_c - where spinodal and binodals become identical). I would recommend reframing this argument.

We cannot refer to percolation threshold as our model is not readily compatible with it. We elaborated and better explained the differences between these models.

It's unlikely, in this reviewer's opinion, that the authors are actually discussing a "first-order" liquid-gas critical point - because saturation concentrations of these proteins can be much higher with temperature and the critical point would thus likely be at much higher concentrations (and ofc temperature). Further, the scaling exponents don't fall into that class naturally. However, if the authors disagree, I would appreciate clear quantitative reasons (including through the scaling exponents in that universality class) and be happy to be convinced to change my mind. As provided, the data does not support this model.

We have now clarified in the manuscript that we do not discuss the liquid-gas critical point.

Reviewer #2:

This is a potentially interesting study addressing a possible scale-invariant log-normal characteristic of droplet size distribution in the phase separation behavior of biomolecular condensates. Some of the data presented are valuable and intriguing. However, as it stands, the validity and utility of this study are uncertain because there are serious deficiencies in the execution and presentation of the authors' results. Many of these shortcomings are fundamental, including a lack of clarity in the basic conceptual framework of the study, insufficient justification of the experimental setup, less-than-conclusive experimental evidence, and inadequate discussion of implications of the authors' findings to future experimental and theoretical studies of biomolecular condensates. Accordingly, this reviewer considers that the manuscript should undergo a major revision to address the following. In particular, the discussion should be significantly expanded by including references mentioned below as well as other references pertinent to the issues raised.

We thank the reviewer for the helpful comments. In the revised version of the manuscript we clarified that we aimed to use a well-established tool – the scaling analysis – to study phase transition and applied to the protein condensation process. This approach offers insight into a universal aspect of protein phase separation, and also provides a practical approach to determine the phase boundary. The observed fat-tailed distribution of protein droplet sizes is not what is normally observed in more standard phase separation systems in the subsaturated phase. Our contribution is not to propose a theoretical model, but rather to report the observation of a scaling behaviour.

(1) The theoretical analysis in this study is based on experimental data on condensed droplet size distributions for FUS and α -synuclein. The size data for FUS droplet is indirect as it relies on the assumption that FUS droplet diameter is proportional to fluorescence intensity of labeled FUS (page 10 of manuscript), with fluorescence data adopted from a previously published work by another group (Kar et al. & Pappu, ref.27). Because fluorescence of a droplet is expected to be dependent upon the condensed-phase concentration of FUS, this proportional relationship, even if it holds, must also be modulated by FUS concentration in the droplet. Moreover, why should fluorescence be proportional to diameter but not the cross-sectional area or volume of the FUS droplet, which would be more intuitive? These issues should be clarified. A new measure by

microscopy is used to determine the size distribution of condensed α -synuclein; but no microscopy image is shown. It is of critical importance that such raw data (for example microscopy images) be presented for the completeness and reproducibility of the experiment because the entire study relies on the soundness of these experimental measurements.

As we mentioned in the article, for the scaling analysis, the droplet dimensions could be assessed in 1D (length), 2D (area) or 3D (volume). For the FUS experiments, we used the data as the authors provided in the original publication (PNAS 2022). For alpha-synuclein, we provided the data in the article.

(2) Despite the authors' claim of a universal scaling relationship, the log-log scatter plots in Figure 1 (page 15 of the manuscript) exhibit significant deviations from linearity at low protein concentrations ($p \rightarrow 0$). Given this fact, is universal scaling really valid? Discussion of this behavior is conspicuously absent (except the statement that these data points are excluded in the fit). In any case, the possible origins of these deviations should be thoroughly discussed so that the regime of universal scaling can be properly delineated.

In general, one would expect the scaling ansatz to be valid close to the phase boundary. It is the feature of the ansatz, that further away from the boundary, deviations are expected because of the decreasing relevance of critical phenomena.

(3) Droplet size distribution most likely depends on the time duration after the preparation of the sample. For α -synuclein, "liquid droplet size characterisation images were captured 10 minutes post-liquid droplet formation" (page 9 of the manuscript). Why 10 minutes? Have the authors tried imaging at different time points and, if so, do the distributions at different time points remain essentially the same? If they are different, what is the criterion for focusing only on a particular time point? Information related to these questions should be provided.

We have now determined the droplet size distribution of alpha-synuclein at different time points, finding that they are not dependent on time within experimental uncertainties (Figure 6 in the revised manuscript).

(4) At least two well-known mechanisms can lead to the time-dependent distribution of liquid droplet sizes: (i) coalescence of droplets in spatial proximity to form a larger droplet, and (ii) Ostwald ripening, i.e., formation of larger droplets concomitant with the dissolution of smaller droplets without fusion of droplets. The implications of these mechanisms on the authors' droplet size distributions should be addressed. Indeed, maintaining a size distribution against these mechanisms in vivo often requires active suppression [Bressloff, Phys Rev E 101, 042804 (2020)] with possible involvement of chemical reactions [Kirschbaum & Zwicker, J R Soc Interface 18, 20210255 (2021)]. These considerations are central to the basic rationale of this study and therefore should be carefully tackled.

These two mechanism of growth are relevant above the critical concentration. Below the critical concentration, which is the regime that we investigated in our work, there is no need of active suppression.

(5) If coalescence and/or Ostwald ripening do occur, given sufficient time after sample preparation, the condensed phase may become a single large "droplet" or a single liquid layer. Does this occur in the authors' experiments?

As we are below the critical concentration, this is unlikely to occur, as indeed supported by the experiments mentioned at point (3).

(6) It is unclear whether the authors aim to address the kinetic phenomenon of liquid droplet formation and evolution or equilibrium properties. The two types of phenomena appear to be conflated in the authors' narrative. Clarification is needed. If this work aims to address timeindependent (or infinite-time) equilibrium properties, how are they expected to be related to droplet size distribution, which most likely is time-dependent?

Our analysis focuses on the equilibrium properties of the droplet size distribution below the critical concentration, and it should guide the proposal of a theoretical model that explains the emergence of scaling. In the introductory part of our manuscript, we proposed a possible scenario that tries to extend the Flory-Huggins's theory to predict a scaling behaviour appropriate to a critical transition. Other scenarios are possible, and our result along with further experiments are needed to arrive at a deeper understanding of protein aggregation.

(7) The relationship between the potentially time-dependent droplet size distribution and equilibrium properties of p_t and p_c (transition and critical concentrations, respectively) should be better spelled out. An added illustrative figure will be helpful.

We are addressing equilibrium properties, not kinetic ones. See also the answers to point 6.

(8) The authors comment that their findings appear to be inconsistent with Flory-Huggins theory because Flory-Huggins "characterizes droplet formation as a consequence of nucleation ..." (page 8 of the manuscript). Here, three issues need detailed clarification: (i) In what way does Flory-Huggins mandate nucleation? (ii) Why are the findings of apparent scale invariance inconsistent with nucleation? (iii) If liquid droplet formations do not arise from nucleation, what physical mechanism(s) is (are) envisioned by the authors to be underpinning the formation of condensed liquid droplets in protein phase separation?

We do agree that the Flory-Huggins theory does not mandate nucleation above the spinodal line. However, we are addressing the equilibrium properties below the critical concentration, so the stable phase is the dilute phase, and there is no nucleation.

(9) Are any of the authors' findings related to finite-system effects of phase separation [see, e.g., Nilsson & Irback, Phys Rev E 101, 022413 (2020)]?

Our experimental system is macroscopic, so we would not expect finite size effects.

(10) Since the authors are using their observation of an apparent scale-invariant droplet size distribution to evaluate phase separation theory, it is important to clarify whether their findings provide any constraint on the shape of coexistence curves (phase diagrams).

We are only reporting the phenomenological observation of a scaling behaviour, so we may not speculate at this stage on the constraints of the coexistence curves. This is indeed an exciting opportunity for future studies.

(11) More specifically, do the authors' findings suggest that the phase diagrams predicted by Flory-Huggins are invalid? Or, are they suggesting that even if the phase diagrams predicted by Flory-Huggins are empirically correct (if verified by experimental testing), they are underpinned by a free energy function different from that of Flory-

Huggins? It is important to answer this question to clarify the implications of the authors' findings on equilibrium phase behaviors and the falsifiability of the implications.

As mentioned above, our main conclusion is that the droplet size distribution follows a scaling behaviour. Our contribution is not to propose a theoretical model, but rather to propose a scaling behaviour that should be accounted for by existing or future theoretical models.

(12) How about the implications of the authors' findings on other theories of protein phase separation that are based on interactions that are different from the short spatial range interactions treated by Flory-Huggins? For instance, it has been observed that whereas the Flory-Huggins-predicted phase diagrams always convex upward, phase diagrams for charged intrinsically disordered proteins with long spatial range Coulomb interactions exhibit a region that concave upward [Das et al., Phys Chem Chem Phys 20, 28558-28574 (2018)]. Can information be provided by the authors' findings regarding apparent scale-invariant droplet size distribution on the underlying interaction driving the protein molecules toward phase separation?

This is an interesting point for future studies about the type of interactions that give rise to the observed scaling behaviour.

(13) Table S1 (page 4) and Table S2 (page 7) are mentioned in the text but these tables are not in the submitted files.

We have added the Supplementary Tables as well as the source files for the figures.

(14) The two systems studied (FUS and α -synuclein) have a single intrinsically disordered protein (IDP) component. It is not clear if the authors expect their claimed scaling relation to be applicable to systems with multiple IDP components and if so, why.

From the data that we have currently analysed, we feel that we may not speculate on this interesting point, leaving it to future studies.

<https://doi.org/10.7554/eLife.94214.2.sa0>