

Reviewed Preprint

v1 • September 1, 2025

Not revised

Reviewed Preprint

v2 • November 24, 2025

Revised by authors

For correspondence:

lmahadev@g.harvard.edu

Competing interests:

No competing interests declared

Reviewing editor:

Saad Jbabdi,
University of Oxford, United Kingdom

© 2025, Yin et al. This article is distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Morphogenesis and morphometry of brain folding patterns across species

Sifan Yin¹, Chunzi Liu¹, Gary PT Choi², Yeonsu Jung¹, Katja Heuer³, Roberto Toro³, L Mahadevan^{1,4,5}✉

¹School of Engineering and Applied Sciences, Harvard University, Cambridge, United States • ²Department of Mathematics, The Chinese University of Hong Kong, Hong Kong, China • ³Institut Pasteur, Université Paris Cité, Unité de Neuroanatomie Appliquée et Théorique, Paris, France • ⁴Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States • ⁵Department of Physics, Harvard University, Cambridge, United States

eLife Assessment

This **important** study presents a cross-species and cross-disciplinary analysis of cortical folding. The authors use a combination of physical gel models, computational simulations, and morphometric analysis, extending prior work in human brain development to macaques and ferrets. The findings support the hypothesis that mechanical forces driven by differential growth can account for major aspects of gyrification. The evidence presented is overall strong and **convincingly** supports the central claims; the findings will be of broad interest in developmental neuroscience.

<https://doi.org/10.7554/eLife.107138.2.sa3>

Abstract

Evolutionary adaptations associated with the formation of a folded cortex in many mammalian brains are thought to be a critical specialization associated with higher cognitive function. The dramatic surface expansion and highly convoluted folding of the cortex during early development is a theme with variations that suggest the need for a comparative study of cortical gyrification. Here, we use a combination of physical experiments using gels, computational morphogenesis, and geometric morphometrics to study the folding of brains across different species. Starting with magnetic resonance images of brains of a newborn ferret, a fetal macaque, and a fetal human, we construct two-layer physical gel brain models that swell superficially in a solvent, leading to folding patterns similar to those seen *in vivo*. We then adopt a three-dimensional continuum model based on differential growth to simulate cortical folding *in silico*. Finally, we deploy a comparative morphometric analysis of the *in vivo*, *in vitro*, and *in silico* surface buckling patterns across species. Our study shows that a simple mechanical instability driven by differential growth suffices to explain cortical folding and suggests that variations in the tangential growth and different initial geometries are sufficient to explain the differences in cortical folding across species.

Introduction

Although not all brains are folded, in many mammals, the folded cerebral cortex is known to be critically important for brain cognitive performance and highly dependent on the hierarchical structure of its morphology, cytoarchitecture, and connectivity [Gautam et al. \(2015\)](#) [Suárez et al. \(2020\)](#) [Pang et al. \(2023\)](#). Brain function is thus related both to the topological structure of neural networks [Bullmore and Bassett \(2011\)](#), as well as the geometry and morphology of the convoluted cortex [Kriegeskorte and Wei \(2021\)](#), both of which serve to enable and constrain neuronal dynamics [Pang et al. \(2023\)](#). Across species, cortical morphologies show a large diversity, as shown in [Fig. 1\(a\)](#) [Takahata et al. \(2012\)](#) [Heuer et al. \(2019\)](#). And within our own species and in model organisms such as the ferret used to study the genetic precursors of

misfolding, cortical folding and misfolding are known to be markers of healthy and pathological neurodevelopment, disease and aging [Hutton et al. \(2009\)](#); [de Moraes et al. \(2024\)](#) (see also **Fig. S1** [Oegema et al. \(2020\)](#); [Choi et al. \(2025\)](#)). Thus a comparative study of cortical folding is essential for understanding brain morphogenesis and functionalization across evolution [Pang et al. \(2023\)](#); [Schwartz et al. \(2023\)](#), during development as well as in pathological situations associated with disease.

The development of cortical morphology involves the coordinated and localized expression of many genes that lead to the migration and differentiation of neural stem and progenitor cells [van der Meer and Kaufmann \(2022\)](#); [Oegema et al. \(2020\)](#); [Del-Valle-Anton and Borrell \(2022\)](#). All these biological processes cooperatively generate an expansion of the cortex relative to the underlying white matter and eventually drive cortical folding [Akula et al. \(2023\)](#). While a range of mechanisms have been proposed in the past for the processes leading to folding [Striedter et al. \(2015\)](#); [Holland et al. \(2015\)](#); [Van Essen \(2020\)](#), over the past decade, theoretical and experimental evidence have converged on the primary determinant of folding as a mechanical instability associated with the formation of a localized crease or sulcus [Hohlfeld and Mahadevan \(2011, 2012\)](#); [Tallinen et al. \(2013\)](#) driven by differential growth, with iterations and variations that qualitatively explain the brain gyrification [Tallinen et al. \(2014, 2016\)](#). However, combining this mechanistic model with a comparative perspective that aims to quantify the variability of folding patterns across species, or linking it to genetic perturbations that change the relative expansion of the cortex remain open questions. In a companion paper [Choi et al. \(2025\)](#), we address the latter using the ferret as a model organism, while in the current study, we address the former question using a combination of physical experiments with gel swelling, numerical simulations of differential growth and geometric morphometrics to compare brain morphogenesis in the ferret, the macaque, and the human. The species were selected as they have very different brain sizes and folding patterns (**Fig. 1(a)**) and thus represent different branches in the evolutionary tree, Carnivora, Old World monkeys, and Hominoidea. Furthermore, in all three species, we have access to a time course of the development of the folds, as shown in **Fig. 1(a)**.

Experiments on swelling gel-brains

To mimic the mechanical basis for brain morphogenesis based on the differential growth of the cortex relative to the white matter, we used the swelling of physical gels that mimic the fetal brain developmental process during post-gestation stages. Previously, this principle has been demonstrated for *Homo sapiens* (human) brain development [Tallinen et al. \(2016, 2014\)](#). To demonstrate that the same principle applies to other species, we constructed physical gels from the fetal brain MRI scans for *Macaca mulatta* (rhesus macaque) and *Mustela furo* (ferret). In brief, a two-layer PDMS gel is constructed from the 3D fetal brain MRI reconstruction and immersed in an organic solvent. Immersion causes solvent imbibition into the surface of the physical gel which swells leading to a compressive strain in the outer layers that causes the surface layer to form convolutions that resemble the folding patterns in the brain cortex layer. Time-lapse images of the gel model to mimic brain folding in *Macaca mulatta* (rhesus macaque) up to G110 are shown in **Fig. 2(a)**, while in **Fig. 2(a)**, we show the initial and final states of swelling to mimic different post-gestation stages corresponding to G85, G110 and G135. A visual inspection of the swollen gels constructed from different post-gestation stages showed qualitatively different folding patterns, indicating the sensitivity of the folds to the initial undulations present on the physical gel surfaces. In **Fig. 2(a)** and Movie S1, we show the results of similar physical gel experiments to mimic brain folding morphogenesis for *Homo sapiens* [Tallinen et al. \(2016\)](#) and *Mustela furo* [Choi et al. \(2025\)](#); in each case the initial condition was based on 3d fetal brain MRIs and the final state was determined qualitatively using the overall volume of the brain relative to the initial state. No attempt was made to vary the swelling ratio of the surface as a function of location, although it is likely that in the different species this was not a constant. To quantitatively describe the folding patterns, the swollen gel surfaces were then scanned and reconstructed using X-ray Computed Tomography (Methods). Movie S2–S4 showed the fetal brain MRI scans and the reconstructed 3D swollen gel surfaces in juxtaposition for *H. sapiens*, *M. furo*, and *M. mulatta*. This paves the way for




Figure 1. The diversity of the cortical morphologies and developmental processes across species.

(a) Phylogenetic relationship of species. Adapted from Takahata et al. (2012) [DOI](#); Heuer et al. (2019) [DOI](#). Typical real brain surfaces of ferret and primates are presented. Color represents mean curvature. Scale bars: 1 cm (estimated from Herculano-Houzel (2009) [DOI](#)). (b) Stained sections of mature brain tissue from ferret, rhesus macaque, and human. Scale bar: 10 mm. Adapted from Rea (2016). (c-e) 3D reconstruction of cortical surfaces of ferret, macaque, and human brains from fetal to adult. (c) Ferret: postnatal day 4, 10, 17 and adult maturation Barnette et al. (2009) [DOI](#). Scale bar: 1 cm; (d) Macaque: gestation day 85, 110, 135 Liu et al. (2020) [DOI](#), and adult maturation Calabrese et al. (2015) [DOI](#). Scale bar: 1 cm; (e) Human: gestation day 175 (week 25), 210 (week 30), 231 (week 33), 273 (week 39), and adult maturation Barnette et al. (2009) [DOI](#). Scale bar: 5 cm.

a quantitative comparison of the results of the physical experiments with those derived from a mechanical theory for brain morphogenesis and those from scans of macaque, ferret and human brains.

Simulations of growing brains

To test the capability of the mechanical model for brain morphogenesis based on differential growth [Tallinen et al. \(2014\)](#) [Budday et al. \(2014\)](#) [Tallinen et al. \(2016\)](#) to explain patterns across species, we perform numerical simulations of the developing brains of ferret, macaque, and human modeled as soft tissues. Here we only consider the simplest homogeneous growth profile which is sufficient to capture the folding formation across different species; regional growth of the cortical layer based on real data from tracking the surface expansion of fetal brains [Garcia et al. \(2018\)](#) [Alenyà et al. \(2022\)](#) [Weickenmeier \(2023\)](#) is a more sophisticated alternative that we do not adopt for reasons of simplicity.

The initial brain models are reconstructed from 3D fetal brain MRI (*Methods*), and assumed to be composed of gray and white matter layers which are considered as hyperelastic materials with differential tangential growth ratio g . A multiplicative decomposition of deformation gradient gives $\mathbf{F} = \mathbf{A} \cdot \mathbf{G}$ with $\mathbf{A}$ the elastic part and $\mathbf{G} = \sqrt{g}\mathbf{I} + (1 - \sqrt{g})\mathbf{n} \otimes \mathbf{n}$ the growth tensor. We adopt a modestly compressible neo-Hookean material with strain energy density

$$W = \frac{\mu}{2} \left[J_A^{-2/3} \text{tr}(\mathbf{A}^T \cdot \mathbf{A} - 3) \right] + \frac{K}{2} (J_A - 1)^2, \quad (1)$$

where $\mathbf{F}$ is the deformation gradient, $J_A = \det \mathbf{A}$, μ is the initial shear modulus, and K is the bulk modulus. Considering modestly compressible material, we assume $K = 5\mu$.

We solve the final shapes of the growing tissues using a custom finite element method [Tallinen et al. \(2016\)](#). All the initial geometries of smooth fetal brains are obtained from open data sources (*Methods*, 3D model reconstruction), and the growth ratio distribution is assumed as a function of the initial location, including the distance to the cortical surface and the presumably non-growing regions. Other parameters, such as the thickness ratio and modulus ratio of gray and white matter, and the temporal changes of growth ratios are assumed (*Methods*, numerical simulations, **Table 2**). The brain models are discretized to tetrahedrons by Netgen. An explicit algorithm is adopted to minimize the total strain energy of the deforming tissues. We adopt a step-wise simulation strategy where the initial geometry of each presumed stress-free state is obtained from real MRI data of earlier-stage fetal brains, instead of using a continuous model where only the initial smooth brain geometry is input [Tallinen et al. \(2016\)](#). The simulated developmental processes of fetal brains are presented in **Fig. 3** and Movie S1.

Morphometric analysis

To verify whether our simulation methods and physical gel models are sufficient to capture the developmental process of fetal brains and reproduce cortical patterns in adult brains, we compare the cortical surfaces of real, simulated, and gel brains across different species. **Fig. 4(a)** presents the hemispherical cortical surfaces of the real, simulated, and gel brains (denoted as $\mathcal{S}_1$, $\mathcal{S}_2$, and $\mathcal{S}_3$, respectively). Left and right symmetry and the comparison of whole brains are presented in **Fig. S3** and S4, Movie S2–S4. Major sulci are extracted and highlighted by hand for further alignment. To analyze the shape differences, we then adapted the parameterization methods in [Choi and Lui \(2015\)](#) [Choi et al. \(2015, 2020\)](#) to map the brain surfaces onto a common disk shape with the major sulci aligned using landmark-matching disk quasi-conformal parameterizations. Denote the three disk parameterization results as $\mathcal{D}_1$, $\mathcal{D}_2$, $\mathcal{D}_3$, as shown in **Fig. 4(a)**. Multiple quantitative measures such as surface area, cortical thickness, curvature, and sulcal depth can be adopted to analyze curved surfaces and to compare different surfaces. Here, we use shape index (SI) and rescaled mean curvature $\tilde{H}$ to describe the similarity among real,

Figure 2. Physical gel model that recapitulates the growth-driven morphogenesis mechanism across phylogeny and developmental stages.

(a) A time-lapse of the physical gel brain mimicking macaque brain development starting from G110. (b) Left views of three physical gels mimicking macaque post gestation day 85, day 110, and day 135 before and after hexane swelling. Scale bar: 1 cm. (c) Comparison of fetal/newborn brain scans and the reconstructed surfaces of swollen physical gels for various species. Scale bars: 1 cm.





Figure 3. Simulations of growing brains of (a) ferret, (b) rhesus macaque, and (c) human.

Starting from smooth fetal/newborn brains, simulations show different gyrification patterns across species. The brains are modeled as soft elastic solids with tangential growth in the gray matter (see *Simulations of growing brains* for details). Initial 3D geometries are taken from the reconstruction of MRI (see *Methods*, 3D model reconstruction). Mechanical parameters of growth ratio and cortical thickness are provided in [Table 2](#). Color from dark to light blue represents shape index (as defined in [Eq. \(2\)](#)) from -1 to 1.



simulated, and gel surfaces. Shape index is a dimensionless and scale-independent surface measure Koenderink and van Doorn (1992). It can be calculated from the mean and Gaussian curvatures by

$$SI = \frac{2}{\pi} \arctan \left(\frac{H}{\sqrt{H^2 - K}} \right). \quad (2)$$

Shape index ranging within $[-1, 1]$ defines a continuous shape change from convex, saddle, to concave shapes. **Fig. S2** shows nine categories of typical curved surfaces. For brain surfaces, we can classify sulcal pits ($-1 < SI < -0.5$), sulcal saddles ($-0.5 < SI < 0$), saddles ($SI = 0$), gyral saddles ($0 < SI < 0.5$) and gyral nodes ($0.5 < SI < 1$). When the shape index equals -1 or 1 , it represents a defect of the curvature tensor with two eigenvalues identical, as shown in **Fig. S3**. The probability of shape index distribution exhibits two peaks, as shown in **Fig. 4(a)**, corresponding to ridge and rut shapes ($SI = \pm 0.5$), where the ridge shape ($SI = 0.5$) is dominating. In contrast, the rescaled mean curvature histogram exhibits a unique peak around 0.2 (**Fig. S3**). The two-peak and unique-peak distributions of adult human brain surfaces have also been presented in previous research Demirci and Holland (2022); Hu et al. (2013). To quantify the similarities between every two brain surfaces, we evaluate the distribution of $I(v)$ differences on the common disk domain at each vertex v . Here $I(v)$ represents either the surface shape index $SI(v)$ or the rescaled mean curvature $\tilde{H}(v)$ defined as

$$\tilde{H}(v) = 2 \frac{H(v) - \min(H)}{\max(H) - \min(H)} - 1. \quad (3)$$

Thus the similarity s of the distributions I_{S_1}, I_{S_2} of the two surfaces S_1, S_2 is then evaluated via

$$s(S_1, S_2) = 1 - \frac{1}{2m^{1/p}} \left\| I_{S_1} - I_{S_2}(g_2^{-1} \circ f_{12} \circ g_1) \right\|_p, \quad (4)$$

where m is the total number of vertices, $g_1: S_1 \rightarrow \mathcal{D}_1$ and $g_2: S_2 \rightarrow \mathcal{D}_2$ are the initial disk conformal parameterizations, f_{12} is the landmark-aligned quasi-conformal map between $\mathcal{D}_1$ and $\mathcal{D}_2$, and $\|\cdot\|_p$ is the vector p -norm:

$$\left\| I_{S_1} - I_{S_2}(g_2^{-1} \circ f_{12} \circ g_1) \right\|_p = \left(\sum_{k=1}^m \left| I_{S_1}(v_k) - I_{S_2}((g_2^{-1} \circ f_{12} \circ g_1)(v_k)) \right|^p \right)^{1/p}. \quad (5)$$

Note that since $I_{S_1} \in [-1, 1]$ and $I_{S_2} \in [-1, 1]$, we have $0 \leq \left\| I_{S_1} - I_{S_2}(g_2^{-1} \circ f_{12} \circ g_1) \right\|_p \leq (2^p m)^{1/p} = 2m^{1/p}$.

Therefore, we always have $0 \leq s(S_1, S_2) \leq 1$.

We calculate the similarity indices with different p -norm: $p = 1$, $p = 2$, and $p = \infty$ of both rescaled mean curvature and shape index. The results are given in **Table 1** and Table S2.

Discussion

In this study, we have explored the mechanisms underlying brain morphogenesis for a few different mammalian species. Using fetal and adult brain MRIs for ferrets, macaques and humans, we carried out physical experiments using swelling gels, combined with a mathematical framework to model the differential growth of the cortex that leads to iterations and variations of a mechanical (sulcification) instability that allowed us to recapitulate the basic morphological development of folds. We then deployed a range of morphometric tools to compare the results of our physical experiments and simulations with 3d scans of real brains, and show that our approaches are qualitatively and quantitatively consistent with experimental observations of brain morphologies.

All together, our study shows that differential growth between the gray matter cortex and white matter bulk provides a minimal physical model to explain the variations in the cortical folding patterns seen in multiple species. More specifically, we see that the overall morphologies are controlled by the relative size of the brain (compared to the cortex), as well as the scaled surface

Figure 4. Comparison among real (δ_1), simulated (δ_2), and gel brains (δ_3) of ferret, rhesus macaque, and human via morphometric analysis.

(a) 3D cortical surfaces of *in vivo*, *in vitro* and *in silico* models. Left brain surfaces are provided here. The symmetry of the left and right halves of the brain surfaces is discussed in Figs. S3 and S4, Movie S2–S4. Three or four major folds of each brain model are highlighted and served as landmarks. The occipital pole region of macaque brains remains smooth in real and simulated brains. (b) The quasi-conformal disk mapping with landmark matching of cortical surfaces on disk (see Sec Morphometric analysis for details). Blue or red curves represent corresponding landmarks. Color represents shape index (SI, as defined in Eq. (2)). Similarity indices of each simulated and gel brain surfaces are presented in Table 1. (c) Histogram of shape index of ferret, macaque, and human. Black, red, and blue dots represent the probability of shape index of real, gel, and simulated surfaces, respectively.




Table 1.

Similarity index evaluated by comparing the shape index of simulated brains (S), swollen gel brain simulacrum (G) and real brain surfaces (R), calculated with vector p-norm $p = 2$, as described in Eq. (4).

Similarity Index	simulation-real	gel-real	simulation-gel
Ferret (left)	0.7632	0.7307	0.7117
Ferret (right)	0.7339	0.7222	0.7019
Macaque (left)	0.7624	0.7608	0.7512
Macaque (right)	0.7559	0.7632	0.7463
Human (left)	0.6947	0.7425	0.7000
Human (right)	0.6909	0.7325	0.7001

expansion rate, both of which can and do have multiple genetic antecedents [Bae et al. \(2014\)](#); [Shinmyo et al. \(2017\)](#); [Qi et al. \(2023\)](#); [Lai et al. \(2003\)](#); [Barresi et al. \(2024\)](#) (see SI, Table S1).

Our results point to some open questions. First, the relationship between physical processes that shape organs and the molecular and cellular processes underlying growth has been the subject of many recent studies, e.g. in the context of gut development [Gill et al. \(2024a,b\)](#), and it would seem natural to expect similar relationships in brain development. There is a growing literature linking genes with brain malformation and pathologies. For example, *GPR56* [Bae et al. \(2014\)](#) and *Cdk5* [Shinmyo et al. \(2017\)](#) can affect progenitors and neurons in migration, *SP0535* [Qi et al. \(2023\)](#) can affect neural proliferation, and *foxp2* can affect neural differentiation [Lai et al. \(2003\)](#); [Barresi et al. \(2024\)](#), all of which change the cortical expansion rate and thickness, consequently leading to brain malformation and pathologies, as listed in Table S1. While a direct relation between gene expression levels and the effective tangential growth rate G and cortical thickness has only been partially resolved, as for example in our companion study on the folding and misfolding of the ferret brain [Choi et al. \(2025\)](#), further studies are needed to address how genetic programs drive cell proliferation, migration, size and shape change that ultimately lead to different cortical morphologies. Second, although our focus has been exclusively on the morphology of the brain, recent studies [Pang et al. \(2023\)](#); [Schwartz et al. \(2023\)](#) are suggestive of a link between cortical geometry and function from both developmental and evolutionary perspectives, and suggest natural questions for future study. Third, despite prescribing a simple spatially homogeneous form for the cortical expansion for fetal brain surfaces of all three species studied, we were able to capture the essential features of the folds and variations therein. The effect of spatio-temporally varying inhomogeneous growth needs to be further investigated by incorporating regional growth of the gray and white matter not only in human brains [Garcia et al. \(2018\)](#); [Weickenmeier \(2023\)](#) based on public datasets [Namburete et al. \(2023\)](#), but also in other species to investigate folding differences across species, inter-individual variability and finally regional differences in folding. More accurate and specific work is expected to focus on these directions. Finally, our physical and computational models along with our morphometric approaches are a promising avenue to pursue in the context of the inverse growth problem, i.e., postulate the fetal brain morphologies from the adult brains. This may one day soon allow us to reconstruct the adult brain geometries from fossil endocasts [de Sousa et al. \(2023\)](#), and eventually provide insights into how a few mutations might have triggered the rapid expansion of the cortex across evolutionary time and led to the highly convoluted human brain that is able to ponder the question of how it might have folded itself.

Methods

3D model reconstruction

Pre-processing

We used a publicly available database for all our 3d reconstructions: fetal macaque brain surfaces are obtained from [Liu et al. \(2020\)](#); newborn ferret brain surfaces are obtained from [Choi et al. \(2025\)](#); and fetal human brain surfaces are obtained from [Tallinen et al. \(2016\)](#). These 2D manifolds of brain surfaces were first normalized by their characteristic lengths $L_0 = L_x/2$ and discretized to triangle meshes by Meshlab [Cignoni et al. \(2008\)](#) and then converted to 3D models and discretized to tetrahedral elements in Netgen Net (2019).

Post-processing

All the numerically calculated brain surfaces and scanned gel brain surfaces were extracted as 3D triangle meshes. These intact cortical surfaces were then dissected to left and right semi-brains and normalized by half of their longitudinal lengths $L = L_x/2$. These surfaces were then checked and fixed to a simply connected open surface to satisfy the requirements for open disk conformal mapping [Choi and Lui \(2015\)](#).

Experimental protocol for gel experiments

The physical gel was constructed following a previous publication Tallinen et al. (2016). In brief, a negative rubber mold was generated with Ecoflex 00-30 from a 3D-printed fetal brain plastic model. The core gel was then generated with SYLGARD 184 at a 1:45 crosslinker: base ratio. Three layers of PDMS gel at a 1:35 crosslinker:base ratio were surface-coated onto the core layer to mimic the cortex layer. Pigments were added to the PDMS mixture for bright-field visualization. To mimic the cortex folding process, the physical gel was immersed in n-hexane. The time-lapse videos were taken with an iPod Touch 7th Gen. To reconstruct the swollen gel surface and analyze the folding patterns, swollen gels were imaged with a Nikon X-Tek HMXST X-ray CT machine. The voxel resolution for all scans was 100 μm . To minimize solvent evaporation during the 30-minute scan, cotton soaked in the solvent was placed inside the container. The container's thin acrylic walls allowed for clear X-ray transmission. The container's thin acrylic walls allowed for clear X-ray transmission. To test the reversibility of the folding pattern formation, the physical gel models were allowed to dry in a laminar flow hood overnight before being immersed in hexane again.

3D reconstruction of swollen gel surface from X-ray CT

The z-stack images obtained from the X-ray CT machine were segmented by a machine-learning-based segmentation toolbox, Labkit, via ImageJ. A classification was created for each swollen gel. Then a 3D surface of the segmented gel image was generated by ImageJ 3D viewer. Further postprocessing was conducted in SOLIDWORKS and MeshLab, including reversing facial normal direction, re-meshing, and surface-preserving Poisson smoothing.

Numerical Simulations

The simulation geometries of ferret, macaque, and human are based on MRI of smooth fetal or newborn brains. For the ferret, we take the P0, P4, P8, and P16 fetal brains as initial shapes of the step-wise growing model; for the macaque, we take G80 and G110 fetal brains as initial shapes of the step-wise growing model. To focus on fold formation, we did not consider the lack of patterning in the relatively smooth regions, such as the Occipital Pole of the macaque; for the human, we take the GW22 fetal brain as the initial shape of the continuous growing model. Small perturbations of the initial geometry typically affect only the minor folds, while the main features of the major folds, such as their orientation, width, and depth, are well conserved across individuals Bohi et al. (2019); Wang et al. (2021). For simplicity, we do not perturb the fetal brain geometry obtained from datasets. Both gray and white matter are considered as neo-Hookean hyperelastic material with the shear modulus distribution

$$\mu(d) = \mu_w + \frac{\mu_g - \mu_w}{1 + \exp[10(d/h - 1)]}, \quad (6)$$

where d is the distance from an arbitrary material point inside the brain to the cortical surface, and h is the approximated cortical thickness assumed to decrease with growth process time t Tallinen et al. (2016). The subscripts 'g' and 'w' represent gray and white matter, respectively. The tangential growth ratio g has a similar spatial distribution

$$g(d) = g_w + \frac{g_g - g_w}{1 + \exp[10(d/h - 1)]}, \quad (7)$$

where g_w and g_g represent the growth ratio at cortical surface and at innermost white matter. The parameters used in simulations are listed in Table 2.

species	model	growth ratio g_g/g_w	stiffness ratio μ_g/μ_w	normalized cortical thickness h/L
Ferret	step-wise	1.8	1	$0.1 - 0.005t$
Macaque	step-wise	1.8	1	$0.1 - 0.05t, 0.1 - 0.1t$
Human	continuous	1.8	1	$0.05 - 0.03t$

Table 2. Parameters for numerical simulations.

Supplementary Text

Malformations of human cortical development

In the main text, we have shown examples of normal cortical development and the resulting folding processes by following fetal and adult animals across a few different species. Defects in neuronal migration disorders may lead to a group of rare brain malformations, the most severe forms being lissencephaly and polymicrogyria, as shown in **Fig. S1(a)** (15). Lissencephaly, or agyria-pachygyria, is characterized by a simplified convolutional pattern where a few broad gyri are separated by rudimentary primary fissures and sulci. Lissencephaly is accompanied by a very thick cortical gray matter layer (**Fig. S1(a)**, middle) which has been confirmed as a critical cause in this malformation (3,18). Polymicrogyria, on the contrary, is an overly convoluted cortex with a reduced cortical thickness and an increasing number of secondary folds (**Fig. S1(a)**, right). Various malformations of cortical development are congenital and genetically heterogeneous diseases which mutations or deletions of genes have been identified. For example, a regional deletion mutation in a regulatory element of GP56 can selectively disrupt the human cortex surrounding the Sylvian fissure bilaterally including “Broca’s area”. **Fig. S1(a)** shows the MRI of polymicrogyria of a noncoding mutation in the GPR56 gene (1). This abnormally thin cortex is folded giving a paradoxical but characteristic thickened appearance. Many other genes have also been found associated with brain malformation and pathologies, such as *foxp2* (2,11), *SP0535* (16) and *Cdk5* (17). These genes affect progenitors and neurons in migration, proliferation, and differentiation which change the cortical expansion rate and thickness, consequently leading to brain malformation and pathologies, as listed in Table S1.

Statistical characteristics of brain folding

To characterize the form of the convolutions on the surface of the brain, we use a dimensionless and scale independent surface measure, the shape index (SI), to quantify 3D cortical morphologies across species. The definition of shape index is given as (9)

$$SI = \frac{2}{\pi} \arctan \left(\frac{H}{\sqrt{H^2 - K}} \right), \quad (S1)$$

or equivalently,

$$SI = \frac{2}{\pi} \arctan \left(\frac{k_{\min} + k_{\max}}{k_{\min} - k_{\max}} \right), \quad (S2)$$

where $k_{\max}$ and $k_{\min}$ are the maximum and minimum curvatures, respectively, $k_{\max, \min} = H \pm \sqrt{H^2 - K}$. The shape index is a continuous number representing the local shape of a surface.

Fig. S2 illustrates a shape index scale divided into nine categories: spherical cup, trough rut, saddle rut, saddle, saddle ridge, ridge, dome and spherical cap, with the color representing the shape index value ranging from -1 to 1.

In the main text, we have shown the shape index distribution of left semi-sphere brains. Here, we present the results of both left and right semi-spheres of ferret, macaque and human brains in **Fig. S3** (top two rows) and **Fig. S4** with comparison among the real (black dots), gel (red dots), and simulated (blue dots) surfaces. The probability of shape index distribution exhibits two peaks,

Gene	Change in geometry or physical properties	Malformations and dysfunctions
<i>foxp2</i> (2, 11)	Reduced gray matter density (thinner cortex) in Broca's area	Polymicrogyria, speech and language disorder
SP0535 (16)	Increased expansion of neural progenitors	improved cognitive ability and working memory
<i>Cdk5</i> (13, 17)	Abnormalities in neuronal migration and organization	Lissencephaly with cerebellar hypoplasia (LCH)
<i>GPR56</i> (1, 12)	Neural progenitors migration	Bilateral frontoparietal polymicrogyria
PAFAH1B1, RELN, TUB1A1, DCX, ARX, NDE1 (10, 19)	Increased cortical thickness	Lissencephaly

Table S1. Gene-related brain properties and malformation.

corresponding to ridge and rut shapes ($SI=\pm 0.5$), where the ridge shape ($SI=0.5$) dominates. In contrast, the rescaled mean curvature histogram exhibits a unique peak around 0.2 (**Fig. S3** [↗](#), the 3rd and 4th rows). These histograms illustrate that shape index and mean curvature exhibit qualitatively different distributions and both of these should be taken into account when describing cortical surface characteristics, consistent with earlier studies (7, 8).

Morphometric analysis

For the morphometric analysis, we developed a landmark-matching disk quasi-conformal parameterization method by extending and combining the methods in (5, 6) and applied it to map the real, simulated, and gel brain surfaces onto the unit disk with major sulci aligned.

More specifically, we denote the real, simulated, and gel brain surfaces as S_1 , S_2 , S_3 respectively, and note that S_1 , S_2 , S_3 are all simply connected open surfaces and hence are topologically equivalent to the unit disk. We then started by applying the disk conformal parameterization method in (6) to map S_1 , S_2 , S_3 onto the unit disk, followed by the Möbius area correction scheme in (4) to further reduce the area distortion of the mappings. Denote the three disk parameterization results as $\mathcal{D}_1$, $\mathcal{D}_2$, $\mathcal{D}_3$.

Next, we followed the idea in (5) to compute landmark-aligned quasi-conformal maps between $\mathcal{D}_1$, $\mathcal{D}_2$, $\mathcal{D}_3$. More explicitly, given certain major sulci identified on each of the three brain surfaces S_1 , S_2 , S_3 , we computed two landmark-matching quasi-conformal maps $f_{13} : \mathcal{D}_1 \rightarrow \mathcal{D}_3$ and $f_{23} : \mathcal{D}_2 \rightarrow \mathcal{D}_3$ that deformed $\mathcal{D}_1$ and $\mathcal{D}_2$ to align the sulci on them with the corresponding sulci positions on $\mathcal{D}_3$. Consequently, we can visualize and compare the folding patterns of the three brain surfaces S_1 , S_2 , S_3 by using the three disk parameterizations $f_{13}(\mathcal{D}_1)$, $f_{23}(\mathcal{D}_2)$, $\mathcal{D}_3$.

Diversity and evolution of cerebral folding in primates

For a surface-based (superficial) comparison of the brain morphologies among different primates from an evolutionary perspective, we take advantage of the fact that biologists have provided plenty of anatomical data on brain geometry, although a systematic study on how these cerebral folding patterns vary across different sizes, initial shapes, and foldedness for human and other primate species is still lacking. Among primates, brains vary enormously from roughly the size of a grape to the size of a grapefruit, and from nearly smooth to dramatically folded; of these, human brains are amongst the most folded, and the largest (relative to body size). These variations in size and form make comparative neuroanatomy a rich resource for investigating common trends that transcend differences between species. We examined 10 primate species in order to cover a wide range of sizes and forms, as shown in **Fig. S5** [↗](#). Using our developed morphometric method, more

investigations on the scaling law of their cortical thickness relative to the surface geometry, folding with respect to size (total surface area) and geometry (i.e. curvature, shape, and sulcal depth), and foldedness (gyrification) would be expected.




Fig. S1. (a) MRI scans showing common malformations of cortical development of human brains. Adapted from (15) with permission. Left: normal brain. Middle: lissencephaly spectrum with agyria–severe pachygyria (arrows). Right: bilateral frontoparietal polymicrogyria with abnormally small gyri and shallow sulci (arrows). Scale bars: 3 cm (estimated from (14)). (b) A noncoding mutation in the GPR56 gene disrupts perisylvian gyri. MRI shows polymicrogyria in the perisylvian area, resulting in a characteristic, thickened appearance. Adapted from (1) with permission.


Table S2.

Similarity index (main text, Eq.(4)) evaluated by rescaled mean curvature of simulated and gel brain surfaces with comparison to the real brain surfaces, calculated with different vector p-norm: $p = 1$, $p = 2$, and $p = \infty$.

Similarity Index p	Simulated			Gel		
	1	2	∞	1	2	∞
Ferret (left)	0.8188	0.7541	1	0.7448	0.6746	1
Ferret (right)	0.7733	0.7005	1	0.7533	0.6887	1
Macaque (left)	0.6886	0.6039	1	0.7658	0.7086	1
Macaque (right)	0.6832	0.5994	1	0.7578	0.6989	1
Human (left)	0.6056	0.5090	1	0.7037	0.6540	1
Human (right)	0.5911	0.4909	1	0.7028	0.6525	1

Fig. S2.

Illustration of shape index scale divided into nine categories: spherical cup, trough rut, saddle rut, saddle, saddle ridge, ridge, dome and spherical cap. The insets are schematics of local curved surfaces. All outward normals pointing upwards.






Fig. S3.

The histogram of shape index SI (top two rows) and rescaled mean curvature $\tilde{H}$ (bottom two rows) of adult cortical surfaces of ferret, macaque and human. Insets are real brain surfaces. Colors represent shape index SI (Eq. (2)) or rescaled mean curvature $\tilde{H}$ (Eq. (3)) in the main text).

Fig. S4. Comparison among real (S_1), simulated (S_2), and gel brains (S_3) of ferret, rhesus macaque, and human via morphometric analysis.

(a) 3D cortical surfaces of *in vivo*, *in silico*, and *in vitro* models. Both left and right cortical surfaces are provided to present the left-right symmetry. (b) The quasi-conformal disk mapping with landmark matching of cortical surfaces on disk. Blue or red curves represent corresponding landmarks. Color represents shape index (SI). Similarity indices of each simulated and gel brain surfaces are presented in Table S2.





Fig. S5. Comparison across ten primate species.

Each species is listed by its common name and scientific name, and accompanied by a picture. Scale bar: 1cm. Color represents the shape index. Pictures are taken from Wikipedia.



Data availability

Reconstructed 3D surface models of fetal and adult brains of macaque and human are available on GitHub at <https://github.com/YinSifan0204/Comparative-brain-morphologies>. Newborn ferret brain surfaces will be made available upon publication of the paper. All other data are included in the article and/or supplementary material.

Acknowledgements

We acknowledge partial financial support from project NeuroWebLab (ANR-19-DATA-0025, K.H., R.T.), DMOBE (ANR-21-CE45-0016), the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 101033485 (K.H. Individual Fellowship), the NSF-ANR grant 2204058 (K.H., R.T., L.M.), the Simons Foundation (L.M.) and the Henri Seydoux Fund (L.M.), and the CUHK Faculty of Science Direct Grant for Research #4053650 (G.P.T.C.).

Additional information

Author contributions

L.M. and R.T. conceived the project; S.Y., C.L., G.P.T.C., and L.M. designed research; S.Y. collected data and performed numerical simulations; C.L. and Y.J. performed gel experiments; S.Y. and G.P.T.C. contributed morphometric analysis; K.H. and R.T. provided adult ferret and primate brain data; S.Y., G.P.T.C., and L.M. wrote the paper with input from all authors; all authors edited the paper; L.M. supervised the project.

Funding

NeuroWebLab (ANR-19-DATA-0025)

- Katja Heuer
- Roberto Toro

DMOBE (ANR-21-CE45-0016)

- Katja Heuer

European Commission

- Katja Heuer

NSF-ANR grant (2204058)

- Katja Heuer
- Roberto Toro
- L Mahadevan

Simons Foundation (SF)

- L Mahadevan

Henri Seydoux Fund

- L Mahadevan

CUHK Faculty of Science Direct Grant for Research (4053650)

- Gary Choi

Author ORCID iDs

Sifan Yin: <http://orcid.org/0000-0002-0296-3981>

Chunzi Liu: <http://orcid.org/0000-0002-3469-3904>

Gary PT Choi: <http://orcid.org/0000-0001-5407-9111>

Yeonsu Jung: <http://orcid.org/0000-0001-7920-3844>

Katja Heuer: <http://orcid.org/0000-0002-7237-0196>
Roberto Toro: <http://orcid.org/0000-0002-6671-858X>
L Mahadevan: <http://orcid.org/0000-0002-5114-0519>

Additional files

Movie S1 [↗](#) Part 1: the swelling processes of physical gel brains of ferret, macaque, and human brains; Part 2: the simulations mimicking the developmental processes of fetal brains of ferret, macaque, and human from smooth surface to the convoluted pattern.

Movie S2 [↗](#) Comparison of the real (gray), gel (pink) and simulated (blue) ferret brains. Color represents shape index.

Movie S3 [↗](#) Comparison of the real (gray), gel (pink) and simulated (blue) macaque brains. Color represents shape index.

Movie S4 [↗](#) Comparison of the real (gray), gel (pink) and simulated (blue) human brains. Color represents shape index.

References

- (2016) Comparisons, 2. <https://ventricular.org/StephenNoctor/comparisons-2/>
- (2019) NetGen. <https://ngsolve.org/>
- Akula SK**, Exposito-Alonso D, Walsh CA (2023) Shaping the brain: The emergence of cortical structure and folding. *Developmental Cell* **58**:2836-2849
- Alenyà M**, Wang X, Lefèvre J, Auzias G, Fouquet B, Eixarch E, Rousseau F, Camara O. (2022) Computational pipeline for the generation and validation of patient-specific mechanical models of brain development. *Brain Multi-physics* **3**:100045
- Bae BI**, Tietjen I, Atabay KD, Evrony GD, Johnson MB, Asare E, Wang PP, Murayama AY, Im K, Lisgo SN *et al.* (2014) Evolutionarily dynamic alternative splicing of GPR56 regulates regional cerebral cortical patterning. *Science* **343**:764-768
- Barnette AR**, Neil JJ, Kroenke CD, Griffith JL, Epstein AA, Bayly PV, Knutsen AK, Inder TE (2009) Characterization of brain development in the ferret via MRI. *Pediatric Research* **66**:80-84
- Barresi M**, Hickmott RA, Bosakhar A, Quezada S, Quigley A, Kawasaki H, Walker D, Tolcos M. (2024) Toward a better understanding of how a gyrified brain develops. *Cerebral Cortex* **34**:bhae055
- Bohi A**, Wang X, Harrach M, Dinomais M, Rousseau F, Lefèvre J. (2019) Global perturbation of initial geometry in a biomechanical model of cortical morphogenesis. In: 2019 41st Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC). pp. 442-445
- Budday S**, Raybaud C, Kuhl E. (2014) A mechanical model predicts morphological abnormalities in the developing human brain. *Scientific Reports* **4**:5644
- Bullmore ET**, Bassett DS (2011) Brain graphs: graphical models of the human brain connectome. *Annual Review of Clinical Psychology* **7**:113-140
- Calabrese E**, Badea A, Coe CL, Lubach GR, Shi Y, Styner MA, Johnson GA (2015) A diffusion tensor MRI atlas of the postmortem rhesus macaque brain. *Neuroimage* **117**:408-416
- Choi GPT**, Leung-Liu Y, Gu X, Lui LM (2020) Parallelizable global conformal parameterization of simply-connected surfaces via partial welding. *SIAM Journal on Imaging Sciences* **13**:1049-1083
- Choi GPT**, Liu C, Yin S, Séjourné G, Smith RS, Walsh CA, Mahadevan L. (2025) Biophysical basis for brain folding and misfolding patterns in ferrets and humans. *eLife* **14**:RP107141 <https://doi.org/10.7554/eLife.107141>
- Choi PT**, Lam KC, Lui LM (2015) FLASH: Fast landmark aligned spherical harmonic parameterization for genus-0 closed brain surfaces. *SIAM Journal on Imaging Sciences* **8**:67-94
- Choi PT**, Lui LM (2015) Fast disk conformal parameterization of simply-connected open surfaces. *Journal of Scientific Computing* **65**:1065-1090

- Cignoni P, Callieri M, Corsini M, Dellepiane M, Ganovelli F, Ranzuglia G, et al. (2008) Meshlab: an open-source mesh processing tool. In: Eurographics Italian Chapter Conference. **2008** Salerno, Italy. pp. 129-136
- Del-Valle-Anton L, Borrell V. (2022) Folding brains: from development to disease modeling. *Physiological Reviews* **102**:511-550
- Demirci N, Holland MA (2022) Cortical thickness systematically varies with curvature and depth in healthy human brains. *Human Brain Mapping* **43**:2064-2084
- Garcia KE, Robinson EC, Alexopoulos D, Dierker DL, Glasser MF, Coalson TS, Ortinau CM, Rueckert D, Taber LA, Van Essen DC et al. (2018) Dynamic patterns of cortical expansion during folding of the preterm human brain. *Proceedings of the National Academy of Sciences* **115**:3156-3161
- Gautam P, Anstey KJ, Wen W, Sachdev PS, Cherbuin N. (2015) Cortical gyrification and its relationships with cortical volume, cortical thickness, and cognitive performance in healthy mid-life adults. *Behavioural Brain Research* **287**:331-339
- Gill HK, Yin S, Lawlor JC, Huycke TR, Nerurkar NL, Tabin CJ, Mahadevan L. (2024) The developmental mechanics of divergent buckling patterns in the chick gut. *Proceedings of the National Academy of Sciences* **121**:e2310992121
- Gill HK, Yin S, Nerurkar NL, Lawlor JC, Lee C, Huycke TR, Mahadevan L, Tabin CJ (2024) Hox gene activity directs physical forces to differentially shape chick small and large intestinal epithelia. *Developmental Cell* **59**:2834-2849
- Herculano-Houzel S. (2009) The human brain in numbers: a linearly scaled-up primate brain. *Frontiers in Human Neuroscience* **3**:857
- Heuer K, Gulban OF, Bazin PL, Osoianu A, Valabregue R, Santin M, Herbin M, Toro R. (2019) Evolution of neocortical folding: A phylogenetic comparative analysis of MRI from 34 primate species. *Cortex* **118**:275-291
- Hohlfeld E, Mahadevan L. (2012) Scale and nature of sulcification patterns. *Physical Review Letters* **109**:025701
- Hohlfeld E, Mahadevan L. (2011) Unfolding the sulcus. *Physical Review Letters* **106**:105702
- Holland MA, Miller KE, Kuhl E. (2015) Emerging brain morphologies from axonal elongation. *Annals of Biomedical Engineering* **43**:1640-1653
- Hu HH, Chen HY, Hung CI, Guo WY, Wu YT (2013) Shape and curvedness analysis of brain morphology using human fetal magnetic resonance images in utero. *Brain Structure and Function* **218**:1451-1462
- Hutton C, Draganski B, Ashburner J, Weiskopf N. (2009) A comparison between voxel-based cortical thickness and voxel-based morphometry in normal aging. *Neuroimage* **48**:371-380
- Koenderink JJ, van Doorn AJ (1992) Surface shape and curvature scales. *Image and Vision Computing* **10**:557-564
- Kriegeskorte N, Wei XX (2021) Neural tuning and representational geometry. *Nature Reviews Neuroscience* **22**:703-718
- Lai CS, Gerrelli D, Monaco AP, Fisher SE, Copp AJ (2003) FOXP2 expression during brain development coincides with adult sites of pathology in a severe speech and language disorder. *Brain* **126**:2455-2462
- Liu Z, Wang X, Newman N, Grant KA, Studholme C, Kroenke CD (2020) Anatomical and diffusion MRI brain atlases of the fetal rhesus macaque brain at 85, 110 and 135 days gestation. *Neuroimage* **206**:116310
- van der Meer D, Kaufmann T. (2022) Mapping the genetic architecture of cortical morphology through neuroimaging: progress and perspectives. *Translational Psychiatry* **12**:447
- de Moraes FHP, Sudo F, Carneiro Monteiro M, de Melo BR, Mattos P, Mota B, Tovar-Moll F. (2024) Cortical folding correlates to aging and Alzheimer's Disease's cognitive and CSF biomarkers. *Scientific Reports* **14**:3222

- Namburete AI**, Papież BW, Fernandes M, Wyburd MK, Hesse LS, Moser FA, Ismail LC, Gunier RB, Squier W, Ohuma EO *et al.* (2023) Normative spatiotemporal fetal brain maturation with satisfactory development at 2 years. *Nature* **623**:106-114
- Oegema R**, Barakat TS, Wilke M, Stouffs K, Amrom D, Aronica E, Bahi-Buisson N, Conti V, Fry AE, Geis T *et al.* (2020) International consensus recommendations on the diagnostic work-up for malformations of cortical development. *Nature Reviews Neurology* **16**:618-635
- Pang JC**, Aquino KM, Oldehinkel M, Robinson PA, Fulcher BD, Breakspear M, Fornito A. (2023) Geometric constraints on human brain function. *Nature* **618**:566-574
- Qi J**, Mo F, An NA, Mi T, Wang J, Qi JT, Li X, Zhang B, Xia L, Lu Y *et al.* (2023) A Human-Specific De Novo Gene Promotes Cortical Expansion and Folding. *Advanced Science* **10**:2204140
- Schwartz E**, Nenning KH, Heuer K, Jeffery N, Bertrand OC, Toro R, Kasprian G, Prayer D, Langs G. (2023) Evolution of cortical geometry and its link to function, behaviour and ecology. *Nature Communications* **14**:2252
- Shinmyo Y**, Terashita Y, Duong TAD, Horiike T, Kawasumi M, Hosomichi K, Tajima A, Kawasaki H. (2017) Folding of the cerebral cortex requires Cdk5 in upper-layer neurons in gyrencephalic mammals. *Cell Reports* **20**:2131-2143
- de Sousa AA**, Beaudet A, Calvey T, Bardo A, Benoit J, Charvet CJ, Dehay C, Gómez-Robles A, Gunz P, Heuer K *et al.* (2023) From fossils to mind. *Communications Biology* **6**:636
- Striedter GF**, Srinivasan S, Monuki ES (2015) Cortical folding: when, where, how, and why?. *Annual Review of Neuroscience* **38**:291-307
- Suárez LE**, Markello RD, Betzel RF, Misic B. (2020) Linking structure and function in macroscale brain networks. *Trends in Cognitive Sciences* **24**:302-315
- Takahata T**, Shukla R, Yamamori T, Kaas JH (2012) Differential expression patterns of striate cortex-enriched genes among old world, new world, and prosimian primates. *Cerebral Cortex* **22**:2313-2321
- Tallinen T**, Biggins JS, Mahadevan L. (2013) Surface sulci in squeezed soft solids. *Physical Review Letters* **110**:024302
- Tallinen T**, Chung JY, Biggins JS, Mahadevan L. (2014) Gyrification from constrained cortical expansion. *Proceedings of the National Academy of Sciences* **111**:12667-12672
- Tallinen T**, Chung JY, Rousseau F, Girard N, Lefèvre J, Mahadevan L. (2016) On the growth and form of cortical convolutions. *Nature Physics* **12**:588-593
- Van Essen DC** (2020) A 2020 view of tension-based cortical morphogenesis. *Proceedings of the National Academy of Sciences* **117**:32868-32879
- Wang X**, Lefèvre J, Bohi A, Harrach MA, Dinomais M, Rousseau F. (2021) The influence of biophysical parameters in a biomechanical model of cortical folding patterns. *Scientific Reports* **11**:7686
- Weickenmeier J**. (2023) Exploring the multiphysics of the brain during development, aging, and in neurological diseases. *Brain Multiphysics* **4**:100068
- Bae Byoung-II**, Tietjen Ian, Atabay Kutay D, Evrony Gilad D, Johnson Matthew B, Asare Ebenezer, Wang Peter P, Murayama Ayako Y, Im Kiho, Lisgo Steven N *et al.* (2014) Evolutionarily dynamic alternative splicing of gpr56 regulates regional cerebral cortical patterning. *Science* **343**:764-768
- Barresi Mikaela**, Hickmott Ryan Alexander, Bosakhar Abdulhameed, Quezada Sebastian, Quigley Anita, Kawasaki Hiroshi, Walker David, Tolcos Mary (2024) Toward a better understanding of how a gyrified brain develops. *Cerebral Cortex* **34**:bhae055
- Budday Silvia**, Raybaud Charles, Kuhl Ellen (2014) A mechanical model predicts morphological abnormalities in the developing human brain. *Scientific Reports* **4**:5644
- Choi Gary P T**, Leung-Liu Yusan, Gu Xianfeng, Lui Lok Ming (2020) Parallelizable global conformal parameterization of simply-connected surfaces via partial welding. *SIAM Journal on Imaging Sciences* **13**:1049-1083

- Choi Pui Tung**, Lam Ka Chun, Lui Lok Ming (2015) Flash: Fast landmark aligned spherical harmonic parameterization for genus-0 closed brain surfaces. *SIAM Journal on Imaging Sciences* **8**:67-94
- Choi Pui Tung**, Lui Lok Ming (2015) Fast disk conformal parameterization of simply-connected open surfaces. *Journal of Scientific Computing* **65**:1065-1090
- Demirci Nagehan**, Holland Maria A (2022) Cortical thickness systematically varies with curvature and depth in healthy human brains. *Human Brain Mapping* **43**:2064-2084
- Hu Hui-Hsin**, Chen Hui-Yun, Hung Chih-I, Guo Wan-Yuo, Wu Yu-Te (2013) Shape and curvedness analysis of brain morphology using human fetal magnetic resonance images in utero. *Brain Structure and Function* **218**:1451-1462
- Koenderink Jan J**, van Doorn Andrea J (1992) Surface shape and curvature scales. *Image and Vision Computing* **10**:557-564
- Koenig Matti**, Dobyns William B, Di Donato Nataliya (2021) Lissencephaly: update on diagnostics and clinical management. *European Journal of Paediatric Neurology* **35**:147-152
- Lai Cecilia SL**, Gerrelli Dianne, Monaco Anthony P, Fisher Simon E, Copp Andrew J (2003) Foxp2 expression during brain development coincides with adult sites of pathology in a severe speech and language disorder. *Brain* **126**:2455-2462
- Luo Rong**, Yang Hye Min, Jin Zhaohui, Halley Dicky JJ, Chang Bernard S, MacPherson Lesley, Brueton Louise, Piao Xianhua (2011) A novel gpr56 mutation causes bilateral frontoparietal polymicrogyria. *Pediatric Neurology* **45**:49-53
- Magen Daniella**, Ofir Ayala, Berger Liron, Goldsher Dorit, Eran Ayelet, Katib Nassser, Nijem Yousif, Vlodavsky Euvgeni, Zur Shay, Behar Doron M *et al.* (2015) Autosomal recessive lissencephaly with cerebellar hypoplasia is associated with a loss-of-function mutation in cdk5. *Human Genetics* **134**:305-314
- Mochida Ganeshwaran H** (2009) Genetics and biology of microcephaly and lissencephaly. *Seminars in pediatric neurology* **16**:120-126
- Oegema Renske**, Barakat Tahsin Stefan, Wilke Martina, Stouffs Katrien, Amrom Dina, Aronica Eleonora, Bahi-Buisson Nadia, Conti Valerio, Fry Andrew E, Geis Tobias *et al.* (2020) International consensus recommendations on the diagnostic work-up for malformations of cortical development. *Nature Reviews Neurology* **16**:618-635
- Qi Jianhuan**, Mo Fan, An Ni A, Mi Tingwei, Wang Jiaxin, Qi Jun-Tian, Li Xiangshang, Zhang Boya, Xia Longkuo, Lu Yingfei *et al.* (2023) A human-specific de novo gene promotes cortical expansion and folding. *Advanced Science* **10**:2204140
- Shinmyo Yohei**, Terashita Yukari, Duong Tung Anh Dinh, Horiike Toshihide, Kawasumi Muneo, Hosomichi Kazuyoshi, Tajima Atsushi, Kawasaki Hiroshi (2017) Folding of the cerebral cortex requires cdk5 in upper-layer neurons in gyrencephalic mammals. *Cell Reports* **20**:2131-2143
- Tallinen Tuomas**, Chung Jun Young, Biggins John S, Mahadevan L (2014) Gyrification from constrained cortical expansion. *Proceedings of the National Academy of Sciences* **111**:12667-12672
- Verrotti Alberto**, Spalice Alberto, Ursitti Fabiana, Papetti Laura, Mariani Rosanna, Castronovo Antonella, Mastrangelo Mario, Iannetti Paola (2010) New trends in neuronal migration disorders. *European Journal of Paediatric Neurology* **14**:1-12

Peer reviews

Reviewer #1 (Public review):

The manuscript by Yin and colleagues addresses a long-standing question in the field of cortical morphogenesis, regarding factors that determine differential cortical folding across species and individuals with cortical malformations. The authors present work based on a computational model of cortical folding evaluated alongside a physical model that makes use of gel swelling to investigate the role of a two-layer model for cortical morphogenesis. The

study assesses these models against empirically derived cortical surfaces based on MRI data from ferret, macaque monkey, and human brains.

The manuscript is clearly written and presented, and the experimental work (physical gel modeling as well as numerical simulations) and analyses (subsequent morphometric evaluations) are conducted at the highest methodological standards. It constitutes an exemplary use of interdisciplinary approaches for addressing the question of cortical morphogenesis by bringing together well-tuned computational modeling with physical gel models. In addition, the comparative approaches used in this paper establish a foundation for broad-ranging future lines of work that investigate the impact of perturbations or abnormalities during cortical development.

The cross-species approach taken in this study is a major strength of the work. However, correspondence across the two methodologies did not appear to be equally consistent in predicting brain folding across all three species. The results presented in Figures 4 (and Figures S3 & S4) show broad correspondence in shape index and major sulci landmarks across all three species. Nevertheless, the results presented for the human brain lack the same degree of clear correspondence for the gel model results as observed in the macaque and ferret. While this study clearly establishes a strong foundation for comparative cortical anatomy across species and the impact of perturbations on individual morphogenesis, further work that fine-tunes physical modeling of complex morphologies, such as that of the human cortex, may help to further understand the factors that determine cortical functionalization and pathologies.

<https://doi.org/10.7554/eLife.107138.2.sa2>

Reviewer #2 (Public review):

This manuscript explores the mechanisms underlying cerebral cortical folding using a combination of physical modelling, computational simulations, and geometric morphometrics. The authors extend their prior work on human brain development (Tallinen et al., 2014; 2016) to a comparative framework involving three mammalian species: ferrets (Carnivora), macaques (Old World monkeys), and humans (Hominoidea). By integrating swelling gel experiments with mathematical differential growth models, they simulate sulcification instability and recapitulate key features of brain folding across species. The authors make commendable use of publicly available datasets to construct 3D models of fetal and neonatal brain surfaces: fetal macaque (ref. [26]), newborn ferret (ref. [11]), and fetal human (ref. [22]).

Using a combination of physical models and numerical simulations, the authors compare the resulting folding morphologies to real brain surfaces using morphometric analysis. Their results show qualitative and quantitative concordance with observed cortical folding patterns, supporting the view that differential tangential growth of the cortex relative to the subcortical substrate is sufficient to account for much of the diversity in cortical folding. This is a very important point in our field, and can be used in the teaching of medical students.

Brain folding remains a topic of ongoing debate. While some regard it as a critical specialization linked to higher cognitive function, others consider it an epiphenomenon of expansion and constrained geometry. This divergence was evident in discussions during the Strüngmann Forum on cortical development (Silver et al., 2019). Though folding abnormalities are reliable indicators of disrupted neurodevelopmental processes (e.g., neurogenesis, migration), their relationship to functional architecture remains unclear. Recent evidence suggests that the absolute number of neurons varies significantly with position-sulcus versus gyrus-with potential implications for local processing capacity (e.g., <https://doi.org/10.1002/cne.25626> ). The field is thus in need of comparative, mechanistic studies like the present one.

This paper offers an elegant and timely contribution by combining gel-based morphogenesis, numerical modelling, and morphometric analysis to examine cortical folding across species. The experimental design - constructing two-layer PDMS models from 3D MRI data and immersing them in organic solvents to induce differential swelling - is well-established in prior literature. The authors further complement this with a continuum mechanics model simulating folding as a result of differential growth, as well as a comparative analysis of surface morphologies derived from in vivo, in vitro, and in silico brains.

Conclusion:

This is a well-executed and creative study that integrates diverse methodologies to address a longstanding question in developmental neurobiology. While a few aspects-such as regional folding peculiarities, sensitivity to initial conditions, and available human data-could be further elaborated, they do not detract from the overall quality and novelty of the work. I enthusiastically support this paper and believe that it will be of broad interest to the neuroscience, biomechanics, and developmental biology communities.

[Editor's note: The reviewers were satisfied with the authors' response. The eLife Assessment was slightly updated to reflect the author's response.]

<https://doi.org/10.7554/eLife.107138.2.sa1>

Author response:

The following is the authors' response to the original reviews

Public Reviews:

Reviewer #1 (Public review):

The manuscript by Yin and colleagues addresses a long-standing question in the field of cortical morphogenesis, regarding factors that determine differential cortical folding across species and individuals with cortical malformations. The authors present work based on a computational model of cortical folding evaluated alongside a physical model that makes use of gel swelling to investigate the role of a two-layer model for cortical morphogenesis. The study assesses these models against empirically derived cortical surfaces based on MRI data from ferret, macaque monkey, and human brains.

The manuscript is clearly written and presented, and the experimental work (physical gel modeling as well as numerical simulations) and analyses (subsequent morphometric evaluations) are conducted at the highest methodological standards. It constitutes an exemplary use of interdisciplinary approaches for addressing the question of cortical morphogenesis by bringing together well-tuned computational modeling with physical gel models. In addition, the comparative approaches used in this paper establish a foundation for broad-ranging future lines of work that investigate the impact of perturbations or abnormalities during cortical development.

The cross-species approach taken in this study is a major strength of the work. However, correspondence across the two methodologies did not appear to be equally consistent in predicting brain folding across all three species. The results presented in Figures 4 (and Figures S3 and S4) show broad correspondence in shape index and major sulci landmarks across all three species. Nevertheless, the results presented for the human brain lack the same degree of clear correspondence for the gel model results as observed in the macaque and ferret. While this study clearly establishes a strong foundation for comparative cortical anatomy across species and the impact of perturbations on individual morphogenesis, further work that fine-tunes physical modeling of complex

morphologies, such as that of the human cortex, may help to further understand the factors that determine cortical functionalization and pathologies.

We thank the reviewer for positive opinions and helpful comments. Yes, the physical gel model of the human brain has a lower similarity index with the real brain. There are several reasons.

First, the highly convoluted human cortex has a few major folds (primary sulci) and a very large number of minor folds associated with secondary or tertiary sulci (on scales of order comparable to the cortical thickness), relative to the ferret and macaque cerebral cortex. In our gel model, the exact shapes, positions, and orientations of these minor folds are stochastic, which makes it hard to have a very high similarity index of the gel models when compared with the brain of a single individual.

Second, in real human brains, these minor folds evolve dynamically with age and show differences among individuals. In experiments with the gel brain, multiscale folds form and eventually disappear as the swelling progresses through the thickness. Our physical model results are snapshots during this dynamical process, which makes it hard to have a concrete one-to-one correspondence between the instantaneous shapes of the swelling gel and the growing human brain.

Third, the growth of the brain cortex is inhomogeneous in space and varying with time, whereas, in the gel model, swelling is relatively homogeneous.

We agree that further systematic work, based on our proposed methods, with more fine-tuned gel geometries and properties, might provide a deeper understanding of the relations between brain geometry, and growth-induced folds and their functionalization and pathologies. Further analysis of cortical pathologies using computational and physical gel models can be found in our companion paper (Choi et al., 2025), also published in eLife:

G. P. T. Choi, C. Liu, S. Yin, G. Séjourné, R. S. Smith, C. A. Walsh, L. Mahadevan, Biophysical basis for brain folding and misfolding patterns in ferrets and humans. eLife, 14, RP107141, 2025. doi:10.7554/eLife.107141

Reviewer# 2 (Public review):

This manuscript explores the mechanisms underlying cerebral cortical folding using a combination of physical modelling, computational simulations, and geometric morphometrics. The authors extend their prior work on human brain development (Tallinen et al., 2014; 2016) to a comparative framework involving three mammalian species: ferrets (Carnivora), macaques (Old World monkeys), and humans (Hominoidea). By integrating swelling gel experiments with mathematical differential growth models, they simulate sulcification instability and recapitulate key features of brain folding across species. The authors make commendable use of publicly available datasets to construct 3D models of fetal and neonatal brain surfaces: fetal macaque (ref. [26]), newborn ferret (ref. [11]), and fetal human (ref. [22]).

Using a combination of physical models and numerical simulations, the authors compare the resulting folding morphologies to real brain surfaces using morphometric analysis. Their results show qualitative and quantitative concordance with observed cortical folding patterns, supporting the view that differential tangential growth of the cortex relative to the subcortical substrate is sufficient to account for much of the diversity in cortical folding. This is a very important point in our field, and can be used in the teaching of medical students.

Brain folding remains a topic of ongoing debate. While some regard it as a critical specialization linked to higher cognitive function, others consider it an epiphenomenon

of expansion and constrained geometry. This divergence was evident in discussions during the Strüngmann Forum on cortical development (Silver et al., 2019). Though folding abnormalities are reliable indicators of disrupted neurodevelopmental processes (e.g., neurogenesis, migration), their relationship to functional architecture remains unclear. Recent evidence suggests that the absolute number of neurons varies significantly with position-sulcus versus gyrus-with potential implications for local processing capacity (e.g., <https://doi.org/10.1002/cne.25626>). The field is thus in need of comparative, mechanistic studies like the present one.

This paper offers an elegant and timely contribution by combining gel-based morphogenesis, numerical modelling, and morphometric analysis to examine cortical folding across species. The experimental design - constructing two-layer PDMS models from 3D MRI data and immersing them in organic solvents to induce differential swelling - is well-established in prior literature. The authors further complement this with a continuum mechanics model simulating folding as a result of differential growth, as well as a comparative analysis of surface morphologies derived from in vivo, in vitro, and in silico brains.

We thank the reviewer for the very positive comments.

I offer a few suggestions here for clarification and further exploration:

Major Comments

(1) Choice of Developmental Stages and Initial Conditions

The authors should provide a clearer justification for the specific developmental stages chosen (e.g., G85 for macaque, GW23 for human). How sensitive are the resulting folding patterns to the initial surface geometry of the gel models? Given that folding is a nonlinear process, early geometric perturbations may propagate into divergent morphologies. Exploring this sensitivity-either through simulations or reference to prior work-would enhance the robustness of the findings.

The initial geometry is one of the important factors that decides the final folding pattern. The smooth brain in the early developmental stage shows a broad consistency across individuals, and we expect the main folds to form similarly across species and individuals.

Generally, we choose the initial geometry when the brain cortex is still relatively smooth. For the human, this corresponds approximately to GW23, as the major folds such as the Rolandic fissure (central sulcus), arise during this developmental stage. For the macaque brain, we chose developmental stage G85, primarily because of the availability of the dataset corresponding to this time, which also corresponds to the least folded.

We expect that large-scale folding patterns are strongly sensitive to the initial geometry but fine-scale features are not. Since our goal is to explain the large-scale features, we expect sensitivity to the initial shape.

Below are some references of other researchers that are consistent with this idea. Figure 4 from Wang et al. shows some images of simulations obtained by perturbing the geometry of a sphere to an ellipsoid. We see that the growth-induced folds mostly maintain their width (wavelength), but change their orientations.

Reference:

Wang, X., Lefèvre, J., Bohi, A., Harrach, M.A., Dinomais, M. and Rousseau, F., 2021. The influence of biophysical parameters in a biomechanical model of cortical folding patterns. *Scientific Reports*, 11(1), p.7686.

Related results from the same group show that slight perturbations of brain geometry, cause these folds also tend to change their orientations but not width/wavelength (Bohi et al., 2019).

Reference:

Bohi, A., Wang, X., Harrach, M., Dinomais, M., Rousseau, F. and Lefèvre, J., 2019, July. Global perturbation of initial geometry in a biomechanical model of cortical morphogenesis. In 2019 41st Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC) (pp. 442-445). IEEE.

Finally, a systematic discussion of the role of perturbations on the initial geometries and physical properties can be seen in our work on understanding a different system, gut morphogenesis (Gill et al., 2024).

We have added the discussion about geometric sensitivity in the section Methods-Numerical Simulations:

“Small perturbations on initial geometry would affect minor folds, but the main features of major folds, such as orientations, width, and depth, are expected to be conserved across individuals [49, 50]. For simplicity, we do not perturb the fetal brain geometry obtained from datasets.”

(2) Parameter Space and Breakdown Points

The numerical model assumes homogeneous growth profiles and simplifies several aspects of cortical mechanics. Parameters such as cortical thickness, modulus ratios, and growth ratios are described in Table II. It would be informative to discuss the range of parameter values for which the model remains valid, and under what conditions the physical and computational models diverge. This would help delineate the boundaries of the current modelling framework and indicate directions for refinement.

Exploring the valid parameter space is a key problem. We have tested a series of growth parameters and will state them explicitly in our revision. In the current version, we chose the ones that yield a relatively high similarity index to the animal brains. More generally, folding patterns are largely regulated by geometry as well as physical parameters, such as cortical thickness, modulus ratios, growth ratios, and inhomogeneity. In our previous work on a different system, gut morphogenesis, where similar folding patterns are seen, we have explored these features (Gill et al., 2024).

Reference:

Gill, H.K., Yin, S., Nerurkar, N.L., Lawlor, J.C., Lee, C., Huycke, T.R., Mahadevan, L. and Tabin, C.J., 2024. Hox gene activity directs physical forces to differentially shape chick small and large intestinal epithelia. *Developmental Cell*, 59(21), pp.2834-2849.

(3) Neglected Regional Features: The Occipital Pole of the Macaque

One conspicuous omission is the lack of attention to the occipital pole of the macaque, which is known to remain smooth even at later gestational stages and has an unusually high neuronal density (2.5× higher than adjacent cortex). This feature is not reproduced in the gel or numerical models, nor is it discussed. Acknowledging this discrepancy-and speculating on possible developmental or mechanical explanations would add depth to the comparative analysis. The authors may wish to include this as a limitation or a target for future work.

Yes, we have added that the omission of the Occipital Pole of the macaque is one of our paper's limitations. Our main aim in this paper is to explore the formation of large-scale

folds, so the smooth region is not discussed. But future work could include this to make the model more complete.

The main text has been modified in Methods, Numerical simulations:

“To focus on fold formation, we did not discuss the relatively smooth region, such as the Occipital Pole of the macaque.”

and also in the caption of Figure 4: “... The occipital pole region of macaque brains remains smooth in real and simulated brains.”

(4) Spatio-Temporal Growth Rates and Available Human Data

The authors note that accurate, species-specific spatio-temporal growth data are lacking, limiting the ability to model inhomogeneous cortical expansion. While this may be true for ferret and macaque, there are high-quality datasets available for human fetal development, now extended through ultrasound imaging (e.g., <https://doi.org/10.1038/s41586-023-06630-3> ). Incorporating or at least referencing such data could improve the fidelity of the human model and expand the applicability of the approach to clinical or pathological scenarios.

We thank the reviewer for pointing out the very useful datasets that exist for the exploration of inhomogeneous growth driven folding patterns. We have referred to this paper to provide suggestions for further work in exploring the role of growth inhomogeneities.

We have referred to this high-quality dataset in our main text, Discussion:

“...the effect of inhomogeneous growth needs to be further investigated by incorporating regional growth of the gray and white matter not only in human brains [29, 31] based on public datasets [45], but also in other species.”

A few works have tried to incorporate inhomogeneous growth in simulating human brain folding by separating the central sulcus area into several lobes (e.g., lobe parcellation method, Wang, PhD Thesis, 2021). Since our goal in this paper is to explain the large-scale features of folding in a minimal setting, we have kept our model simple and show that it is still capable of capturing the main features of folding in a range of mammalian brains.

Reference:

Xiaoyu Wang. Modélisation et caractérisation du plissement cortical. Signal and Image Processing. Ecole nationale supérieure Mines-Télécom Atlantique, 2021. English. 〈NNT : 2021IMTA0248〉 .

(5) Future Applications: The Inverse Problem and Fossil Brains

The authors suggest that their morphometric framework could be extended to solve the inverse growth problem-reconstructing fetal geometries from adult brains. This speculative but intriguing direction has implications for evolutionary neuroscience, particularly the interpretation of fossil endocasts. Although beyond the scope of this paper, I encourage the authors to elaborate briefly on how such a framework might be practically implemented and validated.

For the inverse problem, we could use the following strategies:

- a. Perform systematic simulations using different geometries and physical parameters to obtain the variation in morphologies as a function of parameters.
- b. Using either supervised training or unsupervised training (physics-informed neural networks, PINNs) to learn these characteristic morphologies and classify their dependence on

the parameters using neural networks. These can then be trained to determine the possible range of geometrical and physical parameters that yield buckled patterns seen in the systematic simulations.

c. Reconstruct the 3D surface from fossil endocasts. Using the well-trained neural network, it should be possible to predict the initial shape of the smooth brain cortex, growth profile, and stiffness ratio of the gray and white matter.

As an example in this direction, supervised neural networks have been used recently to solve the forward problem to predict the buckling pattern of a growing two-layer system (Chavoshnejad et al., 2023). The inverse problem can then be solved using machine-learning methods when the training datasets are the folded shape, which are then used to predict the initial geometry and physical properties.

Reference:

Chavoshnejad, P., Chen, L., Yu, X., Hou, J., Filla, N., Zhu, D., Liu, T., Li, G., Razavi, M.J. and Wang, X., 2023. An integrated finite element method and machine learning algorithm for brain morphology prediction. *Cerebral Cortex*, 33(15), pp.9354-9366.

Conclusion

This is a well-executed and creative study that integrates diverse methodologies to address a longstanding question in developmental neurobiology. While a few aspects—such as regional folding peculiarities, sensitivity to initial conditions, and available human data—could be further elaborated, they do not detract from the overall quality and novelty of the work. I enthusiastically support this paper and believe that it will be of broad interest to the neuroscience, biomechanics, and developmental biology communities.

Note: The paper mentions a companion paper [reference 11] that explores the cellular and anatomical changes in the ferret cortex. I did not have access to this manuscript, but judging from the title, this paper might further strengthen the conclusions.

The companion paper (Choi et al., 2025) has also been submitted to eLife and can be found here:

G. P. T. Choi, C. Liu, S. Yin, G. Séjourné, R. S. Smith, C. A. Walsh, L. Mahadevan, Biophysical basis for brain folding and misfolding patterns in ferrets and humans. *eLife*, 14, RP107141, 2025. doi:10.7554/eLife.107141

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

This study was conducted and presented to the highest methodological standards. It is clearly written, and the results are thoroughly presented in the main manuscript and supplementary materials. Nevertheless, I would present the following minor points and comments for consideration by the authors prior to finalizing their work:

We thank the reviewer for positive opinions and helpful comments.

(1) Where did the MRI-based cortical surface data come from? Specifically, it would be helpful to include more information regarding whether the surfaces were reconstructed based on individual- or group-level data. It appears the surfaces were group-level, and, if so, accounting for individual-level cortical folding may be a fruitful direction for future work.

The surface data come from public database, which are stated in the Methods Section. “We used a publicly available database for all our 3d reconstructions: fetal macaque brain surfaces are obtained from Liu et al. (2020); newborn ferret brain surfaces are obtained from Choi et al. (2025); and fetal human brain surfaces are obtained from Tallinen et al. (2016).”

These surfaces are reconstructed based on group-level data. Specifically, the macaque atlas images are constructed for brains at gestational ages of 85 days (G85, $N=18$, 9 females), 110 days (G110, $N=10$, 7 females) and 135 days (G135, $N=16$, 7 females). And yes, future work may focus on individual-level cortical folding, and we expect that more specific results could be found.

(2) One methodological approach for assessing consistency of cortical folding within species might be an evaluation of cross-hemispheric symmetry. I would find this particularly interesting with respect to the gel models, as it could complement the quantification of variation with respect to the computationally derived and real surfaces.

Yes, the cross-hemispheric symmetry comparison can be done by our morphometric analysis method. We have added the results of ferret brain’s left-right symmetry for gel models, simulations, and real surfaces in the supplementary material. A typical conformal mapping figure and the similarity index table are shown here.

(3) Was there a specific reason to reorder the histogram plots in Figure 4c to macaque, ferret, human rather than to maintain the order presented in Figure 4a/b of ferret, macaque, human? I appreciate that this is a minor concern, and all subplots are indeed properly titled, but consistent order may improve clarity.

We have reordered the histogram plots to make all the figure orders consistent.


Reviewer #2 (Recommendations for the authors):

(1) Please consider revising the caption of Figure 1 (or equivalent figures) to explicitly state whether features such as the macaque occipital flatness were reproduced or not.

We thank the reviewer for pointing out the macaque occipital flatness.

real	gel	simu
0.5754	0.7014	0.5868

Author response table 1. Left-right similarity index evaluated by comparing the shape index of ferret brains, calculated with vector P-NORM $p=2$,



Author response image 1. Left-right similarity index of ferret brains

Occipital Pole of the macaque remains relatively smooth in both real brains and computational models. But our main aim in this paper is to explore the large-scale folds formation, so the smooth region is not discussed in depth. But future work could include this to make the model more complete.

(2) Some figures could benefit from clearer labelling to distinguish between in vivo, in vitro, and in silico results.

We have supplemented some texts in panels to make the labelling clearer.

(3) The manuscript would benefit from a short paragraph in the Discussion reflecting on how future incorporation of regional heterogeneities might improve model fidelity.

We have added a sentence in the Discussion Section about improving the model fidelity by considering regional heterogeneities.

“Future more accurate models incorporating spatio-temporal inhomogeneous growth profiles and mechanical properties, such as varying stiffness, would make the folding pattern closer to the real cortical folding. This relies on more in vivo measurements of the brain’s physical properties and cortical expansion.”

(4) Suggestions for improved or additional experiments, data, or analyses.

(5) Clarify and justify the selection of developmental stages: The authors should explain why particular gestational stages (e.g., G85 for macaque, GW23 for human) were chosen as starting points for the physical and computational models. A discussion of how sensitive the folding patterns are to the initial geometry would help assess the robustness of the model. If feasible, a brief sensitivity analysis-varying initial age or surface geometry-would strengthen the conclusions.

The initial geometry is one of the important factors that decides the final folding pattern. The smooth brain in the early developmental stage shows a broad consistency across individuals, and we expect the main folds to form similarly across species and individuals.

Generally, we choose the initial geometry when the brain cortex is still relatively smooth. For the human, this corresponds approximately to GW23, as the major folds such as the Rolandic fissure (central sulcus), arise during this developmental stage. For the macaque brain, we chose developmental stage G85, primarily because of the availability of the dataset corresponding to this time, which also corresponds to the least folded.

We expect that large-scale folding patterns are strongly sensitive to the initial geometry but fine-scale features are not. Since our goal is to explain the large-scale features, we expect sensitivity to the initial shape.

We have added the discussion about geometric sensitivity in the section Methods-Numerical Simulations: “Small perturbations on initial geometry would affect minor folds, but the main features of major folds, such as orientations, width, and depth, are expected to be conserved across individuals [49, 50]. For simplicity, we do not perturb the fetal brain geometry obtained from datasets.”

(6) Explore parameter boundaries more explicitly: The paper would benefit from a clearer account of the ranges of mechanical and geometric parameters (e.g., growth ratios, cortical thickness) for which the model holds. Are there specific conditions under which the physical and numerical models diverge? Identifying breakdown points would help readers understand the model's limitations and applicability.

Exploring the valid parameter space is a key problem. We have tested a series of growth parameters and will state them explicitly in our revision. In the current version, we chose the ones that yield a relatively high similarity index to the animal brains. More generally, folding patterns are largely regulated by geometry as well as physical parameters, such as cortical thickness, modulus ratios, and growth ratios and inhomogeneity. In our previous work on a different system, gut morphogenesis, where similar folding patterns are seen, we have explored these features (Gill et al., 2024).

(7) Address species-specific cortical peculiarities: A striking omission is the flat occipital pole of the macaque, which is not reproduced in the physical or computational models. Given its known anatomical and cellular distinctiveness, this discrepancy warrants discussion. Even if not explored experimentally, the authors could speculate on what developmental or mechanical conditions would be needed to reproduce such regional smoothness.

Please refer to our answer to the public reviewer 2, question (3). From our results, the formation of smooth Occipital Pole might indicate that the spatio-temporal growth rate of gray and white matter are consistent in this region, such that there's no much differential growth.

(8) Consider integration of available human growth data: While the authors note the lack of spatiotemporal growth data across species, such datasets exist for human fetal brain development, including those from MRI and ultrasound studies (e.g., Nature 2023). Incorporating these into the human model-or at least discussing their implications-would enhance biological relevance.

Yes, some datasets for fetal human brains have provided very comprehensive measurements on brain shapes at many developmental stages. This can surely be implemented in our current model by calculating the spatio-temporal growth rate from regional cortical shapes at different stages.

(9) Recommendations for improving the writing and presentation:

a) The manuscript is generally well-written, but certain sections would benefit from more explicit links between the biological phenomena and the modeling framework. For instance, the Introduction and Discussion could more clearly articulate how mechanical principles interface with genetic or cellular processes, especially in the context of evolution and developmental variation.

We have briefly discussed the gene-regulated cellular process and the induced changes of mechanical properties and growth rules in SI, table S1. In the main text, to be clearer, we have added a sentence:

“Many malformations are related to gene-regulated abnormal cellular processes and mechanical properties, which are discussed in SI”

b) The Discussion could better acknowledge limitations and future directions, including regional differences in folding, inter-individual variability, and the model's assumptions of homogeneous material properties and growth.

In the discussion section, we have pointed out four main limitations and open directions based on our current model, including the discussion on spatiotemporal growth and property. To be more complete, we have supplemented other limitations on the regional differences in folding and the interindividual variability. In the main text, we added the following sentence:

“In addition to the homogeneity assumption, we have not investigated the inter-individual variability and regional differences in folding. More accurate and specific work is expected to focus on these directions.”

c) The authors briefly mention the potential for addressing the inverse growth problem. Expanding this idea in a short paragraph - perhaps with hypothetical applications to fossil brain reconstructions-would broaden the paper's appeal to evolutionary neuroscientists.

We have stated general steps in the response to public reviewer 2, question (5).

(10) Minor corrections to the text and figures:

a) Figures:

Label figures more clearly to distinguish between in vivo, in vitro, and in silico brain representations.- Ensure that the occipital pole of the macaque is visible or annotated, especially if it lacks the expected smoothness.

Add scale bars where missing for clarity in morphometric comparisons.

We thank the reviewer for suggestions to improve the readability of our manuscript.

The *in vivo* (real), *in vitro* (gel), and *in silico* (simulated) results are both distinguished by their labels and different color scheme: gray-white for real brain, pink-white for gel model, and blue-white for simulations, respectively.

The occipital pole of the macaque brain remains relatively smooth in our computational model but not in our physical gel model. We have clarified this in the main text: “To focus on fold formation, we did not discuss the relatively smooth region, such as the Occipital Pole of the macaque.”

All the brain models are rescaled to the same size, where the distance between the anterior-most point of the frontal lobe and the posterior-most point of the occipital lobe is two units.

b) Text:

Consider revising figure captions to explicitly mention whether specific regional features (e.g., flat occipital pole) were observed or absent in models.

In Table II (and relevant text), ensure parameter definitions are consistent and explained clearly for across-disciplinary audience.

Add citations to recent human fetal growth imaging work (e.g., ultrasound-based studies) to support claims about available data.

We have added some descriptions of the characters of the folding pattern in the caption of Figure 4, including major folds and smooth regions.

“Three or four major folds of each brain model are highlighted and served as landmarks. The occipital pole region of macaque brains remains smooth in real and simulated brains.”

We have clarified the definition of growth ratio $g_{\text{Msub}}/g_{\text{w}}$ and stiffness ratio $\mu_{\text{g}}/\mu_{\text{w}}$ between gray matter and white matter, and the normalized cortical thickness h/L in Table 2.

We have referred to a high-quality dataset of fetal brain imaging work, the ultrasound-imaging method (Namburete et al. 2023), in our main text, Discussion:

“...the effect of inhomogeneous growth needs to be further investigated by incorporating regional growth of the gray and white matter not only in human brains [29, 31] based on public datasets [45], but also in other species.”

<https://doi.org/10.7554/eLife.107138.2.sa0>