

Reviewed Preprint

v1 • July 24, 2025

Not revised

Reviewed Preprint

v2 • January 23, 2026

Revised by authors

For correspondence:

cepko@genetics.med.harvard.edu

Competing interests:

No competing interests declared

Funding:

See page 57

Reviewing editor:

H Efsun Arda,
National Cancer Institute, United States

© 2025, Tulloch et al. This article is

distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted

use and redistribution provided that

the original author and source are

credited.

Massively parallel reporter assay for mapping gene-specific regulatory regions at single nucleotide resolution

Alastair J Tulloch^{1,3}, Ryan N Delgado^{1,3}, Rinaldo Catta-Preta^{1,3,4,5}, Constance L Cepko^{1,2,3} ✉

¹Department of Genetics, Harvard Medical School, Boston, United States • ²Department of Ophthalmology, Harvard Medical School, Boston, United States • ³Howard Hughes Medical Institute, Harvard Medical School, Boston, United States • ⁴Ocular Genomics Institute, Mass Eye and Ear, Boston, United States • ⁵Department of Ophthalmology, Harvard Medical School, Boston, United States

eLife Assessment

This manuscript presents a **valuable** methodological approach to investigating context-dependent activity of cis-regulatory activity within defined genomic loci. The authors combine a locus-specific massively parallel reporter assay, enabling unbiased and high-coverage profiling of enhancer activity across large genomic regions, with a degenerate reporter assay to identify nucleotides critical for enhancer function. The data supporting the conclusions are **solid**, highlighted by successful identification and characterization of both previously known and new regulatory elements across multiple developmental stages, cell types, and species. While the approach has inherent limitations in sensitivity, and indirect assignment of regulatory elements to target genes, it provides a flexible platform for nominating candidate cis-regulatory elements across defined loci.

<https://doi.org/10.7554/eLife.107565.2.sa4>

Abstract

Precise gene regulation is essential for the development and function of complex tissues, yet comprehensive mapping of cis-regulatory modules (CRMs) remains challenging due to limitations in throughput, resolution, and the ability to assay within specific cell types. Here, we introduce two complementary approaches—a locus-specific massively parallel reporter assay (LS-MPRA) and a degenerate MPRA (d-MPRA)—specifically designed to address some of these shortcomings. LS-MPRA leverages bacterial artificial chromosomes (BACs) to generate high-complexity libraries spanning large genomic regions, enabling unbiased interrogation of millions of DNA fragments potentially relevant for the regulation of a specific gene or set of genes. The d-MPRA employs systematic mutagenesis to resolve the functional architecture of CRMs at nucleotide resolution, thereby nominating critical nucleotides as potential TF binding sites, or for other regulatory roles.

We applied these methods to retinal genes that are stably expressed in differentiated cells of the retina, in rod photoreceptors and in subsets of bipolar interneurons, using both *in vivo* and *ex vivo* preparations of mouse tissue. LS-MPRA recapitulated some of the known CRMs for these genes—such as the proximal promoter region of *Rho*—and identified potentially novel CRMs, including those located within neighboring genes. The method was then applied to a gene that is dynamically expressed in subsets of retinal progenitor cells, *Olig2*, where it identified three distinct CRM regions (Olig2-NR1, NR2, and NR3). D-MPRA and subsequent motif analyses nominated critical TF binding sites within these regions. CUT&RUN experiments confirmed direct binding of these candidates. Moreover, extending LS-MPRA to chick retina and spinal cord demonstrated the applicability of these methods across species and tissues.

Together, the integrated LS-MPRA and d-MPRA strategies provide a robust, high-resolution platform for discovery of the *cis*-regulatory code underlying tissue-specific gene expression. It does not require prior knowledge of potential CRMs, and is quite rapid and straightforward to deploy, using typical molecular biology methods. The fragment size can be scaled to create short CRMs, e.g. for cell type-specific expression within viral vectors. It should enable CRM discovery at a scale and affordability for laboratories wishing to focus on a particular locus or set of loci.

Introduction

The precise regulation of gene expression is fundamental to the development and function of complex tissues. Intricate transcriptional programs govern cell fate decisions and ensure the proper spatial and temporal expression of genes. Central to these programs are *cis*-regulatory modules (CRMs), which integrate combinatorial inputs from transcription factors (TFs) and other regulatory mechanisms to orchestrate gene expression in a context-dependent manner. Despite extensive research identifying individual enhancers and promoters, a comprehensive understanding of the regulatory landscapes that dictate cellular diversity remains elusive. This gap is largely attributable to limitations in conventional assays regarding throughput, resolution, and the reliance on cultured cells.

Massively parallel reporter assays (MPRAs) have emerged as powerful platforms for systematically interrogating enhancer activity (Agarwal et al., 2025 [DOI](#), Arnold et al., 2013 [DOI](#), Inoue & Ahituv, 2015 [DOI](#), Lagunas et al., 2023 [DOI](#), McAfee et al., 2022 [DOI](#), Melnikov et al., 2012 [DOI](#), Shen et al., 2016a [DOI](#)). Most MPRAs rely on predetermined sets of DNA fragments, identified through chromatin, RNA, and/or bioinformatic analyses. They are often broad in scale, in some cases, extending to genome-wide analyses. Constructing such libraries is both time-consuming and resource-intensive, and the selection process can be inherently biased. When an investigator aims to identify CRMs for only one or a few genes, most MPRAs are more complex and expensive than is required, and genome-wide MPRA libraries may lack the resolution needed to detect CRMs for genes that are transiently expressed or limited to small cell populations. CRMs for specific genes might not only lead to an understanding of gene regulation, but they can be used for cell type-specific gene expression for perturbations in specific settings and/or for clinical applications. To provide methods for these latter applications, we developed a locus-specific MPRA (LS-MPRA) that leverages bacterial artificial chromosomes (BACs) to generate highly complex libraries for the unbiased interrogation of millions of fragments surrounding genes of interest (Rabe, 2020 [DOI](#)). Fragment size can be chosen by the investigator, which is especially useful when a small fragment might be required, e.g. for cell type-specific expression in a viral vector. The LS-MPRA library approach not only identifies potential CRMs under homeostatic conditions, but can be run in a perturbed condition, to identify CRMs that might direct a response to e.g. drug treatment. In addition, we devised a degenerate MPRA (d-MPRA) technique to resolve the functional architecture of CRMs at single-nucleotide resolution. By systematically introducing random point mutations into fragments identified in the LS-MPRA, d-MPRA nominates sites required for repression and/or activation of CRMs. Integration of these data with *de novo* motif discovery, single cell RNA sequencing (scRNA-seq) data, and CUT&RUN assays can be used to further analyze the connection between CRM activity and specific transcriptional regulators.

In this study, we first determined that these methods can identify CRMs for stably expressed, cell type-specific genes, previously identified by our laboratory and others. These assays were successfully carried out in the mouse *in vivo* and in explant cultures. We then investigated the regulation of more challenging genes, *Olig2* and *Ngn2*, that are dynamically expressed within subsets of retinal progenitor cells (RPCs) during development. CRMs for *Olig2* were run through the d-MPRA method, where binding sites for known and previously unreported TF regulators of retinal development were identified. Analysis of the patterns of *Olig2* RNA expression and those of these predicted TF regulators, within single cells, ruled in the possibility that the suggested TFs might regulate *Olig2* repression and activation. Binding of the suggested TFs was then carried out and was validated by CUT&RUN assays.

These methods can be launched without any prior knowledge of a gene's architecture, and without synthesizing fragments or producing a library of specific PCR products. They offer a modular framework that can be readily adapted to diverse model systems. The methods are straightforward, requiring only basic molecular biology skills. They are rapid, with library preparation taking 1–2 weeks. Moreover, the methods can be run in a multiplexed fashion, allowing one to combine libraries from several BACs.

Results

LS-MPRA for stably expressed cell type-specific genes

The overall LS-MPRA protocol is shown in Fig. 1A. Fragment libraries from large contiguous genomic regions (~150–250 kb) within a BAC encoding a locus of interest are generated. Essentially random DNA fragments are generated by digestion by the NEBNext® Ultra™ II FS DNA Module enzyme, followed by size selection on an agarose gel. The fragments are then cloned into a barcode library with a synthetic, minimal promoter (TATAA) and GFP, with an intron in the GFP open reading frame (ORF), to allow selection of GFP cDNA vs contaminating plasmid DNA, for some applications. The barcodes are located in the 3'UTR and were used to track DNA fragment activity. Because there are many overlapping fragments from a given region, active genomic regions appear as peaks, where barcode signals from each fragment accumulate. Libraries of at least 6.2×10^6 and as many as 1.3×10^7 fragments were made (Table S3), with each nucleotide in the BAC residing in a different position within at least 5000 fragments and as many as 11,000 fragments. BAC libraries are delivered to the cells of interest, in our case by electroporation into the retina. After incubation, a cDNA library is generated using a RT primer that enables capture of barcoded transcripts, followed by PCR amplification. A gel is then run for isolation of the appropriate sized fragments. The cDNA-derived barcodes are sequenced and mapped to the fragment library. Similar to other MPRA, CRM activity is inferred from cDNA barcode ratio relative to the barcode frequency in the electroporated library.

In order to associate a given DNA fragment with a particular barcode (Fig. 1B), an association library, derived from the original LS-MPRA library, is created and sequenced. This association library is made by excision of the minimal promoter and GFP, followed by ligation (Fig. 1C). This is followed by paired end sequencing, from the barcode at the 3' end and the fragment with a potential CRM from the 5' end. This allows for the association of each fragment with a specific barcode. This baseline sequencing serves two additional purposes. One is normalization of the barcode reads recovered from electroporated tissues or cells to their frequency within the electroporated library, ensuring that barcode enrichments reflect true CRM activity rather than variability in fragment representation. The second is that it allows mapping of each fragment to the BAC and an assessment of library coverage across the BAC.

As an initial test of the LS-MPRA approach, BAC clones containing genes previously analyzed for CRM activity were selected. These genes are stably expressed in specific cell types of the mature retina. They were *Rho* (rhodopsin), expressed in rod photoreceptors, *Cabp5* (calcium-binding protein 5), expressed in subsets of cone bipolar cells and in rod bipolar cells (Haeseleer et al., 2000, Shekhar et al., 2016), *Grm6* (metabotropic glutamate receptor 6), expressed in all ON-bipolar cells (Ueda et al., 1997), and *Vsx2* (visual system homeobox 2; homolog of *Chx10*), expressed in all bipolar cells and some Mueller glial cells (Hatakeyama et al., 2001, Liu et al., 1994, Rowan & Cepko, 2004). LS-MPRA libraries for each of the BACs encoding these genes were made as described above (library characteristics in Table S3 and Supplemental Document A). The libraries were pooled and electroporated into postnatal day 3 (P3) retinas *in vivo* or *ex vivo*, with sacrifice and/or tissue collection 7 days later. This developmental window was chosen as it is the period when rod photoreceptors and bipolar cells are generated from RPCs, and RPCs are readily electroporated, while postmitotic cells are not. The *Rho* library was present at 5% the level of the other libraries as rods are the predominant cell type produced at this time (Young, 1985), and *Rho* is a highly expressed gene. This and all LS-MPRA experiments in this study were




Figure 1. Overview of the LS-MPRA.

(A) Schematic representation of LS-MPRA. BACs containing genomic regions of interest are enzymatically fragmented to generate a high-complexity DNA library. Size selected fragments are cloned into a vector containing a minimal promoter that will drive expression of GFP and a barcode positioned within the 3' untranslated region (UTR). The LS-MPRA library is then electroporated into retinal cells, where CRM activity is inferred by quantifying barcode enrichment in transcribed mRNA. (B) Illustration of fragment-barcoding strategy and necessary elements to sequence, wherein each DNA fragment is uniquely barcoded. (C) Preparation and sequencing of the plasmid library, associating fragment-barcode pairs, and establishing a baseline for barcode abundance, which is used to normalize barcode counts after mapping of barcode-labeled fragments to genomic coordinates. ORF, open reading frame; BC, barcode.

performed using three replicates, with each replicate generated by pooling retinas from several animals. Three retinas were pooled following *in vivo* electroporation, and 3–4 retinas were pooled following *ex vivo* electroporation and culture.

Analysis of the sequences of the fragment-association libraries showed that all regions of the BACs were fairly equally represented when aligned to the mouse mm10 reference genome (Library Coverage plots; Figs. 2 [A](#), 2S [A](#)). When the frequencies of cDNA barcodes representing different genomic regions of each BAC were plotted across the genomic regions, clear signals were observed for all 4 BACs (LS-MPRA plots; Figs. 2 [A](#), 2S [A](#)). The barcode frequency plot for the Rho LS-MPRA library revealed distinct candidate CRM regions (peaks; Fig. 2A [A](#)) in one, or both, of the *in vivo* and *ex vivo* conditions. Several of these regions are conserved across vertebrate and placental mammal species, correspond to areas of open chromatin, as indicated by ATAC-seq data (Lyu et al., 2021 [A](#)), and are annotated as candidate *cis*-regulatory elements in the mouse ENCODE database (not shown; Consortium et al., 2020 [A](#)). Consistent with previous reports identifying the Rho proximal promoter region (PPR) at –176bp to +70bp relative to the *Rho* transcription start site (TSS) (Zack et al., 1991 [A](#)), the *in vivo* Rho LS-MPRA revealed a clear CRM peak at –251bp to +65bp (peak **iii**; Fig. 2A [A](#)). The expanded region of interest (ROI; Fig. 2B [A](#)) clearly demarcates the CRM peak (peak **iii**), which correlates with the ATAC-seq open chromatin profiles. Two additional Rho CRMs known to induce reporter expression *in vitro*—the rhodopsin enhancer region (RER; –1.575 kb to –1.476 kb) and the Crx bound region (CBR; –3.165 kb to –2.693 kb) (Corbo et al., 2010 [A](#), Nie et al., 1996 [A](#))—exhibited minor peaks only in the *ex vivo* LS-MPRA (peaks **iv**, **v**).

In addition to the CRM peaks directly associated with *Rho*, two peaks were identified by the Rho LS-MPRA at approximately –123 kb (peak **i**) and –78 kb (peak **ii**) upstream of the *Rho* TSS. These peaks localize within the putative promoter regions of *Rpl32* (a ribosomal RNA gene) and *Ift122* (encoding an intraflagellar transport protein), respectively, suggesting that the LS-MPRA captured regulatory activity in addition to that of *Rho*. To investigate this possibility, we analyzed publicly available single-cell RNA sequencing (scRNA-seq) data from P8 mouse retina (Clark et al., 2019 [A](#)). In this dataset of 10,200 retinal cells, *Rpl32* was detected in 7,916 cells and *Ift122* in 1,092 cells, while *Rho* was detected in 8,809 cells. The fragment immediately upstream of *Rpl32* was cloned into a plasmid upstream of a TATAA minimal promoter and EGFP reporter, and electroporated *in vivo* into P3 retinas and incubated for 7 days before analysis. GFP expression driven by this fragment was expressed in most electroporated cells (data not shown). Follow up experiments might reveal that this small CRM (237 bp) is active in most tissues, suggesting it would be useful for applications where such breadth of expression driven by a small CRM is desirable. These expression patterns support the notion that some of the CRM activity captured by the Rho LS-MPRA reflects regulatory elements associated with neighboring genes that are co-expressed in retinal cells during this stage of development.

LS-MPRA plots for bipolar cell subtype markers also revealed several relevant peaks. In the Grm6 LS-MPRA, a distinct peak at –8.152kb to –7.857kb relative to the TSS (peak **i**, Fig. 2SA [A](#)) overlapped with a previously identified CRM (–8.126kb to –7.926kb; (Kim et al., 2008 [A](#))), which is moderately conserved across vertebrate and placental mammal species and localizes to an area of open chromatin. In the Vsx2 LS-MPRA, two distinct peaks at –37.617kb to –37.364kb and at –17.833kb to –17.559kb relative to the TSS (peaks **ii**, **iii**; Fig. 2SB [A](#)) overlapped with a Vsx2 enhancer active in RPCs (–37.726kb to –37.117kb (Buenaventura et al., 2018 [A](#)) or –37.778kb to –36.813kb (Honnell et al., 2022 [A](#))) and a Vsx2 enhancer active in bipolar cells (–17.724kb to –17.560kb (Kim et al., 2008 [A](#)) or –19.654kb to –15.836kb (Honnell et al., 2022 [A](#))), respectively. Both peaks are conserved across vertebrate and placental mammal species and localize to areas of open chromatin. However, a broad region previously described as a Vsx2 enhancer (–2.145kb to +87bp; (Rowan & Cepko, 2005 [A](#))) did not align with any peaks in the LS-MPRA. Similarly, no peaks in the Cabp5 LS-MPRA (Fig. 2SC [A](#)) corresponded to the known Cabp5 CRM at –289bp to +156bp relative to the TSS (Kim et al., 2008 [A](#)). These results suggest that while LS-MPRA successfully identified several previously characterized CRMs, some known elements did not exhibit significant activity under these experimental conditions.




Figure 2. Rho LS-MPRA to identify CRMs in the neonatal mouse retina.

(A) LS-MPRA barcode enrichment plots for the Rho locus aligned to a genome browser track from *in vivo* and *ex vivo* experiments (N=3 experimental replicates, combined), and annotated with: (i) known *Rho* regulatory regions, (ii) the coverage of the barcode-fragment association library across the locus, (iii) log₂-transformed base conservation among 60 vertebrate and 40 placental mammal species, (iv) regions of open chromatin in the P8 mouse retina, and (v) RefSeq gene models for the locus. (B) Expanded view of a region of interest from the Rho LS-MPRA plot, showing peaks (i-v) that correspond to known regulatory regions (red bars), genomic conservation, and areas of open chromatin.

LS-MPRA for genes expressed dynamically during retinal development

We are interested in how different RPCs produce different types of retinal cells. Previous work has suggested that there are intrinsic differences among different RPCs over time, and even at one time in development (Cepko, 2014 [↗](#)). We previously found that *Olig2*, a transiently expressed proneural basic helix-loop-helix (bHLH) transcription factor, marks a subset of RPCs at each time in development (Hafler et al., 2012 [↗](#), Shibasaki et al., 2007 [↗](#), Trimarchi et al., 2008 [↗](#)). It is expressed in neurogenic RPCs that divide one or two times, and produce a limited repertoire of cell types among those born at that time. In the embryonic day 13 or 14 (E13–14) retina, cone photoreceptors and horizontal cells, a type of interneuron, are produced. In the postnatal period, *Olig2*⁺ RPCs produce rod photoreceptors and amacrine interneurons. We were interested in the regulators of *Olig2* at these two different times, as different CRMs might lead us to the intrinsic differences between embryonic and postnatal RPCs. The *Olig2* LS-MPRA library was thus generated (library characteristics in Table S3 [↗](#)) and delivered to E14 mouse retinas, which were incubated *ex vivo* for 28 hours. For each of the three experimental replicates, 8 retinas were pooled. Barcode enrichment plots from this library revealed distinct candidate CRM regions (peaks; Fig. 3A [↗](#)), several of which are conserved across vertebrate and placental mammalian species and correspond to areas of open chromatin. Notably, two CRM candidate regions identified by LS-MPRA (Fig. 3B [↗](#)), located at –1.881 kb to –1.700 kb and –85.027 kb to –84.781 kb relative to the *Olig2* TSS, overlap with broader regions previously implicated as potential *Olig2* regulatory elements in mouse embryonic stem cells (Chen et al., 2008 [↗](#)). Other regions with possible regulatory activity (indicated in Fig. 3A [↗](#)), previously identified using *in vitro* assays or in the mouse spinal cord (Fan et al., 2023 [↗](#), Friedli et al., 2010 [↗](#), Sun et al., 2006 [↗](#)), did not exhibit CRM activity in the retina by LS-MPRA.

The peaks at –1.881 kb to –1.700 kb were quite small, so an additional assay was run to determine if they were relevant for *Olig2* regulation. Knock-out mice for *Notch1* have been shown to overproduce cone photoreceptors during the embryonic period, and rod photoreceptors in the neonatal period (Jadhav et al., 2006 [↗](#), Yaron et al., 2006 [↗](#)). Moreover, *Olig2* expression in mouse retinal explants is also transiently upregulated in response to Notch inhibition (Kaufman et al., 2019 [↗](#)). To investigate the precise dynamics of Notch regulation of *Olig2* in the E14 retina, as well as to determine if the small CRMs identified in the LS-MPRA were of relevance to *Olig2* regulation, E14 retinal explants were treated with the γ -secretase inhibitor, LY411575, which inhibits Notch signaling. Notch inhibition was found to increase endogenous *Olig2* RNA levels in E14 retinas over 24 hours of treatment, with a peak at 12 hours (Fig. 4A [↗](#)). Two small *Olig2* CRM candidates, one just 5' (–1.881 kb to –1.700 kb) and one 3' (+6.721 kb to +6.933 kb) of the *Olig2* TSS, also exhibited an increase in activity, also with a peak at 12 hours post Notch inhibition (Fig. 4B [↗](#)). In addition, one CRM within the intron of the *Gm41492* predicted gene, at –84.781 kb to –85.027 kb relative to the *Olig2* TSS, also exhibited an increase in CRM activity with Notch inhibition. The three *Olig2* potential CRMs, now named Notch signaling-responsive (NR) regions 1, 2, and 3, were conserved, and exhibited open chromatin. Reanalysis of one replicate using only unique barcodes also identified these three peaks, including NR3, indicating that barcode collisions did not account for the minor NR3 peak (see Supplemental Document B). Together, these findings suggest that NR1–3 function as CRMs that respond to disruptions in Notch signaling to enhance *Olig2* expression in RPCs that produce cone photoreceptors. More broadly, these experiments demonstrate that the LS-MPRA can be used to rapidly identify CRMs that respond to perturbations that are relevant to the gene under study.

Analysis of cell type-specific CRM activity

To investigate the cellular specificity of the *Olig2* CRMs, NR1, NR2, and NR3, each fragment was cloned individually or as a composite into a plasmid with the endogenous *Olig2* promoter and TSS (–75bp to +51bp) upstream of an EGFP reporter. First, to determine whether the inclusion of the *Olig2* promoter, and/or other sequences within the plasmid, had background transcriptional




Figure 3. Olig2 LS-MPRA to identify candidate CRMs in the developing mouse retina.

(A) Barcode enrichment plot from the Olig2 LS-MPRA, aligned with a genome browser track and annotated with (i) previously described *Olig2* regulatory regions identified in mouse embryonic stem cells, fertilized murine oocytes, a mouse lymphoma cell line, or in mouse ventral neural tube (Chen et al., 2008, Fan et al., 2023, Friedli et al., 2010, Sun et al., 2023), (ii) coverage of the barcode-fragment association library across the locus, (iii) log₂ base conservation across 60 vertebrate or 40 placental mammal species, (iv) regions of open chromatin in the E14 mouse retina, and (v) RefSeq gene models. (B) Two expanded regions of interest from the Olig2 LS-MPRA, showing peaks that align with known regulatory elements, genomic conservation, and open chromatin. Candidate CRMs identified within blue bars.




Figure 4. Dynamic CRM activity and *Olig2* expression following inhibition of Notch signaling.

(A) Bar plot showing relative endogenous *Olig2* RNA expression in the retina over 0–24 hours of treatment with the γ -secretase inhibitor LY411575 or DMSO, normalized to the 0-hour control. Unpaired *t*-tests with Holm’s multiple comparisons correction were performed for each Control vs. Treated timepoint. (B) Barcode enrichment plot from the *Olig2* LS-MPRA, aligned with a genome browser track after 0–28 hours of LY411575 treatment. Notch inhibitor-responsive CRM candidate regions (NR1–3, blue boxes) displayed differential barcode enrichment across the treatment period. Statistical analysis was performed using one-way Welch’s ANOVA with Dunnett’s T3 multiple comparisons test on AUC values, normalized to the 0-hour timepoint. Error bars in (A) and (B) represent SEM. Asterisks indicate statistical significance (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$).

activity, we electroporated backbone plasmids containing GFP alone or GFP under the Olig2 minimal promoter into E14 retinas (Fig. 5A). Minimal and sporadic GFP RNA signal was observed in these controls. This background level was judged to be low enough to allow for assay of specific CRM activity. The Olig2 CRMs were cloned into the backbone plasmid, oriented as they appear in the genome relative to the TSS. These constructs were electroporated into E14 retinas *ex vivo* and incubated for 24 hours before analysis (Fig. 5A).

Electroporated cells were identified by expression of the CAG-BFP co-electroporated control plasmid. Immunolabeling for GFP and Olig2 in transverse retinal sections (Fig. 5B) revealed that all Olig2⁺ electroporated cells, identified by immunohistochemistry (IHC), expressed GFP, also identified by IHC, when driven by NR1, NR2, or the combined NR1-3 composite; additionally, most Olig2⁺ electroporated cells expressed GFP when driven by NR3 alone, albeit at lower levels (Fig. 5C, left column; Table S5). These findings suggest that NR1, NR2, and NR3 contain functional CRMs for Olig2, with NR3 exhibiting very weak CRM activity. However, when the GFP⁺ electroporated cells were examined for Olig2 protein expression using IHC, few GFP⁺ cells demonstrated detectable Olig2 protein (Fig. 5C, middle column; Table S5), and many Olig2⁺ electroporated cells expressed GFP (Fig. 5C, right column; Table S5). This lack of correlation raised two possibilities: either NR1-3 CRM activities were not confined to Olig2⁺ RPCs, and/or the transient nature of Olig2 expression, combined with the stability of EGFP, resulted in a temporal mismatch that limited the detection of co-localized Olig2 protein and GFP protein. This possibility was suggested by the known short half-life of murine Olig2 RNA (~1.66hrs) and protein (~3.5hrs) (Rayon et al., 2020).

We first determined if the short half-lives of Olig2 RNA and protein created a situation where detection of the endogenous Olig2 transcript and protein in the same cell was unlikely. Fluorescent *in situ* hybridization (FISH) for Olig2 transcript, combined with immunolabeling for Olig2 protein, showed that most cells expressed either the transcript or the protein, but not both simultaneously (Fig. 6A). A similar pattern was observed when comparing GFP protein expression and GFP transcript levels in retinas electroporated with the NR1 CRM (Fig. 6B). This observation might indicate that the NR1 CRM activity is also dynamically regulated. To at least partially accommodate this temporal mismatch when assessing the specificity of the Olig2 CRMs, we examined RNA, rather than protein, for Olig2 and GFP, reasoning that RNA might be a more accurate read-out of both the CRM activity and endogenous Olig2 transcription.

FISH for Olig2 and GFP RNA, along with intrinsic GFP fluorescence (Fig. 6C), revealed a pattern similar to the quantification of immunostaining (Fig. 5C). The vast majority of Olig2 RNA⁺ electroporated cells expressed GFP RNA and/or protein when GFP expression was driven by NR1, NR2, or the composite construct (Fig. 6D, left column; Table S6). Approximately 40% of Olig2 RNA⁺ electroporated cells co-expressed GFP when driven by NR3 alone, further supporting its lower but detectable CRM activity. Importantly, a substantial proportion of GFP RNA⁺ cells co-expressed Olig2 RNA when GFP was driven by NR1, NR2, or the composite construct (Fig. 6D, middle-left column; Table S6), and this proportion exceeded that observed when the analysis was limited to GFP protein⁺ cells (Fig. 6D, middle-right column; Table S6). When the expression of GFP RNA or protein was compared in cells that expressed or did not express Olig2 RNA, a greater fraction of Olig2⁺ cells expressed GFP when it was driven by any of the four constructs (Mann-Whitney test with Holm-Šidák method for correcting multiple comparisons, $p < 0.05$ each). These findings indicate a significant enrichment of CRM activities in the Olig2⁺ cells. In addition, these findings suggest that measuring co-localization at the RNA level may provide a more accurate assessment of CRM activity for transiently expressed genes.

D-MPRA for identification of sequences important for CRM activity

To identify nucleotides within potential CRMs that are necessary for their activity, we developed the d-MPRA. Most nucleotides identified are suspected to be within TF binding sites, though other roles, e.g. sites that affect chromatin structure, might also play a role. Libraries with point mutations were created via error-prone PCR amplification of each NR1-3 CRM. Each library was then cloned into a vector backbone. An intraplasmid duplication (IPD) was then made to serve as a




Figure 5. Co-localization of GFP driven by Olig2 CRMs and Olig2 in retinal cells.

(A) Schematic of plasmid constructs containing one or all Notch inhibitor-responsive regions driving GFP expression with the endogenous *Olig2* minimal promoter, alongside the experimental workflow. (B) Representative transverse sections of E14 retinas electroporated with these plasmids, incubated *in vitro* for 24 hours, and stained using IHC for GFP and Olig2. Merged and single-channel images show GFP colocalization with Olig2 (orange arrow) as well as GFP⁺ cells lacking detectable Olig2 expression (yellow arrow). (C) Quantification of GFP and Olig2 co-localization, represented as pie charts showing (i) the percentage of electroporated Olig2⁺ cells that express GFP protein, (ii) the percentage of GFP protein⁺ cells that express Olig2, and (iii) the percentage of electroporated Olig2-cells that express GFP protein in retinas electroporated at E14 with plasmids containing one or all Notch inhibitor-responsive regions driving GFP. Scale bar: 20μm.




Figure 6. Co-localization of CRM-directed RNA and *Olig2* RNA or protein in retinal cells.

(A–C) Representative transverse sections of E14 retinas incubated *in vitro* for 24 hours for analysis of localization patterns of *Olig2* and/or GFP. (A) Sections stained for *Olig2* protein (cyan, yellow arrows) using IHC and *Olig2* RNA (red, magenta arrows) using FISH. A 100% stacked column plot quantifies the overlap between *Olig2* protein and *Olig2* RNA expression. (B) Intrinsic GFP fluorescence and *GFP* RNA driven by *Olig2*-NR1 (yellow and magenta arrows). (C) In retinas electroporated with plasmids containing one or all Notch inhibitor-responsive regions, *Olig2* RNA co-localized with GFP protein only (yellow arrow), *GFP* RNA only (magenta arrow), or both (orange arrow). (D) Pie charts quantifying (column 1) the percentage of electroporated *Olig2* RNA+ cells that expressed GFP, (column 2) the percentage of *GFP* RNA+ cells that expressed *Olig2* RNA, (column 3) the percentage of GFP protein+ cells that expressed *Olig2* RNA, and (column 4) the percentage of *Olig2* RNA-cells that expressed GFP in retinas electroporated. Scale bar: 20µm.

barcode before inserting a GFP reporter into the completed library (Fig. 7A). In the IPD process, the mutant library was nicked on the plus strand upstream and the minus strand downstream of the mutated CRM fragments. Strand-displacing polymerase, lacking exonuclease activity, extended these nicks, linearizing the libraries such that each end carried an exact copy of a particular mutated fragment. The final step involved inserting a minimal promoter and GFP ORF cassette between the two duplicated CRM copies. The 3' copy of the mutated CRM was thus in the 3' UTR, and served as a barcode in the cDNA (Fig. 7A). This configuration allowed the possibility that the 5' and/or 3' mutated CRM could influence transcription of GFP. In addition, it was possible that the 3' fragment could influence the stability of the transcript. To somewhat override this effect, a WPRE sequence was also included in the 3'UTR of all library members. The IPD method was used so that the step of making and sequencing the association library, as was done for the LS-MPRA, was unnecessary.

Mutations reducing positive activity, presumed TF binding that enhances activity, should result in depleted barcode abundance, while those diminishing repressor TF binding should lead to enrichment (Fig. 7B). Similar to LS-MPRAs, the abundance of barcodes of mutated fragments was normalized to their baseline abundance in the d-MPRA plasmid libraries.

The d-MPRA libraries for Olig2 NR1-3 CRMs were constructed with a mutational frequency of ~0.02–0.2% per nucleotide and a complexity of at least 2×10^7 transformants (see Supplemental Document A for mutational rate, co-occurrence across CRMs, and analyses comparing bulk fragments vs singleton mutations). These libraries were electroporated into E14 *ex vivo* retinas, which were incubated for 24 hours before RNA extraction. The 3'UTR region containing transcribed mutated CRMs was reverse transcribed to generate cDNA, PCR amplified, gel purified, indexed, and sequenced. Normalized log₂-transformed values were averaged across replicates, and a sliding window average (window size = 5) was applied to smooth the data and reduce noise. This approach ensured robust comparisons across experimental conditions. The resulting d-MPRA plots for Olig2-NR1 (Fig. 7C), Olig2-NR2 (Fig. 7D), and Olig2-NR3 (Fig. 7E) revealed peaks and troughs, indicating that mutations at multiple sites influence CRM activity. In three independent electroporation experiments for each CRM, the d-MPRA readouts exhibited low variance between experimental replicates. The clear alignment of peaks and troughs with specific nucleotide positions supports the notion that these regions are important for CRM activity.

To identify over-represented sequences in sites corresponding to peaks or troughs in the d-MPRA plots, *de novo* motif discovery analysis, HOMER, was performed. This analysis focused on all three Olig2 CRMs, with an emphasis on NR2, which is proximal to the Olig2 TSS. Across the Olig2-NR2 region, 16 *de novo* motifs were identified (Fig. 8A), several of which overlapped with d-MPRA peaks or troughs. The presence and density of binding motifs in this region are consistent with the open chromatin profile observed in ATAC-seq data. Notably, motifs matching known binding sites for Foxn4, Pax6, Mybl1, and Otx2—recognizable regulators of retinal development (Cvekl & Callaerts, 2017, Farhy et al., 2013, Kaufman et al., 2019, Li et al., 2004, Marquardt et al., 2001, Nishida et al., 2003, Oron-Karni et al., 2008, Remez et al., 2017, Wang et al., 2014)—were detected (Figs. 8B–D). Mybl1 and Otx2 motifs (Motifs 1 and 14, respectively) corresponded to troughs in the d-MPRA plot, whereas the shared motif (Motif 3) for Foxn4 (typically an enhancer) and Pax6 (frequently a repressor) did not align with any prominent peaks or troughs (Fig. 8A). The shared motif for Foxn4 and Pax6 in Olig2-NR2 likely represents a regulatory balance between activation and repression, reflecting the complex interplay of these TFs in fine-tuning Olig2 expression.

To assess whether RPCs that express Olig2 could use these TFs to regulate Olig2, scRNAseq data (Clark et al., 2019) were examined for co-expression with Olig2 at E14. The dataset, originally generated using the 10X Genomics platform, was further processed in Seurat, where filtering, normalization, and dimensionality reduction was carried out to identify distinct retinal cell populations (Fig. 8E). As expected, Olig2 expression was enriched in a subset of neurogenic RPCs (Fig. 8F). Notably, Mybl1, Pax6, Foxn4, and Otx2 were co-expressed with Olig2 in subsets of these cells (Fig. 8G), with 37.7%, 55.5%, 30.7%, and 67.3% of neurogenic RPCs that expressed Olig2 also expressed Mybl1, Pax6, Foxn4, and Otx2, respectively. The expression of Mybl1, Foxn4, and Otx2




Figure 7. Degenerate MPRA to identify functional residues within candidate CRMs.

(A) Schematic of d-MPRA library assembly. Point mutations were introduced into *Olig2* CRM fragments via error-prone PCR, followed by intraplasmid duplication (IPD) to generate constructs with duplicated mutant CRMs flanking a minimal promoter and GFP ORF. The 3' CRM copy, located in the 3' UTR, served as a barcode, with a WPRE sequence included to potentially stabilize transcripts. (B) Conceptual diagram illustrating expected d-MPRA results, showing predicted changes in CRM activity upon disruption of enhancer or repressor binding sites. (C-E) d-MPRA plots displaying $\log_2$ fold changes in mutational frequencies using a 5-base pair sliding window average, normalized across the *Olig2*-NR1 (C), *Olig2*-NR2 (D), and *Olig2*-NR3 (E) regions.




Figure 8. TF binding sites within the Olig2-NR2 CRM.

(A) TF binding motifs predicted using HOMER identified within the *Olig2*-NR2 CRM candidate, aligned with the average d-MPRA plot for this region (from Fig. 7). (B-D) Position frequency matrices of TF binding sites aligned to motifs predicted by HOMER in *Olig2*-NR2: Mybl1 to Motif 1 (B), Foxn4 and Pax6 to Motif 3 (C), and Otx2 to Motif 14 (D). (E) UMAP visualization of scRNA profiles from E14 mouse retinas, with cell types previously identified by marker genes by Clark et al. (2019). (F) Expression of *Olig2* on UMAP. (G) Co-expression (pink) of *Olig2* (blue) with *Mybl1*, *Foxn4*, *Pax6*, or *Otx2* (red) on UMAP. Percentages of respective populations that co-express *Olig2* are indicated (* denotes significance). (H) TF occupancy track aligned to (i) *Olig2* LS-MPRA barcode enrichment after 12-hour LY411575 treatment (replicate of Fig. 4B) and (ii) gene models. Occupancy peaks for Mybl1, Foxn4, and Pax6 align with the Olig2-NR2 CRM candidate (blue box).

were significantly enriched in Olig2⁺ RPCs (Fisher's exact test Odds Ratios [OR]: 11.8 [p<0.0001], 7.1 [p<0.0001], and 8.5 [p<0.0001], respectively), whereas Pax6 (OR: 0.96, p=0.31) was not, perhaps reflecting a role as a repressor. These results support the hypothesis that these factors are functionally intertwined in the regulation of Olig2 expression.

To examine whether the suggested Olig2 regulatory TF's bind to the Olig2-NR2 CRM, a CUT&RUN analysis was carried out. A significant enrichment of Mybl1, Pax6, and Foxn4 binding, with peaks that aligned with the NR2 region (Fig. 8H [8H](#)), provided evidence for their direct regulatory roles via the Olig2-NR2 CRM. Together, these findings suggest that the Olig2-NR2 CRM is a target for multiple retinal developmental regulators, potentially mediating Olig2 expression in neurogenic RPCs.

Using *de novo* motif discovery analysis on Olig2-NR1, 20 motifs were identified (Fig. 8SA [8SA](#)), some of which correspond to binding sites for several retinal TFs, including Sox4, Sox11, Lhx2, Dlx2, Isl1, Foxp1, Mybl1, and Otx2. These factors play critical roles in retinal development: Sox4 and Sox11 are active in neurogenic RPCs and regulate the generation and survival of RGCs (Jiang et al., 2013 [2013](#)); Lhx2 is suggested to influence the progression of RPC competence states (Gordon et al., 2013 [2013](#)); Dlx2 is required for terminal RGC differentiation (de Melo et al., 2005 [2005](#), Zhang et al., 2017 [2017](#)); Isl1 is essential in the gene regulatory network governing RGC development (Mu et al., 2008 [2008](#)); Foxp1 has been implicated in regulating the competence of RPCs to generate early-born retinal cell types (Zhang et al., 2023 [2023](#)); and Mybl1 and Otx2 are known to influence neural progenitor cell cycle regulation (Kaufman et al., 2019 [2019](#)), and photoreceptor and horizontal cell development (Nishida et al., 2003 [2003](#), Wang et al., 2014 [2014](#)), respectively.

In the d-MPRA analysis, specific motifs in Olig2-NR1 aligned with these predicted TF binding sites. For instance, Motifs 13 and 16 matched shared binding sites for Sox4 and Sox11 (Fig. 8SB [8SB](#)); scRNA-seq co-expression data of RPCs (Fig. 8SG [8SG](#)) revealed that 74.1% of Olig2-expressing neurogenic RPCs co-expressed Sox4 (enrichment OR: 5.3 [p<0.0001]) and 89.3% co-expressed Sox11 (enrichment OR: 3.1 [p<0.0001]). In contrast, Motif 11 contained a shared binding site for both Lhx2 and Dlx2 (Fig. 8SC [8SC](#)); among Olig2⁺ neurogenic RPCs, 29.6% co-expressed Lhx2 (enrichment OR: 1.0, p=0.94) and 21.1% co-expressed Dlx2 (enrichment OR: 3.7 [p<0.0001]). Motif 8 matched an Isl1 binding site (Fig. 8SD [8SD](#)) and was associated with only 5.0% co-expression in Olig2⁺ RPCs (Fig. 8SG [8SG](#)) (enrichment OR: 0.33 [p<0.0001]), implying it might suppress Olig2 expression. Motif 18 contained a Foxp1 binding site (Fig. 8SE [8SE](#)), with 40.9% of Olig2⁺ neurogenic RPCs co-expressing Foxp1 (enrichment OR: 0.81 [p<0.0001]), as shown in Fig. 8SG [8SG](#). Additionally, binding sites for Mybl1 and Otx2 were identified in Motifs 2 and 12, respectively (Fig. 8SF [8SF](#)); scRNA-seq data (Fig. 8G [8G](#)) indicate that these TFs may also contribute to the regulation of Olig2 expression via Olig2-NR1. Taken together, and considering the alignment of motifs with d-MPRA peaks and troughs as well as the quantified co-expression data from Figs. 8SG [8SG](#) and 8G [8G](#), these results suggest that Sox4, Sox11, Dlx2, and Isl1 may predominantly function to repress Olig2 expression. Lhx2, Foxp1, Mybl1, and Otx2 are likely to act as activators through the Olig2-NR1 CRM.

The *de novo* motif discovery analysis of Olig2-NR3 identified 19 motifs (Fig. 8SH [8SH](#)), several of which aligned with peaks or troughs in the d-MPRA plot. Among these, four motifs containing known TF binding sites active in the developing retina were associated with troughs, suggesting potential enhancer activity. Motif 13 matched a binding site for Bhlhb5 (encoded by *Bhlhe22*) (Fig. 8SI [8SI](#)), a TF involved in retinal interneuron subtype specification (Feng et al., 2006 [2006](#), Huang et al., 2014 [2014](#)); scRNA-seq co-expression data (Fig. 8SL [8SL](#)) revealed that among Olig2⁺ RPCs, 21.6% co-expressed Bhlhb5 (enrichment OR: 3.5 [p<0.0001]). Motifs 3 and 5 contained binding sites for Ngn2 (Fig. 8SJ [8SJ](#)), a TF expressed in neurogenic RPCs during both early and late retinal development (Hufnagel et al., 2010 [2010](#), Kowalchuk et al., 2018 [2018](#), Ma & Wang, 2006 [2006](#)); 81.4% of Olig2-expressing RPCs co-expressed Ngn2 (Fig. 8SL [8SL](#)) (enrichment OR: 15.4 [p<0.0001]). Motif 15 matched a binding site for Lhx9 (Fig. 8SK [8SK](#)), a TF expressed in early retinal development that is necessary for interneuron subtype specification (Balasubramanian et al., 2014 [2014](#), Balasubramanian et al., 2018 [2018](#)). scRNA-seq analysis (Fig. 8SL [8SL](#)) established that 7.7% of RPCs expressing Olig2 co-expressed Lhx9 (enrichment OR: 3.3 [p<0.0001]). The combined scRNA-seq analysis, d-MPRA, and

motif discovery analysis confirmed that there are likely Bhlhb5, Ngn2, and Lhx9 binding sites in relevant regions of Olig2-NR3, and that they exhibit some degree of co-expression with Olig2, supporting the hypothesis that these TFs may regulate Olig2 expression via the Olig2-NR3 CRM.

The d-MPRA and motif analyses reveal a complex transcriptional network potentially regulating Olig2 expression, with some of the same TFs as candidate regulators across Olig2-NR1, NR2, and NR3 CRMs. Olig2-NR2 emerged as potentially the most important regulatory hub, where motif discovery, scRNA-seq enrichment, and CUT&RUN validation confirmed direct binding of Mybl1, Pax6, and Foxn4. The d-MPRA approach provided a robust, high-resolution method for nominating functionally relevant TF binding sites, with motif-associated peaks and troughs aligning with known retinal regulators. The identification of additional candidates in Olig2-NR1 and NR3 provides candidates to expand our understanding of the gene regulatory network governing Olig2 expression, setting the stage for future *in vivo* studies to dissect their functional roles in retinal development.

LS-MPRAs for Neurogenin-2 in the developing mouse retina

To further test the LS-MPRA in identifying CRMs regulating transiently expressed genes of regulatory importance in the retina, *Ngn2* was analyzed. Ngn2, a proneural bHLH TF, is expressed in a subset of RPCs, those that will produce at least one postmitotic cell (Hufnagel et al., 2010). Though known regulatory regions for Ngn2 have been elucidated in the ventral neural tube, where it also is expressed, their activities in the developing retina were unknown (Henke et al., 2009; Nakazaki et al., 2008; Scardigli et al., 2003; Scardigli et al., 2001; Simmons et al., 2001). The Ngn2 LS-MPRA library (library characteristics in Table S3) was delivered to the E14 mouse retina and processed through the LS-MPRA pipeline, as described for Olig2. The aligned barcode enrichment plot revealed distinct candidates for CRMs (Fig. 9A). Among these, Ngn2-CRM3 was highly conserved across vertebrate and placental mammal species and corresponded to an open chromatin region located -136 bp to +150 bp relative to the *Ngn2* TSS. It also is located in a 4.4kb fragment with regulatory activity in the embryonic mouse spinal cord (Simmons et al., 2001). To investigate the functionality of Ngn2-CRM3 and three additional CRM candidates identified in the LS-MPRA plot (Ngn2-CRM1, CRM2, and CRM4), each fragment was cloned into a plasmid containing a TATAA minimal promoter and EGFP reporter. These constructs were electroporated into E14 retinas and incubated *ex vivo* for 24 hours prior to analysis.

FISH for *Ngn2* and *GFP* RNA, alongside detection of intrinsic GFP fluorescence, in transverse retinal sections (Fig. 9B), demonstrated that Ngn2-CRM2 and Ngn2-CRM3 fragments consistently drove GFP expression in Ngn2+ cells. Co-localization analysis revealed that a substantial majority of *Ngn2* RNA+ electroporated cells expressed *GFP* RNA and/or protein when GFP expression was driven by these two CRMs (Fig. 9C, left column; Table S7). Approximately 57% of *Ngn2* RNA+ cells co-expressed *GFP* RNA when driven by Ngn2-CRM1, suggesting that this fragment may also function as a CRM, albeit less specifically for Ngn2. In contrast, GFP expression driven by Ngn2-CRM4 was sparse, and even fewer cells co-expressed *Ngn2* RNA, indicating that the LS-MPRA peak associated with this region does not correspond to a functional Ngn2 CRM in these conditions.

The proportion of *GFP* RNA+ cells co-expressing *Ngn2* RNA when GFP expression was driven by Ngn2-CRM2 or CRM3 (approximately 21% and 38%, respectively; Fig. 9C, middle-left column; Table S7) was lower compared to the corresponding co-expression analysis for Olig2 CRMs (Fig. 6). Similar to Olig2 CRMs, however, the proportion of *GFP* RNA+ cells driven by any Ngn2-CRM candidate region was consistently greater than that observed when the analysis was restricted to GFP protein+ cells (Fig. 9C, middle-right column; Table S7). These findings suggest that the rapid degradation of transient Ngn2 protein (~30mins half-life, (McDowell et al., 2014; Vosper et al., 2007)) and, likely, *Ngn2* transcript, limits the overlap between Ngn2 and GFP expression.

To address this limitation, the *ex vivo* incubation period was reduced from 24 to 16 hours and focused on Ngn2-CRM3, the most robust CRM candidate. Shortening the incubation period resulted in an increase in the proportion of *GFP* RNA+ cells co-expressing *Ngn2* RNA (38% vs. 49%; Fig. 9C,D, right column; Table S7). Further evidence for this is provided by the concurrent decrease in the proportion of *GFP* RNA+ cells within the population of *Ngn2* RNA-negative cells




Figure 9. Ngn2 LS-MPRA to identify CRM candidates active in mouse retinal cells expressing *Ngn2*.

(A) Ngn2 LS-MPRA barcode enrichment plot aligned with a genome browser track, annotated with (i) known *Ngn2* regulatory regions, (ii) coverage of the barcode-fragment association library, (iii) log₂ base conservation across 60 vertebrate or 40 placental mammal species, (iv) regions of open chromatin in E14 mouse retina, (v) gene models, and (vi) CRM candidate regions 1-4 (blue boxes). (B) Representative transverse sections of E14 retinas incubated for 24 hours *in vitro*, showing localization of *Ngn2* RNA with GFP RNA only (magenta arrow) or both intrinsic GFP and GFP RNA (orange arrows) in retinas electroporated with plasmids containing one of the CRM1-4 regions. (C) Pie charts showing the percentage of *Ngn2* RNA+ cells expressing GFP (column 1), GFP RNA+ cells expressing *Ngn2* RNA (column 2), GFP protein+ cells expressing *Ngn2* RNA (column 3), and *Ngn2* RNA+ cells expressing GFP (column 4) in retinas electroporated at E14 with plasmids containing CRM1-4 regions driving GFP. (D) Co-localization analysis of Ngn2-CRM3 plasmid and GFP following 16-hour incubation *in vitro*. Scale bar: 20µm.

when the incubation period was shortened (69% vs. 39%; Fig. 9C,D, right column; Table S7). This improvement supports the hypothesis that *Ngn2*'s transient expression window and rapid turnover obscured co-expression of *GFP* and *Ngn2* RNA.

LS-MPRAs in chick retina and spinal cord

To determine if LS-MPRAs can be used to identify CRMs across species and tissues, regulatory elements controlling chick *OLIG2* (*cOLIG2*) expression were carried out. Expression in the chick embryonic day 5 (E5) retina, a developmental stage equivalent to the E14 mouse retina, as well as in the chick spinal cord at two time points: E2 and E4, were examined with LS-MPRA. *Olig2* is expressed in spinal cord progenitor cells that give rise to motor neurons and oligodendrocytes (Lu et al., 2000, Zhou et al., 2000). Electroporating the *cOLIG2* LS-MPRA at E2 should capture the transcriptional activity of motor neuron progenitor cells (Hollyday & Hamburger, 1977), while delivery at E4 should capture the activity within oligodendrocyte precursor cells (Soula et al., 2001). The *cOLIG2* LS-MPRA library (library characteristics in Table S3) was electroporated into E5 chick retina explants and incubated *ex vivo* for 24 hours or electroporated into the E2 or E4 spinal cord and incubated *in ovo* for 24 hours or 48 hours, followed by LS-MPRA analysis.

Barcode enrichment plots from these experiments, aligned to the galGal6 chick reference genome, revealed distinct candidate CRM regions in both spinal cord and retina (Fig. 10A). One region, spanning -36.729kb to -36.428kb relative to the *cOLIG2* TSS, is highly conserved across vertebrates and orthologous to the *Olig2*-NR1 CRM identified in the mouse retina (97.3% sequence conservation shared across a 225bp region). Interestingly, this CRM candidate (*cOLIG2*-CRM1) was active in RPCs and in the E4 spinal cord but showed no detectable activity in electroporated E2 spinal cord. In addition to CRM1, two proximal CRM candidates—CRM2 (-264bp to -53bp) and CRM3 (+8.948kb to +9.036kb)—were identified in chick and assessed for their functional relevance to *cOLIG2* expression.

To determine whether *cOLIG2*-CRM1, CRM2, and CRM3 function as CRMs, each fragment was cloned into a plasmid containing the *cOLIG2* minimal promoter and an EGFP reporter. These constructs were electroporated into E5 chick retinas and incubated *ex vivo* for 24 hours prior to analysis. FISH for *cOLIG2* and *GFP* RNA, alongside intrinsic GFP protein expression in transverse retinal sections (Fig. 10B), demonstrated that *cOLIG2*-CRM1 and *cOLIG2*-CRM2 consistently drove GFP expression in *cOLIG2*⁺ cells. Co-localization analysis revealed that the majority of electroporated *cOLIG2* RNA⁺ cells expressed *GFP* RNA and/or protein when GFP expression was driven by these two CRMs (Fig. 10C, left column; Table S8). In contrast, GFP expression driven by *cOLIG2*-CRM3 was nearly absent, indicating that the LS-MPRA peak associated with this region does not correspond to a functional *cOLIG2* CRM in these conditions. Notably, the CRM3 peak overlaps with an *AscI* recognition site, a sequence rarely found in vertebrate genomes, which was used in the LS-MPRA design to facilitate barcode-fragment association (Fig. 1C). This likely led to an under-representation of this region in the association library, thereby generating a false positive due to normalization using the association library flanking the *AscI* site in the BAC.

The proportion of *GFP* RNA⁺ cells co-expressing *cOLIG2* RNA when GFP was driven by *cOLIG2*-CRM1 or CRM2 (Fig. 10C, middle-left column; Table S8) was nearly identical to the proportion of GFP protein⁺ cells (Fig. 10C, middle-right column; Table S8). This suggests a more synchronized developmental window of *cOLIG2* RNA and protein expression in chick compared to mouse (Fig. 6A). However, as observed in the mouse retina, a high proportion of *cOLIG2*-electroporated cells expressed *GFP* RNA (Fig. 10C, right column; Table S8), underscoring the challenges of detecting CRMs for transiently expressed transcription factors.

To further assess CRM activity in spinal cord progenitor cells, the *cOLIG2*-CRM constructs were electroporated into the ventral horn of E2 chick spinal cords (Fig. 10SA) and incubated *in ovo* for 24 hours. FISH for *cOLIG2* and *GFP* RNA in transverse spinal cord sections (Fig. 10SB) revealed that, similar to the chick retina, *cOLIG2*-CRM1 and *cOLIG2*-CRM2 drove *GFP* RNA expression in *cOLIG2*⁺ cells. However, both regions also exhibited GFP expression in *cOLIG2*⁻ cells, suggesting either that their CRM activity is not restricted to *cOLIG2*-expressing progenitors in the spinal cord,




Figure 10. OLIG2 LS-MPRA to identify CRM candidates in chick embryos.

(A) OLIG2 LS-MPRA barcode enrichment plot following electroporation into E5 chick retinal explants, and E2 spinal cords and E4 spinal cords *in ovo*, aligned with a genome browser track and annotated with (i) coverage of the barcode-fragment association library, (ii) $\log_2$ base conservation across 77 vertebrate species, (iii) gene models, and (iv) CRM candidate regions 1-3 (blue boxes). (B) Representative transverse sections of E5 chick retinas incubated for 24 hours *in vitro*, showing localization of chick OLIG2 RNA with GFP RNA only (magenta arrow) or both intrinsic GFP fluorescence and GFP RNA (orange arrows) driven by plasmids containing one of the CRM1-3 regions. (C) Pie charts showing the percentage of OLIG2 RNA+ cells expressing GFP (column 1), GFP RNA+ cells expressing OLIG2 RNA (column 2), GFP protein+ cells expressing OLIG2 RNA (column 3), and OLIG2 RNA-cells expressing GFP (column 4) in retinas electroporated at E5 with plasmids containing CRM1-3 regions driving GFP and visualized by FISH. Scale bar: 20 μ m.

or that low levels of *cOLIG2* RNA are undetectable in some GFP+ cells. No GFP expression was observed from cOLIG2-CRM3, further supporting the conclusion that this fragment does not function as a cOLIG2 CRM.

These findings highlight the ability of LS-MPRAs to identify orthologous CRMs that may regulate gene expression across species and developmental contexts. The conservation of cOLIG2-CRM1 with the mouse *Olig2*-NR1 CRM suggests a shared regulatory mechanism for *Olig2* in vertebrate retinal and oligodendrocyte precursor cells, while the species- and tissue-specific activities of other CRMs underscore the complexity of gene regulation across development and across species.

Olig2 CRMs active in embryonic RPCs are active in postnatal RPCs that produce rod photoreceptors

Thus far, we have primarily focused on the regulatory elements controlling *Olig2* expression in RPCs destined to produce cones and horizontal cells. Later in development, after all cone and horizontal cells have been generated, postnatal neurogenic RPCs express *Olig2* and produce primarily rod photoreceptors (Hafler et al., 2012 [↗](#)). An interesting question concerns whether these two distinct *Olig2*+ RPC populations share the same regulatory mechanisms governing *Olig2* expression. To address this, we electroporated plasmids containing *Olig2*-NR1, NR2, NR3, or all three CRMs, into P0 retinas and incubated them *in vivo* for 24 hours before analysis.

Immunolabeling for GFP and *Olig2* in transverse P1 retinal sections (Fig. 11A [↗](#)) and co-localization analysis revealed that the vast majority of *Olig2*+ electroporated cells expressed GFP when driven by NR1, NR2, NR3, or the combined NR1-3 composite (Fig. 11B [↗](#), left column; Table S9 [↗](#)). These findings indicate that NR1, NR2, and NR3 function as CRMs in RPCs that primarily produce rods in the postnatal retina. Notably, at this postnatal stage, most GFP+ electroporated cells also expressed *Olig2* protein (Fig. 11B [↗](#), middle column; Table S9 [↗](#)), suggesting that the temporal activation of these CRMs is more closely aligned with *Olig2* expression compared to their earlier activity in embryonic RPCs.

Since the majority of RPCs after P0 express *Olig2*, only a small number of *Olig2*-electroporated cells were present in the analysis. Despite their limited representation, most of these cells still exhibited GFP expression (Fig. 11B [↗](#), right column; Table S9 [↗](#)). Similar to the embryonic retina, this could reflect a temporal mismatch in the co-expression of *Olig2* and GFP or indicate a subset of *Olig2*-cells with CRM activity.

Overall, these findings suggest that the same regulatory elements govern *Olig2* expression in embryonic RPCs and the neonatal RPCs, as NR1, NR2, and NR3 serve as CRMs in both populations. The more synchronized co-expression of *Olig2* and GFP in postnatal RPCs further implies that *Olig2* regulation in late RPCs is more tightly coupled to its downstream gene expression programs compared to its earlier, less synchronous expression in early RPCs. This may be a reflection of slower developmental processes, as at least the cell cycle is longer in the later stages of retinal development (Alexiades & Cepko, 1996 [↗](#), Young, 1985 [↗](#)).

Discussion

In this study, we introduce novel LS-MPRA and d-MPRA approaches to search the *cis*-regulatory landscape of key developmental genes. By leveraging BAC-based libraries to survey large genomic regions and employing a degenerate MPRA strategy for nucleotide-resolution mapping, these methods overcome some of the limitations inherent to synthetic MPRA. This approach enables the unbiased identification of candidate CRMs while precisely delineating potential TF binding sites, thereby nominating both activators and repressors that regulate expression of a gene. These methods also offer a rapid and simple route to identify CRMs that can be deployed in applications where driving gene expression in a specific context is required. The scale, expense, and relatively simple molecular methods enable CRM discovery by any laboratory with basic molecular biology skills.




Figure 11. Activity of Olig2 CRM candidates in postnatal retinal cells expressing Olig2.

(A) Representative transverse sections of postnatal retinas electroporated at P0 and incubated *in vivo* for 24 hours, stained with antibodies against Olig2 and GFP to identify co-localized expression (orange arrows). GFP expression was driven by plasmids containing one or more Notch-inhibitor responsive regions (NR1-3). (B) Analysis of GFP and Olig2 co-localization, shown with pie charts depicting the percent of Olig2+ electroporated cells expressing GFP (column 1), the percent of GFP+ cells expressing Olig2 (column 2), and the percent of Olig2-electroporated cells expressing GFP in retinas electroporated at P0 with plasmids containing one or more Notch-inhibitor responsive regions driving GFP. Scale bar: 20µm.

LS-MPRA for stably expressed genes

The LS-MPRA was benchmarked using genes with known CRMs for stably expressed, cell type-specific genes. The experiments were carried out *in vivo* or in intact retinal explants, for discovery of CRMs relevant to authentic gene regulation in specific cell types. In postnatal retinas, CRMs were identified for the Rho PPR as well as some of the known CRMs for Grm6 and Vsx2. Interestingly, while the Rho LS-MPRA consistently detected a clear CRM peak at the PPR in both *in vivo* and *ex vivo* conditions, the previously identified RER and CBR Rho CRMs were seen only in *ex vivo* retinas. Of note, deletions of RER and CBR from the mouse genome showed that these two CRMs are dispensable for Rho expression, while the PPR is required (Sun et al., 2023 [DOI](#)). The LS-MPRA in *ex vivo* retinas displayed a smaller signal to noise ratio around the *Rho* locus, particularly for the RER and CBR regions, but not around other loci on this BAC, perhaps reflecting differences in the requirements for a specific tissue environment or time course for proper *Rho* expression. Another requirement for RER and CBR activity might be the use of the Rho PPR. Both the RER and CBR were shown to enhance expression in constructs that used the Rho promoter (Corbo et al., 2010 [DOI](#)), but neither had been tested using a heterologous promoter, as was used here. In the Vsx2 LS-MPRA, 2 of the 3 known CRMs were readily identified, with those 2 CRMs located quite some distance from the TSS, at -37 and -17 kb. It is unclear why the previously identified, promoter proximal CRM was not captured. One possible explanation is that the relatively short fragment lengths used to construct the libraries may not encompass the broader genomic context required for activation of this proximal CRM. Another potential explanation is again the choice of the promoter, as the Vsx2 promoter was used to identify the promoter proximal CRM in the previous study. The LS-MPRA for two other bipolar cell genes, Grm6 and Cabp5, also showed some positive and some negative results. Grm6 is expressed in the majority of retinal bipolar cells, and our previously described CRM for this gene, at -8kb, was identified by the LS-MPRA in both *ex vivo* and *in vivo* conditions. In contrast, the LS-MPRA failed to capture the known Cabp5 enhancer. It is not clear why this CRM was not identified here. Deeper sequencing of the cDNA libraries and/or using only a single BAC library for Cabp5, rather than the multiplexed format with 4 BACs used here, likely would increase the sensitivity. Sensitivity also should be increased by isolating electroporated cells of interest and/or pharmacologically or genetically increasing gene expression, as we did for Olig2. Finally, the choice of promoter may again play a role as the SV40 promoter was used in the earlier study for identification of the Cabp5 CRM.

LS-MPRA for transiently expressed genes

A focus of our laboratory is the mechanisms of cell fate determination in the retina. Our early study of RNA expression in RPCs showed that some TFs were expressed in subsets of RPCs, at one time and across the period of neurogenesis (Trimarchi et al., 2008 [DOI](#)). We are interested in whether heterogeneity in RPC gene expression accounts for differences in the types of cells produced. Olig2 was identified as expressed in subsets of RPCs. Using a retroviral lineage tracing method to mark clones derived from Olig2 RPCs, we found that they divide only once or twice, and produce a limited repertoire of cell types: cones and horizontal cells in the embryonic period and rods and amacrine cells in the postnatal period. Interestingly, intermingled with the embryonic Olig2+ RPCs are Olig-RPCs, which produce RGCs, the output neurons of the retina. To address how these Olig2+ RPCs might differ from Olig2-RPCs, our goal is to leverage Olig2 regulation. We thus developed the MPRA methods to identify candidates for its regulation, and to see if these regulators might vary over time, in correlation with the different daughter cell types produced at different times.

The LS-MPRA for Olig2 did indeed identify CRMs that lead to potential regulatory sites as well as their cognate TFs. However, it was difficult to validate the specificity of these CRMs, due to the short half-life of endogenous Olig2 RNA and protein. Nonetheless, when tested as single or combined CRMs in individual plasmid constructs, there was activity in a high proportion of Olig2+ cells. CRM activity in Olig2-cells may reflect a mix of cell states, such as those that have recently downregulated Olig2 but retain GFP, RPCs initiating CRM-driven GFP expression prior to detectable *Olig2* levels, or cells in which the CRM functions independently of Olig2 expression. In addition, several technical and biological considerations may contribute to the apparent

disconnect between reporter activity and endogenous Olig2 or Ngn2 expression. Introducing putative CRMs on plasmids removes them from their native genomic context, which may bypass normal chromatin accessibility constraints or local repressive influences and allow activity in cells where the endogenous gene is not expressed. These CRMs may also regulate genes other than Olig2 or Ngn2, reflecting broader or context-dependent roles. Finally, both Olig2 and Ngn2 are likely regulated by many more regulatory elements than the few tested here, so their endogenous expression may not depend solely on these specific CRMs. Despite GFP expression in Olig2-cells, several other lines of evidence increase confidence that Olig2-NR1, NR2, and NR3 are excellent candidates for specific Olig2 regulation. First, in the LS-MPRA, the same CRMs were found across 3 replicates, even though the signal to noise near the Olig2 locus was modest. The modest ratio is likely due to the fact that a modest or small percentage of RPCs express Olig2, reported to be ~14% in a previous study, (Brzezinski et al., 2011 [DOI](#)), or 2.3% in a more recent study (Clark et al., 2019 [DOI](#)). Additionally, and perhaps most importantly, it is expressed only briefly. To boost the signal, to potentially gain confidence in the candidate CRMs identified in the LS-MPRA, we were able to implement a drug manipulation. Notch inhibition greatly increases the production of cones (Jadhav et al., 2006 [DOI](#), Yaron et al., 2006 [DOI](#)), and here we show that this same inhibition leads to greater Olig2 expression, with a peak at 12 hours post inhibition. Importantly, the activity of Olig2-NR1, NR2, and NR3 was upregulated by Notch inhibition with the same magnitude and kinetics as seen for the endogenous Olig2 locus. This experiment shows that the LS-MPRA can be used to not only nominate CRMs for a gene, but to nominate CRMs that respond to perturbation, which might be particularly useful when the experiment is run in the full library format (i.e. all BAC fragments are assayed for a response). Additional evidence that Olig2-NR1, NR2, and NR3 are valid came from the d-MPRA and the follow-on experiments. The d-MPRA showed some of the same potential TF binding sites in more than one CRM, and several of these TFs have been shown to be important in retinal development, including TFs that play a role in cone and horizontal cell development, e.g. Otx2. In keeping with this, scRNAseq data show significant co-expression of Olig2 with the predicted TFs. Finally, CUT&RUN, using retinal tissue, showed binding of the suggested TFs to the CRMs. These data collectively provide support for the legitimacy of Olig2-NR1, NR2, and NR3 as key regulatory regions.

The binding of Mybl1 to Olig2-NR2 and potentially to Olig2-NR1, suggests that it plays a role in cone and/or horizontal cell development, likely as an activator of Olig2 expression. Its role in cone development was suggested previously by a study which tracked RNA expression in cultures induced to produce cones (Kaufman et al., 2019 [DOI](#)). In contrast, Pax6—a master regulator essential for maintaining RPC multipotency and guiding cell fate decisions—has been shown to function as both an activator and a repressor (Cvekl & Callaerts, 2017 [DOI](#), Farhy et al., 2013 [DOI](#), Marquardt et al., 2001 [DOI](#), Oron-Karni et al., 2008 [DOI](#), Remez et al., 2017 [DOI](#)); its binding to Olig2-NR2 may keep Olig2 off in multipotent RPCs, and/or keep Olig2 off in RPCs that produce RGCs. Meanwhile, Foxn4, which is critical for the specification of amacrine and horizontal cells (Li et al., 2004 [DOI](#)), may modulate Olig2 expression for control of the ratio of cone to horizontal cell fates. The overlapping motifs for Pax6 and Foxn4 in Olig2-NR2 may represent a region that balances activation and repression for modulation of early cell fate decisions.

Our results also reveal that different CRMs may be differentially engaged during retinal development. For instance, while Olig2-NR1 appears to integrate inputs from TFs such as Sox4, Sox11, Lhx2, Foxp1, Mybl1, and Otx2, Olig2-NR3 harbors binding sites for TFs including Bhlhb5, Ngn2, and Lhx9. As Sox4 and Sox11 binding sites might represent repressor sites, it is curious that they are co-expressed in 74% and 89% of Olig2+ cells. As they, as well as Lhx2 and Dlx2, play a role in RGC development, these expression patterns and roles in Olig2 regulation might represent a dynamic and complex gene regulatory network governing the production of the early born cell types. Consistent with this idea, and with the role of Bhlhb5 in later-born cell types, its binding site within the Olig2-NR3 may be engaged later in retinal development, considering a subset of bipolar cells exhibit a history of Olig2 expression (Hafler et al., 2012 [DOI](#)), and some mature bipolar cells maintain Olig2 expression. This nuanced and complex regulation underscores the broader utility of LS-MPRA and d-MPRA for discerning CRMs and TFs that are either cell type-specific or shared

across distinct developmental competence states. Future studies employing targeted perturbations, such as CRISPR/Cas9-mediated disruption of specific TF binding sites, will be essential to validate these CRMs and establish the roles that they play in gene regulation.

Considerations for use of LS-MPRA and d-MPRA

There are several issues to consider when choosing a method for CRM identification. Data describing open/active chromatin, conserved genomic sequences, and RNA expression, are increasingly comprehensive, but do not cover all cell types and non-model organisms. However, with the advent of single cell multi-omics, one can anticipate that these gaps will eventually be filled over the period of the next few years. When such data are available, and predict very few CRMs, one can simply PCR a potential CRM from genomic DNA, and test individually. In other cases, these types of data nominate a high number of potential CRMs. Producing many individual PCR products, cloning them into expression vectors, sequencing, introducing into cells, and assaying the results from single constructs, are much more time-consuming and expensive than LS-MPRAs. For example, we have identified potential CRMs for *Otx2*. Using predictions from open chromatin, eRNAs, and DNA conservation, >20 CRMs were nominated (Chan et al., 2020 [DOI](#), Emerson & Cepko, 2011 [DOI](#), Lonfat et al., 2021 [DOI](#)). Assessing each potential CRM and its relevant TF binding sites spanned more than several years. In addition to this level of effort, testing each potential CRM in isolation precludes a quick recognition of the relative strength of each CRM. Speed and expense also can be addressed using LS-MPRA by multiplexing libraries from several genes. Finally, when one needs a relatively small CRM, e.g. for expression in a viral vector, one can choose the fragment size during the BAC library preparation step. CRMs predicted by open chromatin typically do not provide this option.

One critical requirement for identification of CRMs using any method is introduction of constructs into cells. Tissue culture cells offer easy access. However, they often do not recapitulate the endogenous *in vivo* context and are often a single, irrelevant cell type. Even when multiple cell types are present, as in organoids, the correct environment may not be present. This may be exemplified by the difference between CRM identification for the *Rho* gene in the *in vivo* versus *ex vivo* retinal explants. The LS-MPRA from *ex vivo* retinas showed noisier CRM signals, and failed to show as good a signal-to-noise ratio for the required *Rho* CRM, the PPR. This did not seem to hold true for the other *in vivo* versus *ex vivo* comparisons for *Vsx2* and *Grm6*. If one can electroporate *in vivo*, as we do here, the LS-MPRA likely will provide the most reliable results. If electroporation into the cell type of interest is not possible, one can introduce libraries using viral vectors, as was done with an AAV library for identification of CRMs for specific cell types of the central nervous system (Hrvatín et al., 2019 [DOI](#)). Selected genomic regions were chosen bioinformatically, which may introduce a bias, and each predicted element was individually generated by PCR, which is time-consuming and limits library size. Bias may also be introduced by the choice of the promoter or other elements in the expression construct. We chose a simple TATAA promoter to reduce background activity, but as genomic sequences can influence each other, we may have missed some CRMs due to this choice. In addition, while the BAC-based approach enables the survey of large genomic regions, it cannot capture all potential regulatory inputs, and it remains possible that some distal regulatory sequences lie outside of the BACs used here.

When using a CRM as a reagent to gain genetic access to specific cell types, further analysis of its architecture is often unnecessary. However, for those aiming to understand the regulation of a gene of interest, the d-MPRA offers a powerful approach to rapidly pinpoint individual nucleotides that positively or negatively influence CRM activity. Traditional methods, such as introducing single-base mutations or deletions and testing each construct individually, are time-consuming and often yield qualitative, rather than quantitative, results. Repressor sites, in particular, are rarely identified this way. With the growing availability of TF binding site databases, d-MPRA can efficiently highlight key nucleotides and nominate candidate TFs involved in CRM regulation. When multiple CRMs are available, comparing predicted TF binding sites across them can further strengthen TF candidacy. Examining RNA expression data provides additional support. To validate these predictions, TF binding assays performed *in vivo* or *in vitro* can confirm binding to the

endogenous locus. It would also be valuable to integrate LS-MPRA and d-MPRA with analyses of DNA methylation, particularly in regions near genes of interest across stages of retinal development, building on prior studies of methylation in adult tissues associated with disease (Advani et al., 2024 [DOI](#)). Importantly, LS-MPRA and d-MPRA methods are adaptable across species, as demonstrated here in both chick and mouse, and can be applied to different regions of the developing embryo.

Methods

Resource availability

Raw and processed image data are available upon request. All original code and unprocessed sequencing files are available on the Harvard Dataverse (<http://doi.org/10.7910/DVN/TW0ZQL>) (Tulloch, 2025 [DOI](#)). All processed LS-MPRA genome tracks are available to view on a Track Hub in the UCSC Genome Browser (<https://github.com/cattapre/ALAS00> [DOI](#)). Primer sequences are available in Table S1 [DOI](#). Any additional information is available from the lead contact upon request. Plasmids that are required for these methods will be deposited to Addgene.

Experimental model and subject details

All experimental procedures were preapproved by the Harvard Medical Area Standing Committee on Animals (HMA IACUC). Timed and untimed pregnant CD-1 mice used for sequencing and histology were purchased from Charles River Laboratories and were housed in a climate-controlled pathogen free facility. Mouse tissues were collected for sequencing experiments at E14, P3, or P10 and for histological experiments at E14 or P1 as noted within the text. Chick tissues were collected for sequencing experiments at E3, E5, and E6 and for histological experiments at E3 and E5, as noted within the text.

Method details

Electroporation and Culture

For plasmid DNA injections, glass needles were created by pulling Wiretrol II capillaries (Drummond Scientific Company, cat. #5-000-2005) using a needle puller (Sutter Instrument, Model P-97). The glass needles were beveled on two edges with a microgrinder (Narishige, cat. #EG-401). Plasmid DNA with individual or composite enhancers as well as d-MPRA libraries and most LS-MPRA libraries were concentrated to 2.0 µg/µl along with control plasmids containing either bpCAG-mCherry or CAG-BFP concentrated to 0.5 µg/µl. When pooling libraries for multiplexing, LS-MPRAs for Grm6, Vsx2, and Cabp5 were concentrated to 1 µg/µl and the Rho LS-MPRA library was concentrated to 50 ng/µl, with the bpCAG-mCherry control plasmid concentrated to 0.25 µg/µl.

Ex vivo electroporation: perinatal mouse and chick eyes were enucleated from animals and placed in fresh 1×PBS. Using a beveled glass needle, plasmid DNA (prepared in 1×PBS supplemented with 0.01% Fast Green (Millipore Sigma cat. #F7252) was injected into the subretinal space. Injected eyes were either transferred to a modified electroporation chamber (Montana et al., 2011 [DOI](#)) or placed between platinum disk electrode tweezers (Bulldog Bio, cat. #CUY650P2). All retinas were electroporated with five 50 ms pulses at 40 V delivered at 1-second intervals using a NEPA21 type II Nepagene electroporator; electric currents were applied in both directions between electrodes to maximize electroporation efficiency.

Following electroporation, E14 mouse and E5 chick retinas were removed from surrounding tissues and placed in a culture medium consisting of a 1:1 mixture of DMEM:F12, supplemented with 10% FBS, 200 mM L-glutamine (Sigma-Aldrich, cat. #G3126), and 100 U/mL penicillin with 100 mg/mL streptomycin (Invitrogen/Gibco, cat. #15140-122) in microcentrifuge tubes, and cultured on a nutator at 37°C for 16-28 hrs, depending on the experiment. P3 mouse retinas were removed from surrounding tissues and placed directly onto a nucleopore track-etched polycarbonate membrane (Watman, cat. #110410) in a 12-well tissue culture dish, floating on 2 ml explant media consisting of 50% Minimum Essential Medium (Millipore Sigma, cat. #51412C)(v/v), 25% HBSS

(Thermo Fisher Scientific, cat. #14-025-092) (v/v), 25% heat-inactivated horse serum (Thermo Fisher Scientific, cat. #26050-088), 200mM L-glutamine (Sigma-Aldrich, cat. #G3126), 5.75mg/mL glucose, 25mM HEPES (Invitrogen/Gibco, cat. #15630-080), and 100 U/mL penicillin with 100mg/mL streptomycin (Invitrogen/Gibco, cat. #15140-122). The P3 retinas were covered with 20 μ l of explant media, and incubated at 37°C, 5% CO₂ for 7 days; each day, 1ml of media was replaced with new media and 20 μ l of explant media was reapplied on the retina to keep moist.

***In vivo* electroporation**

P0 and P3 mouse pups were anesthetized by cryoanesthesia on ice and injected subcutaneously with 0.012mg/ml buprenorphine (approximately 20 μ l; PAR Pharma). A small skin incision was made by a 30-gauge needle across the eyelid and the eye was exposed. A small drop of 0.5% proparacaine hydrochloride (Bausch & Lomb, cat. #24208-730-06) was placed onto the surface of the eye. A beveled glass needle containing plasmid DNA (prepared in 1 $\times$ PBS supplemented with 0.01% Fast Green, Millipore Sigma cat. #F7252) was inserted through the sclera into the subretinal space. Plasmid DNA was delivered using a Femtojet Express pressure injector (Eppendorf, cat. #920010521) set at 330 Pa for 3 seconds.

The eyelids were then closed with a cotton applicator. Tweezer-type electrodes (Harvard Apparatus, BTX, model 520, 7 mm diameter, cat. #450165) were placed with the positive end slightly above the injected eyelid and the negative electrode on the contralateral side of the head. An electric field was applied using an electroporator (BEX, cat. #CUY21EDIT), delivering five 50 ms pulses at 80 V with 1-second intervals. Eyelids were dried with a cotton applicator to ensure eyelid closure.

***In ovo* electroporation**

Fertilized chick eggs were incubated at 37°C in a humidified chamber until they reached Hamburger and Hamilton (HH) stages 11-13 (E2) or stages 22-24 (E4). A small window was created in the eggshell, and a solution containing plasmid DNA was injected into the neural tube using a beveled glass needle. Platinum chopstick electrodes with 4mm tips (Bulldog Bio, cat. #CUY611P7-4) were positioned flanking the neural tube: at E2, electrodes were placed against the outer membrane to ensure targeted DNA delivery into the ventral neural tube. At E4, due to the natural 90° rotation of the neural tube, a small hole was made in the membrane to enable precise electrode placement on the lateral sides. Electroporation was performed using a NEPA21 Type II Nepagene electroporator with five 50 ms pulses at 25 V delivered at 1-second intervals. After electroporation, the eggs were resealed and incubated for 24–48 hours before tissue collection. Electroporation success was assessed by detecting fluorescence from control plasmids (bpCAG-mCherry or CAG-BFP).

To ensure sequencing replicates contained a sufficient representation of enriched barcodes at each age, pools of at least 8 mouse retinas for each E14 LS-MPRA and d-MPRA experiment, 3 mouse retinas for each P3 LS-MPRA experiment, 3 chick retinas or 1-2 chick spinal cords for each LS-MPRA experiment were used. When indicated and either coupled with *ex vivo* electroporation or *ex vivo* culture alone, the γ -secretase inhibitor LY411575 (MilliporeSigma, cat. #SML0506) was added to a final concentration of 1 μ M. A 10,000 $\times$ stock was prepared at 10 mM in DMSO and stored at –20°C. Just prior to use, the 10 mM stock was diluted 1:10 in 1 $\times$ PBS to yield a 1000 $\times$ working solution; 1 μ l of the working solution was added per 1 ml culture.

RNA Isolation

To isolate RNA for barcode sequencing or to quantify gene expression, up to 8 mouse retinas, 3 chick retinas, or 2 chick spinal cords were pooled and homogenized using 21-gauge needles in 250 μ l TRI Reagent (MilliporeSigma, cat. #T9424). Total RNA was isolated using the Zymo Directzol Microprep Kit (Zymo Research, cat. #R2061), omitting on-column DNase treatment. For LS-MPRA and d-MPRA library preparations, polyA RNA was subsequently enriched using the Dynabeads mRNA Purification Kit (Thermo Fisher Scientific, cat. #61006). The specific cDNA synthesis protocols for each experiment are detailed in following sections.

Retinal Cell Dissociation

Mouse retinas electroporated at E14 or P3 and cultured *ex vivo* were dissociated using papain digestion. For each sample, a papain mix was prepared by combining 315 μ L HBSS, 35 μ L 1M HEPES (pH 7), 20 μ L 50 mM L-cysteine (freshly prepared), 0.4 μ L 0.5 M EDTA (pH 8), 19.6 μ L UltraPure Water, and 10 μ L papain solution (Worthington Biochemical, cat. #LS003126), followed by incubation at 37°C for 15 mins to activate the enzyme. Retinas were incubated in 400 μ L of activated papain mix at 37°C for 7 mins, then centrifuged at 600 $\times$ g for 2.5 mins. The supernatant was removed, and cells were resuspended in 1 mL HBSS/FBS mix (10% FBS in HBSS), centrifuged again, and resuspended in 600 μ L DMEM/BSA mix (4 mg/mL BSA in DMEM). Cells were gently triturated using a fire-polished Pasteur pipette or a P1000 low-retention pipette, centrifuged again at 600 $\times$ g for 2.5 mins, and finally resuspended in 200–1000 μ L of the appropriate solution for downstream applications.

Fixed Tissue Preparation

Retinas electroporated *in vivo* were first dissected from surrounding tissues while retaining the lens in 1 $\times$ PBS. Retinas in *ex vivo* culture were removed from culture media. Spinal cords *in ovo* were dissected from surrounding tissues in 1 $\times$ PBS. All tissues were fixed in 4% paraformaldehyde in 1 $\times$ PBS for 20 mins and rinsed three times in 1 $\times$ PBS for 30 mins. Tissues were transferred to 30% sucrose in 1 $\times$ PBS and gently agitated until fully infiltrated (approximately 30–60 mins). Retinas were equilibrated in a 1:1 mixture of 30% sucrose in 1 $\times$ PBS and Tissue-Tek OCT (VWR, cat. #25608 930), and were then placed in cryomolds (Sakura Finetek, cat. #4565) in this medium and snap frozen on dry ice. Spinal cords were equilibrated in 100% Tissue-Tek OCT prior to snap freezing. Cryosections (20 μ m thick) were air dried for 20 mins at 37°C and stored at –80°C.

Fluorescent in situ Hybridization

For multiplexed HCR™ RNA-FISH, probe sets were manufactured by Molecular Instruments based on the sequences provided: *Olig2* (NM_016967.2), *Neurog2* (XM_030252394), *OLIG2* (NM_001031526.1), and *GFP* (d2eGFP sequence; Lot no. RTA210). Frozen sections on slides were rinsed two times for 5 mins in PBS, pre-hybridized in probe hybridization buffer (Molecular Instruments HCR™ Buffers [v3.0]) for 10 mins at 37°C, and hybridized with 0.4 pmol of each probe set (Molecular Instruments HCR™ Probes [v3.0]) in hybridization buffer O/N at 37°C. Excess probes were removed in serial 15 mins incubations at 37°C in mixtures of 1:0, 3:1, 1:1, 1:3, and 0:1 of probe wash buffer (Molecular Instruments HCR™ Buffers [v3.0]) and 5 $\times$ sodium chloride sodium citrate with 0.1% Tween-20 (SSCT) followed by a 5 min rinse at RT in 5 $\times$ SSCT. Sections were then incubated in amplification buffer (Molecular Instruments HCR™ Buffers [v3.0]) for 30 mins at RT before incubating in a dark humidifying chamber with 6 pmol of h1 and 6pmol of h2 snap-cooled (heated at 95°C for 90s and incubated at RT for 30 mins in the dark) hairpins for each probe set in amplification buffer for 6–18hrs at RT. Alex546- and Alexa647-labeled hairpins were used. Excess hairpins were removed in sequential rinses in 5 $\times$ SSCT at RT for 5 mins, 2 $\times$ 30 mins, and 5 mins and proceeded to immunostaining.

Immunohistochemistry

New frozen sections on slides that were first washed two times for 5 mins in PBSTw (1 $\times$ PBS with 0.1% Tween-20), or immediately following RNA-FISH, were blocked with 2.5% donkey serum (Jackson ImmunoResearch, cat. #017-000-121) in PBSTw for 1hr at RT. Tissues were incubated in primary antibodies O/N at 4°C and with secondary antibodies for 1 hr at RT in blocking solution, rinsed three times with PBSTw between and after antibody incubations, and mounted using Fluoromount-G (SouthernBiotech, cat. # 0100-01). Mouse monoclonal antibodies raised against *Olig2* (1:500, Millipore cat. #MABN50), chicken polyclonal antibodies raised against GFP (1:1000, Abcam, cat. #ab13970), and rabbit polyclonal antibodies raised against BFP (1:200, Abcam, cat. #ab286131) were used. Alexa405-, Alexa488-, and Alexa647-conjugated secondary antibodies (1:200; Jackson ImmunoResearch) raised in donkey against respective primary antibody species as well as DAPI (1:1000, Invitrogen cat. #D1306) were also used.

Microscopy

Samples were imaged with a Yokogawa CSU-W1 single disk (50mm pinhole size) spinning disk confocal unit attached to a motorized Nikon Ti inverted microscope equipped with a Nikon linear-encoded motorized stage with a Mad City Labs 500 mm range Nano-Drive Z piezo insert, an Andor Zyla 4.2 Plus sCMOS camera using a Nikon S Plan Fluor 40x/1.3 DIC H/N2 oil immersion objective lens. The final digital resolution of the image was 0.16 $\mu\text{m}/\text{pixel}$. Fluorescence from (405 nm, 488 nm, 550 nm, and 640 nm) was collected by illuminating the sample with directly modulated solid-state lasers 405 nm diode 100 mW (at the fiber tip) laser line, 488 nm diode 100 mW laser line, 561 nm DPSS 100 mW laser line and 640 nm diode 70 mW laser line in a Toptica iChrome MLE laser combiner, respectively. Signal from each channel was acquired sequentially with hard-coated Chroma ET455/50 nm, Chroma ET525/36 nm, Chroma ET605/52 nm, and Chroma ET705/72 nm emission filters in a filter wheel placed within the scan unit, for blue, green, red, and far-red channels, respectively. All images were captured using the 16-bit dual gain high dynamic range camera mode and no binning. Nikon Elements AR acquisition software was used to acquire the data. Z-stacks were acquired using Piezo Z-device, with the shutter closed during axial movement. Images were acquired by collecting all colors in a Z-step, then each Z-stack before XY-stage movement, with a 15% tile overlap for correct stitching. Data were saved as ND2 files.

Vector Design

Barcoded Stagintbc7

The parent vector, Stagint7, is a minimal pUC19-based construct containing a kanamycin resistance gene, an SV40 polyadenylation site, a multiple cloning site immediately upstream of a minimal TATAA promoter, an intron-containing EGFP open reading frame, and a rabbit β -globin polyadenylation sequence. This vector is based on the previously described Stagia3 vector used for retinal enhancer assays (Billings et al., 2010 [DOI](#)). The parent vector was linearized between the EGFP ORF and polyadenylation sequence via an AscI (NEB, cat. #R0558) recognition sequence, inserting Illumina adapter sequences flanking a recreated AscI recognition sequence, a unique 8-bp library barcode tag for multiplexing (used only for Rho, Grm6, Vsx2, and Cabp5 libraries), and a 24-bp barcode using an Enhanced Gibson Assembly reaction (Rabe & Cepko, 2020 [DOI](#)). Barcodes consisted of either twelve SW base repeats or four VHBD base repeats to prevent high GC content and homopolymer repeats as well as to provide sufficient barcode complexity. The barcoded Stagintbc7 vector used to create LS-MPRA libraries (Rabe, 2020 [DOI](#)) was generated by combining 100 ng of linearized parent vector with 25 nM of each primer pair (see Table S1 [DOI](#)) and, along with the pre-mixed Enhanced Gibson Assembly reaction, incubated at 50°C for 1 hr. The assembled product was purified with a Zymo DNA Clean & Concentrator Kit (Zymo Research, cat. #D4013).

For the creation of the four Stagintbc7 multiplexed libraries, four unique DNA library tags were generated using the DNABarcodes R package (Buschmann, 2017 [DOI](#)) to ensure that all barcodes differed from one another by at least five bases while avoiding homopolymer repeats. The create.dnabarcodes function was used with a Hamming distance metric of 5, and additional filters were applied to exclude triplet repeats, ensure GC content balance, and prevent self-complementarity. The selected library barcode tags included: ATACAGGC, TGTGCAAG, CACGAAGA, and GTGAACCT.

Statadual_WPRE

This vector, used for d-MPRA library construction, is derived from Stagia3 but replaces the minimal TATAA promoter and intron-containing EGFP ORF of Stagint7 with a WPRE sequence.

Assembly of LS-MPRA Libraries

BAC Isolation and Preparation

BACs were first isolated from stab cultures (BACPAC Resources). Cultures were streaked onto chloramphenicol-treated LB agar plates and incubated overnight at 37°C. A single colony was picked and grown in LB media supplemented with chloramphenicol (12.5 $\mu\text{g}/\text{mL}$) in 5 ml overnight, followed by expansion to 500 ml. The culture was pelleted, and BAC DNA was extracted

using a large construct DNA isolation kit with ATP (Qiagen, cat. #12462). DNA concentrations were determined using the Qubit dsDNA HS Assay Kit (ThermoFisher, cats. #Q32851 and # Q32856). For sequencing validation, samples were sent to Plasmidsaurus for large plasmid sequencing. BAC clones used in all experiments included RP23-219M6 (*Rho*), RP23-417M10 (*Grm6*), RP23-127O21 (*Vsx2*), RP24-125H22 (*Cabp5*), CH29-613 (*Olig2*), RP23-182M12 (*Ngn2*), and CH261-60J3 (*OLIG2*).

BAC Fragmentation and Adapter Ligation

BAC DNA (500 ng) was enzymatically digested, end-repaired, and dA-tailed using the NEBNext Ultra II FS DNA Module kit (NEB, cat. #E7810S) with a 20-min enzyme incubation. Digested DNA was purified using SPRIselect beads with a 1.2x left-side size selection ratio. Concentrations were determined using the Qubit dsDNA HS Assay Kit. Double-stranded adapters (Table S1 [↗](#)) were annealed from complementary oligonucleotides and diluted to a molar concentration approximately 100-fold higher than fragments in 1×T4 DNA ligase buffer (NEB, cat. #B0202S). The 5' adapter sequence included an Illumina adapter sequence and a 25-bp homologous sequence for Gibson Assembly insertion into the Stagintbc7 vector, while the 3' adapter sequence included a 23-bp homologous sequence and an *AscI* recognition sequence. Adapters were ligated to fragments using the Blunt/TA Ligase Master Mix (NEB, cat. #M0367), adding a 1:1 mixture of ligase mix and 18 μ l fragments with 2 μ l adapters, incubating at RT for 1 hr. Adapter-ligated DNA was purified using the Zymo DNA Clean & Concentrator Kit and quantified using the Qubit dsDNA HS Assay Kit.

Adapter-ligated Fragment Amplification and Purification

Adapter-ligated fragments were amplified using NEBNext Ultra II Q5 Master Mix (NEB, cat. #M0544) in a 300 μ l reaction with 0.5 μ M primers (Table S1 [↗](#)) complementary to the 5' and 3' adapter ends and at an annealing temperature of 64°C for 15 cycles. Amplified DNA was purified using the Zymo DNA Clean & Concentrator Kit. To ensure proper fragment size selection, adapter-ligated fragments were resolved on a 1% agarose gel, and the desired size range (fragment plus ~90 bp of adapter sequence) was gel purified using Agarose Dissolving Buffer (Zymo Research, cat. #D4001-1-100) and the Zymo DNA Clean & Concentrator Kit.

Cloning into Barcoded Library

The Stagintbc7 barcoded library was linearized with *PacI* (NEB, cat. #R0547) and *EcoRI*-HF (NEB, cat. #R3101). The digested vector was purified via 0.75% agarose gel electrophoresis and gel extraction. An Enhanced Gibson Assembly was performed with 100 ng vector and a 10:1 molar insert:vector ratio at 50°C for 1 hr. The assembled product (Rabe, 2020 [↗](#)) was purified with a Zymo DNA Clean & Concentrator Kit.

Library Transformation and Amplification

All barcoded libraries—Stagintbc7, LS-MPRA, and d-MPRA—were electroporated into DH10 β electrocompetent cells in a pre-cooled 1 mm cuvette (BTX, cat. #45 0134) with the following parameters: 2 kV, 25 μ F, and 200 Ω on a BioRad GenePulser II. Transformed cells were outgrown in 2 ml SOC media at 37°C for 2 hrs, and then grown in 3D culture by embedding bacterial transformations in 0.5% sodium alginate matrices with CaCO₃ and LB to reduce clonal bias inherent in 2D cultures. To prepare 3D cultures (Rabe, 2020 [↗](#)), the transformed bacterial cells grown in SOC were mixed with 12.5 ml of 1% sodium alginate (mixed 1g/100ml LB O/N, then autoclaved) and LB to a volume of 20 ml. Then, 250 μ l of a CaCO₃ suspension (10 g in 100 ml diH₂O; Sigma Aldrich, 310034) and 1000× kanamycin (25 μ L) were combined in 2.5 ml LB, mixed immediately before use, and added to the bacterial-alginate solution. Lastly, 41.7 μ l of 37% HCl in 2.5 ml LB was added for a final volume of 25 ml, and the mixture was gently rocked for 5 minutes before being poured into a sterile 150 mm petri dish. Cultures were incubated at 37°C for 16–20 hours to allow gel formation and outgrowth.

Following incubation, 6.25 ml of 20× SSC + 40 mM EDTA solution (pH 8.0) was added to dissolve the 3D culture, and was gently rocked at RT for 15 min. Bacterial cells were pelleted (6,000×g, 15 min), washed with 25 ml LB, and centrifuged again. The pellet was resuspended in 300 ml LB containing

50 µg/ml kanamycin and 0.05 mg/ml alginate lyase (Sigma Aldrich, cat. #A1603) and incubated at 37°C with shaking (250 RPM) for 6–8 hrs. Cultures were then pelleted for plasmid preparation or storage at –20°C. Plasmids were extracted using a Plasmid Maxi kit (Qiagen, cat. #12163).

To estimate the complexity of transformed libraries, a serial dilution was performed after the transformation and prior to 3D culture. One microliter of the 2 ml culture in SOC was removed and subjected to three sequential dilutions (1:500, 1:5000, and 1:50,000) to represent 1/10,000th, 1/100,000th, and 1/1,000,000th of the total transformation, respectively. For dilution preparation, 1 µl of the culture was added to 499 µl SOC, mixed, and 50 µl of this dilution was transferred to 450 µl SOC, followed by an additional 1:10 dilution under the same conditions. 100 µl of each dilution was plated onto kanamycin-treated LB agar and incubated O/N at 37°C. The number of resulting colonies was used to estimate transformation complexity, with a quantifiable optimal range of 20–200 colonies per plate.

Assembly of d-MPRA Libraries

CRMs of interest were PCR-amplified from their respective BAC clones using primers designed to incorporate 20–30 bp overlapping sequences for Gibson Assembly into the Statadual_WPRE vector (Table S1 [\[link\]](#)). PCR reactions were performed using NEBNext Ultra II Q5 Master Mix under standard conditions, and amplicons were gel-purified using the Zymo DNA Clean & Concentrator kit.

Mutagenesis was performed using the GeneMorph II Random Mutagenesis Kit (Agilent, cat. #200550), following the manufacturer's protocol with 100 pg of input DNA and MutF/R primers (Table S1 [\[link\]](#)), which amplify the entire CRM region and incorporated overlapping sequences. PCR products were gel-purified, and 100–200 ng of each mutagenized fragment was assembled into the EcoRI/PacI-linearized Statadual_WPRE vector via Enhanced Gibson Assembly at a 10:1 insert-to-vector molar ratio. The assembled libraries were electroporated into DH10β electrocompetent cells, expanded in 3D bacterial culture, and purified as previously described.

For intraplasmid duplication (IPD), 3 µg of purified intermediate plasmid libraries were digested with the nicking endonuclease Nt.BspQI (NEB, cat. #R0644) at 50°C for 1 hr, column-purified, and incubated with Bst 2.0 WarmStart Polymerase (8 units, NEB, cat. #M0538) in 1× Isothermal Amplification Buffer supplemented with 6 mM MgSO₄ and 200 µM dNTPs (GoldBio, cat. #D9001) in a 50 µl reaction at 65°C for 45 mins, followed by enzyme inactivation at 80°C for 20 mins. To reduce self-ligation, 2.5 units of Taq DNA polymerase (NEB, cat. #M0273) was added, and the reaction was incubated at 65°C for 20 mins for dA-tailing. The prominent linear band was gel-excised and purified.

The final d-MPRA insert (a minimal TATAA promoter and intron-containing EGFP ORF) was PCR-amplified from Stagint7 with primers designed to incorporate 20–30 bp overlapping sequences for Gibson Assembly into the linearized IPD product (Table S1 [\[link\]](#)), gel-purified, and assembled with 25 fmol of linearized, IPD-processed vector in an Enhanced Gibson Assembly reaction (Rabe, 2020 [\[link\]](#)). The completed d-MPRA libraries were electroporated into DH10β electrocompetent cells, expanded in 3D bacterial culture, and purified as previously described.

CUT&RUN

Experiments were performed using freshly dissociated E14 mouse retinas with the CUTANA™ ChIC/CUT&RUN Kit (EpiCypher, cat. #14-1048) according to the manufacturer's protocol. Mouse monoclonal antibodies raised against Pax6 (Developmental Studies Hybridoma Bank, cat. #AB_528427) and against Foxn4 (Santa Cruz Biotechnologies, cat. #sc-377166) as well as rabbit polyclonal antibodies raised against Mybl1 (Millipore-Sigma, cat. #HPA008791) were used, adding 500ng of the respective antibody to each sample during the antibody binding stage. Indexed libraries were prepared with the NEBNext UltraII DNA Library Prep Kit (NEB, cat. #E7103) and the NEBNext Multiplex Oligos for Illumina Dual Index Sets (NEB, cats. #E7600S or #E7780S). All experiments were validated following Next-Generation sequencing using the SNAP-CUTANA K-MetStat Panel of nucleosome spike-ins in control reactions and the manufacturer's protocol (Appendix 1.6; EpiCypher, cat. #14-1048).

Sequencing Library Preparation

LS-MPRA Fragment–Barcode Association Libraries

To excise the minimal promoter and GFP ORF, enabling BAC fragments to be positioned adjacent to their corresponding barcodes for sequencing, 2–3 µg of the LS-MPRA library was digested with *AscI* and resolved on a 1% agarose gel (Rabe, 2020). The fragment containing both the barcode and insert (~2.1 kb plus the fragment size) was excised and gel-purified. 500 ng of purified fragment was circularized in a 100 µl Quick Ligation reaction (NEB, cat. #M2200) at RT for 5 mins. The ligation product was column-purified and treated with 50 µl exonuclease (Lucigen, cat. #E3101K) to remove residual linear fragments.

Following gel resolution, the smallest and brightest band was excised and gel-purified. Because the ligation product already contained Illumina read primer sequences, 50 ng of purified product was used as a template for index PCR with NEBNext Q5 Ultra II Master and NEBNext Multiplex Oligos for Illumina Dual Index Set for 15 cycles. PCR products were purified via two-sided SPRISelect bead purification (0.5x to 0.9x).

LS-MPRA Barcode Abundance Libraries

To assess barcode abundance in RNA, polyA RNA was reverse-transcribed using the Protoscript II Kit (NEB, cat. #E6560) and a reverse transcription primer at 1 µM specific to all barcoded transcripts (bc7_cdnar; Table S1). The resulting cDNA was purified using the Zymo Oligo Clean & Concentrator Kit (Zymo, cat. #D4060) and then amplified for 30 cycles using NEBNext Q5 Ultra II Master Mix with two primers: one that anneals to a region of the GFP ORF upstream of the intron and the other downstream of the barcode-flanking Illumina adapter sequences [bc7_cdnaf and bc7_cdnar2 (Table S1), respectively], ensuring the intron-containing plasmid DNA was not included. The cDNA-derived amplicons (~356–364 bp) were gel-purified from a 1% agarose gel. Indexed barcode amplicons were generated with 50 ng purified product, NEBNext Q5 Ultra II Master Mix, and NEBNext Multiplex Oligos for Illumina Dual Index Set for 15 cycles, followed by two-sided purification using SPRISelect beads (0.95x–1.2x).

d-MPRA Libraries

Amplicons for d-MPRA mutagenesis analysis were generated from both plasmid libraries and RNA (Rabe, 2020). cDNA was synthesized from polyA RNA using the Protoscript II Kit and the reverse transcription primer at 1 µM specific to all d-MPRA barcoded transcripts [bc7_cdnar_WPRE, (Table S1)], purified with the Zymo Oligo Clean & Concentrator Kit, and amplified for 30 cycles using NEBNext Q5 Ultra II Master Mix with two primers: one that anneals to a region of the GFP ORF upstream of the intron and the other downstream of the barcode and upstream of the WPRE sequence [bc_cdnaf and MutR, respectively (Table S1)]. The resulting cDNA band (~290 bp plus the CRM size) was gel-purified. For plasmid d-MPRA libraries, the corresponding fragment was generated by digesting with *BsrGI* (NEB, cat. #R3575) and *NdeI* (NEB, cat. #R0111), which excises the CRM barcode and WPRE sequence, and gel purifying the product (~760 bp plus the CRM size).

For both plasmid library and RNA-derived amplicons, Illumina adapter sequences were added to each side of the mutated CRM sequence [dualmut_III1F or dualmut_III2F and dualmut_III2R primers (Table S1)]. These fragments, amplified from 50 ng template with NEBNext Q5 Ultra II Master Mix, were gel-purified and subjected to index PCR with NEBNext Multiplex Oligos for Illumina Dual Index Set for 15 cycles. PCR products were purified via two-sided SPRISelect bead purification (0.6x to 0.9x).

Illumina sequencing

All indexed LS-MPRA, d-MPRA, and CUT&RUN libraries were analyzed using an Agilent TapeStation High Sensitivity D1000 assay. Indexed samples were sequenced using 2×150 bp paired end reads on an Illumina NovaSeq instrument at either MedGenome Inc. or Admera Health, with the range of sequencing depth between 20M–50M reads.

Expression Constructs

CRM expression constructs were cloned into the Stagia3 expression vector with an Enhanced Gibson Assembly (Rabe & Cepko, 2020). For all constructs, the vector backbone was amplified from the expression vector using primers compvec_F/R (Table S1). For Olig2 and OLIG2 constructs, the intron-containing EGFP (GFPint) along with the rabbit β -globin polyadenylation sequence was amplified from Stagint7 whereas the insert for the Ngn2 constructs also included the TATAA minimal promoter upstream of EGFP. Olig2 CRMs, the Olig2 minimal promoter, Ngn2 CRMs, OLIG2 CRMs, and the OLIG2 minimal promoter were amplified from their respective BAC clones (all primers in Table S1). Coordinates for all regions are included in Table S2.

For individual constructs, Olig2-NR1 or Olig2-NR2 CRMs were inserted upstream the Olig2 minimal promoter and GFPint, whereas Olig2-NR3 CRM was assembled downstream of the GFPint + polyA sequence. The Olig2 composite construct was assembled accordingly, placing the NR1 and NR2 fragments adjacent to each other and upstream of the Olig2 minimal promoter, GFPint + polyA sequence, and NR3 fragment. Backbone plasmids containing GFPint alone or GFPint under the Olig2 minimal promoter in the absence of CRM fragments were also assembled (primers in Table S1).

The mouse Ngn2-CRM constructs were assembled placing the Ngn2-CRM1, CRM2, or CRM3 fragments upstream of the TATAA minimal promoter and GFPint, while the CRM4 fragment was placed downstream of the GFPint+polyA sequence. The chick OLIG2-CRM constructs were similarly constructed, with OLIG2-CRM1 or CRM2 fragments assembled upstream of the cOLIG2 minimal promoter, while the OLIG2-CRM3 fragment was assembled downstream of the GFPint+polyA sequence.

ddPCR

All ddPCR assays were performed on the Bio-Rad ddPCR system, including the automated droplet generator (Bio-Rad, cat. #1864101). *Olig2* gene expression was determined using Evagreen-based ddPCR with *Hprt* as a normalization control (primers in Table S1). cDNA was generated from isolated RNA with the LunaScript RT SuperMix Kit (NEB, cat. #E3010) from ~2–3 ng of total RNA and purified with the Zymo Oligo Clean & Concentrator kit. 1 ng of cDNA was added to each 22 μ l ddPCR reaction; no-RT controls confirmed the absence of genomic DNA amplification (all assays spanned exon–exon junctions).

Sequencing, quantification, and statistical analyses

LS-MPRA Barcode Association and Quantification Analysis

Paired-end 150-bp sequencing reads were processed to establish barcode-fragment associations and quantify barcode abundances. Two custom shell scripts were used for sequencing data analysis.

Barcode-Fragment Association

For barcode-fragment association, sequencing reads were aligned to the mm10 reference genome using Minimap2 version 2.28 (Li, 2018) and Bowtie2 version 2.5.4 (Langmead & Salzberg, 2012). Adapter sequences were trimmed using bbdut from the BBTools suite version 39.18 (Bushnell et al., 2017), and barcode sequences were extracted from Read 1 using UMI-tools extract version 1.1.6 (Smith et al., 2017). Valid barcode-fragment pairs were identified using UMI-tools group and filtered for library-specific sequences. Genomic coordinates were corrected based on barcode positions, and results were output as BigWig tracks using deepTools bamCoverage version 3.5.5 (Ramirez et al., 2016). Genomic coordinate manipulation, barcode overlap correction, and file conversions were performed using Bedtools version 2.31.1 (Quinlan & Hall, 2010), Samtools version 1.21 (Danecek et al., 2021; Li et al., 2009), and SeqKit version 2.6.1 (Shen et al., 2016b).

An earlier version of the pipeline differed primarily in its alignment and filtering steps. Instead of Minimap2, it used BLAT version 35 (Kent, 2002) to align BAC sequences to the genome, extending flanking regions by 1 kb. Additionally, overlapping barcode-associated reads were

collapsed using R packages GenomicRanges version 1.50.2 (Lawrence et al., 2013), regioneR version 1.30.0 (Gel et al., 2016), and data.table version 1.16.4 (Barrett et al., 2025). Barcode extraction followed a similar approach but lacked the library-specific filtering step included in the updated version. By replacing BLAT with Minimap2 and incorporating stricter filtering, the updated pipeline improved alignment efficiency and accuracy.

Barcode Abundance Quantification

For barcode abundance quantification, extracted barcode sequences were filtered against a whitelist to exclude sequencing errors and ensure accuracy. The whitelist was generated by extracting reverse-complement barcode sequences from a reference FASTA file containing known barcode-fragment associations using the Biostrings R package version 2.66.0 (Pagès et al., 2025). Pairwise Hamming distances were computed using the stringdist R package version 0.9.15 (van der Loo, 2014) to define a mutation threshold. All possible barcode variants within this threshold were enumerated and stored in an RDS file for downstream filtering. Sequencing reads were matched against this whitelist using UMI-tools extract, and barcode counts were normalized to library input using deepTools bamCompare. Normalized BigWig tracks were generated for visualization and analysis.

An earlier version of the pipeline extracted barcodes using bbdduk and SeqKit, without applying Hamming distance-based error correction. Unique barcode counts were merged with barcode-fragment associations, and fragment counts were expanded to match observed barcode frequencies before normalization with bamCompare. While both methods yielded comparable results, the whitelist-based approach provided better error correction and filtering, enhancing reliability in downstream analyses.

All LS-MPRA plots were displayed on the Integrative Genomics Viewer (IGV) (Robinson et al., 2011), and aligned with analyses of genetic conservation from the UCSC Genome Browser database (Perez et al., 2025) as well as with RefSeq genes (O'Leary et al., 2016).

ATAC-seq Analysis

Open chromatin regions were determined based on pseudo-bulk re-analysis of published single-cell ATAC-seq datasets (Lyu et al., 2021), and deposited under accession number GSE181251. Deposited bam files were converted back to FASTQ format using bamtofastq version 1.3.2 from 10XGenomics (Github). Single-cell FASTQ files for E14 and P8 mouse retinas were analyzed using cellranger atac count version 2.0.0 (Satpathy et al., 2019), where reads were filtered and aligned to the mm10 mouse genome, barcode counted, and transposase cut sites identified. Regions of accessible chromatin were detected on a single-cell basis. Bam files contained all the aligned paired-end reads were used to determine pseudo-bulk genome coverage using Deeptools version 3.5 (Ramirez et al., 2016), normalized using RPKM, and filtered for blacklisted (Amemiya et al., 2019), gap and repeat regions, and visualized using IGV (Robinson et al., 2011, Thorvaldsdottir et al., 2013).

CUT&RUN Analysis

Analysis of sequenced samples from the CUT&RUN experiments were analyzed as previously described (Catta-Preta et al., 2025). FASTQ files containing the reads were controlled for quality using FASTQC version 0.11.9 (Andrews, 2010), and further cleaned up of remaining adapter sequences using Trim Galore version 0.6.6 (Krueger, 2015). The resulting pair-ended reads were aligned to the mm10 mouse genome using BWA mem version 0.7.17 (Li & Durbin, 2009), and duplicates removed with Samtools version 1.15.1 (Li et al., 2009). Peaks of enriched immunoprecipitated regions against IgG (negative) control were called using MACS version 2.1.1 (Zhang et al., 2008) with a p-value of 0.05, extension size 200, no lambda baseline correction. Genome coverage from the bam files for all samples was determined using Deeptools version 3.5 (Ramirez et al., 2016), normalized using RPKM, and filtered for blacklisted (Amemiya et al., 2019), gap, and repeat regions. Coverage was visualized using IGV (Robinson et al., 2011, Thorvaldsdottir et al., 2013). Motif enrichment and specific motif presence was determined using Homer version 4.11.1 (Heinz et al., 2010).

D-MPRA Plasmid and Experimental Library Processing

To analyze barcode-fragment associations and quantify changes in mutated fragment (i.e., barcode) frequency in d-MPRA plasmid and experimental libraries, sequencing data were processed through a pipeline involving read merging, alignment, barcode extraction, and statistical analysis of barcode frequencies across CRM sequences.

Plasmid Library Processing

Plasmid control libraries were processed by merging paired-end reads and aligning them to reference sequences using Minimap2 and pysam version 0.22.1 (Bonfield et al., 2021 [DOI](#)). Barcode and fragment associations were then analyzed by parsing Minimap2's cs tags, with mutations identified and quantified using custom Python scripts that leveraged NumPy version 1.26.3 (Harris et al., 2020 [DOI](#)) and pandas version 2.1.4 (McKinney, 2010 [DOI](#)) for data handling. Reads containing deletions within homopolymer regions were excluded based on positions identified from reference FASTA sequences, and reads failing to align fully across the contig were discarded. Monte Carlo simulations were implemented using multiprocessing to perform 10 independent subsampling replicates, and barcode mutation frequencies were normalized to wild-type sequence counts for accurate mutation rate estimation.

Experimental Library Processing

Reads from experimental d-MPRA libraries were merged and aligned similarly to the plasmid workflow, and barcode-fragment associations were identified. An additional filtering step was applied to remove reads that failed to align fully across the contig, reducing false-positive mutation calls.

Mutation Filtering and Statistical Analysis

Both plasmid and experimental workflows applied stringent mutation filtering, discarding deletions occurring in homopolymer regions. Substitutions and deletions were quantified using pysam to parse cs tags from alignments produced by Minimap2, and filtered with logic implemented in numpy and pandas. Barcode frequencies were normalized relative to plasmid control counts using data structures and sampling routines implemented in pandas, numpy, and Python's multiprocessing library. Monte Carlo sampling was used to refine barcode frequency estimates while accounting for sequencing depth. Final mutation statistics, including mean and standard deviation across positions and replicates, were compiled using pandas and json, ensuring accurate and robust quantification of sequence variation. This workflow ensured precise quantification of barcode frequency in both plasmid and experimental d-MPRA libraries while minimizing sequencing artifacts.

Visualization of d-MPRA Profiles

To visualize barcode frequency distributions across CRMs, normalized mutation rates from Monte Carlo-sampled plasmid controls and experimental d-MPRA libraries were compared and plotted. Barcode frequencies corresponding to substitutions and deletions in the experimental datasets were extracted from position-level JSON summaries. These values were normalized to corresponding wild-type-corrected rates from plasmid controls and log₂-transformed to assess relative mutation enrichment or depletion.

To reduce noise and account for local sequence context, the data were smoothed using a five-base sliding window moving average. A baseline correction was applied to each dataset by subtracting the median log₂-transformed value prior to smoothing. For each genomic position, only sites present in both control and experimental datasets were retained for comparison. Edge regions flanking the CRM, specifically, the first 18 and last 14 positions, were excluded from visualization to eliminate non-informative regions required for construct assembly.

For each replicate, baseline-corrected mutation rates were plotted individually alongside the average profile across all replicates. All visualizations were generated in Python using matplotlib version 3.8.2 (Hunter, 2007 [DOI](#)) and numpy.

Quantification of *Olig2* Transcript Levels Using ddPCR

ddPCR was used to quantify the concentration of *Olig2* transcripts in treated and control retinas across multiple time points. Normalized transcript concentrations were determined from ddPCR reactions and analyzed in GraphPad Prism. Statistical comparisons between treated and control groups were performed using unpaired t-tests assuming equal standard deviations, with multiple testing correction applied using the Holm-Šidák method.

Quantification of Barcode Enrichment Using AUC Analysis

CRM barcode enrichment was quantified by calculating the area under the curve (AUC) using BigWig sequencing files. Normalized barcode count values were extracted for the CRM genomic regions, and AUC was computed using the trapezoidal rule. Statistical analysis was performed in GraphPad Prism using Brown-Forsythe and Welch's ANOVA tests, followed by Dunnett's T3 multiple comparisons test to assess significance.

Quantification of Cell Populations Using a Semi-Automatic Colocalization Analysis Toolbox

Fluorescent cell counting was performed using a semi-automatic object-based colocalization analysis (OBCA) toolbox developed by Lunde and Glover (Lunde & Glover, 2020 [DOI](#)). This toolbox comprises two ImageJ plugins, a Microsoft Excel macro, and a MATLAB script, facilitating efficient analysis of complex imaging data. The first ImageJ plugin was employed to enhance visualization of multichannel 3D images, optimizing features relevant to OBCA. The second plugin enabled semi-automatic quantification of colocalization across multiple fluorescence channels. In this study, cells expressing a specific gene transcript or protein, GFP driven by CRM activity, and BFP from an electroporation control plasmid were identified. The toolbox provided counts of cells co-expressing the gene transcript, GFP protein, and/or GFP transcript. These data were used to calculate the percentage of electroporated (BFP+) cells expressing the specific gene transcript that also expressed GFP protein, GFP transcript, or both, for each analyzed section. Additionally, the percentage of GFP+ cells co-expressing the gene transcript was determined.

Motif Analysis

De novo motif analysis was performed on CRM sequences using the HOMER tool (findMotifsGenome.pl) (Heinz et al., 2010 [DOI](#)). The motif search was performed on CRM coordinates in the mm10 genome assembly using a fixed size for each CRM region and a randomly generated reference genome directory. HOMER's built-in TF binding site database, derived from JASPAR, was used to detect known TF binding sites within the discovered motifs. The analysis output was used to identify potential regulatory factors enriched in these enhancer regions. The comprehensive list of potential TF binding sites within each predicted motif of the *Olig2* CRMs is available at the online repository (Tulloch, 2025 [DOI](#)).

Visualization of PFMs

Position frequency matrices (PFMs) for selected TFBSs were parsed and visualized using a custom Python script. PFMs were read into a pandas DataFrame and transposed to organize nucleotide frequencies by position. Rows were normalized to ensure that nucleotide frequencies summed to one at each position. The information content at each position was calculated using Shannon entropy, incorporating a small pseudocount to avoid undefined logarithmic values. Sequence logos were generated using Logomaker (Tareen & Kinney, 2020 [DOI](#)), with bit scores computed by multiplying nucleotide frequencies by their respective information content.

Single-Cell RNA-Seq Data Acquisition and Processing

scRNA-seq data for retinal development across multiple ages were obtained from the published data repository at https://github.com/gofflab/developing_mouse_retina_scRNASeq (Clark et al., 2019 [DOI](#)). The dataset included:

1. a gene expression counts matrix (*10x_mouse_retina_development.mtx*)
2. gene feature annotations (*10x_mouse_retina_development_feature.csv*)

3. sample-level metadata (*10x_mouse_retina_development_phenotype.csv*).

The raw counts matrix was processed and analyzed using Seurat v4.0 (Butler et al., 2018).

Data Preprocessing

The scRNA-seq data were processed in R using Seurat (Hao et al., 2024). The raw counts matrix was loaded with `Read10X()` and converted into a Seurat object. Genes detected in fewer than three cells were removed, and cells with fewer than 200 detected genes or >5% mitochondrial gene content were excluded. To normalize for sequencing depth, the SCTransform workflow was applied, followed by identification of highly variable genes.

Subset Analysis by Developmental Age

To focus on specific developmental stages, cells were subsetted based on “age metadata”. Two ages were subsetted: E14 and P8. Dimensionality reduction, clustering, and Uniform Manifold Approximation and Projection (UMAP) visualization (McInnes et al., 2018) were performed on each subset independently.

Dimensionality Reduction and Clustering

Principal component analysis (PCA) was performed on the top 2,000 variable genes. UMAP was used for visualization, and clustering was conducted using `FindNeighbors()` and `FindClusters()` with a resolution of 0.5 to delineate distinct retinal cell populations. UMAP plots were generated using `DimPlot()`, with cells color-coded by cell type, developmental stage, or gene expression levels.

Gene Expression and Co-Expression Analysis

The expression of individual genes of interest was visualized using `FeaturePlot()`. A continuous color gradient (light grey to blue) was applied to indicate expression levels. To visualize gene co-expression, `FeaturePlot()` was used with `blend = TRUE`, where cells expressing Gene1 appeared in blue, Gene2 in red, and co-expressing cells in pink. The third panel of the blended plot, which represents co-expression, was extracted and saved using `ggsave()`. All visualizations were generated in R using `ggplot2` (Wickham, 2016).

Quantification of Co-Expressing Cells

Expression data for genes of interest were extracted from the RNA assay slot of the Seurat object using `GetAssayData()`. Cells were classified as expressing a gene if their count was greater than zero. The number of cells expressing each gene individually and both genes concurrently was computed. The percentage of Gene1-expressing cells that also expressed Gene2 was calculated as:

Percent Co-Expression

$$= \frac{\text{Cells expressing both genes}}{\text{Total cells expressing Gene1}} \times 100$$

$$= \left(\frac{\text{Cells expressing both genes}}{\text{Total cells expressing Gene1}} \right) \times 100$$

Percent Co-Expression

$$= \frac{\text{Total cells expressing Gene1}}{\text{Cells expressing both genes}} \times 100$$

Statistical Analysis of Gene Co-Expression with *Olig2* from scRNAseq

To assess whether specific genes were significantly enriched or depleted in *Olig2*-expressing cells, Fisher’s exact test for each gene of interest was performed. Gene expression data were obtained from the RNA count matrix of the retinal single-cell dataset. Cells were classified as expressing a given gene if the raw count value was greater than zero.


For each gene, a 2×2 contingency table was constructed, categorizing cells as *Olig2*⁺ or *Olig2*[−] and as expressing or not expressing the target gene. Fisher’s exact test was then applied to each contingency table to determine whether the presence of a given gene was significantly associated with *Olig2* expression. To account for multiple hypothesis testing, the Benjamini-Hochberg false discovery rate (FDR) correction was applied to the resulting p-values. Genes with an adjusted p-

value (FDR q-value) below 0.05 were considered significantly enriched or depleted in *Olig2*-expressing cells. An odds ratio greater than 1 indicated that the gene was enriched in *Olig2*⁺ cells, whereas an odds ratio less than 1 indicated depletion in *Olig2*⁺ cells.


Data availability

All original code and unprocessed sequencing files are available on the Harvard Dataverse (DOI: doi:10.7910/DVN/TW0ZQL). All processed LS-MPRA genome tracks are available to view on a Track Hub in the UCSC Genome Browser (<https://github.com/cattapre/ALAS00>)

Supplemental figures and tables




Supplemental Figure 2S. Grm6, Vsx2, and Cabp5 LS-MPRAs to identify CRMs in the neonatal mouse retina. Barcode enrichment plots from the Grm6 (A), Vsx2 (B), and Cabp5 (C) LS-MPRAs, aligned with genome browser tracks and annotated with (i) known regulatory regions, (ii) coverage of the barcode-fragment association library across the locus, (iii) log₂ base conservation across 60 vertebrate or 40 placental mammal species, (iv) regions of open chromatin in the P8 mouse retina, and (v) RefSeq gene models. Expanded regions of interest (yellow dashed boxes) show peaks (i-iii) or regions that align with known regulatory elements for each gene.




Supplemental Figure 5S. Activity of backbone plasmids containing the *Olig2* minimal promoter and *EGFP*.

(A) Representative transverse sections of E14 retinas incubated *in vitro* for 24 hours show sparse *GFP* RNA and *GFP* fluorescence driven by control plasmids containing either *EGFP* alone or *EGFP* under the *Olig2* minimal promoter. Pie charts of the percentage of electroporated cells with detectable *GFP* expression.



Supplemental Figure 8S. TF Binding sites in Olig2-NR1 and NR3 CRMs.

(A, H) TF binding motifs identified within Olig2-NR1 (A) and Olig2-NR3 (H) CRM candidates, aligned with the average d-MPRA plot (from Fig. 7). (B-F, I-K) Position frequency matrices of transcription factors aligning to Olig2-NR1: Sox4/11 to Motifs 13 and 16 (B), Lhx2 and Dlx2 to Motif 11 (C), Isl1 to Motif 8 (D), Foxp1 to Motif 18 (E), Mybl1 to Motif 2 (F), and Otx2 to Motif 12 (F); or Olig2-NR3: Ngn2 to Motifs 3 and 5 (I), Bhlhe22 to Motif 13 (J), and Lhx9 to Motif 15 (K). (G, L) Co-expression (pink) of Olig2 (blue) with Sox11, Sox4, Lhx2, Dlx2, Isl1, or Foxp1 (red) for Olig2-NR1 (G) and Ngn2, Bhlhe22, or Lhx9 (red) for Olig2-NR3 (L) on UMAP visualization of E14 mouse retinal gene expression. Percentages of respective populations that co-express Olig2 are indicated (* denotes significance).



Supplemental Figure 10S. Activity of OLIG2 CRMs in OLIG2+ cells within embryonic chick spinal cords.

(A) Schematic of an E2 chick transverse spinal cord with the field of view (red outline) shown in B. (B) Representative transverse hemi-sections of ventral E2 chick spinal cords incubated for 24 hours *in ovo*, showing (i) OLIG2 RNA localized with GFP RNA (orange arrow), (ii) OLIG2 expression in GFP-cells (yellow arrows), and (iii) GFP RNA expression in OLIG2-cells (magenta arrows). GFP expression is driven by plasmids containing one of the CRM candidate regions (CRM1-3). Scale bar: 20µm.

Method	Primer Name	Sequence
Stagintbc7 vector design	bc7_ins1	GGCGCGCCaGATCGGAAGAGCACACGTCTGAACTCCAGTCACCCCTTA CTTGACAGCTCGTCCATG
	bc7_ins2	CCGATCtGGCGCGCCSWSWSWSWSWSWSWSWSWaGATCG GAAGAGCGTCGTGTAGGGAAAGAGTGTCCGGCCCCCTTGAGCATC T
	bc7_ins1A	GCCTGTATGGCGCGCCaGATCGGAAGAGCACACGTCTGAACTC CAGTCACCCCTTACTTGTACAGCTCGTCCATG
	bc7_ins1B	CTTGACAGCGCGCCaGATCGGAAGAGCACACGTCTGAACTC CAGTCACCCCTTACTTGTACAGCTCGTCCATG
	bc7_ins1C	TCTTCGTGGCGCGCCaGATCGGAAGAGCACACGTCTGAACTCC AGTCACCCCTTACTTGTACAGCTCGTCCATG
	bc7_ins1D	AGGTTACGGCGCGCCaGATCGGAAGAGCACACGTCTGAACTC CAGTCACCCCTTACTTGTACAGCTCGTCCATG
	bc7_ins2A	CCGATCtGGCGCGCCATACAGGCVHBDVHBDVHBDVHBDVHBD VHBDaGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTCCGGCC CCTTGAGCATCT
	bc7_ins2B	CCGATCtGGCGCGCCTGTGCAAGVHBDVHBDVHBDVHBDVHBD VHBDaGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTCCGGCC CCTTGAGCATCT
	bc7_ins2C	CCGATCtGGCGCGCCACGAAGVHBDVHBDVHBDVHBDVHBD VHBDaGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTCCGGCC CCTTGAGCATCT
	bc7_ins2D	CCGATCtGGCGCGCCGTGAACCTVHBDVHBDVHBDVHBDVHBD VHBDaGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTCCGGCC CCTTGAGCATCT
LS-MPRA Adapters	enh7_5'1S	CGATAGTCGACCAATTGCTCGAGGGTGACTGGAGTTCAGACGTG TGCTCTTCCGATC*t
	enh7_5'2phos	/5Phos/GATCGGAAGAGCACACGTCTGAACTCCAGTCACCCCTCAGCA ATTGGTCGACTATCG
	enh7_3'1phos	/5Phos/GGCGCGCCattaaGCTGGTAGACTgcatCT
	enh7_3'2S	AGatgcAGTCTACCAGCttaaGGCGCGCC*t
Amplify fragments	enh7_ampF	CGATAGTCGACCAATTGCTC
	enh7_ampR	AGatgcAGTCTACCAGC
d-MPRA assembly	Olig2_NR1_mutF	CGATAGTCGACCAATTGCTCGAcagacatcaaatcagtaagccctgttg
	Olig2_NR1_mutR	atcgcAGTCTACCAGCttaaTTGTTAGTGGCGTCTGTCCCG
	Olig2_NR2_mutF	CGATAGTCGACCAATTGCTCGAgagtgtctccacatgccaa
	Olig2_NR2_mutR	atcgcAGTCTACCAGCttaaCGTGTTCCTCCAGCTGGCT
	Olig2_NR3_mutF	CGATAGTCGACCAATTGCTCGActagtgcttgaccccatct
	Olig2_NR3_mutR	atcgcAGTCTACCAGCttaaAGTTGAGGCTTGGGATTCGG
	MutF	CGATAGTCGACCAATTGCTCGA
	MutR	atcgcAGTCTACCAGCttaa
	StatadualA_gfpF	GCTGGTAGACTgcatAcctcaCTATATAATGGAAGCTTG
	StatadualA_gfpR	CCTCGAGCAATTGGTCGACTATCGATTACTTGTACAGCTCGTCCA TGCCG
LS-MPRA BC sequencing	bc7_cdnaR	AAGTCAGATGCTCAAGGGGC
	bc7_cdnaF	GCAGAAGAACGGCATCAAGG
	bc7_cdnaR2	GCCGGACACTCTTTCCCTAC
d-MPRA BC sequencing	bc7_cdnaR_WPRE	AGCAGCGTATCCACATAGCG
	bc7_cdnaF*	GCAGAAGAACGGCATCAAGG
	MutR*	atcgcAGTCTACCAGCttaa
	dualmut_IIIIR1F	ACACTCTTTCCCTACACGACGCTCTTCCGATCtACCAATTGCTCG AGG

Table S1. Primer sequences.

Table S1. (continued)

	dualmut_III2F	ACATCTTTCCCTACACGACGCTCTTCCGATCTACCAATTGCTCGACA
	dualmut_III2R	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCtgcAGTCTACCAGCttaat
Construct vector	compvec F	CAGCTCACTCAAAGGCGGTA
	compvec R	CCTCGAGCAATTGGTCGACTATC
Olig2-NR1 and -NR2 constructs	Olig2_NR1F	GACCAATTGCTCGAGGcagacatacaatcagtaagccctgttg
	Olig2_NR1R	ATGTGCTTTTtgttagtgccgtctgtcccg
	Olig2_NR2F	GACCAATTGCTCGAGGgagtggtctccatcagcc
	Olig2_NR2R	GGAACAATGTGCTTTTcgtgttcccagctggc
	Olig2_minpr-GFPint	AAAAGCACATTGTTCCCGCC
	GFPint_vecR	GCCTTTGAGTGAGCTGgcccctcccatatgtcctcc
Olig2-NR3 construct	Olig2_NR3F	GACATATGGGAGGGCCTAGGTGCTTGGACCCCATC
	Olig2_NR3R	GCCTTTGAGTGAGCTGagttgagcctgggattccg
	Vec GFPintF	gaccaattgctcgaggAAAAGCACATTGTTCCCGCC
	GFPint_R	GCCCTCCCATATGTCCTTCC
Olig2-composite construct	Olig2_NR1F*	GACCAATTGCTCGAGGcagacatacaatcagtaagccctgttg
	Olig2comp_NR1R	GGACCACTCttgttagtgccgtctgtcc
	Olig2comp_NR2F	cactaacaGAGTGGTCTCACATGCCCa
	Olig2_NR2R*	GGAACAATGTGCTTTTcgtgttcccagctggc
	Olig2_minpr-GFPint*	AAAAGCACATTGTTCCCGCC
	GFPint_R*	GCCCTCCCATATGTCCTTCC
	Olig2_NR3F*	GACATATGGGAGGGCCTAGGTGCTTGGACCCCATC
	Olig2_NR3R*	GCCTTTGAGTGAGCTGagttgagcctgggattccg
Backbone control constructs	Vec GFPintF*	gaccaattgctcgaggAAAAGCACATTGTTCCCGCC
	Vec_nominprGFPF	accaattgctcgaggGCCACCATGGTGAGCAAG
	GFPint_R*	GCCCTCCCATATGTCCTTCC
Ngn2-CRM1, -CRM2, and -CRM3 constructs	Ngn2-CRM1_F	ACCAATTGCTCGAGGgaagattggagatgaatagtgtc
	Ngn2-CRM1_R	atcgcAGTCTACCAGCaatgctcacatccctccag
	Ngn2-CRM2_F	ACCAATTGCTCGAGGctcatcagtcaccaactgce
	Ngn2-CRM2_R	atcgcAGTCTACCAGCaatgatgggagtgagag
	Ngn2-CRM3_F	ACCAATTGCTCGAGGactcggccctaatgagc
	Ngn2-CRM3_R	atcgcAGTCTACCAGCttatctaatcagataaaatggtctctaaagctcc
	TATaminpr_F	GCTGGTAGACTgcgatCT
	GFPint_vecR	GCCTTTGAGTGAGCTGgcccctcccatatgtcctcc
Ngn2-CRM4 construct	Ngn2-CRM4_F	GACATATGGGAGGGCcctaaaagtgaagcctggag
	Ngn2-CRM4_R	GCCTTTGAGTGAGCTGcaaggaggcaataaacccccc
	Vec_TATaminpr_F	GACCAATTGCTCGAGGGCTGGTAGACTgcgatCT
	GFPint_R*	GCCCTCCCATATGTCCTTCC
eOLIG2-CRM1 construct	eOLIG2_CRM1F	ACCAATTGCTCGAGGaccaaagattatgtcgttttaaac
	eOLIG2_CRM1R	CTTTATAGCCGCGCCGGGctgctgtaatgtttttatttttctc
	eOLIG2_promF	cccgccgcccgtataaagc
	eOLIG2_promR	CTCACCATGGTGGCggtctcacctgctcggg
	GFPint_F	GCCACCATGGTGAGCAAG
eOLIG2-CRM2 construct	GFPint_vecR*	GCCTTTGAGTGAGCTGGCCCTCCCATATGTCCTTCC
	eOLIG2_CRM2F	ACCAATTGCTCGAGGgttacgcgttacgatggtctgac
	eOLIG2_promR*	CTCACCATGGTGGCggtctcacctgctcggg
	GFPint_F*	GCCACCATGGTGAGCAAG
	GFPint_vecR*	GCCTTTGAGTGAGCTGGCCCTCCCATATGTCCTTCC
eOLIG2-CRM3 construct	eOLIG2_CRM3F	GACATATGGGAGGGCgctccattgtcagcgtg
	eOLIG2_CRM3R	GCCTTTGAGTGAGCTGagccgttaatcccgcc
	eOLIG2-vec_promF	ACCAATTGCTCGAGGcccggcggtataaagc
ddPCR	eOLIG2_promR*	CTCACCATGGTGGCggtctcacctgctcggg
	GFPint_R*	GCCCTCCCATATGTCCTTCC
	Olig2_qF	GCGAGCACCTCAAATCTAATTC
	Olig2_qR	AAAAGATCATCGGGTTCTGGG
	Hprt_qF	CCCCAAAATGGTTAAGGTTGC
	Hprt_qR	AACAAAAGTCTGGCCTGTATCC
	*reused	

Table S2. Genome Coordinates for ROIs.

Genome	Chromosome	Start	End	Description
mm10	chr6	115930332	115930431	Rho PPR (Zack et al., 1991)
	chr6	115931731	115931977	Rho RER (Nie et al., 1996)
	chr6	115928742	115929214	Rho CBR (Corbo et al., 2010)
	chr11	50842558	50842758	Grm6 CRM (Kim et al., 2008)
	chr12	84567682	84569914	Vsx2 2.4 kb promoter CRM (Rowan et al., 2005)
	chr12	84552103	84552267	Vsx2 164 bp enhancer CRM (Kim et al., 2008)
	chr12	84532102	84532711	Vsx2 ECR4 homologue (Buenaventura et al., 2018)
	chr12	84532049	84533013	Vsx2 CRC-SE-mR0-37 (Honnell et al., 2022)
	chr12	84550173	84553991	Vsx2 CRC-SE-mR3-17 (Honnell et al., 2022)
	chr7	13397865	13398310	Cabp5 CRM (Kim et al., 2008)
	chr16	91140523	91140769	Olig2-NR1
	chr16	91223668	91223849	Olig2-NR2
	chr16	91232270	91232482	Olig2-NR3
	chr16	91225474	91225600	Olig2 minimal promoter
	chr16	91235256	91238546	Olig2-K23 (Sun et al., 2006)
	chr16	91139504	91141503	Olig2-ULTRA (Chen et al., 2008)
	chr16	91223550	91225549	Olig2-basal promoter (Chen et al., 2008)
	chr16	91191767	91193766	Olig2-EC2 (Fan et al., 2023)
	chr16	91208209	91211318	Olig2-5F7 (Friedli et al., 2010)
	chr3	127628497	127632832	Ngn2-TgN2 (Simmons et al., 2001)
	chr3	127642407	127642608	Ngn2-CRM1
galGal6	chr3	127605638	127605854	Ngn2-CRM2
	chr3	127632596	127632883	Ngn2-CRM3
	chr3	127572578	127572810	Ngn2-CRM4
galGal6	chr1	106522687	106522790	cOLIG2 minimal promoter
	chr1	106486012	106486309	cOLIG2-CRM1
	chr1	106522473	106522687	cOLIG2-CRM2
	chr1	106531687	106531776	cOLIG2-CRM3

Table S3. Association Library characteristics.

Gene-of-interest	Cabp5	Grm6	Rho	Vsx2	Olig2	Neurog2	cOLIG2
BAC clone	RP23-125H22	RP23-417M10	RP23-219M6	RP23-127O21	CH29-613L21	RP23-182M12	CH261-60J3
Size of clone (bp)	175,529	196,198	205,479	224,092	198,802	176,837	234,491
Sequenced reads	32,993,015	35,980,528	33,804,998	38,542,527	99,453,803	123,039,602	104,836,048
Unique fragments sequenced	6,742,335	6,342,682	6,528,168	5,065,113	7,273,814	9,236,041	11,579,155
Predicted total unique fragments (Vmax)	9,349,334	8,328,615	8,750,444	6,151,434	7,882,427	10,090,748	13,254,886
Estimated under-sequencing (fragments)	27.9%	23.8%	25.4%	17.7%	7.7%	8.5%	12.6%
Unique barcode-fragment pairings	9,283,048	8,115,758	8,339,844	7,121,265	17,931,082	27,383,903	44,774,642
Unique barcodes sequenced	2,745,009	3,299,572	3,434,846	3,541,130	6,494,202	7,894,680	6,859,915
Predicted total unique barcodes (Vmax)	3,033,326	3,694,093	3,902,617	3,991,963	6,911,736	8,419,118	7,213,082
Estimated under-sequencing (barcodes)	9.5%	10.7%	12.0%	11.3%	6.0%	6.2%	4.9%
Mean barcodes per fragment±SD	1.38±0.91	1.28±0.73	1.28±0.73	1.41±0.85	2.47±3.23	2.96±3.51	3.87±4.06
Fragments with 1 BC (%)	5,114,287 (75.9)	5,091,388 (80.3)	5,270,065 (80.7)	3,719,724 (73.4)	3,417,706 (47.0)	3,959,601 (42.9)	3,734,149 (32.2)
Barcode Collisions (%)	1,735,600 (63.2)	1,794,631 (54.4)	1,848,433 (53.8)	1,674,020 (47.3)	3,916,149 (60.3)	5,250,728 (66.5)	5,080,559 (70.1)

Table S4. Profiled Abundance Libraries.

Abundance Library	Fragments observed in RNA	Total fragments in Association Library	Percent profiled
Multiplex-ExVivo-A Cabp5	40,810	6,742,335	0.61%
Multiplex-ExVivo-B Cabp5	32,018		0.47%
Multiplex-ExVivo-C Cabp5	65,346		0.97%
Multiplex-InVivoP10-1 Cabp5	62,482		0.93%
Multiplex-InVivoP10-2 Cabp5	52,526		0.78%
Multiplex-InVivoP10-3 Cabp5	84,138		1.25%
Multiplex-ExVivo-A Grm6	16,902	6,342,682	0.27%
Multiplex-ExVivo-B Grm6	14,886		0.23%
Multiplex-ExVivo-C Grm6	41,249		0.65%
Multiplex-InVivoP10-1 Grm6	23,755		0.37%
Multiplex-InVivoP10-2 Grm6	19,854		0.31%
Multiplex-InVivoP10-3 Grm6	43,001		0.68%
Multiplex-ExVivo-A Rho	26,371	6,528,168	0.40%
Multiplex-ExVivo-B Rho	18,541		0.28%
Multiplex-ExVivo-C Rho	20,949		0.32%
Multiplex-InVivoP10-1 Rho	47,920		0.73%
Multiplex-InVivoP10-2 Rho	37,309		0.57%
Multiplex-InVivoP10-3 Rho	44,261		0.68%
Multiplex-ExVivo-A Vsx2	9,219	5,065,113	0.18%
Multiplex-ExVivo-B Vsx2	8,169		0.16%
Multiplex-ExVivo-C Vsx2	23,762		0.47%
Multiplex-InVivoP10-1 Vsx2	18,753		0.37%
Multiplex-InVivoP10-2 Vsx2	14,380		0.28%
Multiplex-InVivoP10-3 Vsx2	32,321		0.64%
Olig2 LS-MPRA101	400,054	7,273,814	5.50%
Olig2 LS-MPRA202	250,985		3.45%
Olig2 LS-MPRA303	190,997		2.63%
Olig2 4hrA	223,053		3.07%
Olig2 4hrB	210,608		2.90%
Olig2 4hrC	258,610		3.56%
Olig2 12hrA	335,853		4.62%
Olig2 12hrB	323,185		4.44%
Olig2 12hrC	202,155		2.78%
Olig2 28hrA	238,341		3.28%
Olig2 28hrB	218,220		3.00%
Olig2 28hrC	212,975		2.93%
Ngn2-bc101	866,959	9,236,041	9.39%
Ngn2-bc202	754,450		8.17%
Ngn2-bc303	2,751,469		29.79%
cOLIG2 retinaA	2,441,108		21.08%
cOLIG2 retinaB	2,392,209		20.66%
cOLIG2 retinaC	2,557,087		22.08%
cOLIG2 retinaD	2,222,483		19.19%
cOLIG2-E2 SC1	601,082		5.19%
cOLIG2-E2 SC2	788,047		6.81%
cOLIG2-E2 SC3	1,074,630		9.28%
cOLIG2-E2 SC4	1,212,194		10.47%
cOLIG2-E4 SC1	3,719,067		32.12%
cOLIG2-E4 SC2	3,374,676		29.14%
cOLIG2-E4 SC3	2,260,800		19.52%
cOLIG2-E4 SC4	1,725,824		14.90%

Table S5. Quantification of GFP and Olig2 co-localization in embryonic mouse retina.

Plasmid	Electroporated Olig2+ cells that express GFP protein (%)		GFP protein+ cells that express Olig2 (%)		Electroporated Olig2-cells that express GFP protein (%)	
	GFP+	GFP-	Olig2+	Olig2-	GFP+	GFP-
Olig2-NR1	100	0	15.6	84.4	66.8	33.2
Olig2-NR2	100	0	29.7	70.3	48.6	51.4
Olig2-NR3	67.9	32.1	17.0	83.0	50.2	49.8
Olig2-Triple Composite	100	0	21.4	78.6	62.1	37.9

Table S6. Quantification of CRM-directed RNA and Olig2 RNA or protein.

Plasmid	Electroporated Olig2 RNA+ cells that expressed GFP (%)				GFP RNA+ cells that expressed Olig2 RNA (%)		GFP protein+ cells that expressed Olig2 RNA (%)		Olig2 RNA-cells that expressed GFP (%)	
	GFP RNA only	GFP RNA & protein	GFP protein only	GFP-	Olig2 RNA+	Olig2 RNA-	Olig2 RNA+	Olig2 RNA-	GFP+	GFP-
Olig2-NR1	19.3	49.8	30.2	0.64	52.6	47.4	40.6	59.4	57.7	42.3
Olig2-NR2	35.6	45.5	10.9	8.0	64.5	35.5	38.8	61.2	27.9	72.1
Olig2-NR3	34.0	5.1	0.0	60.9	39.6	60.4	35.7	64.3	5.5	94.5
Olig2-Triple Composite	14.2	55.0	25.0	5.8	63.3	36.7	46.0	54.0	63.5	36.5

Table S7. Quantification of CRM-directed RNA and *Ngn2* RNA or protein.

Plasmid	Electroporated <i>Ngn2</i> RNA+ cells that expressed GFP (%)				<i>GFP</i> RNA+ cells that expressed <i>Ngn2</i> RNA (%)		<i>GFP</i> protein+ cells that expressed <i>Ngn2</i> RNA (%)		<i>Ngn2</i> RNA-cells that expressed GFP (%)	
	<i>GFP</i> RNA only	<i>GFP</i> RNA & protein	<i>GFP</i> protein only	<i>GFP</i> -	<i>Ngn2</i> RNA+	<i>Ngn2</i> RNA-	<i>Ngn2</i> RNA+	<i>Ngn2</i> RNA-	<i>GFP</i> +	<i>GFP</i> -
Ngn2-CRM1	22.2	30.6	3.7	43.5	6.6	93.4	6.5	93.5	48.2	51.8
Ngn2-CRM2	7.3	86.4	3.3	2.9	20.7	79.3	19.7	80.3	88.8	11.2
Ngn2-CRM3	32.9	59.6	4.4	3.1	38.5	61.5	32.3	67.7	69.3	30.7
Ngn2-CRM4	9.4	0	0	90.6	11.9	88.1	10.6	89.4	14.8	85.2
Ngn2-CRM3 (reduced ex vivo culture)	24.8	60.2	7.7	7.2	49.2	50.8	44.0	56.0	39.3	60.7

Table S8. Quantification of CRM-directed RNA and *cOLIG2* RNA or protein.


Plasmid	Electroporated <i>cOLIG2</i> RNA+ cells that expressed GFP (%)				<i>GFP</i> RNA+ cells that expressed <i>cOLIG2</i> RNA (%)		<i>GFP</i> protein+ cells that expressed <i>cOLIG2</i> RNA (%)		<i>cOLIG2</i> RNA-cells that expressed GFP (%)	
	<i>GFP</i> RNA only	<i>GFP</i> RNA & protein	<i>GFP</i> protein only	<i>GFP</i> -	<i>cOLIG2</i> RNA+	<i>cOLIG2</i> RNA-	<i>cOLIG2</i> RNA+	<i>cOLIG2</i> RNA-	<i>GFP</i> +	<i>GFP</i> -
cOLIG2-CRM1	5.5	90.3	1.7	2.5	45.7	54.3	45.3	54.7	68.2	31.8
cOLIG2-CRM2	4.1	90.9	3.5	1.5	34.5	65.5	32.5	67.5	74.6	25.4
cOLIG2-CRM3	1.0	0	0	98.9					0.4	99.6


Table S9. Quantification of GFP and Olig2 co-localization in postnatal mouse retina.


Plasmid	Electroporated Olig2+ cells that express GFP protein (%)		<i>GFP</i> protein+ cells that express Olig2 (%)		Electroporated Olig2-cells that express GFP protein (%)	
	<i>GFP</i> +	<i>GFP</i> -	Olig2+	Olig2-	<i>GFP</i> +	<i>GFP</i> -
Olig2-NR1	99.5	0.5	88.0	12.0	68.0	32.0
Olig2-NR2	97.9	2.1	85.3	14.7	51.1	48.9
Olig2-NR3	95.8	4.2	70.8	29.2	56.2	43.8
Olig2-Triple Composite	99.8	0.2	80.9	19.1	65.8	34.2


Supplemental Document A

Association Library Supplemental QC plots










D-MPRA Library Supplemental QC plots


Note: d-MPRA libraries were sequenced with different designated names than those in the main text. For the below supplemental plots:


Olig2_NR1 corresponds with the Olig2-NR2 library in the main text


Olig2_NR2 corresponds with the Olig2-NR3 library in the main text


Olig2_NR3 corresponds with the Olig2-NR1 library in the main text












Supplemental Document B



Acknowledgements

First, we thank Dr. Brian Rabe for developing the original LS-MPRA and d-MPRA methods that formed the basis of this study. We would like to thank Dr. Changhee Lee for discussions related to the study design and for expertise in methodology. As well, we thank Dr. Ryoji Amamoto and Henry Bushnell for expertise in methodology. We would also like to thank Dr. Paula Montero Llopis, Dr. Praju Vikas Anekal, and Dr. Adrienne Wells from the MicRoN imaging core at Harvard Medical School for providing expert advice concerning imaging and analysis. We thank BioRender for assisting in the creation of several schematics (<https://biorender.com/rkf4u51>). We are grateful to the Howard Hughes Medical Institute (C.L.C. and A.J.T.), the National Eye Institute (5K99EY034603, R.N.D. and 5R01EY029771 R.C-P) for support.

Additional information

Funding

Funder	Grant reference number	Author
Howard Hughes Medical Institute (HHMI)		Constance L Cepko Alastair J Tulloch
HHS NIH National Eye Institute (NEI)	5K99EY034603	Ryan Nicholas Delgado
HHS NIH National Eye Institute (NEI)	5R01EY029771	Rinaldo Cattalini Preta

Author ORCID iDs

Alastair J Tulloch: <https://orcid.org/0000-0002-7631-4300>

Constance L Cepko: <https://orcid.org/0000-0002-9945-6387>

References

Advani J, Mehta PA, Hamel AR, Mehrotra S, Kiel C, Strunz T, Corso-Diaz X, Kwicklis M, van Asten F, Ratnapriya R, *et al.* (2024) QTL mapping of human retina DNA methylation identifies 87 gene-epigenome interactions in age-related macular degeneration. *Nat Commun* **15**:1972 [PubMed](#)

Agarwal V, Inoue F, Schubach M, Penzar D, Martin BK, Dash PM, Keukeleire P, Zhang Z, Sohota A, Zhao J, *et al.* (2025) Massively parallel characterization of transcriptional regulatory elements. *Nature* [PubMed](#)

Alexiades MR, Cepko C (1996) Quantitative analysis of proliferation and cell cycle length during development of the rat retina. *Dev Dyn* **205**:293-307 [PubMed](#)

Amemiya HM, Kundaje A, Boyle AP (2019) The ENCODE Blacklist: Identification of Problematic Regions of the Genome. *Sci Rep* **9**:9354 [PubMed](#)

Andrews S. (2010) FASTQC. A quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>

Arnold CD, Gerlach D, Stelzer C, Boryn LM, Rath M, Stark A (2013) Genome-wide quantitative enhancer activity maps identified by STARR-seq. *Science* **339**:1074-7 [PubMed](#)

Balasubramanian R, Bui A, Ding Q, Gan L (2014) Expression of LIM-homeodomain transcription factors in the developing and mature mouse retina. *Gene Expr Patterns* **14**:1-8 [PubMed](#)

Balasubramanian R, Bui A, Dong X, Gan L (2018) Lhx9 Is Required for the Development of Retinal Nitric Oxide-Synthesizing Amacrine Cell Subtype. *Mol Neurobiol* **55**:2922-33 [PubMed](#)

Barrett T, Dowle M, Srinivasan A, Gorecki J, Chirico M, Hocking T, Schwendinger B, Krylov I (2025) data.table: Extension of 'data.frame'. version: 1.17.99 <https://Rdatatable.gitlab.io/data.table>

Billings NA, Emerson MM, Cepko CL (2010) Analysis of thyroid response element activity during retinal development. *PLoS One* **5**:e13739 [PubMed](#)

Bonfield JK, Marshall J, Danecek P, Li H, Ohan V, Whitwham A, Keane T, Davies RM (2021) HTSLib: C library for reading/writing high-throughput sequencing data. *Gigascience* **10** [PubMed](#)

Brzezinski JAt, Kim EJ, Johnson JE, Reh TA (2011) Ascl1 expression defines a subpopulation of lineage-restricted progenitors in the mammalian retina. *Development* **138**:3519-31 [PubMed](#)

Buenaventura DF, Ghinia-Tegla MG, Emerson MM (2018) Fate-restricted retinal progenitor cells adopt a molecular profile and spatial position distinct from multipotent progenitor cells. *Dev Biol* **443**:35-49 [PubMed](#)

Buschmann T (2017) DNABarcodes: an R package for the systematic construction of DNA sample tags. *Bioinformatics* **33**:920-22 [PubMed](#)

Bushnell B, Rood J, Singer E (2017) BBMerge - Accurate paired shotgun read merging via overlap. *PLoS One* **12**:e0185056 [PubMed](#)

Catta-Preta R, Lindtner S, Ypsilanti A, Seban N, Price JD, *et al.* (2025) Combinatorial transcription factor binding encodes cis-regulatory wiring of mouse forebrain GABAergic neurogenesis. *Dev Cell* **60**:288-304 e6 [PubMed](#)

Cepko C (2014) Intrinsically different retinal progenitor cells produce specific types of progeny. *Nat Rev Neurosci* **15**:615-27 [PubMed](#)

Chan CSY, Lonfat N, Zhao R, Davis AE, Li L, Wu MR, Lin CH, Ji Z, Cepko CL, Wang S (2020) Cell type- and stage-specific expression of Otx2 is regulated by multiple transcription factors and cis-regulatory modules in the retina. *Development* **147** [PubMed](#)

Chen CT, Gottlieb DI, Cohen BA (2008) Ultraconserved elements in the Olig2 promoter. *PLoS One* **3**:e3946 [PubMed](#)

Clark BS, Stein-O'Brien GL, Shiao F, Cannon GH, Davis-Marcisak E, Sherman T, Santiago CP, Hoang TV, Rajaii F, James-Esposito RE, *et al.* (2019) Single-Cell RNA-Seq Analysis of Retinal Development Identifies NFI Factors as Regulating Mitotic Exit and Late-Born Cell Specification. *Neuron* **102**:1111-26 e5 [PubMed](#)

Consortium EP, Moore JE, Purcaro MJ, Pratt HE, Epstein CB, *et al.* (2020) Expanded encyclopaedias of DNA elements in the human and mouse genomes. *Nature* **583**:699-710 [PubMed](#)

- Corbo JC**, Lawrence KA, Karlstetter M, Myers CA, Abdelaziz M, Dirkes W, Weigelt K, Seifert M, Benes V, Fritsche LG, *et al.* (2010) CRX ChIP-seq reveals the cis-regulatory architecture of mouse photoreceptors. *Genome Res* **20**:1512-25 [PubMed](#)
- Cvekl A**, Callaerts P (2017) PAX6: 25th anniversary and more to learn. *Exp Eye Res* **156**:10-21 [PubMed](#)
- Danecek P**, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM, *et al.* (2021) Twelve years of SAMtools and BCFtools. *Gigascience* **10** [PubMed](#)
- de Melo J**, Du G, Fonseca M, Gillespie LA, Turk WJ, Rubenstein JL, Eisenstat DD. (2005) Dlx1 and Dlx2 function is necessary for terminal differentiation and survival of late-born retinal ganglion cells in the developing mouse retina. *Development* **132**:311-22 [PubMed](#)
- Emerson MM**, Cepko CL (2011) Identification of a retina-specific Otx2 enhancer element active in immature developing photoreceptors. *Dev Biol* **360**:241-55 [PubMed](#)
- Fan C**, Kim D, An H, Park Y (2023) Identifying an oligodendrocyte enhancer that regulates Olig2 expression. *Hum Mol Genet* **32**:835-46 [PubMed](#)
- Farhy C**, Elgart M, Shapira Z, Oron-Karni V, Yaron O, Menuchin Y, Rechavi G, Ashery-Padan R. (2013) Pax6 is required for normal cell-cycle exit and the differentiation kinetics of retinal progenitor cells. *PLoS One* **8**:e76489 [PubMed](#)
- Feng L**, Xie X, Joshi PS, Yang Z, Shibasaki K, Chow RL, Gan L (2006) Requirement for Bhlhb5 in the specification of amacrine and cone bipolar subtypes in mouse retina. *Development* **133**:4815-25 [PubMed](#)
- Friedli M**, Barde I, Arcangeli M, Verp S, Quazzola A, Zakany J, Lin-Marq N, Robyr D, Attanasio C, Spitz F, *et al.* (2010) A systematic enhancer screen using lentivector transgenesis identifies conserved and non-conserved functional elements at the Olig1 and Olig2 locus. *PLoS One* **5**:e15741 [PubMed](#)
- Gel B**, Diez-Villanueva A, Serra E, Buschbeck M, Peinado MA, Malinverni R (2016) regioneR: an R/Bioconductor package for the association analysis of genomic regions based on permutation tests. *Bioinformatics* **32**:289-91 [PubMed](#)
- Github** (2022) 10XGenomics/bamtofastq: Convert 10x BAM files to the original FASTQs compatible with 10x pipelines v1.4.1. <https://github.com/10XGenomics/bamtofastq>
- Gordon PJ**, Yun S, Clark AM, Monuki ES, Murtaugh LC, Levine EM (2013) Lhx2 balances progenitor maintenance with neurogenic output and promotes competence state progression in the developing retina. *J Neurosci* **33**:12197-207 [PubMed](#)
- Haeseleer F**, Sokal I, Verlinde CL, Erdjument-Bromage H, Tempst P, Pronin AN, Benovic JL, Fariss RN, Palczewski K (2000) Five members of a novel Ca(2+)-binding protein (CABP) subfamily with similarity to calmodulin. *J Biol Chem* **275**:1247-60 [PubMed](#)
- Hafler BP**, Surzenko N, Beier KT, Punzo C, Trimarchi JM, Kong JH, Cepko CL (2012) Transcription factor Olig2 defines subpopulations of retinal progenitor cells biased toward specific cell fates. *Proc Natl Acad Sci U S A* **109**:7882-7 [PubMed](#)
- Hao Y**, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G, Madad S, Fernandez-Granda C, *et al.* (2024) Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat Biotechnol* **42**:293-304 [PubMed](#)
- Harris CR**, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, *et al.* (2020) Array programming with NumPy. *Nature* **585**:357-62 [PubMed](#)
- Hatakeyama J**, Tomita K, Inoue T, Kageyama R (2001) Roles of homeobox and bHLH genes in specification of a retinal cell type. *Development* **128**:1313-22 [PubMed](#)
- Heinz S**, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**:576-89 [PubMed](#)
- Henke RM**, Savage TK, Meredith DM, Glasgow SM, Hori K, Dumas J, MacDonald RJ, Johnson JE (2009) Neurog2 is a direct downstream target of the Ptf1a-Rbpj transcription complex in dorsal spinal cord. *Development* **136**:2945-54 [PubMed](#)

- Hollyday M, Hamburger V (1977) An autoradiographic study of the formation of the lateral motor column in the chick embryo. *Brain Res* **132**:197-208 [PubMed](#)
- Honnell V, Norrie JL, Patel AG, Ramirez C, Zhang J, Lai YH, Wan S, Dyer MA (2022) Identification of a modular super-enhancer in murine retinal development. *Nat Commun* **13**:253 [PubMed](#)
- Hrvatin S, Tzeng CP, Nagy MA, Stroud H, Koutsoumpa C, Wilcox OF, Assad EG, Green J, Harvey CD, Griffith EC, *et al.* (2019) A scalable platform for the development of cell-type-specific viral drivers. *eLife* **8** [PubMed](#)
- Huang L, Hu F, Feng L, Luo XJ, Liang G, Zeng XY, Yi JL, Gan L (2014) Bhlhb5 is required for the subtype development of retinal amacrine and bipolar cells in mice. *Dev Dyn* **243**:279-89 [PubMed](#)
- Hufnagel RB, Le TT, Riesenberger AL, Brown NL (2010) Neurog2 controls the leading edge of neurogenesis in the mammalian retina. *Dev Biol* **340**:490-503 [PubMed](#)
- Hunter JD (2007) Matplotlib: A 2D Graphics Environment. In: Computing in Science & Engineering. **9** pp. 90-95 <https://doi.org/10.1109/MCSE.2007.55>
- Inoue F, Ahituv N (2015) Decoding enhancers using massively parallel reporter assays. *Genomics* **106**:159-64 [PubMed](#)
- Jadhav AP, Mason HA, Cepko CL (2006) Notch 1 inhibits photoreceptor production in the developing mammalian retina. *Development* **133**:913-23 [PubMed](#)
- Jiang Y, Ding Q, Xie X, Libby RT, Lefebvre V, Gan L (2013) Transcription factors SOX4 and SOX11 function redundantly to regulate the development of mouse retinal ganglion cells. *J Biol Chem* **288**:18429-38 [PubMed](#)
- Kaufman ML, Park KU, Goodson NB, Chew S, Bersie S, Jones KL, Lamba DA, Brzezinski JA (2019) Transcriptional profiling of murine retinas undergoing semi-synchronous cone photoreceptor differentiation. *Dev Biol* **453**:155-67 [PubMed](#)
- Kent WJ (2002) BLAT--the BLAST-like alignment tool. *Genome Res* **12**:656-64 [PubMed](#)
- Kim DS, Matsuda T, Cepko CL (2008) A core paired-type and POU homeodomain-containing transcription factor program drives retinal bipolar cell gene expression. *J Neurosci* **28**:7748-64 [PubMed](#)
- Kowalchuk AM, Maurer KA, Shoja-Taheri F, Brown NL (2018) Requirements for Neurogenin2 during mouse postnatal retinal neurogenesis. *Dev Biol* **442**:220-35 [PubMed](#)
- Krueger F (2015) Trim Galore!: A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files.
- Lagunas T Jr, Plassmeyer SP, Fischer AD, Friedman RZ, Rieger MA, Selmanovic D, Sarafinowska S, Sol YK, Kasper MJ, Fass SB, *et al.* (2023) A Cre-dependent massively parallel reporter assay allows for cell-type specific assessment of the functional effects of non-coding elements in vivo. *Commun Biol* **6**:1151 [PubMed](#)
- Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:357-9 [PubMed](#)
- Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ (2013) Software for computing and annotating genomic ranges. *PLoS Comput Biol* **9**:e1003118 [PubMed](#)
- Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**:3094-100 [PubMed](#)
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**:1754-60 [PubMed](#)
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:2078-9 [PubMed](#)
- Li S, Mo Z, Yang X, Price SM, Shen MM, Xiang M (2004) Foxn4 controls the genesis of amacrine and horizontal cells by retinal progenitors. *Neuron* **43**:795-807 [PubMed](#)

- Liu IS, Chen JD, Ploder L, Vidgen D, van der Kooy D, Kalnins VI, McInnes RR (1994) Developmental expression of a novel murine homeobox gene (Chx10): evidence for roles in determination of the neuroretina and inner nuclear layer. *Neuron* **13**:377-93 [PubMed](#)
- Lonfat N, Wang S, Lee C, Garcia M, Choi J, Park PJ, Cepko C (2021) Cis-regulatory dissection of cone development reveals a broad role for Otx2 and Oc transcription factors. *Development* **148** [PubMed](#)
- Lu QR, Yuk D, Alberta JA, Zhu Z, Pawlitzky I, Chan J, McMahon AP, Stiles CD, Rowitch DH (2000) Sonic hedgehog--regulated oligodendrocyte lineage genes encoding bHLH proteins in the mammalian central nervous system. *Neuron* **25**:317-29 [PubMed](#)
- Lunde A, Glover JC (2020) A versatile toolbox for semi-automatic cell-by-cell object-based colocalization analysis. *Sci Rep* **10**:19027 [PubMed](#)
- Lyu P, Hoang T, Santiago CP, Thomas ED, Timms AE, Appel H, Gimmen M, Le N, Jiang L, Kim DW, *et al.* (2021) Gene regulatory networks controlling temporal patterning, neurogenesis, and cell-fate specification in mammalian retina. *Cell Rep* **37**:109994 [PubMed](#)
- Ma W, Wang SZ (2006) The final fates of neurogenin2-expressing cells include all major neuron types in the mouse retina. *Mol Cell Neurosci* **31**:463-9 [PubMed](#)
- Marquardt T, Ashery-Padan R, Andrejewski N, Scardigli R, Guillemot F, Gruss P (2001) Pax6 is required for the multipotent state of retinal progenitor cells. *Cell* **105**:43-55 [PubMed](#)
- McAfee JC, Bell JL, Krupa O, Matoba N, Stein JL, Won H (2022) Focus on your locus with a massively parallel reporter assay. *J Neurodev Disord* **14**:50 [PubMed](#)
- McDowell GS, Hardwick LJ, Philpott A (2014) Complex domain interactions regulate stability and activity of closely related proneural transcription factors. *Biochem Biophys Res Commun* **450**:1283-90 [PubMed](#)
- McInnes L, Healy J, Saul N, Großberger L (2018) UMAP: Uniform Manifold Approximation and Projection. *Journal of Open Source Software* **3**:861 <https://doi.org/10.21105/joss.00861>
- McKinney W (2010) Data Structures for Statistical Computing in Python. In: Proceedings of the 9th Python in Science Conference. Austin, TX: SciPy. pp. 56-61
- Melnikov A, Murugan A, Zhang X, Tesileanu T, Wang L, Rogov P, Feizi S, Gnirke A, Callan CG, Kinney JB, *et al.* (2012) Systematic dissection and optimization of inducible enhancers in human cells using a massively parallel reporter assay. *Nat Biotechnol* **30**:271-7 [PubMed](#)
- Montana CL, Myers CA, Corbo JC (2011) Quantifying the activity of cis-regulatory elements in the mouse retina by explant electroporation. *J Vis Exp* [PubMed](#)
- Mu X, Fu X, Beremand PD, Thomas TL, Klein WH (2008) Gene regulation logic in retinal ganglion cell development: Isl1 defines a critical branch distinct from but overlapping with Pou4f2. *Proc Natl Acad Sci U S A* **105**:6942-7 [PubMed](#)
- Nakazaki H, Reddy AC, Mania-Farnell BL, Shen YW, Ichi S, McCabe C, George D, McLone DG, Tomita T, Mayanil CS (2008) Key basic helix-loop-helix transcription factor genes Hes1 and Ngn2 are regulated by Pax3 during mouse embryonic development. *Dev Biol* **316**:510-23 [PubMed](#)
- Nie Z, Chen S, Kumar R, Zack DJ (1996) RER, an evolutionarily conserved sequence upstream of the rhodopsin gene, has enhancer activity. *J Biol Chem* **271**:2667-75 [PubMed](#)
- Nishida A, Furukawa A, Koike C, Tano Y, Aizawa S, Matsuo I, Furukawa T (2003) Otx2 homeobox gene controls retinal photoreceptor cell fate and pineal gland development. *Nat Neurosci* **6**:1255-63 [PubMed](#)
- O'Leary NA, Wright MW, Brister JR, Ciuffo S, Haddad D, *et al.* (2016) Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res* **44**:D733-45 [PubMed](#)
- Oron-Karni V, Farhy C, Elgart M, Marquardt T, Remizova L, Yaron O, Xie Q, Cvekl A, Ashery-Padan R (2008) Dual requirement for Pax6 in retinal progenitor cells. *Development* **135**:4037-47 [PubMed](#)
- Pagès H, Aboyoun P, Gentleman R, DebRoy S. (2025) Biostrings: Efficient manipulation of biological strings. R package version 2.76.0. <https://bioconductor.org/packages/Biostrings>

- Perez G, Barber GP, Benet-Pages A, Casper J, Clawson H, Diekhans M, Fischer C, Gonzalez JN, Hinrichs AS, Lee CM, *et al.* (2025) The UCSC Genome Browser database: 2025 update. *Nucleic Acids Res* **53**:D1243-D49 [PubMed](#)
- Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**:841-2 [PubMed](#)
- Rabe BA (2020) *Investigation of the Multiple Roles of Notch Signaling in Embryonic Retinal Development Using Novel High-Throughput Techniques*. PhD thesis Cambridge, MA: Harvard University.
- Rabe BA, Cepko C (2020) A Simple Enhancement for Gibson Isothermal Assembly. *bioRxiv* 2020.06.14.150979
- Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dundar F, Manke T. (2016) deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**:W160-5 [PubMed](#)
- Rayon T, Stamataki D, Perez-Carrasco R, Garcia-Perez L, Barrington C, Melchionda M, Exelby K, Lazaro J, Tybulewicz VLJ, Fisher EMC, *et al.* (2020) Species-specific pace of development is associated with differences in protein stability. *Science* **369** [PubMed](#)
- Remez LA, Onishi A, Menuchin-Lasowski Y, Biran A, Blackshaw S, Wahlin KJ, Zack DJ, Ashery-Padan R (2017) Pax6 is essential for the generation of late-born retinal neurons and for inhibition of photoreceptor-fate during late stages of retinogenesis. *Dev Biol* **432**:140-50 [PubMed](#)
- Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP (2011) Integrative genomics viewer. *Nat Biotechnol* **29**:24-6 [PubMed](#)
- Rowan S, Cepko CL (2004) Genetic analysis of the homeodomain transcription factor Chx10 in the retina using a novel multifunctional BAC transgenic mouse reporter. *Dev Biol* **271**:388-402 [PubMed](#)
- Rowan S, Cepko CL (2005) A POU factor binding site upstream of the Chx10 homeobox gene is required for Chx10 expression in subsets of retinal progenitor cells and bipolar cells. *Dev Biol* **281**:240-55 [PubMed](#)
- Satpathy AT, Granja JM, Yost KE, Qi Y, Meschi F, *et al.* (2019) Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion. *Nat Biotechnol* **37**:925-36 [PubMed](#)
- Scardigli R, Baumer N, Gruss P, Guillemot F, Le Roux I. (2003) Direct and concentration-dependent regulation of the proneural gene Neurogenin2 by Pax6. *Development* **130**:3269-81 [PubMed](#)
- Scardigli R, Schuurmans C, Gradwohl G, Guillemot F (2001) Crossregulation between Neurogenin2 and pathways specifying neuronal identity in the spinal cord. *Neuron* **31**:203-17 [PubMed](#)
- Shekhar K, Lapan SW, Whitney IE, Tran NM, Macosko EZ, Kowalczyk M, Adiconis X, Levin JZ, Nemesh J, Goldman M, *et al.* (2016) Comprehensive Classification of Retinal Bipolar Neurons by Single-Cell Transcriptomics. *Cell* **166**:1308-23 e30 [PubMed](#)
- Shen SQ, Myers CA, Hughes AE, Byrne LC, Flannery JG, Corbo JC (2016a) Massively parallel cis-regulatory analysis in the mammalian central nervous system. *Genome Res* **26**:238-55 [PubMed](#)
- Shen W, Le S, Li Y, Hu F. (2016b) SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. *PLoS One* **11**:e0163962 [PubMed](#)
- Shibasaki K, Takebayashi H, Ikenaka K, Feng L, Gan L (2007) Expression of the basic helix-loop-factor Olig2 in the developing retina: Olig2 as a new marker for retinal progenitors and late-born cells. *Gene Expr Patterns* **7**:57-65 [PubMed](#)
- Simmons AD, Horton S, Abney AL, Johnson JE (2001) Neurogenin2 expression in ventral and dorsal spinal neural tube progenitor cells is regulated by distinct enhancers. *Dev Biol* **229**:327-39 [PubMed](#)
- Smith T, Heger A, Sudbery I (2017) UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**:491-99 [PubMed](#)
- Soula C, Danesin C, Kan P, Grob M, Poncet C, Cochard P (2001) Distinct sites of origin of oligodendrocytes and somatic motoneurons in the chick spinal cord: oligodendrocytes arise from Nkx2.2-expressing progenitors by a Shh-dependent mechanism. *Development* **128**:1369-79 [PubMed](#)

- Sun C, Ruzycski PA, Chen S (2023) Rho enhancers play unexpectedly minor roles in Rhodopsin transcription and rod cell integrity. *Sci Rep* **13**:12899 [PubMed](#)
- Sun T, Hafler BP, Kaing S, Kitada M, Ligon KL, Widlund HR, Yuk DI, Stiles CD, Rowitch DH (2006) Evidence for motoneuron lineage-specific regulation of Olig2 in the vertebrate neural tube. *Dev Biol* **292**:152-64 [PubMed](#)
- Tareen A, Kinney JB (2020) Logomaker: beautiful sequence logos in Python. *Bioinformatics* **36**:2272-74 [PubMed](#)
- Thorvaldsdottir H, Robinson JT, Mesirov JP (2013) Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* **14**:178-92 [PubMed](#)
- Trimarchi JM, Stadler MB, Cepko CL (2008) Individual retinal progenitor cells display extensive heterogeneity of gene expression. *PLoS One* **3**:e1588 [PubMed](#)
- Tulloch A (2025) LS-MPRA / d-MPRA Data Repository. Harvard Dataverse. <https://doi.org/10.7910/DVN/TW0ZQL>
- Ueda Y, Iwakabe H, Masu M, Suzuki M, Nakanishi S (1997) The mGluR6 5' upstream transgene sequence directs a cell-specific and developmentally regulated expression in retinal rod and ON-type cone bipolar cells. *J Neurosci* **17**:3014-23 [PubMed](#)
- van der Loo MPJ (2014) The stringdist package for approximate string matching. *The R Journal* <https://doi.org/10.32614/RJ-2014-011>
- Vosper JM, Fiore-Herich CS, Horan I, Wilson K, Wise H, Philpott A (2007) Regulation of neurogenin stability by ubiquitin-mediated proteolysis. *Biochem J* **407**:277-84 [PubMed](#)
- Wang S, Sengel C, Emerson MM, Cepko CL (2014) A gene regulatory network controls the binary fate decision of rod and bipolar cells in the vertebrate retina. *Dev Cell* **30**:513-27 [PubMed](#)
- Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis* New York: Springer-Verlag.
- Yaron O, Farhy C, Marquardt T, Applebury M, Ashery-Padan R (2006) Notch1 functions to suppress cone-photoreceptor fate specification in the developing mouse retina. *Development* **133**:1367-78 [PubMed](#)
- Young RW (1985) Cell differentiation in the retina of the mouse. *Anat Rec* **212**:199-205 [PubMed](#)
- Zack DJ, Bennett J, Wang Y, Davenport C, Klaunberg B, Gearhart J, Nathans J (1991) Unusual topography of bovine rhodopsin promoter-lacZ fusion gene expression in transgenic mouse retinas. *Neuron* **6**:187-99 [PubMed](#)
- Zhang J, Roberts JM, Chang F, Schwakopf J, Vetter ML (2023) Jarid2 promotes temporal progression of retinal progenitors via repression of Foxp1. *Cell Rep* **42**:112237 [PubMed](#)
- Zhang Q, Zagozewski J, Cheng S, Dixit R, Zhang S, de Melo J, Mu X, Klein WH, Brown NL, Wigle JT, *et al.* (2017) Regulation of Brn3b by DLX1 and DLX2 is required for retinal ganglion cell differentiation in the vertebrate retina. *Development* **144**:1698-711 [PubMed](#)
- Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, *et al.* (2008) Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**:R137 [PubMed](#)
- Zhou Q, Wang S, Anderson DJ (2000) Identification of a novel family of oligodendrocyte lineage-specific basic helix-loop-helix transcription factors. *Neuron* **25**:331-43 [PubMed](#)
- Clark BS, Stein-O'Brien GL, Shiao F, Cannon GH, Davis-Marcisak E, Sherman T, Santiago CP, Hoang TV, Rajaii F, James-Esposito RE, *et al.* (2018) Single cell RNA-Seq atlas of the developing murine retina. Github. ID developing_mouse_retina_scRNASeq https://github.com/gofflab/developing_mouse_retina_scRNASeq
- Lyu P, Hoang T, Santiago CP, Thomas ED, Timms AE, Appel H, Gimmen M, Le N, Jiang L, Kim DW, *et al.* (2021) Gene regulatory networks controlling temporal patterning, neurogenesis, and cell fate specification in the mammalian retina. NCBI Gene Expression Omnibus. ID GSE181251 <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE181251>

Peer reviews

Reviewer #1 (Public review):

MPRAs are a high-throughput and powerful tool for assaying the regulatory potential of genomic sequences. However, linking MPRA-nominated regulatory sequences to their endogenous target genes, and identifying the more specific functional regions within these sequences can be challenging. MPRAs that tile a genomic region, and saturation mutagenesis-based MPRAs can help to address these challenges. In this work, Tulloch et al. describe a streamlined MPRA system for the identification and investigation of the regulatory elements surrounding a gene of interest with high resolution. The use of BACs covering a locus of interest to generate MPRA libraries allows for an unbiased, and high-coverage assessment of a particular region. Follow up degenerate MPRAs, where each nucleotide in the nominated sequences are systematically mutated, then can point to key motifs driving their regulatory activity. The authors present this MPRA platform as straightforward, easily customizable, and less time- and resource-intensive than traditional MPRA designs. They demonstrate the utility of their design in the context of the developing mouse retina, where they first use the LS-MPRA to identify active regulatory elements for select retinal genes, followed by d-MPRA which allowed them to dissect the functional regions within those elements and nominate important regulatory motifs. These assays were able to recapitulate some previously known cis-regulatory modules (CRMs), as well as identify some new potential regulatory regions. Follow up experiments assessing co-localization of the gene of interest with the CRM-linked GFP reporter in the target cells, and CUT&RUN assays to confirm transcription factor binding to nominated motifs provided support linking these CRMs to the genes of interest. Overall, this method appears flexible and could be an easy to implement tool for other investigators aiming to study their locus of interest with high resolution.

Strengths:

- (1) The method of fragmenting BACs allows for high, overlapping coverage of the region of interest.
- (2) The d-MPRA method was an efficient way to identify key functional transcription factor motifs, and nominate specific transcription factor-driven regulatory pathways that could be studied further.
- (3) Additional assays like co-expression analyses using the endogenous gene promoter, and use of the Notch inhibitor in the case of *Olig2*, helped correlate the activity of the CRMs to the expression of the gene of interest, and distinguish false positives from the initial MPRA.
- (4) The use of these assays across different time points, tissues, and even species demonstrated that they can be used across many contexts to identify both common and divergent regulatory mechanisms for the same gene.

Weaknesses:

- (1) The LS-MPRA assay most strongly identified promoters, which are not usually novel regulatory elements you would try to discover, and the signal to noise ratio for more TSS-distal, non-promoter regulatory elements was usually high, making it difficult to discriminate lower activity CRMs, like enhancers, from the background. For example, NR2 and NR3 in Figure 3 have very minimal activity peaks (NR3 seems non-existent). The ex vivo data in Figure 2 is similarly noisy. Is there a particular metric or calculation that was or could be used to quantitatively or statistically call a peak above the background? The authors mention in the discussion some adjustments that could reduce the noise, such as increased sequencing depth, which I think is needed to make these initial LS-MPRA results and the benchmarking of this assay more convincing and impactful.

<https://doi.org/10.7554/eLife.107565.2.sa3>

Reviewer #2 (Public review):

Summary:

In this study, Tulloch et al. developed two modified massively parallel reporter assays (MPRAs) and applied them to identify cis-regulatory modules (CRMs) - genomic regions that activate gene expression - controlling retinal gene expression. These CRMs usually function at specific developmental stages and in distinct cell types to orchestrate retinal development. Studying them provides insights into how retinal progenitor cells give rise to various retinal cell types.

The first assay, named locus-specific MPRA (LS-MPRA), tests all genomic regions within 150-300 kb of the gene of interest, rather than relying on previously predicted candidate regulatory elements. This approach reduces potential bias introduced during candidate selection, lowers the cost of synthesizing a library of candidate sequences, and simplifies library preparation. The LS-MPRA libraries were electroporated into mouse retinas in vivo or ex vivo. To benchmark the method, the authors first applied LS-MPRA near stably expressed retinal genes (e.g., *Rho*, *Cabp5*, *Grm6*, and *Vsx2*), and successfully identified both known and novel CRMs. They then used LS-MPRA to identify CRMs in embryonic mouse retinas, near *Olig2* and *Ngn2*, genes expressed in subsets of retinal progenitor cells. Similar experiments were conducted in chick retinas and postnatal mouse retinas, revealing some CRMs with conserved activity across species and developmental stages.

Although the study identified CRMs with robust reporter activity in *Olig2*⁺ or *Ngn2*⁺ cells, the data do not provide sufficient evidence to support the claims that these CRMs regulate *Olig2* or *Ngn2*, rather than other nearby genes, in a cell type-specific manner. For example, the authors propose that three regions (NR1/2/3) regulate *Olig2* specifically in retinal progenitor cells based on: 1) the three regions are close to *Olig2*, 2) increased *Olig2* expression and NR1/2/3 activity upon Notch inhibition, and 3) reporter activity observed in *Olig2*⁺ cells (though also present in many *Olig2*⁻ cells). While these are promising findings, they do not directly support the claims.

The second assay, called degenerate MPRA (d-MPRA), introduces random point mutations into CRMs via error-prone PCR to assess the impact of sequence variations on regulatory activity. This approach was used on NR1/2/3 to identify mutations that alter CRM activity, potentially by influencing transcription factor binding. The authors inferred candidate transcription factors, such as *Mybl1* and *Otx2*, through motif analysis, co-expression with *Olig2* (based on single-cell RNA-seq), and CUR&RUN profiling. While some transcription factors identified in this way overlapped with the d-MPRA results, others did not. This raises questions about how well d-MPRA complements other methods for identifying TF binding sites.

Strengths:

The study introduces two technically robust MPRA protocols that offer advantages over standard methods, such as avoiding reliance on predefined candidate regions, reducing cost and labor, and minimizing selection bias.

The identified regulatory elements and transcription factors contribute to our understanding of gene regulation in retinal development and may have translational potential for cell type-specific gene delivery into developing retinas.

Weakness:

Like other MPRA-based approaches, LS-MPRA mainly tests whether a sequence can drive expression of a reporter gene in given cell type(s). However, this type of assay generally does not show which endogenous gene the sequence regulates. In this study, the evidence supporting gene-specific CRMs is largely correlative. The evidence for cell-type-specific CRMs is also not fully supported (e.g., reporter expression is observed in the intended cell type as well as additional cell types). If further validation in the native genomic context (e.g., CRISPRi of the candidate element followed by RNA-seq across relevant cell types) is out of scope, the manuscript should avoid wording that implies definitive target gene assignment or cell-type specificity.

<https://doi.org/10.7554/eLife.107565.2.sa2>

Reviewer #3 (Public review):

Summary:

Use of reporter assays to understand the regulatory mechanisms controlling gene expression moves beyond simple correlations of cis-regulatory sequence accessibility, evolutionary sequence conservation, and epigenetic status with gene expression, instead quantifying regulatory sequence activity for individual elements. Tulloch et al., provide systematic characterization of two new reporter assay techniques (LS-MPRA and d-MPRA) to comprehensively identify cis-regulatory sequences contained within genomic loci of interest during retinal development. The authors then apply LS-MPRA and d-MPRA to identify putative cis-regulatory sequences controlling *Olig2* and *Ngn2* expression, including potential regulatory motifs that known retinal transcription factors may bind. Transcription factor binding to regulatory sequences is then assessed via CUT&RUN. The broader utility of the techniques are then highlighted by performing the assays across development, across species, and across tissues.

Strengths:

The authors validate the reporter assays on retinal loci for which the regulatory sequences are known (*Rho*, *Vsx2*, *Grm6*, *Cabp5*) mostly confirming known regulatory sequence activity but highlighting either limitations of the current technology or discrepancies of previous reporter assays and known biology. The techniques are then applied to loci of interest (*Olig2* and *Ngn2*) to better understand the regulatory sequences driving expression of these transcription factors across retinal development within subsets of retinal progenitor cells, identifying novel regulatory sequences through comprehensive profiling of the region.

LS-MPRA provides broad coverage of loci of interest

d-MPRA identifies sequence features that are important for cis-regulatory sequence activity.

The authors take into account transcript and protein stability when determining the correlation of putative enhancer sequence activity with target gene expression.

Overall, the manuscript highlights the utility of the techniques to identify novel cis-regulatory sequence contributions to gene expression, including systematic characterizations of sequence motifs conferring activating or repressive functions.

Limitations:

Barcoding strategies have the potential to induce high collision rates (see Table S3) that may lead to misinterpretation of the data and/or high false positive/negative rates.

There are limited robust methods to distinguish differentially active versus inactive CRMs in the LS-MPRA.

<https://doi.org/10.7554/eLife.107565.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public review):

MPRAs are a high-throughput and powerful tool for assaying the regulatory potential of genomic sequences. However, linking MPRA-nominated regulatory sequences to their endogenous target genes and identifying the more specific functional regions within these sequences can be challenging. MPRAs that tile a genomic region, and saturation mutagenesis-based MPRAs, can help to address these challenges. In this work, Tulloch et al. describe a streamlined MPRA system for the identification and investigation of the regulatory elements surrounding a gene of interest with high resolution. The use of BACs covering a locus of interest to generate MPRA libraries allows for an unbiased and high-coverage assessment of a particular region. Follow-up degenerate MPRAs, where each nucleotide in the nominated sequences is systematically mutated, can then point to key motifs driving their regulatory activity. The authors present this MPRA platform as straightforward, easily customizable, and less time- and resource-intensive than traditional MPRA designs. They demonstrate the utility of their design in the context of the developing mouse retina, where they first use the LS-MPRA to identify active regulatory elements for select retinal genes, followed by d-MPRA, which allowed them to dissect the functional regions within those elements and nominate important regulatory motifs. These assays were able to recapitulate some previously known cis-regulatory modules (CRMs), as well as identify some new potential regulatory regions. Follow-up experiments assessing co-localization of the gene of interest with the CRM-linked GFP reporter in the target cells, and CUT&RUN assays to confirm transcription factor binding to nominated motifs, provided support linking these CRMs to the genes of interest. Overall, this method appears flexible and could be an easy-to-implement tool for other investigators aiming to study their locus of interest with high resolution.

Strengths:

- (1) The method of fragmenting BACs allows for high, overlapping coverage of the region of interest.
- (2) The d-MPRA method was an efficient way to identify key functional transcription factor motifs and nominate specific transcription factor-driven regulatory pathways that could be studied further.
- (3) Additional assays like co-expression analyses using the endogenous gene promoter, and use of the Notch inhibitor in the case of *Olig2*, helped correlate the activity of the CRMs to the expression of the gene of interest, and distinguish false positives from the initial MPRA.
- (4) The use of these assays across different time points, tissues, and even species demonstrated that they can be used across many contexts to identify both common and divergent regulatory mechanisms for the same gene.

Weaknesses:

The LS-MPRA assay most strongly identified promoters, which are not usually novel regulatory elements you would try to discover, and the signal-to-noise ratio for more TSS-distal, non-promoter regulatory elements was usually high, making it difficult to

discriminate lower activity CRMs, like enhancers, from the background. For example, NR2 and NR3 in Figure 3 have very minimal activity peaks (NR3 seems non-existent). The ex vivo data in Figure 2 are similarly noisy. Is there a particular metric or calculation that was or could be used to quantitatively or statistically call a peak above the background? The authors mention in the discussion some adjustments that could reduce the noise, such as increased sequencing depth, which I think is needed to make these initial LS-MPRA results and the benchmarking of this assay more convincing and impactful.

Much of the statistical and quantitative data asked for by the Reviewers have been provided in the Revision. However, it is important to note that the types of statistics using peak callers asked for regarding candidate choice will be of limited value. If one is testing a library in a single cell type in vitro, and/or running genome-wide assays, these statistics could aid in the choice of candidates. However, here we are electroporating a complex and dynamic set of cells, with each cell type constituting what can be very different frequencies (e.g. Olig2-expressing cells are <2.4% of cells). This fact alone will give different apparent signal to noise values. In addition, at least for Olig2 and Ngn2, their expression is very transient, suggesting dynamic regulation by what is likely multiple positive and negative CRMs. An additional confound is that the level of expression of each gene that one might test is variable. All of these variables render a statistical prediction of candidates to be less valuable than one might hope, and might lead one to miss those CRMs of interest, particularly those in a small subset of cells. Instead, we suggest that one use one's own level of interest and knowledge in choosing CRM candidates. We provide several examples of experimental, rather than purely statistical, approaches that might help in one's choice of candidates. We used a functional read-out of CRM activity (Notch perturbation), carried out in the context of the entire LS-MPRA library, as one method. Co-expression in single cells of candidate regulators identified by the d-MPRA is another. One can of course use chromatin structure and sequence conservation, as used in many studies of regulatory regions, as other ways to narrow down candidates. The d-MPRA predictions also can be viewed in light of previous genetic studies, i.e. mutations in TFs that effect the cell type of interest or the regulation of the gene of interest, as we were able to do here for CRMs predicted to be regulated by Otx2.

Reviewer #2 (Public review):

Summary:

In this study, Tulloch et al. developed two modified massively parallel reporter assays (MPRAs) and applied them to identify cis-regulatory modules (CRMs) - genomic regions that activate gene expression, controlling retinal gene expression. These CRMs usually function at specific developmental stages and in distinct cell types to orchestrate retinal development. Studying them provides insights into how retinal progenitor cells give rise to various retinal cell types.

The first assay, named locus-specific MPRA (LS-MPRA), tests all genomic regions within 150-300 kb of the gene of interest, rather than relying on previously predicted candidate regulatory elements. This approach reduces potential bias introduced during candidate selection, lowers the cost of synthesizing a library of candidate sequences, and simplifies library preparation. The LS-MPRA libraries were electroporated into mouse retinas in vivo or ex vivo. To benchmark the method, the authors first applied LS-MPRA near stably expressed retinal genes (e.g., Rho, Cabp5, Grm6, and Vsx2), and successfully identified both known and novel CRMs. They then used LS-MPRA to identify CRMs in embryonic mouse retinas, near Olig2 and Ngn2, genes expressed in subsets of retinal progenitor cells. Similar experiments were conducted in chick retinas and postnatal mouse retinas, revealing some CRMs with conserved activity across species and developmental stages.

Although the study identified CRMs with robust reporter activity in Olig2+ or Ngn2+ cells, the data do not provide sufficient evidence to support the claims that these CRMs

regulate *Olig2* or *Ngn2*, rather than other nearby genes, in a cell-type-specific manner. For example, the authors propose that three regions (NR1/2/3) regulate *Olig2* specifically in retinal progenitor cells based on: (1) the three regions are close to *Olig2*, (2) increased *Olig2* expression and NR1/2/3 activity upon Notch inhibition, and (3) reporter activity observed in *Olig2*⁺ cells (though also present in many *Olig2*⁻ cells). While these are promising findings, they do not directly support the claims.

The second assay, called degenerate MPRA (d-MPRA), introduces random point mutations into CRMs via error-prone PCR to assess the impact of sequence variations on regulatory activity. This approach was used on NR1/2/3 to identify mutations that alter CRM activity, potentially by influencing transcription factor binding. The authors inferred candidate transcription factors, such as *Mybl1* and *Otx2*, through motif analysis, co-expression with *Olig2* (based on single-cell RNA-seq), and CUR&RUN profiling. While some transcription factors identified in this way overlapped with the d-MPRA results, others did not. This raises questions about how well d-MPRA complements other methods for identifying transcriptional regulators.

Strengths:

(1) The study introduces two technically robust MPRA protocols that offer advantages over standard methods, such as avoiding reliance on predefined candidate regions, reducing cost and labor, and minimizing selection bias.

(2) The identified regulatory elements and transcription factors contribute to our understanding of gene regulation in retinal development and may have translational potential for cell-type-specific gene delivery into developing retinas.

Weaknesses:

(1) The claims for gene-specific and cell type-specific CRMs would benefit from further validation using complementary approaches, such as CRISPR interference or Prime editing.

The methods that we developed were meant to provide candidates for regulatory elements for a gene of interest. These candidates could be used to further understand the regulation of a gene, a complex and difficult task, especially for dynamically regulated genes in the context of development. These candidates could also, or instead, be used to drive gene expression specifically in a target cell of interest for applications such as gene therapy or perturbations that need this type of specificity. In the first case, to use the candidates to understand the regulation of a gene, one would need to validate the candidates using the types of methods typically employed for this purpose, most rigorously in the *in vivo* genomic context. We did not pursue this level of validation as it would encompass a great deal of work outside the scope of the current study. However, by initially testing loci which have been studied by several groups (as cited in the manuscript, *Rho*, *Grm6*, *Vsx2*, and *Cabp5*), we were able to show that LS-MPRA can identify known CRMs. In the cases of *Rho* and *Vsx2*, previous data have shown the CRMs to be relevant in the genomic context *in vivo*. In addition, two *Vsx2* CRM's identified by LS-MPRA are located at -37 Kb and -17Kb, and the *Grm6* CRM identified by LS-MPRA is at -8Kb. These are the same CRM locations identified previously using classical methods. These data show that the method is capable of identifying distal elements. When one has only one or a few loci of interest, i.e. one does not need to use genome-wide approaches, LS-MPRA is accurate enough to be worth the relatively small effort to identify potential CRMs, even those at some distance from the TSS. However, it is apparent that our methods are not perfect and that the LS-MPRA does not pick up all CRMs. We do not know of a method that has been shown to do so.

Reviewer #3 (Public review):

Summary:

Use of reporter assays to understand the regulatory mechanisms controlling gene expression moves beyond simple correlations of cis-regulatory sequence accessibility, evolutionary sequence conservation, and epigenetic status with gene expression, instead quantifying regulatory sequence activity for individual elements. Tulloch et al., provide a systematic characterization of two new reporter assay techniques (LS-MPRA and d-MPRA) to comprehensively identify cis-regulatory sequences contained within genomic loci of interest during retinal development. The authors then apply LS-MPRA and d-MPRA to identify putative cis-regulatory sequences controlling Olig2 and Ngn2 expression, including potential regulatory motifs that known retinal transcription factors may bind. Transcription factor binding to regulatory sequences is then assessed via CUT&RUN. The broader utility of the techniques is then highlighted by performing the assays across development, across species, and across tissues.

Strengths:

(1) The authors validate the reporter assays on retinal loci for which the regulatory sequences are known (Rho, Vsx2, Grm6, Cabp5) mostly confirming known regulatory sequence activity but highlighting either limitations of the current technology or discrepancies of previous reporter assays and known biology. The techniques are then applied to loci of interest (Olig2 and Ngn2) to better understand the regulatory sequences driving expression of these transcription factors across retinal development within subsets of retinal progenitor cells, identifying novel regulatory sequences through comprehensive profiling of the region.

(2) LS-MPRA provides broad coverage of loci of interest.

(3) d-MPRA identifies sequence features that are important for cis-regulatory sequence activity.

(4) The authors take into account transcript and protein stability when determining the correlation of putative enhancer sequence activity with target gene expression.

Weaknesses:

(1) In its current form, the many important controls that are standard for other MPRA experiments are not shown or not performed, limiting the interpretations of the utility of the techniques. This includes limited controls for basal-promoter activity, limited information about sequence saturation and reproducibility of individual fragments across different barcode sequences, limitations in cloning and assay delivery, and sequencing requirements. Additional quantitative metrics, including locus coverage and number of barcodes/fragments, would be beneficial throughout the manuscript.

We thank the reviewer for these comments and have provided detailed responses to the additional analyses in the subsequent Recommendations section.

(2) There are no statistical metrics for calling a region/sequence 'active'. This is especially important given that NR3 for Olig2 seems to have a small 'peak' and has non-significant activity in Figure 4.

See comments about peak calling in our response to Reviewer #1.

(3) The authors present correlational data for identified cis-regulatory sequences with target gene expression. Additionally, the significance of transcription factor binding to the putative regulatory sequences is not currently tested, only correlated based on previous single-cell RNA-sequencing data. While putative regulatory sequences with

potential mechanisms of regulation are identified/proposed, the lack of validation (and discrepancies with previous literature) makes it hard to decipher the utility of the techniques.

See comments about further validation in our response to Reviewer #2.

(4) While the interpretations that Olig2 mRNA/protein expression is dynamically regulated improved the proportions of cells that co-expressed CRM-regulated GFP and Olig2, alternate explanations (some noted) are just as likely. First, the electroporation isn't specific to Olig2+ progenitors. Also, the tested, short CRM fragments may have activating signals outside of Olig2 neurogenic cells because chromatin conformation, histone modifications, and DNA methylation are not present on plasmids to precisely control plasmid activity. Alternatively, repressive elements that control Olig2 expression are not contained in the reporter vectors.

The electroporation of Olig2 minus and plus cells is an excellent way to determine if a CRM is active in all cells, or only a specific subset, and we therefore consider this the best way to answer the question of specificity. We agree that we were unable to show that all CRM active cells were indeed Olig2-expressing cells. As noted by the Reviewer, we went to some lengths to quantify RNA and protein co-expression, including of endogenous Olig2 protein and RNA. Even with the endogenous RNA and protein, there was a mismatch wherein one infrequently saw the two together in the same cell, which could be predicted from the short half-lives of these molecules. Regarding chromatin, etc., we are intrigued by the proper regulation that we have observed for CRMs that we have previously discovered by plasmid electroporation (e.g. Kim et al. 2008, Matsuda and Cepko, 2004, Wang et al. 2014, Emerson et al. 2013). It is indeed interesting that plasmids can recapitulate proper regulation, without the proper genomic context or chromatin modifications. We have expanded our discussion of these points in the Discussion.

(5) It is unclear as to why the d-MPRA uses a different barcoding strategy, placing a second copy of the cis-regulatory sequence in the 3' UTR. As acknowledged by the author, this will change the transcript stability by changing the 3' UTR sequence. Because of this, comparisons of sequence activity between the LS-MPRA and d-MPRA should not be performed as the experiments are not equivalent.

We had provided a rationale for the different strategies of barcoding in the original submission, and believe it is at the discretion of the experimenter to utilize either strategy for their specific purposes. We agree that comparing activity between different techniques would not be appropriate. The analysis of mutated CRMs using d-MPRA does not utilize data from the LS-MPRA, but is an analysis of relative activity among all mutated d-MPRA constructs.

(6) Furthermore, details of the mutational burden in d-MPRA experiments are not provided, limiting the interpretations of these results.

We have provided detailed responses to the additional analyses in the subsequent Recommendations section and included details of the mutational burden in Supplemental Document A.

(7) Many figures are IGV screenshots that suffer from low resolution. Many figures could be consolidated.

We have increased the resolution of all IGV genome tracks, but believe the content within all figures remains appropriate.

Recommendations for the authors:

Reviewer #1 (Recommendations for the authors):

Suggestions for improving the clarity of the results in the figures:

(1) The pie charts used to show the percentage of overlapping cells in the colocalization analyses were not especially intuitive to read, and although the percentages and any statistical significance were often written in the text, it would've been helpful to have them written in the figures. I would suggest displaying the results in stacked bar plots, possibly like the one shown in Figure 6A, to demonstrate the data more clearly.

We thank the reviewer for the suggestions. Though adding the percentages directly to the pie charts would make the relevant panels too confusing to interpret, we added supplemental tables (Tables S5-S9) with the percentages displayed in all pie charts for readers interested in the precise quantifications.

(2) The scRNA-seq UMAPs showing co-expression of Olig2 with the TFS of interest - it is very hard to see the cells that co-express. I would recommend either having a window zoomed in on the Olig2-expressing cell population to be able to see the co-expression more clearly visually, and/or including a graph demonstrating the percentages of co-expressing cells. These numbers were written in the text, but would be useful to see in the figure.

The resolution of the scRNA-Seq plot has been improved for the visualization of co-expressing cells, which were also brought forward in all UMAP plots to improve clarity. Because of the higher quality images, insets should no longer be necessary. We have also included percentages of co-expression in the figures (Figs. 8 and 8S) and thank the reviewer for the suggestion.

Other minor suggestions/corrections:

(3) Figures 6B and 10S are missing the overlap quantification (in bar or pie charts) like in the other figures.

The quantification for the image in 6B (i.e., GFP fluorescence and GFP RNA) is displayed in 6D for the four Olig2 CRM plasmid constructs. In Fig. 10S, the experiments in early chick ventral neural tube delivered constructs to a very limited number of cells, and quantification of cells would not necessarily represent an accurate number of cells with CRM activity. We therefore decided to show only representative images of CRM activity in this population of cells rather than present a biased count or increase the number of experiments/samples to obtain a robust quantification.

(4) On the second-to-last line of page 10, in the sentence "The d-MPRA approach provided a robust, high resolution method for functionally relevant TF binding sites....", I think you're missing a word between "for" and "functionally". For example, it might be "for identifying..." or "for nominating..."

We have revised the sentence accordingly.

Reviewer #2 (Recommendations for the authors):

Minor suggestions:

(1) Please indicate which mouse reference genome (e.g., mm10) was used in plots such as Figure 2.

We have added text to the relevant sections in the Results (the reference genome was already mentioned in Methods).

(2) In Figures 2 and 2S, the CRMs discussed in the text are not labeled or highlighted, making it unclear which regions are being referenced.

We have labeled peaks with roman numerals in both the figures, legends, and text for clarity and thank the reviewer for the suggestion.

(3) Consider listing the genomic coordinates for the CRMs mentioned in the text, as this information would be especially useful for readers interested in exploring these regions further.

This information was included in Table 2S in the original submission, with all relevant coordinates provided therein.

(4) The d-MPRA plots (e.g., Figure 7C-E) do not clearly show the effects of different nucleotide substitutions. A more informative visualization style can be found in Kircher et al (PMID: 31395865, Fig. 1D) or Deng et al (PMID: 38781390, Fig. 5F).

The precise nucleotide substitutions would be informative to visualize the effects of specific changes. However, we were more interested in how any nucleotide substitution influenced the CRM activity to hone in on relevant TFBS. We therefore believe the current visualization is the most appropriate to accomplish this. However, for some types of future applications, a more informative visualization as noted would be a valuable addition.

(5) It would be extremely helpful to the community if the LS-MPRA data were uploaded to the UCSC genome browser and made accessible via a link.

We have uploaded all LS-MPRA genome tracks to a Track Hub in the UCSC genome browser and provided the appropriate link to access the Hub (<https://github.com/cattapre/ALAS00>) in the methods section.

Reviewer #3 (Recommendations for the authors):

(1) The authors should address the following metrics to showcase the utility of the techniques:

We thank the reviewer for requesting the detailed metrics outlined below. We have addressed all inquiries and included the majority of metrics in the resubmission.

(a) Library size

This should be shown for each library that is generated. It is acknowledged that the complete size of the library is limited by sequencing, and the comprehensiveness of the library will change every time the library is re-prepped. However, metrics of this are not currently provided in a robust manner for each library. "Libraries of at least 7×10^6 and as many as 9×10^7 fragments are made" - vague - how was library complexity established since this seems to be an estimation, how many reads were utilized to estimate library complexity?

We created a new supplemental table (Table S3) that displays the complexity based on sequencing rather than the estimated complexity based on the serial dilutions prior to 3D culture (which was used for the estimates listed in the results). We updated the complexity range in the text as well and thank the reviewer for the suggestion.

Does library size scale proportionally to the BACs of different sizes?

The fragmentation of different BACs with differing sizes does not necessarily alter the size of the library. Library size is primarily determined by the library creation pipeline, with the size selection step of the fragmented BAC and the cloning step that inserts adapter-ligated fragments into the barcoded expression vector being the primary determinants of complexity of plasmid libraries.

(b) Sequence saturation

Can the authors please provide evidence that the libraries have been sequenced to saturation or estimates of the degree of under-sequencing? How many reads does it take to discover a new barcode associated with a new regulatory sequence?

We have provided library characteristics for this in Table S3 and have also generated Sequence Saturation Curves for each association library in Supplemental Document A.

(c) Barcode saturation

How many barcodes are present for each fragment in the libraries? Are most fragments only covered by 1 barcode? The barcoding strategy doesn't prevent the same barcode from being assigned to multiple different fragments, as barcodes are random. What is the incidence of barcode collisions?

We have provided library characteristics for this in Table S3 and have also generated Barcode Saturation Curves for each association library in Supplemental Document A.

Additionally, we tested whether the omission of barcode collisions would affect the output of our LS-MPRA. We reanalyzed one barcode abundance library (one replicate following 12h Notch inhibitor) and filtered the barcodes so that only unique barcodes were analyzed. We were able to replicate all previously identified peaks. Though it is not necessary to filter out barcode collisions, there may be an improvement in signal-to-noise if the sequencing depth of libraries was sufficient (see Supplemental Document B).

(d) Normalization

As performed, fragment activity is normalized by RNA expression compared to the presence of fragments in the library. While this is done for small libraries, for large libraries, this may not be appropriate. For large libraries, every sequence in the library will not be delivered to each cell, and many fragments contained in the library may not be electroporated at all. Ideally, the authors would have sequenced both the RNA and DNA from the electroporations to i) identify the fragment distribution of the library that was successfully electroporated and ii) provide an internal normalization factor across replicate samples. This is especially important if the libraries were ever re-prepped, as the jack-potting or asymmetries in fragment recovery can occur every time the library is re-derived.

We agree with the reviewer's comments about the variability in fragments delivered experimentally, though we also believe the normalization of the libraries is still appropriate. We never needed to re-prepare the libraries as there was sufficient material for many more experiments than were performed. However, should one ever need to re-prepare an LS-MPRA library, all experimental sequencing should be normalized to the respective sequenced association library to account for biased distributions, as the reviewer mentions.

In the absence of these metrics (this would likely require the authors to repeat all experiments and is acknowledged to be outside the scope of revisions), the authors should provide information on the percentage of the library that is profiled in the RNA for each library.

We have provided RNA profiles of all abundance libraries in Table S4. The overall fraction of fragments represented in the RNA pools was lower than that observed in other published MPRA studies. This difference is expected given that most MPRA studies preselect fragments based on chromatin accessibility, transcription factor binding, sequence conservation, or bioinformatically predicted CRMs, thereby enriching for regulatory elements with high activity potential. Our locus-specific MPRA libraries, by contrast, include all fragments across

the targeted genomic region, many of which are likely to be inactive in the tested context. Consequently, only a smaller proportion of fragments show measurable RNA expression.

(e) Fragment sizes

Please provide a density plot or something similar showcasing the size distribution of the libraries generated. Is there any correlation between sequence activity and the size of fragments?

We have generated size distribution plots and correlations between fragment size and activity of all libraries and have included them in Supplemental Document A.

(2) Questions about the statistical validity of results:

(a) What threshold is utilized for calling a sequence as active? This is important as NR3 does not seem to be an element that has significant activity.

See comments about peak calling in prior responses.

(b) A Fisher's exact test using cells from single-cell RNA-sequencing as replicate samples is inappropriate as the cells are i) not from replicate experiments and ii) potentially in different cell states. The proportions of cells across replicate scRNA-seq datasets would be more appropriate.

We thank the reviewer for raising this important point. While we agree that individual cells do not substitute for biological replicates, we believe Fisher's exact test remains appropriate for testing whether gene expression is associated with Olig2 expression within a single scRNA-seq dataset. The test assesses co-occurrence at the level of individual cells, which is valid under the assumption that each cell represents an independent sampling of transcriptional states, even when it is possible that cells are in different states. We use this method as an exploratory tool to identify candidate genes associated with Olig2 expression in this dataset, and in the future, this could also be further validated by comparing the proportions of cells across replicate datasets, as the reviewer mentions.

(3) Discussion of the reporter/Olig2/Ngn2 RNA/protein disconnect needs to be expanded. Some simpler explanations for the presence of GFP in Olig2- and Ngn2- cells, as well as the presence of Olig2 or Ngn2 in GFP- cells, is that (i) these putative CRMs are being introduced to cells in plasmids, taking them out of their native genomic context where they may be inaccessible or repressed and allowing them to drive reporter expression even if their candidate target gene is not endogenously expressed, (ii) these putative CRMs may regulate genes besides just Olig2 or Ngn2, and (iii) Olig2 and Ngn2 are regulated by far more regulatory elements than the 3 or 4 being tested in each reporter assay, so their expression likely does not rely solely on the activity of the few putative CRMs tested.

We have added these points in an expanded discussion in the text.

(4) Problems with figures: Low resolution of many IGV genome tracks, pink 'co-expression' dots are completely indiscernible. Numbers should be listed with the pie charts. BFP expression should be shown since this is being quantified, especially since electroporation efficiency can change across age and/or tissue samples.

We have reconfigured the IGV tracks so that they are higher resolution and have included supplemental tables for the numbers pertaining to the pie charts. For electroporation controls (BFP and RFP), BFP expression is shown in Figs 5S, 6, and 10S and the RFP electroporation control is shown in Fig. 11. Though BFP is sometimes used as a qualifier in the denominator of some of the quantification, displaying its expression, particularly in

combination with three other signals that are already included in most images, provides limited utility.

(5) More information is required to understand the utility of the d-MPRA. Detailed quantification of the number of mutations/fragments needs to be ascertained. When multiple mutations are present, how are the authors controlling for which mutation is affecting activity? What is the coverage of the loci of interest for mutational burden (ie, is every base pair mutated in at least one fragment?). For mutations that increase the activity of the element, are there specific sequence features that increase activity (new motifs generated)?

The d-MPRA platform is a high-throughput assay that seeks to identify putative sub-regions within CRMs nominated by the LS-MPRA, or any other assay. It relies on deep mutational coverage to determine positive and negative regulatory sub-regions of the CRMs. While many reads have multiple mutations, they are broadly co-occurring across the entire fragment (see Supplemental Document A) so as not to create a false linkage between the sites. Every individual site is mutated many times with roughly even coverage across each fragment (see Supplemental Document A), thus allowing us to assess the requirement of each base in contributing to a putative CRM's activity. Comparing d-MPRA plots using bulk fragments or fragments with singleton mutations (Supplemental Document A) yielded almost identical plots for two libraries, and a similar analysis of the third library. Any differences between analysis of fragments with one or more mutations is likely a result of either sequencing depth or the requirement of multiple bases for binding or CRM activation. Follow-up experiments investigating intra-CRM interactions would elucidate such variability. Whether new motifs are generated for any specific substitution is an interesting question, which could be followed up for a CRM of interest. The d-MPRA data that we provide would provide the starting point for such follow-up experiments.

(6) Transcription factors as regulators of CRM-activity.

It is appreciated that the authors validated the binding of transcription factors to NR2. However, this correlative analysis should be further tested in follow-up experiments to highlight novel biology using systems already in place. Potential experiments that could be performed include the following (reagents in hand, or performed in a manner similar to experiments performed by the lab in previous publications):

(a) over-expression of TF using LS-MPRA library.

(b) over-expression of TF using d-MPRA library, showing that mutations in the putative TF binding site disrupt activity compared to non-mutated sequences.

(c) performing TF over-expression using target CRMs, including sequences where the TF binding site is mutated (similar to a small MPRA).

(d) the quantification of target gene expression when i) TF is over-expressed, ii) CRM is activated using CRISPRa, or iii) CRM is inhibited using CRISPRi.

These are all valid follow-up experiments. Please see prior responses we have provided regarding further validation.

Minor points

(1) Please acknowledge that some distal regulatory sequences may be contained outside of the BAC regions. Also, the authors should emphasize the point that the assay is NOT cell-type-specific or specific to regulatory sequences for the gene of interest, but ALL regulatory sequences contained within the locus. The discussion of this with respect to Ift122 and Rpl32 is somewhat confusing.

We have added a sentence in the Discussion addressing possible CRMs outside the BAC coverage. We believe it is implicitly understood that the assay only screens regulatory activity in the BAC, and believe we have addressed this in the manuscript.

If one wishes to use a candidate CRM to drive gene expression in a targeted cell type, one needs to establish specificity. In particular, specificity needs to be established in the context of the vector that is being used. Non-integrated vs integrated vectors, different types of viral vectors with their own confounding regulatory sequences, different types of plasmids and methods of delivery, and copy number can all affect specificity. We provided a double in situ hybridization method for the examination of specificity for some of the novel candidate CRMs. It was quite difficult in the case of Olig2 and Ngn2 as their RNAs and proteins are unstable. We would need to provide further evidence should we wish to use these candidate CRMs for directing expression specifically in Olig2- or Ngn2-expressing cells. We suggest that an investigator can choose the vector and method for establishing specificity depending upon the goals of the application.

(2) I am curious as to why low-resolution, pseudo-bulked single-nucleus ATAC was utilized instead of more comprehensive retina ATAC samples at similar time-points (for example, as available in Al Diri et al., 2017 (E14, E17, P0, P3, P7, P10) samples are all available.

The use of pseudo-bulked single-nucleus ATAC-seq data provided a convenient and consistent comparison to our LS-MPRA results. We agree that incorporating higher-resolution datasets such as those from Al Diri et al. would be valuable for future analyses aimed at linking CRM activity with broader chromatin accessibility dynamics.

<https://doi.org/10.7554/eLife.107565.2.sa0>