

Acidocalcisome-like vacuoles constitute a feedback-controlled phosphate buffering system for the cytosol

Reviewed Preprint

v2 • November 11, 2025

Revised by authors

Reviewed Preprint

v1 • August 22, 2025

Samuel Bru, Lydie Michailat Mayer, Geun-Don Kim, Danye Qiu, Henning Jacob Jessen, Andreas Mayer 

Département d'immunobiologie, Université de Lausanne, Epalinges, Switzerland • Institute of Organic Chemistry, University of Freiburg, Freiburg, Germany • CIBSS – Centre for Integrative Biological Signalling Studies, University of Freiburg, Freiburg, Germany

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

eLife Assessment

This study provides **fundamental** insights into eukaryotic phosphate homeostasis by demonstrating how yeast vacuoles dynamically regulate cytosolic phosphate levels. The conclusions are **convincing**, supported by an elegant combination of in vitro assays and in vivo measurements. This study will be of interest to cell biologists, particularly for those who are working in the field of phosphate metabolism.

<https://doi.org/10.7554/eLife.108181.2.sa3>

Abstract

Cells experience strong variations in the consumption and availability of inorganic phosphate (P_i). Since P_i is an essential macronutrient but excess P_i has negative impacts on nucleotide hydrolysis and metabolism, its concentration must be maintained in a suitable range. Conserved storage organelles, acidocalcisomes, provide this buffering function. We used acidocalcisome-like yeast vacuoles to study how such organelles are set up to for this task. Our combined in vitro and in vivo analyses revealed that their ATP-driven polyphosphate polymerase VTC converts cytosolic P_i into inorganic polyphosphates (polyP), which it transfers into the vacuole lumen. Luminal polyphosphatases immediately hydrolyse this polyP to establish a growing reservoir of vacuolar P_i . Product inhibition by this P_i pool silences the polyphosphatases, caps P_i accumulation, and favours vacuolar polyP storage. Upon cytosolic P_i scarcity, the declining inositol pyrophosphate levels activate the vacuolar P_i exporter Pho91 to replenish cytosolic P_i . In this way, acidocalcisome-like vacuoles constitute a feedback-regulated buffering system for cytosolic P_i , which the cells can switch between P_i accumulation, P_i release, and high-capacity phosphate storage through polyP.

Introduction

Cells actively manage the concentration of P_i in their cytosol because they must strike a balance between conflicting goals (Austin & Mayer, 2020 [\[1\]](#)). On the one hand, P_i is a product of nucleotide-hydrolysing reactions. Its concentration has a significant impact on the free energy that these reactions can provide to drive metabolism. Since, for this reason, excessive P_i concentrations might stall metabolism, we can expect that cytosolic P_i must remain limited. But P_i is also an essential macronutrient. As a major constituent of nucleic acids and phospholipids, and as an important modifier of proteins, carbohydrates and many metabolites, it is consumed in large quantities for anabolic reactions. This can lead to situations where progression of the S-phase of the cell cycle is limited by the P_i uptake capacity of the cells (Bru *et al.*, 2016 [\[2\]](#), 2017 [\[3\]](#); Gillies *et al.*, 1981 [\[4\]](#); Neef & Kladde, 2003 [\[5\]](#)). Rapidly dividing cells, such as yeast cells, remedy this problem by maintaining phosphate stores in the form of inorganic polyP. PolyPs are chains of P_i linked through phosphoric anhydride bonds, which can be stored in acidocalcisome-like organelles (Urech *et al.*, 1978 [\[6\]](#); Okorokov *et al.*, 1980 [\[7\]](#)). Acidocalcisomes are lysosome-related organelles that have been characterised by pioneering work mainly in Trypanosomes, but they have been isolated from a multitude of other organisms (Docampo *et al.*, 2005 [\[8\]](#)). Their defining chemical features are their acidity, high concentrations of polyP, Ca, Mg, Zn, Fe, K, Na, and basic amino acids and they are implicated in osmoregulation and the homeostasis of phosphate and metals (Docampo *et al.*, 2005 [\[8\]](#); Girard-Dias *et al.*, 2023 [\[9\]](#); Docampo, 2024 [\[10\]](#)). Yeast vacuoles share all these properties and are hence good and easily accessible models for studying aspects of acidocalcisomal mechanisms, such as the synthesis and turnover of polyP (Dürr *et al.*, 1979 [\[11\]](#); Huber-Wälichli & Wiemken, 1979 [\[12\]](#); Urech *et al.*, 1978 [\[6\]](#); Austin & Mayer, 2020 [\[1\]](#)). But they have an additional, lysosome-like face, enabling them to assume also hydrolytic degradative functions for the cells (Li & Kane, 2009 [\[13\]](#)). Therefore, we refer to them as acidocalcisome-like organelles.

The polyP stores of vacuoles are accessed by cells when P_i becomes limiting in the environment, or when they face a sudden enhanced P_i consumption, e.g. during metabolic transitions from respiratory growth to fermentation, which requires much more phosphorylated metabolites (Shirahama *et al.*, 1996 [\[14\]](#); Thomas & O'Shea, 2005 [\[15\]](#); Gillies *et al.*, 1981 [\[4\]](#); Nicolay *et al.*, 1982 [\[16\]](#), 1983 [\[17\]](#)). In such cases, cellular polyphosphate content decreases. It has therefore been proposed that polyPs are re-converted into P_i to buffer shortages in cytosolic P_i (Nicolay *et al.*, 1983 [\[16\]](#), 1982 [\[17\]](#)). While this hypothesis is straightforward, we are lacking a coherent concept of how a P_i buffer based on an acidocalcisome-like organelle might work. Pioneering work on these organelles has revealed several characteristic and conserved features, such as their acidity, and their high content of basic amino acids and divalent cations (Docampo & Huang, 2016 [\[18\]](#); Docampo, 2024 [\[10\]](#)). Of direct relevance to P_i homeostasis is the high capacity of acidocalcisome-like organelles for storing polyP. PolyP can greatly vary in length, from two to hundreds of phosphate units. The membrane of acidocalcisome-like organelles can carry polyP polymerases, such as VTC, and P_i transporters, such as Pho91 (Jimenez & Docampo, 2015 [\[19\]](#); Hürlimann *et al.*, 2007 [\[20\]](#); Wang *et al.*, 2015 [\[21\]](#); Huang & Docampo, 2015 [\[22\]](#); Gerasimaite & Mayer, 2016 [\[23\]](#); Müller *et al.*, 2003 [\[24\]](#), 2002 [\[25\]](#)). VTC is a coupled polyP polymerase and translocase (Gerasimaite *et al.*, 2014 [\[26\]](#)), which synthesizes polyP by transferring the ψ -phosphate of cytosolic ATP onto an elongating polyP chain (Hothorn *et al.*, 2009 [\[27\]](#)). PolyP is generated by a catalytic domain in the centre of the Vtc4 subunit of the complex (Hothorn *et al.*, 2009 [\[27\]](#)). The activity of this subunit is controlled through the SPX domains of VTC (Wild *et al.*, 2016 [\[28\]](#)), which may associate into a dimerized inactive state (Pipercevic *et al.*, 2023 [\[29\]](#)). They can be released from this state through the action of inositol pyrophosphates, signalling molecules that accumulate when P_i is abundant in the cytosol and activate P_i storage in the form of polyP. The P_i state of the cytosol is communicated to VTC through a specific inositol pyrophosphate, 1,5-IP₈ (Gerasimaite *et al.*, 2017 [\[30\]](#); Chabert *et al.*, 2023 [\[31\]](#)).

The postulated direct channelling of polyP from the site of its synthesis through the vacuolar membrane (Gerasimaite *et al*, 2014 [\[1\]](#)) is facilitated by the structure of the VTC complex. The catalytic domain of VTC has its exit for polyP right at the entry of a polyP-conducting channel, which is formed through the transmembrane domains of the VTC complex itself (Liu *et al*, 2023 [\[2\]](#); Guan *et al*, 2023 [\[3\]](#); Müller *et al*, 2002 [\[4\]](#), 2003 [\[5\]](#); Gerasimaite *et al*, 2014 [\[1\]](#)). This channel was proposed to exist in an open and closed conformation and to be gated through polyP (Liu *et al*, 2023 [\[2\]](#)). PolyP translocation may be driven by the electrochemical potential across the membrane, which could move the highly negatively charged polyP chain through electrophoresis. This can explain why efficient polyP synthesis depends on proton pumps such as the V-ATPase or H⁺-pumping pyrophosphatases (Gerasimaite & Mayer, 2016 [\[6\]](#); Gerasimaite *et al*, 2014 [\[1\]](#); Freimoser *et al*, 2006 [\[7\]](#); Lemerrier *et al*, 2004 [\[8\]](#)).

The transporter Pho91 was proposed to export vacuolar P_i (Hürlimann *et al*, 2007 [\[9\]](#)). Several properties of yeast Pho91 and its homologs from other organisms are consistent with this view. Its ablation increases vacuolar P_i and polyP content, in yeast as well as in trypanosomes (Hürlimann *et al*, 2007 [\[9\]](#); Jimenez & Docampo, 2015 [\[10\]](#); Farofonova *et al*, 2023 [\[11\]](#)), and it weakly induces the phosphate starvation program in yeast (Pinson *et al*, 2004 [\[12\]](#)), suggesting that it may induce cytosolic P_i scarcity. Furthermore, patch-clamp analyses of Pho91-like channels from yeast, trypanosomes and plants have shown their P_i permeability and the dependence of their directionality on a pH gradient across the membrane (Potapenko *et al*, 2018 [\[13\]](#), 2019 [\[14\]](#); Wang *et al*, 2015 [\[15\]](#)). In the presence of a pH gradient, the Pho91 homolog from rice, OsSpx-MFS3, mediates P_i flux along the proton gradient, consistent with a function in exporting P_i from the acidic lumen of vacuoles (Wang *et al*, 2015 [\[15\]](#)) towards the cytosol.

Acidocalcisome-like organelles also contain polyphosphatases in their lumen (Lander *et al*, 2016 [\[16\]](#); Gerasimaite & Mayer, 2016 [\[6\]](#); Kulakovskaya *et al*, 2021 [\[17\]](#); McCarthy & Downey, 2023 [\[18\]](#)), which can convert polyP back into P_i. In baker's yeast, two vacuolar polyphosphatases are known, Ppn1 and Ppn2 (Gerasimaite & Mayer, 2017 [\[19\]](#); Sethuraman *et al*, 2001 [\[20\]](#); Andreeva *et al*, 2019 [\[21\]](#)). This means that a chain of polyP, when being synthesized by VTC and arriving in the vacuolar lumen, is immediately exposed to hydrolytic enzymes that will degrade it. While this seems at first sight paradoxical, we explored the hypothesis that the co-existence of polyP-synthesizing and polyP-hydrolysing activities might be a key feature conveying to acidocalcisome-like organelles the capacity to buffer cytosolic P_i. That these organelles have a critical role to play in this process is illustrated by observations in yeast, where artificial up- or down-regulation of vacuolar polyP synthesis suffices to drive the cytosol into a state of P_i starvation or P_i excess, respectively (Desfougères *et al*, 2016a [\[22\]](#)). Furthermore, the presence of polyP reserves delays the activation of the transcriptional phosphate starvation response, the PHO pathway (Thomas & O'Shea, 2005 [\[23\]](#)). We hence explored the capacity of isolated yeast vacuoles to interconvert polyP and P_i, and we characterized the roles played by the vacuolar polyphosphatases Ppn1 and Ppn2 and the vacuolar P_i transporter Pho91. Our observations can be combined with previous findings to yield a coherent model of how an acidocalcisome-like organelle can operate as a P_i buffer for the cytosol.

Results

We explored the interplay of VTC, polyphosphatases and Pho91 in the accumulation of polyP and P_i inside vacuoles using an in vitro system with purified organelles. Vacuoles can be isolated in intact form when the cells are gently opened by enzymatic digestion of the cell wall and disruption of the cell membrane by low concentrations of DEAE-dextran (Dürr *et al*, 1975 [\[24\]](#)). Organelles isolated in this way can perform many of their normal cellular functions, such as membrane fusion, membrane fission and autophagy (Michaillat *et al*, 2012 [\[25\]](#); Sattler & Mayer, 2000 [\[26\]](#); D'Agostino & Mayer, 2019 [\[27\]](#); Kunz *et al*, 2004 [\[28\]](#)). They also contain active polyP polymerase (VTC) and active polyphosphatases (Ppn1 and Ppn2) and they can synthesize and import polyP (Gerasimaite *et al*, 2014 [\[1\]](#); Gerasimaite & Mayer, 2017 [\[19\]](#); Gerasimaite *et al*, 2017 [\[29\]](#)).

Vacuolar P_i accumulation depends on polyP synthesis

Purified vacuoles were incubated with an ATP-regenerating system under conditions that allow these organelles to synthesize polyP *in vitro* (Gerasimaite *et al.*, 2014). After different periods of incubation, the organelles were sedimented and solubilized in detergent. Vacuolar accumulation of polyP was assayed through DAPI fluorescence and P_i was measured through malachite green. VTC is stimulated by a variety of inositol pyrophosphates, most efficiently by 1,5-InsP₈ (Gerasimaite *et al.*, 2014, 2017). Since 1,5-InsP₈ is not commercially available, we used saturating concentrations of 5-InsP₇ for our experiments (Pavlovic *et al.*, 2015). This compound stimulates VTC with a higher EC₅₀ but to the same maximal activity as 1,5-InsP₈, and it was more readily available to us for routine experiments. Under the conditions used here, the organelles rapidly and efficiently produced polyP (Fig. 1), in line with previous results (Gerasimaite *et al.*, 2014, 2017). This production was stimulated by 5-InsP₇, reaching 0.7 nmol of phosphate units per μ g of vacuolar protein and min. This means that already within 10 min, wildtype vacuoles produced a mass of polyP equivalent to their total protein content, indicating how efficiently these organelles synthesize polyP. This signal was entirely dependent on VTC, because it was not observed in a mutant lacking the catalytic subunit of VTC (*vtc4Δ*).

Isolated wildtype vacuoles also accumulated P_i with significant efficiency, at an initial rate of at least 0.3 nmol μ g⁻¹ min⁻¹ (Fig. 2A). Vacuoles from *vtc4Δ* cells showed only 8% of this P_i signal. Most of it was not time-dependent and hence may represent a background signal from the organelle preparation. In sum, vacuolar P_i accumulation depended on polyP synthesis through VTC.

The kinetics of polyP and P_i accumulation were remarkably different. Whereas polyP continued to accumulate throughout the entire incubation period, P_i accumulated rapidly in the initial phase but reached a plateau within 30 min (Fig. 2A, B). Thus, although polyP production was essential for P_i accumulation, it could not become the limiting factor for it in the second phase. This is further illustrated by P_i accumulation in strains showing enhanced polyP synthesis, such as overexpressors of VTC5 (Fig. 2A, B) or strains carrying hyperactivating substitutions in the SPX domains of VTC (*vtc3^{K126A} vtc4^{K129A}*, Fig. 2 C, D) (Desfougères *et al.*, 2016b; Wild *et al.*, 2016).

Vacuoles from these strains accumulated polyP at a two to seven times higher initial rate than the wildtype and to higher concentrations. Nevertheless, their accumulation of P_i arrested at the same level as that of wildtype vacuoles, at 5 nmol μ g⁻¹ vacuolar protein. To estimate the corresponding luminal concentration of P_i , we measured the diameters of 100 isolated wildtype vacuoles and counted the number of vacuoles per μ g of vacuolar protein. With an average diameter of 0.8 μ m (Suppl. Fig. 1) and 5×10^7 vacuoles/ μ g of protein we can estimate that P_i accumulation in vacuoles incubated with 5-InsP₇ reached a plateau at a luminal concentration of around 30 mM. *vtc4Δ* vacuoles showed only a very small increase of less than 3 mM in luminal P_i content.

The 30 mM of P_i accumulating in wildtype vacuoles is a substantial concentration, which is in the range of the *in vivo* P_i concentration of 25 mM that was measured by ³¹P-NMR spectroscopy of yeast cells under conditions where vacuolar P_i dominates the signal (Okorokov *et al.*, 1980). Since, upon P_i scarcity, cytosolic P_i drops to 1 mM (Okorokov *et al.*, 1980), a substantial P_i concentration gradient across the vacuolar membrane could drive rapid replenishment of the cytosol from a readily accessible vacuolar pool of P_i .




Fig. 1

VTC- and 5-IP₇-dependent polyP synthesis by isolated vacuoles

Vacuoles were isolated from logarithmic cultures of BY4742 wildtype cells (WT) or from isogenic *vtc4Δ* cells strains. They were incubated in polyphosphate synthesis assays without (A) or (B) in presence of 50 μM 5-InsP₇. At indicated times, aliquots were withdrawn, solubilized in Triton X-100 and polyphosphate was quantified through the polyP-dependent fluorescence of added DAPI. Means ± SD (standard deviation) of three independent experiments are shown. * $p < 0.05$, ** $p < 0.005$ for a paired t-test comparing WT with and without 5-InsP₇ at each time point.




Figure 2.

VTC-dependent accumulation of P_i in isolated vacuoles

Vacuoles were isolated from the indicated, logarithmically growing BY4742 (WT) cells. PolyP overproduction was achieved either through **(A, B)** overexpressing VTC5 from the strong GPD promotor, or **(C,D)** by expressing the hyperactivating *vtc3^{K126A}* and *vtc4^{K129A}* alleles from their native promoters as the sole source of these two proteins. The vacuoles were incubated with an ATP-regenerating system and 50 μM 5-InsP₇ under conditions allowing polyP synthesis. At indicated times, the vacuoles were solubilized with Triton X-100 and P_i was assayed through malachite green. Graphs represent the mean ± SEM of three independent experiments for each strain. * p<0.05, ** p<0.005, *** p<0.001 from a paired t-test comparing each time WT vs VTC5 overexpression or *vtc3^{K126A} vtc4^{K129A}*.

Vacuolar P_i accumulation is limited by feedback inhibition of Ppn1 and Ppn2

We asked why vacuolar P_i accumulation quickly forms a plateau whereas vacuolar polyP continues to accumulate. To this end, we explored the hypothesis that vacuolar polyphosphatases become inhibited when their product, P_i , has accumulated. To test polyphosphatase activity we liberated the polyphosphatases from isolated vacuoles by detergent lysis, followed by incubation with synthetic polyP as a substrate and DAPI as an indicator for polyP. PolyP was efficiently degraded by this extract. The apparent polyPase activity was significantly reduced by 3 mM P_i and efficiently silenced by 30 mM P_i (Fig. 3A). To differentiate the contributions of Ppn1 and Ppn2, we also analysed the respective deletion mutants and performed the assay at higher polyP concentration in the presence Mg^{2+} , which favours Ppn1 activity, or in the presence of Zn^{2+} , which supports activity of Ppn1 and Ppn2 (Gerasimaite & Mayer, 2017). In the presence of Zn^{2+} , the substrate was consumed in less than 3 min (Fig. 3B). Degradation was delayed in vacuoles from *ppn1Δ* or *ppn2Δ* mutants, and it was suppressed in vacuoles from a *ppn1Δ ppn2Δ* mutant, in which both polyphosphatases were ablated. In incubations with only Mg^{2+} instead of Zn^{2+} , which stimulates the activity of Ppn1 much more than that of Ppn2, polyP degradation was slower and genetic ablation of Ppn1 sufficed to stabilize polyP. The addition of 30 mM potassium phosphate, which is equivalent to the maximal P_i accumulation in vacuoles that we observed in Fig. 2, attenuated degradation of polyP 5-to 10-fold, both through Ppn1 (assayed in *ppn2Δ* and in samples without Zn^{2+}) and through Ppn2 (assayed in *ppn1Δ*). That both polyphosphatases are inhibited by P_i at this concentration is consistent with the notion that product inhibition of Ppn1 and Ppn2 might limit the conversion of polyP into P_i in the vacuolar lumen and thus define the maximal concentration of the vacuolar P_i reservoir.

Pho91 limits vacuolar P_i accumulation in an inositol pyrophosphate-dependent manner

To allow vacuoles to function as a P_i buffer for the cytosol, the P_i in the vacuolar lumen should become accessible in a regulated manner. The P_i transporter Pho91 is a prime candidate for mediating regulatable efflux because it is regulated by InsPPs through its SPX domain (Potapenko *et al*, 2018, 2019; Wang *et al*, 2015; Hürlimann *et al*, 2007). Furthermore, an overexpressed GFP fusion was reported to localize to vacuoles (Hürlimann *et al*, 2007; Elbaz-Alon *et al*, 2014), a phenotype that we confirmed (not shown). Since overexpression of membrane proteins in yeast easily leads to their erroneous accumulation in vacuoles, we re-investigated Pho91 localization in the absence of a strong, overexpressing promotor. When we expressed C- or N-terminal Pho91-GFP fusions from the authentic PHO91 promotor, we observed different and complex localization patterns (see Supplementary Figure 2 for examples).

These localization patterns depended on the nature of the fluorescent tag and the linker peptides used to attach it to Pho91, and they often showed significant accumulation in the ER. This suggests that the fluorescent protein tags interfere with intracellular trafficking of Pho91. We hence analysed purified vacuoles to test whether the non-tagged, endogenous Pho91 is a vacuolar protein. Mass spectrometry was used to determine the enrichment of proteins in the vacuolar fraction relative to a whole cell extract (Suppl. Table 1). Peptides from Pho91 were enriched in the purified vacuoles to a similar degree (39-fold) as peptides from vacuolar marker proteins, such as the vacuolar polyP polymerase subunit Vtc3 (35-fold), the vacuolar amino acid transporter Avt3 (48-fold), the v-ATPase subunit Vma9 (46-fold), or the alkaline phosphatase Pho8 (34-fold). By contrast, typical plasma membrane proteins were barely enriched, such as the iron permease Ftr1 (2.5-fold), the P_i importer Pho87 (2.4-fold) or the polyamine importer Tpo5 (2.2-fold) (Supplementary data file 1). This co-enrichment suggests that the major fraction of non-tagged Pho91 indeed resides in the vacuole. In agreement with this, Pho91 engages the AP3-dependent




Figure 3.

Effect of P_i on polyphosphatase activity.

A. P_i titration. Vacuoles were isolated from logarithmically growing wildtype cells (BY4742). The organelles were diluted in polyphosphatase assay reaction buffer, which contained 0.1% Triton X-100 and hence liberated the luminal polyphosphatases. This lysate was incubated with 30 μ M polyP₃₀₀ as a substrate and supplemented with 1 mM $ZnCl_2$ and the indicated concentrations of K- P_i pH 6.8. After the indicated times of incubation, the remaining polyP was quantified through DAPI. The DAPI signal at the beginning of the incubation served as 100% reference. Graphs represent the means $\pm$ SEM of three independent experiments. **B, C.** Differentiation of P_i effects on Ppn1 and Ppn2. Vacuoles were isolated from the indicated, logarithmically growing strains, lysed and used in polyP degradation assays with 300 μ M polyP₃₀₀ as in (A). These assays were performed in the presence or absence of 30 mM K- P_i pH 6.8 and 1 mM $MgCl_2$ (B) or, instead of this, with 1 mM $ZnCl_2$ (C) as cation supporting catalytic activity.

vesicular trafficking pathway, which leads from the Golgi to the vacuole (Eising *et al*, 2022 [↗](#)). In contrast to the other four yeast Pi transporters, Pho84, Pho87, Pho89, and Pho90, which individually suffice to support vigorous growth of yeast, Pho91 was reported to do support slower but significant growth and to fully rescue upon overexpression (Wykoff & O'Shea, 2001 [↗](#); Wykoff *et al*, 2007 [↗](#)). Since this capacity to feed the cells is difficult to reconcile with a vacuolar localisation, unless the cells would feed through fluid phase endocytosis and subsequent export from the vacuole (Klompaker *et al*, 2017 [↗](#)), we reinvestigated this aspect. Instead of relying a down-regulatable PHO84 background, which results in a low but constitutive background expression of the potent Pi importer Pho84 in addition to Pho91, we relied on a more stringent approach, using plasmid shuffling to completely remove PHO89 when PHO91 is brought in. To this end, we generated a quintuple knockout strain lacking these five Pi transporters (Pho84, Pho87, Pho89, Pho90, Pho91), which was kept alive through expression of Pho89 from a URA-based plasmid. Plasmid shuffling allowed to exchange this plasmid against others expressing an individual Pi transporter. Whereas an individual plasma-membrane Pi transporter such as Pho89 supported normal colony formation, confirming previous observations (Wykoff *et al*, 2007 [↗](#); Wykoff & O'Shea, 2001 [↗](#)), we could not confirm that Pho91 supports slower growth. In our plasmid shuffling approach, Pho91 did not support growth, at all (Suppl. Fig. 3 [↗](#)). This supports that Pho91 is indeed a vacuolar Pi transporter, but that C- or N-terminal protein fusions interfere with correct sorting and cannot serve as reliable reporters for localization of this protein.

We tested the impact of Pho91 on Pi accumulation by vacuoles *in vitro*, using the same approach as above (Fig. 4 [↗](#)). Vacuoles lacking Pho91 (*pho91Δ*) accumulated Pi two times faster than the wildtype and the maximal accumulated concentration was two times higher. This is consistent with a function of Pho91 as vacuolar Pi exporter (Hürlimann *et al*, 2007 [↗](#); Potapenko *et al*, 2018 [↗](#); Wang *et al*, 2015 [↗](#)).

The difference between *pho91Δ* and wildtype vacuoles vanished when the vacuoles were incubated in the presence of the inositol pyrophosphate 5-InsP₇. Since, as we showed above, vacuolar Pi accumulation depends on polyP synthesis through VTC, the relative enhancement of Pi accumulation in wildtype vacuoles through 5-InsP₇ could be caused by downregulation of Pho91, which would make the wildtype vacuoles behave similarly as *pho91Δ* vacuoles. Alternatively, 5-InsP₇ might stimulate polyP synthesis in wildtype more than in *pho91Δ* vacuoles. We could rule out the latter explanation based on two observations: An assay of polyP synthesis during the incubation (Fig. 5 [↗](#)) revealed that 5-InsP₇ stimulated polyP synthesis in wildtype and *pho91Δ* vacuoles to similar degrees. Furthermore, as shown above (Fig. 2 [↗](#)), polyP synthesis activity in wildtype cells is not rate-limiting for vacuolar Pi accumulation. Therefore, we attribute the enhancement of Pi accumulation through 5-InsP₇ to Pho91. Upon Pi scarcity, the declining inositol pyrophosphate levels should then activate Pho91 to replenish cytosolic Pi.

Interference with vacuolar polyP turnover provokes cytosolic Pi scarcity

We tested the effects of this postulated cycle of polyP synthesis, polyphosphatase activity and Pho91-mediated Pi export on Pi homeostasis *in vivo*. Pho4-GFP was used as a reporter, because this transcription factor shuttles between nucleus and cytosol. Under cellular Pi scarcity and correspondingly low inositol pyrophosphate levels it is predominantly nuclear, but it shifts to the cytosol under Pi-replete conditions (O'Neill *et al*, 1996 [↗](#); Chabert *et al*, 2023 [↗](#); Desfougères *et al*, 2016a [↗](#); Auesukaree *et al*, 2004 [↗](#)). Pho4-GFP localisation can hence serve as a readout for cytosolic Pi signalling. In wildtype cultures growing logarithmically, cells can experience a transient shortage of Pi during S-phase, when Pi utilization for biosynthesis may exceed the uptake capacity of the cell and consume its polyP stores (Bru *et al*, 2016 [↗](#); Neef & Kladde, 2003 [↗](#); Pondugula *et al*, 2009 [↗](#)). Therefore, even on Pi-rich media, some cells can show nuclear accumulation of Pho4-GFP (Wykoff *et al*, 2007 [↗](#); Vardi *et al*, 2013 [↗](#), 2014 [↗](#)). Since S-phase

Fig. 4

Accumulation of P_i in isolated vacuoles

VTC- and Pho91-dependence. Vacuoles were isolated from the indicated, logarithmically growing strains. The purified organelles were incubated as in [Fig. 1](#), i.e. in a buffer with an ATP-regenerating system that allows the synthesis of polyP, and either (A) without (w/o) or (B) in the presence of 50 μ M 5-IP₇. After the indicated periods of incubation at 27°C, an 80 μ l aliquot was withdrawn, the vacuoles were sedimented by centrifugation, washed and lysed. Released vacuolar P_i was determined by malachite green assay. Graphs represent the mean $\pm$ SEM of three independent experiments. $p < 0.05$ from a paired t-test comparing each time WT versus *pho91* Δ .





Fig. 5

PolyP accumulation in *pho91* mutant vacuoles.

Vacuoles from BY4742 cells (WT) and isogenic *pho91* Δ cells were isolated and incubated under conditions supporting polyP synthesis and P_i accumulation as in [Fig. 4](#), in the absence (w/o) or presence of 50 μ M 5-InsP₇. At the indicated timepoints, aliquots were withdrawn, the vacuoles were lysed in detergent, and polyP was assayed through DAPI fluorescence. Graphs show the means and SEM from three independent experiments. $p > 0.05$ from a paired t-test for all differences between WT and *pho91* Δ .



occupies only around 20% of the yeast cycle (Koren *et al*, 2010 [DOI](#); Donaldson *et al*, 1998 [DOI](#)), this fraction is expected to be correspondingly small. However, it is significant, and it can be clearly quantified by microscopy, which scores individual cells and not the population as an ensemble. We exploited this natural fluctuation of intracellular P_i availability as a sensitising situation to assay how the vacuolar systems influence P_i balance in the cytosol.

We scored the fraction of logarithmically growing cells that show nuclear Pho4-GFP even in P_i -replete medium. This fraction was 7% for wildtype cells (Fig. 6 [DOI](#)). It increased fourfold upon hyperactivation of polyP synthesis, which we achieved by overexpression of the regulatory VTC subunit Vtc5 (Desfougères *et al*, 2016a [DOI](#)). Ablation of polyP synthesis by deletion of the catalytic subunit Vtc4 had the opposite effect and reduced the frequency of cells with nuclear Pho4-GFP by half. Deletion of the vacuolar polyphosphatases Ppn1 and Ppn2 prevents polyP turnover and leads to the accumulation of extremely long polyP chains (Gerasimaite & Mayer, 2017 [DOI](#); Sethuraman *et al*, 2001 [DOI](#)). We may thus expect phosphate to remain fixed in the form of polyP instead of being made available for P_i reflux into the cytosol.

In line with this, the fraction of *ppn1Δ ppn2Δ* double mutants that showed nuclear Pho4-GFP was 3-fold higher than in wildtype. *pho91Δ* cells, in which we expect P_i export from the vacuoles to be impaired, showed two times higher frequency of nuclear Pho4-GFP than wildtype. *pho91Δ vtc4Δ* double mutants showed an even lower frequency of nuclear Pho4-GFP than wildtype, suggesting that the state of P_i starvation that *pho91Δ* favours is dependent on vacuolar polyP accumulation. Cells expressing *pho91*^{K237A} as the sole source of Pho91 also showed a 50% lower frequency of nuclear Pho4-GFP. The *pho91*^{K237A} allele generates an amino acid substitution in the InsPP-binding patch of the Pho91 SPX domain. It mimics the InsPP-free state (Wild *et al*, 2016 [DOI](#)) and hence low- P_i conditions (Chabert *et al*, 2023 [DOI](#)). Collectively, our results are consistent with the notion that loss of InsPP binding activates Pho91 to export P_i from vacuoles to the cytosol, enhancing repression of the PHO pathway.

Since InsPP levels in yeast decline in response to cellular P_i availability (Lonetti *et al*, 2011 [DOI](#); Chabert *et al*, 2023 [DOI](#)), we tested the impact of Pho91, Ppn1 and Ppn2 on the cellular levels of these metabolites using CE-MS, capillary electrophoresis coupled to mass spectrometry (CE-MS) (Qiu *et al*, 2020 [DOI](#), 2023 [DOI](#), 2021 [DOI](#)). The cells were harvested from the same P_i -replete growth conditions as for the microscopic assays above (Fig. 7 [DOI](#)).

The *ppn1Δ ppn2Δ* double mutants showed a decrease by 30 to 50% in the inositol pyrophosphates 1,5-InsP₈, 5-InsP₇ and 1-InsP₇, whereas InsP₆ remained at a similar level as in wildtype. InsPP changes at this scale are functionally significant, because they suffice to trigger the initial phase of the P_i starvation response (Chabert *et al*, 2023 [DOI](#); Kim *et al*, 2023 [DOI](#), 2025 [DOI](#)). This is consistent with the 3-fold increase of cells with nuclear Pho4-GFP (Fig. 6 [DOI](#)). *pho91Δ* cells did not show significant changes for all four metabolites although they showed a partial shift of Pho4-GFP into the nucleus (Fig. 6 [DOI](#)). We attribute this discrepancy to the different nature of the assays. The microscopic assay for Pho4-GFP localization can pick up effects in a fraction of the cells because it offers single cell resolution. Inositol pyrophosphate analysis is an ensemble assay, in which changes in a smaller fraction of the population become diluted through the major pool that does not show the effect. For this reason, a change in inositol pyrophosphates affecting only 10% of the *pho91Δ* cells that show nuclear Pho4-GFP remains undetectable. The microscopic assay suggests that cytosolic P_i scarcity in *ppn1Δ ppn2Δ* affects more cells, perhaps because it is more profound, and it hence becomes detectable even in a whole-population-analysis.




Fig. 6

Impact of the vacuolar polyP/P_i cycle on cytosolic P_i signalling

A) Illustration of nucleo-cytoplasmic relocation of Pho4-GFP in response to P_i availability. Wildtype yeast cells were grown in SC medium under P_i replete conditions. During exponential phase (OD_{600nm}=1), cells were transferred for 30 min to synthetic complete media with 200 μM phosphate (-P_i) or 7.5 mM P_i (+P_i) and imaged by fluorescence microscopy. B) The indicated yeast strains were logarithmically grown over night in SC medium with 7.5 mM P_i, harvested at OD_{600nm}=1 and immediately imaged by fluorescence microscopy as in A. The graph shows the means and SEM of the percentage of cells showing Pho4 predominantly in the nucleus. n=3 independent experiments with 200 cells quantified per sample. ** p<0.005, *** p<0.001, ****p<0.0001 from a paired t-test comparing WT with the mutants.




Fig. 7

Impact of Pho91 and vacuolar polyphosphatases on InsPP levels.

pho91Δ and *ppn1Δppn2Δ* cells, as well as their isogenic wildtypes, were logarithmically grown in P_i-replete SC medium as in Fig. 6. At OD_{600nm}=1 the cells were extracted with perchloric acid as previously described (Wilson *et al.*, 2015) and analysed for the indicated InsPs through capillary electrophoresis coupled to mass spectrometry (CE-MS) as described (Qiu *et al.*, 2023). n=3. * p<0.05 from a paired t-test comparing WT with the mutants.

Discussion

Our observations can be integrated with existing data on the properties of VTC (Wild *et al.*, 2016; Hothorn *et al.*, 2009; Müller *et al.*, 2002; Gerasimaite *et al.*, 2014, 2017; Liu *et al.*, 2023; Guan *et al.*, 2023; Pipercevic *et al.*, 2023) and Pho91 (Hürlimann *et al.*, 2007; Wang *et al.*, 2015; Potapenko *et al.*, 2018) to generate a working model explaining how an acidocalcisome-like organelle such as the yeast vacuole is set up to function as a P_i buffer for the cytosol. Under P_i -replete conditions, high InsPP levels activate VTC to polymerize P_i into polyP and translocate it into the vacuolar lumen. Here, the vacuolar polyphosphatases degrade polyP into P_i , filling the lumen with P_i (Sethuraman *et al.*, 2001; Shi & Kornberg, 2005; Lichko *et al.*, 2010; Gerasimaite & Mayer, 2017). Since the P_i exporter Pho91 is downregulated through InsPP binding to its SPX domain (Potapenko *et al.*, 2018; Hürlimann *et al.*, 2007; Wang *et al.*, 2015), the P_i liberated through polyP hydrolysis accumulates in the vacuoles. Product inhibition of the polyphosphatases attenuates polyP hydrolysis once the vacuolar lumen has reached a P_i concentration above 30 mM. When the cells experience P_i scarcity, InsPP levels decline (Chabert *et al.*, 2023). This activates Pho91 to release P_i from the vacuolar pool into the cytosol and stabilizes cytosolic P_i .

Yeast cells do not only accumulate P_i as a rapidly accessible buffer for the cytosol. Under P_i -replete conditions they accumulate hundreds of millimolar of phosphate in the form of polyP (Urech *et al.*, 1978). In contrast to the vacuolar reserve of P_i , which is presumed to be accessible immediately, mobilizing the polyP store takes minutes to hours (Nicolay *et al.*, 1982; Pondugula *et al.*, 2009; Bru *et al.*, 2016). But the polyP store offers advantages in the form of high capacity - hundreds of millimolar of phosphate units can be stored in the form of polyP - and low osmotic activity of polyP (Dürr *et al.*, 1979). Keeping such a large stock of a critical resource, which is often growth-limiting in nature, is relevant for the cells. In case of phosphate shortage, the vacuolar polyP store can be mobilized to enable the cells to complete the cell cycle and transition into G_0 phase (Spain DCB Facultat de Medicina i Ciències de la Salut Universitat Internacional de Catalunya, 2015; Westenberg *et al.*, 1989; Müller *et al.*, 1992). This can consume substantial amounts of phosphate, because we can estimate that replicating the entire DNA (1.2×10^7 base pairs) immobilizes roughly 1 mM P_i in the cells; and cellular RNA is even 50 times more abundant than DNA (Warner, 1999), accounting for 50 mM phosphate. Phospholipids, which must also be synthesized to complete a cell cycle, fix phosphate in similar amounts (Lange & Heijnen, 2001). Thus, a large polyP store is necessary to guarantee that the cells can finish S-phase upon a shortage of phosphate sources. In accord with this notion, absence of the polyP store impairs cell cycle progression, nucleotide synthesis and induces genome instability (Bru *et al.*, 2016, 2017). Also, a shift from non-fermentable carbon sources to fermentation of glucose leads to a strong requirement for P_i because the activation of glucose uptake and glycolysis depends on large amounts of phosphate-containing sugars and glycolytic intermediates (Nicolay *et al.*, 1982, 1983; Gillies *et al.*, 1981). Shortage of P_i restrains the abundance of these metabolites (Kim *et al.*, 2023).

The properties of the regulatory circuit described above imply an inbuilt switch from vacuolar P_i accumulation to large-scale stocking of vacuolar polyP. P_i -replete conditions generate high cellular InsPP levels. These will not only reduce P_i efflux from the vacuoles through Pho91 and inactivate the vacuolar polyphosphatases, but at the same time stimulate continued polyP synthesis by VTC. Coincidence of these effects will favour storage and high accumulation of phosphate in the form of polyP. Conversely, depletion of the vacuolar P_i reservoir upon P_i scarcity in the medium will activate the vacuolar polyphosphatases. In combination with the downregulation of the polyP polymerase VTC through the decline of InsPPs this will mobilize the large vacuolar polyP reserve once the immediately available vacuolar P_i pool is gradually depleted.




Fig. 8

Working model of acidocalcisome-like vacuoles as P_i buffering systems.

(A) Under P_i -replete conditions, ATP drives the conversion of P_i into polyP and its translocation into the organelle. Here, polyP is degraded by the vacuolar polyphosphatases Ppn1 and Ppn2 to establish a vacuolar pool of free P_i . Feedback inhibition of P_i gradually reduces polyP degradation, enabling the buildup of a vacuolar polyP stock. Red lines and SPX colouring indicate inhibitory action, green colouring stimulation. (B) Cytosolic P_i scarcity decreases InsPP levels, which triggers two compensatory, SPX-controlled effects: The transfer of P_i from the cytosol into vacuoles through VTC ceases; and Pho91-dependent export of P_i from vacuoles is activated. Both measures synergize to stabilize cytosolic P_i . The export of P_i from the vacuole lifts product inhibition on the polyphosphatases Ppn1 and Ppn2 and stimulates a compensatory degradation of polyP.

The concentration of P_i inside vacuoles as a rapidly accessible P_i reserve, and the synthesis of a large polyP stock, come at an energetic cost because the transformation of P_i into polyP requires the formation of phosphoric anhydride bonds (Hothorn *et al*, 2009 [\[1\]](#); Gerasimaite *et al*, 2014 [\[2\]](#)) and vacuolar P_i reaches 30 mM. This exceeds the cytosolic P_i concentration, which was measured through ^{31}P -NMR in a variety of yeasts, yielding values of 5–17 mM (Nicolay *et al*, 1982 [\[3\]](#), 1983 [\[4\]](#)). Cytosolic P_i can also be estimated based on data from several other studies (Auesukaree *et al*, 2004 [\[5\]](#); Hürlimann *et al*, 2009 [\[6\]](#); Theobald *et al*, 1996 [\[7\]](#); Zhang *et al*, 2015 [\[8\]](#); Pinson *et al*, 2004 [\[9\]](#)). Assuming that the cytosolic volume of a BY4741 yeast cell is 40 fL (Chabert *et al*, 2023 [\[10\]](#)), and 1 g of dry weight contains 40×10^9 yeast cells, these studies point to cytosolic values of 10–15 mM in P_i -replete media. Upon P_i starvation, this value rapidly drops up to fivefold, resulting in a strong P_i gradient across the vacuolar membrane (Okorokov *et al*, 1980 [\[11\]](#); Shirahama *et al*, 1996 [\[12\]](#)). To replenish the cytosolic pool under P_i scarcity, Pho91 can exploit not only this P_i concentration gradient, but also the vacuolar electrochemical potential, which was shown to stimulate P_i export through the Pho91 homolog OsSPX-MFS3 from plant vacuoles (Wang *et al*, 2015 [\[13\]](#)).

Vacuolar P_i accumulation is driven indirectly through ATP in two ways. VTC uses ATP as a substrate and transfers the phosphoric anhydride bond of the ψ -phosphate onto a polyP chain (Hothorn *et al*, 2009 [\[1\]](#)). The growing polyP chain exits from the catalytic site directly towards the transmembrane part of VTC (Liu *et al*, 2023 [\[14\]](#); Guan *et al*, 2023 [\[15\]](#)). This transmembrane part likely forms a controlled channel that can guide polyP through the membrane (Liu *et al*, 2023 [\[14\]](#)). Coupled synthesis and translocation require the V-ATPase (Gerasimaite *et al*, 2014 [\[2\]](#)), probably because polyP is highly negatively charged and therefore follows the electrochemical potential across the vacuolar membrane of 180 mV (inside positive) and 1.7 pH units (Kakinuma *et al*, 1981 [\[16\]](#)), which is generated through the proton pumping V-ATPase. Thus, the combination of the VTC complex and vacuolar polyphosphatases can be considered as a P_i pump that is driven by ATP through polyP synthesis and through the electrochemical potential for polyP translocation and P_i export.

It is likely that acidocalcisome- and lysosome-like organelles of other organisms act as buffers for cytosolic P_i similarly as described in our model for yeasts. This notion is supported by the conserved molecular setup of acidocalcisome-like organelles as well as by phenotypic similarities. The acidocalcisomes of Trypanosomes contain VTC, a Pho91 homolog and proton pumps in their membranes, and polyphosphatases in their lumen (Huang & Docampo, 2015 [\[17\]](#); Billington *et al*, 2023 [\[18\]](#); Fang *et al*, 2007 [\[19\]](#); Lander *et al*, 2013 [\[20\]](#); Ulrich *et al*, 2013 [\[21\]](#); Scott *et al*, 1997 [\[22\]](#)). Also the acidocalcisome-like organelles of the alga Chlamydomonas contain such proteins and they accumulate polyP through VTC as a function of the availability of P_i , a proton gradient and metal ions (Zúñiga-Burgos *et al*, 2024 [\[23\]](#); Blaby-Haas & Merchant, 2014 [\[24\]](#); Long *et al*, 2023 [\[25\]](#); Hong-Hermesdorf *et al*, 2014 [\[26\]](#); Aksoy *et al*, 2014 [\[27\]](#); Ruiz *et al*, 2001 [\[28\]](#); Goodenough *et al*, 2019 [\[29\]](#)). Like in yeast, the polyP stores are mobilized upon P_i limitation (Sanz-Luque *et al*, 2020 [\[30\]](#); Plouviez *et al*, 2021 [\[31\]](#)). Drosophila has a potentially lysosome-related compartment, which is acidic, carries V-ATPase and a homolog of the P_i exporter XPR1, impacts cytosolic P_i and diminishes upon P_i starvation (Xu *et al*, 2023 [\[32\]](#)). Mammalian lysosome-like organelles also participate in P_i homeostasis. They can accumulate polyP and take up P_i (Pisoni, 1991 [\[33\]](#); Pisoni & Lindley, 1992 [\[34\]](#)). They carry the P_i exporter XPR1, which interacts with the plasma membrane P_i importer PiT1 to regulate its degradation (Li *et al*, 2024 [\[35\]](#)).

We hence propose that acidocalcisome-like vacuoles may have a general role as feedback-controlled, rapidly accessible P_i buffers for the cytosol, addressing a critical parameter for metabolism. However, given that acidocalcisome-like organelles accumulate not only phosphate but also multiple other metabolites and ions (Docampo, 2024 [\[36\]](#)), they are probably interlinked with cellular metabolism in multiple ways and might form an important hub for its homeostasis.

Materials and Methods

Yeast strains and growth conditions

Saccharomyces cerevisiae cells were grown on yeast extract-peptone-dextrose (YPD: 1% yeast extract, 2% peptone and 2% dextrose) or in synthetic complete (SC) medium from Formedium, supplemented with sodium phosphate as needed). Yeast backgrounds used in this study were BY4741 and BY4742. Genetic manipulations of yeast were performed by homologous recombination according to published procedures and/or transformation with the indicated plasmids (Gietz & Schiestl, 2007 [↗](#); Güldener *et al.*, 1996 [↗](#)). Strains used are listed in **Table 1** [↗](#).

PHO4 localization

Cells transformed with pRS415- P_{PHO4} -PHO4-GFP, a plasmid expressing a PHO4-GFP fusion from the PHO4 promoter (Chabert *et al.*, 2023 [↗](#)), were grown exponentially overnight in SC medium without leucine (SC-Leu). Care was taken that the culture did not grow beyond a density of $OD_{600}=0.7$. At this point, the cells were collected by brief centrifugation (15 sec, 3000 x g) in a tabletop centrifuge and resuspended in 1/10th to 1/20th of their own supernatant. Pho4-GFP localization was immediately checked by microscopy. An aliquot of the cells was washed twice with SC-Leu without P_i and then diluted in SC-Leu with 200 μ M phosphate to $OD_{600}=0.7$. After 30 minutes of incubation in this medium, Pho4-GFP localization was analysed by fluorescence microscopy on a LEICA DMI6000B inverted microscope equipped with a Hamamatsu ORCA-R2 (C10600-10B) camera, an XCite ® series 120Q UV lamp and a Leica 100x 1.4 NA lens.

Vacuole preparation

Vacuoles were purified from yeast cells essentially as described (D'Agostino & Mayer, 2019 [↗](#)). Briefly, yeast cells were grown in 1L of YPD to an OD_{600} of 1.5. 330 ml of cells were collected by centrifugation and resuspended in 50 ml of 30 mM Tris-HCl pH 8.9, 10 mM DTT buffer. Suspensions were incubated in a 30° C water bath for 5 min and collected by centrifugation. The pellet was resuspended in 15 ml of spheroplasting solution (50 mM K-phosphate pH 7.5, 600 mM sorbitol in YPD with 0.2% D-glucose and 3600 U/ml lyticase) and incubated for 25 min at 30°C. Spheroplasts were collected by centrifugation (2500 x g, 3 min) and resuspended in 2.5 ml of 15% Ficoll 400 in PS buffer (10 mM PIPES-KOH pH 6.8, 200 mM sorbitol). 80 μ g of DEAE-dextran were added under gentle mixing. After incubation on ice for 2 min and then at 30 °C for 80 sec, spheroplasts were transferred into Beckman SW41.1 tubes, overlaid with cushions of 8%, 4% and 0% Ficoll 400 in PS buffer, and centrifuged (150'000 x g, 90 min, 2 °C). Vacuoles were collected from the 0-4 % Ficoll interface. Their protein concentration was determined through Bradford assay using BSA as a standard.

Polyphosphatase activity of vacuolar lysates

Polyphosphatase activity was assayed as described previously (Gerasimaite & Mayer, 2017 [↗](#)), with the following modifications. Isolated vacuoles were diluted to a final protein concentration of 0.002 mg/ml in 1 ml of reaction buffer (20 mM PIPES/KOH pH 6.8, 150 mM KCl, 1 mM $ZnCl_2$ or $MgCwWl_2$, 0.1% Triton X-100, 1xPIC, 1 mM PMSF and 30 or 300 μ M polyP₃₀₀) in the presence or absence of various concentrations of KH_2PO_4 and incubated at 27°C. At the indicated times, 80 μ l aliquots were collected and the reaction was stopped by dilution with 160 μ l of stop solution (10 mM PIPES/KOH pH 6.8, 150 mM KCl, 12 mM EDTA pH 8.0, 0.1% Triton X-100, 15 μ M DAPI). Remaining polyP was quantified by measuring polyP-DAPI fluorescence (λ_{exc} . 415 nm, λ_{em} . 550 nm) in a black 96 well plate in a Spectramax Gemini microplate fluorometer (Molecular Devices). A reaction containing boiled vacuoles was used as negative control.

Table 1

Background	Genotype	Source
BY4741	MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0	Euroscarf
BY4742	MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0	Euroscarf
BY4741	<i>vtc4::kanMX</i>	(Müller <i>et al</i> , 2002)
BY4741	<i>pho91::kanMX</i>	This study
BY4742	<i>ppn1::natNT2</i>	(Gerasimaite & Mayer, 2017)
BY4742	<i>ppn2::URA3</i>	Gerasimaite & Mayer, 2017)
BY4741	VTC5 natNT2-GPD1pr	(Desfougères <i>et al</i> , 2016a)
BY4742	VTC3 ^{K126A} VTC4 ^{K129A}	(Wild <i>et al</i> , 2016)
BY4741	<i>pho91::kanMX</i> pRS303-P _{PHO91} -PHO91 (K237A)	This study
BY4741	<i>pho91::kanMX</i> pRS303-P _{PHO91} -PHO91 (Y22F)	This study
BY4741	<i>vtc4::kanMX pho91::hyg</i>	This study
BY4741	pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	<i>vtc4::kanMX</i> pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	<i>pho91::kanMX</i> pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	<i>vtc4::kanMX pho91::hyg</i> pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	VTC5 natNT2-GPD1pr <i>pho91::hyg</i> pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	VTC5 natNT2-GPD1pr pRS415-P _{PHO4} -PHO4-GFP	This study
BY4742	<i>ppn1::natNT2 ppn2::URA3</i> pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	<i>pho91::kanMX</i> pRS303-P _{PHO91} -PHO91 (K237A) pRS415-P _{PHO4} -PHO4-GFP	This study
BY4741	pRS416-P _{ADH} -SPL2 <i>pho91::kanMX</i> pRS303-P _{PHO91} -PHO91 (K237A) pRS415-P _{PHO4} -PHO4-GFP	This study

Polyphosphate synthesis by isolated vacuoles

Polyphosphate synthesis was assayed as described (Gerasimaite *et al.*, 2014 [\[1\]](#)). Isolated vacuoles were diluted to final protein concentration of 0.02 mg/ml on 1 ml of reaction buffer (10 mM PIPES/KOH pH 6.8, 150 mM KCl, 0.5 mM MnCl₂, 200 mM sorbitol) and the reaction was started by adding an ATP regenerating system (1 mM ATP-MgCl₂, 40 mM creatine phosphate and 0.25 mg/ml creatine kinase). The mix was incubated at 27 °C. At different timepoints, 80 µl aliquots were mixed with 160 µl of stop solution (10 mM PIPES/KOH pH 6.8, 150 mM KCl, 200 mM sorbitol, 12 mM EDTA, 0.15% Triton X-100 and 15 µM DAPI). PolyP synthesis was quantified through polyP-DAPI fluorescence (λ_{exc} , 415 nm, λ_{em} , 550 nm) in a black 96 well plate. A calibration curve was prepared using commercial polyP₆₀ as a standard.

Phosphate quantification in isolated vacuoles

Isolated vacuoles were incubated as described for the polyP synthesis assay above. At different time points, 80 µl aliquots were centrifuged (3 min, 2000 x g, 2°C), the pellets were washed with 500 µl of washing solution (10 mM PIPES/KOH pH 6.8, 200 mM Sorbitol, 150 mM KCl) and centrifuged as before. The final pellet was resuspended with 100 µl of lysis buffer (10 mM PIPES/KOH pH 6.8, 200 mM sorbitol, 150 mM KCl, 12 mM EDTA, 0.1 % Triton). Free phosphate was quantified by adding 150 µl of molybdate-malachite green solution (1 mM malachite green, 10 mM ammonium molybdate, 1M HCl) and reading the absorbance at 595 nm in a microplate photometer.

Inositol pyrophosphate synthesis, extraction and quantification

5-InsP7 was synthesized as described (Capolicchio *et al.*, 2013 [\[2\]](#); Wang *et al.*, 2014 [\[3\]](#)). For quantification from cells, InsPPs extraction was performed as described previously (Kim *et al.*, 2023 [\[4\]](#)), with the following modifications. Briefly, 3 ml of yeast cell culture at OD_{600nm}=1 was collected using rapid vacuum-filtration on a polytetrafluoroethylene membrane filter (1.2 µm; Piper Filter GmbH, Germany). After snap freezing in liquid nitrogen, yeast cells on the membrane were resuspended in 400 µL of 1 M perchloric acid and lysed by bead beating (glass beads; 0.25-0.5 mm) for 10 min at 4 °C. After centrifugation at 13,000 rpm for 3 min at 4 °C, the supernatant was transferred into a new tube containing 3 mg of titanium dioxide (TiO₂) beads (GL Sciences, Japan) which had been pre-washed twice with H₂O and 1 M perchloric acid. The sample was gently rotated for 15 min at 4 °C. The TiO₂ beads were collected by centrifugation at 13,000 rpm for 1 min at 4 °C and washed twice using 1 M perchloric acid. After the second washing step, the TiO₂ beads were resuspended in 300 µL of 3 % (v/v) NH₄OH and rotated gently at room temperature. After centrifugation at 13,000 rpm for 1 min, the eluants were transferred into a new tube and dried in SpeedVac (Labogene, Denmark) at 42 °C. InsPPs were measured through capillary electrophoresis coupled to mass spectrometry (CE-MS) as described (Qiu *et al.*, 2021 [\[5\]](#)).


Quantification of the vacuolar proteome

Vacuoles were prepared as described above. During the purification procedure, a sample of the sedimented spheroplasts was withdrawn before DEAE dextran was added. These withdrawn spheroplasts constitute the “whole cell” extract. After the flotation step, the vacuoles withdrawn from the 4%-0% Ficoll interface were used as the vacuole fraction. For each fraction samples with 100 µg protein were precipitated by adding a final concentration 12.5% TCA for 10 min on ice. The proteins were sedimented (12'000 x g, 5 min, room temperature), the supernatant discarded, and the pellets were washed with ice-cold acetone and centrifuged as above twice. The final pellet was dried and dissolved in reducing SDS sample buffer (10% glycerol, 50 mM Tris-HCl pH 6.8, 2 mM DTT, 2% SDS, 0.002% bromophenol blue).

Samples dissolved in Laemmli buffer (50 mM Tris pH 6.8, 10 mM DTT, 2 % SDS, 0.1 % bromophenol blue, 10 % glycerol) were equalized in concentration by dilution in SP3 buffer (2% SDS, 10 mM DTT, 50 mM Tris, pH 7.5). 150 µg of protein per sample were heated at 95 °C for 5 min and cooled down. Reduced cysteine residues were alkylated by adding iodoacetamide (30 mM final) and incubating for 45 min at room temperature in the dark. Digestion was done by the SP3 method (Hughes *et al*, 2019 [\[link\]](#)) using magnetic Sera-Mag Speedbeads (Cytiva 45152105050250, 50 mg/ml). Beads were added at a ratio 10:1 (w:w) to samples, and proteins were precipitated on beads with ethanol (final concentration: 60 %). After 3 washes with 80% ethanol, beads were digested in 50 µl of 100 mM ammonium bicarbonate with 3.0 µg of trypsin (Promega #V5073). After 1 h of incubation at 37 °C, the same amount of trypsin was added to the samples for an additional 1 h of incubation. Supernatants were then recovered, transferred to new tubes, acidified with formic acid (0.5% final concentration), and dried by centrifugal evaporation. To remove traces of SDS, two sample volumes of isopropanol containing 1% TFA were added to the digests, and the samples were desalted on a strong cation exchange (SCX) plate (Oasis MCX; Waters Corp., Milford, MA) by centrifugation. After washing with isopropanol/1%TFA and 2% acetonitrile/0.1% FA, peptides were eluted in 200µl of 80% MeCN, 19% water, 1% (v/v) ammonia, and dried by centrifugal evaporation. Data-dependent LC-MS/MS analyses of samples were carried out on a Fusion Tribrid Orbitrap mass spectrometer (Thermo Fisher Scientific) interfaced through a nano-electrospray ion source to an Ultimate 3000 RSLCnano HPLC system (Dionex). Peptides were separated on a reversed-phase custom packed 45 cm C18 column (75 µm ID, 100Å, Reprosil Pur 1.9 µm particles, Dr. Maisch, Germany) with a 4-90% acetonitrile gradient in 0.1% formic acid at a flow rate of 250 nl/min (total time 140 min). Full MS survey scans were performed at 120'000 resolution. A data-dependent acquisition method controlled by Xcalibur software (Thermo Fisher Scientific) was used that optimized the number of precursors selected (“top speed”) of charge 2+ to 5+ while maintaining a fixed scan cycle of 0.6 s. Peptides were fragmented by higher energy collision dissociation (HCD) with a normalized energy of 32%. The precursor isolation window used was 1.6 Th, and the MS2 scans were done in the ion trap. The m/z of fragmented precursors was then dynamically excluded from selection during 60 s.

Data files were analysed with MaxQuant 1.6.14.0 (Cox & Mann, 2008 [\[link\]](#)) incorporating the Andromeda search engine (Cox *et al*, 2011 [\[link\]](#)). Cysteine carbamidomethylation was selected as fixed modification while methionine oxidation and protein N-terminal acetylation were specified as variable modifications. The sequence databases used for searching were the *S. cerevisiae* reference proteome based on the UniProt database (<https://www.uniprot.org> [\[link\]](#), version of June 6th, 2021, containing 6050 sequences), and a “contaminant” database containing the most usual environmental contaminants and enzymes used for digestion (keratins, trypsin, etc). Mass tolerance was 4.5 ppm on precursors (after recalibration) and 20 ppm on MS/MS fragments. Both peptide and protein identifications were filtered at 1% FDR relative to hits against a decoy database built by reversing protein sequences. The match between runs feature was enabled.


Supplementary figures



Supplementary Figure 1

Isolated vacuoles during polyP synthesis


BJ 4742 (WT) vacuoles were incubated in vitro as in [Fig. 1B](#). After 10 min, 0.1 μM of the fluorophore FM4-64 was added to stain the membranes. The suspension was immediately imaged on a Yokogawa spinning disc microscope with a Gataca LiveSR unit and a 100x 1.45 NA objective. Stacks were taken at 0.2 μm z-distance. The image shows an optical section out of such a stack. To determine vacuole diameters, the stacks were subjected to maximum intensity projection in ImageJ. The largest visible diameter for 100 organelles was measured manually using ImageJ, yielding an average diameter of 0.8 μm with an SD of 0.2 μm Scale bar: 1 μm



Supplementary Figure 2

Effects of N- and C-terminal fluorescent protein tags on the localization of Pho91.

PHO91 was fused with a variety of N- or C-terminal protein tags and peptide spacers as indicated. They were expressed from the endogenous PHO91 promoter or, where indicated, from the overexpressing ADH1 or GPD1 promoters. Cells were logarithmically grown overnight in SC medium, harvested at OD_{600nm} of 1-2, and analysed by fluorescence microscopy.



Supplementary Figure 3

Pho91 cannot replace other P_i transporters to support growth of yeast.

We generated a BY4741 strain with a quintuple deletion of the genes for the known P_i transporters, PHO84, PHO87, PHO89, PHO90 and PHO91 (D5m). These cells were kept alive by expressing the gene for the plasma membrane P_i transporter PHO89 from a URA3-based centromeric (single copy) plasmid (pRS416). PHO91 was expressed from a HIS3-based centromeric plasmid (pRS315). Cells were plated in a dilution series on SC lacking histidine (SC-HIS) to verify that the cells had the HIS3-based PHO91 plasmid, or on SC with 5-fluoro-orotic acid (5-FOA), a drug that forces cells to lose the URA3-based pRS416 and thus to live without PHO89. PHO91 as the sole P_i transporter gene (on SC + 5-FOA) does not allow cells to grow.

Acknowledgements

Mass spectrometry-based proteomics work was performed by the Protein Analysis Facility of the Faculty of Biology and Medicine, University of Lausanne. This study was supported by grants from the SNSF (320030-228119, 31003A_179306 and 310030_204713) and ERC (788442) to AM, by the HFSP (LT000588/2019) to GDK, by the Deutsche Forschungsgemeinschaft (CIBSS, EXC-2189, Project ID 390939984, to HJJ) and the Volkswagen Foundation (VW Momentum Grant 98604 to HJJ).

Additional information

Data availability

Data from the proteomic analysis have been deposited at the PRIDE database under the identifier PXD060102.

Author contributions

AM conceived the study, analysed data and wrote the paper. SB performed in vitro and in vivo experiments with yeast. GDK, LM, GL, and HJJ analysed inositol pyrophosphates. All authors assembled and corrected the manuscript.

Funding

Swiss National Science Foundation (320030-228119)

Swiss National Science Foundation (31003A_179306)

Swiss National Science Foundation (310030_204713)

European Research Council

<https://doi.org/10.3030/788442> 

Deutsche Forschungsgemeinschaft (EXC-2189 Project 390939984)

Volkswagen Foundation (98604)

International Human Frontier Science Program Organization (LT000588/2019)

References

Aksoy M, Pootakham W, Grossman AR (2014) **Critical Function of a *Chlamydomonas reinhardtii* Putative Polyphosphate Polymerase Subunit during Nutrient Deprivation** *Plant Cell* **26**:4214–4229 [Google Scholar](#)

Andreeva N, Ledova L, Ryazanova L, Tomashevsky A, Kulakovskaya T, Eldarov M (2019) **Ppn2 endopolyphosphatase overexpressed in *Saccharomyces cerevisiae*: Comparison with Ppn1, Ppx1, and Ddp1 polyphosphatases** *Biochimie* **163**:101–107 [Google Scholar](#)

Auesukaree C, Homma T, Tochio H, Shirakawa M, Kaneko Y, Harashima S (2004) **Intracellular phosphate serves as a signal for the regulation of the PHO pathway in *Saccharomyces cerevisiae*** *J Biol Chem* **279**:17289–17294 [Google Scholar](#)

Austin S, Mayer A (2020) **Phosphate Homeostasis - A Vital Metabolic Equilibrium Maintained Through the INPHORS Signaling Pathway** *Front Microbiol* **11**:1367 [Google Scholar](#)

Billington K, Halliday C, Madden R, Dyer P, Barker AR, Moreira-Leite FF, Carrington M, Vaughan S, Hertz-Fowler C, Dean S, et al (2023) **Genome-wide subcellular protein map for the flagellate parasite *Trypanosoma brucei*** *Nat Microbiol* **8**:533–547 [Google Scholar](#)

Blaby-Haas CE, Merchant SS (2014) **Lysosome-related organelles as mediators of metal homeostasis** *J Biol Chem* **289**:28129–28136 [Google Scholar](#)

Bru S, Martínez-Laínez JM, Hernández-Ortega S, Quandt E, Torres-Torronteras J, Martí R, Canadell D, Ariño J, Sharma S, Jiménez J, et al (2016) **Polyphosphate is involved in cell cycle progression and genomic stability in *Saccharomyces cerevisiae*** *Mol Microbiol* **101**:367–380 [Google Scholar](#)

Bru S, Samper-Martín B, Quandt E, Hernández-Ortega S, Martínez-Laínez JM, Garí E, Rafel M, Torres-Torronteras J, Martí R, Ribeiro MPC, et al (2017) **Polyphosphate is a key factor for cell survival after DNA damage in eukaryotic cells** *DNA Repair* **57**:171–178 [Google Scholar](#)

Capolicchio S, Thakor DT, Linden A, Jessen HJ (2013) **Synthesis of Unsymmetric Diphospho-Inositol Polyphosphates** *Angew Chem Int Ed* **52**:6912–6916 [Google Scholar](#)

Chabert V, Kim G-D, Qiu D, Liu G, Mayer LM K MJ, Jessen HJ, Mayer A (2023) **Inositol pyrophosphate dynamics reveals control of the yeast phosphate starvation program through 1,5-IP₈ and the SPX domain of Pho81** *eLife* **12**:RP87956 <https://doi.org/10.7554/eLife.87956> | [Google Scholar](#)

Cox J, Mann M (2008) **MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification** *Nat Biotechnol* **26**:1367–1372 [Google Scholar](#)

Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M (2011) **Andromeda: a peptide search engine integrated into the MaxQuant environment** *J Proteome Res* **10**:1794–1805 [Google Scholar](#)

D'Agostino M, Mayer A (2019) **Assay of Lipid Mixing and Fusion Pore Formation in the Fusion of Yeast Vacuoles** *Methods Mol Biol* **1860**:253–262 [Google Scholar](#)

Desfougères Y, Gerasimaitė RU, Jessen HJ, Mayer A (2016a) **Vtc5, a Novel Subunit of the Vacuolar Transporter Chaperone Complex, Regulates Polyphosphate Synthesis and Phosphate Homeostasis in Yeast** *J Biol Chem* **291**:22262–22275 [Google Scholar](#)

Desfougères Y, Neumann H, Mayer A (2016b) **Organelle size control - increasing vacuole content activates SNAREs to augment organelle volume through homotypic fusion** *J Cell Sci* **129**:2817–2828 [Google Scholar](#)

Docampo R (2024) **Advances in the cellular biology, biochemistry, and molecular biology of acidocalcisomes** *Microbiol Mol Biol Rev* **88**:e0004223 [Google Scholar](#)

- Docampo R, Huang G (2016) **Acidocalcisomes of eukaryotes** *Curr Opin Cell Biol* **41**:66–72 [Google Scholar](#)
- Docampo R, Souza W de, Miranda K, Rohloff P, Moreno SNJ (2005) **Acidocalcisomes - conserved from bacteria to man** *Nat Rev Microbiol* **3**:251–261 [Google Scholar](#)
- Donaldson AD, Raghuraman MK, Friedman KL, Cross FR, Brewer BJ, Fangman WL (1998) **CLB5-Dependent Activation of Late Replication Origins in *S. cerevisiae*** *Mol Cell* **2**:173–182 [Google Scholar](#)
- Dürr M, Boller T, Wiemken A (1975) **Polybase induced lysis of yeast spheroplasts. A new gentle method for preparation of vacuoles** *Arch Microbiol* **105**:319–27 [Google Scholar](#)
- Dürr M, Urech K, Boller T, Wiemken A, Schwencke J, Nagy M (1979) **Sequestration of Arginine by Polyphosphate in Vacuoles of Yeast (*Saccharomyces-Cerevisiae*)** *Arch Microbiol* **121**:169–175 [Google Scholar](#)
- Eising S, Esch B, Wälte M, Duarte PV, Walter S, Ungermann C, Bohnert M, Fröhlich F (2022) **A lysosomal biogenesis map reveals the cargo spectrum of yeast vacuolar protein targeting pathways** *J Cell Biol* **221**:e202107148 [Google Scholar](#)
- Elbaz-Alon Y, Rosenfeld-Gur E, Shinder V, Futerman AH, Geiger T, Schuldiner M (2014) **A dynamic interface between vacuoles and mitochondria in yeast** *Dev Cell* **30**:95–102 [Google Scholar](#)
- Fang J, Rohloff P, Miranda K, Docampo R (2007) **Ablation of a small transmembrane protein of *Trypanosoma brucei* (TbVTC1) involved in the synthesis of polyphosphate alters acidocalcisome biogenesis and function, and leads to a cytokinesis defect** *Biochem J* **407**:161–170 [Google Scholar](#)
- Farofonova V, Andreeva N, Kulakovskaya E, Karaginov A, Agaphonov M, Kulakovskaya T (2023) **Multiple effects of the PHO91 gene knockout in *Ogataea parapolyomorpha*** *Folia Microbiol* **73**:1–7 [Google Scholar](#)
- Freimoser FM, Hürlimann HC, Jakob CA, Werner TP, Amrhein N (2006) **Systematic screening of polyphosphate (poly P) levels in yeast mutant cells reveals strong interdependence with primary metabolism** *Genome Biol* **7**:R109 [Google Scholar](#)
- Gerasimaite R, Mayer A (2016) **Enzymes of yeast polyphosphate metabolism: structure, enzymology and biological roles** *Biochem Soc Trans* **44**:234–239 [Google Scholar](#)
- Gerasimaite R, Mayer A (2017) **Ppn2, a novel Zn(2+)-dependent polyphosphatase in the acidocalcisome-like yeast vacuole** *J Cell Sci* **130**:1625–1636 [Google Scholar](#)
- Gerasimaite R, Pavlovic I, Capolicchio S, Hofer A, Schmidt A, Jessen HJ, Mayer A (2017) **Inositol Pyrophosphate Specificity of the SPX-Dependent Polyphosphate Polymerase VTC ACS** *Chem Biol* **12**:648–653 [Google Scholar](#)
- Gerasimaite R, Sharma S, Desfougères Y, Schmidt A, Mayer A (2014) **Coupled synthesis and translocation restrains polyphosphate to acidocalcisome-like vacuoles and prevents its toxicity** *J Cell Sci* **127**:5093–5104 [Google Scholar](#)
- Gietz RD, Schiestl RH (2007) **High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method** *Nat Protoc* **2**:31–34 [Google Scholar](#)

Gillies RJ, Ugurbil K, Hollander JA den, Shulman RG (1981) **³¹P NMR studies of intracellular pH and phosphate metabolism during cell division cycle of *Saccharomyces cerevisiae*** *PNAS* **78**:2125–2129 [Google Scholar](#)

Girard-Dias W, Augusto I, Fernandes TVA, Pascutti PG, Souza W de, Miranda K (2023) **A spatially resolved elemental nanodomain organization within acidocalcisomes in *Trypanosoma cruzi*** *Proc National Acad Sci* **120**:e2300942120 [Google Scholar](#)

Goodenough U, Heiss AA, Roth R, Rusch J, Lee J-H (2019) **Acidocalcisomes: Ultrastructure, Biogenesis, and Distribution in Microbial Eukaryotes** *Protist* **170**:287–313 [Google Scholar](#)

Guan Z, Chen J, Liu R, Chen Y, Xing Q, Du Z, Cheng M, Hu J, Zhang W, Mei W, et al (2023) **The cytoplasmic synthesis and coupled membrane translocation of eukaryotic polyphosphate by signal-activated VTC complex** *Nat Commun* **14**:718 [Google Scholar](#)

Güldener U, Heck S, Fiedler T, Beinhauer J, Hegemann JH (1996) **A new efficient gene disruption cassette for repeated use in budding yeast** *Nucleic Acids Res* **24**:2519–2524 [Google Scholar](#)

Hong-Hermesdorf A, Miethke M, Gallaher SD, Kropat J, Dodani SC, Chan J, Barupala D, Domaille DW, Shirasaki DI, Loo JA, et al (2014) **Subcellular metal imaging identifies dynamic sites of Cu accumulation in *Chlamydomonas*** *Nat Chem Biol* **10**:1034–1042 [Google Scholar](#)

Hothorn M, Neumann H, Lenherr ED, Wehner M, Rybin V, Hassa PO, Uttenweiler A, Reinhardt M, Schmidt A, Seiler J, et al (2009) **Catalytic core of a membrane-associated eukaryotic polyphosphate polymerase** *Science* **324**:513–516 [Google Scholar](#)

Huang G, Docampo R (2015) **Proteomic analysis of acidocalcisomes of *Trypanosoma brucei* uncovers their role in phosphate metabolism, cation homeostasis, and calcium signaling** *Commun Integr Biol* **8**:e1017174 [Google Scholar](#)

Huber-Wälchli V, Wiemken A (1979) **Differential extraction of soluble pools from the cytosol and the vacuoles of yeast (*Candida utilis*) using DEAE-dextran** *Arch Microbiol* **120**:141–149 [Google Scholar](#)

Hughes CS, Moggridge S, Müller T, Sorensen PH, Morin GB, Krijgsveld J (2019) **Single-pot, solid-phase-enhanced sample preparation for proteomics experiments** *Nat Protoc* **14**:68–85 [Google Scholar](#)

Hürlimann HC, Pinson B, Stadler-Waibel M, Zeeman SC, Freimoser FM (2009) **The SPX domain of the yeast low-affinity phosphate transporter Pho90 regulates transport activity** *EMBO Rep* **10**:1003–1008 [Google Scholar](#)

Hürlimann HC, Stadler-Waibel M, Werner TP, Freimoser FM (2007) **Pho91 Is a vacuolar phosphate transporter that regulates phosphate and polyphosphate metabolism in *Saccharomyces cerevisiae*** *Mol Biol Cell* **18**:4438–4445 [Google Scholar](#)

Jimenez V, Docampo R (2015) **TcPho91 is a contractile vacuole phosphate sodium symporter that regulates phosphate and polyphosphate metabolism in *Trypanosoma cruzi*** *Mol Microbiol* **97**:911–925 [Google Scholar](#)

Kakinuma Y, Ohsumi Y, Anraku Y (1981) **Properties of H⁺-translocating adenosine triphosphatase in vacuolar membranes of *Saccharomyces cerevisiae*** *J Biol Chem* **256**:10859–63 [Google Scholar](#)

Kim G-D, Liu G, Qiu D, Leo MGD, Gopaldass N, Hermes J, Timmer J, Saiardi A, Mayer A, Jessen HJ (2025) **Pools of Independently Cycling Inositol Phosphates Revealed by Pulse Labeling with 18O-Water** *J Am Chem Soc* [Google Scholar](#)

Kim G-D, Qiu D, Jessen HJ, Mayer A (2023) **Metabolic Consequences of Polyphosphate Synthesis and Imminent Phosphate Limitation** *Mbio: e* :102–23 [Google Scholar](#)

Klompmaaker SH, Kohl K, Fasel N, Mayer A (2017) **Magnesium uptake by connecting fluid-phase endocytosis to an intracellular inorganic cation filter** *Nature communications* **8**:1879 [Google Scholar](#)

Koren A, Soifer I, Barkai N (2010) **MRC1-dependent scaling of the budding yeast DNA replication timing program** *Genome Res* **20**:781–790 [Google Scholar](#)

Kulakovskaya TV, Andreeva NA, Ledova LA, Ryazanova LP, Trilisenko LV, Eldarov MA (2021) **Enzymes of Polyphosphate Metabolism in Yeast: Properties, Functions, Practical Significance** *Biochemistry Mosc* **86**:S96–S108 [Google Scholar](#)

Kunz JB, Schwarz H, Mayer A (2004) **Determination of Four Sequential Stages during Microautophagy in Vitro *** *J Biol Chem* **279**:9987–9996 [Google Scholar](#)

Lander N, Cordeiro C, Huang G, Docampo R (2016) **Polyphosphate and acidocalcisomes** *Biochem Soc Trans* **44**:1–6 [Google Scholar](#)

Lander N, Ulrich PN, Docampo R (2013) **Trypanosoma brucei vacuolar transporter chaperone 4 (TbVtc4) is an acidocalcisome polyphosphate kinase required for in vivo infection** *J Biol Chem* **288**:34205–34216 [Google Scholar](#)

Lange HC, Heijnen JJ (2001) **Statistical reconciliation of the elemental and molecular biomass composition of Saccharomyces cerevisiae** *Biotechnol Bioeng* **75**:334–344 [Google Scholar](#)

Lemercier G, Espiau B, Ruiz FA, Vieira M, Luo S, Baltz T, Docampo R, Bakalara N (2004) **A Pyrophosphatase Regulating Polyphosphate Metabolism in Acidocalcisomes Is Essential for Trypanosoma brucei Virulence in Mice*** *J Biol Chem* **279**:3420–3425 [Google Scholar](#)

Li SC, Kane PM (2009) **The yeast lysosome-like vacuole: endpoint and crossroads** *Biochim Biophys Acta* **1793**:650–663 [Google Scholar](#)

Li X, Kirkpatrick RB, Wang X, Tucker CJ, Shukla A, Jessen HJ, Wang H, Shears SB, Gu C (2024) **Homeostatic coordination of cellular phosphate uptake and efflux requires an organelle-based receptor for the inositol pyrophosphate IP8** *Cell Rep* **43**:114316 [Google Scholar](#)

Lichko LP, Kulakovskaya TV, Kulaev IS (2010) **Properties of partially purified endopolyphosphatase of the yeast Saccharomyces cerevisiae** *Biochemistry Mosc* **75**:1404–1407 [Google Scholar](#)

Liu W, Wang J, Comte-Miserez V, Zhang M, Yu X, Chen Q, Jessen HJ, Mayer A, Wu S, Ye S (2023) **Cryo-EM structure of the polyphosphate polymerase VTC reveals coupling of polymer synthesis to membrane transit** *Embo J* [Google Scholar](#)

Lonetti A, Szigyarto Z, Bosch D, Loss O, Azevedo C, Saiardi A (2011) **Identification of an evolutionarily conserved family of inorganic polyphosphate endopolyphosphatases** *J Biol Chem* **286**:31966–74 [Google Scholar](#)

- Long H, Fang J, Ye L, Zhang B, Hui C, Deng X, Merchant SS, Huang K (2023) **Structural and functional regulation of Chlamydomonas lysosome-related organelles during environmental changes** *Plant Physiol* [Google Scholar](#)
- McCarthy L, Downey M (2023) **The emerging landscape of eukaryotic polyphosphatases** *Febs Lett* [Google Scholar](#)
- Michaillat L, Baars TL, Mayer A (2012) **Cell-free reconstitution of vacuole membrane fragmentation reveals regulation of vacuole size and number by TORC1** *Mol Biol Cell* **23**:881–895 [Google Scholar](#)
- Müller J, Westenberg B, Boller T, Wiemken A (1992) **Synthesis and degradation of polyphosphate in the fission yeast Schizosaccharomyces pombe: mutations in phosphatase genes do not affect polyphosphate metabolism** *FEMS Microbiol Lett* **71**:151–156 [Google Scholar](#)
- Müller O, Bayer MJ, Peters C, Andersen JS, Mann M, Mayer A (2002) **The Vtc proteins in vacuole fusion: coupling NSF activity to V(0) trans-complex formation** *EMBO J* **21**:259–269 [Google Scholar](#)
- Müller O, Neumann H, Bayer MJ, Mayer A (2003) **Role of the Vtc proteins in V-ATPase stability and membrane trafficking** *J Cell Sci* **116**:1107–1115 [Google Scholar](#)
- Neef DW, Kladde MP (2003) **Polyphosphate loss promotes SNF/SWI- and Gcn5-dependent mitotic induction of PHO5** *Mol Cell Biol* **23**:3788–3797 [Google Scholar](#)
- Nicolay K, Scheffers WA, Bruinenberg PM, Kaptein R (1982) **Phosphorus-31 nuclear magnetic resonance studies of intracellular pH, phosphate compartmentation and phosphate transport in yeasts** *Arch Microbiol* **133**:83–89 [Google Scholar](#)
- Nicolay K, Scheffers WA, Bruinenberg PM, Kaptein R (1983) **In vivo 31P NMR studies on the role of the vacuole in phosphate metabolism in yeasts** *Arch Microbiol* **134**:270–275 [Google Scholar](#)
- Okorokov LA, Lichko LP, Kulaev IS (1980) **Vacuoles: main compartments of potassium, magnesium, and phosphate ions in Saccharomyces carlsbergensis cells** *J Bacteriol* **144**:661–5 [Google Scholar](#)
- O'Neill EM, Kaffman A, Jolly ER, O'Shea EK (1996) **Regulation of PHO4 nuclear localization by the PHO80-PHO85 cyclin-CDK complex** *Science* **271**:209–212 [Google Scholar](#)
- Pavlovic I, Thakor DT, Bigler L, Wilson MSC, Laha D, Schaaf G, Saiardi A, Jessen HJ (2015) **Prometabolites of 5-Diphospho-myo-inositol Pentakisphosphate** *Angew Chem Int Ed Engl* **54**:9622–9626 [Google Scholar](#)
- Pinson B, Merle M, Franconi J-M, Daignan-Fornier B (2004) **Low affinity orthophosphate carriers regulate PHO gene expression independently of internal orthophosphate concentration in Saccharomyces cerevisiae** *J Biol Chem* **279**:35273–35280 [Google Scholar](#)
- Pipercevic J, Kohl B, Gerasimaite R, Comte-Miserez V, Hostachy S, Müntener T, Agustoni E, Jessen HJ, Fiedler D, Mayer A, et al (2023) **Inositol pyrophosphates activate the vacuolar transport chaperone complex in yeast by disrupting a homotypic SPX domain interaction** *Nat Commun* **14**:2645 [Google Scholar](#)

Pisoni RL (1991) **Characterization of a phosphate transport system in human fibroblast lysosomes** *J Biol Chem* **266**:979–985 [Google Scholar](#)

Pisoni RL, Lindley ER (1992) **Incorporation of [32P]orthophosphate into long chains of inorganic polyphosphate within lysosomes of human fibroblasts** *J Biol Chem* **267**:3626–3631 [Google Scholar](#)

Plouviez M, Fernández E, Grossman AR, Sanz-Luque E, Sells M, Wheeler D, Guieysse B (2021) **Responses of *Chlamydomonas reinhardtii* during the transition from P-deficient to P-sufficient growth (the P-overplus response): The roles of the vacuolar transport chaperones and polyphosphate synthesis** *J Phycol* **57**:988–1003 [Google Scholar](#)

Pondugula S, Neef DW, Voth WP, Darst RP, Dhasarathy A, Reynolds MM, Takahata S, Stillman DJ, Kladde MP (2009) **Coupling phosphate homeostasis to cell cycle-specific transcription: mitotic activation of *Saccharomyces cerevisiae* PHO5 by Mcm1 and Forkhead proteins** *Mol Cell Biol* **29**:4891–4905 [Google Scholar](#)

Potapenko E, Cordeiro CD, Huang G, Docampo R (2019) **Pyrophosphate Stimulates the Phosphate-Sodium Symporter of *Trypanosoma brucei* Acidocalcisomes and *Saccharomyces cerevisiae* Vacuoles** *mSphere* **4**:57 [Google Scholar](#)

Potapenko E, Cordeiro CD, Huang G, Storey M, Wittwer C, Dutta AK, Jessen HJ, Starai VJ, Docampo R (2018) **5-Diphosphoinositol Pentakisphosphate (5-IP7) Regulates Phosphate Release from Acidocalcisomes and Yeast Vacuoles** *J Biol Chem: jbc.RA* **118**:005884 [Google Scholar](#)

Qiu D, Eisenbeis VB, Saiardi A, Jessen HJ (2021) **Absolute Quantitation of Inositol Pyrophosphates by Capillary Electrophoresis Electrospray Ionization Mass Spectrometry** *J Vis Exp* [Google Scholar](#)

Qiu D, Gu C, Liu G, Ritter K, Eisenbeis VB, Bittner T, Gruzdev A, Seidel L, Bengsch B, Shears SB, et al (2023) **Capillary electrophoresis mass spectrometry identifies new isomers of inositol pyrophosphates in mammalian tissues** *Chem Sci* **14**:658–667 [Google Scholar](#)

Qiu D, Wilson MS, Eisenbeis VB, Harmel RK, Riemer E, Haas TM, Wittwer C, Jork N, Gu C, Shears SB, et al (2020) **Analysis of inositol phosphate metabolism by capillary electrophoresis electrospray ionization mass spectrometry** *Nat Commun* **11**:6035 [Google Scholar](#)

Ruiz FA, Marchesini N, Seufferheld M, Govindjee & Docampo R (2001) **The polyphosphate bodies of *Chlamydomonas reinhardtii* possess a proton-pumping pyrophosphatase and are similar to acidocalcisomes** *J Biol Chem* **276**:46196–46203 [Google Scholar](#)

Sanz-Luque E, Saroussi S, Huang W, Akkawi N, Grossman AR (2020) **Metabolic control of acclimation to nutrient deprivation dependent on polyphosphate synthesis** *Sci Adv* **6**:eabb5351 [Google Scholar](#)

Sattler T, Mayer A (2000) **Cell-free reconstitution of microautophagic vacuole invagination and vesicle formation** *J Cell Biol* **151**:529–538 [Google Scholar](#)

Scott DA, Docampo R, Dvorak JA, Shi S, Leapman RD (1997) **In situ compositional analysis of acidocalcisomes in *Trypanosoma cruzi*** *Journal of Biological Chemistry* **272**:28020–28029 [Google Scholar](#)

- Sethuraman A, Rao NN, Kornberg A (2001) **The endopolyphosphatase gene: essential in *Saccharomyces cerevisiae*** *PNAS* **98**:8542–8547 [Google Scholar](#)
- Shi X, Kornberg A (2005) **Endopolyphosphatase in *Saccharomyces cerevisiae* undergoes post-translational activations to produce short-chain polyphosphates** *FEBS Lett* **579**:2014–2018 [Google Scholar](#)
- Shirahama K, Yazaki Y, Sakano K, Wada Y, Ohsumi Y (1996) **Vacuolar Function in the Phosphate Homeostasis of the Yeast *Saccharomyces cerevisiae*** *Plant Cell Physiol* **37**:1090–1093 [Google Scholar](#)
- Spain DCB Facultat de Medicina i Ciències de la Salut Universitat Internacional de Catalunya, Barcelona Catalonia, Jimenez J, Bru S, Ribeiro M, Clotet J (2015) **Live fast, die soon: cell cycle progression and lifespan in yeast cells** *Microb Cell* **2**:62 [Google Scholar](#)
- Theobald U, Mohns J, Rizzi M (1996) **Determination of in-vivo cytoplasmic orthophosphate concentration in yeast** *Biotechnol Tech* **10**:297–302 [Google Scholar](#)
- Thomas MR, O'Shea EK (2005) **An intracellular phosphate buffer filters transient fluctuations in extracellular phosphate levels** *Proc Natl Acad Sci USA* **102**:9565–9570 [Google Scholar](#)
- Ulrich PN, Lander N, Kurup SP, Reiss L, Brewer J, Soares-Medeiros LC, Miranda K, Docampo R (2013) **The Acidocalcisome Vacuolar Transporter Chaperone 4 Catalyzes the Synthesis of Polyphosphate in Insect-stages of *Trypanosoma brucei* and *T. cruzi*** *J Eukaryot Microbiol* **61**:155–165 [Google Scholar](#)
- Urech K, Dürr M, Boller T, Wiemken A, Schwencke J (1978) **Localization of Polyphosphate in Vacuoles of *Saccharomyces-Cerevisiae*** *Arch Microbiol* **116**:275–278 [Google Scholar](#)
- Vardi N, Levy S, Assaf M, Carmi M, Barkai N (2013) **Budding Yeast Escape Commitment to the Phosphate Starvation Program Using Gene Expression Noise** *Curr Biol* [Google Scholar](#)
- Vardi N, Levy S, Gurvich Y, Polacheck T, Carmi M, Jaitin D, Amit I, Barkai N (2014) **Sequential feedback induction stabilizes the phosphate starvation response in budding yeast** *Cell Rep* **9**:1122–1134 [Google Scholar](#)
- Wang C, Yue W, Ying Y, Wang S, Secco D, Liu Y, Whelan J, Tyerman SD, Shou H (2015) **Rice SPX-Major Facility Superfamily3, a Vacuolar Phosphate Efflux Transporter, Is Involved in Maintaining Phosphate Homeostasis in Rice** *Plant Physiol* **169**:2822–2831 [Google Scholar](#)
- Wang H, Godage HY, Riley AM, Weaver JD, Shears SB, Potter BVL (2014) **Synthetic Inositol Phosphate Analogs Reveal that PPIP5K2 Has a Surface-Mounted Substrate Capture Site that Is a Target for Drug Discovery** *Chem Biol* **21**:689–699 [Google Scholar](#)
- Warner JR (1999) **The economics of ribosome biosynthesis in yeast** *Trends Biochem Sci* **24**:437–440 [Google Scholar](#)
- Westenberg B, Boller T, Wiemken A (1989) **Lack of arginine-storage and polyphosphate-storage pools in a vacuole-deficient mutant (end1) of *Saccharomyces-cerevisiae*** *FEBS Lett* **254**:133–136 [Google Scholar](#)
- Wild R, Gerasimaite R, Jung J-Y, Truffault V, Pavlovic I, Schmidt A, Saiardi A, Jessen HJ, Poirier Y, Hothorn M, et al (2016) **Control of eukaryotic phosphate homeostasis by inositol**

polyphosphate sensor domains *Science* **352**:986–990 [Google Scholar](#)

Wilson MSC, Bulley SJ, Pisani F, Irvine RF, Saiardi A (2015) **A novel method for the purification of inositol phosphates from biological samples reveals that no phytate is present in human plasma or urine** *Open Biol* **5**:150014 [Google Scholar](#)

Wykoff DD, O’Shea EK (2001) **Phosphate transport and sensing in *Saccharomyces cerevisiae*** *Genetics* **159**:1491–1499 [Google Scholar](#)

Wykoff DD, Rizvi AH, Raser JM, Margolin B, O’Shea EK (2007) **Positive feedback regulates switching of phosphate transporters in *S. cerevisiae*** *Mol Cell* **27**:1005–1013 [Google Scholar](#)

Xu C, Xu J, Tang H-W, Ericsson M, Weng J-H, DiRusso J, Hu Y, Ma W, Asara JM, Perrimon N (2023) **A phosphate-sensing organelle regulates phosphate and tissue homeostasis** *Nature* **617**:1–9 [Google Scholar](#)

Zhang J, Sassen T, Pierick A ten, Ras C, Heijnen JJ, Wahl SA (2015) **A fast sensor for in vivo quantification of cytosolic phosphate in *Saccharomyces cerevisiae*** *Biotechnol Bioeng* **112**:1033–1046 [Google Scholar](#)

Zúñiga-Burgos T, Saiardi A, Camargo-Valero MA, Baker A (2024) **Phosphate overplus response in *Chlamydomonas reinhardtii*: Polyphosphate dynamics to monitor phosphate uptake and turnover** *Algal Res* **81**:103589 [Google Scholar](#)

Lydie Michailat Mayer (2025) **Enrichment of proteins in purified *Saccharomyces cerevisiae* vacuoles relative to a whole cell extract**. PRIDE ID PXD060102 <https://www.ebi.ac.uk/pride/archive/projects/PXD060102>

Author information

Samuel Bru*

Département d’immunobiologie, Université de Lausanne, Epalinges, Switzerland

*Present address: Department of Biomedical Sciences, Faculty of Medicine and Health Sciences, Universitat Internacional de Catalunya. 08195 Barcelona, Spain.

Lydie Michailat Mayer

Département d’immunobiologie, Université de Lausanne, Epalinges, Switzerland

Geun-Don Kim

Département d’immunobiologie, Université de Lausanne, Epalinges, Switzerland

Danye Qiu

Institute of Organic Chemistry, University of Freiburg, Freiburg, Germany, CIBSS – Centre for Integrative Biological Signalling Studies, University of Freiburg, Freiburg, Germany

Henning Jacob Jessen

Institute of Organic Chemistry, University of Freiburg, Freiburg, Germany
ORCID iD: [0000-0002-1025-9484](https://orcid.org/0000-0002-1025-9484)

Andreas Mayer

Département d'immunobiologie, Université de Lausanne, Epalinges, Switzerland
ORCID iD: [0000-0001-6131-313X](https://orcid.org/0000-0001-6131-313X)

For correspondence: andreas.mayer@unil.ch

Editors

Reviewing Editor

Johannes Herrmann

University of Kaiserslautern, Kaiserslautern, Germany

Senior Editor

Felix Campelo

Universitat Pompeu Fabra, Barcelona, Spain

Reviewer #1 (Public review):

The manuscript by Bru et al. focuses on the role of vacuoles as a phosphate buffering system for yeast cells. The authors describe here the crosstalk between the vacuole and the cytosol using a combination of in vitro analyses of vacuoles and in vivo assays. They show that the luminal polyphosphatases of the vacuole can hydrolyze polyphosphates to generate inorganic phosphate, yet they are inhibited by high concentrations. This balances the synthesis of polyphosphates against the inorganic phosphate pool. Their data further show that the Pho91 transporter provides a valve for the cytosol as it gets activated by a decline in inositol pyrophosphate levels. The authors thus demonstrate how the vacuole functions as a phosphate buffering system to maintain a constant cytosolic inorganic phosphate pool.

This is a very consistent and well-written manuscript with a number of convincing experiments, where the authors use isolated vacuoles and cellular read-out systems to demonstrate the interplay of polyphosphate synthesis, hydrolysis, and release. The beauty of this system the authors present is the clear correlation between product inhibition and the role of Pho91 as a valve to release Pi to the cytosol to replenish the cytosolic pool. I find the paper overall an excellent fit.

Comments on Revision:

The authors have addressed all my concerns.

<https://doi.org/10.7554/eLife.108181.2.sa2>

Reviewer #3 (Public review):

Bru et al. investigated how inorganic phosphate (Pi) is buffered in cells using *S. cerevisiae* as a model. Pi is stored in cells in the form of polyphosphates in acidocalcisomes. In *S. cerevisiae*, the vacuole, which is the yeast lysosome, also fulfills the function of Pi storage organelle. Therefore, yeast is an ideal system to study Pi storage and mobilization.

They can recapitulate in their previously established system, using isolated yeast vacuoles, findings from their own and other groups. They integrate the available data and propose a working model of feedback loops to control the level of Pi on the cellular level.

This is a solid study, in which the biological significance of their findings is not entirely clear. The data analysis and statistical significance need to be improved and included, respectively.

The manuscript would have benefited from rigorously testing the model, which would also have increased the impact of the study.

<https://doi.org/10.7554/eLife.108181.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public review):

The manuscript by Bru et al. focuses on the role of vacuoles as a phosphate buffering system for yeast cells. The authors describe here the crosstalk between the vacuole and the cytosol using a combination of in vitro analyses of vacuoles and in vivo assays. They show that the luminal polyphosphatases of the vacuole can hydrolyse polyphosphates to generate inorganic phosphate, yet they are inhibited by high

concentrations. This balances the synthesis of polyphosphates against the inorganic phosphate pool. Their data further show that the Pho91 transporter provides a valve for the cytosol as it gets activated by a decline in inositol pyrophosphate levels. The authors thus demonstrate how the vacuole functions as a phosphate buffering system to maintain a constant cytosolic inorganic phosphate pool.

This is a very consistent and well-written manuscript with a number of convincing experiments, where the authors use isolated vacuoles and cellular read-out systems to demonstrate the interplay of polyphosphate synthesis, hydrolysis, and release. The beauty of this system the authors present is the clear correlation between product inhibition and the role of Pho91 as a valve to release Pi to the cytosol to replenish the cytosolic pool. I find the paper overall an excellent fit and only have a few issues, including:

(1) Figure 3: The authors use in their assays 1 mM ZnCl₂ or 1mM MgCl₂. Is this concentration in the range of the vacuolar luminal ion concentration? Did they also test the effect of Ca²⁺, as this ion is also highly concentrated in the lumen?

The concentrations inside vacuoles reach those values. However, given that polyP can chelate divalent metal ions, what would matter are the concentrations of free Zn²⁺ or Mg²⁺ inside the organelle. These are not known. This is not critical since we use those two conditions only as a convenient tool to differentiate Ppn1 and Ppn2 activity in vitro. In our initial characterisation of Ppn2 (10.1242/jcs.201061), we had also tested Mn, Co, Ca, Ni, Cu. Only Zn and Co supported activity. Ca did not. Andreeva et al. (10.1016/j.biochi.2019.06.001) reached similar conclusions and extended our results.

(2) Regarding the concentration of 30 mM K-PI, did the authors also use higher and lower concentrations? I agree that there is inhibition by 30 mM, but they cannot derive conclusions on the luminal concentration if they use just one in their assay. A titration is necessary here.

The concentration of 30 mM was not chosen arbitrarily. It is the luminal Pⁱ concentration that the vacuoles reached through polyP synthesis and hydrolysis when they entered a plateau of luminal Pⁱ. We consider this as an upper limit because polyP kept increasing which luminal Pⁱ did not. Thus, there is no physiological motivation for trying higher values. We have nevertheless added a titration to the revised version (new Fig. 3A).

(3) What are the consequences on vacuole morphology if the cells lack Pho91?

We had not observed significant abnormalities during a screen of the genome-wide deletion collection of yeast (10.1371/journal.pone.0054160), nor in other experiments with *pho91* mutants, which we have not included in this manuscript due to a lack of effect.

(4) Discussion: The authors do not refer to the effect of calcium, even though I would expect that the levels of the counterion should affect the phosphate metabolism. I would appreciate it if they would extend their discussion accordingly.

The situation is much more complex because Ca^{2+} is not the only counterion. Major pools of counterions (up to hundreds of mM) are constituted by vacuolar lysine, arginine, polyamines, Mg, Zn etc. Their interplay with polyP is probably complex and worth to be treated in a dedicated project. If we wanted to limit the discussion of this complexity not to the simple statement that it is not understood, which is not very useful, we would have to engage in a lot of speculation. We feel that this would make the discussion lose focus and not contribute concrete insights.

(5) I would appreciate a brief discussion on how phosphate sensing and control are done in human cells. Do they use a similar lysosomal buffer system?

Mammalian cells have their Pi exporter XPR1 mainly on a lysosome-like compartment (10.1016/j.celrep.2024.114316). Whether and how it functions there for Pi export from the cytosol is not entirely clear. We have addressed this situation in the revised discussion section.

Reviewer #2 (Public review):

Summary:

This manuscript presents a well-conceived and concise study that significantly advances our understanding of polyphosphate (polyP) metabolism and its role in cytosolic phosphate (Pi) homeostasis in a model unicellular eukaryote. The authors provide evidence that yeast vacuoles function as dynamic regulatory buffers for Pi homeostasis, integrating polyP synthesis, storage, and hydrolysis in response to cellular metabolic demands. The work is methodologically sound and offers valuable insights into the conserved mechanisms of phosphate regulation across eukaryotes.

Strengths:

The results demonstrate that the vacuolar transporter chaperone (VTC) complex, in conjunction with luminal polyphosphatases (Ppn1/Ppn2) and the Pi exporter Pho91, establishes a finely tuned feedback system that balances cytosolic Pi levels. Under Pi-replete conditions, inositol pyrophosphates (InsPPs) promote polyP synthesis and storage while inhibiting polyP hydrolysis, leading to vacuolar Pi accumulation.

Conversely, Pi scarcity triggers InsPP depletion, activating Pho91-mediated Pi export and polyP mobilization to sustain cytosolic phosphate levels. This regulatory circuit ensures metabolic flexibility, particularly during critical processes such as glycolysis, nucleotide synthesis, and cell cycle progression, where phosphate demand fluctuates dramatically.

From my viewpoint, one of the most important findings is the demonstration that vacuoles act as a rapidly accessible Pi reservoir, capable of switching between storage (as polyP) and release (as free Pi) in response to metabolic cues. The energetic cost of polyP synthesis-driven by ATP and the vacuolar proton gradient-highlights the evolutionary importance of this buffering system. The study also draws parallels between yeast vacuoles and acidocalcisomes in other eukaryotes, such as Trypanosoma and

Chlamydomonas, suggesting a conserved role for these organelles in phosphate homeostasis.

Weaknesses:

While the manuscript is highly insightful, referring to yeast vacuoles as "acidocalcisome-like" may warrant further discussion. Canonical acidocalcisomes are structurally and chemically distinct (e.g., electrondense, in most cases spherical, and not routinely subjected to morphological changes, and enriched with specific ions), whereas yeast vacuoles have well-established roles beyond phosphate storage. A comment on this terminology could strengthen the comparative analysis and avoid potential confusion in the field.

Yeast vacuoles show all major chemical features of acidocalcisomes. They are acidified, contain high concentrations of Ca, polyP (which make them electron-dense, too), other divalent ions, such as Mg, Zn, Mn etc, and high concentrations of basic amino acids. Thus, they clearly have an acidocalcisome-like character. In addition, they have hydrolytic, lysosomelike functions and, depending on the strain background, they can be larger than acidocalcisomes described e.g. in protists. We have elaborated on this point in the introduction of the revised version.

Reviewer #3 (Public review):

*Bru et al. investigated how inorganic phosphate (Pi) is buffered in cells using *S. cerevisiae* as a model. Pi is stored in cells in the form of polyphosphates in acidocalcisomes. In *S. cerevisiae*, the vacuole, which is the yeast lysosome, also fulfills the function of Pi storage organelle. Therefore, yeast is an ideal system to study Pi storage and mobilization.*

They can recapitulate in their previously established system, using isolated yeast vacuoles, findings from their own and other groups. They integrate the available data and propose a working model of feedback loops to control the level of Pi on the cellular level.

This is a solid study, in which the biological significance of their findings is not entirely clear. The data analysis and statistical significance need to be improved and included, respectively. The manuscript would have benefited from rigorously testing the model, which would also have increased the impact of the study.

It is not clear to us what the reviewer would see as a more rigorous test of the model.

Reviewer #1 (Recommendations for the authors):

(1) Figure 2: Why do the authors label the blue curve in A and B as BY and in C and D as WT? Is this a different genetic background they used here? This should be specified in the legend.

No, it is the same background. The figures had been reshuffled before submission and we overlooked to replace "BY" by "WT". This has been corrected. Now we consistently use WT in all figures

(2) Figure 4 has different scaling for the two panels, which should be labeled as A and B. I am aware that the authors do this for comparison, but it is rather confusing at first glance. I recommend having them at the same scale.

We chose this representation on two separate scales because this figure shall primarily illustrate that the shift between pho91^Δ and WT curves vanishes in the presence of IP7. We

now highlight in the figure legend that the scales are different to avoid confusion.

(3) *Figure 8: I would appreciate a model with normal and low Pi concentrations in comparison, as this is what the authors worked out.*

We have modified the figure. It now compares Pi-rich and Pi-limited scenarios.

(4) *Minor issue: Wouldn't it make more sense to show the molar concentration in the Figures rather than the nmol of Pi/ug of protein? I am aware that this would require information on the vacuole volume rather than the reaction volume, and the authors do this calculation later on.*

It depends. We often chose this representation because it illustrates the price to pay (metabolic input in terms of protein that must be dedicated to this task) to sequester a certain quantity of P^i . But, as we provide the corresponding P^i concentration in the text, this information is accessible to the reader, too.

Reviewer #2 (Recommendations for the authors):

As stated above in the weaknesses section, while functional parallels exist, canonical acidocalcisomes are structurally and chemically distinct, typically smaller, electron-dense, and enriched with cations. Whereas yeast vacuoles are larger, multifunctional organelles with well-established roles beyond phosphate storage. Explicitly addressing these differences would strengthen the comparative framework and prevent potential confusion in interpreting the evolutionary relationships between these organelles.

We agree to some degree, which is the reason why we refer to vacuoles as acidocalcisome-like organelles. In fact, vacuoles share virtually all defining chemical traits of acidocalcisomes. They just have a second functional domain as hydrolytic, lysosome-like organelles. Given the plasticity of endo-lysosomal compartments, and acidocalcisomes belong to this group because of their biogenesis through the AP3 pathway, this is not shocking to us. But the reviewer's comment made us realize that it is better to explicitly address this point. We have added a section to the introduction to do this.

Reviewer #3 (Recommendations for the authors):

(1) *Page 8: It is unclear why the authors only estimated the Pi concentration in wild-type vacuoles. This should also be done for vacuoles from other strains.*

This information is inherent in Figure 2. PolyP hyperaccumulating strains show the same plateau as the wildtype, meaning that they also reach around 30 mM luminal Pi concentration, whereas vtc4^Δ vacuoles reach only around 1/10th of that increase, indicating that they remain at 3 mM. We mention this now in the text.

(2) *The attempts of the localization of Pho91 through tagging are not satisfactory. The author described different localizations for Pho91 depending on whether it was tagged on the N- or C-terminus or when Nterminally tagged and overexpressed using two strong promoters. While it is not uncommon that proteins show different localization patterns, depending on where the tag is inserted, it is possible that one of the tags would reflect the localization of the endogenous protein. There is an easy way to test this, in particular when Pho91 is endogenously tagged. pho91Δ has reported phenotypes such as abnormal vacuolar morphology or increased autophagy. They could also measure PI content in vacuoles. The authors could compare the phenotypes of the endogenously tagged strains with WT and a pho91Δ strain.*

Indeed, the attempts to localise the protein through fluorescent tags are unsatisfactory, in our hands as in the hands of others. We would not have created a series of many different tagged versions (we present only a selection of these in the manuscript) if the creation of a faithful reporter for Pho91 localisation were so straightforward. Expression from the endogenous promoter yields quite low signals (which is why others have overexpressed their GFP fusion from strong promoters). But overexpression brings at least a significant part of the protein to the cell surface, where it can then function as Pi importer and suffice to restore much of the maximal Pi uptake capacity that genuine plasma membrane transporters provide and support normal growth of the cells (Wykoff & O'Shea, 2001). But the localisation pattern of Pho91-GFP, likewise overexpressed from a strong promoter, does not reflect this plasma membrane localisation (see the references that the reviewer mentioned under (3)). The published overexpressed GFP-fusions localise only to the vacuole, suggesting that even in this case the GFP tag may create an artefact. Therefore, we went through a large variety of Pho91 gene fusions, which led us to the conclusion that the protein is very sensitive to tags at both ends and that fusion proteins hence are unlikely to reliably report the correct location of the protein. Given this, we resorted to quantitative proteomics to clarify the issue. This quantitative experiment goes beyond previously published proteomics analyses that the reviewer mentions under (3), which found the protein in the vacuolar fraction but did not calculate the enrichment factors, which is crucial.

A strong phenotype of abnormal vacuolar morphology is not apparent in our cultures.

(3) Moreover, Pho91 has been identified as a component enriched in vacuolar-mitochondria contact sites (vCLAMP), and this localization was confirmed with GFP-Pho91 (PMID: 25026036). Likewise, PMID: 35175277 also detected Pho91 by mass spectrometry as a vacuolar protein and showed endogenously tagged GFP-Pho91 on the vacuole (co-staining with Vph1). The authors may request the strains from the authors of these papers and use them for their experiments. PMID: 17804816, the oldest of the three reports (from 2007) reports a GFP-Pho91 under either TEF or ADH promoter that localizes to the vacuole. They also showed that the fusion protein is functional. These and other experiments led them to conclude that Pho91 exports phosphate from the vacuolar lumen to the cytoplasm.

We have now included these references. As argued above, we have analysed also the strains from PMID17804816. The observed clear localisation of the fusion protein to vacuoles is only visible upon overexpression, not upon expression from the endogenous locus. Apparently also this construct is unlikely to report Pho91 localisation reliably (though, by chance, overexpression leads it to the correct location). Thus, we maintain our conclusion that C- or N-terminally GFP-tagged versions of Pho91 are unreliable tools for localising the protein.

(4) The impact of pho91Δ on Pho4-GFP nuclear localization is modest at best (increase from 5% of cells showing Pho4-GFP in the nucleus in WT vs 10% in pho91Δ), and only somewhat stronger in ppn1Δ/ppn2Δ. This means 90% of pho91Δ cells do not respond, and Pho4-GFP stays cytoplasmic. It is unclear how the author can derive a meaningful conclusion from these data. Moreover, are these data really supporting the model, or do these data rather indicate that there are additional factors/pathways needed? What is the biological significance of the marginal increase from 5% to 10% of cells that would respond? What happens to the cells that cannot respond? Will they die or at least have a growth disadvantage? It would be useful to provide some functional studies.

We should have explained the nature of the assay better. The experiment exploits the fact that dividing yeast cells transiently fall into a state of Pi scarcity during S-phase. Since S-phase is less than a quarter of the cell cycle, only a small fraction of the cells transiently activates the PHO pathway. These cannot be well characterised by ensemble assays, but microscopy

circumvents this background of the whole population and picks them up very clearly, allowing to quantify them. We have adapted the respective chapter in the results section to improve the description of this experiment.

(5) The quantification of the data is suboptimal, as in most assays the mean and standard error of the mean (SEM) are given. SEM is not really appropriate in these cases because it gives only the error of the mean and not of the entire data. Therefore, the standard deviation (SD) is needed, which reports on the variability of the data, and which is usually much larger than the SEM. Using the SD, would also allow the authors to do proper statistical analysis, which is missing entirely in this manuscript.

SEM also comprises the variability of the data. It is linked with the SD ($SEM = SD / \sqrt{n}$), but SEM also considers the number of the experiments n . The main goal is to compare the means, and SEM is an appropriate and frequently used tool for this because it illustrates how well the arithmetic mean may estimate the true mean of the population. Therefore, we kept the SEM but have added tests of significance for the differences shown.

(6) Statistical testing in Figure 7 is essential as the effects are very small. Again, are these changes big enough for a biologically meaningful response? The authors should at least discuss this.

Our previous time course analyses of InsPP dynamics, performed under comparable conditions as in this study, showed that InsP8 decreases by around 50% in the first 30 min after transfer to Pi starvation (DOI: <https://doi.org/10.7554/eLife.87956>) and that this decline is already sufficient to trigger the PHO starvation program, as assessed by Pho4-GFP translocation into the nucleus. Thus, a 50% decrease, which is observed in *ppn1* *ppn2* mutants, is functionally significant. We have now also evaluated statistical significance in Fig. 7, which is given for the 50% reduction of InsP8 and 1-InsP7 in *ppn1* *ppn2*.

Minor points:

(1) There are a number of smaller edits (use of italic or better the absence thereof, lacking information in the reference list, and some typos).

Thank you. We have corrected those.

(2) The exact n should be given in the Figure legend.

Corrected.

(3) Page 8, line 8: it would be nice to have a picture of the wild-type vacuoles and what you measured.

We now present a sample image in the new Suppl. Fig. 1.

(4) PMID: 11779791 showed already that Pho91 cannot rescue the absence of the plasma membrane Pi transporters. This study should be at least cited.

This is not quite correct. The study that the reviewer mentions showed that Pho91 supports slower growth and the authors concluded that "A synthetic lethal phenotype was observed when (all) five phosphate transporters were inactivated...". We had cited the same group and the same first author, just using their later study (Wykoff et al., 2007) that had recapitulated the results from PMID11779791 and showed in addition quite good growth of the PHO91 expressing strain on YPD (Suppl. Fig. 2). We had obtained the strains from this group. In

reproducing their experiments, we noticed that the growth of Pho91 that these authors had observed is due to incomplete repression of Pho84. They had overexpressed Pho84 from a galactose inducible promotor to generate a background with a regulatable Pi transporter. This trick allowed them to conveniently manipulate the strain and reduce (but not abolish) Pho84 expression by transferring the cells from galactose to glucose for their experiments. Therefore, we chose a more rigorous plasmid shuffling strategy to test the individual P_i transporter, which allows an assessment without the leaky background expression of Pho84 on glucose. In contrast to O'Shea and colleagues, we observed zero growth of a strain expressing only PHO91. We have revised the results section to make this discrepancy more evident and provide a better motivation for our experiment.

| (5) *It would be nice to see the actual data in Figure 6; not only a quantification.*

We illustrate the phenotype of nuclear Pho4-GFP in panel A. Showing all the images necessary to appreciate the differences between the strains would require including many dozens of images into the figure, which would not be useful.

<https://doi.org/10.7554/eLife.108181.2.sa0>