

Pronounced expression of extracellular matrix proteoglycans regulated by Wnt pathway underlies the parallel evolution of lip hypertrophy in East African cichlids

Reviewed Preprint

v2 • March 21, 2025

Revised by authors

Reviewed Preprint

v1 • September 5, 2024

Nagatoshi Machii, Ryo Hatashima, Tatsuya Niwa, Hideki Taguchi, Ismael A Kimirei, Hillary DJ Mrosso, Mitsuto Aibara, Tatsuki Nagasawa, Masato Nikaido 

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan • Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, Yokohama, Japan • Tanzania Fisheries Research Institute (TAFIRI), Dar es Salaam, United Republic of Tanzania • Tanzania Fisheries Research Institute (TAFIRI), Mwanza Fisheries Research Center, Mwanza, United Republic of Tanzania

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

Cichlid fishes have attracted attention from a wide range of biologists because of their extensive species diversification at the ecological and phenotypic levels. In this **important** study, the authors have partially revealed the mechanism behind lip thickening in cichlid fishes, which has evolved independently across three lakes in Africa. To explore this phenomenon, the authors used histological comparison, proteomics, and transcriptomics, all of which are well suited for their objectives. With **compelling** evidence, this contribution provides insights into parallel evolution in polygenic traits and holds significant value for the field.

<https://doi.org/10.7554/eLife.99160.2.sa2>

Abstract

Cichlid fishes inhabiting the East African Great Lakes, Victoria, Malawi, and Tanganyika, are textbook examples of parallel evolution, as they have acquired similar traits independently in each of the three lakes during the process of adaptive radiation. In particular, “hypertrophied lip” has been highlighted as a prominent example of parallel evolution. However, the underlying molecular mechanisms remain poorly understood. In this study, we conducted an integrated comparative analysis between the hypertrophied and normal lips of cichlids across three lakes based on histology, proteomics, and transcriptomics. Histological and proteomic analyses revealed that the hypertrophied lips were characterized by enlargement of the proteoglycan-rich layer, in which versican and periostin proteins were abundant. Transcriptome analysis revealed that the expression of extracellular matrix-related genes, including collagens, glycoproteins and proteoglycans, was higher in hypertrophied lips,

regardless of their phylogenetic relationships. In addition, the genes in Wnt signaling pathway, which is involved in promoting proteoglycan expression, was highly expressed in both the juvenile and adult stages of hypertrophied lips. Our comprehensive analyses showed that hypertrophied lips of the three different phylogenetic origins can be explained by similar proteomic and transcriptomic profiles, which may provide important clues into the molecular mechanisms underlying phenotypic parallelisms in East African cichlids.

Introduction

Parallel evolution is evidence of natural selection that drives adaptive solutions to the environment (Losos, 2011). Acquisition of similar traits in distantly related lineages can be observed in multiple lineages, such as the loss of eyes in cave fish, patterns of wing colors in *Heliconius* butterflies, and body armor of sticklebacks (Krishnan and Rohner, 2017; Stern, 2013). Since it is unlikely that such a complex phenotype was acquired repeatedly by neutral evolution alone, the same direction of natural selection among species is likely to drive the parallel phenotypic change (Losos, 2011; Stern, 2013). Therefore, research on parallel evolution enables us to compare the commonalities and differences in the mechanisms behind the acquisition of novel traits and elucidate how adaptation occurs in similar environments.

East African cichlids offer one of the best opportunities to examine the molecular basis of parallel evolution (Kocher et al., 1993; Muschick et al., 2012). There are hundreds of endemic cichlid species in the East African Great Lakes, Victoria, Malawi, and Tanganyika, as a result of explosive adaptive radiation (Salzburger, 2018; Santos et al., 2023). As a result of independent adaptations in each lake, parallel evolution is often observed among lakes, such as that of the nose snout (Duenser et al., 2023), jaw morphology, and body shape (Kocher et al., 1993; Muschick et al., 2012). To reveal their evolutionary mechanisms, genetic bases of the phenotypes have been investigated. For example, the parallel evolution of body stripes in cichlids of Lake Victoria and Lake Malawi was explained by relatively simple changes in pigmental cell arrangement caused by mutations in the cis-regulatory element of agouti-related peptide 2 (*agrp2*) (Kratochwil et al., 2018). Although this parallel evolution of body stripe patterns can be explained by a single gene, most traits are likely to be polygenic, hindering elucidation of the parallel evolution mechanism.

Cichlid fishes have acquired a huge diversity in their oral morphologies, reflecting ecological and trophic specializations (Albertson and Kocher, 2006; Kocher, 2004; Lecaudey et al., 2021, 2019). Hypertrophied lips, one of the most prominent examples of oral diversity, have been independently acquired in cichlids inhabiting each of the East African Great Lakes, as well as in the Nicaraguan crater lakes and South American rivers (Henning et al., 2017; Kautt et al., 2020; Masonick et al., 2023, 2022). Several hypotheses have been proposed for the adaptive roles of hypertrophied lips, such as for effectively sucking prey out from narrow crevices (Lukas et al., 2015), reducing mechanical stress during foraging in rocky habitats (Fryer, 1972; Greenwood, 1974), and facilitating prey detection by providing an enlarged area for taste receptors (Arnégard and Snoeks, 2001; Oliver and Arnégard, 2010). Thus, hypertrophied lips are considered an adaptive trait for a specific trophic niche, making them an ideal model for understanding the mechanism of parallel evolution.

The molecular mechanisms underlying hypertrophied lips have been investigated in various lineages of cichlids. For example, transcriptome comparisons between hypertrophied and normal lips of cichlids in Lake Tanganyika (Colombo et al., 2013) and South American crater lakes (Manousaki et al., 2013) have identified some differentially expressed genes (DEGs) as candidates for lip hypertrophy; however, their primary regulators remain to be elucidated. In addition, quantitative trait locus (QTL) mapping (Henning et al., 2017) and genome-wide association study (GWAS) (Masonick et al., 2023) have been conducted to elucidate the genomic loci responsible for hypertrophied lips in the cichlids of Lake Victoria and Lake Malawi. These

genome-wide studies concluded that lip hypertrophy was a polygenic trait involving several candidate genes. However, the molecular mechanism underlying lip hypertrophy has not yet been elucidated.

In this study, we comprehensively compared the hypertrophied and normal lips of cichlids across all East African Great Lakes at various biological levels using histology, proteomics, and transcriptomics. As a result, we showed that a novel key pathway commonly involved in the formation of hypertrophied lips, providing insight into a better understanding of the molecular basis of a typical example of parallel evolution.

Materials and Methods

Animals

Three species of Victoria cichlids (*Haplochromis chilotes* with hypertrophied lips; *H. sauvagei*, and *H. pyrrhocephalus* with normal lips), four species of Malawi cichlids (*Placidochromis milomo* with hypertrophied lips; *Maylandia lombardoi*, *M. zebra*, and *Labidochromis caeruleus* with normal lips), and two species of Taganyika cichlids (*Lobochilotes labiatus* with hypertrophied lips; *Tropheus moorii* with normal lips) were used in the experiments. *H. chilotes*, *H. sauvagei*, and *H. pyrrhocephalus* were collected from the Mwanza gulf of Lake Victoria during an exploration that was led by the Nikaido Laboratory in 2018. The other fish were obtained from a commercial supplier. Fish were maintained and bred at 27°C on a 12/12 h light/dark cycle. One to 12 individuals were kept in a plastic tank (40 cm × 25 cm × 36 cm) and fed food pellets 2-3 times a day. The sample information is provided in Table S1. All experiments were conducted in accordance with the Institutional Animal Experiment Committee of the Tokyo Institute of Technology.

Histological observation

Cichlids were anesthetized with 0.02% ethyl p-aminobenzoate. The lower lips were then dissected, along with the lower jaw, and were fixed in 4% paraformaldehyde (PFA, Fujifilm-Wako, Chuo-ku, Osaka, Japan) in phosphate-buffered saline (PBS) at 4°C for 24 h. Tissues were demineralized in 0.34 M ethylenediaminetetraacetic acid (EDTA, Fujifilm-Wako, Chuo-ku, Osaka, Japan) in 100 mM, pH7.0 tris (hydroxymethyl) aminomethane (Tris, Fujifilm-Wako, Chuo-ku, Osaka, Japan) buffer at 4°C for 24 h. After demineralization, tissues were treated with 20% sucrose/PBS at 4°C for 24 h. Tissues were then embedded in Tissue-Tek O.C.T. compound (Sakura Finetek, Chuo-ku, Tokyo, Japan) and frozen with liquid nitrogen. Embedded blocks were sliced in the sagittal plane into 20 µm sections and placed on a coated glass slide (MAS-01, Matsunami, Bunkyo-ku, Tokyo, Japan). Sections were stored at −80°C until they were used for experiments. To observe the histological structure, hematoxylin–eosin (HE) staining, Alcian blue (AB) staining, and Van Gieson (VG) staining were conducted. Mayer's hematoxylin solution (Fujifilm-Wako, Chuo-ku, Osaka, Japan), Eosin Y solution (Fujifilm-Wako, Chuo-ku, Osaka, Japan), ALCIAN BLUE 8GX (Sigma-Aldrich, St. Louis, MO, USA), Van Gieson solution P (Fujifilm-Wako, Chuo-ku, Osaka, Japan), and Van Gieson solution F (Fujifilm-Wako, Chuo-ku, Osaka, Japan) were used for each staining. Experiments were performed according to the manufacturer's protocol and previous studies (Arnegard and Snoeks, 2001 [↗](#); Oliver and Arnegard, 2010 [↗](#)). For AB staining, the pH was adjusted to 2.5 with 3% acetic acid (Fujifilm-Wako, Chuo-ku, Osaka, Japan). For VG staining, Van Gieson solution P and Van Gieson solution F were prepared at a ratio of 7.55:1. The stained area was calculated using ImageJ version 1.5.1 (Schneider et al., 2012 [↗](#)) after trimming the image to include only the lips. The images were captured using a Zeiss Axioplan fluorescence microscope and a Zeiss Axiocam 503 color CCD camera (Carl Zeiss, Oberkochen, Germany). Images were level-corrected, contrast adjusted, and tiled using Adobe Photoshop CC 2022.

Chondroitinase digestion

Chondroitinase ABC from *Proteus vulgaris* (Sigma-Aldrich, St. Louis, MO, USA) was used for chondroitinase digestion. 2 mUnits/0.1% bovine serum albumin solution BSA in 20 mmol/L Tris-HCl pH 7.7 were prepared as the enzyme solution. In the experiment, the sections were first treated with 100 μ L of 0.1% BSA solution for 5 min. Next, 100 μ L of enzyme solution, or BSA solution for the control, was added. During the reaction, the sections were placed in a moist chamber. The enzymatic reaction was performed at 37°C for 1 h. After the reaction, the samples were washed with water for 15 min, and then AB staining was examined.

Identification of extracellular matrix (ECM)-related genes in cichlids

To compare the transcriptome of cichlid lips, cichlid extracellular matrix (ECM)-related genes were identified using the data from a previous study (Hynes and Naba, 2012 [DOI](#)). The amino acid sequences of human, mouse, and zebrafish ECM-related genes registered in Human GENCODE 41, Mouse GENCODE M30, Ensembl, and UniProtKB were used as queries, and those of *M. zebra* and *O. niloticus* were identified from their protein sequences registered in Ensembl and Uniprot using OrthoFinder2 v.2.5.4. The longest amino acid sequence of each isoform was selected to represent the gene. After OrthoFinder2 analysis, ortholog correspondences were corrected manually, if necessary. A list of genes is provided in Table S2.

Sample preparation for proteomic analysis

Proteomic analysis was conducted on the lips of cichlids, *H. chilotes* (hypertrophied) and *H. sauvagei* (normal), from Lake Victoria. The lips of *H. chilotes* were dissected in two half anterior (tip) and half posterior (base), which are analyzed separately. For proteomics analysis of pellet fractions, dissected samples were homogenized with a pestle in 1% sodium dodecyl sulfate in PBS supplemented with a protease inhibitor cocktail (cOmplete™ mini, EDTA free, Roche, Basel, Switzerland). The resultant lysate was centrifuged at 20,000 $\times$ g for 15 min, and the supernatant was extracted. A fourfold volume of methanol, an equal volume of chloroform, and a threefold volume of distilled water were added to the supernatants, and the suspension was vortexed vigorously for 1 min after each addition. After vortexing, the supernatants were centrifuged at 15,000 $\times$ g for 2 min and the upper water layer was removed. A threefold volume of methanol was added, and the samples were centrifuged at 15,000 $\times$ g for 2 min. The supernatant was removed by aspiration and the pellet fraction was obtained.

Subsequently, sample preparation for LC-MS/MS was performed according to a previous study (Masuda et al. 2009), with some modifications. Proteins in the pellet fraction were solubilized in phase transfer surfactant (PTS) solution (12 mM sodium deoxycholate, 12 mM sodium lauryl sulfate, 100 mM Tris-HCl, pH 9.0), and the protein concentration was determined using a Pierce BCA Protein Assay Kit (Thermo Fisher Scientific, Waltham, MA). Then, 50 μ g of protein in 50 μ L of PTS was prepared for the following procedure. First, the proteins were reduced by treatment with 5 mM dithiothreitol (DTT) at room temperature for 30 min, and then alkylated with 50 mM iodoacetamide in the dark at room temperature for 30 min. The reaction was stopped by a fivefold dilution with 50 mM ammonium bicarbonate. Denatured proteins in the solution were digested into peptide fragments in two steps: first, incubation with 0.5 μ g of Trypsin/Lys-C Mix (Promega, Madison, WI) at room temperature for 3 h, followed by incubation with 1.0 μ g of Trypsin/Lys-C Mix at 37°C overnight. After digestion, an equal volume of ethyl acetate and 0.5% trifluoroacetic acid (final concentration) were added and the mixture was shaken vigorously for 2 min. The mixture was then centrifuged at 15,700 $\times$ g for 2 min and the upper ethyl acetate layer was removed. The solvent was dried using a centrifugal evaporator. The residual pellet was redissolved in 600 μ L of 0.1% TFA and 2% acetonitrile and then desalted using a handmade StageTip composed of an SDB-XC Empore Disk (3 M, Maplewood, MN, USA). The solution was

applied to a StageTip equilibrated with 0.1% TFA and 2% acetonitrile, washed with 0.1% TFA and 2% acetonitrile, and eluted with 0.1% TFA and 80% acetonitrile. After the desalting, the solvent was dried using a centrifugal evaporator, and the residual peptides were redissolved in 120 μ L of 0.1% TFA and 2% acetonitrile. The solution was centrifuged at $20,000 \times g$ for 5 min, and the supernatant was collected and subjected to LC–MS/MS measurement.

LC–MS/MS measurement and data processing

LC–MS/MS measurements and SWATH–MS acquisition (Gillet et al., 2012) were performed using an Eksigent NanoLC Ultra and TripleTOF 4600 tandem-mass spectrometer or an Eksigent nanoLC 415 and TripleTOF 6600 mass spectrometer (AB Sciex, USA). The trap column used for nanoLC was a 5.0 mm $\times$ 0.3 mm ODS column (L-column2, CERI, Bunkyo-ku, Tokyo, Japan), and the separation column was a 12.5 cm $\times$ 75 μ m capillary column packed with 3 μ m C18-silica particles (Nikkyo Technos, Bunkyo-ku, Tokyo, Japan). The flow rate of nanoLC was 300 nL/min, and peptides were separated by applying a 10–40% linear acetonitrile gradient over 70 min in the presence of 0.1% formic acid. Data were acquired in DIA/SAWTH (Gillet et al., 2012) mode controlled by Analyst software (version 1.7, AB Sciex, USA) with the following settings: 15 Da fixed width window in 350–1100 m/z (50 cycles), 45 msec acquisition time, and 100–1600 fragment ion m/z range. The measurements were performed three times for each sample. Quantitative values were calculated by DIA–NN software (version 1.8.1) with default settings (Demichev et al., 2020). The library for DIA/SWATH quantification was generated using DIA–NN with the amino acid sequences of *M. zebra* total proteins obtained from UniProtKB. The P-value and fold change were calculated by Python libraries (scipy.stats and pandas) using the proteins detected in all three measurements for three parts of the lips. Quantitative values were scaled by log10 before statistical tests (Welch's t-test, two sided).

RNA-seq analysis

RNA was extracted from dissected lower lips using TRI Reagent (Molecular Research Center, Inc.). Three species of Lake Victoria cichlids (*Haplochromis chilotes* [n = 4], *H. sauvagei* [n = 4], *H. pyrrhocephalus* [n = 2]), three species of Lake Malawi cichlids (*Placidochromis milomo* [n = 5], *Maylandia zebra* [n = 2], *Labidochromis caeruleus* [n = 5]), and two species of Lake Tanganyika cichlids (*Lobochilotes labiatus* [n = 4], *Tropheus moorii* [n = 4]) were used for RNA-seq (Table S1). The extracted total RNA was sequenced at 100-bp paired-end reads on a Illumina NovaSeq 6000 by Macrogen Japan Corp., using a TruSeq stranded mRNA Library Prep Kit (Illumina). The raw FASTQ data were qualified using fastp version 0.20.1 (Chen et al., 2018) in default option. The high-quality reads were mapped to *M. zebra* genome (Conte and Kocher, 2015) using STAR version 2.7.10a (Dobin et al., 2013) and quantified using rsem-calculate-expression version 1.3.1 (Li and Dewey, 2011). Principal component analysis (PCA) was conducted using Python scikit-learn package. Low-expression genes (average transcripts per kilobase million (TPM) below 5.0) were removed, and expression rates were normalized by Z score. The resulting 12,231 genes were used for PCA analysis. For PCA of ECM-related genes, we used 186 genes as inputs, which were extracted in the same manner. DEGs were detected using the TCC package (Sun et al., 2013) with the TMM-edgeR-edgeR combination. Enrichment analysis was conducted using DAVID version 2023q3 (Huang et al., 2009a, 2009b) after the gene IDs of *M. zebra* were converted to human gene IDs using ensembl biomart. The annotation terms used for functional annotation clustering were collected from biological process (Gene Ontology), KEGG, and Reactome.

Results

Comparative histology of hypertrophied and normal lips among cichlids from the East African Great Lakes

We compared the histological structures of hypertrophied and normal lips among cichlids from Lake Victoria, Lake Malawi, and Lake Tanganyika. For an accurate comparison, we first observed serial sections of the lower lips with the jaws to determine the midline of the lips in each species, which was defined by the shape of the lower dentary bone (**Figure S1** [↗](#)). Based on this definition, we compared the distribution of collagens and proteoglycans, the main ECM components, at the midline of the lower lips using VG stain (red) and AB stain (blue). Hypertrophied lips of cichlids from all lakes generally consisted of two mutually exclusive structures in the hypodermis: a collagen-rich layer at the base and a proteoglycan-rich layer at the tip. Moreover, comparison of hypertrophied lips and normal lips showed that the area of the proteoglycan-rich (AB positive) layer was notably enlarged in hypertrophied lips, which was statistically significant in cichlids from Lake Victoria (**Figure 1** [↗](#); **Figure S2** [↗](#)).

Proteoglycans and their related proteins were abundant in the hypertrophied lips

To explore the proteins associated with lip hypertrophy, we used shotgun proteomics to compare the protein profiles of hypertrophied lips and normal lips in Lake Victoria cichlids. Using LC-MS/MS in DIA/SWATH acquisition mode, we detected 2,720 proteins in total from three types of lip samples: the tip of hypertrophied lips (proteoglycan-rich loose connective tissue), the base of hypertrophied lips (collagen-rich loose connective tissue), and normal lips. First, we observed few differences between the tip of hypertrophied lips and the base of the hypertrophied lips (**Figure S3** [↗](#)).

On the other hand, we identified 138 differentially accumulated proteins between the tip of hypertrophied and normal lips (133 proteins upregulated and 5 proteins downregulated in the hypertrophied lip) under the criteria (P -value < 0.05; log₂ fold change > 1) (**Figure 2A** [↗](#)). Compared to normal lips, tip of hypertrophied lips is abundant in proteoglycans and their related proteins such as versican a/b (vcan), brevican (bcan), periostin (postn), hyaluronan and proteoglycan link protein 1 (hapln1), and hyaluronan binding protein 2 (happ2), which were highly accumulated in hypertrophied lips with statistical significance (**Figures 2B-D** [↗](#)). The results of the above proteomic analyses were consistent with the histological observation that proteoglycan-rich loose connective tissue was enlarged in hypertrophied lips compared to that in normal lips.

Chondroitin sulfate proteoglycans such as vcan are the main component of hypertrophied lips

Next, we performed glycosaminoglycan digestion experiments to examine whether loose connective tissue was composed of the proteoglycans, which were detected as highly accumulated proteins in hypertrophied lips. Among the proteoglycans, we focused on vcan, which showed a clear difference in protein amount between hypertrophied and normal lips. Vcan is a high-molecular-weight core protein (approximately 370 kDa) that is glycosylated with chondroitin sulfate (Iozzo and Schaefer, 2015 [↗](#); Wight et al., 2020 [↗](#)). Therefore, we stained sections of hypertrophied lips with AB stain after the treatment with chondroitinase, which specifically digests glycosylated chondroitin sulfate. As a result, AB staining in loose connective tissue became drastically weak in treated sections, which was common among lakes (**Figures 2E-G** [↗](#)). Our data suggested that the main components of enlarged loose connective tissue in hypertrophied lips are chondroitin sulfate proteoglycans such as vcan.




Fig. 1

Histological comparisons of hypertrophied and normal lips of cichlids among lakes, focusing on proteoglycan-rich loose connective tissue (A,B). Examples of typical cichlids with hypertrophied lips (A, *Haplochromis chilotes*) and normal lips (B, *H. sauvagei*). The lower jaw and lip are highlighted by black squares. (C) Phylogenetic relationships among cichlids from Lake Victoria, Lake Malawi, and Lake Tanganyika. (D–O) The midline of the lips, based on the lower dentary bone of the lower jaw (marked by asterisks), in cichlids from Lake Victoria (D,E, *H. chilotes* with hypertrophied lips; F,G, *H. sauvagei* with normal lips), Lake Malawi (H,I, *Placidochromis milomo* with hypertrophied lips; J,K, *Maylandia lombardoi* with normal lips), and Lake Tanganyika (L,M, *Lobochilotes labiatus* with hypertrophied lips; N,O, *Tropheus moorii* with normal lips) were sectioned in sagittal, and stained with AB: Alcian blue (D, H, L, F, J, and N), or VG: Van Gieson (E, I, M, F, J, and N). Each section is shown with the dorsal side up and the anterior side to the right. The area surrounded by dotted lines shows proteoglycan-rich tissue (stained dense blue by AB) or collagen-rich tissue (stained dense red by VG). Each species showed mutually exclusive staining by AB and VG, and AB-positive (proteoglycan-rich) tissues were commonly enlarged in hypertrophied lips among the cichlids from all three lakes with statistical significance. Scale bars are 1mm.




Fig. 2

Chondroitin sulfate proteoglycans accumulate in hypertrophied lips. (A) Volcano plot of proteomics comparison between the tips of hypertrophied and normal lips. Statistical significance was examined using Welch's t-test. Yellow plots highlight highly accumulated proteins in hypertrophied lips, and blue plots highlight highly accumulated proteins in normal lips with the criteria of P-value < 0.05 & log2 fold change > 1. (B,C) The top 20 proteins highly accumulated in normal (B) and hypertrophied lips (C) are shown along with log2 fold change relative to normal lips. (D) The relative amount of each proteoglycan-related protein. MS2 intensity indicates the intensity of peptides obtained from LC-MS/MS and DIA/SWATH analysis using DIA-NN software by comparing three parts of the lip: the base of hypertrophied lips (HLip), the tip of HLip, and normal lips (NLip). Welch's t-test was used to examine statistical significance. (E–G) Digestion of chondroitin sulfate by treatment with (upper panels) and without (lower panels) chondroitinase in hypertrophied lips of cichlids from Lake Victoria (E), Lake Malawi (F), and Lake Tanganyika (G). Each section is shown with the dorsal side up and the anterior side to the right. Scale bars are 1mm.

Convergent expression in ECM-related genes

Given that proteomic analysis is not well suited for detecting high-turnover proteins (Decaris et al., 2014), we additionally conducted transcriptomic comparisons between hypertrophied and normal lips across Lake Victoria, Lake Malawi and Lake Tanganyika. PCA with all expressed genes clustered according to each lake, reflecting genetic closeness (Figure 3A). Next, we compared the gene expression of ECM-related genes, for which the proteomics profiles were clearly distinct between hypertrophied and normal lips. A PCA plot was drawn using only ECM-related genes in cichlids, which were identified based on a previous study (Hynes and Naba, 2012) (Figure 3B). In contrast to the PCA plot of all expressed genes, that of ECM-related genes reflected differences of lip morphology (hypertrophied and normal), regardless of the genetic closeness. To evaluate the results of these PCAs, we calculated the average silhouette (AS) score, which assesses the cohesiveness of the cluster. In the PCA of all genes, the AS score of the cluster by lake (0.1601) was higher than that of the cluster by lip morphology (0.0849). In contrast, in the PCA of ECM-related genes, the AS score of the cluster by lip morphology (0.1798) was higher than that of the cluster by lake (0.0774). These results indicate that hypertrophied lips are characterized by the expression of ECM-related genes.

Next, we compared the expression of each ECM-related gene between hypertrophied and normal lips of cichlids from the three lakes (Figures 3C, D). As a result, several ECM-related genes showed higher expression in hypertrophied lips than in normal lips, such as type-I collagen fibrils (col1a1a, col1a1b, and col1a2), hyaluronan-binding proteoglycans (vcanb, hapln1a/b, and hapln3), periostins (postna/b), small leucine-rich proteoglycans (dcn, lum, and aspn), and microfibril-related genes (mfap2, emilin1b, emilin2a, emilin2b, and emilin3b). Consistent with the proteomics comparison, some proteoglycans and their related proteins (vcanb, hapln1a, and postna) showed significantly higher expression in hypertrophied lips than in normal lips. In conclusion, we observed convergent expression of ECM-related genes in hypertrophied lips of cichlids among the East African Great Lakes.

High expression of ECM-related genes in the juvenile and adult stages of hypertrophied lips

Next, we performed transcriptome analysis by comparing juvenile and adult cichlids from Lake Victoria to examine the possible expression changes in ECM-related genes during development (Figure 4A). In the adult stage, the expression profiles of ECM-related genes between hypertrophied and normal lips showed obvious differences (low correlation value 0.75; orange square in Figure 4B), which was consistent with previous analyses (Figure 3B). In the juvenile stage, the difference was weaker than in adult stage (0.85; blue square in Figure 4B). Interestingly, the difference in the expression profiles between the juvenile and adult stages of hypertrophied lips (0.90; Figure 4B black square) was lower than that of normal lips (0.83; Figure 4B black dotted square). To examine the genes contributing to the differences in expression profiles, we performed hierarchical clustering analysis for the relative expression levels of each ECM-related gene (Figure 4C, D). The analyses consistently showed the higher expression of proteoglycans and their related genes (e.g. postna, vcanb, and hapln1b) in hypertrophied lips at both the juvenile and adult stages. These results indicated that the accumulation of proteoglycans, which leads to the lip hypertrophy, begins at the juvenile stage.

The Wnt pathway-related genes are highly expressed in hypertrophied lips

To explore the molecular pathways involved in lip hypertrophy, we performed an enrichment analysis using DAVID (Huang et al., 2009a, 2009b) against DEGs obtained from the transcriptome. As a result of enrichment analysis, DEGs were categorized in the canonical and




Fig. 3

Hypertrophied lips of cichlids are characterized by higher expression of ECM-related genes. Transcriptomic analysis using three species of Lake Victoria cichlids (*H. chilotes* (Hch) with hypertrophied lips; *H. sauvagei* (Hsa) and *H. pyrrhocephalus* (Hpy) with normal lips), three species of Lake Malawi cichlids (*Placidochromis milomo* (Pmi) with hypertrophied lips; *Maylandia zebra* (Mze) and *Labidochromis caeruleus* (Lca) with normal lips), and two species of Lake Tanganyika cichlids (*Lobochilotes labiatus* (Lla) with hypertrophied lips; *Tropheus moorii* (Tmo) with normal lips). (A, B) PCA plots of all expressed genes showed differences depending on the lakes (A), while those of ECM-related genes showed differences depending on lip morphology: hypertrophied or normal (B). (C) Heatmap of ECM-related gene expression between hypertrophied and normal lips. The expression rate (TPM) normalized by Z score of each gene is compared for Lake Victoria (yellow; Hch, Hsa and Hpy), Lake Malawi (blue; Pmi, Lca and Mze), and Lake Tanganyika (red; Lla and Tmo). The intensity of the relative expression rate is shown in the bottom right of the figure. (D) Cumulative fold change of the ECM-related genes comparing hypertrophied and normal lips. The gene symbol corresponds to **Figure 3C**. The color represents the lake information. The gray dotted line represents the value at which higher expression was observed in the hypertrophied lips.




Fig. 4

Comparison of juvenile and adult cichlids to examine gene expression changes during the development of hypertrophied lips. (A) The lips of the cichlids used in this analysis are shown with the dorsal side up and the anterior side to the right: the juvenile and adult stages of *Haplochromis chilotes* (Hch, representative of hypertrophied lips: HLip) and *H. sauvagei* (Hsa, representative of normal lips: NLip) from Lake Victoria. Hypertrophied lips at juvenile stage are not notably enlarged. Scale bars are 0.5 mm. (B) Spearman correlation matrix of the ECM-related gene expression rate (TPM) normalized by Z score. The raw correlation values are indicated on the right side and the average correlation values within each stage and species are indicated on the left side. The orange frame highlights the correlations between the hypertrophied and normal lips at the adult stage, and the blue frame highlights those at the juvenile stage. The black frame highlights the correlations between juvenile and adult samples of each species. (C) Cluster heatmap of the ECM-related gene expression rate (TPM) normalized by Z score in each gene. Each gene is clustered by similarity of expression rates. The intensity of the relative expression rate is shown on the bottom left of the figure. (D) The log10 cumulative fold change of the ECM-related genes comparing hypertrophied and normal lips. The gene symbol corresponds to **Figure 4C**. The positive values indicate high expression in hypertrophied lips, whereas the negative values indicate low expression in hypertrophied lips. The colors of the bars indicate the stage of the samples.

noncanonical Wnt signaling pathways, suggesting that high expression of genes in the Wnt signaling pathway is likely to be involved in the hypertrophied lips of juvenile and adult fish. Comparison of the developmental stages revealed differences in the expression patterns between the canonical and noncanonical Wnt pathways: the genes of the canonical Wnt pathway were highly expressed in both stages (**Figures 5C–I** [↗](#)), whereas those of the non-canonical pathway were highly expressed only in the adult stage (**Figures 5J–N** [↗](#)). We further compared the expression patterns of these Wnt signaling pathways in cichlids from Lake Malawi and Lake Tanganyika. Notably, the expression of genes for the Wnt pathway tend to be higher in the hypertrophied-lip cichlids in Lake Malawi and Lake Tanganyika, although the differences were not as distinct as those found in cichlids from Lake Victoria. These results were consistent with those of previous studies that showed that the expression of ECM-related genes, such as *vcan*, was promoted by the Wnt signaling pathway (Rahmani et al., 2005 [↗](#)). Continuous activation of the Wnt signaling pathway throughout all developmental stages may lead to lip hypertrophy via the deposition of ECM-related proteins.

Discussion

In this study, we investigated the differences between hypertrophied and normal lips using histology, proteomics, and transcriptomics. Histological comparisons revealed that cichlids with hypertrophied lips from all three East African Great Lakes exhibited an enlargement of the proteoglycan-rich loose connective tissue. By integrating proteomics and transcriptomics, we identified several proteins that were significantly accumulated in the hypertrophied lips. Among them, proteoglycans and their related proteins such as *vcan* and *postn*, could be a primary factor that explains lip hypertrophy. Interestingly, these proteins are known to be involved in the formation of human keloids, a skin disease characterized by an overabundance of ECM in the dermis (Deng et al., 2021 [↗](#); Naitoh et al., 2005 [↗](#)). It has been discussed that excessive deposition of *vcan* retains large amounts of water in the sulfate or carbonyl moieties, contributing to the increase in keloid volume (Yagi et al., 2013 [↗](#)). In addition, *postn* is involved in fibrosis through its ability to interact with multiple ECM proteins. For example, upregulation of *postn* leads to strengthening of the ECM by crosslinking with collagens and excessive deposition of ECM proteins by differentiating quiescent fibroblasts into activated myofibroblasts (Ashley et al., 2017 [↗](#); Yamaguchi et al., 2013 [↗](#)). In cichlid, *vcan* and *postn* may have a similar function, leading to hypertrophied loose connective tissue.

The Wnt signaling pathway was also elevated in both the adult and juvenile stages of hypertrophied lips. The Wnt signaling pathway is known to be a key regulator of the expression of ECM-related genes; for example, *vcan* is directly regulated by the Wnt/*b-catenin* and *tcf4* signaling pathways, leading to *vcan* upregulation (Rahmani et al., 2005 [↗](#)). Additionally, *Wnt5a*, a noncanonical pathway, plays a critical role in dermal condensation during mouse and sugar glider development (Feigin et al., 2023 [↗](#)). Cellular-level localization of gene expression for the canonical and non-canonical Wnt pathway is required for future examination of the involvement of the Wnt signaling pathway in lip hypertrophy.

From a genomic perspective, several studies have investigated the genetic basis of hypertrophied lip cichlids (Masonick et al., 2023 [↗](#); Nakamura et al., 2021 [↗](#)). Importantly, some Wnt pathway-related genes (*tcf4* and *daam2*) and ECM-related genes (*postna*, *col12a1a*, and *col12a1b*) have been found to be under positive selection in cichlids with hypertrophied lips of Lake Victoria (see Nakamura et al., 2021 [↗](#) Table S3). For future research, examining whether these genes are under selection in other lakes is crucial to understand the genetic mechanisms underlying the parallel evolution of hypertrophied lips.




Fig. 5

Wnt pathway-related genes are highly expressed in hypertrophied lips of juvenile and adult cichlids. (A, B) Enrichment analysis of highly expressed genes in the hypertrophied lips. Labels are according to the top annotation in DAVID annotation clusters. Asterisks indicate statistically significant annotation clusters (FDR < 0.05). The Wnt-related annotations are highlighted in yellow. (C–N) Expression of the specific Wnt pathway-related genes. (C, J) The size of the dots represents the log fold change, and the color of the dots represents the P-value. Blue dots represent genes that are highly expressed in hypertrophied lips (HLip), and red dots represent genes that are highly expressed in normal lips (NLip). (D–I, K–N) Box plots show the expression rate (TPM) of Wnt pathway-related genes. Yellow boxes represent genes that are highly expressed in hypertrophied lips and blue boxes represent genes that are highly expressed in normal lips, respectively. Mann–Whitney U-test was used to examine the statistical significance (*: P-value < 0.05). (C–I) The canonical Wnt pathway-related genes, and (J–N) show the non-canonical Wnt pathway-related genes.




Fig. 6

The trend of higher expression of Wnt-related genes in hypertrophied lips. (A) Box plots of expression rate (TPM) of Wnt related genes of hypertrophied lips (HLip: yellow) and normal lips (NLip: blue). Mann-Whitney U-test was used to examine the statistical significance (*: P-value < 0.05, **: P-value < 0.01). (B) Illustration of the canonical Wnt pathway. DEGs are marked in the bold.

Although the parallel evolution of hypertrophied lips in cichlids has received considerable attention in evolutionary biology, the commonality and diversity underlying their molecular basis remain poorly understood. This might be primarily due to the fact that many genetic factors are possibly involved in the dynamic organization of hypertrophied lips. We uncovered the apparent relationships between hypertrophied lips and the expression profiles of ECM proteins, in particularly proteoglycans. The trends for the overall expression of ECM-related genes were similar across hypertrophied lip species, but we rarely observed a specific gene that was commonly expressed at high or low levels in all three examples of hypertrophied lips across all East African Great Lakes. Furthermore, although we focused primarily on the relationship between the Wnt signaling pathway and lip hypertrophy, several other pathways may be involved in the development of hypertrophied lips. For example, our enrichment analysis in adult cichlids identified VEGF-related pathways, which could contribute to lip hypertrophy by increasing vascularization and nutrient supply to the lip tissue. In addition, previous quantitative trait locus (QTL) analysis by [Henning et al. \(2017\)](#) [↗](#) concluded that lip hypertrophy is likely influenced by numerous loci with small additive effects. These lines of data imply that although enlargement of proteoglycan-rich loose connective tissue is common in hypertrophied lips, the developmental pathways to accomplish this are diverse in each lake. The pathways responsible for lip hypertrophy therefore require further investigation. In conclusion, our approach of integrating histology, proteomics and transcriptomics revealed complex evolutionary pathways for hypertrophied lips in cichlids of the East African Great Lakes, providing an opportunity to further understand the molecular mechanism underlying the parallel evolution of polygenic traits, such as jaw morphology or body shape.

Data availability

RNA-seq short reads and raw proteome data are stored at NCBI and jPOST ([Okuda et al., 2017](#) [↗](#)) (Accession: PRJDB3401, PRJDB17757, and JPST003042/PXD051571).

Acknowledgements

This study was supported by JSPS KAKENHI (23KJ0951 to N. M., 17H04606, 20KK0167 to M.N.) and JST SPRING (JPMJSP2106). Computations were partially performed on the NIG supercomputer at ROIS National Institute of Genetics. The authors thank the Open Research Facilities for Life Science and Technology, Tokyo Institute of Technology for LC-MS/MS analysis (for technical assistance). The authors thank the Hongoh laboratory members. Nagatoshi Machii thank my family, Nagashi Machii, Mitsue M. and Sawa M.

Additional files

Supplementary Table 1-5. Table S1. Basic information of the cichlids used in this study. Table S2. List of ECM-related genes of cichlids identified in this study. The classification of ECM-related genes (the column of *matrisome_annotation*) follows a previous study ([Hynes and Naba, 2012](#)). Table S3. All differentially accumulated proteins detected by LC-MS/MS under the criteria of $P\text{-value} < 0.05$, $\log_2 \text{fold change} > 1$. Table S4. Results of enrichment analysis of highly expressed genes in hypertrophied lips of juvenile and adult cichlids. The annotation terms were clustered by DAVID annotation clustering. Table S5. All differentially expressed genes between hypertrophied lips and normal lips. We compared adult *Victoria* cichlids (*H. chilotes* with hypertrophied lips and *H. sauvagei* with normal lips), juvenile *Victoria* cichlids (*H.*

Fig S1

The midlines of the lips were defined by the shape of the lower dentary bone. Hematoxylin–eosin stain (HE stain) was used to examine serial sections of the lower lips and lower jaws of cichlids from Lake Victoria (A, *H. chilotes* with hypertrophied lips; B, *H. sauvagei* with normal lips), Lake Malawi (C, *Placidochromis milomo* with hypertrophied lips; D, *Maylandia lombardoi* with normal lips), and Lake Tanganyika (E, *Lobochilotes labiatus* with hypertrophied lips; F, *Tropheus moorii* with normal lips). Each section is shown with the dorsal side up and the anterior side to the left. The lower dentary bone became smaller as it approached the midline, and just at the midline, the bone was composed of cartilage. Scale bars are 1mm.





Fig. S2

The relative area of the proteoglycan-rich layer was significantly enlarged in the hypertrophied lips of Victoria cichlids. The stained area percentage of alcian blue stain (AB stain; A) and Van Gieson stain (VG stain; B) relative to the whole-lip area was calculated using ImageJ software. *H. chilotes* with hypertrophied lips (yellow box) and *H. sauvagei* with normal lips (blue box) were used in the experiment. Each plot is an average value calculated by three sections near the midline of the lips. Mann–Whitney U-test was used to examine the statistical significance (*: P-value < 0.05, **: P-value < 0.01).






Fig. S3

Volcano plot of proteomics comparison between the tip of hypertrophied lips and the base of hypertrophied lips (A) and the base of hypertrophied lips and normal lips (B). Statistical significance was examined by Welch's t-test. Differentially accumulated proteins in each comparison are highlighted in the black plot with the following criteria: P-value < 0.05, log2 fold change > 1.

chilotes and *H. sauvagei*), Malawi cichlids (*Placidochromis milomo* with hypertrophied lips and *Labidochromis caeruleus* with normal lips), and Taganyika cichlids (*Lobochilotes labiatus* with hypertrophied lips and *Tropheus moorii* with normal lips). [↗](#)

References

1. Albertson RC, Kocher TD 2006) **Genetic and developmental basis of cichlid trophic diversity** *Heredity* **97**:211–221
2. Arnegard ME, Snoeks J 2001) **New Three-Spotted Cichlid Species with Hypertrophied Lips (Teleostei: Cichlidae) from the Deep Waters of Lake Malaŵi/Nyasa Africa**. *The American Society of Ichthyologists and Herpetologists* [https://doi.org/10.1643/0045-8511\(2001\)001\[0705:NTSCSW\]2.0.CO](https://doi.org/10.1643/0045-8511(2001)001[0705:NTSCSW]2.0.CO)
3. Ashley SL, Wilke CA, Kim KK, Moore BB 2017) **Periostin regulates fibrocyte function to promote myofibroblast differentiation and lung fibrosis** *Mucosal Immunol* **10**:341–351
4. Chen S, Zhou Y, Chen Y, Gu J 2018) . **fastp: an ultra-fast all-in-one FASTQ preprocessor** *Bioinformatics* **34**:i884–i890
5. Colombo M, Diepeveen ET, Muschick M, Santos ME, Indermaur A, Boileau N, Barluenga M, Salzburger W 2013) **The ecological and genetic basis of convergent thick-lipped phenotypes in cichlid fishes** *Mol Ecol* **22**:670–684
6. Conte MA, Kocher TD 2015) **An improved genome reference for the African cichlid, *Metriaclicma zebra*** *BMC Genomics* **16**:724
7. Decaris ML, Gatmaitan M, FlorCruz S, Luo F, Li K, Holmes WE, Hellerstein MK, Turner SM, Emson CL 2014) **Proteomic analysis of altered extracellular matrix turnover in bleomycin-induced pulmonary fibrosis** *Mol Cell Proteomics* **13**:1741–1752
8. Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M 2020) **DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput** *Nat Methods* **17**:41–44
9. Deng C-C, Hu Y-F, Zhu D-H, Cheng Q, Gu J-J, Feng Q-L, Zhang L-X, Xu Y-P, Wang D, Rong Z, Yang B 2021) **Single-cell RNA-seq reveals fibroblast heterogeneity and increased mesenchymal fibroblasts in human fibrotic skin diseases** *Nat Commun* **12**:3709
10. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR 2013) **STAR: ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21
11. Duenser A, Singh P, Lecaudey LA, Sturmbauer C, Albertson RC, Gessl W, Ahi EP 2023) **Conserved Molecular Players Involved in Human Nose Morphogenesis Underlie Evolution of the Exaggerated Snout Phenotype in Cichlids** *Genome Biol Evol* **15** <https://doi.org/10.1093/gbe/evad045>
12. Feigin CY, Moreno JA, Ramos R, Mereby SA, Alivisatos A, Wang W, van Amerongen R, Camacho J, Rasweiler JJ, Behringer RR, Ostrow B, Plikus MV, Mallarino R 2023) **Convergent deployment of ancestral functions during the evolution of mammalian flight membranes** *Sci Adv* **9**:eade7511
13. Fryer G 1972) **The cichlid fishes of the Great Lakes of Africa** *Their Biology and Evolution* **641**

14. Gillet LC, Navarro P, Tate S, Röst H, Selevsek N, Reiter L, Bonner R, Aebersold R (2012) **Targeted Data Extraction of the MS/MS Spectra Generated by Data-independent Acquisition: A New Concept for Consistent and Accurate Proteome Analysis** *Mol Cell Proteomics* **11**:O111.016717
15. Greenwood PH (1974) **Cichlid fishes of Lake Victoria, East Africa: the biology and evolution of a species flock**. *Bull Brit Mus (Nat Hist) Zool Suppl* **6**:1–134
16. Henning F, Machado-Schiaffino G, Baumgarten L, Meyer A (2017) **Genetic dissection of adaptive form and function in rapidly speciating cichlid fishes** *Evolution* **71**:1297–1312
17. Huang DW, Sherman BT, Lempicki RA (2009a) **Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists** *Nucleic Acids Res* **37**:1–13
18. Huang DW, Sherman BT, Lempicki RA (2009b) **Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources** *Nat Protoc* **4**:44–57
19. Hynes RO, Naba A (2012) **Overview of the matrisome--an inventory of extracellular matrix constituents and functions** *Cold Spring Harb Perspect Biol* **4**:a004903
20. Iozzo RV, Schaefer L (2015) **Proteoglycan form and function: A comprehensive nomenclature of proteoglycans** *Matrix Biol* **42**:11–55
21. Kautt AF, Kratochwil CF, Nater A, Machado-Schiaffino G, Olave M, Henning F, Torres-Dowdall J, Härer A, Hulsey CD, Franchini P, Pippel M, Myers EW, Meyer A (2020) **Contrasting signatures of genomic divergence during sympatric speciation** *Nature* **588**:106–111
22. Kocher T, Conroy JA, McKaye KR, Richard SJ (1993) **Similar Morphologies of Cichlid Fish in Lakes Tanganyika and Malawi Are Due to Convergence** *Mol Phylogenet Evol* **2**:158–165
23. Kocher TD (2004) **Adaptive evolution and explosive speciation: the cichlid fish model** *Nat Rev Genet* **5**:288–298
24. Kratochwil CF, Liang Y, Gerwin J, Woltering JM, Urban S, Henning F, Machado-Schiaffino G, Hulsey CD, Meyer A (2018) **Agouti-related peptide 2 facilitates convergent evolution of stripe patterns across cichlid fish radiations** *Science* **362**:457–460
25. Krishnan J, Rohner N (2017) **Cavefish and the basis for eye loss** *Philos Trans R Soc Lond B Biol Sci* **372** <https://doi.org/10.1098/rstb.2015.0487>
26. Lecaudey LA, Singh P, Sturmbauer C, Duenser A, Gessl W, Ahi EP (2021) **Transcriptomics unravels molecular players shaping dorsal lip hypertrophy in the vacuum cleaner cichlid, *Gnathochromis permaxillaris*** *BMC Genomics* **22**:506
27. Lecaudey LA, Sturmbauer C, Singh P, Ahi EP (2019) **Molecular mechanisms underlying nuchal hump formation in dolphin cichlid, *Cyrtocara moorii*** *Sci Rep* **9**:20296
28. Li B, Dewey CN (2011) **RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome** *BMC Bioinformatics* **12**:323
29. Losos JB (2011) **Convergence, adaptation, and constraint** *Evolution* **65**:1827–1840
30. Lukas B, Gonzalo M-S, Frederico H, Axel M (2015) **What big lips are good for: on the adaptive function of repeatedly evolved hypertrophied lips of cichlid fishes** *Biol J Linn Soc Lond*

31. Manousaki T, Hull PM, Kusche H, Machado-Schiaffino G, Franchini P, Harrod C, Elmer KR, Meyer A (2013) **Parsing parallel evolution: ecological divergence and differential gene expression in the adaptive radiations of thick-lipped Midas cichlid fishes from Nicaragua** *Mol Ecol* **22**:650–669
32. Masonick P, Meyer A, Darrin Hulsey C (2023) **A Kiss of Deep Homology: Partial Convergence in the Genomic Basis of Hypertrophied Lips in Cichlid Fish and Human Cleft Lip** *Genome Biol Evol* **15** <https://doi.org/10.1093/gbe/evad072>
33. Masonick P, Meyer A, Hulsey CD (2022) **Phylogenomic Analyses Show Repeated Evolution of Hypertrophied Lips Among Lake Malawi Cichlid Fishes** *Genome Biol Evol* **14** <https://doi.org/10.1093/gbe/evac051>
34. Muschick M, Indermaur A, Salzburger W (2012) **Convergent evolution within an adaptive radiation of cichlid fishes** *Curr Biol* **22**:2362–2368
35. Naitoh M, Kubota H, Ikeda M, Tanaka T, Shirane H, Suzuki S, Nagata K (2005) **Gene expression in human keloids is altered from dermal to chondrocytic and osteogenic lineage** *Genes Cells* **10**:1081–1091
36. Nakamura H, Aibara M, Kajitani R, Mrosso HDJ, Mzighani SI, Toyoda A, Itoh T, Okada N, Nikaido M (2021) **Genomic Signatures for Species-Specific Adaptation in Lake Victoria Cichlids Derived from Large-Scale Standing Genetic Variation** *Mol Biol Evol* **38**:3111–3125
37. Okuda S, Watanabe Y, Moriya Y, Kawano S, Yamamoto T, Matsumoto M, Takami T, Kobayashi D, Araki N, Yoshizawa AC, Tabata T, Sugiyama N, Goto S, Ishihama Y (2017) **jPOSTrepo: an international standard data repository for proteomes** *Nucleic Acids Res* **45**:D1107–D1111
38. Oliver MK, Arnegard ME (2010) **A new genus for *Melanochromis labrosus*, a problematic Lake Malawi cichlid with hypertrophied lips (Teleostei: Cichlidae)** *Ichthyol Explor Freshwaters* **21**:209–232
39. Rahmani M, Read JT, Carthy JM, McDonald PC, Wong BW, Esfandiarei M, Si X, Luo Z, Luo H, Rennie PS, McManus BM (2005) **Regulation of the Versican Promoter by the β -Catenin-T-cell Factor Complex in Vascular Smooth Muscle Cells*** *J Biol Chem* **280**:13019–13028
40. Salzburger W (2018) **Understanding explosive diversification through cichlid fish genomics** *Nat Rev Genet* **19**:705–717
41. Santos ME, Lopes JF, Kratochwil CF (2023) **East African cichlid fishes** *Evodevo* **14**:1
42. Schneider CA, Rasband WS, Eliceiri KW (2012) **NIH Image to ImageJ: 25 years of image analysis** *Nat Methods* **9**:671–675
43. Stern DL (2013) **The genetic causes of convergent evolution** *Nat Rev Genet* **14**:751–764
44. Sun J, Nishiyama T, Shimizu K, Kadota K (2013) **TCC: an R package for comparing tag count data with robust normalization strategies** *BMC Bioinformatics* **14**:219
45. Wight TN, Kang I, Evanko SP, Harten IA, Chang MY, Pearce OMT, Allen CE, Frevert CW (2020) **Versican-A Critical Extracellular Matrix Regulator of Immunity and Inflammation** *Front Immunol* **11**:512

46. Yagi Y, Muroga E, Naitoh M, Isogai Z, Matsui S, Ikehara S, Suzuki S, Miyachi Y, Utani A (2013) **An ex vivo model employing keloid-derived cell-seeded collagen sponges for therapy development** *J Invest Dermatol* **133**:386–393
47. Yamaguchi Y, Ono J, Masuoka M, Ohta S, Izuhara K, Ikezawa Z, Aihara M, Takahashi K (2013) **Serum periostin levels are correlated with progressive skin sclerosis in patients with systemic sclerosis** *Br J Dermatol* **168**:717–725

Author information

Nagatoshi Machii

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan
ORCID iD: [0009-0003-8703-339X](https://orcid.org/0009-0003-8703-339X)

Ryo Hatashima

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan
ORCID iD: [0000-0003-2734-1027](https://orcid.org/0000-0003-2734-1027)

Tatsuya Niwa

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan, Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, Yokohama, Japan
ORCID iD: [0000-0002-1330-8974](https://orcid.org/0000-0002-1330-8974)

Hideki Taguchi

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan, Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, Yokohama, Japan
ORCID iD: [0000-0002-6612-9339](https://orcid.org/0000-0002-6612-9339)

Ismael A Kimirei

Tanzania Fisheries Research Institute (TAFIRI), Dar es Salaam, United Republic of Tanzania
ORCID iD: [0000-0002-1101-5262](https://orcid.org/0000-0002-1101-5262)

Hillary DJ Mrosso

Tanzania Fisheries Research Institute (TAFIRI), Mwanza Fisheries Research Center, Mwanza, United Republic of Tanzania
ORCID iD: [0000-0003-1924-7268](https://orcid.org/0000-0003-1924-7268)

Mitsuto Aibara

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan
ORCID iD: [0000-0001-8526-7449](https://orcid.org/0000-0001-8526-7449)

Tatsuki Nagasawa

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan
ORCID iD: [0000-0002-5759-2524](https://orcid.org/0000-0002-5759-2524)

Masato Nikaido

School of Life Science and Technology, Tokyo Institute of Technology, Tokyo, Japan

ORCID iD: [0000-0002-2430-6974](https://orcid.org/0000-0002-2430-6974)

For correspondence: mnikaido@life.isct.ac.jp

Editors

Reviewing Editor

Vincent Lynch

University at Buffalo, State University of New York, Buffalo, United States of America

Senior Editor

George Perry

Pennsylvania State University, University Park, United States of America

Reviewer #1 (Public review):

Summary:

Machii et al. reported a possible molecular mechanism underlying the parallel evolution of lip hypertrophy in African cichlids. The multifaceted approach taken in this manuscript is highly valued, as it uses histology, proteomics, and transcriptomics to reveal how phylogenetically distinct thick-lips have evolved in parallel. Findings from histology and proteomics connected to wnt signaling through the transcriptome are very exciting.

Strengths:

There is consistency between the results and it is possible to make a strong argument from the results.

Comments on revised version:

The issues I pointed out in the previous review have been carefully answered, and all issues have been addressed. The main points of the manuscript are clear, and the conclusions are easy to understand. The enlarged lips are a notable example of convergent evolution in African cichlids.

<https://doi.org/10.7554/eLife.99160.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer 1:

Weaknesses:

The authors do not discuss based on genomic information; the genomes of the cichlids from the three lakes have been decoded and are therefore available. However, indeed, the species in Lake Tanganyika and Lake Malawi/Victoria are genetically distant from each other, so a comparative genome analysis would not have yielded the results presented here. I recommend adding such a discussion to the Discussion.

We appreciate your comment. We added the discussion regarding the genomic aspect of parallel evolution.

Line 386-393: “From a genomic perspective, several studies have investigated the genetic basis of hypertrophied lip cichlids (Masonick et al., 2023; Nakamura et al., 2021). Importantly, some Wnt pathway-related genes (tcf4 and daam2) and ECM-related genes (postna, col12a1a, and col12a1b) have been found to be under positive selection in cichlids with hypertrophied lips of Lake Victoria (see Nakamura et al., 2021 Table S3). For future research, examining whether these genes are under selection in other lakes is crucial to understand the genetic mechanisms underlying the parallel evolution of hypertrophied lips.”

Minor comments:

Line 30, the Wnt --> the genes in Wnt

We appreciate your comment. According to the comment, we corrected the sentence.

Line 30: “the Wnt signaling pathway” -> “the genes in Wnt signaling pathway”

Line 42-44, "It is considered that the same direction of natural selection drives phenotypic changes among species since it is unlikely that these complex phenotypes have been acquired repeatedly just by neutral evolution". How about "Since it is unlikely that such a complex phenotype was acquired repeatedly by neutral evolution alone, the same direction of natural selection among species is likely to drive the parallel phenotypic change."?

We agree with your suggestion and correct the sentence of our manuscript.

Line 42-44: “It is considered that the same direction of natural selection drives phenotypic changes among species since it is unlikely that these complex phenotypes have been acquired repeatedly just by neutral evolution”

“Since it is unlikely that such a complex phenotype was acquired repeatedly by neutral evolution alone, the same direction of natural selection among species is likely to drive the parallel phenotypic change”

Line 60, polygenic --> likely to be polygenic

We appreciate your comment. Indeed, it is better to weaken the wording.

Line 60: “most traits are polygenic” -> “most traits are likely to be polygenic”

Line 91, the Wnt --> the genes in Wnt

We appreciate your correction. Last paragraph of introduction has been corrected according to the suggestion of Reviewer 2 (Q1).

Line 230, NovaSeq --> Illumina NovaSeq

We appreciate your correction.

Line 222: “NovaSeq 6000” -> “Illumina NovaSeq 6000”

Line 231 "mRNA Library Prep Kit". Please add a company name.

We appreciate your correction. We added company's information.

Line 223: “a TruSeq stranded mRNA Library Prep Kit.” -> “a TruSeq stranded mRNA Library Prep Kit (Illumina)”

Line 267, as for the tip of hypertrophied lips, could you add and point out which part is the tip?

We dissected hypertrophied lips in two half anterior and half posterior. We added the sentence in the materials and methods section.

Line 156-158: “The lips of *H. chilotes* were analyzed separately for the base and tip.” -> “The lips of *H. chilotes* were dissected in two half anterior (tip) and half posterior (base), which are analyzed separately.”

Line 272, “133 proteins upregulated and 5 proteins downregulated” in hypertrophied lip or normal lip?

We appreciate your correction. We added the sentence as follows.

Line 264: “133 proteins upregulated and 5 proteins downregulated”

“133 proteins upregulated and 5 proteins downregulated in the hypertrophied lip”

Line 274, “hypertrophied lips” means tip of hypertrophied lips?

We appreciate your correction. We corrected the sentence as follows.


Line 266: “hypertrophied lips are abundant” -> “tip of hypertrophied lips is abundant”

Line 277, Did you perform multiple testing correction for statistical significance?


We appreciate your comment about multiple testing corrections. We did not apply multiple testing corrections in our “exploratory” analysis of proteomics not to miss biologically important candidates in a limited sample size (n=3). We calculated the multiple corrected p-value in the Benjamini Hochberg method (Author response image 1, right). The result suggested that almost the same proteoglycans and its related proteins as we focused on are highly accumulated in the hypertrophied lips in milder conditions (significance level of 0.1).

Author response image 1.

No correction of p-value (the same as the Figure 2A)



Correction of p-value with Benjamini Hochberg



Thus, our main conclusions remain unchanged even with correction applied, however, the overall balance of the volcano plot is not visually appealing (Author response image 1, right).

It is important to note that we selected the Top 20 proteins based on fold change rather than statistical significance. In addition, our proteomic findings show consistency with our

histological and transcriptome data, providing the biological validation from various aspects. While we understand the potential benefits of multiple testing correction, our current approach without multiple testing still offers valuable and fair data to propose hypothesis on the molecular mechanisms of lip hypertrophy in cichlids. Therefore, we want to use original figure without multiple testing. We greatly appreciate the understanding of the reviewer.

Line 349-351, "The results of the enrichment analysis suggested that the genes that were categorized into both canonical and non-canonical Wnt signaling pathways, were highly expressed in the hypertrophied lips of juvenile and adult cichlids."

The wnt category was enriched by analyzing the highly expressed genes, so isn't it natural that the wnt category is highly expressed?

Did you mean to say as in the following sentence?

"Enrichment of genes categorized in the canonical and noncanonical Wnt signaling pathways suggested that high expression of genes in the Wnt signaling pathway is likely to be involved in the hypertrophied lips of juvenile and adult fish."

Thank you for your comments. We corrected our manuscript as follows.

Line 341-344: "The results of the enrichment analysis suggested that the genes that were categorized into both canonical and non-canonical Wnt signaling pathways, were highly expressed in the hypertrophied lips of juvenile and adult cichlids."

"As a result of enrichment analysis, DEGs were categorized in the canonical and noncanonical Wnt signaling pathways, suggesting that high expression of genes in the Wnt signaling pathway is likely to be involved in the hypertrophied lips of juvenile and adult fish."

Line 403-404, "several other pathways may be involved in the development of hypertrophied lips". Do you have any evidence?

We appreciate your comment regarding possible evidence for the involvement of multiple pathways in hypertrophied lip development. Our statement was based on two main points:

(1) While we highlighted the Wnt pathway because this pathway is known to increase proteoglycan expression, we cannot exclude the possibility of the involvement of other pathways. For instance, our enrichment analysis in adult cichlids identified VEGF-related pathways, which could contribute to lip hypertrophy by increasing vascularization and nutrient supply to the lip tissue.

(2) Previous quantitative trait locus (QTL) analysis by Henning et al. (2017) concluded that lip hypertrophy is likely influenced by numerous loci with small additive effects. This indicates that lip hypertrophy is a complex phenotype consisted of multiple genetic factors, some which probably correspond to different molecular pathways.

Given these points, we draw a conclusion that emphasize the importance of Wnt pathway while also recognizing the potential cooperative interaction of multiple pathways in developing lip hypertrophy. Without confusing the two statements, we corrected our manuscript as follows.

Line 398-412: "We uncovered the apparent relationships between hypertrophied lips and the expression profiles of ECM proteins, in particularly proteoglycans. The trends for the overall expression of ECM-related genes were similar across hypertrophied lip species, but we rarely observed a specific gene that was commonly expressed at high or low levels in all three examples of hypertrophied lips across all East African Great Lakes. Furthermore, although we focused primarily on the relationship between the Wnt signaling pathway and lip

hypertrophy, several other pathways may be involved in the development of hypertrophied lips. These findings imply that although enlargement of proteoglycan-rich loose connective tissue is common in hypertrophied lips, the developmental pathways to accomplish this are diverse in each lake.”

“We uncovered the apparent relationships between hypertrophied lips and the expression profiles of ECM proteins, in particularly proteoglycans. The trends for the overall expression of ECM-related genes were similar across hypertrophied lip species, but we rarely observed a specific gene that was commonly expressed at high or low levels in all three examples of hypertrophied lips across all East African Great Lakes. Furthermore, although we focused primarily on the relationship between the Wnt signaling pathway and lip hypertrophy, several other pathways may be involved in the development of hypertrophied lips. For example, our enrichment analysis in adult cichlids identified VEGF-related pathways, which could contribute to lip hypertrophy by increasing vascularization and nutrient supply to the lip tissue. In addition, previous quantitative trait locus (QTL) analysis by Henning et al. (2017) concluded that lip hypertrophy is likely influenced by numerous loci with small additive effects. These lines of data imply that although enlargement of proteoglycan-rich loose connective tissue is common in hypertrophied lips, the developmental pathways to accomplish this are diverse in each lake.”

Reviewer 2:

Minor comments:

Last paragraph of Introduction: Remove the results of this study.

We appreciate your suggestion. We remove the specialized results from the last paragraph.

“In this study, we comprehensively compared the hypertrophied lips of cichlids across all East African Great Lakes using histology, proteomics, and transcriptomics. Histological and proteomic analyses revealed a distinct microstructure of hypertrophied lips compared to normal lips, and primary candidate proteins were identified. Transcriptome analysis at different developmental stages showed that the genes in Wnt signaling pathway was highly expressed in cichlids with hypertrophied lips at both the juvenile and adult stages. It is noteworthy that the distinct expression profiles observed in the proteome and transcriptome analyses of hypertrophied lips were similar among cichlids from each of the East African Great Lakes. The present study, which integrates comprehensive analyses for cichlids from all East African Great Lakes, provides insight for a better understanding of the molecular basis of a typical example of parallel evolution.”

Line 87-91: “In this study, we comprehensively compared the hypertrophied and normal lips of cichlids across all East African Great Lakes at various biological levels using histology, proteomics, and transcriptomics. As a result, we showed that a novel key pathway commonly involved in the formation of hypertrophied lips, providing insight into a better understanding of the molecular basis of a typical example of parallel evolution.”

Line 156: Italicize the scientific names.

We appreciate your correction.

Line 148: “*M. zebra* and *O. niloticus*” -> “*M. zebra* and *O. niloticus*”

Line 261: Remove the period after "Victoria."

We appreciate your correction.

Line 253: “Lake Victoria. (Figure 1; Figure S2).” -> “Lake Victoria (Figure 1; Figure S2).”

| *Line 416: Remove the period after "tissue."*

We appreciate your correction.

Line 420: “tissue. (A,B)” -> “tissue (A,B)”

| *Line 646: Probably "the anterior side to the left."*

We apologize for our mistake. As you commented, the anterior side is left. We corrected our manuscript as follows.

Line 648: “the anterior side to the right” -> “the anterior side to the left”

| *Fig. S2: Based on Fig. 1, the VG stained area appears larger in the Hypertrophied lip species; however, it is the opposite in Fig. S2.*

We appreciate your comments. This is because we calculated the ratio of the VG-stained area to the whole lip area. While the absolute VG-stained area is larger in hypertrophied lips, the proportion of the VG-stained area relative to the total lip area is smaller. This correction using entire area allows us to simply compare the degree of lip hypertrophy among species.

<https://doi.org/10.7554/eLife.99160.2.sa0>